

**(12) PATENT
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 200145098 B2
(10) Patent No. 783874**

(54) Title
Uncharged antisense oligonucleotides targeted to bacterial 16S and 23S PRNAS and their uses

(51)⁷ International Patent Classification(s)
C12N 015/11

(21) Application No: **200145098** (22) Application Date: **2000.11.29**

(87) WIPO No: **WO01/42457**

(30) Priority Data

(31) Number **60/168150** (32) Date **1999.11.29** (33) Country **US**

(43) Publication Date : **2001.06.18**

(43) Publication Journal Date : **2001.08.30**

(44) Accepted Journal Date : **2005.12.15**

(71) Applicant(s)
AVI BioPharma, Inc.

(72) Inventor(s)
Patrick L. Iversen

(74) Agent/Attorney
F.B. Rice and Co.,Level 23,200 Queen Street,MELBOURNE Vic 3000

(56) Related Art
WO 1998/003533

AU 200145098

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
14 June 2001 (14.06.2001)

PCT

(10) International Publication Number
WO 01/42457 A3

(51) International Patent Classification⁷: C12N 15/11. (74) Agents: GORTHEY, LeeAnn et al.; Iota Pi Law Group, A61K 31/7125, A61P 31/04 P.O. Box 60850, Palo Alto, CA 94306-0850 (US).

(21) International Application Number: PCT/US00/42391

(81) Designated States (national): AU, CA, JP, KR.

(22) International Filing Date:

29 November 2000 (29.11.2000)

(84) Designated States (regional): European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR).

(25) Filing Language:

English

Published:

— with international search report

(26) Publication Language:

English

(88) Date of publication of the international search report:
31 January 2002

(30) Priority Data:


60/168,150 29 November 1999 (29.11.1999) US

(71) Applicant: AVI BIOPHARMA, INC. [US/US]; Suite 200, 4575 S.W. Research Way, Corvallis, OR 97333 (US).

(72) Inventor: IVERSEN, Patrick, L.; 3902 N.W. Fair Oaks Place, Corvallis, OR 97330 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: UNCHARGED ANTISENSE OLIGONUCLEOTIDES TARGETED TO BACTERIAL 16S AND 23S PRNAS AND THEIR USES



(57) Abstract: The invention relates to compositions comprising oligomers antisense to bacterial 16S or 23S rRNA and capable of selectively modulating the biological activity thereof, and methods for their use. More particularly, the invention relates to antisense oligomers directed to 16S or 23S rRNA found in one or more particular bacteria, or generally conserved among bacteria in general, and to pharmaceutical compositions and methods of treatment comprising the same.

WO 01/42457 A3

ANTISENSE ANTIBACTERIAL METHOD AND COMPOSITIONField of the Invention

5 The present invention relates to oligonucleotide compositions antisense to bacterial 16S and 23S rRNA and methods for use of such compositions in the treatment of bacterial infection in a mammal.

References

Agrawal, S. *et al.*, *Proc. Natl. Acad. Sci. USA* **87**(4):1401-5 (1990).

10 Ardhammar, M. *et al.*, *J. Biomolecular Structure & Dynamics* **17**(1):33-40 (Aug 1999).

Attia, S.A. *et al.*, *Antisense & Nucleic Acid Drug Dev.* **8** (3):207-14 (1998).

Bennett, M.R. *et al.*, *Circulation* **92**(7):1981-1993 (1995).

Bonham, M.A. *et al.*, *Nucleic Acids Res.* **23**(7):1197-1203 (1995).

Boudvillain, M. *et al.*, *Biochemistry* **36**(10):2925-31 (1997).

15 Cross, C.W. *et al.*, *Biochemistry* **36**(14):4096-107 (Apr 8 1997).

Dagle, J.M. *et al.*, *Nucleic Acids Research* **28**(10):2153-7 (May 15 2000).

Ding, D. *et al.*, *Nucleic Acids Research* **24**(2):354-60 (Jan 15 1996).

Egholm, M. *et al.*, *Nature* **365**(6446):566-8 (Oct 7 1993).

Felgner *et al.*, *Proc. Nat. Acad. Sci. USA* **84**:7413 (1987).

20 Gait, M.J.; Jones, A.S. and Walker, R.T., *J. Chem. Soc. Perkin I*, 1684-86 (1974).

Gee, J.E. *et al.*, *Antisense & Nucleic Acid Drug Dev.* **8**:103-111 (1998).

Good, L. and Nielsen, P.E., *Proc. Nat. Acad. Sci. USA* **95**:2073-2076 (1998).

Huie, E.M. *et al.*, *J. Org. Chem.* **57**:4569 (1992).

Jones, A.S., MacCross, M. and Walker, R.T., *Biochem. Biophys. Acta* **365**:365-377 (1973).

25 Lesnikowski, Z.J. *et al.*, *Nucleic Acids Research* **18**(8):2109-15 (Apr 25 1990).

Matteucci, M., *Tetrahedron Lett.* **31**:2385-88 (1990).

McElroy, E.B. *et al.*, *Bioorg. Med. Chem. Lett.* **4**:1071 (1994).

Mertes, M.P. and Coates, E.A., *J. Med. Chem.* **12**:154-157 (1969).

Miller, P.S. *et al.*, in: *Antisense Research Applications*, Crooke, S.T. and Lebleu, B., Eds.,

30 CRC Press, Boca Raton, FL, p. 189. (1993).

Olgive, K.K. and Cormier, J.F., *Tetrahedron Lett* **26**:4159-4162 (1986).

Rahman, M.A. *et al.*, *Antisense Res Dev* **1**(4):319-27 (1991).

Roughton, A.L. *et al.*, *J. Am. Chem. Soc.* **117**:7249 (1995).

Stein, D. *et al.*, *Antisense & Nucleic Acid Drug Dev.* **7**(3):151-7 (Jun 1997); see also

35 Summerton, J. *et al.*, *Antisense & Nucleic Acid Drug Dev.* **7**(2):63-70 (Apr 1997).

Toulme, J.J. *et al.*, *Biochimie* 78(7):663-73 (1996).

Vasseur, J.J. *et al.*, *J. Am. Chem. Soc.* 114:4006 (1992).

Background of the Invention

5 Currently, there are several types of antibiotics in use against bacterial pathogens, with a variety of anti-bacterial mechanisms. Beta-lactam antibiotics, such as penicillin and cephalosporin, act to inhibit the final step in peptidoglycan synthesis. Glycopeptide antibiotics, including vancomycin and teichoplanin, inhibit both transglycosylation and transpeptidation of muramyl-pentapeptide, again interfering with peptidoglycan synthesis. Other well-known 10 antibiotics include the quinolones, which inhibit bacterial DNA replication, inhibitors of bacterial RNA polymerase, such as rifampin, and inhibitors of enzymes in the pathway for production of tetrahydrofolate, including the sulfonamides.

Some classes of antibiotics act at the level of protein synthesis. Notable among these are the 15 aminoglycosides, such as kanamycin and gentamycin. These compounds target the bacterial 30S ribosome subunit, preventing the association with the 50S subunit to form functional ribosomes. Tetracyclines, another important class of antibiotics, also target the 30S ribosome subunit, acting by preventing alignment of aminoacylated tRNA's with the corresponding mRNA codon. Macrolides and lincosamides, another class of antibiotics, inhibit bacterial synthesis by binding to 20 the 50S ribosome subunit, and inhibiting peptide elongation or preventing ribosome translocation.

25 Despite impressive successes in controlling or eliminating bacterial infections by antibiotics, the widespread use of antibiotics both in human medicine and as a feed supplement in poultry and livestock production has led to drug resistance in many pathogenic bacteria. Antibiotic resistance mechanisms can take a variety of forms. One of the major mechanisms of resistance to beta lactams, particularly in Gram-negative bacteria, is the enzyme beta-lactamase, which renders the antibiotic inactive. Likewise, resistance to aminoglycosides often involves an enzyme capable of 30 inactivating the antibiotic, in this case by adding a phosphoryl, adenyl, or acetyl group. Active efflux of antibiotics is another way that many bacteria develop resistance. Genes encoding efflux proteins, such as the tetA, tetG, tetL, and tetK genes for tetracycline efflux, have been identified. A bacterial target may develop resistance by altering the target of the drug. For example, the so-called penicillin binding proteins (PBPs) in many beta-lactam resistant bacteria are altered to inhibit the critical antibiotic binding to the target protein. Resistance to tetracycline may involve, in addition to enhanced efflux, the appearance of cytoplasmic proteins capable of competing with ribosomes for binding to the antibiotic. Where the antibiotic acts by inhibiting a bacterial enzyme, such as for sulfonamides, point mutations in the target enzyme may confer resistance.

The appearance of antibiotic resistance in many pathogenic bacteria, in many cases involving multi-drug resistance, has raised the specter of a pre-antibiotic era in which many bacterial pathogens are simply untreatable by medical intervention. There are two main factors that could contribute to this scenario. The first is the rapid spread of resistance and multi-resistance genes across bacterial strains, species, and genera by conjugative elements, the most important of which are self-transmissible plasmids. The second factor is a lack of current research efforts to find new types of antibiotics, due in part to the perceived investment in time and money needed to find new antibiotic agents and bring them through clinical trials, a process that may require a 20-year research effort in some cases.

10 In addressing the second of these factors, some drug-discovery approaches that may accelerate the search for new antibiotics have been proposed. For example, efforts to screen for and identify new antibiotic compounds by high-throughput screening have been reported, but to date no important lead compounds have been discovered by this route.

15 Several approaches that involve antisense agents designed to block the expression of bacterial resistance genes or to target cellular RNA targets, such as the rRNA in the 30S ribosomal subunit, have been proposed (Good *et al.*, 1998; Rahman *et al.*, 1991). In general, these approaches have been marginally successful, presumably because of poor uptake of the antisense agent (*e.g.*, Summerton *et al.*, 1997), or the requirement that the treated cells show high permeability for antibiotics (Good *et al.*, 1998).

20 There is thus a growing need for new antibiotics that (i) are not subject to the principal types of antibiotic resistance currently hampering antibiotic treatment of bacteria, (ii) can be developed rapidly and with some reasonable degree of predictability as to target-bacteria specificity, (iii) can also be designed for broad-spectrum activity, (iv) are effective at low doses, meaning, in part, that they are efficiently taken up by wild-type bacteria or even bacteria that have reduced permeability for antibiotics, and (v) show few side effects.

Summary of the Invention

30 In one aspect, the invention provides an antibacterial compound, consisting of a substantially uncharged antisense oligomer containing from 8 to 40 nucleotide subunits, including a targeting nucleic acid sequence at least 10 nucleotides in length which is complementary to a bacterial 16S or 23S rRNA nucleic acid sequence. Each of the subunits comprises a 5- or 6-membered ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base in the bacterial nucleic acid sequence. Adjacent subunits are joined by uncharged linkages selected from the group consisting of: uncharged phosphoramidate, phosphorodiamidate, carbonate, carbamate, amide, phosphotriester, alkyl phosphonate, siloxane, sulfone, sulfonamide.

sulfamate, thioformacetyl, and methylene-N-methylhydroxylamino, or by charged linkages selected from the group consisting of phosphate, charged phosphoramidate and phosphorothioate. The ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1, preferably at least 5:1, and more preferably at least 8:1. In one embodiment, the oligomer is fully uncharged.

5 Preferably, the oligomer is able to hybridize with the bacterial sequence at a T_m substantially greater than the T_m of a duplex composed of a corresponding DNA and the same bacterial sequence. Alternatively, the oligomer is able to hybridize with the bacterial sequence at a T_m substantially greater than 37°C, preferably greater than 50°C, and more preferably in the range of 60-80°C.

10 In one embodiment, the oligomer is a morpholino oligomer. The uncharged linkages, and, in one embodiment, all of the linkages, in such an oligomer are preferably selected from the group consisting of the structures presented in Figures 2A through 2D. Particularly preferred are phosphorodiamidate-linked oligomers, as represented at Figure 2B, where X=NR₂, R being hydrogen or methyl, Y=O, and Z=O.

15 The length of the oligomer is preferably 12 to 25 subunits. In one embodiment, the oligomer is a phosphorodiamidate-linked morpholino oligomer having a length of 15 to 20 subunits, and more preferably 17-18 subunits.

20 In selected embodiments, the targeting sequence is a broad spectrum sequence selected from the group consisting of SEQ ID NOs: 15, 16, and 21-25. In other embodiments, the targeting sequence is complementary to a Gram-positive bacterial 16S rRNA consensus sequence, e.g., SEQ ID NOs: 27-28, or is complementary to a Gram-negative bacterial 16S rRNA consensus sequence, e.g. SEQ ID NOs: 29-30.

Other targeting sequences can be used for treatment of an infection produced by various organisms, for example:

25 (a) *E. coli*, where the sequence is selected from the group consisting of SEQ ID NO:32 and SEQ ID NO:35;

(b) *Salmonella typhimurium*, where the sequence is selected from the group consisting of SEQ ID NO:18 and SEQ ID NO:36;

30 (c) *Pseudomonas aeruginosa*, where the sequence is selected from the group consisting of SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42 and SEQ ID NO:43;

(d) *Vibrio cholera*, where the sequence is selected from the group consisting of SEQ ID NO:45, SEQ ID NO:46 and SEQ ID NO:47;

(e) *Neisseria gonorrhoea*, where the sequence is selected from the group consisting of SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50 and SEQ ID NO:51;

(f) *Staphylococcus aureus*, where the sequence is selected from the group consisting of SEQ ID NO:53, SEQ ID NO:54 and SEQ ID NO:55;

(g) *Mycobacterium tuberculosis*, where the sequence is selected from the group consisting of SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58 and SEQ ID NO:59;

5 (h) *Helicobacter pylori*, where the sequence is selected from the group consisting of SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62 and SEQ ID NO:63;

(i) *Streptococcus pneumoniae*, where the sequence is selected from the group consisting of SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66 and SEQ ID NO:67;

10 (j) *Treponema palladium*, where the sequence is selected from the group consisting of SEQ ID NO:69, SEQ ID NO:70 and SEQ ID NO:71;

(k) *Chlamydia trachomatis*, where the sequence is selected from the group consisting of SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74 and SEQ ID NO:75;

(l) *Bartonella henselae*, where the sequence is selected from the group consisting of SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78 and SEQ ID NO:79;

15 (m) *Hemophilis influenza*, where the sequence is selected from the group consisting of SEQ ID NO:81, SEQ ID NO:82 and SEQ ID NO:83;

(n) *Shigella dysenteriae*, where the sequence is presented as SEQ ID NO:88; or

(o) *Enterococcus faecium*, where the sequence is presented as SEQ ID NO: 92.

20 In other embodiments, the targeting sequence is an antisense oligomer sequence selected from one of the following groups, for use in treatment of an infection produced by:

(a) *E. coli*, *Salmonella typhimurium* and *Shigella dysenteriae*, where the sequence is selected from the group consisting of SEQ ID NO:15, SEQ ID NO:16, SEQ ID NO:33, SEQ ID NO:34, SEQ ID NO:38, SEQ ID NO:39 and SEQ ID NO:86 and SEQ ID NO:87;

25 (b) *E. coli*, *Salmonella typhimurium* and *Hemophilis influenza*, where the sequence is presented as SEQ ID NO:31;

(c) *E. coli* and *Shigella dysenteriae*, where the sequence is presented as SEQ ID NO:17;

(d) *E. coli*, *Salmonella typhimurium*, *Shigella dysenteriae*, *Hemophilis influenza* and *Vibrio cholera*, where the sequence is presented as SEQ ID NO:44;

30 (e) *Staphylococcus aureus* and *Bartonella henselae*, where the sequence is presented as SEQ ID NO:52;

(f) *Salmonella typhimurium*, *Hemophilis influenza* and *Treponema palladium*, where the sequence is presented as SEQ ID NO:68; or

(g) *E. coli*, *Salmonella typhimurium*, *Shigella dysenteriae*, *Hemophilis influenza* and *Neisseria gonorrhoea*, where the sequence is presented as SEQ ID NO:84.

In a related aspect, the invention provides a method of treating a bacterial infection in a human or mammalian animal subject, by administering to the subject, in a pharmaceutically effective amount, a substantially uncharged antisense oligomer as described above. Various selected embodiments of the oligomer and the target sequence are as described above. Preferably, the antisense oligomer is administered in an amount and manner effective to result in a peak blood concentration of at least 200-400 nM antisense oligomer. The method can be used, for example, for treating bacterial infections of the skin, wherein administration is by a topical route, or for use in treating a bacterial respiratory infection, wherein administration is by inhalation.

10 In a further related aspect, the invention provides a livestock and poultry food composition containing a food grain supplemented with a subtherapeutic amount of an antibacterial compound, said compound consisting of a substantially uncharged antisense oligomer as described above.

15 Also contemplated is, in a method of feeding livestock and poultry with a food grain supplemented with subtherapeutic levels of an antibiotic, an improvement in which the food grain is supplemented with a subtherapeutic amount of an antibacterial compound of the type described above.

20 According to the invention there is also provided an antibacterial compound consisting of a substantially uncharged antisense oligomer containing from 10 to 40 nucleotide subunits, each of said subunits comprising a morpholino ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base, said base-pairing moieties including a targeting nucleic acid sequence at least 10 nucleotides in length which is complementary to a bacterial 16S or 23S rRNA nucleic acid sequence,

25 wherein adjacent subunits are joined by uncharged linkages selected from the group consisting of uncharged phosphoramidate and phosphorodiamidate, or by charged phosphoramidate linkages, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1.

30 According to the invention there is also provided a method of treating a bacterial infection in a human or mammalian animal subject, comprising administering to the subject, in a pharmaceutically effective amount, a substantially uncharged antisense oligomer containing from 10 to 40 nucleotide subunits, each of said subunits comprising a morpholino ring supporting a base-pairing moiety effective to bind by

Watson-Crick base pairing to a respective nucleotide base, said base-pairing moieties including a targeting nucleic acid sequence at least 10 nucleotides in length which is complementary to a bacterial 16S or 23S rRNA nucleic acid sequence, wherein

wherein adjacent subunits are joined by uncharged linkages selected from the 5 group consisting of uncharged phosphoramidate and phosphorodiamidate, or by charged phosphoramidate linkages, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1.

According to the invention there is also provided a livestock and poultry food composition containing a food grain supplemented with a subtherapeutic amount of an

10 antibacterial compound, said compound consisting of a substantially uncharged antisense oligomer containing from 10 to 40 nucleotide subunits, each of said subunits comprising a morpholino ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base, said base-pairing including a targeting nucleic acid sequence at least 10 nucleotides in length which is 15 complementary to a bacterial 16S or 23S rRNA nucleic acid sequence,

wherein adjacent subunits are joined by uncharged linkages selected from the group consisting of uncharged phosphoramidate and phosphorodiamidate, or by charged phosphoramidate linkages, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1.

20 These and other desires and features of the invention will become more fully apparent when the following detailed description is read in conjunction with the accompanying figures and examples.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated 25 element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is solely for the purpose of providing a context for the present invention. It is not to be taken as an admission that any or all of 30 these matters form part of the prior art base or were common general knowledge in the field relevant to the present invention as it existed before the priority date of each claim of this application.

Brief Description Of The Figures

35 Figure 1 shows several preferred morpholino-type subunits having 5-atom (A), six-atom (B) and seven-atom (C-D) linking groups suitable for forming polymers ;

Figures 2A-D show the repeating subunit segment of exemplary morpholino oligonucleotides, designated A through D, constructed using subunits A-D, respectively, of Figure 1.

5 Figures 3A-3G show examples of uncharged linkage types in oligonucleotide analogs;

Figure 4 depicts the results of a study on the effect of a phosphorodiamidate morpholino antisense oligomer (PMO) designated VRE-2 (SEQ ID NO: 92) (see Table 10), targeted against an *Enterococcus faecium* 16S rRNA sequence, alone or in

10 combination with 50 μ M of an oligomer antisense to c-myc (SEQ ID NO: 139), on bacterial colony formation in *E. coli*, presented as percent viability ;

Figure 5 depicts the results of a study on the effect of various concentrations of a PMO having SEQ ID NO: 15 (broad spectrum ; see Table 2A), targeted against a bacterial 16S rRNA consensus sequence, on the bacterial colony formation in *E. coli*,

15 presented as percent inhibition of colony formation;

Figure 6 depicts the results of a study wherein PMO oligomers targeting various different regions of *Enterococcus faecium* 16S rRNA, designated AVI-1-23-22, -32, -45, -33, -34, -44, -35 and -36 (SEQ ID NOS: 92, 102, 115, 103, 104, 114, 105, and 106), indicated in the figure as 22, 23, 45, 33, 34, 44, 35 and 36, respectively, were added at 1 μ M to vancomycin-resistant *Enterococcus faecium* (VRE) cultures, with the results presented as percent viability;

5 Figure 7 depicts the results of a study wherein PMO oligomers targeting various different regions of *Enterococcus faecium* 23S rRNA, designated AVI-1-23-46, -47, -48, -49 and -50 (SEQ ID NOS: 116-120), indicated in the figure as 46, 47, 48, 49 and 50, respectively, were added at 1 μ M to vancomycin-resistant *Enterococcus faecium* cultures, with the results presented as percent viability;

10 Figure 8 depicts the results of a study on the effect of 1 μ M of PMOs of various lengths targeted against the 16S rRNA of a vancomycin-resistant *Enterococcus faecium* bacterial strain on viability of the bacteria (percent viability, reported as percent of untreated control). The PMO sequences corresponding to the oligomer lengths are shown in Table 12, which illustrates antisense targeting of 16S rRNA in VRE, reported as percent inhibition (100 - percent of untreated control);

15 Figure 9 depicts the results of a study on the effect of 1 μ M PMO targeted against *Enterococcus faecium* 16S rRNA, designated VRE-2, AVI 1-23-22 (SEQ ID NO: 92), on bacterial colony formation in VRE, presented as percent viability (percent of control) as determined on days 1 through 6; and

20 Figures 10A-B depict the results of a study on the effect of 1 μ M of a PMO targeted against *Enterococcus faecium* 16S rRNA (SEQ ID NO: 92), alone or in combination with (A) 3 μ M vancomycin, or (B) 3 μ M ampicillin, on growth of VRE, with the results reported as percent viability.

25 **Detailed Description of the Invention**

I. **Definitions**

The terms below, as used herein, have the following meanings, unless indicated otherwise:

As used herein, the term "16S ribosomal RNA", also termed "16S rRNA", refers to RNA which is part of the structure of a ribosome and is involved in the synthesis of proteins.

30 The term "polynucleotide" as used herein refers to a polymeric molecule having a backbone which supports bases capable of hydrogen bonding to typical polynucleotides, where the polymer backbone presents the bases in a manner to permit such hydrogen bonding in a sequence specific fashion between the polymeric molecule and a typical polynucleotide (e.g., single-stranded RNA, double-stranded RNA, single-stranded DNA or double-stranded DNA). "Polynucleotides" include

polymers with nucleotides which are an N- or C-glycoside of a purine or pyrimidine base, and polymers containing non-standard nucleotide backbones, for example, backbones formed using phosphorodiamidate morpholino chemistry, polyamide linkages (e.g., peptide nucleic acids or PNAs) and other synthetic sequence-specific nucleic acid molecules.

5 As used herein, the terms "antisense oligonucleotide" and "antisense oligomer" are used interchangeably and refer to a sequence of nucleotide bases and a subunit-to-subunit backbone that allows the antisense oligomer to hybridize to a target nucleic acid (e.g., RNA) sequence by Watson-Crick base pairing, to form a nucleic acid:oligomer heteroduplex within the target sequence. The oligomer may have exact sequence complementarity to the target sequence or near complementarity. In one exemplary application, such an antisense oligomer may block or inhibit the function of 16S or 23S rRNA containing a given target sequence, may bind to a double-stranded or single stranded portion of the 16S or 23S rRNA target sequence, may inhibit mRNA translation and/or protein synthesis, and may be said to be "directed to" a sequence with which it specifically hybridizes.

10 15 As used herein, an oligonucleotide or antisense oligomer "specifically hybridizes" to a target polynucleotide if the oligomer hybridizes to the target under physiological conditions, with a T_m substantially greater than 37°C, preferably at least 50°C, and typically 60°C-80°C or higher. Such hybridization preferably corresponds to stringent hybridization conditions. At a given ionic strength and pH, the T_m is the temperature at which 50% of a target sequence hybridizes to a complementary polynucleotide.

20 25 Polynucleotides are described as "complementary" to one another when hybridization occurs in an antiparallel configuration between two single-stranded polynucleotides. A double-stranded polynucleotide can be "complementary" to another polynucleotide, if hybridization can occur between one of the strands of the first polynucleotide and the second. Complementarity (the degree that one polynucleotide is complementary with another) is quantifiable in terms of the proportion (i.e., the percentage) of bases in opposing strands that are expected to form hydrogen bonds with each other, according to generally accepted base-pairing rules.

As used herein, the term "consensus sequence", relative to 16S or 23S rRNA sequences, refers to a sequence which is common to or shared by a particular group of organisms. The 30 consensus sequence shows the nucleic acid most commonly found at each position within the polynucleotide. For example, a Gram-negative bacterial 16S or 23S rRNA consensus sequence is common to Gram-negative bacteria and generally not found in bacteria that are not Gram-negative.

As used herein, the term "conserved", relative to 16S or 23S rRNA sequences, also refers to a sequence which is common to or shared by a particular group of organisms (e.g., bacteria).

A "subunit" of an oligonucleotide or oligonucleotide analog refers to one nucleotide (or nucleotide analog) unit of the oligomer. The term may refer to the nucleotide unit with or without the attached intersubunit linkage, although, when referring to a "charged subunit", the charge typically resides within the intersubunit linkage (e.g. a phosphate or phosphorothioate linkage).

5 As used herein, a "morpholino oligomer" refers to a polymeric molecule having a backbone which supports bases capable of hydrogen bonding to typical polynucleotides, wherein the polymer lacks a pentose sugar backbone moiety, and more specifically lacks a ribose backbone linked by phosphodiester bonds which is typical of nucleotides and nucleosides, but instead contains a ring nitrogen with coupling through the ring nitrogen. A typical "morpholino" 10 oligonucleotide is composed of morpholino subunit structures of the form shown in Fig. 1A-1D, where (i) the structures are linked together by phosphorous-containing linkages, one to three atoms long, joining the morpholino nitrogen of one subunit to the 5' exocyclic carbon of an adjacent subunit, and (ii) B is a purine or pyrimidine base-pairing moiety effective to bind, by 15 base-specific hydrogen bonding, to a base in a polynucleotide.

15 As used herein, the term "PMO" refers to a phosphorodiamidate morpholino oligomer, as further described below, wherein the oligomer is a polynucleotide of about 8-40 bases in length, preferably 12-25 bases in length. This preferred aspect of the invention is illustrated in Fig. 2B, where the two subunits are joined by a phosphorodiamidate linkage.

20 As used herein, a "nuclease-resistant" oligomeric molecule (oligomer) is one whose backbone is not susceptible to nuclease cleavage of a phosphodiester bond. Exemplary nuclease resistant antisense oligomers are oligonucleotide analogs such as phosphorothioate and phosphate-amine DNA (pnDNA), both of which have a charged backbone, and methyl phosphonate and phosphoramidate- or phosphorodiamidate-linked morpholino oligonucleotides, which have uncharged backbones.

25 A "2'-O-allyl (or alkyl) modified oligonucleotide" is an oligoribonucleotide in which the 2' hydroxyl is converted to an allyl or alkyl ether, respectively. The alkyl ether is typically a methyl ether.

30 "Alkyl" refers to a fully saturated acyclic monovalent radical containing carbon and hydrogen, which may be branched or a straight chain. Examples of alkyl groups are methyl, ethyl, n-butyl, t-butyl, n-heptyl, and isopropyl. "Lower alkyl" refers to an alkyl radical of one to six carbon atoms, and preferably one to four carbon atoms, as exemplified by methyl, ethyl, isopropyl, n-butyl, isobutyl, and t-butyl.

35 As used herein, a first sequence is an "antisense sequence" with respect to a second sequence if a polynucleotide with a first sequence specifically binds to, or specifically hybridizes with, a polynucleotide which has a second sequence, under physiological conditions.

As used herein, a "base-specific intracellular binding event involving a target RNA" refers to the specific binding of an oligomer to a target RNA sequence inside a cell. The base specificity of such binding is sequence specific. For example, a single-stranded polynucleotide can specifically bind to a single-stranded polynucleotide that is complementary in sequence.

5 As used herein, "nuclease-resistant heteroduplex" refers to a heteroduplex formed by the binding of an antisense oligomer to its complementary target, such that the heteroduplex is resistant to *in vivo* degradation by ubiquitous intracellular and extracellular nucleases.

As used herein, the term "broad spectrum bacterial sequence", with reference to bacterial 16S rRNA, refers to an oligonucleotide of the invention which is antisense to some segment of 10 most if not all of the bacterial 16S rRNA sequences described herein. A corresponding definition applies to bacterial 23S rRNA. Exemplary broad spectrum bacterial sequences described herein include the antisense oligomers presented as SEQ ID NO:21, SEQ ID NO:22 and SEQ ID NO:23, which are antisense to an *Escherichia coli* (*E. coli*), *Salmonella thyphimurium* (*S. thyphi*), *Pseudomonas aeruginosa* (*P. aeruginosa*), *Vibrio cholera*, *Neisseria gonorrhoea* (*N. gonorrhoea*), 15 *Staphylococcus aureus* (*Staph. aureus*), *Mycobacterium tuberculosis* (*Myco. tubercul.*), *Helicobacter pylori* (*H. pylori*), *Streptococcus pneumoniae* (*Strep. pneumoniae*), *Treponema palladium* (*Treponema pallad.*), *Chlamydia trachomatis* (*Chlamydia trach.*), *Bartonella henselae* (*Bartonella hens.*), *Hemophilis influenza* (*H. influenza*) and *Shigella dysenteriae* (*Shigella dys.*) 16S rRNA sequence (see Table 5A), and SEQ ID NOs 24-25, which are antisense to the 16s rRNA of the 20 majority of these organisms (see Table 5B).

As used herein, the term "narrow spectrum bacterial sequence", with respect to 16S bacterial rRNA, refers to an oligonucleotide of the invention which is antisense to particular, but not most or all, bacterial 16S rRNA sequences described herein. Again, a corresponding definition applies to bacterial 23S rRNA. A narrow spectrum bacterial sequence may be specific to one or more 25 different bacteria, e.g., an antisense oligomer which is antisense to *E. coli*, *S. thyphi* and *Shigella dys.* 16S rRNA, but not the other bacterial 16S rRNA sequences described herein, as exemplified by SEQ ID NO:31; or an antisense oligomer which is antisense to the *E. coli* 16S rRNA sequence, but not the other bacterial 16S rRNA sequences described herein, as exemplified by SEQ ID NO:32.

30 As used herein, the term "modulating expression" relative to oligonucleotides refers to the ability of an antisense oligomer to either enhance or reduce the expression of a given protein by interfering with the expression or translation of RNA.

As used herein, "effective amount" relative to an antisense oligomer refers to the amount of antisense oligomer administered to a mammalian subject, either as a single dose or as part of a

series of doses, that is effective to inhibit a biological activity, *e.g.*, expression of a selected target nucleic acid sequence.

As used herein, "treatment" of an individual or a cell is any type of intervention provided as a means to alter the natural course of the individual or cell. Treatment includes, but is not limited to, administration of a pharmaceutical composition, and may be performed either prophylactically or subsequent to the initiation of a pathologic event or contact with an etiologic agent.

As used herein, the term "improved therapeutic outcome", relative to a patient diagnosed as infected with a particular bacteria, refers to a slowing or diminution in the growth of the bacteria and/or a decrease in, or elimination of, detectable symptoms typically associated with infection by that particular bacteria.

II. Antisense Oligomers: Selection Criteria

Antisense compounds employed in the invention preferably meet several criteria of structure and properties, considered in the subsections below.

15 A. Base Sequence and Length

The antisense compound has a base sequence targeted against a selected RNA target sequence. The region of complementarity with the target RNA sequence may be as short as 10-12 bases, but is preferably 13-20 bases, and more preferably 17-20 bases, in order to achieve the requisite binding T_m , as discussed below.

20 In some cases, the target for modulation of the activity of 16S rRNA using the antisense oligomers of the invention is a sequence in a double stranded region of the 16s rRNA, such as the peptidyl transferase center, the alpha-sarcin loop or the mRNA binding region of the 16S rRNA sequence. In other cases, the target for modulation of gene expression is a sequence in a single stranded region of bacterial 16S or 23S rRNA. The target may be a consensus sequence for 25 bacterial 16S or 23S rRNAs in general, a sequence common to the 16s or 23S rRNA of one or more types of bacteria (*e.g.*, Gram positive or Gram negative bacteria), or specific to a particular 16S or 23S rRNA sequence.

30 The oligomer may be 100% complementary to the bacterial RNA target sequence, or it may include mismatches, *e.g.*, to accommodate variants, as long as the heteroduplex formed between the oligomer and bacterial RNA target sequence is sufficiently stable to withstand the action of cellular nucleases and other modes of degradation which may occur *in vivo*. Mismatches, if present, are less destabilizing toward the end regions of the hybrid duplex than in the middle. The number of mismatches allowed will depend on the length of the oligomer, the percentage of G:C base pairs in the duplex and the position of the mismatch(es) in the duplex, according to well 35 understood principles of duplex stability. Although such an antisense oligomer is not necessarily

100% complementary to the bacterial RNA target sequence, it is effective to stably and specifically bind to the target sequence such that a biological activity of the nucleic acid target, e.g., expression of bacterial protein(s) is modulated.

5 Oligomers as long as 40 bases may be suitable, where at least the minimum number of bases, e.g., 10-15 bases, are complementary to the target RNA sequence. In general, however, facilitated or active uptake in cells is optimized at oligomer lengths less than about 30, preferably less than 25, and more preferably 20 or fewer bases. For PMO oligomers, described further below, an optimum balance of binding stability and intake generally occurs at lengths of 17-18 bases.

10 **B. Duplex Stability (Tm)**

The oligomer must form a stable hybrid duplex with the target sequence. Preferably, the oligomer is able to hybridize to the target RNA sequence with a Tm substantially greater than the Tm of a duplex composed of a corresponding DNA and the same target RNA sequence. The antisense oligomer will have a binding Tm, with respect to a complementary-sequence RNA, of greater than body temperature and preferably greater than 50°C. Tm's in the range 60-80°C or greater are preferred. The Tm of an antisense compound with respect to complementary-sequence RNA may be measured by conventional methods, such as those described by Hames *et al.*, Nucleic Acid Hybridization, IRL Press 1985, pp.107-108. According to well known principles, the Tm of an oligomer compound, with respect to a complementary-base RNA hybrid, can be increased by increasing the length (in basepairs) of the heteroduplex. At the same time, for purposes of optimizing cell transport, it may be advantageous to limit the size of the oligomer. For this reason, compounds that show high Tm (50°C or greater) at a length of 15-20 bases or less will be preferred over those requiring 20+ bases for high Tm values.

25 Increasing the ratio of C:G paired bases in the duplex is also known to generally increase in the Tm of an oligomer compound. Studies in support of the invention suggest that maximizing the number of C bases in the antisense oligomer is particularly favorable.

C. Uptake by Cells

30 In order to achieve adequate intracellular levels, the antisense oligomer must be actively taken up by cells, meaning that the compound is taken up by facilitated or active transport, if administered in free (non-complexed) form, or is taken by an endocytotic mechanism if administered in complexed form.

When the antisense compound is administered in complexed form, the complexing agent typically is a polymer, e.g., a cationic lipid, polypeptide, or non-biological cationic polymer, having an opposite charge to a net charge on the antisense compound. Methods of forming complexes, including bilayer complexes, between anionic oligonucleotides and cationic lipid or

other polymer components are well known. For example, the liposomal composition Lipofectin® (Felgner *et al.*, 1987), containing the cationic lipid DOTMA (N-[1-(2,3-dioleyloxy)propyl]-N,N,N-trimethylammonium chloride) and the neutral phospholipid DOPE (dioleyl phosphatidyl ethanolamine), is widely used. After administration, the complex is taken up by cells through an 5 endocytotic mechanism, typically involving particle encapsulation in endosomal bodies. The ability of the antisense agent to resist cellular nucleases promotes survival and ultimate delivery of the agent to the cell cytoplasm.

In the case where the agent is administered in free form, the agent should be substantially uncharged, meaning that a majority of its intersubunit linkages are uncharged at physiological pH. 10 Experiments carried out in support of the invention indicate that a small number of net charges, e.g., 1-2 for a 15- to 20-mer oligomer, can enhance cell uptake of certain oligomers with substantially uncharged backbones. The charges may be carried on the oligomer itself, e.g., in the backbone linkages, or may be terminal charged-group appendages. Preferably, the number of charged linkages is no more than one charged linkage per four uncharged linkages.

15 An oligomer may also contain both negatively and positively charged backbone linkages, as long as two opposite charges are substantially offsetting, and preferably do not include runs of more than 3-5 consecutive subunits of either charge. For example, the oligomer may have a given number of anionic linkages, e.g. phosphorothioate or N3'→P5' phosphoramidate linkages, and a comparable number of cationic linkages, such as N,N-diethylenediamine phosphoramidates (Dagle). The net charge is preferably neutral or at most 1-2 net charges per oligomer, as above.

20 In addition to being substantially or fully uncharged, the antisense agent is preferably a substrate for a membrane transporter system (i.e. a membrane protein or proteins) capable of facilitating transport or actively transporting the oligomer across the cell membrane. This feature may be determined by one of a number of tests, as follows, for oligomer interaction or cell 25 uptake.

25 A first test assesses binding at cell surface receptors, by examining the ability of an oligomer compound to displace or be displaced by a selected charged oligomer, e.g., a phosphorothioate oligomer, on a cell surface. The cells are incubated with a given quantity of test oligomer, which is typically fluorescently labeled, at a final oligomer concentration of between about 10-300 nM. 30 Shortly thereafter, e.g., 10-30 minutes (before significant internalization of the test oligomer can occur), the displacing compound is added, in incrementally increasing concentrations. If the test compound is able to bind to a cell surface receptor, the displacing compound will be observed to displace the test compound. If the displacing compound is shown to produce 50% displacement at a concentration of 10X the test compound concentration or less, the test compound is considered 35 to bind at the same recognition site for the cell transport system as the displacing compound.

A second test measures cell transport, by examining the ability of the test compound to transport a labeled reporter, e.g., a fluorescence reporter, into cells. The cells are incubated in the presence of labeled test compound, added at a final concentration between about 10-300 nM. After incubation for 30-120 minutes, the cells are examined, e.g., by microscopy, for intracellular label. The presence of significant intracellular label is evidence that the test compound is transported by facilitated or active transport.

A third test relies on the ability of certain antisense compounds to effectively inhibit bacterial growth when targeted against bacterial 16S or 23S rRNA. Studies carried out in support of the present invention show that the inhibition requires active or facilitated transport across bacterial cell membranes. The test compound is prepared with a target 16S sequence that has been demonstrated to be effective in inhibiting bacterial growth. For example, SEQ ID NOS: 1-3 herein are representative sequences against *E. coli* 16S rRNA. The compound is added to the growing bacterial culture at increasing concentrations, typically between 10nM and 1 mM. The ability to inhibit bacterial growth is measured from number of cell colonies cell counts at 24-72 hours after addition of the test compound. Compounds which can produce a 50% inhibition at a concentration of between about 100-500 nM or lower are considered to be good candidates for active transport.

As shown by the data in Fig. 4, 500 nM of PMO antisense oligomer targeted against VRE (vancomycin-resistant *Enterococcus*) 16s rRNA, having SEQ ID NO: 92, inhibited growth in VRE by about 50%. It was also observed that addition of a comparatively large concentration (50 μ M) of a nontarget sequence PMO (antisense to *c-myc*; SEQ ID NO: 139) essentially nullified this effect, suggesting that the transport mechanism has a finite capacity.

D. mRNA Resistance to RNaseH

Two general mechanisms have been proposed to account for inhibition of expression by antisense oligonucleotides. (See e.g., Agrawal *et al.*, 1990; Bonham *et al.*, 1995; and Boudvillain *et al.*, 1997). In the first, a heteroduplex formed between the oligonucleotide and mRNA is a substrate for RNaseH, leading to cleavage of the mRNA. Oligonucleotides belonging, or proposed to belong, to this class include phosphorothioates, phosphotriesters, and phosphodiesters (unmodified "natural" oligonucleotides). However, because such compounds would expose mRNA in an oligomer:RNA duplex structure to proteolysis by RNaseH, and therefore loss of duplex, they are suboptimal for use in the present invention. A second class of oligonucleotide analogs, termed "steric blockers" or, alternatively, "RNaseH inactive" or "RNaseH resistant", have not been observed to act as a substrate for RNaseH, and are believed to act by sterically blocking target RNA nucleocytoplasmic transport, splicing or translation. This class includes methylphosphonates (Toulme *et al.*, 1996), morpholino oligonucleotides, peptide nucleic acids

(PNA's), 2'-O-allyl or 2'-O-alkyl modified oligonucleotides (Bonham, 1995), and N3'→P5' phosphoramidates (Gee, 1998; Ding).

5 A test oligomer can be assayed for its ability to protect mRNA against RNaseH by forming an RNA:oligomer duplex with the test compound, then incubating the duplex with RNaseH under a standard assay conditions, as described in Stein *et al.* After exposure to RNaseH, the presence or absence of intact duplex can be monitored by gel electrophoresis or mass spectrometry.

10 In testing an oligomer for suitability in the present invention, each of the properties detailed above is preferably met. It is recognized that the "substantially uncharged" feature is inherently met where the linkages are uncharged, and the target-sequence complementarity is achieved by base-sequence design. Thus, an oligomer is preferably tested as to its (i) Tm with respect to target RNA at a duplex length preferably between 12-20 basepairs, (ii) ability to be transported across cell membranes by active or facilitated transport, and (iii) ability to prevent RNA proteolysis by RNaseH in duplex form.

15 The antibacterial effectiveness of a given antisense oligomer may be further evaluated by screening methods known in the art. For example, the oligomer may be incubated with a bacterial culture *in vitro* and the effect on the target 16S RNA evaluated by monitoring (1) heteroduplex formation with the target sequence and/or non-target sequences, using procedures known to those of skill in the art, *e.g.*, an electrophoretic gel mobility assay; (2) the amount of 16S mRNA, as determined by standard techniques such as RT-PCR or Northern blot; (3) the 20 amount of bacterial protein production, as determined by standard techniques such as ELISA or Western blotting; or (4) the amount of bacterial growth *in vitro* for both bacteria known to have the 16S rRNA sequence targeted by a particular antisense oligomer and bacteria not predicted to have the target 16S rRNA sequence.

25 Candidate antisense oligomers may also be evaluated, according to well known methods, for acute and chronic cellular toxicity, such as the effect on protein and DNA synthesis as measured via incorporation of ³H-leucine and ³H-thymidine, respectively. In addition, various control oligonucleotides, *e.g.*, one or more control oligonucleotides such as sense, nonsense or scrambled antisense sequences, or sequences containing mismatched bases, are generally included in the evaluation process, in order to confirm the specificity of binding of candidate antisense oligomers. 30 The results of such tests allow discrimination of specific effects of antisense inhibition of gene expression from indiscriminate suppression. (See, *e.g.* Bennett *et al.*, 1995). Sequences may be modified as needed to limit non-specific binding of antisense oligomers to non-target sequences, *e.g.*, by changing the length or the degree of complementarity to the target sequence.

III. Uncharged Oligonucleotide Analogs

Examples of uncharged linkages that may be used in oligonucleotide analogs of the invention are shown in Figs. 3A-3G. (As noted below, a small number of charged linkages, e.g. charged phosphoramidate or phosphorothioate, may also be incorporated into the oligomers.) The uncharged linkages include carbonate (3A, R=O) and carbamate (3A, R=NH₂) linkages, (Mertes; Gait); alkyl phosphonate and phosphotriester linkages (3B, R=alkyl or -O-alkyl) (Miller; Lesnikowski); amide linkages (3C); sulfones (3D, R₁, R₂ = CH₂) (Roughten); sulfonamides (3D, R₁=NH, R₂=CH₂ or vice versa) (McElroy); sulfamates (3D, R₁, R₂ = NH) (Huie); and a thioformacetyl linkage (3E) (Matteucci; Cross). The latter is reported to have enhanced duplex and triplex stability with respect to phosphorothioate antisense compounds (Cross). Also reported are the 3'-methylene-N-methylhydroxyamino compounds of structure 3F (Vasseur). In Figs. 3A-3G, B represents a purine or pyrimidine base-pairing moiety effective to bind, by base-specific hydrogen bonding, to a base in a polynucleotide, preferably selected from adenine, cytosine, guanine and uracil. The linkages join nucleotide subunits, each consisting of a 5- or 6-membered ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base in the bacterial nucleic acid sequence. These subunits may comprise, for example, ribose rings, as in native nucleic acids, or morpholino rings, as described further below.

PNAs (peptide nucleic acids) are analogs of DNA in which the backbone is structurally homomorphous with a deoxyribose backbone, consisting of N-(2-aminoethyl) glycine units to which pyrimidine or purine bases are attached. PNAs containing natural pyrimidine and purine bases hybridize to complementary oligonucleotides obeying Watson-Crick base-pairing rules, and mimic DNA in terms of base pair recognition (Egholm *et al.*, 1993). However, PNA antisense agents have been observed to display slow membrane penetration in cell cultures, possibly due to poor uptake (transport) into cells. (See, e.g., Ardhhammar, M. *et al.*, 1999).

Oligomeric ribonucleotides substituted at the 2'-oxygen show high RNA binding affinities and, in comparison to unsubstituted ribonucleotides, reduced sensitivity to endogenous nucleases. Methyl- substituted ribonucleotides are reported to provide greater binding affinity and cellular uptake than those having larger 2'-oxygen substituents (e.g. ethyl, propyl, allyl, or pentyl).

One preferred oligomer structure employs morpholino-based subunits bearing base-pairing moieties, joined by uncharged linkages as outlined above. Especially preferred is a substantially uncharged morpholino oligomer such as illustrated by the phosphorodiamidate-linked compound shown in Fig. 3G. Morpholino oligonucleotides, including antisense oligomers, are detailed, for example, in co-owned U.S. Patent Nos. 5,698,685, 5,217,866, 5,142,047, 5,034,506, 5,166,315, 5,185,444, 5,521,063, and 5,506,337, all of which are expressly incorporated by reference herein. Desirable chemical properties of the morpholino-based subunits are the ability to be

linked in a oligomeric form by stable, uncharged backbone linkages, the ability of the polymer so formed to hybridize with a complementary-base target nucleic acid, including target RNA, with high T_m, even with oligomers as short as 10-14 bases, the ability of the oligomer to be actively transported into mammalian cells, and the ability of the oligomer:RNA heteroduplex to resist 5 RNase degradation.

Exemplary backbone structures for antisense oligonucleotides of the invention include the morpholino subunit types shown in Figs. 1A-D, each linked by an uncharged, phosphorous-containing subunit linkage. In these figures, the X moiety pendant from the phosphorous may be any of the following: fluorine; an alkyl or substituted alkyl; an alkoxy or substituted alkoxy; a 10 thioalkoxy or substituted thioalkoxy; or, an unsubstituted, monosubstituted, or disubstituted nitrogen, including cyclic structures. Alkyl, alkoxy and thioalkoxy preferably include 1-6 carbon atoms, and more preferably 1-4 carbon atoms. Monosubstituted or disubstituted nitrogen 15 preferably refers to lower alkyl substitution, and the cyclic structures are preferably 5- to 7-membered nitrogen heterocycles optionally containing 1-2 additional heteroatoms selected from oxygen, nitrogen, and sulfur. Z is sulfur or oxygen, and is preferably oxygen.

Fig. 1A shows a phosphorous-containing linkage which forms the five atom repeating-unit backbone shown in Fig. 2A, where the morpholino rings are linked by a 1-atom phosphoamide linkage.

Subunit B in Fig. 1B is designed for 6-atom repeating-unit backbones, as shown in Fig. 2B. 20 In Fig. 1B, the atom Y linking the 5' morpholino carbon to the phosphorous group may be sulfur, nitrogen, carbon or, preferably, oxygen. The X and Z moieties are as defined above. Particularly preferred morpholino oligonucleotides include those composed of morpholino subunit structures of the form shown in Fig. 2B, where X=NH₂ or N(CH₃)₂, Y=O, and Z=O.

Subunits C-D in Figs. 1C-D are designed for 7-atom unit-length backbones as shown for 25 structures in Figs. 2C and D. In Structure C, the X moiety is as in Structure B, and the moiety Y may be methylene, sulfur, or preferably oxygen. In Structure D, the X and Y moieties are as in Structure B. In all subunits depicted in Figures 1 and 2, each P_i and P_j is a purine or pyrimidine base-pairing moiety effective to bind, by base-specific hydrogen bonding, to a base in a polynucleotide, and is preferably selected from adenine, cytosine, guanine and uracil.

30 As noted above, the substantially uncharged oligomer may advantageously include a limited number of charged linkages, e.g. up to about 1 per every 5 uncharged linkages. In the case of the morpholino oligomers, such a charged linkage may be a linkage as represented by any of Figs. 2A-D, preferably Fig. 2B, where X is oxide (-O⁻) or sulfide (-S⁻).

35 The antisense compounds of the invention can be synthesized by stepwise solid-phase synthesis, employing methods detailed in the references cited above. The sequence of subunit

additions will be determined by the selected base sequence (see Sections IID and IV below). In some cases, it may be desirable to add additional chemical moieties to the oligomer compounds, e.g. to enhance the pharmacokinetics of the compound or to facilitate capture or detection of the compound. Such a moiety may be covalently attached, typically to the 5'- or 3'-end of the 5 oligomer, according to standard synthesis methods. For example, addition of a polyethyleneglycol moiety or other hydrophilic polymer, e.g., one having 10-100 polymer subunits, may be useful in enhancing solubility. One or more charged groups, e.g., anionic charged groups such as an organic acid, may enhance cell uptake. A reporter moiety, such as fluorescein or a radiolabeled group, may be attached for purposes of detection. Alternatively, the 10 reporter label attached to the oligomer may be a ligand, such as an antigen or biotin, capable of binding a labeled antibody or streptavidin. In selecting a moiety for attachment or modification of an oligomer antisense, it is generally of course desirable to select chemical compounds of groups that are biocompatible and likely to be tolerated by a subject without undesirable side effects.

15 **IV. Exemplary Bacterial Targets**

Escherichia coli (*E. coli*) is a Gram negative bacteria that is part of the normal flora of the gastrointestinal tract. There are hundreds of strains of *E. coli*, most of which are harmless and live in the gastrointestinal tract of healthy humans and animals. Currently, there are four 20 recognized classes of enterovirulent *E. coli* (the "EEC group") that cause gastroenteritis in humans. Among these are the enteropathogenic (EPEC) strains and those whose virulence mechanism is related to the excretion of typical *E. coli* enterotoxins. Such strains of *E. coli* can cause various diseases including those associated with infection of the gastrointestinal tract and urinary tract, septicemia, pneumonia, and meningitis. Antibiotics are not effective against some strains and do not necessarily prevent recurrence of infection.

25 For example, *E. coli* strain 0157:H7 is estimated to cause 10,000 to 20,000 cases of infection in the United States annually (Federal Centers for Disease Control and Prevention). Hemorrhagic colitis is the name of the acute disease caused by *E. coli* O157:H7. Preschool children and the elderly are at the greatest risk of serious complications. *E. coli* strain 0157:H7 was recently reported as the cause of death of four children who ate under cooked hamburgers from a fast-food 30 restaurant in the Pacific Northwest.

Salmonella typhimurium are Gram negative bacteria which cause various conditions that range clinically from localized gastrointestinal infections and gastroenteritis (diarrhea, abdominal cramps, and fever) to enteric fevers (including typhoid fever) which are serious systemic illnesses. *Salmonella* infection also causes substantial losses of livestock.

Typical of Gram-negative bacilli, the cell wall of *Salmonella spp.* contains a complex lipopolysaccharide (LPS) structure that is liberated upon lysis of the cell and may function as an endotoxin, which contributes to the virulence of the organism.

Contaminated food is the major mode of transmission for non-typhoidal salmonella infection, 5 due to the fact that *Salmonella* survive in meats and animal products that are not thoroughly cooked. The most common animal sources are chickens, turkeys, pigs, and cows, in addition to numerous other domestic and wild animals. The epidemiology of typhoid fever and other enteric fevers caused by *Salmonella spp.* is associated with water contaminated with human feces.

Vaccines are available for typhoid fever and are partially effective; however, no vaccines are 10 available for non-typhoidal *Salmonella* infection. Non-typhoidal salmonellosis is controlled by hygienic slaughtering practices and thorough cooking and refrigeration of food. Antibiotics are indicated for systemic disease, and Ampicillin has been used with some success. However, in patients under treatment with excessive amounts of antibiotics, patients under treatment with immunosuppressive drugs, following gastric surgery, and in patients with hemolytic anemia, 15 leukemia, lymphoma, or AIDS, *Salmonella* infection remains a medical problem.

Pseudomonas spp. are motile, Gram-negative rods which are clinically important because 20 they are resistant to most antibiotics, and are a major cause of hospital acquired (nosocomial) infections. Infection is most common in: immunocompromised individuals, burn victims, individuals on respirators, individuals with indwelling catheters, IV narcotic users and individuals with chronic pulmonary disease (e.g., cystic fibrosis). Although infection is rare in healthy individuals, it can occur at many sites and lead to urinary tract infections, sepsis, pneumonia, pharyngitis, and numerous other problems, and treatment often fails with greater significant mortality.

Vibrio cholerae is a Gram negative rod which infects humans and causes cholera, a disease 25 spread by poor sanitation, resulting in contaminated water supplies. *Vibrio cholerae* can colonize the human small intestine, where it produces a toxin that disrupts ion transport across the mucosa, causing diarrhea and water loss. Individuals infected with *Vibrio cholerae* require rehydration either intravenously or orally with a solution containing electrolytes. The illness is generally self-limiting; however, death can occur from dehydration and loss of essential electrolytes. Antibiotics 30 such as tetracycline have been demonstrated to shorten the course of the illness, and oral vaccines are currently under development.

Neisseria gonorrhoeae is a Gram negative coccus, which is the causative agent of the common sexually transmitted disease, gonorrhea. *Neisseria gonorrhoeae* can vary its surface 35 antigens, preventing development of immunity to reinfection. Nearly 750,000 cases of gonorrhea are reported annually in the United States, with an estimated 750,000 additional unreported cases

annually, mostly among teenagers and young adults. Ampicillin, amoxicillin, or some type of penicillin used to be recommended for the treatment of gonorrhea. However, the incidence of penicillin-resistant gonorrhea is increasing, and new antibiotics given by injection, e.g., ceftriaxone or spectinomycin, are now used to treat most gonococcal infections.

5 *Staphylococcus aureus* is a Gram positive coccus which normally colonizes the human nose and is sometimes found on the skin. *Staphylococcus* can cause bloodstream infections, pneumonia, and surgical-site infections in the hospital setting (i.e., nosocomial infections). *Staph. aureus* can cause severe food poisoning, and many strains grow in food and produce exotoxins. 10 *Staphylococcus* resistance to common antibiotics, e.g., vancomycin, has emerged in the United States and abroad as a major public health challenge both in community and hospital settings. Recently a vancomycin-resistant *Staph. aureus* isolate has also been identified in Japan.

15 *Mycobacterium tuberculosis* is a Gram positive bacterium which is the causative agent of tuberculosis, a sometimes crippling and deadly disease. Tuberculosis is on the rise globally and is the leading cause of death from a single infectious disease (with a current death rate of three million people per year). It can affect several organs of the human body, including the brain, the kidneys and the bones; however, tuberculosis most commonly affects the lungs.

20 In the United States, approximately ten million individuals are infected with *Mycobacterium tuberculosis*, as indicated by positive skin tests, with approximately 26,000 new cases of active disease each year. The increase in tuberculosis (TB) cases has been associated with HIV/AIDS, homelessness, drug abuse and immigration of persons with active infections. Current treatment programs for drug-susceptible TB involve taking two or four drugs (e.g., isoniazid, rifampin, pyrazinamide, ethambutol or streptomycin) for a period of from six to nine months, because all of the TB germs cannot be destroyed by a single drug. In addition, the observation of drug-resistant and multiple drug resistant strains of *Mycobacterium tuberculosis* is on the rise.

25 *Helicobacter pylori* (*H. pylori*) is a micro-aerophilic, Gram negative, slow-growing, flagellated organism with a spiral or S-shaped morphology which infects the lining of the stomach. *H. pylori* is a human gastric pathogen associated with chronic superficial gastritis, peptic ulcer disease, and chronic atrophic gastritis leading to gastric adenocarcinoma. *H. pylori* is one of the most common chronic bacterial infections in humans and is found in over 90% of patients with active gastritis. 30 Current treatment includes triple drug therapy with bismuth, metronidazole, and either tetracycline or amoxicillin, which eradicates *H. pylori* in most cases. Problems with triple therapy include patient compliance, side effects, and metronidazole resistance. Alternate regimens of dual therapy which show promise are amoxicillin plus metronidazole or omeprazole plus amoxicillin.

35 *Streptococcus pneumoniae* is a Gram positive coccus and one of the most common causes of bacterial pneumonia as well as middle ear infections (otitis media) and meningitis. Each year in

the United States, pneumococcal diseases account for approximately 50,000 cases of bacteremia; 3,000 cases of meningitis; 100,000-135,000 hospitalizations; and 7 million cases of otitis media. Pneumococcal infection causes an estimated 40,000 deaths annually in the United States. Children less than 2 years of age, adults over 65 years of age, persons of any age with underlying medical conditions, including, e.g., congestive heart disease, diabetes, emphysema, liver disease, sickle cell, HIV, and those living in special environments, e.g., nursing homes and long-term care facilities, are at highest risk for infection.

5 Drug-resistant *S. pneumoniae* strains have become common in the United States, with many penicillin-resistant pneumococci also resistant to other antimicrobial drugs, such as erythromycin or trimethoprim-sulfamethoxazole.

10 *Treponema pallidum* is a spirochete which causes syphilis. *T. pallidum* is exclusively a pathogen which causes syphilis, yaws and non-venereal endemic syphilis or pinta. *Treponema pallidum* cannot be grown in vitro and does replicate in the absence of mammalian cells. The initial infection causes an ulcer at the site of infection; however, the bacteria move throughout the 15 body, damaging many organs over time. In its late stages, untreated syphilis, although not contagious, can cause serious heart abnormalities, mental disorders, blindness, other neurologic problems, and death.

12 Syphilis is usually treated with penicillin, administered by injection. Other antibiotics are available for patients allergic to penicillin, or who do not respond to the usual doses of penicillin. In all stages of syphilis, proper treatment will cure the disease, but in late syphilis, damage already done to body organs cannot be reversed.

15 *Chlamydia trachomatis* is the most common bacterial sexually transmitted disease in the United States, and it is estimated that 4 million new cases occur each year. The highest rates of infection are in 15 to 19 year olds. Chlamydia is a major cause of non-gonococcal urethritis (NGU), cervicitis, bacterial vaginitis, and pelvic inflammatory disease (PID). Chlamydia infections may have very mild symptoms or no symptoms at all; however, if left untreated, Chlamydia infections can lead to serious damage to the reproductive organs, particularly in women. Antibiotics such as azithromycin, erythromycin, ofloxacin, amoxicillin or doxycycline are typically prescribed to treat *Chlamydia* infection.

20 *Bartonella henselae*. Cat Scratch Fever (CSF) or cat scratch disease (CSD) is a disease of humans acquired through exposure to cats, caused by a Gram negative rod originally named *Rochalimaea henselae*, and currently known as *Bartonella henselae*. Symptoms include fever and swollen lymph nodes. CSF is generally a relatively benign, self-limiting disease in people; however, infection with *Bartonella henselae* can produce distinct clinical symptoms in 25 immunocompromised people, including acute febrile illness with bacteremia, bacillary

angiomatosis, peliosis hepatitis, bacillary splenitis, and other chronic disease manifestations such as AIDS encephalopathy.

The disease is treated with antibiotics, such as doxycycline, erythromycin, rifampin, penicillin, gentamycin, ceftriaxone, ciprofloxacin, and azithromycin.

5 *Haemophilus influenzae* (*H. influenza*) is a family of Gram negative bacteria; six types of which are known, with most *H. influenza*-related disease caused by type B, or "HIB". Until a vaccine for HIB was developed, HIB was a common causes of otitis media, sinus infections, bronchitis, the most common cause of meningitis, and a frequent culprit in cases of pneumonia, septic arthritis (joint infections), cellulitis (infections of soft tissues), and pericarditis (infections of 10 the membrane surrounding the heart). The *H. influenza* type B bacterium is widespread in humans and usually lives in the throat and nose without causing illness. Unvaccinated children under age 5 are at risk for HIB disease. Meningitis and other serious infections caused by *H. influenza* infection can lead to brain damage or death.

15 *Shigella dysenteriae* (*Shigella dys.*) is a Gram negative rod which causes dysentary. In the colon, the bacteria enter mucosal cells and divide within mucosal cells, resulting in an extensive inflammatory response. *Shigella* infection can cause severe diarrhea which may lead to dehydration and can be dangerous for the very young, very old or chronically ill. *Shigella dys.* forms a potent toxin (shiga toxin), which is cytotoxic, enterotoxic, and neurotoxic and acts as a inhibitor of protein synthesis. Resistance to antibiotics such as ampicillin and TMP-SMX has 20 developed; however, treatment with newer, more expensive antibiotics such as ciprofloxacin, norfloxacin and enoxacin, remains effective.

25 *Enterococcus faecium*. *Enterococci* are a component of the normal flora of the gastrointestinal and female urogenital tracts, however, recent studies indicate that pathogenic *Enterococci* can be transmitted directly in the hospital setting. (See, e.g., Boyce, *et al.*, *J Clin Microbiol* 32, 1148-53, 1994) *Enterococci*, have been recognized as a cause of nosocomial infection and some strains are resistant to multiple antimicrobial drugs. The most common *Enterococci*-associated nosocomial infections are urinary tract infections, post-surgical infections and bacteremia (Murray BE, *Clin Microbiol* 3, 46-65, Rev. 1990; Moellering RC Jr., *Clin Infect Dis* 14, 1173-8, 1992; Schaberg DR *et al.*, *Am J Med* 91(Suppl 3B), 72S-75S, 1991).

30 Vancomycin has been used extensively to treat *Enterococcus* infection since the late 1970s. Recently, a rapid increase in the incidence of infection and colonization with vancomycin-resistant enterococci (VRE) has been reported. The observed resistance is of concern due to (1) the lack of effective antimicrobial therapy for VRE infections because most VRE are also resistant to drugs previously used to treat such infections, *i.e.*, penicillin and aminoglycosides (CDC. MMWR 35 42:597-9, 1993; Handwerger, *et al.*, *Clin Infect Dis* 16, 750-5, 1993); and (2) the possibility that

the vancomycin-resistant genes present in VRE can be transferred to other gram-positive microorganisms.

Resistance to vancomycin and other glycopeptide antibiotics has been associated with the synthesis of a modified cell-wall precursor, terminating in D-lactate which has a lower affinity for antibiotics such as vancomycin.

5 *Listeria* is a genus of Gram-positive, motile bacteria found in human and animal feces.

Listeria monocytogenes causes such diseases as meningoencephalitis and meningitis. In cattle and sheep, listeria infection causes encephalitis and spontaneous abortion.

10 Veterinary applications. A healthy microflora in the gastro- intestinal tract of livestock is of vital importance for health and corresponding production of associated food products. As with humans, the gastrointestinal tract of a healthy animal contains numerous types of bacteria (i.e., *E. coli*, *Pseudomonas aeruginosa* and *Salmonella spp.*), which live in ecological balance with one another. This balance may be disturbed by a change in diet, stress, or in response to antibiotic or other therapeutic treatment, resulting in bacterial diseases in the animals generally caused by 15 bacteria such as *Salmonella*, *Campylobacter*, *Enterococci*, *Tularemia* and *E. coli*. Bacterial infection in these animals often necessitates therapeutic intervention, which has treatment costs as well being frequently associated with a decrease in productivity.

20 As a result, livestock are routinely treated with antibiotics to maintain the balance of flora in the gastrointestinal tract. The disadvantages of this approach are the development of antibiotic resistant bacteria and the carry over of such antibiotics into resulting food products for human consumption.

V. Exemplary 16S rRNA Antisense Oligomers

25 In one embodiment, the antisense oligomers of the invention are designed to hybridize to a region of a bacterial 16S rRNA nucleic acid sequence under physiological conditions, with a T_m substantially greater than 37°C, e.g., at least 50°C and preferably 60°C-80°C. The oligomer is designed to have high binding affinity to the nucleic acid and may be 100% complementary to the 16S rRNA nucleic acid target sequence, or it may include mismatches, as further described above.

30 In various aspects, the invention provides an antisense oligomer having a nucleic acid sequence effective to stably and specifically bind to a target sequence selected from the group consisting of 16S rRNA sequences which have one or more of the following characteristics: (1) a sequence found in a double stranded region of a 16s rRNA, e.g., the peptidyl transferase center, the alpha-sarcin loop and the mRNA binding region of the 16S rRNA sequence; (2) a sequence found in a single stranded region of a bacterial 16s rRNA; (3) a sequence specific to a particular

strain of a given species of bacteria, *i.e.*, a strain of *E. coli* associated with food poisoning; (4) a sequence specific to a particular species of bacteria; (5) a sequence common to two or more species of bacteria; (6) a sequence common to two related genera of bacteria (*i.e.*, bacterial genera of similar phylogenetic origin); (7) a sequence generally conserved among Gram-negative bacterial 16S rRNA sequences; (6) a sequence generally conserved among Gram-positive bacterial 16S rRNA sequences; or (7) a consensus sequence for bacterial 16S rRNA sequences in general.

Exemplary bacteria and associated GenBank Accession Nos. for 16S rRNA sequences are provided in Table 1, below.

Table 1

Organism	GenBank Reference for 16S rRNA	SEQ ID NO:
<i>Escherichia coli</i>	X80725	1
<i>Salmonella thyphimurium</i>	U88545	2
<i>Pseudomonas aeruginosa</i>	AF170358	3
<i>Vibrio cholera</i>	AF118021	4
<i>Neisseria gonorrhoea</i>	X07714	5
<i>Staphylococcus aureus</i>	Y15856	6
<i>Mycobacterium tuberculosis</i>	X52917	7
<i>Helicobacter pylori</i>	M88157	8
<i>Streptococcus pneumoniae</i>	AF003930	9
<i>Treponema palladium</i>	AJ010951	10
<i>Chlamydia trachomatis</i>	D85722	11
<i>Bartonella henselae</i>	X89208	12
<i>Hemophilis influenza</i>	M35019	13
<i>Shigella dysenteriae</i>	X96966	14

10

It will be understood that one of skill in the art may readily determine appropriate targets for antisense oligomers, and design and synthesize antisense oligomers using techniques known in the art. Targets can be identified by obtaining the sequence of a target 16S or 23S nucleic acid of interest (*e.g.* from GenBank) and aligning it with other 16S or 23S nucleic acid sequences using, for example, the MacVector 6.0 program, a ClustalW algorithm, the BLOSUM 30 matrix, and default parameters, which include an open gap penalty of 10 and an extended gap penalty of 5.0 for nucleic acid alignments. An alignment may also be carried out using the Lasergene99 MegAlign Multiple Alignment program with a ClustalW algorithm run under default parameters.

For example, given the 16s rRNA sequences provided in Table 1 and other 16s rRNA sequences available in GenBank, one of skill in the art can readily align the 16s rRNA sequences of interest and determine which sequences are conserved among one or more different bacteria, and those which are specific to one or more particular bacteria. A similar alignment can be performed on 23S rRNA sequences.

As an illustration, the 16S rRNA sequences from the organisms shown in Table 1 were aligned using the Lasergene 99 MegAlign Multiple Alignment program, with a ClustalW algorithm and default parameters. Tables 2-5 show exemplary oligomers antisense to 16S rRNA of these bacterial species, including sequences targeting individual bacteria, multiple bacteria, and broad spectrum sequences. These oligomers were derived from the sequences in Table 1 and from the alignment performed as described above. As the Tables show, a number of sequences were 5 conserved among different organisms.

Exemplary oligomers antisense to *E. coli* 16S rRNA (SEQ ID NO:32 and SEQ ID NO:35) were designed based on the sequence found at GenBank Accession No. X80725. Further 10 exemplary oligomers antisense to *E. coli* 16S rRNA and one or more other bacterial 16S rRNA sequences are provided in Table 2A.

Exemplary oligomers antisense to *Salmonella thyphimurium* 16S rRNA (SEQ ID NO:18 and SEQ ID NO:36) were designed based on the sequence found at GenBank Accession No. U88545. Further exemplary oligomers antisense to *S. thyphi*, 16S rRNA and one or more other bacterial 16S 15 rRNA sequences are provided in Table 2A.

Exemplary oligomers antisense to *Pseudomonas aeruginosa* 16S rRNA (SEQ ID NO:40, SEQ ID NO:41, SEQ ID NO:42 and SEQ ID NO:43) were designed based on the sequence found at GenBank Accession No. AF170358.

Exemplary oligomers antisense to *Vibrio cholera* 16S rRNA (SEQ ID NO:45, SEQ ID NO:46 20 and SEQ ID NO:47) were designed based on the sequence found at GenBank Accession No. AF118021. A further exemplary oligomer, antisense to *Vibrio cholera* 16S rRNA and other bacterial 16S rRNA sequences (SEQ ID NO:44), is provided in Table 2A.

25 **Table 2A. BACTERIAL 16S rRNA SEQUENCES AND ANTISENSE OLIGOMERS**

Organism	GenBank Reference	Native sequence	Antisense oligomer
<i>E. coli</i> (NS-1) <i>Shigella dys.</i>	X80725 nt 446-466 X96966 nt 436-456	GAGTAAAGTTAAT ACCTTTGC	GCAAAGGTATTAA CTTTACT (SEQ ID NO:17)
<i>E. coli</i> (BS-1) <i>S. thyphi</i> <i>Shigella dys.</i>	X80725 nt 1270-1290 U88545 nt 1282-1302 X96966 nt 1263-1283	TCATAAAGTGCCT CGTAGTCC	GGACTACGACGCA CTTTATGAG (SEQ ID NO:15)
<i>E. coli</i> <i>S. thyphi</i> <i>H. influenza</i>	X80725 nt 1-21 U88545 nt 10-30 M35019 nt 10-30	AGTTTGATCATGG CTCAGATT	AATCTGAGCCATG ATCAAAT (SEQ ID NO:31)
<i>E. coli</i>	X80725 nt 173-193	ACGTCCGAAGCAC AAAGAGGG	CCCTCTTGTGCT TGCACGCT (SEQ ID NO:32)
<i>E. coli</i> <i>S. thyphi</i> <i>Shigella dys.</i>	X80725 nt 643-663 U88545 nt 652-672 X96966 nt 653-673	TTGAGTCTCGTAG AGGGGGGT	ACCCCCCTCTACG AGACTCAA (SEQ ID NO:33)

<i>E. coli</i> <i>S. thyphi</i> <i>Shigella dys.</i>	X80725 nt 823-843 U88545 nt 832-852 X96966 nt 813-833	GGTTGTGCCCTTG AGGCGTGG	CCACGCCCTCAAGG GCACAACC (SEQ ID NO:34)
<i>E. coli</i>	X80725 nt 991-1011	CGGAAGTTTCAG AGATGAGA	TCTCATCTCTGAA AACTTCCG (SEQ ID NO:35)
<i>S. thyphi</i> (NS-2)	U88545 nt 455-475	GTTGTGGTTAATA ACCGCAGC	GCTGCGGTTATTAA ACCACAAC (SEQ ID NO:18)
<i>S. thyphi.</i> (BS-2) <i>E. coli</i> <i>Shigella dys.</i>	U88545 nt 1261-1281 X80725 nt 1252-1272 X96966 nt 1242-1262	CCTCGCGAGAGCA AGCGGACC	GGTCCGCTTGCTC TCGCGAGG (SEQ ID NO:16)
<i>S. thyphi.</i>	U88545 nt 1-21	AAATTGAAGAGTT TGATCATG	CATGATCAAACTC TTCAATT (SEQ ID NO:36)
<i>S. thyphi.</i> <i>Shigella dys.</i>	U88545 nt 181-201 X96966 nt 162-182	ACGTCGCAAGACC AAAGAGGG	CCCTCTTGGTCT TGCACGT (SEQ ID NO:37)
<i>S. thyphi.</i> <i>E. coli</i> <i>Shigella dys.</i>	U88545 nt 652-672 X80725 nt 643-663 X96966 nt 633-653	TGAGTCTCGTAGA GGGGGGTA	TACCCCCCTCTAC GAGACTCA (SEQ ID NO:38)
<i>S. thyphi.</i> <i>E. coli</i> <i>Shigella dys.</i>	U88545 nt 832-852 X80725 nt 823-843 X96966 nt 813-833	GTTGTGCCCTTG GGCGTGGC	GCCACGCCCTCAAG GGCACAAAC (SEQ ID NO:39)
<i>P. aeruginosa</i>	AF170358 nt 1-21	ATGAAGAGGGCTT GCTCTCTG	CAGAGAGCAAGC CCTCTTCAT (SEQ ID NO:40)
<i>P. aeruginosa</i>	AF170358 nt 107-127	CGTCCTACGGGAG AAAGCAGG	CCTGCTTTCTCCC GTAGGACG (SEQ ID NO:41)
<i>P. aeruginosa</i>	AF170358 nt 578-598	AGAGTATGGCAGA GGGTGGTG	CACCAACCCTCTGC CATACTCT (SEQ ID NO:42)
<i>P. aeruginosa</i>	AF170358 nt 758-778	TTGGGATCCTTGA GATCTTAG	CTAAGATCTCAAG GATCCCAA (SEQ ID NO:43)
<i>Vibrio cholera</i> <i>E. coli</i> <i>H. influenza</i> <i>S. thyphi.</i> <i>Shigella dys.</i>	AF118021 nt 1-21 X80725 nt 19-39 M35019 nt 26-46 U88545 nt 18-48 X96966 nt 9-29	ATTGAACGCTGGC GGCAGGCC	GGCCTGCCGCCAG CGTTCAAT (SEQ ID NO:44)
<i>Vibrio cholera</i>	AF118021 nt 157-177	ATGTTTACGGACC AAAGAGGG	CCCTCTTGGTCC GTAAACAT (SEQ ID NO:45)
<i>Vibrio cholera</i>	AF118021 nt 625-645	GCTAGAGTCTTGT AGAGGGGG	CCCCCTCTACAAG ACTCTAGC (SEQ ID NO:46)
<i>Vibrio cholera</i>	AF118021 nt 805-825	GAGGTTGTGACCT ARAGTCGT	ACGACTYTAGGTC ACAACCTC (SEQ ID NO:47)

1: Approximate nucleotide locations

Exemplary oligomers antisense to *Neisseria gonorrhoea* 16S rRNA (SEQ ID NO:48, SEQ ID NO:49, SEQ ID NO:50 and SEQ ID NO:51) were designed based on the sequence found at GenBank Accession No. X07714. These are shown in Table 2B, below.

Exemplary oligomers antisense to *Staphylococcus aureus* 16S rRNA (SEQ ID NO:53, SEQ ID NO:54 and SEQ ID NO:55) were designed based on the sequence found at GenBank Accession No. Y15856. A further exemplary oligomer, antisense to a *Staph. aureus* 16S rRNA and a *Bartonella henselae* 16S rRNA sequence (SEQ ID NO:52), is provided in Table 2B, below.

Exemplary oligomers antisense to *Mycobacterium tuberculosis* 16S rRNA (SEQ ID NO:56, SEQ ID NO:57, SEQ ID NO:58 and SEQ ID NO:59) were designed based on the sequence found at GenBank Accession No. X52917.

Exemplary oligomers antisense to *Helicobacter pylori* 16S rRNA (SEQ ID NO:60, SEQ ID NO:61, SEQ ID NO:62 and SEQ ID NO:63) were designed based on the sequence found at GenBank Accession No. M88157.

Exemplary oligomers antisense to *Streptococcus pneumoniae* 16S rRNA (SEQ ID NO:64, SEQ ID NO:65, SEQ ID NO:66 and SEQ ID NO:67) were designed based on the sequence found at GenBank Accession No. AF003930.

Exemplary oligomers antisense to *Treponema palladium* 16S rRNA (SEQ ID NO:69, SEQ ID NO:70 and SEQ ID NO:71) were designed based on the sequence found at GenBank Accession No. AJ010951. A further exemplary oligomer, antisense to *Treponema palladium* 16S rRNA and other 16S rRNA sequences (SEQ ID NO:68), is provided in Table 2B, below.

Table 2B. BACTERIAL 16s rRNA SEQUENCES AND ANTISENSE OLIGOMERS

Organism	GenBank Reference	Native sequence	Antisense oligomer
<i>N. gonorrhoea</i>	X07714 nt 1-21	TGAACATAAGAGT TTGATCCT	AGGATCAAACCTTTATGTTCA (SEQ ID NO:48)
<i>N. gonorrhoea</i>	X07714 nt 183-203	CGTCTTGAGAGGG AAAGCAGG	CCTGCTTCCCTCTCAAGACG (SEQ ID NO:49)
<i>N. gonorrhoea</i>	X07714 nt 654-674	CGAGTGTGTCAGA GGGAGGTG	CACCTCCCTCTGACACACTCG (SEQ ID NO:50)
<i>N. gonorrhoea</i>	X07714 nt 834-854	TTGGGCAACTTGA TTGCTTGG	CCAAGCAATCAAGTTGCCAA (SEQ ID NO:51)
<i>Staph. aureus</i>	Y15856 nt 1-21	CTGGCTCAGGATG AACGCTGG	CCAGCGTTCATCCTGAGCCAG (SEQ ID NO:52)
<i>Bartonella hens</i>	X89208 nt 3-23		
<i>Staph. aureus</i>	Y15856 nt 163-183	ATATTTGAACCG CATGGTTC	GAACCATGCGGTTCAAAATAT (SEQ ID NO:53)
<i>Staph. aureus</i>	Y15856 nt 640-660	CTTGAGTGCAGAA GAGGAAAG	CTTCCTCTTCTGCACTCAAG (SEQ ID NO:54)
<i>Staph. aureus</i>	Y15857 nt 447-466		ATGTGCACAGTTACTTACAC avi ref no. 23
<i>Staph. aureus</i>	Y15857 nt 1272-1291		CTGAGAACAACTTTATGGGA avi ref no. 24

<i>Staph. aureus</i>	Y15856 nt 819-839	GTGTTAGGGGGTT TCCGCC	GGGGCGGAAACCCCTAACAC (SEQ ID NO:55)
<i>Myco. tubercul.</i>	X52917 nt 1-21	GGCGCGTGTCTTA ACACATGC	GCATGTGTTAACGACGCCGCC (SEQ ID NO:56)
<i>Myco. tubercul.</i>	X52917 nt 138-158	GGACCACGGGATG CATGTCTT	AAGACATGCATCCGTGGTCC (SEQ ID NO:57)
<i>Myco. tubercul.</i>	X52917 nt 604-624	AGAGTACTGCAGG GGAGACTG	CAGTCTCCCTGCAGTACTCT (SEQ ID NO:58)
<i>Myco. tubercul.</i>	X52917 nt 784-804	TGGGTTTCTTCCT TGGGATC	GATCCAAGGAAGGAAACCCA (SEQ ID NO:59)
<i>H. pylori</i>	M88157 nt 1-21	TTTATGGAGAGTT TGATCCTG	CAGGATCAAACCTCCATAAA (SEQ ID NO:60)
<i>H. pylori</i>	M88157 nt 181-201	ACTCCTACGGGG AAAGATT	AAATCTTCCCCGTAGGAGT (SEQ ID NO:61)
<i>H. pylori</i>	M88157 nt 613-633	AGAGTGTGGGAGA GGTAGGTG	CACCTACCTCTCCCACACTCT (SEQ ID NO:62)
<i>H. pylori</i>	M88157 nt 794-814	TTGGAGGGCTTAG TCTCTCCA	TGGAGAGACTAAGCCCTCCAA (SEQ ID NO:63)
<i>Strep. pneumoniae</i>	AF003930 nt 1-21	ATTTGATCCTGGC TCAGGACG	CGTCCTGAGCCAGGATCAAAT (SEQ ID NO:64)
<i>Strep. pneumoniae</i>	AF003930 169-189	AGAGTGGATGTTG CATGACAT	ATGTCATGCAACATCCACTCT (SEQ ID NO:65)
<i>Strep. pneumoniae</i>	AF003930 646-666	TTGAGTGCAAGAG GGGAGAGT	ACTCTCCCTCTTGCACCAA (SEQ ID NO:66)
<i>Strep. pneumoniae</i>	AF003930 826-846	GTTAGACCTTTC CGGGGTTT	AAACCCCGGAAGGGTCTAAC (SEQ ID NO:67)
<i>Treponema pallad.</i>	AJ010951 nt 1-21	AGAGTTGATCAT GGCTCAGA	TCTGAGCCATGATCAAACCTCT (SEQ ID NO:68)
<i>S. typhi.</i>	U88545 nt 8-28		
<i>H. influenza</i>	M35019 nt 8-28		
<i>Treponema pallad.</i>	AJ010951 nt 173-193	ACTCAGTGCTTCA TAAGGGGT	ACCCCTTATGAAGCACTGAGT (SEQ ID NO:69)
<i>Treponema pallad.</i>	AJ010951 nt 651-671	TTGAATTACGGAA GGGAAACT	AGTTTCCCTCCGTAATTCAA (SEQ ID NO:70)
<i>Treponema pallad.</i>	AJ010951 nt 831-851	GTTGGGGCAAGAG CTTCAGTG	CACTGAAGCTTGCACCAA (SEQ ID NO:71)

2 Approximate nucleotide locations

Exemplary oligomers antisense to *Chlamydia trachomatis* 16S rRNA (SEQ ID NO:72, SEQ ID NO:73, SEQ ID NO:74 and SEQ ID NO:75) were designed based on the sequence found at GenBank Accession No. D85722. These are shown in Table 2C, below.

Exemplary oligomers antisense to *Bartonella henselae* 16S rRNA (SEQ ID NO:76, SEQ ID NO:77, SEQ ID NO:78 and SEQ ID NO:79) were designed based on the sequence found at GenBank Accession No. X89208.

Exemplary oligomers antisense to *Hemophilis influenza* 16S rRNA (SEQ ID NO:81, SEQ ID NO:82 and SEQ ID NO:83) were designed based on the sequence found at GenBank Accession No. M35019. A further exemplary oligomer, antisense to a *H. influenza* 16S rRNA sequence and a *Salmonella typhimurium* 16S rRNA sequence (SEQ ID NO:80), is provided in Table 2C, below.

An exemplary oligomer antisense to *Shigella dysenteriae* 16S rRNA (SEQ ID NO:88) was designed based on the sequence found at GenBank Accession No. X96966. Further exemplary antisense oligomers antisense to *Shigella dys* 16S rRNA and one or more other bacterial 16S rRNA sequences are provided in Table 2C.

5

Table 2C. BACTERIAL 16S rRNA SEQUENCES AND ANTISENSE OLIGOMERS

Organism	GenBank Reference	Native sequence	Antisense oligomer
<i>Chlamydia trach.</i>	D85722 nt 1-21	CTGAGAATTGATCTTGGTTC	GAACCAAGATCAAATTCTCAG (SEQ ID NO:72)
<i>Chlamydia trach.</i>	D85722 nt 176-196	ATATTGGGCATC CGAGTAAC	GTTACTCGGATGCCAAATAT (SEQ ID NO:73)
<i>Chlamydia trach.</i>	D85722 nt 658-678	AGAGGGTAGATG GAGAAAAGG	CCTTTCTCCATCTACCCCTCT (SEQ ID NO:74)
<i>Chlamydia trach.</i>	D85722 nt 838-858	TGGATGGTCTCA ACCCCATCC	GGATGGGGTTGAGACCATCCA (SEQ ID NO:75)
<i>Bartonella hens.</i>	X89208 nt 1-21	TCCTGGCTCAGG ATGAACGCT	AGCGTTCATCCTGAGCCAGGA (SEQ ID NO:76)
<i>Bartonella hens.</i>	X89208 nt 149-169	CGTCCTACTGGA GAAAGATT	AAATCTTCTCCAGTAGGACG (SEQ ID NO:77)
<i>Bartonella hens.</i>	X89208 nt 581-601	TGAGTATGGAAG AGGTGAGT	CACTCACCTCTCCATACTCA (SEQ ID NO:78)
<i>Bartonella hens.</i>	X89208 nt 761-781	TTGGGTGGTTTAC TGCTCAGT	ACTGAGCAGTAAACCACCCAA (SEQ ID NO:79)
<i>H. influenza</i>	M35019 nt 2-21	AATTGAAGAGTT	CATGATCAAACCTTTCAATTN (SEQ ID NO:80)
<i>S. thyphi.</i>	U88545 nt 2-21	TGATCATG	
<i>H. influenza</i>	M35019 nt 180-200	TATTATCGGAAG ATGAAAGTG	CACTTCATCTCCGATAATA (SEQ ID NO:81)
<i>H. influenza</i>	M35019 nt 649-669	AACTAGAGTACT TTAGGGAGG	CCTCCCTAAAGTACTCTAGTT (SEQ ID NO:82)
<i>H. influenza</i>	M35019 nt 829-849	GGGGGTGGGGTT TAACTCTG	CAGAGTTAAACCCCAACCCCC (SEQ ID NO:83)
<i>Shigella dys.</i>	X96966 nt 1-21	TGGCTCAGATTG AACGCTGGC	GCCAGCGTTCAATCTGAGCCA (SEQ ID NO:84)
<i>E. coli</i>	X80725 nt 11-31		
<i>S. thyphi.</i>	X96966 nt 20-40		
<i>N. gonorrhoea</i>	X07714 nt 21-41		
<i>H. influenza</i>	M35019 nt 20-40		
<i>Shigella dys.</i>	X96966 nt 162-182	ACGTCGCAAGAC	CCCTCTTGGTCTTGCACGT (SEQ ID NO:85)
<i>S. thyphi.</i>	X96966 nt 181-201	CAAAGAGGG	
<i>Shigella dys.</i>	X96966 nt 633-653	TGAGTCTCGTAG	TACCCCCCTCTACGAGACTCA (SEQ ID NO:86)
<i>E. coli</i>	X80725 nt 644-664	AGGGGGGTA	
<i>S. thyphi.</i>	X96966 nt 652-672		
<i>Shigella dys.</i>	X96966 nt 813-833	GTTGTGCCCTTGA	GCCACGCCTCAAGGGCACAAC (SEQ ID NO:87)
<i>E. coli</i>	X80725 nt 824-844	GGCGTGGC	

<i>S. thypi.</i>	X96966 nt 832-852		
<i>Shigella dys.</i>	X96966 nt 983-1003	GAACCTTGTAGA GATACGAGG	CCTCGTATCTCTACAAGGTT (SEQ ID NO:88)

3 Approximate nucleotide locations

Exemplary Gram-positive bacterial targets include, but are not limited to, *Staphylococcus aureus*, *Mycobacterium tuberculosis* and *Streptococcus pneumoniae*.

5 Exemplary oligomer sequences antisense to Gram-positive bacterial 16S rRNA sequences are exemplified in Table 3 by the sequences presented as SEQ ID NO:27 and SEQ ID NO:28, with the bacterial 16s rRNAs to which the exemplary antisense oligomers are targeted indicated in Table 3 as "+" and those which are not targeted indicated as "-".

10 **Table 3. GRAM POSITIVE 16s rRNA SEQUENCES AND ANTISENSE OLIGOMERS**

Organism	SEQUENCE	AACTACGTGCCAGC AGCCGCG	TCGTGAGATGTTGG GTTAAGT
	ANTISENSE	CGCGGCTGCTGGCA CGTAGTT	ACTTAACCCAACATC TCACGA
<i>Staph aureus</i>	Y15856	+	+
<i>Myco. tubercul.</i>	X52917	+	+
<i>Strep. pneumoniae</i>	AF003930	+	+
<i>E. coli</i>	X80725	-	-
<i>S. thypi</i>	U88545	-	-
<i>P. aeruginosa</i>	AF170358	-	+
<i>Vibrio cholera</i>	AF118021	-	-
<i>N. gonorrhoea</i>	X07714	+	+
<i>H. pylori</i>	M88157	-	+
<i>Treponema pallad.</i>	AJ010951	-	-
<i>Chlamydia trach.</i>	D85722	-	-
<i>Bartonella hens</i>	X89208	-	+
<i>H. influenza</i>	M35019	-	-
<i>Shigella dys.</i>	X96966	-	-

4 Based on nucleotides 497-517 of GenBank Y15856, designated SEQ ID NO: 27

5 Based on nucleotides 1064-1084 of GenBank Y15856, designated SEQ ID NO: 28

15 Exemplary Gram-negative bacterial targets include, but are not limited to, *E. coli*, *Salmonella thymurium*, *Pseudomonas aeruginosa*, *Vibrio cholera*, *Neisseria gonorrhoea*, *Helicobacter pylori*, *Bartonella henselae*, *Hemophilis Influenza* and *Shigella dysenteriae*.

20 Exemplary oligomer sequences antisense to Gram-negative bacterial 16S rRNA sequences are exemplified in Table 4 by the sequences presented as SEQ ID NO:29 and SEQ ID NO:30, with the bacterial 16s rRNAs to which the exemplary antisense oligomers are targeted indicated in Table 4 as "+" and those which are not targeted indicated as "-".

Table 4. GRAM NEGATIVE 16s rRNA SEQUENCES AND ANTISENSE OLIGOMERS

Organism	SEQUENCE	TCGGAATTACTGGGC GTAAA	CCGCCCCGTACACCAT GGGAGT
	ANTISENSE	TTTACGCCAGTAATT CCGA	ACTCCCATGGTGTGACG GGCGG
<i>E. coli</i>	X80725	+	+
<i>S. thyphi</i>	U88545	+	+
<i>P. aeruginosa</i>	AF170358	+	+
<i>Vibrio cholera</i>	AF118021	+	+
<i>N. gonorrhoea</i>	X07714	+	+
<i>Staph aureus</i>	Y15856	-	-
<i>Myco. tubercul.</i>	X52917	-	-
<i>H. pylori</i>	M88157	-	+
<i>Strep. pneumoniae</i>	AF003930	-	-
<i>Treponema pallad.</i>	AJ010951	-	+
<i>Chlamydia trach.</i>	D85722	-	+
<i>Bartonella hens</i>	X89208	-	+
<i>H. influenza</i>	M35019	-	+
<i>Shigella dys.</i>	X96966	+	+

6 Based on nucleotides 546-566 of GenBank X80725, designated SEQ ID NO: 29

7 Based on nucleotides 1389-1409 of GenBank X80725, designated SEQ ID NO: 30

5 Exemplary bacterial targets for broad spectrum antisense oligomers include, but are not limited to, *E. coli*, *Salmonella* *thyphimurium*, *Pseudomonas aeruginosa*, *Vibrio cholera*, *Neisseria gonorrhoea*, *Helicobacter pylori*, *Bartonella henselae*, *Hemophilis Influenza*, *Shigella dysenteriae*, *Staphylococcus aureus*, *Mycobacterium tuberculosis*, *Streptococcus pneumoniae*, *Treponema palladium* and *Chlamydia trachomatis*. (See Table 1.)

10 Exemplary broad spectrum antisense oligomers are presented in Tables 5A and 5B as SEQ ID NOs:21-25, with the bacterial 16s rRNAs to which the exemplary antisense oligomers are targeted indicated in Tables 5A and 5B as "+" and those which are not targeted indicated as "-".

Table 5A. BROAD SPECTRUM ANTISENSE OLIGONUCLEOTIDE SEQUENCES

Organism	SEQUENCE	AGACTCCTACGG GAGGCAGCA	CGTGCCAGCAGC CGCGGTAAAT	AACAGGATTAG ATACCCCTGGT
	ANTISENSE	TGCTGCCTCCCGT AGGAGTCT	ATTACCGCGGCT GCTGGCACG	ACCAGGGTATC TAATCCTGTT
<i>E. coli</i>	X80725	+	+	+
<i>S. thyphi</i>	U88545	+	+	+
<i>P. aeruginosa</i>	AF170358	+	+	+
<i>Vibrio cholera</i>	AF118021	+	+	+
<i>N. gonorrhoea</i>	X07714	+	+	+
<i>Staph. aureus</i>	Y15856	+	+	+
<i>Myco. tubercul.</i>	X52917	+	+	+
<i>H. pylori</i>	M88157	+	+	+
<i>Strep. pneumoniae</i>	AF003930	+	+	+
<i>Treponema pallad.</i>	AJ010951	+	+	+

<i>Chlamydia trach.</i>	D85722	+	+	+
<i>Bartonella hens</i>	X89208	+	+	+
<i>H. influenza</i>	M35019	+	+	+
<i>Shigella dys.</i>	X96966	+	+	+

8: based on nucleotides 327-347 of GenBank No. X80725, designated SEQ ID NO:21

9: based on nucleotides 504-524 of GenBank No. X80725, designated SEQ ID NO:22

10: based on nucleotides 781-801 of GenBank No. X80725, designated SEQ ID NO:23

5

Table 5B. BROAD SPECTRUM ANTISENSE OLIGONUCLEOTIDE SEQUENCES

Organism	SEQUENCE	GCACAAGCGGTGGA GCATGTG	ATGTTGGGTTAAGT CCCGCAA
	ANTISENSE	CACATGCTCCACCG CTTGTGC	TTGCGGGACTTAAC CCAACAT
<i>E. coli</i>	X80725	+	+
<i>S. typhi</i>	U88545	+	+
<i>P. aeruginosa</i>	AF170358	+	-
<i>Vibrio cholera</i>	AF118021	+	+
<i>N. gonorrhoea</i>	X07714	-	+
<i>Staph aureus</i>	Y15856	+	+
<i>Myco. tubercul.</i>	X52917	-	+
<i>H. pylori</i>	M88157	-	+
<i>Strep. pneumoniae</i>	AF003930	+	+
<i>Treponema pallad.</i>	AJ010951	+	-
<i>Chlamydia trach.</i>	D85722	-	-
<i>Bartonella hens</i>	X89208	+	+
<i>H. influenza</i>	M35019	+	+
<i>Shigella dys.</i>	X96966	+	+

11: based on nucleotides 924-944 of GenBank No. X80725, designated SEQ ID NO:24

12: based on nucleotides 1072-1092 of GenBank No. X80725, designated SEQ ID NO:25.

10

VI. Inhibitory Activity of Antisense Oligomers

A. Effect of Antisense Oligomers to Bacterial 16S rRNA on Bacterial Growth

The effect of PMO antisense oligomers on bacterial culture viability was tested using the protocol described below; see "Bacterial Cultures" in Materials and Methods. Briefly, test oligonucleotides, diluted in phosphate buffered saline (PBS), are added to the freshly inoculated bacterial cultures; the cultures are incubated at 37°C overnight, e.g., 6 to 26 hours, diluted, and plated on agar plates; colonies are counted 16-24 hours later. Non-selective bacterial growth media, e.g., agar containing nutrients appropriate to the type of bacteria being cultured, are utilized, as generally known in the art.

20 The viability of bacteria following overnight culture with a test oligomer is based on the number of bacterial colonies in antisense oligomer-treated cultures relative to untreated or

nonsense treated cultures. An exemplary nonsense control is an oligomer antisense to c-myc, having the sequence presented as SEQ ID NO: 139.

5 A1. Inhibition of *Salmonella typhimurium* with a Conserved-Sequence Oligomer Antisense to 16S rRNA. Two strains of *Salmonella typhimurium* (1535 and 1538) were inoculated into
broth media, as described in Materials and Methods, below. An oligomer antisense to a 16S
rRNA sequence conserved amongst *E. coli*, *S. typhimurium* and *S. dysenteriae* ("BS-1"; SEQ ID
NO:15) was added to a final concentration of 1 μ M and the tube placed in an incubator at 37°C for
6 to 16 hours. At the end of the incubation, the broth was spread onto plates, incubated overnight
for 16 to 24 hours and colonies counted. The data, shown in Table 6, provides evidence that
10 *Salmonella typhimurium* is inhibited by a 16S rRNA antisense oligomer based on a 16S rRNA
sequence which is conserved amongst *E. coli*, *S. typhimurium* and *S. dysenteriae*.

Table 6. Effect of Broad Spectrum Antisense on *Salmonella typhimurium*

Strain (culture time)	Control (colonies)	1 μ M AS to 16S rRNA (colonies)	% Inhibition
1535 (6 hours)	217	141	35
1535 (16 hours)	214	52	76
1538 (6 hours)	824	664	19
1538 (16 hours)	670	133	80

15 A2. Effect of Antisense Oligomers to Bacterial 16S rRNA on Growth Of *E. coli*.

The effect of PMO antisense oligomers on inhibition of *E. coli* was evaluated, using a procedure such as described above, by adding an antisense oligomer targeting particular 20-22 nucleotide portions of the *E. coli* 16S rRNA sequence found at GenBank Accession No. X80725 to individual *E. coli* cultures. Each antisense oligomer was incubated at a 1 μ M concentration with *E. coli* bacteria for 16 hours, the cultures were diluted and plated on agar plates, and colonies were counted 16-24 hours later. The results, shown in Table 7, indicate that PMO antisense oligomers targeting *E. coli* 16S rRNA inhibited growth of colonies by up to 60%, with oligomers targeting various regions throughout the 16S rRNA sequence observed to be effective.

25 **Table 7. *E. coli* 16s rRNA Targeting Study**

AVI Ref. No.	Location	Antisense sequence (5'→3')	SEQ ID NO.	Percent Inhibition	S.E.	Repeats (n)
9	1263-1283	GCA CTT TAT GAG GTC CGC TTG	19	59.8	3.4	8
15	1272-1293	GGA CTA CGA CGC ACT TTA TGA G	15	19.5	7.4	9
16	1252-1272	GGT CCG CTT GCT CTC GCG AGG	16	21.5	11	9
17	446-466	GCA AAG GTA TTA ACT TTA CTC	17	66	3.3	14

27	1-20	ATC TGA GCC ATG ATC AAA CT	97	55.2	9.7	5
28	301-320	TGT CTC AGT TCC AGT GTT GC	98	35	7.2	8
29	722-741	GTC TTC GTC CAG GGG GCC GC	99	52.5	4	7
30	1021-1040	CAC CTG TCT CAC GGT TCC CG	100	56	8.4	5
31	1431-1450	CGC CCT CCC GAA GTT AAG CT	101	43	13	5

Figure 5 depicts the results of a study on the effect of various concentrations of the PMO having SEQ ID NO: 15 (broad spectrum) targeted against a bacterial 16S rRNA consensus sequence on the bacterial colony formation in *E. coli*, presented as percent inhibition of colony formation. As the figure shows, about 70% inhibition was achieved at about 0.1 μ M PMO.

A3. Inhibition of *Staphylococcus aureus* and *Pseudomonas aeruginosa* with Oligomers Antisense to 16S rRNA.

Tables 8 and 9 show the effect of oligomers targeting 16S rRNA, at a concentration of 1 μ M, on bacterial growth in *Staphylococcus aureus* and *Pseudomonas aeruginosa*. In a typical experiment, antisense oligomers targeting particular 22-nucleotide portions of the *Staphylococcus aureus* and *Pseudomonas aeruginosa* 16S rRNA sequences, found at GenBank Accession Nos. Y15857 and Z76651, respectively, were incubated with the respective bacteria at a concentration of 1 μ M for 16 hours. Growth of *S. aureus* was inhibited by up to 25%, and growth of *P. aeruginosa* was inhibited by up to about 53%.

Table 8. *Staphylococcus aureus* 16s rRNA Targeting Study

AVI Ref. No.	Location	Antisense sequence (5'→3')	SEQ ID NO	Percent Inhibition	S.E.	n =
23	447-466	ATG TGC ACA GTT ACT TAC AC	93	2.5	8.6	2
24	1272-1291	CTG AGA ACA ACT TTA TGG GA	94	25.3	11	2

Table 9. *Pseudomonas aeruginosa* 16s rRNA Targeting Study

AVI Ref. No.	Location	Antisense sequence (5'→3')	SEQ ID NO:	Percent Inhibition	S.E.	n =
25	447-466	TTA TTC TGT TGG TAA CGT CA	95	37.3	9.8	3
26	1272-1291	CG AGT TGC AGA CTG CGA TC	96	52.7	7.1	3

Inhibition of *Listeria* was also demonstrated by a corresponding anti-16S PMO. A very low dose (about 30 nM) of the PMO gave about 40% inhibition.

5 A4. Effect of Antisense Oligomers to Bacterial rRNA on Growth Of Vancomycin-Resistant *Enterococcus faecium* (VRE)

(a) Bacterial 16S rRNA Targets

The effect of PMO antisense oligomers on the growth of VRE was evaluated, using the method described above, by adding antisense PMO's targeting numerous 16S rRNA sequences to cultures of VRE and incubating at a concentration of 1 μ M for 16 hours. The results shown in 10 Table 10 and in Figure 6 indicate that inhibition ranged from about 48% to about 70%, averaging about 60%, with no significant differences in effectiveness seen among the oligomers tested. (The nucleotide symbol "M" in the sequences represents methyl cytidine.)

15 Figure 6 illustrates the effect of a broad spectrum PMO on VRE colony formation. The oligomer designated SEQ ID NO: 114 is considered broad spectrum, targeted to a region conserved in all of the bacteria listed in Table 5A, above. This oligomer targets approximately the same region as that targeted by SEQ ID NO: 23, which is shown in Table 5A. As can be seen from the data in Table 10, this oligomer was similar in effectiveness to a "narrow spectrum" oligomer specific to *Enterococcus*, SEQ ID NO: 115.

20 Also included were several oligomers specific to 16s rRNA of other organisms (*E. coli*, *S. aureus*, and *P. aeruginosa*). These oligomers had no inhibitory effect on VRE.

Table 10. Targeting Study in *Enterococcus faecium*.

PMO Source	GenBank Acc. No.	Location	Antisense Sequence (5'→3')	SEQ ID	Percent Inhibition	S.E.	n =
VRE	Y18294	447-466	GAT GAA CAG TTA CTC TCA TC	91	61.7	2.7	3
VRE	Y18294	1272-1291	ACT GAG AGA AGC TTT AAG AG	92	59.7	5.1	6
VRE	Y18294	1-20	GGC ACG CCG CCA GCG TTC G	102	56.7	7.8	3
VRE	Y18294	300-319	TGT CTC AGT CCC AAT GTG GC	103	53.7	1.0	3
VRE	Y18294	721-740	GTT ACA GAC CAG AGA GCC GC	104	69.7	3.0	3
VRE	Y18294	1022-1041	CAC CTG TCA CTT TGC CCC CG	105	47.9	10.1	3
VRE	Y18294	1438-1456	GGC GGC TGG CTC CAA AAG G	106	58.5	3.2	3
VRE	Y18294	776-795	GAC TAC CAG GGT ATC TAA TC	114	62.2	5.5	3
VRE	Y18294	194-213	CAG CGA CAC CCG AAA GCC CC	115	70.1	3.3	3
<i>S. aureus</i>	See Table 8		CTG AGA ACA ACT TTA TGG GA	94	24	8.8	3
<i>P. aeruginosa</i>	See Table 9		TCG AGT TGC AGA CTG CGA TC	96	26	11.6	3
<i>E. coli</i>	See Table 7		GCA AAG GTA TTA ACT TTA CTC	17	17	22.4	3
<i>E. coli</i>	See Table 7		GCA CTT TAT GAG GTC CGC TTG	19	9	10	3
VRE	Y18294	0077-95	CAC CCG TTC GCC ACT CCT C	107	45.1	6.1	3
VRE	Y18294	0895-914	TCA ATT CCT TTG AGT TTC AA	108	31.8	15.3	3
VRE	Y18294	1263-1291	GCA ATC CGC ACT GAG AGA AGC TTT AAG AG	109	39.1	11.4	6

VRE	Y18294	1268-1291	C CGC ACT GAG AGA AGC TTT AAG AG	110	50.1	5.5	6
VRE	Y18294	1275-1291	GAG AGA AGC TTT AAG AG	111	61.5	3.3	6
VRE	Y18294	1277-1291	G AGA AGC TTT AAG AG	112	46.3	5	6
VRE	Y18294	1282-1291	A AGC TTT AAG AG	113	39.5	8.2	6
VRE	Y18294	1274-1291	T GAG AGA AGC TTT AAG AG	121	57.2	4.8	3
VRE	Y18294	1273-1291	CT GAG AGA AGC TTT AAG AG	122	54.4	2.7	3
VRE	Y18294	196-213	GCG ACA CCC GAA AGC GCC	123	59.0	5.3	6
VRE	Y18294	723-740	TAC AGA CCA GAG AGC CGC	124	63.3	4.9	9
VRE	Y18294	197-213	CGA CAC CCG AAA GCG CC	125	63.6	3.7	9
VRE	Y18294	195-213	A GCG ACA CCC GAA AGC GCC	126	60.6	4.8	12
VRE	Y18294	196-213	CG ACA CCC GAA AGC GCC A	127	58.9	5.6	9
VRE	Y18294	197-213	MG AMA MMA GAA AGM GMM	128	60.3	4.5	9
VRE	Y18294	723-740	TAM AGA MMA GAG AGM MGM	129	56.9	3.9	9
VRE	Y18294	1162-1177	MMM MAM MTT MTT MMG G	130	56.1	3.7	9
VRE	Y18294	1345-1363	CAC CGC GGC GTG CTG ATC C	131	64.0	3.9	6
VRE	Y18294	1162-1177	CCC CAC CTT CCT CCG G	132	70.2	1.6	3
VRE	Y18294	916-933	CCG CTT GTG CGG GCC CCC	133	66.8	4.3	3
VRE	Y18294	1345-1362	CAC CGC GGC GTG CTG ATC	134	71.4	11.3	3
VRE	Y18294	1345-1361	CAC CGC GGC GTG CTG AT	135	57.3	3.8	3
VRE	Y18294	1346-1364	ACC GCG GCG TGC TGA TCC	136	75.0	4.4	3
VRE	Y18294	1344-1360	CCG CGG CGT GCT GAT CC	137	66.3	3.5	3
VRE	Y18294	1346-1363	ACC GCG GCG TGC TGA TC	138	63.8	2.2	3

M represents methyl cytidine.

A dose-response study was also conducted using different concentrations of the oligomer having SEQ ID NO: 92. About 70% inhibition was achieved at 1-10 μ M, about 50% at 0.1 μ M, about 20% at 0.01 μ M, and about 12% at 1 nM.

5 (b) Bacterial 23S rRNA Targets

In a related experiment, also using vancomycin-resistant *Enterococcus faecium* (VRE) as the target bacteria, the effect of PMO antisense oligomers targeting 23S rRNA sequences on bacterial growth was evaluated, using the method described above. In individual assays, antisense PMO's targeting VRE 23S rRNA sequences were added to cultures of VRE and incubated at a 10 concentration of 1 μ M for 16 hours. The data in Table 11, below, represented graphically in Fig. 7, shows that antisense targeting of 23S rRNA in VRE was successful in inhibiting bacterial growth. Locations refer to GenBank Acc. No. X79341.

Table 11. VRE 23S rRNA Targeting Study

Ref. No.	Location	Antisense Sequence (5'→3')	SEQ ID NO:	Percent Inhibition	S.E. (N=3)
46	20-39	GTG CCA AGG CAT CCA CCG TG	116	61.9	4.6
47	679-698	CAT ACT CAA ACG CCC TAT TC	117	46.8	6.6
48	1462-1480	CCT TAG CCT CCT GCG TCC C	118	47.6	7.5
49	2060-2079	GGG GTC TTT CCG TCC TGT CG	119	67.0	5.7
50	2881-2900	CGA TCG ATT AGT ATC AGT CC	120	63.0	10.5

B. Effect of Length of Antisense Oligomer on Inhibition of VRE

The procedure used to obtain the data shown in Table 10, above, was repeated using different-length versions (SEQ ID NOs: 109-113) of the anti-16S rRNA oligomer having SEQ ID NO: 92, ranging from a 12-mer (SEQ ID NO: 113) to a 29-mer (SEQ ID NO:109). Results are 5 given in Table 12, below.

As shown in Table 12 and Fig. 8, the optimum length in this study was in the 17- to 20-mer range. Further studies confirmed that oligomers with a length of from 17 to 20 nucleotide subunits, and more preferably 17-18 subunits, are generally preferred. The results suggest that shorter oligomers, such as 12-mers, may have insufficient binding affinity, and that longer 10 oligomers, such as the 29-mer, are less easily transported into cells.

Table 12. Antisense Targeting of 16S rRNA in VRE

Ref. No.	length	Antisense sequence (5'→3')	SEQ ID NO:	Percent Inhibition	SE	n =
39	29mer	GCA ATC CGC ACT GAG AGA AGC TTT AAG AG	109	29.1	11.4	6
40	24mer	C CGC ACT GAG AGA AGC TTT AAG AG	110	51.1	5.5	6
22	20mer	ACT GAG AGA AGC TTT AAG AG	92	59.7	5.2	6
41	17mer	GAG AGA AGC TTT AAG AG	111	61.5	3.3	6
42	15mer	G AGA AGC TTT AAG AG	112	46.3	5.0	6
43	12mer	A AGC TTT AAG AG	113	39.5	8.2	6

C. Antisense PMO Resistance Study in VRE

15 The 20-mer anti-16S rRNA antisense oligomer referred to above (SEQ ID NO: 92) was used in a resistance study with VRE. After each day of incubation (concn. 1 μ M), three colonies were picked and retreated with oligomer to test for resistance. As shown in Table 13, below, and in Fig. 9, viability increased somewhat at four days but then dropped again at five and six days. Tests carried out to twelve days (data not shown) showed no evidence that resistance to the 20 oligomer had developed.

Table 13. Resistance Study with anti-16S rRNA (SEQ ID NO: 92) in VRE

Day	Percent Inhibition	S.E. (n=3)
1	41.8	5.2
2	49.6	2.7
3	51.8	12.3
4	19.2	11.9
5	34.1	10.9
6	47.2	12.0

D. Combination Therapy with Antibiotic Drugs

Enterococcus faecium was treated with vancomycin alone and in combination with 1.0 μ M antisense PMO targeted to VRE 16S rRNA (SEQ ID NO: 92). Inhibition was greatly increased by addition of the PMO, as shown in Figure 10A, and the organisms were completely eliminated at 3 μ M vancomycin and 1 μ M PMO. The results show that use of an antisense PMO targeted to VRE 16S rRNA together with vancomycin results in an enhanced anti-bacterial effect relative that of vancomycin alone.

A similar study was conducted with vancomycin resistant *Enterococcus faecium* (VRE), treated with ampicillin alone and in combination with 1.0 μ M of the same antisense PMO (see Fig. 10B). Again, essentially complete inhibition was achieved by the combination at 3 μ M ampicillin. Similar to the results obtained for vancomycin, the combination of an antisense PMO targeted to VRE 16S rRNA and ampicillin resulted in an enhanced anti-bacterial effect.

VII. In Vivo Administration Of Antisense Oligomers

In another aspect, the invention is directed to slowing or limiting bacterial infection *in vivo* in a mammal, and/or decreasing or eliminating detectable symptoms typically associated with infection by that particular bacteria. In general, a therapeutically effective amount of an antisense oligonucleotide-containing pharmaceutical composition is administered to a mammalian subject, in a manner effective to inhibit the activity of a 16S rRNA.

The antisense oligonucleotides of the invention and pharmaceutical compositions containing them are useful for inhibiting bacterial infection *in vivo* in a mammal, and for inhibiting or arresting the growth of bacteria in the host. In other words, the bacteria may be decreased in number or eliminated, with little or no detrimental effect on the normal growth or development of the host.

In some cases, the antisense oligomer will inhibit the growth of bacteria in general. In other cases, the antisense oligomer will be specific to one or more particular types of bacteria, *e.g.* a particular genus, species or strain.

It will be understood that the *in vivo* efficacy of such an antisense oligomer in a subject using the methods of the invention is dependent upon numerous factors including, but not limited to, (1) the target sequence; (2) the duration, dose and frequency of antisense administration; and (3) the general condition of the subject.

The efficacy of an *in vivo* administered antisense oligomer of the invention on inhibition or elimination of the growth of one or more types of bacteria may be determined by *in vitro* culture or microscopic examination of a biological sample (tissue, blood, etc.) taken from a subject prior

to, during and subsequent to administration of the antisense oligomer. (See, for example, Pari, G.S. *et al.*, *Antimicrob. Agents and Chemotherapy* 39(5):1157-1161, 1995; Anderson, KP *et al.*, *Antimicrob. Agents and Chemotherapy* 40(9):2004-2011, 1996.)

5 A. Treating Subjects

Effective delivery of the antisense oligomer to the target RNA is an important aspect of the methods of the invention. In accordance with the invention, such routes of antisense oligomer delivery include, but are not limited to, various systemic routes, including oral and parenteral routes, *e.g.*, intravenous, subcutaneous, intraperitoneal, and intramuscular, as well as inhalation, transdermal and topical delivery. The appropriate route may be determined by one of skill in the art, as appropriate to the condition of the subject under treatment.

For example, an appropriate route for delivery of an antisense oligomer in the treatment of a bacterial infection of the skin is topical delivery, while delivery of an antisense oligomer in the treatment of a bacterial respiratory infection is by inhalation.

15 Additional exemplary embodiments include oral delivery of an antisense oligomer directed to bacterial 16S or 23S rRNA for treatment of a urinary tract infection or sepsis and IV delivery for treatment of sepsis.

It is appreciated that methods effective to deliver the oligomer to the site of bacterial infection or to introduce the oligonucleotide into the bloodstream are contemplated.

20 Transdermal delivery of antisense oligomers may be accomplished by use of a pharmaceutically acceptable carrier adapted for topical administration. One example of morpholino oligomer delivery is described in PCT patent application WO 97/40854, incorporated herein by reference.

25 In one aspect of the invention, an antisense oligomer directed to bacterial 16S or 23S rRNA is delivered by way of a catheter, microbubbles, a heart valve coated or impregnated with oligomer, a Hickman catheter or a coated stent.

30 In one preferred embodiment, the oligomer is a morpholino oligomer, contained in a pharmaceutically acceptable carrier, and delivered orally. In a further aspect of this embodiment, a morpholino antisense oligonucleotide is administered at regular intervals for a short time period, *e.g.*, daily for two weeks or less. However, in some cases the antisense oligomer is administered intermittently over a longer period of time.

Typically, one or more doses of antisense oligomer are administered, generally at regular intervals, for a period of about one to two weeks. Preferred doses for oral administration are from about 1 mg oligomer/patient to about 25 mg oligomer/patient (based on a weight of 70 kg). In some cases, doses of greater than 25 mg oligomer/patient may be necessary. For IV

administration, the preferred doses are from about 0.5 mg oligomer/patient to about 10 mg oligomer/patient (based on an adult weight of 70 kg).

The antisense compound is generally administered in an amount and manner effective to result in a peak blood concentration of at least 200-400 nM antisense oligomer.

5 In general, the method comprises administering to a subject, in a suitable pharmaceutical carrier, an amount of an antisense agent effective to inhibit the biological activity of a bacterial 16S or 23S rRNA target sequence of interest.

10 It follows that a morpholino antisense oligonucleotide composition may be administered in any convenient vehicle which is physiologically acceptable. Such an oligonucleotide composition may include any of a variety of standard pharmaceutically accepted carriers employed by those of ordinary skill in the art. Examples of such pharmaceutical carriers include, but are not limited to, saline, phosphate buffered saline (PBS), water, aqueous ethanol, emulsions such as oil/water emulsions, triglyceride emulsions, wetting agents, tablets and capsules. It will be understood that the choice of suitable physiologically acceptable carrier will vary dependent upon the chosen 15 mode of administration.

15 In some instances liposomes may be employed to facilitate uptake of the antisense oligonucleotide into cells. (See, e.g., Williams, S.A., *Leukemia* 10(12):1980-1989, 1996; Lappalainen *et al.*, *Antiviral Res.* 23:119, 1994; Uhlmann *et al.*, ANTISENSE OLIGONUCLEOTIDES: A NEW THERAPEUTIC PRINCIPLES, *Chemical Reviews*, Volume 90, No. 4, pages 544-584, 1990; 20 Gregoriadis, G., Chapter 14, Liposomes, *Drug Carriers in Biology and Medicine*, pp. 287-341, Academic Press, 1979). Hydrogels may also be used as vehicles for antisense oligomer administration, for example, as described in WO 93/01286. Alternatively, the oligonucleotides may be administered in microspheres or microparticles. (See, e.g., Wu GY and Wu CH, *J. Biol. Chem.* 262:4429-4432, 1987.)

25 Sustained release compositions are also contemplated within the scope of this application. These may include semipermeable polymeric matrices in the form of shaped articles such as films or microcapsules.

30 In one aspect of the method, the subject is a human subject, typically a subject diagnosed as having a localized or systemic bacterial infection.

35 In another aspect, the condition of the patient may dictate prophylactic administration of an antisense oligomer of the invention, *i.e.*, a patient who (1) is immunocompromised; (2) is a burn victim; (3) has an indwelling catheter; (4) is about to undergo or has recently undergone surgery, etc.

35 In another application of the method, the subject is a livestock animal, *e.g.*, a chicken, turkey, pig, cow or goat, etc, and the treatment is either prophylactic or therapeutic.

In addition, the methods of the invention are applicable to treatment of any condition wherein inhibiting or eliminating the growth of bacteria would be effective to result in an improved therapeutic outcome for the subject under treatment.

It will be understood that an effective *in vivo* treatment regimen using the antisense oligonucleotides of the invention will vary according to the frequency and route of administration, as well as the condition of the subject under treatment (*i.e.*, prophylactic administration versus administration in response to localized or systemic infection). Accordingly, such *in vivo* therapy will generally require monitoring by tests appropriate to the particular type of bacterial infection under treatment and a corresponding adjustment in the dose or treatment regimen in order to achieve an optimal therapeutic outcome.

B. Monitoring Treatment

The efficacy of a given therapeutic regimen involving the methods described herein may be monitored, *e.g.*, by general indicators of infection, such as complete blood count (CBC), nucleic acid detection methods, immunodiagnostic tests or bacterial culture.

Identification and monitoring of bacterial infection generally involves one or more of (1) nucleic acid detection methods; (2) serological detection methods, *i.e.*, conventional immunoassay; (3) culture methods; and (4) biochemical methods. Such methods may be qualitative or quantitative.

DNA probes may be designed based on publicly available bacterial nucleic acid sequences, and used to detect target genes or metabolites (*i.e.*, toxins) indicative of bacterial infection, which may be specific to a particular bacterial type, *e.g.*, a particular species or strain, or common to more than one species or type of bacteria (*i.e.*, Gram positive or Gram negative bacteria). In addition, nucleic amplification tests (*e.g.*, PCR) may be used in such detection methods.

Serological identification may be accomplished using a bacterial sample or culture isolated from a biological specimen, *e.g.*, stool, urine, cerebrospinal fluid, blood, etc. Immunoassay for the detection of bacteria is generally carried out by methods routinely employed by those of skill in the art, *e.g.*, ELISA or Western blot.

In general, procedures and/or reagents for immunoassay of bacterial infections are routinely employed by those of skill in the art. In addition, monoclonal antibodies specific to particular bacterial strains or species are often commercially available.

Culture methods may be used to isolate and identify particular types of bacteria, by employing techniques including, but not limited to, aerobic versus anaerobic culture, and growth and morphology under various culture conditions.

Exemplary biochemical tests include Gram stain (Gram, 1884; Gram positive bacteria stain dark blue, and Gram negative stain red), enzymatic analyses (*i.e.*, oxidase, catalase positive for *Pseudomonas aeruginosa*), and phage typing.

It will be understood that the exact nature of such diagnostic, and quantitative tests as well as other physiological factors indicative of bacterial infection will vary dependent upon the bacterial target, the condition being treated and whether the treatment is prophylactic or therapeutic.

In cases where the subject has been diagnosed as having a particular type of bacterial infection, the status of the bacterial infection is also monitored using diagnostic techniques typically used by those of skill in the art to monitor the particular type of bacterial infection under treatment.

The antisense oligomer treatment regimen may be adjusted (dose, frequency, route, etc.), as indicated, based on the results of immunoassays, other biochemical tests and physiological examination of the subject under treatment.

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications may be made without departing from the invention.

MATERIALS AND METHODS

Standard recombinant DNA techniques were employed in all constructions, as described in Ausubel, FM, *et al.*, in CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley and Sons, Inc., Media, PA, 1992 and Sambrook J, *et al.*, in MOLECULAR CLONING: A LABORATORY MANUAL, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, Vol. 2, 1989), both of which are expressly incorporated by reference herein.

Plasmid. The plasmid used for studies in support of the present invention was engineered using pCi-Neo mammalian expression vector (Promega), by inserting 36 bases of the c-myc target region along with the coding region for firefly luciferase into the vector in the polylinker downstream from the T7 promoter. The A from the ATG of codon No. 1 of luciferase was removed by *in vitro* mutagenesis, leaving the ATG that is present in the c-myc sequence in frame with the reporter. The plasmid, pCiNeo(myc)luc δ A, also contained the b-lactamase gene coding for antibiotic resistance and was transformed into *Escherichia Coli* DHS.

Bacterial Cultures. In evaluating the effectiveness of antisense oligonucleotides of the invention, approximately 3 ml bacterial cultures were aliquoted into plastic snap cap tubes from a 45 ml starting culture in Luria-Bertani (LB) Broth containing 4.5 mg of Ampicillin and a single bacterial colony taken from a freshly streaked LB agar plate containing 100 μ g/mL ampicillin. The test oligomer diluted in phosphate buffered saline (PBS) was added to the cultures, incubated

at 37°C for a specific time, e.g., 16 or 26 hours with shaking at 210 rpm, then placed on ice for 15 minutes.

Culture staining microscopy and colony scanning. Bacterial plate counts require that a measured volume of material be added to agar either by the pour plate or spread plate technique. If 5 the original sample has a large number of bacteria, dilutions are prepared and plated. The plates are incubated and the number of colony-forming units (CFU) reflect the viable organisms in the sample.

The colonies may be counted manually using a microscope, however, it is preferred that an automatic colony counter be employed (e.g., as offered by Bioscience International, Rockville, MD). Bacterial cultures are stained in accordance with standard Gram staining protocols. The 10 stained bacterium are visualized using a Nikon Optiphot-2 upright microscope, with images magnified 1000X using the combination of an 100X oil immersion lens and the 10X magnification of the camera. The camera used to capture the images is a Nikon N8008S. The images are taken using bright field microscopy with a 4 second exposure on a setting 5 light output. A preferred film was Kodak Gold 400 ASA. After developing, the images are scanned using a Microtek Scan Maker 15 4, then cropped using Adobe PhotoShop.

Sequence Listing Table

Description	SEQ ID NO.
<i>E. coli</i> GenBank Accession No: X80725	1
<i>Salmonella typhimurium</i> GenBank Accession No: U88545	2
<i>Pseudomonas aeruginosa</i> GenBank Accession No: AF170358	3
<i>Vibrio cholera</i> GenBank Accession No: AF118021	4
<i>Staphylococcus aureus</i> GenBank Accession No: Y15856	6
<i>Mycobacterium tuberculosis</i> GenBank Accession No: X52917	7
<i>Helicobacter pylori</i> GenBank Accession No: M88157	8
<i>Streptococcus pneumoniae</i> GenBank Accession No: AF003930	9
<i>Treponema palladium</i> GenBank Accession No: AJ010951	10
<i>Chlamydia trachomatis</i> GenBank Accession No: D85722	11
<i>Bartonella henselae</i> GenBank Accession No: X89208	12
<i>Hemophilis influenza</i> GenBank Accession No: M35019	13
<i>Shigella dysenteriae</i> GenBank Accession No: X96966	14
0-1-23-15 (BS-1; Table 2A)	15
5'- GGA CTA CGA CGC ACT TTA TGA G -3' (22-mer)	
0-1-23-16 (BS-2; Table 2A)	16
5'- GGT CCG CTT GCT CTC GCG AGG -3' (21-mer)	
0-1-23-17 (NS-1; Table 2A)	17
5'-GCA AAG GTA TTA ACT TTA CTC-3' (21-mer)	
0-1-23-18 (NS-2; Table 2A)	18
5'- GCT GCG GTT ATT AAC CAC AAC -3' (21-mer)	
0-1-23-9 (<i>E. coli</i> 16S)	19
5'-GCA CTT TAT GAG GTC CGC TTG-3' (21-mer)	
TGCTGCCTCCCGTAGGAGTCT	Table 2A-broad
ATTACCGCGGCTGCTGGCACG	Table 2A-broad
ACCAGGGTATCTAATCCTGTT	Table 2A-broad
CACATGCTCCACCGCTTGTGC	Table 2B-broad
TTGCGGGACTTAACCCAACAT	Table 2B-broad
CGCGGCTGCTGGCACGTAGTT	Table 3-Gram positive
ACTTAACCCAACATCTCACGA	Table 3-Gram positive
TTTACGCCAGTAATTCCGA	Table 4-Gram negative
ACTCCCATGGTGTGACGGGCGG	Table 4-Gram negative
AATCTGAGCCATGATCAAAC	Table 2A
CCCTCTTGTGCTTGCACGT	Table 2A
ACCCCCCTCTACGAGACTCAA	Table 2A
CCACGCCTCAAGGGCACACC	Table 2A
TCTCATCTCTGAAAACCTCCG	Table 2A
CATGATCAAACCTTCAATT	Table 2A
CCCTCTTGGTCTTGCACGT	Table 2A
TACCCCCCTCTACGAGACTCA	Table 2A
GCCACGCCTCAAGGGCACACC	Table 2A
CAGAGAGCAAGCCCTTTCAT	Table 2A
CCTGCTTCTCCCGTAGGACG	Table 2A
CACCAACCCTCTGCCATACTCT	Table 2A

CTAAGATCTCAAGGATCCAA	Table 2A	43
GGCCTGCCGCCAGCGTCAAT	Table 2A	44
CCCTCTTGGTCCGTAAACAT	Table 2A	45
CCCCCTCTACAAGACTCTAGC	Table 2A	46
ACGACTRTAGGTACAACCTC	Table 2A	47
AGGATCAAACCTTATGTTCA	Table 2B	48
CCTGCTTCCCTCTCAAGACG	Table 2B	49
CACCTCCCTGTGACACACTG	Table 2B	50
CCAAGCAATCAAGTTGCCAA	Table 2B	51
CCAGCGTTCATCCTGAGCCAG	Table 2B	52
GAACCATGCGGGTCAAAATAT	Table 2B	53
CTTCCTCTTCTGACTCAAG	Table 2B	54
GGGGCGGAAACCCCCCTAACAC	Table 2B	55
GCATGTGTTAACGACGCCGCC	Table 2B	56
AAGACATGCATCCCCTGGTCC	Table 2B	57
CAGTCTCCCTGCAGTACTCT	Table 2B	58
GATCCAAGGAAGGAAACCCA	Table 2B	59
CAGGATCAAACCTCTCCATAAA	Table 2B	60
AAATCTTCCCCCTAGGAGT	Table 2B	61
CACCTACCTCTCCCACACTCT	Table 2B	62
TGGAGAGACTAAGCCCTCCAA	Table 2B	63
CGTCCGTAGCCAGGATCAAAT	Table 2B	64
ATGTCATGCAACATCCACTCT	Table 2B	65
ACTCTCCCTTTGACTCAA	Table 2B	66
AAACCCCGGAAAGGGTCTAAC	Table 2B	67
TCTGAGCCATGATCAAACCT	Table 2B	68
ACCCCTTATGAAGCACTGAGT	Table 2B	69
AGTTTCCCTCCGTAACTCAA	Table 2B	70
CACTGAAGCTTGGCCCAAC	Table 2B	71
GAACCAAGATCAAATTCTCAG	Table 2C	72
GTTACTCGGATGCCAAATAT	Table 2C	73
CCTTTCTCCATCTACCCCT	Table 2C	74
GGATGGGGTTGAGACCATCCA	Table 2C	75
AGCGTTCATCCTGAGCCAGGA	Table 2C	76
AAATCTTCTCCAGTAGGACG	Table 2C	77
CACTCACCTCTCCATACTCA	Table 2C	78
ACTGAGCAGTAAACCCACCCAA	Table 2C	79
CATGATCAAACCTTCAATTN	Table 2C	80
CACTTCATCTCCGATAATA	Table 2C	81
CCTCCCTAAAGTACTCTAGTT	Table 2C	82
CAGAGTTAACCCCCAACCCCC	Table 2C	83
GCCAGCGTTCAATCTGAGCCA	Table 2C	84
CCCTCTTGGTCTTGCACGT	Table 2C	85
TACCCCCCTACGAGACTCA	Table 2C	86
GCCACGCCCTCAAGGGCACAAC	Table 2C	87
CCTCGTATCTCTACAAGGTT	Table 2C	88
CCC CAT CAT TAT GAG TGA TGT GC	AVI-1-23-19	89
TCA TTA TGA G GTG ACC CCA	AVI-1-23-20	90
GAT GAA CAG TTA CTC TCA TC	AVI-1-23-21	91

ACT GAG AGA AGC TTT AAG AG	AVI-1-23-22	92
ATG TGC ACA GTT ACT TAC AC	AVI-1-23-23	93
CTG AGA ACA ACT TTA TGG GA	AVI-1-23-24	94
TTA TTC TGT TGG TAA CGT CA	AVI-1-23-25	95
CG AGT TGC AGA CTG CGA TC	AVI-1-23-26	96
ATC TGA GCC ATG ATC AAA CT	AVI-1-23-27	97
TGT CTC AGT TCC AGT GTT GC	AVI-1-23-28	98
GTC TTC GTC CAG GGG GCC GC	AVI-1-23-29	99
CAC CTG TCT CAC GGT TCC CG	AVI-1-23-30	100
CGC CCT CCC GAA GTT AAG CT	AVI-1-23-31	101
GGC ACG CCG CCA GCG TTC G	AVI-1-23-32	Table 10
TGT CTC AGT CCC AAT GTG GC	AVI-1-23-33	Table 10
GTT ACA GAC CAG AGA GCC GC	AVI-1-23-34	Table 10
CAC CTG TCA CTT TGC CCC CG	AVI-1-23-35	Table 10
GGC GGC TGG CTC CAA AAG G	AVI-1-23-36	Table 10
CAC CCG TTC GCC ACT CCT C	AVI-1-23-37	Table 10
TCA ATT CCT TTG AGT TTC AA	AVI-1-23-38	Table 10
GCA ATC CGC ACT GAG AGA AGC TTT AAG AG	AVI-1-23-39	Table 10
C CGC ACT GAG AGA AGC TTT AAG AG	AVI-1-23-40	Table 10
GAG AGA AGC TTT AAG AG	AVI-1-23-41	Table 10
G AGA AGC TTT AAG AG	AVI-1-23-42	Table 10
A AGC TTT AAG AG	AVI-1-23-43	Table 10
GAC TAC CAG GGT ATC TAA TC	AVI-1-23-44	Table 10
CAG CGA CAC CCG AAA GCC CC	AVI-1-23-45	Table 10
GTG CCA AGG CAT CCA CCG TG	AVI-1-23-46	Table 11
CAT ACT CAA ACG CCC TAT TC	AVI-1-23-47	Table 11
CCT TAG CCT CCT GCG TCC C	AVI-1-23-48	Table 11
GGG GTC TTT CCG TCC TGT CG	AVI-1-23-49	Table 11
CGA TCG ATT AGT ATC AGT CC	AVI-1-23-50	Table 11
T GAG AGA AGC TTT AAG AG	AVI-1-23-63	Table 10
CT GAG AGA AGC TTT AAG AG	AVI-1-23-66	Table 10
GCG ACA CCC GAA AGC GCC	AVI-1-23-67	Table 10
TAC AGA CCA GAG AGC CGC	AVI-1-23-68	Table 10
CGA CAC CCG AAA GCG CC	AVI-1-23-69	Table 10
A GCG ACA CCC GAA AGC GCC	AVI-1-23-70	Table 10
CG ACA CCC GAA AGC GCC A	AVI-1-23-71	Table 10
MG AMA MMM GAA AGM GMM	AVI-1-23-72	Table 10
TAM AGA MMA GAG AGM MGM	AVI-1-23-73	Table 10
MMM MAM MTT MTT MMG G	AVI-1-23-74	Table 10
CAC CGC GGC GTG CTG ATC C	AVI-1-23-75	Table 10
CCC CAC CTT CCT CCG G	AVI-1-23-76	Table 10
CCG CTT GTG CGG GCC CCC	AVI-1-23-77	Table 10
CAC CGC GGC GTG CTG ATC	AVI-1-23-78	Table 10
CAC CGC GGC GTG CTG AT	AVI-1-23-79	Table 10
ACC GCG GCG TGC TGA TCC	AVI-1-23-80	Table 10
CCG CGG CGT GCT GAT CC	AVI-1-23-81	Table 10
ACC GCG GCG TGC TGA TC	AVI-1-23-82	Table 10
5'- ACG TTG AGG GGC ATC GTC GC-3'	AVI 1-22-126	AS to c-myc
		139

EDITORIAL NOTE

APPLICATION NUMBER – AU 45098/01

The following sequence listing pages 51a – 78 are part of the description. The claims pages follow on pages 47 – 51.

SEQUENCE LISTING

<110> AVI BioPharma, Inc.

<120> Antisense Antibacterial Method and Composition

<130> 0450-0032.41

<140> PCT/US00/42391
<141> 2000-11-29<150> US 60/168,150
<151> 1999-11-29

<160> 139

<170> FastSEQ for Windows Version 4.0

<210> 1
<211> 1450

<212> DNA

<213> Escherichia coli

<220>

<221> misc_feature

<222> 392, 852, 853

<223> n = A,T,C or G

<400> 1

agtttgatca	tggctcagat	tgaacgctgg	cgccaggcct	aacacatgca	agtgcAACGG	60
taacaggaag	cagcttgctg	cttgcgtgac	gagttggcgg	cgggtggat	atgtctggga	120
aactgcctga	tggaggggg	taactactgg	aaacggtagc	taataccgca	taacgtcgca	180
agcacaaaga	gggggaccc	tttggccttt	ggccatggat	gtgcccagat	gggatttagct	240
atagggtggg	gttacggctc	acctaggcga	cgatccctag	ctgggtctgag	aggatgacca	300
gcaacactgg	aactgagaca	cggtccagac	tcctacggg	ggcagcagt	ggaaatattg	360
cacaatgggc	gcaaggctga	tgcagccat	cnccgtgtat	gaagaaggcc	ttccgggttgt	420
aaagtacttt	cagcggggag	gaaggggagta	aagttataac	ctttgctcat	tgacgttacc	480
cgagaagaa	gcaccggcta	actccgtgcc	agcagcccg	gtatatacgga	gggtgcaagc	540
gttaatcgga	attactgggc	gtaaagcgca	cgcaggcggt	ttgttaagtc	agatgtgaaa	600
tccccgggct	caacctggga	actgcacatcg	atactggca	gtttgagct	cgtagagggg	660
ggtagaattc	cagggttagc	ggtgaaatgc	gttagagatct	ggaggaatac	cggtggcgaa	720
ggcggccccc	tggacgaaga	ctgacgctca	ggtgcgaaag	cgtggggagc	aaacaggatt	780
agataccctg	gttagtccacg	ccgttaaacga	tgtgcacttg	gagggtgtgc	ccttgaggcg	840
tggctcccg	anntaacgcg	ttaagtgcac	cgccctggga	gtacggccgc	aagttaaaaa	900
ctcaaatgaa	ttgacgggg	ccgcacaaag	ggtggagcat	gtgggttaat	tgcatgcac	960
gcaagaacc	ttacctggtc	ttgacatcca	cggaagttt	cagagatgag	aatgtgcctt	1020
cgggaaccgt	gagacaggtg	ctgcatggct	gtcgtcagct	cgtgttgta	aatgtgggt	1080
taagtcccg	aacgacgcga	acccttattcc	tttggttgcca	gcccgtccgc	cggaactca	1140
aaggagactg	ccagtataaa	actggaggaa	ggtggggatg	acgtcaagtc	atcatggccc	1200
ttacgaccag	ggctacacac	gtgctacaat	ggcgcataca	aagagaagcg	acctcgcgag	1260
agcaagcgga	cctcataaaag	tgcgtcgtag	tccggattgg	agtctgcac	tgcactccat	1320
gaagtccgaa	tcgtcgat	tcgtggatca	gaatgccacg	gtgaaatacg	tcccgccct	1380
tgtacacacc	gccccgtcaca	ccatgggagt	gggttgcaaa	agaagtaggt	agcttaactt	1440
cgggagggcg						1450

<210> 2

<211> 1541

<212> DNA

<213> Salmonella thyphimurium

<400> 2

aaattgaaga	gtttgatcat	ggctcagatt	gaacgctggc	ggcaggccta	acacatgcaa	60
gtcgaacgg	aacaggaaac	agcttgctct	ttgctgacga	gtggcggacg	ggtagataat	120
gtctggaaa	ctgcctgtat	gagggggata	actactggaa	acgggtggcta	ataccgcata	180
acgtcgcaag	accaaagagg	gggaccttcg	ggccctttgc	catcgatgt	gcccagatgg	240
gattagctag	taggtgggt	aacggctcac	ctaggcgcac	atccctagct	ggtctgagag	300
gatgaccagc	cacactgaag	ctgaagcacg	gtccagactc	ctacggagg	cagcagtggg	360

0450-0032.41-SEQLIST-corr

gaatatgtca	caatgggcgc	aagcctgatg	cagccatgcc	gcgtgtatga	agaaggcctt	420
cgggtttaaa	agtactttca	gcggggagga	agggtttgtg	gttaataacc	gcagcaattg	480
acgttacccg	cagaagaagc	accggctaac	tccgtgccag	cagccgcgtt	aatacggagg	540
gtgcaagcgt	taatcggaat	tactgggcgt	aaagcgcacg	caggcggttt	gttaagtca	600
atgtgaaatc	cccggtctca	acctgggaac	tgcacatctgtat	actggcaagc	ttgagttctcg	660
tagagggggg	tagaatttcca	ggtgttagcgg	tgaaatgcgt	agagatctgg	aggaataccg	720
gtggcgaagg	cggccccctg	gacgaagact	gacgctcagg	tgcgaaagcg	tggggagcaa	780
acaggattag	atacccttgg	agtccacgccc	gtaaacgatc	tctacttgg	ggttgtgccc	840
ttgaggcgtg	gcttccggag	ctaacgcgtt	aagtagagtg	cttggggaggt	acggccgcaa	900
ggttaaaact	caaataaatt	gacggggggcc	cgcacaagcg	gtggagcatg	tggttttaatt	960
cgtatgcacg	cgaagaacct	tacctggct	tgcacatccac	agaacttcc	agagatgaga	1020
ttgtgccttc	gggaactgtg	agacagggtgc	tgcattggctg	tcgtcagtc	gtgttgtgaa	1080
atgttgggtt	aagtcccgca	acgagcgc	cccttataatc	ttgttgccag	cgggtccggcc	1140
gggaactcaa	aggagactgc	cagtataaa	ctggaggaag	gtggggatga	cgtcaagtca	1200
tcatggccct	tacgaccagg	gctacacacg	tgctacaatg	gcgcatacaa	agagaagcga	1260
cctcgcgaga	gcaagcggac	ctcataaaagt	gcgtcgtagt	ccggatttgg	gtctgcaact	1320
cgactccatg	aagtccgtat	cgctagtaat	cgtggatcag	aatgccacgg	tgaatacgtt	1380
cccgggcctt	gtacacaccg	cccgtcacac	catgggatgt	ggttgcaaaa	gaagtaggt	1440
gcttaacctt	cgggagggcg	cttaccactt	tgtgattcat	gactgggtg	aagtcgtaac	1500
aaggttaaccg	tagggaaacc	tgcgggttgg	tcacccctt	a		1541

<210> 3

<211> 1467

<212> DNA

<213> *Pseudomonas aeruginosa*

<400> 3

atgaagaggg	cttgctctct	gattcagcgg	cggacgggtg	agtaatgcct	aggaatctgc	60
ctgatagtgg	gggacaacgt	ttcgaagga	acgctaatac	cgcatacgtc	ctacgggaga	120
aagcagggga	ccttcgggcc	ttgcgtatc	agatgagccct	aggtcgatt	agctagttgg	180
tgaggttaacg	gcttaccaag	gcgacgatcc	gtaactggtc	tgagaggatg	atcagtacaca	240
ctggaaactga	gacacggtcc	agactctac	gggaggcagc	agtgggaaat	attggaaat	300
gggcgaaagc	ctgatccagc	catgcccgt	gtgtgaagaa	ggtctcgga	ttgtaaagca	360
ctttaagttg	ggaggaaggg	cattaaaccta	atacggttagt	gttttgcgt	taccgcacaga	420
ataaggcccg	gcttaacttcg	tgccagcagc	cgcgttaata	cgaagggtgc	aagcgttaat	480
cggaattact	gggcgtaaag	cgcgttagg	tggtttgtta	agttaatgt	gaaagcccg	540
ggctcaacct	gggaaactgca	tccaaaactg	gcaagctaga	gtatggcaga	gggtgggtgaa	600
atttcctgtg	tagcggtgaa	atgcgtagat	ataggaaggg	acaccagtgg	cgaaggcgcac	660
cacctgggct	aatactgaca	ctgaggtgcg	aaagcgtggg	gagcaaacag	gattagatac	720
cctggtagtc	cacccgtaa	acgatgtcga	ctagccgttg	ggatccttga	gatcttagtgc	780
gcgcagctaa	cgcattaaatg	cgaccgcctg	gggagttacgg	ccgctagtt	aaaactctaa	840
tgaattgacg	ggggcccgca	caagcgttgg	agcatgtggt	ttaattcga	gcaacgcgaa	900
gaaccttacc	aggccttgcac	atgcagagaa	cttccagaa	atggattgt	gccttcgggaa	960
actctgacac	agggtgtcga	tggctgtcgt	cagctcgtgt	cgtgagatgt	tgggttaagt	1020
cccgtaacga	gcgcacccct	tgtcccttagt	taccagcaca	ttaaggtggg	cactctaagg	1080
agactgcgg	tgacaaacccg	gaggaaagggt	gggatgcacgt	caagtcatca	ttggcccttac	1140
ggcctgggct	acacacgtgc	tacaatgtc	ggtacaaagg	gttgcacaagc	cgcgagggtgg	1200
agctaattcc	ataaaacccga	tcgtatgtcc	gatcgcagtc	tgcaactcga	ctgcgtgaag	1260
tcggaatcgc	tagtaatcgt	gaatcagaat	gtcacgggtga	atacgttccc	gggccttgcg	1320
cacaccgccc	gtcacacccat	gggagttgggt	tgttccagaa	gtagctagtc	taaccttcgg	1380
ggggacgggtt	accacggagg	tattcatgac	tggggtgaag	tcgtaacaag	gtagccgtag	1440
gggaacctqc	qgctqgatca	cctcctt				1467

<210> 4

<211> 1500

<212> DNA

<213> *vibrio cholera*

<400> 4

attgaacgct

cttttgaaga

gggataaacag ttggaaaacg

gctcaccaaag	gcgacgatcc	ctagctggtt	tgagaggatg	atcagccaca	cttggaaactga	300
gacacggtcc	agactcctac	gggaggcagc	agtggggaaat	attgcacaat	gggcgcgaagc	360
ctgtatgcagc	catgccgcgt	gtgtgaagaa	ggccttcggg	ttgttaagca	ctttcagcag	420
tgaggaaggt	tggtgtcgta	atagcgtatc	aatttgacgt	tagctgcaga	agaagcaccg	480
gctaactccg	tgccagcagc	cgcggtaata	cgagggtgc	gagcgttaat	cggaattact	540
gggcgtaaag	cgcattcgagg	cggtttgtta	agcaagatgt	gaaagccccg	ggctcaacct	600

0450-0032.41-SEQLIST-corr

gggaaccgca	ttttgaactg	gcaggctaga	gtcttgtaga	gggggtttaga	atttcagggt	660
tagcggtgaa	atgcgttagag	atctgaagga	ataccggtgg	cgaaggcgc	cccctggaca	720
aagactgacg	ctcagatgcg	aaagcgtgg	gagcaaacag	gattagatac	cctggtagtc	780
cacgctgtaa	acgatgtcta	cttggaggt	gtgacctrara	gtcgtggctt	tcggagctaa	840
cgcgttaagt	agaccgcctg	gggagtagcgg	tcgcaagatt	aaaactcaa	tgaattgacg	900
ggggcccgca	caagcggtgg	agcatgtgg	ttaattcgat	gcaacgcga	gaaccttacc	960
tactcttgac	atccagagaa	gccgaaagag	attttgggt	gccttcggg	actctgagac	1020
aggtgtcgca	tggctgtcgt	cagctcgtgt	tgtgaatgt	tgggttaagt	cccgcaacga	1080
gcgcacccct	tatccctgtt	tgcgcagcg	taatgtcggg	aactccagg	agactgccgg	1140
tgataaaaccg	gaggaagggt	gggacgacgt	caagtcata	tggcccttac	gatggggct	1200
acacacgtgc	tacaatggca	tatacagagg	gcagcgaggc	cgcgagggtt	agcgaatccc	1260
agaaagtatg	tcgtatgtcc	gatcggagtc	tgcaactcga	ctccgtgaag	tcggaatcgc	1320
tagtaatcgt	gaatcagaat	gtcacggtga	atacgttccc	gggccttga	cacaccgccc	1380
gtcacacccat	gggagttggc	tgcaccagaa	gtagatagct	taaccttccg	gagggcgttt	1440
accacggtgt	ggttcatgac	tgggtgaag	tcgtaacaag	gtagccctag	gggaacctgg	1500

<210> 5

<211> 1544

<212> DNA

<213> Neisseria gonorrhoea

<400> 5

•••••tgaacataag	agtttgatcc	tggctcagat	tgaacgctgg	cggcatgctt	tacacatgca	60
•••••agtcggacgg	cagcacaggg	aagcttgctt	ctcggtggc	gagtggcgaa	cgggtgagta	120
•••••acatatcgga	acgtacccgg	tagcggggga	taactgatcg	aaagatcagc	taataccgca	180
•••••tagtcttga	gagggaaagc	aggggacctt	cgggccttgc	gctatccgag	cggccgatat	240
•••••ctgattagct	ggttggcggg	gttaaaggccc	accaaggcga	cgatcgttag	cgggtctgag	300
•••••aggatgatcc	gccacactgg	gactgagaca	cggcccgac	tcctacggg	ggcagcagtg	360
•••••ggaaattttg	gacaatggc	gcaagcctga	tccagccatg	ccgcgtgtct	gaagaaggcc	420
•••••ttcgggttgt	aaaggacttt	tgtcagggaa	gaaaaggctg	ttgccaatat	cggccggccga	480
•••••tgacggtacc	tgaagaataa	gcaccggcta	actacgtgcc	agcagccg	gtaatacg+a	540
•••••gggtgcgagc	gttaatcgga	attactgggc	gtaaagcggg	cgcagacgt	tacttaagca	600
•••••ggatgtgaaa	tcccccggct	caaccgggga	actgcgttct	gaactgggt	actcgagtgt	660
•••••gtcagaggg	ggttggattc	cacgtgttagc	agtgaatgc	gtagagatgt	ggaggaatac	720
•••••cgatggcgaa	ggcagccccc	tggataaca	ctgacgttca	tgtccgaaag	cgtgggtagc	780
•••••aaacaggagg	agataccctg	gtatgtccac	cccttaaaca	tgtcaattag	ctgttggca	840
•••••acttgattgc	tttgttagcgt	agctaacgcg	tgaaattgac	cgcctgggg	gtacggtcgc	900
•••••aagattaaaa	ctcaaaggaa	ttgacgggg	ccgcacaa	cggtggatga	tgtgattaa	960
•••••ttcgatgcaa	cgcgaagaac	cttacctgg	tttgacatgt	gcggaaatct	ccggagacgg	1020
•••••aggagtgcct	tcgggagccg	taacacagg	gctgcattgc	tgtcgtcagc	tcgtgtcg	1080
•••••agatgttggg	ttaagtcccg	caacgagcgc	aacccttgc	attagttgc	atcattcggt	1140
•••••tgggcactct	aatgagactg	ccggtgacaa	gccggaggaa	ggtggggatg	acgtcaagtc	1200
•••••ctcatggccc	ttatgaccag	ggcttcacac	gtcatacaat	gtcgtgtaca	gagggtagcc	1260
•••••aagccgcgag	gcggagccaa	tctcacaaa	ccgatcgtag	tccggattgc	actctgcaac	1320
•••••tcgagtgcat	gaagtcggaa	tcgctagtaa	tcgcaggta	gcatactgcg	gtaaatacgt	1380
•••••tcccgggtct	tgtacacacc	gcccgtcaca	ccatgggagt	gggggatacc	agaagtaggt	1440
•••••agggtaaccg	caaggagtcc	gcttaccacg	gtatgttca	tgactgggt	gaagtcgtaa	1500
•••••caaggtagcc	gtaggggaac	ctgcggctgg	atcacccct	ttct		1544

<210> 6

<211> 1484

<212> DNA

<213> Staphylococcus aureus

<400> 6

ctggctcagg	atgaacgctg	gcggcgtgcc	taatacatgc	aagtcgagcg	aacggacgag	60
aagcttgctt	ctctgtatgtt	agcggcggac	gggtgagtaa	cacgtggata	acctacctat	120
aagactggaa	taacttcggg	aaacggcggac	taataccaga	taatattttt	aaccgcattgg	180
ttcaaaatgt	aaagacggtc	ttgctgtac	ttatagatgg	atccgcgtc	cattagctag	240
ttggtaatgt	aacggcttac	caaggcaacg	atgcatacg	gacctgagag	ggtgtatcgk	300
cacactggaa	ctgagacacg	gtccagactc	ctacgggagg	cagcgttgg	gaatttccg	360
caatgggcga	aaggctgacg	gagcaacg	gcgtgagtga	tgaaggctt	cgatcgtaa	420
aactctgtt	ttagggaa	acatatgtgt	aagtaactgt	gcacatctt	acggtaaccta	480
atcagaaaagc	cacggctaac	tacgtgccc	cagccgcgt	aatacgtagg	tggcaagcgt	540
tatccggaaat	tattgggcgt	aaagcgcgcg	taggcggttt	ttyaagtctg	atgtgaaagc	600
ccacggctca	accgtggagg	gtcattggaa	actggaaaac	ttgagtgcag	aagagggaaag	660
tggaattcca	tgtgttagcgg	tgaaatgcgc	agagatatgg	aggaacacca	gtggcgaagg	720
cgactttctg	gtctgttaact	gacgcgtatg	tgcgaaagcg	tgggtatcaa	acaggattag	780
atacccttgtt	agtccacg	gtaaacacgat	agtgtatgt	tttaggggt	ttccggccc	840

0450-0032.41-SEQLIST-corr

tagtgctgca	gctaacgcac	taagcactcc	gcctggggag	tacgaccgca	aggttgaaac	900
tcaaaggaaat	tgacggggac	ccgcacaagc	ggtggagcat	gtggtttaat	tcgaagcaac	960
gcgaagaacc	ttaccaaatac	ttgacatcct	ttgacaactc	tagagataga	gccttcccct	1020
tcgggggaca	aagtgcacagg	tggtgcatgg	ttgtcgctcg	ctcgtgtcg	gagatgttgg	1080
gttaagtccc	gcaacgagcg	caacccttaa	gcttagttgc	catcattaag	ttggcactc	1140
taagttact	gcccgtgaca	aaccggagga	aggtggggat	gacgtcaat	catcatcccc	1200
cttatgattt	gggctacaca	cgtgctacaa	tggacaatac	aaagggcagc	gaaaccggaga	1260
ggtcaagcaa	atcccataaa	gttgtctca	gttcggattt	tagtctgcaa	ctcgactaca	1320
tgaagctgga	atcgctagta	atcgtagatc	agcattctac	ggtgaatacg	ttcccgggtc	1380
ttgtacacac	cggccgtcac	accacgagag	ttttaacac	ccgaagccgg	tggagtaacc	1440
ttttaggagc	tagccgtcga	aggtgggaca	aatgatttggg	gtga		1484

<210> 7

<211> 1464

<212> DNA

<213> Mycobacterium tuberculosis

<400> 7

ggccgcgtgc	ttaacacatg	caagtcgaac	ggaaaggct	cttcggagat	actcgagtgg	60
cgaacgggtg	agtaacacgt	gggtgatctg	ccctgcactt	cgggataagc	ctggaaact	120
gggtctaata	ccggatagga	ccacggatg	catgtcttgc	ggtggaaagc	gctttagcg	180
tgtggatga	gcccgggcc	tatcagcttgc	ttgggggggt	gacggcctac	caaggcgacg	240
acgggtagcc	ggcctgagag	ggtgtccggc	cacactggga	ctgagatacg	gcccagactc	300
ctacgggagg	cagcagtggg	aatatttgc	caatgggcgc	aagcctgatg	cagcgacgcc	360
gcgtggggga	tgacggctt	cgggttgtaa	acctcttca	ccatcgacga	aggtccgggt	420
tctctcgat	tgacgttagg	tggagaagaa	gcaccggcca	actacgtgcc	agcagccg	480
gtaatacgt	gggtgcgagc	gttgtccgg	attactgggc	gtaaagagct	cgtaggttgt	540
ttgtcgcgtt	gttcgtaaa	tctcacggct	taactgtgag	cgtgcggg	atacgggcag	600
actagagttac	tgcaggggag	acttggattc	ctgggttagc	ggtggaaatgc	gcagatatca	660
ggaggaacac	cgggtggc	ggcgggttctc	tgggcagtaa	ctgacgctga	ggagcgaaag	720
cgtggggagc	gaacaggatt	agatccctgt	gtatccacg	ccgtaaacgg	tgggtactag	780
gtgtgggtt	ccttccctgg	gatccgtg	gtagtaac	cattaatgc	ccgccttgg	840
gagtagcc	gcaaggctaa	aactcaaagg	aatttgcggg	ggccgcaca	agcggcgag	900
catgtggatt	aattcgatgc	aacgcgaaga	accttacctg	ggtttgacat	gcacaggacg	960
cgtctagaga	taggcgttcc	cttgcggct	gtgtcgaggt	ggtgcacggc	tgtcgacgc	1020
tcgtgtcg	agatgttgg	ttaagtcccg	caacgagcgc	aacccttgc	tcatgttgc	1080
agcacgtat	ggtggggact	cgtgagagac	tgccgggtc	aactcggagg	aagggtggga	1140
tgacgtcaag	tcatcatg	ccttatgtcc	agggcttac	acatgctaca	atggccggta	1200
caaaggctg	cgtgcgcgc	agtttaagcg	aatccctaaa	agccgtctc	agttcgatc	1260
gggtctgca	actcgacccc	gtgaagtgc	agtcgttagt	aatcgacat	cagcaacgc	1320
gcgtgtata	cgttcccg	ccttgcac	accgcgcgtc	acgtcatgaa	agtcgttaac	1380
accgaagcc	agtggctaa	ccctcgggag	ggagctgtcg	aagggtggat	cggcgattgg	1440
gacgaagtgc	taacaaggt	gccc				1464

<210> 8

<211> 1450

<212> DNA

<213> Helicobacter pylori

<220>

<221> misc_feature

<222> 912, 927, 928, 929, 930, 1083

<223> n = A,T,C or G

<400> 8

tttatggaga	gtttgatcct	ggctcagagt	gaacgctggc	ggcgtgccta	atacatgca	60
gtcgaacgat	gaagcttcta	gcttgcata	gtgctgat	gtggcgcacg	ggtgagtaac	120
gcataaggta	tgtgcctt	agtttggat	agccatttgg	aacgatgatt	aataccagat	180
actcctacgg	gggaaagatt	tatcgctaag	agatcagct	atgtcctatc	agcttgggt	240
taaggtaatg	gcttaccaag	gtatgacgg	gtatccggcc	ttaggggtg	aacggacaca	300
ctggaaactg	gacacggtcc	agactccat	gggaggcagc	agtagggat	attgctcaat	360
gggggaaacc	ctgaagcagc	aacgcgcgt	ggaggatgaa	ggttttagga	ttgtaaactc	420
cttttggtag	agaagataat	gacggtatct	aacgaataag	caccggctaa	ctccgtgcca	480
gcagccgcgg	taatacggag	ggtgcaagcg	ttactcgaa	tcactggcg	taaagagcgc	540
gtaggcggga	tagtcagtca	gggtgtaaat	cctatggctt	aaccatagaa	ctgcatttga	600
aactactatt	ctagagtgt	ggagaggtag	gtggatttct	tgggttaggg	gtaaaatccg	660
tagagatcaa	gaggaatact	cattgcgaag	gcaacatttgc	tgacgctgt		720
tgcgttaaaag	cgtggggagc	aaacaggatt	agatccctg	gtagtccacg	ccctaaacga	780
tggatgctag	ttgttggagg	gcttagtctc	tccagtaatg	caacgtacgc	attaagcatc	840

0450-0032.41-SEQLIST-corr

ccgcctgggg	agtacggtcg	caagattaaa	actcaaagga	atagacgggg	acccgcacaa	900
gcggtgagc	angtgttta	attcgannnn	acacgaagaa	ccttacctag	gcttgcacatt	960
gagagaatcc	gctagaataa	gtggagtgtc	tagttgcta	gaccttgaaa	acaggtgctg	1020
cacggctgtc	gtcagctgt	gtcgtgagat	gttgggttaa	gtcccgcac	gagcgaacc	1080
ccnttctta	gttgctaaca	ggttatgctg	agaactctaa	ggatactgcc	tccgtaaagg	1140
ggaggaaggt	ggggacgacg	tcaagtcatc	atggccctta	cgcctaggc	tacacacgtg	1200
ctacaatggg	gtgcacaaag	agaagcaata	ctgtgaagtg	gagccaatct	tcaaaacacc	1260
tctcagttcg	gattgttaggc	tgcaactcgc	ctgcatgaag	ctggaatcgc	tagtaatcgc	1320
aaatcagcca	tgttgcggtg	aatacgttcc	cgggtcttgc	actcaccgc	cgtcacacca	1380
tgggagttgt	gtttgcctta	agttaggtg	ctaaattggc	tactgcccac	ggcacacaca	1440
gcgactgggg						1450

<210> 9

<211> 1515

<212> DNA

<213> *Streptococcus pneumoniae*

<400> 9

atttgatcct	ggctcaggac	gaacgctggc	ggcgtgccta	atacatgca	gtagaacgct	60
gaaggaggag	cttgcttctc	tggatgagtt	gcgaacgggt	gagtaacgcg	tagtaacct	120
gcctggtagc	gggggataac	tattggaaac	gataactaat	accgcataag	agtggatgtt	180
gcatgacatt	tgcttaaaag	gtgcacttgc	atcaactacca	gatggacact	cgtgttatta	240
gctagttgtt	ggggtaacgg	ctcaccaagg	cgacgataca	tagccgac	gagagggtga	300
tcggccacac	tgggacttag	acacgkccca	gactcttacg	ggaggcagca	gtagggaaatc	360
ttcggcaatg	gacggaaatc	tgaccgagca	acgcgcgtg	agtgaagaag	gttttcggat	420
cgtaaagctc	tgttgtaa	gaagaacgag	tgtgagagtg	gaaagttcac	actgtgacgg	480
tatcttacca	gaaaggacg	gctaactacg	tgccagcagc	cgccgtaaata	cgttagtccc	540
gagcgttgc	cggatttatt	gggcgttaaag	cgagcgcagg	cggttagata	agtctgaagt	600
taaaggctgt	ggcttaacca	tagtaggctt	tggaaactgt	ttaactttag	tgcaagaggg	660
gagagtggaa	ttccatgtgt	agcggtaaa	tgcgtagata	tatggagaa	caccgggtggc	720
gaaagcggct	ctctggcttg	taactgacgc	tgaggctcga	aagcgtgggg	agcaaacagg	780
attagatacc	ctggtagtcc	acgctgtaaa	cgatgagtg	taggtgttag	acccttccg	840
gggttagtg	ccgtagctaa	cgcattaagc	actccgcctg	gggagtagca	ccgcaagggtt	900
gaaactcaaa	ggaatttgac	ggggcccgca	caagcggtgg	agcatgtgtt	ttaattcgaa	960
gcaacgcgaa	gaaccttacc	aggtcttgc	atccctctga	ccgctctaga	gatagagttt	1020
tccctcggg	cagagggtac	aggtgggtca	tggtgtcgt	cagctgtgt	cgtgagatgt	1080
tgggttaagt	cccgcaacga	gcbcaccc	tatttttagt	tgccatcatt	caggggca	1140
ctctagcgag	actgcccgt	ataaaaccgg	ggaaggtggg	gatgacgtca	aatcatcatg	1200
cccttatga	cctggctac	acacgtgcta	caatggctgg	tacaacgagt	cgcaagccgg	1260
tgacggcaag	ctaatctt	aaagccagtc	tcagttcga	ttttaggtgt	caactcgcct	1320
acatgaagtc	ggaatcgcta	gtaatcgccg	atcagcacgc	cgcgtgaat	acgtcccccgg	1380
gccttgtaca	caccggccgt	cacaccacga	gagttttaa	caccgaagt	cggtgaggta	1440
accctaagga	gccagccgccc	taaggtggg	tagatgattg	gggtgaagtc	gtaacaaggt	1500
cagccgtttg	ggaga					1515

<210> 10

<211> 1544

<212> DNA

<213> *Treponema palladium*

<400> 10

agagtttgat	catggctcag	aacgaacgct	ggcggcgcgt	cttaagcatg	caagtcgaac	60
ggcaagagag	gagcttgctt	cttcctaga	gtggcggact	ggtgaggaac	acgtgggtaa	120
tctaccctta	agatggggat	agctgctaga	aatagcaggt	aataccgaat	atactcagtg	180
cttcataagg	ggtatttggg	aaaggaagct	acggcttcgc	ttgaggatga	gcttgcgtcc	240
cattagctag	ttgggtgaggt	aaaggcccac	caaggcagc	atgggtatcc	ggcctgagag	300
ggtgatcrga	cacattggga	ctgagatacg	gccccaaactc	ctacgggagg	cagcagctaa	360
gaatattccg	caatggacgg	aagtctgacg	gagcgaacgc	gcgtggatga	agaaggctga	420
aaagttgtaa	aatccctttt	ttgtatga	ataagggtga	gagggatgc	tcatctgtat	480
acgtaatcg	acgataatacg	cccggttaat	tacgtccag	cagccgcgt	aacacgtaa	540
gggcgagcgt	tgttcgaaat	tattttggcgt	aaagggtatg	taggcgttgc	tgtaagcctg	600
atgtaaatc	ctggggctta	accccagaat	agcatgggt	actgtgtac	ttgaattacg	660
gaagggaaac	tggaaatcca	agttgtgggg	tggaaatctgt	agatattttg	aagaacaccg	720
gtggcgaagg	cggtttctg	gcccataatt	gacgctgaga	tgcgaaatgt	tggggatcga	780
acaggattag	atacccttgg	agtccacacc	gtaaacgtat	tacacttagt	gttggggccaa	840
gagttcagt	gccaagcaa	acgcgataag	tgtaccgcct	ggggagttatg	cccgcaagggg	900
tgaaactcaa	aggaatgtac	ggggggccgc	acaagcgtg	gagcatgtgg	tttaattcga	960
tggtagcgt	ggaaccttac	ctgggtttga	catctagtag	aaggctttag	agataaggcc	1020
ggtagcaat	accctgctag	acagggtctg	catggctgtc	gtcagctgt	gccgtgagggt	1080

0450-0032.41-SEQLIST-corr

gttgggttaa	gtcccgcaat	gagcgcaacc	cctactgcca	gttactaaca	ggtaaagctt	1140
gaggactctg	gcggactgc	cgatgacaaa	tcggaggaag	gtggggatga	cgtcaagtc	1200
tcatggccct	tatgtccagg	gctacacacg	tgctacaatg	gttgcataaa	agcgaagcaa	1260
gaccgttaagg	tggagcaagc	cgcaaaaaag	caatcgtagt	tcggattgaa	gtctgaaact	1320
cgacttcatg	aagttgaaat	cgctagtaat	cgcgcacac	cacggcgcgg	tgaatacggt	1380
ccggggcctt	gtacacaccg	cccgtcacac	catccgagtt	gggggtaccc	gaagtcgctt	1440
gtctaaccctg	caaaggagga	cggtgccgaa	ggtacgctt	gtaaggaggg	tgaagtcgta	1500
acaaggtagc	cgtaccggaa	ggtgccgctg	gatcacctcc	ttaa		1544

<210> 11
<211> 1548

<212> DNA

<213> Chlamydia trachomatis

<400> 11	ctgagaattt	gatcttggtt	cagattgaac	gctggcggcg	tggatgaggc	atgcaagtcg	60
	aacgggca	ttgtttcgcc	aattgtttag	tggcggaaagg	tttagtaatg	catagataat	120
	tttgccttaa	cttggaaata	acgggtggaa	acggccgcta	ataccgaat	tggcgatatt	180
	tgggcattcg	agtaacgtta	aagaaggggg	tcttaggacc	tttcggtaaa	gggagagtc	240
	atgtatatac	agctagttgg	ttgggttaaaag	gcctaccaag	gctatgacgt	ctaggcggat	300
	tgagagattt	gccgccaaca	ctgggactga	gacactgccc	agactccat	gggaggctgc	360
	agtcgagaat	ctttcgcaat	ggacggaaat	ctgacgaagc	gacgcccgt	gtgtatgaa	420
	ggctctaggg	ttgttaagca	ctttcgctt	ggaataagag	aagacggtt	ataccgctg	480
	gatttgagcg	taccaggtaa	agaaggcaccg	gctaactccg	tgccagcagc	tgcgtaata	540
	cggaggggtc	tagcgtaat	cggtttatt	ggccgtaaaag	gccgtgtagg	cgaaaaggta	600
	agttagttgt	caaagatcg	ggctcaaccc	cgagtcggca	tctaatacta	tttttctaga	660
	gggttagatgg	agaaaaggga	atttcacgt	tagcgtgaa	atgcgtat	atgttggaa	720
	acaccagtgg	cgaaggcgt	tttctaattt	atacctgacg	ctaaggcgcg	aaagcaaggg	780
	gagcaaaacag	gattagatac	cctggtagt	cttgcgtaa	acgatgcata	tttgatgtgg	840
	atgttctcaa	ccccatccgt	tgcggagct	acgcgttaag	tatgcccct	gaggataca	900
	ctcgcaaggg	tgaactcaa	agaatttgc	gggggcccgc	acaaggatgt	gagcatgtgg	960
	ttaatttca	tgcaacgcga	aggaccttac	ctgggtttga	catgtatat	accgcggcag	1020
	aaatgtcg	ttccgcagg	acatatacac	aggtgctgca	tggctgtcg	cagctcgtgc	1080
	cgtgaggtgt	tgggttaagt	cccgcaacga	gcgcacccct	tatcggtat	tgcagact	1140
	taggggtggg	actctaaca	gactgcctt	gttaaccagg	aggaaggcga	ggatgacgtc	1200
	aagttagcat	ggccctttag	cccagggcga	cacacgtct	acaatggca	gtacagaagg	1260
	tagcaagatc	gtgagatgg	gcaatccctc	aaagtggcc	ccagttcgg	ttttagtct	1320
	caactcgact	acatgaagtc	ggaatttcta	gtatggct	gtcagccata	acgcgtgaa	1380
	tacgttcccg	ggccttgtac	acaccgccc	tcacatcat	ggagttgtt	ttaccttaag	1440
	tcgttgactc	aacccgcaag	gagagaggcg	cccaagggt	ggctgtat	tagatgaag	1500
	tcgttacaag	gtagccctac	cggaagggtt	ggctggatca	cctcctt		1548

<210> 12
<211> 1466

<212> DNA

<213> Bartonella henselae

<220>							
<221> misc_feature							
<222> 1311							
<223> n = A,T,C or G							
<400> 12							
tcctggctca	ggatgaacgc	tggcggcagg	cttaacacat	gcaagtcgag	cgcactcatt		60
tagagtggc	ggcagacggg	ttagtaacgc	gtggaaatct	acccttttct	acggaataac		120
acagagaaat	ttgtgctaat	accgtatacg	tcctacttgg	gaaagattt	tcggagaagg		180
atgagccgc	gttggattag	ctagttggt	aggtaaaggc	tcaccaaggc	gacgatccat		240
agctggctcg	agaggatgt	cagccacact	gggactgaga	cacggcccg	actcctacgg		300
gaggcagcag	tggggatat	tggacaatgg	gggcaacccct	gatcgat	tgccgcgtga		360
gtatgtgggg	ccctagggtt	gtaaagctt	ttcaccgggt	aatcgat	cggttaaccgg		420
agaagaagcc	ccggctaact	tcgtgcacgc	agccgcgtt	atacgaaagg	ggctagcg		480
gttcggattt	actgggcgt	aagcgtatgt	aggcggat	ttaagtca	ggtgaaatcc		540
cagggctcaa	ccctggaaact	gcctttgata	ctggatatct	tgatgtat	agaggtgagt		600
ggaatttccg	gtgttagaggt	aaaattcgta	gatatttgg	ggaacaccag	tggcgaaggc		660
ggctcactgg	tccattactg	acgctgaggt	gcaaaagcgt	gggggagaaa	caggattaga		720
taccctggta	gtccacgc	taaacatgt	atgttagcc	ttgggtgtt	tactgtct		780
tggcgcacgt	aacgcattaa	acattccgccc	tggggagtt	ggtcgcaaga	ttaaaactca		840
aagaatttga	cggggccccg	cacaagggt	ggagcatgt	ttttaattcg	aagcaacgcg		900
cagaaccccta	ccagcccttgc	acatcccgat	cgcggaaat	ggagacaccc	tccttcgtt		960

0450-0032.41-SEQLIST-corr

cggctggatc	ggagacaggt	gctgcattgc	tgtcgctcagc	tcgtgtcg	agatgttggg	1020
ttaagtcccg	caacgagcgc	aaccctcg	cttagttgc	agcattcagt	tggcactct	1080
agggggactg	ccgggtataa	gccgagagga	aggtggggat	gacgtcaagt	cctcatggcc	1140
cttacgggc	gggtacaca	cgtgctacaa	tggttgtac	agtgggcagc	gagatcgcaa	1200
ggtcgagcta	atctccaaaa	gccatctca	ttcggattgc	actctgcaac	tcgagtgc	1260
gaagttggaa	tcgctagtaa	tcgtggatca	gcatgctacg	gtgaatacgt	nccggggct	1320
tgtacacacc	gcccgctaca	ccatgggagt	tggttttacc	cgaagggtct	gtctaaccg	1380
caaggaggca	ggttaaccacg	gttagggtc	cgactggggt	gaagtcgtaa	caaggttagcc	1440
gtagggaaacc	tgcggctgga	tcacct				1466

<210> 13

<211> 1487

<212> DNA

<213> *Hemophilis influenza*

<220>

<221> misc_feature

<222> 1, 373, 541, 629, 930, 1182, 1387

<223> n = A,T,C or G

<400> 13

naattgaaga	gtttgatcat	ggctcagatt	gaacgctggc	ggcaggctta	acacatgcaa	60
gtcgaacgg	agcaggagaa	agttgc	cttgcgtac	agtggcggac	gggtgagtaa	120
tgcttggaa	tctggcttat	ggagggggat	aacgacggga	aactgtcgt	aataccgcgt	180
attatcgaa	gatgaaagt	cgggactgag	aggccgc	ccataggat	agcccaagt	240
ggatttagta	gttgggggg	taatgccta	ccaaggctgc	gatctctag	tggcttgaga	300
ggatgaccag	ccacactgga	actgagacac	gttccagact	cctacgggag	gcagcagtgg	360
ggaatattgc	gcnatgggg	gaacccgtac	gcagccatgc	cgcgtgaat	aagaaggcct	420
tcgggttga	aagtcttc	ggtattgagg	aagggtat	tgttaatag	acatcaaatt	480
gacgttaaat	acagaagaag	caccggctaa	ctccgtgc	gcagccgc	taatacggag	540
ngtgcagcg	ttaatcgaa	taactggcg	taaaggggac	gcaggcggtt	atthaagtga	600
ggtgtgaaag	ccccgggctt	aacctggg	ttgcattca	gactgggtaa	ctagagact	660
ttagggaggg	gtagaattcc	acgtgtac	gtgaaatgc	tagagatgt	gaggaatacc	720
gaaggcgaag	gcagccctt	ggaatgtac	tgacgctcat	gtgcgaaagc	gtggggagca	780
aacaggatta	gataccctgg	tagtccacgc	tgttaacgct	gtcgatttgg	gggttgggg	840
ttaactctgg	caccgttagc	taacgtgata	aatcgaccgc	ctggggagta	cggccgcaag	900
gttaaaactc	aaatgaatt	acggggccn	gcacaagcgg	tggagcatgt	ggttaattc	960
gatgcaacgc	gaagaacctt	acctactct	gacatcctaa	gaagagctca	gagatgagct	1020
tgtgccttcg	ggaactttaga	gacaggtgct	gcatgctgt	cgtcagctcg	tgttgtgaaa	1080
tgttgggtt	agtccgc	cgagcgaac	ccttac	tgttgc	gacttgg	1140
ggaactcaa	ggagactg	agtgttaaa	tggagaaagg	tnggat	gtcaagt	1200
catggccctt	acgatgtgg	ctacacacgt	gctacaatgg	cgtatacaga	gggaagcga	1260
gctgcgagg	ggagcgaatc	tcataaaagta	cgtctaagtc	cggattggag	tctgcact	1320
gactccatga	agtcggaatc	gctagtaatc	gcaatcaga	atgtcg	gaatacg	1380
ccgggcntt	tacacaccgc	ccgtcacacc	atgggatgg	gttgc	taccacagg	1440
cttaacctt	tggaggg	gttaccacgg	atgat	actgggg	aaatgat	1487

<210> 14

<211> 1487

<212> DNA

<213> *Shigella dysenteriae*

<400> 14

tggctcagat	tgaacgctgg	cgccaggcc	aacacatgc	agtcgaacgg	taacagaaag	60
cagcttgc	tttgcgtac	agtggcggac	gggtgagtaa	tgtctgggaa	actgcctgat	120
ggagggggat	aactactg	aacggtagct	aataccgc	aacgtcg	gaccaaagag	180
ggggacctt	gggccttgc	ccatggat	tgcccagat	ggattagct	gtagg	240
taacggctc	cctaggcgac	gatccctag	tggtctgaga	ggatgacc	ccacactg	300
actgagacac	gttccagact	cctacgggag	gcagcgt	ggaatatt	acaatggcg	360
caaggctgat	gcagccatgc	cgcgttat	aagaaggc	tcgggtt	aatactt	420
agcggggagg	aagggat	atgttac	tttgcatt	gacgttac	gcagaaga	480
cacccgctaa	ctccgtgc	gcagccg	taatacgg	ggtcaag	ttaatcg	540
ttactggcg	taaagcgc	gcaggcg	tgttaat	gatgt	cccccggc	600
aacctggaa	ctgcatt	tactgg	cttgc	gtagagg	gtagaatt	660
aggtgtac	gtgaaatgc	tagatct	gaggat	ggtgg	gcggcccc	720
ggacaaac	tgacgctc	gtgc	gtgggg	gacaggat	gatacc	780
tagtccacgc	ctgttac	gtgcact	aggttgc	cttgcagg	ggcttcc	840
gctaacgcgt	taagtgc	gcctggg	tacggcc	aggtt	aaatgaat	900
tgacggggc	ccgcaca	gttg	gttg	at	tcgatgc	960

0450-0032.41-SEQLIST-corr

ttacctggtc ttgacatcca cagaacctt tagagatacg agggtgccctt cgggaactgt	1020
gagacagggt ctgcgtcgact cgtgttgcgt aatgttgggt taagtccgc	1080
aacgagcgca acccttatcc tttgttgcctt gcggtccggc cgggaactca aaggagactg	1140
ccagtataaa actggaggaa ggtggggatg acgtcaagtc atcatggccc ttacgaccag	1200
ggctacacac gtgctacaat ggcgcataca aagagaagcg acctcgcgag agcaagcgaa	1260
cctcataaaag tgcgtcgtag tccggattgg agtctgcaac tcgactccat gaagtccgaa	1320
tcgctagtaa tcgtggatca gaatgtcagc gtgaatacgt tcccgccct ttacacacc	1380
gcccgtcaca ccatgggagt gggttgcaaa agaagtaggt agcttaacct tcgggagggc	1440
gcttaccact ttgtgattca tgactggggt gaagtctgtaa caaggta	1487

<210> 15

<211> 22

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 15

ggactacgac gcactttatg ag

22

<210> 16

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 16

ggtccgcttg ctctcgcgag g

21

<210> 17

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 17

gcaaaggat taactttact c

21

<210> 18

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 18

gctgcggta ttaaccacaa c

21

<210> 19

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 19

gcactttatg aggtccgctt g

21

<210> 20

<211> 4

<212> DNA

<213> Unknown

<220>
<223> No sequence is present

<400> 20
000

<210> 21
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 21
tgctgcctcc cgttaggagtc t 21

<210> 22
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 22
attaccgcgg ctgctggcac g 21

<210> 23
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 23
accagggtat ctaatccctgt t 21

<210> 24
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 24
cacatgctcc accgcttgtg c 21

<210> 25
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 25
ttgcgggact taacccaaca t 21

<210> 26
<211> 4
<212> DNA
<213> Unknown

<220>
<223> No sequence is present

<400> 26	
000	
<210> 27	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 27	
cgccggctgct ggcacgtgt t	21
<210> 28	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
.....	
<400> 28	
acttaaccctt acatctcactt a	21
.....	
<210> 29	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
.....	
<220>	
<223> antisense oligomer	
<400> 29	
tttacgccccca gtaattccga	20
.....	
<210> 30	
<211> 22	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
.....	
<400> 30	
actccccatgg tgtgacgggc gg	22
.....	
<210> 31	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 31	
aatctgagcc atgatcaaac t	21
.....	
<210> 32	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 32	
ccctctttgt gcttgcgacg t	21

<210> 33		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 33		
accccccctct acgagactca a		21
<210> 34		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 34		
ccacgcctca agggcacaac c		21
.....	<210> 35	
.....	<211> 21	
.....	<212> DNA	
.....	<213> Artificial Sequence	
.....	<220>	
.....	<223> antisense oligomer	
.....	<400> 35	
.....	tctcatctct gaaaacttcc g	21
.....	<210> 36	
.....	<211> 21	
.....	<212> DNA	
.....	<213> Artificial Sequence	
.....	<220>	
.....	<223> antisense oligomer	
.....	<400> 36	
.....	catgatcaaa ctcttcaatt t	21
.....	<210> 37	
.....	<211> 21	
.....	<212> DNA	
.....	<213> Artificial Sequence	
.....	<220>	
.....	<223> antisense oligomer	
.....	<400> 37	
.....	ccctctttgg tcttgcgacg t	21
<210> 38		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 38		
taccccccctc tacgagactc a		21
<210> 39		
<211> 21		

0450-0032.41-SEQLIST-corr

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 39
gccacgcctc aaggcacaa c 21

<210> 40
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 40
cagagagcaa gccctttca t 21

<210> 41
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 41
cctgctttct cccgttaggac g 21

<210> 42
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 42
caccacccctc tgccatactc t 21

<210> 43
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 43
ctaagatctc aaggatccca a 21

<210> 44
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 44
ggctgtccgc cagcgttcaa t 21

<210> 45
<211> 21
<212> DNA
<213> Artificial Sequence

<220>		
<223> antisense oligomer		
<400> 45		
ccctcttgg tccgtaaaca t		21
<210> 46		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 46		
ccccctctac aagactctag c		21
<210> 47		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
.....<223> antisense oligomer		
.....<400> 47		
.....acgactytag gtcacaacct c		21
.....<210> 48		
.....<211> 21		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
.....<400> 48		
.....aggatcaaac tcttatgttc a		21
.....<210> 49		
.....<211> 21		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
.....<400> 49		
.....cctgcttcc ctctcaagac g		21
.....<210> 50		
.....<211> 21		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
.....<400> 50		
.....cacccccc tgacacactc g		21
.....<210> 51		
.....<211> 21		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		

<400> 51
 ccaagcaatc aagttgccca a
 <210> 52
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 52
 ccagcggtca tcctgagcca g

<210> 53
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 53
 gaaccatgcg gttcaaaata t

<210> 54
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 54
 ctttcctctt ctgcactcaa g

<210> 55
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 55
 ggggcggaaa cccccctaaca c

<210> 56
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 56
 gcatgtgtta agcacgcccgc c

<210> 57
 <211> 21
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 57
 aagacatgca tcccggtggtc c

<210> 58		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 58		
cagtctcccc tgcagtactc t		21
<210> 59		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 59		
gatcccaagg aaggaaaccc a		21
<210> 60		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 60		
caggatcaaa ctctccataaa a		21
<210> 61		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 61		
aaatcttcc cccgttaggag t		21
<210> 62		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 62		
cacctacctc tccccacactc t		21
<210> 63		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 63		
tggagagact aagccctcca a		21
<210> 64		
<211> 21		
<212> DNA		

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 64

cgtcctgagc caggatcaaa t

21

<210> 65

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 65

atgtcatgca acatccactc t

21

<210> 66

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 66

actctccct cttgcactca a

21

<210> 67

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 67

aaaccccgga aagggtctaa c

21

<210> 68

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 68

tctgagccat gatcaaactc t

21

<210> 69

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 69

acccttatg aagcactgag t

21

<210> 70

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

0450-0032.41-SEQLIST-corr

<223> antisense oligomer		
<400> 70		
agtttccctt ccgtaattca a		21
<210> 71		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 71		
cactgaagct cttgccccaa c		21
<210> 72		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
.....		
<400> 72		
gaaccaagat caaattctca g		21
.....		
<210> 73		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
.....		
<220>		
<223> antisense oligomer		
<400> 73		
gttactcgga tgcccaaata t		21
.....		
<210> 74		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
.....		
<220>		
<223> antisense oligomer		
.....		
<400> 74		
ccttttctcc atctaccctc t		21
.....		
<210> 75		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
.....		
<220>		
<223> antisense oligomer		
<400> 75		
ggatggggtt gagaccatcc a		21
.....		
<210> 76		
<211> 21		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 76		

agcgttcatc ctgagccagg a

<210> 77
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 77
aaatctttct ccagtaggac g

<210> 78
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 78
cactcacctc ttccatactc a

21

.....<210> 79
.....<211> 21
.....<212> DNA
.....<213> Artificial Sequence

.....<220>
.....<223> antisense oligomer

....<400> 79
....<actgagcagt aaaccaccca a

21

.....<210> 80
.....<211> 21
.....<212> DNA
.....<213> Artificial Sequence

.....<220>
.....<223> antisense oligomer

.....<221> misc_feature
.....<222> 21
.....<223> n = A,T,C or G

<400> 80
catgatcaaa ctcttcaatt n

21

.....<210> 81
.....<211> 21
.....<212> DNA
.....<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 81
cactttcatc ttccgataat a

21

<210> 82
<211> 21
<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 82	21
cctccctaaa gtactctagt t	
<210> 83	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 83	21
cagagttaaa ccccaacccc c	
<210> 84	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 84	21
gccagcggttc aatctgagcc a	
<210> 85	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 85	21
ccctctttgg tcttgcgacg t	
<210> 86	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 86	21
taccccccctc tacgagactc a	
<210> 87	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 87	21
gccacgcctc aagggcacaa c	
<210> 88	
<211> 21	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 88	21
cctcgttatct ctacaaggtt c	

<210> 89		
<211> 23		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 89		
ccccatcatt atgagtatgatg tgc		23
<210> 90		
<211> 19		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 90		
tcattatgatg gtgacccca		19
<210> 91		
<211> 20		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 91		
gatgaacagt tactctcatc		20
<210> 92		
<211> 20		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 92		
actgagagaa gctttaagag		20
<210> 93		
<211> 20		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 93		
atgtgcacag ttacttacac		20
<210> 94		
<211> 20		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 94		
ctgagaacaa ctttatggga		20
<210> 95		
<211> 20		
<212> DNA		

0450-0032.41-SEQLIST-corr

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 95

ttattctgtt ggtaacgtca

20

<210> 96

<211> 19

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 96

cgagttgcag actgcgatc

19

<210> 97

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 97

atctgagcca tcatcaaact

20

<210> 98

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 98

tgtctcgtt ccagtgttgc

20

<210> 99

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 99

gttttcgtcc agggggccgc

20

<210> 100

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer

<400> 100

caccctgtctc acgggtttcccg

20

<210> 101

<211> 20

<212> DNA

<213> Artificial Sequence

<220>

<223> antisense oligomer	
<400> 101	
cgccctcccg aagttaagct	20
<210> 102	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 102	
ggcacgccgc cagcgttcg	19
<210> 103	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 103	
tgtctcagtc ccaatgtggc	20
<210> 104	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 104	
gttacagacc agagagccgc	20
<210> 105	
<211> 20	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	,
<400> 105	
cacctgtcac tttgcccccg	20
<210> 106	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 106	
ggcggctggc tccaaaagg	19
<210> 107	
<211> 19	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 107	

caccggttcg ccactcctc
 <210> 108
 <211> 20
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 108
 tcaattcctt tgagttcaa

<210> 109
 <211> 29
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> antisense oligomer

<400> 109
 gcaatccgaa ctgagagaag ctttaagag

.....<210> 110
<211> 24
<212> DNA
<213> Artificial Sequence

.....<220>
<223> antisense oligomer

.....<400> 110
 ccgaactgag agaagcttta agag

.....<210> 111
<211> 17
<212> DNA
<213> Artificial Sequence

.....<220>
<223> antisense oligomer

.....<400> 111
 gagagaagct ttaagag

.....<210> 112
<211> 15
<212> DNA
<213> Artificial Sequence

.....<220>
<223> antisense oligomer

.....<400> 112
 gagaagcttt aagag

.....<210> 113
<211> 12
<212> DNA
<213> Artificial Sequence

.....<220>
<223> antisense oligomer

.....<400> 113
 aagctttaag ag

<210> 114

```

<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligomer
<400> 114
gactaccagg gtatctaatc 20
<210> 115
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligomer
<400> 115
cagcgacacc cgaaagcgcc 20
<210> 116
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligomer
<400> 116
gtgccaaggc atccaccgtg 20
<210> 117
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligomer
<400> 117
catactcaaa cgccctattc 20
<210> 118
<211> 19
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligomer
<400> 118
ccttagcctc ctgcgtccc 19
<210> 119
<211> 20
<212> DNA
<213> Artificial Sequence
<220>
<223> antisense oligomer
<400> 119
ggggtcttcc cgtcctgtcg 20
<210> 120
<211> 20
<212> DNA
<213> Artificial Sequence

```

<220>		
<223> antisense oligomer		
<400> 120		
cgatcgatta gtatcagtcc		20
<210> 121		
<211> 18		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 121		
tgagagaagc tttaagag		18
<210> 122		
<211> 19		
<212> DNA		
<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
.....<400> 122		
.....ctgagagaag cttaagag		19
.....<210> 123		
.....<211> 18		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
<400> 123		
gacacacccg aaagcgcc		18
.....<210> 124		
.....<211> 18		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
<400> 124		
tacagaccag agagccgc		18
.....<210> 125		
.....<211> 17		
.....<212> DNA		
.....<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 125		
cgacacccga aagcgcc		17
<210> 126		
<211> 19		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		

<400> 126		
agcgacacccc gaaagcgcc		19
<210> 127		
<211> 18		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 127		
cgacacccga aagcgctt		18
<210> 128		
<211> 17		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
.....<400> 128		
.....mgamammmga aagmgmm		17
.....<210> 129		
.....<211> 18		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
.....<400> 129		
.....tamagammag agagmmgm		18
.....<210> 130		
.....<211> 16		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
.....<400> 130		
.....mmmmammmttm mtmmgg		16
.....<210> 131		
.....<211> 19		
.....<212> DNA		
.....<213> Artificial Sequence		
.....<220>		
.....<223> antisense oligomer		
<400> 131		
caccgcggcg tgctgatcc		19
<210> 132		
<211> 16		
<212> DNA		
<213> Artificial Sequence		
<220>		
<223> antisense oligomer		
<400> 132		
ccccacacctc ctccgg		16

<210> 133	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 133	
ccgcttgtgc gggccccc	18
<210> 134	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 134	
caccgcggcg tgctgatc	18
•••• <210> 135	
<211> 17	
<212> DNA	
<213> Artificial Sequence	
•••• <220>	
<223> antisense oligomer	
•••• <400> 135	
caccgcggcg tgctgat	17
•••• <210> 136	
<211> 18	
<212> DNA	
<213> Artificial Sequence	
•••• <220>	
<223> antisense oligomer	
•••• <400> 136	
accgcggcgt gctgatcc	18
•••• <210> 137	
<211> 17	
<212> DNA	
<213> Artificial Sequence	
•••• <220>	
<223> antisense oligomer	
•••• <400> 137	
ccgcggcgtg ctgatcc	17
<210> 138	
<211> 17	
<212> DNA	
<213> Artificial Sequence	
<220>	
<223> antisense oligomer	
<400> 138	
accgcggcgt gctgatc	17
<210> 139	
<211> 20	

0450-0032.41-SEQLIST-corr

<212> DNA
<213> Artificial Sequence

<220>
<223> antisense oligomer

<400> 139
acgttgaggg gcatcgtcgc

20

.....

.....

.....

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. An antibacterial compound consisting of a substantially uncharged antisense oligomer containing from 10 to 40 nucleotide subunits, each of said subunits comprising a morpholino ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base, said base-pairing moieties including a targeting nucleic acid sequence at least 10 nucleotides in length which is complementary to a bacterial 16S or 23S rRNA nucleic acid sequence,
5 wherein adjacent subunits are joined by uncharged linkages selected from the group consisting of uncharged phosphoramidate and phosphorodiamidate, or by charged phosphoramidate linkages, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1.
- 10 2. The compound of claim 1, wherein said oligomer is able to hybridize with the bacterial sequence at a T_m substantially greater than the T_m of a duplex composed of a corresponding DNA and the same bacterial sequence.
- 15 3. The compound of claim 1 or claim 2, wherein said oligomer is able to hybridize with the bacterial sequence at a T_m substantially greater than 37°C.
- 20 4. The compound of any preceding claim, wherein the uncharged linkages are selected from the group consisting of the structures presented in Figures 2A through 2D, where:
25 X is fluorine, an alkyl or substituted alkyl, an alkoxy or substituted alkoxy, a thioalkoxy or substituted thioalkoxy, or an unsubstituted, monosubstituted, or disubstituted nitrogen;
Z is sulfur or oxygen;
Y, in Figs. 2B and 2D, is sulfur, nitrogen, carbon or oxygen, and in Fig. 30 2C is methylene, sulfur, or oxygen;
and each P_i and P_j is a purine or pyrimidine base-pairing moiety effective to bind, by base-specific hydrogen bonding, to a base in a polynucleotide.
- 35 5. The compound of claim 4, wherein each uncharged linkage is a phosphorodiamidate linkage as represented at Figure 2B, where $X=NR_2$, where R is hydrogen or methyl, $Y=O$, and $Z=O$.

6. The compound of any preceding claim, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 8:1.

7. The compound of any preceding claim, wherein the antisense oligomer has a length of from 12 to 25 subunits.

5

8. The compound of any preceding claim, wherein the region of complementarity with the target RNA sequence has a length of from 13 to 20 bases.

10 9. The compound of any preceding claim, wherein the targeting sequence is selected from the group consisting of SEQ ID NOs : 15, 16, and 21-25.

15

10. The compound of any preceding claim, where the targeting sequence is complementary to a Gram-positive bacterial 16S rRNA consensus sequence or a Gram-negative bacterial 16S rRNA consensus sequence.

11. The compound of claim 10, where the targeting sequence is selected from the group consisting of SEQ ID NOs : 27-30.

20 12. The compound of any one of claims 1 to 9, wherein the targeting sequence is SEQ ID NO: 92.

13. A method of treating a bacterial infection in a human or mammalian animal subject, comprising administering to the subject, in a pharmaceutically effective amount, a substantially uncharged antisense oligomer containing from 10 to 40 nucleotide subunits, each of said subunits comprising a morpholino ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base, said base-pairing moieties including a targeting nucleic acid sequence at least 10 nucleotides in length which is complementary to a bacterial 16S or 23S rRNA nucleic acid sequence,

25

30

35

wherein adjacent subunits are joined by uncharged linkages selected from the group consisting of uncharged phosphoramidate and phosphorodiamidate, or by charged phosphoramidate linkages, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1.

14. The method of claim 13, wherein said oligomer is able to hybridize with the bacterial sequence at a T_m substantially greater than 37°C.

15. The method of claim 13 or claim 14, wherein the uncharged linkages are selected from the group consisting of the structures presented in Figures 2A through 2D, where:

X is fluorine, an alkyl or substituted alkyl, an alkoxy or substituted alkoxy, a thioalkoxy, or substituted thioalkoxy, or an unsubstituted, monosubstituted, or disubstituted nitrogen;

Z is sulfur or oxygen;

Y in Figs, 2B and 2D, is sulfur, nitrogen, carbon or oxygen, and in Fig. 2C is methylene, sulfur, or oxygen;

and each P_i and P_j is a purine or pyrimidine base-pairing moiety effective to bind, by base-specific hydrogen bonding, to a base in a polynucleotide.

16. The method of claim 15, wherein each uncharged linkage is a phosphorodiamidate linkage as represented at Figure 2B, where X=NR₂, where R is hydrogen or methyl, Y=O, and Z=O.

17. The method of any one of claims 13 to 16, where the antisense oligomer has a length of from 12 to 25 bases.

18. The method of any one of claims 13 to 17, where the region of complementarity with the target RNA sequence has a length of from 13 to 20 bases.

19. The method of any one of claims 13 to 18, wherein the targeting sequence is selected from the group consisting of SEQ ID NOs : 15, 16 and 21-25.

20. The method of any one of claims 13 to 19, where the targeting sequence is complementary to a Gram-positive bacterial 16S rRNA consensus sequence or a Gram-negative bacterial 16S rRNA consensus sequence.

21. The method of claim 20, where the targeting sequence is selected from the group consisting of SEQ ID NOs: 27-30.

22. The method of any one of claims 13 to 19, for use in treatment of an infection produced by *E. coli*, *Salmonella thyphimurium*, *Pseudomonas aeruginosa*, *Vibrio*

cholera, Neisseria gonorrhoea, Helicobacter pylori, Bartonella henselae, Hemophilis Influenza, Shigella dysenteriae, Staphylococcus aureus, Mycobacterium tuberculosis, Streptococcus pneumoniae, Treponema palladium and Chlamydia trachomatis, where the antisense oligomer has a sequence selected from the group consisting of SEQ ID NOS: 21-25.

5

10

15

20

25

30

35

23. The method of any one of claims 13 to 22, wherein the antisense oligomer is administered in an amount and manner effective to result in a peak blood concentration of at least 200-400 nM antisense oligomer.
24. The method of any one of claims 13 to 22 for treating bacterial infections of the skin, wherein said administering is by a topical route.
25. The method of any one of claims 13 to 22 for use in treating a bacterial respiratory infection, wherein said administering is by inhalation.
26. A livestock and poultry food composition containing a food grain supplemented with a subtherapeutic amount of an antibacterial compound, said compound consisting of a substantially uncharged antisense oligomer containing from 10 to 40 nucleotide subunits, each of said subunits comprising a morpholino ring supporting a base-pairing moiety effective to bind by Watson-Crick base pairing to a respective nucleotide base, said base-pairing including a targeting nucleic acid sequence at least 10 nucleotides in length which is complementary to a bacterial 16S or 23S rRNA nucleic acid sequence,
wherein adjacent subunits are joined by uncharged linkages selected from the group consisting of uncharged phosphoramidate and phosphorodiamidate, or by charged phosphoramidate linkages, wherein the ratio of uncharged linkages to charged linkages in the oligomer is at least 4:1.
27. The composition of claim 28, wherein each linkage is a phosphorodiamidate linkage as represented at Figure 2B, where $X = NR_2$, where R is hydrogen or methyl, Y=0, and Z=0.
28. The composition of claim 26 or claim 27, wherein the antisense oligomer has a length of from 12 to 25 bases.

29. The composition of any one of claims 26 to 28, wherein the region of complementarity with the target RNA sequence has a length of 13 to 20 bases.

5 30. The composition of any one of claims 26 to 29, wherein the targeting sequence is selected from the group consisting of SEQ ID NOs : 15, 16, 21-25 and 27-30.

31. A compound according to claim 1, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

10 32. A method according to claim 13, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.

33. A composition according to claim 26, substantially as herein described with reference to any one or more of the Examples and/or accompanying Figures.





15

Dated this twenty-first day of June 2005





AVI BioPharma, Inc.
Patent Attorneys for the Applicant:

F B RICE & CO








1/11

**Fig. 1A****Fig. 1B****Fig. 1C****Fig. 1D**

2/11

**Fig. 2A****Fig. 2B****Fig. 2C****Fig. 2D**

3 / 11



4 / 11






Fig. 4


5/11

**Fig. 5**

6/11

**Fig. 6**

7/11

**Fig. 7**

8 / 11




Fig. 8

9/11






Fig. 9

10/11



11/11

**Fig. 10B**