



(51) International Patent Classification:

C07D 401/14 (2006.01) *A61P 35/00* (2006.01)
A61K 31/4439 (2006.01)

(21) International Application Number:

PCT/EP2011/072640

(22) International Filing Date:

13 December 2011 (13.12.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/422,422 13 December 2010 (13.12.2010) US

(71) Applicant (for all designated States except US): **NO-VARTIS AG** [CH/CH]; Lichtstrasse 35, CH-4056 Basel (CH).

(72) Inventors; and

(75) **Inventors/Applicants (for US only): STRAUB, Christopher Sean** [US/US]; Novartis Institutes for BioMedical Research, Inc., 250 Massachusetts Avenue, Cambridge, Massachusetts 02139 (US). **CHEN, Zhuoliang** [CN/US]; Novartis Institutes for BioMedical Research, Inc., 250 Massachusetts Avenue, Cambridge, Massachusetts 02139 (US).

(74) Agent: **STRANG, Andrea**; Novartis Pharma AG, Patent Department, CH-4002 Basel (CH).

(81) **Designated States** (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) **Designated States** (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

Published:

— with international search report (Art. 21(3))

(54) Title: DIMERIC IAP INHIBITORS



(57) **Abstract:** The present invention provides compounds of formula M-L-M' (where M and M' are each independently a monomeric moiety of Formula (I) and L is a linker). The dimeric compounds have been found to be effective in promoting apoptosis in rapidly dividing cells. (I)

DIMERIC IAP INHIBITORS

FIELD OF THE INVENTION

The present invention relates to dimeric compounds that act as inhibitors of the
5 Inhibitor of Apoptosis Proteins (IAPs), as well as pharmaceutical compositions thereof,
methods of their use, and methods for their manufacture.

BACKGROUND

Programmed cell death plays a critical role in regulating cell number and in
10 eliminating stressed or damaged cells from normal tissues. Indeed, the network of
apoptotic signaling mechanisms inherent in most cell types provides a major barrier to
the development and progression of human cancer. Since most commonly used
radiation and chemo-therapies rely on activation of apoptotic pathways to kill cancer
cells, tumor cells which are capable of evading programmed cell death often become
15 resistant to treatment.

Apoptosis signaling networks are classified as extrinsic when mediated by death
receptor-ligand interactions or intrinsic when mediated by cellular stress and
mitochondrial permeabilization. Both pathways ultimately converge on individual
caspases, cysteine-aspartic proteases. Once activated, caspases cleave a number of
20 cell death-related substrates, effecting destruction of the cell.

Tumor cells have devised a number of strategies to circumvent apoptosis. One
recently reported molecular mechanism involves the overexpression of members of the
IAP (Inhibitor of Apoptosis Protein) family. IAPs sabotage apoptosis by directly
interacting with and neutralizing caspases. The prototype IAPs, XIAP and cIAP have
25 three functional domains referred to as BIR 1, 2 & 3 domains. The BIR3 domain
interacts directly with caspase 9 and inhibits its ability to bind and cleave its natural
substrate, procaspase 3.

A proapoptotic mitochondrial protein, Smac (also known as DIABLO), can
neutralize XIAP and/or cIAP by binding to a peptide binding pocket (Smac binding site)
30 on the surface of BIR3 thereby precluding interaction with caspase 9. Binding of
peptides derived from Smac has also been reported to trigger autocatalytic
polyubiquitination and subsequent proteasome-mediated degradation of cIAP1. The
present invention relates to therapeutic molecules that bind to the Smac binding pocket
thereby promoting apoptosis in rapidly dividing cells. Such therapeutic molecules are
35 useful for the treatment of proliferative diseases, including cancer.


SUMMARY

The present invention provides compounds of formula M-L-M' that have been
found to be effective in promoting apoptosis in rapidly dividing cells. Advantageously,

the compounds of the present invention are selectively more toxic to abnormal cells *e.g.* cells that are proliferating more rapidly than normal cells, particularly in human tumor or cancer cells. Accordingly, the compounds of the present invention are useful in the treatment of diseases and conditions characterized by cell proliferation.

5 In each of the embodiments below, M and M' are preferably both the same.

In one embodiment of the present invention, a compound of formula M-L-M' is provided where M and M' is each independently a monomeric moiety of Formula (I)



(I)

10 wherein,

R¹ is (C₁-C₄)alkyl, deuterated methyl, or hydrogen;

R² is (C₁-C₄)alkyl or hydrogen;

R³ is (C₁-C₄)alkyl or hydrogen, or

15 R¹ or R² along with the nitrogen to which R¹ or R² is attached is taken together with R³ to form an aziridinyl, azetidiny, pyrrolidinyl, or piperidinyl;

R⁴ is

(i) (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, (C₃-C₆)cycloalkyl, phenyl, a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each
20 independently selected from O, N or S, or

(ii) R^{4a}-(C₁-C₆)alkylene, where R^{4a} is (C₃-C₆)cycloalkyl, phenyl, a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each independently selected from O, N or S,

25 where said R⁴ and said R^{4a} are optionally substituted with 1 to 3 substituents selected from halo, hydroxyl, -SH, -CO₂H, (C₁-C₄)alkyl, halo-substituted(C₁-C₄)alkyl, (C₁-C₄)alkoxy, (C₁-C₄)alkyl-S-, -SO₂-, -NH₂ or -NO₂, and where 1 of the ring members of said cycloalkyl and said heterocycle moieties are optionally replaced with oxo or thione;

A, B, and D are CR⁵, and E is N,

30 A, B and E are CR⁵ and D is N,

A, D and E are CR⁵, and B is N,

B, D and E are CR⁵, and A is N,





A and B are both N, and D and E are both CR⁵,


A and E are both N, and B and D are both CR⁵, or

B and E are both N, and A and D are both CR⁵, where R⁵ are each independently

5 selected from H, F, -CH₃ or -CF₃;

V is selected from moieties of Formula (1a), (1b), (1c), (1d), (1e), (1f), (1g), (1h), (1i), (1j), or (1k)





where * indicates the point of attachment to L,

R^{6a} and R^{6b} are each independently H, (C₁-C₃)alkyl, Cl, or CN, where at least one of R^{6a} and R^{6b} is H or (C₁-C₃)alkyl;

W is a bond or (C₁-C₄)alkylene;

R^7 are each independently halo, -CH₃, -CF₃, -CN or -OCH₃;

p is 0, 1, 2, 3 or 4 for formula (1a) or (1e),

p is 0, 1, 2 or 3 for formula (1b), (1c), (1d), (1f), (1i) or (1j),

p is 0, 1 or 2 for formula (1g) or (1h);

L is -NR⁸-X¹-NR⁸-, where each R⁸ is independently selected from H, (C₁-C₄)alkyl, or halo-substituted(C₁-C₄)alkyl and X¹ is

(i) a bond,

(ii) (C₁-C₁₀)alkylene, (C₂-C₁₀)alkenylene, (C₂-C₁₀)alkynylene, ((C₁-C₁₀)alkylene)-(O(C₁-C₆)alkylene)_q-, or (C₁-C₁₀)alkylene-NH(C₁-C₆)alkylene, where q is 0, 1 or 2,

(iii) phenylene, naphthylene, fluorenylene, 9H-fluoren-9-onylene, 9,10-dihydroanthracenylenylene, anthracen-9,10-dionylene, a partially or fully saturated (C₃-C₈)cycloalkylene, a 5- to 7-membered heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S, or N, or a 5- to 10-membered heteroarylene containing 1 to 3 heteroatoms each independently selected from O, S or N, where said phenylene is optionally fused to a (C₅-C₆)cycloalkyl,

(iv) (phenylene)-G-(phenylene), where G is a bond, O, S, -NH-, -N=N-, -S=S-, -SO₂-, (C₁-C₆)alkylene, (C₂-C₆)alkenylene, (C₂-C₁₀)alkynylene, (C₃-C₆)cycloalkylene, a 5- to 6-membered heteroaryl containing 1 to 3 heteroatoms each independently selected from O, S or N, or a 5- to 6-membered partially or fully saturated heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S or N, and where said phenylene is optionally fused to a phenyl,

(v) ((C₁-C₆)alkylene)_r-Z¹-((C₁-C₆)alkylene)_s, or ((C₁-C₆)alkenylene)_r-Z¹-((C₁-C₆)alkenylene)_s, where r and s are each independently 0, 1, or 2; and Z¹ is -O-, -N=N-, (C₃-C₆)cycloalkylene, phenylene, bisphenylene, a 5- to 6-

5 membered partially or fully saturated heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S or N, or a 5- to 6-membered heteroarylene containing 1 to 3 heteroatoms each independently selected from O, S or N, where said heteroarylene and said heterocyclene are optionally fused to a phenyl, phenylene, a 5- to 6-membered partially or fully saturated heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S or N, or a 5- to 6-membered heteroarylene containing 1 to 3 heteroatoms each independently selected from O, S or N, or

10 (vi) (C₁-C₂₀)alkylene or -NH-((C₁-C₂₀)alkylene)-NH-, where said alkylene contains 1 to 6 oxygen atoms interspersed within the alkylene chain and optionally 1 to 2 phenylene groups interspersed within the alkylene chain; or X¹ is optionally taken together with one or both R⁸ groups along with the nitrogen to which the R⁸ group is attached to form a 4- to 14-membered
15 heterocyclene, (4- to 6-membered heterocyclyl)-(C₁-C₆)alkylene-(4- to 6-membered heterocyclyl), or *bis*-(4- to 6-membered heterocyclene, where said heterocyclene and said heterocyclyl moieties optionally contain 1 to 3 additional heteroatoms selected from O, S and N, and X¹ and R⁸ are optionally substituted with oxo or 1 to 3 substituents each independently
20 selected from hydroxyl or (C₁-C₄)alkyl;

where said group (ii) moieties of X¹ are each independently substituted with one or more fluoro atoms, or 1 to 2 substituents each independently selected from halo, oxo, amino, phenyl, naphthyl, (C₃-C₆) cycloalkyl, or 5- to 6-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, where said phenyl, said
25 cycloalkyl, and said heterocycle are optionally substituted with 1 to 3 substituents each independently selected from halo, (C₁-C₄)alkyl, or trifluoromethyl,

where said group (iii) and (iv) moieties of X¹ are optionally substituted with 1 to 4 substituents each independently selected from (C₁-C₄)alkyl, (C₁-C₄)alkoxy, halo, amino, -OH, benzyl, or a fused 5- to 6-membered cycloalkyl, where said (C₁-C₄)alkyl, said (C₁-
30 C₄)alkoxy, and said fused cycloalkyl are optionally substituted with 1 to 3 substituents selected from halo, or (C₁-C₄)alkyl,

where said group (v) moieties of X¹ are optionally substituted with 1 to 3 substituents each independently selected from halo, hydroxy, oxo, amino, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, or phenyl; or a pharmaceutically acceptable salt thereof.

35 In one particular embodiment,

R¹ is (C₁-C₄)alkyl, or deuterated methyl;

R² is hydrogen;

R^3 is (C_1-C_4) alkyl;

R^4 is

(i) (C_1-C_{10}) alkyl, (C_3-C_6) cycloalkyl, phenyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or





5 (ii) $R^{4a}-(C_1-C_6)$ alkylene, where R^{4a} is (C_3-C_6) cycloalkyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each independently selected from O, N or S,

10 where said R^4 and said R^{4a} are optionally substituted with 1 to 3 substituents selected from halo, or (C_1-C_4) alkoxy and where 1 of the ring members of said cycloalkyl and said heterocycle moieties are optionally replaced with oxo;



A, B, and D are CR^5 , and E is N, or A, B and E are CR^5 and D is N, where R^5 are each independently selected from H, or F;


V is selected from moieties of Formula (1a), (1b), (1e), (1f), (1g), (1i), (1j), or (1k)

15



20





where * indicates the point of attachment to L,

R^{6a} and R^{6b} are H;

5 W is a bond or (C_1-C_4) alkylene;

R^7 is halo;

p is 0 or 1 for formula (1a) or (1e),

p is 0 or 1 for formula (1b), (1f), (1i) or (1j),

p is 0 or 1 for formula (1g);

10 L is $-NR^8-X^1-NR^8-$, where each R^8 is independently selected from H, (C_1-C_4) alkyl, or halo-substituted (C_1-C_4) alkyl and X^1 is

(i) a bond,

(ii) (C_1-C_{10}) alkylene, (C_2-C_{10}) alkynylene, or $((C_1-C_{10})$ alkylene)- $(O(C_1-C_6)$ alkylene) $_q-$, where q is 0, 1 or 2,

15 (iii) phenylene, naphthylene, or a fully saturated (C_3-C_8) cycloalkylene,

(iv) (phenylene)-G-(phenylene), where G is a bond, O, $-SO_2-$, (C_1-C_6) alkylene,


(v) $((C_1-C_6)$ alkylene) $_r-Z^1-((C_1-C_6)$ alkylene) $_s$, where r and s are each independently 0, 1, or 2; and Z^1 is $-O-$, or

(vi) (C_1-C_{20}) alkylene or $-NH-((C_1-C_{20})$ alkylene)- $NH-$, where said alkylene contains 1 to 6 oxygen atoms interspersed within the alkylene chain;

20 or X^1 is optionally taken together with one or both R^8 groups along with the nitrogen to which the R^8 group is attached to form a 4- to 14-membered heterocyclene;

or a pharmaceutically acceptable salt thereof.


25 In another particular embodiment, M and M' are the same and each are a monomeric moiety of Formula (II)



(II)

or a pharmaceutically acceptable salt thereof.

In a preferred embodiment, R^1 is methyl or deuterated methyl; R^2 is H; R^3 is methyl; R^4 is isopropyl or cyclohexyl; R^{6a} is H; p is 0; W is a bond; L is $-\text{NH}-(\text{CH}_2)_3-(\text{O}-\text{CH}_2\text{CH}_2)_4-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_3-(\text{O}-\text{CH}_2\text{CH}_2)_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_3-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_3-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-(\text{CH}_2)_2-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_2-(\text{O}-\text{CH}_2\text{CH}_2)_2-\text{O}-(\text{CH}_2)_2-\text{NH}-$, $-\text{NH}-((\text{C}_1-\text{C}_{12})\text{alkylene})-\text{NH}-$, $-\text{NH}-\text{CH}_2-(\text{phenylene})-\text{CH}_2-\text{NH}-$, $-\text{NH}-\text{CH}_2-(\text{phenylene})-(\text{phenylene})-\text{CH}_2-\text{NH}-$, $-\text{NH}-(\text{cyclohexylene})-\text{NH}-$,



or a pharmaceutically acceptable salt thereof.

Representative compounds include: (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(4,7,10,13,16-pentaoxanonadecane-1,19-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))bis(propane-3,1-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(ethane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(biphenyl-4,4'-diylbis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(undecane-1,11-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

1-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(3-(3-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)-propoxy)propyl)-1H-indole-3-carboxamide;

(S,S,S)-N,N'-(butane-1,4-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(5-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)pentyl)-1H-indole-3-carboxamide;

(S,S,S)-N,N'-(hexane-1,6-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-((1S,4S)-4-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)cyclohexyl)-1H-indole-3-carboxamide;

(2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(2,6-diazaspiro[3.3]heptane-2,6-diylbis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide);

(2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(piperazine-1,4-diylbis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide); and

(S,S,S)-N,N'-(1,4-phenylene-bis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

or a pharmaceutically acceptable salt thereof.

One preferred compound is (S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)-pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

5 Another preferred compound is (S,S,S)-N,N'-(ethane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

Yet another preferred compound is 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(3-(3-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)propoxy)propyl)-1H-indole-3-carboxamide; or a pharmaceutically acceptable salt thereof.

Another preferred compound is (S,S,S)-N,N'-(butane-1,4-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

Yet another preferred compound is (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

Another preferred compound is (S,S,S)-N,N'-(4,7,10,13,16-pentaoxanonadecane-1,19-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

Yet another preferred compound is (S,S,S)-N,N'-(3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

Another preferred compound is (S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

Yet another preferred compound is (S,S,S)-N,N'-(hexane-1,6-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.


Another preferred compound is (2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(piperazine-1,4-diylbis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide); or a pharmaceutically acceptable salt thereof.

Yet another preferred compound is 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(5-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)pentyl)-1H-indole-3-carboxamide; or a pharmaceutically

5 acceptable salt thereof.


Another preferred compound is (S,S,S)-N,N'-(1,4-phenylene-bis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

10 Another particular embodiment of the above embodiments is where M and M' are the same and each are a monomeric moiety of Formula (III)



or a pharmaceutically acceptable salt thereof.

15 Preferably, R¹ is methyl or deuterated methyl; R² is H; R³ is methyl; R⁴ is isopropyl or cyclohexyl; R^{6a} and R^{6b} are both H; p is 0; W is a bond; and L is -NH-(CH₂)₃-(O-CH₂CH₂)₄-O-(CH₂)₃-NH-, -NH-(CH₂)₃-(O-CH₂CH₂)₂-O-(CH₂)₃-NH-, -NH-(CH₂)₃-O-CH₂CH₂-O-(CH₂)₃-NH-, -NH-(CH₂)₃-O-(CH₂)₃-NH-, -NH-(CH₂)₂-O-CH₂CH₂-O-(CH₂)₂-NH-, -NH-(CH₂)₂-(O-CH₂CH₂)₂-O-(CH₂)₂-NH-, -NH-((C₁-C₁₂)alkylene)-NH-,
 20 -NH-CH₂-(phenylene)-CH₂-NH-, -NH-CH₂-(phenylene)-(phenylene)-CH₂-NH-,
 -NH-(cyclohexylene)-NH-,



or a pharmaceutically acceptable salt thereof.

25 Preferably, L is -NH-(CH₂)₃-NH-, -NH-(CH₂)₃-O-(CH₂)₂-O-(CH₂)₃-NH-, -NH-(cyclohexylene)-NH-, or -NH-CH₂-(phenylene)-CH₂-NH-; or a pharmaceutically acceptable salt thereof.

Representative compounds include:

(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy)))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

5 (S,S,S)-N,N'-(1,4-phenylenebis-(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

(S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

10 (S,S,S)-N,N'-((1S,2S)-cyclohexane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); and

(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy)))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(trideuteromethylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

A preferred compound is (S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy)))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

Another preferred compound is (S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy)))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(trideuteromethylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

25 Yet another preferred compound is (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

In another aspect of the present invention, a pharmaceutical composition is provided which comprises any one of the compound described above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent or excipient. The pharmaceutical composition may further comprise at least one additional pharmaceutical agent (described herein below). In particular, the at least one additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist. More particularly, the at least one additional pharmaceutical agent is paclitaxel.

In yet another aspect of the present invention, a method for treating a disease, disorder, or condition associated with the over expression of an IAP in a subject is

provided which comprises the step of administering to a subject in need to such treatment a therapeutically effective amount of any one of the compounds described above, or a pharmaceutically acceptable salt thereof.

In yet another aspect, a method for treating a disease, disorder, or condition mediated by IAPs is provided which comprises the step of administering to a subject in need of such treatment a therapeutically effective amount of any one of the compounds described above, or a pharmaceutically acceptable salt thereof.

In yet another aspect, the use of any one of the compounds described above is provided for inducing or enhancing apoptosis in a tumor or cancer cell.

Any one of the compounds described above may be used in therapy.

Also described is the use of any one of the compounds described above in the manufacture of a medicament for the treatment of a disease, disorder or condition mediated by IAPs.

In another aspect, the use of any one of the compounds described above is provided for the treatment of a disease, disorder or condition associated with the overexpression of one or more IAPs.

In yet another aspect, a method for treating a disease, disorder, or condition mediated by IAPs is provided which comprises the step(s) of administering to a patient in need of such treatment

(i) a compound as defined above, or a pharmaceutically acceptable salt thereof; and

(ii) at least one additional pharmaceutical agent (described herein below).

In particular, the additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist.

More particularly, the additional pharmaceutical agent is paclitaxel.


The compound, or pharmaceutical acceptable salt thereof, and the additional pharmaceutical agent may be administered simultaneously or sequentially.

In yet another aspect, a method for treating a disease, disorder, or condition mediated by IAP is provided which comprises the step of administering to a patient in need of such treatment a pharmaceutical composition comprising any one of the compounds described above, or a pharmaceutically acceptable salt thereof, and a pharmaceutical acceptable carrier. The method composition may further comprise at least one additional pharmaceutical agent (described herein below). In particular, the additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist. More particularly, the additional pharmaceutical agent is paclitaxel.

In yet another aspect, a method for treating a disease, disorder, or condition mediated by IAPs is provided which comprises the step(s) of administering to a patient in need of such treatment

- (i) a first composition comprising any one of the compounds described above, or a pharmaceutically acceptable salt thereof, and a pharmaceutical carrier; and
- (ii) a second composition comprising at least one additional pharmaceutical agent and a pharmaceutical carrier. In particular, the additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist. More particularly, the additional pharmaceutical agent is a paclitaxel. The first composition and the second composition may be administered simultaneously or sequentially.

Another aspect of the invention are useful intermediates. For example, a compound having Formula (I-1d) or (I-1e)







wherein

- R is a carboxy-protecting group;
R¹ is (C₁-C₄)alkyl, or deuterated methyl;
R² is hydrogen or an amino-protecting group;
20 R³ is (C₁-C₄)alkyl;
R⁴ is
(i) (C₁-C₁₀)alkyl, (C₃-C₆)cycloalkyl, phenyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or
(ii) R^{4a}-(C₁-C₆)alkylene, where R^{4a} is (C₃-C₆)cycloalkyl, or a 3- to 7-membered
25 heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S,
or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each independently selected from O, N or S,

where said R⁴ and said R^{4a} are optionally substituted with 1 to 3 substituents selected from halo, or (C₁-C₄)alkoxy and where 1 of the ring members of said cycloalkyl and said heterocycle moieties are optionally replaced with oxo;

A, B, and D are CR⁵, and E is N, or A, B and E are CR⁵ and D is N, where R⁵ are each independently selected from H, or F;

V is selected from moieties of Formula (1a), (1b), (1e), (1f), (1g), (1i), (1j), or (1k)



10

where * indicates the point of attachment to -OH or OR,

R^{6a} and R^{6b} are H;

W is a bond or (C₁-C₄)alkylene;

R^7 is halo;

p is 0 or 1 for formula (1a) or (1e),

15

p is 0 or 1 for formula (1b), (1f), (1i) or (1j), and


p is 0 or 1 for formula (1g).

Preferably, R^1 is methyl or deuterated methyl; R^2 is an amino-protecting group; R^3 is methyl; R^4 is isopropyl or cyclohexyl; R^{6a} is H; p is 0; and W is a bond.

In one embodiment, A , B , and D are CR^5 , and E is N , where R^5 are each independently selected from H, or F.

- 5 In another embodiment, A , B and E are CR^5 and D is N , where R^5 are each independently selected from H, or F.

In a particular embodiment, V is a moiety of Formula (1a)




(1a).

- 10 A preferred intermediate is a compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid ethyl ester.

Another preferred intermediate is a compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid.

- 15

In another particular embodiment, V is a moiety of Formula (1f)



(1f).

- 20 A preferred intermediate is a compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-D₃-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester.

Another preferred intermediate is a compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-D₃-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid.

- 25 Yet another preferred intermediate is a compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester.

Another preferred intermediate is a compound which is 1-[5-((S)-1-((S)-2-((S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid.

Definitions

5 As used herein, the term "alkyl" refers to a hydrocarbon moiety of the general formula C_nH_{2n+1} . The alkane group may be straight or branched. For example, the term "(C₁-C₁₀)alkyl" refers to a monovalent, straight, or branched aliphatic group containing 1 to 10 carbon atoms (e.g., methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl, s-butyl, t-butyl, n-pentyl, 1-methylbutyl, 2-methylbutyl, 3-methylbutyl, neopentyl, 3,3-dimethylpropyl, 10 hexyl, 2-methylpentyl, heptyl, and the like). Similarly, the alkyl portion (i.e., alkyl moiety) of an alkoxy have the same definition as above. When indicated as being "optionally substituted", the alkane radical or alkyl moiety may be unsubstituted or substituted with one or more substituents (generally, one to three substituents except in the case of halogen substituents such as perchloro or perfluoroalkyls). "Halo-substituted alkyl" refers 15 to an alkyl group having at least one halogen substitution.

The term "alkenyl" refers to an alkyl moiety containing at least one unsaturation in the alkyl group. The alkenyl group may be straight or branched. For example, vinyl, prop-1-enyl, prop-2-enyl, allenyl, 2-methylprop-2-enyl, 3-methylbut-2-enyl, butadienyl, and the like.

20 The term "alkynyl" refers to an alkyl moiety containing at least one triple bond. The alkynyl group may be straight or branched. For example, $CH_3-C\equiv C-$, $H-C\equiv C-CH_2-$, $CH_3-C\equiv C-CH_2-$, $H-C\equiv C-CH(CH_3)-$, $H-C\equiv C-CH_2CH_2-$, $H-C\equiv C-CH(CH_3)CH_2-$, $H-C\equiv C-CH_2-C\equiv C-CH_2-$, and the like.

The term "alkylene" or "alkylenyl" refers to an alkyl moiety where the moiety 25 contains two binding sites. The alkylene group may be straight (e.g., $-(CH_2)-$, $-(CH_2)_2-$, $-(CH_2)_3-$, or branched (e.g., $-CH(CH_3)-$, $-C(CH_3)_2-$, $-CH_2CH(CH_3)-$, $-CH(CH_3)-CH_2-$, $-C(CH_3)_2-CH_2-$, etc.). Suitable alkylene moieties are the same as those described above for alkyl except with two binding sites instead of just one.

30 The term "alkenylene" or "alkenylenyl" refers to an alkenyl moiety containing two binding sites. For example, $-CH_2-CH=CH-CH_2-$, $-CH=CH-CH=CH-$, and the like. Suitable alkenylene moieties are the same as those described above for alkenyl except with two binding sites instead of just one.


The term "alkynylene" or "alkynylenyl" refers to an alkynyl moiety containing two 35 binding sites. For example, $-CH_2-C\equiv C-CH_2-$. Suitable alkynylene moieties are the same as those described above for alkynyl except with two binding sites instead of just one.

The term "aryl" refers to aromatic moieties having a single (e.g., phenyl) or a fused ring system (e.g., naphthalene, anthracene, phenanthrene, etc.). A typical aryl group is a 6- to 14-membered aromatic carbocyclic ring(s). A fused aromatic ring system may also include a phenyl fused to a partially or fully saturated cycloalkyl. For example,
 5 2,3-dihydroindenyl, 1,2,3,4-tetrahydronaphthalenyl, 1,2-dihydronaphthalenyl, 2,3-dihydronaphthalenyl, 9,10-dihydroanthracenyl, fluorenyl, and the like.


The term "arylene" refers to a carbocyclic aromatic moiety having two binding sites. Suitable arylenes include those groups described above for an aryl moiety except with two binding sites rather than one. For example, 1,2-phenylene, 1,3-phenylene, 1,4-phenylene, 1,3-naphthylene, 1,4-naphthylene, 1,5-naphthylene, 1,6-naphthylene, 1,7-naphthylene, 2,3-naphthylene, 2,4-naphthylene, 2,5-naphthylene, 2,6-naphthylene, 2,7-naphthylene, 3,4-naphthylene, 3,5-naphthylene, 3,6-naphthylene, 3,7-naphthylene, etc. The two binding sites on the fused arylene system may be on the same ring or different rings.

15 The term "partially or fully saturated cycloalkyl" refers to a carbocyclic ring which is fully hydrogenated (e.g., cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, etc.) or partially hydrogenated (e.g., cyclopropenyl, cyclobutenyl, cyclopentyl, cyclopenta-1,3-dienyl, cyclohexenyl, cyclohexa-1,3-dienyl, cyclohexa-1,4-dienyl, etc.). The carbocyclic ring may be a single ring (as described above), a bicyclic ring (e.g.,
 20 octahydropentalenyl, bicyclo[1.1.1]pentanyl, bicyclo[2.1.1]hexanyl, bicyclo[2.1.1]hex-2-enyl, bicyclo[2.2.1]hept-2-enyl, bicyclo[2.2.1]heptanyl, bicyclo[2.2.2]octanyl, bicyclo[2.2.2]oct-2-enyl, bicyclo[2.2.2]octa-2,5-dienyl, etc.) or a spiral ring (e.g., spiro[2.2]pentanyl, etc.), and the like.

The term "partially or fully saturated cycloalkylene" refers to a carbocyclic ring
 25 having either no unsaturation in the ring (fully hydrogenated) or at least one unsaturation (partially hydrogenated) without being aromatic and contains two binding sites. Suitable ring systems include those described above for a partially or fully saturated cycloalkyl except having two bind sites instead of one. For example, 1,2-cyclopropyl, 1,2-cycloprop-1-enyl, 1,2-cyclobutyl, 1,3-cyclobutyl, 1,2-cyclobut-1-enyl, 3,4-cyclobut-1-enyl,
 30 3,5-cyclopent-1-enyl, 1,4-cyclopenta-1,3-dienyl, 1,5-cyclopenta-1,3-dienyl, 1,2-cyclopenta-1,3-dienyl, 1,3-cyclopenta-1,3-dienyl, etc. The carbocyclic ring may be a single ring, a bicyclic ring, fused ring (e.g., decahydronaphthalene), or a spiral ring where the two binding sites on the bicyclic ring and spiral ring may be on the same ring or different rings. See, e.g., the illustration below.




Same ring



Different ring

The term “partially or fully saturated heterocycle” refers to a nonaromatic ring that is either partially or fully hydrogenated and may exist as a single ring, bicyclic ring
 5 (including fused rings) or a spiral ring. Unless specified otherwise, the heterocyclic ring is generally a 3- to 14-membered ring containing 1 to 3 heteroatoms (preferably 1 or 2 heteroatoms) independently selected from sulfur, oxygen and/or nitrogen. Partially saturated or fully saturated heterocyclic rings include groups such as epoxy, aziridinyl, azetidiny, tetrahydrofuranyl, dihydrofuranyl, dihydropyridinyl, pyrrolidinyl, imidazolidinyl,
 10 imidazoliny, 1H-dihydroimidazolyl, hexahydropyrimidinyl, piperidinyl, piperazinyl, pyrazolidinyl, 2H-pyranyl, 4H-pyranyl, 2H-chromenyl, oxazinyl, morpholino, thiomorpholino, tetrahydrothienyl, tetrahydrothienyl, 1,4,7-triazonane, diazepanyl, 1,1-dioxide, oxazolidinyl, thiazolidinyl, octahydropyrrolo[3,2-b]pyrrolyl, decahydro-2,7-naphthyridinyl, and the like. A partially saturated heterocyclic ring also includes groups
 15 wherein the heterocyclic ring is fused to an aryl or heteroaryl ring (e.g., 2,3-dihydrobenzofuranyl, indolinyl (or 2,3-dihydroindolyl), 2,3-dihydrobenzothiophenyl, 2,3-dihydrobenzothiazolyl, 1,2,3,4-tetrahydroquinolinyl, 1,2,3,4-tetrahydroisoquinolinyl, 5,6,7,8-tetrahydropyrido[3,4-b]pyrazinyl, and the like). Examples of spiral rings include 2,6-diazaspiro[3.3]heptanyl, 2,7-diazaspiro[4.4]nonanyl, 3-azaspiro[5.5]undecanyl, 3,9-
 20 diazaspiro[5.5]undecanyl, and the like.

The term “partially or fully saturated heterocyclene” refers to a partially or fully saturated heterocyclic ring (as described above) except having two binding sites instead of one. The heterocyclene ring may be a single ring, a bicyclic ring, or a spiral ring where the two binding sites on the bicyclic ring (including fused rings) and spiral ring may
 25 be on the same ring or different rings. See, e.g., the illustration below.



Same ring



Different ring

The term “heteroaryl” refers to aromatic moieties containing at least one heteroatom (e.g., oxygen, sulfur, nitrogen or combinations thereof) within a 5- to 10-

membered aromatic ring system (e.g., pyrrolyl, pyridyl, pyrazolyl, indolyl, indazolyl, thienyl, furanyl, benzofuranyl, oxazolyl, imidazolyl, tetrazolyl, triazinyl, pyrimidyl, pyrazinyl, thiazolyl, purinyl, benzimidazolyl, quinolinyl, isoquinolinyl, benzothiophenyl, benzoxazolyl, 1H-benzo[d][1,2,3]triazolyl, and the like.). The heteroaromatic moiety may
5 consist of a single or fused ring system. A typical single heteroaryl ring is a 5- to 6-membered ring containing one to three heteroatoms independently selected from oxygen, sulfur and nitrogen and a typical fused heteroaryl ring system is a 9- to 10-membered ring system containing one to four heteroatoms independently selected from oxygen, sulfur and nitrogen. The fused heteroaryl ring system may consist of two
10 heteroaryl rings fused together or a heteroaryl fused to an aryl (e.g., phenyl).

The term "heteroarylene" refers to a heteroaryl having two binding sites instead of one. Suitable heteroarylene groups include those described above for heteroaryl having two binding sites instead of one.

Unless specified otherwise, the term "compounds of the present invention" refers
15 to dimeric Compounds of Formula (M-L-M'), and salts thereof, as well as all stereoisomers (including diastereoisomers and enantiomers), rotamers, tautomers and isotopically labeled compounds (including deuterium substitutions), as well as inherently formed moieties (e.g., polymorphs, solvates and/or hydrates). For purposes of this invention, solvates and hydrates are generally considered compositions.

20

DETAILED DESCRIPTION

The present invention provides compounds and pharmaceutical formulations thereof that are useful in the treatment of diseases, conditions and/or disorders in which the inhibition of apoptosis contributes to disease pathogenesis.

25 Compounds of the present invention may be synthesized by synthetic routes that include processes analogous to those well-known in the chemical arts, particularly in light of the description contained herein. The starting materials are generally available from commercial sources such as Aldrich Chemicals (Milwaukee, Wis.) or are readily prepared using methods well known to those skilled in the art (e.g., prepared by methods
30 generally described in Louis F. Fieser and Mary Fieser, Reagents for Organic Synthesis, v. 1-19, Wiley, New York (1967-1999 ed.), or Beilsteins Handbuch der organischen Chemie, 4, Aufl. ed. Springer-Verlag, Berlin, including supplements (also available via the Beilstein online database)).

For illustrative purposes, the reaction schemes depicted below provide potential
35 routes for synthesizing the compounds of the present invention as well as key intermediates. For a more detailed description of the individual reaction steps, see the Examples section below. Those skilled in the art will appreciate that other synthetic

routes may be used to synthesize the inventive compounds. Although specific starting materials and reagents are depicted in the schemes and discussed below, other starting materials and reagents can be easily substituted to provide a variety of derivatives and/or reaction conditions. In addition, many of the compounds prepared by the methods

5 described below can be further modified in light of this disclosure using conventional chemistry well known to those skilled in the art.


In the preparation of compounds of the present invention, protection of remote functionality (e.g., primary or secondary amino, or carboxyl groups) of intermediates may be necessary. The need for such protection will vary depending on the nature of the

10 remote functionality and the conditions of the preparation methods. Suitable amino-protecting groups (NH-Pg) include acetyl, trifluoroacetyl, t-butoxycarbonyl (BOC), benzyloxycarbonyl (CBz) and 9-fluorenylmethylenoxycarbonyl (Fmoc). Suitable carboxyl protecting groups (C(O)O-Pg) include alkyl esters (e.g., methyl, ethyl or t-butyl), benzyl esters, silyl esters, and the like. The need for such protection is readily

15 determined by one skilled in the art. For a general description of protecting groups and their use, see T. W. Greene, Protective Groups in Organic Synthesis, John Wiley & Sons, New York, 1991.

Scheme 1 (below) describes a potential route for producing dimeric compounds of Formula M-L-M', wherein M and M' are a monomeric moiety of Formula (I). The

20 monomeric units may be the same or different; however, generally they are the same.



Scheme I

The desired V group having a protected carboxyl group (e.g., ester) is coupled to the starting material (SM-1), where the pyrrolidine nitrogen is protected (e.g., p-methoxyphenethyl (PMPE)), using standard transition metal catalyzed (e.g., copper-catalyzed) coupling of aryl halides well-known to those of skill in the art. Suitable V-OR compounds for Formula (1a) which are commercially available or readily prepared from literature preparations include: indole-3-acetic acid methyl ester, methyl 5,6-dichloro-3-indoleacetate, 2-methyl-5-fluoro-1H-indol-3-yl-acetic acid methyl ester, methyl (5-chloro-

2-methyl-1H-indol-3-yl)acetate, ethyl 2-(5-cyano-2-methyl-1H-indol-3-yl)acetate, 2-methyl-1H-indole-3-acetic acid methyl ester, ethyl 2-(5-methoxy-1H-indol-3-yl)acetate, ethyl 2-(2,5-dimethyl-1H-indol-3-yl)acetate, (4-fluoro-1H-indol-3-yl)-acetic acid ethyl ester, 3-(1H-indol-3-yl)-propionic acid methyl ester, 4-(1H-indol-3-yl)-butanoic acid methyl ester (see, e.g., Jackson, R.; Manske, R. *J. Am. Chem. Soc.* **1930**, 52 (12), 5029-5035), 5-methoxy-2-methyl-1H-indole-3-carboxylic acid ethyl ester, methyl indole-3-carboxylate, 2-methyl-1H-indole-3-carboxylic acid methyl ester, 5-methylindole-3-carboxylic acid methyl ester, 4,5-difluoro-2-methylindole-3-carboxylic acid ethyl ester, 4-fluoro-2-methylindole-3-carboxylic acid ethyl ester, 6-fluoro-2-methylindole-3-carboxylic acid ethyl ester, 5-fluoro-2-methylindole-3-carboxylic acid ethyl ester, 5-methoxy-1H-indole-3-carboxylic acid methyl ester, 5-chloro-1H-indole-3-carboxylic acid methyl ester, 5-fluoro-1H-indole-3-carboxylic acid methyl ester, 6-fluoro-1H-indole-3-carboxylic acid methyl ester, 2-chloro-1H-indole-3-carboxylic acid methyl ester, 4,5,6,7-tetrafluoro-1H-indole-3-carboxylic acid methyl ester, 6-methyl-1H-indole-3-carboxylic acid methyl ester, 7-methyl-1H-indole-3-carboxylic acid methyl ester, 6-chloro-1H-indole-3-carboxylic acid methyl ester, 6-cyano-1H-indole-3-carboxylic acid methyl ester, 6-methoxy-1H-indole-3-carboxylic acid methyl ester, 4-methoxyindole-3-carboxylic acid methyl ester, 7-cyano-1H-indole-3-carboxylic acid methyl ester, 7-methoxyindole-3-carboxylic acid methyl ester, 4-methyl-1H-indole-3-carboxylic acid methyl ester, 4-fluoro-1H-indole-3-carboxylic acid methyl ester, 4-chloro-1H-indole-3-carboxylic acid methyl ester, 4-cyano-1H-indole-3-carboxylic acid methyl ester, 5-cyano-1H-indole-3-carboxylic acid methyl ester, 7-fluoro-1H-indole-3-carboxylic acid methyl ester, 7-chloro-1H-indole-3-carboxylic acid methyl ester, 5,7-dichloro-1H-indole-3-carboxylic acid ethyl ester, 7-methoxy-2-methyl-1H-indole-3-carboxylic acid methyl ester, 6-(trifluoromethyl)-1H-indole-3-carboxylic acid methyl ester, 7-(trifluoromethyl)-1H-indole-3-carboxylic acid methyl ester, 2-ethyl-1H-indole-3-carboxylic acid methyl ester, and 7-methoxy-1H-indole-3-carboxylic acid methyl ester (see, e.g., Prashad, M.; La Vecchia, L.; Prasad, K.; Repic, O., *Syn. Comm.* **1995**, 25 (1), 95-100)).

Suitable V-OR compounds for Formula (1b) which are commercially available or readily prepared from literature preparations include: methyl 6-(trifluoromethyl)-1H-pyrrolo[2,3-b]pyridine-3-carboxylate, methyl 1H-pyrrolo[2,3-b]pyridine-3-carboxylate, 5-(trifluoromethyl)-7-azaindole-3-carboxylic acid methyl ester, 5-methyl-7-azaindole-3-carboxylic acid methyl ester, 5-fluoro-7-azaindole-3-carboxylic acid methyl ester, 5-chloro-7-azaindole-3-carboxylic acid methyl ester, ethyl 4-chloro-1H-pyrrolo[2,3-b]pyridine-3-carboxylate, 1H-pyrrolo[2,3-b]pyridine-3-acetic acid methyl ester, and (4-chloro-1H-pyrrolo[2,3-b]pyridin-3-yl)-acetic acid methyl ester.

Suitable V-OR compounds for Formula (1c) which are commercially available or readily prepared from literature preparations include: 4-fluoro-6-azaindole-3-carboxylic acid methyl ester and 1H-pyrrolo[2,3-c]pyridine-3-carboxylic acid methyl ester (see, e.g., Moeller, K.; Sus, O., *Liebigs Ann. Der. Chem.*, **1956**, 599 (3), 233-236).

- 5 Suitable V-OR compounds for Formula (1d) which are commercially available or readily prepared from literature preparations include: methyl 1H-pyrrolo[3,2-b]pyridine-3-carboxylate, methyl 6-cyano-1H-pyrrolo[3,2-b]pyridine-3-carboxylate, methyl 6-chloro-1H-pyrrolo[3,2-b]pyridine-3-carboxylate, methyl 5-(trifluoromethyl)-1H-pyrrolo[3,2-b]pyridine-3-carboxylate, 1H-pyrrolo[3,2-b]pyridine-3-carboxylic acid ethyl ester, 5-methoxy-1H-pyrrolo[3,2-b]pyridine-3-carboxylic acid ethyl ester, ethyl 2-(1H-pyrrolo[3,2-b]pyridin-3-yl)acetate and ethyl 2-(5-methoxy-1H-pyrrolo[3,2-b]pyridin-3-yl)acetate.

Suitable V-OR compounds for Formula (1e) which are commercially available or readily prepared from literature preparations include: ethyl 2-methyl-2,3-dihydro-indole-3-carboxylate and 2,3-dihydro-1H-indole-3-carboxylic acid methyl ester.

- 15 Suitable V-OR compounds for Formula (1f) which are commercially available or readily prepared from literature preparations include: ethyl indole-4-acetate, 7-chloro-4-indole carboxylic acid methyl ester, methyl indole-4-carboxylate, methyl 3-cyanoindole-4-carboxylate, 6-fluoro indole-4-methylcarboxylate, and 6-chloro indole-4-methylcarboxylate.

- 20 Suitable V-OR compounds for Formula (1g) which are commercially available or readily prepared from literature preparations include: methyl 1H-pyrrolo[2,3-b]pyridine-4-carboxylate, methyl 5-chloro-1H-pyrrolo[2,3-b]pyridine-4-carboxylate, 3-chloro-7-azaindole-4-carboxylic acid methyl ester, 6-chloro-7-azaindole-4-carboxylic acid methyl ester, 5-fluoro-7-azaindole-4-carboxylic acid methyl ester, 6-fluoro-7-azaindole-4-carboxylic acid methyl ester, and tert-butyl 2-(1H-pyrrolo[2,3-b]pyridin-4-yl)acetate.

A suitable V-OR compound for Formula (1h) which is commercially available or readily prepared from literature preparations includes methyl 1H-pyrrolo[3,2-c]pyridine-4-carboxylate.

- 30 A suitable V-OR compound for Formula (1i) which is commercially available or readily prepared from literature preparations includes 1H-benzimidazole-4-carboxylic acid methyl ester.

A suitable V-OR compound for Formula (1j) which is commercially available or readily prepared from literature preparations includes 2,3-dihydro-1H-indole-4-carboxylic acid methyl ester.

- 35 A suitable V-OR compound for Formula (1k) which is commercially available or readily prepared from literature preparations includes methyl 1H-benzo[d][1,2,3]triazole-4-carboxylate.

One of skill in the art would know how to make further modifications of these compounds using conventional chemistry.


The pyrrolidine nitrogen of Intermediate (I-1a) can then be de-protected to produce Intermediate (I-1b) using procedures suitable for the particular protecting group used. For example, a PMPE protecting group can be removed by treating with trifluoroacetic acid at elevated temperatures. Once deprotected, the pyrrolidine nitrogen can then be coupled with the desired amino-protected amino acid derivative (HO-C(O)-C(CR⁴)-NH-Pg, such as Boc-L-valine, 2-(Boc-amino)-2-cyclohexylacetic acid, 2-(Boc-amino)-2-morpholinoacetic acid, 2-(Boc-amino)-2-*tert*-butylacetic acid, 2-(Boc-amino)-2-(tetrahydro-2H-pyran-4-yl) acetic acid, 2-(Boc-amino)-2-phenyl acetic acid, 2-(Boc-amino)-2-(4-hydroxycyclohexyl) acetic acid, 2-(Boc-amino)-3-methylpentanoic acid, 2-(Boc-amino)-3-hydroxy-3-methylbutanoic acid, and 2-(Boc-amino)-2-(4,4-difluorocyclohexyl)acetic acid) using standard peptide coupling conditions. Common activating agents for the coupling reaction include carbodiimides (e.g., dicyclohexylcarbodiimide (DCC), 1-Ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC) and diisopropylcarbodiimide (DIC)), triazoles (e.g., 1-hydroxy-benzotriazole (HOBt), 1-hydroxy-7-aza-benzotriazole (HOAt)), (2-(7-Aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate) (HATU), and O-Benzotriazole-N,N',N'-tetramethyl-uronium-hexafluoro-phosphate (HBTU), triazines (e.g., 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM)). Once the coupling is complete, then the amino acid protecting group can be removed so that a second amino acid (e.g., HO-C(O)-C(R³)-N(R¹)R², when R² is H, then an amino-protecting group, such as Boc, is used) can be added using the same or different standard peptide coupling conditions.

Alternatively, the two amino acid derivatives can be coupled prior to condensing onto the deprotected pyrrolidine nitrogen of Intermediate (I-1b).

The carboxylic acid protecting group on the "V" moiety is then removed using procedures appropriate for the particular protecting group utilized to form the corresponding carboxylic acid. Once the carboxylic acid protecting group is removed using standard hydrolysis conditions (e.g., lithium hydroxide in methanol and water), then two monomeric units can be linked using a desired diamino linker to produce the dimeric compound (M-L-M'). The two unprotected monomeric carboxylic acids coupled to the diamino compound can be the same or different. When a protecting group is used in the R² position, then the amino-protecting group may be removed using conditions commensurate with the particular amino-protecting group used to provide compounds of Formula (M-L-M') where R² is H in one or both monomeric units.

Suitable diamino linker compounds ($H-NR^8-X^1-NR^8-H$) which are commercially available or readily prepared from literature preparations include 2,6-diazaspiro[3.3]heptane; 2,2-dimethylpropane-1,3-diamine; 4,7,10,13,16-pentaoxanonadecane-1,19-diamine; 3,3'-oxydipropan-1-amine; 2,2'-(ethane-1,2-diylbis(oxy))diethanamine; 3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))dipropan-1-amine; 2,2'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))diethanamine; 3,3'-(ethane-1,2-diylbis(oxy))dipropan-1-amine; propane-1,3-diamine; butane-1,4-diamine; 4-[2-(4-aminophenyl)ethynyl]aniline; 1,4-bis(3-aminophenyl)butadiyne; 1,4-diamino-2-butyne; hex-3-yne-2,5-diamine; hexa-2,4-diyne-1,6-diamine (see, e.g., Jeon, J. H.; Sayre, L. M., *Biochem. Biophys. Res. Commun.* **2003**, 304(4), 788-794); N^1, N^4 -diethylbut-2-yne-1,4-diamine; (*E*)- N^1, N^4 -diethylbut-2-ene-1,4-diamine; *cis*-octahydro-pyrrolo[3,4-*c*]pyridine; 1,1'-ethylenedipiperazine; 1,5-diethyl-3,7-diaza-bicyclo[3.3.1]nonan-9-one; 1-ethyl-5-methyl-3,7-diaza-bicyclo[3.3.1]nonan-9-ol; 1-ethyl-5-methyl-3,7-diaza-bicyclo[3.3.1]nonan-9-one; 4,10-diaza-12-crown-4-ether; 1,5,9-triazacyclododecane; 1,5-dimethyl-3,7-diaza-bicyclo[3.3.1]nonan-9-ol; 4,4-bipiperidine; 1,5-dimethyl-3,7-diaza-bicyclo[3.3.1]nonan-9-one; 1,5-dimethyl-3,7-diazabicyclo[3.3.1]nonane; 2,8-diazaspiro[5,5]undecane; decahydro-2,7-naphthyridine; 1,4,7-triazacyclononane; 6,6-dimethyl-1,4-diazepane; (*S*)-2,7-diazaspiro[4.4]nonane; *cis*-octahydro-pyrrolo[3,4-*c*]pyridine; 1,5-diazacyclooctane; 6-methyl-[1,4]diazepane; 3,7-diazabicyclo[3.3.0]octane; homopiperazine; 2,6-diazaspiro[3.3]heptane; piperazine; (3a*S*,7a*R*)-octahydro-pyrrolo[2,3-*c*]p; (3a*R*,7a*S*)-octahydro-pyrrolo[2,3-*c*]p; 1-(furan-2-yl)-*N*-(piperidin-4-ylmethyl)methanamine; 2,2,2-trifluoro-*N*-(pyrrolidin-3-ylmethyl)ethanamine; *N*-((morpholin-2-yl)methyl)ethanamine; methyl-morpholin-2-ylmethyl-amine; methyl-piperidin-4-ylmethyl-amine; ethyl-pyrrolidin-3-ylmethyl-amine; methyl-pyrrolidin-3-ylmethyl-amine; *N*-methyl-3-azetidinemethanamine; and (2,3-dihydro-1*H*-pyrrolo[2,3-*b*]pyridin-5-yl)methanamine. Those of skill in the art will know how to make modifications to the literature preparations or commercial compounds to make additional derivatives.

Scheme 2 below provides an alternative route for making the dimeric compounds (M-L-M').



Scheme 2

Intermediate (I-1a) can alternatively be linked prior to condensing the diamino acid sidechain to the monomeric units. The carboxylic acid protecting group on the "V" moiety is removed using procedures appropriate for the particular protecting group utilized to form the corresponding carboxylic acid ($R=H$). Once the carboxylic acid protecting group is removed using standard hydrolysis conditions (e.g., lithium hydroxide in methanol and water), then two monomeric units (I-1a) can be linked using a desired diamino linker to produce the dimeric intermediate (I-2a). The two unprotected monomeric carboxylic acids coupled to the diamino compound can be the same or different. The pyrrolidine nitrogens of Intermediate (I-2a) can then be de-protected to produce Intermediate using procedures suitable for the particular protecting group used. Once deprotected, the pyrrolidine nitrogens can then be coupled with the desired amino-protected amino acid derivative (followed by condensation with a second amino acid derivative) or diamino acid derivative using the procedures described above in Scheme I to produce the dimer compound M-L-M'.

The dimeric compounds may be isolated and used as the compound *per se* or as its salt. As used herein, the terms "salt" or "salts" refers to an acid addition or base addition salt of a compound of the invention. "Salts" include in particular "pharmaceutical

acceptable salts". The term "pharmaceutically acceptable salts" refers to salts that retain the biological effectiveness and properties of the compounds of this invention and, which typically are not biologically or otherwise undesirable. In many cases, the compounds of the present invention are capable of forming acid and/or base salts by virtue of the

5 presence of amino and/or carboxyl groups or groups similar thereto.

Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids, e.g., acetate, aspartate, benzoate, besylate, bromide/hydrobromide, bicarbonate/carbonate, bisulfate/sulfate, camphorsulfonate, chloride/hydrochloride, chlortheophyllonate, citrate, ethandisulfonate, fumarate, gluceptate, gluconate, glucuronate, hippurate, hydroiodide/iodide, isethionate, lactate, lactobionate, laurylsulfate, malate, maleate, malonate, mandelate, mesylate, methylsulphate, naphthoate, napsylate, nicotinate, nitrate, octadecanoate, oleate, oxalate, palmitate, pamoate, phosphate/hydrogen phosphate/dihydrogen phosphate, polygalacturonate, propionate, stearate, succinate, sulfosalicylate, tartrate, tosylate and trifluoroacetate salts.

Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like.

Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, toluenesulfonic acid, sulfosalicylic acid, and the like.

Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases.

25 Inorganic bases from which salts can be derived include, for example, ammonium salts and metals from columns I to XII of the periodic table. In certain embodiments, the salts are derived from sodium, potassium, ammonium, calcium, magnesium, iron, silver, zinc, and copper; particularly suitable salts include ammonium, potassium, sodium, calcium and magnesium salts.

30 Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like. Certain organic amines include isopropylamine, benzathine, choline, diethanolamine, diethylamine, lysine, meglumine, piperazine and tromethamine.

35 The pharmaceutically acceptable salts of the present invention can be synthesized from a parent compound, a basic or acidic moiety, by conventional chemical methods. Generally, such salts can be prepared by reacting free acid forms of these

compounds with a stoichiometric amount of the appropriate base (such as Na, Ca, Mg, or K hydroxide, carbonate, bicarbonate or the like), or by reacting free base forms of these compounds with a stoichiometric amount of the appropriate acid. Such reactions are typically carried out in water or in an organic solvent, or in a mixture of the two.

- 5 Generally, use of non-aqueous media like ether, ethyl acetate, ethanol, isopropanol, or acetonitrile is desirable, where practicable. Lists of additional suitable salts can be found, e.g., in "Remington's Pharmaceutical Sciences", 20th ed., Mack Publishing Company, Easton, Pa., (1985); and in "Handbook of Pharmaceutical Salts: Properties, Selection, and Use" by Stahl and Wermuth (Wiley-VCH, Weinheim, Germany, 2002).

- 10 Any formula given herein is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. Isotopically labeled compounds have structures depicted by the formulas given herein except that one or more atoms are replaced by an atom having a selected atomic mass or mass number. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of
- 15 hydrogen, carbon, nitrogen, oxygen, phosphorous, fluorine, and chlorine, such as ^2H , ^3H , ^{11}C , ^{13}C , ^{14}C , ^{15}N , ^{18}F , ^{31}P , ^{32}P , ^{35}S , ^{36}Cl , ^{125}I respectively. The invention includes various isotopically labeled compounds as defined herein, for example those into which radioactive isotopes, such as ^3H , ^{13}C , and ^{14}C , are present. Such isotopically labelled compounds are useful in metabolic studies (with ^{14}C), reaction kinetic studies (with, for
- 20 example ^2H or ^3H), detection or imaging techniques, such as positron emission tomography (PET) or single-photon emission computed tomography (SPECT) including drug or substrate tissue distribution assays, or in radioactive treatment of patients. In particular, an ^{18}F or labeled compound may be particularly desirable for PET or SPECT studies. Isotopically labeled compounds of this invention can generally be prepared by
- 25 carrying out the procedures disclosed in the schemes or in the examples and preparations described below by substituting a readily available isotopically labeled reagent for a non-isotopically labeled reagent.

- Further, substitution with heavier isotopes, particularly deuterium (i.e., ^2H or D) may afford certain therapeutic advantages resulting from greater metabolic stability, for
- 30 example increased in vivo half-life, reduced dosage requirements, reduced cyp inhibition (competitive or time dependent) or an improvement in therapeutic index. For example, substitution with deuterium may modulate undesirable side effects of the undeuterated compound, such as competitive cyp inhibition, time dependent cyp inactivation, etc. It is understood that deuterium in this context is regarded as a substituent in a compound of
- 35 the present invention (including both the monomeric and linker moieties of the dimer). The concentration of such a heavier isotope, specifically deuterium, may be defined by the isotopic enrichment factor. The term "isotopic enrichment factor" as used herein

means the ratio between the isotopic abundance and the natural abundance of a specified isotope. If a substituent in a compound of this invention is denoted deuterium (e.g., deuterated methyl or $-CD_3$), such compound has an isotopic enrichment factor for each designated deuterium atom of at least 3500 (52.5% deuterium incorporation at
5 each designated deuterium atom), at least 4000 (60% deuterium incorporation), at least 4500 (67.5% deuterium incorporation), at least 5000 (75% deuterium incorporation), at least 5500 (82.5% deuterium incorporation), at least 6000 (90% deuterium incorporation), at least 6333.3 (95% deuterium incorporation), at least 6466.7 (97% deuterium incorporation), at least 6600 (99% deuterium incorporation), or at least 6633.3
10 (99.5% deuterium incorporation).

Isotopically-labeled compounds of present invention can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Examples and Preparations using an appropriate isotopically-labeled reagents in place of the non-labeled reagent previously
15 employed.

Pharmaceutically acceptable solvates in accordance with the invention include those wherein the solvent of crystallization may be isotopically substituted, e.g. D_2O , d_6 -acetone, d_6 -DMSO.

It will be recognized by those skilled in the art that the compounds of the present
20 invention may contain chiral centers and as such may exist in different isomeric forms. As used herein, the term "isomers" refers to different compounds that have the same molecular formula but differ in arrangement and configuration of the atoms. Also as used herein, the term "an optical isomer" or "a stereoisomer" refers to any of the various stereo isomeric configurations which may exist for a given compound of the present invention
25 and includes geometric isomers. It is understood that a substituent may be attached at a chiral center of a carbon atom. Therefore, the invention includes enantiomers, diastereomers or racemates of the compound.

"Enantiomers" are a pair of stereoisomers that are non- superimposable mirror images of each other. A 1:1 mixture of a pair of enantiomers is a "racemic" mixture. The
30 term is used to designate a racemic mixture where appropriate.

"Diastereoisomers" are stereoisomers that have at least two asymmetric atoms, but which are not mirror-images of each other. The absolute stereochemistry is specified according to the Cahn- Ingold- Prelog R-S system. When a compound is a pure enantiomer the stereochemistry at each chiral carbon may be specified by either *R* or *S*.
35 Resolved compounds whose absolute configuration is unknown can be designated (+) or (-) depending on the direction (dextro- or levorotatory) which they rotate plane polarized light at the wavelength of the sodium D line. Certain of the compounds described herein

contain one or more asymmetric centers or axes and may thus give rise to enantiomers, diastereomers, and other stereoisomeric forms that may be defined, in terms of absolute stereochemistry, as (*R*)- or (*S*)-.

Unless specified otherwise, the compounds of the present invention are meant to
5 include all such possible isomers, including racemic mixtures, optically pure forms and intermediate mixtures. Optically active (*R*)- and (*S*)- isomers may be prepared using chiral synthons or chiral reagents, or resolved using conventional techniques. If the compound contains a double bond, the substituent may be *E* or *Z* configuration. If the compound contains a disubstituted cycloalkyl, the cycloalkyl substituent may have a *cis*-
10 or *trans*-configuration. All tautomeric forms are also intended to be included.

Compounds of the invention that contain groups capable of acting as donors and/or acceptors for hydrogen bonds may be capable of forming co-crystals with suitable co-crystal formers. These co-crystals may be prepared from compounds of the present invention by known co-crystal forming procedures. Such procedures include grinding,
15 heating, co-subliming, co-melting, or contacting in solution compounds of the present invention with the co-crystal former under crystallization conditions and isolating co-crystals thereby formed. Suitable co-crystal formers include those described in WO 2004/078163. Hence the invention further provides co-crystals comprising a compound of the present invention.

20 Compounds of the present invention have been found to induce or enhance apoptosis and therefore useful in the treatment of cancer. Consequently, a compound of the present invention may be used in the manufacture of a medicament for the treatment of diseases, conditions or disorders associated with the overexpression of an IAP in a subject (or mammal, preferably a human), inducing apoptosis in a tumor or cancer cell,
25 inhibiting the binding of an IAP protein to a caspase protein, or sensitizing a tumor or cancer cell to an apoptotic signal. In the process, a compound of the present invention may also induce the degradation of individual or multiple IAPs in cells (specifically cIAP1, cIAP2 and/or XIAP), and may induce expression of TNF α in some cells.

The compounds of the present invention are typically used as a pharmaceutical
30 composition (e.g., a compound of the present invention and at least one pharmaceutically acceptable carrier). As used herein, the term "pharmaceutically acceptable carrier" includes generally recognized as safe (GRAS) solvents, dispersion media, surfactants, antioxidants, preservatives (e.g., antibacterial agents, antifungal agents), isotonic agents, salts, preservatives, drug stabilizers, buffering agents (e.g.,
35 maleic acid, tartaric acid, lactic acid, citric acid, acetic acid, sodium bicarbonate, sodium phosphate, and the like), and the like and combinations thereof, as would be known to those skilled in the art (see, for example, Remington's Pharmaceutical Sciences, 18th

Ed. Mack Printing Company, 1990, pp. 1289- 1329). Except insofar as any conventional carrier is incompatible with the active ingredient, its use in the therapeutic or pharmaceutical compositions is contemplated. For purposes of this invention, solvates and hydrates are considered pharmaceutical compositions comprising a compound of
5 the present invention and a solvent (i.e., solvate) or water (i.e., hydrate).

The formulations may be prepared using conventional dissolution and mixing procedures. For example, the bulk drug substance (i.e., compound of the present invention or stabilized form of the compound (e.g., complex with a cyclodextrin derivative or other known complexation agent)) is dissolved in a suitable solvent in the presence of
10 one or more of the excipients described above. The compound of the present invention is typically formulated into pharmaceutical dosage forms to provide an easily controllable dosage of the drug and to give the patient an elegant and easily handleable product.

The pharmaceutical composition (or formulation) for application may be packaged in a variety of ways depending upon the method used for administering the drug.
15 Generally, an article for distribution includes a container having deposited therein the pharmaceutical formulation in an appropriate form. Suitable containers are well-known to those skilled in the art and include materials such as bottles (plastic and glass), ampoules, plastic bags, metal cylinders, and the like. The container may also include a tamper-proof assemblage to prevent indiscreet access to the contents of the package. In
20 addition, the container has deposited thereon a label that describes the contents of the container. The label may also include appropriate warnings.

The pharmaceutical composition comprising a therapeutically effective amount of a compound of the present invention is generally formulated for use as a parenteral administration. The pharmaceutical compositions (e.g., intravenous (iv) formulation) can
25 be subjected to conventional pharmaceutical operations such as sterilization and/or can contain conventional inert diluents, or buffering agents, as well as adjuvants, such as preservatives, stabilizers, wetting agents, emulsifiers and buffers well known to those of skill in the art.

In certain instances, it may be advantageous to administer the compound of the
30 present invention in combination with at least one additional pharmaceutical (or therapeutic) agent (e.g., an anti-cancer agent or adjunct therapy typically used in chemotherapy). The compound of the present invention may be administered either simultaneously with, or before or after, one or more other therapeutic agent(s). Alternatively, the compound of the present invention may be administered separately, by
35 the same or different route of administration, or together in the same pharmaceutical composition as the other agent(s).

Suitable additional anti-cancer agents include

(i) Taxane anti-neoplastic agents such as Cabazitaxel (1-hydroxy-7 β ,10 β -dimethoxy-9-oxo-5 β ,20-epoxytax-11-ene-2 α ,4,13 α -triyl-4-acetate-2-benzoate-13-[(2R,3S)-3-[[*tert*-butoxy]carbonyl]amino]-2-hydroxy-3-phenylpropanoate), larotaxel ((2 α ,3 ξ ,4 α ,5 β ,7 α ,10 β ,13 α)-4,10-bis(acetyloxy)-13-((2R,3S)-3-[[*tert*-butoxycarbonyl]amino]-2-hydroxy-3-phenylpropanoyl)oxy)-1-hydroxy-9-oxo-5,20-epoxy-7,19-cyclotax-11-en-2-yl benzoate) and paclitaxel;

(ii) Vascular Endothelial Growth Factor (VEGF) receptor inhibitors and antibodies such as Bevacizumab (sold under the trademark Avastin® by Genentech/Roche), axitinib, (N-methyl-2-[[3-[(E)-2-pyridin-2-ylethenyl]-1H-indazol-6-yl]sulfanyl]benzamide, also known as AG013736, and described in PCT Publication No. WO 01/002369), Brivanib Alaninate ((S)-((R)-1-(4-(4-Fluoro-2-methyl-1H-indol-5-yloxy)-5-methylpyrrolo[2,1-f][1,2,4]triazin-6-yloxy)propan-2-yl)2-aminopropanoate, also known as BMS-582664), motesanib (N-(2,3-dihydro-3,3-dimethyl-1H-indol-6-yl)-2-[(4-pyridinylmethyl)amino]-3-pyridinecarboxamide, and described in PCT Publication No. WO 02/066470), pasireotide (also known as SOM230, and described in PCT Publication No. WO 02/010192), and sorafenib (sold under the tradename Nexavar®);

(iii) Tyrosine kinase inhibitors such as Erlotinib hydrochloride (sold under the trademark Tarceva® by Genentech/Roche), Linifanib (N-[4-(3-amino-1H-indazol-4-yl)phenyl]-N'-(2-fluoro-5-methylphenyl)urea, also known as ABT 869, available from Genentech), sunitinib malate (sold under the tradename Sutent® by Pfizer), bosutinib (4-[(2,4-dichloro-5-methoxyphenyl)amino]-6-methoxy-7-[3-(4-methylpiperazin-1-yl)propoxy]quinoline-3-carbonitrile, also known as SKI-606, and described in US Patent No. 6,780,996), dasatinib (sold under the tradename Sprycel® by Bristol-Myers Squibb), armala (also known as pazopanib, sold under the tradename Votrient® by GlaxoSmithKline), and imatinib and imatinib mesylate (sold under the tradenames Gilvec® and Gleevec® by Novartis);

(iv) Bcr/Abl kinase inhibitors such as nilotinib hydrochloride (sold under the tradename Tasigna® by Novartis);

(v) DNA Synthesis inhibitors such as Capecitabine (sold under the trademark Xeloda® by Roche), gemcitabine hydrochloride (sold under the trademark Gemzar® by Eli Lilly and Company), and nelarabine ((2R,3S,4R,5R)-2-(2-amino-6-methoxy-purin-9-yl)-5-(hydroxymethyl)oxolane-3,4-diol, sold under the tradenames Arranon® and Atriance® by GlaxoSmithKline);

(vi) Antineoplastic agents such as oxaliplatin (sold under the tradename Eloxatin® by Sanofi-Aventis and described in US Patent No. 4,169,846);

(vii) Epidermal growth factor receptor (EGFR) inhibitors such as Gefitinib (sold under the tradename Iressa®), N-[4-[(3-Chloro-4-fluorophenyl)amino]-7-[(3"S)-

tetrahydro-3-furanyl]oxy]-6-quinazolinyl]-4(dimethylamino)-2-butenamide, sold under the tradename Tovok® by Boehringer Ingelheim), cetuximab (sold under the tradename Erbitux® by Bristol-Myers Squibb), and panitumumab (sold under the tradename Vectibix® by Amgen);

5 (viii) Pro-apoptotic receptor agonists (PARAs) such as Dulanermin (also known as AMG-951, available from Amgen/Genentech);

(ix) PI3K inhibitors such as 4-[2-(1H-Indazol-4-yl)-6-[[4-(methylsulfonyl)piperazin-1-yl]methyl]thieno[3,2-d]pyrimidin-4-yl]morpholine (also known as GDC 0941 and described in PCT Publication Nos. WO 09/036082 and WO 09/055730), and 2-Methyl-2-
10 [4-[3-methyl-2-oxo-8-(quinolin-3-yl)-2,3-dihydroimidazo[4,5-c]quinolin-1-yl]phenyl]propionitrile (also known as BEZ 235 or NVP-BEZ 235, and described in PCT Publication No. WO 06/122806);

(x) BCL-2 inhibitors such as 4-[4-[[2-(4-chlorophenyl)-5,5-dimethyl-1-cyclohexen-1-yl]methyl]-1-piperazinyl]-N-[[4-[[[(1R)-3-(4-morpholinyl)-1-
15 [(phenylthio)methyl]propyl]amino]-3-[(trifluoromethyl)sulfonyl]phenyl]-sulfonyl]benzamide (also known as ABT-263 and described in PCT Publication No. WO 09/155386);

(xi) Topoisomerase I inhibitors such as Irinotecan (sold under the trademark Camptosar® by Pfizer), topotecan hydrochloride (sold under the tradename Hycamtin® by GlaxoSmithKline);

20 (xii) Topoisomerase II inhibitors such as etoposide (also known as VP-16 and Etoposide phosphate, sold under the tradenames Toposar®, VePesid® and Etopophos®), and teniposide (also known as VM-26, sold under the tradename Vumon®);

(xiii) CTLA-4 inhibitors such as Tremelimumab (IgG2 monoclonal antibody
25 available from Pfizer, formerly known as ticilimumab, CP-675,206), and ipilimumab (CTLA-4 antibody, also known as MDX-010, CAS No. 477202-00-9);

(xiv) Histone deacetylase inhibitors (HDI) such as Voninostat (sold under the tradename Zolinza® by Merck) and Panobinostat (N-hydroxy-3-[4-[[[2-(2-methyl-1H-indol-3-yl)ethyl]amino]methyl]phenyl]-(2E)-2-Propenamide described in PCT Publication
30 No. 02/0022577 or US Patent No. 7,067,551);

(xv) Alkylating agents such as Temozolomide (sold under the tradenames Temodar® and Temodal® by Schering-Plough/Merck), dactinomycin (also known as actinomycin-D and sold under the tradename Cosmegen®), melphalan (also known as L-PAM, L-sarcosylsin, and phenylalanine mustard, sold under the tradename Alkeran®),
35 altretamine (also known as hexamethylmelamine (HMM), sold under the tradename Hexalen®), carmustine (sold under the tradename BiCNU®), bendamustine (sold under the tradename Treanda®), busulfan (sold under the tradenames Busulfex® and

Myleran®), carboplatin (sold under the tradename Paraplatin®), lomustine (also known as CCNU, sold under the tradename CeeNU®), cisplatin (also known as CDDP, sold under the tradenames Platinol® and Platinol®-AQ), chlorambucil (sold under the tradename Leukeran®), cyclophosphamide (sold under the tradenames Cytoxan® and
 5 Neosar®), dacarbazine (also known as DTIC, DIC and imidazole carboxamide, sold under the tradename DTIC-Dome®), altretamine (also known as hexamethylmelamine (HMM) sold under the tradename Hexalen®), ifosfamide (sold under the tradename Ifex®), procarbazine (sold under the tradename Matulane®), mechlorethamine (also known as nitrogen mustard, mustine and mechloroethamine hydrochloride, sold under
 10 the tradename Mustargen®), streptozocin (sold under the tradename Zanosar®), and thiotepa (also known as thiophosphoamide, TESPA and TSPA, sold under the tradename Thioplex®;

(xvi) Anti-tumor antibiotics such as doxorubicin (sold under the tradenames Adriamycin® and Rubex®), bleomycin (sold under the tradename lenoxane®),
 15 daunorubicin (also known as dauorubicin hydrochloride, daunomycin, and rubidomycin hydrochloride, sold under the tradename Cerubidine®), daunorubicin liposomal (daunorubicin citrate liposome, sold under the tradename DaunoXome®), mitoxantrone (also known as DHAD, sold under the tradename Novantrone®), epirubicin (sold under the tradename Ellence™), idarubicin (sold under the tradenames Idamycin®, Idamycin
 20 PFS®), and mitomycin C (sold under the tradename Mutamycin®);

(xvii) Anti-mitotic agents such as Docetaxel (sold under the tradename Taxotere® by Sanofi-Aventis);

(xviii) Proteasome inhibitors such as Bortezomib (sold under the tradename Velcade®);

25 (xix) Plant Alkaloids such as Paclitaxel protein-bound (sold under the tradename Abraxane®), vinblastine (also known as vinblastine sulfate, vincalkeboplastine and VLB, sold under the tradenames Alkaban-AQ® and Velban®), vincristine (also known as vincristine sulfate, LCR, and VCR, sold under the tradenames Oncovin® and Vincasar Pfs®), vinorelbine (sold under the tradename Navelbine®), and paclitaxel (sold under the
 30 tradenames Taxol and Onxal™);

(xx) Glucocorticosteroids such as Hydrocortisone (also known as cortisone, hydrocortisone sodium succinate, hydrocortisone sodium phosphate, and sold under the tradenames Ala-Cort®, Hydrocortisone Phosphate, Solu-Cortef®, Hydrocort Acetate® and Lanacort®), dexamethazone ((8S,9R,10S,11S,13S,14S,16R,17R)-9-fluoro-11,17-
 35 dihydroxy-17-(2-hydroxyacetyl)-10,13,16-trimethyl-6,7,8,9,10,11,12,13,14,15,16,17-dodecahydro-3H-cyclopenta[a]phenanthren-3-one), prednisolone (sold under the tradenames Delta-Cortel®, Orapred®, Pediapred® and Prelone®), prednisone (sold

under the tradenames Deltasone®, Liquid Red®, Meticorten® and Orasone®), and methylprednisolone (also known as 6-Methylprednisolone, Methylprednisolone Acetate, Methylprednisolone Sodium Succinate, sold under the tradenames Duralone®, Medralone®, Medrol®, M-Prednisol® and Solu-Medrol®);

- 5 (xxi) Tumor necrosis factor–related apoptosis-inducing ligand (TRAIL, also referred to as Apo2 Ligand) receptor agonists such as TRAIL antibodies (e.g., Adecatumumab, Belimumab, Cixutumumab, Conatumumab, Figitumumab, Iratumumab, Lexatumumab, Lucatumumab, Mapatumumab, Necitumumab, Ofatumumab, Olaratumab, Panitumumab, Pritumumab, Pritumumab, Robatumumab, Votumumab, 10 Zalutumumab, and TRAIL (referred to as anti-DR-5) antibodies described in US Patent No. 7,229,617 and PCT Publication No. WO2008/066854, incorporated herein by reference), and recombinant TRAIL (e.g., Dulanermin (also known as AMG 951 (rhApo2L/TRAIL))); and

- (xxii) Tumor-vascular disrupting agents such as Vadimezan (5,6-dimethyl-9-oxo- 15 9H-Xanthene-4-acetic acid described in US Patent No. 5,281,620).

A preferred anti-cancer agent for use in combination with a compound of the present invention is paclitaxel.

- Another preferred anti-cancer agent for use in combination with a compound of the present invention is a PI3K inhibitor (e.g., 2-Methyl-2-[4-[3-methyl-2-oxo-8-(quinolin- 20 3-yl)-2,3-dihydroimidazo[4,5-c]quinolin-1-yl]phenyl]propionitrile).

Another preferred anti-cancer agent for use in combination with a compound of the present invention is a TRAIL (or anti-DR-5) antibody or recombinant TRAIL.

- Suitable therapeutic agents for adjunct therapy include steroids, anti-inflammatory agents, anti-histamines, antiemetics, and other agents well-known to those of skill in art 25 for use in improving the quality of care for patients being treated for the diseases, conditions, or disorders described herein.

- The compound of the present invention or pharmaceutical composition thereof for use in humans is typically administered intravenously *via* infusion at a therapeutic dose of less than or equal to about 100 mg/kg, 75 mg/kg, 50 mg/kg, 25 mg/kg, 10 mg/kg, 7.5 30 mg/kg, 5.0 mg/kg, 3.0 mg/kg, 1.0 mg/kg, 0.5 mg/kg, 0.05 mg/kg or 0.01 mg/kg, but preferably not less than about 0.0001 mg/kg. The dosage may depend upon the infusion rate at which the formulation is administered. In general, the therapeutically effective dosage of a compound, the pharmaceutical composition, or the combinations thereof, is dependent on the species of the subject, the body weight, age and individual condition, 35 the disorder or disease or the severity thereof being treated. A physician, pharmacist, clinician or veterinarian of ordinary skill can readily determine the effective amount of

each of the active ingredients necessary to prevent, treat or inhibit the progress of the disorder or disease.

The above-cited dosage properties are demonstrable *in vitro* and *in vivo* tests using advantageously mammals, e.g., mice, rats, dogs, monkeys or isolated organs, tissues and preparations thereof. The compounds of the present invention can be applied *in vitro* in the form of solutions, e.g., aqueous solutions, and *in vivo* either enterally, parenterally, advantageously intravenously, e.g., as a suspension or in aqueous solution. The dosage *in vitro* may range between about 10^{-3} molar and 10^{-9} molar concentrations.

In general, a therapeutically effective amount of a compound of the present invention is administered to a patient in need of treatment. The term "a therapeutically effective amount" of a compound of the present invention refers to an amount of the compound of the present invention that will elicit the biological or medical response of a subject, for example, reduction or inhibition of an enzyme or a protein activity, or ameliorate symptoms, alleviate conditions, slow or delay disease progression, or prevent a disease, etc.

In one non-limiting embodiment, the term "a therapeutically effective amount" refers to the amount of a compound of the present invention, when administered to a subject, is effective to (1) at least partially alleviate, inhibit, prevent and/or ameliorate a condition, a disorder or a disease mediated by IAP, or characterized by normal or abnormal activity of such IAP mediation or action; or (2) enhance programmed cancerous cell death (apoptosis). Preferably, when administered to a cancer cell, or a tissue, or a non-cellular biological material, or a medium, the compound of the present invention is effective to at least partially increase or enhance apoptosis. Not to be bound by any particular mechanism, a compound of the present may inhibit the binding of IAP protein to a caspase protein and/or may initiate degradation of XIAP, cIAP1 and/or cIAP2, directly or indirectly.

In one embodiment, a method for inhibiting the binding of an IAP protein to a caspase protein is provided which comprises contacting the IAP protein with a compound of the present invention.

In another embodiment, a method of inducing apoptosis in a tumor or cancer cell is provided which comprises introducing into the cell, a compound of the present invention.

In yet another embodiment, a method of sensitizing a tumor or cancer cell to an apoptotic signal is provided which comprises introducing into the cell a compound of the present invention.

In yet another embodiment, a method for treating a disease, disorder, or condition associated with the overexpression of an IAP in a mammal, is provided which comprises administering to the mammal an effective amount of a compound of the present invention.

5 In yet another embodiment, a method for treating cancer in a mammal is provided which comprises administering to a mammal in need of such treatment an effective amount of a compound of the present invention.

As used herein, the term "subject" refers to an animal. Typically the animal is a mammal. A subject also refers to for example, primates (e.g., humans, male or female),
10 cows, sheep, goats, horses, dogs, cats, rabbits, rats, mice, fish, birds and the like. In certain embodiments, the subject is a primate. Preferably, the subject is a human.

As used herein, the term "inhibit", "inhibition" or "inhibiting" refers to the reduction or suppression of a given condition, symptom, or disorder, or disease, or a significant decrease in the baseline activity of a biological activity or process.

15 As used herein, the term "treat", "treating" or "treatment" of any disease or disorder, refers (i) to ameliorating the disease or disorder (i.e., slowing or arresting or reducing the development of the disease or at least one of the clinical symptoms thereof); (ii) to alleviating or ameliorating at least one physical parameter including those which may not be discernible by the patient; or (iii) to preventing or delaying the onset or
20 development or progression of the disease or disorder. In general, the term "treating" or "treatment" describes the management and care of a patient for the purpose of combating the disease, condition, or disorder and includes the administration of a compound of the present invention to prevent the onset of the symptoms or complications, alleviating the symptoms or complications, or eliminating the disease,
25 condition or disorder.

As used herein, a subject is "in need of" a treatment if such subject would benefit biologically, medically or in quality of life from such treatment (preferably, a human).

Another aspect of the invention is a product comprising a compound of the present invention and at least one other therapeutic agent (or pharmaceutical agent) as a
30 combined preparation for simultaneous, separate or sequential use in therapy to enhance apoptosis.

In the combination therapies of the invention, the compound of the present invention and the other therapeutic agent may be manufactured and/or formulated by the same or different manufacturers. Moreover, the compound of the present invention and
35 the other therapeutic (or pharmaceutical agent) may be brought together into a combination therapy: (i) prior to release of the combination product to physicians (e.g. in the case of a kit comprising the compound of the invention and the other therapeutic

agent); (ii) by the physician themselves (or under the guidance of the physician) shortly before administration; (iii) in the patient themselves, e.g. during sequential administration of the compound of the invention and the other therapeutic agent.

Accordingly, the invention provides the use of a compound of the present
5 invention for treating a disease or condition by inhibiting IAPs (or enhancing apoptosis), wherein the medicament is prepared for administration with another therapeutic agent. The invention also provides for the use of another therapeutic agent, wherein the medicament is administered as a combination of a compound of the present invention with the other therapeutic agent.

10 Embodiments of the present invention are illustrated by the following Examples. It is to be understood, however, that the embodiments of the invention are not limited to the specific details of these Examples, as other variations thereof will be known, or apparent in light of the instant disclosure, to one of ordinary skill in the art.

EXAMPLES

15 Unless specified otherwise, starting materials are generally available from commercial sources such as Aldrich Chemicals Co. (Milwaukee, Wis.), Lancaster Synthesis, Inc. (Windham, N.H.), Acros Organics (Fairlawn, N.J.), Maybridge Chemical Company, Ltd. (Cornwall, England), Tyger Scientific (Princeton, N.J.), and AstraZeneca Pharmaceuticals (London, England).

20 The following abbreviations used herein below have the corresponding meanings:

Boc: *tert*-butoxycarbonyl (N-protecting group)

DIEA: *N,N*-Diisopropylethylamine (also known as Hunig's base)

DMF: Dimethylformamide

DMTMM: 4-(4,6-Dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride

25 HATU: 2-(1H-7-Azabenzotriazol-1-yl)--1,1,3,3-tetramethyl uronium
hexafluorophosphate Methanaminium

HBTU: O-Benzotriazole-*N,N,N',N'*-tetramethyl-uronium-hexafluoro-phosphate

HOBt: *N*-Hydroxybenzotriazole


NMP: *N*-Methyl-2-pyrrolidone

30 TFA: Trifluoroacetic acid

3-Bromo-5-{(S)-1-[(R)-1-(4-methoxy-phenyl)-ethyl]-pyrrolidin-2-yl}-pyridine was prepared as described in PCT Publication No. WO 2008/016893 A1 on page 33.


Example 1

Preparation of (S,S,S)-N,N'-(3,3'-(Ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide) (1A):



5 (1A)

Preparation of Intermediate 1-(5-((S)-1-[(R)-1-(4-Methoxy-phenyl)-ethyl]-pyrrolidin-2-yl)-pyridin-3-yl)-1H-indole-4-carboxylic acid ethyl ester (I-1A-1a):




(I-1A-1a)

10 A mixture of 3-bromo-5-[(S)-1-[(R)-1-(4-methoxy-phenyl)-ethyl]-pyrrolidin-2-yl]-pyridine (4.0 g, 11 mmol), 1H-indole-4-carboxylic acid ethyl ester (6.3 g, 33 mmol), K₂CO₃ (2.3 g, 16.6 mmol), copper iodide (0.42 g, 2.2 mmol) in NMP (22 mL) was heated in a microwave tube at 190°C for 30 minutes. The resulting mixture was cooled to room temperature filtered through celite and concentrated under reduced pressure to give an
15 oil. This oil was purified by silica gel column chromatography (30% ethyl acetate/heptane) to give the titled compound as a yellow solid (2.2 g, 42%).

¹H NMR (400 MHz, chloroform-*d*) δ ppm 8.82 (br. s., 2 H), 7.97 (d, *J*=7.6 Hz, 1 H), 7.65 (d, *J*=8.2 Hz, 1 H), 7.51 (s, 1 H), 7.39 (s, 2 H), 7.25 (t, *J*=7.8 Hz, 1 H), 7.10 (d, *J*=8.6 Hz, 2 H), 6.65 (d, *J*=8.6 Hz, 2 H), 4.50 (q, *J*=7.2 Hz, 2 H), 3.77 (dd, *J*=8.8, 5.7 Hz, 1 H), 3.52 - 3.67 (m, 4 H), 3.24 (t, *J*=7.2 Hz, 1 H), 2.51 - 2.67 (m, 1 H), 2.28-2.32 (m, 1 H), 1.92 - 2.06 (m, 1 H), 1.79 - 1.90 (m, 1 H), 1.62 - 1.79 (m, 1 H), 1.51 (t, *J*=7.1 Hz, 3 H), 1.36 (d, *J*=6.6 Hz, 3 H); LC-MS calculated for C₂₉H₃₂N₃O₃ 470.2, found 470.4 (ESI *m/e* [M + H⁺]); *t_R* 1.78 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

25 Preparation of Intermediate 1-((S)-5-Pyrrolidin-2-yl-pyridin-3-yl)-1H-indole-4-carboxylic acid ethyl ester (I-1A-1b):




(I-1A-1b)

A solution of 1-(5-[(S)-1-[(R)-1-(4-methoxy-phenyl)-ethyl]-pyrrolidin-2-yl]-pyridin-3-yl)-1H-indole-4-carboxylic acid ethyl ester (I-1A-1a: 5.0 g, 11 mmol) in TFA (15 mL) was heated in a microwave tube at 100°C for 30 minutes. The resulting mixture was concentrated under reduced pressure to give a di-TFA salt of the titled compound as a red oil (3.2 g) which was used directly in the next step without purification.

¹H NMR (400 MHz, chloroform-*d*) δ ppm 9.80 - 10.03 (br. s, 1 H), 9.40 - 9.80 (br. s, 1 H), 8.88 - 9.02 (m, 1 H), 8.72 - 8.85 (m, 1 H), 8.45 (s, 1 H), 7.91 (m, 1 H), 7.60-7.72 (m, 1 H), 7.33 - 7.45 (m, 2 H), 7.22 - 7.33 (m, 1 H), 4.75 - 4.95 (m, 1 H), 4.37 (q, *J*=7.1 Hz, 2 H), 3.63 - 3.80 (m, 1 H), 3.36 - 3.59 (m, 2 H), 2.45 - 2.63 (m, 1 H), 2.27 - 2.41 (m, 2 H), 2.08 - 2.26 (m, 1 H), 1.37 (t, *J*=7.1 Hz, 3 H); LC-MS calculated for C₂₀H₂₂N₃O₂ 336.2, found 336.4 (ESI *m/e* [M + H⁺]); *t_R* 1.09 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

Preparation of Intermediate 1-{5-[(S)-1-[(S)-2-tert-Butoxycarbonylamino-3-methyl-butyryl]-pyrrolidin-2-yl]-pyridin-3-yl}-1H-indole-4-carboxylic acid ethyl ester (IA-1c):




(I-1A-1c)

To a solution of 1-((S)-5-pyrrolidin-2-yl-pyridin-3-yl)-1H-indole-4-carboxylic acid ethyl ester di-TFA salt (I-1A-1b: 2 g, 4.5 mmol) in DMF (20 mL) was added DIEA (10.4 mL, 60 mmol), Boc-L-valine (1.3 g, 6 mmol), O-(benzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HBTU) (2.5 g, 6.6 mmol), and HOBT (1.0 g, 6.6 mmol). The mixture was stirred at room temperature overnight and then diluted with EtOAc (200 mL) and washed with water (2 x 100 mL). The aqueous layer was back extracted with EtOAc (50 mL) and the combined organic layer was washed with water (2 x 25 mL). The combined organic phase was dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give an oil. This oil was purified by silica gel column chromatography (5-50% EtOAc/heptane) to give the titled compound as a light yellow foam (1.8 g, 94%).

¹H NMR (400 MHz, chloroform-*d*) δ ppm 8.65 - 8.78 (m, 1 H), 8.52 (s, 1 H), 8.00 (d, *J*=7.5 Hz, 1 H), 7.59 - 7.77 (m, 2 H), 7.38 - 7.46 (m, 2 H), 7.29 (t, *J*=7.9 Hz, 1 H), 5.08

- 5.36 (m, 2 H), 4.49 (q, $J=7.1$ Hz, 2 H), 4.35-4.44 (m, 1 H), 3.92 - 4.09 (m, 1 H), 3.65 - 3.88 (m, 1 H), 2.38 - 2.59 (m, 1 H), 1.93 - 2.17 (m, 4 H), 1.49 (t, $J=7.1$ Hz, 3 H), 1.45 (s, 9 H), 0.99 (d, $J=6.7$ Hz, 3 H), 0.86 (d, $J=6.8$ Hz, 3 H); LC-MS calculated for $C_{30}H_{39}N_4O_5$ 535.3, found 535.5 (ESI m/e [$M + H^+$]); t_R 1.67 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

Preparation of Intermediate 1-[5-((S)-1-((S)-2-((S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester (I-1A-1d):




(I-1A-1d)

A solution of 1-[5-((S)-1-((S)-2-*tert*-butoxycarbonylamino-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester (I-1A-1c: 2.5 g, 4.7 mmol) in TFA (2 mL) and CH_2Cl_2 (2 mL) was stirred at room temperature for 18 hours. More TFA (3.6 mL) was added. The mixture was stirred for 3 hours and concentrated under reduced pressure. To the resulting oily residue was added anhydrous DMF, DIEA (7.6 mL, 44 mmol), Boc-L-valine (0.89 g, 4.4 mmol), HBTU (1.8 g, 4.8 mmol), and HOBT (0.74 g, 4.9 mmol). The mixture was stirred at room temperature overnight. To the reaction mixture was added more HBTU (0.83 g, 2.2 mmol), Boc-Me-Alanine (0.18 g, 0.87 mmol), and DIEA (2.3 mL, 13.1 mmol). Stirring continued at room temperature overnight. The reaction mixture was then diluted with EtOAc (200 mL) and washed with water (2 x 100 mL). The aqueous layer was extracted with EtOAc (50 mL) and the combined organic layers were washed with water again (2 x 25 mL). The organic layer was dried over anhydrous Na_2SO_4 and concentrated under reduced pressure to give an oil. This oil was purified by silica gel column chromatography (0-100% EtOAc/heptane) to give the titled compound as a light yellow foam (1.7 g, 58% for 2 steps).

1H NMR (400 MHz, chloroform- d) δ ppm 8.71 (s, 1 H), 8.52 (s, 1 H), 8.10-7.90 (m, 1 H), 7.60 - 7.70 (m, 2 H), 7.43-7.35 (m, 2 H), 7.20-7.33 (m, 1 H), 6.75-6.56 (br, d, 1H), 5.29 - 5.25 (m, 1 H), 4.71 - 4.67 (m, 1 H), 4.43 - 4.54 (m, 2 H), 3.98 - 4.08 (m, 1 H), 3.67 - 3.87 (m, 3 H), 2.80 (s, 3 H), 2.49 - 2.41 (m, 1 H), 1.91 - 2.22 (m, 4 H), 1.43 - 1.53 (m, 12 H), 1.35 (d, $J=7.0$ Hz, 3 H), 0.96 (d, $J=6.0$ Hz, 3 H), 0.86 (d, $J=6.0$ Hz, 3 H); LC-MS calculated for $C_{34}H_{46}N_5O_6$ 620.3, found 620.5 (ESI m/e [$M + H^+$]); t_R 1.63 minutes

(Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

5 Preparation of Intermediate 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid (I-1A-1e)



(I-1A-1e)

To a solution of 1-[5-((S)-1-((S)-2-[(S)-2-(tert-butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester (I-1A-1d: 1.2 g, 1.9 mmol) in THF (5 mL) and MeOH (1.7 mL) was added a solution of LiOH (0.080 g, 3.5 mmol) in water (1.7 mL). The mixture was stirred at room temperature for 5 hours. More LiOH (0.023 g, 0.97 mmol) was added and stirring continued for overnight. The reaction mixture was quenched with water and concentrated under reduced pressure to remove THF. The remaining mixture was diluted with water and neutralized with 1 M HCl dropwise until pH = 4. The white precipitate formed was filtered and dried in vacuo to give the titled compound as a white powder (1.0 g, 87%).

¹H NMR (400 MHz, chloroform-*d*) δ ppm 8.66 - 8.79 (m, 1 H), 8.42 - 8.58 (m, 1 H), 8.06 (d, *J*=7.5 Hz, 1 H), 7.62 - 7.79 (m, 2 H), 7.44 (s, 2 H), 7.32 (t, *J*=7.8 Hz, 1 H), 6.60-6.90 (br, m, 1 H), 5.12 - 5.35 (m, 1 H), 4.64 - 4.81 (m, 2 H), 3.98 - 4.11 (m, 1 H), 3.77 - 3.87 (m, 1 H), 2.80 (br. s., 3 H), 2.37 - 2.54 (m, 1 H), 1.93 - 2.19 (m, 4 H), 1.47 (s, 9 H), 1.35 (d, *J*=7.0 Hz, 3 H), 0.95 (d, *J*=6.0 Hz, 3 H), 0.86 (d, *J*=6.5 Hz, 3 H); LC-MS calculated for C₃₂H₄₂N₅O₆ 592.3, found 592.5 (ESI *m/e* [M + H⁺]); *t_R* 1.35 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

Preparation of the title Compound 1A:

To a solution of 1-[5-((S)-1-((S)-2-[(S)-2-(tert-butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid (I-1A-1e: 0.20 g, 0.34 mmol) in THF (4 mL) was added DIEA (0.14 mL, 0.80 mmol),

followed by O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.13 g, 0.35 mmol) at 0°C. The mixture was stirred at 0°C for 15 minutes. 3,3'-(Ethane-1,2-diylbis(oxy))dipropylamine (0.028 g, 0.16 mmol) was added. The resulting mixture was warmed up to room temperature and stirred overnight.

5 It was concentrated under reduced pressure and the residue was dissolved in MeOH (3 mL) and filtered consecutively through SiliaPrep propylsulfonic acid (SCX-2) resin and SiliaPrep carbonate resin to remove amine materials, excess carboxylic acid and peptide couplings by-products, respectively. Both resins were washed again with MeOH (6 mL) and the combined filtrate was concentrated under reduced pressure to give a yellow oil

10 (alternatively, the reaction mixture was quenched with saturated NaHCO₃ (15 mL) and extracted with EtOAc (3x50 mL). The combined organic phase was washed with saturated citric acid (15 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give an oil). To this oil was added CH₂Cl₂ (2 mL) and TFA (2 mL). The mixture was stirred at room temperature for 3 hours, concentrated under reduced

15 pressure, and purified by preparative HPLC (10-100% CH₃CN/H₂O over 10 minutes with 0.1% TFA, Sunfire C18 OBD 50x50 mm column with flow rate of 60 mL/minute). The product-containing fractions were concentrated in genevac overnight to give an oil. This oil was dissolved in MeOH (3 mL) and filtered through a StratoSpheres SPE PL-HCO₃ MP-Resin and the cartridge was washed with MeOH (6 mL). The combined filtrate was


20 concentrated under reduced pressure to afford the free base form of the titled compound as a white foam. To this foam (0.090 g, 0.080 mmol) in EtOAc (2 mL) was added citric acid (0.031 g, 0.16 mmol) in MeOH (2 mL). The resulting mixture was stirred at room temperature for 3 hours and concentrated under reduced pressure to give a glassy residue. This residue was dissolved in minimum amount of water (1 mL) and lyophilized

25 to give the citrate salt (2.25 eq. by ¹H NMR) of the titled compound as a white powder (0.12 g, 47% in three steps).

¹H NMR (400 MHz, MeOD) δ ppm 8.60 - 8.74 (m, 2 H), 8.34 - 8.57 (m, 2 H), 7.84 - 7.95 (m, 2 H), 7.55 - 7.77 (m, 4 H), 7.44 - 7.54 (m, 2 H), 7.21 - 7.34 (m, 2 H), 7.07 - 7.16 (m, 2 H), 5.16 - 5.56 (m, 2 H), 4.54 - 4.64 (m, 2 H), 4.00 - 4.14 (m, 2 H), 3.88 - 3.99 (m, 2 H), 3.78 - 3.88 (m, 2 H), 3.64 (s, 4 H), 3.55 - 3.62 (m, 4 H), 3.43 - 3.53 (m, 4 H), 2.62 (s, 6 H), 2.37 - 2.53 (m, 2 H), 1.96 - 2.23 (m, 8 H), 1.79 - 1.94 (m, 4 H), 1.41 - 1.53 (m, 6 H), 0.93 - 1.06 (m, 6 H), 0.81 - 0.91 (m, 6 H), citrate: 2.69-2.75 (m, 4.5 H), 2.78-2.83 (m, 4.5H), LC-HRMS calculated for C₆₂H₈₃N₁₂O₈ 1123.6457, found 1123.6478 (ESI *m/e* [M + H⁺]); *t_R* 2.98 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 35 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

The compounds below were prepared using procedures analogous to those described above for the preparation of Example 1A with the appropriate starting materials.


- 5 Preparation of (S,S,S)-N,N'-(1,4-phenylenebis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide) (1B):



- 10 The TFA salt of the titled compound was prepared as a white powder (0.017 g, 20% in two steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl-(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxylic acid (l-1A-1e: 0.10 g, 0.17 mmol), 1,4-phenylenedimethanamine (0.011 g, 0.08 mmol) and 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide (EDC) hydrochloride (0.034 g, 0.18 mmol) in DMF following the general procedure described above for the preparation of Example 1A.

- ¹H NMR (400 MHz, MeOD) δ ppm 8.75 (br. s., 2 H), 8.41 - 8.61 (m, 2 H), 8.03 (br. s., 2 H), 7.66 - 7.82 (m, 2 H), 7.61 - 7.79 (m, 2 H), 7.58 (d, J=7.0 Hz, 2 H), 7.44 (s, 4 H), 7.22-7.32 (m, 2 H), 7.17 (br. s., 2 H), 5.19 (t, J=6.3 Hz, 2 H), 4.66 (s, 4 H), 4.61 (d, J=6.5 Hz, 2 H), 4.05 - 4.16 (m, 2 H), 3.81 - 4.02 (m, 4 H), 2.67 (s, 6 H), 2.41 - 2.59 (m, 2 H), 1.89 - 2.26 (m, 8 H), 1.49 (d, J=7.0 Hz, 6 H), 0.97 (d, J=6.5 Hz, 6 H), 0.91 (d, J=7.0 Hz, 6 H); LC-HRMS calcd for C₆₂H₇₅N₁₂O₆ 1083.5933, found 1083.5878 (ESI m/e [M + H⁺]); t_R 3.13 min (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 99% by UV 254/214 nm.


Preparation of (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide) (1C):



The citrate salt (3 equivalents) of the titled compound was prepared as a white powder (0.011 g, 9% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxylic acid (l-1A-1e: 0.10 g, 0.17 mmol), propane-1,3-diamine (0.006 g, 0.081 mmol) and HATU (0.067 g, 0.18 mmol) following the general procedure described above for the preparation of Example 1A.

¹H NMR (400 MHz, MeOD) δ ppm 8.68-8.55 (m, 2 H), 8.53-8.35 (m, 2 H), 7.93-7.83 (m, 2 H), 7.55 - 7.74 (m, 6 H), 7.32-7.22 (m, 2 H), 7.16 - 7.25 (m, 2 H), 5.05 - 5.22 (m, 2 H), 4.62 (d, *J*=6.5 Hz, 2 H), 4.22 (dd, *J*=6.0, 2.0 Hz, 2 H), 4.03 - 4.16 (m, 2 H), 3.82 - 3.94 (m, 2 H), 3.50-3.65 (m, 4 H), 2.67 (s, 6 H), 2.40 - 2.57 (m, 2 H), 1.95 - 2.24 (m, 8 H), 1.53-1.72 (m, 2 H), 1.48 - 1.50 (m, 6 H), 0.96 (d, *J*=7.5 Hz, 6 H), 0.91 (d, *J*=6.5 Hz, 6 H), citrate: 3.57 - 3.77 (m, 6 H), 3.84 - 4.01 (m, 6 H); LC-HRMS calculated for C₅₇H₇₃N₁₂O₆ 1021.5776, found 1021.5788 (ESI *m/e* [M + H⁺]); *t_R* 2.87 min (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 95% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-((1S,2S)-cyclohexane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide)(1D):




The TFA salt of the titled compound was prepared as a white powder (0.005 g, 6% in two steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)-propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxylic acid (1-1A-1e: 0.10 g, 0.17 mmol), (1R,2R)-cyclohexane-1,2-diamine (0.009 g, 0.081 mmol) and HATU (0.068 g, 0.18 mmol) following general procedure described above for the preparation of Example 1A.

¹H NMR (400 MHz, MeOD) δ ppm 8.47 - 8.66 (m, 4 H), 7.99 - 8.13 (m, 4 H), 7.79 - 7.91 (m, 2 H), 7.37 (d, *J*=8.1 Hz, 2 H), 7.19 (t, *J*=7.3 Hz, 2 H), 7.08-7.15 (m, 2 H), 5.17 (d, *J*=5.6 Hz, 2 H), 4.59 (d, *J*=6.6 Hz, 2 H), 4.01 - 4.14 (m, 4 H), 3.81 - 3.95 (m, 4 H), 2.67 (s, 6 H), 2.37 - 2.53 (m, 2 H), 1.83 - 2.24 (m, 12 H), 1.60-1.80 (m, 2H), 1.48 (d, *J*=7.1 Hz, 6 H), 1.30-1.40 (m, 2 H), 0.91 - 1.05 (m, 6 H), 0.88 (d, *J*=6.6 Hz, 6 H); LC-
HRMS calculated for C₆₀H₇₇N₁₂O₆ 1061.6089, found 1061.6088 (ESI *m/e* [M + H⁺]); *t_R* 3.22 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 99% by UV 254/214 nm.


Example 2

Preparation of (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2A):



(2A)

Preparation of Intermediate 1-(5-{(S)-1-[(R)-1-(4-Methoxy-phenyl)-ethyl]-pyrrolidin-2-yl}-pyridin-3-yl)-1H-indole-3-carboxylic acid ethyl ester (I-2A-2a):



(I-2A-2a)


5

To a suspension of 3-bromo-5-{(S)-1-[(R)-1-(4-methoxy-phenyl)-ethyl]-pyrrolidin-2-yl}-pyridine (2.0 g, 5.5 mmol), 1H-Indole-3-carboxylic acid ethyl ester (1.3 g, 6.6 mmol), copper iodide (0.11 g, 0.55 mmol), K₃PO₄ (2.5 g, 11.6 mmol) in toluene (5 mL) was added (1S,2S)-N1,N2-dimethylcyclohexane-1,2-diamine (0.16 g, 1.1 mmol). After being flushed with N₂, the resulting mixture was heated at 110°C overnight, cooled to room temperature, filtered through celite and concentrated under reduced pressure to give an oil. This oil was purified by silica gel column chromatography using 30-50% EtOAc/heptane to give the titled compound as a light yellow foam (2.0 g, 77%).

¹H NMR (400 MHz, chloroform-*d*) δ ppm 8.43 (br. s., 2 H), 8.19 (d, *J*=8.0 Hz, 1 H), 7.84 (s, 1 H), 7.42 (s, 1 H), 7.14 - 7.32 (m, 3 H), 6.97 (d, *J*=8.3 Hz, 2 H), 6.53 (d, *J*=8.3 Hz, 2 H), 4.36 (q, *J*=7.2 Hz, 2 H), 3.65 (dd, *J*=8.9, 5.6 Hz, 1 H), 3.56 (s, 3 H), 3.52 (d, *J*=6.6 Hz, 1 H), 3.11 - 3.24 (m, 1 H), 2.46 - 2.61 (m, 1 H), 2.20 (m, 1 H), 1.88 (m, 1 H), 1.71 - 1.82 (m, 1 H), 1.53 - 1.69 (m, 1 H), 1.38 (t, *J*=7.1 Hz, 3 H), 1.29 (d, *J*=6.6 Hz, 3 H); LC-MS calculated for C₂₉H₃₂N₃O₃ 470.2, found 470.4 (ESI *m/e* [M + H⁺]); *t_R* 1.79 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

Preparation of Intermediate 1-((S)-5-Pyrrolidin-2-yl-pyridin-3-yl)-1H-indole-3-carboxylic acid ethyl ester (I-2b):

25




(I-2A-2b)

In a 20 mL microwave vial was added 1-((S)-5-Pyrrolidin-2-yl-pyridin-3-yl)-1H-indole-3-carboxylic acid ethyl ester (I-2A-2a: 1.5 g, 3.19 mmol) and trifluoroacetic acid (7 mL, 91 mmol). The vial was capped, and microwaved at 100°C for 35 minutes. The trifluoroacetic acid was removed under reduced pressure then the resulting oil was taken up in CH₂Cl₂ (50 mL) and excess saturated aqueous NaHCO₃ was added until the solution was slightly basic. The layers were partitioned and the organic layer was washed with brine, then dried over Na₂SO₄. The drying agent was filtered off, and the filtrate was concentrated to give an amber oil as the crude product (1.14 g) which was used in the subsequent step without further purification. LC-MS calculated for C₂₀H₂₂N₃O₂ 336.2, found 336.3 (ESI *m/e* [M + H⁺]); *t_R* 1.18 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 5 mM ammonium formate, at 2 mL/minute over 2 minutes, 40°C column temperature).

¹H NMR (400 MHz, chloroform-*d*) δ ppm 8.76 (br. s, 2 H), 8.27 (d, *J*=7.5 Hz, 1 H), 8.09 (s, 1 H), 8.03 (s, 1 H), 7.52 (d, *J*=7.5 Hz, 1 H), 7.30-7.41 (m, 2 H), 7.10-7.18 (m, 1 H), 6.72-7.02 (m, 1 H), 4.43 (q, *J*=7.0 Hz, 2 H), 3.26 (br. s, 1 H), 3.18 (br. s, 1 H), 2.39 (br. s, 1 H), 1.94-2.19 (m, 2 H), 1.83-1.91 (m, 2 H), 1.46 (t, *J*=7.0 Hz, 3 H).

Preparation of Intermediate 1-[5-((S)-1-[(S)-2-[(S)-2-(tert-butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)]-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid ethyl ester (I-2A-2c):



(I-2A-2c)

To a solution of 1-((S)-5-pyrrolidin-2-yl-pyridin-3-yl)-1H-indole-3-carboxylic acid ethyl ester (I-2A-2b: 0.25 g, 0.75 mmol) and (S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoic acid (0.26 g, 0.86 mmol, prepared from (S)-2-amino-3-methyl-butyl acid methyl ester and (S)-2-(tert-

butoxycarbonyl-methyl-amino)-propionic acid *via* amide coupling followed ester hydrolysis) in THF (41 mL) at 0°C was added 4-(4, 6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (DMTMM) (0.25 g, 0.89 mmol). The mixture was warmed to room temperature, stirred for 3 hours, then quenched with water (20 mL) and


5 concentrated under reduced pressure to remove THF. The remaining mixture was extracted with EtOAc (3 x 40 mL) and the organic layers were combined, dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give an oil. This oil was purified by silica gel column chromatography using 90-100% EtOAc/heptane to give the titled compound as a white solid (0.30 g, 65%).

10 ¹H NMR (400 MHz, MeOD) δ ppm 8.71 (br. s., 1 H), 8.59 (br. s., 1 H), 8.13 - 8.25 (m, 2 H), 7.83 - 8.01 (m, 1 H), 7.44 - 7.71 (m, 1 H), 7.21 - 7.38 (m, 2 H), 5.13 - 5.27 (m, 1 H), 4.49 - 4.72 (m, 1 H), 4.34 - 4.44 (m, 2 H), 4.27 - 4.36 (m, 1 H), 4.12-4.30 (m, 1 H), 3.90-4.00 (m, 1 H). 2.85 (s, 3 H), 2.40 - 2.63 (m, 1 H), 1.94 - 2.25 (m, 4 H), 1.47-1.55 (m, 12 H), 1.30-1.44 (m, 3 H), 1.15-0.80 (m, 6 H); LC-MS calculated for C₃₄H₄₆N₅O₆ 620.3, found 620.7 (ESI *m/e* [M + H⁺]); *t_R* 1.68 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

15

Preparation of Intermediate 1-[5-((S)-1-{(S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-buteryl}-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid

20 (I-2A-2d):



(I-2A-2d)

To a solution of 1-[5-((S)-1-{(S)-2-[(S)-2-(tert-butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-buteryl}-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid ethyl ester (I-2A-2c: 0.29 g, 0.46 mmol) in THF (1 mL) and MeOH (0.3 mL) was added a

25 solution of LiOH (0.035 g, 0.80 mmol) in water (0.3 mL). The resulting mixture was stirred at room temperature for 5 hours. More LiOH (0.065 g, 2.70 mmol) was added and stirring continued for overnight. The reaction mixture was then concentrated under reduced pressure to remove all organic solvents, then diluted with water. To this mixture

30 1 M HCl was added dropwise until pH = 4. The white precipitate formed was filtered and dried *in vacuo* to give the titled compound as a white powder (0.19 g, 70%).

¹H NMR (400 MHz, chloroform-*d*) δ ppm 8.69 - 8.84 (m, 1 H), 8.50 - 8.66 (m, 1 H), 8.21 - 8.38 (m, 1 H), 8.02 - 8.12 (m, 1 H), 7.63 - 7.79 (m, 1 H), 7.47 (d, *J*=8.5 Hz, 1 H), 7.31 - 7.44 (m, 2 H), 6.91 (br. s., 1 H), 5.19 - 5.39 (m, 1 H), 4.54 - 4.87 (m, 2 H), 4.02 - 4.15 (m, 1 H), 3.69 - 3.91 (m, 1 H), 2.83 (s, 3 H), 2.40 - 2.60 (m, 1 H), 1.91 - 2.25 (m, 4 H), 1.49 (s, 9 H), 1.37 (d, *J*=7.0 Hz, 3 H), 0.97 (d, *J*=6.5 Hz, 3 H), 0.89 (d, *J*=7.0 Hz, 3 H); LC-MS calculated for C₃₂H₄₂N₅O₆ 592.3, found 592.7 (ESI *m/e* [M + H⁺]); *t_R* 1.54 minutes (Insertsil C-8-3, 3 X 3 mm: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 2 mL/minute over 2 minutes).

10 Preparation of Compound 2A:


To a solution of 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)-amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol) in THF (2 mL) was added DIEA (0.070 mL, 0.40 mmol), followed by O-(7-azabenzotriazol-1-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU) (0.068 g, 0.18 mmol) at 0°C. The mixture was stirred at 0°C for 15 minutes, then propane-1,3-diamine (0.006 g, 0.081 mmol) was added. The resulting mixture was warmed to room temperature and stirred overnight. It was concentrated under reduced pressure and the residue was dissolved in MeOH (3 mL) and filtered through SiliaPrep propylsulfonic acid (SCX-2) resin and consecutively
20 through SiliaPrep carbonate resin to remove amine materials and excess carboxylic acid and peptide couplings by-products, respectively. Both resins were washed again with MeOH (6 mL) and the combined filtrate was concentrated under reduced pressure to give a yellow oil (alternatively, the reaction mixture was quenched with saturated NaHCO₃ (15 mL) and extracted with EtOAc (3x50 mL). The organic layers were
25 combined then washed with satd. citric acid (15 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give an oil). To this oil was added CH₂Cl₂ (2 mL) and TFA (2 mL). The mixture was stirred at room temperature for 3 hours, concentrated under reduced pressure, and purified by preparative HPLC (10-100% CH₃CN/H₂O over 10 minutes with 0.1% TFA, Sunfire C18 OBD 50x50 mm column with
30 flow rate of 60 mL/min). The product-containing fractions were concentrated to give an oil. This oil was dissolved in MeOH (3 mL) and filtered through a column of StratoSpheres SPE PL-HCO₃ MP-Resin and the cartridge was washed with MeOH (6 mL). The combined filtrate was concentrated under reduced pressure and to afford the free base form of the titled compound as white foam. To this foam (0.042 g, 0.041 mmol)
35 in EtOAc (5 mL) was added citric acid (0.016 g, 0.082 mmol) in MeOH (2 mL). The resulting mixture was stirred at room temperature for 3 hours and concentrated under reduced pressure to give a glassy residue. This residue was dissolved in minimum

amount of water (1 mL) and lyophilized to give the citrate salt (2 equivalents) of the titled compound as a white powder (0.043 g, 38 % in 3 steps).

¹H NMR (400 MHz, MeOD) δ ppm 8.61 - 8.73 (m, 2 H), 8.46 - 8.59 (m, 2 H), 8.19 - 8.30 (m, 2 H), 8.05 - 8.13 (m, 2 H), 7.87 - 7.97 (m, 2 H), 7.43 - 7.62 (m, 2 H), 7.21 - 7.34 (m, 4 H), 5.10 - 5.21 (m, 2 H), 4.51 - 4.65 (m, 2 H), 4.03 - 4.15 (m, 2 H), 3.89 - 3.97 (m, 2 H), 3.76 - 3.86 (m, 2 H), 3.48 - 3.72 (m, 4 H), 2.61 (s, 6 H), 2.41 - 2.53 (m, 2 H), 1.93 - 2.24 (m, 10 H), 1.41 - 1.52 (m, 6 H), 0.92 - 1.05 (m, 6 H), 0.85 - 0.92 (m, 6 H), citrate: 2.66-2.74 (m, 4 H), 2.75-2.82 (m, 4 H); LC-HRMS calculated for C₅₇H₇₃N₁₂O₆ 1021.5776, found 1021.5778 (ESI *m/e* [M + H⁺]); *t_R* 3.06 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

The following compounds were prepared using procedures analogous to those described above for the preparation of Example 2A with the appropriate starting materials.

Preparation of (S,S,S)-N,N'-(4,7,10,13,16-pentaoxanonadecane-1,19-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2B):



(2B)


The citrate salt (2 eq) of the titled compound was prepared as a white powder (0.037 g, 28% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl-
 5 (methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (I-2A-2d: 0.10 g, 0.17 mmol), 4,7,10,13,16-pentaoxanonadecane-1,19-diamine (0.025 g, 0.081 mmol) and HATU (0.068 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.65 - 8.80 (m, 2 H), 8.46 - 8.60 (m, 2 H), 8.14
 10 - 8.32 (m, 2 H), 8.06-8.13 (m, 2 H), 7.90 - 8.04 (m, 2 H), 7.43 - 7.65 (m, 2 H), 7.19 - 7.34 (m, 4 H), 5.12 - 5.24 (m, 2 H), 4.52 - 4.64 (m, 2 H), 4.01 - 4.15 (m, 2 H), 3.88 - 3.99 (m, 2 H), 3.78 - 3.88 (m, 2 H), 3.54 - 3.63 (m, 12 H), 3.41 - 3.51 (m, 12 H), 2.62 (s, 6 H), 2.40 - 2.52 (m, 2 H), 1.95 - 2.22 (m, 8 H), 1.81 - 1.94 (m, 4 H), 1.41 - 1.53 (m, 6 H), 0.92 - 1.05 (m, 6 H), 0.83 - 0.91 (m, 6 H), citrate: 2.68-2.76 (m, 4 H), 2.76-2.84 (m, 4 H); LC-HRMS

calculated for $C_{68}H_{95}N_{12}O_{11}$ 1255.7243, found 1255.7240 (ESI m/e $[M + H^+]$); t_R 3.20 minutes (Inertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

5

Preparation of (S,S,S)-N,N'-(3,3'-(2,2'-oxybis(ethane-2,1-diyl))bis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2C):



(2C)


10

The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.045 g, 36% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-

yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), 3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))dipropan-1-amine (0.018 g, 0.081 mmol) and HATU (0.068 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.63 - 8.77 (m, 2 H), 8.44 - 8.59 (m, 2 H), 8.10 - 8.27 (m, 2H), 8.00 - 8.07 (m, 2 H), 7.88 - 7.98 (m, 2 H), 7.42 - 7.63 (m, 2 H), 7.17 - 7.34 (m, 4 H), 5.10 - 5.22 (m, 2 H), 4.52 - 4.65 (m, 2 H), 4.02 - 4.16 (m, 2 H), 3.88 - 3.98 (m, 2 H), 3.77 - 3.87 (m, 2 H), 3.52 - 3.64 (m, 12 H), 3.43 - 3.50 (m, 4 H), 2.61 (s, 6 H), 2.40 - 2.51 (m, 2 H), 1.95 - 2.23 (m, 8 H), 1.79 - 1.92 (m, 4 H), 1.41 - 1.52 (m, 6 H), 0.92 - 1.05 (m, 6 H), 0.85 - 0.91 (m, 6 H), citrate: 2.68-2.76 (m, 4 H), 2.76-2.84 (m, 4 H); LC-HRMS calculated for C₆₄H₈₇N₁₂O₉ 1167.6719, found 1167.6741 (ESI *m/e* [M + H⁺]); *t_R* 3.16 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

15 Preparation of (S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2D):




(2D)

20 The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.052 g, 43% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-

carboxylic acid (I-2A-2d: 0.10 g, 0.17 mmol), 2,2'-(ethane-1,2-diylbis(oxy))diethanamine (0.012 g, 0.081 mmol) and HATU (0.068 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.59 - 8.72 (m, 2 H), 8.42 - 8.56 (m, 2 H), 8.07 - 8.25 (m, 2 H), 7.93 - 8.05 (m, 2 H), 7.81 - 7.94 (m, 2 H), 7.37 - 7.56 (m, 2 H), 7.13 - 7.30 (m, 4 H), 5.10 - 5.21 (m, 2 H), 4.54 - 4.70 (m, 2 H), 4.02 - 4.15 (m, 2 H), 3.84 - 3.96 (m, 2 H), 3.66 - 3.79 (m, 10 H), 3.43 - 3.66 (m, 4 H), 2.81 (s, 6 H), 2.25 - 2.50 (m, 2 H), 1.93 - 2.20 (m, 8 H), 1.36 (s, 6 H), 0.90 - 1.03 (m, 6 H), 0.84 - 0.90 (m, 6 H), citrate: 2.67-2.74 (m, 4 H), 2.75-2.88 (m, 4 H); LC-HRMS calculated for C₆₀H₇₉N₁₂O₈ 1095.6144, found 1095.6090 (ESI *m/e* [M + H⁺]); *t_R* 3.02 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 97% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2E):




The citrate salt (2.4 equivalents) of the titled compound was prepared as a white powder (0.647 g, 52% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (I-2A-2d: 0.929 g, 0.157 mmol), 3,3'-(ethane-1,2-diylbis(oxy))dipropan-1-amine (0.132 g, 0.748 mmol), HATU (0.625 g, 1.645 mmol) and HOBT (0.222 g, 1.645 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.63 - 8.80 (m, 2 H), 7.95-8.58 (m, 2 H), 8.07 - 8.30 (m, 2 H), 7.97 - 8.05 (m, 2 H), 7.80-7.95 (m, 2 H), 7.40- 7.65 (m, 2 H), 7.17 - 7.40 (m, 4 H), 5.12 - 5.60 (m, 2 H), 4.30-4.65 (m, 2 H), 4.00 - 4.20 (m, 2 H), 3.80-3.98 (m, 4

H), 3.60-3.70 (m, 4 H), 3.55-3.60 (m, 4 H), 3.38-3.50 (m, 4 H), 2.65 (s, 6 H), 2.35 - 2.55 (m, 2 H), 1.88 - 2.24 (m, 8 H), 1.78-1.89 (m, 4 H), 1.37 - 1.58 (m, 6 H), 0.75 - 1.32 (m, 12 H), citrate: 2.71-2.75 (m, 4.8 H), 2.80-2.84 (m, 4.8 H); LC-HRMS calculated for $C_{62}H_{83}N_{12}O_8$ 1123.6457, found 1123.6494 (ESI m/e $[M + H^+]$); t_R 3.35 minutes (100 X 3 mm 3um C18 column: mobile phase: 5-95% acetonitrile/water with 5 mM ammonium formate, at 1 mL/minute over 10 minutes); purity 100% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-(ethane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2F):




The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.059 g, 53% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), ethane-1,2-diamine (0.005 g, 0.081 mmol), HBTU (0.068 g, 0.18 mmol) and HOBT (0.027 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

1H NMR (400 MHz, MeOD) δ ppm 8.62 - 8.75 (m, 2 H), 8.45 - 8.60 (m, 2 H), 8.14 - 8.32 (m, 2 H), 8.04 - 8.14 (m, 2 H), 7.85 - 7.98 (m, 2 H), 7.42 - 7.62 (m, 2 H), 7.13 - 7.36 (m, 4 H), 5.10 - 5.24 (m, 2 H), 4.51 - 4.65 (m, 2 H), 4.01 - 4.16 (m, 2 H), 3.87 - 3.97 (m, 2 H), 3.77 - 3.87 (m, 2 H), 3.62 - 3.75 (m, 2 H), 2.62 (s, 6 H), 2.41 - 2.52 (m, 2 H),

1.90 - 2.24 (m, 8 H), 1.41 - 1.52 (m, 6 H), 0.91 - 1.05 (m, 6 H), 0.83 - 0.91 (m, 6 H),
citrate: 2.68-2.75 (m, 4 H), 2.76-2.84 (m, 4 H); LC-HRMS calculated for $C_{56}H_{71}N_{12}O_6$
1007.5620, found 1007.5570 (ESI m/e $[M + H]^+$); t_R 3.04 minutes (Insertsil ODS3, 100 X
3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1
5 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-(biphenyl-4,4'-diylbis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2G):




10

The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.12 g, 48% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.20 g, 0.34 mmol), biphenyl-4,4'-diyl dimethanamine (0.034 g, 0.16 mmol) and HATU (0.13 g, 0.35 mmol) following general procedure described above for the preparation of Example 2A.

1H NMR (400 MHz, MeOD) δ ppm 8.69-8.80 (m, 2 H), 8.40-8.51 (m, 2 H), 8.18 - 8.34 (m, 2 H), 8.11 (s, 2 H), 7.80-8.00 (m, 2 H), 7.59 (d, $J=8.0$ Hz, 5 H), 7.41 - 7.54 (m, 5
20 H), 7.25 - 7.37 (m, 4 H), 5.17-5.60 (m, 2 H), 4.65 (s, 4 H), 4.51 - 4.60 (m, 2 H), 4.06 (t, $J=6.5$ Hz, 2 H), 3.78 - 3.97 (m, 2 H), 3.56 - 3.74 (m, 2 H), 2.51 (s, 6 H), 2.40 - 2.50 (m, 2 H), 1.92 - 2.22 (m, 8 H), 1.40 (d, 6 H), 0.93 (d, $J=6.5$ Hz, 6 H), 0.87 (d, $J=7.0$ Hz, 6 H),
citrate: 2.65-2.72 (m, 4 H), 2.73-2.80 (m, 4 H); LC-HRMS calcd for $C_{68}H_{79}N_{12}O_6$

1159.6246, found 1159.6204 (ESI m/e $[M + H^+]$); t_R 3.60 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

- 5 Preparation of (S,S,S)-N,N'-(undecane-1,11-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2H):




(2H)

- 10 The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.088 g, 71% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), undecane-1,11-diamine (0.018 g, 0.097 mmol) and HATU (0.068 g, 0.18 mmol) following general procedure
- 15 described above for the preparation of Example 2A.

- 1H NMR (400 MHz, MeOD) δ ppm 8.65 - 8.78 (m, 2 H), 8.46 - 8.59 (m, 2 H), 8.13 - 8.28 (m, 2 H), 8.02 - 8.12 (m, 2 H), 7.90 - 7.98 (m, 2 H), 7.45 - 7.64 (m, 2 H), 7.21 - 7.36 (m, 4 H), 5.13 - 5.23 (m, 2 H), 4.51 - 4.65 (m, 2 H), 4.02 - 4.15 (m, 2 H), 3.89 - 3.97 (m, 2 H), 3.76 - 3.88 (m, 2 H), 3.37 - 3.45 (m, 4 H), 2.62 (s, 6 H), 2.40 - 2.55 (m, 2 H),
- 20 1.95 - 2.23 (m, 8 H), 1.58 - 1.72 (m, 4 H), 1.46 (d, $J=6.5$ Hz, 6 H), 1.30 - 1.54 (m, 14 H), 0.93 - 1.10 (m, 6 H), 0.81 - 0.90 (m, 6 H), citrate: 2.68-2.76 (m, 4 H), 2.76-2.85 (m, 4 H); LC-HRMS calculated for $C_{65}H_{89}N_{12}O_6$ 1133.7023, found 1133.6960 (ESI m/e $[M + H^+]$); t_R 3.73 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95%

acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 min); purity 98% by UV 254/214 nm.

Preparation of 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-
 5)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(3-(3-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-
(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-
carboxamido)propoxy)propyl)-1H-indole-3-carboxamide) (2I):




(2I)

10 The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.073 g, 62% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), 3,3'-oxydipropyl-1-amine (0.011 g, 0.081 mmol) and HATU (0.067 g, 0.18 mmol) following general procedure
 15 described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.59-8.75 (m, 2 H), 8.40-8.55 (ms, 2 H), 8.08 - 8.28 (m, 2 H), 8.05 (s, 2 H), 7.89 (s, 2 H), 7.43 (d, J=8.0 Hz, 2 H), 7.11 - 7.30 (m, 4 H), 5.13-5.55 (m, 2 H), 4.51 - 4.63 (m, 2 H), 4.01 - 4.15 (m, 2 H), 3.74 - 3.99 (m, 4 H), 3.47 - 3.69 (m, 8 H), 2.62 (s, 6 H), 2.38 - 2.51 (m, 2 H), 2.00 - 2.21 (m, 4 H), 1.85 - 2.01 (m, 8
 20 H), 1.45 (d, J=7.0 Hz, 6 H), 0.94 (d, J=7.0 Hz, 6 H), 0.89 (d, J=7.0 Hz, 6 H), citrate: 2.68-2.76 (m, 4 H), 2.76-2.84 (m, 4 H); LC-HRMS calculated for C₆₀H₇₉N₁₂O₇ 1079.6195, found 1079.6162 (ESI m/e [M + H⁺]); t_R 3.11 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 min); purity 99% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-(butane-1,4-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2J):



(2J)


5

The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.096 g, 36% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.26 g, 0.44 mmol), butane-1,4-diamine (0.026 g, 0.21 mmol) and HATU (0.18 g, 0.46 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.68-8.80 (m, 2 H), 8.40-8.60 (m, 2 H), 8.15-8.30 (m, 2 H), 8.09 (s, 2 H), 7.93 (s, 2 H), 7.44 - 7.62 (m, 2 H), 7.17 - 7.35 (m, 4 H), 5.18-5.60 (m, 2 H), 4.50 - 4.63 (m, 2 H), 4.00 - 4.14 (m, 2 H), 3.79 - 3.97 (m, 2 H), 3.60 - 3.74 (m, 2 H), 3.50 (br. s., 4 H), 2.53 (s, 6 H), 1.94 - 2.21 (m, 10 H), 1.80 (br. s., 4 H), 1.39 (d, *J*=6.5 Hz, 6 H), 0.94 (d, *J*=6.5 Hz, 6 H), 0.88 (d, *J*=6.5 Hz, 6 H), citrate: 2.65-2.73 (m, 4 H), 2.73-2.80 (m, 4 H); LC-HRMS calculated for C₅₈H₇₅N₁₂O₆ 1035.5933, found 1035.5942 (ESI *m/e* [M + H⁺]); *t_R* 3.13 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

20

Preparation of 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(5-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)pentyl)-1H-indole-3-carboxamide (2K):




(2K)

The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.070 g, 60% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), pentane-1,5-diamine (0.008 g, 0.081 mmol), HBTU (0.068 g, 0.18 mmol) and HOBT (0.027 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.64-8.80 (m, 2 H), 8.40-8.60 (m, 2 H), 8.10-8.30 (m, 2 H), 8.05 (s, 2 H), 7.91 (s, 2 H), 7.46-7.70 (m, 2 H), 7.07 - 7.35 (m, 4 H), 5.10 - 5.24 (m, 2 H), 4.51 - 4.64 (m, 2 H), 4.06 (t, *J*=6.5 Hz, 2 H), 3.78 - 3.97 (m, 4 H), 3.46-3.60 (m, 4 H), 2.62 (s, 6 H), 2.39 - 2.54 (m, 2 H), 1.93 - 2.22 (m, 8 H), 1.68 - 1.82 (m, 4 H), 1.50-1.70 (m, 2 H), 1.46 (d, *J*=6.5 Hz, 6 H), 0.95 (d, *J*=6.5 Hz, 6 H), 0.90 (d, *J*=7.0 Hz, 6 H), citrate: 2.68-2.76 (m, 4 H), 2.77-2.84 (m, 4 H); LC-HRMS calculated for C₅₉H₇₇N₁₂O₆ 1049.6089, found 1049.6039 (ESI *m/e* [*M* + *H*⁺]); *t_R* 3.13 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-(hexane-1,6-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2L):

5




The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.080 g, 34% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.20 g, 0.34 mmol), hexane-1,6-diamine (0.019 g, 0.16 mmol) and HATU (0.14 g, 0.36 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.68-8.85 (m, 2 H), 8.50-8.60 (m, 2 H), 8.18-8.30 (m, 2 H), 8.09 (s, 2 H), 7.85-8.00 (m, 2 H), 7.41 - 7.64 (m, 2 H), 7.14 - 7.34 (m, 4 H), 5.11 - 5.26 (m, 2 H), 4.58 (d, *J*=7.0 Hz, 2 H), 4.06-4.20 (m, 2 H), 3.79 - 3.97 (m, 2 H), 3.55-3.70 (m, 2 H), 3.50-3.40 (m, 4 H), 2.52 (s, 6 H), 2.42 - 2.50 (m, 2 H), 1.93 - 2.24 (m, 8 H), 1.70 (t, *J*=6.3 Hz, 4 H), 1.53 (br. s., 4 H), 1.38 (d, *J*=7.0 Hz, 6 H), 0.91 - 1.05 (m, 6 H), 0.88 (d, *J*=7.0 Hz, 6 H), citrate: 2.66-2.73 (m, 4 H), 2.74-2.81 (m, 4 H); LC-HRMS calculated for C₆₀H₇₉N₁₂O₆ 1063.6246, found 1063.6235 (ESI *m/e* [M + H⁺]); *t_R* 3.29 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95%

acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

5 Preparation of 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-
butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-((1S,4S)-4-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-
(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-
carboxamido)cyclohexyl)-1H-indole-3-carboxamide (2M):




(2M)

10 The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.041 g, 36% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), *trans*-cyclohexane-1,4-diamine (0.009 g, 0.081 mmol) and HATU (0.068 g, 0.18 mmol) in THF (2 mL) and

CH₂Cl₂ (2 mL) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, DMSO-*d*₆) δ ppm 8.68 - 8.80 (m, 2 H), 8.52 - 8.64 (m, 2 H), 8.20 - 8.38 (m, 2 H), 7.98 - 8.08 (m, 2 H), 7.87 - 7.99 (m, 2 H), 7.40 - 7.59 (m, 2 H), 7.16 - 7.32 (m, 4 H), 5.04 - 5.19 (m, 2 H), 4.38-4.70 (m, 2 H), 4.00 - 4.20 (m, 2 H), 3.90 - 3.98 (m, 4 H), 3.80 - 3.90 (m, 2 H), 2.65 (s, 6 H), 2.40-2.60 (m, 2 H), 1.89 - 2.30 (m, 12 H), 1.70 - 1.83 (m, 4 H), 1.23 - 1.37 (m, 6 H), 0.83 - 1.05 (m, 6 H), 0.71 - 0.81 (m, 6 H), citrate: 2.44-2.53 (m, 4 H), 2.53-2.62 (m, 4 H); LC-HRMS calculated for C₆₀H₇₇N₁₂O₆ 1061.6089, found 1061.6086 (ESI *m/e* [M + H⁺]); *t*_R 3.24 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

Preparation of (2*S*,2'*S*)-*N,N'*-((2*S*,2'*S*)-1,1'-((2*S*,2'*S*)-2,2'-(5,5'-(3,3'-(2,6-diazaspiro[3.3]heptane-2,6-diylbis(oxomethylene))bis(1*H*-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide) (2*N*):




The citrate salt (2 eq) of the titled compound was prepared as a white powder (0.076 g, 66% in three steps) from 1-(5-((*S*)-1-((*S*)-2-((*S*)-2-(tert-butoxycarbonyl-(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1*H*-indole-3-carboxylic acid (1-2A-2d: 0.10 g, 0.17 mmol), 2,6-diazaspiro[3.3]heptane (0.008 g, 0.08 mmol) and HATU (0.067 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

¹H NMR (400 MHz, MeOD) δ ppm 8.66 - 8.81 (m, 2 H), 8.42 - 8.61 (m, 2 H), 8.16 - 8.29 (m, 2 H), 7.90 - 8.11 (m, 4 H), 7.43 - 7.64 (m, 2 H), 7.19 - 7.37 (m, 4 H), 5.13 -

5.23 (m, 2 H), 4.65-4.85 (m, 4 H), 4.54 - 4.65 (m, 2 H), 4.33 - 4.52 (m, 4 H), 4.03 - 4.16 (m, 2 H), 3.88 - 4.00 (m, 2 H), 3.77 - 3.88 (m, 2 H), 2.62 (s, 6 H), 2.40 - 2.55 (m, 2 H), 1.92 - 2.25 (m, 8 H), 1.40 - 1.54 (m, 6 H), 0.93 - 1.08 (m, 6 H), 0.80 - 0.92 (m, 6 H), citrate: 2.68-2.77 (m, 4 H), 2.77-2.87 (m, 4 H); LC-HRMS calculated for $C_{59}H_{73}N_{12}O_6$ 1045.5776, found 1045.5781 (ESI m/e [$M + H^+$]); t_R 3.02 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 96% by UV 254/214 nm.

Preparation of (2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(piperazine-1,4-diylbis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide) (2O):




(2O)

The titled compound was prepared as a white powder (0.005 g, 6% in two steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)-propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (I-2A-2d: 0.10 g, 0.17 mmol), piperazine (0.007 g, 0.08 mmol) and HATU (0.068 g, 0.18 mmol) following general procedure described above for the preparation of Example 2A.

1H NMR (400 MHz, MeOD) δ ppm 8.69 - 8.82 (m, 2 H), 8.45 - 8.61 (m, 2 H), 7.96 - 8.10 (m, 2 H), 7.92 (s, 2 H), 7.75 - 7.88 (m, 2 H), 7.48 - 7.68 (m, 2 H), 7.19 - 7.39 (m, 4 H), 5.13 - 5.26 (m, 2 H), 4.51 - 4.66 (m, 2 H), 4.09-4.15 (m, 2 H), 3.65 - 4.01 (m, 12 H), 2.67 (s, 6 H), 2.42 - 2.57 (m, 2 H), 1.93 - 2.25 (m, 8 H), 1.49 (d, $J=7.1$ Hz, 6 H), 0.96 - 1.08 (m, 6 H), 0.91 (d, $J=6.6$ Hz, 6 H); LC-HRMS calculated for $C_{58}H_{73}N_{12}O_6$ 1033.5776, found 1033.5748 (ESI m/e [$M + H^+$]); t_R 3.11 minutes (Insertsil ODS3, 100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

Preparation of (S,S,S)-N,N'-(1,4-phenylenebis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide) (2P):




5

The citrate salt (2 equivalents) of the titled compound was prepared as a white powder (0.082 g, 33% in three steps) from 1-(5-((S)-1-((S)-2-((S)-2-(tert-butoxycarbonyl(methyl)amino)propanamido)-3-methylbutanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxylic acid (I-2A-2d: 0.10 g, 0.17 mmol), 1,4-phenylene-
 10 dimethanamine (0.023 g, 0.17 mmol) and HATU (0.14 g, 0.37 mmol) following general procedure described above for the preparation of Example 2A.


¹H NMR (400 MHz, MeOD) δ ppm 8.66 - 8.79 (m, 2 H), 8.47 - 8.62 (m, 2 H), 8.17 - 8.36 (m, 2 H), 8.08 - 8.14 (m, 2 H), 7.91 - 8.02 (m, 2 H), 7.45 - 7.68 (m, 2 H), 7.37 -
 15 7.45 (m, 4 H), 7.23 - 7.36 (m, 4 H), 5.18 (t, *J*=6.8 Hz, 2 H), 4.53-4.69 (m, 6 H), 4.02 - 4.14 (m, 2 H), 3.85 - 4.01 (m, 4 H), 2.66 (s, 6 H), 2.41 - 2.55 (m, 2 H), 1.95 - 2.22 (m, 8 H), 1.48 (d, *J*=7.0 Hz, 6 H), 0.95 (d, *J*=7.0 Hz, 6 H), 0.89 (d, *J*=6.5 Hz, 6 H), citrate: 2.74-2.85 (m, 4 H), 2.85-2.96 (m, 4 H); LC-HRMS calculated for C₆₂H₇₅N₁₂O₆ 1083.5933, found 1083.5977 (ESI *m/e* [*M* + *H*⁺]); *t_R* 3.25 minutes (Insertsil ODS3, 100 X 3 mm C18
 20 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 7.75 minutes); purity 100% by UV 254/214 nm.

Example 3

Preparation of (S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(trideuteromethylamino)propanamido)butanoyl)-pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide) (3A):




Preparation of Intermediate 1-[5-((S)-1-((S)-2-((S)-2-(tert-Butoxycarbonyl-D₃-methylamino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester (I-3A-3a):



The titled compound was prepared as yellow powder (0.600 g, 68% in two steps) from 1-[5-[(S)-1-((S)-2-*tert*-butoxycarbonylamino-3-methyl-butyl)-pyrrolidin-2-yl]-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester (**I-1A-1c**: 0.75 g, 1.40 mmol), CH₂Cl₂ (2 mL), and TFA (2 mL) following the general procedure described above for the preparation of Example 1A. To the resulting oily residue was added anhydrous THF (10 mL), DIPEA (0.6 mL), Boc-Me-d₃-alanine (0.26 g, 1.27 mmol), and DMTMM (0.41 g, 1.52 mmol) at 0°C. The mixture was stirred for 3 hours at 0°C, then worked-up following the general procedure described above for the preparation of Example 1A.

¹H NMR (400 MHz, dichloromethane-*d*) δ ppm 8.79 - 8.63 (m, 1 H), 8.60 - 8.44 (m, 1 H), 8.04 - 7.93 (m, 1 H), 7.81 - 7.63 (m, 2 H), 7.51 - 7.48 (m, 1 H), 7.42 - 7.38 (m, 1 H), 7.36 - 7.28 (m, 1 H), 6.86 - 6.45 (br, 1H), 5.31 - 5.18 (m, 1 H), 4.87 - 4.64 (m, 1 H), 4.48 (dd, *J*=7.0 Hz, 2 H), 4.17-4.09 (m, 1 H), 4.04-3.91 (m, 1 H), 3.87 - 3.77 (m, 1 H), 2.56 - 2.39 (m, 1 H), 2.21 - 1.83 (m, 4 H), 1.48 (s, 9 H), 1.38 - 1.30 (m, 3 H), 1.26 (t, *J*=7.0 Hz, 3 H), 0.97 d, *J*=7.0 Hz, 3 H), 0.82 (d, *J*=7.0 Hz, 3 H); LCMS calculated for C₃₄H₄₂D₃N₅O₆ 622.77, found 623.4 (ESI *m/e* [M + H⁺]).

Preparation of Intermediate 1-[5-((S)-1-{(S)-2-[{(S)-2-(tert-Butoxycarbonyl-D₃-methyl-amino)-propionylamino]-3-methyl-butyl-yl)-pyrrolidin-2-yl]-pyridin-3-yl]-1H-indole-4-carboxylic acid (I-3A-3b):



(1-3A-3b)

The titled compound was prepared as a white powder (0.420 g, 73%) from 1-[5-((S)-1-((S)-2-[(S)-2-(*tert*-butoxycarbonyl-D₃-methyl-amino)-propionylamino]-3-methylbutyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid ethyl ester (I-3A-3a: 0.60 g, 0.96 mmole), and LiOH (0.81 g, 19.27 mmole) following general procedure described above for the preparation of Example 1A.

¹H NMR (400 MHz, CD₃OD) δ ppm 8.80 - 8.64 (m, 1 H), 8.58 - 8.38 (m, 1 H), 8.00 - 7.85 (m, 2 H), 7.74 (d, J = 8.0 Hz, 1 H), 7.64 (d, J = 3.0 Hz, 1 H), 7.36 (d, J = 3.0 Hz, 1 H), 7.31 (dd, J = 8.0 Hz, 1 H), 5.56 - 5.10 (m, 1 H), 4.71 - 4.25 (m, 2 H), 4.16 - 3.61 (m, 2 H), 2.64 - 2.38 (m, 1 H), 2.24 - 1.85 (m, 4 H), 1.58 - 1.23 (m, 12 H), 1.10 - 0.80 (m, 6 H); LCMS calcd for C₃₂H₃₈D₃N₅O₆ 594.72, found 595.3 (ESI *m/e* [M + H⁺]).

Preparation of Example 3A:

To a solution of 1-[5-((S)-1-((S)-2-(*tert*-butoxycarbonyl-D₃-methyl-amino)-propionylamino]-3-methyl-butyryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-carboxylic acid (1-3A-3b: 0.42 g, 0.71 mmol) in DMF (4 mL) was added O-(7-azabenzotriazol-1-yl)-N,N,N', N'-tetramethyluronium hexafluorophosphate (HATU) (0.28 g, 0.74 mmol) and 1-hydroxybenzotriazole hydrate (HOBt · H₂O) and was stirred for 10 minutes. 3,3'-(Ethane-1,2-diylbis(oxy))dipropan-1-amine (0.059 g, 0.33 mmol) was added, followed by DIPEA (0.3 mL). The resulting mixture was stirred for three hours. The reaction mixture was quenched with water and extracted with EtOAc. The combined organic was washed with brine and dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to give an oil. To this oil was added CH₂Cl₂ (2 mL) and TFA (2 mL) following general procedure described above for the preparation of Example 1A to give the citrate salt (2.4

equivalents by ^1H NMR) of the titled compound as a white powder (0.25 g, 53% in three steps).

^1H NMR (400 MHz, MeOD) δ ppm 8.73 - 8.60 (m, 2 H), 8.54 - 8.39 (m, 2 H), 7.98 - 7.87 (m, 2 H), 7.75 - 7.55 (m, 4 H), 7.52 - 7.44 (m, 2 H), 7.32 - 7.21 (m, 2 H), 7.15 - 7.09 (m, 2 H), 5.55 - 5.10 (m, 2 H), 4.63 - 4.32 (m, 2 H), 4.16 - 3.99 (m, 2 H), 3.98 - 3.77 (m, 4 H), 3.63 (s, 4 H), 3.61 - 3.56 (m, 4 H), 3.52 - 3.44 (m, 4 H), 2.53 - 2.38 (m, 2 H), 2.22 - 1.91 (m, 8 H), 1.90 - 1.81 (m, 4 H), 1.55 - 1.43 (m, 6 H), 1.04 - 0.82 (m, 12 H), citrate: 2.87 - 2.68 (m, 9.6 H), LCHRMS calculated for $\text{C}_{62}\text{H}_{76}\text{D}_6\text{N}_{12}\text{O}_8$ 1129.6833, found 1129.6884 (ESI m/e $[\text{M} + \text{H}^+]$); t_R 3.13 minutes (100 X 3 mm C18 column: mobile phase: 5-95% acetonitrile/water with 0.1% formic acid, at 1 mL/minute over 10 minutes); purity 100% by UV 254 nm and 97% by UV 214 nm.

PHARMACOLOGICAL DATA

The compounds described herein above were profiled using a cellular assay (using Panc3.27 tumor cells) and a binding assay to determine the competition between the compounds of the present invention and smac7mer peptide for XIAP-BIR3 and clAP1-BIR3 binding groove occupancy.

Cellular Assay - Treatment of Panc3.27 tumor cells with dimeric IAP antagonists

On day one adherent Panc3.27 cells are plated into two 96-well, clear, flat bottom plates. All wells in row A contain 90uL of media. All wells in rows B-G contain a total volume of 90uL per well and 4000 cells per well for Panc3.27 cell lines. Plates are then incubated overnight for 18 hours at 37°C, 5% CO_2 .

On day two cells are treated with the compounds of formula M-L-M'. Treatments are done in triplicate. The compounds are first serially diluted in DMSO and then added to media giving a final concentration of 0.2% DMSO when added to cells. Cells are treated with 10uL of serially diluted compounds of formula M-L-M' at a final concentration of 1000nM, 200nM, 40nM, 8nM, 1.6nM, 0.32nM, 0.06nM, 0.013nM, 0.0026nM, and one untreated well. Plate two is used as a time zero plate.

To measure cell viability 50uL of Cell Titer Glo (CTG) solution is added to row A, media only and B, cells and media. CTG is purchased from Promega Corporation catalog number G7573. The solution is prepared according to manufacturer's instructions. CTG measures the amount of ATP released from viable cells that is proportional to the number of cells in each well. After incubating for ten minutes with CTG plates are measured on a luminescent reader at 700nm wavelength. Read time is approximately one second per well for a 96-well plate.

On day five 50uL of CTG is added to plate one, rows A-G, incubated for 10 minutes at room temperature and read on a luminescent reader. Raw data is adjusted to account for the time zero plate as well as background noise. Triplicate values are averaged and percent control growth is calculated. Percent control growth is calculated using the following logical test: If well read data point (a) is greater than time zero data point (t=0), then $100 * [(a) - (t=0)] / [(72\text{hour total growth}) - (t=0)]$, OR $100 * [(a) - (t=0)] / [(t=0)]$. Data is represented by line graph with the concentration of compound on the x axis and percent control growth on the y axis.

The results are presented in Table 1 below.

Binding Assay

The present method includes utility of a Surface plasmon resonance (SPR)-based biosensor (Biacore™, GE Healthcare, Uppsala, Sweden) to examine competition between the compounds of the present invention and smac7mer peptide for XIAP-BIR3 and cIAP1-BIR3 binding groove occupancy.

Biacore™ utilizes the phenomenon of surface plasmon resonance (SPR) to detect and measure binding interactions. In a typical Biacore experiment, one of the interacting molecules is immobilized on a flexible dextran matrix while the interacting partner is flowed over the derivatized surface. A binding interaction results in an increase in mass on the sensor surface and a corresponding direct change in the refractive index of the medium in the vicinity of the sensor surface. Changes in refractive index or signal are recorded in resonance units (R.U.) Signal changes due to association and dissociation of complexes are monitored in a non-invasive manner, continuously and in real-time, the results of which are reported in the form of a sensorgram.

Solution inhibition assay format:

Biacore™ T100 (GE Healthcare, Uppsala, Sweden) was used to conduct all experiments reported herein. Sensor surface preparation and interaction analyses were performed at 25°C. Buffer and Biacore reagents were purchased from GE Healthcare. Running buffer containing 10mM Hepes, pH7.4, 150mM sodium chloride, 1.25mM Dithiothreitol, 2% Dimethyl sulfoxide and 0.05% polysorbate 20 was utilized throughout all experiments.

Biotinylated smac7mer peptide was diluted to 10nM in running buffer and captured onto a sensor surface pre-derivatized with streptavidin (sensor chip SA) towards peptide surface densities in the range 40 – 100 R.U. Peptide captured surfaces were blocked with 500µM PEO₂-Biotin (Thermo Scientific). A blank flowcell was similarly blocked with PEO₂-biotin and served as a reference flowcell in the competition assay.

Interaction analyses were performed by first equilibrating each compound within a six point seven fold compound dilution series in the range 1 μ M to 0.06nM with either 100nM XIAP-BIR3 or 6nM cIAP1-BIR3 for at least one hour during instrument start-up procedures. Protein compound mixtures were then injected over reference and

5 smac7mer peptide surfaces in series for 60 seconds at a flow-rate of 60 μ L/min. Surface regeneration was performed at the end of each analysis cycle by a 30 second injection of 10mM Glycine, pH 2.5, 1M Sodium Chloride, 0.05% polysorbate 20. Additionally, control compound samples and control XIAP-BIR3 or cIAP1-BIR3 samples were prepared and run at regular intervals to monitor surface and assay performance.

10 Data analyses were carried out using Biacore™ T100 evaluation software v2.0 to validate assay quality. Binding level report points were plotted versus logarithmic compound concentration values and analyzed in Graphpad prism 5 *via* non-linear regression using a one-site competition model. EC50 values were generated and used as a measure of inhibitor potency.

15 The results are presented in Table 1 below.

Table 1

Ex No	Structural Name	XIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	CIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	PANC proliferation IC50 [nmol l ⁻¹]
1A	(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide), citrate	34.5 - 56.9	0.24 - 0.57	< 0.0026 - 0.83
1B	(S,S,S)-N,N'-(1,4-phenylenebis-(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide), trifluoroacetate	56.8 - 57.0	1.22 - 1.23	< 0.0026

Ex No	Structural Name	XIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	CIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	PANC proliferation IC50 [nmol l ⁻¹]
1C	(S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide), citrate	70.7 - 71.3	1.66 - 1.71	< 0.0026 - 0.14
1D	(S,S,S)-N,N'-((1S,2S)-cyclohexane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide), trifluoroacetate	76.0 - 76.7	1.20 - 1.22	2.61 - 2.75
2A	(S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	49.8 - 52.2	0.60 - 1.10	0.14 - 2.86
2B	(S,S,S)-N,N'-(4,7,10,13,16-pentaoxanonadecane-1,19-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	51.2 - 59.4	0.51 - 1.04	0.02 - 0.21
2C	(S,S,S)-N,N'-(3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))bis(propane-3,1-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	42.0 - 79.9	0.97 - 0.98	0.10 - 0.27

Ex No	Structural Name	XIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	CIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	PANC proliferation IC50 [nmol l ⁻¹]
2D	(S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	33.3 - 53.3	0.37 - 4.29	1.41 - 2.22
2E	(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	47.0 - 54.0	0.52 - 1.83	0.26 - 0.44
2F	(S,S,S)-N,N'-(ethane-1,2-diylbis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	23.5 - 55.2	0.21 - 0.86	0.08 - 0.28
2G	(S,S,S)-N,N'-(biphenyl-4,4'-diylbis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	47.9 - 51.6	2.24 - 2.27	< 0.0026
2H	(S,S,S)-N,N'-(undecane-1,11-diylbis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	60.3 - 60.8	1.00 - 1.02	< 0.0026

Ex No	Structural Name	XIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	CIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	PANC proliferation IC50 [nmol l ⁻¹]
2I	1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(3-(3-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)propoxy)propyl)-1H-indole-3-carboxamide), citrate	19.7 - 20.8	0.50 - 0.51	0.18 - 1.21
2J	(S,S,S)-N,N'-(butane-1,4-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	19.3 - 19.5	1.13 - 1.15	0.18 - 1.21
2K	1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(5-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)pentyl)-1H-indole-3-carboxamide, citrate	45.2 - 45.3	1.82 - 1.83	0.25 - 1.65
2L	(S,S,S)-N,N'-(hexane-1,6-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	25.5 - 25.8	1.71 - 1.73	< 0.0026

Ex No	Structural Name	XIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	CIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	PANC proliferation IC50 [nmol l ⁻¹]
2M	1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-((1S,4S)-4-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)cyclohexyl)-1H-indole-3-carboxamide, citrate	26.1 - 26.4	0.70 - 0.71	< 0.0026
2N	(2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(2,6-diazaspiro[3.3]heptane-2,6-diyl)bis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide), citrate	32.6 - 33.3	0.57 - 0.68	7.04 - 8.37
2O	(2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(piperazine-1,4-diyl)bis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide), trifluoroacetate	39.0	2.17 - 2.18	0.54 - 1.37
2P	(S,S,S)-N,N'-(1,4-phenylene-bis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide), citrate	70.9 - 71.0	1.21 - 1.23	< 0.0026

Ex No	Structural Name	XIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	CIAP-BIR3 Binding Biacore (Competitive) EC50 [nmol l ⁻¹]	PANC proliferation IC50 [nmol l ⁻¹]
3A	(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(trideuteromethylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide), citrate	44.7	0.59	0.12 - 0.17


In vivo combination studies were performed as follows. Cohorts of mice (n=8) were treated with (1) Cohort A: Example 1A (1.8 mg/kg, 1x per week), (2) Cohort B: Paclitaxel (12 mg/kg, 1x per week), or (3) Cohort C: a combination of Example 1A (1.8 mg/kg, 1x per week) + Paclitaxel (12 mg/kg, 1x per week). Studies typically initiated dosing when xenograft tumors were 200mm³ and lasted 2-3 weeks. Positive combination activity occurs when neither single agent treatment alone results in tumor regression (i.e. increase in tumor volume is not inhibited in treated animals) but the combination results in tumor regression (i.e. tumor volume decreases in treated animals).

For Cohort A tumor stasis was achieved and for Cohort B tumor growth control was >50%, while tumor regression was observed for Cohort C (combination of Example 1A with Paclitaxel).

CLAIMS

What is claimed is:

1. A compound of Formula M-L-M', wherein M and M' are each independently a
5 monomeric moiety of Formula (I)



(I)

wherein,





- R^1 is (C₁-C₄)alkyl, deuterated methyl, or hydrogen;
10 R^2 is (C₁-C₄)alkyl or hydrogen;
 R^3 is (C₁-C₄)alkyl or hydrogen, or
 R^1 or R^2 along with the nitrogen to which R^1 or R^2 is attached is taken together
with R^3 to form an aziridinyl, azetidiny, pyrrolidinyl, or piperidinyl;
 R^4 is
15 (i) (C₁-C₁₀)alkyl, (C₂-C₁₀)alkenyl, (C₂-C₁₀)alkynyl, (C₃-C₆)cycloalkyl, phenyl, a 3- to
7-membered heterocycle containing 1 to 3 heteroatoms each independently selected
from O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each
independently selected from O, N or S, or
(ii) R^{4a} -(C₁-C₆)alkylene, where R^{4a} is (C₃-C₆)cycloalkyl, phenyl, a 3- to 7-
20 membered heterocycle containing 1 to 3 heteroatoms each independently selected from
O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each
independently selected from O, N or S,
where said R^4 and said R^{4a} are optionally substituted with 1 to 3 substituents
selected from halo, hydroxyl, -SH, -CO₂H, (C₁-C₄)alkyl, halo-substituted(C₁-C₄)alkyl, (C₁-
25 C₄)alkoxy, (C₁-C₄)alkyl-S-, -SO₂-, -NH₂ or -NO₂, and where 1 of the ring members of said
cycloalkyl and said heterocycle moieties are optionally replaced with oxo or thione;
A, B, and D are CR⁵, and E is N,
A, B and E are CR⁵ and D is N,
A, D and E are CR⁵, and B is N,
30 B, D and E are CR⁵, and A is N,
A and B are both N, and D and E are both CR⁵,

A and E are both N, and B and D are both CR⁵, or


B and E are both N, and A and D are both CR⁵, where R⁵ are each independently selected from H, F, -CH₃ or -CF₃;

V is selected from moieties of Formula (1a), (1b), (1c), (1d), (1e), (1f), (1g), (1h),

5 (1i), (1j), or (1k)



10



where * indicates the point of attachment to L,

R^{6a} and R^{6b} are each independently H, (C₁-C₃)alkyl, Cl, or CN, where at least one of R^{6a} and R^{6b} is H or (C₁-C₃)alkyl;

W is a bond or (C₁-C₄)alkylene;

R^7 are each independently halo, -CH₃, -CF₃, -CN or -OCH₃;

p is 0, 1, 2, 3 or 4 for formula (1a) or (1e),

p is 0, 1, 2 or 3 for formula (1b), (1c), (1d), (1f), (1i) or (1j),

p is 0, 1 or 2 for formula (1g) or (1h);

L is -NR⁸-X¹-NR⁸-, where each R⁸ is independently selected from H, (C₁-C₄)alkyl, or halo-substituted(C₁-C₄)alkyl and X¹ is

(i) a bond,

(ii) (C₁-C₁₀)alkylene, (C₂-C₁₀)alkenylene, (C₂-C₁₀)alkynylene, ((C₁-C₁₀)alkylene)-(O(C₁-C₆)alkylene)_q-, or (C₁-C₁₀)alkylene-NH(C₁-C₆)alkylene, where q is 0, 1 or 2,

(iii) phenylene, naphthylene, fluorenylene, 9H-fluoren-9-onylene, 9,10-dihydroanthracenylenylene, anthracen-9,10-dionylene, a partially or fully saturated (C₃-C₈)cycloalkylene, a 5- to 7-membered heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S, or N, or a 5- to 10-membered heteroarylene containing 1 to 3 heteroatoms each independently selected from O, S or N, where said phenylene is optionally fused to a (C₅-C₆)cycloalkyl,

(iv) (phenylene)-G-(phenylene), where G is a bond, O, S, -NH-, -N=N-, -S=S-, -SO₂-, (C₁-C₆)alkylene, (C₂-C₆)alkenylene, (C₂-C₁₀)alkynylene, (C₃-C₆)cycloalkylene, a 5- to 6-membered heteroaryl containing 1 to 3 heteroatoms each independently selected from O, S or N, or a 5- to 6-membered partially or fully saturated heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S or N, and where said phenylene is optionally fused to a phenyl,

(v) ((C₁-C₆)alkylene)_r-Z¹-((C₁-C₆)alkylene)_s, or ((C₁-C₆)alkenylene)_r-Z¹-((C₁-C₆)alkenylene)_s, where r and s are each independently 0, 1, or 2; and Z¹ is -O-, -N=N-, (C₃-C₆)cycloalkylene, phenylene, bisphenylene, a 5- to 6-

- 5 membered partially or fully saturated heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S or N, or a 5- to-6-membered heteroarylene containing 1 to 3 heteroatoms each independently selected from O, S or N, where said heteroarylene and said heterocyclene are optionally fused to a phenyl, phenylene, a 5- to 6-membered partially or fully saturated heterocyclene containing 1 to 3 heteroatoms each independently selected from O, S or N, or a 5- to-6-membered heteroarylene containing 1 to 3 heteroatoms each independently selected from O, S or N, or
- 10 (vi) (C₁-C₂₀)alkylene or -NH-((C₁-C₂₀)alkylene)-NH-, where said alkylene contains 1 to 6 oxygen atoms interspersed within the alkylene chain and optionally 1 to 2 phenylene groups interspersed within the alkylene chain; or X¹ is optionally taken together with one or both R⁸ groups along with the nitrogen to which the R⁸ group is attached to form a 4- to 14-membered
- 15 heterocyclene, (4- to 6-membered heterocyclyl)-(C₁-C₆)alkylene-(4- to 6-membered heterocyclyl), or *bis*-(4- to 6-membered heterocyclene, where said heterocyclene and said heterocyclyl moieties optionally contain 1 to 3 additional heteroatoms selected from O, S and N, and X¹ and R⁸ are optionally substituted with oxo or 1 to 3 substituents each independently
- 20 selected from hydroxyl or (C₁-C₄)alkyl;
- where said group (ii) moieties of X¹ are each independently substituted with one or more fluoro atoms, or 1 to 2 substituents each independently selected from halo, oxo, amino, phenyl, naphthyl, (C₃-C₆) cycloalkyl, or 5- to 6-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, where said phenyl, said
- 25 cycloalkyl, and said heterocycle are optionally substituted with 1 to 3 substituents each independently selected from halo, (C₁-C₄)alkyl, or trifluoromethyl,
- where said group (iii) and (iv) moieties of X¹ are optionally substituted with 1 to 4 substituents each independently selected from (C₁-C₄)alkyl, (C₁-C₄)alkoxy, halo, amino, -OH, benzyl, or a fused 5- to 6-membered cycloalkyl, where said (C₁-C₄)alkyl, said (C₁-
- 30 C₄)alkoxy, and said fused cycloalkyl are optionally substituted with 1 to 3 substituents selected from halo, or (C₁-C₄)alkyl,
- where said group (v) moieties of X¹ are optionally substituted with 1 to 3 substituents each independently selected from halo, hydroxy, oxo, amino, (C₁-C₄)alkyl, (C₁-C₄)alkoxy, or phenyl;
- 35 or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1, wherein

R^1 is (C₁-C₄)alkyl, or deuterated methyl;

R^2 is hydrogen;

R^3 is (C₁-C₄)alkyl;

R^4 is







5 (i) (C₁-C₁₀)alkyl, (C₃-C₆)cycloalkyl, phenyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or

(ii) R^{4a} -(C₁-C₆)alkylene, where R^{4a} is (C₃-C₆)cycloalkyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each independently
10 selected from O, N or S,


where said R^4 and said R^{4a} are optionally substituted with 1 to 3 substituents selected from halo, or (C₁-C₄)alkoxy and where 1 of the ring members of said cycloalkyl and said heterocycle moieties are optionally replaced with oxo;

A, B, and D are CR⁵, and E is N, or A, B and E are CR⁵ and D is N, where R⁵ are
15 each independently selected from H, or F;

V is selected from moieties of Formula (1a), (1b), (1e), (1f), (1g), (1i), (1j), or (1k)



20



where * indicates the point of attachment to L,

R^{6a} and R^{6b} are H;

5 W is a bond or (C_1-C_4) alkylene;

R^7 is halo;

p is 0 or 1 for formula (1a) or (1e),

p is 0 or 1 for formula (1b), (1f), (1i) or (1j),

p is 0 or 1 for formula (1g);

10 L is $-NR^8-X^1-NR^8-$, where each R^8 is independently selected from H, (C_1-C_4) alkyl, or halo-substituted (C_1-C_4) alkyl and X^1 is

(i) a bond,

(ii) (C_1-C_{10}) alkylene, (C_2-C_{10}) alkynylene, or $((C_1-C_{10})$ alkylene)- $(O(C_1-C_6)$ alkylene) $_q-$, where q is 0, 1 or 2,

15 (iii) phenylene, naphthylene, or a fully saturated (C_3-C_8) cycloalkylene,

(iv) (phenylene)- G -(phenylene), where G is a bond, O, $-SO_2-$, (C_1-C_6) alkylene,

(v) $((C_1-C_6)$ alkylene) $_r-Z^1-((C_1-C_6)$ alkylene) $_s$, where r and s are each independently 0, 1, or 2; and Z^1 is $-O-$, or


(vi) (C_1-C_{20}) alkylene or $-NH-((C_1-C_{20})$ alkylene)- $NH-$, where said alkylene contains 1 to 6 oxygen atoms interspersed within the alkylene chain;

20 or X^1 is optionally taken together with one or both R^8 groups along with the nitrogen to which the R^8 group is attached to form a 4- to 14-membered heterocyclene;

or a pharmaceutically acceptable salt thereof.

25


3. The compound of Claim 1 or 2, wherein M and M' are the same and each are a monomeric moiety of Formula (II)



(II)

or a pharmaceutically acceptable salt thereof.

- 5 4. The compound of Claim 3 wherein
 R^1 is methyl or deuterated methyl;
 R^2 is H;
 R^3 is methyl;
 R^4 is isopropyl or cyclohexyl;
10 R^{6a} is H;
 p is 0;
 W is a bond;
 L is $-\text{NH}-(\text{CH}_2)_3-(\text{O}-\text{CH}_2\text{CH}_2)_4-\text{O}-(\text{CH}_2)_3-\text{NH}-$,
 $-\text{NH}-(\text{CH}_2)_3-(\text{O}-\text{CH}_2\text{CH}_2)_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_3-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$,
15 $-\text{NH}-(\text{CH}_2)_3-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-(\text{CH}_2)_2-\text{NH}-$,
 $-\text{NH}-(\text{CH}_2)_2-(\text{O}-\text{CH}_2\text{CH}_2)_2-\text{O}-(\text{CH}_2)_2-\text{NH}-$, $-\text{NH}-((\text{C}_1-\text{C}_{12})\text{alkylene})-\text{NH}-$,
 $-\text{NH}-\text{CH}_2-(\text{phenylene})-\text{CH}_2-\text{NH}-$, $-\text{NH}-\text{CH}_2-(\text{phenylene})-(\text{phenylene})-\text{CH}_2-\text{NH}-$,
 $-\text{NH}-(\text{cyclohexylene})-\text{NH}-$,



- 20 or a pharmaceutically acceptable salt thereof.

5. A compound selected from the group consisting of
 (S,S,S) -N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-
(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-
25 carboxamide);

(S,S,S)-N,N'-(4,7,10,13,16-pentaoxanonadecane-1,19-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(ethane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(biphenyl-4,4'-diylbis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

(S,S,S)-N,N'-(undecane-1,11-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(3-(3-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)-propoxy)propyl)-1H-indole-3-carboxamide;

(S,S,S)-N,N'-(butane-1,4-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(5-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)pentyl)-1H-indole-3-carboxamide;

(S,S,S)-N,N'-(hexane-1,6-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);

1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-((1S,4S)-4-(1-(5-((S)-1-((S)-3-

methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)cyclohexyl)-1H-indole-3-carboxamide;

- (2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(2,6-diazaspiro[3.3]heptane-2,6-diylbis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide);
- (2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(piperazine-1,4-diylbis(oxomethylene))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide); and
- (S,S,S)-N,N'-(1,4-phenylene-bis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide);
- or a pharmaceutically acceptable salt thereof.

6. A compound which is (S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy))bis(ethane-2,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.


7. A compound which is (S,S,S)-N,N'-(ethane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

8. A compound which is 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(3-(3-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)propoxy)propyl)-1H-indole-3-carboxamide; or a pharmaceutically acceptable salt thereof.

9. A compound which is (S,S,S)-N,N'-(butane-1,4-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.


10. A compound which is (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.

11. A compound which is (S,S,S)-N,N'-(4,7,10,13,16-pentaoxanonadecane-1,19-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.
- 5 12. A compound which is (S,S,S)-N,N'-(3,3'-(2,2'-oxybis(ethane-2,1-diyl)bis(oxy)))bis(propane-3,1-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.
- 10 13. A compound which is (S,S,S)-N,N'-(2,2'-(ethane-1,2-diylbis(oxy)))bis(ethane-2,1-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.
- 15 14. A compound which is (S,S,S)-N,N'-(hexane-1,6-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.
- 20 15. A compound which is (2S,2'S)-N,N'-((2S,2'S)-1,1'-((2S,2'S)-2,2'-(5,5'-(3,3'-(piperazine-1,4-diylbis(oxomethylene)))bis(1H-indole-3,1-diyl))bis(pyridine-5,3-diyl))bis(pyrrolidine-2,1-diyl))bis(3-methyl-1-oxobutane-2,1-diyl))bis(2-(methylamino)propanamide); or a pharmaceutically acceptable salt thereof.
- 25 16. A compound which is 1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-N-(5-(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamido)pentyl)-1H-indole-3-carboxamide; or a pharmaceutically acceptable salt thereof.
- 30 17. A compound which is (S,S,S)-N,N'-(1,4-phenylene-bis(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)-butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-3-carboxamide); or a pharmaceutically acceptable salt thereof.
- 35 18. The compound of Claim 1 or 2, wherein M and M' are the same and each are a monomeric moiety of Formula (III)



or a pharmaceutically acceptable salt thereof.

- 5 19. The compound of Claim 18 wherein
 R^1 is methyl or deuterated methyl;
 R^2 is H;
 R^3 is methyl;
 R^4 is isopropyl or cyclohexyl;
10 R^{6a} and R^{6b} are both H;
 p is 0;
 W is a bond; and
 L is $-\text{NH}-(\text{CH}_2)_3-(\text{O}-\text{CH}_2\text{CH}_2)_4-\text{O}-(\text{CH}_2)_3-\text{NH}-$,
 $-\text{NH}-(\text{CH}_2)_3-(\text{O}-\text{CH}_2\text{CH}_2)_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_3-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$,
15 $-\text{NH}-(\text{CH}_2)_3-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_2-\text{O}-\text{CH}_2\text{CH}_2-\text{O}-(\text{CH}_2)_2-\text{NH}-$,
 $-\text{NH}-(\text{CH}_2)_2-(\text{O}-\text{CH}_2\text{CH}_2)_2-\text{O}-(\text{CH}_2)_2-\text{NH}-$, $-\text{NH}-((\text{C}_1-\text{C}_{12})\text{alkylene})-\text{NH}-$,
 $-\text{NH}-\text{CH}_2-(\text{phenylene})-\text{CH}_2-\text{NH}-$, $-\text{NH}-\text{CH}_2-(\text{phenylene})-(\text{phenylene})-\text{CH}_2-\text{NH}-$,
 $-\text{NH}-(\text{cyclohexylene})-\text{NH}-$,



- 20 or a pharmaceutically acceptable salt thereof.

20. The compound of Claim 19 wherein L is $-\text{NH}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{CH}_2)_3-\text{O}-(\text{CH}_2)_2-\text{O}-(\text{CH}_2)_3-\text{NH}-$, $-\text{NH}-(\text{cyclohexylene})-\text{NH}-$, or $-\text{NH}-\text{CH}_2-(\text{phenylene})-\text{CH}_2-\text{NH}-$;
or a pharmaceutically acceptable salt thereof.

25

21. A compound selected from the group consisting of

(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

(S,S,S)-N,N'-(1,4-phenylenebis-(methylene))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

(S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

(S,S,S)-N,N'-((1S,2S)-cyclohexane-1,2-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); and

(S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(trideuteromethylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide);

or a pharmaceutically acceptable salt thereof.

22. A compound which is (S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)-propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

23. A compound which is (S,S,S)-N,N'-(3,3'-(ethane-1,2-diylbis(oxy))bis(propane-3,1-diyl))bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(trideuteromethylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

24. A compound which is (S,S,S)-N,N'-(propane-1,3-diyl)bis(1-(5-((S)-1-((S)-3-methyl-2-((S)-2-(methylamino)propanamido)butanoyl)pyrrolidin-2-yl)pyridin-3-yl)-1H-indole-4-carboxamide); or a pharmaceutically acceptable salt thereof.

25. A pharmaceutical composition comprising a compound of any one of the preceding claims, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable carrier, diluent or excipient.

26. The pharmaceutical composition of Claim 25 further comprising at least one additional pharmaceutical agent.

27. The pharmaceutical composition of Claim 26 wherein said at least one additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist.

5

28. The pharmaceutical composition of Claim 26 wherein said at least one additional pharmaceutical agent is paclitaxel.

29. A method for treating a disease, disorder, or condition associated with the overexpression of an IAP in a subject comprising the step of administering to a subject in need to such treatment a therapeutically effective amount of a compound according to Claims 1 through 24, or a pharmaceutically acceptable salt thereof.

30. A method for treating a disease, disorder, or condition mediated by IAPs comprising the step of administering to a subject in need of such treatment a therapeutically effective amount of a compound according to Claims 1 through 24, or a pharmaceutically acceptable salt thereof.

31. The use of a compound as defined in any one of Claims 1 through 24 for inducing or enhancing apoptosis in a tumor or cancer cell.

32. A compound according to any one of Claims 1 to 24, for use as a medicament.

33. Use of a compound according to any one of Claim 1 through 24 in the manufacture of a medicament for the treatment of a disease, disorder or condition mediated by IAPs.

34. Use of a compound according to any one of Claims 1 through 24 for the treatment of a disease, disorder or condition associated with the overexpression of an IAPs.

35. A method for treating a disease, disorder, or condition mediated by IAPs comprising the steps of administering to a patient in need of such treatment
(i) a compound according to claims 1 to 24 or a pharmaceutically acceptable salt thereof; and
(ii) at least one additional pharmaceutical agent.

36. The method of Claim 35 wherein said additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist.

5 37. The method of Claim 35 wherein said additional pharmaceutical agent is paclitaxel.

38. The method of Claim 35, 36 or 37 wherein said compound, or pharmaceutical acceptable salt thereof, and said additional pharmaceutical agent are administered
10 simultaneously.

39. The method of Claim 35, 36 or 37 wherein said compound, or pharmaceutical acceptable salt thereof, and said additional pharmaceutical agent are administered sequentially.

15

40. A method for treating a disease, disorder, or condition mediated by IAP comprising the step of administering to a patient in need of such treatment a pharmaceutical composition comprising a compound according to Claims 1 through 24, or a pharmaceutically acceptable salt thereof, and a pharmaceutical acceptable carrier.

20

41. The method of Claim 40 wherein said composition further comprises at least one additional pharmaceutical agent.

42. The method of Claim 41 wherein said additional pharmaceutical agent is
25 paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist.

43. The method of Claim 41 wherein said additional pharmaceutical agent is paclitaxel.

30

44. A method for treating a disease, disorder, or condition mediated by IAPs comprising the steps of administering to a patient in need of such treatment

(i) a first composition comprising a compound according to Claims 1 through 24, or a pharmaceutically acceptable salt thereof, and a pharmaceutical carrier; and

35 (ii) a second composition comprising at least one additional pharmaceutical agent and a pharmaceutical carrier.

45. The method of Claim 44 wherein said additional pharmaceutical agent is paclitaxel, a PI3K inhibitor, a topoisomerase inhibitor, a Trail antibody, recombinant Trail, or a Trail receptor agonist.


5 46. The method of Claim 44 wherein said additional pharmaceutical agent is a paclitaxel.

47. The method of Claim 44, 45 or 46 wherein said first composition and said second composition are administered simultaneously.

10

48. The method of Claim 44, 45 or 46 wherein said first composition and said second composition are administered sequentially.

49. A compound having Formula (I-1d) or (I-1e)



wherein

R is a carboxy-protecting group;

R¹ is (C₁-C₄)alkyl, or deuterated methyl;

20 R^2 is hydrogen or an amino-protecting group;

R³ is (C₁-C₄)alkyl;

 $\mathbb{R}^4$ is


(i) (C₁-C₁₀)alkyl, (C₃-C₆)cycloalkyl, phenyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or

25 (ii) R^{4a}-(C₁-C₆)alkylene, where R^{4a} is (C₃-C₆)cycloalkyl, or a 3- to 7-membered heterocycle containing 1 to 3 heteroatoms each independently selected from O, N or S, or a 5- to 9-membered heteroaryl containing 1 to 3 heteroatoms each independently selected from O, N or S.




where said R⁴ and said R^{4a} are optionally substituted with 1 to 3 substituents
30 selected from halo, or (C₁-C₄)alkoxy and where 1 of the ring members of said cycloalkyl
and said heterocycle moieties are optionally replaced with oxo;

A, B, and D are CR⁵, and E is N, or A, B and E are CR⁵ and D is N, where R⁵ are each independently selected from H, or F;

V is selected from moieties of Formula (1a), (1b), (1e), (1f), (1g), (1i), (1j), or (1k)



5



10

where * indicates the point of attachment to -OH or -OR,

R^{6a} and R^{6b} are H;

W is a bond or (C₁-C₄)alkylene;

R⁷ is halo;


p is 0 or 1 for formula (1a) or (1e),

p is 0 or 1 for formula (1b), (1f), (1i) or (1j), and

15


p is 0 or 1 for formula (1g).

50. The compound of Claim 49 wherein
 R^1 is methyl or deuterated methyl;
 5 R^2 is an amino-protecting group;
 R^3 is methyl;
 R^4 is isopropyl or cyclohexyl;
 R^{6a} is H;
 p is 0; and
 10 W is a bond.
51. The compound of Claim 49 or 50, wherein A, B, and D are CR^5 , and E is N, where R^5 are each independently selected from H, or F.
- 15 52. The compound of Claim 49 or 50, wherein A, B and E are CR^5 and D is N, where R^5 are each independently selected from H, or F.
53. The compound of Claim 49, 50, 51 or 52, wherein V is a moiety of Formula (1a)



(1a).

54. A compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid ethyl ester.
- 25 55. A compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-amino)-propionylamino]-3-methyl-butyl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-3-carboxylic acid.
- 30 56. The compound of Claim 49, 50, 51 or 52, wherein V is a moiety of Formula (1f)



(1f).

57. A compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-D₃-methyl-
5 amino)-propionylamino]-3-methyl-buteryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-
carboxylic acid ethyl ester.

58. A compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-D₃-methyl-
amino)-propionylamino]-3-methyl-buteryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-
10 carboxylic acid.

59. A compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-
amino)-propionylamino]-3-methyl-buteryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-
carboxylic acid ethyl ester.

15

60. A compound which is 1-[5-((S)-1-((S)-2-[(S)-2-(tert-Butoxycarbonyl-methyl-
amino)-propionylamino]-3-methyl-buteryl)-pyrrolidin-2-yl)-pyridin-3-yl]-1H-indole-4-
carboxylic acid.

20

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2011/072640

A. CLASSIFICATION OF SUBJECT MATTER

INV. C07D401/14 A61K31/4439 A61P35/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2008/045905 A1 (NOVARTIS AG [CH]; CHAREST MARK G [US]; CHEN CHRISTINE HIU-TUNG [US]; C) 17 April 2008 (2008-04-17) page 2, line 8 - page 5, line 4 -----	1-60
A	WO 2009/155709 A1 (AEGERA THERAPEUTICS INC [CA]; LAURENT ALAIN [CA]; PROULX MELANIE [CA];) 30 December 2009 (2009-12-30) paragraphs [0007], [0144] -----	1-60
A,P	WO 2011/104266 A1 (NOVARTIS AG [CH]; STRAUB CHRISTOPHER SEAN [US]; CHEN ZHUOLIANG [US]; P) 1 September 2011 (2011-09-01) page 1, line 34 - page 6, line 23 -----	1-60



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

13 January 2012

Date of mailing of the international search report

25/01/2012

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Uselli, Ambrogio

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2011/072640

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2008045905	A1	17-04-2008	
		AU 2007307763 A1	17-04-2008
		CA 2666112 A1	17-04-2008
		CN 101595121 A	02-12-2009
		EP 2102229 A1	23-09-2009
		JP 2010506847 A	04-03-2010
		KR 20090065548 A	22-06-2009
		RU 2009117701 A	20-11-2010
		US 2011015232 A1	20-01-2011
		WO 2008045905 A1	17-04-2008
WO 2009155709	A1	30-12-2009	
		AU 2009261919 A1	30-12-2009
		CA 2728933 A1	30-12-2009
		EP 2310402 A1	20-04-2011
		US 2011171171 A1	14-07-2011
		WO 2009155709 A1	30-12-2009
WO 2011104266	A1	01-09-2011	
		US 2011206690 A1	25-08-2011
		UY 33236 A	30-09-2011
		WO 2011104266 A1	01-09-2011