



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2583750 C 2015/11/24

(11)(21) **2 583 750**

(12) **BREVET CANADIEN
CANADIAN PATENT**

(13) **C**

(86) **Date de dépôt PCT/PCT Filing Date:** 2005/10/19
(87) **Date publication PCT/PCT Publication Date:** 2006/04/27
(45) **Date de délivrance/Issue Date:** 2015/11/24
(85) **Entrée phase nationale/National Entry:** 2007/04/16
(86) **N° demande PCT/PCT Application No.:** US 2005/037584
(87) **N° publication PCT/PCT Publication No.:** 2006/044962
(30) **Priorités/Priorities:** 2004/10/19 (US60/619,999);
2005/06/10 (US60/689,192)

(51) **Cl.Int./Int.Cl. A01K 67/027** (2006.01),
C12N 15/00 (2006.01), **C12N 15/87** (2006.01),
C12N 15/873 (2010.01), **C12N 5/073** (2010.01),
C12Q 1/68 (2006.01)

(72) **Inventeurs/Inventors:**
POUEYMIROU, WILLIAM, US;
DECHIARA, THOMAS M., US;
AUERBACH, WOJTEK, US;
FRENDEWEY, DAVID, US;
VALENZUELA, DAVID M., US

(73) **Propriétaire/Owner:**
REGENERON PHARMACEUTICALS, INC., US

(74) **Agent:** BLAKE, CASSELS & GRAYDON LLP

(54) **Titre : METHODE DE PRODUCTION D'UN ANIMAL HOMOZYGOTE POUR UNE MODIFICATION GENETIQUE**
(54) **Title: METHODS FOR GENERATING A RODENT HAVING A GENETIC MODIFICATION**

(57) **Abrégé/Abstract:**

Methods of generating modified embryos and mammals by introduction of donor cells into an early stage embryo are provided, such that the resulting embryo and animal generated therefrom has a significant or complete contribution to all tissues from the donor cells and is capable of transmitting the donor cell DNA.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 April 2006 (27.04.2006)

PCT

(10) International Publication Number
WO 2006/044962 A1

(51) International Patent Classification:

A01K 67/027 (2006.01) C12N 5/06 (2006.01)
C12N 15/87 (2006.01)

(74) Agent: VALETA, Gregg; Regeneron Pharmaceuticals, Inc., 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US).

(21) International Application Number:

PCT/US2005/037584

(22) International Filing Date: 19 October 2005 (19.10.2005)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/619,999 19 October 2004 (19.10.2004) US
60/689,192 10 June 2005 (10.06.2005) US

(71) Applicant (for all designated States except US): **REGENERON PHARMACEUTICALS, INC.** [US/US]; 777 Old Saw Mill River Road, Tarrytown, NY 10591 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **POUEYMIROU, William** [US/US]; 94 County Center road, White Plains, NY 10607 (US). **DECHIARA, Thomas, M.** [US/US]; 300 Croton Lake Road, Katonah, NY 10536 (US). **AUERBACH, Wojtek** [CA/US]; 320 South Pleasant Avenue, Ridgewood, NJ 07450 (US). **FRENDEWEY, David** [US/US]; 564 First Avenue, New York, NY 10016 (US). **VALENZUELA, David, M.** [US/US]; 529 Giordano Drive, Yorktown Heights, NY 10598 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: METHOD FOR GENERATING AN ANIMAL HOMOZYGOUS FOR A GENETIC MODIFICATION

(57) Abstract: Methods of generating modified embryos and mammals by introduction of donor cells into an early stage embryo are provided, such that the resulting embryo and animal generated therefrom has a significant or complete contribution to all tissues from the donor cells and is capable of transmitting the donor cell DNA.

WO 2006/044962 A1

METHODS FOR GENERATING A RODENT HAVING A GENETIC MODIFICATION

BACKGROUND

Field of the Invention

[0001] The invention is directed to improved methods for generating an animal homozygous for a genetic modification. More particularly, the present invention is directed to methods for increasing the efficiency of generating animals capable of transmitting a desired modification to subsequent offspring, for generating animals having high genetic material contribution from donor cells, and for increasing the efficiency of generating homozygous animals carrying the desired genetic modification.

Description of the Related Art

[0002] Methods for modifying eukaryotic donor cells are known in the art. See, for example, U.S. Patent No. 6,586,251. Methods for generating transgenic animals are also known in the art. A proposed method for improving the efficiency of generating animals capable of transmitting a desired modification in TT2 ES cells is described in Yagi et al. (1993) *Analytical Biochemistry* 214:70-76.

BRIEF SUMMARY OF THE INVENTION

[0003] The invention is based in part on the realization that introduction of a modified cell into an early stage host embryo substantially increases the genetic material and cellular contribution of the donor cells to the host embryo, such that the animal produced has increased fraction of cells from the donor and thus has an increased likelihood of transmitting the modification through the germline. Further, the methods of the invention allow a broader range of donor cells to be used than was practical with prior art methods. The methods of the instant invention thus reduce the number of animals and breedings necessary to generate an animal homozygous for a modification.

[0004] In a first aspect, the invention features a method for generating a modified mammal, comprising: (a) introducing a eukaryotic donor cell into an early stage embryo; and (b) introducing the embryo of (a) into a surrogate mother for gestation, wherein a modified mammal is generated. The early stage embryo is a pre-morula stage cell. In a more specific

embodiment, the early stage embryo is an 8-cell embryo. The early stage embryo may be derived from any strain. In one embodiment, an inbred embryo, e.g., C57BL/6, is used as host embryo. In another embodiment, an outbred embryo, e.g., Swiss Webster, is used as a host embryo.

[0005] The eukaryotic donor cell is a stem cell. More specifically, the stem cell is an embryonic stem (ES) cell or an ES-like cell. ES cells from any suitable ES cell line can be used in the method of the invention. In one embodiment, the ES cell is derived from an inbred strain, e.g., 129, C57BL/6, or BalbC. In another embodiment, the ES cell is derived from a hybrid strain, e.g., C57BL/6 x 129. In a specific embodiment, the cell is a mouse ES cell. However, the method of the invention may be practiced with cells derived from other mammals, e.g., a rat ES or ES-like cell. In one embodiment, the ES or ES-like cell is a modified cell comprising a genetic modification. In a more specific embodiment, the modification may arise spontaneously, may be random, or may result from experimental manipulation, for example, by the introduction of a foreign nucleic acid. A foreign nucleic acid may be introduced by any method known to the art, e.g., by homologous recombination.

[0006] In a second aspect, the invention features a method for generating a mammal homozygous for a genetic modification, comprising: (a) introducing a female eukaryotic donor cell into an early stage host embryo, wherein the eukaryotic donor cell is heterozygous for the genetic modification; (b) maintaining the embryo of (a) in culture for further development; (c) introducing the embryo of (b) into a surrogate mother for gestation, wherein a modified female mammal is generated; (a') introducing a male eukaryotic donor cell into an early stage host embryo, wherein the eukaryotic donor cell is heterozygous for the genetic modification; (b') maintaining the embryo of (a') in culture for further development; (c') introducing the embryo of (b') into a surrogate mother for gestation, wherein a modified male mammal is generated; and (d) breeding the sexually mature female and male mammal of (c and c'), wherein a mammal homozygous for the genetic modification is generated.

[0007] In one embodiment, the female eukaryotic donor cell of step (a) is derived from the same cell line of the male eukaryotic donor cell of step (a'). In a more specific embodiment, the female eukaryotic donor cell of step (a) is an XO cell and the modified female mammal of step (c) is an XO mammal.

[0008] The male or female eukaryotic donor cell is a stem cell, preferably a heterozygous ES or ES-like cell. In one embodiment, the heterozygous ES cell is generated by (i) obtaining a large cloned genomic fragment containing a DNA sequence of interest; (ii) using bacterial homologous recombination to genetically modify the large cloned genomic fragment of (i) to create a large targeting vector for use in the eukaryotic donor cells (LTVEC, or large targeting vector); (iii) introducing the LTVEC of (ii) into the eukaryotic donor cells to modify an endogenous gene or chromosomal locus in the eukaryotic donor cells; and (iv) using a quantitative assay to detect modification of allele in the eukaryotic donor cells of (iii) to identify those eukaryotic donor cells in which the endogenous gene or chromosomal locus has been genetically modified.

[0009] In various embodiments, the culture conditions of step (b) and/or (b') allow the embryo to develop to a post-morula stage or further. In one embodiment, the culture conditions of step (b) and/or (b') are provided by a growth factor-conditioned culture medium. In a preferred embodiment, the growth factor is a protein of the Wnt family. In one embodiment, the Wnt protein is added to the culture medium directly. In another embodiment, the Wnt protein is produced by mouse L-cell. In yet another embodiment, the culture medium further comprises a leukemia inhibitor factor (LIF).

[0010] In one specific embodiment, a blastocyst stage embryo of (b) and/or (b') is introduced into a surrogate mother for gestation and further development.

[0011] The eukaryotic donor cell may be introduced into an early stage embryo by any method known to the art. In one embodiment, the eukaryotic donor cell is introduced under the zona pellucida of the early stage embryo. Other methodologies include removal of the zona pellucida membrane. In a more specific embodiment, the eukaryotic donor cell may be introduced under the zona pellucida by injection or other methods that create an opening in the zona pellucida. In one specific embodiment, an opening is created in the zona pellucida by a laser. In one embodiment, the laser comprises an Infrared Solid State Diode laser. In another embodiment, the eukaryotic donor cell is introduced into the host embryo by cell fusion methods.

[0012] In one embodiment, a modified mammal comprising more than 90% donor cell-derived cells is bred in step (d). In a preferred embodiment, the modified mammal comprises more than 95% donor cell-derived cells. More preferably, the modified mammal comprises 100% donor cell-derived cells.

[0013] In a third aspect, the invention features a method of generating an embryo homozygous for a genetic modification, comprising: (a) introducing a female eukaryotic donor cell into an early stage host embryo, wherein the female eukaryotic donor cell is heterozygous for the genetic modification; (b) maintaining the embryo of (a) in culture for further development; (c) introducing the embryo of (b) into a surrogate mother for gestation, wherein a modified female mammal is generated; (a') introducing a male eukaryotic donor cell into an early stage host embryo, wherein the male eukaryotic donor cell is heterozygous for the genetic modification; (b') maintaining the embryo of (a') in culture for further development; (c') introducing the embryo of (b') into a surrogate mother for gestation, wherein a modified male mammal is generated; and (d) breeding the modified female and the modified male mammal of (c and c'), wherein an embryo homozygous for the genetic modification is generated.

[0014] In a fourth aspect, the invention features a method of generating a mammal homozygous for a genetic modification, comprising: (a) introducing a eukaryotic donor cell into an early stage embryo, wherein the eukaryotic donor cell is homozygous for the genetic modification; (b) maintaining the embryo of (a) in culture for further development; (c) introducing the embryo of (b) into a surrogate mother for gestation, wherein a mammal homozygous for the genetic modification is generated.

[0015] In one embodiment, the method for generating the eukaryotic donor cell homozygous for the genetic modification comprises a gene conversion, targeting both alleles of the same gene, or targeting either an X- or a Y-linked gene in a male ES cell.

[0016] In a fifth aspect, the invention features a method of generating an embryo homozygous for genetic modification, comprising: (a) introducing a eukaryotic donor cell into an early stage embryo, wherein the eukaryotic donor cell is homozygous for the genetic modification; and (b) culturing the early stage embryo of step (a), wherein an embryo homozygous for the genetic modification is generated.

[0017] In a sixth aspect, the invention features a method of increasing the relative contribution of genetic material from a donor eukaryotic donor cell to a host embryo, comprising: (a) introducing a eukaryotic donor cell into an early stage embryo; and (b) culturing the early stage embryo of step (a) under conditions that allow the embryo to develop to a post-morula stage or further.

[0018] In one embodiment, the culture conditions of step (b) allow the embryo to develop to a blastocyst. In another embodiment, the culture conditions of step (b) allow the embryo to develop to a gastrula or beyond. In various embodiments, the culture conditions of step (b) allow the embryo to develop to a post-morula stage or further. In one embodiment, the culture conditions of step (b) is provided by a growth factor-conditioned culture medium. In a preferred embodiment, the growth factor is a protein of the Wnt family. In one embodiment the Wnt protein is added to the culture medium directly. In another embodiment, the Wnt protein is produced by L-cells. In yet another embodiment, the culture medium further comprises a leukemia inhibitor factor (LIF).

[0019] In a seventh aspect, the invention features a method of generating a mammal having an increased relative contribution of genetic material from a donor eukaryotic donor cell to a host embryo, comprising: (a) introducing a eukaryotic donor cell into an early stage embryo; (b) culturing the early stage embryo of step (a) under conditions that allow the embryo to develop to a post-morula stage or further; and (c) introducing the embryo of (b) into a surrogate mother for gestation, wherein a mammal having an increased fraction of cells derived from the eukaryotic donor cell is generated.

[0020] In an eighth aspect, the invention features a method of generating a mammal capable of transmitting a genetic modification, comprising: (a) introducing a eukaryotic donor cell into an early stage embryo, wherein the eukaryotic donor cell carries a genetic modification; (b) culturing the early stage embryo of step (a) under conditions which allow the embryo to develop to a post-morula stage or further; and (c) introducing the embryo of (b) into a surrogate mother for gestation, wherein a mammal capable of transmitting the genetic modification is generated.

[0021] In a ninth aspect, the invention features a genetically modified mammal generated by a method of the invention. In a preferred embodiment, 90% or more of the cells of the mammal are donor cell-derived cells; more preferably, 95% or more of the cells of the mammal are donor cell-derived cells; even more preferably, 99% or more of the cells of the mammal are donor cell-derived cells. In one embodiment, 100% of the cells of the mammal are donor cell-derived cells.

[0022] In a tenth aspect, the invention features a genetically modified embryo generated by a method of the invention. In a preferred embodiment, 90% or more of the cells of the embryo are donor cell-derived cells; more preferably, 95% or more of the cells are donor cell-derived cells; even more preferably, 99% or more of the cells are donor cell-derived cells. In one embodiment, 100% of the cells of the embryo are donor cell-derived cells.

[0023] In an eleventh aspect, the invention features a culture medium for maintaining and/or growing an early stage embryo, wherein the culture medium comprises a growth factor.

[0024] In one embodiment, the growth factor is a protein of the Wnt family. The Wnt family protein is selected from the group consisting of Wnt1, Wnt3a, Wnt2, Wnt2b, Wnt3, Wnt4, Wnt5a, Wnt6, Wnt7a, Wnt8a, Wnt9a, Wnt9b, Wnt10a, Wnt10b, Wnt11, and Wnt16. In a preferred embodiment, the Wnt protein is Wnt3a. In one embodiment, the Wnt protein is a recombinant protein. In another embodiment, the culture medium is conditioned by a Wnt-producing mouse L-cell.

[0025] In one embodiment, the culture medium further comprises a maintenance agent. In one embodiment, the maintenance agent is a leukemia inhibitor factor (LIF).

[0025a] Thus, in one aspect, the invention provides a method for generating a rodent embryo homozygous for a genetic modification, comprising:

- (a) incorporating a rodent donor cell into a pre-morula embryo of a rodent, wherein the donor cell is an embryonic stem cell (ES cell) or an ES-like cell from an inbred rodent, and is homozygous for the genetic modification; and,
- (b) culturing the early stage host embryo of (a) to the blastocyst stage;

(c) introducing the embryo of (b) into a surrogate rodent mother for gestation, wherein a rodent embryo homozygous for the genetic modification is generated having at least 90% cellular contribution from the rodent donor cell.

[0025b] In another aspect, the invention provides a method for generating a rodent embryo heterozygous for a genetic modification, comprising:

- (a) incorporating a rodent donor cell into a pre-morula embryo of a rodent, wherein the rodent donor cell is an embryonic stem cell (ES cell) or an ES-like cell from an inbred rodent; and is heterozygous for the genetic modification;
- (b) culturing the early stage host embryo of (a) to the blastocyst stage;
- (c) gestating the embryo of (b) in a surrogate rodent mother for gestation, wherein a rodent embryo heterozygous for the genetic modification is obtained having at least 90% cellular contribution from the rodent donor cell.

[0025c] In another aspect, the invention provides a method for generating a mouse homozygous for a genetic modification, comprising:

- (a) introducing a female mouse embryonic stem (ES) donor cell into a pre-morula diploid mouse host embryo, wherein the ES donor cell is heterozygous for the genetic modification and is obtained from an inbred strain;
- (b) culturing the embryo of (a) to the blastocyst stage;
- (c) introducing the embryo of (b) into a surrogate mother for gestation, wherein a modified female mouse is generated having at least 90% cellular contribution from the female donor cell;
- (d) introducing a male mouse ES donor cell into a pre-morula diploid mouse host embryo, wherein the ES donor cell is heterozygous for the genetic modification and is obtained from an inbred strain;
- (e) culturing the embryo of (d) to the blastocyst stage;
- (f) introducing the embryo of (e) into a surrogate mother for gestation, wherein a modified male mouse is generated; and,
- (g) breeding a sexually mature modified male mouse and a sexually mature modified female mouse of (c and f), wherein a mouse homozygous for the genetic modification is generated.

[0025d] In another aspect, the invention provides a method for making a mouse embryo homozygous for a genetic modification, comprising:

- (a) incorporating a female mouse ES donor cell into a first pre-morula diploid mouse host embryo, wherein the female mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
- (b) culturing the first embryo to the blastocyst stage;
- (c) gestating the embryo of (b) in a surrogate mother, wherein upon gestation a modified female mouse is obtained having at least 90% cellular contribution from the female mouse ES donor cell;
- (d) incorporating a male mouse ES donor cell into a second pre-morula diploid mouse host embryo, wherein the male mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
- (e) culturing the embryo of (d) to the blastocyst stage;
- (f) gestating the embryo of (e) in a surrogate mother, wherein upon gestation a modified male mouse is obtained having at least 90% cellular contribution from the male mouse ES donor cell; and,
- (g) breeding the modified female mouse of (c) with the modified male mouse of (f) to obtain a mouse embryo homozygous for the genetic modification.

[0025e] In another aspect, the invention provides a method for making a mouse embryo heterozygous for a genetic modification, comprising:

- (a) incorporating a mouse ES donor cell into a pre-morula diploid mouse host embryo, wherein the mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
- (b) culturing the embryo to the blastocyst stage;
- (c) gestating the embryo of (b) in a surrogate mother, wherein upon gestation a modified mouse embryo is obtained having at least 90% cellular contribution from the mouse ES donor cell.

[0025f] In another aspect, the invention provides a method for making a mouse heterozygous for a genetic modification, comprising:

- (a) incorporating a mouse ES donor cell into a pre-morula diploid mouse host embryo, wherein the mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
- (b) culturing the embryo to the blastocyst stage;
- (c) gestating the embryo of (b) in a surrogate mother, wherein upon gestation a modified mouse is obtained having at least 90% cellular contribution from the mouse ES donor cell.

[0025g] In another aspect, the invention provides a method of generating a mouse wherein at least 90% of the cells in the mouse comprise a homozygous genetic modification, the method comprising:

- (a) incorporating a donor mouse embryonic stem (ES) cell from an inbred strain into a pre-morula diploid mouse host embryo, wherein the donor ES cell is homozygous for the genetic modification;
- (b) culturing the pre-morula host embryo of (a) to the blastocyst stage; introducing the embryo of (b) into a surrogate mother for gestation, and,
- (c) generating a mouse having at least a 90% cellular contribution from the donor ES cell, wherein at least 90% of the cells in the mouse are homozygous for the genetic modification.

[0025h] In another aspect, the invention provides method of generating a mouse embryo wherein at least 90% of the cells in the mouse embryo comprise a homozygous genetic modification, the method comprising:

- (a) incorporating a donor mouse ES cell from an inbred strain into a pre-morula diploid mouse host embryo, wherein the donor ES cell is homozygous for the genetic modification; and,
- (b) culturing the pre-morula host embryo of (a), wherein a mouse embryo comprising at least a 90% cellular contribution from the donor ES cell is generated and wherein at least 90% of the cells in the mouse embryo are homozygous for the genetic modification.

[0025i] In another aspect, the invention provides a method for making a mouse having a genetic modification, comprising:

- (a) incorporating a donor mouse ES cell comprising a genetic modification into a mouse host pre-morula stage embryo to obtain a mouse embryo comprising a donor ES cell;
- (b) growing the embryo in a medium conditioned by mouse fibroblasts that produce a protein that is a member of the Wnt family of proteins; and,
- (c) gestating the embryo comprising the donor ES cell in a surrogate mother to obtain a mouse progeny.

[0025j] In another aspect, the invention provides a method for generating a mouse embryo comprising cells that are homozygous for a genetic modification, comprising:

- (a) introducing mouse donor cells into a pre-morula host mouse embryo, wherein the donor cells comprise ES cells or ES-like cells from an inbred mouse, and are homozygous for the genetic modification; and
- (b) culturing the pre-morula host embryo of (a) to the blastocyst stage, wherein at least 90% of the cells of a mouse that develops from the blastocyst are derived from the donor cells.

[0025k] In another aspect, the invention provides a method for generating a mouse embryo comprising cells that are heterozygous for a genetic modification, comprising:

- (a) introducing mouse donor cells into a pre-morula host mouse embryo, wherein the mouse donor cells are ES cells or an ES-like cells from an inbred mouse; and are heterozygous for the genetic modification; and
- (b) culturing the pre-morula host embryo of (a) to the blastocyst stage, wherein at least 90% of the cells of a mouse that develops from the blastocyst are derived from the donor cells.

[0026] Other objects and advantages will become apparent from a review of the ensuing detailed description.

DETAILED DESCRIPTION

[0027] Before the methods, constructs and transgenic animals of the present invention are described, it is to be understood that this invention is not limited to particular methods, constructs, transgenic animals, and experimental conditions described, as such all may vary. It

is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0028] As used in this specification and in the appended claims, the singular forms “a”, “an” and “the” include plural references unless the context clearly dictates otherwise, e.g., “a cell” includes a plurality of cells. Thus, for example, a reference to “a method” includes one or more methods, and/or steps of the type described herein and/or which will become apparent to those persons skilled in the art upon reading this disclosure.

[0029] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, the preferred methods, constructs and materials are now described.

Definitions

[0030] The term “embryonic stem (ES)-like cell” includes a cell which, upon introduction into an embryo, can contribute to any tissue of the developing embryo.

[0031] The terms “increased contribution”, “higher relative percentage” and the like, include an improved genetic material contribution of a donor eukaryotic donor cell to an organism resulting from further development of a modified early stage host embryo. The method of the invention provides the means of increasing the probability that cells introduced into the host embryo contribute to all tissues, including germline tissues, of the generated animal.

[0032] By “gene knockout” as used herein is meant a genetic modification resulting from the disruption of the genetic information encoded at a chromosomal locus. By “gene knockin” as used herein is meant a genetic modification resulting from replacement of genetic information encoded at a chromosomal locus with a different DNA sequence or insertion of foreign genetic information at a chromosomal locus. By “knockout animal” as used herein is meant an animal in which a significant proportion of the animal’s cells harbor a gene knockout. By “knockin animal”

as used herein is meant an animal in which a significant proportion of the animal's cells harbor a genetic knockin.

General Description

[0033] One of the desired components of a transgenic animal study is generating a genetically modified transgenic animal capable of transmitting the genetic modification to progeny, i.e., a transgenic animal comprising the genetic modification in its germline. Current methods of creating such a transmission-capable transgenic animal tend to be inefficient in terms of resources and time expenditures. For example, to generate a genetically modified transgenic animal capable of transmitting the genetic modification to progeny, a modified ES cell heterozygous for a desired genetic modification is injected into a recipient blastocyst embryo, and the recipient embryo is implanted into a surrogate mother for gestation and birth of transgenic progeny. The resulting transgenic progeny are chimeric because some of the progeny's tissues are derived from the injected ES cells while other of the progeny's tissues are derived from the recipient embryo cells. Because of this chimerism, the injected ES cells comprising the genetic modification may or may not form germline tissues in the progeny and be capable of transmitting the genetic modification to the next generation. To determine whether a chimera is capable of transmitting the genetic modification, the chimera must be bred to another animal that does not carry the same genetic modification to establish if the desired modification is transmitted to the resulting progeny (F1). Detection of the desired genetic modification in the F1 progeny of the cross between the chimera and the other animal establishes that the chimera carries the desired genetic modification in its germline and is capable of transmitting the modification to its progeny (germline transmission). Typically, approximately 50% of chimeras exhibit germline transmission. Coat color is the most frequently used marker of the extent of ES cell contribution to the chimera and the transmission of ES cell genetic content to the F1 progeny.

[0034] The current need to generate an F1 generation to determine if the chimera is capable of transmitting the genetic modification is inefficient and costly in terms of time and the costs of breeding and maintaining F1 progeny. One method of improving the efficiency of the process for generating transgenic animals is provided by the instant invention which allows introduction into pre-morula embryos of cells that generate animals having an increased genetic material contribution of the exogenous cells relative to the results obtained when donor cells are

introduced into later stage embryos, e.g., blastocysts. As a result, a much higher percentage of the chimeras are germline transmitters. In some instances, about 100% of the chimeras are germline transmitters and thus these chimeras can transmit ES cell materials to their offspring.

[0035] Introducing donor cells into early stage host embryos, e.g., 8-cell embryos, provides several important benefits over the current methods, which teach the use of later stage host embryos, e.g., blastocysts. As shown in Example 1 below, the number of early stage embryos harvested from a donor mother (e.g., female BL/6 mouse) is higher than the number of later stage embryos harvested. Thus, fewer pregnant female mice are needed as donors, decreasing the cost of obtaining and maintaining pregnant female mice.

[0036] Further, as shown below, donor cells may be introduced into a smaller number of early stage host embryos than later stage embryos to generate the same number of chimeric animals, reducing the time and cost of introducing donor cells into embryos, e.g., when introduction is by microinjection, the amount of time spent on microinjections is greatly reduced.

[0037] The instant invention also allows culturing the donor cell-containing host embryo to post-morula stages, e.g., to a blastocyst stage or a gastrula stage, before being introduced into a surrogate mother for gestation. Since *in vitro* culturing conditions are more favorable for the donor cells than for the host embryo, the resulting embryo has a higher content of donor cell-derived cells compared to the method by which the morula stage host embryo is introduced into a surrogate mother for gestation.

[0038] An important improvement provided by the instant invention is that the number of animals generated that are capable of transmitting donor DNA is increased substantially with the use of early stage host embryos, such that an entire generation of breeding is eliminated. This is a significant practical improvement with important commercial implications.

[0039] A method known in the art that allows an entire generation of breeding to be eliminated employs tetraploid embryos as recipients of the modified donor cells. As the tetraploid cells of the recipient embryo are incapable of contributing to the tissue of the developing animal, the animals that are born are completely derived from the modified donor cells. If the resulting animals do not have a genetic modification that affects fertility, they will all be able to transmit

the desired genetic modification to their progeny upon mating. This process is laborious and inefficient, producing only a small fraction of live birth from hybrid ES cell lines. Injections of cells under the zona pellucida of pre-morula stage diploid embryos produces increased survival and generation of larger number of completely or almost completely ES cell derived live animals. Both male and females can be obtained by this method.

[0040] The method of the invention can be applied to introduce inbred ES cells into outbred recipient embryos through microinjection.

[0041] Other features of the invention will become apparent in the course of the following descriptions of exemplary embodiments which are given for illustration of the invention and are not intended to be limiting thereof.

EXAMPLES

[0042] The following examples are put forth so as to provide those of ordinary skill in the art with examples of how to make and use the methods, compositions and animals of the invention, and are not intended to limit the scope of the invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperature, etc.) but some experimental deviations are to be expected as is known to one of skill in the art. Unless indicated otherwise, parts are parts by weight, molecular weight is average molecular weight, temperature is in degrees Centigrade, and pressure is at or near atmospheric.

Example 1. Generation of Germline Transmitting Animals

[0043] Mouse embryos for injection were obtained from natural matings where the morning of the plug was designated as 0.5 days p.c. (post coitus). Oviducts were collected from plugged females at 2.5 days p.c. and were flushed with Dulbecco's medium to harvest the 8-cell embryos. The embryos were maintained in this medium for culture (37°C in a 5% CO₂ incubator) and microinjection procedures.

[0044] The injection of the 8-cell embryos was performed using standard microinjection techniques except that prior to introduction of the ES cells, a hole was generated in the zona pellucida of the 8-cell embryo using an XY Clone laser system according to the manufacturer's

specifications. The injection needle is inserted through this hole and 8 to 10 ES cells are deposited into the perivitelline space.

[0045] The injected embryos were transferred to a culture dish KSOM+AA and cultured overnight to the blastocyst stage. The surviving embryos were transferred as blastocysts to surrogate females in the afternoon of the next day (3.5 days p.c.).

Example 2. Generation of Homozygous DLL4 Knockout Mouse.

[0046] Homologous DLL4 knockout ES cells were microinjected into an 8-cell mouse embryo as described above. Injected embryos were cultured to blastocyst stage and were transferred to a surrogate female for gestation. All DLL4 knockout embryos died during gestation. The cause of death was identical to that observed for embryos produced by conventional breeding of heterozygous mice. Observation of the null DLL4 phenotype in the F0 generation avoided the two generations of breeding that would normally be required to generate DLL4 null mice.

Example 3. Generation of Highly and Fully ES Cell-Derived F0 Mice with Genetic Modifications

[0047] Genetically modified male F1H4 ES cells were microinjected into either 8-cell or blastocyst stage C57BL/6 embryos. The microinjected blastocyst embryos were transferred to a surrogate female for gestation immediately after the injection. The microinjected 8-cell embryos were further cultured in KSOM+AA culture medium to blastocyst stage and were then transferred to a surrogate female for gestation. The percentages of ES cell contribution were estimated by the coat color of the male F0 mice. Results are summarized in Table 1. As shown, when genetically modified ES cells were microinjected into 8-cell embryos, all of the F0 mice are ES cell-derived. On the other hand, when genetically modified ES cells were microinjected into blastocyst stage embryos, none of the male F0 mice were fully ES cell-derived and only about half of the male F0 mice (2 out of 4 and 4 out of 7 for 494B-F5 and 1218X-E2, respectively) have more than 90% ES-derived cells.

Table 1

ES Cell Line	Embryo		No. of Pups		No. of Male F0 Mice with Various Percentage of ES Cell Contribution*				
	Stage	No.	Total	Male	<50%	50% to 80%	90%	>90%	100%
494B-F5	8-Cell	33	11	11	0	0	0	0	11
494B-F5	Blastocyst	20	5	4	0	0	2	2	0
1218X-E2	8-Cell	50	6	6	0	0	0	0	6
1218X-E2	Blastocyst	36	7	4	0	0	0	4	0

* Percentages of ES cell contribution were estimated by the coat color of F0 mice.

[0048] More ES cell lines were used to test the effect of the host embryo stage (8-cell vs. blastocyst) on the ES cell contribution in the resulting male F0 mice. Similar results were obtained and are summarized in Table 2 (* Percentages of ES cell contribution were estimated by the coat color of F0 mice).

Table 2

ES Cell Line	Embryo		No. of Pups		No. of Male F0 Mice with Various Percentage of ES Cell Contribution*				
	Stage	No.	Total	Male	<50%	50% to 80%	90%	>90%	100%
F1H4 Parental	8-Cell	120	34	30	3	0	0	0	27
F1H4 Parental	Blastocyst	125	24	20	5	1	2	12	0
609B-G2	8-Cell	50	17	15	2	0	0	0	13
609B-G2	Blastocyst	50	21	16	1	0	1	14	0
698B-A8	8-Cell	25	5	4	1	0	0	1	2
698B-A8	Blastocyst	25	9	6	3	3	0	0	0
639C-A9	8-Cell	37	12	11	1	0	0	0	10
639C-A9	Blastocyst	25	7	7	0	0	0	7	0
619A-A1	8-Cell	25	12	11	1	0	0	0	10
619A-A1	Blastocyst	25	7	6	0	0	0	6	0
576A-E11	8-Cell	25	12	10	1	0	0	0	9

576A-E11	Blastocyst	25	21	16	2	1	1	12	0
----------	------------	----	----	----	---	---	---	----	---

Example 4. Generation of Highly and Fully ES Cell-Derived Female F0 Mice

[0049] Genetically modified XO ES cells (648B-H12 clone) were microinjected into either 8-cell or blastocyst stage C57BL/6 embryos. The microinjected blastocyst embryos were transferred to a surrogate female for gestation immediately after the injection. The microinjected 8-cell embryos were further cultured in KSOM+AA culture medium to blastocyst stage and were then transferred to a surrogate female for gestation. The percentages of ES cell contribution in the female F0 mice were estimated by their coat color. As shown in Table 3, when genetically modified ES XO ES cells were microinjected into 8-cell embryos, all of the chimeras were females. When genetically modified ES cells were microinjected into blastocyst stage embryos, only 7 out of a total of 16 for 648B-H12 and 11 out of 16 for 648C-H1 F0 mice were female (* Percentages of ES cell contribution were estimated by the coat color of F0 mice).

Table 3

XO Clone	SW Embryo		Pups	No. of Chimera		No. of Female F0 mice with Various Percentage of ES Cell Contribution*		
	Stage	No.		Total	Female	<90%	>90%	100%
648B-H12	8-Cell	50	13	13	13	0	0	13
648B-H12	Blastocyst	50	28	16	7	1	6	0
648C-H1	8-Cell	50	6	6	6	0	0	6
648C-H1	Blastocyst	50	27	16	11	1	10	0

Example 5. Generation of Highly and Fully ES Cell-Derived F0 Mice Using Outbred and Inbred Host Embryos

[0050] Unmodified male inbred (C57BL/6) or hybrid (F1H4) ES cells were microinjected into either 8-cell stage or blastocyst stage Swiss Webster (SW) embryos. The microinjected blastocyst embryos were transferred to a surrogate female for gestation immediately after the injection. The microinjected 8-cell embryos were further cultured in KSOM+AA culture medium to blastocyst stage and were then transferred to a surrogate female for gestation. Results are

summarized in Table 4 (* Percentages of ES cell contribution were estimated by the coat color of F0 mice).

Table 4

	SW Embryo		No. of Pups		No. of Male F0 Mice with Various Percentage of ES Cell Contribution*		
ES Cells	Stage	No.	Total	Male	<90%	>90%	100%
F1H4	8-Cell	140	46	46	34	4	8
F1H4	Blastocyst	50	11	11	11	0	0
C57/BL6.2	8-Cell	75	19	6	0	0	6
C57/BL6.2	Blastocyst	10	7	6	6	0	0

[0051] Similar results were obtained when male inbred 129 (CJ7) and Balb/c ES cells were microinjected into either 8-cell or blastocyst stage inbred C56BL/6 host embryos. Results are summarized in Table 5 (* Percentages of ES cell contribution were estimated by the coat color of F0 mice).

Table 5

	C56BL/6 Embryo		No. of Pups		No. of Male F0 Mice with Various Percentage of ES Cell Contribution*		
ES Cells	Stage	No.	Total	Male	<90%	>90%	100%
129 (CJ7)	8-Cell	57	8	8	3	0	5
129 (CJ7)	Blastocyst	50	18	14	4	10	0
Balb/c	8-Cell	50	11	11	5	0	6
Balb/c	Blastocyst	57	8	1	1	0	0

Example 6. Highly and Fully ES Cell-Derived F0 Mice are Germline Transmission Competent

[0052] F1H4-derived male F0 mice having more than 90% cells derived from the ES cells (more than 90% ES cell contribution) were used to test germline transmission competency. Using male ES cells, sexually mature male F0 mice were bred with sexually mature females. The coat color of the offspring was used as a marker for the evaluation of germline transmission

competency. Results are summarized on Table 6. When 8-cell stage embryos were used as host embryos, over 95% (51 out of 53) of the F0 males generated by 8-cell microinjection exhibited 100% germline transmission competency. When blastocyst stage embryos were used as host embryos, only about 64% (57 out of 89) of the F0 males exhibited 100% germline transmission competency. In addition, about 21% (19 out of 89) of the F0 males generated by blastocyst microinjection did not produce any offspring compared to less than 4% (2 out of 53) of the F0 males generated by 8-cell microinjection with no offspring (**Percentages of germline transmission competency were estimated by the coat color of the corresponding offspring).

Table 6

Stage of Host Embryo	No. of Male F0 Bred	No. of Male F0 Shown Various Percentage of Germline Transmission Competency**			No. of Male F0 with No Offspring
		100%	Partial	Zero	
8-Cell	53	51	0	0	2
Blastocyst	89	57	9	4	19

Example 7. Effect of Post-Microinjection Culture Medium on the Quality of F0 Mice

[0053] Modified male C57BL/6 and F1H4 ES cells were microinjected into 8-cell stage Swiss Webster (SW) embryos. The microinjected embryos were cultured in different culture media to blastocyst stage and were then transferred to a surrogate female for gestation. Three different culture media were used: (1) KSOM, mouse embryo culture medium; (2) LIF, ES cell culture medium containing LIF; (3) Wnt3, Wnt3 conditioned ES cell culture medium. Results are summarized in Table 7. Wnt3 conditioned ES medium was produced as following: (i) mouse L-cells were plated into a T75 flask in medium made of high glucose DMEM, 10% FBS, and L-glutamine, and were incubated at 37°C, with 5% CO₂; (ii) when cell density reached 100% confluency, 10% of the cells were re-plated into another T75 flask; (iii) the culture medium was collected until cell density reached confluency again (about 3 days after re-plating); and finally, (iv) the collected culture medium was mixed with equal volume of ES cell culture medium without LIF but with serum replacement (* Percentages of ES cell contribution were estimated by the coat color of F0 mice).

Table 7

ES Cells	Culture Medium	No. Embryo Transferred	No. of Pups		No. of Male F0 Mice with Various Percentage of ES Cell Contribution*	
			Total	Male	<100%	100%
F1H4	KSOM	140	55	45	37	8
F1H4	LIF	40	8	8	3	5
F1H4	Wnt3	40	9	9	2	7
C57BL/6.2	KSOM	75	16	16	10	6
C57BL/6.2	LIF	38	10	10	5	5
C57BL/6.2	Wnt3	38	11	11	0	11

[0054] Similar results were obtained when male modified F1H4 cells were microinjected into 8-cell C57BL/6 ES embryos.

Example 8. Generation of Animals Homozygous for a Genetic Modification from Heterozygous ES cells.

[0055] Male ES cells heterozygous for a desired genetic modification are microinjected into an 8-cell mouse embryo as described in Example 1. Female ES cells derived from the same male ES cell line and heterozygous for the same genetic modification are microinjected into another 8-cell mouse embryo as described above. Both embryos are cultured to the blastocyst stage and are transferred to a surrogate female for gestation. The resulting germline transmitting male and female F0 mice are bred to obtain progeny homozygous for the desired genetic modification. Two pairs of mice were bred. A total of 39 pups were born and 9 of these pups were homozygous for the genetic modification.

WE CLAIM:

1. A method for generating a rodent homozygous for a genetic modification, comprising:
 - (a) incorporating a rodent donor cell into a pre-morula host embryo of a rodent, wherein the donor cell is an embryonic stem cell (ES cell) or an ES-like cell from an inbred rodent, and is homozygous for the genetic modification;
 - (b) culturing the pre-morula host embryo of (a) to the blastocyst stage; and,
 - (c) introducing the embryo of (b) into a surrogate rodent mother for gestation,wherein the rodent homozygous for the genetic modification is generated having at least 90% cellular contribution from the rodent donor cell.
2. The method of claim 1, wherein the rodent donor cell homozygous for the genetic modification is generated by a process selected from the group consisting of gene conversion, targeting both alleles of the same gene, and targeting either an X- or a Y-linked gene in the ES cell.
3. A method for generating a rodent heterozygous for a genetic modification, comprising:
 - (a) incorporating a rodent donor cell into a pre-morula host embryo of a rodent, wherein the rodent donor cell is an embryonic stem cell (ES cell) or an ES-like cell from an inbred rodent; and is heterozygous for the genetic modification;
 - (b) culturing the pre-morula embryo of (a) to the blastocyst stage;
 - (c) gestating the embryo of (b) in a surrogate rodent mother for gestation,wherein the rodent heterozygous for the genetic modification is obtained having at least 90% cellular contribution from the rodent donor cell.
4. The method according to any one of claims 1 to 4, wherein the pre-morula host embryo of the rodent is a diploid embryo.
5. The method according to any one of claims 1 to 4, wherein the pre-morula host embryo is from an inbred strain.
6. The method according to any one of claims 1 to 4, wherein the pre-morula host embryo is from an outbred strain.

7. The method according to any one of claims 1 to 6, wherein the pre-morula host embryo is an 8-cell stage embryo.
8. The method according to any one of claims 1 to 7, wherein the pre-morula host embryo comprises a zona pellucida, and wherein the rodent donor cell is introduced into the host embryo through an opening in the zona pellucida.
9. The method according to claim 8, wherein the opening is created by a laser.
10. The method according to any one of claims 1 to 9, wherein, prior to step (a), the rodent donor cell is generated by:
 - (i) obtaining a cloned genomic fragment containing a DNA sequence of interest;
 - (ii) using bacterial homologous recombination to genetically modify the cloned genomic fragment of (i) to create a targeting vector (LTVEC) for use in the rodent donor cells;
 - (iii) introducing the LTVEC of (ii) into the rodent donor cells to modify an endogenous gene or a chromosomal locus in the cells; and
 - (iv) using a quantitative assay to detect modification of allele in the rodent donor cells of (iii) to identify those rodent donor cells in which the endogenous gene or the chromosomal locus has been genetically modified.
11. The method according to any one of claims 1 to 10, wherein the culture of step (b) or (e) is conditioned by a growth factor.
12. The method of claim 11, wherein the growth factor is a protein of the Wnt family.
13. The method of claim 12, wherein the protein of the Wnt family is Wnt3a.
14. The method according to any one of claims 1 to 13, wherein the rodent is a mouse.

15. A method for generating a mouse homozygous for a genetic modification, comprising:
- (a) introducing a female mouse embryonic stem (ES) donor cell into a first pre-morula diploid mouse host embryo, wherein the female mouse ES donor cell is heterozygous for the genetic modification and is obtained from an inbred strain;
 - (b) culturing the embryo of (a) to the blastocyst stage;
 - (c) introducing the embryo of (b) into a first surrogate mother for gestation, wherein a modified female mouse is generated having at least 90% cellular contribution from the female donor cell; and,
 - (d) introducing a male mouse ES donor cell into a second pre-morula diploid mouse host embryo, wherein the male mouse ES donor cell is heterozygous for the genetic modification and is obtained from an inbred strain;
 - (e) culturing the embryo of (d) to the blastocyst stage;
 - (d) introducing the embryo of (e) into a second surrogate mother for gestation, wherein a modified male mouse is generated; and,
 - (e) breeding a sexually mature modified male mouse and a sexually mature modified female mouse of (c) and (f), wherein a mouse homozygous for the genetic modification is generated.
16. The method of claim 15, wherein the female mouse ES donor cell of step (a) is obtained from the same cell line of the male mouse ES donor cell of step (d).
17. The method of claim 15, wherein the female mouse ES donor cell of step (a) is an XO cell and the modified female mouse of step (c) is an XO mouse.
18. The method of claim 15, wherein the first and the second pre-morula stage diploid mouse host embryo is an 8-cell embryo.
19. The method of claim 15, wherein the inbred strain is 129, C57BL/6, or BalbC.
20. The method of claim 15, wherein the genetically modified male and female mouse ES donor cell are generated by:
- (i) obtaining a cloned genomic fragment containing a DNA sequence of interest;

- (ii) using bacterial homologous recombination to genetically modify the cloned genomic fragment of (i) to create a targeting vector (LTVEC) for use in the mouse ES donor cells; and,
- (iii) introducing the LTVEC of (ii) into the mouse ES donor cells to modify an endogenous gene or chromosomal locus in the cells; and,
- (iv) using a quantitative assay to detect modification of an allele in the mouse ES donor cells of (iii) to identify those donor cells in which an endogenous gene or chromosomal locus has been genetically modified.

21. The method of claim 15, wherein the first and the second pre-morula diploid mouse host embryo is obtained from an inbred strain.

22. The method of claim 21, wherein the inbred strain is C57BL/6.

23. The method of claim 15, wherein the mouse host embryo is obtained from an outbred strain.

24. The method of claim 23, wherein the outbred strain is Swiss Webster.

25. The method of claim 15, wherein the culture of step (b) or (e) is conditioned by a growth factor.

26. The method of claim 25, wherein the growth factor is a protein of the Wnt family.

27. The method of claim 26, wherein the protein of the Wnt family is Wnt3a.

28. The method of claim 15, wherein the ES donor cell is introduced into the mouse host embryo through an opening in the zona pellucida.

29. The method of claim 28, wherein the opening is created by a laser.

30. A method for making a mouse embryo homozygous for a genetic modification, comprising:
- (a) incorporating a female mouse ES donor cell into a first pre-morula diploid mouse host embryo, wherein the female mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
 - (b) culturing the first embryo to the blastocyst stage
 - (c) gestating the embryo of (b) in a first surrogate mother, wherein upon gestation a modified female mouse is obtained having at least 90% cellular contribution from the female mouse ES donor cell;
 - (d) incorporating a male mouse ES donor cell into a second pre-morula diploid mouse host embryo, wherein the male mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
 - (e) culturing the embryo of (d) to the blastocyst stage;
 - (f) gestating the embryo of (e) in a second surrogate mother, wherein upon gestation a modified male mouse is obtained having at least 90% cellular contribution from the male mouse ES donor cell; and,
 - (g) breeding the modified female mouse of (c) with the modified male mouse of (f) to obtain a mouse embryo homozygous for the genetic modification.
31. The method of claim 30, wherein the first and the second pre-morula host embryos are 8-cell stage embryos.
32. The method of claim 31, wherein the first and the second pre-morula host embryos are from an inbred strain.
33. The method of claim 31, wherein the first and the second pre-morula host embryos are from an outbred strain.
34. The method of claim 31, wherein the first and the second pre-morula host embryos each comprise a zona pellucida, and the ES donor cell is incorporated through an opening in the zona pellucida.
35. The method of claim 34, wherein the opening is produced by a laser.

36. A method for making a mouse embryo heterozygous for a genetic modification, comprising:
- (a) incorporating a mouse ES donor cell into a pre-morula diploid mouse host embryo, wherein the mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
 - (b) culturing the host embryo of (a) to the blastocyst stage; and,
 - (c) gestating the host embryo of (b) in a surrogate mother, wherein upon gestation a modified mouse embryo is obtained having at least 90% cellular contribution from the mouse ES donor cell.
37. The method of claim 36, wherein the pre-morula host embryo is an 8-cell stage embryo.
38. The method of claim 36, wherein the pre-morula host embryo is from an inbred strain.
39. The method of claim 36, wherein the pre-morula host embryo is from an outbred strain.
40. The method of claim 365, wherein the pre-morula embryo comprises a zona pellucida, and the mouse ES donor cell is incorporated through an opening in the zona pellucida.
41. The method of claim 40, wherein the opening is produced by a laser.
42. A method for making a mouse heterozygous for a genetic modification, comprising:
- (a) incorporating a mouse ES donor cell into a pre-morula diploid mouse host embryo, wherein the mouse ES donor cell is heterozygous for the genetic modification and is from an inbred strain;
 - (b) culturing the host embryo to the blastocyst stage; and,
 - (c) gestating the host embryo of (b) in a surrogate mother, wherein upon gestation a modified mouse is obtained having at least 90% cellular contribution from the mouse ES donor cell.
43. The method of claim 42, wherein the pre-morula host embryo is an 8-cell stage embryo.

44. The method of claim 42, wherein the pre-morula host embryo is from an inbred strain.
45. The method of claim 42, wherein the pre-morula host embryo is from an outbred strain.
46. The method of claim 42, wherein the pre-morula diploid mouse host embryo comprises a zona pellucida, and the mouse ES donor cell is incorporated through an opening in the zona pellucida.
47. The method of claim 46, wherein the opening is produced by a laser.
48. A method of generating a mouse wherein at least 90% of the cells in said mouse comprise a homozygous genetic modification, the method comprising:
- (a) incorporating a donor mouse embryonic stem (ES) cell from an inbred strain into a pre-morula diploid mouse host embryo, wherein the donor ES cell is homozygous for the genetic modification;
 - (b) culturing the pre-morula host embryo of (a) to the blastocyst stage;
 - (c) introducing the embryo of (b) into a surrogate mother for gestation, and,
 - (d) generating a mouse having at least a 90% cellular contribution from the donor ES cell, wherein at least 90% of the cells in said mouse are homozygous for the genetic modification.
49. The method of claim 48, wherein the donor mouse ES cell homozygous for the genetic modification is generated by a process selected from the group consisting of gene conversion, targeting both alleles of the same gene, and targeting either an X- or a Y-linked gene in an ES cell.
50. The method of claim 48, wherein the pre-morula host embryo is an 8-cell stage embryo.
51. The method of claim 48, wherein the pre-morula host embryo is from an inbred strain.
52. The method of claim 48, wherein the pre-morula host embryo is from an outbred strain.

53. The method of claim 48, wherein the pre-morula host embryo of (a) comprises a zona pellucida, and the donor mouse embryonic stem cell is incorporated through an opening in the zona pellucida.
54. The method of claim 53, wherein the opening is made by a laser.
55. A method of generating a mouse embryo wherein at least 90% of the cells in said mouse embryo comprise a homozygous genetic modification, the method comprising:
- (a) incorporating a donor mouse ES cell from an inbred strain into a pre-morula diploid mouse host embryo, wherein the donor ES cell is homozygous for the genetic modification; and,
 - (b) culturing the pre-morula host embryo of (a), wherein a mouse embryo comprising at least a 90% cellular contribution from the donor ES cell is generated and wherein at least 90% of the cells in said mouse embryo are homozygous for the genetic modification.
56. The method of claim 55, wherein the donor mouse ES cell homozygous for the genetic modification is generated by a process selected from the group consisting of gene conversion, targeting both alleles of the same gene, and targeting either an X- or a Y-linked gene in an ES cell.
57. The method of claim 55, wherein the pre-morula host embryo is an 8-cell stage embryo.
58. The method of claim 55, wherein the pre-morula host embryo is from an inbred strain.
59. The method of claim 55, wherein the pre-morula host embryo is from an outbred strain.
60. The method of claim 55, wherein the pre-morula host embryo of (a) comprises a zona pellucida, and the donor mouse embryonic stem cell is incorporated through an opening in the zona pellucida.
61. The method of claim 60, wherein the opening is made by a laser.

62. The method of claim 48, wherein at least 95% of the cells of the mouse are donor ES cell-derived.
63. The method of claim 48, wherein at least 99% of the cells of the mouse are donor ES cell-derived.
64. The method of claim 55, wherein at least 95% of the cells of the generated embryo are donor ES cell-derived.
65. The method of claim 55, wherein at least 99% of the cells of the generated embryo are donor ES cell-derived.
66. A method for making a mouse having a genetic modification, comprising:
 (a) incorporating a donor mouse ES cell comprising a genetic modification into a mouse host pre-morula stage embryo to obtain a mouse embryo comprising a donor ES cell;
 (b) growing the mouse embryo comprising the donor ES cell in a medium conditioned by mouse fibroblasts that produce a protein that is a member of the Wnt family of proteins; and,
 (c) gestating the mouse embryo comprising the donor ES cell in a surrogate mother to obtain a mouse that develops from the mouse embryo comprising the donor ES cell.
67. The method according to claim 65, wherein the contribution of the donor mouse ES cell to the mouse that develops from the mouse embryo is 100%.
68. The method according to claim 65, wherein the method is performed a plurality of times to obtain a plurality of mice that develops from the mouse embryo, and wherein at least 40% of the plurality of mice that develops from the mouse embryo display a contribution of the donor mouse ES cell that is 100%.
69. A method for generating a mouse embryo comprising cells that are homozygous for a genetic modification, comprising:

(a) introducing mouse donor cells into a pre-morula host mouse embryo, wherein the donor cells comprise ES cells or ES-like cells from an inbred mouse, and are homozygous for the genetic modification; and

(b) culturing the pre-morula host embryo of (a) to the blastocyst stage, wherein at least 90% of the cells of a mouse that develops from the blastocyst are derived from the donor cells.

70. The method of claim 69, further comprising introducing the embryo of (b) into a surrogate mouse mother for gestation.

71. The method of claim 69 or 70, wherein the mouse donor cells homozygous for the genetic modification are generated by a process selected from the group consisting of gene conversion, targeting both alleles of the same gene, and targeting either an X- or a Y-linked gene in the ES cells or ES-like cells.

72. The method according to any one of claims 69 to 71, wherein (i) at least 95% of the cells of a mouse that develops from the blastocyst are derived from the donor cells, or (ii) at least 99% of the cells of a mouse that develops from the blastocyst are derived from the donor cells.

73. A method for generating a mouse embryo comprising cells that are heterozygous for a genetic modification, comprising:

(a) introducing mouse donor cells into a pre-morula host mouse embryo, wherein the mouse donor cells are ES cells or an ES-like cells from an inbred mouse; and are heterozygous for the genetic modification; and

(b) culturing the pre-morula host embryo of (a) to the blastocyst stage, wherein at least 90% of the cells of a mouse that develops from the blastocyst are derived from the donor cells.

74. The method of claim 73, further comprising gestating the embryo of (b) in a surrogate mouse mother.

75. The method according to claim 73 or 74, wherein (i) at least 95% of the cells of a mouse that develops from the blastocyst are derived from the donor cells, or (ii) at least 99% of the cells of a mouse that develops from the blastocyst are derived from the donor cells.

76. The method according to any one of claims 69 to 75, wherein the pre-morula host mouse embryo is: (i) a diploid embryo; and/or (ii) from an inbred strain or an outbred strain.

77. The method according to any one of claims 69 to 76, wherein the pre-morula host mouse embryo is an 8-cell stage embryo.

78. The method according to any one of claims 69 to 77, wherein the pre-morula host mouse embryo comprises a zona pellucida, and wherein the mouse donor cells are introduced into the host mouse embryo through an opening in the zona pellucida; optionally wherein the opening is created by a laser.

79. The method according to any one of claims 69 to 78, wherein the genetically modified donor cells are generated by:

- (a) obtaining a cloned genomic fragment containing a DNA sequence of interest;
- (b) using bacterial homologous recombination to genetically modify the cloned genomic fragment of (a) to create a targeting vector (LTVEC) for use in the donor cells ;
- (c) introducing the LTVEC of (b) into the donor cells to modify an endogenous gene or a chromosomal locus in the cells; and
- (d) using a quantitative assay to detect modification of allele in the donor cells of (c) to identify those donor cells in which the endogenous gene or chromosomal locus has been genetically modified.

80. The method according to any one of claims 69 to 79, wherein the donor ES cells are from an inbred mouse strain selected from the group consisting of 129, C57BL/6, and BalbC.

81. The method according to any of claims 69 to 80, wherein the host mouse embryo comprises an inbred embryo from strain C57BL/6 or an outbred embryo from strain Swiss Webster.

82. The method according to any one of claims 69 to 81, wherein the culture of step (b) is conditioned by a growth factor, optionally a protein of the Wnt family; preferably Wnt3a.

83. The method according to any one of claims 69 to 82 further comprising gestation of the genetically modified mouse embryo to generate a genetically modified mouse, wherein when the donor cells are homozygous for the genetic modification, at least 90% of the cells in the genetically modified mouse are homozygous for the genetic modification.

84. The method according to any one of claims 69 to 82 further comprising gestation of the genetically modified mouse embryo to generate a genetically modified mouse, wherein when the donor cells are heterozygous for the genetic modification at least 90% of the cells in the genetically modified mouse are heterozygous for the genetic modification.

85. The method according to claim 83, wherein: (a) at least 95% of the cells in the genetically modified mouse are homozygous for the genetic modification; or (b) at least 99% of the cells in the genetically modified mouse are homozygous for the genetic modification.

86. The method according to claim 84, wherein:

(a) at least 95% of the cells in the genetically modified mouse are heterozygous for the genetic modification, or

(b) at least 99% of the cells in the genetically modified mouse are heterozygous for the genetic modification.

87. The method according to claim 83, wherein at least 90% of the germ cells in the genetically modified mouse are homozygous for the genetic modification:

88. The method according to claim 83, wherein at least 90% of the germ cells in the genetically modified mouse are heterozygous for the genetic modification.

89. The method according to any one of claims 83 to 88, wherein the donor cells are female XO cells and the genetically modified mouse is an XO mouse.

90. The method of claim 42 or 48, wherein in step (b) the host embryo is cultured in a medium that comprises a Wnt family protein produced by mouse fibroblasts.
91. The method according to claim 90, wherein the contribution of the donor mouse ES cell to the mouse that develops from the mouse embryo is 100%.
92. The method according to claim 90, wherein the method is performed a plurality of times to obtain a plurality of mice that develops from the mouse embryo, and wherein at least 40% of the plurality of mice that develops from the mouse embryo display a contribution of the donor mouse ES cell that is 100%.