



(43) International Publication Date  
12 November 2015 (12.11.2015)

(10) International Publication Number  
**WO 2015/171430 A1**

- (51) **International Patent Classification:**  
*C07F 5/04* (2006.01) *C07F 5/02* (2006.01)
- (21) **International Application Number:**  
PCT/US2015/028613
- (22) **International Filing Date:**  
30 April 2015 (30.04.2015)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**  
61/988,690 5 May 2014 (05.05.2014) US
- (71) **Applicant:** REMPEX PHARMACEUTICALS, INC.  
[US/US]; 3033 Science Park Rd., Suite 200, San Diego,  
CA 92121 (US).
- (72) **Inventors:** HECKER, Scott; 4802 Rancho Viejo Dr., Del  
Mar, CA 92014 (US). BOYER, Serge; 5112 Castle Hills  
Drive, San Diego, CA 92109 (US).
- (74) **Agent:** DELANEY, Karoline, A.; Knobbe Martens Olson  
& Bear, LLP, 2040 Main Street, 14th Floor, Irvine, CA  
92614 (US).

- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

**Published:**

— with international search report (Art. 21(3))

(54) **Title:** SYNTHESIS OF BORONATE SALTS AND USES THEREOF

(57) **Abstract:** Disclosed herein are boronate intermediates in the synthesis of antimicrobial compounds and the use and preparation thereof. Some embodiments relate to crystalline boronate salt derivatives and their use in the synthesis of therapeutic compounds.



WO 2015/171430 A1

## SYNTHESIS OF BORONATE SALTS AND USES THEREOF

### BACKGROUND OF THE INVENTION

#### Field of the Invention

[0001] The present application relates to certain compounds and to methods for the preparation of certain compounds that can be used in the fields of chemistry and medicine. More specifically, the present application relates to intermediates and methods in the synthesis of boronic acid antimicrobial compounds.

#### Description of the Related Art

[0002] Antibiotics have been effective tools in the treatment of infectious diseases during the last half-century. From the development of antibiotic therapy to the late 1980s there was almost complete control over bacterial infections in developed countries. However, in response to the pressure of antibiotic usage, multiple resistance mechanisms have become widespread and are threatening the clinical utility of anti-bacterial therapy. The increase in antibiotic resistant strains has been particularly common in major hospitals and care centers. The consequences of the increase in resistant strains include higher morbidity and mortality, longer patient hospitalization, and an increase in treatment costs.

[0003] Various bacteria have evolved  $\beta$ -lactam deactivating enzymes, namely,  $\beta$ -lactamases, that counter the efficacy of the various  $\beta$ -lactam antibiotics.  $\beta$ -lactamases can be grouped into 4 classes based on their amino acid sequences, namely, Ambler classes A, B, C, and D. Enzymes in classes A, C, and D include active-site serine  $\beta$ -lactamases, and class B enzymes, which are encountered less frequently, are Zn-dependent. These enzymes catalyze the chemical degradation of  $\beta$ -lactam antibiotics, rendering them inactive. Some  $\beta$ -lactamases can be transferred within and between various bacterial strains and species. The rapid spread of bacterial resistance and the evolution of multi-resistant strains severely limits  $\beta$ -lactam treatment options available.

[0004] The increase of class D  $\beta$ -lactamase-expressing bacterium strains such as *Acinetobacter baumannii* has become an emerging multidrug-resistant threat. *A. baumannii* strains express A, C, and D class  $\beta$ -lactamases. The class D  $\beta$ -lactamases such as the OXA families are particularly effective at destroying carbapenem type  $\beta$ -lactam antibiotics, e.g.,

imipenem, the active carbapenems component of Merck's Primaxin® (Montefour, K.; *et al.* Crit. Care Nurse 2008, 28, 15; Perez, F. *et al.* Expert Rev. Anti Infect. Ther. 2008, 6, 269; Bou, G.; Martinez-Beltran, J. Antimicrob. Agents Chemother. 2000, 40, 428. 2006, 50, 2280; Bou, G. *et al.* J. Antimicrob. Agents Chemother. 2000, 44, 1556). This has imposed a pressing threat to the effective use of drugs in that category to treat and prevent bacterial infections. Indeed the number of catalogued serine-based  $\beta$ -lactamases has exploded from less than ten in the 1970s to over 300 variants. These issues fostered the development of five "generations" of cephalosporins. When initially released into clinical practice, extended-spectrum cephalosporins resisted hydrolysis by the prevalent class A  $\beta$ -lactamases, TEM-1 and SHV-1. However, the development of resistant strains by the evolution of single amino acid substitutions in TEM-1 and SHV-1 resulted in the emergence of the extended-spectrum  $\beta$ -lactamase (ESBL) phenotype.


**[0005]** New  $\beta$ -lactamases have recently evolved that hydrolyze the carbapenem class of antimicrobials, including imipenem, biapenem, doripenem, meropenem, and ertapenem, as well as other  $\beta$ -lactam antibiotics. These carbapenemases belong to molecular classes A, B, and D. Class A carbapenemases of the KPC-type predominantly in *Klebsiella pneumoniae* but now also reported in other *Enterobacteriaceae*, *Pseudomonas aeruginosa* and *Acinetobacter baumannii*. The KPC carbapenemase was first described in 1996 in North Carolina, but since then has disseminated widely in the US. It has been particularly problematic in the New York City area, where several reports of spread within major hospitals and patient morbidity have been reported. These enzymes have also been recently reported in France, Greece, Sweden, United Kingdom, and an outbreak in Germany has recently been reported. Treatment of resistant strains with carbapenems can be associated with poor outcomes.

**[0006]** Another mechanism of  $\beta$ -lactamase mediated resistance to carbapenems involves combination of permeability or efflux mechanisms combined with hyper production of beta-lactamases. One example is the loss of a porin combined in hyperproduction of ampC beta-lactamase results in resistance to imipenem in *Pseudomonas aeruginosa*. Efflux pump over expression combined with hyperproduction of the ampC  $\beta$ -lactamase can also result in resistance to a carbapenem such as meropenem.

[0007] Thus, there is a need for improved  $\beta$ -lactamase inhibitors and efficient methods for making these improved  $\beta$ -lactamase inhibitors.

## SUMMARY OF THE INVENTION

[0008] Some embodiments of the present application provide a compound of Formula (I) or a salt thereof:



[0009] In some embodiments,  $n$  is 0 or 1.

[0010] In some embodiments,  $Y_1$  is O or  $N^+R_5R_6$ .

[0011] In some embodiments,  $Y_2$  is O or  $NR_{10}$ .

[0012] In some embodiments,  $R_1$  and  $R_2$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl, or  $R_1$  and  $R_2$  together with the atoms to which they are attached, form =O.

[0013] In some embodiments,  $R_3$  is selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl, or  $R_3$  and  $R_5$ , together with the atoms to which they are attached, form a heteroaryl ring; or  $R_3$  and  $R_4$  together with the atoms to which they are attached, form =O.

[0014] In some embodiments,  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl.

[0015] In some embodiments,  $R_9$  is selected from the group consisting of optionally substituted  $C_1$ - $C_{12}$  alkyl, optionally substituted  $C_3$ - $C_8$  cycloalkyl, optionally substituted  $C_2$ - $C_{12}$  alkenyl, optionally substituted  $C_2$ - $C_{12}$  alkynyl, optionally substituted aryl and optionally substituted heteroaryl.

[0016] In some embodiments,  $R_1$  is H. In some embodiments,  $R_1$  is optionally substituted phenyl. In some embodiments,  $R_1$  is optionally substituted  $C_{1-4}$  alkyl.

[0017] In some embodiments,  $R_2$  is H. In some embodiments,  $R_2$  is optionally substituted phenyl. In some embodiments,  $R_2$  is optionally substituted  $C_{1-4}$  alkyl.

[0018] In some embodiments,  $R_1$  and  $R_2$  together with the atoms to which they are attached, form =O. In some embodiments,  $R_3$  and  $R_4$  together with the atoms to which

they are attached, form =O. In some embodiments, R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O and wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

**[0019]** In some embodiments, R<sub>3</sub> is H. In some embodiments, R<sub>3</sub> is optionally substituted phenyl. In some embodiments, R<sub>3</sub> is C<sub>1-4</sub> optionally substituted alkyl. In some embodiments, R<sub>1</sub> and R<sub>3</sub> together with the atoms to which they are attached form carbocyclic ring. In some embodiments, R<sub>3</sub> and R<sub>5</sub>, together with the atoms to which they are attached, form a heteroaryl ring.

**[0020]** In some embodiments, R<sub>4</sub> is H. In some embodiments, R<sub>4</sub> is optionally substituted phenyl. In some embodiments, R<sub>4</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0021]** In some embodiments, R<sub>5</sub> is H. In some embodiments, R<sub>5</sub> is optionally substituted phenyl. In some embodiments, R<sub>5</sub> is optionally substituted C<sub>1-4</sub> alkyl.


**[0022]** In some embodiments, R<sub>6</sub> is H. In some embodiments, R<sub>6</sub> is optionally substituted phenyl. In some embodiments, R<sub>6</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0023]** In some embodiments, R<sub>10</sub> is H. In some embodiments, R<sub>10</sub> is optionally substituted phenyl. In some embodiments, R<sub>10</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0024]** In some embodiments, R<sub>9</sub> is optionally substituted C<sub>1-C12</sub> alkyl. In some embodiments, R<sub>9</sub> is t-butyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>3-C8</sub> cycloalkyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2-C12</sub> alkenyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2-C12</sub> alkynyl. In some embodiments, R<sub>9</sub> is optionally substituted aryl. In some embodiments, R<sub>9</sub> is optionally substituted heteroaryl.


**[0025]** In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with OMe. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with phenyl. In some embodiments, each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen. In some embodiments, each phenyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with halogen.

**[0026]** In some embodiments, the compound of Formula (I) is selected from the group consisting of:




[0027] In some embodiments,  $M^+$  is a cation selected from the group consisting of lithium, sodium, potassium, calcium, ammonium, triethylammonium, and aluminum.


[0028] Some embodiments of the present application provides a method of making a compound of Formula (I), or a salt thereof:




[0029] In some embodiments, the method of making comprises the steps of: protecting the primary hydroxy group of the compound of Formula (A)




to form a compound of Formula (B):




reacting a compound of Formula (B) with a compound of Formula (C):




to form a compound of Formula (D):




deprotecting a compound of Formula (D) to form a compound of Formula (E):



reacting a compound of Formula (E) with a complexing agent of Formula (F):



to form a compound of Formula (I):



[0030] In some embodiments,  $n$  is 0 or 1.

[0031] In some embodiments,  $Y_1$  is O or  $N^+R_5R_6$ .

[0032] In some embodiments,  $Y_2$  is O or  $NR_{10}$ .

[0033] In some embodiments,  $R_1$  and  $R_2$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl, or  $R_1$  and  $R_2$  together with the atoms to which they are attached, form =O.

[0034] In some embodiments,  $R_3$  is selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl, or  $R_3$  and  $R_5$ , together with the atoms to which they are attached, form a heteroaryl ring; or  $R_3$  and  $R_4$  together with the atoms to which they are attached, form =O.

[0035] In some embodiments,  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl.

[0036] In some embodiments,  $R_9$  is selected from the group consisting of optionally substituted  $C_1$ - $C_{12}$  alkyl, optionally substituted  $C_3$ - $C_8$  cycloalkyl, optionally substituted  $C_2$ - $C_{12}$  alkenyl, optionally substituted  $C_2$ - $C_{12}$  alkynyl, optionally substituted aryl and optionally substituted heteroaryl.

[0037] In some embodiments,  $R_1$  is H. In some embodiments,  $R_1$  is optionally substituted phenyl. In some embodiments,  $R_1$  is optionally substituted  $C_{1-4}$  alkyl.

[0038] In some embodiments,  $R_2$  is H. In some embodiments,  $R_2$  is optionally substituted phenyl. In some embodiments,  $R_2$  is optionally substituted  $C_{1-4}$  alkyl.

[0039] In some embodiments,  $R_1$  and  $R_2$  together with the atoms to which they are attached, form =O. In some embodiments,  $R_3$  and  $R_4$  together with the atoms to which they are attached, form =O. In some embodiments,  $R_1$  and  $R_2$  together with the atoms to

which they are attached, form =O and wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

**[0040]** In some embodiments, R<sub>3</sub> is H. In some embodiments, R<sub>3</sub> is optionally substituted phenyl. In some embodiments, R<sub>3</sub> is C<sub>1-4</sub> optionally substituted alkyl. In some embodiments, R<sub>1</sub> and R<sub>3</sub> together with the atoms to which they are attached form carbocyclic ring. In some embodiments, R<sub>3</sub> and R<sub>5</sub>, together with the atoms to which they are attached, form a heteroaryl ring.

**[0041]** In some embodiments, R<sub>4</sub> is H. In some embodiments, R<sub>4</sub> is optionally substituted phenyl. In some embodiments, R<sub>4</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0042]** In some embodiments, R<sub>5</sub> is H. In some embodiments, R<sub>5</sub> is optionally substituted phenyl. In some embodiments, R<sub>5</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0043]** In some embodiments, R<sub>6</sub> is H. In some embodiments, R<sub>6</sub> is optionally substituted phenyl. In some embodiments, R<sub>6</sub> is optionally substituted C<sub>1-4</sub> alkyl.


**[0044]** In some embodiments, R<sub>10</sub> is H. In some embodiments, R<sub>10</sub> is optionally substituted phenyl. In some embodiments, R<sub>10</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0045]** In some embodiments, R<sub>9</sub> is optionally substituted C<sub>1-C12</sub> alkyl. In some embodiments, R<sub>9</sub> is t-butyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>3-C8</sub> cycloalkyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2-C12</sub> alkenyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2-C12</sub> alkynyl. In some embodiments, R<sub>9</sub> is optionally substituted aryl. In some embodiments, R<sub>9</sub> is optionally substituted heteroaryl.

**[0046]** In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with OMe. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with phenyl. In some embodiments, each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen. In some embodiments, each phenyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with halogen.

**[0047]** In some embodiments, the compound of Formula (I) is selected from the group consisting of:





**[0048]** In some embodiments, M<sup>+</sup> is a cation selected from the group consisting of lithium, sodium, potassium, calcium, ammonium, triethylammonium, and aluminum.

**[0049]** In some embodiments, PG is a hydroxyl protecting group. In some embodiments, PG is a trialkylsilyl group. In some embodiments, PG is trimethylsilyl.

**[0050]** In some embodiments, R<sub>11</sub> is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkenyl, C<sub>3</sub>-C<sub>12</sub> cycloalkynyl, C<sub>3</sub>-C<sub>12</sub> heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl.

**[0051]** In some embodiments, X is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, and C<sub>2</sub>-C<sub>4</sub> alkynyl.

**[0052]** In some embodiments, R<sub>12</sub> is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkenyl, C<sub>3</sub>-C<sub>12</sub> cycloalkynyl, C<sub>3</sub>-C<sub>12</sub> heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl.


**[0053]** In some embodiments, R<sub>11</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkenyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkynyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> heterocyclyl. In some embodiments, R<sub>11</sub> is optionally substituted aryl. In some embodiments, R<sub>11</sub> is optionally substituted heteroaryl. In some embodiments, R<sub>11</sub> is optionally substituted arylalkyl. In some embodiments, R<sub>11</sub> is optionally substituted heteroarylalkyl. In some embodiments, R<sub>11</sub> is optionally substituted (cycloalkyl)alkyl.

**[0054]** In some embodiments, X is optionally substituted C<sub>1</sub>-C<sub>4</sub> alkyl. In some embodiments, X is optionally substituted C<sub>2</sub>-C<sub>4</sub> alkenyl. In some embodiments, X is optionally substituted C<sub>2</sub>-C<sub>4</sub> alkynyl.


**[0055]** In some embodiments, R<sub>12</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl. In some embodiments, R<sub>12</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl. In some embodiments, R<sub>12</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl. In some embodiments, R<sub>12</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkyl. In some embodiments, R<sub>12</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkenyl. In some embodiments, R<sub>12</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkynyl. In some embodiments, R<sub>12</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> heterocyclyl. In some embodiments, R<sub>12</sub> is optionally substituted aryl. In some embodiments, R<sub>12</sub> is optionally substituted heteroaryl. In some embodiments, R<sub>12</sub> is optionally substituted arylalkyl. In some embodiments, R<sub>12</sub> is optionally substituted heteroarylalkyl. In some embodiments, R<sub>12</sub> is optionally substituted (cycloalkyl)alkyl.

**[0056]** In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with OMe. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with phenyl. In some embodiments, each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen. In some embodiments, each phenyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with halogen.


**[0057]** Some embodiments of the present application provides a method of making a compound of Formula (D):




**[0058]** In some embodiments, the method of making comprises the steps of: reacting a compound of Formula (B):



with a compound of Formula (C):



to form a compound of Formula (D):



**[0059]** In some embodiments, R<sub>9</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl. In some embodiments, R<sub>9</sub> is t-butyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>3</sub>-C<sub>8</sub> cycloalkyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl. In some embodiments, R<sub>9</sub> is optionally substituted aryl. In some embodiments, R<sub>9</sub> is optionally substituted heteroaryl.

**[0060]** In some embodiments, R<sub>11</sub> is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkenyl, C<sub>3</sub>-C<sub>12</sub> cycloalkynyl, C<sub>3</sub>-C<sub>12</sub> heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl.

**[0061]** In some embodiments, X is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>2</sub>-C<sub>4</sub> alkenyl, and C<sub>2</sub>-C<sub>4</sub> alkynyl.

**[0062]** In some embodiments, R<sub>12</sub> is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkenyl, C<sub>3</sub>-C<sub>12</sub> cycloalkynyl, C<sub>3</sub>-C<sub>12</sub> heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl.

**[0063]** In some embodiments, R<sub>11</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkenyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkynyl. In some embodiments, R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> heterocyclyl. In some embodiments, R<sub>11</sub> is optionally substituted aryl. In some embodiments, R<sub>11</sub> is optionally substituted heteroaryl. In some embodiments, R<sub>11</sub> is optionally substituted arylalkyl. In some embodiments, R<sub>11</sub> is

optionally substituted heteroarylalkyl. In some embodiments,  $R_{11}$  is optionally substituted (cycloalkyl)alkyl.


**[0064]** In some embodiments, X is optionally substituted  $C_1$ - $C_4$  alkyl. In some embodiments, X is optionally substituted  $C_2$ - $C_4$  alkenyl. In some embodiments, X is optionally substituted  $C_2$ - $C_4$  alkynyl.

**[0065]** In some embodiments,  $R_{12}$  is optionally substituted  $C_1$ - $C_{12}$  alkyl. In some embodiments,  $R_{12}$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl. In some embodiments,  $R_{12}$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl. In some embodiments,  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkyl. In some embodiments,  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkenyl. In some embodiments,  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkynyl. In some embodiments,  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  heterocyclyl. In some embodiments,  $R_{12}$  is optionally substituted aryl. In some embodiments,  $R_{12}$  is optionally substituted heteroaryl. In some embodiments,  $R_{12}$  is optionally substituted arylalkyl. In some embodiments,  $R_{12}$  is optionally substituted heteroarylalkyl. In some embodiments,  $R_{12}$  is optionally substituted (cycloalkyl)alkyl.

**[0066]** In some embodiments, each  $C_{1-4}$  alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl. In some embodiments, each  $C_{1-4}$  alkyl is optionally substituted with OMe. In some embodiments, each  $C_{1-4}$  alkyl is optionally substituted with phenyl. In some embodiments, each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen. In some embodiments, each phenyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with halogen.


**[0067]** In some embodiments, PG is a hydroxyl protecting group. In some embodiments, PG is a trialkylsilyl group. In some embodiments, PG is trimethylsilyl.

**[0068]** Some embodiments of the present application provides a method of making a compound of Formula (I), or a salt thereof:




**[0069]** In some embodiments, the method of making comprises the steps of:


reacting a compound of Formula (E):



with a complexing agent of Formula (F):



to form a compound of Formula (I):



**[0070]** In some embodiments,  $n$  is 0 or 1.

**[0071]** In some embodiments,  $Y_1$  is O or  $N^+R_5R_6$ .

**[0072]** In some embodiments,  $Y_2 = O$  or  $NR_{10}$ .

**[0073]** In some embodiments,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl.

**[0074]** In some embodiments, independently two germinal  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  together with the atoms to which they are attached, form  $=O$ .

**[0075]** In some embodiments,  $R_9$  is optionally substituted  $C_1$ - $C_{12}$  alkyl, optionally substituted  $C_3$ - $C_8$  cycloalkyl, optionally substituted  $C_2$ - $C_{12}$  alkenyl, optionally substituted  $C_2$ - $C_{12}$  alkynyl, optionally substituted aryl or optionally substituted heteroaryl.

**[0076]** In some embodiments,  $R_1$  is H. In some embodiments,  $R_1$  is optionally substituted phenyl. In some embodiments,  $R_1$  is optionally substituted  $C_{1-4}$  alkyl.

**[0077]** In some embodiments,  $R_2$  is H. In some embodiments,  $R_2$  is optionally substituted phenyl. In some embodiments,  $R_2$  is optionally substituted  $C_{1-4}$  alkyl.

**[0078]** In some embodiments,  $R_1$  and  $R_2$  together with the atoms to which they are attached, form  $=O$ .

**[0079]** In some embodiments,  $R_3$  is H. In some embodiments,  $R_3$  is optionally substituted phenyl. In some embodiments,  $R_3$  is optionally substituted  $C_{1-4}$  alkyl.

**[0080]** In some embodiments, R<sub>4</sub> is H. In some embodiments, R<sub>4</sub> is optionally substituted phenyl. In some embodiments, R<sub>4</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0081]** In some embodiments, R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O. In some embodiments, R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O and wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

**[0082]** In some embodiments, R<sub>5</sub> is H. In some embodiments, R<sub>5</sub> is optionally substituted phenyl. In some embodiments, R<sub>5</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0083]** In some embodiments, R<sub>6</sub> is H. In some embodiments, R<sub>6</sub> is optionally substituted phenyl. In some embodiments, R<sub>6</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0084]** In some embodiments, R<sub>10</sub> is H. In some embodiments, R<sub>10</sub> is optionally substituted phenyl. In some embodiments, R<sub>10</sub> is optionally substituted C<sub>1-4</sub> alkyl.

**[0085]** In some embodiments, R<sub>9</sub> is optionally substituted C<sub>1-C12</sub> alkyl. In some embodiments, R<sub>9</sub> is t-butyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>3-C8</sub> cycloalkyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2-C12</sub> alkenyl. In some embodiments, R<sub>9</sub> is optionally substituted C<sub>2-C12</sub> alkynyl. In some embodiments, R<sub>9</sub> is optionally substituted aryl. In some embodiments, R<sub>9</sub> is optionally substituted heteroaryl.

**[0086]** In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with OMe. In some embodiments, each C<sub>1-4</sub> alkyl is optionally substituted with phenyl. In some embodiments, each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen. In some embodiments, each phenyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with halogen.

#### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

**[0087]** Numerous references are cited herein. The references cited herein, including the U.S. patents cited herein, are each to be considered incorporated by reference in their entirety into this specification.

**[0088]** Embodiments of the invention include, but are not limited to, methods for the preparation of various compounds and intermediates, and the compounds and intermediates themselves. In some embodiments, one or more substituents, one or more compounds, or groups of compounds can be specifically excluded in any one or more of the methods or compounds as described more fully below.

**[0089]** Where the compounds disclosed herein have at least one chiral center, they may exist as individual enantiomers and diastereomers or as mixtures of such isomers, including racemates. Separation of the individual isomers or selective synthesis of the individual isomers is accomplished by application of various methods which are well known to practitioners in the art. Unless otherwise indicated, all such isomers and mixtures thereof are included in the scope of the compounds disclosed herein. Furthermore, compounds disclosed herein may exist in one or more crystalline or amorphous forms. Unless otherwise indicated, all such forms are included in the scope of the compounds disclosed herein including any polymorphic forms. In addition, some of the compounds disclosed herein may form solvates with water (i.e., hydrates) or common organic solvents. Unless otherwise indicated, such solvates are included in the scope of the compounds disclosed herein. Additionally, in some embodiments, the compounds disclosed herein can form oligomers and other higher-order polymers.

**[0090]** The skilled artisan will recognize that some structures described herein may be resonance forms or tautomers of compounds that may be fairly represented by other chemical structures, even when kinetically; the artisan recognizes that such structures may only represent a very small portion of a sample of such compound(s). Such compounds are considered within the scope of the structures depicted, though such resonance forms or tautomers are not represented herein.

**[0091]** Isotopes may be present in the compounds described. Each chemical element as represented in a compound structure may include any isotope of said element. For example, in a compound structure a hydrogen atom may be explicitly disclosed or understood to be present in the compound. At any position of the compound that a hydrogen atom may be present, the hydrogen atom can be any isotope of hydrogen, including but not limited to hydrogen-1 (protium) and hydrogen-2 (deuterium). Thus, reference herein to a compound encompasses all potential isotopic forms unless the context clearly dictates otherwise.

Definitions

**[0092]** “Solvate” refers to the compound formed by the interaction of a solvent and a compound described herein or salt thereof. Suitable solvates are pharmaceutically acceptable solvates including hydrates.

**[0093]** The term “pharmaceutically acceptable salt” refers to salts that retain the biological effectiveness and properties of a compound and, which are not biologically or otherwise undesirable for use in a pharmaceutical. In many cases, the compounds disclosed herein are capable of forming acid and/or base salts by virtue of the presence of amino and/or carboxyl groups or groups similar thereto. Pharmaceutically acceptable acid addition salts can be formed with inorganic acids and organic acids. Inorganic acids from which salts can be derived include, for example, hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid, and the like. Organic acids from which salts can be derived include, for example, acetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, p-toluenesulfonic acid, salicylic acid, and the like. Pharmaceutically acceptable base addition salts can be formed with inorganic and organic bases. Inorganic bases from which salts can be derived include, for example, sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum, and the like; particularly preferred are the ammonium, potassium, sodium, calcium and magnesium salts. Organic bases from which salts can be derived include, for example, primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines, basic ion exchange resins, and the like, specifically such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, and ethanolamine. Many such salts are known in the art, as described in WO 87/05297, Johnston et al., published September 11, 1987 (incorporated by reference herein in its entirety).

**[0094]** As used herein, “C<sub>a</sub> to C<sub>b</sub>” or “C<sub>a-b</sub>” in which “a” and “b” are integers refer to the number of carbon atoms in the specified group. That is, the group can contain from “a” to “b”, inclusive, carbon atoms. Thus, for example, a “C<sub>1</sub> to C<sub>4</sub> alkyl” or “C<sub>1-4</sub> alkyl”



group refers to all alkyl groups having from 1 to 4 carbons, that is,  $\text{CH}_3$ -,  $\text{CH}_3\text{CH}_2$ -,  $\text{CH}_3\text{CH}_2\text{CH}_2$ -,  $(\text{CH}_3)_2\text{CH}$ -,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{CH}_2$ -,  $\text{CH}_3\text{CH}_2\text{CH}(\text{CH}_3)$ - and  $(\text{CH}_3)_3\text{C}$ -.

**[0095]** The term “halogen” or “halo,” as used herein, means any one of the radio-stable atoms of column 7 of the Periodic Table of the Elements, *e.g.*, fluorine, chlorine, bromine, or iodine, with fluorine and chlorine being preferred.

**[0096]** As used herein, “alkyl” refers to a straight or branched hydrocarbon chain that is fully saturated (*i.e.*, contains no double or triple bonds). The alkyl group may have 1 to 20 carbon atoms (whenever it appears herein, a numerical range such as “1 to 20” refers to each integer in the given range; *e.g.*, “1 to 20 carbon atoms” means that the alkyl group may consist of 1 carbon atom, 2 carbon atoms, 3 carbon atoms, *etc.*, up to and including 20 carbon atoms, although the present definition also covers the occurrence of the term “alkyl” where no numerical range is designated). The alkyl group may also be a medium size alkyl having 1 to 9 carbon atoms. The alkyl group could also be a lower alkyl having 1 to 4 carbon atoms. The alkyl group may be designated as “ $\text{C}_{1-4}$  alkyl” or similar designations. By way of example only, “ $\text{C}_{1-4}$  alkyl” indicates that there are one to four carbon atoms in the alkyl chain, *i.e.*, the alkyl chain is selected from the group consisting of methyl, ethyl, propyl, isopropyl, n-butyl, iso-butyl, sec-butyl, and t-butyl. Typical alkyl groups include, but are in no way limited to, methyl, ethyl, propyl, isopropyl, butyl, isobutyl, tertiary butyl, pentyl, hexyl, and the like.

**[0097]** As used herein, “alkoxy” refers to the formula  $-\text{OR}$  wherein R is an alkyl as is defined above, such as “ $\text{C}_{1-9}$  alkoxy”, including but not limited to methoxy, ethoxy, n-propoxy, 1-methylethoxy (isopropoxy), n-butoxy, iso-butoxy, sec-butoxy, and tert-butoxy, and the like.

**[0098]** As used herein, “alkylthio” refers to the formula  $-\text{SR}$  wherein R is an alkyl as is defined above, such as “ $\text{C}_{1-9}$  alkylthio” and the like, including but not limited to methylmercapto, ethylmercapto, n-propylmercapto, 1-methylethylmercapto (isopropylmercapto), n-butylmercapto, iso-butylmercapto, sec-butylmercapto, tert-butylmercapto, and the like.

**[0099]** As used herein, “alkenyl” refers to a straight or branched hydrocarbon chain containing one or more double bonds. The alkenyl group may have 2 to 20 carbon atoms, although the present definition also covers the occurrence of the term “alkenyl” where

no numerical range is designated. The alkenyl group may also be a medium size alkenyl having 2 to 9 carbon atoms. The alkenyl group could also be a lower alkenyl having 2 to 4 carbon atoms. The alkenyl group may be designated as “C<sub>2-4</sub> alkenyl” or similar designations. By way of example only, “C<sub>2-4</sub> alkenyl” indicates that there are two to four carbon atoms in the alkenyl chain, i.e., the alkenyl chain is selected from the group consisting of ethenyl, propen-1-yl, propen-2-yl, propen-3-yl, buten-1-yl, buten-2-yl, buten-3-yl, buten-4-yl, 1-methyl-propen-1-yl, 2-methyl-propen-1-yl, 1-ethyl-ethen-1-yl, 2-methyl-propen-3-yl, buta-1,3-dienyl, buta-1,2,-dienyl, and buta-1,2-dien-4-yl. Typical alkenyl groups include, but are in no way limited to, ethenyl, propenyl, butenyl, pentenyl, and hexenyl, and the like.

**[0100]** As used herein, “alkynyl” refers to a straight or branched hydrocarbon chain containing one or more triple bonds. The alkynyl group may have 2 to 20 carbon atoms, although the present definition also covers the occurrence of the term “alkynyl” where no numerical range is designated. The alkynyl group may also be a medium size alkynyl having 2 to 9 carbon atoms. The alkynyl group could also be a lower alkynyl having 2 to 4 carbon atoms. The alkynyl group may be designated as “C<sub>2-4</sub> alkynyl” or similar designations. By way of example only, “C<sub>2-4</sub> alkynyl” indicates that there are two to four carbon atoms in the alkynyl chain, i.e., the alkynyl chain is selected from the group consisting of ethynyl, propyn-1-yl, propyn-2-yl, butyn-1-yl, butyn-3-yl, butyn-4-yl, and 2-butynyl. Typical alkynyl groups include, but are in no way limited to, ethynyl, propynyl, butynyl, pentynyl, and hexynyl, and the like.

**[0101]** As used herein, “heteroalkyl” refers to a straight or branched hydrocarbon chain containing one or more heteroatoms, that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur, in the chain backbone. The heteroalkyl group may have 1 to 20 carbon atom, although the present definition also covers the occurrence of the term “heteroalkyl” where no numerical range is designated. The heteroalkyl group may also be a medium size heteroalkyl having 1 to 9 carbon atoms. The heteroalkyl group could also be a lower heteroalkyl having 1 to 4 carbon atoms. The heteroalkyl group may be designated as “C<sub>1-4</sub> heteroalkyl” or similar designations. The heteroalkyl group may contain one or more heteroatoms. By way of example only, “C<sub>1-4</sub> heteroalkyl” indicates that there are one to four carbon atoms in the heteroalkyl chain and additionally one or more heteroatoms in the backbone of the chain.

**[0102]** As used herein, “alkylene” means a branched, or straight chain fully saturated di-radical chemical group containing only carbon and hydrogen that is attached to the rest of the molecule via two points of attachment (i.e., an alkanediyl). The alkylene group may have 1 to 20 carbon atoms, although the present definition also covers the occurrence of the term alkylene where no numerical range is designated. The alkylene group may also be a medium size alkylene having 1 to 9 carbon atoms. The alkylene group could also be a lower alkylene having 1 to 4 carbon atoms. The alkylene group may be designated as “C<sub>1-4</sub> alkylene” or similar designations. By way of example only, “C<sub>1-4</sub> alkylene” indicates that there are one to four carbon atoms in the alkylene chain, i.e., the alkylene chain is selected from the group consisting of methylene, ethylene, ethan-1,1-diyl, propylene, propan-1,1-diyl, propan-2,2-diyl, 1-methyl-ethylene, butylene, butan-1,1-diyl, butan-2,2-diyl, 2-methyl-propan-1,1-diyl, 1-methyl-propylene, 2-methyl-propylene, 1,1-dimethyl-ethylene, 1,2-dimethyl-ethylene, and 1-ethyl-ethylene.

**[0103]** As used herein, “alkenylene” means a straight or branched chain di-radical chemical group containing only carbon and hydrogen and containing at least one carbon-carbon double bond that is attached to the rest of the molecule via two points of attachment. The alkenylene group may have 2 to 20 carbon atoms, although the present definition also covers the occurrence of the term alkenylene where no numerical range is designated. The alkenylene group may also be a medium size alkenylene having 2 to 9 carbon atoms. The alkenylene group could also be a lower alkenylene having 2 to 4 carbon atoms. The alkenylene group may be designated as “C<sub>2-4</sub> alkenylene” or similar designations. By way of example only, “C<sub>2-4</sub> alkenylene” indicates that there are two to four carbon atoms in the alkenylene chain, i.e., the alkenylene chain is selected from the group consisting of ethenylene, ethen-1,1-diyl, propenylene, propen-1,1-diyl, prop-2-en-1,1-diyl, 1-methyl-ethenylene, but-1-enylene, but-2-enylene, but-1,3-dienylene, buten-1,1-diyl, but-1,3-dien-1,1-diyl, but-2-en-1,1-diyl, but-3-en-1,1-diyl, 1-methyl-prop-2-en-1,1-diyl, 2-methyl-prop-2-en-1,1-diyl, 1-ethyl-ethenylene, 1,2-dimethyl-ethenylene, 1-methyl-propenylene, 2-methyl-propenylene, 3-methyl-propenylene, 2-methyl-propen-1,1-diyl, and 2,2-dimethyl-ethen-1,1-diyl.

**[0104]** The term “aromatic” refers to a ring or ring system having a conjugated pi electron system and includes both carbocyclic aromatic (e.g., phenyl) and heterocyclic

aromatic groups (e.g., pyridine). The term includes monocyclic or fused-ring polycyclic (i.e., rings which share adjacent pairs of atoms) groups provided that the entire ring system is aromatic.

**[0105]** As used herein, “aryl” refers to an aromatic ring or ring system (i.e., two or more fused rings that share two adjacent carbon atoms) containing only carbon in the ring backbone. When the aryl is a ring system, every ring in the system is aromatic. The aryl group may have 6 to 18 carbon atoms, although the present definition also covers the occurrence of the term “aryl” where no numerical range is designated. In some embodiments, the aryl group has 6 to 10 carbon atoms. The aryl group may be designated as “C<sub>6-10</sub> aryl,” “C<sub>6</sub> or C<sub>10</sub> aryl,” or similar designations. Examples of aryl groups include, but are not limited to, phenyl, naphthyl, azulenyl, and anthracenyl.

**[0106]** As used herein, “aryloxy” and “arylthio” refers to RO- and RS-, in which R is an aryl as is defined above, such as “C<sub>6-10</sub> aryloxy” or “C<sub>6-10</sub> arylthio” and the like, including but not limited to phenyloxy.

**[0107]** An “aralkyl” or “arylalkyl” is an aryl group connected, as a substituent, via an alkylene group, such as “C<sub>7-14</sub> aralkyl” and the like, including but not limited to benzyl, 2-phenylethyl, 3-phenylpropyl, and naphthylalkyl. In some cases, the alkylene group is a lower alkylene group (i.e., a C<sub>1-4</sub> alkylene group).

**[0108]** As used herein, “heteroaryl” refers to an aromatic ring or ring system (i.e., two or more fused rings that share two adjacent atoms) that contain(s) one or more heteroatoms, that is, an element other than carbon, including but not limited to, nitrogen, oxygen and sulfur, in the ring backbone. When the heteroaryl is a ring system, every ring in the system is aromatic. The heteroaryl group may have 5-18 ring members (i.e., the number of atoms making up the ring backbone, including carbon atoms and heteroatoms), although the present definition also covers the occurrence of the term “heteroaryl” where no numerical range is designated. In some embodiments, the heteroaryl group has 5 to 10 ring members or 5 to 7 ring members. The heteroaryl group may be designated as “5-7 membered heteroaryl,” “5-10 membered heteroaryl,” or similar designations. Examples of heteroaryl rings include, but are not limited to, furyl, thienyl, phthalazinyl, pyrrolyl, oxazolyl, thiazolyl, imidazolyl, pyrazolyl, isoxazolyl, isothiazolyl, triazolyl, thiadiazolyl, pyridinyl, pyridazinyl,

pyrimidinyl, pyrazinyl, triazinyl, quinolinyl, isoquinlinyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, indolyl, isoindolyl, and benzothienyl.

**[0109]** A “heteroaralkyl” or “heteroarylalkyl” is heteroaryl group connected, as a substituent, via an alkylene group. Examples include but are not limited to 2-thienylmethyl, 3-thienylmethyl, furylmethyl, thienylethyl, pyrrolylalkyl, pyridylalkyl, isoxazolylalkyl, and imidazolylalkyl. In some cases, the alkylene group is a lower alkylene group (i.e., a C<sub>1-4</sub> alkylene group).

**[0110]** As used herein, “carbocyclyl” means a non-aromatic cyclic ring or ring system containing only carbon atoms in the ring system backbone. When the carbocyclyl is a ring system, two or more rings may be joined together in a fused, bridged or spiro-connected fashion. Carbocyclyls may have any degree of saturation provided that at least one ring in a ring system is not aromatic. Thus, carbocyclyls include cycloalkyls, cycloalkenyls, and cycloalkynyls. The carbocyclyl group may have 3 to 20 carbon atoms, although the present definition also covers the occurrence of the term “carbocyclyl” where no numerical range is designated. The carbocyclyl group may also be a medium size carbocyclyl having 3 to 10 carbon atoms. The carbocyclyl group could also be a carbocyclyl having 3 to 6 carbon atoms. The carbocyclyl group may be designated as “C<sub>3-6</sub> carbocyclyl” or similar designations. Examples of carbocyclyl rings include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cyclohexenyl, 2,3-dihydro-indene, bicycle[2.2.2]octanyl, adamantyl, and spiro[4.4]nonanyl.

**[0111]** A “(carbocyclyl)alkyl” is a carbocyclyl group connected, as a substituent, via an alkylene group, such as “C<sub>4-10</sub> (carbocyclyl)alkyl” and the like, including but not limited to, cyclopropylmethyl, cyclobutylmethyl, cyclopropylethyl, cyclopropylbutyl, cyclobutylethyl, cyclopropylisopropyl, cyclopentylmethyl, cyclopentylethyl, cyclohexylmethyl, cyclohexylethyl, cycloheptylmethyl, and the like. In some cases, the alkylene group is a lower alkylene group.

**[0112]** As used herein, “cycloalkyl” means a fully saturated carbocyclyl ring or ring system. Examples include cyclopropyl, cyclobutyl, cyclopentyl, and cyclohexyl.

**[0113]** As used herein, “cycloalkenyl” means a carbocyclyl ring or ring system having at least one double bond, wherein no ring in the ring system is aromatic. An example is cyclohexenyl.

**[0114]** As used herein, “heterocyclyl” means a non-aromatic cyclic ring or ring system containing at least one heteroatom in the ring backbone. Heterocyclyls may be joined together in a fused, bridged or spiro-connected fashion. Heterocyclyls may have any degree of saturation provided that at least one ring in the ring system is not aromatic. The heteroatom(s) may be present in either a non-aromatic or aromatic ring in the ring system. The heterocyclyl group may have 3 to 20 ring members (i.e., the number of atoms making up the ring backbone, including carbon atoms and heteroatoms), although the present definition also covers the occurrence of the term “heterocyclyl” where no numerical range is designated. The heterocyclyl group may also be a medium size heterocyclyl having 3 to 10 ring members. The heterocyclyl group could also be a heterocyclyl having 3 to 6 ring members. The heterocyclyl group may be designated as “3-6 membered heterocyclyl” or similar designations. In preferred six membered monocyclic heterocyclyls, the heteroatom(s) are selected from one up to three of O, N or S, and in preferred five membered monocyclic heterocyclyls, the heteroatom(s) are selected from one or two heteroatoms selected from O, N, or S. Examples of heterocyclyl rings include, but are not limited to, azepinyl, acridinyl, carbazolyl, cinnolinyl, dioxolanyl, imidazolynyl, imidazolidinyl, morpholinyl, oxiranyl, oxepanyl, thiepanyl, piperidinyl, piperazinyl, dioxopiperazinyl, pyrrolidinyl, pyrrolidonyl, pyrrolidionyl, 4-piperidonyl, pyrazolynyl, pyrazolidinyl, 1,3-dioxynyl, 1,3-dioxanyl, 1,4-dioxynyl, 1,4-dioxanyl, 1,3-oxathianyl, 1,4-oxathiinyl, 1,4-oxathianyl, 2*H*-1,2-oxazinyl, trioxanyl, hexahydro-1,3,5-triazinyl, 1,3-dioxolyl, 1,3-dioxolanyl, 1,3-dithiolyl, 1,3-dithiolanyl, isoxazolynyl, isoxazolidinyl, oxazolynyl, oxazolidinyl, oxazolidinonyl, thiazolynyl, thiazolidinyl, 1,3-oxathiolanyl, indolynyl, isoindolynyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, tetrahydro-1,4-thiazinyl, thiamorpholinyl, dihydrobenzofuranyl, benzimidazolidinyl, and tetrahydroquinoline.

**[0115]** A “(heterocyclyl)alkyl” is a heterocyclyl group connected, as a substituent, via an alkylene group. Examples include, but are not limited to, imidazolynylmethyl and indolynylethyl.

**[0116]** As used herein, “oxo” refers to =O.

**[0117]** As used herein, “acyl” refers to  $-C(=O)R$ , wherein R is hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and

5-10 membered heterocyclyl, as defined herein. Non-limiting examples include formyl, acetyl, propanoyl, benzoyl, and acryl.

**[0118]** An “O-carboxy” group refers to a “-OC(=O)R” group in which R is selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0119]** A “C-carboxy” group refers to a “-C(=O)OR” group in which R is selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein. A non-limiting example includes carboxyl (i.e., -C(=O)OH).

**[0120]** A “cyano” group refers to a “-CN” group.

**[0121]** A “cyanato” group refers to an “-OCN” group.

**[0122]** An “isocyanato” group refers to a “-NCO” group.

**[0123]** A “thiocyanato” group refers to a “-SCN” group.

**[0124]** An “isothiocyanato” group refers to an “-NCS” group.

**[0125]** A “sulfinyl” group refers to an “-S(=O)R” group in which R is selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0126]** A “sulfonyl” group refers to an “-SO<sub>2</sub>R” group in which R is selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0127]** An “S-sulfonamido” group refers to a “-SO<sub>2</sub>NR<sub>A</sub>R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0128]** An “N-sulfonamido” group refers to a “-N(R<sub>A</sub>)SO<sub>2</sub>R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0129]** An “O-carbamyl” group refers to a “-OC(=O)NR<sub>A</sub>R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl,

C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0130]** An “N-carbamyl” group refers to an “-N(R<sub>A</sub>)C(=O)OR<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0131]** An “O-thiocarbamyl” group refers to a “-OC(=S)NR<sub>A</sub>R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0132]** An “N-thiocarbamyl” group refers to an “-N(R<sub>A</sub>)C(=S)OR<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0133]** A “C-amido” group refers to a “-C(=O)NR<sub>A</sub>R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0134]** An “N-amido” group refers to a “-N(R<sub>A</sub>)C(=O)R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein.

**[0135]** An “amino” group refers to a “-NR<sub>A</sub>R<sub>B</sub>” group in which R<sub>A</sub> and R<sub>B</sub> are each independently selected from hydrogen, C<sub>1-6</sub> alkyl, C<sub>2-6</sub> alkenyl, C<sub>2-6</sub> alkynyl, C<sub>3-7</sub> carbocyclyl, C<sub>6-10</sub> aryl, 5-10 membered heteroaryl, and 5-10 membered heterocyclyl, as defined herein. A non-limiting example includes free amino (i.e., -NH<sub>2</sub>).

**[0136]** An “aminoalkyl” group refers to an amino group connected via an alkylene group.


**[0137]** An “alkoxyalkyl” group refers to an alkoxy group connected via an alkylene group, such as a “C<sub>2-8</sub> alkoxyalkyl” and the like.




**[0138]** As used herein, a substituted group is derived from the unsubstituted parent group in which there has been an exchange of one or more hydrogen atoms for another atom or group. Unless otherwise indicated, when a group is deemed to be “substituted,” it is meant that the group is substituted with one or more substituents independently selected from C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkenyl, C<sub>1</sub>-C<sub>6</sub> alkynyl, C<sub>1</sub>-C<sub>6</sub> heteroalkyl, C<sub>3</sub>-C<sub>7</sub> carbocyclyl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), C<sub>3</sub>-C<sub>7</sub>-carbocyclyl-C<sub>1</sub>-C<sub>6</sub>-alkyl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), 5-10 membered heterocyclyl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), 5-10 membered heterocyclyl-C<sub>1</sub>-C<sub>6</sub>-alkyl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), aryl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), aryl(C<sub>1</sub>-C<sub>6</sub>)alkyl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), 5-10 membered heteroaryl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), 5-10 membered heteroaryl(C<sub>1</sub>-C<sub>6</sub>)alkyl (optionally substituted with halo, C<sub>1</sub>-C<sub>6</sub> alkyl, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> haloalkyl, and C<sub>1</sub>-C<sub>6</sub> haloalkoxy), halo, cyano, hydroxy, C<sub>1</sub>-C<sub>6</sub> alkoxy, C<sub>1</sub>-C<sub>6</sub> alkoxy(C<sub>1</sub>-C<sub>6</sub>)alkyl (i.e., ether), aryloxy, sulfhydryl (mercapto), halo(C<sub>1</sub>-C<sub>6</sub>)alkyl (e.g., -CF<sub>3</sub>), halo(C<sub>1</sub>-C<sub>6</sub>)alkoxy (e.g., -OCF<sub>3</sub>), C<sub>1</sub>-C<sub>6</sub> alkylthio, arylthio, amino, amino(C<sub>1</sub>-C<sub>6</sub>)alkyl, nitro, O-carbamyl, N-carbamyl, O-thiocarbamyl, N-thiocarbamyl, C-amido, N-amido, S-sulfonamido, N-sulfonamido, C-carboxy, O-carboxy, acyl, cyanato, isocyanato, thiocyanato, isothiocyanato, sulfinyl, sulfonyl, and oxo (=O). Wherever a group is described as “optionally substituted” that group can be substituted with the above substituents.

**[0139]** It is to be understood that certain radical naming conventions can include either a mono-radical or a di-radical, depending on the context. For example, where a substituent requires two points of attachment to the rest of the molecule, it is understood that the substituent is a di-radical. For example, a substituent identified as alkyl that requires two points of attachment includes di-radicals such as -CH<sub>2</sub>-, -CH<sub>2</sub>CH<sub>2</sub>-, -CH<sub>2</sub>CH(CH<sub>3</sub>)CH<sub>2</sub>-, and the like. Other radical naming conventions clearly indicate that the radical is a di-radical such as “alkylene” or “alkenylene.”

**[0140]** When two R groups are said to form a ring (e.g., a carbocyclyl, heterocyclyl, aryl, or heteroaryl ring) “together with the atom to which they are attached,” it is meant that the collective unit of the atom and the two R groups are the recited ring. The ring is not otherwise limited by the definition of each R group when taken individually. For example, when the following substructure is present:




and R<sup>1</sup> and R<sup>2</sup> are defined as selected from the group consisting of hydrogen and alkyl, or R<sup>1</sup> and R<sup>2</sup> together with the nitrogen to which they are attached form a heteroaryl, it is meant that R<sup>1</sup> and R<sup>2</sup> can be selected from hydrogen or alkyl, or alternatively, the substructure has structure:




where ring A is a heteroaryl ring containing the depicted nitrogen.

**[0141]** Similarly, when two “adjacent” R groups are said to form a ring “together with the atoms to which they are attached,” it is meant that the collective unit of the atoms, intervening bonds, and the two R groups are the recited ring. For example, when the following substructure is present:




and R<sup>1</sup> and R<sup>2</sup> are defined as selected from the group consisting of hydrogen and alkyl, or R<sup>1</sup> and R<sup>2</sup> together with the atoms to which they are attached form an aryl or carbocyclyl, it is meant that R<sup>1</sup> and R<sup>2</sup> can be selected from hydrogen or alkyl, or alternatively, the substructure has structure:



where A is an aryl ring or a carbocyclyl containing the depicted double bond.

**[0142]** Wherever a substituent is depicted as a di-radical (*i.e.*, has two points of attachment to the rest of the molecule), it is to be understood that the substituent can be attached in any directional configuration unless otherwise indicated. Thus, for example, a

substituent depicted as  $-AE-$  or  includes the substituent being oriented such that the A is attached at the leftmost attachment point of the molecule as well as the case in which A is attached at the rightmost attachment point of the molecule.

**[0143]** “Leaving group,” or “LG,” as used herein refers to any atom or moiety that is capable of being displaced by another atom or moiety in a chemical reaction. More specifically, in some embodiments, “leaving group” refers to the atom or moiety that is displaced in a nucleophilic substitution reaction. In some embodiments, “leaving groups” are any atoms or moieties that are conjugate bases of strong acids. Examples of suitable leaving groups include, but are not limited to, tosylates and halogens. Non-limiting characteristics and examples of leaving groups can be found, for example in Organic Chemistry, 2d ed., Francis Carey (1992), pages 328-331; Introduction to Organic Chemistry, 2d ed., Andrew Streitwieser and Clayton Heathcock (1981), pages 169-171; and Organic Chemistry, 5th ed., John McMurry (2000), pages 398 and 408; all of which are incorporated herein by reference for the limited purpose of disclosing characteristics and examples of leaving groups.

**[0144]** Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which the embodiments belong. Although any methods and materials similar or equivalent to those described herein can also be used in the practice or testing of the embodiments, the preferred methods and materials are now described. All publications mentioned herein are incorporated herein by reference to disclose and describe the methods and/or materials in connection with which the publications are cited.

### Protecting Groups

**[0145]** In some circumstances, a chemical reaction may need to be performed selectively at one reactive site in a multifunctional compound. One such method that is useful for accomplishing such selectivity is to temporarily block one or more reactive sites in the multifunctional compound with a protective group. Such a method is often referred to as “protecting” the functional group. Many protecting groups are known in the art. See, e.g., Greene et al., *Protective Groups in Organic Synthesis*, Third Ed. (John Wiley & Sons, Inc. 1999), herein incorporated by reference in its entirety; Wutz et al., *Greene’s Protective Groups in Organic Synthesis*, Fourth Ed. (John Wiley & Sons, Inc. 2007), herein incorporated by reference in its entirety. When more than one reactive site in a multifunctional compound requires protecting, or when a compound is prepared that will possess more than one protected functional group, it is important to use orthogonal protecting groups. Protecting groups are orthogonal if they are susceptible to selective removal.

**[0146]** In some embodiments, it may be necessary to protect one or more functional groups so as to prevent their interference in the desired reaction. For example, it may be necessary to protect one or more functional groups such as amines, carboxylic acids, and/or hydroxyl groups.

**[0147]** Suitable protecting groups for protecting amines include: carbamates such as alkyl carbamates including methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, pentyl, neopentyl, hexyl, heptyl, nonnyl, decanyl, and configurational isomers thereof; 9-fluorenylmethyl; 9-(2-sulfo)fluorenylmethyl; 9-(2,7-dibromo)fluorenylmethyl; 17-tetrabenzo[a,c,g,i]fluorenylmethyl; 2-chloro-3-indenylmethyl; benz[f]inden-3-ylmethyl; 2,7-di-t-butyl[9-(10,10-dioxo-10,10,10,10-tetrahydrothioxanthyl)]methyl; 1,1-dioxobenzo[b]thiophene-2-ylmethyl; substituted ethyl carbamates such as 2,2,2-trichloroethyl; 2-trimethylsilylethyl; 2-phenylethyl; 1-(1-adamantyl)-1-methylethyl; 2-chloroethyl; 1,1-dimethyl-2-haloethyl; 1,1-dimethyl,2,2-dibromoethyl; 1,1-dimethyl-2,2,2-trichloroethyl; 1-methyl-1-(4-biphenyl)ethyl; 1-(3,5-di-t-butylphenyl)-1-methylethyl; 2-(2'- and 4'-pyridyl)ethyl; N-(2-pivaloylamino)-1,1-dimethylethyl; 2-[(2-nitrophenyl)dithio]-1-phenylethyl; 2-(N,N-dicyclohexylcarboxamido)ethyl; t-butyl; 1-adamantyl; 2-adamantyl; vinyl; allyl; 1-isopropylallyl; cinnamyl; 4-nitrocinnamyl; 3-(3'-pyridyl)prop-2-enyl; 8-quinolyl; N-hydroxypiperidinyl; alkydithio; benzyl; p-methoxybenzyl; p-nitrobenzyl; p-

bromobenzyl; p-chlorobenzyl; 2,4-dichlorobenzyl; 4-methylsulfinylbenzyl; 9-anthrylmethyl; diphenylmethyl; 2-methylthioethyl; 2-methylsulfonyl; 2-(p-toluenesulfonyl)ethyl; [2-(1,3-dithianyl)]methyl; 4-methylthiophenyl; 2,4-dimethylthiophenyl; 2-phosphonioethyl; 1-methyl-1-(triphenylphosphonio)ethyl; 1,1-dimethyl-2-cyanoethyl; 2-dansylethyl; 2-(4-nitrophenyl)ethyl; 4-phenylacetoxybenzyl; 4-azidobenzyl; 4-azidomethoxybenzyl; m-chloro-p-acyloxybenzyl; p-(dihydroxyboryl)benzyl; 5-benzisoxazolylmethyl; 2-(trifluoromethyl)-6-chromonylmethyl; m-nitrophenyl; 3,5-dimethoxybenzyl; 1-methyl-1-(3,5-dimethoxyphenyl)ethyl;  $\alpha$ -methylnitropiperonyl; o-nitrobenzyl; 3,4-dimethoxy-6-nitrobenzyl; phenyl(o-nitrophenyl)methyl; 2-(2-nitrophenyl)ethyl; 6-nitroveratryl; 4-methoxyphenacyl; 3',5'-dimethoxybenzoin; phenothiazinyl-(10)-carbonyl derivatives; N'-p-toluenesulfonylaminocarbonyl; N'-phenylaminothiocarbonyl; t-amyl; S-benzyl thiocarbamate; butynyl; p-cyanobenzyl; cyclobutyl; cyclohexyl; cyclopentyl; cyclopropylmethyl; p-dicyloxybenzyl; diisopropylmethyl; 2,2-dimethoxycarbonylvinyl; o-(N',N'-dimethylcarboxamido)benzyl; 1,1-dimethyl-3-(N',N'-dimethylcarboxamido)propyl; 1,1-dimethylpropynyl; di(2-pyridyl)methyl; 2-furanylmethyl; 2-iodoethyl; isobornyl; isobutyl; isonicotinyl; p-(p'-methoxyphenylazo)benzyl; 1-methylcyclobutyl; 1-methylcyclohexyl; 1-methyl-1-cyclopropylmethyl; 1-methyl-1-(p-phenylazophenyl)ethyl; 1-methyl-1-phenylethyl; 1-methyl-1-(4'-pyridyl)ethyl; phenyl; p-(phenylazo)benzyl; 2,4,6-tri-t-butylphenyl; 4-(trimethylammonium)benzyl; 2,4,6-trimethylbenzyl; and other similar carbamates; amides, including, but not limited to, formyl, acetyl, chloroacetyl, trichloroacetyl, trifluoroacetyl, phenylacetyl, propionyl, 3-phenylpropionyl, 4-pentenoyl, picolinoyl, 3-pyridylcarboxamide, benzoylphenylalanyl, benzoyl, p-phenylbenzoyl, amides whose cleavage is induced by nitro group reduction, such as o-nitrophenylacetyl, o-nitrophenoxycetyl, 3-(o-nitrophenyl)propionyl, 2-methyl-2-(o-nitrophenoxycetyl)propionyl, 3-methyl-3-nitrobutyryl, o-nitrocinnamoyl, o-nitrobenzoyl, and 3-(4-t-butyl-2,6-dinitrophenyl)-2,2-dimethylpropionyl; amides whose cleavage is induced by release of an alcohol, such as o-(benzoyloxymethyl)benzoyl, (2-acetoxymethyl)benzoyl, 2-[(t-butyl)diphenylsiloxy)methyl]benzoyl, 3-(3',6'-dioxo-2',4',5'-trimethylcyclohexa-1',4'-diene-3,3-dimethylpropionyl, and o-hydroxy-trans-cinnamoyl; amides whose cleavage is induced by other chemical reactions, such as 2-methyl-2-(o-phenylazophenoxy)propionyl, 4-chlorobutyryl, acetoacetyl, 3-(p-hydroxyphenyl)propionyl, (N'-

dithiobenzyloxycarbonylamino)acetyl, N-acetylmethionine, and 4,5-diphenyl-3-oxazolin-2-one; cyclic imide derivatives such as N-phthaloyl, N-tetrachlorophthaloyl, N-4-nitrophthaloyl, N-dithiasuccinoyl, N-2,3-diphenylmaleoyl, N-2,5-dimethylpyrrolyl, N-2,5-bis(triisopropylsiloxy)pyrrolyl, N-1,1,4,4-tetramethyldisilylazacyclopentane adduct, N-1,1,3,3-tetramethyl-1,3-disilaisoindolyl, 5-substituted 1,3-dimethyl-1,3,5-triazacyclohexan-2-one, 5-substituted 1,3-dibenzyl-1,3,5-triazacyclohexan-2-one, 1-substituted 3,5-dinitro-4-pyridonyl, and 1,3,5-dioxazinyl; N-alkyl and N-aryl derivatives, such as N-methyl, N-t-butyl, N-allyl, N-[2-(trimethylsilyl)ethoxy]methyl, N-3-acetoxypropyl, N-cyanomethyl, N-(1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl), N-2,4-dimethoxybenzyl, N-2-azanorbornenyl, N-2,4-dinitrophenyl, quaternary ammonium salts, N-benzyl, N-4-methoxybenzyl, N-2,4-dimethoxybenzyl, N-2-hydroxybenzyl, N-diphenylmethyl, N-bis(4-methoxyphenyl)methyl, N-5-dibenzosuberyl, N-triphenylmethyl, N-(4-methoxyphenyl)diiphenylmethyl, N-9-phenylfluorenyl, N-ferrocenylmethyl, and N-2-picolylamine N'-oxide; imine derivatives, such as N-1,1-dimethylthiomethylene, N-benzylidene, N-p-methoxybenzylidene, N-diphenylmethylene, N-[(2-pyridyl)mesityl]methylene, N-(N',N'-dimethylaminomethylene), N-(N',N'-dibenzylaminomethylene), N-(N'-t-butylaminomethylene), N,N'-isopropylidene, N-p-nitrobenzylidene, N-salicylidene, N-5-chlorosalicylidene, N-(5-chloro-2-hydroxyphenyl)phenylmethylene, N-cyclohexylidene, and N-t-butylidene; enamine derivatives, such as N-(5,5-dimethyl-3-oxo-1-cyclohexenyl), N-2,7-dichloro-9-fluorenylmethylene, N-2-(4,4-dimethyl-2,6-dioxocyclohexylidene)ethyl, N-4,4,4-trifluoro-3-oxo-1-butenyl, and N-1-isopropyl-4-nitro-2-oxo-3-pyrrolin-3-yl; and N-heteroatom derivatives such as N-metal, N-borane, N-diphenylborinic acid, N-diethylborinic acid, N-difloroborinic acid, N,N'-3,5-bis(trifluoromethyl)phenylboronic acid, N-[phenyl(pentacarbonylchromium-or-tungsten)]carbonyl, N-copper chelates, N-zinc chelates, and 18-crown-6 derivatives, N-N derivatives such as N-nitro, N-nitroso, N-oxide, and triazene derivatives, N-P derivatives such as N-diphenylphosphinyl, N-dimethylthiophosphinyl, N-diphenylthiophosphinyl, N-dialkylphosphoryl, N-dibenzylphosphoryl, N-diphenylphosphoryl, and iminotriphenylphosphorane derivatives, N-Si derivatives, N-sulfenyl derivatives such as N-benzenesulfonyl, N-o-nitrobenzenesulfonyl, N-2,4-dinitrobenzenesulfonyl, N-pentachlorobenzenesulfonyl, N-2-nitro-4-methoxybenzenesulfonyl, N-triphenylmethylsulfenyl, N-1-(2,2,2-trifluoro-1,1-

diphenyl)ethylsulfenyl, and N-3-nitro-2-pyridinesulfenyl, and/or N-sulfonyl derivatives such as N-p-toluenesulfonyl, N-benzenesulfonyl, N-2,3,6-trimethyl-4-methoxybenzenesulfonyl, N-2,4,6-trimethoxybenzenesulfonyl, N-2,6-dimethyl-4-methoxybenzenesulfonyl, N-pentamethylbenzenesulfonyl, N-2,3,5,6-tetramethyl-4-methoxybenzenesulfonyl, N-4-methoxybenzenesulfonyl, N-2,4,6-trimethylbenzenesulfonyl, N-2,6-dimethoxy-4-methylbenzenesulfonyl, N-3-methoxy-4-t-butylbenzenesulfonyl, N-2,2,5,7,8-pentamethylchroman-6-sulfonyl, N-2-nitrobenzenesulfonyl, N-4-nitrobenzenesulfonyl, N-2,4-dinitrobenzenesulfonyl, N-benzothiazole-2-sulfonyl, N-methanesulfonyl, N-2-(trimethylsilyl)ethanesulfonyl, N-9-anthracenesulfonyl, N-4-(4',8'-dimethoxynaphthylmethyl)benzenesulfonyl, N-benzylsulfonyl, N-trifluoromethylsulfonyl, N-phenacysulfonyl, and N-t-butylsulfonyl.

**[0148]** Suitable protecting groups for carboxylic acids include: esters such as enzymatically cleavable esters including heptyl, 2-N-(morpholino)ethyl, choline, (methoxyethoxy)ethyl, methoxyethyl; alkyl esters such as methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, pentyl, neopentyl, hexyl, heptyl, nonnyl, decanyl, and configurational isomers thereof; substituted methyl esters such as 9-fluorenylmethyl, methoxymethyl, methylthiomethyl, tetrahydropyranyl, tetrahydrofuranlyl, methoxyethoxymethyl, 2-(trimethylsilyl)ethoxymethyl, benzyloxymethyl, pivaloyloxymethyl, phenylacetoxymethyl, triisopropylsilylmethyl, cyanomethyl, acetol, phenacetyl, p-bromophenacetyl,  $\alpha$ -methylphenacetyl, p-methoxyphenacetyl, desyl, carboamidomethyl, p-azobenzenecarboxamidomethyl, N-phthalidimidomethyl; 2-substituted ethyl esters such as 2,2,2-trichloroethyl, 2-haloethyl,  $\omega$ -chloroalkyl, 2-(trimethylsilyl)ethyl, 2-methylthioethyl, 1,3-dithianyl-2-methyl, 2-(p-nitrophenylsulfenyl)ethyl, 2-(p-toluenesulfonyl)ethyl, 2-(2'-pyridyl)ethyl, 2-(p-methoxyphenyl)ethyl, 2-(diphenylphosphino)ethyl, 1-methyl-1-phenylethyl, 2-(4-acetyl-2-nitrophenyl)ethyl, 2-cyanoethyl, 3-methyl-3-pentyl, dicyclopropylmethyl, 2,4-dimethyl-3-pentyl, cyclopentyl, cyclohexyl, allyl, methallyl, 2-methylbut-3-en-2-yl, 3-methylbut-2-(prenyl), 3-buten-1-yl, 4-(trimethylsilyl)-2-buten-1-yl, cinnamyl,  $\alpha$ -methylcinnamyl, prop-2-ynyl, phenyl; 2,6-dialkylphenyl esters such as 2,6-dimethylphenyl, 2,6-diisopropylphenyl, 2,6-di-t-butyl-4-methylphenyl, 2,6-di-t-butyl-4-methoxyphenyl, p-(methylthio)phenyl, pentafluorophenyl, benzyl; substituted benzyl esters such as triphenylmethyl, diphenylmethyl, bis(o-

mitrophenyl)methyl, 9-anthrylmethyl, 2-(9,10-dioxo)anthrylmethyl, 5-dibenzosuberyl, 1-pyreneylmethyl, 2-(trifluoromethyl)-6-chromonylmethyl, 2,4,6-trimethylbenzyl, p-bromobenzyl, o-nitrobenzyl, p-nitrobenzyl, p-methoxybenzyl, 2,6-dimethoxybenzyl, 4-(methylsulfinyl)benzyl, 4-sulfobenzyl, 4-azidomethoxybenzyl, 4-{N-[1-(4,4-dimethyl-2,6-dioxocyclohexylidene)-3-methylbutyl]amino}benzyl, piperonyl, 4-picolyl, polymer supported p-benzyl; silyl esters such as trimethylsilyl, triethylsilyl, t-butyl dimethylsilyl, i-propyldimethylsilyl, phenyldimethylsilyl, di-t-butylmethylsilyl, triisopropylsilyl; activated esters such as thiol esters; oxazoles; 2-alkyl-1,3-axazoline; 4-alkyl-5-oxo-1,3-oxazolidine; 2,2-bistrifluoromethyl-4-alkyl-5-oxo-1,3-oxazolidine; 5-alkyl-4-oxo-1,3-dioxolane; dioxanones; ortho esters; pentaaminocobalt(III) complexes; and stannyl esters such as triethylstannyl and tri-n-butylstannyl; amides such as N,N-dimethyl, pyrrolidinyl, piperidinyl, 5,6-dihydrophenanthridinyl, o-nitroanilide, N-7-nitroindolyl, N-8-nitro-1,2,3,4-tetrahydroquinolyl, 2-(2-aminophenyl)acetaldehyde dimethyl acetal amide, and polymer supported p-benzenesulfonamide; hydrazides such as N-phenyl, N,N'-diisopropyl; and tetraalkylammonium salts such as methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, pentyl, neopentyl, hexyl, heptyl, nonnyl, decanyl, and configurational isomers thereof.

**[0149]** Suitable protecting groups for hydroxyl groups include: silyl ethers such as trimethylsilyl, triethylsilyl, triisopropylsilyl, dimethylisopropylsilyl, diethylisopropylsilyl, dimethylthexylsilyl, 2-norbornyldimethylsilyl, t-butyl-dimethylsilyl, t-butyl diphenylsilyl, tribenzylsilyl, tri-p-xylylsilyl, triphenylsilyl, diphenylmethylsilyl, di-t-butylmethylsilyl, bis(t-butyl)-1-pyrenylmethoxysilyl, tris(trimethylsilyl)silyl; silyl; (2-hydroxystyryl)dimethylsilyl; (2-hydroxystyryl)diisopropylsilyl, t-butylmethoxyphenylsilyl, t-butoxydiphenylsilyl, 1,1,3,3-tetraisopropyl-3-[2-(triphenylmethoxy)ethoxy]disiloxane-1-yl, fluorous silyl; C<sub>1-10</sub>alkyl ethers such as methyl, ethyl, propyl, isopropyl, butyl, tert-butyl, sec-butyl, pentyl, neopentyl, hexyl, heptyl, nonnyl, decanyl, and configurational isomers thereof; substituted methyl ethers such as methoxymethyl, methylthiomethyl, (phenyldimethylsilyl)methoxymethyl, benzyloxymethyl, p-methoxybenzyloxymethyl, [(3,4-dimethoxybenzyl)oxy]methyl, p-nitrobenzyloxymethyl, o-nitrobenzyloxymethyl, [(R)-1-(2-nitrophenyl)ethoxy]methyl, (4-methoxyphenoxy)methyl, guaiacolmethyl, t-butoxymethyl, 4-pentenylloxymethyl, siloxymethyl, 2-methoxyethoxymethyl, 2-cyanoethoxymethyl, 2,2,2-trichloroethoxymethyl, bis(2-chloroethoxy)methyl, 2-(trimethylsilyl)ethoxymethyl, methoxymethyl, O-Bis(2-



acetoxyethoxy)methyl, tetrahydropyranyl, fluorous tetrahydropyranyl, 3-bromotetrahydropyranyl, tetrahydrothiopyranyl, 1-methoxycyclohexyl, 4-methoxytetrahydropyranyl, 4-methoxytetrahydrothiopyranyl, 4-methoxytetrahydrothiopyranyl S,S-dioxide, 1-[(2-chloro-4-methyl)phenyl]-4-methoxypiperidin-4-yl, 1-(2-fluorophenyl)-4-methoxypiperidin-4-yl, 1-(4-chlorophenyl)-4-methoxypiperidin-4-yl, 1,4-dioxan-2-yl, tetrahydrofuranyl, tetrahydrothiofuranyl, and 2,3,2a,4,5,6,7,7a-octahydro-7,8,8-trimethyl-4,7-methanobenzofuran-2-yl; substituted ethyl ethers such as 1-ethoxyethyl, 1-(2-chloroethoxy)ethyl, 2-hydroxyethyl, 2-bromoethyl, 1-[2-(trimethylsilyl)ethoxy]ethyl, 1-methyl-1-methoxyethyl, 1-methyl-1-benzyloxyethyl, 1-methyl-1-benzyloxy-2-fluoroethyl, 1-methyl-1-phenoxyethyl, 2,2,2-trichloroethyl, 1,1-dianisyl-2,2,2-trichloroethyl, 1,1,1,3,3,3-hexafluoro-2-phenylisopropyl, 1-(2-cyanoethoxy)ethyl, 2-trimethylsilylethyl, 2-(benzylthio)ethyl, 2-(phenylselenyl)ethyl, t-butyl, allyl, prenyl, cinnamyl, 2-phenallyl, propargyl, p-chlorophenyl, p-methoxyphenyl, p-nitrophenyl, 2,4-dinitrophenyl, 2,3,5,6-tetrafluoro-4-(trifluoromethyl)phenyl; benzyl; substituted benzyl ethers such as p-methoxybenzyl, 3,4-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, pentadienyl nitrobenzyl, pentadienyl nitropiperonyl, halobenzyl, 2,6-dichlorobenzyl, 2,4-dichlorobenzyl, p-cyanobenzyl, p-phenylbenzyl, 2,6-difluorobenzyl, fluorous benzyl, 4-fluorousalkoxybenzyl, trimethylsilylxylyl, 2-phenyl-2-propyl (Cumyl), p-acylaminobenzyl, p-azidobenzyl, 4-azido-3-chlorobenzyl, 2-trifluoromethylbenzyl, 4-trifluoromethylbenzyl, p-(methylsulfinyl)benzyl, p-silylbenzyl, 4-acetoxybenzyl, 4-(2-trimethylsilyl)ethoxymethoxybenzyl, 2-naphthylmethyl, 2-picolyl, 4-picolyl, 3-methyl-2-picolyl N-oxide, 2-quinolinylmethyl, 6-methoxy-2-(4-methylphenyl)-4-quinolinemethyl, 1-pyrenylmethyl, diphenylmethyl, 4-methoxydiphenylmethyl, 4-phenyldiphenylmethyl, p,p'-dinitrobenzhydryl, 5-dibenzosuberyl, triphenylmethyl,  $\alpha$ -naphthyl diphenylmethyl, p-methoxyphenyldiphenylmethyl, di(p-methoxyphenyl)phenylmethyl, tri(p-methoxyphenyl)methyl, 4-(4'-bromophenacyloxy)phenyldiphenylmethyl, 4,4',4''-tris(4,5-dichlorophthalimidophenyl)methyl, 4,4',4''-tris(levulinoyloxyphenyl)methyl, 4,4',4''-tris(benzoyloxyphenyl)methyl, 4,4'-dimethoxy-3''-[N-imidazolylmethyl]]trityl, 4,4'-dimethoxy-3''-[N-imidazolylethyl]carbamoyle]trityl, 1,1-bis(4-methoxyphenyl)-1'-pyrenylmethyl, 4-(17-tetrabenzo[a,c,g,i]fluorenylmethyl)-4,4''-dimethoxytrityl, 9-anthryl, 9-(9-phenyl)xanthenyl, 9-phenylthioxanthyl, 9-(9-phenyl-10-oxo)anthryl, 1,3-benzodithiolan-

2-yl, 4,5-bis(ethoxycarbonyl)-[1,3]-dioxolan-2-yl, benzisothiazolyl S,S-dioxido; C<sub>1-10</sub>alkyl esters such as formyl, acetyl, propionyl, isopropionyl, butyryl, tert-butyryl, sec-butyryl, pentanoyl, neopentanoyl, hexanoyl, heptanoyl, nonanoyl, decanoyl, and configurational isomers thereof, esters such as benzoylformate, chloroacetate, dichloroacetate, trichloroacetate, trifluoroacetate, methoxyacetate, triphenylmethoxyacetate, phenoxyacetate, p-chlorophenoxyacetate, phenylacetate, polymer supported p-phenylacetate, diphenylacetate, bisfluorous chain type propanoyl, nicotinate, 3-phenylpropionate, 4-pentenoate, 4-oxopentanoate, 4,4-(ethylenedithio)pentanoate, 5-[3-bis(4-methoxyphenyl)hydroxymethylphenoxy]levulinate, pivaloate, 1-adamantoate, crotonate, 4-methoxycrotonate, benzoate, p-phenylbenzoate, 2,4,6-trimethylbenzoate, picolinate, nicotinate, 4-bromobenzoate, 2,5-difluorobenzoate, p-nitrobenzoate, 2,6-dichloro-4-methylphenoxyacetate, 2,6-dichloro-4-(1,1,3,3-tetramethylbutyl)phenoxyacetate, 2,4-bis(1,1-dimethylpropyl)phenoxyacetate, chlorodiphenylacetate, isobutyrate, monosuccinoate, 2-methyl-2-butenolate, (E)-2-methyl-2-butenolate, (Z)-2-methyl-2-butenolate, o-(methoxycarbonyl)benzoate, polymer supported p-benzoate,  $\alpha$ -naphthoate, nitrate, alkyl N,N,N',N'-tetramethylphosphorodiamidate, 2-chlorobenzoate, 3',5'-dimethoxybenzoin, N-phenylcarbamate, borate, dimethylphosphinothioyl, 2,4-dinitrophenylsulfenate, and photolabile esters; carbonates, including methyl, methoxymethyl, 9-fluorenylmethyl, ethyl, 2,2,2-trichloroethyl, 1,1-dimethyl-2,2,2-trichloroethyl, 2-(trimethylsilyl)ethyl, 2-(triphenylsulfonyl)ethyl, 2-(triphenylphosphonia)ethyl, isobutyl, vinyl, allyl, p-nitrophenyl, benzyl, p-methoxybenzyl, 3,3-dimethoxybenzyl, o-nitrobenzyl, p-nitrobenzyl, and silyl esters; carbonates cleaved by  $\beta$ -elimination such as 2-dansylethyl, 2-(4-nitrophenyl)ethyl, 2-(2,4-dinitrophenyl)ethyl, 2-cyano-1-phenylethyl, S-benzyl thiocarbonate, 4-ethoxy-1-mapthyl, and methyl dithiocarbonate, carbonates cleaved with assisted cleavage such as 2-iodobenzoate, 4-azidobutyrate, 4-nitro-4-methylpentanoate, o-(dibromomethyl)benzoate, 2-formylbenzenesulfonate, 2-(methylthiomethoxy)ethyl, 4-(methylthiomethoxymethyl)butyrate, 2-(methylthiomethoxymethyl)benzoate, 2-(chloroacetoxymethyl)benzoate, 2-[(2-chloroacetoxy)ethyl]benzoate, 2-[2-(benzyloxy)ethyl]benzoate, 2-[2-(4-methoxybenzyloxy)ethyl]benzoate; and sulfonates such as sulfate, allylsulfate, C<sub>1-10</sub>alkyl sulfonates such as methyl, ethyl, propyl, isopropyl, butyl,


tert-butyl, sec-butyl, pentyl, neopentyl, hexyl, heptyl, nonnonyl, decanyl, and configurational isomers thereof, benzylsulfonate, tosylate, and 2-[(4-nitrophenyl)ethyl]sulfonate.

## Protection and Deprotection Reactions

**[0150]** Reagents, solvents, and reaction conditions useful for protecting amines, carboxylic acids, and alcohols are well-known in the art. Likewise, reagents, solvents, and reaction conditions useful for deprotecting amines, carboxylic acids, and alcohols are well known in the art. See, e.g., Greene et al., *Protective Groups in Organic Synthesis*, Third Ed. (John Wiley & Sons, Inc. 1999), herein incorporated by reference in its entirety; Wutz et al., *Greene's Protective Groups in Organic Synthesis*, Fourth Ed. (John Wiley & Sons, Inc. 2007), herein incorporated by reference in its entirety. While references have been made to specific reagents, solvents, and reaction conditions in the schemes described above, it is readily envisioned that equivalent reagents, solvents, and reaction conditions may be utilized to protect and deprotect amines, carboxylic acids, and alcohols.

### Intermediate Compounds

**[0151]** Some embodiments disclosed herein include intermediates in the synthetic methods described herein, including a compound having the structure of Formula (I):



or a salt thereof.

**[0152]** In some embodiments, R<sub>1</sub> and R<sub>2</sub> are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted C<sub>1-4</sub> alkyl, or R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O.

**[0153]** In some embodiments, R<sub>3</sub> is selected from the group consisting of H, optionally substituted phenyl and optionally substituted C<sub>1-4</sub> alkyl, R<sub>3</sub> and R<sub>5</sub> together with the atoms to which they are attached form heteroaryl ring, or R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

**[0154]** In some embodiments,  $R_4$  is selected from the group consisting of H, optionally substituted phenyl, optionally substituted  $C_{1-4}$  alkyl, or  $R_3$  and  $R_4$  together with the atoms to which they are attached, form  $=O$ .

**[0155]** In some embodiments,  $R_5$ ,  $R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl.

**[0156]** In some embodiments,  $R_9$  is selected from the group consisting of optionally substituted  $C_1-C_{12}$  alkyl, optionally substituted  $C_3-C_8$  cycloalkyl, optionally substituted  $C_2-C_{12}$  alkenyl, optionally substituted  $C_2-C_{12}$  alkynyl, optionally substituted aryl and optionally substituted heteroaryl.

**[0157]** In some embodiments, independently, two germinal  $R_1$ ,  $R_2$ ,  $R_3$ , and  $R_4$  together with the atoms to which they are attached, form  $=O$ .

**[0158]** In some embodiments,  $Y_1$  is O. In some embodiments,  $Y_1$  is  $N^+R_5R_6$ .

**[0159]** In some embodiments,  $Y_2$  is O. In some embodiments,  $Y_2$  is  $NR_{10}$ .

**[0160]** In some embodiments,  $n$  is 0. In some embodiments,  $n$  is 1.

**[0161]** In some embodiments,  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ , and  $R_6$ , are hydrogen.

**[0162]** In some embodiments,  $R_1$  is optionally substituted phenyl. In some embodiments,  $R_1$  is optionally substituted  $C_1-C_4$  alkyl. In some embodiments,  $R_2$  is optionally substituted phenyl. In some embodiments,  $R_2$  is optionally substituted  $C_1-C_4$  alkyl. In some embodiments  $R_1$  and  $R_2$ , together with the atoms to which they are attached, form  $=O$ . In some embodiments  $R_1$  and  $R_3$ , together with the atoms to which they are attached, form a carbocyclic ring

**[0163]** In some embodiments,  $R_3$  is optionally substituted phenyl. In some embodiments,  $R_3$  is optionally substituted  $C_1-C_4$  alkyl. In some embodiments,  $R_4$  is optionally substituted phenyl. In some embodiments,  $R_4$  is optionally substituted  $C_1-C_4$  alkyl. In some embodiments,  $R_5$  is optionally substituted phenyl. In some embodiments,  $R_5$  is optionally substituted  $C_1-C_4$  alkyl. In some embodiments,  $R_6$  is optionally substituted phenyl. In some embodiments,  $R_6$  is optionally substituted  $C_1-C_4$  alkyl. In some embodiments  $R_3$  and  $R_5$ , together with the atoms to which they are attached, form a heteroaryl ring.

**[0164]** In some embodiments,  $R_9$  is optionally substituted  $C_1-C_{12}$  alkyl. In some embodiments,  $R_9$  is t-butyl. In some embodiments,  $R_9$  is optionally substituted  $C_3-C_8$


cycloalkyl. In some embodiments,  $R_9$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl. In some embodiments,  $R_9$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl. In some embodiments,  $R_9$  is optionally substituted aryl. In some embodiments,  $R_9$  is optionally substituted heteroaryl.

**[0165]** In some embodiments,  $R_{10}$  is H. In some embodiments,  $R_{10}$  is optionally substituted phenyl. In some embodiments,  $R_{10}$  is optionally substituted  $C_{1-4}$  alkyl.

**[0166]** In some embodiments, each  $C_{1-4}$  alkyl is optionally substituted with one or more substituents independently selected from the group consisting of, OMe, and phenyl. In some embodiments, each  $C_{1-4}$  alkyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen.

**[0167]** In some embodiments, each phenyl is optionally substituted with OMe. In some embodiments, each phenyl is optionally substituted with halogen.

**[0168]** Some specific embodiments of the compound described herein have the following structures:



**[0169]** In some embodiments,  $M^+$  is a metal cation. In some embodiments,  $M^+$  is a non-metal cation. In some embodiments,  $M^+$  and the compound of Formula (I) are in a 1:1 ratio. In some embodiments,  $M^+$  and the compound of Formula (I) are in a ratio between 1:10 and 10:1. Non-limiting examples of  $M^+$  include cations of lithium, sodium, potassium, calcium, ammonia, triethylamine, and aluminum.

**[0170]** Intermediate compounds of Formula (I) have the advantageous property of being crystalline, thus facilitating purification. More particularly, crystalline intermediates of Formula (I) are advantageous because no chromatography is required to remove impurities and excess reagents. For example, metal catalysts and other borane reagents would normally necessitate silica gel chromatography prior to entering the next step in the synthesis of therapeutic boronate beta-lactamase inhibitors. In some embodiments, intermediate

compounds of Formula (I) are filtered, washed with suitable solvent, and carried onto the next synthetic step without further purification. In some embodiments, suitable solvents include dimethyl ether, diethyl ether, diisopropyl ether, MTBE, ACN, THF, DCM, chloroform, ethyl acetate, pentanes, hexanes, heptanes, petroleum ether, benzene, toluene, trifluorotoluene, acetone, 2-Me-THF, CPME, 2-butanone, 1,2-dichloroethane, dioxane, pyridine, o-xylene, m-xylene, p-xylene, or any combination thereof.

#### Methods of Preparation

**[0171]** The compounds disclosed herein may be synthesized by methods described below, or by modification of these methods. Ways of modifying the methodology include, among others, temperature, solvent, reagents etc., known to those skilled in the art. In general, during any of the processes for preparation of the compounds disclosed herein, it may be necessary and/or desirable to protect sensitive or reactive groups on any of the molecules concerned. This may be achieved by means of conventional protecting groups, such as those described in *Protective Groups in Organic Chemistry* (ed. J.F.W. McOmie, Plenum Press, 1973); and P.G.M. Green, T.W. Wutts, *Protecting Groups in Organic Synthesis* (3rd ed.) Wiley, New York (1999), which are both hereby incorporated herein by reference in their entirety. The protecting groups may be removed at a convenient subsequent stage using methods known from the art. Synthetic chemistry transformations useful in synthesizing applicable compounds are known in the art and include e.g. those described in R. Larock, *Comprehensive Organic Transformations*, VCH Publishers, **1989**, or L. Paquette, ed., *Encyclopedia of Reagents for Organic Synthesis*, John Wiley and Sons, **1995**, which are both hereby incorporated herein by reference in their entirety. The routes shown and described herein are illustrative only and are not intended, nor are they to be construed, to limit the scope of the claims in any manner whatsoever. Those skilled in the art will be able to recognize modifications of the disclosed syntheses and to devise alternate routes based on the disclosures herein; all such modifications and alternate routes are within the scope of the claims. Unless otherwise indicated, substituent variables in the following schemes have the same definitions as elsewhere in this application.

**[0172]** An exemplary but non-limiting general synthetic scheme for preparing the intermediate compound of Formula (I) is shown in Scheme 1, below. Unless otherwise

indicated, the variable definitions are as above for Formula (I). This process starts with  $\beta$ -hydroxy ester (A), where  $R_9$  is selected from the group consisting of optionally substituted  $C_1$ - $C_{12}$  alkyl, optionally substituted  $C_3$ - $C_8$  cycloalkyl, optionally substituted  $C_2$ - $C_{12}$  alkenyl, optionally substituted  $C_2$ - $C_{12}$  alkynyl, optionally substituted aryl and optionally substituted heteroaryl.

**[0173]** Compound (A) is treated with a base and PG-LG, where PG is a hydroxyl protecting group and LG is a leaving group, to append a protecting group to the alcohol moiety of (A), providing Compound (B). In some embodiments, suitable bases include, but are not limited to, sodium hydride, potassium hydride, triethylamine, diisopropylethylamine, triisopropylamine, piperidine, imidazole, t-butyl magnesium chloride, DBU, DABCO, and potassium t-butoxide. In some embodiments, PG is a trialkylsilyl group, such as a trimethylsilyl or t-butyldimethylsilyl. In some embodiments, PG is an acetal, such as O-methoxymethyl ether or O-tetrahydropyran. In some embodiments, PG is an ether, such as benzyl ether, p-methoxybenzyl ether, or methyl ether. In some embodiments, suitable leaving groups include Cl, Br, I, methanesulfonyl, p-toluenesulfonyl, trifluoromethanesulfonyl, p-fluorophenyl, pentafluorophenyl, or p-nitrosulfonyl.

**[0174]** Compound (B) is subsequently treated with borane (C) and a catalyst, to form boronate (D). In some embodiments, X is selected from the group consisting of substituted or unsubstituted variants of the following:  $C_1$ - $C_4$  alkyl,  $C_2$ - $C_4$  alkenyl, and  $C_2$ - $C_4$  alkynyl. In some embodiments,  $R_{11}$  is selected from the group consisting of substituted or unsubstituted variants of the following:  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl,  $C_2$ - $C_{12}$  alkynyl,  $C_3$ - $C_{12}$  cycloalkyl,  $C_3$ - $C_{12}$  cycloalkenyl,  $C_3$ - $C_{12}$  cycloalkynyl,  $C_3$ - $C_{12}$  heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl. In some embodiments,  $R_{12}$  is selected from the group consisting of substituted or unsubstituted variants of the following:  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl,  $C_2$ - $C_{12}$  alkynyl,  $C_3$ - $C_{12}$  cycloalkyl,  $C_3$ - $C_{12}$  cycloalkenyl,  $C_3$ - $C_{12}$  cycloalkynyl,  $C_3$ - $C_{12}$  heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl. In some embodiments, suitable catalysts include ligand-bound transition metals such as rhodium, iridium, ruthenium, cobalt, nickel, palladium, titanium, and zirconium. In some embodiments, suitable ligands include triphenylphosphine, carbon monoxide, chloride, BINAP, CPhos, Xantphos, 1,5-cyclooctadiene, 1,2-Bis(diphenylphosphino)ethane, 1,2-

Bis(diphenylphosphino)propane, 1,2-Bis(diphenylphosphino)butane, 1,1'-Bis(diphenylphosphino)ferrocene, and any combination thereof.

**[0175]** Compound (D) is subsequently exposed to deprotection conditions to remove the protecting group, followed by oxidation, affording cyclic boronic acid (E). In some embodiments, suitable deprotection conditions include treatment with acid, such as HF, HCl, HBr, p-toluenesulfonic acid, methanesulfonic acid, acetic acid, or trifluoroacetic acid. In some embodiments, suitable deprotection conditions with acid include aqueous acid. In some embodiments, suitable deprotection conditions include treatment with base, such as NaOH, KOH, sodium hydride, potassium hydride, t-butyl magnesium chloride, LDA, and potassium t-butoxide. In some embodiments, suitable deprotection conditions include treatment with a reducing agent, such as H<sub>2</sub>, Ni°, Pd°, Na/NH<sub>3</sub>. In some embodiments, suitable deprotection conditions include treatment with an oxidizing agent, such NaIO<sub>4</sub>. In some embodiments, suitable deprotection conditions include hydrogenation, for example with Pd/C and H<sub>2</sub> or Raney nickel and H<sub>2</sub>. In some embodiments, suitable deprotection conditions include hydride abstraction, for example, with DDQ or Ph<sub>3</sub>CBF<sub>4</sub>. In some embodiments, suitable deprotection conditions include treatment with a Lewis acid, such as BBr<sub>3</sub> or Me<sub>3</sub>SiI. In some embodiments, suitable deprotection conditions include treatment with a nucleophile, such as EtSNa or LiI. In some embodiments, suitable deprotection conditions include combinations of the above listed embodiments (e.g aqueous acid followed or together with NaIO<sub>4</sub> oxidation).

**[0176]** Compound (E) is then treated with complexing group (F) to form a compound of Formula (I). In some embodiments, Compound (E) and complexing group (F) are dissolved in a suitable solvent or combination of solvents. In some embodiments, a suitable solvent is, for example, dimethyl ether, diethyl ether, diisopropyl ether, MTBE, ACN, THF, DCM, chloroform, ethyl acetate, pentanes, hexanes, heptanes, petroleum ether, benzene, toluene, trifluorotoluene, acetone, 2-Me-THF, CPME, 2-butanone, 1,2-dichloroethane, dioxane, pyridine, , o-xylene, m-xylene, p-xylene, or any combination thereof.


**[0177]** In some embodiments, the solvent is an ethereal solvent such as dimethyl ether, diethyl ether, diisopropyl ether, MTBE or dioxane. In some embodiments, the solvent is a mixture of an ethereal solvent such as dimethyl ether, diethyl ether, diisopropyl ether,



MTBE or dioxane and another solvent selected from heptane, hexanes, ethyl acetate, toluene or ACN. In some embodiments the ethereal solvent is MTBE. In some embodiments, the ethereal solvent is dioxane. In some embodiments, solvent is a mixture of MTBE and ACN. In some embodiments, the solvent is a mixture of MTBE, ACN and heptane. In some embodiments, the solvent is a mixture of MTBE and dioxane. In some embodiments, the ratio of MTBE to ACN is between 5 to 1 and 10 to 1. In some embodiments the ratio of MTBE to ACN is between 6 to 1 and 7 to 1.

**[0178]** In some embodiments,  $n$  is 0 or 1. In some embodiments,  $Y_1$  is O. In some embodiments,  $Y_1$  is  $N^+R_5R_6$ . In some embodiments,  $Y_2$  is O. In some embodiments,  $Y_2$  is  $NR_{10}$ . In some embodiments,  $R_1, R_2, R_3, R_4, R_5, R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl. In some embodiments, Compound (F) is ethanolamine. In some embodiments, Compound (F) is glycine. In some embodiments, Compound (F) is oxalic acid. In some embodiments, Compound (F) is glycolic acid.

#### Scheme 1



**[0179]** The intermediate compound of Formula (I) can be crystallized and isolated. One skilled in the art will recognize that specific crystallization conditions will

depend, in part upon the identities of the particular substituents in each molecule, for example R<sub>1</sub>, R<sub>2</sub>, R<sub>3</sub>, R<sub>4</sub>, R<sub>5</sub>, R<sub>6</sub>, and/or R<sub>9</sub>. Crystallization occurs in a suitable solvent or solvent system at a particular temperature or range of temperatures. In some embodiments, suitable solvents include dimethyl ether, diethyl ether, diisopropyl ether, MTBE, ACN, THF, DCM, chloroform, ethyl acetate, pentanes, hexanes, heptanes, petroleum ether, benzene, toluene, trifluorotoluene, acetone, 2-Me-THF, CPME, 2-butanone, 1,2-dichloroethane, dioxane, pyridine, , o-xylene, m-xylene, p-xylene, or any combination thereof. In some embodiments, suitable temperatures include no higher than -25°C, no higher than -20°C, no higher than -15°C, no higher than -10°C, no higher than -5°C, no higher than 0°C, no higher than 5°C, no higher than 10°C, no higher than 15°C, no higher than 20°C, no higher than 25°C, no higher than 30°C.

**[0180]** In some embodiments, the compound of Formula (I) can be crystallized in a mixture of acetonitrile and MTBE. In some embodiments this mixtures is at least 5% MTBE, at least 10% MTBE, at least 15% MTBE, at least 20% MTBE, at least 25% MTBE, at least 30% MTBE, at least 35% MTBE, at least 40% MTBE, at least 45% MTBE, at least 50% MTBE, at least 55% MTBE, at least 60% MTBE, at least 65% MTBE, at least 70% MTBE, at least 75% MTBE, at least 80% MTBE, at least 85% MTBE, at least 90% MTBE, or at least 95% MTBE. In some embodiments, the compound of Formula (I) heptane is added to the mixture of MTBE and ACN. In some embodiments, these mixtures of MTBE and ACN contain at least 5% heptane, 10% heptane, at least 15% heptane, at least 20% heptane, at least 25% heptane, at least 30% heptane, at least 35% heptane, at least 40% heptane, at least 45% heptane, at least 50% heptane, at least 55% heptane, at least 60% heptane, at least 65% heptane, at least 70% heptane, at least 75% heptane, at least 80% heptane, at least 85% heptane, at least 90% heptane, or at least 95% heptane. The intermediate compound of Formula (I) can be used to synthesize therapeutic boronate beta-lactamase inhibitors. For example, an exemplary but non-limiting general synthetic scheme for preparing therapeutic boronate compounds from compounds of Formula (I) is shown in Scheme 2, below.

**[0181]** Unless otherwise indicated, the variable definitions are as above for Formula (I). The process starts with decomplexation and protection of the boronate moiety.

In some embodiments, decomplexation and protection can be accomplished concomitantly or in separate individual steps.

**[0182]** In some embodiments, the decomplexation and protection of the boronate is accomplished using a concomitant process which entails mixing the compound of Formula (I) and the boronate protecting group in a biphasic solvent system, one aqueous/hydrophilic solvent and an organic solvent. Non-limiting examples include water, with or without acid or base, and MTBE,  $\text{CH}_2\text{Cl}_2$ , ethyl acetate, 2-Me-THF, CPME, and diisopropyl ether. This reaction provides protected boronate (I-B).

**[0183]** In some embodiments, the complexing moiety (F) is then removed by phase partitioning, and the protected boronate is isolated as a solution. In some embodiments, the decomplexation and protection of the boronate is accomplished using a step-wise process. In some embodiments, compound of Formula (I) is dissolved in water with or without an acid and the decomplexed boronate is extracted with an organic solvent not miscible with water, such as MTBE,  $\text{CH}_2\text{Cl}_2$ , ethyl acetate, 2-Me-THF, CPME, diisopropyl ether. In some embodiments, the protecting-boronate reagent is then added to the organic solution of the decomplexed boronate and the protected boronate (I-B) is then isolated as a solution.

**[0184]** In some embodiments the acid is HCl or  $\text{H}_2\text{SO}_4$ . In some embodiments, the organic solvent is MTBE or  $\text{CH}_2\text{Cl}_2$ . In some embodiments, the organic solvent is MTBE. In some embodiments, the organic solvent is  $\text{CH}_2\text{Cl}_2$ . In some embodiments, the boronate protecting group is a compound of Formula G. In some embodiments, n is 0 or 1. In some embodiments,  $\text{R}_1$ ,  $\text{R}_2$ ,  $\text{R}_3$  and  $\text{R}_4$  are independently selected from the group consisting of H, optionally substituted phenyl, optionally substituted carbocyclyl and optionally substituted  $\text{C}_{1-4}$  alkyl, or  $\text{R}_1$  and  $\text{R}_3$  together with the atoms to which they are attached form a carbocyclic ring.

**[0185]** In some embodiments, n is 0. In some embodiments, n is 1.

**[0186]** In some embodiments,  $\text{R}_1$ , and  $\text{R}_3$ , are phenyl and  $\text{R}_2$ , and  $\text{R}_4$  are hydrogen. In some embodiments,  $\text{R}_1$ , and  $\text{R}_3$ , are isopropyl and  $\text{R}_2$ , and  $\text{R}_4$  are hydrogen. In some embodiments,  $\text{R}_1$ , and  $\text{R}_3$ , are cyclohexyl and  $\text{R}_2$ , and  $\text{R}_4$  are hydrogen.

**[0187]** In some embodiments, the boronate-protecting group is pinanediol, pinacol, 1,2-dicycloethanediol, 1,2-diisopropylethanediol, 1,2-diphenylethanediol. In some embodiments, the boronate-protecting group is pinanediol. In some embodiments, the

boronate-protecting group is (+)-pinanediol. In some embodiments, the boronate-protecting group is (-)-pinanediol.


**[0188]** In some embodiments, the compound of Formula I-B is then treated with a base and PG-LG, where PG is a hydroxyl protecting group and LG is a leaving group, to append a protecting group to the alcohol moiety of (I-B), providing Compound (I-C). In some embodiments, suitable bases include, but are not limited to, sodium hydride, potassium hydride, triethylamine, diisopropylethylamine, triisopropylamine, piperidine, imidazole, t-butyl magnesium chloride, DBU, DABCO, and potassium t-butoxide. In some embodiments, PG is a trialkylsilyl group, such as a trimethylsilyl or t-butyl dimethylsilyl. In some embodiments, PG is an acetal, such as O-methoxymethyl ether or O-tetrahydropyran. In some embodiments, PG is an ether, such as benzyl ether, p-methoxybenzyl ether, or methyl ether. In some embodiments, suitable leaving groups include Cl, Br, I, methanesulfonyl, p-toluenesulfonyl, trifluoromethanesulfonyl, p-fluorophenyl, pentafluorophenyl, or p-nitrosulfonyl.

**[0189]** Halomethylation of compound of Formula (I-C), at the carbon adjacent to the boron atom affords Compound I-D. In some embodiments, halomethylation is accomplished by treatment with n-butyllithium and dichloromethane. In some embodiments, halomethylation is accomplished with dibromomethane and lithium diisopropylamide or lithium hexamethyldisilazide. In some embodiments, subsequent treatment of Compound I-D with an amine nucleophile displaces the halogen to form an amino group at the alpha position. Exemplary nucleophiles include, but are not limited to lithium hexamethyldisilazide and ammonia. Subsequent reaction with a carboxylic acid containing moiety under amide coupling conditions provides Compound I-E. In some embodiments, amide bond coupling conditions include treatment with EDCI, HOBt, HBTU, HATU, PyBOP, PyBrop, or DCC. In some embodiments,  $R_{13}$  is  $C_{1-9}$  alkyl. In some embodiments,  $R_{13}$  is  $C_{1-9}$ alkyl- $R_{14}$ . In some embodiments,  $R_{14}$  is substituted or unsubstituted aryl. In some embodiments,  $R_{14}$  is substituted or unsubstituted heteroaryl. In some embodiments,  $R_{14}$  is substituted or unsubstituted carbocyclyl. In some embodiments,  $R_{14}$  is substituted or unsubstituted heterocyclyl. In some embodiments,  $R_{14}$  is thiophene-2-yl.

**[0190]** De-esterification and de-complexation of Compound I-E provides therapeutic boronate I-F. In some embodiments, de-esterification and de-complexation are

accomplished in one step. In some embodiments, de-esterification and de-complexation are accomplished in more than one step. In some embodiments, de-esterification and de-complexation are accomplished by treatment with aqueous acid, dioxane, and a boronic acid acid. In some embodiments, the acid is hydrochloric or sulfuric acid. In some embodiments, the acid is hydrochloric acid. In some embodiments, the acid is sulfuric acid. In some embodiments, the boronic acid is isobutyl boronic acid, phenylboronic acid or boric acid. In some embodiments, the boronic acid is isobutylboronic acid. In some embodiments, the boronic acid is phenylboronic acid. In some embodiments, the boronic acid is boric acid. In some embodiments, the de-esterification and de-complexation are accomplished by treatment with sulfuric acid, dioxane and boric acid. In some embodiments, the de-esterification and de-complexation are accomplished by treatment with hydrochloric acid, dioxane and isobutylboronic acid.


Scheme 2



## EXAMPLES

Example 1 - Synthesis of Intermediate Compound 10

[0191] The compound of Formula 10 was synthesized as shown in Scheme 3, below:

Scheme 3(R)-t-butyl 3-(trimethylsilyloxy)-pent-4-enoate (7)

[0192] Chlorotrimethylsilane (4.6 mL, 36.3 mmol, 1.25 eq) was added to a solution of (R)-t-butyl 3-hydroxy-pent-4-enoate (**1**, 5 g, 29 mmol) and triethylamine (5.3 mL, 37.3 mmol, 1.3 eq) in dichloromethane (25 mL) keeping the temperature below 30 °C. After completion of the addition, the white heterogeneous mixture was stirred at rt for 20 minutes (TLC, GC, note 2) then quenched with MeOH (352 µL, 0.3 eq). After stirring at rt for 5 minutes, the white heterogeneous reaction mixture was diluted with heptane (25 mL). The salts were filtered off and rinsed with heptane (2 x 10 mL). The combined turbid filtrates were washed with a saturated solution of NaHCO<sub>3</sub> (2 x 25 mL) and concentrated to dryness. The residual oil was azeotroped with heptane (25 mL) to give a colorless oil that was used immediately.

(S)-t-butyl 3-(trimethylsilyloxy)-5-(4,4,5,5-tetramethyl-[1,3,2]dioxaborolan-2-yl)-pentanoate (8)

[0193] A solution of bis-diphenylphosphino-ethane (46.3 mg, 0.2 mol%) and [Ir(COD)Cl]<sub>2</sub> (39 mg, 0.1 mol%) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added to a refluxing solution of crude TMS-protected pentenoate **7**. Pinacol borane (9.3 mL, 1.1 eq) was added to the

refluxing solution. After stirring at reflux for 3 h, the reaction mixture was cooled to room temperature, concentrated to dryness and taken up in heptane (50 mL). The insolubles were filtered over Celite and rinse with heptane (10 mL).


Ethanolamine-boronic acid salt (10)

**[0194]** A mixture of fully protected boronate **8** (5.0 g, 13.4 mmol), 0.5 N HCl (5 mL) and acetone (0.5 mL) was stirred vigorously at room temperature, providing intermediate **9**. After complete consumption of the starting material, a solution of NaIO<sub>4</sub> (3.44 g, 1.2 eq) in water (15 mL) was added slowly keeping the temperature <30 °C. Upon the completion of the addition (30 min), the reaction mixture was allowed to cool to room temperature. After consumption of all pinacol, MTBE (5 mL) was added. After stirring at room temperature for 10 min, the white solids were filtered off and rinsed with MTBE (2 x 5 mL). The filtrate was partitioned and the aqueous layer was extracted with MTBE (10 mL). The combined organic extracts were washed sequentially with a 0.1 M NaHSO<sub>3</sub> solution (2 x 5 mL), a saturated NaHCO<sub>3</sub> solution (5 mL) and brine (5 mL). The organic layer was concentrated to dryness. The residue was taken up in MTBE (15 mL) and the residual salts filtered off. The filtrate was concentrated to dryness and the residue was taken up in MTBE (10 mL) and acetonitrile (1.7 mL). Ethanolamine (0.99 mL, 1.1 eq) was added. After stirring at room temperature for 1 hour, the heterogeneous mixture was stirred at 0 °C. After stirring at 0 °C for 2 hours, the solids were collected by filtration, rinsed with MTBE (2 x 5 mL), air dried then dried under high vacuum to give Compound **10** as a white granular powder.

Example 2 – Preparation of Beta-Lactamase Inhibitor (15)

**[0195]** The compound of Formula 15 was synthesized as shown in Scheme 4 below:

Scheme 4



#### Synthesis of pinanediol boronate (**12**)

**[0196]** Ethanolammonium boronate **11** (15 g, 61.7 mmol) and pinanediol (10.5 g, 61.7 mmol, 1 eq) were suspended in MTBE (75 mL). Water (75 mL) was added and the yellow biphasic heterogeneous mixture was stirred at room temperature. After stirring for 2 hours at room temperature, some pinanediol was still present and stirring was continued overnight. The layers were separated and the organic layer was washed with brine, concentrated under reduced pressure and azeotroped with MTBE (2 x 30 mL). The residual oil was taken up in dichloromethane (40 mL). In another flask, TBSCl (11.6 g, 77.1 mmol, 1.25 eq) was added to a solution of imidazole (9.66 g, 141.9 mmol, 2.3 eq) in dichloromethane (25 mL). The white slurry was stirred at room temperature. After 5 minutes, the solution of pinanediol boronate was added to the white slurry and the flask was rinsed with dichloromethane (2 x 5 mL). The heterogeneous reaction mixture was heated at reflux temperature. After stirring at reflux for 8 hours, the reaction mixture was cooled to 30 °C and TMSCl (330 µL) was added. After stirring 30 minutes at 30 °C, MeOH (15 mL) was added. After stirring at room temperature overnight, the reaction mixture was washed sequentially



with 0.5 N HCl (115 mL), 0.5 N HCl (60 mL) and saturated NaHCO<sub>3</sub> (90 mL). The organic layer was concentrated under reduced pressure and azeotroped with heptane (150 mL) to give **12** as a yellow oil (27.09 g, 94.1%) which was used without purification.

#### Synthesis of chloroboronate (**13**)

[0197] A solution of n-butyllithium (2.5 M in hexane, 29.6 mL, 74.1 mmol, 1.3 eq) was added to THF (100 mL) at -80 °C. The resulting solution was cooled to -100 °C. A solution of dichloromethane (14.6 mL, 228 mmol, 4 eq) in THF (25 mL) was added via syringe pump on the sides of the flask keeping the temperature < -95 °C. During the second half of the addition a precipitate starts to appear which became thicker with the addition of the remaining dichloromethane solution. After stirring between -100 and -95 °C for 30 min, a solution of **12** (26.59 g, 57 mmol) in THF (25 mL) was added by syringe pump on the sides of the flask while maintaining the batch temperature < -95 °C to give a clear yellow solution. After stirring between -100 and -95 °C for 30 min, a solution of zinc chloride (1 M in ether, 120 mL, 120 mmol, 2.1 eq) was added keeping the temperature < -70 °C. The reaction mixture was then warmed to room temperature (at about -18 °C the reaction mixture became turbid/heterogeneous). After stirring at room temperature for 2 hours, the reaction mixture was cooled to 15 °C and quenched with 1 N HCl (100 mL). The layers were separated and the organic layer was washed sequentially with 1 N HCl (100 mL) and water (2 x 100 mL), concentrated to oil and azeotroped with heptane (3 x 150 mL) to provide **13** as a yellow oil (30.03 g, 102%) which was used without purification.

#### Synthesis of (**14**)

[0198] LiHMDS (1 M in THF, 63 mL, 62.7 mmol, 1.1 eq) was added to a solution of **13** (29.5 g, 57 mmol) in THF (60 mL) while maintaining the batch temperature at < -65 °C. After stirring at -78 °C for 2 hours, additional LiHMDS (5.7 mL, 0.1 eq) was added to consume the remaining starting material. After stirring at -78 °C for 30 minutes, the tan reaction mixture was warmed to room temperature. After stirring at room temperature for one hour, the solution of silylated amine was added via cannula to a solution of HOBt ester of 2-thienylacetic acid in acetonitrile at 0 °C (the solution of HOBt ester was prepared by adding EDCI (16.39 g, 85.5 mmol, 1.5 eq) to a suspension of recrystallized 2-thienylacetic acid

(9.73 g, 68.4 mmol, 1.2 eq) and HOBT.H<sub>2</sub>O (11.35 g, 74.1 mmol, 1.3 eq) in acetonitrile (10 mL) at 0 °C. The clear solution was stirred at 0 °C for 30 minutes prior to the addition of the silylated amine). After stirring at 0 °C for one hour, the heterogeneous reaction mixture was placed in the fridge overnight. Saturated aqueous sodium bicarbonate (80 mL) and heptane (80 mL) were added, and after stirring 30 minutes at room temperature, the layers were separated. The organic layer was washed with saturated aqueous sodium bicarbonate (2 x 80 mL) and filtered through Celite. The filtrate was concentrated under reduced pressure and the tan oil was azeotroped with heptane (3 x 110 mL). The residue was taken up in heptane (60 mL) and seeds were added. After stirring at room temperature for one hour, the reaction mixture became heterogeneous. After stirring 4 hours at 0 °C, the solids were collected by filtration and washed with ice cold heptane (3 x 20 mL), air dried then dried under high vacuum to give **14** as an off white powder (10.95 g, 31%).

#### Synthesis of (15)

[0199] A mixture of **14** (10 g, 16.1 mmol), boric acid (1.3 g, 20.9 mmol, 1.3 eq), dioxane (20 mL), and 1 M sulfuric acid (10 mL) was heated at 75 °C. After stirring 7 hours at 75 °C, the cooled reaction mixture was diluted with water (10 mL) and MTBE (30 mL) and the residual mixture was cooled to 0 °C. The pH was adjusted to 5.0 with a solution of 2 N NaOH (14 mL). The layers were separated and the aqueous layer was extracted with MTBE (2 x 30 mL) then concentrated to dryness. The residue was taken up in water (10 mL) and the solution was filtered through a 0.45 µm GMF syringe filter. The flask and filter were rinsed with water (7.5 mL). The pH of the filtrate was lowered to 4.0 with 2 M H<sub>2</sub>SO<sub>4</sub> and seeds (5 mg) were added. After stirring at room temperature for 10 minutes, the pH was lowered to 1.9 over 1 hour with 2 M H<sub>2</sub>SO<sub>4</sub> (total volume 3.5 mL). After stirring at room temperature for 2 hours, the solids were collected by filtration. The filtrate was recirculated twice to rinse the flask and the cake was washed with water (2 x 12 mL), air dried then dried under high vacuum to give **15** as a white powder (3.63 g, 76%).

#### Conclusion

[0200] While the invention has been described with reference to the specific embodiments thereof, it should be understood by those skilled in the art that various changes

may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. This includes embodiments which do not provide all of the benefits and features set forth herein. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the present invention. All such modifications are intended to be within the scope of the claims appended hereto. Accordingly, the scope of the invention is defined only by reference to the appended claims.



9. The compound of any one of Claims 1 to 7, wherein R<sub>1</sub> is optionally substituted phenyl.

10. The compound of any one of Claims 1 to 7, wherein R<sub>1</sub> is optionally substituted C<sub>1-4</sub> alkyl.

11. The compound of any one of Claims 1 to 10, wherein R<sub>2</sub> is H.

12. The compound of any one of Claims 1 to 10, wherein R<sub>2</sub> is optionally substituted phenyl.

13. The compound of any one of Claims 1 to 10, wherein R<sub>2</sub> is optionally substituted C<sub>1-4</sub> alkyl.

14. The compound of any one of Claims 1 to 7, wherein R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O.

15. The compound of any one of Claims 1 to 14, wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

16. The compound of any one of Claims 1 to 7, wherein R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O and wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

17. The compound of any one of Claims 1 to 14, wherein R<sub>3</sub> is H.

18. The compound of any one of Claims 1 to 14, wherein R<sub>3</sub> is optionally substituted phenyl.

19. The compound of any one of Claims 1 to 14, wherein R<sub>3</sub> is C<sub>1-4</sub> optionally substituted alkyl.

20. The compound of any one of Claims 1 to 7 and 11 to 13, wherein R<sub>1</sub> and R<sub>3</sub> together with the atoms to which they are attached form carbocyclic ring.


21. The compound of any one of Claims 1 to 14, wherein R<sub>3</sub> and R<sub>5</sub>, together with the atoms to which they are attached, form a heteroaryl ring.
22. The compound of any one of Claims 1 to 21, wherein R<sub>4</sub> is H.
23. The compound of any one of Claims 1 to 21, wherein R<sub>4</sub> is optionally substituted phenyl.
24. The compound of any one of Claims 1 to 21, wherein R<sub>4</sub> is optionally substituted C<sub>1-4</sub> alkyl.
25. The compound of any one of Claims 1 to 20 and 22 to 24, wherein R<sub>5</sub> is H.
26. The compound of any one of Claims 1 to 20 and 22 to 24, wherein R<sub>5</sub> is optionally substituted phenyl.
27. The compound of any one of Claims 1 to 20 and 22 to 24, wherein R<sub>5</sub> is optionally substituted C<sub>1-4</sub> alkyl.
28. The compound of any one of Claims 1 to 27, wherein R<sub>6</sub> is H.
29. The compound of any one of Claims 1 to 27, wherein R<sub>6</sub> is optionally substituted phenyl.
30. The compound of any one of Claims 1 to 27, wherein R<sub>6</sub> is optionally substituted C<sub>1-4</sub> alkyl.
31. The compound of any one of Claims 1 to 30, wherein R<sub>10</sub> is H.
32. The compound of any one of Claims 1 to 30, wherein R<sub>10</sub> is optionally substituted phenyl.
33. The compound of any one of Claims 1 to 30, wherein R<sub>10</sub> is optionally substituted C<sub>1-4</sub> alkyl.

34. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl.
35. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is t-butyl.
36. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is optionally substituted C<sub>3</sub>-C<sub>8</sub> cycloalkyl.
37. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl.
38. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl.
39. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is optionally substituted aryl.
40. The compound of any one of Claims 1 to 33, wherein R<sub>9</sub> is optionally substituted heteroaryl.
41. The compound of any one of Claims 1 to 7, 10, 13, 19, 24, 27, 30 and 33, wherein each C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl.
42. The compound of any one of Claims 1 to 7, 10, 13, 19, 24, 27, 30, and 33, wherein each C<sub>1-4</sub> alkyl is optionally substituted with OMe.
43. The compound of any one of Claims 1 to 7, 10, 13, 19, 24, 27, 30 and 33, wherein each C<sub>1-4</sub> alkyl is optionally substituted with phenyl.
44. The compound of any one of Claims 1 to 7, 9, 12, 18, 23, 26, 29 and 32, wherein each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen.

45. The compound of any one of Claims 1 to 7, 9, 12, 18, 23, 26, 29 and 32, wherein each phenyl is optionally substituted with OMe.

46. The compound of any one of Claims 1 to 7, 9, 12, 18, 23, 26, 29 and 32, wherein each phenyl is optionally substituted with halogen.

47. The compound of Claim 1, having a structure selected from the group consisting of:




wherein:


$M^+$  is a cation selected from the group consisting of lithium, sodium, potassium, calcium, ammonium, triethylammonium, and aluminum.

48. A method of making a compound of Formula (I), or a salt thereof, comprising the steps of:

(a) protecting the primary hydroxy group of the compound of Formula (A):




to form a compound of Formula (B):




(b) reacting a compound of Formula (B) with a compound of Formula (C):







to form a compound of Formula (D):




(c) deprotecting a compound of Formula (D) to form a compound of Formula (E):



(d) reacting a compound of Formula (E) with a complexing agent of Formula (F):



to form a compound of Formula (I):



wherein:

PG is a hydroxyl protecting group;

n is 0 or 1;

$Y_1$  is O or  $N^+R_5R_6$

$Y_2$  is O or  $NR_{10}$ ;

$R_1$  and  $R_2$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl, or  $R_1$  and  $R_2$  together with the atoms to which they are attached, form =O;

$R_3$  is selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl, or  $R_1$  and  $R_3$  together with the atoms to which they are attached form an aryl or heteroaryl ring; or  $R_3$  and  $R_4$  together with the atoms to which they are attached, form =O;

$R_4$ ,  $R_5$ ,  $R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl;

$R_9$  is optionally substituted  $C_1$ - $C_{12}$  alkyl, optionally substituted  $C_3$ - $C_8$  cycloalkyl, optionally substituted  $C_2$ - $C_{12}$  alkenyl, optionally substituted  $C_2$ - $C_{12}$  alkynyl, optionally substituted aryl or optionally substituted heteroaryl;

$R_{11}$  is selected from the group consisting of substituted or unsubstituted variants of the following:  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl,  $C_2$ - $C_{12}$  alkynyl,  $C_3$ - $C_{12}$  cycloalkyl,  $C_3$ - $C_{12}$  cycloalkenyl,  $C_3$ - $C_{12}$  cycloalkynyl,  $C_3$ - $C_{12}$  heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl, and

X is selected from the group consisting of substituted or unsubstituted variants of the following:  $C_1$ - $C_4$  alkyl,  $C_2$ - $C_4$  alkenyl, and  $C_2$ - $C_4$  alkynyl; and

$R_{12}$  is selected from the group consisting of substituted or unsubstituted variants of the following:  $C_1$ - $C_{12}$  alkyl,  $C_2$ - $C_{12}$  alkenyl,  $C_2$ - $C_{12}$  alkynyl,  $C_3$ - $C_{12}$  cycloalkyl,  $C_3$ - $C_{12}$  cycloalkenyl,  $C_3$ - $C_{12}$  cycloalkynyl,  $C_3$ - $C_{12}$  heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl.

49. The method of Claim 48, wherein PG is a trialkylsilyl group.

50. The method of any one of Claims 48 to 49, wherein PG is trimethylsilyl.

51. The method of any one of Claims 48 to 50, wherein n is 0.

52. The method of any one of Claims 48 to 50, wherein n is 1.

53. The method of any one of Claims 46 to 52, wherein  $Y_1$  is O.
54. The method of any one of Claims 46 to 52, wherein  $Y_1$  is  $N^+R_5R_6$ .
55. The method of any one of Claims 48 to 54, wherein  $Y_2$  is O.
56. The method of any one of Claims 48 to 54, wherein  $Y_2$  is  $NR_{10}$ .
57. The method of any one of Claims 48 to 56, wherein  $R_1$  is H.
58. The method of any one of Claims 48 to 56, wherein  $R_1$  is optionally substituted phenyl.
59. The method of any one of Claims 48 to 56, wherein  $R_1$  is optionally substituted  $C_{1-4}$  alkyl.
60. The method of any one of Claims 48 to 59, wherein  $R_2$  is H.
61. The method of any one of Claims 48 to 59, wherein  $R_2$  is optionally substituted phenyl.
62. The method of any one of Claims 48 to 59, wherein  $R_2$  is optionally substituted  $C_{1-4}$  alkyl.
63. The method of any one of Claims 48 to 56, wherein  $R_1$  and  $R_2$  together with the atoms to which they are attached, form  $=O$ .
64. The method of any one of Claims 48 to 63, wherein  $R_3$  and  $R_4$  together with the atoms to which they are attached, form  $=O$ .
65. The method of any one of Claims 48 to 56, wherein  $R_1$  and  $R_2$  together with the atoms to which they are attached, form  $=O$  and wherein  $R_3$  and  $R_4$  together with the atoms to which they are attached, form  $=O$ .
66. The method of any one of Claims 48 to 65, wherein  $R_3$  is H.

67. The method of any one of Claims 48 to 63, wherein  $R_3$  is optionally substituted phenyl.

68. The method of any one of Claims 48 to 63, wherein  $R_3$  is  $C_{1-4}$  optionally substituted alkyl.

69. The method of any one of Claims 48 to 56, wherein  $R_1$  and  $R_3$  together with the atoms to which they are attached form a carbocyclic ring.

70. The method of any one of Claims 48 to 65, wherein,  $R_3$  and  $R_5$ , together with the atoms to which they are attached, form a heteroaryl ring.

71. The method of any one of Claims 48 to 70, wherein  $R_4$  is H.

72. The method of any one of Claims 48 to 70, wherein  $R_4$  is optionally substituted phenyl.

73. The method of any one of Claims 48 to 69, wherein  $R_4$  is optionally substituted  $C_{1-4}$  alkyl.

74. The method of any one of Claims 48 to 69 and 71-73, wherein  $R_5$  is H.

75. The method of any one of Claims 48 to 69 and 71-73, wherein  $R_5$  is optionally substituted phenyl.

76. The method of any one of Claims 48 to 69 and 71-73, wherein  $R_5$  is optionally substituted  $C_{1-4}$  alkyl.

77. The method of any one of Claims 48 to 76, wherein  $R_6$  is H.

78. The method of any one of Claims 48 to 76, wherein  $R_6$  is optionally substituted phenyl.

79. The method of any one of Claims 48 to 76, wherein  $R_6$  is optionally substituted  $C_{1-4}$  alkyl.

80. The method of any one of Claims 48 to 79, wherein  $R_{10}$  is H.
81. The method of any one of Claims 48 to 79, wherein  $R_{10}$  is optionally substituted phenyl.
82. The method of any one of Claims 48 to 79, wherein  $R_{10}$  is optionally substituted  $C_{1-4}$  alkyl.
83. The method of any one of Claims 48 to 82, wherein  $R_9$  is optionally substituted  $C_1$ - $C_{12}$  alkyl.
84. The method of any one of Claims 48 to 82, wherein  $R_9$  is t-butyl.
85. The method of any one of Claims 48 to 82, wherein  $R_9$  is optionally substituted  $C_3$ - $C_8$  cycloalkyl.
86. The method of any one of Claims 48 to 82, wherein  $R_9$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl.
87. The method of any one of Claims 48 to 82, wherein  $R_9$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl.
88. The method of any one of Claims 48 to 82, wherein  $R_9$  is optionally substituted aryl.
89. The method of Claim 48, wherein  $R_9$  is optionally substituted heteroaryl.
90. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_1$ - $C_{12}$  alkyl.
91. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl.
92. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl.

93. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkyl.
94. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkenyl.
95. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkynyl.
96. The method of Claim 48, wherein  $R_{11}$  is optionally substituted  $C_3$ - $C_{12}$  heterocyclyl.
97. The method of Claim 48, wherein  $R_{11}$  is optionally substituted aryl.
98. The method of Claim 48, wherein  $R_{11}$  is optionally substituted heteroaryl.
99. The method of Claim 48, wherein  $R_{11}$  is optionally substituted arylalkyl.
100. The method of Claim 48, wherein  $R_{11}$  is optionally substituted heteroarylalkyl.
101. The method of Claim 48, wherein  $R_{11}$  is optionally substituted (cycloalkyl)alkyl.
102. The method of Claim 48, wherein X is optionally substituted  $C_1$ - $C_4$  alkyl.
103. The method of Claim 48, wherein X is optionally substituted  $C_2$ - $C_4$  alkenyl.
104. The method of Claim 48, wherein X is optionally substituted  $C_2$ - $C_4$  alkynyl.
105. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_1$ - $C_{12}$  alkyl.
106. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl.
107. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl.

108. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkyl.

109. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkenyl.

110. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkynyl.

111. The method of Claim 48, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  heterocyclyl.

112. The method of Claim 48, wherein  $R_{12}$  is optionally substituted aryl.

113. The method of Claim 48, wherein  $R_{12}$  is optionally substituted heteroaryl.

114. The method of Claim 48, wherein  $R_{12}$  is optionally substituted arylalkyl.

115. The method of Claim 48, wherein  $R_{12}$  is optionally substituted heteroarylalkyl.

116. The method of Claim 48, wherein  $R_{12}$  is optionally substituted (cycloalkyl)alkyl.

117. The method of any one of Claims 48 to 56, 59, 62, 68, 73, 76, 79, 82, and 102, wherein each  $C_{1-4}$  alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl.

118. The method of any one of Claims 48 to 56, 59, 62, 68, 73, 76, 79, 82, and 102, wherein each  $C_{1-4}$  alkyl is optionally substituted with OMe.


119. The method of any one of Claims 48 to 56, 59, 62, 68, 73, 76, 79, 82, and 102, wherein each  $C_{1-4}$  alkyl is optionally substituted with phenyl.

120. The method of any one of Claims 48 to 56, 58, 61, 67, 72, 75, 78, and 81, wherein each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen.


121. The method of any one of Claims 48 to 56, 58, 61, 67, 72, 75, 78, and 81, wherein each phenyl is optionally substituted with OMe.

122. The method of any one of Claims 48 to 56, 58, 61, 67, 72, 75, 78, and 81, wherein each phenyl is optionally substituted with halogen.


123. A method of making a compound of Formula (D) comprising the steps of: reacting a compound of Formula (B):



with a compound of Formula (C):



to form a compound of Formula (D)



wherein PG is a hydroxyl protecting group, and

wherein R<sub>9</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl, optionally substituted C<sub>3</sub>-C<sub>8</sub> cycloalkyl, optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl, optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl, optionally substituted aryl or optionally substituted heteroaryl;



R<sub>11</sub> is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkenyl, C<sub>3</sub>-C<sub>12</sub> cycloalkynyl, C<sub>3</sub>-C<sub>12</sub> heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl, and

X is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, and C<sub>2</sub>-C<sub>12</sub> alkynyl; and

R<sub>12</sub> is selected from the group consisting of substituted or unsubstituted variants of the following: C<sub>1</sub>-C<sub>12</sub> alkyl, C<sub>2</sub>-C<sub>12</sub> alkenyl, C<sub>2</sub>-C<sub>12</sub> alkynyl, C<sub>3</sub>-C<sub>12</sub> cycloalkyl, C<sub>3</sub>-C<sub>12</sub> cycloalkenyl, C<sub>3</sub>-C<sub>12</sub> cycloalkynyl, C<sub>3</sub>-C<sub>12</sub> heterocyclyl, aryl, heteroaryl, arylalkyl, heteroarylalkyl, and (cycloalkyl)alkyl.

124. The method of Claim 123, wherein PG is a trialkylsilyl group.

125. The method of any one of Claims 123 to 124, wherein PG is trimethylsilyl.

126. The method of any one of Claims 123 to 125, wherein R<sub>9</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl.

127. The method of any one of Claims 123 to 126, wherein R<sub>9</sub> is t-butyl.

128. The method of any one of Claims 123 to 125, wherein R<sub>9</sub> is optionally substituted C<sub>3</sub>-C<sub>8</sub> cycloalkyl.

129. The method of any one of Claims 123 to 125, wherein R<sub>9</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl.

130. The method of any one of Claims 123 to 125, wherein R<sub>9</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl.

131. The method of any one of Claims 123 to 125, wherein R<sub>9</sub> is optionally substituted aryl.

132. The method of any one of Claims 123 to 125, wherein R<sub>9</sub> is optionally substituted heteroaryl.

133. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>1</sub>-C<sub>12</sub> alkyl.

134. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkenyl.

135. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>2</sub>-C<sub>12</sub> alkynyl.

136. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkyl.

137. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkenyl.

138. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> cycloalkynyl.

139. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted C<sub>3</sub>-C<sub>12</sub> heterocyclyl.

140. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted aryl.

141. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted heteroaryl.

142. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted arylalkyl.

143. The method of any one of Claims 123 to 132, wherein R<sub>11</sub> is optionally substituted heteroarylalkyl.

144. The method of any one of Claims 123 to 132, wherein  $R_{11}$  is optionally substituted (cycloalkyl)alkyl.

145. The method of any one of Claims 123 to 144, wherein X is optionally substituted  $C_1$ - $C_4$  alkyl.

146. The method of any one of Claims 123 to 144, wherein X is optionally substituted  $C_2$ - $C_4$  alkenyl.

147. The method of any one of Claims 123 to 144, wherein X is optionally substituted  $C_2$ - $C_4$  alkynyl.

148. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_1$ - $C_{12}$  alkyl.

149. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl.

150. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl.

151. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkyl.

152. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkenyl.

153. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  cycloalkynyl.

154. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted  $C_3$ - $C_{12}$  heterocyclyl.

155. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted aryl.


156. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted heteroaryl.

157. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted arylalkyl.


158. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted heteroarylalkyl.

159. The method of any one of Claims 123 to 147, wherein  $R_{12}$  is optionally substituted (cycloalkyl)alkyl.


160. A method of making a compound of Formula (I), or a salt thereof, comprising the steps of: reacting a compound of Formula (E):



with a complexing agent of Formula (F):



to form a compound of Formula (I):



wherein

n is 0 or 1,

$Y_1$  is O or  $N^+R_5R_6$

$Y_2 = O$  or  $NR_{10}$ ,

$R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$ ,  $R_5$ ,  $R_6$ , and  $R_{10}$  are independently selected from the group consisting of H, optionally substituted phenyl and optionally substituted  $C_{1-4}$  alkyl; or

wherein independently two germinal  $R_1$ ,  $R_2$ ,  $R_3$ ,  $R_4$  together with the atoms to which they are attached, form  $=O$ ; and

wherein  $R_9$  is optionally substituted  $C_1$ - $C_{12}$  alkyl, optionally substituted  $C_3$ - $C_8$  cycloalkyl, optionally substituted  $C_2$ - $C_{12}$  alkenyl, optionally substituted  $C_2$ - $C_{12}$  alkynyl, optionally substituted aryl or optionally substituted heteroaryl.

161. The method of Claim 160, wherein n is 0.

162. The method of Claim 160, wherein n is 1.

163. The method of any one of Claims 160 to 162, wherein  $Y_1$  is O.

164. The method of any one of Claims 160 to 162, wherein  $Y_1$  is  $N^+R_5R_6$ .

165. The method of any one of Claims 160 to 164, wherein  $Y_2$  is O.

166. The method of any one of Claims 160 to 164, wherein  $Y_2$  is  $NR_{10}$ .

167. The method of any one of Claims 160 to 166, wherein  $R_1$  is H.

168. The method of any one of Claims 160 to 166, wherein  $R_1$  is optionally substituted phenyl.

169. The method of any one of Claims 160 to 166, wherein  $R_1$  is optionally substituted  $C_{1-4}$  alkyl.

170. The method of any one of Claims 160 to 169, wherein  $R_2$  is H.

171. The method of any one of Claims 160 to 169, wherein R<sub>2</sub> is optionally substituted phenyl.

172. The method of any one of Claims 160 to 169, wherein R<sub>2</sub> is optionally substituted C<sub>1-4</sub> alkyl.

173. The method of any one of Claims 160 to 169, wherein R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O.

174. The method of any one of Claims 160 to 173, wherein R<sub>3</sub> is H.

175. The method of any one of Claims 160 to 173, wherein R<sub>3</sub> is optionally substituted phenyl.

176. The method of any one of Claims 160 to 173, wherein R<sub>3</sub> is C<sub>1-4</sub> optionally substituted alkyl.

177. The method of any one of Claims 160 to 176, wherein R<sub>4</sub> is H.

178. The method of any one of Claims 160 to 176, wherein R<sub>4</sub> is optionally substituted phenyl.

179. The method of any one of Claims 160 to 173, wherein R<sub>4</sub> is optionally substituted C<sub>1-4</sub> alkyl.

180. The method of any one of Claims 160 to 173, wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

181. The method of any one of Claims 160 to 166, wherein R<sub>1</sub> and R<sub>2</sub> together with the atoms to which they are attached, form =O and wherein R<sub>3</sub> and R<sub>4</sub> together with the atoms to which they are attached, form =O.

182. The method of any one of Claims 160 to 181, wherein R<sub>5</sub> is H.

183. The method of any one of Claims 160 to 181, wherein R<sub>5</sub> is optionally substituted phenyl.

184. The method of any one of Claims 160 to 181, wherein  $R_5$  is optionally substituted  $C_{1-4}$  alkyl.

185. The method of any one of Claims 160 to 184, wherein  $R_6$  is H.

186. The method of any one of Claims 160 to 184, wherein  $R_6$  is optionally substituted phenyl.

187. The method of any one of Claims 160 to 184, wherein  $R_6$  is optionally substituted  $C_{1-4}$  alkyl.

188. The method of any one of Claims 160 to 187, wherein  $R_{10}$  is H.

189. The method of any one of Claims 160 to 187, wherein  $R_{10}$  is optionally substituted phenyl.

190. The method of any one of Claims 160 to 187, wherein  $R_{10}$  is optionally substituted  $C_{1-4}$  alkyl.

191. The method of any one of Claims 160 to 190, wherein  $R_9$  is optionally substituted  $C_1$ - $C_{12}$  alkyl.

192. The method of any one of Claims 160 to 191, wherein  $R_9$  is t-butyl.

193. The method of any one of Claims 160 to 190, wherein  $R_9$  is optionally substituted  $C_3$ - $C_8$  cycloalkyl.

194. The method of any one of Claims 160 to 190, wherein  $R_9$  is optionally substituted  $C_2$ - $C_{12}$  alkenyl.

195. The method of any one of Claims 160 to 190, wherein  $R_9$  is optionally substituted  $C_2$ - $C_{12}$  alkynyl.

196. The method of any one of Claims 160 to 190, wherein R<sub>9</sub> is optionally substituted aryl.

197. The method of any one of Claims 160 to 190, wherein R<sub>9</sub> is optionally substituted heteroaryl.

198. The method of any one of Claims 160 to 166, 169, 172, 176, 179, 184, 187, and 190, wherein each C<sub>1-4</sub> alkyl is optionally substituted with one or more substituents independently selected from the group consisting of OMe, and phenyl.

199. The method of any one of Claims 160 to 166, 169, 172, 176, 179, 184, 187, and 190, wherein each C<sub>1-4</sub> alkyl is optionally substituted with OMe.

200. The method of any one of Claims 160 to 166, 169, 172, 176, 179, 184, 187, and 190, wherein each C<sub>1-4</sub> alkyl is optionally substituted with phenyl.

201. The method of any one of Claims 160 to 166, 168, 171, 175, 178, 183, 186, and 189, wherein each phenyl is optionally substituted with one to three substitutions independently selected from the group consisting of OMe, and halogen.

202. The method of any one of Claims 160 to 166, 168, 171, 175, 178, 183, 186, and 189, wherein each phenyl is optionally substituted with OMe.

203. The method of any one of Claims 160 to 166, 168, 171, 175, 178, 183, 186, and 189, wherein each phenyl is optionally substituted with halogen.



## INTERNATIONAL SEARCH REPORT

 International application No.  
**PCT/US2015/028613**

## A. CLASSIFICATION OF SUBJECT MATTER

**C07F 5/04 (2006.01) C07F 5/02 (2006.01)**

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

PATENTSCOPE: applicant, inventor search.

CA: structure searches based on Formulae (1) and (D)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	Documents are listed in the continuation of Box C	



Further documents are listed in the continuation of Box C



See patent family annex

* "A"	Special categories of cited documents: document defining the general state of the art which is not considered to be of particular relevance	"T"	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E"	earlier application or patent but published on or after the international filing date	"X"	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L"	document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y"	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O"	document referring to an oral disclosure, use, exhibition or other means	"&"	document member of the same patent family
"P"	document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search 17 August 2015	Date of mailing of the international search report 17 August 2015
<b>Name and mailing address of the ISA/AU</b>  AUSTRALIAN PATENT OFFICE PO BOX 200, WODEN ACT 2606, AUSTRALIA Email address: pct@ipaustalia.gov.au	<b>Authorised officer</b>  Geoffrey Peters AUSTRALIAN PATENT OFFICE (ISO 9001 Quality Certified Service) Telephone No. 0262832184

INTERNATIONAL SEARCH REPORT		International application No.
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		PCT/US2015/028613
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2012/021455 A1 (REMPEX PHARMACEUTICALS, INC.) 16 February 2012 Whole Document.	1-203
A	WO 2013/122888 A2 (REMPEX PHARMACEUTICALS, INC.) 22 August 2013 Whole Document.	1-203

INTERNATIONAL SEARCH REPORT		International application No.	
Information on patent family members		PCT/US2015/028613	
This Annex lists known patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.			
Patent Document/s Cited in Search Report		Patent Family Member/s	
Publication Number	Publication Date	Publication Number	Publication Date
WO 2012/021455 A1	16 February 2012	WO 2012021455 A1	16 Feb 2012
		AU 2011289615 A1	21 Mar 2013
		AU 2011289615 B2	07 May 2015
		CA 2807546 A1	16 Feb 2012
		CL 2013000399 A1	23 Aug 2013
		CN 103180328 A	26 Jun 2013
		CO 6680667 A2	31 May 2013
		EP 2603514 A1	19 Jun 2013
		JP 2013535502 A	12 Sep 2013
		KR 20130099923 A	06 Sep 2013
		MX 2013001517 A	29 Apr 2013
		NZ 607354 A	30 Jan 2015
		RU 2013104951 A	20 Sep 2014
		SG 187757 A1	28 Mar 2013
		US 2012040932 A1	16 Feb 2012
		US 8680136 B2	25 Mar 2014
		US 2013345172 A1	26 Dec 2013
WO 2013/122888 A2	22 August 2013	WO 2013122888 A2	22 Aug 2013
		EP 2814483 A2	24 Dec 2014
		US 2015119363 A1	30 Apr 2015
End of Annex			