

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2007264682 B2**

(54) Title  
**Method of treatment of age-related macular degeneration**

(51) International Patent Classification(s)  
**A61K 38/17 (2006.01) A61K 38/20 (2006.01)**  
**A61K 31/00 (2006.01)**

(21) Application No: **2007264682** (22) Date of Filing: **2007.06.28**

(87) WIPO No: **WO08/001380**

(30) Priority Data

(31) Number **60/806,041** (32) Date **2006.06.28** (33) Country **US**

(43) Publication Date: **2008.01.03**  
(44) Accepted Journal Date: **2012.09.06**

(71) Applicant(s)  
**Yeda Research & Development Co. Ltd**

(72) Inventor(s)  
**Butovsky, Oleg;Eisenbach-Schwartz, Michal**

(74) Agent / Attorney  
**FB Rice, Level 23 44 Market Street, Sydney, NSW, 2000**

(56) Related Art  
**US 2005/0118180 A1 (RUBINFELD, J) 2 June 2005**  
**US 2006/0057110 A1 (EISEN BACH-SCHWARTZ, M. AND COHEN, A.) 16 March 2006**  
**US 2003/0004099 A1 (EISEN BACH-SCHWARTZ, M., ET AL) 2 January 2003**  
**WO 2004/060265 A2 (YEDA RESEARCH AND DEVELOPMENT CO. LTD) 22 July 2004**  
**US 2005/0220803 A1 (EISEN BACH-SCHWARTZ, M., AND MIZRAHI, T.) 6 October 2005**  
**US 2004/0092435 A1 (PEYMAN, G.A.) 13 May 2004**  
**US 2004/0248802 A1 (EISEN BACH-SCHWARTZ, M., ET AL.) 9 December 2004**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



PCT

(43) International Publication Date  
3 January 2008 (03.01.2008)

(10) International Publication Number  
WO 2008/001380 A2

(51) International Patent Classification:  
A61K 38/20 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(21) International Application Number:  
PCT/IL2007/000798

(22) International Filing Date: 28 June 2007 (28.06.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/806,041 28 June 2006 (28.06.2006) US

(71) Applicant (for all designated States except US): YEDA RESEARCH AND DEVELOPMENT CO. LTD. [IL/IL]; at the Weizmann Institute of Science, P.O. Box 95, 76100 Rehovot (IL).

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): EISEN-BACH-SCHWARTZ, Michal [IL/IL]; New Scientist Housing, Alley H, House #4, The Weizmann Institute of Science, 76100 Rehovot (IL). BUTOVSKY, Oleg [IL/IL]; 9/1 Alkalay Street, 76304 Rehovot (IL).

(74) Agent: BEN-AMI, Paulina; P.O Box 94, 76100 Rehovot (IL).

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 2008/001380 A2

(54) Title: METHOD OF TREATMENT OF AGE-RELATED MACULAR DEGENERATION

(57) **Abstract:** The invention provides methods and compositions for treatment of age-related macular degeneration, which comprises causing T cells that produce IL-4 to accumulate in the eye by administration of an agent such as Copolymer-1, IL-4, cells activated with IL-4, IL-13 or up to 20 ng/ml IFN- $\gamma$ , or a pathogenic self-antigen associated with a T-cell-mediated specific autoimmune disease of the eye or a peptide derived therefrom, and any combination of such agents.

## METHOD OF TREATMENT OF AGE-RELATED MACULAR DEGENERATION

### 5 FIELD OF THE INVENTION

The present invention relates to methods and compositions for treatment of age-related macular degeneration.

**Abbreviations:** A $\beta$ , amyloid  $\beta$ -peptide; AD, Alzheimer's disease; AMD, age-related macular degeneration; APCs, antigen-presenting cells; BrdU, 5-bromo-2'-deoxyuridine; CNS, central nervous system; COP-1, copolymer 1; GFP, green fluorescent protein; IB-4, *Bandeiraea simplicifolia* isolectin B4; IGF, insulin-like growth factor; IL, interleukin; MG, microglia; MHC-II, class II major histocompatibility complex; MWM, Morris water maze; NPCs, neural stem/progenitor cells; RGCs, retinal ganglion cells; Tg, transgenic; Th, T-helper; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ .

### BACKGROUND OF THE INVENTION

#### Age-related macular degeneration

Age-related macular degeneration (AMD) is a disease affecting the macular region of the eye, which is the area in the retina where the sharp vision is obtained. Macular degeneration is caused by the deterioration of the central portion of the retina, the inside back layer of the eye that records the images we see and sends them via the optic nerve from the eye to the brain. The retina's central portion, known as the macula, is responsible for focusing central vision in the eye, and it controls our ability to read, drive a car, recognize faces or colors, and see objects in fine detail.

AMD is the leading cause of irreversible blindness among the elderly in industrialized nations, and its prevalence increases in the population over the age of 60 (Klein et al., 1992; Mitchell et al., 1995). Numerous attempts have been made to understand the etiology of the disease, its pathophysiology and factors involved in the 5 progression of the disease. A common early sign of AMD is the buildup of drusen, tiny yellow or white fat globules and extracellular material in the retina of the eye or on the optic nerve head. Drusen occurs as hard drusen (small, solid deposits that seem harmless) or larger deposits of soft drusen with indistinct borders. Soft drusen accumulating between the retinal pigment epithelium (RPE) and Bruch's membrane 10 force these two structures apart.

Most people over 40 have a small amount of hard drusen, which can join to form soft drusen in AMD cases. However, not all soft drusen come from hard drusen.

There are two types of macular degeneration: the dry or atrophic type, and the wet or hemorrhagic type. The dry form of AMD, which constitutes 80% of all AMD 15 patients, is characterized by the appearance of drusen. The presence of drusen is considered to be a pre-existing factor associated with the progression of the disease to either advanced dry AMD or wet AMD.

### **Alzheimer's disease**

20 Alzheimer's disease (AD) is an age-related progressive neurodegenerative disorder characterized by memory loss and severe cognitive decline (Hardy & Selkoe, 2002). The clinical features are manifested morphologically by excessive accumulation of extracellular aggregations of amyloid  $\beta$ -peptide (A $\beta$ ) in the form of amyloid plaques in the brain parenchyma, particularly in the hippocampus and cerebral cortex, leading 25 to neuronal loss (Selkoe, 1991). In addition, in most mouse models of Alzheimer's disease the neurogenesis that normally occurs throughout life in the hippocampus of the adult brain is disrupted (Haughey et al., 2002). In Alzheimer patients, like in

transgenic mice (PDGF-APPSw, Ind), some increase in neurogenesis takes place but is apparently not sufficient to overcome the disease (Jin et al., 2004a, b).

### **The similarity between AMD and Alzheimer's disease**

5       AMD and Alzheimer's disease are both chronic neurodegenerative disorders that affect a substantial proportion of elderly persons. Characteristic of these disorders is the irreversible loss of function, for which there is no cure. The degeneration occurring in AMD and Alzheimer's disease may, to some extent, have a common pathogenesis (Klaver et al., 1999). Although the etiology of both AMD and  
10      Alzheimer's disease is largely unknown, the pathogeneses of the two diseases show some striking similarities. In AMD, early histopathological manifestations are extracellular drusen deposits and basal laminar deposits (Hageman & Mullins, 1999). These lesions contain lipids, glycoproteins and glycosaminoglycans, which are presumably derived from a degenerating neuroretina (Kliffen et al., 1995).  
15      Accumulation of these deposits is associated with loss of photoreceptors and subsequent deterioration of macular function (Holz et al., 1994). As noted above, an early pathologic hallmark in Alzheimer's disease is the presence of extracellular senile plaques (Selkoe, 1991). These plaques are composed of many components, including small peptides generated by proteolytic cleavage of a family of transmembrane  
20      polypeptides known as amyloid precursor proteins. Two peptides that are widely regarded as major contributors to the pathology of Alzheimer's disease are known as amyloid- $\beta$  (A $\beta$ ) peptides. Shared components of amyloid deposits and drusen include proteins such as vitronectin, amyloid P, apolipoprotein E, and even the A $\beta$  peptides and amyloid oligomers that are associated with amyloid plaques in Alzheimer's disease  
25      (Luibl et al., 2006; Mullins et al., 2000; Yoshida et al., 2005).

      The A $\beta$  peptides present in Alzheimer's disease activate microglial cells to produce potentially neurotoxic substances such as reactive oxygen and nitrogen species, proinflammatory cytokines, complement proteins, and other inflammatory

mediators that bring about neurodegenerative changes (Akiyama et al., 2000). The inflammatory response that has been associated with Alzheimer's disease often involves CD11b<sup>+</sup> activated microglia, representing the innate arm of the immune system in the central nervous system (CNS) (Streit, 2004). CD11b<sup>+</sup> microglia were 5 reported to be associated with age-related normal human brain (Streit, 2004), and it is possible that such microglia are the ones that contribute both to age-related cognitive loss and to impaired neurogenesis (Monje et al., 2003). CD11b have also been found in patients with Alzheimer's disease (Akiyama & McGeer, 1990). Moreover, inflammatory mediators are present in amyloid deposits as well as in drusen, 10 suggesting a possible common role for the inflammatory pathway in AMD and Alzheimer's disease (Hageman et al., 2001). A role for local inflammation in drusen biogenesis suggests that it is analogous to the process that occurs in Alzheimer's disease, where accumulation of extracellular plaques and deposits elicits a local chronic inflammatory response that exacerbates the effects of primary pathogenic stimuli 15 (Akiyama et al., 2000).

### **Microglial activation in neurodegeneration**

Microglia are bone marrow-derived glial cells, which are present within all layers of the adult human retina (Penfold et al., 1991). Several types are present which 20 may be associated with neurons or with blood vessels, and some of these are antigen-presenting cells (APCs) (Penfold et al., 1991; Provis, 2001). The nature of microglial activation, either beneficial or harmful, in damaged neural tissue depends on how microglia interpret the threat (Butovsky et al., 2005). Although the presence of microglial cells in normal undamaged neural tissue has been debated for years, it is 25 now an accepted fact (Nimmerjahn et al., 2005), including their presence in the eye. The role of microglia in inflammatory processes is controversial. On the one hand, participation of microglia in inflammatory process of the eye can stimulate mature retinal ganglion cells (RGCs) to regenerate their axons (Yin et al., 2003). On the other

hand, the role of microglia in neurodegenerative processes may be detrimental to the neuronal tissue. Roque et al (1999) showed that microglial cells release soluble product(s) that induce degeneration of cultured photoreceptor cells. This controversy may be explained by the contradicting reports regarding the presence of antigen-presenting cells, which are crucial factors of an antigen-specific cell-mediated immune response. Immunological responses in neural retinal microglia are related to early pathogenic changes in retinal pigment epithelium pigmentation and drusen formation. Activated microglia may also be involved in rod cell death in AMD and late-onset retinal degeneration. A recent study has proposed that microglia, activated by primary rod cell death, migrate to the outer nuclear layer, remove rod cell debris and may kill adjacent cone photoreceptors (Gupta et al., 2003).

Like blood-derived macrophages, microglia exhibit scavenging of extracellular deposits, and phagocytosis of abnormal amyloid deposits in Alzheimer's disease. Such microglia, while efficiently acting as phagocytic cells, cause neuronal death by the secretion of mediators like tumor necrosis factor alpha (TNF- $\alpha$ ) (Butovsky et al., 2005), and thus, while acting as phagocytic cells (Frenkel et al., 2005), they are apparently not efficient enough to fight off the Alzheimer's disease symptoms. In contrast to these resident microglia, microglia derived from the bone marrow of matched wild-type mice can effectively remove plaques (Simard et al., 2006). Moreover, an absence of normally functioning macrophages lead to the development of clinical AMD (Ambati et al., 2003). Thus, AMD, like Alzheimer's disease, illustrates a disease in which scavenging of abnormal deposits inevitably induces self-perpetuation of disease progression mediated by the phagocytic cell themselves (Gupta et al., 2003).

## 25 **Protective autoimmunity**

Some years ago our group formulated the concept of 'protective autoimmunity' (Moalem et al., 1999). Both pro-inflammatory and anti-inflammatory cytokines were found to be critical components of a T cell-mediated beneficial autoimmune response,

provided that the timing and the intensity of the T-cell activity was suitably controlled (Butovsky et al., 2005; Shaked et al., 2004), and depending on the nature of the disease (Schwartz et al., 2006). According to our concept, an uncontrolled autoimmunity leads to the commonly known condition of autoimmune diseases associated with 5 overwhelmed activation of microglia (Butovsky et al., 2006a), as will be discussed below. The beneficial effect of the autoreactive T cells was found to be exerted via their ability to induce the CNS-resident microglia to adopt a phenotype capable of presenting antigens (Butovsky et al., 2001; Butovsky et al., 2005; Schwartz et al., 2006; Butovsky et al., 2006a; Shaked et al., 2004), expressing growth factors 10 (Butovsky et al., 2005; Butovsky et al., 2006a;b), and buffering glutamate (Shaked et al., 2005).

In attempting to boost the efficacy of the protective autoreactive T cells, we tested many compounds in the search for a safe and suitable antigen for neuroprotection. We then suggested to use glatiramer acetate, also known as 15 Copolymer 1 or Cop-1 (Kipnis et al., 2000; Avidan et al., 2004; Angelov et al., 2003), a synthetic 4-amino-acid copolymer known to be safe and currently used as a treatment for multiple sclerosis by a daily administration regimen (Copaxone®, Teva Pharmaceutical Industries Ltd., Israel). In our studies we have demonstrated its low-affinity cross-reaction with a wide range of CNS autoantigens. Because the affinity of 20 cross-reaction is low, the Cop-1-activated T cells, after infiltrating the CNS, have the potential to become locally activated with little or no attendant risk of autoimmune disease (Kipnis et al., 2000).

A single injection of Cop-1 is protective in acute models of CNS insults (Kipnis et al., 2000; Avidan et al., 2004; Kipnis. & Schwartz, 2002), while in chronic models 25 occasional boosting is required for a long-lasting protective effect (Angelov et al., 2003). In the rat model of chronically high intraocular pressure, vaccination with Cop-1 significantly reduces RGC loss even if the pressure remains high. It should be noted that the vaccination does not prevent disease onset, but can slow down its progression

by controlling the local extracellular environment of the nerve and retina, making it less hostile to neuronal survival and allowing the RGCs to be better able to withstand the stress (Schori et al., 2001; Benner et al., 2004; Kipnis & Schwartz, 2002; Kipnis et al., 2000).

5 For chronic conditions occasional boosting is needed. For example, in a model of chronically elevated intraocular pressure, weekly administration of adjuvant-free Cop-1 was found to result in neuroprotection (Bakalash et al., 2005). The neuroprotective effect of Cop-1 has been attributed in part to production of brain-derived neurotrophic factor (BDNF) (Kipnis et al., 2004b; Ziemssen et al., 2002).

10

#### **Aggregated A $\beta$ induces toxicity on resident microglia and impairs cell renewal**

Recent studies performed in our laboratory suggested that microglia exposed to aggregated A $\beta$ , although effective in removing plaques, are toxic to neurons and impair neural cell renewal (Butovsky et al., 2006a); these effects are reminiscent of the response of microglia to invading microorganisms (as exemplified by their response to LPS) (Butovsky et al., 2005; Schwartz et al., 2006). Such activities are manifested by increased production of TNF- $\alpha$ , down-regulation of insulin-like growth factor (IGF-I), inhibition of the ability to express class II major histocompatibility complex (MHC-II) proteins and CD11c (a marker of dendritic cells) and thus to act as antigen-presenting cells (APCs), and failure to support neural tissue survival and renewal ((Butovsky et al., 2006a; Butovsky et al., 2005; Butovsky et al., 2006b). Further, we found that when microglia encounter aggregated  $\beta$ -amyloid, their ability to remove these aggregates without exerting toxic effects on neighboring neurons or impairing neurogenesis depends upon their undergoing a phenotype switch. A switch in microglial phenotype might take place via a local dialog between microglia and T-cells, which is mediated by T cell-derived cytokines such as interleukin (IL)-4. Addition of IL-4, a cytokine derived from T-helper (Th)-2 cells, to microglia activated by aggregated A $\beta$  can

reverse the down-regulation of IGF-I expression, the up-regulation of TNF- $\alpha$  expression, and the failure to act as APCs (Butovsky et al., 2005). The significance of microglia for in-vivo neural cell renewal was demonstrated by enhanced neurogenesis in the rat dentate gyrus after injection of IL-4-activated microglia 5 intracerebroventricularly and by the presence of IGF-I-expressing microglia in the dentate gyrus of rats kept in an enriched environment (Ziv et al., 2006). In rodents with acute or chronic EAE, injection of IL-4-activated microglia into the cerebrospinal fluid resulted in increased oligodendrogenesis in the spinal cord and improved clinical symptoms. The newly formed oligodendrocytes were spatially associated with 10 microglia expressing MHC-II and IGF-I (Butovsky et al., 2006c).

#### **In both Alzheimer's disease and AMD there are systemic components**

Our first observation that systemic immune cells (in the form of T cells directed to certain self-antigens) can protect injured neurons from death came from studies in 15 rodents showing that passive transfer of T cells specific to myelin basic protein reduces the loss of RGCs after a traumatic optic nerve injury (Moalem et al., 1999). We found that these T cells are also effective when directed to either cryptic or pathogenic epitopes of myelin basic protein, as well as to other myelin antigens or their epitopes (Mizrahi et al., 2002). These findings raised a number of critical questions. For 20 example, are myelin antigens capable of protecting the nervous system from any type of acute or chronic insult? Is the observed neuroprotective activity of immune cells merely an anecdotal finding reflecting our experimental conditions, or does it point to the critical participation of the immune system in fighting off injurious conditions in the visual system and in the CNS in general? If the latter, does it mean that 25 neurodegenerative diseases are systemic diseases? If so, can this finding be translated into a systemic therapy that would protect the brain, the eye, and the spinal cord?

In a series of experiments carried out over the last few years we have learned, firstly, that protective T cell response is a physiologically evoked response that might

not be sufficient in severe insults or might not always be properly controlled. Moreover, we discovered that the specificity of such protective T cells depends on the site of the insult. Thus, for example, the protective effect of vaccination with myelin-associated antigens is restricted to injuries of the white matter, i.e., to myelinated axons 5 (Mizrahi et al., 2002; Avidan et al., 2004; Schori et al., 2001). If the insult is to the retina, which contains no myelin, myelin antigens have no effect. Secondly, we observed that the injury-induced response of T cells reactive to specific self-antigens residing in the site of stress (eye or brain) is a spontaneous physiological response (Yoles et al., 2001). We then sought to identify the phenotype of the beneficial 10 autoimmune T cells and to understand what determines the balance between a beneficial (neuroprotective) outcome of the T cell-mediated response to a CNS injury and a destructive effect causing autoimmune disease. We also examined ways of translating the beneficial response into a therapy for glaucoma. We found that in immune deficient animals the number of surviving RGCs following an insult in the 15 eye, the spinal cord or the brain is significantly lower than in matched controls with an intact immune system, suggesting that the ability to withstand insult to the CNS depends on the integrity of the immune system and specifically on specific population within the immune system; those that recognize the site-specific self-antigens. Interestingly, the use of steroids caused significant loss of RGCs (Bakalash et al., 20 2003).

#### **T cells specific to antigens residing in the site of damage help clean and heal**

In order to be protective, the anti-self T cells should home to the site of damage and be locally activated. This is why only those antigens that are being presented at the 25 site of lesion can be used for the vaccination. Once activated, the T cells provide a source of cytokines and growth factors that shape the resident eye sentinels cells – the microglia, so as to make them active defensible cells that the eye can tolerate. Namely, such activated microglia can take up glutamate, remove debris and produce growth

factors while refraining from production of agents that are part of their killing mechanism (e.g. TNF- $\alpha$ ) to which the eye, like the brain, has a low tolerance (Butovsky et al., 2005; Butovsky et al., 2001; Barouch & Schwartz, 2002; Moalem et al., 2000; Shaked et al., 2005). Such T cells are constitutively controlled by 5 physiologically existing regulatory T cells that are themselves amenable to control upon need (Kipnis et al., 2004a; Kipnis et al., 2002).

Reference is made to copending International Patent Application No. PCT/IL2007/001379 entitled "Activated myeloid cells for promoting tissue repair and detecting damaged tissue" filed by applicant at the Israel PCT Receiving Office 10 (RO/IL) on the same date, the contents thereof being explicitly excluded from the scope of the present invention.

Citation of any document herein is not intended as an admission that such document is pertinent prior art, or considered material to the patentability of any claim of the present application. Any statement as to content or a date of any document is 15 based on the information available to applicant at the time of filing and does not constitute an admission as to the correctness of such a statement.

Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general 20 knowledge in the field relevant to the present disclosure as it existed before the priority date of each claim of this application.

## **SUMMARY OF THE INVENTION**

It is a preferred aim the present invention to provide an immune-based therapy 25 for age-related macular degeneration.

In one aspect, the present invention provides a method for treatment of age-related macular degeneration, which comprises causing T cells that produce IL-4 to accumulate in the eye of a patient in need by administering a single active agent

2007264682

08 Aug 2012

selected from the group consisting of Copolymer 1 or T cells activated against Copolymer 1 to said individual, thereby halting or delaying progress of the macular degeneration.

In another aspect, the invention provides a use of a single active agent that causes T cells that produce IL-4 to accumulate in the eye, wherein the single active agent is selected from the group consisting of Copolymer 1 or T cells activated against Copolymer 1, for the preparation of a medicament for treatment of age-related macular degeneration.

#### **BRIEF DESCRIPTION OF THE DRAWINGS**

**Figs. 1A-1D** demonstrate that IL-4 can counteract the adverse effect of aggregated A $\beta$  on microglial toxicity and promotion of neurogenesis in adult mouse neural progenitor cells. (**Fig. 1A**) *In-vitro* treatment paradigm. (**Fig. 1B**) Representative confocal microscopic images of neural progenitor cells (NPCs) expressing green

fluorescent protein (GFP) and  $\beta$ III-T (neuronal marker), co-cultured for 10 days without microglia (MG, control), or with untreated microglia, or with microglia that were pre-activated by aggregated  $A\beta_{(1-40)}$  (5  $\mu$ M) ( $MG_{(A\beta_{1-40})}$ ) for 48 h and subsequently activated with  $IFN-\gamma$  (10 ng/ml) ( $MG_{(A\beta_{1-40}/IFN\gamma)}$ ) or with IL-4 (10 ng/ml) 5 ( $MG_{(A\beta_{1-40}/IL-4)}$ ) or with both  $IFN-\gamma$  (10 ng/ml) and IL-4 (10 ng/ml) ( $MG_{(A\beta_{1-40}/IFN\gamma+IL-4)}$ ). Note, aggregated  $A\beta$  induced microglia to adopt an amoeboid morphology, but after IL-4 was added they exhibited a ramified structure. (Fig. 1C) Separate confocal images 10 of NPCs co-expressing GFP and  $\beta$ III-T adjacent to  $CD11b^+$  microglia. (Fig. 1D) Quantification of cells double-labeled with GFP and  $\beta$ III-T (expressed as a percentage 15 of  $GFP^+$  cells) obtained from confocal images. Results are of three independent experiments in replicate cultures; bars represent means  $\pm$  SEM. Asterisks above bars denote the significance of differences relative to untreated (control) NPCs ( $^*P < 0.05$ ;  $^{***}P < 0.001$ ; two-tailed Student's t-test). Horizontal lines with  $P$  values above them show differences between the indicated groups (ANOVA).

15 **Figs. 2A-2L** show that Cop-1 vaccination leads to reduction in  $\beta$ -amyloid and counteracts loss of hippocampal neurons in the brains of transgenic Alzheimer's disease mice: key role of microglia. (Fig. 2A) Representative confocal microscopic images of brain hippocampal slices from non-transgenic (Tg), untreated-Tg-Alzheimer's disease (AD), and Cop-1-vaccinated Tg-AD mice stained for NeuN (mature neurons) and human  $A\beta$ . The non-Tg mouse shows no staining for human  $A\beta$ . The untreated-Tg-AD mouse shows an abundance of extracellular  $A\beta$  plaques, whereas 20 in the Cop-1-treated Tg-AD mouse  $A\beta$ -immunoreactivity is low. Weak NeuN $^+$  staining is seen in the hippocampal CA1 and dentate gyrus regions of the untreated-Tg-AD mouse relative to its non-Tg littermate, whereas NeuN $^+$  staining in the Cop-1- 25 vaccinated Tg-AD mouse is almost normal. (Fig. 2B) Staining for activated microglia using anti- $CD11b$  antibodies. Images at low and high magnification show a high incidence of microglia double-stained for  $A\beta$  and  $CD11b$  in the CA1 and dentate gyrus

regions of the hippocampus of an untreated-Tg-AD mouse, but only a minor presence of CD11b<sup>+</sup> microglia in the Cop-1-vaccinated Tg-AD mouse. Arrows indicate areas of high magnification, shown below. (Fig. 2C) CD11b<sup>+</sup> microglia, associated with an A $\beta$  plaque, strongly expressing TNF- $\alpha$  in an untreated-Tg-AD mouse. (Fig. 2D) Staining 5 for MHC-II (a marker of antigen presentation) in a cryosection taken from a Cop-1-vaccinated Tg-AD mouse in an area that stained positively for A $\beta$  shows a high incidence of MHC-II<sup>+</sup> microglia and almost no TNF- $\alpha$ <sup>+</sup> microglia. (Fig. 2E) All MHC-II<sup>+</sup> microglia in a brain area that stained positively for A $\beta$  (arrowheads) in a Cop-1-vaccinated Tg-AD mouse co-express CD11c (a marker of dendritic cells), but only a 10 few CD11c<sup>+</sup>/MHC-II<sup>+</sup> microglia are seen in a corresponding area in the brain of an untreated-Tg-AD mouse. (Fig. 2F) MHC-II<sup>+</sup> microglia in a Cop-1-vaccinated Tg-AD mouse co-expresses IGF-I. (Fig. 2G) CD3<sup>+</sup> T cells are seen in close proximity to an A $\beta$ -plaque and (Fig. 2H) are associated with MHC-II<sup>+</sup> microglia. Boxed area shows 15 high magnification of an immunological synapse between a T cell (CD3<sup>+</sup>) and a microglial cell expressing MHC-II. (Fig. 2I) Histogram showing the total number of A $\beta$ -plaques (in a 30- $\mu$ m hippocampal slice). (Fig. 2J) Histogram showing staining for A $\beta$ -immunoreactivity. Note the significant differences between Cop-1-vaccinated Tg-AD and untreated-Tg-AD mice, verifying the decreased presence of A $\beta$ -plaques in the 20 vaccinated mice. (Fig. 2K) Histogram showing a marked reduction in cells stained for CD11b, indicative of activated microglia and inflammation, in the Cop-1-vaccinated Tg-AD mice relative to untreated-Tg-AD mice. Note the increase in CD11b<sup>+</sup> microglia with age in the non-Tg littermates. (Fig. 2L) Histogram showing significantly more CD3<sup>+</sup> cells associated with an A $\beta$ -plaque in Cop-1-vaccinated Tg-AD mice than in 25 untreated-Tg-AD mice. Quantification of CD3<sup>+</sup> cells was analyzed from 30–50 plaques of each mouse tested in this study. Error bars indicate means  $\pm$  SEM. \* $P$  < 0.05, \*\*\* $P$  < 0.001 versus non-Tg littermates (Student's *t*-test). The  $P$  values indicated in the figure

represent a comparison of the groups as analyzed by ANOVA. All of the mice in this study were included in the analysis (6–8 sections per mouse).

**Figs. 3A-3C** show that Cop-1 vaccination induces microglia to express CD11c. (Fig. 3A) CD11b<sup>+</sup> microglia co-expressing CD11c surround an A $\beta$ -plaque in Cop-1-vaccinated Tg-AD mice. All of the CD11c-expressing microglia are co-labeled for CD11b. Separate confocal channel is shown in right panel. (Fig. 3B) Histograms showing the number of CD11b<sup>+</sup> cells associated with A $\beta$ -plaque. (Fig. 3C) Histograms showing quantification of CD11c<sup>+</sup> cells as a percentage of the total number of CD11b<sup>+</sup> and CD11c<sup>+</sup> cells associated with an A $\beta$ -plaque. For this analysis, cells were counted surrounding 30–50 plaques in each mouse tested. Error bars represent means  $\pm$  SEM. Asterisks above bars denote the significance of differences between the groups ( $^{**}P < 0.01$ ;  $^{***}P < 0.001$ ; two-tailed Student's *t*-test).

**Figs. 4A-4D** show that Cop-1 vaccination induces microglia to express CD11c: role of IL-4. (Fig. 4A) IL-4-activated microglia (MG<sub>(IL-4)</sub>) induce CD11c expression in a primary culture of mouse microglia 5 days after activation. Untreated microglia (MG<sub>(-)</sub>) express hardly any CD11c. (Fig. 4B) Effect of IL-4 (in terms of morphology and CD11c expression) on microglia pretreated for 3 days with aggregated A $\beta$ <sub>(1-40)</sub> (MG<sub>(A $\beta$ )</sub>) and assessed 10 days later compared to IL-4 treatment for 10 days without pre-exposure to A $\beta$ . Note that dendritic-like morphology was adopted upon addition of IL-4 to the A $\beta$ -pretreated microglia only, whereas CD11c expression was induced by IL-4 both with and without A $\beta$  pretreatment. (Fig. 4C) Quantitative analysis of microglial expression of CD11c<sup>+</sup> microglia (expressed as a percentage of IB-4-labeled microglia) and of CD11c intensity per cell, both expressed as a function of time in culture with or without IL-4. (Fig. 4D) Quantitative analysis of CD11c expression (calculated as a percentage of IB-4-labeled microglia) by the cultures shown in (Fig. 4B). Results are of three independent experiments in replicate cultures; bars represent

means  $\pm$  SEM. Asterisks above bars denote the significance of differences relative to untreated microglia at each time point ( $^{***}P < 0.001$ ; two-tailed Student's *t*-test).

**Figs. 5A-5B** show engulfment of aggregated A $\beta$  by activated microglia. Microglia were treated with IL-4 (10 ng/ml) 24 h after seeding (MG<sub>(IL-4)</sub>) or were left untreated for 48 h (MG<sub>(-)</sub>). The media were then replaced by a labeling medium (DMEM containing 10 mg/ml bovine serum albumin), and aggregated A $\beta$ <sub>(1-40)</sub> was added (5  $\mu$ g/ml) for 1 h. Following incubation the cultures were fixed and immunostained with antibodies directed to human A $\beta$  and co-stained for microglia (IB-4). (**Fig. 5A**) Confocal photomicrographs. (**Fig. 5B**) Quantitative analysis expressed as intensity per cell. Results of one of two experiments, each containing eight replicates (20–30 cells per replicate) per group are presented (means  $\pm$  SD).

**Figs. 6A-6E** depict enhanced neurogenesis induced by Cop-1 vaccination in the hippocampal dentate gyri of adult transgenic (Tg) AD mice. Three weeks after the first Cop-1 vaccination, mice in each experimental group were injected i.p. with 5-bromo-2'-deoxyuridine (BrdU) twice daily for 2.5 days. Three weeks after the last injection their brains were excised and the hippocampi analyzed for BrdU, DCX (a marker of early differentiation of the neuronal lineage), and NeuN (a marker of mature neurons). (**Figs. 6A-6C**) Histograms showing quantification of the proliferating cells (BrdU $^{+}$ ). (**Fig. 6A**) Newly formed mature neurons (BrdU $^{+}$ /NeuN $^{+}$ ) (**Fig. 6B**), and all pre-mature (DCX $^{+}$ -stained) neurons (**Fig. 6C**). Numbers of BrdU $^{+}$ , BrdU $^{+}$ /NeuN $^{+}$  and DCX $^{+}$  cells per dentate gyrus (DG), calculated from six equally spaced coronal sections (30  $\mu$ m) from both sides of the brains of all the mice tested in this study. Error bars represent means  $\pm$  SEM. Asterisks above bars denote the significance of differences relative to non-Tg littermates ( $^{**}P < 0.01$ ;  $^{***}P < 0.001$ ; two-tailed Student's *t*-test). Horizontal lines with *P* values above them show differences between the indicated groups (ANOVA). (**Fig. 6D**) Representative confocal microscopic images of the dentate gyrus showing immunostaining for BrdU/DCX/NeuN in a Cop-1-vaccinated Tg-AD mouse

and in a non-Tg littermate relative to that in an untreated-Tg-AD mouse. (Fig. 6E) Branched DCX<sup>+</sup> cells are found near MHC-II<sup>+</sup> microglia located in the subgranular zone (SGZ) of the hippocampal dentate gyrus of a Cop-1-vaccinated Tg-AD mouse.

Figs. 7A-7B show that Cop-1 vaccination counteracts cognitive decline in 5 transgenic (Tg) AD mice. Hippocampus-dependent cognitive activity was tested in the Morris water maze (MWM). (Figs. 7A-7B) Cop-1-vaccinated Tg-AD mice (*diamond*;  $n = 6$ ) showed significantly better learning/memory ability than untreated-Tg-AD mice (*square*;  $n = 7$ ) during the acquisition and reversal. Untreated-Tg-AD mice showed consistent and long-lasting impairments in spatial memory tasks. In contrast, 10 performance of the MWM test by the Cop-1-vaccinated Tg-AD mice was rather similar, on average, to that of their age-matched naïve non-Tg littermates (*triangle*;  $n = 6$ ) (3-way ANOVA, repeated measures: groups, df (2,16),  $F = 22.3$ ,  $P < 0.0002$ ; trials, df (3,48),  $F = 67.9$ ,  $P < 0.0001$ ; days, df (3,48),  $F = 3.1$ ,  $P < 0.035$ , for the acquisition phase; and groups, df (2,16),  $F = 14.9$ ,  $P < 0.0003$ ; trials, df (3,48),  $F = 21.7$ ,  $P < 15 0.0001$ ; days, df (1,16),  $F = 16.9$ ,  $P < 0.0008$ , for the reversal phase).

Figs. 8A-8D show a time course of CD11c expression in microglia activated by IFN- $\gamma$  and IL-4. (Fig. 8A) Microglia were treated with IFN- $\gamma$  (10 ng/ml; MG<sub>(IFN- $\gamma$ )</sub>) or IL-4 (10 ng/ml; MG<sub>(IL-4)</sub>) for 1, 3, 5, 10 and 18 days. Untreated microglia (MG<sub>(-)</sub>) were used as controls. (Fig. 8B) Confocal images of microglia, identified by staining for 20 IB4, immunolabeled for  $\beta$ III-T, and CD11c after 5 days of treatment. MG<sub>(-)</sub> did not express CD11c. After exposure to IFN- $\gamma$  or IL-4 the microglia expressed CD11c and exhibited their characteristic morphology. (Fig. 8C) Co-expression of  $\beta$ III-T and CD11c in microglia activated with IFN- $\gamma$  (10 ng/ml) for 5 days (IB4/ $\beta$ III-T/CD11c). Note, confocal channels are presented separately. (Fig. 8D) Quantitative analysis of the 25 numbers of CD11c<sup>+</sup> microglia (expressed as a percentage of IB4<sup>+</sup> (microglia marker) cells) were examined in all treatments at all time points. Results are of four independent experiments with duplicate or triplicate wells; bars represent means  $\pm$  SEM. Asterisks above bars denote the significance of differences relative to MG<sub>(-)</sub> (\* $P$

< 0.05; \*\* $P$  < 0.01; \*\*\* $P$  < 0.001; two-tailed Student's  $t$ -test). Horizontal lines with  $P$  values above them show differences between the indicated groups (ANOVA).

## DETAILED DESCRIPTION OF THE INVENTION

5 In searching for a prospect for a T-cell-based vaccination for treatment of AMD, the following considerations may be relevant.

In general, the brain, like the retina, is considered to be immune privileged in the sense that the blood-brain barrier resists passive deposition of antibodies and reduces the recruitment of antigen-specific lymphocytes (Streilein et al., 1992).

10 Paradoxically, antigen-specific immunity might actually function to protect against degenerative diseases. Recently, it was shown in a Alzheimer's disease mouse model that immunization with the abnormal amyloid, or passive administration of antibodies against the abnormal protein, greatly reduced the quantity of deposition in the brain of the genetically modified mice, and improved their performance in laboratory tests of 15 memory and cognitive function (e.g. Morgan et al., 2000). The mechanisms are unclear, but may be related to enhanced phagocytosis, neutralization of toxic molecules, or interference with amyloid fibril aggregation. In Alzheimer's disease however, patients may be immunologically tolerant to amyloid, preventing protective autoimmunization to the abnormally processed protein and thus developing 20 autoimmune encephalomyelitis (Furlan et al., 2003).

Studies from our the laboratory over the last few years have shown that recovery from CNS injury is critically dependent on the well-controlled activity of T cells directed to specific CNS autoantigens (Moalem et al., 1999; Yoles et al., 2001; Kipnis et al., 2002). After homing to the site of damage, these autoreactive T cells evidently 25 regulate microglia in a way that renders them supportive of neuronal survival and neural tissue repair (Butovsky et al., 2005; Schwartz et al., 2006; Butovsky et al., 2001; Shaked et al., 2005).

Our results herein argue in favor of the use of a myelin-related antigen such as Cop-1, but not an A $\beta$  peptide, as a T cell-based therapy for AMD. Even if any T cells expressing T-cell receptors for drusen-associated peptides such as A $\beta$  were to home to the site of a CNS lesion (Monsonego et al., 2006), it is unlikely that they would 5 encounter their relevant APCs there, and they would therefore not be able to become locally activated.

On the other hand, myelin-presenting microglia, with which myelin-specific T cells can readily hold a dialog, are likely to be present at the damaged sites. Myelin-related antigens, or antigens that are weakly cross-reactive with myelin (such as Cop-10 1), are therefore likely to be the antigens of choice for therapeutic vaccination (Avidan et al., 2004). T cells activated by these antigens will then home to the CNS and, upon 15 encountering their relevant APCs there, become locally activated to supply the cytokines and growth factors in order to switch the phenotype of the harmful microglia (activated by aggregated A $\beta$ ; Butovsky et al., 2005) into microglia with dendritic-like characteristics. The resulting immunological synapse between T cells and microglia will then create a supportive niche for cell renewal by promoting neurogenesis from the pool of adult stem cells (Butovsky et al., 2006a).

Our results indicate that T cells constitute the immune-based therapy of choice for AMD. This does not preclude the potential benefit of antibodies as a supplementary 20 therapy as shown in animal model of Alzheimer disease with antibodies against A $\beta$  peptide (Bard et al., 2000). Moreover, the T cells can function as a mini-factory capable of producing a variety of compounds, including cytokines and neurotrophic factors (Ziemssen et al., 2002). Above all, they represent a physiological system of maintenance and repair that might help to counteract the age-related conditions leading 25 to brain senescence.

In developing the immune-based therapy for AMD according to the present invention we thus took into consideration the lessons from neurodegenerative diseases with similar pathogenicity.

Formation of extracellular deposits consisting of misfolded protein is the hallmark of many neurodegenerative diseases. The accumulation of amyloid in drusen and Alzheimer's disease and the presence of activated microglia as inflammatory mediators in both neurodegenerative conditions, suggests a possible common chronic inflammatory pathway in AMD and Alzheimer's disease. In the context of Alzheimer's disease we have recently demonstrated that aggregated  $\beta$ -amyloid (A $\beta$ ) activates microglia to acquire a phenotype which is reminiscence of that activated by microorganisms. Although such microglia/macrophage can act as phagocytic cells their overall activity is cytotoxic and can hardly be tolerated by the brain. As a result, microglia activated by A $\beta$  rather than help the suffering tissue further contribute to the chaos. In the present invention, we explore possible immune-based therapy to modulate microglia activity in Alzheimer's disease and AMD with the target of maintaining their phagocytic activity while conferring their ability to support cell survival and renewal.

As shown herein, aggregated A $\beta$  induces microglia to become cytotoxic and block neurogenesis from adult rodent neural progenitor cells (NPCs). Addition of IL-4, a cytokine derived from T-helper (Th)2 cells, to microglia activated by A $\beta$  can reverse the impediment, the down-regulation of IGF-I, the up-regulation of TNF- $\alpha$ , and the failure to act as APCs. Using Alzheimer's disease double-transgenic mice expressing mutant human genes encoding presenilin 1 and chimeric mouse/human amyloid precursor protein, we show that switching of microglia phenotype into professional APCs producing IGF-I, achieved here by a T cell-based vaccination with Copolymer-1, resulted in reduction of amyloid loads and the induction of neuronal survival and neurogenesis.

The present invention thus relates to a method for treatment of age-related macular degeneration (AMD), which comprises causing T cells that produce IL-4 to accumulate in the eye of an individual in need by administering an agent selected from the group consisting of Copolymer 1 or T cells activated against Copolymer 1 to said

individual, thereby halting or delaying progress of the macular degeneration. This effect can be affected by several self antigens and cytokine activated cells.

Herein disclosed are agents that can cause T cells producing IL-4 to accumulate in the eye include, without being limited to:

- (i) Copolymer-1, a Copolymer-1-related-peptide, or a Copolymer 1-related polypeptide;
- 5 (ii) IL-4;
- (iii) dendritic cells, monocytes, bone marrow-derived myeloid cells or peripheral blood mononuclear cells activated by IL-4 ;
- (iv) genetically engineered cells that produce IL-4;
- 10 (v) bone marrow-derived myeloid cells or peripheral blood-derived myeloid cells activated with IL-13 or with up to 20 ng/ml IFN- $\gamma$ ;
- (vi) a pathogenic self-antigen associated with a T-cell-mediated specific autoimmune disease of the eye;
- 15 (vii) a peptide which sequence is comprised within the sequence of said pathogenic self-antigen of (vi) or a peptide obtained by modification of said peptide, which modification consists in the replacement of one or more amino acid residues of the peptide by different amino acid residues (hereinafter "modified peptide"), said modified peptide still being capable of recognizing the T-cell receptor recognized by the parent peptide but with less affinity;
- 20 (viii) a nucleotide sequence encoding a pathogenic self-antigen of (vi) or a peptide or a modified peptide of (vii) ;
- (ix) T cells activated by an agent of (i), (vi) or (vii); and
- (x) any combination of (i) – (ix).

In one preferred embodiment, the agent is Copolymer 1, a Copolymer 1-related peptide or a Copolymer 1-related polypeptide.

For the purpose of the present invention, "Copolymer 1 or a Copolymer 1-related peptide or polypeptide" is intended to include any peptide or polypeptide,

including a random copolymer that cross-reacts functionally with MBP and is able to compete with MBP on the MHC class II in the antigen presentation.

The Cop 1 or a Cop 1-related peptide or polypeptide is represented by a random copolymer consisting of a suitable ratio of a positively charged amino acid such as 5 lysine or arginine, in combination with a negatively charged amino acid (preferably in a lesser quantity) such as glutamic acid or aspartic acid, optionally in combination with a non-charged neutral amino acid such as alanine or glycine, serving as a filler, and optionally with an amino acid adapted to confer on the copolymer immunogenic properties, such as an aromatic amino acid like tyrosine or tryptophan. Such 10 copolymers are disclosed, for example, in WO 00/05250, the entire contents of which are herewith incorporated herein by reference.

More specifically, the Copolymer 1 or a Copolymer 1-related peptide or polypeptide is a copolymer selected from the group consisting of random copolymers comprising one amino acid selected from each of at least three of the following groups: 15 (a) lysine and arginine; (b) glutamic acid and aspartic acid; (c) alanine and glycine; and (d) tyrosine and tryptophan. The amino acids may be L- or D-amino acids or mixtures thereof. The present invention contemplates the use of copolymers containing both D- and L-amino acids, as well as copolymers consisting essentially of either L- or D-amino acids.

20 In one embodiment of the invention, the copolymer contains four different amino acids, each from a different one of the groups (a) to (d).

In a more preferred embodiment, the agent is Copolymer 1, composed of a mixture of random polypeptides consisting essentially of the amino acids L-glutamic acid (E), L-alanine (A), L-tyrosine (Y) and L-lysine (K) in an approximate ratio of 25 1.5:4.8:1:3.6, having a net overall positive electrical charge and of a molecular weight from about 2 KDa to about 40 KDa.

In one preferred embodiment, the Cop 1 has average molecular weight of about 2 KDa to about 20 KDa, more preferably of about 4,7 KDa to about 13 K Da, still more

preferably of about 4 KDa to about 8.6 KDa, of about 5 KDa to 9 KDa, or of about 6.25 KDa to 8.4 KDa. In another preferred embodiment, the Cop 1 has average molecular weight of about 13 KDa to about 20 KDa, more preferably of about 13,5 KDa to about 18 KDa, with an average of about 15 KDa to about 16 KDa, preferably of 5 16kDa. Other average molecular weights for Cop 1, lower than 40 KDa, are also encompassed by the present invention. Copolymer 1 of said molecular weight ranges can be prepared by methods known in the art, for example by the processes described in U.S. Patent No. 5,800,808, the entire contents of which are hereby incorporated by reference in the entirety. The Copolymer 1 may be a polypeptide comprising from 10 about 15 to about 100, preferably from about 40 to about 80, amino acids in length.

In one more preferred embodiment of the invention, the agent is Cop 1 in the form of its acetate salt known under the generic name glatiramer acetate or its trade name Copaxone® (a trademark of Teva Pharmaceutical Industries Ltd., Petach Tikva, Israel). As used herein in the application, the terms "Cop 1", "Copolymer 1", 15 "glatiramer acetate" and "GA" are used interchangeably.

In another embodiment, the Copolymer 1-related peptide is a random copolymer of 4 amino acids in which one or more of the following substitutions is made: aspartic acid for glutamic acid, glycine for alanine, arginine for lysine, and tryptophan for tyrosine, which is expected to have the same activity of Copolymer.

20 In another embodiment of the invention, the Cop 1-related peptide or polypeptide is a copolymer of three different amino acids each from a different one of three groups of the groups (a) to (d). These copolymers are herein referred to as terpolymers. In one embodiment, the terpolymer contains tyrosine (Y), alanine (A), and lysine (K), hereinafter designated YAK, in which the average molar fraction of the 25 amino acids can vary: Y, A and K can be present in a mole fraction of about 0.05-0.250, 0.3 - 0.6; and 0.1-0.5, respectively, more preferably, the molar ratios of Y, A and K are about 0.10:0.54:0.35, respectively. It is possible to substitute arginine for lysine, glycine for alanine, and/or tryptophan for tyrosine.

In another embodiment, the terpolymer contains tyrosine (Y), glutamic acid (E), and lysine (K), hereinafter designated YEK, in which the average molar fraction of the amino acids can vary: E, Y and K can be present in a mole fraction of about 0.005 - 0.300, 0.005-0.250, and 0.3-0.7, respectively. More preferably, the molar ratios of E, Y and K are about 0.26:0.16:0.58, respectively. It is possible to substitute aspartic acid for glutamic acid, arginine for lysine, and/or tryptophan for tyrosine.

In another embodiment, the terpolymer contains lysine (K), glutamic acid (E), and alanine (A), hereinafter designated KEA, in which the average molar fraction of the amino acids can vary: E, A and K can be present in a mole fraction of about 0.005-0.300, 0.005-0.600, and 0.2 - 0.7, respectively. More preferably, the molar ratios of E, A and K are about 0.15:0.48:0.36, respectively. It is possible to substitute aspartic acid for glutamic acid, glycine for alanine, and/or arginine for lysine.

In a further embodiment, the terpolymer contains tyrosine (Y), glutamic acid (E), and alanine (A0, hereinafter designated YEA, in which the average molar fraction of the amino acids can vary: Y, E and A can be present in a mole fraction of about 0.005-0.250, 0.005-0.300, and 0.005-0.800, respectively. More preferably, the molar ratios of E, A, and Y are about 0.21: 0.65:0.14, respectively. It is possible to substitute tryptophan for tyrosine, aspartic acid for glutamic acid, and/or glycine for alanine.

The average molecular weight of the terpolymers YAK, YEK, KEA and YEA can vary between about 2 KDa to 40 KDa, preferably between about 3 KDa to 35 KDa, more preferably between about 5 KDa to 25 KDa.

Copolymer 1 and related peptides and polypeptides may be prepared by methods known in the art, for example, by the process disclosed in U.S. Patent 3,849,550. The molecular weight of the copolymers can be adjusted during polypeptide synthesis or after the copolymers have been made. To adjust the molecular weight during polypeptide synthesis, the synthetic conditions or the amounts of amino acids are adjusted so that synthesis stops when the polypeptide reaches the approximate length that is desired. After synthesis, polypeptides with the desired molecular weight

can be obtained by any available size selection procedure, such as chromatography of the polypeptides on a molecular weight sizing column or gel, and collection of the molecular weight ranges desired. The copolymers can also be partially hydrolyzed to remove high molecular weight species, for example, by acid or enzymatic hydrolysis, 5 and then purified to remove the acid or enzymes.

In one embodiment, the copolymers with a desired molecular weight may be prepared by a process, which includes reacting a protected polypeptide with hydrobromic acid to form a trifluoroacetyl-polypeptide having the desired molecular weight profile. The reaction is performed for a time and at a temperature that is 10 predetermined by one or more test reactions. During the test reaction, the time and temperature are varied and the molecular weight range of a given batch of test polypeptides is determined. The test conditions that provide the optimal molecular weight range for that batch of polypeptides are used for the batch. Thus, a trifluoroacetyl-polypeptide having the desired molecular weight profile can be 15 produced by a process, which includes reacting the protected polypeptide with hydrobromic acid for a time and at a temperature predetermined by test reaction. The trifluoroacetyl-polypeptide with the desired molecular weight profile is then further treated with an aqueous piperidine solution to form a low toxicity polypeptide having the desired molecular weight. In one embodiment, a test sample of protected 20 polypeptide from a given batch is reacted with hydrobromic acid for about 10-50 hours at a temperature of about 20-28°C. The best conditions for that batch are determined by running several test reactions. For example, in one embodiment, the protected polypeptide is reacted with hydrobromic acid for about 17 hours at a temperature of about 26°C.

25 As binding motifs of Cop 1 to MS-associated HLA-DR molecules are known (Fridkis-Hareli et al, 1999), polypeptides derived from Cop 1 having a defined sequence can readily be prepared and tested for binding to the peptide binding groove of the HLA-DR molecules as described in the Fridkis-Hareli et al (1999) publication.

Examples of such peptides are those disclosed in WO 00/05249 and WO 00/05250, the entire contents of which are hereby incorporated herein by reference, and include the peptides of SEQ ID NOs. 1-32 hereinbelow.

SEQ ID NO:1	AAAYAAAAAKAAAA
SEQ ID NO:2	AEKYAAAAAKAAAA
SEQ ID NO:3	AKEYAAAAAKAAAA
SEQ ID NO:4	AKKYAAAAAKAAAA
SEQ ID NO:5	AEAYAAAAAKAAAA
SEQ ID NO:6	KEAYAAAAAKAAAA
SEQ ID NO:7	AEEYAAAAAKAAAA
SEQ ID NO:8	AAEYAAAAAKAAAA
SEQ ID NO:9	EKAYAAAAAKAAAA
SEQ ID NO:10	AAKYEA AAAKAAAA
SEQ ID NO:11	AAKYAEAAA AKAAAA
SEQ ID NO:12	EAAYAAAAAKAAAA
SEQ ID NO:13	EKKYAAAAAKAAAA
SEQ ID NO:14	EAKYAAAAAKAAAA
SEQ ID NO:15	AEKYAAAAAA AAAAAA
SEQ ID NO:16	AKEYAAAAAA AAAAAA
SEQ ID NO:17	AKKYEA AAAA AAAAAA
SEQ ID NO:18	AKKYAEAAA AAAAAA
SEQ ID NO:19	AEAYKAAA AAAAAA
SEQ ID NO:20	KEAYAAAAAA AAAAAA
SEQ ID NO:21	AEEYKAAA AAAAAA
SEQ ID NO:22	AAEYKAAA AAAAAA
SEQ ID NO:23	EKAYAAAAAA AAAAAA
SEQ ID NO:24	AAKYEA AAAA AAAAAA
SEQ ID NO:25	AAKYAEAAA AAAAAA
SEQ ID NO:26	EKKYAAAAAA AAAAAA
SEQ ID NO:27	EAKYAAAAAA AAAAAA
SEQ ID NO:28	AEYAKAAA AAAAAA
SEQ ID NO:29	AEKAYAAA AAAAAA
SEQ ID NO:30	EKYAAAAAA AAAAAA
SEQ ID NO:31	AYKAEEAAA AAAAAA
SEQ ID NO:32	AKYAEAAA AAAAAA

Such peptides of SEQ ID Nos: 1-32 and other similar peptides derived from Cop 1 would be expected to have similar activity as Cop 1. Such peptides, and other similar peptides, are also considered to be within the definition of Cop 1-related peptides or polypeptides and their use is encompassed by the present invention as well as other 5 synthetic amino acid copolymers such as the random four-amino acid copolymers described by Fridkis-Hareli et al., 2002 (as candidates for treatment of multiple sclerosis), namely copolymers (14-, 35- and 50-mers) containing the amino acids phenylalanine, glutamic acid, alanine and lysine (poly FEAK), or tyrosine, phenylalanine, alanine and lysine (poly YFAK), and any other similar copolymer to be 10 discovered that can be considered a universal antigen similar to Cop 1.

In another preferred embodiment of the present invention, the agent that causes T cells that produce IL-4 to accumulate in the eye is IL-4, preferably human recombinant IL-4, that can be administered nasally.

In a further embodiment, the agent is IL-4 activated dendritic cells, IL-4 15 activated monocytes, IL-4 activated bone marrow-derived myeloid cells or IL-4 activated peripheral blood mononuclear cells (PBMCs). In this regard, IL-13 have the same effect as IL-4, because it is well established in the field of cytokines that IL-4 and IL-13 can utilize a common receptor and share many actions such as B-cell activation and suppression of Th-1 cells.

20 In an additional embodiment, the agent is bone marrow-derived myeloid cells or peripheral blood-derived myeloid cells activated with IL-13 or with a narrow concentration range of IFN- $\gamma$ , more preferably up to 20 ng/ml IFN- $\gamma$ .

In another embodiment, the agent is a mammalian pathogenic self-antigen 25 associated with a T-cell-mediated specific autoimmune disease of the eye such as, but not limited to, a pathogenic uveitogenic antigen selected from mammalian interphotoreceptor retinoid-binding protein (IRBP), S-antigen (S-Ag), or rhodopsin. The mammalian uveitogenic antigen is preferably a human IRBP (SEQ ID NO: 33) or

a bovine IRBP (SEQ ID NO: 34), a human S-Ag (SEQ ID NO: 35) or a bovine S-Ag (SEQ ID NO: 36), or human rhodopsin (SEQ ID NO: 37).

In still another embodiment, the agent is a peptide which sequence is comprised within the sequence of said pathogenic self-antigen; a peptide obtained by modification of said peptide, which modification consists in the replacement of one or more amino acid residues of the peptide by different amino acid residues (hereinafter "modified peptide"), said modified peptide still being capable of recognizing the T-cell receptor recognized by the parent peptide but with less affinity; or a nucleotide sequence encoding said pathogenic self-antigen, said peptide or said modified peptide.

In one embodiment, the agent is: (a) a peptide which sequence is comprised within the sequence of bovine IRBP (SEQ ID NO: 34); (b) a modified peptide as defined above obtained by modification of the peptide of (a); or (c) a nucleotide sequence encoding human or bovine IRPB, a peptide of (a), or a modified peptide of (b).

The peptide which sequence is comprised within the sequence of bovine IRBP (SEQ ID NO: 34) may be the peptide R16 (sequence 1177–1191 of IRBP), ADGSSWEGVGVVPDV (SEQ ID NO:38); the peptide PTARSVGAADGSSWEGVGVVPDV (SEQ ID NO:39); or the peptide HVDDTDLYLTIPTARSVGAADGS (SEQ ID NO:40).

In another embodiment, the agent is: (a) a peptide which sequence is comprised within the sequence of bovine S-Ag (SEQ ID NO:36); (b) a modified peptide as defined above obtained by modification of the peptide of (a); or (c) a nucleotide sequence encoding human or bovine S-Ag, a peptide of (a), or a modified peptide of (b).

The peptide (a) which sequence is comprised within the sequence of bovine S-Ag may be the peptide G-8 (sequence 347–354 of S-Ag) of the sequence TSSEVATE (SEQ ID NO:41); the peptide M-8 (sequence 307–314 of S-Ag), DTNLASST (SEQ ID NO:42; or the peptides of the sequences:

DTNLASSTIIKEGIDKTV (SEQ ID NO:43);  
VPLLANNRRGIALDGKIKHE (SEQ ID NO:44);  
TSSEVATEVPFRLMHPQPED (SEQ ID NO:45);  
SLTKTLTVPLLANNRRG (SEQ ID NO:46);  
5 SLTRTLTLLPLLANNRERAG (SEQ ID NO:47);  
KEGIDKTVMGILVSYQIKVKL (SEQ ID NO:48); and  
KEGIDRTVLGILVSYQIKVKL (SEQ ID NO:49).

The modified peptide (c) derived from bovine S-Ag may be the G-8 analog, TSSEAATE (SEQ ID NO:50) or the M-8 analog, DTALASST (SEQ ID NO:51).

10 In another embodiment, the agent is a nucleotide sequence encoding a pathogenic self-antigen associated with a T-cell-mediated specific autoimmune disease of the eye, or a peptide or a modified peptide derived therefrom as defined herein.

15 In a further embodiment, the agent is T cells activated by Copolymer 1, or by a pathogenic self-antigen associated with a T-cell-mediated specific autoimmune disease of the eye, a peptide or a modified peptide derived therefrom as defined herein.

In yet a further embodiment, the agent is any combination of the agents defined above.

20 The invention further relates to the use of an agent selected from the agents defined herein for treatment of age-related macular degeneration or for the manufacture of a medicament for treatment of age-related macular degeneration.

When the agent is activated cells as described above, the cells can be preferably autologous or they can be from a matched donor and are preferably administered intravenously.

25 In one preferred embodiment, the agent is T cells activated by Copolymer 1, which can be prepared close to the administration step or cell banks can be established to store Copolymer 1-sensitized T cells for treatment of individuals at a later time, as needed. Autologous T cells may be obtained from the individual and allogeneic or semi-allogeneic T cells may be obtained from a bank of stored T cells of each of the most

common MHC-class II types are present. The patient is preferably treated with its autologous stored T cells, but if autologous T cells are not available, then cells should be used which share an MHC type II molecule with the patient, and these would be expected to be operable in that patient. The T cells are preferably stored in an activated 5 state after exposure to Copolymer 1. However, the T cells may also be stored in a resting state and activated with Copolymer 1 once they are thawed and prepared for use.

The T cell lines are prepared in any way that is well known in the art. The cell 10 lines of the bank are preferably cryopreserved. Once the cells are thawed, they are preferably cultured prior to injection in order to eliminate non-viable cells. During this culturing, the T cells can be activated or reactivated using the Copolymer 1 antigen as used in the original activation. Alternatively, activation may be achieved by culturing in the presence of a mitogen, such as phytohemagglutinin (PHA) or concanavalin A (preferably the former). This will place the cells into an even higher state of activation.

15 The bone marrow-derived myeloid cells for use in the present invention can be obtained from autologous or donor's peripheral blood or bone marrow and processed by techniques well-known in the art. The donor should be a matched, namely HLA-matched, donor. Once obtained, the myeloid cells may be cultured until they multiply to the level needed for transplant back into the patient and are then activated with the 20 cytokine (IL-4, IL-13 or IFN- $\gamma$ ) for the time necessary to upregulate CD11c expression. For example, activation with up to 20 ng/ml IFN- $\gamma$  may take 2-3 days until the peak of CD11c expression is reached.

In the examples, CD11c $^{+}$  microglia cells are described. Microglia are immune 25 cells of the CNS that are derived from myeloid progenitor cells, which come from the bone marrow. Thus, microglia are the resident CNS cells whereas the bone marrow-derived myeloid cells are the infiltrating cells.

Pharmaceutical compositions/medicaments for use in accordance with the present invention may be formulated in conventional manner using one or more

pharmaceutically/physiologically acceptable carriers or excipients, depending on the agent used. The carrier(s) must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

Methods of administration and doages will depend on the agent used and include, but are not limited to, parenteral, e.g., intravenous, intraperitoneal, intramuscular, subcutaneous, mucosal, e.g., oral, intranasal, buccal, vaginal, rectal, or intraocular, intrathecal, topical and intradermal routes, with or without adjuvant. Administration can be systemic or local.

The dosage of the agent to be administered will be determined by the physician according to the agent, the age of the patient and stage of the disease. For example, for Cop 1, the dosage may be chosen from a range of 1-80 mg, preferably 20 mg, although any other suitable dosage is encompassed by the invention. The treatment can be carried out by administration of repeated doses at suitable time intervals, according to the stage of the disease, the age and condition of the patient. In one embodiment, Cop 1 may be administered daily. In another preferred embodiment, the administration may be made according to a regimen suitable for immunization, for example, at least once a week, once a month or at least once every 2 or 3 months, or less frequently, but any other suitable interval between the immunizations is envisaged by the invention according to the condition of the patient.

When the agent is genetically engineered cells that produce IL-4, the cells are preferably engineered bone marrow-derived dendritic cells (DCs) that express IL-4, which may be obtained as described by Morita et al., 2001. Since DCs are specialized APCs that migrate from the periphery to lymphoid tissues, where they activate and regulate T cells, genetic modification of DCs to express immunoregulatory cytokines such as IL-4 provides a new immunotherapeutic strategy for treatment of AMD and other diseases.

Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated

element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

The invention will now be illustrated by the following non-limiting examples.

## EXAMPLES

### Materials and Methods

5 (i) *Animals.* Neonatal (P0-P1) mice, inbred adult male C57Bl/6J mice (8-10 weeks) were supplied by the Animal Breeding Center, Weizmann Institute of Science, Rehovot, Israel.

10 (ii) *Neural progenitor cell culture.* Coronal sections (2 mm thick) of tissue containing the subventricular zone of the lateral ventricle were obtained from the brains of adult C57Bl/6J mice. The tissue was minced and then incubated for digestion at 37°C, 5% CO<sub>2</sub> for 45 min in Earle's balanced salt solution containing 0.94 mg/ml papain (Worthington, Lakewood, NJ) and 0.18 mg/ml of L-cysteine and EDTA. After 15 centrifugation at 110 × g for 15 min at room temperature, the tissue was mechanically dissociated by pipette trituration. Cells obtained from single-cell suspensions were plated (3500 cells/cm<sup>2</sup>) in 75-cm<sup>2</sup> Falcon tissue-culture flasks (BD Biosciences, San Diego, CA), in neural stem/progenitor cell (NPC)-culturing medium [Dulbecco's modified Eagles's medium (DMEM)/F12 medium (Gibco/Invitrogen, Carlsbad, CA) containing 2 mM L-glutamine, 0.6% glucose, 9.6 µg/ml putrescine, 6.3 ng/ml progesterone, 5.2 ng/ml sodium selenite, 0.02 mg/ml insulin, 0.1 mg/ml transferrin, 2 µg/ml heparin (all from Sigma-Aldrich, Rehovot, Israel), fibroblast growth factor-2 20 (human recombinant, 20 ng/ml), and epidermal growth factor (human recombinant, 20 ng/ml; both from Peprotech, Rocky Hill, NJ)]. Spheres were passaged every 4-6 days and replated as single cells. Green fluorescent protein (GFP)-expressing NPCs were obtained as previously described (Pluchino et al., 2003).

25 (iii) *Primary microglial culture.* Brains from neonatal (P0-P1) C57Bl/6J mice were stripped of their meninges and minced with scissors under a dissecting microscope (Zeiss, Stemi DV4, Germany) in Leibovitz-15 medium (Biological Industries, Kibbutz Beit Ha-Emek, Israel). After trypsinization (0.5% trypsin, 10 min, 37°C/5% CO<sub>2</sub>), the tissue was triturated. The cell suspension was washed in culture

medium for glial cells [DMEM supplemented with 10% fetal calf serum (FCS; Sigma-Aldrich, Rehovot), L-glutamine (1 mM), sodium pyruvate (1 mM), penicillin (100 U/ml), and streptomycin (100 mg/ml)] and cultured at 37°C/5% CO<sub>2</sub> in 75-cm<sup>2</sup> Falcon tissue-culture flasks (BD Biosciences) coated with poly-D-lysine (PDL) (10 mg/ml; 5 Sigma-Aldrich, Rehovot) in borate buffer (2.37 g borax and 1.55 g boric acid dissolved in 500 ml sterile water, pH 8.4) for 1 h, then rinsed thoroughly with sterile, glass-distilled water. Half of the medium was changed after 6 h in culture and every 2<sup>nd</sup> day thereafter, starting on day 2, for a total culture time of 10–14 days. Microglia were shaken off the primary mixed brain glial cell cultures (150 rpm, 37°C, 6 h) with 10 maximum yields between days 10 and 14, seeded (10<sup>5</sup> cells/ml) onto PDL-pretreated 24-well plates (1 ml/well; Corning, New York, NY), and grown in culture medium for microglia [RPMI-1640 medium (Sigma-Aldrich, Rehovot) supplemented with 10% FCS, L-glutamine (1 mM), sodium pyruvate (1 mM), β-mercaptoethanol (50 mM), penicillin (100 U/ml), and streptomycin (100 mg/ml)]. The cells were allowed to 15 adhere to the surface of a PDL-coated culture flask (30 min, 37°C/5% CO<sub>2</sub>), and non-adherent cells were rinsed off.

(iv) *Immunocytochemistry and Immunohistochemistry.* Primary antibodies: *Bandeiraea simplicifolia* isolectin B4 (IB-4; 1:50; Sigma-Aldrich, Rehovot); mouse anti-β-tubulin (anti-βIII-T) isoform C-terminus antibodies (1:500; Chemicon, 20 Temecula, CA), rat anti-CD11b (MAC1; 1:50; BD-Pharmingen, Franklin Lakes, NJ), hamster anti-CD11c (1:100; eBioscience, San Diego, CA), rat anti-MHC-II Abs (clone IBL-5/22; 1:50), mouse anti-Aβ (human amino-acid residues 1–17; clone 6E10; Chemicon), rat anti-BrdU (1:200; Oxford Biotechnology, Kidlington, Oxfordshire, UK), goat anti-doublecortin (anti-DCX) (1:400; Santa Cruz Biotechnology, Santa Cruz, 25 CA), mouse anti-neuronal nuclear protein (NeuN; 1:200; Chemicon), goat anti-IGF-I Abs (1:20; R&D Systems), goat anti-TNF-α Abs (1:100; R&D Systems), rabbit anti-CD3 polyclonal Abs (1:100; DakoCytomation, CA). Secondary antibodies: FITC-

conjugated donkey anti-goat, Cy-3-conjugated donkey anti-mouse, and Cy-3- or Cy-5-conjugated donkey anti-rat, biotin-conjugated anti-hamster antibody and Cy-3- or Cy-5-conjugated streptavidin antibody (all from Jackson ImmunoResearch).

Cover slips from co-cultures of NPCs and mouse microglia were washed with  
5 PBS, fixed as described above, treated with a permeabilization/blocking solution containing 10% FCS, 2% bovine serum albumin, 1% glycine, and 0.1% Triton X-100 (Sigma-Aldrich, Rehovot), and stained with a combination of mouse anti- $\beta$ -tubulin (anti- $\beta$ III-T) isoform C-terminus antibodies (1:500; Chemicon, Temecula, CA), rat anti-CD11b (MAC1; 1:50; BD-Pharmingen, Franklin Lakes, NJ) and hamster anti-  
10 CD11c (1:100; eBioscience, San Diego, CA). To capture the microglia FITC- or Cy3-conjugated *Bandeiraea simplicifolia* isolectin B4 (IB-4; 1:50; Sigma-Aldrich, Rehovot) was used. To detect expression of human A $\beta$  anti-A $\beta$  (human amino-acid residues 1–17) (mouse, clone 6E10; Chemicon) was used.

For BrdU staining, sections were washed with PBS and incubated in 2N HCl at  
15 37°C for 30 min. Sections were blocked for 1 h with blocking solution [PBS containing 20% normal horse serum and 0.1% Triton X-100, or PBS containing mouse immunoglobulin blocking reagent obtained from Vector Laboratories (Burlingame, CA)].

For immunohistochemistry, tissue sections were treated with a  
20 permeabilization/blocking solution containing 10% FCS, 2% bovine serum albumin, 1% glycine, and 0.05% Triton X-100 (Sigma-Aldrich, St. Louis). Tissue sections were stained overnight at 4°C with specified combinations of the following primary antibodies: rat anti-BrdU (1:200; Oxford Biotechnology, Kidlington, Oxfordshire, UK), goat anti-doublecortin (anti-DCX) (1:400; Santa Cruz Biotechnology, Santa Cruz, CA),  
25 and mouse anti-neuronal nuclear protein (anti-NeuN) (1:200; Chemicon). Secondary antibodies were FITC-conjugated donkey anti-goat, Cy-3-conjugated donkey anti-mouse, and Cy-3- or Cy-5-conjugated donkey anti-rat (1:200; Jackson

ImmunoResearch, West Grove, PA). CD11b (MAC1; 1:50; BD-Pharmingen) or FITC-conjugated IB-4 was used for labeling of microglia. Anti-MHC-II Abs (rat, clone IBL-5/22; 1:50) was used to detect expression of cell-surface MHC-II proteins. To detect expression of CD11c hamster anti-CD11c (1:100; eBioscience, San Diego, CA) was 5 used. Anti-A $\beta$  (human amino-acid residues 1–17) (mouse, clone 6E10; Chemicon) was used to detect expression of human A $\beta$ . Expression of IGF-I was detected by goat anti-IGF-I Abs (1:20; R&D Systems). Expression of TNF- $\alpha$  was detected by goat anti-TNF- $\alpha$  Abs (1:100; R&D Systems). T cells were detected with anti-CD3 polyclonal 10 Abs (rabbit, 1:100; DakoCytomation, CA). Propidium iodide (1  $\mu$ g/ml; Molecular Probes, Invitrogen, Carlsbad, CA), was used for nuclear staining.

Control sections (not treated with primary antibody) were used to distinguish specific staining from staining of nonspecific antibodies or autofluorescent components. Sections were then washed with PBS and cover-slipped in polyvinyl alcohol with diazabicyclo-octane as anti-fading agent.

15 **(v) Transgenic mice.** Nineteen adult double-transgenic APP<sub>K595N, M596L</sub> + PS1<sub>ΔE9</sub> mice of the B6C3-Tg (APPswe, PSEN1dE9) 85Dbo/J strain (Borchelt et al., 1997) were purchased from The Jackson Laboratory (Bar Harbor, ME) and were bred and maintained in the Animal Breeding Center of The Weizmann Institute of Science. All animals were handled according to the regulations formulated by the Weizmann 20 Institute's Animal Care and Use Committee, and all experiments and procedures were approved by the Weizmann Institute's Animal Care and Use Committee.

**(vi) Genotyping.** All mice used in this experiment were genotyped for the presence of the transgenes by PCR as previously described (Jankowsky et al., 2004).

25 **(vii) Reagents.** Recombinant mouse IFN- $\gamma$  and IL-4 were obtained from R&D Systems (Minneapolis, MN).  $\beta$ -amyloid peptide [fragment 1–40 (A $\beta$ <sub>1–40</sub>)] was purchased from Sigma-Aldrich, St. Louis, MO. The A $\beta$  peptide was dissolved in

endotoxin-free water, and A $\beta$  aggregates were formed by incubation of A $\beta$ , as described (Butovsky et al., 2005).

5 *(viii) Copolymer-1 vaccination.* Each mouse was subcutaneously injected five times with a total of 100  $\mu$ g of high-molecular-weight (TV-5010 DS, from batch no. 486220205; Teva Pharmaceutical Industries, Petach Tikva, Israel) emulsified in 200  $\mu$ l PBS, from experimental day 0 until day 24, twice during the first week and once a week thereafter.

10 *(ix) Behavioral testing.* Spatial learning/memory was assessed by performance on a hippocampus-dependent visuo-spatial learning task in the Morris water maze (MWM) and carried out as described (Lichtenwalner et al., 2001).

15 *(x) Administration of 5-bromo-2'-deoxyuridine and tissue preparation.* The cell-proliferation marker 5-bromo-2'-deoxyuridine (BrdU) was dissolved by sonication in phosphate-buffered saline (PBS) and injected intraperitoneally (i.p.) into each mouse (50 mg/kg body weight; 1.25 mg BrdU in 200  $\mu$ l PBS). Starting from experimental day 22 after the first Cop-1 vaccination, BrdU was injected i.p. twice daily, every 12 h for 2.5 days, to label proliferating cells. Three weeks after the first BrdU injection the mice were deeply anesthetized and perfused transcardially, first with PBS and then with 4% paraformaldehyde. The whole brain was removed, postfixed overnight, and then equilibrated in phosphate-buffered 30% sucrose. Free-floating 30- $\mu$ m sections were 20 collected on a freezing microtome (Leica SM2000R) and stored at 4°C prior to immunohistochemistry.

25 *(xi) Co-culturing of neural progenitor cells and microglia.* Cultures of treated or untreated microglia were washed twice with fresh NPC-differentiation medium (same as the culture medium for NPCs but without growth factors except for 0.02 mg/ml insulin and with 2.5% FCS) to remove all traces of the tested reagents, then incubated on ice for 15 min and shaken at 350 rpm for 20 min at room temperature. Microglia were removed from the flasks and immediately co-cultured ( $5 \times 10^4$

cells/well) with NPCs ( $5 \times 10^4$  cells/well) for 10 days on cover slips coated with Matrigel™ (BD Biosciences) in 24-well plates, in the presence of NPC-differentiation medium. The cultures were then fixed with 2.5% paraformaldehyde in PBS for 30 min at room temperature and stained for neuronal and glial markers.

5 **(xii) Quantification and stereological counting procedure.** A Zeiss LSM 510 confocal laser scanning microscope ( $\times 40$  magnification) was used for microscopic analysis. For experiments *in vitro* fields of  $0.053 \text{ mm}^2$  ( $n = 8-16$  from at least two different cover slips) were scanned for each experimental group. For each marker, 500–1000 cells were sampled. Cells co-expressing GFP and βIII-T were counted.

10 For *in-vivo* experiments, the numbers of A $\beta$  plaques and CD11b $^+$  microglia in the hippocampus were counted at 300- $\mu\text{m}$  intervals in 6–8 coronal sections (30  $\mu\text{m}$ ) from each mouse. Neurogenesis in the dentate gyrus was evaluated by counting of premature neurons (DCX $^+$ ), proliferating cells (BrdU $^+$ ), and newly formed mature neurons (BrdU $^+$ /NeuN $^+$ ) in six coronal sections (370  $\mu\text{m}$  apart) per mouse brain. To obtain an 15 estimate of the total number of labeled cells per dentate gyrus, the total number of cells counted in the selected coronal sections from each brain was multiplied by the volume index (the ratio between the volume of the dentate gyrus and the total combined volume of the selected sections). Specificity of BrdU+/NeuN+ co-expression was assayed using the confocal microscope (LSM 510) in optical sections at 1- $\mu\text{m}$  intervals.

20 Quantification of CD3 $^+$ , CD11b $^+$  and CD11c $^+$  cells were analyzed from 30–50 A $\beta$ -plaques of each mouse tested in this study. Cell counts, numbers of A $\beta$  plaques, plaque areas, and intensity of NeuN staining per unit area in the dentate gyrus were evaluated automatically using Image-Pro Plus 4.5 software (Media Cybernetics, Carlsbad, CA).

25 **(xiii) Statistical analysis.** MWM behavior scores were analyzed using 3-way ANOVA. Treatment group and trial block were used as sources of variation to evaluate the significance of differences between mean scores during acquisition trial blocks in the MWM. When the *P*-value obtained was significant, a pairwise Fisher's least-

significant-difference multiple comparison test was run to determine which groups were significantly different.

The *in-vitro* results were analyzed by two-tailed unpaired Student's *t*-test and by the Tukey–Kramer multiple comparisons test (ANOVA) and are expressed as means ±

5 SEM. Results *in vivo* were analyzed by two-tailed unpaired Student's *t*-test or 1-way ANOVA and are expressed as means ± SEM.

**Example 1. Aggregated  $\beta$ -amyloid induces microglia to express a phenotype that blocks neurogenesis, and the blocking is counteracted by IL-4.**

10 Previous in vitro findings from our laboratory have suggested that the microglia found in association with inflammatory and neurodegenerative diseases (e.g. microglia activated by LPS or by aggregated  $A\beta_{(1-40)}$ ) have an impaired ability to present antigen, whereas IL-4-activated microglia, shown to be associated with neural tissue survival, express MHC-II, produce IGF-I, and decrease TNF- $\alpha$  expression (Butovsky et al.,

15 2005). Here we first examined whether  $A\beta$ -activated microglia block neurogenesis, and if so, whether T cell-derived cytokines can counteract the inhibitory effect. To this end we co-cultured green fluorescent protein (GFP)-expressing neural stem/progenitor cells (NPCs) with microglia that had been pre-incubated for 48 h in their optimal growth

medium (Butovsky et al., 2005) in the presence or absence of the aggregated  $A\beta$  peptide 1–40 ( $A\beta_{(1-40)}$ ; 5  $\mu$ M) and subsequently treated for an additional 48 h with

20 IFN- $\gamma$  (10 ng/ml) or IL-4 (10 ng/ml) or IL-4 together with IFN- $\gamma$  (10 ng/ml). The choice of  $A\beta_{(1-40)}$  rather than  $A\beta_{(1-42)}$  and its concentration was based on our previous demonstration that this compound induces cytotoxic activity in microglia (Butovsky et al., 2005). Growth media and cytokine residues were then washed off the co-cultured

25 microglia, and each of the treated microglial preparations was freshly co-cultured with dissociated adult subventricular zone-derived NPC spheres (Butovsky et al., 2006a) on coverslips coated with Matrigel™ in the presence of differentiation medium (Butovsky

et al., 2006a) (**Fig. 1A**). Expression of GFP by NPCs confirmed that any differentiating neurons seen in the cultures were derived from the NPCs rather than from contamination of the primary microglial culture. After 10 days, we could discern a few GFP-positive NPCs expressing the neuronal marker  $\beta$ III-T in microglia-free cultures 5 (control). In co-cultures of NPCs with microglia previously activated by incubation with IFN- $\gamma$  (10 ng/ml; MG<sub>(IFN- $\gamma$ )</sub>) a dramatic increase in numbers of GFP<sup>+</sup>/ $\beta$ III-T<sup>+</sup> cells was seen. On the contrary, microglia activated by aggregated A $\beta$ <sub>(1-40)</sub> (MG<sub>(A $\beta$ 1-40)</sub>) 10 blocked neurogenesis and decreased the number of NPCs. The addition of IFN- $\gamma$  to A $\beta$ -activated microglia (MG<sub>(A $\beta$ 1-40/IFN- $\gamma$ )</sub>), failed to reverse their negative effect on 15 neurogenesis. In contrast, the addition of IL-4 (10 ng/ml) to microglia pretreated with aggregated A $\beta$ <sub>(1-40)</sub> (MG<sub>(A $\beta$ 1-40/IL-4)</sub>) partially counteracted the adverse effect of the aggregated A $\beta$  on NPCs survival and differentiation, with the result that these 20 microglia were able to induce NPCs to differentiate into neurons. However, when IFN- $\gamma$  was added in combination with IL-4 (MG<sub>(A $\beta$ 1-40/IFN- $\gamma$ +IL-4)</sub>), their effect in 15 counteracting the negative activity of the A $\beta$ -activated microglia on NPC survival and differentiation was stronger than the effect of IL-4 alone (**Fig. 1B**). We verified that in all cases the  $\beta$ III-T<sup>+</sup> cells also expressed GFP (**Fig. 1C**). This finding is particularly interesting in view of our earlier demonstration that the order in which threatening 25 stimuli are presented to the microglia critically affects the ability of these cells to withstand them (Butovsky et al., 2005). The quantitative analysis presented in **Fig. 1D** summarizes the data shown in **Fig. 1B**, and in addition shows that differentiation in the presence of untreated microglia occurred only to a small extent. Notably, no  $\beta$ III-T<sup>+</sup> cells were seen in microglia cultured without NPCs (Butovsky et al., 2006a).

**Example 2. T-cell-based vaccination with copolymer-1 modulates immune activity of microglia, eliminates  $\beta$ -amyloid plaque formation, and induces neurogenesis.**

The above findings prompted us to examine whether a T cell-based vaccination would alter the default microglial phenotype in Alzheimer's disease and hence lead to plaque removal and neurogenesis. The antigen we chose for the vaccination was Cop-1 (Teitelbaum et al., 1996), shown by us to be weakly cross-reactive with a wide-range of CNS autoantigens and, depending on the regimen, to be neuroprotective under conditions of both acute and chronic neurodegeneration (Kipnis et al., 2000; Schori et al., 2001; Angelov et al., 2003). We examined the effect of Cop-1 in Tg-AD mice, suffering from learning/memory impairment and an accumulation of aggregated A $\beta$  plaques deposited mainly in the cortex and the hippocampus, both characteristic features of early-onset familial Alzheimer's disease (Borchelt et al., 1997). The regimen for Cop-1 administration was similar to that used to evoke neuroprotection in a model of chronic elevation of intraocular pressure (Bakalash et al., 2005).

We verified the presence of both transgenes in each mouse by PCR amplification of genomic DNA. Tg-AD mice aged approximately 8 months were then vaccinated subcutaneously with Cop-1 ( $n = 6$ ) twice during the first week and once a week thereafter. Age-matched untreated Tg-AD mice ( $n = 7$ ) and non-Tg littermates ( $n = 6$ ) served, respectively, as untreated-Tg and wild-type controls. Seven weeks after the first Cop-1 injection all the mice were euthanized and analyzed. Staining of brain cryosections from Tg-AD mice with antibodies specific to human A $\beta$  disclosed numerous plaques in the untreated-Tg-AD mice but very few in the Tg-AD mice vaccinated with Cop-1 (Fig. 2A). No plaques were seen in their respective non-Tg littermates (Fig. 2A).

The above results, coupled with the *in-vitro* findings, prompted us to look for changes in microglial features in the vaccinated Tg-AD mice. Plaques in the untreated-Tg-AD mice were found to be associated with the abundant appearance of CD11b $^{+}$  microglia (Fig. 2A and Fig 2B) expressing TNF- $\alpha$  (Fig. 2C). Fewer CD11b $^{+}$  microglia

were detectable in the Cop-1-vaccinated Tg-AD mice (**Fig. 2A**). It is important to note that the CD11b<sup>+</sup> microglia in the untreated-Tg-AD mice showed relatively few ramified processes **Fig. 2C**). Staining with anti-MHC-II antibodies disclosed that in the Cop-1-vaccinated Tg-AD mice most of the microglia adjacent to residual A $\beta$  plaques 5 expressed MHC-II, and hardly any of them expressed TNF- $\alpha$  (**Fig. 2D**), whereas in the untreated-Tg-AD mice hardly any microglia expressed MHC-II (**Fig. 2E**), suggesting that their ability to function as APCs is limited. All of the MHC-II<sup>+</sup> cells were co-labeled with IB-4 (data not shown), verifying their identification as microglia. The 10 dendritic-like morphology (**Fig. 2D**) of the MHC-II<sup>+</sup> microglia seen in the Cop-1-vaccinated Tg-AD mice encouraged us to examine whether they express the characteristic marker of dendritic cells, namely CD11c. CD11c<sup>+</sup> microglia in untreated 15 Tg-AD mice were only rarely found in association with A $\beta$ <sup>+</sup> plaques, whereas any residual A $\beta$ -stained plaques seen in the Cop-1-vaccinated mice were surrounded by MHC-II<sup>+</sup>/CD11c<sup>+</sup> microglia (**Fig. 2E**). Notably, these CD11c<sup>+</sup> microglia were also 20 positively stained for CD11b (**Fig. 3A**); in addition, they were loaded with A $\beta$ , indicative of their engulfment of this peptide (**Fig. 2E**). Quantitative analysis revealed that the number of CD11b<sup>+</sup> cells associated with A $\beta$ -plaque significantly decreased in the Cop-1-vaccinated Tg-AD mice (**Fig. 3B**), and that as a result of the vaccination 87 % of the CD11b<sup>+</sup> became CD11b<sup>+</sup>/CD11c<sup>+</sup> cells relative to 25 % in the untreated Tg- 25 AD mice (**Fig. 3C**).

In view of our recent finding that MHC-II<sup>+</sup> microglia (which are activated by IL-4) abundantly express IGF-I (Butovsky et al., 2005; Butovsky et al., 2006a;b), we examined IGF-I expression in the vaccinated Tg-AD mice. MHC-II<sup>+</sup> microglia in these mice were indeed found to express IGF-I (**Fig. 2F**). Staining for the presence of T cells, 25 identified by anti-CD3 antibodies, revealed that unlike in the untreated-Tg-AD mice, in the Cop-1-vaccinated Tg-AD mice there were numerous T cells associated with A $\beta$ -plaques (**Fig. 2G**). Moreover, most of the T cells in the Cop-1 vaccinated Tg-AD mice

were found to be located close to MHC-II<sup>+</sup> microglia. Any A $\beta$ -immunoreactivity detected in those mice appeared to be associated with the MHC-II<sup>+</sup> microglia, suggesting the occurrence of an immune synapse between these microglia and CD3<sup>+</sup> T cells (Fig. 2H). Quantitative analysis confirmed the presence of significantly fewer 5 plaques in the Cop-1-vaccinated Tg-AD mice than in the untreated-Tg-AD mice (Fig. 2I), and showed that the area occupied by the plaques was significantly smaller in the vaccinated Tg-AD mice than in their age-matched untreated counterparts (Fig. 2J). In addition, significantly fewer CD11b<sup>+</sup> microglia (Fig. 2K) and significantly more T 10 cells associated with A $\beta$ -plaque were observed in the Cop-1-vaccinated Tg-AD mice than in the corresponding groups of untreated-Tg-AD mice (Fig. 2L).

On the basis of our previous findings, we suspected that the switch from a CD11b<sup>+</sup>/CD11c<sup>-</sup>/IGF-1<sup>-</sup> to a CD11b<sup>+</sup>/CD11c<sup>+</sup>/IGF-1<sup>+</sup> microglial phenotype in the Cop-1 vaccinated Tg-AD mice might be attributable to IL-4. We examined this possibility *in vitro*. Staining of 5-day microglia cultures with the CD11c marker showed that 15 CD11c was hardly expressed at all by untreated, but was abundantly expressed by microglia activated by IL-4 (Fig. 4A). Moreover, IL-4, even if only added 3 days after the microglia were exposed to A $\beta$ , was able to induce them to express CD11c (Fig. 4B). Differential activation of the microglia was also reflected in morphological differences: microglia activated by A $\beta$  exhibited amoeboid morphology, whereas the 20 rounded shape of the CD11c<sup>+</sup> microglia was reminiscent of dendritic cells (Fig. 4B). Most importantly, the amoeboid morphology of the A $\beta$ -stained microglia was reversible on addition of IL-4, when they again took on the morphological appearance of dendritic-like cells (Fig. 4B). The various treatments applied to the microglia did not affect their expression of CD11b, suggesting that they did not lose their CD11b 25 characteristics when they took on the expression of CD11c (Fig. 4B). Quantitative analysis of CD11c expression, assessed by the number of CD11c<sup>+</sup> cells and the intensity of their staining as a function of time in culture, revealed that soon after seeding (day 0) untreated microglia expressed low levels of CD11c, which gradually

disappeared (**Fig. 4C**). In contrast, the expression of CD11c induced by IL-4 was not transient. Quantification of the ability of IL-4 to induce CD11c expression even after the microglia were pretreated with A $\beta$  is shown in **Fig. 4D**.

5 The correlation between the phenotype that was found to be induced by the Cop-1 vaccination and the IL-4 effect on microglia *in vitro* prompted us to examine the ability of IL-4-activated microglia to phagocytize aggregated A $\beta$ <sub>(1-40)</sub>. Quantitative comparison (by intracellular staining) of immunoreactive A $\beta$  engulfed by IL-4-treated and untreated microglia indicated that IL-4 did not interfere with the ability of microglia to engulf A $\beta$  (**Fig. 5**).

10 The observed effects of IL-4 on the expression of CD11c, MHC-II, and TNF- $\alpha$  prompted us to examine whether the Cop-1-vaccinated Tg-AD mice would show increased neurogenesis *in vivo*. Three weeks before tissue excision all mice had been injected with the proliferating-cell marker BrdU, making it possible to detect new neurons. Quantitative analysis of additional sections from the same areas of the 15 hippocampal dentate gyrus disclosed significantly more BrdU $^+$  cells in the Cop-1-vaccinated Tg-AD mice (**Fig. 6A**) than in their untreated-Tg counterparts. In addition, compared to the numbers of newly formed mature neurons (BrdU $^+$ /NeuN $^+$ ) in their respective non-Tg littermates the numbers were significantly lower in the untreated Tg-AD group, but were similar in the Cop-1-vaccinated Tg-AD group; indicating that 20 the neurogenesis capacity had been at least partially restored by the Cop-1 vaccination (**Fig. 6B**). Analysis of corresponding sections for DCX, a useful marker for analyzing the absolute number of newly generated pre-mature neurons in the adult dentate gyrus (Rao et al., 2004), disclosed that relative to the non-Tg littermates there were significantly fewer DCX $^+$  cells in the dentate gyri of untreated-Tg-AD mice, and 25 slightly but significantly more in the dentate gyri of Tg-AD mice vaccinated with Cop-1 (**Fig. 6C**). Confocal micrographs illustrate the differences in the numbers of BrdU $^+$ /NeuN $^+$  cells, and in the numbers of DCX $^+$  cells and their dendritic processes, between non-Tg littermates, untreated-Tg-AD mice, and Cop-1-vaccinated Tg-AD

mice (**Fig. 6D**). The results showed that neurogenesis was indeed significantly more abundant in the Cop-1-treated Tg-AD mice than in the untreated-Tg-AD mice. Interestingly, however, in both untreated and Cop-1-vaccinated Tg-AD mice the processes of the DCX-stained neurons in the subgranular zone of the dentate gyrus 5 were short, except in those Cop-1-vaccinated Tg-AD mice in which the DCX<sup>+</sup> cells were located adjacent to MHC-II<sup>+</sup> microglia (**Fig. 6E**).

## Discussion

We showed here that vaccination of Tg-AD mice with Cop-1 reduced plaque 10 formation and attenuated cognitive decline. Labeling of activated microglia with anti-CD11b antibodies disclosed that staining was heavy in the untreated-Tg-AD mice and significantly less intense in the age-matched Cop-1-vaccinated Tg-AD mice. The decrease in numbers of CD11b<sup>+</sup> microglia in Cop-1-vaccinated mice could be an outcome of a reduction in A $\beta$ -plaques, whose deposition in the brain had led to the 15 microglial activation in the first place.

The role of microglia in Alzheimer's disease and other neurodegenerative diseases has been intensively investigated over the last few years, with apparently conflicting results (Streit, 2004). The detection of some CD11b<sup>+</sup> microglia in aged wild-type mice that are healthy is in line with the reported age-related increase in 20 activated microglia in the normal human brain (Streit, 2004). It is possible that such microglia are the ones that contribute both to age-related cognitive loss and to impaired neurogenesis (Monje et al., 2003). CD11b were found also in patients with Alzheimer's disease (Akiyama & McGeer, 1990). Although these microglia are phagocytic (Frenkel et al., 2005), they are apparently not efficient enough to fight off the 25 Alzheimer symptoms. In contrast, and in line with our present study, microglia derived from the bone marrow of matched wild-type mice can effectively remove plaques (Simard et al., 2006). These microglia express higher protein levels than are required for antigen presentation and might therefore be more effective phagocytes than the

resident microglia. On the basis of our present results, we suggest that the microglia that are needed to support brain maintenance and fight Alzheimer's disease are dendritic-like cells (CD11b<sup>+</sup>/CD11c<sup>+</sup>). The CD11c<sup>+</sup> microglia might engage in a dialog with T cells that can help to fight off adverse conditions by promoting the buffering of 5 excessive A $\beta$  and supporting both neuronal survival (Butovsky et al., 2005) and neural renewal (Butovsky et al., 2006b). In view of our earlier finding that IL-4, but not IFN- $\gamma$ , can alter the phenotype of A $\beta$ -committed microglia (Butovsky et al., 2005), it seems likely that the MHC-II<sup>+</sup> microglia found adjacent to A $\beta$  plaques in the present study were activated by IL-4. This likelihood is further supported by two of the *in-vitro* 10 findings of the present study: first, that the A $\beta$ -induced blockage of neurogenesis was partially counteracted by IL-4, either alone or in combination with IFN- $\gamma$ , but not by IFN- $\gamma$  alone; and secondly, that the MHC-II<sup>+</sup>/CD11c<sup>+</sup> microglia which were seen in close proximity to the residual A $\beta$  plaques in the Cop-1-vaccinated Tg-AD mice also 15 expressed IGF-I. Taken together, these findings reinforce the contention that the terms 'beneficial' and 'harmful' cannot be applied in a generalized way to microglial activity (Schwartz et al., 2006).

IL-4 has often been described as an anti-inflammatory cytokine (Chao et al., 1993). Our results strongly argue against this perception and suggest instead that IL-4 20 activates microglia to adopt a phenotype that seems to acquire a different morphology and a different activity from those of the innately activated microglia or of the activated microglia commonly seen in Alzheimer's disease or in MS. In the latter disease, unlike in Alzheimer's disease, the microglia appear to be overwhelmed by an onslaught of adaptive immunity (Butovsky et al., 2006a). Interestingly, it seems that IL-4 is capable of restoring a favorable activated phenotype even after the microglia have already 25 exhibited phenotypic characteristics of aggregated A $\beta$  ((Butovsky et al., 2005), and the present study), or been overwhelmed by IFN- $\gamma$  (Butovsky et al., 2006a). Another interesting finding is that LPS and A $\beta$  exhibit similar patterns of MAPK activation in

microglia (Pyo et al., 1998). IL-4 can attenuate a MAPK pathway activated by LPS, an effect evidently associated with serine/threonine phosphatase activity (Iribarren et al., 2003). The latter phenomenon might indeed serve as a molecular mechanism underlying the present finding that IL-4 attenuates the detrimental effect of A $\beta$ -activated microglia. Accordingly, we suggest that the activity of IL-4 should not be regarded as anti-inflammatory, but as immunomodulatory.

Based on our *in-vitro* findings in connection with the effect of IL-4 on microglial phenotype, and the ability of the regimen chosen in the present study to evoke a T-cell response with an IL-4 bias, we suggest that the observed beneficial 10 effect of Cop-1 vaccination on the Tg-AD mice in this study was a result of the evoked T-cell effect on their microglial phenotype.

**Example 3. Copolymer-1 vaccination counteracts cognitive decline in Alzheimer's disease.**

15 Two weeks before the end of the experiment, all mice were tested in a Morris water maze (MWM) for cognitive activity, as reflected by their performance of a hippocampus-dependent spatial learning/memory task. The MWM performance of the untreated-Tg-AD mice was significantly worse, on average, than that of their age-matched non-Tg littermates (Figs. 7A-7B). However, the performance of Cop-1-vaccinated Tg-AD mice was superior to that of the untreated-Tg-AD mice and did not 20 differ significantly from that of the non-Tg-AD mice, suggesting that the Cop-1 vaccination had prevented further cognitive loss. Differences in cognitive performance were manifested in both the acquisition (Fig. 7A) and the reversal tasks (Fig. 7B).

25 The vaccinated mice in this study demonstrated attenuated cognitive loss (tested in a Morris water maze) and increased neurogenesis. These two aspects of hippocampal plasticity are apparently related to the presence of IGF-I and cognitive activity and cell renewal (Butovsky et al., 2006a;b). Reported observations in Tg-AD mice housed in an

enriched environment also support a link between mechanisms associated with neurogenesis (Ziv et al., 2006) and with plaque reduction (Lazarov et al., 2005).

Because aggregated A $\beta$  evidently interferes with the ability of microglia to engage in dialog with T cells, its presence in the brain can be expected to cause loss of cognitive ability and impairment of neurogenesis. Homing of CNS-autoreactive T cells to the site of disease or damage in such cases is critical, but will be effective only if those T cells can counterbalance the destructive activity of the aggregated A $\beta$ . As shown here, IFN- $\gamma$  by itself is impotent against the activity of microglia that are already committed to an aggregated A $\beta$  phenotype, but is effective when added together with IL-4. Thus, the results of this study strongly suggest that the occurrence of neurogenesis in the adult hippocampus depends on well controlled local immune activity associated with microglial production of growth factors such as IGF-I and BDNF (Ziv et al., 2006). In line with this notion is the reported finding that neurogenesis is impaired in animals treated with LPS (Monje et al., 2003), shown to impair microglial production of IGF-I and induce microglial secretion of TNF- $\alpha$  (Butovsky et al., 2005).

**Example 4. IFN- $\gamma$ -activated myeloid cells and their uses for promoting tissue repair, detection of and delivery of drugs to damaged tissue.**

We have shown that when IFN- $\gamma$  was added in combination with IL-4 their effect in counteracting the negative activity of A $\beta$ -activated microglia on NPC survival and differentiation was stronger than the effect of IL-4 alone (Fig. 1B). However, our data demonstrates that IFN- $\gamma$  alone, similarly to IL-4 but transiently, induces CD11c if the cells are treated with IFN- $\gamma$  in a narrow concentration range of up to 20 ng/ml, induces neuroprotection (Butovsky et al., 2005) and is able to support neurogenesis from neural stem/progenitor cells (Butovsky et al., 2006b). CD11c is upregulated and reach the peak at 2-3 days after IFN- $\gamma$  activation in microglia (Figs. 8A-8D) or bone-

marrow-derived myeloid cells (data not shown). Thus, upregulation of CD11c on bone marrow derived myeloid cells by IFN- $\gamma$  will allow the cells to reach injured sites and induce beneficial effect.

5 **Example 5. Weekly vaccination with Copaxone as a potential therapy for dry Age-related Macular Degeneration**

The results of the previous examples showed that vaccination of Tg-AD mice with Cop-1 in a regimen previously found to lead to neuroprotection resulted in a microglial phenotype switch from CD11b $^+$  to CD11b $^+$ /CD11c $^+$ , and that this was 10 correlated with plaque removal, neurogenesis, and attenuated cognitive loss. The beneficial effect was attributed to a phenotype switch of the infiltrating microglia as well as to recruitment of blood-borne monocytes.

Based on these findings and on the many features common between Alzheimer's disease and AMD, we hypothesized that in AMD, similar to Alzheimer's 15 disease, weekly Cop-1 treatment would likely lead to clearing of drusen and this, subsequently, may restrain the progression of dry to wet AMD. It should be emphasized that Cop-1 given in different regimens results in different therapies; daily treatment has no beneficial effects in paradigms of neurodegenerative diseases, but is effective in suppressing inflammatory disease in patients like multiple sclerosis. 20 Therefore, we decided to embark on a clinical trial with AMD patients in a protocol of weekly administration.

The natural fate of drusen in our population of patients was examined by analysis of fundus photographs of unenrolled and untreated dry AMD patients. These patients comprised an observational group.

25 The effect of Cop-1 (Copaxone®, Teva) on drusen was examined during a prospective, pilot, randomized, double-masked, placebo-controlled, comparative trial. Patients over 50 years of age with intermediate dry AMD in both eyes were enrolled.

Enrolled patients were randomly (ratio 2 to 1) treated either with subcutaneous injections of Copaxone (20 mg) or placebo injections on a weekly basis for a period of 12 weeks. Complete eye examination, along with fundus photography, was performed at baseline, 6- and 12-week visits. The primary outcome was the change in total drusen area and it was calculated using Image-Pro software. Analysis of patients from the observational group (17 eyes) showed an increase of 25.2% in total drusen area.

At 12 week, 8 eyes treated with Copaxone showed a reduction in total drusen area from baseline of 53.6% (range, 5-89%). Two eyes receiving placebo injections demonstrated reduction of 0.6% in total drusen area.

This study thus shows a potential beneficial effect for Copaxone for treating AMD. Due to the limited numbers of the enrolled patients, no statistical significant conclusion can be drawn. Yet, the results justify and encourage to extend and continue this study for fighting off drusens. In light of the underlying mechanism of weekly treatment with Copaxone in animal models of neurodegenerative diseases, its effect on AMD might be beyond the drusens' elimination in protecting neurons and promoting repair.

## REFERENCES

Akiyama, H., Barger, S., Barnum, S., Bradt, B., Bauer, J., Cole, GM., Cooper, NR, Eikelenboom, P., Emmerling, M., Fiebich, BL., Finch, CE, Frautschy, S., Griffin, WS., Hampel, H., Hull, M. *et al.* Inflammation and Alzheimer's disease.

5 *Neurobiol Aging* **21**, 383-421 (2000).

Akiyama, H & McGeer, PL. Brain microglia constitutively express beta-2 integrins. *J Neuroimmunol* **30**, 81-93 (1990).

Ambati, J. et al. An animal model of age-related macular degeneration in senescent Ccl-2- or Ccr-2-deficient mice. *Nat Med* **9**, 1390-7 (2003).

10 Angelov, DN, Waibel, S, Guntinas-Lichius, O, Lenzen, M, Neiss, WF, Tomov, TL, Yoles, E, Kipnis, J, Schori, H, Reuter, A, Ludolph A, and Schwartz, M. Therapeutic vaccine for acute and chronic motor neuron diseases: Implications for amyotrophic lateral sclerosis. *Proc Natl Acad Sci U S A* **100**, 4790-5 (2003).

Avidan, H, Kipnis, J, Butovsky, O, Caspi, RR & Schwartz, M. Vaccination 15 with autoantigen protects against aggregated beta-amyloid and glutamate toxicity by controlling microglia: effect of CD4+CD25+ T cells. *Eur J Immunol* **34**, 3434-45 (2004).

Bakalash, S, Shlomo, GB, Aloni, E, Shaked, I, Wheeler, L, Ofri, R & Schwartz, M. T cell-based vaccination for morphological and functional 20 neuroprotection in a rat model of chronically elevated intraocular pressure. *J Mol Med* **83**, 904-16 (2005).

Bakalash, S, Kessler, A, Mizrahi, T, Nussenblatt, R & Schwartz, M. Antigenic specificity of immunoprotective therapeutic vaccination for glaucoma. *Invest. Ophthalmol. Vis. Sci.* **44**, 3374-81 (2003).

25 Bard, F, Cannon, C, Barbour, R, Burke, RL, Games, D, Grajeda, H, Guido, T, Hu, K, Huang, J, Johnson-Wood, K. *et al.* Peripherally administered antibodies against amyloid beta-peptide enter the central nervous system and reduce pathology in a mouse model of Alzheimer disease. *Nat Med* **6**, 916-9 (2000).

Barouch, R. & Schwartz, M. Autoreactive T cells induce neurotrophin production by immune and neural cells in injured rat optic nerve: implications for protective autoimmunity. *Faseb J* 16, 1304-6 (2002).

5 Borchelt, DR, Ratovitski, T, van Lare, J, Lee, MK, Gonzales, V, Jenkins, NA, Copeland, NG, Price, DL & Sisodia, SS. Accelerated amyloid deposition in the brains of transgenic mice co-expressing mutant presenilin 1 and amyloid precursor proteins. *Neuron* 19, 939-45 (1997).

10 Butovsky, O., M. Koronyo-Hamaoui, G. Kunis, E. Ophir, G. Landa, H. Cohen & M. Schwartz. Glatiramer acetate fights against Alzheimer's disease by inducing dendritic-like microglia expressing insulin-like growth factor-1. 2006 Jul 24 [Epub ahead of print] (2006a)

15 Butovsky, O, Ziv, Y, Schwartz, A, Landa, G, Talpalar, AE, Pluchino, S, Martino, G & Schwartz, M. Microglia activated by IL-4 or IFN-gamma differentially induce neurogenesis and oligodendrogenesis from adult stem/progenitor cells. *Mol Cell Neurosci* 31, 149-60 (2006b).

Butovsky, O, Landa, G, Kunis, G, Ziv, Y, Avidan, H, Greenberg, N, Schwartz, A, Smirnov, I, Pollack, A, Jung, S & Schwartz, M. Induction and blockage of oligodendrogenesis by differently activated microglia in an animal model of multiple sclerosis. *J Clin Invest* 116 :905-15 (2006c).

20 Butovsky, O, Talpalar, AE, Ben-Yaakov, K & Schwartz, M. Activation of microglia by aggregated beta-amyloid or lipopolysaccharide impairs MHC-II expression and renders them cytotoxic whereas IFN-gamma and IL-4 render them protective. *Mol Cell Neurosci* 29, 381-93 (2005).

25 Butovsky, O, Hauben, E & Schwartz, M. Morphological aspects of spinal cord autoimmune neuroprotection: colocalization of T cells with B7-2 (CD86) and prevention of cyst formation. *Faseb J.* 15, 1065-7 (2001).

Chao, CC, Molitor, TW & Hu, S. Neuroprotective role of IL-4 against activated microglia. *J Immunol* 151, 1473-81 (1993).

Frenkel, D, Maron, R, Burt, DS & Weiner, HL. Nasal vaccination with a proteosome-based adjuvant and glatiramer acetate clears beta-amyloid in a mouse model of Alzheimer disease. *J Clin Invest* 115, 2423-2433 (2005).

5 Furlan, R. et al. Vaccination with amyloid-beta peptide induces autoimmune encephalomyelitis in C57/BL6 mice. *Brain* 126, 285-91 (2003).

Gupta, N, Brown, KE & Milam, AH. Activated microglia in human retinitis pigmentosa, late-onset retinal degeneration, and age-related macular degeneration. *Exp Eye Res* 76, 463-71 (2003).

10 Hageman, GS. & Mullins, RF. Molecular composition of drusen as related to substructural phenotype. *Mol Vis* 5, 28 (1999).

Hageman, GS. et al. An integrated hypothesis that considers drusen as biomarkers of immune-mediated processes at the RPE-Bruch's membrane interface in aging and age-related macular degeneration. *Prog Retin Eye Res* 20, 705-32 (2001).

15 Hardy, J. & Selkoe, DJ. The amyloid hypothesis of Alzheimer's disease: progress and problems on the road to therapeutics. *Science* 297, 353-6 (2002).

Haughey, NJ, Nath, A, Chan, SL, Borchard, AC, Rao, MS & Mattson, MP. Disruption of neurogenesis by amyloid beta-peptide, and perturbed neural progenitor cell homeostasis, in models of Alzheimer's. *J Neurochem* 83, 1509-24 (2002).

Holz, FG. et al. Bilateral macular drusen in age-related macular degeneration. Prognosis and risk factors. *Ophthalmology* 101, 1522-8 (1994).

Iribarren, P, Cui, YH, Le, Y, Ying, G, Zhang, X, Gong, W & Wang, JM. *J Immunol* 171, 5482-8 (2003).

25 Jankowsky, J. L., Fadale, D. J., Anderson, J., Xu, G. M., Gonzales, V., Jenkins, N. A., Copeland, N. G., Lee, M. K., Younkin, L. H., Wagner, S. L. et al. Mutant presenilins specifically elevate the levels of the 42 residue beta-amyloid peptide in vivo: evidence for augmentation of a 42-specific gamma secretase. *Hum Mol Genet* 13, 159-70 (2004).

Jin, K, Galvan, V, Xie, L, Mao, XO, Gorostiza, OF, Bredesen, DE & Greenberg, DA. Enhanced neurogenesis in Alzheimer's disease transgenic (PDGF-APPsw,Ind) mice. *Proc Natl Acad Sci USA* 101, 13363-7 (2004a).

5 Jin, K, Peel, AL, Mao, XO, Xie, L, Cottrell, BA, Henshall, DC & Greenberg, DA. Increased hippocampal neurogenesis in Alzheimer's disease. *Proc Natl Acad Sci USA* 101, 343-7 (2004b).

10 Kerschensteiner, M, Gallmeier, E, Behrens, L, Leal, VV, Misgeld, T, Klinkert, WE, Kolbeck, R, Hoppe, E, Oropesa-Wekerle, RL, Bartke, I. *et al.* Activated human T cells, B cells, and monocytes produce brain-derived neurotrophic factor in vitro and in inflammatory brain lesions: a neuroprotective role of inflammation? *J. Exp. Med.* 189, 865-70 (1999).

15 Kipnis, J. & Schwartz, M. Dual action of glatiramer acetate (Cop-1) in the treatment of CNS autoimmune and neurodegenerative disorders. *Trends Mol Med* 8, 319-23 (2002).

Kipnis, J, Avidan, H, Caspi, RR & Schwartz, M. Dual effect of CD4+CD25+ regulatory T cells in neurodegeneration: Pro- and anti-inflammatory cytokines determine microglial activity. *Proc. Natl. Acad. Sci. U S A* 101 Suppl 2, 14663-14669 (2004a).

20 Kipnis, J., Cohen, H., Cardon, M., Ziv, Y. & Schwartz, M. (2004b) *Proc Natl Acad Sci USA* 101, 8180-5.

Kipnis, J, Mizrahi, T, Hauben, E, Shaked, I, Shevach, E & Schwartz, M. Neuroprotective autoimmunity: naturally occurring CD4+CD25+ regulatory T cells suppress the ability to withstand injury to the central nervous system. *Proc Natl Acad Sci USA* 99, 15620-5 (2002).

25 Kipnis, J, Yoles, E, Porat, Z, Cohen, A, Mor, F, Sela, M, Cohen, IR & Schwartz, M. T cell immunity to copolymer 1 confers neuroprotection on the damaged optic nerve: possible therapy for optic neuropathies. *Proc. Natl. Acad. Sci. USA* 97, 7446-51 (2000).

30 Klaver, C.C. *et al.* Is age-related maculopathy associated with Alzheimer's Disease? The Rotterdam Study. *Am J Epidemiol* 150, 963-8 (1999).

Klein R, Klein BE, Linton KL. Prevalence of age-related maculopathy. The Beaver Dam Eye Study. *Ophthalmology* 99:933–43 (1992).

Kliffen, M., de Jong, PT & Luider, TM. Protein analysis of human maculae in relation to age-related maculopathy. *Lab Invest* 73, 267-72 (1995).

5 Lazarov, O., Robinson, J., Tang, Y. P., Hairston, I. S., Korade-Mirnics, Z., Lee, V. M., Hersh, L. B., Sapolisky, R. M., Mirnics, K. & Sisodia, S. S. *Cell* 120, 701-13 (2005).

Lichtenwalner, R. J., Forbes, M. E., Bennett, S. A., Lynch, C. D., Sonntag, W. E. & Riddle, D. R. (2001) *Neuroscience* 107, 603-13.

10 Luibl, V. et al. Drusen deposits associated with aging and age-related macular degeneration contain nonfibrillar amyloid oligomers. *J Clin Invest* 116, 378-85 (2006).

15 Mitchell P, Smith W, Attebo K, Wang JJ. Prevalence of age-related maculopathy in Australia. The Blue Mountains Eye Study. *Ophthalmology* 102:1450–60 (1995).

Mizrahi, T, Hauben, E & Schwartz, M. The tissue-specific self-pathogen is the protective self-antigen: the case of uveitis. *J. Immunol.* 169, 5971-7 (2002).

Moalem, G. et al. Production of neurotrophins by activated T cells: implications for neuroprotective autoimmunity. *J. Autoimmun.* 15, 331-45 (2000).

20 Moalem, G, Leibowitz-Amit, R, Yoles, E, Mor, F, Cohen, IR & Schwartz, M. Autoimmune T cells protect neurons from secondary degeneration after central nervous system axotomy. *Nat. Med.* 5, 49-55 (1999).

Monje, ML, Toda, H. & Palmer, TD. Inflammatory blockade restores adult hippocampal neurogenesis. *Science* 302, 1760-5 (2003).

25 Monsonego, A, Imitola, J, Petrovic, S, Zota, V, Nemirovsky, A, Baron, R, Fisher, Y, Owens, T & Weiner, H. L. A $\beta$ -induced meningoencephalitis is IFN- $\gamma$ -dependent and is associated with T cell-dependent clearance of A $\beta$  in a mouse model of Alzheimer's disease. *Proc Natl Acad Sci U S A*, 103:5048-53 (2006).

Morgan, D. et al. A beta peptide vaccination prevents memory loss in an animal model of Alzheimer's disease. *Nature* 408, 982-5 (2000).

5 Morita Y, Yang J, Gupta R, Shimizu K, Shelden EA, Endres J, Mulé JJ, McDonagh KT, Fox DA. Dendritic cells genetically engineered to express IL-4 inhibit murine collagen-induced arthritis. *J Clin Invest* 107(10):1275-84.(2001)

Mullins, RF, Russell, SR, Anderson, DH & Hageman, GS. Drusen associated with aging and age-related macular degeneration contain proteins common to extracellular deposits associated with atherosclerosis, elastosis, amyloidosis, and dense deposit disease. *Faseb J* 14, 835-46 (2000).

10 Nimmerjahn, A, Kirchhoff, F & Helmchen, F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 308, 1314-8 (2005).

15 Penfold, PL, Madigan, MC & Provis, JM. Antibodies to human leucocyte antigens indicate subpopulations of microglia in human retina. *Vis Neurosci* 7, 383-8 (1991).

Pluchino, S, Quattrini, A, Brambilla, E, Gritti, A, Salani, G, Dina, G, Galli, R, Del Carro, U, Amadio, S, Bergami, A. et al. Injection of adult neurospheres induces recovery in a chronic model of multiple sclerosis. *Nature* 422, 688-94 (2003).

20 Provis, JM. Development of the primate retinal vasculature. *Prog Retin Eye Res* 20, 799-821 (2001).

Pyo, H., Jou, I., Jung, S., Hong, S. & Joe, E H *Neuroreport* 9, 871-4 (1998).

Rao, M. S. & Shetty, A. K. *Eur J Neurosci* 19, 234-46 (2004).

25 Roque, RS, Rosales, AA, Jingjing, L, Agarwal, N & Al-Ubaidi, MR. Retina-derived microglial cells induce photoreceptor cell death in vitro. *Brain Res* 836, 110-9 (1999).

Schori, H, Kipnis, J, Yoles, E, WoldeMussie, E, Ruiz, G, Wheeler, LA & Schwartz, M. Vaccination for protection of retinal ganglion cells against death from glutamate cytotoxicity and ocular hypertension: implications for glaucoma. *Proc. 30 Natl. Acad. Sci. USA* 98, 3398-403 (2001).

Schwartz, M, Butovsky, O, Bruck, W & Hanisch, UK. Microglial phenotype: is the commitment reversible? *Trends Neurosci* 29, 68-74 (2006).

Selkoe, D J. The molecular pathology of Alzheimer's disease. *Neuron* 6, 487-98 (1991).

5 Shaked, I, Porat, Z, Gersner, R, Kipnis, J. & Schwartz, M. Early activation of microglia as antigen-presenting cells correlates with T cell-mediated protection and repair of the injured central nervous system. *J Neuroimmunol* 146, 84-93 (2004).

Shaked, I, Tchoresh, D, Gersner, R, Meiri, G, Mordechai, S, Xiao, X, Hart, RP. & Schwartz, M. Protective autoimmunity: interferon-gamma enables microglia 10 to remove glutamate without evoking inflammatory mediators. *J Neurochem* 92, 997-1009 (2005).

Simard, AR, Soulet, D, Gowing, G, Julien, JP & Rivest, S. Bone marrow-derived microglia play a critical role in restricting senile plaque formation in Alzheimer's disease. *Neuron* 49, 489-502 (2006).

15 Streilein, JW, Wilbanks, GA & Cousins, SW. Immunoregulatory mechanisms of the eye. *J Neuroimmunol* 39, 185-200 (1992).

Streit, WJ. Microglia and Alzheimer's disease pathogenesis. *J Neurosci Res* 77, 1-8 (2004).

Teitelbaum, D., Fridkis-Hareli, M., Arnon, R. & Sela, M. *J Neuroimmunol* 20 64, 209-17 (1996).

20 Yin, Y. et al. Macrophage-derived factors stimulate optic nerve regeneration. *J Neurosci*. 23, 2284-93 (2003).

Yoles, E, Hauben, E, Palgi, O, Agranov, E, Gothilf, A, Cohen, A, Kuchroo, V, Cohen, IR, Weiner, H & Schwartz, M. Protective autoimmunity is a 25 physiological response to CNS trauma. *J Neurosci* 21, 3740-8 (2001).

Yoshida, T. et al. The potential role of amyloid beta in the pathogenesis of age-related macular degeneration. *J Clin Invest* 115, 2793-800 (2005).

Ziemssen, T, Kumpfel, T, Klinkert, WE, Neuhaus, O & Hohlfeld, R. Glatiramer acetate-specific T-helper 1- and 2-type cell lines produce BDNF:

implications for multiple sclerosis therapy. Brain-derived neurotrophic factor. *Brain* 125, 2381-91 (2002).

5 Ziv, Y, Ron, N, Butovsky, O, Landa, G, Sudai, E, Greenberg, N, Cohen, H, Kipnis, J & Schwartz, M. Immune cells contribute to the maintenance of neurogenesis and spatial learning abilities in adulthood. *Nat Neurosci* 9, 268-75 (2006).

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:-

1. A method for treatment of age-related macular degeneration, which comprises causing T cells that produce IL-4 to accumulate in the eye of an individual in need by administering a single active agent selected from the group consisting of Copolymer 1 or T cells activated against Copolymer 1 to said individual, thereby halting or delaying progress of the macular degeneration.

2. The method according to claim 1, wherein said agent is Copolymer 1.

3. The method according to claim 1, wherein said agent is T cells activated by Copolymer 1.

4. Use of a single active agent that causes T cells that produce IL-4 to accumulate in the eye, wherein the single active agent is selected from the group consisting of Copolymer 1 or T cells activated against Copolymer 1, for the preparation of a medicament for treatment of age-related macular degeneration.

5. The use according to claim 4, wherein said agent is Copolymer 1.

6. The use according to claim 4, wherein said agent is T cells activated by Copolymer 1.

1/15




Fig. 1A




Fig. 1B

SUBSTITUTE SHEET (RULE 26)

2/15




Fig. 1C




Fig. 1D

3/15




Fig. 2A




Fig. 2B






Fig. 2C

4/15

**Fig. 2D****Fig. 2E**

SUBSTITUTE SHEET (RULE 26)

5/15




Fig. 2F

Fig. 2G




Fig. 2H

6/15




Fig. 2I




Fig. 2J




Fig. 2K




Fig. 2L

7/15




Fig. 3A




Fig. 3B




Fig. 3C

8/15




Fig. 4A




Fig. 4B

9/15




Fig. 4C




Fig. 4D

10/15




Fig. 5A




Fig. 5B

11/15




Fig. 6A






Fig. 6B

**Fig. 6C****Fig. 6E**

13/15




Fig. 7A




Fig. 7B

14/15




Fig. 8A




Fig. 8B



Fig. 8C

15/15




Fig. 8D

YEDA-068 PCT 27.06.07 MS.ST25  
SEQUENCE LISTING

<110> YEDA RESEARCH AND DEVELOPMENT CO. LTD  
Eisenbach-Schwartz, Michal  
Butovsky, Oleg

<120> METHOD OF TREATMENT OF AGE-RELATED MACULAR DEGENERATION

<130> YEDA-068 PCT

<150> US 60/806,041  
<151> 2006-06-28

<160> 51

<170> PatentIn version 3.3

<210> 1  
<211> 15  
<212> PRT  
<213> Artificial

<220>

<223> .

<400> 1

Ala Ala Ala Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 2  
<211> 15  
<212> PRT  
<213> Artificial

<220>

<223> .

<400> 2

Ala Glu Lys Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 3  
<211> 15  
<212> PRT  
<213> Artificial

<220>

<223> .

<400> 3

Ala Lys Glu Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 4  
<211> 15  
<212> PRT  
<213> Artificial

<220>

<223> .

<400> 4

Ala Lys Lys Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 5  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 5

Ala Glu Ala Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 6  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 6

Lys Glu Ala Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 7  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 7

Ala Glu Glu Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 8  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 8

Ala Ala Glu Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 9  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 9

Glu Lys Ala Tyr Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 10

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 10

Ala Ala Lys Tyr Glu Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 11

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 11

Ala Ala Lys Tyr Ala Glu Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 12

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 12

Glu Ala Ala Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 13

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 13

Glu Lys Lys Tyr Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

<210> 14

<211> 15

<212> PRT

<213> Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 14

Glu Ala Lys Tyr Ala Ala Ala Ala Ala Ala Lys Ala Ala Ala Ala  
1 5 10 15

&lt;210&gt; 15

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 15

Ala Glu Lys Tyr Ala  
1 5 10 15

&lt;210&gt; 16

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 16

Ala Lys Glu Tyr Ala Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

&lt;210&gt; 17

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 17

Ala Lys Lys Tyr Glu Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

&lt;210&gt; 18

&lt;211&gt; 15

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 18

Ala Lys Lys Tyr Ala Glu Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

&lt;210&gt; 19

&lt;211&gt; 15

&lt;212&gt; PRT

<213> Artificial

<220>

<223> .

<400> 19

Ala Glu Ala Tyr Lys Ala 15  
1 5 10 15

<210> 20

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 20

Lys Glu Ala Tyr Ala 15  
1 5 10 15

<210> 21

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 21

Ala Glu Glu Tyr Lys Ala Ala Ala Ala Ala Ala Ala Ala 15  
1 5 10 15

<210> 22

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 22

Ala Ala Glu Tyr Lys Ala Ala Ala Ala Ala Ala Ala Ala 15  
1 5 10 15

<210> 23

<211> 15

<212> PRT

<213> Artificial

<220>

<223> .

<400> 23

Glu Lys Ala Tyr Ala Ala Ala Ala Ala Ala Ala Ala 15  
1 5 10 15

<210> 24

<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 24

Ala Ala Lys Tyr Glu Ala Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

<210> 25  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 25

Ala Ala Lys Tyr Ala Glu Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

<210> 26  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 26

Glu Lys Lys Tyr Ala Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

<210> 27  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 27

Glu Ala Lys Tyr Ala Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

<210> 28  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 28

Ala Glu Tyr Ala Lys Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

YEDA-068 PCT 27.06.07 MS.ST25

<210> 29  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 29

Ala Glu Lys Ala Tyr Ala  
1 5 10 15

<210> 30  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 30

Glu Lys Tyr Ala  
1 5 10 15

<210> 31  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 31

Ala Tyr Lys Ala Glu Ala Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

<210> 32  
<211> 15  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 32

Ala Lys Tyr Ala Glu Ala Ala Ala Ala Ala Ala Ala Ala Ala  
1 5 10 15

<210> 33  
<211> 1247  
<212> PRT  
<213> Homo sapiens

<400> 33

Met Met Arg Glu Trp Val Leu Leu Met Ser Val Leu Leu Cys Gly Leu  
1 5 10 15

YEDA-068 PCT 27.06.07 MS.ST25

Ala Gly Pro Thr His Leu Phe Gln Pro Ser Leu Val Leu Asp Met Ala  
 20 25 30

Lys Val Leu Leu Asp Asn Tyr Cys Phe Pro Glu Asn Leu Leu Gly Met  
 35 40 45

Gln Glu Ala Ile Gln Gln Ala Ile Lys Ser His Glu Ile Leu Ser Ile  
 50 55 60

Ser Asp Pro Gln Thr Leu Ala Ser Val Leu Thr Ala Gly Val Gln Ser  
 65 70 75 80

Ser Leu Asn Asp Pro Arg Leu Val Ile Ser Tyr Glu Pro Ser Thr Pro  
 85 90 95

Glu Pro Pro Pro Gln Val Pro Ala Leu Thr Ser Leu Ser Glu Glu Glu  
 100 105 110

Leu Leu Ala Trp Leu Gln Arg Gly Leu Arg His Glu Val Leu Glu Gly  
 115 120 125

Asn Val Gly Tyr Leu Arg Val Asp Ser Val Pro Gly Gln Glu Val Leu  
 130 135 140

Ser Met Met Gly Glu Phe Leu Val Ala His Val Trp Gly Asn Leu Met  
 145 150 155 160

Gly Thr Ser Ala Leu Val Leu Asp Leu Arg His Cys Thr Gly Gly Gln  
 165 170 175

Val Ser Gly Ile Pro Tyr Ile Ile Ser Tyr Leu His Pro Gly Asn Thr  
 180 185 190

Ile Leu His Val Asp Thr Ile Tyr Asn Arg Pro Ser Asn Thr Thr Thr  
 195 200 205

Glu Ile Trp Thr Leu Pro Gln Val Leu Gly Glu Arg Tyr Gly Ala Asp  
 210 215 220

Lys Asp Val Val Val Leu Thr Ser Ser Gln Thr Arg Gly Val Ala Glu  
 225 230 235 240

Asp Ile Ala His Ile Leu Lys Gln Met Arg Arg Ala Ile Val Val Gly  
 245 250 255

Glu Arg Thr Gly Gly Ala Leu Asp Leu Arg Lys Leu Arg Ile Gly  
 260 265 270

Glu Ser Asp Phe Phe Phe Thr Val Pro Val Ser Arg Ser Leu Gly Pro  
 275 280 285

YEDA-068 PCT 27.06.07 MS-ST25  
 Leu Gly Gly Gly Ser Gln Thr Trp Glu Gly Ser Gly Val Leu Pro Cys  
 290 295 300  
 Val Gly Thr Pro Ala Glu Gln Ala Leu Glu Lys Ala Leu Ala Ile Leu  
 305 310 315 320  
 Thr Leu Arg Ser Ala Leu Pro Gly Val Val His Cys Leu Gln Glu Val  
 325 330 335  
 Leu Lys Asp Tyr Tyr Thr Leu Val Asp Arg Val Pro Thr Leu Leu Gln  
 340 345 350  
 His Leu Ala Ser Met Asp Phe Ser Thr Val Val Ser Glu Glu Asp Leu  
 355 360 365  
 Val Thr Lys Leu Asn Ala Gly Leu Gln Ala Ala Ser Glu Asp Pro Arg  
 370 375 380  
 Leu Leu Val Arg Ala Ile Gly Pro Thr Glu Thr Pro Ser Trp Pro Ala  
 385 390 395 400  
 Pro Asp Ala Ala Ala Glu Asp Ser Pro Gly Val Ala Pro Glu Leu Pro  
 405 410 415  
 Glu Asp Glu Ala Ile Arg Gln Ala Leu Val Asp Ser Val Phe Gln Val  
 420 425 430  
 Ser Val Leu Pro Gly Asn Val Gly Tyr Leu Arg Phe Asp Ser Phe Ala  
 435 440 445  
 Asp Ala Ser Val Leu Gly Val Leu Ala Pro Tyr Val Leu Arg Gln Val  
 450 455 460  
 Trp Glu Pro Leu Gln Asp Thr Glu His Leu Ile Met Asp Leu Arg His  
 465 470 475 480  
 Asn Pro Gly Gly Pro Ser Ser Ala Val Pro Leu Leu Leu Ser Tyr Phe  
 485 490 495  
 Gln Gly Pro Glu Ala Gly Pro Val His Leu Phe Thr Thr Tyr Asp Arg  
 500 505 510  
 Arg Thr Asn Ile Thr Gln Glu His Phe Ser His Met Glu Leu Pro Gly  
 515 520 525  
 Pro Arg Tyr Ser Thr Gln Arg Gly Val Tyr Leu Leu Thr Ser His Arg  
 530 535 540  
 Thr Ala Thr Ala Ala Glu Glu Phe Ala Phe Leu Met Gln Ser Leu Gly  
 545 550 555 560

YEDA-068 PCT 27.06.07 MS.ST25  
 Trp Ala Thr Leu Val Gly Glu Ile Thr Ala Gly Asn Leu Leu His Thr  
 565 570 575  
  
 Arg Thr Val Pro Leu Leu Asp Thr Pro Glu Gly Ser Leu Ala Leu Thr  
 580 585 590  
  
 Val Pro Val Leu Thr Phe Ile Asp Asn His Gly Glu Ala Trp Leu Gly  
 595 600 605  
  
 Gly Gly Val Val Pro Asp Ala Ile Val Leu Ala Glu Glu Ala Leu Asp  
 610 615 620  
  
 Lys Ala Gln Glu Val Leu Glu Phe His Gln Ser Leu Gly Ala Leu Val  
 625 630 635 640  
  
 Glu Gly Thr Gly His Leu Leu Glu Ala His Tyr Ala Arg Pro Glu Val  
 645 650 655  
  
 Val Gly Gln Thr Ser Ala Leu Leu Arg Ala Lys Leu Ala Gln Gly Ala  
 660 665 670  
  
 Tyr Arg Thr Ala Val Asp Leu Glu Ser Leu Ala Ser Gln Leu Thr Ala  
 675 680 685  
  
 Asp Leu Gln Glu Val Ser Gly Asp His Arg Leu Leu Val Phe His Ser  
 690 695 700  
  
 Pro Gly Glu Leu Val Val Glu Glu Ala Pro Pro Pro Pro Ala Val  
 705 710 715 720  
  
 Pro Ser Pro Glu Glu Leu Thr Tyr Leu Ile Glu Ala Leu Phe Lys Thr  
 725 730 735  
  
 Glu Val Leu Pro Gly Gln Leu Gly Tyr Leu Arg Phe Asp Ala Met Ala  
 740 745 750  
  
 Glu Leu Glu Thr Val Lys Ala Val Gly Pro Gln Leu Val Arg Leu Val  
 755 760 765  
  
 Trp Gln Gln Leu Val Asp Thr Ala Ala Leu Val Ile Asp Leu Arg Tyr  
 770 775 780  
  
 Asn Pro Gly Ser Tyr Ser Thr Ala Ile Pro Leu Leu Cys Ser Tyr Phe  
 785 790 795 800  
  
 Phe Glu Ala Glu Pro Arg Gln His Leu Tyr Ser Val Phe Asp Arg Ala  
 805 810 815  
  
 Thr Ser Lys Val Thr Glu Val Trp Thr Leu Pro Gln Val Ala Gly Gln  
 820 825 830

YEDA-068 PCT 27.06.07 MS.ST25

Arg Tyr Gly Ser His Lys Asp Leu Tyr Ile Leu Met Ser His Thr Ser  
 835 840 845

Gly Ser Ala Ala Glu Ala Phe Ala His Thr Met Gln Asp Leu Gln Arg  
 850 855 860

Ala Thr Val Ile Gly Glu Pro Thr Ala Gly Gly Ala Leu Ser Val Gly  
 865 870 875 880

Ile Tyr Gln Val Gly Ser Ser Pro Leu Tyr Ala Ser Met Pro Thr Gln  
 885 890 895

Met Ala Met Ser Ala Thr Thr Gly Lys Ala Trp Asp Leu Ala Gly Val  
 900 905 910

Glu Pro Asp Ile Thr Val Pro Met Ser Glu Ala Leu Ser Ile Ala Gln  
 915 920 925

Asp Ile Val Ala Leu Arg Ala Lys Val Pro Thr Val Leu Gln Thr Ala  
 930 935 940

Gly Lys Leu Val Ala Asp Asn Tyr Ala Ser Ala Glu Leu Gly Ala Lys  
 945 950 955 960

Met Ala Thr Lys Leu Ser Gly Leu Gln Ser Arg Tyr Ser Arg Val Thr  
 965 970 975

Ser Glu Val Ala Leu Ala Glu Ile Leu Gly Ala Asp Leu Gln Met Leu  
 980 985 990

Ser Gly Asp Pro His Leu Lys Ala Ala His Ile Pro Glu Asn Ala Lys  
 995 1000 1005

Asp Arg Ile Pro Gly Ile Val Pro Met Gln Ile Pro Ser Pro Glu  
 1010 1015 1020

Val Phe Glu Glu Leu Ile Lys Phe Ser Phe His Thr Asn Val Leu  
 1025 1030 1035

Glu Asp Asn Ile Gly Tyr Leu Arg Phe Asp Met Phe Gly Asp Gly  
 1040 1045 1050

Glu Leu Leu Thr Gln Val Ser Arg Leu Leu Val Glu His Ile Trp  
 1055 1060 1065

Lys Lys Ile Met His Thr Asp Ala Met Ile Ile Asp Met Arg Phe  
 1070 1075 1080

Asn Ile Gly Gly Pro Thr Ser Ser Ile Pro Ile Leu Cys Ser Tyr  
 1085 1090 1095

YEDA-068 PCT 27.06.07 MS.ST25

Phe Phe Asp Glu Gly Pro Pro Val Leu Leu Asp Lys Ile Tyr Ser  
 1100 1105 1110

Arg Pro Asp Asp Ser Val Ser Glu Leu Trp Thr His Ala Gln Val  
 1115 1120 1125

Val Gly Glu Arg Tyr Gly Ser Lys Lys Ser Met Val Ile Leu Thr  
 1130 1135 1140

Ser Ser Val Thr Ala Gly Thr Ala Glu Glu Phe Thr Tyr Ile Met  
 1145 1150 1155

Lys Arg Leu Gly Arg Ala Leu Val Ile Gly Glu Val Thr Ser Gly  
 1160 1165 1170

Gly Cys Gln Pro Pro Gln Thr Tyr His Val Asp Asp Thr Asn Leu  
 1175 1180 1185

Tyr Leu Thr Ile Pro Thr Ala Arg Ser Val Gly Ala Ser Asp Gly  
 1190 1195 1200

Ser Ser Trp Glu Gly Val Gly Val Thr Pro His Val Val Val Pro  
 1205 1210 1215

Ala Glu Glu Ala Leu Ala Arg Ala Lys Glu Met Leu Gln His Asn  
 1220 1225 1230

Gln Leu Arg Val Lys Arg Ser Pro Gly Leu Gln Asp His Leu  
 1235 1240 1245

<210> 34

<211> 551

<212> PRT

<213> Bos taurus

<400> 34

Val Leu Pro Gly Gln Leu Gly Tyr Leu Arg Phe Asp Ala Met Ala Glu  
 1 5 10 15

Leu Glu Thr Val Lys Ala Val Gly Pro Gln Leu Val Gln Leu Val Trp  
 20 25 30

Gln Lys Leu Val Asp Thr Ala Ala Leu Val Val Asp Leu Arg Tyr Asn  
 35 40 45

Pro Gly Ser Tyr Ser Thr Ala Val Pro Leu Leu Cys Ser Tyr Phe Phe  
 50 55 60

Glu Ala Glu Pro Arg Arg His Leu Tyr Ser Val Phe Asp Arg Ala Thr  
 65 70 75 80

Ser Arg Val Thr Glu Val Trp Thr Leu Pro His Val Thr Gly Gln Arg  
 Page 12

85 YEDA-068 PCT 27.06.07 MS.ST25  
 90 95

Tyr Gly Ser His Lys Asp Leu Tyr Val Leu Val Ser His Thr Ser Gly  
 100 105 110

Ser Ala Ala Glu Ala Phe Ala His Thr Met Gln Asp Leu Gln Arg Ala  
 115 120 125

Thr Ile Ile Gly Glu Pro Thr Ala Gly Gly Ala Leu Ser Val Gly Ile  
 130 135 140

Tyr Gln Val Gly Ser Ser Ala Leu Tyr Ala Ser Met Pro Thr Gln Met  
 145 150 155 160

Ala Met Ser Ala Ser Thr Gly Glu Ala Trp Asp Leu Ala Gly Val Glu  
 165 170 175

Pro Asp Ile Thr Val Pro Met Ser Val Ala Leu Ser Thr Ala Arg Asp  
 180 185 190

Ile Val Thr Leu Arg Ala Lys Val Pro Thr Val Leu Gln Thr Ala Gly  
 195 200 205

Lys Leu Val Ala Asp Asn Tyr Ala Ser Pro Glu Leu Gly Val Lys Met  
 210 215 220

Ala Ala Glu Leu Ser Gly Leu Gln Ser Arg Tyr Ala Arg Val Thr Ser  
 225 230 235 240

Glu Ala Ala Leu Ala Glu Leu Leu Gln Ala Asp Leu Gln Val Leu Ser  
 245 250 255

Gly Asp Pro His Leu Lys Thr Ala His Ile Pro Glu Asp Ala Lys Asp  
 260 265 270

Arg Ile Pro Gly Ile Val Pro Met Gln Ile Pro Ser Pro Glu Val Phe  
 275 280 285

Glu Asp Leu Ile Lys Phe Ser Phe His Thr Asn Val Leu Glu Gly Asn  
 290 295 300

Val Gly Tyr Leu Arg Phe Asp Met Phe Gly Asp Cys Glu Leu Leu Thr  
 305 310 315 320

Gln Val Ser Glu Leu Leu Val Glu His Val Trp Lys Lys Ile Val His  
 325 330 335

Thr Asp Ala Leu Ile Val Asp Met Arg Phe Asn Ile Gly Gly Pro Thr  
 340 345 350

Ser Ser Ile Ser Ala Leu Cys Ser Tyr Phe Phe Asp Glu Gly Pro Pro  
 Page 13

355 YEDA-068 PCT 27.06.07 MS.ST25  
 360 365

Ile Leu Leu Asp Lys Ile Tyr Asn Arg Pro Asn Asp Ser Val Ser Glu  
 370 375 380

Leu Trp Thr Leu Ser Gln Leu Glu Gly Glu Arg Tyr Gly Ser Lys Lys  
 385 390 395 400

Ser Met Val Ile Leu Thr Ser Thr Leu Thr Ala Gly Ala Ala Glu Glu  
 405 410 415

Phe Thr Tyr Ile Met Lys Arg Leu Gly Arg Ala Leu Val Ile Gly Glu  
 420 425 430

Val Thr Ser Gly Gly Cys Gln Pro Pro Gln Thr Tyr His Val Asp Asp  
 435 440 445

Thr Asp Leu Tyr Leu Thr Ile Pro Thr Ala Arg Ser Val Gly Ala Ala  
 450 455 460

Asp Gly Ser Ser Trp Glu Gly Val Gly Val Val Pro Asp Val Ala Val  
 465 470 475 480

Pro Ala Glu Ala Ala Leu Thr Arg Ala Gln Glu Met Leu Gln His Thr  
 485 490 495

Pro Leu Arg Ala Arg Arg Ser Pro Arg Leu His Gly Arg Arg Lys Gly  
 500 505 510

His Gln Gln Gln Ser Gln Gly Arg Ala Gly Ser Leu Gly Arg Asn Gln  
 515 520 525

Gly Val Val Arg Pro Glu Val Leu Thr Glu Ala Pro Ser Gly Gln Lys  
 530 535 540

Arg Gly Leu Leu Gln Cys Gly  
 545 550

<210> 35  
 <211> 405  
 <212> PRT  
 <213> Homo sapiens

<400> 35

Met Ala Ala Ser Gly Lys Thr Ser Lys Ser Glu Pro Asn His Val Ile  
 1 5 10 15

Phe Lys Lys Ile Ser Arg Asp Lys Ser Val Thr Ile Tyr Leu Gly Asn  
 20 25 30

Arg Asp Tyr Ile Asp His Val Ser Gln Val Gln Pro Val Asp Gly Val  
 35 40 45

YEDA-068 PCT 27.06.07 MS.ST25

Val Leu Val Asp Pro Asp Leu Val Lys Gly Lys Lys Val Tyr Val Thr  
50 55 60

Leu Thr Cys Ala Phe Arg Tyr Gly Gln Glu Asp Ile Asp Val Ile Gly  
65 70 75 80

Leu Thr Phe Arg Arg Asp Leu Tyr Phe Ser Arg Val Gln Val Tyr Pro  
85 90 95

Pro Val Gly Ala Ala Ser Thr Pro Thr Lys Leu Gln Glu Ser Leu Leu  
100 105 110

Lys Lys Leu Gly Ser Asn Thr Tyr Pro Phe Leu Leu Thr Phe Pro Asp  
115 120 125

Tyr Leu Pro Cys Ser Val Met Leu Gln Pro Ala Pro Gln Asp Ser Gly  
130 135 140

Lys Ser Cys Gly Val Asp Phe Glu Val Lys Ala Phe Ala Thr Asp Ser  
145 150 155 160

Thr Asp Ala Glu Glu Asp Lys Ile Pro Lys Lys Ser Ser Val Arg Leu  
165 170 175

Leu Ile Arg Lys Val Gln His Ala Pro Leu Glu Met Gly Pro Gln Pro  
180 185 190

Arg Ala Glu Ala Ala Trp Gln Phe Phe Met Ser Asp Lys Pro Leu His  
195 200 205

Leu Ala Val Ser Leu Asn Lys Glu Ile Tyr Phe His Gly Glu Pro Ile  
210 215 220

Pro Val Thr Val Thr Val Thr Asn Asn Thr Glu Lys Thr Val Lys Lys  
225 230 235 240

Ile Lys Ala Phe Val Glu Gln Val Ala Asn Val Val Leu Tyr Ser Ser  
245 250 255

Asp Tyr Tyr Val Lys Pro Val Ala Met Glu Glu Ala Gln Glu Lys Val  
260 265 270

Pro Pro Asn Ser Thr Leu Thr Lys Thr Leu Thr Leu Leu Pro Leu Leu  
275 280 285

Ala Asn Asn Arg Glu Arg Arg Gly Ile Ala Leu Asp Gly Lys Ile Lys  
290 295 300

His Glu Asp Thr Asn Leu Ala Ser Ser Thr Ile Ile Lys Glu Gly Ile  
305 310 315 320

YEDA-068 PCT 27.06.07 MS.ST25

Asp Arg Thr Val Leu Gly Ile Leu Val Ser Tyr Gln Ile Lys Val Lys  
 325 330 335

Leu Thr Val Ser Gly Phe Leu Gly Glu Leu Thr Ser Ser Glu Val Ala  
 340 345 350

Thr Glu Val Pro Phe Arg Leu Met His Pro Gln Pro Glu Asp Pro Ala  
 355 360 365

Lys Glu Ser Tyr Gln Asp Ala Asn Leu Val Phe Glu Glu Phe Ala Arg  
 370 375 380

His Asn Leu Lys Asp Ala Gly Glu Ala Glu Glu Gly Lys Arg Asp Lys  
 385 390 395 400

Asn Asp Val Asp Glu  
 405

<210> 36

<211> 404

<212> PRT

<213> Bos taurus

<400> 36

Met Lys Ala Asn Lys Pro Ala Pro Asn His Val Ile Phe Lys Lys Ile  
 1 5 10 15

Ser Arg Asp Lys Ser Val Thr Ile Tyr Leu Gly Lys Arg Asp Tyr Ile  
 20 25 30

Asp His Val Glu Arg Val Glu Pro Val Asp Gly Val Val Leu Val Asp  
 35 40 45

Pro Glu Leu Val Lys Gly Lys Arg Val Tyr Val Ser Leu Thr Cys Ala  
 50 55 60

Phe Arg Tyr Gly Gln Glu Asp Ile Asp Val Met Gly Leu Ser Phe Arg  
 65 70 75 80

Arg Asp Leu Tyr Phe Ser Gln Val Gln Val Phe Pro Pro Val Gly Ala  
 85 90 95

Ser Gly Ala Thr Thr Arg Leu Gln Glu Ser Leu Ile Lys Lys Leu Gly  
 100 105 110

Ala Asn Thr Tyr Pro Phe Leu Leu Thr Phe Pro Asp Tyr Leu Pro Cys  
 115 120 125

Ser Val Met Leu Gln Pro Ala Pro Gln Asp Val Gly Lys Ser Cys Gly  
 130 135 140

YEDA-068 PCT 27.06.07 MS.ST25

Val Asp Phe Glu Ile Lys Ala Phe Ala Thr His Ser Thr Asp Val Glu  
145 150 155 160

Glu Asp Lys Ile Pro Lys Lys Ser Ser Val Arg Leu Leu Ile Arg Lys  
165 170 175

Val Gln His Ala Pro Arg Asp Met Gly Pro Gln Pro Arg Ala Glu Ala  
180 185 190

Ser Trp Gln Phe Phe Met Ser Asp Lys Pro Leu Arg Leu Ala Val Ser  
195 200 205

Leu Ser Lys Glu Ile Tyr Tyr His Gly Glu Pro Ile Pro Val Thr Val  
210 215 220

Ala Val Thr Asn Ser Thr Glu Lys Thr Val Lys Lys Ile Lys Val Leu  
225 230 235 240

Val Glu Gln Val Thr Asn Val Val Leu Tyr Ser Ser Asp Tyr Tyr Ile  
245 250 255

Lys Thr Val Ala Ala Glu Glu Ala Gln Glu Lys Val Pro Pro Asn Ser  
260 265 270

Ser Leu Thr Lys Thr Leu Thr Leu Val Pro Leu Leu Ala Asn Asn Arg  
275 280 285

Glu Arg Arg Gly Ile Ala Leu Asp Gly Lys Ile Lys His Glu Asp Thr  
290 295 300

Asn Leu Ala Ser Ser Thr Ile Ile Lys Glu Gly Ile Asp Lys Thr Val  
305 310 315 320

Met Gly Ile Leu Val Ser Tyr Gln Ile Lys Val Lys Leu Thr Val Ser  
325 330 335

Gly Leu Leu Gly Glu Leu Thr Ser Ser Glu Val Ala Thr Glu Val Pro  
340 345 350

Phe Arg Leu Met His Pro Gln Pro Glu Asp Pro Asp Thr Ala Lys Glu  
355 360 365

Ser Phe Gln Asp Glu Asn Phe Val Phe Glu Glu Phe Ala Arg Gln Asn  
370 375 380

Leu Lys Asp Ala Gly Glu Tyr Lys Glu Glu Lys Thr Asp Gln Glu Ala  
385 390 395 400

Ala Met Asp Glu

YEDA-068 PCT 27.06.07 MS.ST25

<210> 37  
<211> 348  
<212> PRT  
<213> Homo sapiens

&lt;400&gt; 37

Met Asn Gly Thr Glu Gly Pro Asn Phe Tyr Val Pro Phe Ser Asn Ala  
1 5 10 15

Thr Gly Val Val Arg Ser Pro Phe Glu Tyr Pro Gln Tyr Tyr Leu Ala  
20 25 30

Glu Pro Trp Gln Phe Ser Met Leu Ala Ala Tyr Met Phe Leu Leu Ile  
35 40 45

Val Leu Gly Phe Pro Ile Asn Phe Leu Thr Leu Tyr Val Thr Val Gln  
50 55 60

His Lys Lys Leu Arg Thr Pro Leu Asn Tyr Ile Leu Leu Asn Leu Ala  
65 70 75 80

Val Ala Asp Leu Phe Met Val Leu Gly Gly Phe Thr Ser Thr Leu Tyr  
85 90 95

Thr Ser Leu His Gly Tyr Phe Val Phe Gly Pro Thr Gly Cys Asn Leu  
100 105 110

Glu Gly Phe Phe Ala Thr Leu Gly Gly Glu Ile Ala Leu Trp Ser Leu  
115 120 125

Val Val Leu Ala Ile Glu Arg Tyr Val Val Val Cys Lys Pro Met Ser  
130 135 140

Asn Phe Arg Phe Gly Glu Asn His Ala Ile Met Gly Val Ala Phe Thr  
145 150 155 160

Trp Val Met Ala Leu Ala Cys Ala Ala Pro Pro Leu Ala Gly Trp Ser  
165 170 175

Arg Tyr Ile Pro Glu Gly Leu Gln Cys Ser Cys Gly Ile Asp Tyr Tyr  
180 185 190

Thr Leu Lys Pro Glu Val Asn Asn Glu Ser Phe Val Ile Tyr Met Phe  
195 200 205

Val Val His Phe Thr Ile Pro Met Ile Ile Ile Phe Phe Cys Tyr Gly  
210 215 220

Gln Leu Val Phe Thr Val Lys Glu Ala Ala Ala Gln Gln Gln Glu Ser  
225 230 235 240

YEDA-068 PCT 27.06.07 MS.ST25  
 Ala Thr Thr Gln Lys Ala Glu Lys Glu Val Thr Arg Met Val Ile Ile  
 245 250 255

Met Val Ile Ala Phe Leu Ile Cys Trp Val Pro Tyr Ala Ser Val Ala  
 260 265 270

Phe Tyr Ile Phe Thr His Gln Gly Ser Asn Phe Gly Pro Ile Phe Met  
 275 280 285

Thr Ile Pro Ala Phe Phe Ala Lys Ser Ala Ala Ile Tyr Asn Pro Val  
 290 295 300

Ile Tyr Ile Met Met Asn Lys Gln Phe Arg Asn Cys Met Leu Thr Thr  
 305 310 315 320

Ile Cys Cys Gly Lys Asn Pro Leu Gly Asp Asp Glu Ala Ser Ala Thr  
 325 330 335

Val Ser Lys Thr Glu Thr Ser Gln Val Ala Pro Ala  
 340 345

<210> 38  
 <211> 15  
 <212> PRT  
 <213> Artificial

<220>  
 <223> peptide R16  
 <400> 38

Ala Asp Gly Ser Ser Trp Glu Gly Val Gly Val Val Pro Asp Val  
 1 5 10 15

<210> 39  
 <211> 23  
 <212> PRT  
 <213> Artificial

<220>  
 <223> .

<400> 39

Pro Thr Ala Arg Ser Val Gly Ala Ala Asp Gly Ser Ser Trp Glu Gly  
 1 5 10 15

Val Gly Val Val Pro Asp Val  
 20

<210> 40  
 <211> 23  
 <212> PRT  
 <213> Artificial

<220>  
 <223> .

YEDA-068 PCT 27.06.07 MS.ST25

&lt;400&gt; 40

His Val Asp Asp Thr Asp Leu Tyr Leu Thr Ile Pro Thr Ala Arg Ser  
1 5 10 15Val Gly Ala Ala Asp Gly Ser  
20

&lt;210&gt; 41

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; peptide G-8

&lt;400&gt; 41

Thr Ser Ser Glu Val Ala Thr Glu  
1 5

&lt;210&gt; 42

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; ;

&lt;400&gt; 42

Asp Thr Asn Leu Ala Ser Ser Thr  
1 5

&lt;210&gt; 43

&lt;211&gt; 18

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 43

Asp Thr Asn Leu Ala Ser Ser Thr Ile Ile Lys Glu Gly Ile Asp Lys  
1 5 10 15

Thr Val

&lt;210&gt; 44

&lt;211&gt; 22

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 44

Val Pro Leu Leu Ala Asn Asn Arg Glu Arg Arg Gly Ile Ala Leu Asp  
1 5 10 15

YEDA-068 PCT 27.06.07 MS.ST25

Gly Lys Ile Lys His Glu  
20

<210> 45  
<211> 20  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 45

Thr Ser Ser Glu Val Ala Thr Glu Val Pro Phe Arg Leu Met His Pro  
1 5 10 15

Gln Pro Glu Asp  
20

<210> 46  
<211> 20  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 46

Ser Leu Thr Lys Thr Leu Thr Leu Val Pro Leu Leu Ala Asn Asn Arg  
1 5 10 15

Glu Arg Arg Gly  
20

<210> 47  
<211> 20  
<212> PRT  
<213> Artificial

<220>  
<223> .

<400> 47

Ser Leu Thr Arg Thr Leu Thr Leu Leu Pro Leu Leu Ala Asn Asn Arg  
1 5 10 15

Glu Arg Ala Gly  
20

<210> 48  
<211> 21  
<212> PRT  
<213> Artificial

<220>  
<223> .

YEDA-068 PCT 27.06.07 MS.ST25

&lt;400&gt; 48

Lys Glu Gly Ile Asp Lys Thr Val Met Gly Ile Leu Val Ser Tyr Gln  
1 5 10 15Ile Lys Val Lys Leu  
20

&lt;210&gt; 49

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; .

&lt;400&gt; 49

Lys Glu Gly Ile Asp Arg Thr Val Leu Gly Ile Leu Val Ser Tyr Gln  
1 5 10 15Ile Lys Val Lys Leu  
20

&lt;210&gt; 50

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; PEPTIDE 8 ANALOG

&lt;400&gt; 50

Thr Ser Ser Glu Ala Ala Thr Glu  
1 5

&lt;210&gt; 51

&lt;211&gt; 8

&lt;212&gt; PRT

&lt;213&gt; Artificial

&lt;220&gt;

&lt;223&gt; M-8

&lt;400&gt; 51

Asp Thr Ala Leu Ala Ser Ser Thr  
1 5