

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
31 December 2008 (31.12.2008)

PCT

(10) International Publication Number
WO 2009/002918 A2

(51) International Patent Classification:
A61K 48/00 (2006.01)

(21) International Application Number:
PCT/US2008/067871

(22) International Filing Date: 23 June 2008 (23.06.2008)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/929,335 22 June 2007 (22.06.2007) US

(71) Applicant (for all designated States except US): NUCLEONICS, INC. [US/US]; 702 Electronic Drive, Horsham, PA 19044 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): MCCALLUS, Daniel [US/US]; P.O. Box 596, 312 Gumbes Road, Oaks, PA 19456 (US). PACHUK, Catherine [US/US]; 400 Foxcroft Drive, Blue Bell, PA 19422 (US).

(74) Agents: HAYMAN, Mark et al.; Cooley Godward Kronish LLP, 777 6th Street, N.W., Suite 1100, Washington, DC 20001 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- without international search report and to be republished upon receipt of that report
- with sequence listing part of description published separately in electronic form and available upon request from the International Bureau

(54) Title: HEPATITIS C DSRNA EFFECTOR MOLECULES, EXPRESSION CONSTRUCTS, COMPOSITIONS, AND METHODS OF USE




Fig. 1

(57) Abstract: The present invention provides agents, compositions, constructs and methods for silencing HCV polynucleotides, as well as methods and compositions for treating or preventing HCV infection in a mammalian cell. In one aspect, the present invention provides an agent or composition comprising at least one double-stranded RNA effector molecule or complex. The double-stranded RNA effector molecule or complex comprises: (1) a sequence of at least 19 nucleotides having at least 90% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), HCV Conserved Region 5 (SEQ ID NO: 4), (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), ATR-4 (SEQ ID NO: 89); and (2) its complementary sequence. In another aspect, the present invention provides a construct suitable for replication in a host cell, and/or suitable for expression of an RNA molecule or complex of the invention in vitro or in vivo. In a third aspect, the present invention provides a method for silencing HCV RNA in a mammalian cell, which comprises administering to the mammalian cell an agent, composition, or construct of the invention in a manner and amount effective to silence HCV RNA in the cell. In a related aspect, the invention provides a method for treating or preventing HCV infection in a patient, comprising administering to the patient an effective amount of an agent, composition, or construct of the invention as described herein.

WO 2009/002918 A2

HEPATITIS C dsRNA EFFECTOR MOLECULES, EXPRESSION CONSTRUCTS, COMPOSITIONS, AND METHODS OF USE

5

PRIORITY

This application claims priority to US Provisional Application No. 60/929,335, filed June 22, 2007, which is hereby incorporated by reference in its entirety.

10

FIELD OF THE INVENTION

The present invention relates to nucleic acid-based therapeutics for treating or preventing Hepatitis C Virus (HCV) infection, and particularly RNAi-based therapeutics.

15

BACKGROUND OF THE INVENTION

Hepatitis C is an RNA virus containing a single-stranded positive-sense RNA genome of about 9,600 nts. The genes for the viral structural and non-structural proteins are flanked by 5' and 3' untranslated regions (UTRs), which are essential for genome replication. For example, the 5' UTR contains an internal ribosome entry site (IRES) which is indispensable for the initiation of HCV polyprotein translation. HCV has been classified into six major genotypes, each comprising further subgroups, which differ in their sequence homology by more than 30%. The distribution of these genotypes differs geographically. For example, genotypes 1a and 1b are the most prevalent genotypes found within the U.S., while genotypes 2 and 3 are more prevalent in other countries.

20

Because of the sequence variability of HCV, the development of vaccines and therapeutic drugs, including RNAi-based therapeutics, that would be active against the majority of viruses, must take advantage of the rare conserved epitopes and sequences found among the viral genotypes and quasispecies. In fact, the mutability of HCV is such that even within an infected individual, the HCV virus exists as a swarm of variants or "quasispecies" of a predominant type rather than as a single entity.

25

30

Thus, to apply a gene-silencing-based strategy to the treatment or prevention of HCV infection, it is necessary to identify sufficiently conserved stretches of nucleotide sequence in this highly mutable virus. That is, since RNA interference is a sequence-specific effect, therapeutic or prophylactic RNAi molecules must be specific for HCV target sequences, despite the fact that hepatitis C viral genomes are highly variable. While HCV target sequence conservation is an important consideration in the design of sequence-specific anti-HCV prophylactic or therapeutic modalities such as RNAi or antisense, e.g., some of the highly conserved regions of the HCV genome such as the 5' UTR are known to be highly structured, while some regions of the viral genome are present in the infected cell in association with proteins which make them largely inaccessible to antisense or RNAi. The lack of a readily available HCV animal model and problems with various HCV cell culture models, e.g., the absence or deficiencies in viral infection or replication models, have hindered the development of anti-HCV pharmaceuticals of all types.

Despite well over a decade of research efforts, there are no vaccines available for HCV. As a consequence, the rate of new HCV infections around the world is extremely high. The WHO estimates that globally 170 million individuals carry chronic HCV infections and that new infections are established at a rate of 3 to 4 million annually.

Chronic HCV infection induces liver inflammation, causing progressive liver disease that can lead to cirrhosis and hepatocellular carcinoma (liver cancer). Chronic HCV infection becomes established in 75%-85% of individuals experiencing an initial infection, and HCV-related liver failure is the most common indication cited for liver transplantation in the U.S. Chronic HCV infection in its early stages may cause only mild non-specific symptoms, such as fatigue, or be completely asymptomatic, leaving many infected individuals unaware that they carry a dangerous chronic infection.

Current therapies for HCV infection, which may include a 6 to 12 month regimen of pegylated interferon and ribavirin, can lead to a cure in a minority of patients. Response rates vary by HCV genotype, with genotype 2 and 3 patients exhibiting a 76% response rate to the current standard therapy while patients infected with genotype 1a and 1b having only a 46% response rate. Unfortunately, genotype 1 accounts for 60% of global infections and is the

dominant strain in the U.S., Japan, and Western Europe. Complicating genotype 1 resistance to ribavarin and interferon is the fact that both drugs have side effect profiles that can require dose reduction or discontinuance of therapy when patients experience side effects. Further complicating patient outcomes is the fact that patients who fail an initial treatment regimen rarely respond favorably to a subsequent round of treatment with interferon and ribavarin.

Clinicians who treat HCV patients are hopeful that current and future research programs will yield options that improve the response rate for genotype 1 patients, which is currently less than 50% using ribavarin and interferon. New treatment options that have a more tolerable side effect profile would improve patient compliance and enable more patients to complete a full course of therapeutic intervention.

There remains a need for treatment options for HCV-exposed or infected patients, including for highly conserved nucleic acid-based molecules, including double-stranded RNAs and constructs encoding dsRNAs, capable of inhibiting the replication of HCV in mammalian cells. Such nucleic acid based anti-HCV therapeutic agents have the potential to improve patient response rates to therapy, improve adverse event profile, and eliminate or significantly delay the development of drug resistant escape mutant virus.

SUMMARY OF THE INVENTION

The present invention provides agents, compositions, constructs, and methods for silencing HCV polynucleotides, as well as methods and compositions capable of inhibiting HCV replication and for treating or preventing HCV infection in a mammalian cell.

In one aspect, the present invention provides an agent or composition for silencing HCV RNA in a cell. In one aspect, the agent or composition inhibits HCV viral replication and antigen expression in a mammalian cell, preferably a human cell. In this aspect, the agent or composition comprises at least one double-stranded RNA effector molecule or complex. The double-stranded RNA effector molecule or complex comprises: (1) a sequence of at least 19, 20, or 21 consecutive nucleotides having at least 90%, 95%, or 100% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), HCV Conserved Region 5 (SEQ ID NO: 4),

active target region (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89); and (2) its complementary sequence. Preferably the dsRNA effector molecule will include (1) a sequence of 19 to 29, 19 to 25, 20 to 25, 21 to 25, or 21 to 23 consecutive nucleotides within
5 HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), HCV Conserved Region 5 (SEQ ID NO: 4), active target region (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89); and (2) its complementary sequence. The effector molecule may optionally form a stem-loop structure, with the sequence of at least 19
10 nucleotides and its complementary sequence being connected via a loop sequence, thereby providing, for example, short-hairpin RNAs (shRNAs) suitable for RNAi-based HCV therapeutics. In one aspect, multiple different of such dsRNA effector molecules of the invention, e.g., two, three, four, five or more, are administered to or expressed concomitantly in a mammalian cell, to eliminate or
15 substantially delay the emergence of drug resistant viral escape mutants.

In another aspect, the present invention provides a construct suitable for replication in a host cell, and/or suitable for expression of an RNA molecule of the invention *in vitro* or *in vivo*. The construct of the invention encodes at least one RNA effector molecule of the invention, which may be operably linked to a
20 promoter sequence, such as an RNA Polymerase I, RNA polymerase II, or RNA polymerase III promoter sequence as described herein. In one aspect, multiple such dsRNA effector molecules of the invention, e.g., hairpin dsRNA molecules, are encoded by a single expression construct under the control of one or more of such promoters.

25 In a third aspect, the present invention provides a method for silencing or inhibiting the replication of HCV, including inhibition of HCV RNA and/or HCV antigen expression, in a mammalian cell. In this aspect, the method of the invention comprises administering to the mammalian cell an agent, composition, or construct of the invention in a manner and amount effective to inhibit HCV
30 replication and/or HCV RNA or antigen expression in the cell. In a related aspect, the present invention provides a method for treating or preventing HCV infection in a patient, comprising administering to the patient an effective amount and regimen of an agent, composition, or construct of the invention as described herein.

The present invention further provides methods for preparing the dsRNA effector molecules, compositions, and constructs of the invention.

DESCRIPTION OF THE DRAWINGS

5 **Figure 1** shows a number of highly active shRNAs which map to four active target regions (ATR-1, ATR-2, ATR-3, and ATR-4) within the HCV 5' UTR.

Figure 2 is a plasmid diagram of the multi-cistronic plasmid QJ, which expresses four different active shRNAs, HCV 5'-21-61, HCV 5'-21-94, HCV 5'-21-124, and HCV 5'-21-135.

10 **Figure 3** is a plasmid diagram of a mono-cistronic plasmid HCV 5' 21-61, which expresses the shRNA HCV 5' 21-61 from the 7SK 4A promoter.

DETAILED DESCRIPTION OF THE INVENTION

To identify sequences that are most conserved among HCV genomes
15 worldwide, a bioinformatic analysis was conducted. There are 93 complete genomes published in GenBank version 134.0 and these were compared for the identification of sequences of 19 nts or greater that are >95% conserved, and which could potentially serve as target sites for small inhibitory RNAs (siRNAs) and short-hairpin RNAs (shRNAs). The following sequences were identified
20 within the HCV 5'UTR, and are shown with respect to GenBank Accession ID AB047639 (SEQ ID NO: 1).

HCV Conserved region 1: nts 35-102 of AB047639 (SEQ ID NO: 2).

5'- atcactcccctgtgaggaactactgtcttcacgcagaaagcgcctagccatggcgtagtatgagtgt-3'

25

HCV Conserved region 2: nts 119-176 of AB047639 (SEQ ID NO: 3).

5'-ccccccctcccgggagagccatagtggtctgcggaaccggtgagtacaccggaattgc-3'

HCV Conserved region 5: nts 270-338 of AB047639 (SEQ ID NO: 4).

5'-gcgaaaggccttggtactgcctgatagggcgcttgcgagtgccccgggaggtctcgtagaccgtgca-3'

Four highly conserved and highly active target regions (ATR), preferred for some applications, were identified:

5 ATR-1: 5'-CCCTGTGAGGAACTACTGTCTTCACGCAGAA-3' (SEQ ID NO: 86), mapping to nucleotide coordinates 42 to 76 of GenBank Accession No. AB047639, found within Conserved Region 1 (SEQ ID NO: 2).

ATR-2: 5'-TCCCGGGAGAGCCATAGTGGTCTGCGGAA-3' (SEQ ID NO: 87), mapping to nucleotide coordinates 126 to 154 of GenBank Accession No. AB047639, found within Conserved Region 2 (SEQ ID NO: 3).

ATR-3: 5'-CGAAAGGCCTTGTGGTACTGC-3', (SEQ ID NO: 88) mapping to nucleotide coordinates 271 to 297 of GenBank Accession No. AB047639, found within Conserved Region 5 (SEQ ID NO: 4).

15 ATR-4: 5'-TGCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCA-3', (SEQ ID NO: 89) mapping to nucleotide coordinates 305 to 338 of GenBank Accession No. AB047639, found within Conserved Region 5 (SEQ ID NO: 4).

In this context, the present invention provides agents, compositions, constructs and methods for silencing HCV polynucleotides, as well as for treating or preventing HCV infection in a mammalian cell.

20 Effector RNA Molecules and Complexes

In one aspect, the present invention provides an agent for silencing HCV RNA in a cell. The agent comprises at least one double-stranded RNA effector molecule or complex, which comprises: (1) a sequence of at least 19, 20, or 21 consecutive nucleotides having at least about 90% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), HCV Conserved Region 5 (SEQ ID NO: 4), ATR-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89) and (2) its complementary sequence.

30 In this context, the nucleotide "t" in SEQ ID NOS: 1, 2, 3, 4, etc. is considered to be identical with "u," which would take the place of "t" in the corresponding RNA sequence. Thus, throughout this application it will be

understood that where RNA sequences are described for convenience with respect to encoding or corresponding DNA sequences, "t" will be replaced by "u" in the RNA sequence.

In certain embodiments, the at least 19 nucleotides of the dsRNA effector molecule or complex has at least about 95% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), HCV Conserved Region 5 (SEQ ID NO: 4), ATR-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89), such as at least about 96%, 97%, 98%, 99%, or about 100% identity. Identity between two nucleotide sequences may be determined using any suitable algorithm known in the art, such as Tatusova et al., Blast 2 sequences - a new tool for comparing protein and nucleotide sequences, *FEMS Microbiol Lett.* 174:247-250 (1999).

In some embodiments, the at least 19 nucleotides of the dsRNA effector molecule or complex has a sequence selected from within nucleotides 42-76 (ATR-1), 126-154 (ATR-2), 271 to 297 (ATR-3), 305 to 338 (ATR-4) or 271-338 of SEQ ID NO: 1 (GenBank Accession ID AB047639). For example, the dsRNA effector molecule or complex may comprise at least 19 nucleotides selected from within nucleotides 305-338 of SEQ ID NO: 1.

The dsRNA effector molecules and complexes may have double-stranded regions that vary somewhat in length, so long as the effector molecule or complex is effective for silencing the target polynucleotide, that is, the length of the double-stranded region is generally sufficient to trigger RNAi-mediated degradation of the target sequence. For example, the agent or composition of the invention may comprise at least one double-stranded RNA effector molecule or complex containing a double-stranded region of from 19 or 21 base pairs to about 30 or 40 base pairs, or from 21 base pairs to about 27 base pairs. In certain embodiments, the double-stranded RNA effector molecule or complex contains a double-stranded region of about 21 or about 27 base pairs.

In accordance with the invention, the double-stranded RNA molecules and complexes of the invention may exist in a denatured or substantially denatured form. Alternatively, the double-stranded RNA molecules and complexes may exist in a double-stranded conformation, or a substantially double-stranded

conformation, or a partially double-stranded conformation, at least with respect to the regions of complementarity. Generally, the RNA molecules and complexes of the invention are capable of forming double-stranded structures under physiological conditions (e.g., intracellular conditions), where these structures are sufficient to trigger RNAi-mediated gene silencing.

In some embodiments, the double-stranded RNA effector molecule or complex has a double-stranded region that comprises, or consists essentially of (or consists of) in one strand, a sequence selected from SEQ ID NOS: 5-42 as disclosed herein. Such sequences, which correspond to HCV Conserved Regions 1, 2, and 5, are listed in Table 3.

The double-stranded RNA effector molecules of the invention comprise a region of self-complementarity such that nucleotides in one segment of the molecule base pair with nucleotides in another segment of the molecule (e.g., stem-loop or hairpin structure as described herein). In contrast, the double-stranded RNA effector duplexes or complexes of the invention include at least two separate polynucleotide strands that have a region of complementarity to each other. The double-stranded RNA complexes (i.e., duplexes) may be fully complementary, that is, may contain no single stranded regions, such as single stranded ends. In other embodiments, the double-stranded RNA complex contains short single-stranded ends, such as single-stranded 5' or 3' ends of from about 1 to about 5 nucleotides (e.g., 1, 2, or 3 nucleotides).

In certain embodiments, the effector molecule is a short hairpin dsRNA (shRNA) or a microRNA. A "shRNA" (short-hairpin RNA) is an RNA molecule of less than approximately 200 or 100 nucleotides, such as about 70 nucleotides or less, in which at least one stretch of nucleotides (e.g., at least about 19 nucleotides) is base paired with a complementary sequence located on the same RNA molecule and separated from the complementary sequence by an unpaired region. These single-stranded hairpin regions form a single-stranded loop between the stem structure created by the two regions of base complementarity. The length and nucleotide sequence of the loop sequence is not narrowly critical, and may range, for example, from about 4 to 5 to about 20 nucleotides in length, or from about 7 to about 10 nucleotides in length. For example, the loop sequence may be about 9 nucleotides in length. An exemplary loop sequence is 5'- agagaactt - 3' (SEQ ID NO: 43). The loop may vary considerably in length

and sequence, although loop sequences which assume significant secondary structure are to be avoided, as are poly T (e.g., T₄ to T₅ or more) sequences, which might trigger premature termination of transcription for effector molecules expressed by polymerase III promoters. In addition to a "stem" region which
5 comprises the identified homologous and complementary HCV sequences, in certain embodiments the hairpin molecule and/or the expression vector encoding the hairpin RNA will include additional 5' and/or 3' sequences, including in some embodiments 5' and/or 3' flanking sequences as well as loop sequences derived from miRNAs. See e.g., US 2004/0053411, the teaching of which is hereby
10 incorporated by reference.

Exemplary shRNAs in accordance with the invention have a sequence selected from SEQ ID NOS: 44-81. The complementary strand may also, optionally, have from one to five uracil nucleotides at its 3'-end, which may correspond to transcribed transcription termination sequences.

15 In addition to single shRNAs, the invention includes dual or bi-finger and multi-finger hairpin dsRNAs, in which the RNA molecule comprises two or more of such stem-loop structures, each separated by a single-stranded spacer region. In some embodiments such two or more stem-loop structures may be encoded by an expression construct and operably linked to a single promoter. Thus, the
20 hairpin dsRNA may be a single hairpin dsRNA or a bi-fingered, or multi-fingered dsRNA hairpin as described in PCT/US03/033466 or WO 04/035766, or a partial or forced hairpin structure as described in WO 2004/011624, the disclosures of which are hereby incorporated by reference. In these embodiments, the multi-finger hairpin RNA contains two, three, or four double stranded regions, each
25 corresponding to an HCV target sequence independently selected from SEQ ID NOS: 5-42, such as SEQ ID NOS: 11, 19, 22, and 33. In these multi-finger hairpin RNAs the stem loop sequences may be independently selected from SEQ ID NOS: 44-81.

HCV target sequences (as described above) may also be combined into
30 other RNA structures suitable for RNAi-based therapeutics, including a single stem loop structure with multiple double-stranded regions separated by mismatch region(s). Such RNA structures are disclosed in US Application No. 10/531,349, which was published as US 2006/0035344 on February 16, 2006, which is hereby incorporated by reference. In these embodiments, the multi-target

effector molecule may comprise two, three, or four double-stranded regions each corresponding to an HCV target sequence (as described herein) independently selected from SEQ ID NOS: 5-42, such as SEQ ID NO: 11, 19, 22, and 33.

The RNA molecules and complexes of the invention may be composed
5 purely or predominately of ribonucleotides found naturally in RNA (A, U, C, G), or may contain chemically modified nucleotides. Exemplary chemically modified nucleotides include phosphorothioate internucleotide linkages, 2'-deoxyribonucleotides, 2'-O-methyl ribonucleotides, 2'-deoxy-2'-fluoro ribonucleotides, "universal base" nucleotides, "acyclic" nucleotides, 5-C-methyl
10 nucleotides, and terminal glyceryl and/or inverted deoxy abasic residue incorporation. These, as well as other chemical modifications, support RNAi-mediated gene silencing while having superior serum stability.

RNA Compositions

The invention further provides compositions containing from two to five
15 double-stranded RNA effector molecules or complexes, such as from two to five double-stranded RNA effector molecules or complexes as described above. For example, the two to five double-stranded RNA effector molecules (e.g., shRNAs) or complexes may each comprise: (1) a sequence of at least 19 nucleotides having at least 90% identity with a nucleotide sequence within Conserved Region
20 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4); ATR-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89), and (2) its complementary sequence. Where the two to five double-stranded RNAs are each shRNAs, the shRNAs each further comprise a loop sequence as described above.

25 Thus, the composition of the invention may contain a plurality of double-stranded RNA effector molecules, each having a double-stranded region that comprises (or consists essentially of) a sequence independently selected from SEQ ID NOS: 5-42, and its complementary sequence. An exemplary composition may have: a first dsRNA effector molecule comprising SEQ ID NO:
30 11 and its complementary sequence; a second dsRNA effector molecule comprising SEQ ID NO: 19 and its complementary sequence; a third dsRNA effector molecule comprising SEQ ID NO: 22 and its complementary sequence; and a fourth dsRNA effector molecule comprising SEQ ID NO: 33 and its

complementary sequence. One, two, three, or four of these double stranded RNA effector molecules may be in the form of short-hairpin RNAs, that is, the sense and anti-sense strands may be connected by a short loop sequence as described herein. Thus, in these embodiments, the composition of the invention
5 may comprise from two to five (e.g., four) double-stranded RNA effector molecules independently selected from SEQ ID NOS: 44-81. For example, the composition may comprise the four shRNAs represented by the sequences: SEQ ID NO: 50, SEQ ID NO: 58, SEQ ID NO: 61, and SEQ ID NO: 72. In some
10 embodiments the plurality of anti-HCV RNA effector molecules are expressed from a plasmid or viral vector within a human cell.

Alternatively still, the composition of the invention may contain a plurality of double-stranded RNA effector duplexes or complexes, each having a double-stranded region that comprises (or consists essentially of) a sequence
15 independently selected from SEQ ID NOS: 5-42, and its complementary sequence. For example: a first RNA effector complex may comprise, in one RNA strand, the nucleotide sequence of SEQ ID NO: 11; a second RNA effector complex may comprise, in one strand, the nucleotide sequence of SEQ ID NO:
20 19; a third RNA effector complex may comprise, in one strand, the nucleotide sequence of SEQ ID NO: 22; and a fourth RNA effector complex may comprise, in one strand, the nucleotide sequence of SEQ ID NO: 33. Such RNA molecules are base-paired in the complex with (or are capable of base-pairing with) a
complementary or partially complementary RNA molecule in the complex, that is an RNA molecule having a sequence complementary to SEQ ID NOS: 11, 19, 22,
and 33, respectively.

25 Constructs

In another aspect, the present invention provides a construct encoding at least one RNA molecule or complex of the invention. The construct may be, for example, a plasmid or viral vector. Such constructs may be expression
30 constructs suitable for expression of the encoded RNA *in vitro* or *in vivo*, or may otherwise be suitable for replication of the construct in a host cell, such as a prokaryotic or eukaryotic host cell, including bacteria, yeast, and mammalian including human host cells. The construct may include an origin of replication, mechanism for selection (e.g., antibiotic resistance gene) as well as elements to

facilitate removal of the RNA-encoding sequence for sub-cloning into additional constructs or vectors as may be desired, such as conveniently placed restriction endonuclease cleavage sites (e.g., flanking the RNA-encoding sequence(s)).

In some embodiments, the construct is an expression construct containing a DNA segment that encodes an RNA molecule of the invention, with the DNA segment being operably linked to a promoter to drive expression of the RNA molecule. An "expression construct" is any double-stranded DNA or double-stranded RNA designed to produce an RNA of interest. The invention includes expression constructs in which one or more of the promoters is not in fact operably linked to a polynucleotide sequence to be transcribed, but instead is designed for efficient insertion of an operably-linked polynucleotide sequence to be transcribed by the promoter, for instance by way of one or more restriction cloning sites in operative association with the one or more promoters.

Transfection or transformation of the expression construct into a recipient cell allows the cell to express an RNA effector molecule encoded by the expression construct. An expression construct may be a genetically engineered plasmid, virus, recombinant virus, or an artificial chromosome derived from, for example, a bacteriophage, adenovirus, adeno-associated virus, retrovirus, lentivirus, poxvirus, or herpesvirus. Expression vectors for use with the invention contain sequences from bacteria, viruses or phages. Such vectors include chromosomal, episomal and virus-derived vectors, e.g., vectors derived from bacterial plasmids, bacteriophages, yeast episomes, yeast chromosomal elements, and viruses; as well as vectors derived from combinations thereof, such as those derived from plasmid and bacteriophage genetic elements, cosmids and phagemids. Exemplary vectors are double-stranded DNA phage vectors and double-stranded DNA viral vectors.

In certain embodiments, the expression construct of the invention is a plasmid, such as a plasmid suitable for delivery to and/or for RNA expression in a mammalian cell.

The construct of the invention encodes at least one double-stranded RNA effector molecule, which comprises: (1) a sequence of at least 19 nucleotides having at least about 90% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3),

HCV Conserved Region 5 (SEQ ID NO: 4); ATR-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89), and (2) its complementary sequence.

In certain embodiments, the at least 19 nucleotides have at least about 5 95% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), HCV Conserved Region 5 (SEQ ID NO: 4), ATR-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89), such as at least about 96%, 97%, 98%, 99%, or about 100% identity. Identity between two nucleotide sequences may be 10 determined using any suitable algorithm known in the art, such as Tatusova et al., Blast 2 sequences - a new tool for comparing protein and nucleotide sequences, *FEMS Microbiol Lett.* 174:247-250 (1999).

In some embodiments, the at least one RNA effector molecule encoded by the construct or vector has a sequence selected from within nucleotides 42-76, 15 126-154, or 271-338 of SEQ ID NO: 1 (GenBank Accession ID AB047639). For example, the construct may encode an RNA effector molecule comprising at least 19 nucleotides selected from within nucleotides 305-338 of SEQ ID NO: 1.

The expression construct of the invention may encode RNA effector molecules having double-stranded regions that vary somewhat in length, so long 20 as the effector molecule is effective for silencing the target polynucleotide. For example, the expression construct may encode at least one double-stranded RNA effector molecule containing a double-stranded region of from 19 or 21 base pairs to about 30 or 40 base pairs, or from 21 to about 27 base pairs. In certain embodiments, the double-stranded RNA effector molecule contains a double- 25 stranded region of about 21 or about 27 base pairs.

In some embodiments, the construct (e.g., expression vector) encodes one or more double-stranded RNA effector molecules, each having a double-stranded region that comprises, or consists essentially of (or consists of) in one strand, a sequence selected from SEQ ID NOS: 5-42. Such sequences are listed in Table 30 3. In some embodiments, the expression vector encodes at least one double stranded RNA effector molecule that comprises: 1) a sequence selected from SEQ ID NO: 9, 11, 19, 22, 26, 31, 32, and 33; 2) the complement of said sequence, and optionally, 3) a sequence linking 1) and 2). In some

embodiments, the expression vector will encode at least two, three, four, or more of such dsRNA effector molecules. In some embodiments the expression vector will encode at least four different dsRNA effector molecules comprising, in double stranded conformation, SEQ ID NOS: 11, 19, 22, and 33; SEQ ID NOS: 9, 11, 31, and 33; SEQ ID NOS: 11, 19, 31, and 33; SEQ ID NOS: 19, 22, 32, and 33; SEQ ID NOS: 11, 19, 22, and 33; or SEQ ID NOS: 26, 31, 32, and 33.

The encoded double stranded RNA effector molecule(s) may be composed of two separate complementary, or partially complementary RNA molecules (e.g., expressed from separate expression cassettes on the same or different vectors), or may alternatively be in the form of a single RNA molecule, such as a short-hairpin RNA. In the latter embodiment, the sense and anti-sense polynucleotides are connected via a loop sequence, which is generally single-stranded. The length of the loop sequence is not narrowly critical, and may range, for example, from 4 to 20 nucleotides in length, or from 7 to 10 nucleotides in length, or longer if desired, e.g., 30nt, 40nt, etc. For example, the loop sequence may be about 9 nucleotides in length. An exemplary loop sequence is 5'- agagaactt - 3' (SEQ ID NO: 43). The loop sequence may also vary considerably, so long as structure-forming complementarity is avoided, and, in the case of RNA polymerase III promoter transcribed sequences, poly T termination sequences are avoided.

In accordance with this aspect of the invention, exemplary expression constructs encode double-stranded RNA effector molecule(s) having a sequence selected from SEQ ID NOS: 44-81. The complementary strand may also, optionally have from 1 to 5 uracil nucleotides at its 3'-end, representing transcribed RNA pol III termination signal nucleotides. The expressed dsRNA effector molecule may also optionally include one to five or more additional 5' nucleotides, which may vary depending on the transcription start site.

The expression construct may be engineered to encode multiple, e.g., three, four, five or more RNA molecules, such as the RNA molecules described herein, including short hairpin dsRNAs. The expression construct may include a plurality of promoters, e.g., RNA polymerase I, II or III promoters, mitochondrial promoters, etc., operable in the host mammalian cell. Each promoter may be operably linked to a sequence encoding one or more RNA molecules of the invention, followed by an appropriate termination sequence. In addition to

targeting highly conserved HCV sequences, the ability to co-deliver two, three, four, five or more different RNA effector molecules radically reduces the ability of the virus to develop escape mutants. While "cocktail" pharmaceutical preparations including multiple active components can be formulated, dsRNA
5 expression constructs provide an attractive delivery vehicle for accomplishing such co-delivery of a plurality of different antiviral effector molecules by a single pharmaceutical entity. The manufacturing and regulatory advantages of such an approach are readily apparent.

In certain embodiments, the expression construct encodes two or more
10 RNAs of the invention, such as 2, 3, 4, 5, or more double stranded RNA molecules, such as shRNAs. Thus, the expression construct may encode double stranded RNAs, such as shRNA hairpins, specific for one or more of HCV Conserved Regions 1, 2 or 5, or ATR-1, ATR-2, ATR-3, or ATR-4.

In these embodiments, the invention provides one or more expression
15 constructs, which collectively encode from two to five or more double-stranded RNA effector molecules, such as from two to five or more double-stranded RNA effector molecules as described herein. In certain embodiments, each expression construct encodes two, three, or four double-stranded RNA effector molecules, and preferably shRNAs. The encoding sequences, which may each
20 be operably linked to a promoter, may each encode: (1) a sequence of at least 19 nucleotides having at least 90% identity with a nucleotide sequence within HCV Conserved Region 1 (SEQ ID NO: 2), HCV Conserved Region 2 (SEQ ID NO: 3), or HCV Conserved Region 5 (SEQ ID NO: 4), ATR-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89); (2) its
25 complementary sequence; and optionally (3) a loop sequence.

For example, the expression construct of the invention may encode a plurality of double-stranded RNA effector molecules having double-stranded regions comprising (or consisting essentially of) a sequence independently selected from SEQ ID NOS: 5-42 and its complementary sequence. In one
30 embodiment, the construct encodes: a first effector molecule comprising or consisting essentially of (or consisting of) the nucleotide sequence of SEQ ID NO: 11, a second effector molecule comprising or consisting essentially of (or consisting of) the nucleotide sequence of SEQ ID NO: 19, a third effector molecule comprising or consisting essentially of (or consisting of) the nucleotide

sequence of SEQ ID NO: 22, and a fourth effector molecule comprising or consisting essentially of (or consisting of) the nucleotide sequence of SEQ ID NO: 33. One, two, three, or four of these double stranded RNA effector molecules may be in the form of short-hairpin RNAs, that is, having sense and anti-sense strands connected by a short loop sequence as described herein. Thus, in these
5 embodiments, the expression construct of the invention may encode a plurality of dsRNA effector molecules independently selected from SEQ ID NOS: 44-81. The construct may encode two, three, or four double-stranded RNA effector molecules represented by the sequences: SEQ ID NO: 50, SEQ ID NO: 58, SEQ
10 ID NO: 61, and SEQ ID NO: 72.

The expression construct may encode an RNA molecule of the invention and its complementary stand separately, that is from separate promoters (e.g., separate expression cassettes). In this embodiment, the double-stranded molecule is produced intracellularly upon expression of the RNAs and
15 subsequent hybridization. Typically, with expressed interfering RNA (eiRNA), the dsRNA is expressed in the first transfected cell from an expression vector. In such a vector, the sense strand and the antisense strand of the dsRNA may be transcribed from the same nucleic acid sequence using e.g., two convergent promoters at either end of the nucleic acid sequence or separate promoters
20 transcribing either a sense or antisense sequence. Alternatively, two plasmids can be cotransfected, with one of the plasmids designed to transcribe one strand of the dsRNA while the other is designed to transcribe the other strand.

In certain embodiments, the construct is an expression construct suitable for expression (e.g., transcription) of the encoded RNA *in vitro* or *in vivo*. In these
25 embodiments, the construct encodes an RNA molecule of the invention operably linked to a promoter sequence. Suitable promoter sequences suitable for run-off transcription *in vitro* are known, and include T3 and T7 promoters. In other embodiments, the expression construct encodes an RNA molecule of the invention operably linked to a promoter suitable for expression (e.g., transcription)
30 of the RNA in a mammalian cell. The mammalian cell may be in culture, or may be a patient's cell, e.g., a patient afflicted with or at risk of acquiring HCV. Promoters suitable for RNA expression in a mammalian cell are known, and include those described in WO 2006/033756, which is hereby incorporated by reference. For example, the construct of the invention may contain a DNA

sequence encoding an RNA molecule of the invention operably linked to an RNA polymerase III promoter, a 7SK promoter, an H1 promoter, or a U6 promoter. Further, where the construct encodes a plurality of RNA molecules, each independently controlled by its own promoter, the promoters may be the same or different. In one aspect, an expression construct may comprise two, three, four or more 7SK 4A promoters as described in WO 2006/033756. Additional promoters may be selected from an RNA polymerase I promoter, an RNA polymerase II promoter, a T7 polymerase promoter, an SP6 polymerase promoter, a tRNA promoter, and a mitochondrial promoter.

10 Generally, vector-directed expression of short RNA effector molecules including short hairpin dsRNAs is most efficient when under the control of a mammalian promoter that the host cell naturally employs for expression of small RNA molecules. These promoters comprise the family of RNA Polymerase III promoters, including Type 1, Type 2, and Type 3 RNA Polymerase III promoters. Prototypical examples of promoters in each class are found in genes encoding 5s RNA (Type 1), various transfer RNAs (Type 2) and U6 small nuclear RNA (Type 3). Another promoter family (transcribed by RNA Polymerase I) is also dedicated in the cell to transcription of small structural RNAs; however, this family may be less diverse in sequence than the RNA Polymerase III promoters. Finally, RNA Polymerase II promoters are used in the transcription of the protein-coding messenger RNA molecules, as distinguished from the small structural and regulatory RNA mentioned above. The majority of promoter systems known in the art utilize RNA Polymerase II promoters, which may not be preferred for production of small RNAs. An exception may be shRNAs expressed by RNA polymerase II or III promoters in a miRNA context as taught in e.g. USSN 25 10/429,249 and PCT/US2007/81103, which is hereby incorporated by reference. RNA polymerase III promoter-based vectors containing one promoter have been described in the art (see, e.g., U.S. Patent No. 5,624,803, Noonberg et al., "In vivo oligonucleotide generator, and methods of testing the binding affinity of triplex forming oligonucleotides derived therefrom"), and a description of U6-based vector systems can be found in Lee et al., Nat. Biotechnol. 20:500-05 (2002). Yu et al., Proc. Natl. Acad. Sci. USA 99:6047-52 (2002), describe an expression system for short duplex siRNAs comprising a T7 and U6 promoter. Miyagishi and Taira, Nat. Biotechnol. 20:497-500 (2002), describe expression

plasmids for short duplex siRNAs comprising expression cassettes containing tandem U6 promoters, each transcribing either the sense or the antisense strand of an siRNA, which are then annealed to form duplex siRNAs.

Where it is desired to deliver short dsRNAs, multiple RNA polymerase III promoter expression constructs (as taught in WO 06/033756, which is hereby incorporated by reference), may be used in accordance with the invention. The multiple RNA polymerase III promoters may be utilized in conjunction with promoters of other classes, including RNA polymerase I promoters, RNA polymerase II promoters, etc. Preferred in some applications are the Type III RNA pol III promoters including U6, H1, and 7SK, which exist in the 5' flanking region, include TATA boxes, and lack internal promoter sequences. A preferred 7SK promoter is the 7SK 4A promoter variant taught in WO 06/033756, the nucleotide sequence of which is hereby incorporated by reference. In such expression constructs each promoter may be designed to control expression of an independent RNA expression cassette, e.g., a shRNA expression cassette. Such multiple RNA polymerase III promoter expression constructs are suitable for expression of multiple, e.g., three, four, five, or more anti-HCV shRNA effector molecules of the invention. Each dsRNA effector molecule, e.g., hairpin dsRNA, may be transcribed from its own promoter or one or more promoters may be engineered to each transcribe a single RNA strand encoding a series or "gang" of two, three or more shRNA molecules separated by single-stranded regions. RNA Pol III promoters may be especially beneficial for expression of small engineered RNA transcripts, because RNA Pol III termination occurs efficiently and precisely at a short run of thymine residues in the DNA coding strand, without other protein factors. T₄ and T₅ are the shortest Pol III termination signals in yeast and mammals, with oligo (dT) terminators longer than T₅ being rare in mammals. Accordingly, the multiple polymerase III promoter expression constructs of the invention may also include an appropriate oligo (dT) termination signal, i.e., a sequence of 4, 5, 6 or more Ts, operably linked 3' to each RNA Pol III promoter cassette in the DNA coding strand. That is, a DNA sequence encoding one or more RNA effector molecules, e.g., a dsRNA hairpin or RNA stem-loop structure to be transcribed, is inserted between the Pol III promoter and the termination signal.

The invention provides means for delivering to a host cell sustained amounts of 2, 3, 4, 5, or more different antiviral dsRNA hairpin molecules (e.g., specific for 2, 3, 4, 5, or more different viral sequences or elements), in a genetically stable mode, so as to inhibit viral replication while preventing, decreasing, or delaying generation of viral escape mutants, and without evoking a dsRNA stress response. In accordance with this aspect, each dsRNA hairpin may be expressed from an expression construct, and controlled by e.g. an RNA polymerase III promoter. In some such embodiments a single RNA polymerase promoter, e.g. a pol II or pol III promoter, may express a plurality of dsRNA hairpins. In some such embodiments the dsRNA hairpins will be present within 5' and 3' flanking miRNA sequences.

Thus, the expression constructs of the invention provide a convenient means for delivering a multi-drug regimen comprising several different RNAs of the invention to a cell or tissue of a host vertebrate organism, thereby potentiating the anti-viral activity, and reducing the likelihood that multiple independent mutational events will produce resistant virus. This provides an important advantage in countering viral variation both within human and animal host populations and temporally within a host due to mutation events.

In certain embodiments, the expression construct contains at least two expression cassettes, each expression cassette directing the expression of a shRNA independently selected from SEQ ID NOS: 44-81, such as SEQ ID NOS: 50, 58, 61, and 72. Each expression cassette may independently comprise at least one promoter, e.g., an RNA polymerase III promoter selected from a U6 promoter, a 7SK promoter, an H1 promoter, and an MRP promoter. For example, each expression cassette may comprise a 7SK promoter driving the expression of at least one double-stranded RNA effector molecule, and an RNA pol III termination signal.

Pharmaceutical Compositions

The RNAs and constructs of the invention may be formulated as pharmaceutical compositions, comprising, in addition to effective amounts of the RNA(s) or construct(s) necessary to produce the desired biological effect, pharmaceutically acceptable carriers. Such carriers may comprise, for example, agents for facilitating the transfection of mammalian cells, which are well known.

Exemplary transfection agents and compositions, which may be used in accordance with the present invention, are Lipofectamine2000™ (Invitrogen, Carlsbad, Cal.), as well as those reviewed and described in US 2006/0084617 published April 20, 2006, which is hereby incorporated by reference in its entirety.

5 See also, the methods and compositions for delivery of nucleic acids as taught in in WO 2006/033756, the teaching of which is hereby incorporated herein in its entirety. Nucleic acids such as shRNAs may also be delivered to distal organs such as the liver by transfecting skeletal muscle cells (e.g., injection, injection/electroporation, or hydrodynamic vessel delivery) with an expression

10 vector as taught in USSN 11/935,925 (PCT/US2007/83805), the teaching of which is hereby incorporated by reference in its entirety.

In various embodiments, the pharmaceutical composition includes about 1 ng to about 20 mg of nucleic acid, e.g., RNA, DNA, plasmids, viral vectors, recombinant viruses, or mixtures thereof (as described above), which provide the

15 desired amounts of the nucleic acid molecules. In some embodiments, the composition contains about 10 ng to about 10 mg of nucleic acid, about 0.1 mg to about 500 mg, about 1 mg to about 350 mg, about 25 mg to about 250 mg, or about 100 mg of nucleic acid. Those of skill in the art of clinical pharmacology can readily arrive at appropriate dosing schedules with routine experimentation.

20 Other suitable carriers include, but are not limited to, saline, buffered saline, dextrose, water, glycerol, ethanol, and combinations thereof. The pharmaceutical composition generally contains one or more pharmaceutically acceptable additives suitable for the selected route and mode of administration. These compositions may be administered by, without limitation, any parenteral

25 route including intravenous (IV) or intra-arterial (IA) (including hydrodynamic delivery methods in which increasing intravessel pressure increases transfection of the surrounding cells), intramuscular (IM), intramuscular/electroporation, subcutaneous (SC), intradermal, intraperitoneal, intrathecal, as well as topically, orally, and by mucosal routes of delivery such as intranasal, inhalation, rectal,

30 vaginal, buccal, and sublingual. Generally, the pharmaceutical compositions of the invention are prepared for administration to mammalian subjects including primates and humans, and are in the form of liquids, including sterile, non-pyrogenic liquids for injection, emulsions, powders, aerosols, tablets, capsules, enteric coated tablets, or suppositories.

Methods for Inhibiting Expression of Target Polynucleotides

The invention further provides a method for silencing an HCV RNA in a mammalian cell. As used herein, the term silencing of an HCV RNA refers to reducing the abundance of the target polynucleotide in the cell, or in a patient
5 treated with a sequence-specific anti-HCV agent of the invention, whether via RNA stability and/or via replication rate of the HCV polynucleotide, including inhibitory effects on production, levels or persistence of positive and/or negative strand HCV RNA, including various qualitative and quantitative measures of HCV viral load, e.g., quantitative PCR, transcription-mediated amplification (TMA), and
10 branched DNA assay (bDNA). HCV RNA silencing also includes inhibitory effects on viral protein expression. The target polynucleotide (e.g., HCV RNA) may be reduced by about 1/2, 1/5, 1/10, 1/100 as compared to a positive control cell (e.g., a cell infected with HCV). In some embodiments, the target polynucleotide is undetectable in cells transfected or patients treated with an agent, composition
15 (including pharmaceutical composition), or construct of the invention).

In this aspect, the invention comprises administering to a mammalian cell, such as a primate or human cell infected with HCV, at least one agent, composition, or construct of the invention, as described previously herein, including small-hairpin RNAs and including the encoding expression constructs.
20 The cell in such embodiments may be a cell culture, or may be an animal model or patient. In these embodiments, the invention provides a method of reducing levels of HCV RNA in a cell either *in vitro* or *in vivo*, as well as methods for reducing an HCV titer *in vitro* or *in vivo*. In a preferred aspect, the agent is capable of reducing HCV replication in a human hepatocyte in an *in vitro* HCV
25 infection/replication model such as the recently developed infectious cell culture systems, which more closely reflect actual human HCV infection. The JFH-1 HCV clone, utilized in the Examples below, which is able to replicate efficiently in Huh7 cells and can secrete infectious viral particles, represents one such improved model. See e.g., Zhong et al., Robust hepatitis C virus infection *in vitro*,
30 *PNAS*, 102, No. 26, pp 9294-9299, (2005). See also Yi M, Villanueva RA, Thomas DL, Wakita T, Lemon SM. Production of infectious genotype 1a hepatitis C virus (Hutchinson strain) in cultured human hepatoma cells. *Proc Natl Acad Sci USA* 2006; 103(7): 2310-2315. In accordance with these embodiments, the double-stranded RNA effector molecules may be introduced into the cell by

transforming or transfecting a cell with an expression construct of the invention, or alternatively by directly introducing the double stranded RNA.

The present invention further provides a method for treating, ameliorating or preventing HCV infection in a patient (e.g., a patient having, or at risk of acquiring an HCV infection), comprising administering to said patient, including primates such as humans, an effective amount of an agent, composition, or construct of the invention, suitable for triggering RNAi-mediated degradation of the patient. Thus, the method of the invention reduces viral replication in infected cells, and in certain embodiments, eliminated the infection. The presence of HCV RNA in the blood is considered to be an indication that the virus is actively replicating (reproducing and infecting new cells). In some embodiments, HCV viral load is reduced (by 10%, 20%, 50%, 75%, 90% or more) in an infected individual administered a sequence-specific dsRNA of the invention. In some aspects, viral load will be reduced from high to low levels (less than 2 million copies/mL), desirably to undetectable levels. It is thought that individuals with viral load below 400,000 IU/mL respond better to therapeutic agents than those with higher levels of virus.

The present invention may, in certain embodiments, employ the methods disclosed in USSN 11/935,925 (PCT/US2007/83805), "*In Vivo* Delivery of Double Stranded RNA to a Target Cell", which is hereby incorporated by reference in its entirety. Specifically, delivery into skeletal muscle cells of expression constructs encoding dsRNA(s) may result in targeted inhibition of gene expression in other organs and tissues of the body such as liver. The intramuscular delivery may be achieved in a variety of ways, including needle or needleless IM injection, IM injection/electroporation, and intravascular/hydrodynamic delivery. Without being bound by any particular theory, delivery of dsRNA to distal tissues such as liver cells, for example, may be mediated by extracellular vesicles (exovesicles) containing expressed dsRNA or injected siRNA or shRNA that bud from the surface of transfected muscle cells.

The HCV treated by the method of the invention (e.g., targeted by the agents, compositions, and constructs described herein) may be any genotype, subtype, or quasispecies, including genotypes 1a, 1b, 2a, 2b, 3a, 4 and 5, and combinations thereof. In certain embodiments, the HCV infection is non-responsive to interferon-based therapy, and/or other nucleoside analogs such as

ribavirin, or other antivirals, making the methods of the invention particularly desirable. In other embodiments, the agents, compositions, and constructs are administered before, during, or after interferon therapy, where necessary to control or eliminate infection.

5 In certain embodiments, the agents, compositions, and constructs of the invention are administered so as to inhibit viral replication without evoking a dsRNA stress response. Further, unlike traditional antiviral agents, the method of the present invention may reduce the likelihood that multiple independent
10 mutational events will produce resistant virus, thereby providing an important advantage in countering viral variation both within human and animal host populations and temporally within a host due to mutation events.

 Some dsRNA sequences, possibly in certain cell types and through certain delivery methods, may result in an interferon response. The methods of the invention may be performed so as not to trigger an interferon/PKR response,
15 for instance by using shorter dsRNA molecules between 20 to 25 base pairs, by expressing dsRNA molecules intracellularly, or by using other methods known in the art. See US Published Application 20040152117, which is herein incorporated by reference. For instance, one of the components of an interferon response is the induction of the interferon-induced protein kinase PKR. To
20 prevent an interferon response, interferon and PKR responses may be silenced in the transfected and target cells using a dsRNA species directed against the mRNAs that encode proteins involved in the response. Alternatively, interferon response promoters are silenced using dsRNA, or the expression of proteins or transcription factors that bind interferon response element (IRE) sequences is
25 abolished using dsRNA or other known techniques.

 By "under conditions that inhibit or prevent an interferon response or a dsRNA stress response" is meant conditions that prevent or inhibit one or more interferon responses or cellular RNA stress responses involving cell toxicity, cell death, an anti-proliferative response, or a decreased ability of a dsRNA to carry
30 out a PTGS event. These responses include, but are not limited to, interferon induction (both Type 1 and Type II), induction of one or more interferon stimulated genes, PKR activation, 2'5'-OAS activation, and any downstream cellular and/or organismal sequelae that result from the activation/induction of one or more of these responses. By "organismal sequelae" is meant any

effect(s) in a whole animal, organ, or more locally (e.g., at a site of injection) caused by the stress response. Exemplary manifestations include elevated cytokine production, local inflammation, and necrosis. Desirably the conditions that inhibit these responses are such that not more than 95%, 90%, 80%, 75%,
5 60%, 40%, or 25%, and most desirably not more than 10% of the cells undergo cell toxicity, cell death, or a decreased ability to carry out a PTGS event, compared to a cell not exposed to such interferon response inhibiting conditions, all other conditions being equal (e.g., same cell type, same transformation with the same dsRNA).

10 Apoptosis, interferon induction, 2'5'-OAS activation/induction, PKR induction/activation, anti-proliferative responses, and cytopathic effects are all indicators for the RNA stress response pathway. Exemplary assays that can be used to measure the induction of an RNA stress response as described herein include a TUNEL assay to detect apoptotic cells, ELISA assays to detect the
15 induction of alpha, beta and gamma interferon, ribosomal RNA fragmentation analysis to detect activation of 2'5'-OAS, measurement of phosphorylated eIF2a as an indicator of PKR (protein kinase RNA inducible) activation, proliferation assays to detect changes in cellular proliferation, and microscopic analysis of cells to identify cellular cytopathic effects. See, e.g., US Published Application
20 20040152117, which is herein incorporated by reference.

Methods for Making Agents and Compositions

The dsRNA molecules and constructs of the invention may be made using conventional molecular biology techniques, including standard gene cloning and
25 *in vitro* RNA synthesis protocols. Such methods are well known, and are described in Sambrook et al, Molecular Cloning: A Laboratory Manual, 2nd edition, Cold Spring Harbor Laboratory Press (1989); and Ausubel et al., ed. Current Protocols in Molecular Biology, John Wiley & Sons, Inc. (1987)). See also the methods taught in US 6,143,527, "Chain reaction cloning using a bridging oligonucleotide and DNA ligase".

30 The following examples are provided to describe and illustrate the present invention. As such, they should not be construed to limit the scope of the invention. Those in the art will well appreciate that many other embodiments also

fall within the scope of the invention, as it is described hereinabove and in the claims.

EXAMPLES

5 The following Examples are provided for illustration only.

 All possible 21-mer and 27-mer expressed shRNAs were constructed based on HCV Conserved Regions 1 (SEQ ID NO: 2), 2 (SEQ ID NO: 3) and 5 (SEQ ID NO: 4). Thus, for shRNAs targeting Conserved Region 1, twenty-six 21-
10 21-mers and sixteen 27-mers were constructed; for Conserved Region 2, thirty-nine 21-mers and thirty-three 27-mers were constructed; and for Conserved Region 5, twenty-eight 21-mers and sixteen 27-mers were constructed.

Example 1: Silencing HCV replication in a viral infection cell culture model

 A human liver-derived cell line such as the Huh7 cell line is transfected with an eiRNA plasmid expressing short-hairpin RNA (shRNA) molecules that
15 comprise sequences homologous and complementary to the identified conserved sequences in the HCV genome. Following internalization of the plasmid into hepatocytes and nuclear localization, transcription of the eiRNA plasmid from one or more RNA pol III promoters results in production of the HCV-specific shRNAs.

 The transfected cells are then infected with HCV JFH-1 virus. The JFH-1
20 clone is able to replicate efficiently in Huh7 cells and can secrete infectious viral particles. See e.g., Zhong et al., Robust hepatitis C virus infection *in vitro*, *PNAS*, 102, No. 26, pp 9294-9299, (2005).

 Using this model, cells were transfected with various eiRNA constructs and then infected with HCV virus. The cells were then monitored for loss of HCV
25 replication as described below.

 The following is an example of an experiment that was performed using eiRNA vectors encoding HCV sequences derived from GenBank accession number ABO47639. HCV sequences in these described eiRNA vectors were highly conserved sequences identified as described elsewhere herein. The
30 particular eiRNA backbone vector used for this experiment contained a 7SK promoter to drive expression of the encoded RNAs. Each vector encoded only one shRNA. The shRNA coding sequence was followed by an RNA pol III

termination sequence. Sequences of the 7SK promoter, RNA pol III termination signal, and encoded shRNAs are all shown at the end of the example. Similar vectors containing U6 promoters (another RNA pol III type 3 promoter) and RNA pol III termination signals are commercially available such as the “siLentGene-2 Cloning Systems” vector from Promega, Inc., Madison, Wis. One of ordinary skill

5

Experimental Procedure: Transfection.

Huh7 cells cultured in DMEM medium were seeded into 96-well plates at a density of 4×10^3 cells/well. All transfections were performed the day after cell seeding using Lipofectamine2000™ (Invitrogen, Carlsbad, Cal.) according to the manufacturer's directions. In this experiment, cells were transfected with 200 ng

10

of the eiRNA plasmids and were infected 12 hours later with HCV infectious viral clone JFH-1 at a multiplicity of infection (MOI) of 0.1ffu/cell. DNA was held constant/transfection at 200 ng. In experiments where less than 200 ng of the eiRNA plasmid was transfected, the remainder of the DNA is made up by including an inert plasmid DNA, pGL2-Basic (Promega, Madison Wis.) in amounts that brought the total DNA in the transfection to 200 ng. Prior to transfection, medium was removed from the cells and the cells washed with phosphate-buffered saline (PBS), followed by addition of 100 uL of DMEM

15

containing 10% Fetal Bovine Serum (FBS). The DNA/Lipofectamine2000 transfection mix was added to the cells and these were incubated at 37°C for 12 hours. The medium containing the transfection mix was then removed and the cells were washed once with PBS. At this point, 100 uL of DMEM containing 10% FBS was added to the cells. Infection with JFH-1 then commenced with the

20

addition of the virus at a MOI of 0.1 ffu/cell. All transfections were carried out in triplicate. Control groups were pGL2-basic alone (200 ng of pGL2-basic), a green fluorescent protein (GFP) plasmid instead of the eiRNA plasmid (200ng pGFP), and a group that was untransfected but was infected with JFH-1 at the same MOI.

30 Monitoring cells for loss of HCV replication.

At 48 hours following transfection, cells were monitored for the loss or reduction in HCV replication by measuring viral RNA by quantitative reverse-transcription PCR (qRT-PCR). The cells were lysed using Total RNA Lysis Buffer

(Applied Biosystems, Foster City, CA) and the RNA was harvested using an ABI Prism 6100 Nucleic Acid Prep Station (Applied Biosystems). HCV RNA was quantified by qRT-PCR using a SYBR-green labeled probe. The PCR primers used to amplify a 75 bp product corresponding the 5' UTR of the genotype 2a HCV genome (GenBank AB047639) were 5'-TCTGCGGAACCGGTGAGTA-3' (sense) (SEQ ID NO: 82) and 5'-TCAGGCAGTACCACAAGGC-3' (antisense) (SEQ ID NO: 83). The PCR primers used to amplify a 225 bp product of the human GAPDH coding sequence (GenBank NM002046) were 5'-GAAGGTGAAGGTCGGAGTC -3' (sense) (SEQ ID NO: 84) and 5'-GAAGATGGTGATGGGATTTTC -3' (antisense) (SEQ ID NO: 85). HCV and GAPDH transcript levels were determined relative to a standard curve comprised of serial dilutions of plasmid containing the HCV cDNA or the human GAPDH gene. HCV copies per ug of total cellular RNA were normalized to GAPDH transcript levels using a modified comparative threshold cycle method 2^{-ddCt} (2^{-ddCt}); where ddCt is the difference between the Average GAPDH and the GAPDH of the individual sample. Fold reduction in HCV copy numbers was calculated as: (average number of HCV copies per ug total cellular RNA in control transfected samples) ÷ (average number of HCV copies per ug total cellular RNA in eiRNA transfected samples).

20 Results:

A number of the shRNA plasmids transfected led to a decrease in HCV replication as compared to negative controls. The results in Table 1 are presented as the average of three plates and are shown as both fold-inhibition and the equivalent percent reduction.

25 A total of 135 eiRNAs (plasmid expressed shRNAs) containing 21-mer double-stranded stem sequences and 118 eiRNAs containing 27-mer double-stranded stem sequences were tested in the assay. The eiRNAs showing a 4-fold or greater inhibition are listed in Table 1 along with a representative selection of eiRNAs showing less than 4-fold inhibition. The eiRNA vectors encode the HCV sequences listed in Table 2. The sequences of the shRNAs are shown as well as the map coordinates of the sense sequence (found 3' to the underlined 9 nt loop sequence) relative to the HCV JFH-1 clone, GenBank accession number ABO47639 (SEQ ID NO: 1). The sequences of the encoded shRNAs include an antisense HCV sequence followed by the loop sequence (underlined in Table 1)

followed by a second HCV sequence, which is the complement to the first HCV sequence. (It will be understood that either the antisense or sense sequence may be located 5' to the loop sequence, i.e., 5'-antisense-loop-sense-3' or 5'-sense-loop-antisense-3'.) The loop structure does not need to be a fixed sequence or length, and several loop sequences were used with no significant impact on the functioning of the eiRNA construct. The second HCV sequence is followed by a series of T residues, e.g., 1, 2, 3, or more Ts, preferably at least 4 or 5 Ts, that function as the termination signal for RNA pol III. These T nucleotides are not included in Table 2.

10

Table 1.

<u>eiRNA</u>	<u>Fold inhibition</u>	<u>Percent reduction</u>
HCV 5' 21-2	1	0
HCV 5' 21-16	1	0
HCV 5' 21-50	1	0
HCV 5'-21-55	5	80
HCV 5'-21-56	5	80
HCV 5'-21-57	5	80
HCV 5'-21-61	7	85.7
HCV 5' 21-63	4	75
HCV 5' 21-70	1	0
HCV 5' 21-73	1	0
HCV 5' 21-88	5	80
HCV 5' 21-89	5	80
HCV 5' 21-90	4	75
HCV 5' 21-92	7	85.7
HCV 5' 21-94	9	88.9
HCV 5'-21-122	4	75
HCV 5'-21-123	4	75
HCV 5'-21-124	7	85.7

<u>eiRNA</u>	<u>Fold inhibition</u>	<u>Percent reduction</u>
HCV 5'-21-125	5	80
HCV 5'-21-126	8	87.5
HCV 5'-21-127	7	85.7
HCV 5'-21-128	8	87.5
HCV 5'-21-129	4	75
HCV 5'-21-130	6	83.3
HCV 5'-21-131	3	67
HCV 5'-21-133	12	91.7
HCV 5'-21-134	9	88.9
HCV 5'-21-135	10	90
HCV 5' 27-1	1	0
HCV 5' 27-8	4	75
HCV 5' 27-9	4	75
HCV 5' 27-12	1	0
HCV 5'-27-16	8	87.5
HCV 5' 27-45	1	0
HCV 5'-27-53	4	75
HCV5'-27-73	1	0
HCV 5' 27-111	1	0

NUCL-029/01WO

<u>Sequence (5' antisense-loop-sense 3')</u>	<u>nucleotide coordinates</u> <u>sense sequence relative</u> <u>GenBank No. AB047639</u>
TAGTTCCTCACAGGGGAGTGA <u>agagaactt</u> TCACTCCCCTGTGAGGAACTA	36-56
CCGGTCCGCAGACCACTATG <u>agagaactt</u> CATAGTGGTCTGCGGAACCGG	50-70
TATGGCTCTCCCGGAGGGG <u>agagaactt</u> CCCCCTCCCGGAGAGCCATA	121-141
ACCACTATGGCTCTCCCGGA <u>agagaactt</u> TCCCGGAGAGCCATAGTGGT	126-146
GACCACTATGGCTCTCCCGG <u>agagaactt</u> CCCGGAGAGCCATAGTGGTC	127-147
AGACCACTATGGCTCTCCCGG <u>agagaactt</u> CCGGGAGAGCCATAGTGGTCT	128-148
CCGCAGACCACTATGGCTCTC <u>agagaactt</u> GAGAGCCATAGTGGTCTGCGG	132-152

NUCL-029/01WO

Sequence (5' antisense-loop-sense 3')	<u>nucleotide coordinates</u> <u>sense sequence relative</u> GenBank No. AB047639
TTCCGCAGACCACTATGGCTC <u>agagaactt</u> GAGCCATAGTGGTCTGCGGAA	134-154
TCACCGGTTCCGCAGACCACT <u>agagaactt</u> AGTGGTCTGCGGAACCGGTGA	141-161
TACTACCGGTTCCGCAGACC <u>agagaactt</u> GGTCTGCGGAACCGGTGAGTA	144-164
GCAGTACCACAAGGCCTTTC <u>agagaactt</u> CGAAAGGCCTTGTGGTACTGC	271-291
GGCAGTACCACAAGGCCTTTC <u>agagaactt</u> GAAAGGCCTTGTGGTACTGCC	272-292
AGGCAGTACCACAAGGCCTTTC <u>agagaactt</u> AAAGGCCTTGTGGTACTGCCT	274-294
TCAGGCAGTACCACAAGGCCT <u>agagaactt</u> AGGCCTTGTGGTACTGCCTGA	275-295
TATCAGGCAGTACCACAAGGC <u>agagaactt</u> GCCTTGTGGTACTGCCTGATA	277-297

NUCL-029/01WO

<u>Sequence (5' antisense-loop-sense 3')</u>	<u>nucleotide coordinates</u> <u>sense sequence relative</u> <u>GenBank No. AB047639</u>
AGACCTCCCGGGGCACTCGCA <u>agagaactt</u> TGCGAGTGCCCCGGGAGGTCT	305-325
GAGACCTCCCGGGGCACTCGC <u>agagaactt</u> GCGAGTGCCCCGGGAGGTCTC	306-326
CGAGACCTCCCGGGGCACTCG <u>agagaactt</u> CGAGTGCCCCGGGAGGTCTCG	307-327
ACGAGACCTCCCGGGGCACTC <u>agagaactt</u> GAGTGCCCCGGGAGGTCTCGT	308-328
TACGAGACCTCCCGGGGCACT <u>agagaactt</u> AGTGCCCCGGGAGGTCTCGTA	309-329
CTACGAGACCTCCCGGGGCAC <u>agagaactt</u> GTGCCCCGGGAGGTCTCGTAG	310-330
TCTACGAGACCTCCCGGGGCA <u>agagaactt</u> TGCCCCGGGAGGTCTCGTAGA	311-331

NUCL-029/01WO

<u>Sequence (5' antisense-loop-sense 3')</u>	<u>nucleotide coordinates</u> <u>sense sequence relative</u> <u>GenBank No. AB047639</u>
GTCTACGAGACCTCCCGGGG <u>Cagagaactt</u> GCCCCGGGAGGTCTCGTAGAC	312-332
GGTCTACGAGACCTCCCGGGG <u>agagaactt</u> CCCCGGGAGGTCTCGTAGACC	313-333
CGGTCTACGAGACCTCCCGGG <u>agagaactt</u> CCCCGGGAGGTCTCGTAGACCG	314-334
ACGGTCTACGAGACCTCCCGG <u>agagaactt</u> CCGGGAGGTCTCGTAGACCGT	315-335
CACGGTCTACGAGACCTCCC <u>agagaactt</u> CGGGAGGTCTCGTAGACCGTG	316-336
GCACGGTCTACGAGACCTCCC <u>agagaactt</u> GGGAGGTCTCGTAGACCGTGC	317-337
TGCACGGTCTACGAGACCTCC <u>agagaactt</u> GGAGGTCTCGTAGACCGTGCA	318-338
GACAGTAGTTCCTCACAGGGGAGTGAT <u>agagaactt</u> ATCACTCCCCTGTGAGGAACTACTGTC	35-61

NUCL-029/01WO

<u>Sequence (5' antisense-loop-sense 3')</u>	<u>nucleotide coordinates</u> <u>sense sequence relative</u> <u>GenBank No. AB047639</u>
GCGTGAAGACAGTAGTTCCTCACAGG <u>Gagagaactt</u> CCCTGTGAGGAACTACTGTCTTCACGC	42-68
TGGTGAAGACAGTAGTTCCTCACAGG <u>Gagagaactt</u> CCCTGTGAGGAACTACTGTCTTCACGCA	43-69
TTCTGCGTGAAGACAGTAGTTCCTCAC <u>Gagagaactt</u> GTGAGGAACTACTGTCTTCACGCAGAA	46-72
CGCTTTCTGCGTGAAGACAGTAGTTC <u>Gagagaactt</u> GAACTACTGTCTTCACGCAGAAAGCG	50-76
GACCACTATGGCTCTCCCGGGAGGGG <u>Gagagaactt</u> CCCCCTCCCGGAGAGCCATAGTGGTC	122-148
GTTCCGAGACCACTATGGCTCTCCCG <u>Gagagaactt</u> CGGAGAGCCATAGTGGTCTGCGGAAC	129-155
CAATCCGGTGTACTCACCGGTTCCGC <u>Gagagaactt</u> GCGGAACCGGTGAGTACACCGGAATTG	149-175

NUCL-029/01WO

	<u>nucleotide coordinates</u> <u>sense sequence relative</u>
<u>Sequence (5' antisense-loop-sense 3')</u>	<u>GenBank No. AB047639</u>
TCTACGAGACCTCCCGGGGCACTCGCA <u>agagaact</u> TGCGAGTGCCCCGGGAGGTCTCGTAGA	305-331

Table 3.

<u>eiRNA</u>	<u>Sense sequence</u>
HCV5'-21-2 (SEQ ID NO: 5)	tcactcccctgtgaggaacta
HCV5'-21-16 (SEQ ID NO: 6)	ggaactactgtcttcacgcag
HCV5'-21-50 (SEQ ID NO: 7)	ccccctcccgaggagagccata
HCV5'-21-55 (SEQ ID NO: 8)	tccgggagagccatagtggt
HCV5'-21-56 (SEQ ID NO: 9)	ccgggagagccatagtggtc
HCV5'-21-57 (SEQ ID NO: 10)	ccgggagagccatagtggtct
HCV5'-21-61 (SEQ ID NO: 11)	gagagccatagtggtctgcgg
HCV5'-21-63 (SEQ ID NO: 12)	gagccatagtggtctgcggaa
HCV5'-21-70 (SEQ ID NO: 13)	agtggtctgcggaaccggtga
HCV5'-21-73 (SEQ ID NO: 14)	ggtctgcggaaccggtgagta
HCV5'-21-88 (SEQ ID NO: 15)	cgaaaggccttggtactgc
HCV5'-21-89 (SEQ ID NO: 16)	gaaaggccttggtactgcc
HCV5'-21-90 (SEQ ID NO: 17)	aaaggccttggtactgcct
HCV5'-21-92 (SEQ ID NO: 18)	aggccttggtactgcctga

<u>eiRNA</u>	<u>Sense sequence</u>
HCV5'-21-94 (SEQ ID NO: 19)	gccttggtactgcctgata
HCV5'-21-122 (SEQ ID NO: 20)	tcgagtgccccgggaggctct
HCV5'-21-123 (SEQ ID NO: 21)	gcgagtgccccgggaggctctc
HCV5'-21-124 (SEQ ID NO: 22)	cgagtgccccgggaggctctcg
HCV5'-21-125 (SEQ ID NO: 23)	gagtgccccgggaggctctcgt
HCV5'-21-126 (SEQ ID NO: 24)	agtccccgggaggctctcgta
HCV5'-21-127 (SEQ ID NO: 25)	gtgccccgggaggctctcgtag
HCV5'-21-128 (SEQ ID NO: 26)	tgccccgggaggctctcgtaga
HCV5'-21-129 (SEQ ID NO: 27)	gccccgggaggctctcgtagac
HCV5'-21-130 (SEQ ID NO: 28)	ccccgggaggctctcgtagacc
HCV5'-21-131 (SEQ ID NO: 29)	cccgggaggctctcgtagaccg
HCV5'-21-132 (SEQ ID NO: 30)	ccgggaggctctcgtagaccgt
HCV5'-21-133 (SEQ ID NO: 31)	cgggaggctctcgtagaccgtg
HCV5'-21-134 (SEQ ID NO: 32)	gggaggctctcgtagaccgtgc
HCV5'-21-135	ggaggctctcgtagaccgtgca

<u>siRNA</u>	<u>Sense sequence</u>
(SEQ ID NO: 33)	
HCV5'-27-1 SEQ ID NO: 34)	atcactcccctgtgaggaaactactgtc
HCV5'-27-8 (SEQ ID NO 35)	ccctgtgaggaaactactgtcttcacgc
HCV5'-27-9 (SEQ ID NO: 36)	cctgtgaggaaactactgtcttcacgca
HCV5'-27-12 (SEQ ID NO: 37)	gtgaggaaactactgtcttcacgcagaa
HCV5'-27-16 (SEQ ID NO: 38)	ggaactactgtcttcacgcagaaagcg
HCV5'-27-45 (SEQ ID NO: 39)	ccccctccgggagagccatagtggtc
HCV5'-27-53 (SEQ ID NO: 40)	cgggagagccatagtggctgcggaac
HCV5'-27-73 (SEQ ID NO: 41)	gcggaaccgggtgagtacaccggaattg
HCV5'-27-111 (SEQ ID NO: 42)	tgcgagtgtccccgggagggtctcgtaga

Preferred for use in anti-HCV dsRNA effector molecules of the invention are the following conserved and actively inhibitory HCV sequences:

HCV-5-21-56, SEQ ID NO: 9, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID. NO: 48;

HCV-5-21-61, SEQ. ID NO: 11, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO: 50;

HCV-5-21-90, SEQ. ID NO: 17, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO: 56;

HCV-5-21-94, SEQ. ID NO: 19, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO: 58;

HCV-5-21-124, SEQ. ID NO:22, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO: 61;

5 HCV-5-21-128, SEQ. ID NO: 26, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO: 65;

HCV-5-21-133, SEQ. ID NO:31, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO:70;

10 HCV-5-21-134, SEQ. ID NO: 32, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO:71;

HCV-5-21-135, SEQ. ID NO:33, and its complement, which may be utilized as a duplex dsRNA, or as an shRNA, such as that of SEQ. ID NO:72;

In preferred embodiments, two, three, four, five or more of the above-identified anti-HCV dsRNA effector molecules are administered concomitantly to a human cell infected with HCV. In some aspects, an expression construct or constructs encoding such a plurality of dsRNA effector molecules is/are administered concomitantly to a human cell infected with HCV. In some aspects said two or more dsRNA effector molecules will comprise in double-stranded conformation two or more sequences selected from SEQ ID NOS: 9, 11, 17, 19, 20 22, 26, 31, 32, and 33. In some aspects said two or more dsRNA effector molecules will comprise two or more sequences selected from SEQ ID NOS: 48, 50, 56, 58, 61, 65, 70, 71, and 72.

The actively inhibitory HCV sequences map to four conserved and highly active target regions (ATR) (**see Figure 1**):

25 ATR-1: 5'-CCCTGTGAGGAACTACTGTCTTCACGCAGAA-3' (SEQ ID NO: 86), mapping to nucleotide coordinates 42 to 76 of GenBank Accession No. AB047639, found within Conserved Region 1 (SEQ ID NO: 2).

30 ATR-2: 5'-TCCCGGGAGAGCCATAGTGGTCTGCGGAA-3' (SEQ ID NO: 87), mapping to nucleotide coordinates 126 to 154 of GenBank Accession No. AB047639, found within Conserved Region 2 (SEQ ID NO: 3).

ATR-3: 5'-CGAAAGGCCTTGTGGTACTGC-3' (SEQ ID NO: 88), mapping to nucleotide coordinates 271 to 297 of GenBank Accession No. AB047639, found within Conserved Region 5 (SEQ ID NO: 4).

ATR-4: 5'-TGCGAGTGCCCCGGGAGGTCTCGTAGACCGTGCA3' (SEQ ID NO: 89), mapping to nucleotide coordinates 305 to 338 of GenBank Accession No. AB047639, found within Conserved Region 5 (SEQ ID NO: 4).

Example 2: Silencing HCV replication using a plasmid expressing multiple eiRNA molecules.

10 Plasmids expressing multiple eiRNAs (shRNAs) (known as multi-cistronic plasmids) can be used to inhibit HCV replication in a cell culture model (as described in Example 1). In this example, sequences encoding four different shRNA molecules were sequentially cloned into a single plasmid expression vector, each operably linked to an individual RNA polymerase III promoter and
15 terminator. Each shRNA contained a different sequence homologous and complementary to the HCV genome that was previously shown to inhibit HCV replication. By expressing several different dsRNA molecules within an HCV infected cell or a human cell capable of HCV infection, the eiRNA expression vector represents a multi-drug regimen having an enhanced ability to inhibit HCV
20 replication and to prevent the development during therapy of HCV escape mutants.

We cloned the genes for four HCV-suppressive shRNAs into a single plasmid, each operably linked to a different RNA polymerase III promoter and terminator. The four shRNAs were 1) shRNA SEQ ID NO: 50, which includes
25 HCV 5'-21-61 (SEQ ID NO: 11) and its complement; 2) shRNA SEQ ID NO: 58, which includes HCV-5'-21-94 (SEQ. ID NO: 19) and its complement; 3) shRNA SEQ. ID NO: 61, which includes HCV 5'-21-124 (SEQ. ID NO: 22) and its complement; and 4) shRNA SEQ. ID NO: 72, which includes HCV 5'21-135 (SEQ. ID NO: 33) and its complement. (See HCV multi-cistronic plasmid HCV-
30 QJ, **Figure 2**). Each shRNA sequence was expressed from a different copy of the RNA pol III promoter 7SK 4A, although one or more other RNA pol III type 3 promoters such as 7SK, H1 and/or U6 could also be used. It may be desirable in some circumstances to select two, three, four or more different promoters to

express two, three, four or more different dsRNAs of the invention. The quad-cistron plasmid was constructed essentially as described in WO 2006/0033756 and the 7SK 4A promoter utilized is also described therein. The eiRNAs were transcribed following intracellular uptake of the plasmid.

5 In the same manner as HCV-QJ (described above), other multi-cistronic plasmids (e.g., including quad-cistron plasmids HVC-QF, HCV-QG, HCV-QH, and HCV-QK), were constructed using different combinations of eiRNAs (anti-HCV shRNAs described herein) that had each been shown to be suppressive of HCV replication in their original mono-cistronic forms (see Table 4 below).

10 Such a mono-cistronic HCV eiRNA expression plasmid (HCV 5' 21-61 eiRNA plasmid, which expresses shRNA SEQ ID NO: 50, which includes HCV 5'-21-61 (SEQ ID NO: 11) and its complement) is illustrated in **Figure 3**.

Note that any number of combinations, e.g., two, three, four, five or more different suppressive eiRNAs, can constitute the multi-cistronic eiRNA plasmids of the invention. While a combination of two, three, four or more of such different dsRNA effector molecules (e.g., duplex and/or hairpin dsRNAs) may be administered concomitantly to a mammalian cell as a "cocktail" of exogenously generated RNAs, it is desirable in some aspects to express them within a target mammalian cell from a multi-cistronic plasmid as described herein.

20

Table 4.

Multi-cistronic plasmid

shRNA sequence and corresponding HCV sequence

HCV-QF	HCV 5' 21-56 (SEQ. ID. NO: 9) and its complement, expressed as shRNA SEQ ID NO: 48
	HCV 5' 21-135 (SEQ. ID NO:33) and its complement, expressed as shRNA SEQ. ID NO:72
	HCV 5' 21-133 (SEQ. ID NO:31) and its complement, expressed as shRNA SEQ. ID NO:70
	HCV 5' 21-61 (SEQ. ID NO: 11) and its complement, expressed

as shRNA SEQ. ID NO: 50

HCV-QG

HCV-5-21-94 (SEQ. ID NO: 19) and its complement, expressed as shRNA SEQ. ID NO: 58

HCV 5' 21-135 (SEQ. ID NO:33) and its complement, expressed as shRNA SEQ. ID NO:72

HCV 5' 21-133 (SEQ. ID NO:31) and its complement, expressed as shRNA SEQ. ID NO:70

HCV 5' 21-61 (SEQ. ID NO: 11) and its complement, expressed as shRNA SEQ. ID NO: 50

HCV-QH

HCV-5-21-94 (SEQ. ID NO: 19) and its complement, expressed as shRNA SEQ. ID NO: 58

HCV-5-21-134 (SEQ. ID NO: 32) and its complement, expressed as shRNA SEQ. ID NO:71

HCV-5-21-124 (SEQ. ID NO:22) and its complement, expressed as shRNA, SEQ. ID NO: 61

HCV 5' 21-135 (SEQ. ID NO:33) and its complement, expressed as shRNA SEQ. ID NO:72

HCV-QJ

HCV-5-21-94 (SEQ. ID NO: 19) and its complement, expressed as shRNA SEQ. ID NO: 58

HCV 5' 21-61 (SEQ. ID NO: 11) and its complement, expressed as shRNA SEQ. ID NO: 50

HCV-5-21-124 (SEQ. ID NO:22)and its complement, expressed as shRNA, SEQ. ID NO: 61

HCV 5' 21-135 (SEQ. ID NO: 33) and its complement, expressed as shRNA SEQ. ID NO:72

HCV-QK	HCV-5-21-128 (SEQ. ID NO: 26) and its complement, expressed as shRNA SEQ. ID NO: 65
	HCV-5-21-134 (SEQ. ID NO: 32) and its complement, expressed as shRNA SEQ. ID NO:71
	HCV 5' 21-133 (SEQ. ID NO: 31) and its complement, expressed as shRNA SEQ. ID NO: 70
	HCV 5' 21-135 (SEQ. ID NO: 33) and its complement, expressed as shRNA SEQ. ID NO:72

The methods that were utilized for transfection of the multi-cistronic plasmids into Huh7 cells, the subsequent infection of these cells with HCV and the quantification of HCV mRNA (as a measure of the suppression of HCV replication) were carried out as described in Example 1 above.

Results.

The suppressive activities that were observed for each of the multi-cistronic plasmids are shown in Table 5. Not only did the use of multiple different dsRNA effector molecules produce a very high level of inhibition of HCV replication, we expect that use of such combinations of different sequence-specific inhibitors will have enhanced ability to prevent the development of HCV escape mutants during therapy.

Table 5.

Multi-cistronic plasmid	Fold-inhibition	Percent reduction
HCV-QF	12	91.7
HCV-QG	11	90.9
HCV-QH	20	95
HCV-QJ	28	96.4
HCV-QK	25	96

All publications, patents and patent applications discussed herein are incorporated herein by reference. While in the foregoing specification this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be
5 apparent to those skilled in the art that the invention is susceptible to additional embodiments and that certain of the details described herein may be varied considerably without departing from the basic principles of the invention.

WHAT IS CLAIMED IS:

1. A method for silencing an hepatitis C virus (HCV) RNA in a mammalian cell comprising administering to said cell at least one double-stranded RNA effector molecule or complex, comprising: (1) a sequence of at least 19 consecutive nucleotides having at least 90% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), or Conserved Region 5 (SEQ ID NO: 4); active target region (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89); and (2) its complementary sequence.
2. The method of claim 1, wherein the sequence of at least 19 consecutive nucleotides has at least 95% or 100% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), or Conserved Region 5 (SEQ ID NO: 4), active target region (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89).
3. The method of claim 1 or 2, wherein the sequence of at least 19 nucleotides is selected from within nucleotides 42-76, 126-154, or 271-297 of SEQ ID NO: 1 (GenBank accession number AB047639).
4. The method of claim 1, 2, or 3, wherein the sequence of at least 19 nucleotides is selected from within nucleotides 305-338 of SEQ ID NO: 1 (GenBank accession number AB047639).
5. The method of any one of claims 1-4, wherein the double-stranded RNA effector molecule or complex contains a double-stranded region of from 19 to 30 base pairs.
6. The method of any one of claims 1-4, wherein the double-stranded RNA effector molecule or complex contains a double-stranded region of from 21 to 27 base pairs.
7. The method of any one of claims 1-4, wherein the double-stranded RNA effector molecule or complex contains a double-stranded region of about 21 or about 27 base pairs.
8. The method of any one of claims 1-7, wherein the sequence of at least 19 nucleotides comprises or consists essentially of a sequence selected from SEQ

ID NOS: 5-42, more preferably from SEQ ID NOS. 9, 11, 17, 19, 22, 26, 31, 32, and 33.

9. The method of any one of claims 1-8, wherein the sequence of at least 19 nucleotides and its complementary sequence are connected via a loop sequence.

5 10. The method of claim 9, wherein the loop sequence is from 5 to 20 nucleotides in length.

11. The method of claim 9, wherein the loop sequence is from 7 to 10 nucleotides in length.

10 12. The method of claim 9, wherein the loop sequence is about 9 nucleotides in length.

13. The method of claim 1, wherein at least one double-stranded RNA effector molecule has a sequence selected from SEQ ID NOS: 44-81, more preferably from SEQ ID NOS: 48, 50, 56, 58, 61, 65, 70, 71, and 72.

15 14. The method of any one of claims 1-13, wherein from two to five double-stranded RNA effector molecules or complexes are administered.

15. The method of claim 14, wherein the two to five double-stranded RNA effector molecules each comprise: (1) a sequence of at least 19 nucleotides having at least 90% to 100% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3),
20 Conserved Region 5 (SEQ ID NO: 4); (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), ATR-4 (SEQ ID NO: 89); SEQ ID NO: 9; SEQ, ID NO: 11; SEQ ID NO: 19; SEQ ID NO: 22; SEQ ID NO: 26; SEQ ID NO: 31; SEQ ID NO: 32; or SEQ ID NO: 33; and (2) its complementary sequence; and (3) a loop sequence.

25 16. The method of claim 15, wherein at least four double-stranded RNA effector molecules are administered, wherein: 1) a first effector molecule comprises the nucleotide sequence of SEQ ID 11, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 19, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 22, and a fourth effector
30 molecule comprises the nucleotide sequence of SEQ ID NO: 33; or 2) a first effector molecule comprises the nucleotide sequence of SEQ ID 9, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 11, a third

effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33, or 3) a first effector molecule comprises the nucleotide sequence of SEQ ID 11, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 19, 5 a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33; or 4) a first effector molecule comprises the nucleotide sequence of SEQ ID 19, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 22, a third effector molecule comprises the nucleotide sequence of SEQ ID 10 NO: 32, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33; or 5) a first effector molecule comprises the nucleotide sequence of SEQ ID 26, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 32, and a fourth effector molecule comprises the nucleotide 15 sequence of SEQ ID NO: 33 .

17. The method of claim 16, wherein the four double-stranded RNA effector molecules are selected from SEQ. ID. NOS: 48, 50, 56, 58, 61, 65, 70, 71, and 72; or more preferably comprise the sequences: SEQ ID NO: 50, SEQ ID NO: 58, SEQ ID NO: 61, and SEQ ID NO: 72; or sequences: SEQ ID NO: 48, SEQ ID NO: 20 50, SEQ ID NO: 70, and SEQ ID NO: 72; or sequences SEQ ID NO: 50, SEQ ID NO: 58, SEQ ID NO: 70, and SEQ ID NO: 72; or sequences SEQ ID NO: 58, SEQ ID NO: 61, SEQ ID NO: 71, and SEQ ID NO: 72; or SEQ ID NO: 65, SEQ ID NO: 70, SEQ ID NO: 71, and SEQ ID NO: 72.18 The method of any one of claims 1-17, wherein said administering is accomplished by providing to the 25 mammalian cell at least one expression construct capable of expressing the double-stranded RNA effector molecule(s).

19. The method of claim 18, wherein the expression construct contains at least two expression cassettes, each expression cassette directing the expression of at least one double-stranded RNA effector molecule.

20. The method of claim 19, wherein each expression cassette independently comprises at least one RNA polymerase III promoter selected from a U6 promoter, a 7SK promoter, an H1 promoter, and an MRP promoter.

21. The method of claim 19, wherein each expression cassette comprises a 7SK promoter driving the expression of one double-stranded RNA effector molecule, and an RNA pol III termination signal.
22. The method of any one of claims 1-21, wherein the mammalian cell is a human cell.
23. An agent for silencing hepatitis C Virus (HCV) RNA, comprising at least one double-stranded RNA effector molecule or complex that comprises: (1) a sequence of at least 19 nucleotides having at least 90% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4); (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), ATR-4 (SEQ ID NO: 89); SEQ ID NO: 9; SEQ, ID NO: 11; SEQ ID NO: 19; SEQ ID NO: 22; SEQ ID NO: 26; SEQ ID NO: 31; SEQ ID NO: 32; or SEQ ID NO: 33; and (2) its complementary sequence.
24. The agent of claim 23, wherein the sequence of at least 19 nucleotides has at least 95% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4); (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), ATR-4 (SEQ ID NO: 89); SEQ ID NO: 9; SEQ, ID NO: 11; SEQ ID NO: 19; SEQ ID NO: 22; SEQ ID NO: 26; SEQ ID NO: 31; SEQ ID NO: 32; or SEQ ID NO: 33.
25. The agent of claim 23 or 24, wherein the sequence of at least 19 nucleotides is selected from within nucleotides 42-76, 126-154, or 271-297 of SEQ ID NO: 1 (GenBank accession number AB047639).
26. The agent of claim 23, 24, or 25, wherein the sequence of at least 19 nucleotides is selected from within nucleotides 305-338 of SEQ ID NO: 1 (GenBank accession number AB047639).
27. The agent of any one of claims 23-26, wherein the double-stranded RNA effector molecule or complex contains a double-stranded region of from 19 to 30 base pairs.

28. The agent of any one of claims 23-26, wherein the double-stranded RNA effector molecule or complex contains a double-stranded region of from 21 to 27 base pairs.
29. The agent of any one of claims 23-26, wherein the double-stranded RNA effector molecule or complex contains a double-stranded region of about 21 or about 27 base pairs.
30. The agent of any one of claims 23-29, wherein the sequence of at least 19 nucleotides comprises a sequence selected from SEQ ID NOS: 5-42, more preferably a sequence selected from SEQ ID NOS. 9, 11, 19, 22, 26, 31, 32, or 33.
31. The agent of any one of claims 23-30, wherein the sequence of at least 19 nucleotides and its complementary sequence are connected via a loop sequence.
32. The agent of claim 31, wherein the loop sequence is from 5 to 20 nucleotides in length.
33. The agent of claim 32, wherein the loop sequence is from 7 to 10 nucleotides in length.
34. The agent of claim 33, wherein the loop sequence is about 9 nucleotides in length.
35. The agent of claim 23, wherein at least one double-stranded RNA effector molecule has a sequence selected from SEQ ID NOS: 44-81, more preferably a sequence selected from SEQ ID NOS: 48, 50, 56, 58, 61, 65, 70, 71, and 72.
36. The agent of any one of claims 23-35, comprising from two to five double-stranded RNA effector molecules.
37. The agent of claim 36, wherein the two to five double-stranded RNA effector molecules each comprise: (1) a sequence of at least 19 nucleotides having at least 90% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4); (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), ATR-4 (SEQ ID NO: 89); SEQ ID NO: 9; SEQ, ID NO: 11; SEQ ID NO: 19; SEQ ID NO: 22; SEQ ID NO: 26; SEQ ID NO: 31; SEQ ID NO: 32; or SEQ ID NO: 33; and (2) its complementary sequence; and (3) a loop sequence.

38. The agent of claim 36 or 37, wherein at least four double-stranded RNA effector molecules are administered, wherein: 1) a first effector molecule comprises the nucleotide sequence of SEQ ID 11, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 19, a third effector molecule
5 comprises the nucleotide sequence of SEQ ID NO: 22, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33; or 2) a first effector molecule comprises the nucleotide sequence of SEQ ID 9, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 11, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, and a
10 fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33, or 3) a first effector molecule comprises the nucleotide sequence of SEQ ID 11, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 19, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO:
15 33; or 4) a first effector molecule comprises the nucleotide sequence of SEQ ID 19, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 22, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 32, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33; or 5) a first effector molecule comprises the nucleotide sequence
20 of SEQ ID 26, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 32, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33 . .

39. The agent of claim 38, wherein the four double-stranded RNA effector
25 molecules are selected from SEQ. ID. NOS: 48, 50, 56, 58, 61, 65, 70, 71, and 72; or more preferably comprise the sequences: SEQ ID NO: 50, SEQ ID NO: 58, SEQ ID NO: 61, and SEQ ID NO: 72; or sequences: SEQ ID NO: 48, SEQ ID NO: 50, SEQ ID NO: 70, and SEQ ID NO: 72; or sequences SEQ ID NO: 50, SEQ ID NO: 58, SEQ ID NO: 70, and SEQ ID NO: 72; or sequences SEQ ID NO: 58,
30 SEQ ID NO: 61, SEQ ID NO: 71, and SEQ ID NO: 72; or SEQ ID NO: 65, SEQ ID NO: 70, SEQ ID NO: 71, and SEQ ID NO: 72.

40. A construct for silencing a hepatitis C virus (HCV) RNA, the construct encoding at least one double-stranded RNA effector molecule or complex wherein the effector molecule or complex comprises: (1) a sequence of at least

19 nucleotides having at least 90% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4); active target region (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89); and (2) its complementary sequence.

41. The construct of claim 40, wherein the sequence of at least 19 nucleotides has at least 95% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4), active target region (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), or ATR-4 (SEQ ID NO: 89);.

42. The construct of claim 40 or 41, wherein the sequence of at least 19 nucleotides is selected from within nucleotides 42-76, 126-154, or 271-297 of SEQ ID NO: 1 (GenBank accession number AB047639).

43. The construct of claim 40, 41, or 42, wherein the sequence of at least 19 nucleotides is selected from within nucleotides 305-338 of SEQ ID NO: 1 (GenBank accession number AB047639).

44. The construct of any one of claims 40-43, wherein the double-stranded RNA effector molecule contains a double-stranded region of from 19 to 30 base pairs.

45. The construct of any one of claims 40-43, wherein the double-stranded RNA effector molecule contains a double-stranded region of from 21 to 27 base pairs.

46. The construct of any one of claims 40-43, wherein the double-stranded RNA effector molecule contains a double-stranded region of about 21 or about 27 base pairs.

47. The construct of any one of claims 40-46, wherein the sequence of at least 19 nucleotides comprises a sequence selected from SEQ ID NOS: 5-42, more preferably a sequence selected from SEQ ID NOS. 9, 11, 19, 22, 26, 31, 32, or 33.

30 .

48. The construct of any one of claims 40-47, wherein the sequence of at least 19 nucleotides and its complementary sequence are connected via a loop sequence.
49. The construct of claim 48, wherein the loop sequence is from 5 to 20 nucleotides in length.
50. The construct of claim 48, wherein the loop sequence is from 7 to 10 nucleotides in length.
51. The construct of claim 48, wherein the loop sequence is about 9 nucleotides in length.
52. The construct of claim 40, wherein at least one double-stranded RNA effector molecule has a sequence selected from SEQ ID NOS: 44-81, more preferably a sequence selected from SEQ ID NOS: 48, 50, 56, 58, 61, 65, 70, 71, and 72.
53. The construct of any one of claims 40-52, wherein the expression construct encodes from two to five double-stranded RNA effector molecules.
54. The construct of claim 53, wherein the two to five double-stranded RNA effector molecules each comprise: (1) a sequence of at least 19 nucleotides having at least 90% identity with a nucleotide sequence within Conserved Region 1 (SEQ ID NO: 2), Conserved Region 2 (SEQ ID NO: 3), Conserved Region 5 (SEQ ID NO: 4); (ATR)-1 (SEQ ID NO: 86), ATR-2 (SEQ ID NO: 87), ATR-3 (SEQ ID NO: 88), ATR-4 (SEQ ID NO: 89); SEQ ID NO: 9; SEQ, ID NO: 11; SEQ ID NO: 19; SEQ ID NO: 22; SEQ ID NO: 26; SEQ ID NO: 31; SEQ ID NO: 32; or SEQ ID NO: 33; (2) its complementary sequence; and (3) a loop sequence.
55. The construct of claim 54, wherein the expression construct encodes at least four double-stranded RNA effector molecules, wherein: 1) a first effector molecule comprises the nucleotide sequence of SEQ ID 11, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 19, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 22, and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33; or 2) a first effector molecule comprises the nucleotide sequence of SEQ ID 9, a second effector molecule comprises the nucleotide sequence of SEQ ID NO: 11, a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 31, and a

fourth effector molecule comprises the nucleotide sequence of SEQ ID NO: 33, or
3) a first effector molecule comprises the nucleotide sequence of SEQ ID 11, a
second effector molecule comprises the nucleotide sequence of SEQ ID NO: 19,
a third effector molecule comprises the nucleotide sequence of SEQ ID NO: 31,
5 and a fourth effector molecule comprises the nucleotide sequence of SEQ ID NO:
33; or 4) a first effector molecule comprises the nucleotide sequence of SEQ ID
19, a second effector molecule comprises the nucleotide sequence of SEQ ID
NO: 22, a third effector molecule comprises the nucleotide sequence of SEQ ID
NO: 32, and a fourth effector molecule comprises the nucleotide sequence of
10 SEQ ID NO: 33; or 5) a first effector molecule comprises the nucleotide sequence
of SEQ ID 26, a second effector molecule comprises the nucleotide sequence of
SEQ ID NO: 31, a third effector molecule comprises the nucleotide sequence of
SEQ ID NO: 32, and a fourth effector molecule comprises the nucleotide
sequence of SEQ ID NO: 33.

15 56. The construct of claim 55, wherein the four double-stranded RNA effector
molecules are selected from SEQ. ID. NOS: 48, 50, 56, 58, 61, 65, 70, 71, and
72; or more preferably comprise the sequences: SEQ ID NO: 50, SEQ ID NO: 58,
SEQ ID NO: 61, and SEQ ID NO: 72; or sequences: SEQ ID NO: 48, SEQ ID NO:
50, SEQ ID NO: 70, and SEQ ID NO: 72; or sequences SEQ ID NO: 50, SEQ ID
20 NO: 58, SEQ ID NO: 70, and SEQ ID NO: 72; or sequences SEQ ID NO: 58,
SEQ ID NO: 61, SEQ ID NO: 71, and SEQ ID NO: 72; or sequences SEQ ID NO:
65, SEQ ID NO: 70, SEQ ID NO: 71, and SEQ ID NO: 72.57. The construct
of claim 40, wherein the expression construct contains at least two expression
cassettes, each expression cassette directing the expression of at least one
25 double-stranded RNA effector molecule.

58. The construct of claim 57, wherein each expression cassette
independently comprises at least one RNA polymerase III promoter selected from
a U6 promoter, a 7SK promoter, an H1 promoter, and an MRP promoter.

59. The construct of claim 58, wherein each expression cassette comprises a
30 7SK promoter driving the expression of one double-stranded RNA effector
molecule, and an RNA pol III termination signal.

60. A method of treating or preventing HCV infection in a patient, comprising administering to said patient, an effective amount of an agent of composition according to any one of claims 23-59 to said patient.




Fig. 1




Fig. 2




Fig. 3