



HU000031060T2

(19) **HU**(11) Lajstromszám: **E 031 060**(13) **T2****MAGYARORSZÁG**
Szellemi Tulajdon Nemzeti Hivatala**EURÓPAI SZABADALOM**
SZÖVEGÉNEK FORDÍTÁSA

(21) Magyar ügyszám: **E 06 784009** (51) Int. Cl.: **A01N 63/04** (2006.01)
(22) A bejelentés napja: **2006. 08. 04.** **A01H 17/00** (2006.01)
A01P 3/00 (2006.01)
(96) Az európai bejelentés bejelentési száma: **A01N 436/53** (2006.01)
EP 20060784009
(97) Az európai bejelentés közzétételi adatai: (86) A nemzetközi (PCT) bejelentési szám:
EP 1916890 A1 **2007. 02. 22.** **PCT/NZ 06/000202**
(97) Az európai szabadalom megadásának meghirdetési adatai: (87) A nemzetközi közzétételi szám:
EP 1916890 B1 **2016. 09. 07.** **WO 07021200**

(30) Elsőbbségi adatok: 54160605 2005. 08. 16. NZ	(73) Jogosult(ak): Grasslanz Technology Limited, Palmerston North (NZ)
(72) Feltaláló(k): ROLSTON, Maurice, Philip, Christchurch RD 2 (NZ) SIMPSON, Wayne, Roydon, Palmerston North (NZ)	(74) Képviselő: Danubia Szabadalmi és Jogi Iroda Kft Sár és Társai Ügyvédi Iroda, Budapest

(54) **Füendofita kiterjesztett tulajdonságai**

Az európai szabadalom ellen, megadásának az Európai Szabadalmi Közlönyben való meghirdetésétől számított kilenc hónapon belül, felszólalást lehet benyújtani az Európai Szabadalmi Hivatalnál. (Európai Szabadalmi Egyezmény 99. cikk(1))

A fordítást a szabadalmas az 1995. évi XXXIII. törvény 84/H. §-a szerint nyújtotta be. A fordítás tartalmi helyességét a Szellemi Tulajdon Nemzeti Hivatala nem vizsgálta.

(19)



(11)

EP 1 916 890 B1

(12)

EUROPEAN PATENT SPECIFICATION

(45) Date of publication and mention of the grant of the patent:
07.09.2016 Bulletin 2016/36

(51) Int Cl.:
A01N 63/04 ^(2006.01) **A01H 17/00** ^(2006.01)
A01N 43/653 ^(2006.01) **A01P 3/00** ^(2006.01)

(21) Application number: **06784009.0**

(86) International application number:
PCT/NZ2006/000202

(22) Date of filing: **04.08.2006**

(87) International publication number:
WO 2007/021200 (22.02.2007 Gazette 2007/08)

(54) GRASS ENDOPHYTE ENHANCED ATTRIBUTES

ERWEITERTE ATTRIBUTE VON GRASENDOPHYTEN

ATTRIBUTS AMELIORES D'HERBE ENDOPHYTE

(84) Designated Contracting States:
AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC NL PL PT RO SE SI SK TR

(56) References cited:
WO-A-02/13616 WO-A-2004/106487
US-A- 6 111 170 US-A1- 2005 150 024

(30) Priority: **16.08.2005 NZ 54160605**

(43) Date of publication of application:
07.05.2008 Bulletin 2008/19

(73) Proprietor: **Grasslanz Technology Limited Palmerston North (NZ)**

(72) Inventors:
• **ROLSTON, Maurice, Philip Christchurch RD 2 (NZ)**
• **SIMPSON, Wayne, Roydon Palmerston North (NZ)**

(74) Representative: **Vogel, Andreas et al Bals & Vogel**
Universitätsstrasse 142
44799 Bochum (DE)

- **ROLSTON M P ET AL: "Tolerance of AR1 Neotyphodium endophyte to fungicides used in perennial ryegrass seed production" PROCEEDINGS OF THE NEW ZEALAND PLANT PROTECTION CONFERENCE, XX, XX, vol. 55, 1 January 2002 (2002-01-01), pages 322-326, XP008089558**
- **TIMPER P. ET AL.: 'Response of Pratylenchus spp. in tall fescue infected with different strains of the fungal endophyte Neotyphodium coenophialum' NEMATOTOLOGY vol. 7, no. 1, 2005, pages 105 - 110, XP008077066**
- **HILL N.S. ET AL.: 'Endophyte viability in seeding tall fescue treated with fungicides' CROP SCIENCE vol. 40, no. 5, 2000, pages 1490 - 1491, XP003008972**
- **SIEGEL M.R. ET AL.: 'A fungal endophyte of tall fescue: evaluation of control methods' PHYTOPATHOLOGY vol. 74, no. 8, 1984, pages 937 - 941, XP008077069**

EP 1 916 890 B1

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

Description**TECHNICAL FIELD**

5 [0001] The invention relates to grass endophyte enhanced attributes. More specifically, the invention relates to an endophyte with improved fungicide resistance and stability when in combination with a cultivated plant or part thereof.

BACKGROUND ART

10 [0002] It is known from New Zealand patent specification 233,083 that synthetic combinations of endophyte/herbage cultivars can be made which are resistant to pests and can be less toxic to grazing animals than naturally occurring combinations. These selected properties may be achieved by selecting synthetic combinations which produce relatively high levels of peramine and relatively low levels of or no lolitrem B in ryegrass.

15 [0003] US 6,111,170 (the '170 patent), describes findings that selected endophytes of the genus *Neotyphodium* form stable synthetic combinations with tall fescue hosts (*Festuca arundinacea*). The combinations described have improved resistance to invertebrate pests as compared to tall fescue cultivars that do not contain such endophytes. The particular strains of endophyte described in the '170 patent include AR501, AR502, AR510, AR542, AR572, AR577 and AR584. This patent does not however describe, nor suggest other attributes of these endophytes and their combinations beyond improved pest resistance and reduced toxicity.

20 [0004] Russian patent abstract, RU2201678C2 describes a biopreparation for protecting plants against fungus and bacterial diseases. The biopreparation is based upon live cells and spores of endophytic bacterium *Bacillus subtilis*. This bacterium has a high antagonistic activity to agents of fungus and bacterial diseases in vegetable, grain and fruit cultures. No description is made regarding use of the bacterium in grasses. It is also not obvious from this abstract that the bacterium described is resistant to fungicides. The description merely makes reference to the bacterium preventing

25 fungal infection of the plant in effect acting as a fungicide itself.
 [0005] Meriaux *et al* describes the viability of ryegrass (*Lolium perenne*) seed infected with *Neotyphodium* endophyte after treatment with fungicides. The fungicides used were prochloraz (prochloraz), triticonazole, bitertanol, and fluquinconazole administered at two doses or heat treated (oven treatment at 60 or 80°C). Seed germination is described as being effected for treatment with prochloraz (1 g/kg), triticonazole (5 and 10 g/kg) and heat treatment at 80°C. No reference is made or suggested towards endophyte viability or viability (seed or endophyte) when combined with cultivated

30 grasses. It is the inventor's experience that seed viability and endophyte viability are two very different issues with seeds often retaining viability for longer than endophytes.
 [0006] Rolston *et al* describes fungicide treatment effects on AR1 endophyte and ryegrass combinations. AR1 is a *Neotyphodium* species endophyte that has a different alkaloid profile to AR584 in that AR1 does not produce loline alkaloids and may produce terpendoles differently to AR584. The publication notes that AR1 endophyte viability was not affected by the fungicide treatments including triazole and strobilurin fungicides. It is the inventor's experience that it is not obvious that a property noted for one endophyte in one type of plant would also exhibit the same properties with another endophyte or plant. Anecdotal evidence exists showing that endophyte strain and plant species may result in very different results. One reason for the variation is that different endophytes give different attributes to grass endophyte

35 combinations. One reason for the variation is that different endophytes have different levels and types of alkaloid profile. As a result different endophytes in combination with different grasses have variable properties. In addition, different species of grass also influence properties of combinations with endophytes.
 [0007] Saiga *et al* describes the effects of *Neotyphodium* endophytes on plant characteristics including fungicide soak treatments. The fungicide, benomyl (benzimidazole group) is described as being effective in killing endophyte in both perennial ryegrass and tall fescue. Triforine (amide group) fungicide was effective in tall fescue and effective in some perennial ryegrass plants. Thiophanate-methyl (benzimidazole precursor fungicide) had little effect on endophyte in perennial ryegrass and had no effect on endophyte in tall fescue. The aim of the publication method is to remove endophytes altogether from the grass. As a result, particularly strong doses of fungicides are used and the grass is soaked in fungicide. No disclosure is made towards identification of endophytes with higher resistance to fungicides, or

40 disclosure regarding stability.
 [0008] Hill *et al* describes tall fescue cultivars containing endophytes including *Neotyphodium coenophialum*. The abstract discusses how to maintain viable endophyte in seedling plants by applying chloroneb (aromatic fungicide) and terrazole (thiazole group) fungicides during the first 7 to 21 days post germination. Hill *et al* does not identify endophytes with resistance to fungicides or identify AR584 as exhibiting any particular resistance. Further, the time period of analysis described is only post germination and only in seedling plants.

45 [0009] Park *et al* describes development of a microbial fungicide that utilises endophytic bacteria for the control of cucumber and red pepper anthracnoses caused by *Colletotrichum orbiculare* and *C. coccodes* and *C. acutatum*. Among 18 strains isolated, a bacterial strain EB215 isolated from cucumber roots displayed the most potent antifungal activity

against *Colletotrichum* species. No mention is made regarding improved fungicide resistance due to endophyte, nor is usage in terms of grasses such as tall fescue.

[0010] Bluett *et al* describes a trial comparing various properties between ryegrass (*Lolium perenne*) infected with wild type endophyte, AR1 endophyte or no endophyte. AR1-infected ryegrass was found to produce similar pasture yields as wild endophyte-infected ryegrass, while offering small improvements in milk yield with no incidence of ryegrass staggers noted in grazing animals. No discussion is made on the endophyte effect with regard to fungicide treatment, endophyte stability or to the endophyte AR584 or tall fescue grass cultivars.

[0011] Timper, P.; Gates, R. N.; Bouton J. H-'Response of *Pratylenchus* spp. In tall fescue infected with different strains of the fungal endophyte *Neotyphodium coenophialum*' *Nematology*, Vol. 7 No 1, Pages 105-110 (2005) disclose that a non-ergot producing endophyte strain of *Neotyphodium* AR584 maintains some resistance to nematode pests. Timper et al are entirely silent regarding any relationship between endophyte viability and plant/endophyte fungicide resistance.

[0012] US200515002 is the US national phase entry of WO2002013616. This document describes grass having improved stress tolerance and lower animal toxicity when infected with certain isolates of *Neotyphodium* fungi.

[0013] WO2004106487 describes an endophyte or endophyte culture of *Neotyphodium lolii* which in combination with ryegrass does not cause typical symptoms of ryegrass toxicosis in grazing animals and which produces janthitrem epoxide compounds.

[0014] It should be appreciated by a person skilled in the art that an endophyte plant combination that has increased fungicide resistance would be of benefit to enhance existing turf and forage seed crop management practices. In addition, increased stability during transport would also be an advantage.

[0015] It is an object of the present invention to address the foregoing problems or at least to provide the public with a useful choice.

[0016] It is acknowledged that the term 'comprise' may, under varying jurisdictions, be attributed with either an exclusive or an inclusive meaning. For the purpose of this specification, and unless otherwise noted, the term 'comprise' shall have an inclusive meaning - i.e. that it will be taken to mean an inclusion of not only the listed components it directly references, but also other non-specified components or elements. This rationale will also be used when the term 'comprised' or 'comprising' is used in relation to one or more steps in a method or process.

[0017] Further aspects and advantages of the present invention will become apparent from the ensuing description which is given by way of example only.

DISCLOSURE OF INVENTION

[0018] For the purposes of this specification, the term 'endophyte' or grammatical variations thereof refer to fungi of *Neotyphodium* spp. living within cultivated grasses or axenic culture medium.

[0019] The term 'cultivated grasses' or grammatical variations thereof refer to varieties of grasses that have been created or selected intentionally and maintained through cultivation.

[0020] The term 'synthetic grass cultivar' refers to the grass cultivar being produced through selective breeding techniques including selection and development from an uncultivated population. For example, a synthetic grass cultivar refers to where -

- reproducible units are from a cross-pollinated crop. This could encompass clones or inbred grasses for example;
- materials used are selected from their performance in combining ability or progeny tests;
- the cultivar is constituted by random inter-mating of the units;
- the units are maintained so that the synthetic can be reconstituted.

[0021] The term 'combination' or grammatical variations thereof refer to the combination of an endophyte and culturally improved grass varieties each of which have been isolated from nature, but the combination of which does not exist in nature.

The term 'plant or a part thereof' refers to the plant in entirety or selected portions of the plant during the plant life cycle such as the plant seed, shoots, roots, flowers, stems and the like.

The term 'retain' or grammatical variations thereof in relation to endophytes refers to the ability of the endophyte to maintain a level of viability in combination with a host plant, or part thereof, at a level acceptable for a given circumstance or set of circumstances over a given period of time and under a given set of conditions. It should be appreciated by those skilled in the art that acceptable percentages for viability may vary depending on the generation or class of seed to which it relates as well as by varying industry standards.

The term 'viable' or grammatical variations thereof refer to the endophyte being capable of living, developing, or multiplying under favorable conditions in symbiotic relationship with a host plant or in axenic culture.

The term 'treatment with fungicide' refers to the combination being sprayed, powdered, infused or other methods of

administration, by a fungicide containing composition.

[0022] The term 'fungicide' refers to substances that destroy or inhibit growth of fungi.

[0023] According to a further aspect of the present invention there is provided a method of retaining endophyte viability in a plant or a part thereof after treatment with fungicide, produced by a combination a grass cultivar and endophyte combination, wherein the endophyte is AR584 endophyte (AGAL deposit no. NM98/04676 dated 12 May 1998).

[0024] The inventor's have unexpectedly found that AR584 endophyte exhibits an improved ability to resist fungicide treatments. It is envisaged that this attribute will have significant commercial benefit in farming. By way of example, standard farm turf or seed crop management calls for spraying of fungicide treatments to the turf or seed crop to avoid foliar and soil borne fungal diseases. A problem with existing grasses and grass endophyte combinations is that, fungicides commonly used in turf and seed crop management and control of unwanted fungal pathogens also tend to significantly reduce endophyte viability. As a result, the favourable properties noted for endophyte and grass combinations may be reduced or negated altogether. AR584 however has been noted by the inventors to have a resistance to standard commercial fungicides to an extent that means a farmer can spray to avoid foliar diseases and unwanted soil borne fungal pathogens and still retain adequate endophyte viability.

[0025] Preferably, at least a 50% or greater endophyte viability is retained after fungicide treatment.

[0026] Preferably, the fungicides used in treatment have systemic characteristics.

[0027] Preferably, the fungicides used in treatment have foliar disease control characteristics.

[0028] Preferably, the fungicide used in treatment has activity against soil borne fungicidal pathogens.

[0029] Preferably, fungicides used in treatment are selected from the triazole group, strobilurin group or combinations of these two fungicide groups.

[0030] Preferably, at least 60% or greater endophyte viability is retained after being subjected to treatment with triazole based fungicide.

[0031] Preferably, the combination retains at least 65% or greater endophyte viability, more preferably, greater than 80% viability, when subjected to a 50:50wt/wt combination of triazole based fungicide and strobilurin based fungicide.

[0032] Preferably, the fungicide treatment is applied to the grass cultivar and endophyte combination at standard rates used in control of foliar diseases.

[0033] Preferably, triazole based fungicide treatment is applied as a spray at a rate of 1000 ml/ha. This is envisaged to be the equivalent to 250g epoxiconazole/ha.

[0034] Preferably, strobilurin based fungicide treatment is applied as a spray at a rate of 1000 ml/ha. This is envisaged to be the equivalent to 250g azoxystrobin/ha.

[0035] Preferably, the fungicide treatment is applied at early flowering of the grass cultivar and endophyte combination.

[0036] Preferably, the fungicide treatment is applied to the combination at early flowering and re-applied again at late flowering of the combination.

[0037] In preferred embodiments, the fungicide is applied as a spray although this should not be seen as limiting as the fungicide may be applied by other methods such as immersion, powders, dusting, gels or combinations thereof. It is envisaged that this could be applied to the plant or part thereof.

[0038] According to an aspect of the present invention there is provided a method of retaining endophyte viability in a plant or a part thereof after treatment with fungicide, produced by a combination of a grass cultivar and endophyte combination, wherein the endophyte is AR584 endophyte (AGAL deposit no. NM98/04676 dated 12 May 1998) wherein the combination retains viable AR584 endophyte therein after the combination is subjected to temperatures of between 15°C and 55°C for a time period of at least 3 days.

[0039] Preferably the combination retains at least 60% or greater endophyte viability when the combination is subjected to a temperature of up to 40°C and a relative humidity of up to 100% for at least a time period of 3 days.

[0040] Preferably, the plant or a part thereof produced by the combination of the endophyte and grass cultivar combination as described above is further characterised by retaining endophyte viability when the grass and endophyte combination is subjected to temperatures above 20°C.

[0041] Preferably, the AR584 endophyte in the combination retains at least a 50% or greater endophyte viability.

[0042] Preferably, the endophyte viability is retained when the endophyte and plant or part thereof combination is subjected to temperatures of 35°C to 45°C.

[0043] Preferably, endophyte viability is retained when the endophyte and plant or part thereof combination is subjected to temperatures of 40°C.

[0044] Preferably, endophyte viability is retained when the endophyte and plant or part thereof combination is subjected to a temperature increase as described over on a continuous basis. Preferably, endophyte viability is retained when the temperature is increased over time periods of at least 3 days.

[0045] Preferably, the plant or a part thereof produced by the endophyte and grass cultivar combination is further characterised by retaining endophyte viability when the grass and endophyte combination is subjected to a relative humidity of 70% or greater.

[0046] Preferably, the endophyte viability is retained when the combination is subjected to a relative humidity of 70%

or greater for at least 3 days.

[0047] Preferably, the plant or a part thereof produced by the endophyte and grass cultivar combination is able to maintain viability when subjected to relative humidity of up to 100%. Preferably, endophyte viability is retained when the endophyte and plant or part thereof combination is subjected to a humidity increase as described over on a continuous basis. Preferably, endophyte viability is retained when the humidity is increased over time periods of at least 3 days.

[0048] Preferably, at least 75% or greater endophyte viability is retained.

[0049] Preferably, the combination:

- (a) retains at least 60% or greater endophyte viability when the combination is subjected to a temperature of up to 40°C and a relative humidity of up to 100% for at least a time period of 3 days; and,
- (b) retains at least a 50% or greater endophyte viability after treatment with fungicide.

[0050] The inventors have also unexpectedly found that AR584 endophyte has a surprisingly high tolerance to elevated temperatures and humidities allowing it to remain viable within a plant or part thereof for extended periods of time in such conditions. This attribute is of commercial benefit where plants or parts thereof are transported as few special treatments such as refrigeration are required therefore also reducing the cost of transport.

[0051] Preferably, the plant or part thereof is the entire plant or selected portions of the plant. It is envisaged that this may encompass the plant seed(s), shoot(s), leaves, stem(s), flower(s), root(s) and combinations thereof.

[0052] Preferably, the plant or part thereof is a seed or seeds.

[0053] Preferably, the grass cultivar is selected from one or more tall fescue plant varieties.

[0054] Preferably, this may be selected from the group including: Grasslands Flecha, Grasslands Advance, Kentucky 31, Georgia 5, Jesup, Jackal, Quantum, and combinations thereof.

[0055] Preferably, the combination is produced by inoculating the grass cultivar with an axenic culture of the endophyte.

[0056] Preferably, the combination is produced by crossing the endophyte/grass cultivar combination with an endophyte free grass cultivar to form a grass cultivar infected with endophyte. Preferably, the grass cultivar is a synthetic grass cultivar.

[0057] It should be appreciated from the above description that there is provided new methods for AR584 strain endophyte. In particular, the plant or part thereof produced by the combination of AR584 endophyte and a grass cultivar has significant new and unexpected advantages including an enhanced level of resistance to fungicides commonly used to prevent foliar diseases in grass. This attribute means that a farmer can apply standard fungicides without need for any special handling.

[0058] The plant or part thereof also has a significantly improved stability compared to other existing endophyte / grass cultivar combinations with respect to retaining endophyte viability when the plant or part thereof is subjected to extremes of temperature and humidity. One major benefit from this attribute is in transport, for example of seeds, where viability may be reduced using existing endophyte / grass combinations due to poor handling practices or extremes in temperature and humidity experienced in certain climates and countries. One such example may be when quantities of seed are transported in the holds of ships across varying climatic zones in uncontrolled storage environments.

BRIEF DESCRIPTION OF DRAWINGS

[0059] Further aspects of the present invention will become apparent from the following description which is given by way of example only and with reference to the accompanying drawings in which:

Figure 1 is a graph comparing endophyte viability levels of AR584, AR542 and AR502 in combination with two different grass cultivars with five different fungicide treatments;

Figure 2 is a graph comparing endophyte viability levels of AR584, AR542 and AR502 in combination with two different grass cultivars with either 0 days or 3 days accelerated aging (AA); and,

Figure 3 is a graph comparing viable endophyte levels for AR542 and AR584 in two different grass cultivars treated with three different fungicides treatments.

BEST MODES FOR CARRYING OUT THE INVENTION

[0060] The invention methods is now described with reference to trials completed by the inventors.

TRIAL 1

Introduction

5 [0061] The trial was undertaken to determine the tolerance of three endophytes (AR502, AR542 and AR584) to fungicides commonly used in grass seed production for the control of foliar diseases.

[0062] The fungicides evaluated belong to two different chemical groups. The first group relate to triazole fungicides including:

- 10
- Opus® with active ingredient epoxiconazole and
 - Folicur® with active ingredient tebuconazole.

[0063] The second group relate to the strobilurin fungicides including Amistar® with active agent azoxystrobin.

15 [0064] It is the inventor's experience that the above chemical groups account for over 90% of the fungicides used in tall fescue grass production and management.

[0065] The accelerated aging (AA) process is a high temperature (40°C) and high relative humidity (RH) (100% RH) treatment of seed to simulate long term storage effects. Based on prior trials, accelerated aging is a useful and accurate predictor of long term natural storage stability.

20 Methods

Field Trial

25 [0066] Plants of two tall fescue cultivars 'Jesup' and 'Georgia 5' were inoculated with endophyte and transplanted into trial plots. Each plot consisted of 4 plants of each cultivar, in rows 1.0 m apart. Treatments were replicated three times in a randomised block design.

Fungicides

30 [0067] Five fungicide treatments were evaluated as follows:

1. Control

- 35
- Nil fungicide;

2. Opus®

- Epoxiconazole 250 g active ingredient per litre, triazole group sprayed at the label rate of 1000 ml/ha;

40 3. Amistar®

- Azoxystrobin 250 g active ingredient per hectare, strobilurin group

45 4. Opus® and Amistar®

- Opus® at 500 ml/ha (half label rate) and Amistar® 500 ml/ha and applied as a tank mix;

5. Folicur®

50 1. Tebuconazole 430 g active ingredient per litre; triazole group at the label rate of 440 ml/ha.

[0068] The doses used were the approximate recommended label rates for use of the above products. It should be noted that growers commonly use less than the label rate particularly if they plan to apply the fungicide more than once. As a result, the dosing rates used in this trial present a significant challenge to endophyte viability.

55 [0069] Fungicides were applied twice during the trial to replicate normal turf practice. Applications were made at Zadocks Growth Score (GS) 69 (early flowering/anthesis) and at Zadocks growth score (GS) 71 (caryopsis water ripe). This timing of application was as per those recommended to growers and coincides with the period of most disease pressure in crops.

Accelerated aging

[0070] Seeds were suspended on a fine mesh above water enclosed within plastic sealed containers and placed in an incubator set to maintain a temperature of approximately 40°C and with the seed chamber controlled to 100% relative humidity.

Endophyte Assessment by Grow-out test

[0071] Six replicates (each of 48 seedlings) consisting of duplicates from each treatment, were raised in peat-based trays and grown in a greenhouse for at least 6 weeks before blotting.

Endophyte detection

[0072] Cut tillers from the seedlings after 6 weeks growth were blotted on nitro-cellulose paper and developed using an antibody/antigen reaction.

[0073] The development process is completed by immersing nitrocellulose membrane sheets in blocking solution for at least two hours at room temperature. The blocking solution is composed of 2.42g of Tris (hydroxymethyl) methylamine, 2.92g of NaCl, 5g of non-fat milk powder and 10ml HCl (1 M) in 1 L of water that has been passed through a reverse osmosis (RO) system. The pH of the solution is 7.5. During the immersion, any unbound site prior to the application of primary antibody is blocked.

[0074] After removing the blocking solution, 25ml of fresh blocking solution per sheet is added along with 25µl of primary antibody per sheet. The sheets are soaked overnight at 4°C and shaken to keep the solution moving over the sheets. The next day, the blocking solution and primary antibody are removed and the sheets are rinsed several times with fresh blocking solution. Afterwards, the sheets are soaked and shaken for two hours at room temperature in 25ml of fresh blocking solution with 12.5µl of secondary antibody added. An enzyme is conjugated to this secondary antibody. Subsequently, blocking solution and the secondary antibody are decanted off and the sheets are rinsed by fresh blocking solution.

[0075] In order to develop the colour that allows endophyte identification, the nitrocellulose membranes are immersed into chromogen. This product is prepared by dissolving 75mg of Fast Red in 12.5 ml of Tris buffer (For one litre of RO water, the Tris buffer is composed of 24.2g of Tris (hydroxymethyl) methylamine (pH: 8.2)) per sheet and by dissolving separately 12.5mg of naphthol as-mx phosphate in 12.5 ml Tris buffer per sheet and these solutions combined. The sheets are incubated on shaker for 15 minutes. At the end of the 15 minutes, the chromogen is removed and the sheets are rinsed in water that has been passed through a reverse osmosis system. The sheets are then ready to be read.

Statistical Analysis

[0076] Data analysis was undertaken using a statistical package - GenStat. Version 7.

Results**Endophyte effect**

[0077] The average viable endophyte level of AR584 endophyte was significantly higher ($P < 0.001$) than AR542 and AR502 for both fungicide treatment and accelerated aging experiments. AR584 averaged 81% viable endophyte after all treatments, while AR542 and AR502 averaged 60 and 56% respectively (Table 1).

[0078] The three endophytes were also noted as behaving differently between the main treatment variables: endophyte x cultivar; endophyte x accelerated aging (AA) and endophyte x fungicide (Table 1).

Table 1. Viable endophyte comparison of three endophyte strains (AR584; AR542 and AR502). Each value is averaged across two cultivars, two accelerated aging treatments, five fungicides and six repetitions).

Endophyte	E+%
AR584	81
AR542	60
AR502	56
LSD 5%	5.1
F probability	

(continued)

Endophyte	E+%
Endophyte	<0.001
Endophyte x cultivar	0.005
Endophyte x accelerated aging	<0.001
Endophyte x fungicide	<0.001

10 **Fungicide effect**

[0079] The two fungicides significantly reduced the level of viable endophyte (means of all other treatments) from 92% in the nil fungicide treatment to 61% for the Opus@ and Amistar® mix; and 38% for the Opus@ only treatment (Table 2).

[0080] The data points to epoxiconazole as being the component most significantly reducing viable endophyte levels.

15 **Table 2.** Effect of fungicides on viable endophytes (data means averaged across two endophytes, two cultivars and six repetitions).

Fungicide	E+%
nil	81
Amistar®	76
Folicur®	70
Opus@ + Amistar®	61
Opus@	45
LSD 5%	6.6
F probability	
Fungicide	<0.001
Fungicide x endophyte	<0.001

30 **Endophyte and Fungicide Interaction**

[0081] The endophyte and fungicide interaction as shown in Table 1 is further broken down in Table 3 below.

[0082] It is the inventor's unexpected findings that the interaction appears to arise because AR584 is significantly more tolerant of the two fungicides compared to the control and other endophyte combinations tested.

[0083] With the Opus@ and Amistar® mixture treatment, AR584 had at least a 50% higher endophyte viability compared to other endophytes (73% viable endophyte compared to 49% in AR542). For Opus@ only treatments, AR584 had more than three times the level of viable endophyte compared to AR542. The interaction is shown in more detail in Figure 1.

40 **Table 3.** AR584 and AR542 viable endophyte for fungicide treatments (means of two cultivars accelerated aging treatments and six repetitions).

Fungicide	Endophyte		
	AR584	AR542	AR502
	Endophyte Viability [%]		
Nil	93	91	58
Amistar	93	73	63
Folicur	89	70	51
Opus + Amistar	73	49	46
Opus	60	17	61
LSD 5%		11.3	

55 **Cultivar Effect**

[0084] There was a small cultivar effect, with Georgia 5 cultivar having higher endophyte levels than Jesup cultivar following fungicide treatments (Table 4). The cultivar and endophyte interaction was statistically significant.

Table 4. Viable endophyte for two cultivars (means averaged for three endophytes, five fungicides and six repetitions).

Cultivar	Endophyte Viability [%]
Georgia 5	68
Jesup	63
LSD 5%	5
F probability	
Cultivar	0.02
Cultivar x Endophyte	0.005

Accelerated Aging

[0085] Three days of accelerated aging were tested to determine endophyte viability levels for AR584, AR542 and AR502 (Table 5).

[0086] The results found show that AR502 appears to be very sensitive to the accelerated aging treatment (Table 5 and Figure 2) and AR584 has a significantly improved resistance to the elevated temperature and humidity.

Table 5. Effect of 0 or 3 days accelerated aging (AA) on viable endophyte (means of two cultivars, five fungicides and 6 reps).

Endophyte	Days accelerated aging	
	0 AA	3 AA
AR584	89	74
AR542	61	60
AR502	79	33
LSD 5%	10	

[0087] Time periods of longer than 3 days were also tested and gave proportionally similar results as above however, with reduced rates of endophyte viability.

Conclusion

[0088] The data strongly supports the contention that AR584 has unique properties compared to other endophytes in retaining seed viability characteristics when plants are treated with fungicides during seed development. AR584 was found to be more robust to fungicides of different chemical activities, especially to the fungicide "Opus@".

[0089] AR584 was also found to have increased resistance to extremes in heat and humidity when compared to other endophytes.

TRIAL 2

Introduction

[0090] This trial was done to supplement the results shown in trial one. The tolerance of two endophytes, AR584 and AR542 in two tall fescue cultivars, 'Jesup' and 'Georgia 5' when treated with three different fungicide treatments was tested.

[0091] The fungicide treatments tested included:

1. Nil treatment
2. Opus only applied at 1.0 L/ha; and
3. a mixture of Opus@ and Amistar@ applied at 0.5 + 0.5 L/ha

[0092] The accelerated ageing (AA) process was also adapted. As discussed above in trial 1, AA was undertaken by incubating seeds at a high temperature of 40°C and 100% Relative Humidity (RH).

Method**Field Trial**

5 **[0093]** A field trial was conducted on two tall fescue cultivars, 'Jesup' and 'Georgia 5' that were inoculated with the two endophytes AR584 and AR542. The field trial was carried out as described above in Trial one, wherein the plants of were inoculated with endophyte and then transplanted into trial plots. Each plot consisted of 4 plants of each cultivar, in rows 1.0 m apart. Treatments were replicated three times in a randomised block design.

10 **Fungicides**

[0094] Three fungicide treatments where trialed as outlined below:

1. Nil treatment;
- 15 2. Opus@ only (O), applied at 1.0 L/ha; and
3. a mixture of Opus@ applied at 0.5 L/ha and Amistar® (O+A) applied at 0.5 L/ha.

[0095] Again, the doses used were the approximately the recommended label rates for use of the above products. The timing of the fungicide application was as per those recommended to growers and coincides with the period of most disease pressure in crops.

20 **[0096]** The accelerated ageing, Endophyte Assessment by Grow-out test, Endophyte detection and Statistical Analysis was also carried out as outlined above in Trial 1.

Results

25

Endophyte and Fungicide effect

[0097] Figure 3 and Table 6 shows the viability of each endophyte when treated with the various fungicide treatments. As shown, AR584 had 81% viability when treated with a mixture of Opus@ and Amistar® (O+A) and 64% viability when treated with Opus@ (O) only in the Georgia 5 cultivar. The endophyte viability for the Jesup cultivar was 70% and 69% for the respective treatments. In comparison, AR542 had 63% and 24% viability in the Georgia 5 cultivar and 35% and 11% viability in the Jesup cultivar.

[0098] These results show that there was a clear difference in endophyte viability of both endophytes after the treatment of the various fungicides.

35

Table 6. The percentage of viable endophyte for AR542 and AR584 in two cultivars (Georgia 5 and Jesup) when treated with: no fungicide, Opus@ + Amistar® (O+A) fungicide or Opus@ (O) fungicide alone.

Cultivar	Endophyte	Fungicide treatment		
		nil	O+A	Opus
Georgia 5	AR542	91	63	24
Georgia 5	AR584	92	81	64
Jesup	AR542	90	35	11
Jesup	AR584	93	70	69
LSD 5%			11.6	

[0099] When the averages of the percentage viability of the different endophytes are compared across the treatments and cultivars, the difference between AR584 and AR542 as shown in Table 7, is highly significant. The data shows that AR584 endophyte has a higher viability of 76% compared to AR542, which had 54% viability across the different cultivar and fungicide treatments.

50

Table 7. Endophyte for data averaged over the different cultivar, fungicide and accelerated aging.

Endophyte	E+%
AR584	76
AR542	54

55

EP 1 916 890 B1

(continued)

Endophyte	E+%
LSD 5%	13
F prob	<0.001

[0100] There was no difference in the level of viable endophyte for both AR548 and AR542 endophytes for nil fungicide treatment. However, when the fungicide Opus@ was applied, there was a significant interaction of P<0.001. As shown in Table 8 below, when AR584 endophyte inoculated plants were treated with Opus@, the viable endophyte level (average of nil and 3 day AA) dropped to 60%. In contrast AR542 for the same treatment was 17%.

Table 8. Percentage viability of endophyte for AR584 and AR542 endophyte and treatment of Opus@ fungicide.

	nil	Opus
AR584	93	60
AR542	91	17
LSD 5%	9.2	
F prob	<0.001	

Cultivar effect

[0101] Table 9 below shows that there was also a significant interaction (P=0.045) between the endophyte, cultivar and fungicide. This interaction is the result of the Opus@ fungicide treatment, which considerably dropped the level of viable endophyte. The percentage result of this was less in Georgia cultivar in comparison to the Jesup cultivar.

[0102] However, this difference between AR584 endophyte and the AR542 endophyte in comparison to the cultivar type was comparable over both treatments.

Table 9. Shows the interaction between the various Endophyte, Cultivar and Fungicide.

	Fungicide	AR584 nil	Opus	AR542 nil	Opus
Cultivar	Georgia 5	93	64	91	24
	Jesup	93	55	91	11
LSD 5%	18.4				
F prob	0.045				

Conclusion

[0103] This trial showed that the AR584 endophyte was more robust than AR542 endophyte when treated with various fungicide treatments. As shown, the AR584 endophyte had an improved tolerance to the various fungicide treatments, when compared to AR542 endophyte. This relationship was the same in both tall fescue cultivars tested.

[0104] This robustness is particularly evident when tall fescue combined with AR584 endophyte was exposed to Opus@, a triazole rust control fungicide.

TRIAL 3

[0105] To test whether or not endophytes can be transferred to other species of grass besides Jesup and Georgia 5, results are now included for a different endophyte, AR1 showing that this endophyte can be inoculated into cultivars 'Grasslands Advance' type grass and Kentucky 31 tall fescues.

[0106] Inoculation data for inter-species inoculation of AR1 (*Neotyphodium lolii*) into Tall Fescue *(*Schedonorus phoenix*) and Meadow Fescue (*Festuca pratensis*) hosts. (*previously *Festuca arundinacea*)

Tall Fescue:

plant	fungus	pos	neg	dead	% positive live
949	AR1	19	24	20	44

EP 1 916 890 B1

(continued)

plant	fungus	pos	neg	dead	% positive live
K31	AR1	26	53	37	33
949 = four replicates of 10 - 18 plants K31 = seven replicates of 10 - 19 plants					

Meadow Fescue:

plant	fungus	pos	neg	dead	% positive live
J103	AR1	2	5	13	29
J110	AR1	3	27	37	10
J103 = one replicate of 20 plants J110 = six replicates of 4 - 2- plants					

[0107] It should be appreciated from the above that there is provided an improved endophyte which in combination with a plant cultivar, addresses problems associated with current turf management practice and transport issues.

[0108] Aspects of the present invention have been described by way of example only and it should be appreciated that modifications and additions may be made thereto without departing from the scope thereof.

Claims

1. A method of retaining endophyte viability in a plant or a part thereof produced by a combination a grass cultivar and endophyte combination, wherein the endophyte is AR584 endophyte (AGAL deposit no. NM98/04676 dated 12 May 1998), comprising treating the combination with fungicide.
2. The method as claimed in claim 1 wherein the AR584 endophyte in the combination retains at least a 50% or greater endophyte viability after treatment with fungicide.
3. The method as claimed in claim 1 or claim 2 wherein the fungicide used in treatment has systemic characteristics, foliar disease control characteristics, or activity against soil borne fungicidal pathogens.
4. The method as claimed in any one of claims 1 to 3 wherein the fungicide used in treatment is selected from: the triazole group, strobilurin group or combinations of these two fungicide groups.
5. The method as claimed in any one of claims 1 to 4 wherein at least 65% or greater endophyte viability is retained when the combination is treated with 50:50wt/wt combination of triazole based fungicide and strobilurin based fungicide.
6. The method as claimed in claims 1 to 5 wherein the fungicide used in treatment is triazole based and applied to the combination as a spray at a rate of 1000 ml/ha.
7. The method as claimed in any one of claims 1 to 5 wherein the fungicide used in treatment is strobilurin based and applied as a spray at a rate of 1000 ml/ha.
8. The method as claimed in any one of claims 1 to 7 wherein the fungicide used in treatment is applied to the combination at early flowering of the combination, or is applied at early flowering and re-applied again at late flowering to the combination.
9. The method as claimed in any one of claims 1 to 8 wherein the fungicide is applied to the combination in the form

selected from: a spray, an immersion, a powder, a gel, and combinations thereof.

- 5
10. A method as claimed in any one of claims 1 to 9 comprising subjecting the combination to temperatures of between 15°C and 55°C for a time period of at least 3 days.
11. A method as claimed in claim 10, wherein the combination is produced by inoculating the grass cultivar with an axenic culture of the endophyte.
- 10
12. The method as claimed in claim 10 or 11 wherein the AR584 endophyte in the combination retains at least a 50% or greater endophyte viability.
13. The method as claimed in any one of claims 10 to 12 wherein the endophyte viability is retained when the combination of endophyte and plant or part thereof is subjected to temperatures of 35°C to 45°C.
- 15
14. The method as claimed in any one of claims 10 to 13 wherein endophyte viability is retained when the combination is subjected to a relative humidity of 70% or greater for least 3 days.
- 20
15. A method as claimed in any one of claims 1 to 9, wherein the combination retains at least 60% or greater endophyte viability when the combination is subjected to a temperature of up to 40°C and a relative humidity of up to 100% for at least a time period of 3 days.
16. The method as claimed in claim 15 wherein at least 75% or greater endophyte viability is retained.
- 25
17. A method as claimed in any one of claims 1 to 9, comprising subjecting the combination to a temperature of up to 40°C and a relative humidity of up to 100% for at least a time period of 3 days, wherein the combination retains at least 60% or greater endophyte viability, when the combination is subjected to the temperature of up to 40°C and the relative humidity of up to 100% for at least a time period of 3 days, wherein the combination retains at least a 50% or greater endophyte viability after treatment with fungicide.
- 30
18. The method as claimed in any one of claims 1 to 17 wherein the plant or part thereof is selected from: seed(s), shoot(s), stem(s), flower(s) or root(s) and combinations thereof.
19. The method as claimed in any one of claims 1 to 18 wherein the grass cultivar is a tall fescue plant variety.
- 35
20. The method as claimed in any one of claims 1 to 19 wherein the grass cultivar is a synthetic grass cultivar.

Patentansprüche

- 40
1. Verfahren zum Bewahren der Endophyten-Lebensfähigkeit in einer Pflanze oder einem Teil derselben, die durch eine Kombination aus einer Grassorte und einer Endophytenkombination erzeugt wird, wobei der Endophyt der AR584-Endophyt ist (AGAL-Deposit-Nr. NM98/04676 mit Datum vom 12. Mai 1998), das die Behandlung der Kombination mit Fungizid umfasst.
- 45
2. Verfahren nach Anspruch 1, wobei der AR584-Endophyt in der Kombination eine Lebensfähigkeit von mindestens 50 % oder mehr nach einer Behandlung mit Fungizid bewahrt.
3. Verfahren nach Anspruch 1 oder 2, wobei das Fungizid, das bei der Behandlung verwendet wird, systemische Merkmale, Merkmale zur Kontrolle von Blattkrankheiten oder eine Aktivität gegen bodenbedingte fungizide Pathogene hat.
- 50
4. Verfahren nach einem der Ansprüche 1 bis 3, wobei das Fungizid, das bei der Behandlung verwendet wird, unter folgenden ausgewählt wird: Triazol-Gruppe, Strobilurin-Gruppe oder Kombinationen dieser beiden Fungizid-Gruppen.
- 55
5. Verfahren nach einem der Ansprüche 1 bis 4, wobei eine Lebensfähigkeit des Endophyten von mindestens 65 % oder mehr bewahrt wird, wenn die Kombination mit einer Kombination von 50:50 Gew-% eines Fungizids auf Triazol-Basis und einem Fungizid auf Strobilurin-Basis behandelt wird.

EP 1 916 890 B1

6. Verfahren nach einem der Ansprüche 1 bis 5, wobei das Fungizid, das bei der Behandlung verwendet wird, auf Triazol basiert und auf die Kombination als Spray mit einer Rate von 1000 ml/ha angewendet wird.
- 5 7. Verfahren nach einem der Ansprüche 1 bis 5, wobei das Fungizid, das bei der Behandlung verwendet wird, auf Strobilurin basiert und als Spray mit einer Rate von 1000 ml/ha angewendet wird.
8. Verfahren nach einem der Ansprüche 1 bis 7, wobei das Fungizid, das bei der Behandlung verwendet wird, auf die Kombination bei der frühen Blüte der Kombination angewendet wird oder bei der frühen Blüte angewendet und bei der späten Blüte auf die Kombination wieder angewendet wird.
- 10 9. Verfahren nach einem der Ansprüche 1 bis 8, wobei das Fungizid auf die Kombination in der Form angewendet wird, die aus Folgendem ausgewählt wird: ein Spray, eine Immersion, ein Pulver, ein Gel und Kombinationen derselben.
- 15 10. Verfahren nach einem der Ansprüche 1 bis 9, umfassend das Einwirken von Temperaturen zwischen 15 °C und 55 °C für eine Zeitdauer von mindestens 3 Tagen auf die Kombination.
11. Verfahren nach Anspruch 10, wobei die Kombination durch Impfen der Grassorte mit einer axenischen Kultur des Endophyten erzeugt ist.
- 20 12. Verfahren nach Anspruch 10 oder 11, wobei der AR584-Endophyt in der Kombination eine Lebensfähigkeit des Endophyten von mindestens 50 % oder mehr bewahrt.
13. Verfahren nach einem der Ansprüche 10 bis 12, wobei die Lebensfähigkeit des Endophyten bewahrt wird, wenn die Kombination aus Endophyt und Pflanze oder einem Teil derselben Temperaturen von 35 °C bis 45 °C ausgesetzt wird.
- 25 14. Verfahren nach einem der Ansprüche 10 bis 13, wobei die Lebensfähigkeit des Endophyten bewahrt wird, wenn die Kombination einer relativen Feuchtigkeit von 70 % oder mehr über mindestens 3 Tage ausgesetzt wird.
- 30 15. Verfahren nach einem der Ansprüche 1 bis 9, wobei die Kombination eine Lebensfähigkeit des Endophyten von mindestens 60 % oder mehr bewahrt, wenn die Kombination einer Temperatur von bis zu 40 °C und einer relativen Feuchte von bis zu 100 % für mindestens eine Zeitdauer von 3 Tagen ausgesetzt wird.
- 35 16. Verfahren nach Anspruch 15, wobei die Lebensfähigkeit des Endophyten von mindestens 75 % oder mehr bewahrt wird.
17. Verfahren nach einem der Ansprüche 1 bis 9, umfassend das Einwirken einer Temperatur von bis zu 40 °C und einer relativen Feuchte von bis zu 100 % über eine Zeitdauer von mindestens 3 Tagen auf die Kombination, wobei die Kombination eine Lebensfähigkeit des Endophyten von mindestens 60 % oder mehr bewahrt, wenn die Kombination der Temperatur von bis zu 40 °C und der relativen Feuchte von bis zu 100 % über eine Zeitdauer von mindestens 3 Tagen ausgesetzt wird, wobei die Kombination eine Entwicklungsfähigkeit des Endophyten von mindestens 50 % oder mehr nach einer Behandlung mit Fungizid bewahrt.
- 40 18. Verfahren nach einem der Ansprüche 1 bis 17, wobei die Pflanze oder ein Teil derselben aus Folgendem ausgewählt wird: Saat(en), Trieb(e), Stängel, Blüte(n) oder Wurzel(n) und Kombinationen derselben.
19. Verfahren nach einem der Ansprüche 1 bis 18, wobei die Grassorte eine große Schwingelpflanzenvarietät ist.
- 50 20. Verfahren nach einem der Ansprüche 1 bis 19, wobei die Grassorte eine synthetische Grassorte ist.

Revendications

- 55 1. Méthode de conservation de la viabilité des endophytes dans une plante ou une partie de celle-ci produite par une combinaison d'un cultivar de graminée et d'une combinaison d'endophytes, dans laquelle l'endophyte est l'endophyte AR584 (dépôt AGAL n° NM98/04676 daté du 12 mai 1998), comprenant le traitement de la combinaison par un fongicide.

EP 1 916 890 B1

2. Méthode selon la revendication 1, dans laquelle l'endophyte AR584 dans la combinaison conserve au moins 50% ou plus de viabilité des endophytes après traitement par un fongicide.
- 5 3. Méthode selon la revendication 1 ou la revendication 2, dans laquelle le fongicide utilisé dans le traitement possède des caractéristiques systémiques, des caractéristiques de contrôle de maladies foliaires, ou une activité contre les pathogènes fongicides du sol.
- 10 4. Méthode selon l'une quelconque des revendications 1 à 3, dans laquelle le fongicide utilisé dans le traitement est choisi parmi : le groupe des triazoles, le groupe des strobilurines ou des combinaisons de ces deux groupes de fongicides.
- 15 5. Méthode selon l'une quelconque des revendications 1 à 4, dans laquelle au moins 65% ou plus de la viabilité des endophytes est conservée lorsque la combinaison est traitée par une combinaison de 50:50 p/p d'un fongicide à base de triazole et d'un fongicide à base de strobilurine.
- 20 6. Méthode selon les revendications 1 à 5, dans laquelle le fongicide utilisé dans le traitement est à base de triazole et est appliqué à la combinaison sous forme de pulvérisation à un taux de 1000 ml/ha.
- 25 7. Méthode selon l'une quelconque des revendications 1 à 5, dans laquelle le fongicide utilisé dans le traitement est à base de strobilurine et est appliqué sous forme de pulvérisation à un taux de 1000 ml/ha.
- 30 8. Méthode selon l'une quelconque des revendications 1 à 7, dans laquelle le fongicide utilisé dans le traitement est appliqué à la combinaison à la floraison précoce de la combinaison, ou est appliqué à la floraison précoce et réappliqué de nouveau à la floraison tardive à la combinaison.
- 35 9. Méthode selon l'une quelconque des revendications 1 à 8, dans laquelle le fongicide est appliqué à la combinaison sous une forme choisie parmi : une pulvérisation, une immersion, une poudre, un gel, et des combinaisons de ceux-ci.
- 40 10. Méthode selon l'une quelconque des revendications 1 à 9, comprenant la soumission de la combinaison à des températures comprises entre 15°C et 55°C pendant une période de temps d'au moins 3 jours.
- 45 11. Méthode selon la revendication 10, dans laquelle la combinaison est produite par l'inoculation du cultivar de graminée par une culture axénique de l'endophyte.
- 50 12. Méthode selon la revendication 10 ou 11, dans laquelle l'endophyte AR584 dans la combinaison conserve au moins 50% ou plus de viabilité des endophytes.
- 55 13. Méthode selon l'une quelconque des revendications 10 à 12, dans laquelle la viabilité des endophytes est conservée lorsque la combinaison des endophytes et de la plante ou d'une partie de celle-ci est soumise à des températures allant de 35°C à 45°C.
14. Méthode selon l'une quelconque des revendications 10 à 13, dans laquelle la viabilité des endophytes est conservée lorsque la combinaison est soumise à une humidité relative de 70% ou plus pendant au moins 3 jours.
15. Méthode selon l'une quelconque des revendications 1 à 9, dans laquelle la combinaison conserve au moins 60% ou plus de viabilité des endophytes lorsque la combinaison est soumise à une température allant jusqu'à 40°C et une humidité relative allant jusqu'à 100% pendant au moins une période de temps de 3 jours.
16. Méthode selon la revendication 15, dans laquelle au moins 75% ou plus de la viabilité des endophytes est conservée.
17. Méthode selon l'une quelconque des revendications 1 à 9, comprenant la soumission de la combinaison à une température allant jusqu'à 40°C et une humidité relative allant jusqu'à 100% pendant au moins une période de temps de 3 jours, où la combinaison conserve au moins 60% ou plus de viabilité des endophytes, lorsque la combinaison est soumise à la température allant jusqu'à 40°C et l'humidité relative allant jusqu'à 100% pendant au moins une période de temps de 3 jours, où la combinaison conserve au moins 50% ou plus de viabilité des endophytes après traitement par un fongicide.
18. Méthode selon l'une quelconque des revendications 1 à 17, dans laquelle la plante ou une partie de celle-ci est

EP 1 916 890 B1

choisie parmi : une ou plusieurs graines, une ou plusieurs pousses, une ou plusieurs tiges, une ou plusieurs fleurs ou une ou plusieurs racines, et des combinaisons de celles-ci.

5 19. Méthode selon l'une quelconque des revendications 1 à 18, dans laquelle le cultivar de graminée est une variété de fétuque élevée.

10 20. Méthode selon l'une quelconque des revendications 1 à 19, dans laquelle le cultivar de graminée est un cultivar de graminée synthétique.

10

15

20

25

30

35

40

45

50

55




FIGURE 1




FIGURE 2




FIGURE 3

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- US 6111170 A [0003]
- RU 2201678 C2 [0004]
- US 200515002 B [0012]
- WO 2002013616 A [0012]
- WO 2004106487 A [0013]

Non-patent literature cited in the description

- **TIMPER, P. ; GATES, R. N. ; ; BOUTON J. H.** Response of *Pratylenchus* spp. In tall fescue infected with different strains of the fungal endophyte *Neotyphodium coenophialum*. *Nematology*, 2005, vol. 7 (1), 105-110 [0011]

FÜENDOFITA KITERJESZTETT TULAJDONSÁGAI

Szabadalmi igénypontok

1. Eljárás növényben vagy annak egy részében lévő endofita-életképesség fenntartásához, melyet fü kultivár és endofita kombináció kombinálása révén állítunk elő, ahol az endofita AR584 endofita (NM98/04676 számú AGAL letét, 1998. május 12.) és amely eljárás a kombináció fungiciddel való kezelését tartalmazza.

2. Az 1. igénypont szerinti eljárás, ahol a kombinációban lévő AR584 endofita, a fungiciddel való kezelés után, legalább 50%-os vagy nagyobb endofita-életképességet tart fenn.

3. Az 1. vagy a 2. igénypont szerinti eljárás, ahol a kezeléshez használt fungicid szisztémás tulajdonságokkal, levélbetegség szabályozásának tulajdonságaival vagy talajban lévő fungicid patogének elleni aktivitással rendelkezik.

4. Az 1-3. igénypontok bármelyike szerinti eljárás, ahol a kezeléshez használt fungicidet a következő csoportok közül választjuk ki: triazol csoport, strobilurin csoport vagy ennek a két fungicid csoportnak a kombinációja.

5. Az 1-4. igénypontok bármelyike szerinti eljárás, ahol a legalább 65%-os vagy nagyobb endofita életképességet akkor tartjuk fenn, mikor a kombinációt triazol alapú fungicid és strobilurin alapú fungicid 50:50 tömeg%-os kombinálásával kezeljük.

6. Az 1-5. igénypontok bármelyike szerinti eljárás, ahol a kezeléshez használt fungicid triazol alapú és a kombinációhoz 1000 ml/ha arányú sprayként alkalmazzuk.

7. Az 1-5. igénypontok bármelyike szerinti eljárás, ahol a kezeléshez használt fungicid strobilurin alapú és 1000 ml/ha arányú sprayként alkalmazzuk.

8. Az 1-7. igénypontok bármelyike szerinti eljárás, ahol a kezeléshez használt fungicidet a kombinációhoz a kombináció korai virágzása esetén alkalmazzuk vagy korai virágzás esetén, és késő virágzás esetén pedig ismételtén alkalmazzuk a kombinációhoz.

9. Az 1-8. igénypontok bármelyike szerinti eljárás, ahol a kombinációhoz alkalmazott fungicidet a következő formák közül választjuk ki: spray, immerzió, por, gél és ezeknek a kombinációja.

10. Az 1-9. igénypontok bármelyike szerinti eljárás, amely tartalmazza a kombináció kitételét 15°C és 35°C közötti hőmérsékletek, legalább 3 napos időtartamon keresztül.



11. A 10. igénypont szerinti eljárás, ahol a kombinációt a fű kultivár endofita parazitamentes kultúrájával való oltása révén állítjuk elő.

12. A 10. vagy a 11. igénypont szerinti eljárás, ahol a kombinációban lévő AR584 endofita legalább 50%-os vagy nagyobb endofita-életképességet tart fenn.

13. A 10-12. igénypontok bármelyike szerinti eljárás, ahol az endofita életképesség fennmarad, amikor endofita és növény vagy annak egy részének a kombinációját 35°C-45°C közötti hőmérsékleten tartjuk.

14. A 10-13. igénypontok bármelyike szerinti eljárás, ahol az endofita életképesség fennmarad, amikor a kombinációt 70%-os vagy nagyobb relatív páratartalomnak tesszük ki, legalább 3 napra.

15. Az 1-9. igénypontok bármelyike szerinti eljárás, ahol a kombináció legalább 60%-os vagy nagyobb endofita életképességet tart fenn, amikor a kombinációt legfeljebb 40°C-os hőmérsékletnek és legfeljebb 100%-os relatív páratartalomnak tesszük ki, legalább 3 napra.

16. A 15. igénypont szerinti eljárás, ahol legalább 75%-os vagy nagyobb endofita életképesség marad fenn.

17. Az 1-9. igénypontok bármelyike szerinti eljárás, amely tartalmazza a kombináció kitételét legfeljebb 40°C-os hőmérsékletnek és legfeljebb 100%-os relatív páratartalomnak, legalább 3 napig, ahol a kombináció legalább 60%-os vagy nagyobb endofita életképességet tart fenn, amikor a kombinációt legfeljebb 40°C-os hőmérsékletnek és legfeljebb 100%-os relatív páratartalomnak tesszük ki legalább 3 napra, ahol a kombináció legalább 50%-os vagy nagyobb endofita-életképességet tart fenn fungiciddel való kezelés után.

18. Az 1-17. igénypontok bármelyike szerinti eljárás, ahol a növényt vagy annak egy részét a következők közül választjuk ki: mag(ok), hajtás(ok), szár(ak) vagy gyökér(ek) és ezeknek a kombinációja.

19. Az 1-18. igénypontok bármelyike szerinti eljárás, ahol a fű kultivár nádképző csekesz növényfajta.

20. Az 1-19. igénypontok bármelyike szerinti eljárás, ahol a fű kultivár egy szintetikus fű kultivár.