(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau

(43) International Publication Date
21 April 2005 (21.04.2005)

PCT

(10) International Publication Number
WO 2005/035507 A2

(51) International Patent Classification[7]: C07D 239/00

(21) International Application Number:
PCT/US2004/033430

(22) International Filing Date: 8 October 2004 (08.10.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/510,804 10 October 2003 (10.10.2003) US

(71) Applicant *(for all designated States except US)*: **BAYER PHARMACEUTICALS CORPORATION** [US/US]; 400 Morgan Lane, West Haven, Connecticut 06516 (US).

(72) Inventors; and

(75) Inventors/Applicants *(for US only)*: **DIXON, Julie, A.** [US/US]; 40 Ridgewood Road, Bethany, Connecticut 06524 (US). **NAGARATHNAM, Dhanapalan** [US/US]; 52 Virginia Rail Drive, Bethany, Connecticut 06524 (US). **ZHANG, Lei** [CN/US]; 45 Tanglewood Drive, Hamden, Connecticut 06518 (US). **WANG, Chunguang** [CN/US]; 27 Timberwood Trail, Hamden, Connecticut 06514 (US). **YI, Lin** [CN/US]; 58 Magnolia Road, Milford, Connecticut 06460 (US). **CHEN, Yuanwei** [US/US]; 15 Blue Ridge Lane, North Haven, Connecticut 06473 (US). **CHEN, Jianqing** [CA/US]; 117 Frederick Street, Apt. #2-L, New Haven, Connecticut 06515 (US). **BEAR, Brian** [US/US]; 373 Acorn Lane, Milford, Connecticut 06460 (US). **BRANDS, Michael** [DE/US]; 35 Deer Pond Trail, Hamden, Connecticut 06518 (US). **HILLISCH, Alexander** [AT/DE]; Zum Teller Hof 44, 42553 Velbert (DE). **BIERER, Donald** [US/US]; 46 Hilltop Road, Bethany, Connecticut 06524 (US). **WANG, Ming** [US/US]; 32 Milford Hunt Lane, Milford, Connecticut 06460 (US). **FU, Wenlang** [CN/US]; 12133 Town Walk Drive, Hamden, Connecticut 06518 (US). **HENTEMANN, Martin, F.** [US/US]; 80 Morris Street, Hamden, Connecticut 06517 (US). **BULLION, Ann-Marie** [US/US]; 104 D East Broadway, Milford, Connecticut 06460 (US).

(74) Agents: **GREENMAN, Jeffrey, M.** et al.; Bayer Pharmaceuticals Corporation, 400 Morgan Lane, West Haven, Connecticut 06516 (US).

(81) Designated States *(unless otherwise indicated, for every kind of national protection available)*: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.


(84) Designated States *(unless otherwise indicated, for every kind of regional protection available)*: ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE,*

[Continued on next page]

(54) Title: PYRIMIDINE DERIVATIVES FOR TREATMENT OF HYPERPROLIFERATIVE DISORDERS



(I)

(57) Abstract: Pyrimidine derivatives of formula (I) in which J and Y represent aromatic or heteroaromatic rings; R^2, G, G', and G" represent substituent groups and R^{2a} represents H or halogen; L represents a linking group; and M represents CH or N. Pharmaceutical compositions containing these compounds, and methods of using these compounds in treatment of hyperproliferative diseases such as cancer are also disclosed and claimed.

WO 2005/035507 A2

BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)

— *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii)) for all designations*

— *of inventorship (Rule 4.17(iv)) for US only*

Published:

— *without international search report and to be republished upon receipt of that report*

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Pyrimidine Derivatives for Treatment of Hyperproliferative Disorders

FIELD:

This application relates to small molecule heterocyclic pharmaceuticals, and more particularly, to amino-substituted pyrimidine derivatives having cytotoxic activity.

BACKGROUND:

Nitrogen-containing heterocycles such as pyrimidine derivatives have been disclosed in patent and non-patent publications as having a variety of pharmaceutical properties and utilities. Several such publications are listed below.

WO 03/062225 (Bayer) relates to pyrimidine derivatives as rho-kinase inhibitors, and their use in treatment of rho-kinase mediated conditions including cancer.

WO 2001/87845 (Fujisawa) relates to N-containing heterocyclic compounds having 5-HT antagonistic activity. These compounds are stated as being useful for treating or preventing central nervous system disorders.

WO 95/10506 (Du Pont Merck) relates to 1N-alkyl-N-arylpyrimidinamines and derivatives thereof, which are stated to inhibit the corticopropin releasing factor (CRF) peptide and to be useful for treatment of psychiatric disorders and neurological diseases.

WO 2004/048365 (Chiron) relates to 2,4,6-trisubstituted pyrimidines as phosphotidylinositol (PI) 3-kinase inhibitors and their use in treatment of cancer.

WO 2004/000820 (Cellular Genomics) relates to N-containing heterocycles and other compounds as kinase modulators, and their use in treatment of numerous kinase-associated disorders including cancer.


WO 01/62233 (Hoffmann La Roche) relates to nitrogen-containing heterocycles and their use in treatment of diseases modulated by the adenosine receptor.

US 2004/0097504 (Vertex) relates to nitrogen-containing heterocycles useful in treatment of various protein kinase-mediated disorders.

The pharmaceutical field is always interested in identifying new pharmaceutically active compounds. Such materials are the subject of the present application.

Compounds of the invention

In a first embodiment, this invention relates to compounds of Formula (I)



(I)

wherein

R^1 represents H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^2 represents $(C_1\text{-}C_3)$alkyl, cyclopropyl, $O(C_1\text{-}C_3)$alkyl, or NR^3R^4

wherein R^3 and R^4 are H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^{2a} represents H or halogen;


M represents CH or N;

L represents a carbonyl group, O, NR^5 , CR^6R^7 , or $(C_2\text{-}C_3)$alkylenyl which is optionally substituted up to twice by groups independently selected from halogen and OH; wherein

R^5 is H or $(C_1\text{-}C_3)$alkyl; and


R^6 and R^7 are independently H, CH_3, halogen, or OH;

J represents an aromatic or heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

wherein R^8 represents H or $(C_1\text{-}C_3)$alkyl;

G" represents a substituent selected from the group consisting of $(C_1\text{-}C_3)$alkyl, cyclopropyl, $O(C_1\text{-}C_3)$alkyl, halogen, CF_3, CN and CO_2R^9 ;

wherein

R^9 represents H or $(C_1\text{-}C_3)$alkyl; and

m represents the number of substituents G", and is 0, 1, or 2;

G represents a substituent located on ring J;

G' represents a substituent located on ring Y;

n represents the number of substituents G; and

n' represents the number of substituents G' ;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,

2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G11, to a maximum total of 3 substituents on rings J and Y, and

3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12-G37;

and subject to the further provisos

4) when J is phenyl, G is other than OH or alkylthio; and when J is phenyl or pyridyl, n is 1, 2, or 3;

5) when J is phenyl, and G is G4 shown below, then R^2 is NR^3R^4;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1\text{-}C_4)$alkyl which optionally is substituted up to two times by $O(C_1\text{-}C_2)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_5)$alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3 ;

G6) $NHC(O)(C_1\text{-}C_3)$alkyl ;

G7) $NHSO_2(C_1\text{-}C_3)$alkyl ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,

$NR^{12}R^{13}$ wherein


R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula



wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,


or by

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G9) $(CH_2)_a$-$NR^{19}R^{20}$ wherein

R^{19} and R^{20} are independently H, $(C_1$-$C_5)$alkyl, or $(C_3$-$C_6)$cycloalkyl, or may be joined to form a saturated 5-6-membered N-containing ring; and

the subscript "a" is an integer of 1-4;

G10)  wherein

Q' is O or NR^{21} ;

R^{21} is H, $(C_1$-$C_3)$alkyl, or cyclopropyl; and

the subscript "b" is an integer of 1-3;

G11) $CH_2NR^{22}(CH_2)_cOCH_3$ wherein

R^{22} is H, $(C_1$-$C_3)$alkyl, or cyclopropyl; and

the subscript "c" is an integer of 2-4;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2$-$C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1$-$C_3)$alkyl;

G13) CN ;

G14) NO_2 ;

G15) cyclopropyl ;


G16) OR^{27}, wherein


R^{27} represents phenyl or benzyl;

G17) $S(C_1$-$C_3)$alkyl;

G18) $CH=CH-(CH_2)_{1-3}-OR^5$; wherein

R^5 represents H or (C_1-C_3)alkyl;

G19)  ;


G20)  ;

G21) $C(O)NR^{28}R^{29}$, wherein

R^{28} and R^{29} are independently selected from

H,

cyclopropyl, provided that both R^{28} and R^{29} are not simultaneously cyclopropyl,


, provided that this group does not constitute both R^{28} and R^{29} simultaneously,

and

(C_1-C_3)alkyl which is optionally substituted up to two times by OH;

or

R^{28} and R^{29} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by OH, or by (C_1-C_3)alkyl which in turn is optionally substituted up to two times by OH or $O(C_1-C_3)$alkyl;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is

H,


cyclopropyl, or

$(C_1$-$C_3)$alkyl which is optionally substituted once by halogen, OH, or $O(C_1$-$C_3)$alkyl;

G23) O-$(CH_2)_d$-$NR^{31}R^{32}$ wherein

R^{31} and R^{32} are independently H, $(C_1$-$C_3)$alkyl, or cyclopropyl, or may be joined to form a saturated 5-6-membered N-containing ring; and


the subscript "d" is an integer of 2-4;

G24)  wherein

the subscript "e" is an integer of 2-3; and

Q''' is O or NR^{33} ; and

R^{33} is H, $(C_1$-$C_3)$alkyl, or cyclopropyl;

G25)  wherein

Q^{iv} is O or NR^{34} ; and

R^{34} is H, $(C_1$-$C_3)$alkyl, or cyclopropyl;

G26) $C(O)NR^{35}(CH_2)_fOR^{36}$ wherein

R^{35} is H, $(C_1$-$C_3)$alkyl, or cyclopropyl;

R^{36} is $(C_1$-$C_6)$alkyl optionally substituted up to two times by halogen, OH, or $O(C_1$-$C_3)$alkyl, and

the subscript "f" is an integer of 2-4;

G27) CO_2R^{37} wherein

R^{37} is H or $(C_1$-$C_3)$alkyl;

G28) phenyl, which is optionally substituted by up to 2 groups selected from halogen, $(C_1$-$C_3)$alkyl, OR^{38}, CN, CF_3, and $NR^{39}R^{40}$

wherein

R^{38} represents H or $(C_1$-$C_3)$alkyl; and

R^{39} and R^{40} represent H or $(C_1$-$C_3)$alkyl;

G29) $NR^{41}SO_2NR^{42}R^{43}$ wherein

R^{41} represents H, or $(C_1\text{-}C_4)$alkyl, and

R^{42} and R^{43} independently represent H, CH_3, or $(C_2\text{-}C_3)$alkyl which may optionally be substituted once by -OH or $NR^{44}R^{45}$, wherein

R^{44} and R^{45} independently represent H or $(C_1\text{-}C_3)$alkyl;

G30) $OC(O)\text{-}CH_2\text{-}NR^{46}R^{47}$ wherein

R^{46} and R^{47} independently represent H, $(C_1\text{-}C_3)$alkyl, or CO_2(t-butyl), provided that R^{46} and R^{47} are not both simultaneously CO_2(t-butyl);

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl;

G32) $C(O)\text{-}(C_1\text{-}C_3)$alkyl;

G33) $(CH_2)_g\text{-}N(R^{55})\text{-}C(O)\text{-}R^{56}$ wherein

g represents 1, 2, or 3;


R^{55} represents H or $(C_1\text{-}C_3)$alkyl;

R^{56} represents

$(C_1\text{-}C_3)$alkyl optionally substituted up to two times by OR^{57} or $NR^{58}R^{59}$, wherein

R^{57} represents H or $(C_1\text{-}C_3)$alkyl, and

R^{58} and R^{59} each represents H or $(C_1\text{-}C_3)$alkyl,

or R^{56} represents  wherein

R^{60} represents halogen, $(C_1\text{-}C_3)$alkyl, $O(C_1\text{-}C_3)$alkyl, CN, OH, CF_3, or $NR^{61}R^{62}$, wherein

R^{61} and R^{62} represent H or $(C_1\text{-}C_3)$alkyl; and

h represents 0, 1, or 2;


G34) $(CH_2)_i\text{-}N(R^{63})\text{-}C(O)\text{-}NR^{64}R^{65}$ wherein

i represents 1, 2, or 3;

R^{63} represents H or $(C_1\text{-}C_3)$alkyl;


R^{64} and R^{65} each represents H or $(C_1\text{-}C_3)$alkyl;

or

R^{64} and R^{65} may be joined to form  wherein

Q^V represents CH_2, O or NR^{66} wherein

R^{66} represents H or $(C_1\text{-}C_3)$alkyl;

G35)  wherein

j represents 1, 2, or 3;

R^{67}represents H or $(C_1\text{-}C_3)$alkyl; and

R^{68} represents H or $(C_1\text{-}C_3)$alkyl;

G36) $(CH_2)_k\text{-}N(R^{69})\text{-}SO_2\text{-}R^{70}$ wherein

k represents 1, 2, or 3;

R^{69} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{70} represents $(C_1\text{-}C_4)$alkyl, or phenyl which is optionally substituted up to perhalo by halogen or up to three times by OR^{71}, CN, CF_3, or $NR^{72}R^{73}$, wherein

R^{71} represents H or $(C_1\text{-}C_3)$alkyl; and


R^{72} and R^{73} each represents H or $(C_1\text{-}C_3)$alkyl;

G37) $CH{=}CH\text{-}(CH_2)_{1\text{-}3}\text{-}NR^{74}R^{75}$ wherein

R^{74} and R^{75} represent H or $(C_1\text{-}C_3)$alkyl;

or a pharmaceutically acceptable salt, solvate, solvate of a salt, or stereoisomer thereof.

In a second embodiment, the invention relates to a compound of the above first embodiment, having the structure




(I)

wherein the definitions of the variables are as given above in the first embodiment, except that in this second embodiment

R^1 represents H;


M represents CH;

J represents a heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,

2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G5 and G8, to a maximum total of 3 substituents on rings J and Y, and

3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12, G13, G22, G29, and G31;

and subject to the further proviso

4) when J is pyridyl, n is 1, 2, or 3;

and proviso 5 does not apply;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1\text{-}C_4)$alkyl which optionally is substituted up to two times by $O(C_1\text{-}C_2)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_5)$alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,

or by


$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G13) CN ;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is

H,

cyclopropyl, or

$(C_1\text{-}C_3)$alkyl which is optionally substituted once by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G29) $NR^{41}SO_2NR^{42}R^{43}$ wherein

R^{41}represents H, or $(C_1\text{-}C_4)$alkyl, and

R^{42} and R^{43} independently represent H, CH_3, or $(C_2\text{-}C_3)$alkyl which may optionally be substituted once by -OH or $NR^{44}R^{45}$, wherein

R^{44} and R^{45} independently represent H or $(C_1\text{-}C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein


R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

In a third embodiment the invention relates to a compound of the above second embodiment, having the structure



(I)

wherein the definitions of the variables are as given above in the second embodiment, except that in this third embodiment

R^1 represents H;

R^2 represents $O(C_1\text{-}C_3)$alkyl or NR^3R^4

wherein R^3 and R^4 are H or $(C_1\text{-}C_3)$alkyl;

R^{2a} represents H;

L represents O or CR^6R^7 wherein

R^6 and R^7 are independently H, CH_3 , or OH;

G" represents a substituent selected from the group consisting of $O(C_1\text{-}C_3)$alkyl, halogen, and CF_3;

n and n' are independently 0 or 1, and provisos 1-3 do not apply;

G and G' moieties are independently selected from the group consisting of:

G1) Cl or F;

G2) $O(C_1\text{-}C_3)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_3)$alkyl, which is optionally substituted up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and


R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G13) CN ;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is H or $(C_1\text{-}C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1-3}$-CO_2H ,

$O(C_2$-$C_4)$alkyl,

$(CH_2)_{1-4}$-$NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1$-$C_3)$alkyl,


or

$CH(R^{52})$-$NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1-4}$-NH_2, CH_2OH, $CH(CH_3)OH$, or $(C_1$-$C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1$-$C_3)$alkyl.

In a fourth embodiment, the invention relates to a compound of the above first embodiment, having the structure




(I)

wherein the definitions of the variables are as given above in the first embodiment, except that in this fourth embodiment

R^1 represents H;


M represents CH;

J represents a heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,

2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G5 and G8, to a maximum total of 3 substituents on rings J and Y, and

3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12, G21, G25, G26, and G31;

and subject to the further proviso

4) when J is pyridyl, n is 1, 2, or 3;

and proviso 5 does not apply;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1-C_4)$alkyl which optionally is substituted up to two times by $O(C_1-C_2)$alkyl;

G3) OH ;

G4) (C_1-C_5)alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,

or by

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein


R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G21) $C(O)NR^{28}R^{29}$, wherein

R^{28} and R^{29} are independently selected from

H,

cyclopropyl, provided that both R^{28} and R^{29} are not simultaneously cyclopropyl,




, provided that this group does not constitute both R^{28} and R^{29} simultaneously,

and

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by OH;

or

R^{28} and R^{29} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by OH, or by $(C_1\text{-}C_3)$alkyl which in turn is optionally substituted up to two times by OH or $O(C_1\text{-}C_3)$alkyl;

G25)  wherein

Q^{iv} is O or NR^{34} ; and

R^{34} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

G26) $C(O)NR^{35}(CH_2)_fOR^{36}$ wherein

R^{35} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^{36} is $(C_1\text{-}C_6)$alkyl optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl, and

the subscript "f" is an integer of 2-4; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl,


or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

In a fifth embodiment, the invention relates to a compound of the above fourth embodiment, having the structure



(I)

wherein the definitions of the variables are as given above in the fourth embodiment, except that in this fifth embodiment

R^1 represents H;

R^2 represents $O(C_1\text{-}C_3)$alkyl or NR^3R^4

wherein R^3 and R^4 are H or $(C_1\text{-}C_3)$alkyl;

R^{2a} represents H;

L represents O or CR^6R^7, wherein

R^6 and R^7 are independently H, CH_3, or OH;

G" represents a substituent selected from the group consisting of $O(C_1\text{-}C_3)$alkyl, halogen, and CF_3;

n and n' are independently 0 or 1, and provisos 1-3 do not apply;

G and G' moieties are independently selected from the group consisting of:

G1) Cl or F;

G2) $O(C_1\text{-}C_3)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_3)$alkyl, which is optionally substituted up to three times by halogen;

G5) OCF_3;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;


G21) $C(O)NR^{28}R^{29}$, wherein

R^{28} and R^{29} are independently selected from

H

and

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by OH;

G25)  wherein

Q^{iv} is O or NR^{34}; and

R^{34} is H or $(C_1\text{-}C_3)$alkyl;

G26) $C(O)NR^{35}(CH_2)_fOR^{36}$ wherein

R^{35} is H or $(C_1\text{-}C_3)$alkyl;

R^{36} is $(C_1\text{-}C_6)$alkyl optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl, and

the subscript "f" is an integer of 2-4; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein


R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

In a sixth embodiment, the invention relates to a compound of the above first embodiment, having the structure




(I)

wherein the definitions of the variables are as given above in the first embodiment, except that in this sixth embodiment

R^1 represents H;


M represents CH;

J represents an aromatic or heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,

2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G5 and G8, to a maximum total of 3 substituents on rings J and Y, and

3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12, G22, and G31;

and subject to the further proviso

4) when J is pyridyl, n is 1, 2, or 3;

and proviso 5 does not apply;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1\text{-}C_4)$alkyl which optionally is substituted up to two times by $O(C_1\text{-}C_2)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_5)$alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,


or by

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is

H,

cyclopropyl, or

$(C_1\text{-}C_3)$alkyl which is optionally substituted once by halogen, OH, or $O(C_1\text{-}C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl,


or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

In a seventh embodiment, the invention relates to a compound of the above sixth embodiment, having the structure



(I)

wherein the definitions of the variables are as given above in the sixth embodiment, except that in this seventh embodiment

R^1 represents H;

R^2 represents $O(C_1\text{-}C_3)$alkyl, or NR^3R^4

wherein R^3 and R^4 are H or $(C_1\text{-}C_3)$alkyl;

R^{2a} represents H;

L represents O or CR^6R^7, wherein

R^6 and R^7 are independently H, CH_3, or OH;

G" represents a substituent selected from the group consisting of $O(C_1\text{-}C_3)$alkyl, halogen, and CF_3 ;

n and n' are independently 0 or 1, and provisos 1-3 do not apply;

G and G' moieties are independently selected from the group consisting of:

G1) Cl or F;

G2) $O(C_1\text{-}C_3)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_3)$alkyl, which is optionally substituted up to three times by halogen;

G5) OCF_3;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein


Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is H or $(C_1\text{-}C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

Pharmaceutically acceptable salts of these compounds as well as commonly used prodrugs of these compounds such as, for example, O-acyl derivatives of invention compounds which contain hydroxy groups, ester derivatives of invention compounds which contain carboxyl groups, and amide derivatives of invention compounds which contain amino groups, are also within the scope of the invention.

It is to be understood that:

1) in compounds of the invention in which an alkyl moiety may bear substituents such as amino, hydroxyl, alkoxy, and halogen groups, a single carbon atom of this alkyl moiety may not simultaneously bear two groups independently selected from amino, hydroxyl, and alkoxy; and where this alkyl moiety bears a halogen, it may not simultaneously also bear an amino, hydroxyl, or alkoxy substituent.

2) in compounds of the invention in which any moiety is defined in terms of a numerical range of atoms and this moiety is further permitted to bear up to a certain number

of substituents, if the total number of substituents possible exceeds the number of valences available for moieties at the lower end of the defined numerical range of atoms, then the number of substituents is limited to the number of available valences. For example, if a (C_1-C_3)alkyl moiety if defined as optionally bearing up to three halogens and up to two other defined substituents, this means that a C_1-alkyl group could bear up to three substituents (the number of available valences), all of which could be halogen, but no more than two of which could be other defined substituent groups.

The compounds of Formula (I) may contain one or more asymmetric centers, depending upon the location and nature of the various substituents desired. Asymmetric carbon atoms may be present in the (*R*) or (*S*) configuration. Preferred isomers are those with the absolute configuration which produces the compound of Formula (I) with the more desirable biological activity. In certain instances, asymmetry may also be present due to restricted rotation about a given bond, for example, the central bond adjoining two aromatic rings of the specified compounds.

It is intended that all isomers (including enantiomers and diastereomers), either by nature of asymmetric centers or by restricted rotation as described above, as separated, pure or partially purified isomers or racemic mixtures thereof, be included within the scope of the instant invention. The purification of said isomers and the separation of said isomeric mixtures may be accomplished by standard techniques known in the art.

The terms identified above have the following meaning throughout:

The term "optionally substituted" means that the moiety so modified may have from none to up to at least the highest number of substituents indicated. The substituent may replace any H atom on the moiety so modified as long as the replacement is chemically possible and chemically stable. When there are two or more substituents on any moiety, each substituent is chosen independently of any other substituent and can, accordingly, be the same or different.

The term "halogen" means an atom selected from Cl, Br, F, and I.

The terms "(C_1-C_2)alkyl," "(C_1-C_3)alkyl" "(C_1-C_4)alkyl" "(C_1-C_5)alkyl," and "(C_1-C_6)alkyl" mean linear or branched saturated carbon groups having from about 1 to about 2, about 3, about 4, about 5 or about 6 C atoms, respectively. Such groups include, but are not limited to, methyl, ethyl, *n*-propyl, isopropyl, *sec*-butyl, *n*-hexyl, and the like.

The term "alkylenyl" means a divalent linear or branched saturated carbon chain, usually having from about 1 to about 3 carbon atoms in this application. Such chains

include, but are not limited to methylene (-CH_2-), ethylenyl (-CH_2CH_2)-, and propylenyl (-$CH_2CH_2CH_2$-) and the like.

The term "(C_3-C_6)cycloalkyl" means a saturated monocyclic alkyl group of from about 3 to about 6 carbon atoms and includes such groups as cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and the like.

Pharmaceutical compositions

The invention also relates to pharmaceutical compositions comprising at least one of the compounds of the invention, or a salt or prodrug thereof, in a pharmaceutically acceptable carrier.

Method of treating hyperproliferative disorders

The present invention also relates to a method of using the compounds described above, including salts, prodrugs, and corresponding pharmaceutical compositions thereof, to treat mammalian hyperproliferative disorders. This method comprises administering to a patient an amount of a compound of this invention, or a pharmaceutically acceptable salt or prodrug thereof, which is effective to treat the patient's hyperproliferative disorder. A patient, for the purpose of this invention, is a mammal, including a human, in need of treatment for a particular hyperproliferative disorder. A pharmaceutically effective amount of a compound or composition is that amount which produces a desired result or exerts an influence on the particular hyperproliferative disorder being treated.

Hyperproliferative disorders include but are not limited to solid tumors, such as cancers of the breast, respiratory tract, brain, reproductive organs, digestive tract, urinary tract, eye, liver, skin, head and neck, thyroid, parathyroid and their distant metastases. These disorders also include lymphomas, sarcomas, and leukemias.

Examples of breast cancer include, but are not limited to invasive ductal carcinoma, invasive lobular carcinoma, ductal carcinoma in situ, and lobular carcinoma in situ.

Examples of cancers of the respiratory tract include, but are not limited to small-cell and non-small-cell lung carcinoma, as well as bronchial adenoma and pleuropulmonary blastoma.

Examples of brain cancers include, but are not limited to brain stem and hypophtalmic glioma, cerebellar and cerebral astrocytoma, medulloblastoma, ependymoma, as well as neuroectodermal and pineal tumor.

Tumors of the male reproductive organs include, but are not limited to prostate and testicular cancer. Tumors of the female reproductive organs include, but are not limited to endometrial, cervical, ovarian, vaginal, and vulvar cancer, as well as sarcoma of the uterus.

Tumors of the digestive tract include, but are not limited to anal, colon, colorectal, esophageal, gallbladder, gastric, pancreatic, rectal, small-intestine, and salivary gland cancers.

Tumors of the urinary tract include, but are not limited to bladder, penile, kidney, renal pelvis, ureter, and urethral cancers.

Eye cancers include, but are not limited to intraocular melanoma and retinoblastoma.

Examples of liver cancers include, but are not limited to hepatocellular carcinoma (liver cell carcinomas with or without fibrolamellar variant), cholangiocarcinoma (intrahepatic bile duct carcinoma), and mixed hepatocellular cholangiocarcinoma.

Skin cancers include, but are not limited to squamous cell carcinoma, Kaposi's sarcoma, malignant melanoma, Merkel cell skin cancer, and non-melanoma skin cancer.

Head-and-neck cancers include, but are not limited to laryngeal / hypopharyngeal / nasopharyngeal / oropharyngeal cancer, and lip and oral cavity cancer.

Lymphomas include, but are not limited to AIDS-related lymphoma, non-Hodgkin's lymphoma, cutaneous T-cell lymphoma, Hodgkin's disease, and lymphoma of the central nervous system.

Sarcomas include, but are not limited to sarcoma of the soft tissue, osteosarcoma, malignant fibrous histiocytoma, lymphosarcoma, and rhabdomyosarcoma.

Leukemias include, but are not limited to acute myeloid leukemia, acute lymphoblastic leukemia, chronic lymphocytic leukemia, chronic myelogenous leukemia, and hairy cell leukemia.

These disorders have been well characterized in humans, but also exist with a similar etiology in other mammals, and can be treated by administering pharmaceutical compositions of the present invention.

The utility of the compounds of the present invention can be illustrated, for example, by their activity *in vitro* in the *in vitro* tumor cell proliferation assay described below. The link between activity in tumor cell proliferation assays *in vitro* and anti-tumor activity in the clinical setting has been well established in the art. For example, the therapeutic utility of taxol (Silvestrini et al. *Stem Cells* 1993, 11(6), 528-35), taxotere (Bissery et al. *Anti Cancer Drugs* 1995, 6(3), 339), and topoisomerase inhibitors (Edelman et al. *Cancer Chemother.*

Pharmacol. 1996, 37(5), 385-93) was demonstrated with the use of *in vitro* tumor proliferation assays.

In this application, where the plural form is used for compounds, salts, and the like, this is taken to mean also a single compound, salt, or the like.

Salts are especially the pharmaceutically acceptable salts of compounds of formula I such as, for example, acid addition salts, preferably with organic or inorganic acids, from compounds of formula I with a basic nitrogen atom. Suitable inorganic acids are, for example, halogen acids such as hydrochloric acid, sulfuric acid, or phosphoric acid. Suitable organic acids are, for example, carboxylic, phosphonic, sulfonic, or sulfamic acids, for example acetic acid, propionic acid, octanoic acid, decanoic acid, dodecanoic acid, glycolic acid, lactic acid, -hydroxybutyric acid, gluconic acid, glucosemonocarboxylic acid, fumaric acid, succinic acid, adipic acid, pimelic acid, suberic acid, azeiaic acid, malic acid, tartaric acid, citric acid, glucaric acid, galactaric acid, amino acids, such as glutamic acid, aspartic acid, *N*-methylglycine, acetytaminoacetic acid, *N*-acetylasparagine or *N*-acetylcysteine, pyruvic acid, acetoacetic acid, phosphoserine, 2- or 3-glycerophosphoric acid.

The compounds of the invention may be administered orally, dermally, parenterally, by injection, by inhalation or spray, or sublingually, rectally or vaginally in dosage unit formulations. The term 'administered by injection' includes intravenous, intraarticular, intramuscular, subcutaneous and parenteral injections, as well as use of infusion techniques. Dermal administration may include topical application or transdermal administration. One or more compounds may be present in association with one or more non-toxic pharmaceutically acceptable carriers and if desired, other active ingredients.

Compositions intended for oral use may be prepared according to any suitable method known to the art for the manufacture of pharmaceutical compositions. Such compositions may contain one or more agents selected from the group consisting of diluents, sweetening agents, flavoring agents, coloring agents and preserving agents in order to provide palatable preparations.

Tablets contain the active ingredient in admixture with non-toxic pharmaceutically acceptable excipients which are suitable for the manufacture of tablets. These excipients may be, for example, inert diluents, such as calcium carbonate, sodium carbonate, lactose, calcium phosphate or sodium phosphate; granulating and disintegrating agents, for example, corn starch, or alginic acid; and binding agents, for example magnesium stearate, stearic acid or talc. The tablets may be uncoated or they may be coated by known techniques to delay

disintegration and adsorption in the gastrointestinal tract and thereby provide a sustained action over a longer period. For example, a time delay material such as glyceryl monostearate or glyceryl distearate may be employed. These compounds may also be prepared in solid, rapidly released form.

Formulations for oral use may also be presented as hard gelatin capsules wherein the active ingredient is mixed with an inert solid diluent, for example, calcium carbonate, calcium phosphate or kaolin, or as soft gelatin capsules wherein the active ingredient is mixed with water or an oil medium, for example peanut oil, liquid paraffin or olive oil.

Aqueous suspensions containing the active materials in admixture with excipients suitable for the manufacture of aqueous suspensions may also be used. Such excipients are suspending agents, for example sodium carboxymethylcellulose, methylcellulose, hydroxypropyl-methylcellulose, sodium alginate, polyvinylpyrrolidone, gum tragacanth and gum acacia; dispersing or wetting agents may be a naturally-occurring phosphatide, for example, lecithin, or condensation products of an alkylene oxide with fatty acids, for example polyoxyethylene stearate, or condensation products of ethylene oxide with long chain aliphatic alcohols, for example heptadecaethyleneoxycetanol, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol such as polyoxyethylene sorbitol monooleate, or condensation products of ethylene oxide with partial esters derived from fatty acids and hexitol anhydrides, for example polyethylene sorbitan monooleate. The aqueous suspensions may also contain one or more preservatives, for example ethyl, or *n*-propyl, *p*-hydroxybenzoate, one or more coloring agents, one or more flavoring agents, and one or more sweetening agents, such as sucrose or saccharin.

Dispersible powders and granules suitable for preparation of an aqueous suspension by the addition of water provide the active ingredient in admixture with a dispersing or wetting agent, suspending agent and one or more preservatives. Suitable dispersing or wetting agents and suspending agents are exemplified by those already mentioned above. Additional excipients, for example, sweetening, flavoring and coloring agents, may also be present.

The compounds may also be in the form of non-aqueous liquid formulations, e.g., oily suspensions which may be formulated by suspending the active ingredients in a vegetable oil, for example arachis oil, olive oil, sesame oil or peanut oil, or in a mineral oil such as liquid paraffin. The oily suspensions may contain a thickening agent, for example beeswax, hard paraffin or cetyl alcohol. Sweetening agents such as those set forth above,

and flavoring agents may be added to provide palatable oral preparations. These compositions may be preserved by the addition of an anti-oxidant such as ascorbic acid.

Pharmaceutical compositions of the invention may also be in the form of oil-in-water emulsions. The oil phase may be a vegetable oil, for example olive oil or arachis oil, or a mineral oil, for example liquid paraffin or mixtures of these. Suitable emulsifying agents may be naturally-occurring gums, for example gum acacia or gum tragacanth, naturally-occurring phosphatides, for example soy bean, lecithin, and esters or partial esters derived from fatty acids and hexitol anhydrides, for example sorbitan monooleate, and condensation products of the said partial esters with ethylene oxide, for example polyoxyethylene sorbitan monooleate. The emulsions may also contain sweetening and flavoring agents.

Syrups and elixirs may be formulated with sweetening agents, for example glycerol, propylene glycol, sorbitol or sucrose. Such formulations may also contain a demulcent, a preservative and flavoring and coloring agents.

The compounds may also be administered in the form of suppositories for rectal or vaginal administration of the drug. These compositions can be prepared by mixing the drug with a suitable non-irritating excipient which is solid at ordinary temperatures but liquid at the rectal or vaginal temperature and will therefore melt in the rectum or vagina to release the drug. Such materials include cocoa butter and polyethylene glycols.

Compounds of the invention may also be administered transdermally using methods known to those skilled in the art (see, for example: Chien; "Transdermal Controlled Systemic Medications"; Marcel Dekker, Inc.; 1987. Lipp et al. WO 94/04157 3Mar94). For example, a solution or suspension of a compound of Formula I in a suitable volatile solvent optionally containing penetration enhancing agents can be combined with additional additives known to those skilled in the art, such as matrix materials and bacteriocides. After sterilization, the resulting mixture can be formulated following known procedures into dosage forms. In addition, on treatment with emulsifying agents and water, a solution or suspension of a compound of Formula I may be formulated into a lotion or salve.

Suitable solvents for processing transdermal delivery systems are known to those skilled in the art, and include lower alcohols such as ethanol or isopropyl alcohol, lower ketones such as acetone, lower carboxylic acid esters such as ethyl acetate, polar ethers such as tetrahydrofuran, lower hydrocarbons such as hexane, cyclohexane or benzene, or halogenated hydrocarbons such as dichloromethane, chloroform, trichlorotrifluoroethane, or trichlorofluoroethane. Suitable solvents may also include mixtures one or more materials

selected from lower alcohols, lower ketones , lower carboxylic acid esters, polar ethers, lower hydrocarbons, halogenated hydrocarbons.

Suitable penetration enhancing materials for transdermal delivery systems are known to those skilled in the art, and include, for example, monohydroxy or polyhydroxy alcohols such as ethanol, propylene glycol or benzyl alcohol, saturated or unsaturated C_8-C_{18} fatty alcohols such as lauryl alcohol or cetyl alcohol, saturated or unsaturated C_8-C_{18} fatty acids such as stearic acid, saturated or unsaturated fatty esters with up to 24 carbons such as methyl, ethyl, propyl, isopropyl, *n*-butyl, *sec*-butyl isobutyl *tert*-butyl or monoglycerin esters of acetic acid, capronic acid, lauric acid, myristinic acid, stearic acid, or palmitic acid, or diesters of saturated or unsaturated dicarboxylic acids with a total of up to 24 carbons such as diisopropyl adipate, diisobutyl adipate, diisopropyl sebacate, diisopropyl maleate, or diisopropyl fumarate. Additional penetration enhancing materials include phosphatidyl derivatives such as lecithin or cephalin, terpenes, amides, ketones, ureas and their derivatives, and ethers such as dimethyl isosorbid and diethyleneglycol monoethyl ether. Suitable penetration enhancing formulations may also include mixtures one or more materials selected from monohydroxy or polyhydroxy alcohols, saturated or unsaturated C_8-C_{18} fatty alcohols, saturated or unsaturated C_8-C_{18} fatty acids, saturated or unsaturated fatty esters with up to 24 carbons, diesters of saturated or unsaturated dicarboxylic acids with a total of up to 24 carbons, phosphatidyl derivatives, terpenes, amides, ketones, ureas and their derivatives, and ethers.

Suitable binding materials for transdermal delivery systems are known to those skilled in the art and include polyacrylates, silicones, polyurethanes, block polymers, styrene-butadiene coploymers, and natural and synthetic rubbers. Cellulose ethers, derivatized polyethylenes, and silicates may also be used as matrix components. Additional additives, such as viscous resins or oils may be added to increase the viscosity of the matrix.

For all regimens of use disclosed herein for compounds of Formula I, the daily oral dosage regimen will preferably be from 0.01 to 200 mg/Kg of total body weight. The daily dosage for administration by injection, including intravenous, intramuscular, subcutaneous and parenteral injections, and use of infusion techniques will preferably be from 0.01 to 200 mg/Kg of total body weight. The daily rectal dosage regimen will preferably be from 0.01 to 200 mg/Kg of total body weight. The daily vaginal dosage regimen will preferably be from 0.01 to 200 mg/Kg of total body weight. The daily topical dosage regimen will preferably be from 0.1 to 200 mg administered between one to four times daily. The transdermal concentration will preferably be that required to maintain a daily dose of from 0.01 to 200


mg/Kg. The daily inhalation dosage regimen will preferably be from 0.01 to 10 mg/Kg of total body weight.

It will be appreciated by those skilled in the art that the particular method of administration will depend on a variety of factors, all of which are considered routinely when administering therapeutics. It will also be understood, however, that the specific dose level for any given patient will depend upon a variety of factors, including, but not limited to the activity of the specific compound employed, the age of the patient, the body weight of the patient, the general health of the patient, the gender of the patient, the diet of the patient, time of administration, route of administration, rate of excretion, drug combinations, and the severity of the condition undergoing therapy. It will be further appreciated by one skilled in the art that the optimal course of treatment, i.e., the mode of treatment and the daily number of doses of a compound of Formula I or a pharmaceutically acceptable salt thereof given for a defined number of days, can be ascertained by those skilled in the art using conventional treatment tests.

GENERAL PREPARATIVE METHODS

The compounds of the invention have the general chemical structure shown below and may be prepared by the use of known chemical reactions and procedures. The particular process to be utilized in the preparation of the compounds of this invention depends upon the specific compound desired. Such factors as the selection of the specific J and Y moieties, as well as the specific substituents possible at various locations on the molecule, all play a role in the path to be followed in the preparation of the specific compounds of this invention. Those factors are readily recognized by one skilled in the art.


Nevertheless, the following general preparative methods are presented to aid the reader in synthesizing the compounds of the invention, with more detailed particular examples being presented below in the experimental section describing the working examples.




(I)

All variable groups of these methods are as described in the generic description if they are not specifically defined below. When a variable group or substituent with a given symbol (i.e. G, G', M) is used more than once in a given structure, it is to be understood that each of these groups or substituents may be independently varied within the range of definitions for that symbol.

Within these general methods the variable Z is equivalent to the moiety

Z =  in which each variable group or substituent is allowed to vary independently within the limits defined for that symbol.

Within these general methods the variable E is equivalent to the moiety

E =  in which each variable group or substituent is allowed to vary independently within the limits defined for that symbol.


It is recognized that compounds of the invention with each claimed optional functional group cannot be prepared with each of the below-listed methods. Within the scope of each method optional substituents are used which are stable to the reaction conditions, or the functional groups which may participate in the reactions are present in protected form where necessary, and the removal of such protective groups is completed at appropriate stages by methods well known to those skilled in the art.

Additional compounds of formula (I) may be prepared from other formula (I) compounds by elaboration of functional groups present. Such elaboration includes, but is

not limited to, hydrolysis, reduction, oxidation, alkylation, acylation, esterification, amidation and dehydration reactions. Such transformations may in some instances require the use of protecting groups by the methods disclosed in T. W. Greene and P.G.M. Wuts, *Protective Groups in Organic Synthesis*; Wiley: New York, (1999), and incorporated herein by reference. Such methods would be initiated after synthesis of the desired compound or at another place in the synthetic route that would be readily apparent to one skilled in the art.


General Method A - Invention compounds of formula **5** in which Z and E are as defined above, may be conveniently prepared according to a reaction sequence as shown in General Method "A". Thus, amidine or guanidine **1** and β-ketoester **2** are either obtained from commercial sources or made by one skilled in the art according to published procedures (amidine **1**: Granik et al *Russ Chem. Rev.* **1983**, *52*, 377-393; β-ketoester **2**: Tabuchi, H. et al. *Synlett* **1993**, *(9)*, 651-2). Amidine or guanidine **1** is treated with β-ketoester **2** in a refluxing mixed solvent such as alcohol and toluene or benzene to furnish pyrimidinone intermediate **3**. The alcohol is typically a lower molecular weight alcohol such as ethanol, isopropanol, *n*-propanol, *n*-butanol, *iso*-butanol, or *t*-butanol. Compound **3** is treated with a chlorinating agent such as phosphorous oxychloride, thionyl chloride or phosphorous pentachloride to yield chloropyrimidine intermediate **4**. Intermediate **4** is reacted with a nucleophile of formula NHR^1Z in a refluxing solvent such as alcohol, water, DMF, DMA, acetonitrile, acetone, dioxane or DMSO to furnish the invention compound of formula **5** [formula (I), where R^{2a} is H]. Such reactions can also be done in a melt free of solvent or in a solvent catalyzed by acids such as HCl, H_2SO_4 or bases such as but not limited to triethylamine, Cs_2CO_3, K_2CO_3, Na_2CO_3, K_3PO_4, Na_3PO_4, NaOH, KOH, NaH, $NaNH_2$, KNH_2, or a sodium or potassium alkoxide or 1,8-diazobicyclo[5.4.0]undec-7-ene (DBU). Invention compounds of formula **5a** [(I) where R^{2a} is Cl, Br or I] can be prepared from compounds of formula **5** by halogenation with Cl_2, Br_2, or I_2. Invention compounds of formula **5a** [(I) where R^{2a} is F] can be prepared from the formula (I) compounds where R^{2a} is Cl, Br or I by a nucleophilic substitution reaction using a fluoride source, e.g., KF.

General Method A



General Method B - Compounds of formula **5** in which R^1, R^2, Z and E are as previously defined can also be prepared via an alternative reaction sequence outlined in General Method "B" below. Thus, dichloropyrimidine **8**, which is either commercially available or can be made by one skilled in the art according to published procedures (Bagli, J. et al, *J. Med. Chem.* **1988**, *31(4)*, 814-23), is reacted with a nucleophile of formula NHR^1Z in a solvent such as alcohol, water, DMF or DMSO to furnish intermediate **9**. Such condensations can also be done in a solvent catalyzed by acids such as HCl, H_2SO_4 or an aforementioned base. Compound **9** is reacted with a boronic acid or ester of formula $E\text{-}B(OR')_2$ where R' is H, alkyl or two R' may form a ring, under standard Suzuki coupling conditions (such as $Pd(PPh_3)_4$ or $PdCl_2(dppf)\cdot CH_2Cl_2$ /base/solvent) to provide invention compound **5**.

General Method B




R' = H, alkyl or two R' may form a ring

General Method C - Invention compounds of formula **5** in which R^1, R^2 and E are as defined above, and Z, L' are defined below can also be prepared via an alternative

reaction sequence outlined in General Method "C" below. Thus, intermediate **4** is reacted with a nucleophile of formula **6** using aforementioned conditions (General Method A) to furnish intermediate **10**. Compound **10** is treated with an aromatic intermediate of formula **7** in an aprotic solvent and base (such as bases in General **Method A**) to furnish invention compounds of formula **5**.


General Method C



X = halogen, OTf, OMs, OTs
L' = O, NR^5

General Method D - Invention compounds of formula **13** in which R^1, R^2 and Z are as defined above, and R^D is G2, G12, G23, G24, G30, or benzyl can also be prepared via a reaction sequence as shown in General Method "D" below. Thus, demethylation of intermediate **11** (General method A or B or C) employing standard conditions (such as BBr_3, Me_3SiI, $AlCl_3$/EtSH etc.) provides intermediate **12**. Subsequently, compound **12** can then undergo alkylation, acylation, or sulfamylation to introduce the R^D substituent and provide the compound of formula **13**. Standard reaction conditions for these transformations can be used, i.e., a reagent of formula R^D-halo in the presence a base. In addition, O-alkylation can be accomplished using a Mitsunobu reaction (i.e., DEAD/PPh_3) to provide invention compound **13** where R^D is alkyl.


General Method D



11 **12** **13**

General Method E - Invention compounds of formula **16** and **17** in which R^1, R^2, G, G", m, n, and E are as defined above, and M' is CH or N, can be prepared via a reaction sequence as shown in General Method "E" below. Thus, the cyano group of intermediate **14** can be hydrolyzed and the resulting carboxylic acid can be coupled with an amine such as $NHR^{28}R^{29}$, a piperdine, or morpholine, under standard conditions to provide compound **16** where G^{E-1} is G21, G25 or G26. Invention compound **17** can be prepared by reduction of the amide **16** with $LiAlH_4$ or BH_3, followed by optional sulfonylation or acylation. Alternatively, compound **17** can be prepared by alkylation or reductive amination of amine **15**, which is prepared by reduction of **14** by a reducing agent such as H_2/Pd on C in acetic acid.


General Method E



14 15 16 17

General Method F - Invention compounds of formula **17b** can be prepared by displacement of the halo substituent on the compound of formula **17a** with a sulfur, nitrogen or oxygen nucleophile, represented by G^{F-1}-H, e.g., a thiol, ammonia, a mono or dialkylamine, water or an optionally substituted alcohol, in the optional presence of a base such as triethylamine, Cs_2CO_3, K_2CO_3, Na_2CO_3, K_3PO_4, Na_3PO_4, NaOH, KOH, NaH, $NaNH_2$, KNH_2, or a sodium or potassium alkoxide or 1,8-diazobicyclo[5.4.0]undec-7-ene (DBU). Thus are prepared compounds of formula (I) in which G^{F-1} is selected from G2, G3, G8, G16, G17, G22, G23, and G24. In addition, compounds of formula **17c** may be prepared by acylation or sulfonylation of the compounds of formula **17b** where at least one H may be replaced, using appropriate reagents such as acyl halides or alkylsulfonyl halides, generally in the presence of a base. Thus are prepared compounds of formula (I) in which G^{F-2} is selected from G12, G29, G30 and G31.

General Method F




General Methods (a-e) for Preparation of Intermediate NHR^1Z

Method a - The compounds of formula **18** in which M, G, G", m and n are as defined above, M' is independently CH or N, and L' is O or NR^5 can be conveniently prepared as shown in Method **a** below. Generally, the intermediate **18** may be prepared by an aromatic substitution reaction of intermediate **7** and intermediate **6**. Thus, aniline or aminopyridine **6** is treated with an aromatic intermediate of formula **7** in an aprotic solvent such as DMF, DMA, acetonitrile, acetone, dioxane or DMSO and base to furnish the intermediate of formula **18** (when X = OTf, OMs, OTs see ref. Sammes, P. et al. *J. Chem. Soc. Perkin Trans 1*, **1988**, *(12)*, 3229-31). Compounds of formula **18a** can be obtained


through reductive amination of **18** with an aldehyde under reductive amination conditions such as $NaBH_4$, $NaBH_3CN$, or $NaBH(OAc)_3$.

Method a




Method b - Alternatively, compounds of formula **18b,** in which M, G, G", m and n are as defined above, M' is independently CH or N, and L' is O, NR^5 or CH_2, can be conveniently prepared as shown in Method **b** below. Thus, the aromatic intermediate of formula **20** is deprotonated with an aforementioned base or LDA, *n*-BuLi, *t*-BuLi in an aprotic solvent, followed by reaction with intermediate **19** to furnish the intermediate of formula **21**. The nitro group of compound **21** can be reduced by one skilled in the art according to published procedures such as catalytic hydrogenation, Fe/HOAc and $SnCl_2$ to provide intermediate **18b**.

Method b




Method c The 4-substituted aniline compound of formula **25**, **26** and **27** in which G, G", m and n are as defined above, P' is a protecting group, M' is independently CH or N, and R^6 is H or (C_1-C_3)alkyl can be prepared via a reaction sequence as outlined in Method **c** below. Thus, intermediate **22** is treated with acyl chloride **23** under Friedel-Crafts acylation conditions (Lewis acid such as $AlCl_3$) to furnish the intermediate of formula **24**. Compound **24** can be converted to aniline **25** by Grignard reaction with R^6MgBr or reduction with $LiAlH_4$ followed by deprotection. Aniline **26** can be obtained by reduction of the carbonyl group of **24** by methods such as but not limited to N_2H_4/OH^-, $Pd/C/H_2$, Et_3SiH/Lewis acid, or $NaBH_4$/Lewis acid (see ref. Ono, A. et al, *Synthesis*, **1987**, (8), 736-8) or alternatively by formation of a dithiane and subsequent desulfuration with Raney Nickel. In some instances, deprotection of aniline may be necessary to obtain **26**. By deprotection of the amino group of compound **24**, the aniline intermediate **27** can also be obtained.

Method c




Method d The 3-substituted aniline compounds **30**, **30a** and **31** in which G, G", m and n are as defined above, M' is independently CH or N, and R^6 is H or (C_1-C_3)alkyl can be prepared conveniently via a reaction sequence as shown in Method **d** below. Thus, nitration of intermediate **28** employing standard nitration conditions such as but not limited to HNO_3/H_2SO_4, or $NaNO_3/HCl$ furnishes intermediate **29**. Reduction of **29** with a reducing agent such as $SnCl_2$, Fe/HOAc, or catalytic hydrogenation provides aniline **30**. Additionally, compound **29** can be converted to aniline **30a** by treatment with R^6MgBr or reduction with $LiAlH_4$ followed by the above-mentioned reduction conditions. Aniline **31** can be obtained by reduction of the carbonyl group by a method such as but not limited to $N_2H_4/NaOH$, Pd-C/H_2, Et_3SiH/Lewis acid, or $NaBH_4$/Lewis acid (see ref. Ono, A. et al, *Synthesis*, **1987**, (8), 736-8) or alternatively by formation of dithiane and subsequent desulfuration with Raney Nickel. In some instances, reduction of the nitro group by an aforementioned method may be necessary to obtain aniline **31**.

Method d



Method e The compounds of formula **36** and **37** in which M, G, G", m, n, R^{10} and R^{11} are as defined above and R^e is G2, G16, G23, and G24, can be prepared conveniently via a reaction sequence as shown in Method **e** below. Thus, intermediate pyridine **32** is oxidized by a reagent such as *m*-CPBA, H_2O_2, $CH_3C(O)OOH$, or $CF_3C(O)OOH$ to the *N*-oxide, followed by chlorination with a chlorinating agent such as phosphorous oxychloride, thionyl chloride or phosphorous pentachloride to yield chloropyridine **33**. Compound **33** can be converted to aniline **36** by treatment with alcohol in the presence of base such as NaH, followed by reduction of the nitro group with a reducing agent such as $SnCl_2$, Fe/H^+, or catalytic hydrogenation. Treatment of compound **33** with amine $HNR^{10}R^{11}$ followed by reduction of the nitro group of resulting compound **34** with the above mentioned reagents provides compound **37**.

Method e



By using these above described methods, the compounds of the invention may be prepared. The following specific examples are presented to further illustrate the invention described herein, but they should not be construed as limiting the scope of the invention in any way.

Abbreviations and Acronyms

A comprehensive list of the abbreviations utilized by organic chemists of ordinary skill in the art appears in the first issue of each volume of the *Journal of Organic Chemistry*; this list is typically presented in a table entitled Standard List of Abbreviations. The abbreviations contained in said list, and all abbreviations utilized by organic chemists of ordinary skill in the art are hereby incorporated by reference.

For purposes of this invention, the chemical elements are identified in accordance with the Periodic Table of the Elements, CAS version, Handbook of Chemistry and Physics, 67th Ed., 1986-87.

More specifically, when the following abbreviations are used throughout this disclosure, they have the following meaning:

2X	two times
3X	three times

$AlMe_3$	trimethylaluminum
Boc	*t*-butoxycarbonyl
n-BuLi	butyllithium
t-BuOK	potassium *t*-butoxide
calcd	calculated
Celite®	diatomaceous earth filtering agent, registered trademark of Celite Corp.
CD_3OD	methanol-d_4
$CHCl_3$-*d*	chloroform-*d*
d	doublet
DBU	1,8-diazobicyclo[5.4.0]undec-7-ene
DCC	dicyclohexylcarbodiimide
DEAD	diethylazodicarboxylate
DIBAH	diisobutylaluminum hydride
DIEA	diisopropylethylamine
DMA	dimethylacetamide
DMAP	4-dimethylaminopyridine
DME	dimethoxyethane
DMF	*N,N*-dimethylformamide
DMSO	dimethylsulfoxide
DMSO-d_6	dimethylsulfoxide-d_6
EDCI	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
EtSH	ethanethiol
EtOAc	ethyl acetate
EtOH	ethanol
Et_3SiH	triethylsilane
h	hour(s)
HATU	*O*-(7-azabenzotriazol-1-yl)-*N,N,N',N'*-tetramethyluronium hexafluorophosphate
Hex	hexanes
^1H NMR	proton nuclear magnetic resonance
HOAc	acetic acid
HPLC	high performance liquid chromatography
LC-MS	liquid chromatography / mass spectroscopy
LDA	lithium diisopropylamide

LiHMDS	lithium hexamethyldisilazide
m	multiplet
m-CPBA	3-chloroperoxybenzoic acid
MeOH	methanol
min	minute(s)
Me_3SiI	trimethylsilyl iodide
MS ES	mass spectroscopy with electrospray
$NaBH(OAc)_3$	sodium triacetoxyborohydride
OMs	O-methanesulfonyl (mesylate)
OTs	O-*p*-toluenesulfononyl (tosyl)
OTf	O-trifluoroacetyl (triflyl)
Pd/C	palladium on carbon
$Pd_2(dba)_3$	tris(dibenzylideneacetone)dipalladium(0)
$Pd(PPh_3)_4$	tetrakis(triphenylphosphine)palladium(0)
$PdCl_2(dppf)\cdot CH_2Cl_2$	[1,1'-bis(diphenylphosphino)ferrocene]dichloropalladium(II) complex with dichloromethane
RT	retention time
rt	room temperature
R_f	TLC Retention factor
s	singlet
t	triplet
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC	thin layer chromatography

General Analytical Procedures

The structure of representative compounds of this invention were confirmed using the following procedures.

Electron impact mass spectra (EI-MS) were obtained with a Hewlett Packard 5989A mass spectrometer equipped with a Hewlett Packard 5890 Gas Chromatograph with a J & W DB-5 column (0.25 uM coating; 30 m x 0.25 mm). The ion source was maintained at 250 °C and spectra were scanned from 50-800 amu at 2 sec per scan.

High pressure liquid chromatography-electrospray mass spectra (LC-MS) were obtained using either a:

(A) Hewlett-Packard 1100 HPLC equipped with a quaternary pump, a variable wavelength detector set at 254 nm, a YMC pro C-18 column (2 x 23 mm, 120A), and a Finnigan LCQ ion trap mass spectrometer with electrospray ionization. Spectra were scanned from 120-1200 amu using a variable ion time according to the number of ions in the source. The eluents were A: 2% acetonitrile in water with 0.02% TFA and B: 2% water in acetonitrile with 0.018% TFA. Gradient elution from 10% B to 95% over 3.5 min at a flowrate of 1.0 mL/min is used with an initial hold of 0.5 min and a final hold at 95% B of 0.5 min. Total run time is 6.5 min.

or

(B) Gilson HPLC system equipped with two Gilson 306 pumps, a Gilson 215 Autosampler, a Gilson diode array detector, a YMC Pro C-18 column (2 x 23 mm, 120 A), and a Micromass LCZ single quadrupole mass spectrometer with z-spray electrospray ionization. Spectra were scanned from 120-800 amu over 1.5 seconds. ELSD (Evaporative Light Scattering Detector) data is also acquired as an analog channel. The eluents were either A: 2% acetonitrile in water with 0.02% TFA or B: 2% water in acetonitrile with 0.018% TFA. Gradient elution from 10% B to 90% over 3.5 min at a flowrate of 1.5 mL/min is used with an initial hold of 0.5 min and a final hold at 90% B of 0.5 min. Total run time is 4.8 min. An extra switching valve is used for column switching and regeneration.

Routine one-dimensional NMR spectroscopy is performed on 400 MHz Varian Mercury-plus spectrometers. The samples were dissolved in deuterated solvents obtained from Cambridge Isotope Labs, and transferred to 5 mm ID Wilmad NMR tubes. The spectra were acquired at 293 K. The chemical shifts were recorded on the ppm scale and were referenced to the appropriate solvent signals, such as 2.49 ppm for DMSO-d_6, 1.93 ppm for CD_3CN-d_3, 3.30 ppm for CD_3OD 5.32 ppm for CD_2Cl_2-d_2 and 7.26 ppm for $CHCl_3$-d for ^1H spectra.

General HPLC Purification Method


Preparative reversed-phase HPLC chromatography was accomplished using a Gilson 215 system, typically using a YMC Pro-C18 AS-342 (150 x 20 mm I.D.) column. Typically, the mobile phase used was a mixture of (A) H_2O containing 0.1% TFA, and (B) acetonitrile. A typical gradient was:

Time [min]	A: %	B: %	Flow [mL/min]
0.50	90.0	10.0	1.0
11.00	0.0	100.0	1.0
14.00	0.0	100.0	1.0
15.02	100.0	0.0	1.0

Experimental Examples

Preparation of chloropyrimidine amine intermediates


Intermediate 1A: Preparation of 4-chloro-6-phenylpyrimidin-2-amine



A suspension of guanidine carbonate (3.60 g, 20 mmol) in ethanol (120 mL) and toluene (20 mL) was refluxed under nitrogen for 1 h, during which time about 50 mL of solvent was removed by distillation. After the mixture was cooled to 45 °C, ethyl 3-oxo-3-phenylpropanoate (7.68 g, 40 mmol) was added and the solution was heated at reflux overnight. The desired product precipitated as a white solid during the reaction. Water (50 mL) was added to the reaction and the mixture was refluxed for an additional 30 min. After cooling to rt, the mixture was neutralized with 1N HCl and placed in the refrigerator for 6 h. The solid was filtered, washed with water followed by ether and dried at 60 °C under vacuum to give the product as white solid (6.45 g, 86%). **MS** ES: 188 $(M+H)^+$, calcd 188; RT = 0.91 min; **TLC** (CH_2Cl_2/ 2M NH_3 in MeOH 95/5) R_f = 0.10.


A mixture of the above product (6.0 g, 32 mmol) and $POCl_3$ (100 mL) was heated at reflux for 1 h. The majority of the $POCl_3$ was removed in vacuo and the residue was diluted with EtOAc and poured over an ice/saturated $NaHCO_3$ solution. The aqueous layer was extracted with EtOAc and the combined organic layers were washed with brine, dried (Na_2SO_4), and concentrated. The crude organic concentrate was re-crystallized from EtOAc/ether to give the product **1A** as an off-white powder (2.8 g, 43%). **MS** ES: 206 $(M+H)^+$, calcd 206; RT = 2.49 min; **TLC** (CH_2Cl_2/ 2M NH_3 in MeOH 95/5) R_f = 0.72. (Reference 1: H. L. Skulnick, S. D. Weed, E. E. Edison, H. E. Renis, W. Wierenga, and D. A. Stringfellow, *J. Med. Chem.* 1985, 28, 1854-1869).

Intermediate 1B: Preparation of 4-chloro-6- (2-furyl) pyrimidin-2-amine




The (2-furyl) pyrimidin-2-amine intermediate **1B** was prepared by an analogous method to that described for **1A,** starting from guanidine carbonate and ethyl 3-(2-furyl)-3-oxopropanoate. **MS** ES: 196 $(M+H)^+$, calcd 196, RT = 2.13 min.

Intermediate 1C: Preparation of 4-chloro-6- (3-furyl) pyrimidin-2-amine




The (3-furyl) pyrimidin-2-amine intermediate **1C** was prepared by an analogous method to that described for **1A,** starting from guanidine carbonate and ethyl 3-(3-furyl)-3-oxopropanoate. **MS** ES: 196 $(M+H)^+$, calcd 196, RT = 2.04 min.

Intermediate 1D: Preparation of 4-chloro-6-(2-thienyl) pyrimidin-2-amine



Step 1: Preparation of ethyl 3-oxo-3- (2-thienyl) propanoate




A solution of 2,2-dimethyl-1,3-dioxane-4,6-dione (12 g, 83.26 mmol) and thiophene-2-carboxylic acid (8.97 g, 70.0 mmol) and DMAP (17.10 g, 140 mmol) in methylene chloride (100 mL) was cooled in an ice bath and treated with a solution of DCC (15.88 g, 76.96 mmol) in methylene chloride (50 mL). The reaction was stirred at rt for 2 h. The resulting precipitate was filtered and the filtrate was concentrated and re-dissolved in EtOH (400 mL). To this solution was added *p*-toluenesulfonic acid (32 g) and the reaction mixture was refluxed for 1 h. The solvent was removed in vacuo to afford the crude organic concentrate which was dissolved in ethyl acetate (1000 mL) and washed with water (300

mL). The organic layer was washed with saturated aqueous sodium bicarbonate (200 mL), 1N hydrochloric acid (200 mL), saturated aqueous sodium chloride, dried (Na_2SO_4), and concentrated. The residue was purified using silica gel column chromatography (0-7% ethyl acetate in hexane) to furnish the desired product as a colorless oil (3.67 g, 27%). **MS** ES 199 $(M+H)^+$, calcd 199; RT = 2.12 min; **TLC** (25% ethyl acetate in hexane) R_f = 0.50.


Step 2: Preparation of title compound (2-thienyl) pyrimidin-2-amine **1D**

(2-Thienyl) pyrimidin-2-amine **1D** was prepared by an analogous method to that described for **1A**, starting form guanidine carbonate and ethyl 3-oxo-3-(2-thienyl) propanoate. **MS** ES: 212 (M+H), calcd 212, RT = 2.42 min; **TLC** (20% EtOAc-80% hexane): R_f = 0.29.

Intermediate 1E: Preparation of 4-chloro-6-(3-methoxyphenyl)pyrimidin-2-amine



Step 1: Preparation of ethyl 3-oxo-3-(3-methoxyphenyl) propanoate




This material is prepared by a method analogous to that described for preparation of ethyl 3-oxo-3-(2-thienyl)propanoate in preparation of **1D**, starting from 2,2-dimethyl-1,3-dioxane-4,6-dione and 3-methoxybenzoic acid.


Step 2: Preparation of the title compound

1E is prepared by a method analogous to that described for **1A**, starting from guanidine carbonate and ethyl 3-oxo-3-(3-methoxyphenyl) propanoate.

Intermediate 1F: Preparation of 4-chloro-6-(4-methoxyphenyl)pyrimidin-2-amine



Step 1: Preparation of ethyl 3-oxo-3-(4-methoxyphenyl) propanoate




This material is prepared by a method analogous to that described for preparation of ethyl 3-oxo-3-(2-thienyl)propanoate in preparation of **1D**, starting from 2,2-dimethyl-1,3-dioxane-4,6-dione and 4-methoxybenzoic acid.


Step 2: Preparation of title compound

1F is prepared by a method analogous to that described for **1A**, starting from guanidine carbonate and ethyl 3-oxo-3-(4-methoxyphenyl) propanoate.

Intermediate 1G: Preparation of 4-chloro-6-[4-(trifluoromethyl) phenyl] pyrimidin-2-amine



Step 1: Preparation of ethyl 3-oxo-3-[4-(trifluoromethyl) phenyl]propanoate




This material is prepared by a method analogous to that described for preparation of ethyl 3-oxo-3-(2-thienyl)propanoate in preparation of **1D**, starting from 2,2-dimethyl-1,3-dioxane-4,6-dione and 4-(trifluoromethyl)benzoic acid.


Step 2: Preparation of title compound

This material is prepared by a method analogous to that described for **1A**, starting from guanidine carbonate and 3-oxo-3-[4-(trifluoromethyl) phenyl]propanoate.

Intermediate 1H: Preparation of 4-chloro-6-(4-fluorophenyl)pyrimidin-2-amine



Step 1: Preparation of ethyl 3-(4-fluorophenyl)-3-oxopropanoate




This material is prepared by a method analogous to that described for preparation of ethyl 3-oxo-3-(2-thienyl)propanoate in preparation of **1D**, starting from 2,2-dimethyl-1,3-dioxane-4,6-dione and 4-fluorobenzoic acid.

Step 2: Preparation of title compound

1H is prepared by a method analogous to that described for **1A**, starting from guanidine carbonate and the product from Step 1, ethyl 3-(4-fluorophenyl)-3-oxopropanoate.

Preparation of substituted aniline intermediates

Intermediate 2A: Preparation of {4-[(2-ethylpyridin-4-yl) oxy] phenyl} amine




To a –78 °C solution of diisopropylamine (12.1 mL, 86.2 mmol) in THF (20 mL) was added a solution of *n*-BuLi in hexanes (1.60 M, 26.9 mL, 43.0 mmol) dropwise over 5 min. The mixture was stirred for 30 min, then a solution of 4-chloro-picoline (5.00 g, 39.2 mmol) in THF (20 mL) was added slowly over 30 min. The reaction mixture was warmed to -60 °C and stirred for 30 min. after which time a solution of methyl iodide (2.44 mL, 39.2 mmol) in 10 mL THF was added over a 20 min period. The reaction was stirred for 30 min at -60 °C and 1.5 h at –30 °C. The reaction was quenched by pouring the mixture into cold brine. The mixture was extracted with dichloromethane. The organic layers were dried (sodium sulfate) and concentrated. Vacuum distillation of the residue (10 mm Hg, 70-80 °C) furnished 5 g of a 4.5:1 mixture of the desired 2-ethyl-4-chloropyridine and the isopropyl analog.

A well stirred, degassed solution of *t*-BuOK (5.43 g, 44.5 mmol), 4-aminophenol (4.16 g, 38.2 mmol) and 2-ethyl-4-chloropyridine (4.5 g, 32 mmol, contains 20% isopropyl analog) in dimethylacetamide (100 mL) was heated at 100 °C for 30 h. The reaction mixture was cooled to rt and concentrated in vacuo. The residue was partitioned between dichloromethane (200 mL) and 0.1 N NaOH (200 mL). The organic phase was washed with 0.1 N NaOH, dried (Na_2SO_4), and concentrated in vacuo. The crude oil was purified by silica gel chromatography (20% EtOAc to 60% EtOAc in hexanes) to provide 3.22 g of the


desired ethyl compound **2A** and 465 mg of the isopropyl analog. **MS** ES: 215 $(M+H)^+$, calcd 215, RT = 0.19 min.

Intermediate 2B: Preparation of {4-[(2-methylpyridin-4-yl) oxy] phenyl} amine




{4-[(2-Methylpyridin-4-yl) oxy] phenyl} amine **(2B)** was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from 4-aminophenol and 4-chloro-2-methylpyridine, **MS** ES: 201 $(M+H)^+$, calcd 201, RT = 1.01 min.

Intermediate 2C: Preparation of 4-(3-aminophenoxy) pyridine-2-carboxamide




3-Aminophenol (18.12 g, 0.17 mmol) and potassium *t*-butoxide (12.07 g, 0.17 mmol) were suspended in *N,N*-dimethylformide (350 mL) and stirred at rt for 30 min. 2-Amido-4-chloropyridine (20 g, 0.13 mmol) was added and the mixture was stirred at 90 °C overnight. The reaction mixture was concentrated in vacuo. The residue was partitioned between ethyl acetate and water. The aqueous layer was extracted with ethyl acetate. The organic layers were combined, dried (Na_2SO_4) and evaporated to dryness. The crude tan solid was recrystallized from ethyl acetate to afford 10.5 g (27%) of the desired product **2C**. **MS** ES: 230 $(M+H)^+$, calcd 230, RT = 1.29 min.

Intermediate 2D: Preparation of 4-(3-aminophenoxyamino] phenoxy)-*N*-methyl-pyridine-2-carboxamide




Aniline **2D** was prepared by a procedure described in WO 00/42012 (Bayer Corporation, ω-Carboxyaryl Substitued Diphenyl Ureas as RAF kinase Inhibitors), starting from 3-aminophenol and 4-chloro-2-(*N*-methylamido)pyridine. **MS** ES: 244 $(M+H)^+$, calcd 244, RT = 1.51 min.

Intermediate 2E: Preparation of {4-[(3, 5-difluoropyridin-4-yl) oxy] phenyl} amine




{4-[(3, 5-Difluoropyridin-4-yl) oxy] phenyl} amine **(2E)** was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from 4-aminophenol and 3, 4, 5-trifluoropyridine, **MS** ES: 223 $(M+H)^+$, calcd 223, RT = 0.50 min.

Intermediate 2F: Preparation of 4-(4-aminophenoxy)-*N*-methylpyridine-2-carboxamide




4-(4-Aminophenoxy)-*N*-methylpyridine-2-carboxamide **(2F)** was prepared by a procedure described in WO 00/42012 (Bayer Corporation, ω-Carboxyaryl Substitued Diphenyl Ureas as RAF kinase Inhibitors), starting from 4-aminophenol and 4-chloro-2-(*N*-methylamido)pyridine **MS** ES: 244 $(M+H)^+$, calcd 244, RT = 1.16 min.

Intermediate 2G: Preparation of 4-(4-amino-3-fluorophenoxy) pyridine-2-carbonitrile




4-(4-Amino-3-fluorophenoxy) pyridine-2-carbonitrile **(2G)** was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from 4-amino-3-fluorophenol and 4-chloro-2-cyanopyridine, **MS** ES: 230 $(M+H)^+$, calcd 230, RT = 2.85 min.

Intermediate 2H: Preparation of 4-(4-amino-2-fluorophenoxy) pyridine-2-carbonitrile




4-(4-Amino-2-fluorophenoxy) pyridine-2-carbonitrile **(2H)** was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from 4-amino-2-fluorophenol and 4-chloro-2-cyanopyridine, **MS** ES: 230 $(M+H)^+$, calcd 230, RT = 2.18 min.

Intermediate 2I: Preparation of 4-(4-aminophenoxy) pyridine-2-carbonitrile




4-(4-Aminophenoxy) pyridine-2-carbonitrile **(2I)** was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from 4-aminophenol and 4-chloro-2-cyanopyridine, **MS** ES: 212 $(M+H)^+$, calcd 212, RT = 1.23 min.

Intermediate 2J: Preparation of {3-fluoro-4-[(2-methylpyridin-4-yl)oxy]phenyl}-amine



{3-Fluoro-4-[(2-methylpyridin-4-yl)oxy]phenyl}-amine **(2J)** was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from 4-amino-2-fluorophenol and 4-chloro-2-cyanopyridine, **MS** ES: 219 $(M+H)^+$, calcd 219, RT = 1.07 min.


Intermediate 2K: Preparation of [4-(4-methoxyphenoxy) phenyl] amine



1-Fluoro-4-nitrobenzene (7.76 g, 55.0 mmol) and potassium carbonate (12.0 g, 86.8 mmol) were suspended in anhydrous DMF (100 mL) and stirred at 125 °C for 2 h. 4-Methoxyphenol (6.21 g, 50.0 mmol) was added and the mixture was stirred vigorously at 125 °C for 4 h. After cooling to rt, the reaction mixture was poured into ice-water (1000 mL) and stirred vigorously for 30 min. The resulting yellow solid was collected by vacuum filtration and washed with water to give 11.7 g of the nitro intermediate which was dried in vacuo overnight. This nitro intermediate (8.00 g, 32.6 mmol) was suspended in ethanol (180 mL) and added to a flask charged with 10% Pd/C (0.35 g). The reaction mixture was


flushed with hydrogen gas three times and then stirred at rt under hydrogen atmosphere overnight. The catalyst was removed by filtration and the filtrate was concentrated. The resulting precipitate was collected by vacuum filtration to give a white solid product (6.76 g, 96%). **MS** ES 216 $(M+H)^+$, calcd 216, RT = 1.24 min; **TLC** (25% ethyl acetate-hexane) R_f = 0.18.

Intermediate 2L: Preparation of 4-[4-amino-3-(trifluoromethyl)phenoxy] pyridine-2-carbonitrile




This material is prepared by a method analogous to that described for preparation of **2C**, starting from 4-amino-3-(trifluoromethyl) phenol and 4-chloro-2-cyanopyridine.

Intermediate 2M: Preparation of {4-[(2-methylpyrimidin-4-yl)oxy]phenyl}amine




This material is prepared by a method analogous to that described for preparation of **2C**, starting from 4-amino-phenol and 4-chloro-2-methylpyrimidine.

Intermediate 2N: Preparation of 4-(4-aminophenoxy)-N-(2-{[*tert*-butyl(dimethyl)silyl]oxy}ethyl)pyridine-2-carboxamide




4-(4-Aminophenoxy)-*N*-(2-{[*tert*-butyl(dimethyl)silyl]oxy}ethyl)pyridine-2-carboxamide was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide (Intermediate **2C**), starting from 4-aminophenol and N-(2-{[*tert*-butyl(dimethyl)silyl]oxy}ethyl)chloropyridine-2-carboxamide **MS** ES: 388 $(M+H)^+$, calcd 388, RT = 3.60 min.

Intermediate 2O: Preparation of 4-(4-fluoro-benzyl)-phenylamine




Step 1. Preparation of (4-fluoro-phenyl)-(4-nitrophenyl)-methanone



To a solution of 4-nitrobenzoyl chloride (2.3 g, 13 mmol) in nitroethane (20 mL) was added aluminum chloride (3.5 g, 26 mmol) followed by fluorobenzene (1.2 mL, 13 mmol). The mixture was stirred at rt for 4 h, then quenched carefully with 6M HCl. The reaction mixture was washed with dilute aqueous NaOH and brine, dried over sodium sulfate, filtered, and concentrated in vacuo to provide the crude product as a light yellow solid. The solid was purified by recrystallization from hexanes to give (4-fluoro-phenyl)- (4-nitrophenyl)-methanone (2.0 g, 65%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.41-8.32 (m, 5H), 7.90 (m, 1H), 7.84 (m, 1H), 7.20 (m, 1H).

Step 2. Preparation of 1-fluoro-4(4-nitrobenzyl)benzene




To a solution of (4-fluoro-phenyl)-(4-nitrophenyl)-methanone (2.0 g, 8.2 mmol) in dichloromethane (16 mL) at 0 °C was added trifluromethanesulfonic acid (1.4 mL, 16 mmol) in dichloromethane (16 mL). A solution of triethylsilane (2 mL, 12 mmol) in dichloromethane (16 mL) was subsequently added dropwise, resulting in an exotherm. After 5 min, additional trifluoromethanesulfonic acid (1.4 mL, 16 mmol) was added, followed by triethylsilane (2.0 mL, 12 mmol). The reaction mixture was stirred at rt for 2h, then poured into cold saturated sodium bicarbonate and extracted several times with dichloromethane. The combined organic extracts were dried over sodium sulfate and concentrated in vacuo. Purification by column chromatography eluting with 0-10% ethyl acetate in hexanes, gave the desired product as a white solid (260 mg, 14%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.14 (m, 2H), 7.31 (m, 2H), 7.12 (m, 2H), 7.01 (m, 2H), 4.06 (s, 2H).

Step 3. Preparation of the title compound.


To a solution of the product prepared in Step 2 (260 mg, 1.1 mmol) in ethanol (4 mL) and water (1.2 mL) was added iron powder (188 mg, 3.40 mmol) and ammonium chloride (36 mg, 0.70 mmol). The reaction was stirred at 85 °C for 2 h, cooled to rt, and filtered through Celite®. The filtrate was concentrated then diluted in dichloromethane, washed with water, and dried over sodium sulfate. The combined organic layers were concentrated in vacuo to afford 4-(4-fluoro-benzyl)-phenylamine as light brown oil which crystallized

upon standing (150 mg, 67%). **^{1}H NMR** ($CHCl_3$-*d*) δ 7.11 (m, 2H), 6.95 (m, 4H), 6.62 (m, 2H), 3.85 (s, 2H), 3.59 (br s, 2H).

Intermediate 2P: Preparation of 4-(2-trifluoromethyl-pyridin-4-ylmethyl)-phenylamine




Step 1. Preparation of (4-nitro-phenyl)-(2-trifluoromethyl-pyridin-4-yl)-acetic acid ethyl ester



To a solution of ethyl (4-nitrophenyl)acetate (760 mg, 3.6 mmol) in DMF (10 mL) was added 60% sodium hydride (145 mg, 3.6 mmol). The deep purple reaction mixture was stirred at rt for 30 min, then 4-fluoro-2-trifluoromethyl-pyridine (500 mg, 3.0 mmol) was added. After heating at 70 °C for 2 h, the mixture was poured onto ice water and extracted with ethyl acetate. The organic layers were washed with water and brine, then dried over sodium sulfate and concentrated in vacuo. The residue was purified by column chromatography, eluting with 10-30% ethyl acetate in hexanes, to give (4-nitro-phenyl)-(2-trifluoromethyl-pyridin-4-yl)-acetic acid ethyl ester as a viscous yellow oil (440 mg, 41%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.70 (d, J = 5.1 Hz, 1H), 8.23 (m, 2H), 7.63 (m, 1H), 7.50 (m, 2H), 7.44 (dd, J = 5.0, 1.6 Hz, 1H), 5.15 (s, 1H), 4.27 (q, J = 7.0 Hz, 2H), 1.30 (t, J = 7.1 Hz, 3H).

Step 2. Preparation of 4-(4-nitrobenzyl)-2-(trifluoromethyl)pyridine




To a solution of the product prepared in Step 1 (440 mg, 1.24 mmol) in methanol (13 mL) containing a drop of water, was added powdered LiOH (36 mg, 1.5 mmol) and the mixture was stirred at rt overnight. The mixture was concentrated to remove the methanol, diluted in dichloromethane, and washed with water. The combined organic extracts were dried over sodium sulfate, concentrated in vacuo, and purified by column chromatography eluting with 10-25% ethyl acetate in hexanes to give 4-(4-nitrobenzyl)-2-(trifluoromethyl)pyridine as a light yellow solid (100 mg, 29%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.65 (d, J = 4.7 Hz, 1H), 8.21 (m, 2H), 7.49 (s, 1H), 7.35 (m, 2H), 7.28 (m, 1H), 4.18 (s, 2H).


Step 3. Preparation of the title compound

10% Degussa Pd on carbon (15 mg, 0.14mmol) was flushed with nitrogen then diluted in ethanol (2 mL). A solution of 4-(4-nitrobenzyl)-2-(trifluoromethyl)pyridine (100 mg, 0.35 mmol) in ethanol (2 mL) and pyridine (14 mg, 0.18 mmol) was subsequently added, and the mixture was flushed again with nitrogen prior to placing a hydrogen balloon on the flask. The mixture was stirred at rt overnight then filtered through Celite® and concentrated. The residue was dissolved in ethyl acetate and filtered through a silica gel plug, eluting with 50-100% ethyl acetate in hexanes, to give 4-(2-trifluoromethyl-pyridin-4-ylmethyl)-phenylamine as a clear colorless oil (76 mg, 85%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.57 (d, J = 5.0 Hz, 1H), 7.47 (s, 1H), 7.26 (m, 1H), 6.93 (m, 2H), 6.65 (m, 2H), 3.92 (s, 2H), 3.57 (br s, 2H).

Intermediate 2Q: Preparation of 4-(4-amino-benzyl)-pyridine-2-carbonitrile



Step 1. Preparation of 4-(4-nitro-benzyl)-pyridine-2-carbonitrile




To a solution of 4-(4-nitro-benzyl)-pyridine 1-oxide (1.0 g, 4.3 mmol) in dichloromethane (9 mL) was added trimethylsilyl cyanide (2.3 mL, 17 mmol). After 5 min, benzoyl chloride (1.0 mL, 8.7 mmol) was added dropwise and the mixture was stirred at rt for an additional 30 min. Water (10 mL) was carefully added, followed by solid potassium carbonate (2.1 g). After 30 min, the aqueous phase was extracted with dichloromethane and the combined organic layers were dried over sodium sulfate and concentrated in vacuo. Purification of the residue by column chromatography, eluting with 5-25% ethyl acetate in hexanes, gave an orange oil. This oil was subsequently triturated with toluene to afford 4-(4-nitro-benzyl)-pyridine-2-carbonitrile as a tan solid (353 mg, 34%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.63 (d, J = 4.8 Hz, 1H), 8.21 (m, 2H), 7.49 (m, 1H), 7.32 (m, 3H), 4.15 (s, 2H).

Step 2. Preparation of the title compound

10% Degussa Pd on carbon (40 mg, 0.38 mmol) was flushed with nitrogen then diluted in ethanol (5 mL). 4-(4-Nitro-benzyl)-pyridine-2-carbonitrile (250 mg, 1.05 mmol) in ethanol (5 mL) and pyridine (42 mg, 0.52 mmol) was subsequently added, and the mixture was flushed again with nitrogen prior to placing a hydrogen balloon on the flask. The mixture was stirred at rt overnight then filtered through Celite® and concentrated. The


residue was dissolved in ethyl acetate and filtered through a silica gel plug, eluting with 50-100% ethyl acetate in hexanes, to give 4-(4-amino-benzyl)-pyridine-2-carbonitrile (134 mg, 61%). **^{1}H NMR** ($CHCl_3$-*d*) δ 8.55 (d, J = 5.2 Hz, 1H), 7.46 (s, 1H), 7.30 (d, J = 4.8 Hz, 1H), 6.93 (d, J = 8.3 Hz, 2H), 6.66 (d, J = 8.2 Hz, 2H), 3.91 (s, 2H).

Intermediate 2R: Preparation of 4-(4-aminophenoxy)-2-chloropyridine




4-(4-Aminophenoxy)-2-chloropyridine was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide (**2C**), starting from 4-aminophenol and 2,4-dichloropyridine **MS** ES: 221 $(M+H)^+$, calcd 221, RT = 0.32 min.

Intermediate 2S: Preparation of 4-(2-chloro-pyridin-4-ylmethyl)-phenylamine



4-(2-Chloro-pyridin-4-ylmethyl)-phenylamine was prepared by a method analogous to that described for 4-(2-trifluoromethyl-pyridin-4-ylmethyl)-phenylamine (**Intermediate 2P**), starting from ethyl (4-nitrophenyl) acetate and 2-chloro-4-nitro-pyridine. **^{1}H NMR** ($CHCl_3$-*d*) δ 8.23 (dd, J = 5.1, 0.5 Hz, 1H), 7.11 (m, 1H), 7.01 (m, 1H), 6.95 (m, 2H), 6.65 (m, 2H), 3.83 (s, 2H).


Intermediate 2T: Preparation of 4-[(4-bromopyridin-2-yl)oxy]aniline



A solution of 4-aminophenol (1.86 g, 17.05 mmol) in anhydrous DMF was added to a suspension of potassium *t*-butoxide (2.10 g, 18.75 mmol) in DMF. The mixture was stirred at rt for 1 h. 4-Bromo-2-fluoropyridine (3.00 g, 17.05 mmol) was added into the reaction mixture and it was heated at 90 °C with stirring for 20 h. It was cooled down to rt and 100 ml of water was slowly added to quench the reaction. The reaction mixture was concentrated in vacuum to provide a residue which was extracted with EtOAc (3 X) and washed with water (3 X). The organic layer was dried ($MgSO_4$) and concentrated to give the crude product, which was purified by flash chromatography (Hexane:EtOAc=6:4) to provide 1.02 g (23%)


of the intermediate **2T** as a yellow solid. **MS** ES 265 $(M+H)^+$, calc. 265, RT = 2.52 min; **TLC** (Hexane/EtOAc=6/4) $R_f = 0.26$.

Intermediate 2U: Preparation of 4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}aniline




A cold (-5 °C), de-gassed solution of 4-aminophenol (41.6 g, 0.38 mol) in *N,N*-dimethylacetamide (250 mL) was treated with potassium *tert*-butoxide and stirred while warming to 20 °C. A solution containing 4-fluoro-2-trifluoromethylpyridine (60 g, 0.36 mol) in dimethylacetamide (150 mL) was slowly added and the mixture was stirred at 25 °C for 18 h. The reaction mixture was then concentrated in vacuo and the residue was added to vigorously stirred water (1 L). The precipitated solids were collected by suction filtration and washed with isopropanol/ether (1:1) followed by ether and hexane. The yellow tan solids were dried to afford 72.8 g (79%) of product. **^{1}H NMR** (DMSO-d_6) δ 5.20 (s, 2H, -NH_2), 6.62 (m, 2H), 6.86 (m, 2H), 7.04 (dd, 1H, *J*=5.6, 2.4 Hz), 7.24 (d, 1H, *J*=2.4 Hz), 8.54 (d, 1H, 5.7 Hz). **MS** ES 255 $(M+H)^+$, calcd 255, RT=1.66 min.

Intermediate 2V: Preparation of methyl 4-(4-aminophenoxy)pyridine-2-carboxylate



Step 1. Synthesis of methyl 4-chloropyridine-2-carboxylate HCl salt



Anhydrous DMF (10.0 mL) was slowly added to $SOCl_2$ (300 mL) at 40-48 °C. The solution was stirred at for 10 min., then picolinic acid (100 g, 812 mmol) was added over 30 min. The resulting solution was heated at 72 °C (vigorous SO_2 evolution) for 16 h to generate a yellow solid. The resulting mixture was cooled to rt, diluted with toluene (500 mL) and concentrated to 200 mL. The toluene addition/concentration process was repeated twice. The resulting nearly dry residue was filtered, and the solids were washed with toluene (50 mL) and dried under high vacuum for 4 h to afford 4-chloropyridine-2-carbonyl chloride HCl salt as an off-white solid (27.2 g, 16%). This material was set aside.


The red filtrate from above was added to MeOH (200 mL) at a rate which kept the internal temperature below 55 °C. The contents were stirred at rt for 45 min, cooled to 5 °C and treated with Et_2O (200 mL) dropwise. The resulting solids were filtered, washed with Et_2O (200 mL) and dried under reduced pressure at 35 °C to provide methyl 4-chloropyridine-2-carboxylate HCl salt as a white solid (110 g, 65%): mp 108-112 °C; **^{1}H-NMR** (DMSO- d_6) δ 3.88 (s, 3H); 7.82 (dd, J=5.5, 2.2 Hz, 1H); 8.08 (d, J=2.2 Hz, 1H); 8.68 (d, J=5.5 Hz, 1H); 10.68 (br s, 1H); **MS** ES 172 $(M+H)^+$ calcd 172.

Step 2. Preparation of the title compound

Methyl 4-(4-aminophenoxy)pyridine-2-carboxylate was prepared by a method analogous to that described for 4-(3-aminophenoxy) pyridine-2-carboxamide **(2C)**, starting from the product of step 1 and 4-aminophenol.

Preparation of Invention Compounds

Example 1: Preparation of N^4-{4-[(2-ethylpyridin-4-yl) oxy]phenyl}-6-phenyl-pyrimidine-2, 4-diamine




Chloropyrimidine **1A** (75 mg, 0.35 mmol) and aniline **2A** (72 mg, 0.35 mmol) were suspended in water (2 mL) containing concentrated hydrochloric acid (0.1 mL) and stirred at 100 °C for 17 h. After cooling to rt, the mixture was neutralized with 1 N aqueous sodium hydroxide and stirred for 20 min. The precipitate was collected by filtration and purified by silica gel column chromatography (0-5% methanol-methylene chloride) to afford 43 mg (32%) of the title compound as a yellow solid. **^{1}H NMR** (DMSO-d_6) δ 9.34 (s, 1H), 8.31 (d, 5.7 Hz, 1H), 7.91-7.93 (m, 2H), 7.86 (d, J = 8.8 Hz, 2H), 7.45-7.47 (m, 3H), 7.07-7.09 (m, 2H), 6.76 (d, J = 2.3 Hz, 1H), 6.68 (dd, J = 5.6 Hz, 1.3 Hz, 1H), 6.49 (s, 1H), 6.37 (b, 2H), 2.69 (q, J = 7.6 Hz, 2H), 1.19 (t, J = 7.4 Hz, 3H), **MS** ES 384 $(M+H)^+$, calcd 384, RT = 1.87 min; **TLC** (5/95 v/v methanol-methylene chloride) R_f= 0.41. The reaction mixture can also be purified by preparative HPLC using an elution gradient from 15% to 85 %

acetonitrile in water containing 0.1 % TFA over 15 min with Phenomenex Luna 5 μ C18 150 x 30 mm column to provide the title compound as its TFA salt.

By dissolving the title compound in an appropriate solvent such as MeOH or dioxane, addition of either 1 N HCl or 1 N methanesulfonic acid, and filtration, the corresponding HCl or methanesulfonate salt is isolated.


By using the method described for Example 1, and by substituting appropriate starting materials, **Examples 60-68 and 87** were similarly prepared.

Example 2: Preparation of N^4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6 phenylpyrimidine-2,4-diamine



Stating from chloropyrimidine **1A** and aniline **2B**, this compound was prepared by a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.31 (s, 1H), 8.25 (d, J = 5.7 Hz, 1H), 7.89-7.92 (m, 2H), 7.83-7.86 (m, 2H), 7.42-7.48 (m, 3H), 7.04 (d, J = 8.9 Hz, 2H), 6.73 (d, J = 2.4 Hz, 1H), 6.67-6.69 (m, 1H), 6.47 (s, 1H), 6.34 (s, 2H), 2.37 (s, 3H); **MS** ES: 370 $(M+H)^+$, calcd 370, RT = 1.41 min; **TLC** (5/95 methanol-methylene chloride) R_f = 0.33.

Example 3: Preparation of 4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy} pyridine-2-carboxamide




Starting from chloropyrimidine **1A** and aniline **2C**, this material was prepared using a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.43 (s, 1H), 8.50 (d, J = 5.6 Hz, 1H), 8.12 (s, 1H), 7.88-7.90 (m, 2H), 7.75 (t, J = 2.2 Hz, 1H), 7.70

(s, 1H), 7.60 (d, J = 8.4 Hz, 1H), 7.43-7.46 (m, 4H), 7.38 (t, J = 8 Hz, 1H), 7.19 (dd, J = 5.5 Hz, 1.5 Hz, 1H), 6.75 (dd, J = 7.5 Hz, 1.0 Hz, 1H), 6.48 (s, 1H), 6.38 (b, 2H); **MS** ES 399 $(M+H)^+$, calcd 399, RT = 2.64 min; **TLC** (5/95 methanol-methylene chloride) R_f=0.27.

By using the method described for **Example 3,** and by substituting appropriate starting materials, **Example 69-70** were similarly prepared.


Example 4: Preparation of 4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-*N*-methylpyridine-2-carboxamide



Starting from chloropyrimidine **1A** and aniline **2D**, this material was prepared using the method similar to **Example 1.** **^{1}H NMR** (DMSO-d_6) δ 9.43 (s, 1H), 8.76-8.79 (m, 1H), 8.50 (d, J = 5.8 Hz, 1H), 7.88-7.90 (m, 2H), 7.75 (t, J = 2.0 Hz, 1H), 7.59 (d, J = 7.9 Hz, 1H), 7.42-7.47 (m, 4H), 7.38 (t, J = 8.0 Hz, 1H), 7.17-7.19 (m, 1H), 6.75 (dd, J = 8.0 Hz, 1.0 Hz, 1H), 6.48 (s, 1H), 6.38 (s, 2H), 2.78 (d, J = 4.8 Hz, 3H); **MS** ES 413 $(M+H)^+$, calcd 413, RT = 2.13 min; **TLC** (5/95 methanol-methylene chloride) R_f = 0.31.

By using the method described for **Example 4**, and by substituting appropriate starting materials, **Examples 71-74** were similarly prepared.


Example 5: Preparation of N^4-{4-[(3,5-difluoropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine



Staring from chloropyrimidine **1A** and aniline **2E**, this compound was prepared by a method analogous to that described for **Example 1.** **^{1}H NMR** (DMSO-d_6) δ 9.24 (s, 1H),


8.63 (s, 2H), 7.88-7.90 (m, 2H), 7.74 (d, J = 9.0 Hz, 2H), 7.43-7.46 (m, 3H), 7.03 (d, J = 9.0 Hz, 2H), 6.44 (s, 1H), 6.31 (b, 2H); **MS** ES 392 $(M+H)^+$, calcd 392, RT = 2.27 min.

Example 6: Preparation of 4-(4-amino-3-fluorophenoxy)pyridine-2-carbonitrile hydrochloride




Starting from chloropyrimide **1A** and aniline **2G**, this material was prepared by a method analogous to that described for **Example 1**. After the reaction was complete, the solid was filtered and washed with MeOH to provide the title compound. **^{1}H NMR** (DMSO-d_6) δ 12.99 (s, broad, 1H, 10.51 (s, broad, 1H) 8.63 (d, J = 6.0 Hz, 1H), 8.07 (s, broad, 1H), 7.88 (m, 2H), 7.79 (d, J = 2.4 Hz, 1H), 7.66 (m, 3H), 7.45 (dd, J = 11.2 Hz, 2.4 Hz, 1H), 7.30 (d, J = 3.2 Hz, 1H), 7.16 (dd, J = 8.8 Hz, 1.6 Hz, 1H), 6.82 (s, broad, 1H); **MS** ES: 399 $(M+H)^+$, calcd 399, RT = 2.24 min.

Example 7: N^4-{3-fluoro-4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine



Starting from chloropyrimidine **1A** and aniline **2H**, this material was prepared by a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.54 (s, 1H), 8.24-8.30 (m, 2H), 7.19 (dd, J = 7.6 Hz, 2.4 Hz, 2H), 7.44-7.48 (m, 3H), 7.23 (t, J = 9.2 Hz, 1H), 6.70-6.76 (m, 2H), 6.49 (b, 2H), 2.41 (s, 3H); **MS** ES 388 $(M+H)^+$, calcd 388, RT = 1.70 min.


Example 8: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy}pyridine-2-carbonitrile



Starting from chloropyrimidine **1A** and aniline **2I**, this material was prepared by a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 12.73 (b, 1H), 10.85 (b, 2H), 8.56 (d, J = 6.0 Hz, 1H), 7.94-7.96 (m, 2H), 7.83-7.85 (m, 2H), 7.71 (d, J = 2.4 Hz, 1H), 7.63-7.67 (m, 3H), 7.29 (d, J = 8.8 Hz, 2H), 7.18-7.21 (m, 1H), 6.65 (s, 1H); **MS** ES 381 $(M+H)^+$, calcd 381, RT = 2.22 min.


By using the method described for **Example 8,** and by substituting appropriate starting materials, **Example 75** was similarly prepared.

Example 9: Preparation of 4-(3-{[2-amino-6-(3-furyl)pyrimidin-4-yl]amino}phenoxy)-*N*-methylpyridine-2-carboxamide




Starting from chloropyrimidine **1C** and aniline **2D**, this material was prepared by a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.37 (s, 1H), 8.77 (d, J = 5.0 Hz, 1H), 8.50 (d, J = 5.0 Hz, 1H), 8.11 (s, 1H), 7.71-7.74 (m, 2H), 7.56-7.59 (m, 1H), 7.43 (d, J = 4.0 Hz, 1H), 7.36 (t, J = 8.0 Hz, 1H), 7.16-7.18 (m, 1H), 6.79-6.80 (m, 1H), 6.72-6.73 (m, 1H), 6.29 (s, 2H), 6.22 (b, 1H), 2.78 (d, J = 5.0 Hz, 3H); **MS** ES 403 $(M+H)^+$, calcd 403, RT = 1.99 min; **TLC** (5/95 methanol-methylene chloride) R_f= 0.27.

Example 10: Preparation of 4-(4-{[2-amino-6-(3-furyl)pyrimidin-4-yl]amino}-phenoxy)-*N*-methylpyridine-2-carboxamide




Starting from chloropyrimidine **1C** and aniline **2F**, this material was prepared by the method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.31 (s, 1H), 8.76 (d, J = 5.0 Hz, 1H), 8.47 (d, J = 6.0 Hz, 1H), 8.12 (s, 1H), 7.85 (d, J = 7.2 Hz, 2H), 7.74 (s, 1H), 7.36 (d, J = 3.0 Hz, 1H), 7.10-7.14 (m, 3H), 6.81 (s, 1H), 6.18 (s, 2H), 6.23 (s, 1H), 2.78 (d, J = 5.0 Hz, 3H); **MS** ES 403 $(M+H)^+$, calcd 403, RT = 1.94 min; **TLC** (5/95 methanol-methylene chloride) R_f= 0.26.

Example 11: Preparation of N^4-[4-(4-nitrophenoxy)phenyl]-6-phenylpyrimidine-2,4 diamine




Starting from chloropyrimidine **1A** and [4-(4-nitrophenoxy) phenyl] amine, this material was prepared by a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.36 (s, 1H), 8.23 (d, J = 9.2 Hz, 2H), 7.87-7.93 (m, 4H), 7.43-7.48 (m, 3H), 7.08-7.12 (m, 4H), 6.49 (s, 1H), 6.37 (b, 2H); **MS** ES 400 $(M+H)^+$, calcd 400, RT = 3.01 min; **TLC** (5/95 methanol-methylene chloride) R_f = 0.67.

Example 12: Preparation of N^4-[4-(4-chlorophenoxy)phenyl]-6-phenylpyrimidine-2,4-diamine




Starting from chloropyrimidine **1A** and [4-(4-chlorophenoxy) phenyl] amine, this material was prepared by a method analogous to that described for **Example 1**. **1H NMR** (DMSO-d_6) δ 9.24 (s, 1H), 7.90 (dd, J = 7.8 Hz, 1.8 Hz, 2H), 7.79 (d, J = 8.8 Hz, 2H), 7.45 (m, 3H), 7.39 (d, J = 8.4 Hz, 2H), 6.99 (m, 4H), 6.46 (s, 1H), 6.31 (s, 2H); **MS** ES 389 $(M+H)^+$, calcd 389, RT = 2.78 min; **TLC** (CH_2Cl_2/ 2M NH_3 in MeOH 95/5) R_f = 0.33

Example 13: Preparation of N^4-[4-(4-methoxyphenoxy)phenyl]-6-phenylpyrimidine-2,4-diamine




Starting from chloropyrimidine **1A** and aniline **2K**, this material was prepared by a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.15 (s, 1H), 7.90 (dd, J = 9.6 Hz, 1.6 Hz, 2H), 7.70 (m, 2H), 7.44 (m, 3H), 6.93 (m, 4H), 6.88 (d, J = 8.8 Hz, 2H), 6.43 (s, 1H), 6.27 (s, 2H), 3.72 (s, 3H); **MS** ES 385 $(M+H)^+$, calcd 385, RT = 2.48 min.

Example 14: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]-2-fluorophenoxy}pyridine-2-carbonitrile




This material is prepared by a method analogous to that described in **Example 1**, starting from **2H** and **1A**.

Example 15: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]-3 (trifluoromethyl) phenoxy} pyridine-2-carbonitrile




This material is prepared by a method analogous to that described in **Example 1**, starting from **2L** and **1A**.

Example 16: Preparation of N^4-{4-[(2-methylpyrimidin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine




This material is prepared by a method analogous to that described in **Example 1**, starting from **2M** and **1A**.

Example 17: Preparation of N^4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-[4-(2-pyrrolidin-1-ylethoxy)phenyl]pyrimidine-2,4-diamine




Step 1: Preparation of 6-(4-methoxyphenyl)-N^4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine




This material is prepared by a method analogous to that described for **Example 1**, starting from **1F** and **2B**.

Step 2: Preparation of 4-[2-amino-6-({4-[(2-methylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenol



The intermediate from Step 1 above is treated with BBr_3 in methylene chloride at 0 °C for 12 h. After work-up and purification by a published procedure (J. F. W. McOmie and D. E. West, *Org. Synth., Collect.* Vol. V, 412 (1973)), the desired compound is obtained.

Step 3: Preparation of 6-[4-(2-bromoethoxy)phenyl]-N^4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine




To a solution of Step 2 product (1 equiv) in DMF is added 1,2- dibromoethane (1 equiv) and K_2CO_3 (3 equiv). The mixture is refluxed overnight. After cooling to rt, the mixture is diluted with EtOAc and washed sequentially with 1N NaOH, water and brine. The organic layer is dried (Na_2SO_4) and concentrated to afford a crude product which is to be used in next step without further purification.

Step 4: Preparation of the title compound

A mixture of the product from Step 3 (1 equiv), pyrrolidine (2 equiv) and K_2CO_3 (8 equiv) in DMF is stirred at 65 °C overnight. The solvent is removed and the residue is dissolved in EtOAc. The organic solution is washed with water, dried, and evaporated to dryness. The residue is purified by chromatography on a silica column to afford the title compound.


By using the method described for Example 17, and by substituting appropriate starting materials, **Examples 76-77** are similarly prepared.

Example 18: Preparation of 4-[4-({2-amino-6-[4-(2-piperidin-1-ylethoxy) phenyl] pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile




This is prepared by a method analogous to that described for **Example 17**, starting from **1F**, **2I** and using piperidine in step 4.

Example19: Preparation of methyl 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxylate



Starting from chloropyrimidine **1A** and aniline **2V**, this material was prepared using a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.37 (s, 1H), 8.52 (d, 1H), 7.85 (m, 4H), 7.41 (m, 4H), 7.16 (m, 3H), 6.45 (s, 1H), 6.36 (s, 2H), 3.79 (s, 3H); **MS** ES 414 $(M+H)^+$, calcd 414, RT = 2.16 min.


Example 20: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy} pyridine-2-carboxylic acid



A solution containing the 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy}pyridine-2-carbonitrile (20 g, 0.05 mol, **Example 8**) in concentrated sulfuric acid (150 mL) was heated at 70 °C for 12 h. The reaction mixture was then cooled to -40 °C and water (30 mL) was added, followed by heating at 70 °C for 12 h. The solution was cooled to rt and poured into vigorously stirred ice water (2 L) and stirring was continued for 2 h. The solids were then collected by suction filtration, washed with water (500 mL) and dried by air suction. The slightly damp material was then dissolved in a minimum volume of hot (90 °C) *N,N*-dimethylformamide and triethylamine was added until the mixture tested slightly acidic. The cooled solution was then poured into ice water (2 L), stirred for 0.5-1 h and the precipitated material was collected by suction filtration. The filter cake was washed with water, followed by isopropanol, diethyl ether, and finally hexane. Air-drying sequentially afforded the carboxylic acid as an off-white solid, 18.5 g (90%). **^{1}H NMR** (DMSO- d_6) δ 9.40 (s, 1H). 8.53 (d, 1H, J=5.8 Hz), 7.90 (m, 4H), 7.46 (m, 3H), 7.40 (d, 1H, *J*=7.1 Hz), 7.16 (m, 1H,), 7.13 (d, 2H, *J*=9.1 Hz), 6.50 (s, 1H), 6.40 (s, 2H), 3.30 (br s, 1H), **MS** ES 400 $(M+H)^+$, calcd 400, RT = 1.71 min.

The HCl salt of the title compound, **(Example 78)** .was prepared by addition of **Example 20** to a 1N HCl.


Example 21: Preparation of N^4-(4-{[2-(morpholin-4-ylcarbonyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine



To a solution of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy} pyridine-2-carboxylic acid (**Example 20**, 0.15 g, 0.38 mmol) in dry DMA (3 mL) was added HATU


(0.14 g, 0.38 mmol) and DIEA (0.15 g, 1.13 mmol). The solution was stirred at rt for 0.5 h, followed by addition of morpholine (0.16 g, 1.88 mmol). The resulting solution was stirred at rt overnight, followed by prep-HPLC separation to give 77 mg (35%) pure product. **^{1}H NMR** (DMSO- d_6) δ 10.79 (s, 1H), 8.40 (s, 1H), 7.85 (m, 2H), 7.74 (m, 2H), 7.60 (m, 3H), 7.21 (m, 2H), 7.00 (m, 2H), 6.59 (s, 1H), 3.49 (m, 8H); **MS** ES 469 $(M+H)^+$, calcd 469.

Example 22: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-*N,N*-dimethylpyridine-2-carboxamide



This material is prepared by a method analogous to that described for **Example 21**, starting from 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxylic acid and dimethylamine.


Example 23: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-*N*-(2-methoxyethyl)pyridine-2-carboxamide




This material was prepared by a method analogous to that described for **Example 21**, starting from 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy}pyridine-2-carboxylic acid and 2-methoxyethylamine.

By using the method described for **Example 23**, and by substituting appropriate starting materials, **Examples 79-82** were similarly prepared.

Example 24: Preparation of 4-[4-({2-amino-6-[4-(trifluoromethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]-*N*-(2-methoxyethyl)pyridine-2-carboxamide



Step 1: Preparation of 4-[4-({2-amino-6-[4-(trifluoromethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxylic acid




This material is prepared by methods analogous to that described for **Example 1** and **Example 20**, starting from **2I** and **1G.**

Step 2: Preparation of the title compound


This material is prepared by a method analogous to that described for **Example 21** starting from 2-methoxyethylamine and 4-[4-({2-amino-6 (trifluoromethyl)phenyl] pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxylic acid.

Example 25: Preparation of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-*N*-(2-methoxyethyl)-*N*-methylpyridine-2-carboxamide




This material is prepared by a method analogous to that described for **Example 21**, starting from 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxylic acid and 2-methoxyethyl-*N*-methyl amine.

Example 26: Preparation of N^4-[4-({2-[(4-methylpiperazin-1-yl)carbonyl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine



This material is prepared by a method analogous to that described for **Example 21**, starting from 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy} pyridine-2-carboxylic acid (**Example 20**) and 1-methylpiperizine. **^{1}H NMR** (DMSO-d_6) δ 10.85 (s, 1H), 10.19 (s, 1H), 8.42 (d, 1H), 7.90 (m, 2H), 7.74 (m, 3H), 7.59 (m, 4H), 7.22 (m, 3H), 7.06 (m, 2H), 6.60 (s,1H), 4.51 (m, 1H), 4.08 (m, 1H), 3.45 (m, 3H), 3.17 (m, 3H), 2.78 (s, 3H); **MS** ES 482 $(M+H)^+$, calcd 482, RT = 1.86 min.


Example 27: Preparation of N^4-{4-[(2-{[(2-methoxyethyl)amino]methyl}pyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine




A solution of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl) amino]phenoxy}-*N*-(2-methoxyethyl)pyridine-2-carboxamide from **Example 23** (50 mmol) in anhydrous THF (50 mL) is added in portions to a pre-cooled in ice-bath solution of lithium aluminum hydride (100 mmol, 1.0 M in THF) in anhydrous THF (150 mL). The reaction is stirred at 0 °C for 30 min until evolution of hydrogen subsides. The reaction mixture is refluxed under nitrogen for 48 h. The mixture is brought to 5-10 °C and carefully quenched with water (3.8 mL), 15% NaOH (3.8 mL) and water (12 mL). The mixture is extracted with EtOAc and the organic layer is dried and concentrated to give a crude product which is purified by chromatography on a silica column to give the title compound.

(Reference: *Org. Synth. Collect.*, 1988, Vol. VI, 382-385)

Example 28: Preparation of 6-(4-fluorophenyl)-N^4-(4-{[2-(piperidin-1-ylcarbonyl) pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine




Step 1: Preparation of 4-(4-{[2-amino-6-(4-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxylic acid



This material is prepared by a method analogous to that described for **Examples 1** and **20** starting from **1H** and **2I.**

Step 2: Preparation of 6-(4-methoxyphenyl)-N^4-(4-{[2-(morpholin-4-ylcarbonyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine




This material is prepared by a method analogous to that described for **Example 21,** starting from 6-(4-fluorophenyl)-N^4-[4-(pyridin-4-yloxy) phenyl] pyrimidine-2,4-diamine and piperidine.


Step 3: Preparation of the title compound

This material is prepared by a method analogous to that described for **Example 27,** starting from 6-(4-methoxyphenyl)-N^4-(4-{[2-(morpholin-4-ylcarbonyl)pyridin-4-yl]oxy} phenyl)pyrimidine-2,4-diamine.

Example 29: Preparation of 6-(4-methoxyphenyl)- N^4-(4-{[2-(morpholin-4-ylmethyl) pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine




Step 1: Preparation of 4-(4-{[2-amino-6-(4-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxylic acid



This material is prepared by a method analogous to that described for **Examples 1** and **20**, starting from **1F** and **2I**.

Step 2: Preparation of 6-(4-methoxyphenyl)-N^4-(4-{[2-(morpholin-4-ylcarbonyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine




This material is prepared by a method analogous to that described for **Example 21**, starting from the product from step 1.


Step 3: Preparation of title compound

This material is prepared by a method analogous to that described for **Example 27**, starting from the product of step 2.

Example 30: Preparation of N^2-ethyl-6-(3-methoxy-phenyl)- N^4-[4-(2-trifluoromethyl-pyridin-4-yloxy)-phenyl]-pyrimidine-2,4-diamine




Step 1: Preparation of 2,4-dichloro-6-(3-methoxy-phenyl)-pyrimidine



Trichloropyrimidine (11.83 g, 64.49 mmol) was added to a solution of 3-methoxyphenylboronic acid (9.8 g, 64.49 mmol) in a solvent mixture of ethanol (30 mL), toluene (30 mL) and 2M aqueous sodium bicarbonate (96.7 mL) at rt. The resulting mixture was degassed under vacuum for several min before the flask was purged with nitrogen. Dichloro[1,1'-bis(diphenylphosphino)ferrocene]palladium (II)dichloromethane adduct (2.4 g, 3.22 mmol) was added and the resulting mixture was heated for 3 h at 50 °C. The cooled reaction mixture was filtered through a silica gel pad and the pad was washed with acetone. The filtrated was evaporated under reduced pressure. The crude material was purified by column chromatography eluting with a gradient of 0 to 45% ethyl acetate/hexanes to give 2,4-dichloro-6-(3-methoxy-phenyl)-pyrimidine as a white solid (14.4 g, 65.4%). **MS** ES 255 $(M+H)^+$, calcd 255, RT = 3.35 min.

Step 2: Preparation of [2-chloro-6-(3-methoxy-phenyl)-pyrimidin-4-yl]-[4-(2-trifluoromethyl-pyridin-4-yloxy)-phenyl]-amine




2,4-Dichloro-6-(3-methoxy-phenyl)-pyrimidine (1.0 g, 3.92 mmol) and 4-(2-trifluoromethyl-pyridin-4-yloxy)-phenylamine **Intermediate 2U** (1.0 g, 3.92 mmol) were suspended in a mixture of isopropanol/water 2:8 (40 mL). The reaction mixture was heated at reflux for 24 h at which point the TLC showed a completed reaction. The reaction mixture was filtered with a fritted glass funnel. The crude residue was purified by HPLC eluting with

a gradient of 0 to 71% acetonitrile/water containing 0.1% TFA in both solvents. The TFA salt of [2-chloro-6-(3-methoxy-phenyl)-pyrimidin-4-yl]-[4-(2-trifluoromethyl-pyridin-4-yloxy)-phenyl]-amine was obtained as a yellow oil which solidified on standing. (926 mg, 50.1%). **MS** ES 473 $(M+H)^+$, calcd 473, RT = 3.98 min.

Step 3: Preparation of the title compound: N^2-ethyl-6-(3-methoxy-phenyl)- N^4-[4-(2-trifluoromethyl-pyridin-4-yloxy)-phenyl]-pyrimidine-2,4-diamine

[2-Chloro-6-(3-methoxy-phenyl)-pyrimidin-4-yl]-[4-(2-trifluoromethyl-pyridin-4-yloxy)-phenyl]-amine (100 mg, 0.21 mmol) and ethylamine (2M THF, 1 mL) were dissolved in *n*-butanol (3 mL) and the reaction mixture was heated at 120 °C overnight. The reaction mixture were evaporated under vacuum, and the crude residue was purified by HPLC eluting with a gradient of 10 to 85% acetonitrile/water containing 0.1% TFA in both solvents. The TFA salt of N^2-Ethyl-6-(3-methoxy-phenyl)- N^4-[4-(2-trifluoromethyl-pyridin-4-yloxy)-phenyl]-pyrimidine-2,4-diamine (13.9 mg, 11%) was obtained as a beige solid. **^{1}H NMR** (acetone-d_6) δ 10.36 (br, 1H), 10.06 (Br, 1H), 8.62 (d, J = 6 Hz, 1H), 8.00 (br, 1H), 7.53-7.51 (m, 1H), 7.47-7.31 (m, 2H), 7.19-7.12 (m, 3H), 6.66 (s, 1H), 3.94 (s, 3H), 3.59-3.56 (m, 2H), 1.31 (t, J = 7 Hz, 3H). **MS** ES 482 $(M+H)^+$, calcd 482, RT = 2.83 min.


Example 31: Preparation of 4-[4-(2-amino-5-bromo-6-phenyl-pyrimidin-4-ylamino)-phenoxy]-pyridine-2-carbonitrile



To a solution of 4-[4-(2-amino-6-phenyl-pyrimidin-4-ylamino)-phenoxy]-pyridine-2-carbonitrile (**Example 8,** 200 mg, 0.53 mmol) and sodium acetate (146.6 mg, 459.8 mmol) in acetic acid (4 mL) at rt was added bromine (84 mg, 0.53 mmol). The reaction was allowed to stand for 2 h after which time dichloromethane (20 mL) was added followed by water (20 mL). The phases were separated and the organic layer was washed with a saturated aqueous bicarbonate solution. The combined organic extracts were dried over $MgSO_4$ and then evaporated under vacuum. The crude material was purified by column chromatography eluting with a gradient of 0 to 60% AcOEt/Hexanes to give 4-[4-(2-Amino-5-bromo-6-phenyl-pyrimidin-4-ylamino)-phenoxy]-pyridine-2-carbonitrile as an orange solid (200 mg,

83%). **^{1}H NMR** (DMSO d_6) δ 8.58 (d, J = 5 Hz, 1H, 8.50 (Br, 1H), 7.90-7.86 (m, 2H), 7.67 (d, J = 2Hz, 1H), 7.54-7.51 (m, 2H), 7.47-7..41 (m, 3H), 7.20-7.16 (m, 3H), 6.57 (Br, 2H). **MS** ES 459 $(M+H)^+$, calcd 459, RT = 2.85 min.


Example 32: Preparation of N^4-{4-[2-(2-morpholin-4-yl-ethoxy)-pyridin-4-yloxy]-phenyl}-6-phenyl-pyrimidine-2,4-diamine



N^4-[4-(2-chloro-pyridin-4-yloxy)-phenyl]-6-phenyl-pyrimidine-2,4-diamine (**Example 48,** 75 mg, 0.19 mmol) was dissolved in toluene (1.5 mL). 2-Morpholin-4-yl-ethanol (61 mg, 0.46 mmol), powdered KOH (22 mg, 0.38 mmol), and 18-crown-6 (20 mg, 0.08 mmol) were subsequently added. The mixture was stirred at 90 °C overnight, after which time it was diluted with water and extracted with both ethyl acetate and dichloromethane. The combined organic extracts were concentrated and the residue was purified by prep HPLC to give the title compound (14 mg, 15%). **^{1}H NMR** (DMSO-d_6) δ 10.79 (br s, 1H), 9.99 (br s, 1H), 8.08 (d, J = 5.8 Hz, 1H), 7.90 (m, 2H), 7.77 (dd, J = 7.6, 2.0 Hz, 2H), 7.64 (m, 3H), 7.21 (d, J = 8.9 Hz, 2H), 6.69 (dd, J = 5.8, 2.1 Hz, 1H), 6.60 (s, 1H), 6.22 (d, J = 2.2 Hz, 1 H), 4.57 (t, J = 5.1 Hz, 2H), 3.96 (m, 2H), 3.68 (m, 2H), 3.50 (m, 4H), 3.15 (m, 2H); **MS** ES: 485 $(M+H)^+$, calcd 485, RT = 1.96 min.


By using the method described for **Example 32**, and by substituting appropriate starting materials, **Examples 83-86** were similarly prepared.

Example 33: Preparation of 6-phenyl-N^4-[4-(2-trifluoromethyl-pyridin-4-ylmethyl)-phenyl]-pyrimidine-2,4-diamine




Starting from chloropyrimidine **1A** and aniline **2P**, this material was prepared using a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 12.90 (br s, 1H), 10.82 (br s, 1H), 8.64 (d, J = 5.1 Hz, 1H), 7.83 (m, 3 H),. 7.74 (m, 2H), 7.64 (m, 4H), 7.31 (d, J = 8.3 Hz, 2H), 6.67 (br s, 1 H), 4.11 (s, 2H); **MS** ES: 422 $(M+H)^+$, calcd 422, RT = 2.51 min.

Example 34: Preparation of N^4-[4-(2-chloro-pyridin-4-ylmethyl)-phenyl)-6-phenyl-pyrimidine-2,4-diamine



Starting from chloropyrimidine **1A** and aniline **2S**, this material was prepared using a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 12.80 (br s, 1H), 10.74 (br s, 1H), 8.29 (dd, J = 5.4, 0.6 Hz, 1H), 7.83 (m, 2H), 7.67 (m, 2H), 7.63 (m, 3H), 7.41 (m, 1H), 7.30 (m, 3H), 6.61 (br s, 1H), 3.98 (s, 2H); **MS** ES: 388 $(M+H)^+$, calcd 388, RT = 2.38 min.


Example 35: Preparation of 4-[4-(2-amino-6-phenyl-pyrimidin-4-ylamino)-benzyl]-pyridine-2-carbonitrile



Starting from chloropyrimidine **1A** and **Intermediate 2Q**, this material was prepared using a method analogous to that described for **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 12.74 (br s, 1H), 10.73 (br s, 1H), 8.63 (d, J = 4.9 Hz, 1H), 7.96 (s, 1H), 7.82 (m, 2H), 7.72 (m, 2H), 7.61 (m, 4H), 7.30 (d, J = 7.9 Hz, 2H), 6.61 (s, 1H), 4.05 (s, 2H); **MS** ES: 379 $(M+H)^+$, calcd 379, RT = 2.35 min.


By using the method described for **Example 35**, and by substituting appropriate starting materials, **Examples 88-91** were similarly prepared.

Example 36: Preparation of N^4-[4-(2-aminomethyl-pyridin-4-ylmethyl)-phenyl]-6-phenyl-pyrimidine-2,4-diamine




10% Degussa Pd on carbon (15 mg, 0.14 mmol) was flushed with nitrogen then diluted in methanol (1 mL). 4-[4-(2-amino-6-phenyl-pyrimidin-4-ylamino)-benzyl]-pyridine-2-carbonitrile (**Example 35,** 90 mg, 0.24 mmol) in methanol (2 mL) and concentrated HCl (0.03 mL) were subsequently added, and the mixture was flushed again with nitrogen prior to placing a hydrogen balloon on the flask. The mixture was stirred at rt for 3 h then filtered through Celite® and concentrated. The residue was purified by prep HPLC to give N^4-[4-(2-aminomethyl-pyridin-4-ylmethyl)-phenyl]-6-phenyl-pyrimidine-2,4-diamine (10 mg, 11%). **^{1}H NMR** (CD_3OD) δ 8.50 (d, J = 5.3 Hz, 1H), 7.76 (m, 4H), 7.63 (m, 3H), 7.28 (m, 4H), 6.53 (s, 1H), 4.22 (s, 2H), 4.05 (s, 2H); **MS** ES: 383 $(M+H)^+$, calcd 383, RT = 1.84 min.

Example 37: Preparation of 6-phenyl-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine




Starting from chloropyrimidine **1A** and 4-(2-trifluoromethyl-pyridin-4-yloxy)-phenylamine **2U**, the title compound was prepared using a method analogous to that described for **Example 1.** **^{1}H NMR** (CD_3OD) δ ppm 6.59 (1 H, s), 7.16 (1 H, dd, J = 5.6, 2.4 Hz), 7.29 - 7.33 (2 H, m), 7.42 (1 H, d, J = 2.4 Hz), 7.61 - 7.67 (3 H, m), 7.75 - 7.79 (2 H, m), 7.94 (2 H, s), 8.63 (1 H, d, J = 5.7 Hz), 10.79 (1 H, s); **MS** ES 424 $(M+H)^+$, calcd 424, RT = 2.48 min.

Example 38: Preparation of N^4-(4-{[1-oxido-2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine



To a solution of 6-phenyl- N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine (0.10 g, 0.24 mmol, **Example 37**) in $CHCl_3$, *m*-CPBA (77%, 0.053 g, 0.24 mmol) was added and the mixture was stirred at rt overnight. Solvent was removed in vacuo, and the residue was taken up in DMF and purified by prep-HPLC to provide 11 mg of an off-white solid (11%). **^{1}H NMR** (DMSO-d_6) δ10.20 (s, 1H), 8.64 (d, J = 5.7 Hz, 1H), 7.93 (s, 1H), 7.86 – 7.92 (m, 2H), 7.52 – 7.56 (m, 2H), 7.41 – 7.49 (m, 5H), 7.30 – 7.39 (m, 2H), 7.24 (dd, J = 5.7 Hz, 1H), 6.71 (s, 1H). **MS** ES 440 $(M+H)^+$, calcd 440, RT = 2.97 min.

Example 39: N^4-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine




A mixture containing of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy}pyridine-2-carbonitrile (3.2 g, 8.4 mmol, **Example 8**) and 10% palladium on carbon catalyst (0.75 g, Degussa) in glacial acetic acid (100 mL) was shaken on a Parr hydrogenation apparatus (3 atm H_2) until hydrogen consumption ceased. The suspension was filtered through diatomaceous earth and the filtrate was concentrated in vacuo. The residue was dissolved in *N,N*-dimethylformamide and treated with triethylamine until basic, then was added to vigorously stirred ice water. The precipitated solids were collected by suction filtration and washed with water, isopropanol, diethyl ether and finally hexane. The product was dried by air suction to afford a tan powder, 2.36 g (73%). **^{1}H NMR** (DMSO-d_6) δ ppm 9.39 (s, 1H), 8.30 (m, 1H), 7.89 (m, 4H), 7.44 (m, 3H), 7.05 (m, 2H), 6.96 (dm, 1H),

6.69 (dm, 1H), 6.51 (s, 1H), 6.36 (s, 2H), 4.16 (d, 0.5H, *J*=5.8 Hz, $C\underline{H}_2NH_2$), 3.73 (s, 1.5H, $C\underline{H}_2NH_2$), 3.28 (br s, 2H, NH_2), **MS** ES 385 $(M+H)^+$, calcd 385 RT=1.75 min.

The TFA salt (**Example 92**) was obtained by preparative HPLC of the above reaction mixture.


Example 40: Preparation of *N*-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]-phenoxy}pyridin-2-yl)methyl]methanesulfonamide



Methanesulfonyl chloride (0.040 mL, 0.52 mmol) was added to a solution of N^4-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine (**Example 39**, 0.20 g, 0.52 mmol) and DMAP (0.064 g, 0.52 mmol) in pyridine (8.0 mL) at 0 °C. The mixture was allowed to warm to rt and was stirred overnight. The mixture was concentrated in vacuo and the residue was taken up in DMF and purified by prep-HPLC to provide 82 mg of an off-white solid (27%). **^{1}H NMR** (CD_3OD) δ ppm 8.57 (1 H, d, *J*=6.8 Hz), 8.00 (2 H, s), 7.77 - 7.80 (2 H, m), 7.63 (3 H, d, *J*=7.4 Hz), 7.42 (1 H, d, *J*=2.5 Hz), 7.29 - 7.33 (3 H, m), 6.58 (1 H, s), 4.54 (2 H, s), 3.04 (3 H, s); **MS** ES 463 $(M+H)^+$, calcd 463, RT = 1.17 min.


By using the method described for **Example 40,** and by substituting appropriate starting materials, **Examples 93-98** were similarly prepared.

Example 41: Preparation of *N*-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-4-fluorobenzamide



N^4-(4-{[2-(Aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine (50 mg, 0.13 mmol, **Example 39**) and 4-fluorobenzoyl chloride (20.6 mg, 0.13 mmol) were suspended in THF (1 mL) and stirred at rt for 24 h. TLC and LC-MS indicated the reaction was completed. The mixture was extracted with EtOAc and washed with 1N aqueous sodium hydroxide solution (2X) and H_2O (3X). The organic layer was dried and concentrated to give 68 mg of the crude product. The residue was purified by Prep-TLC (CH_3OH:EtOAc=2:8) to obtain 38 mg (58%) of the title product as a yellowish oil. **^{1}H NMR** (CD_3OD) δ 8.34 (d, 1H), 7.82 (m, 4H), 7.78 (d, 2H), 7.44 (m, 3H), 7.15 (t, 2H), 7.02 (d, 2H), 6.85 (d, 1H), 6.82 (s, 1H), 6.42 (s, 1H), 4.59 (s, 2H); **MS** ES 507 $(M+H)^+$, calcd 507, RT = 2.50 min; **TLC** (MeOH/EtOAc=20/80) R_f = 0.57.


Example 42: Preparation of N'-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-*N,N*-diethylurea



N^4-(4-{[2-(Aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine (50 mg, 0.13 mmol, **Example 39**) and diethylcarbamyl chloride (20.6 mg, 0.13 mmol) were suspended in THF (1 mL) and stirred at rt for 24 h. TLC and LC-MS indicated that the reaction was complete. The mixture was extracted with EtOAc and washed with 1N aqueous sodium hydroxide solution (2X) and H_2O (3X). The organic layer was dried and concentrated to give 72 mg of the crude product. The residue was purified by Prep-TLC (CH_3OH:EtOAc=2:8) to obtain 40 mg (63%) of the title product as a yellowish oil. **^{1}H NMR** (CD_3OD) δ 8.28 (d, 1H), 7.84 (m, 3H), 7.81 (s, 1H), 7.48 (m, 3H), 7.05 (d, 2H), 6.84 (m, 1H), 6.75 (s, 1H), 6.44 (s, 1H), 4.39 (s, 2H), 3.23 (m, 4H), 1.20 (t, 6H); **MS** ES 484 $(M+H)^+$, calcd 484, RT = 2.37 min; **TLC** (MeOH/EtOAc = 20/80) R_f = 0.4

By using the method described for **Example 42**, and by substituting appropriate starting materials, **Examples 99-104** were similarly prepared.


Example 43: Preparation of N^4-[4-({4-[(2S)-(+)-2-(methoxymethyl)pyrrolidin-1-yl]pyridin-2-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine



N^4-{4-[(4-Bromopyridin-2-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine (75 mg, 0.17 mmol, was prepared by the method of **Example 1** using and **Intermediates 2T** and **1A** as starting materials. N^4-[4-(4-bromo-pyridin-2-yloxy)-phenyl]-6-phenyl-pyrimidine-2,4-diamine, was then combined with (*S*)-(+)-2-(methoxymethyl)pyrrolidine (99.5 mg, 0.86 mmol) in a 5-mL reaction flask and heated at 108 °C with stirring for 24 h. TLC and LC-MS indicated that the reaction was complete. After cooling to rt, the reaction mixture was extracted with EtOAc and washed with 1N aqueous sodium hydroxide solution (2X) and H_2O (3X). The organic layer was dried and concentrated to give 75 mg of the crude product. The residue was purified by Prep-TLC (CH_3OH:EtOAc=2:8) to obtain 32 mg (40%) of the title product as a yellowish oil. **^{1}H NMR** (CD_3OD) δ 7.82 (m, 2H), 7.70 (m, 3H), 7.41 (m, 3H), 7.04 (d, 2H), 6.41 (m, 2H), 5.94 (s, 1H), 3.86 (s, 1H), 3.38 (m, 2H), 3.30 (m, 3H), 3.18 (m, 1H), 2.01 (m, 4H), 1.30 (m, 1H); **MS** ES 469 $(M+H)^+$, calcd 469, RT = 1.9 min; **TLC** (EtOAc) R_f = 0.2.

By using the method described for **Example 43**, and by substituting appropriate starting materials, **Examples 105-109** were similarly prepared.

Example 44: Preparation of N^4-[4-({2-[(isopropylamino)methyl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine




Acetone (11.51 mg, 0.20 mmol), N^4-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine (80 mg, 0.21 mmol, **Example 39**) and titanium (IV) methoxide (68.2 mg, 0.40 mmol) were suspended in CH_2Cl_2 (5 mL) and stirred at rt for 24 h. Sodium triacetoxyborohydride (105 mg, 0.50 mmol) was added into the reaction mixture and

the mixture was stirred at rt for another 24 h. The mixture was filtered through a Celite® pad and washed with CH_2Cl_2. A small amount of Celite® was added to the filtrate and 5 mL of water was added to quench the reaction. After it was stirred for 20 min, the CH_2Cl_2 was removed in vacuo. The residue was taken up in ethyl acetate and washed with 1N NaOH (2X) and water (3X). The organic layer was concentrated and purified by Prep-TLC (MeOH) to give 36 mg (42.2%) of the title product as a white solid. **^{1}H NMR** (CD_3OD) δ 8.35 (d, 1H), 7.82 (d, 2H), 7.00 (d, 2H), 7.44 (m, 3H), 7.08 (m, 2H), 7.01 (s, 1H), 6.82 (m, 1H), 6.46 (s, 1H), 3.81 (s, 2H), 2.80 (m, 1H), 1.10 (d, 6H); **MS** ES 427 $(M+H)^+$, calcd 427, RT = 2.59 min; **TLC** (MeOH) R_f = 0.38.


By using the method described for **Example 44**, and by substituting appropriate starting materials, **Examples 110-111** were similarly prepared.

Example 45: Preparation of 4-[4-(2-Amino-6-phenylpyrimidin-4-ylamino)phenoxy]pyridine-2-carboxylic acid (2-hydroxyethyl)amide




Chloropyrimidine **1A** (0.2 g, 0.97 mmol) and **Intermediate 2N** (0.38 g, 0.97 mmol), were suspended in *n*-butanol (5 mL) and heated at 80 °C for 12 h. LC-MS indicated that the reaction was complete. KF was then added to the reaction mixture and heating was continued at 80 °C for 5 h. The solvent was removed by rotary evaporation, and the residue was treated with 10% sodium carbonate and extracted with EtOAc (20 mL x 3). The organic extracts were combined, washed with water and brine, dried over magnesium sulfate, and evaporated to afford a solid that was washed with methanol to give pure product, 0.19 g (44%). **^{1}H NMR** (DMSO-d_6) δ 9.36 (s, 1H), 8.62 (t, 1H), 8.45 (d, 1H), 7.88 (m, 4H), 7.47 (m, 3H), 7.38 (d, 1H), 7.11 (m, 3H), 6.45 (s, 1H), 6.34 (s, 2H), 3.49 (t, 2H), 3.36 (t, 2H); **MS** ES 443 $(M+H)^+$ calcd 443.

Example 46: Preparation of 6-phenyl-N^4-{4-[2-(1*H*-tetrazol-5-yl)pyridin-4-yloxy]phenyl}pyrimidine-2, 4-diamine




A mixture of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carbonitrile (0.20 g, 0.53 mmol, prepared in **Example 8**), sodium azide (0.051 g, 0.79 mmol), and triethylamine (0.11 g, 0.79 mmol) in toluene (15 mL) was heated at 100 °C for 2 days. The mixture was then treated with cold water. The solid was collected by filtration, washed with water and methanol to give pure product, 0.14 g (63%). **^{1}H NMR** (DMSO-d_6) δ 9.51 (s, 1H), 8.62 (d, 1H), 7.94 (m, 4H), 7.55 (d, 1H), 7.48 (m, 3H), 7.19 (m, 3H), 6.48 (m, 3H); **MS** ES 424 $(M+H)^+$ calc 424.

Example 47: Preparation of N^4-{4-[2-(4,5-Dihydro-1*H*-imidazol-2-yl)pyridin-4-yloxy]phenyl}-6-phenylpyrimidine-2,4-diamine



A mixture of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carbonitrile (0.2 g, 0.53 mmol, prepared in **Example 8**), ethylene diamine (0.095 g, 1.58 mmol), and sulfur (0.05 g, 1.58 mmol) in DMF (3 mL) was heated at 80 °C for 3 days. The solvent was then removed by evaporation under reduced pressure. The residue was purified by preparative HPLC followed by preparative TLC (EtOAc:NH_4OH = 99:2) to afford pure product, 0.01 g (5%). **^{1}H NMR** (DMSO-d_6) δ 9.39 (s, 1H), 8.41 (d, 1H), 7.84 (m, 4H), 7.42 (m, 3H), 7.37 (d, 1H), 7.09 (m, 3H), 6.95 (s, 1H), 6.46 (s, 1H), 6.37 (s, 2H); **MS** ES 424 $(M+H)^+$ calc 424.


Example 48: Preparation of N^4-[4-(2-Chloro-pyridin-4-yloxy)-phenyl]-6-phenyl-pyrimidine-2,4-diamine



This compound was prepared by reaction of **1A** with **2R** by the method described in **Example 1**. **^{1}H NMR** (DMSO-d_6) δ 9.40 (s, 1H), 8.22 (d, 1H), 7.85 (m, 4H), 7.45 (m, 3H), 7.11 (m, 2H), 6.88 (m, 2H), 6.46 (s, 1H), 6.36 (s, 2H); **MS** ES 390 $(M+H)^+$, calcd 390, RT = 2.27 min.

By using the method described for **Example 48**, and by substituting appropriate starting materials, **Examples 113-118** were similarly prepared.


Example 49: Preparation of (*S*)-N^4-{4-[2-(2-Methoxymethylpyrrolidin-1-yl)pyridin-4-yloxy]phenyl}-6-phenylpyrimidine-2,4-diamine



A mixture of N^4-{4-[(2-chloropyridin-4-yl)oxy]phenyl}- 6-phenylpyrimidine-2,4-diamine (0.15 g, 0.38 mmol, prepared in **Example 48**) and (*S*)-(+)-2-(methoxymethyl)pyrrolidine (2 mL) was heated at 80 °C for 3 days. The mixture was cooled to rt and separated by preparative HPLC directly. The desired fractions were combined, neutralized by 10% sodium carbonate, and extracted with EtOAc (3X). The extracts were combined, dried over magnesium sulfate, and evaporated to furnish pure product, 0.04 g (22%). **^{1}H NMR** (DMSO-d_6) δ 9.24 (s, 1H), 7.89 (m, 3H), 7.78 (m, 2H), 7.42 (m, 3H), 7.00 (m, 2H), 6.44 (s, 1H), 6.34 (s, 2H), 6.10 (m, 1H), 5.85 (d, 1H), 4.06 (m, 2H), 3.40 (m, 1H), 3.21 (s, 3H), 3.08 (m, 4H), 1.82 (m, 4H); **MS** ES 469 $(M+H)^+$ calc 469.

By using the method described for **Example 49**, and by substituting appropriate starting materials, **Examples 119-147** were similarly prepared.


Example 50: Preparation of 2-amino-*N*-(4-{2-amino-6-[4-(2-trifluoromethylpyridin-4-yloxy)phenylamino]pyrimidin-4-yl}phenyl)-3-hydroxypropionamide




To a solution of (4S)-3-(tert-butoxycarbonyl)-2,2-dimethyl-1,3-oxazolidine-4-carboxylic acid (0.56 g, 2.3 mmol) in dry *N,N*-dimethylacetamide (10 mL) was added HATU (0.11 g, 2.87 mmol) and DIEA (0.742 g, 5.75 mmol). After the reaction solution was stirred at rt for 1 h, 6-(4-aminophenyl)-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine (**Example 165,** 0.84 g, 1.92 mmol) was added. The solution was stirred for an additional 24 h at rt and separated by preparative HPLC to afford a solid intermediate, which was treated with methanol (10 mL) and concentrated HCl (0.5 mL) for 12 h at rt. The resulting mixture was diluted with DMSO and purified by HPLC to give a solid. The solid was stirred with saturated sodium bicarbonate and EtOAc for 2 h. The organic layer was separated, washed with water and brine, dried over magnesium sulfate, and evaporated to afford 0.433 g (43%) of pure product. **^{1}H NMR** (DMSO-d_6) δ 9.34 (s, 1H), 8.59 (d, 1H), 7.88 (m, 4H), 7.76 (d, 2H), 7.37 (s, 1H), 7.16 (m, 3H), 6.45 (s, 1H), 6.36 (s, 2H), 4.88 (t, 1H), 3.56 (m, 2H), 3.39 (m, 1H); **MS** ES 526 $(M+H)^+$ calcd 526.

By using the method described for **Example 50**, and by substituting appropriate starting materials, **Examples 148-152** were similarly prepared.

Example 51: Preparation of 4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenol



Step 1: Preparation of 6-chloro-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine




A stirred solution containing 2-amino-4,6-dichloropyrimidine (17.74 g, 0.11 mol) and 4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}aniline (**Intermediate 2U**) in water (400 mL), isopropanol (100 mL) and concentrated hydrochloric acid (5 mL) was heated at 65 °C for 18 h. The reaction was cooled to 0 °C and the yellow-tan solid was collected by suction filtration and washed with water. The filtered product was dissolved in hot *N,N*-dimethylformamide (90 °C) and triethylamine was slowly added until the solution was slightly basic (pH~8). The solution was then cooled to rt, added to vigorously stirred ice water (1.2 L), and stirring was continued for 1 h. The tan solids were collected by suction filtration and washed sequentially with water , isopropanol, diethyl ether and finally hexane. The material was dried by air suction to afford a light tan solid, 28.8 g (77%). **^{1}H NMR** (DMSO- d_6) δ ppm 9.46 (s, 1H), 8.59 (d, 1H, *J*=5.6 Hz), 7.81 (d, 2H, *J*=9.0), 7.37 (d, 1H, *J*=2.4 Hz), 7.17 (d, 2H, *J*=9.0 Hz), 7.11 (dd, 1H, *J*= 2.6, 5.6 Hz), 6.78 (s, 2H), 6.00 (s, 1H). **MS** ES 382 $(M+H)^+$, calcd 382 RT=2.93 min.

Step 2: Preparation of the title compound

To a 8 mL microwave tube was added 6-chloro-*N*^{4}-(4-{[2-(trifluoromethyl)- pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine (0.2 g, 0.52 mmol), boronic ester (0.17 g, 0.79 mmol), $PdCl_2$dppf-CH_2Cl_2 complex (0.023 g, 0.03 mmol), potassium carbonate (0.18 g, 1.3 mmol), *N,N*-dimethylacetamide (3 mL), and water (1 mL). The mixture was degassed, flushed with nitrogen, and heated at 150 °C for 15 min in a microwave reactor. The mixture was filtered, and the filtrate was separated by prep HPLC. The desired fractions were combined, basified, and extracted with EtOAc (3X). The EtOAc extracts were then washed with water and brine, dried over magnesium sulfate, and evaporated to give 45 mg (20%) pure product. **^{1}H NMR** (DMSO-d_6) δ 9.75 (s, 1H), 9.22 (s, 1H), 8.58 (d, 1H), 7.86 (d, 2H), 7.76 (d, 2H), 7.35 (s, 1H), 7.12 (m, 3H), 6.78 (m, 2H), 6.39(s, 1H), 6.20 (s, 2H); **MS** ES 440 $(M+H)^+$, calcd 440.

By using the method described for **Example 51**, and by substituting appropriate starting materials, **Examples 153-184** were similarly prepared.


Example 52: Preparation of sulfamic acid, 4-{2-amino-6-[4-(2-trifluoromethylpyridin-4-yloxy)phenylamino]pyrimidin-4-yl}phenyl ester



To neat chlorosulfonyl isocyanate (0.166 g, 1.37 mmol) was added dropwise formic acid (97%, 0.63 g, 1.37 mmol) while cooling in ice-water bath. The mixture was stirred at rt until gas evolution ceased. The resulting sulfamoyl chloride was added to a solution of 4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl) amino] pyrimidin-4-yl}phenol (0.06 g, 0.14 mmol, **Example 51)** in dry *N,N*-dimethylacetamide at 0 °C. The reaction mixture was then stirred at rt for 12 h. The solution was diluted with methanol and separated by preparative HPLC. The desired fractions were combined, basified with saturated sodium carbonate, and extracted with EtOAc (3X). The extracts were combined, washed with brine, dried over magnesium sulfate, and evaporated to give desired product, 0.025 g (35%). **^{1}H NMR** (DMSO-d_6) δ 9.42 (s, 1H), 8.59 (d, 1H), 8.09 (d, 2H), 7.98 (m, 2H), 7.85 (m, 2H), 7.35 (m, 3H), 7.10 (m, 3H), 6.42 (s, 1H), 6.36 (s, 2H); **MS** ES 519 $(M+H)^+$ calcd 519.

By using the method described for **Example 52**, and by substituting appropriate starting materials, **Examples 185-186** and **332** were similarly prepared. In addition, by selection of the appropriate starting materials, this method can be used for the preparation of Examples **326-331,** and **333-334**

Example 53: Preparation of N^4-[4-(2-aminopyridin-4-yloxy)phenyl]-6-phenylpyrimidine-2,4-diamine



To a 8-mL vial was added $Pd_2(dba)_3$ (0.028 g, 0.03 mmol), 2-dicyclohexylphosphinobiphenyl (0.025 g, 0.070 mmol), and N^4-{4-[(2-chloropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine (0.20 g, 0.51 mmol, **Example 48**). The vial

was sealed, evacuated, and back filled with nitrogen. THF was then added via syringe, followed by addition of LiHMDS (1M in THF, 0.72 mL, 0.72 mmol). The mixture was heated at 65 °C overnight. The mixture was then cooled to rt, treated with 1N HCl, and stirred for 12 h at rt. The mixture was then neutralized using 1N NaOH and extracted with methylene chloride (10 mL x 3). The organic extracts were combined, washed with brine, dried over magnesium sulfate, and purified by preparative HPLC to furnish 0.035 g of the desired product (18%). **^{1}H NMR** (CD_3OD) δ 7.96 (d, 2H), 7.79 (m, 3H), 7.60 (m, 3H), 7.22 (m, 2H), 6.61 (d, 1H), 6.58 (s, 1H), 6.20 (s, 1H); **MS** ES 371 $(M+H)^+$ calcd 371.

Example 54: High-Speed Analoging (HSA) Synthesis Method A

To a mixture of 1 equivalent of the chloropyrimidine (100 mg, e.g. compound **9 General Method B**), 2 equivalent of boronic acid (e.g. **General Method B**), and 0.06 equivalent of $PdCl_2$(dppf) CH_2Cl_2 complex in 2.3 mL anhydrous N,N-dimethylacetamide in a 5 mL microwave reaction vessel was added 3.1 equivalent of 2 M K_2CO_3 aqueous solution. After the resulting mixture was degassed for 10 min using N_2, the vial was sealed and heated at 150 °C for 20 min in a microwave reactor. The reaction mixture was filtered, and the filtrate was purified by pre-HPLC eluting with 15% to 85 % acetonitrile using a Phenomenex Luna 5 μ C18 150 x 30 mm column to provide the final product.

By using the appropriate starting materials, the method described for **Example 54**, was utilized for the preparation of **Examples 112** and **187-219**.

Example 55: High-Speed Analoging (HSA) Synthesis Method B

To a mixture of 1 equivalent of the chloropyrimidine (100 mg, e.g. compound **9 General Method B**), 2 equivalents of boronic acid (**General Method B**), and 0.06 equivalent of $PdCl_2$(dppf) CH_2Cl_2 complex in 2.3 mL anhydrous *N,N*-dimethylacetamide in a 8 mL microwave reaction vessel was added 3.1 equivalent of 2M K_2CO_3 aqueous solution. After the resulting mixture was degassed for 10 min using N_2, the vial was sealed and heated at 140 °C for 20 min in a microwave reactor. The reaction mixture was filtered, and the filtrate was purified by pre-HPLC eluting with 15% to 85 % acetonitrile using a Phenomenex Luna 5 μ C18 150 x 30 mm column to provide the final product.


By using the appropriate starting materials, the method described for **Example 55**, was utilized for the preparation of **Examples 220-279**

Example 56: High-Speed Analoging (HSA) Synthesis Method C


A mixture of 1 equivalent chloropyrimidine (e.g. compound **9 General Method B)**, 2 equivalents of boronic acid **(General Method B)**, and 0.1 equivalent of $PdCl_2(dppf)$-CH_2Cl_2 complex in 2.5 mL anhydrous *N,N*-dimethylacetamide and 0.5 mL of 2 M K_2CO_3 in water in a 5 mL microwave reaction vessel under nitrogen was heated at 140 °C for 20 min in the personal microwave reactor. The reaction mixture was filtered, and the filtrate was purified by pre-HPLC eluting with 15% to 85 % acetonitrile containing 0.1%TFA using a Phenomenex Luna 5 μ C18 150 x 30 mm column to provide the final product.

By using the appropriate starting materials, the method described for **Example 56,** was utilized for the preparation of **Examples 280-325**

Example 57: Preparation of 4-[2-amino-6-({4-[(2-chloropyridin-4yl)oxy]phenyl} amino) pyrimidin-4-yl]phenol



Step 1: Preparation of 4-(2-amino-6-chloropyrimidin-4-yl)phenyl *tert*-butyl carbonate




To a mixture of 2-amino-4,6-dichloropyrimidine (1.5 g, 9.15 mmol), *t*-butyl-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)phenyl carbonate (2.9 g, 9.15 mmol), $PdCl_2$dppf CH_2Cl_2 complex (0.45 g, 0.55 mmol), and DME (14 mL) was added a solution of potassium carbonate (3.2 g, 22.9 mmol) in water (4mL). The mixture was then degassed, flushed with nitrogen and heated at 80 °C overnight. The organic layer was separated, washed with water and brine, dried over magnesium sulfate, and evaporated. The residue

was purified by column (2% MeOH:50% hexane:48% EtOAc) to afford 0.64 g (22%) desired product. **MS** ES 322 $(M+H)^+$, calcd 322, RT = 3.37 min.

Step 2: Preparation of title compound


This material was prepared by a method analogous to that described for **Example 1**, starting from the product from 4-(2-amino-6-chloropyrimidin-4-yl)phenyl *tert*-butyl carbonate and **Intermediate 2R**. **^{1}H NMR** (DMSO-d_6) δ 9.79 (s, 1H), 9.25 (s, 1H), 8.22 (d, 1H), 7.86 (m, 2H), 7.74 (m, 2H), 7.09 (m, 2H), 6.95 (m, 2H), 6.80 (m, 2H), 6.39 (s, 1H), 6.23 (s, 2H); **MS** ES 406 $(M+H)^+$, calcd 406, RT = 2.74 min.

Example 58: Preparation of (3E)-4-(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy}pyridin-2-yl)but-3-en-1-ol



N^4-{4-[(2-chloropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine (0.10 g, 0.26 mmol, **Example 48**), K_2CO_3 (0.089 g, 0.64 mmol), and DMA (2.5 mL) were placed into a small microwave vial. The mixture was degassed for 10 min before (3E)-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)but-3-en-1-ol (0.10 g, 0.33 mmol) and Pd(dppf)Cl_2 CH_2Cl_2 complex (0.012 g, 0.020 mmol) were added. The mixture was heated at 150 °C for 20 min in a microwave reactor. The mixture was cooled, filtered, and purified by prep-HPLC. Concentration of the desired fractions gave 0.016 g of the title compound (10%). **MS** ES: 426 $(M+H)^+$, calcd 426, RT = 1.90 min.

Example 59: Preparation of (4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy} pyridin-2-yl)methanol trifluoroacetate)



To a cloudy solution of 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino] phenoxy} pyridine-2-carboxylic acid (748 mg, 1.87 mmol, **Example 20**) in anhydrous DMF (50 mL) at rt was added carbonyldiimidazole (456 mg, 2.81 mmol). The white suspension was stirred at 80 °C overnight, concentrated to a volume of 10 mL, and diluted with anhydrous THF (7 mL). The reaction mixture was cooled to 0 °C and water (10 mL) was added. The mixture was vigorously stirred as $NaBH_4$ (142 mg, 3.75 mmol) was added and was allowed to warm from 0 °C to rt over 2 h before it was quenched with conc. HCl (1 mL) in an ice bath. After stirring for 15 min, the mixture was slowly added to a stirred solution of sat. $NaHCO_3$ (20 mL) at 0 °C. After stirring for 30 min, it was extracted with EtOAc (3 x 100 mL). The combined organic layers were dried over Na_2SO_4, filtered, and concentrated to give an off-white gum (420 mg, 85% pure). The crude material (100 mg) was purified by prep HPLC purification to give 37 mg (40% yield) of the title compound as a colorless gum. **^{1}H NMR** (DMSO-d_6) δ 10.8 (s, 1H), 8.55 (d, 1H), 7.90 (m, 2H), 7.75 (m, 2H), 7.60 (m, 3H), 7.35 (d, 2H), 7.20 (m, 2H), 6.60 (s, 1H), 4.60 (s, 2H); **MS** ES 386 $(M+H)^+$ calcd 386, RT = 1.73 min.

Examples of the compounds of the invention are summarized in Table 1 below. It is to be noted that in some of the IUPAC names, "N4*" is to be understood as a symbol for "N^4".

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
1		N^4-{4-[(2-ethylpyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	384 (1.87)
2		N^4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	370 (1.41)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
3		4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxamide	399.4 (2.74)
4		4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N-methylpyridine-2-carboxamide	413 (2.13)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
5		N^4-{4-[(3,5-difluoropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	392 (2.27)
6		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]-3-fluorophenoxy}pyridine-2-carbonitrile hydrochloride	399.3 (2.06)
7		N4*-{3-fluoro-4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	388 (1.70)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
8		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carbonitrile	381 (2.22)
9		4-(3-{[2-amino-6-(3-furyl)pyrimidin-4-yl]amino}phenoxy)-N-methylpyridine-2-carboxamide	403 (1.99)
10		4-(4-{[2-amino-6-(3-furyl)pyrimidin-4-yl]amino}phenoxy)-N-methylpyridine-2-carboxamide	403 (1.94)

Table 1










Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
11		N4*-[4-(4-nitrophenoxy)phenyl]-6-phenylpyrimidine-2,4-diamine	400 (3.01)
12		N4*-[4-(4-chlorophenoxy)phenyl]-6-phenylpyrimidine-2,4-diamine	389 (2.78)
13		N4*-[4-(4-methoxyphenoxy)phenyl]-6-phenylpyrimidine-2,4-diamine	385 (2.48)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
14		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]-2-fluorophenoxy}pyridine-2-carbonitrile	
15		4-[4-[(2-amino-6-phenylpyrimidin-4-yl)amino]-3-(trifluoromethyl)phenoxy]pyridine-2-carbonitrile	
16		N^4-{4-[(2-methylpyrimidin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	

106

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
17		N^4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-[4-(2-pyrrolidin-1-ylethoxy)phenyl]pyrimidine-2,4-diamine	
18		4-[4-({2-amino-6-[4-(2-piperidin-1-ylethoxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	
19		methyl 4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxylate	414.2 (2.16)

107

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
20		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxylic acid	400 (1.71)
21		N4*-(4-{[2-(morpholin-4-ylcarbonyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine trifluoroacetate	469.2 (2.13)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
22		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N,N-dimethylpyridine-2-carboxamide	
23		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N-(2-methoxyethyl)pyridine-2-carboxamide	457.4 (2.17)
24		4-[4-({2-amino-6-[4-(trifluoromethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]-N-(2-methoxyethyl)pyridine-2-carboxamide	

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
25		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]benzyl}-N-(2-methoxyethyl)-N-methylpyridine-2-carboxamide	
26		N4*-[4-({2-[(4-methylpiperazin-1-yl)carbonyl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine trifluoroacetate	482.2 (1.86)
27		N^4-{4-[(2-{[(2-methoxyethyl)amino]methyl}pyridin-4-yl)oxy]phenyl}6-phenylpyrimidine-2,4-diamine	

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
28		6-(4-fluorophenyl)-N^4-(4-{[2-(piperidin-1-ylmethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	
29		6-(4-methoxyphenyl)-N^4-(4-{[2-(morpholin-4-ylmethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
30		N^2-ethyl-6-(3-methoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine trifluoroacetate	482 (2.83)
31		4-{4-[(2-amino-5-bromo-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carbonitrile	459.3 (2.85)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
32		N4*-(4-{[2-(2-morpholin-4-ylethoxy)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	485.1 (1.96)
33		6-phenyl-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]methyl}phenyl)pyrimidine-2,4-diamine	422.3 (2.51)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
34		N4*-{4-[(2-chloropyridin-4-yl)methyl]phenyl}-6-phenylpyrimidine-2,4-diamine	388.3 (2.38)
35		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]benzyl}pyridine-2-carbonitrile	379.3 (2.35)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
36		N4*-(4-{[2-(aminomethyl)pyridin-4-yl]methyl}phenyl)-6-phenylpyrimidine-2,4-diamine	383.3 (1.84)
37		6-phenyl-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	424.3 (2.48)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
38		N4*-(4-{[1-oxido-2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	440.3 (2.97)
39		N4*-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	385.2 (1.80)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
40		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]methanesulfonamide trifluoroacetate	463.3 (1.17)
41		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-4-fluorobenzamide	507.7 (2.5)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
42		N'-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-N,N-diethylurea	484.4 (2.37)
43		N4*-[4-({4-[2-(methoxymethyl)pyrrolidin-1-yl]pyridin-2-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	469.4 (1.90)
44		N4*-[4-({2-[(isopropylamino)methyl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	427.1 (2.59)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
45		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N-(2-hydroxyethyl)pyridine-2-carboxamide	443.3 (2.19)
46		6-phenyl-N4*-(4-{[2-(1H-tetrazol-5-yl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	424.1 (1.84)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
47		N4*-(4-{[2-(4,5-dihydro-1H-imidazol-2-yl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	424.2 (0.81)
48		N4*-{4-[(2-chloropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	390.3 (2.40)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
49	Chiral	N4*-[4-({2-[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	469.3 (1.36)
50		N-(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)serinamide	536.1 (2.14)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
51		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenol	440.3 (1.56)
52		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl sulfamate	519.1 (2.64)
53		N4*-{4-[(2-aminopyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	371.4 (1.90)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
57		4-[2-amino-6-({4-[(2-chloropyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenol	406.4 (2.74)
58		(3E)-4-(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)but-3-en-1-ol trifluoroacetate (salt)	426.3 (1.90)
59		(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methanol trifluoroacetate	386.2 (1.73)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
60		ethyl 2-[(2-amino-6-phenylpyrimidin-4-yl)amino]-5-[(5-chloropyridin-3-yl)oxy]benzoate	462.2 (2.53)
61		6-phenyl-N4*-[4-(pyrimidin-5-yloxy)phenyl]pyrimidine-2,4-diamine	357.3 (2.07)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
62		N^4-[4-(4-aminophenoxy)phenyl]-6-phenylpyrimidine-2,4-diamine	370.3 (2.08)
63		N4*-{4-[(4-bromopyridin-2-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	436.1 (2.81)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
64		N4*-{4-[(2-fluoropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine trifluoroacetate	374.7 (2.52)
65		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-ol	372.9 (2.34)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
66		N4*-{4-[(2-chloropyridin-4-yl)oxy]-2-fluorophenyl}-6-phenylpyrimidine-2,4-diamine	408.3 (1.42)
67		N4*-(2-fluoro-4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	442.3 (2.54)
68		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}benzonitrile	380.4 (2.39)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
69		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxamide	399.3 (2.54)
70		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N-methylpyridine-2-carboxamide	413.3 (2.21)
71		4-(4-{[2-amino-6-(4-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)-N-methylpyridine-2-carboxamide	431.3 (2.25)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
72		4-(3-{[2-amino-6-(4-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)-N-methylpyridine-2-carboxamide	431.5 (2.15)
73		4-(4-{[2-amino-6-(4-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	417.3 (2.07)
74		4-(3-{[2-amino-6-(4-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	417.3 (2.09)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
75		4-(4-{[2-amino-6-(4-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	399.4 (2.21)
76		4-(3-{[2-amino-6-(3-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)-N-methylpyridine-2-carboxamide	443.4 (2.18)
77		4-(3-{[2-amino-6-(4-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)-N-methylpyridine-2-carboxamide	443.4 (2.19)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
78		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxylic acid hydrochloride	400.1 (1.67)
79		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N-1-azabicyclo[2.2.2]oct-3-ylpyridine-2-carboxamide trifluoroacetate	508.3 (1.99)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
80		4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-N-(2,3-dihydroxypropyl)pyridine-2-carboxamide	473.2 (2.11)
81	Chiral	N4*-{4-[(2-{[(2S)-2-(methoxymethyl)pyrrolidin-1-yl]carbonyl}pyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine trifluoroacetate	497.3 (2.23)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
82	 Chiral	(3R)-1-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)carbonyl]pyrrolidin-3-ol trifluoroacetate (salt)	469.4 (1.98)
83		N4*-{4-[(2-butoxypyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	428.3 (2.56)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
84		N^4-(4-{[2-(2-methoxyethoxy)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	430.3 (2.24)
85		N^4-[4-({2-[2-(dimethylamino)ethoxy]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	443.1 (1.94)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
86		N4*-{4-[(2-ethoxypyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	400.3 (2.24)
87		1-[4-({5-[(2-amino-6-phenylpyrimidin-4-yl)amino]pyridin-2-yl}oxy)phenyl]ethanone	398.2 (2.29)
88		N^{4*}-[4-(4-aminobenzyl)phenyl]-6-phenylpyrimidine-2,4-diamine	368.3 (1.91)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
89		N^4-[4-(4-methylbenzyl)phenyl]-6-phenylpyrimidine-2,4-diamine	367.4 (2.78)
90		N^4-[4-(4-methoxybenzyl)phenyl]-6-phenylpyrimidine-2,4-diamine	398.2 (2.29)
91		N^4-[4-(4-fluorobenzyl)phenyl]-6-phenylpyrimidine-2,4-diamine	371.4 (2.68)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
92		N4*-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine trifluoroacetate	385.2 (1.86)
93		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-1-methyl-1H-imidazole-4-sulfonamide trifluoroacetate	529.3 (1.15)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
94		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]ethanesulfonamide trifluoroacetate	477.4 (1.89)
95		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]propane-2-sulfonamide trifluoroacetate	491.4 (2.06)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
96		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]benzenesulfonamide trifluoroacetate	525.4 (2.20)
97		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-4-methoxybenzenesulfonamide trifluoroacetate	555.4 (2.21)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
98		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-4-fluorobenzenesulfonamide trifluoroacetate	543.4 (2.24)
99		N'-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-N,N-dimethylurea	457.1 (2.52)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
100		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]morpholine-4-carboxamide	498.4 (2.37)
101		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]urea trifluoroacetate	428.3 (0.72)
102		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-N,N',N'-trimethylurea	470.1 (1.88)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
103		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-N',N'-diethyl-N-methylurea	498.2 (2.04)
104		N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-N-methylmorpholine-4-carboxamide	512.2 (1.88)
105		N4*-{4-[(4-morpholin-4-ylpyridin-2-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	441.4 (1.27)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
106		6-phenyl-N4*-{4-[(4-piperidin-1-ylpyridin-2-yl)oxy]phenyl}pyrimidine-2,4-diamine	439.4 (1.93)
107		N4*-(4-{[4-(4-methylpiperazin-1-yl)pyridin-2-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	454.2 (2.22)
108		2,2'-[(2-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-4-yl)imino]diethanol	459.3 (2.38)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
109		N4*-[4-({4-[bis(2-methoxyethyl)amino]pyridin-2-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	488.0 (2.32)
110		N4*-{4-[(2-{[(1-ethylpropyl)amino]methyl}pyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	455.2 (2.47)
111		N4*-[4-({2-[(cyclohexylamino)methyl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	467.1 (2.42)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
112		6-(3,4-dichlorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	492.3 (3.29)
113		N4*-{3-[(2-chloropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	390.3 (2.37)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
114		N4*-{4-[(2-chloropyridin-4-yl)oxy]phenyl}-6-(4-fluorophenyl)pyrimidine-2,4-diamine	408.3 (1.04)
115		N4*-{4-[(2-chloropyridin-4-yl)oxy]phenyl}-6-(3-nitrophenyl)pyrimidine-2,4-diamine	435.2 (1.27)
116		6-(4-aminophenyl)-N4*-{4-[(2-chloropyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	405.3 (1.60)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
117		N4*-{4-[(2-chloropyridin-4-yl)oxy]-3-fluorophenyl}-6-phenylpyrimidine-2,4-diamine	408.2 (1.72)
118		6-(3-aminophenyl)-N4*-{4-[(2-chloropyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	405.3 (1.05)
119		6-phenyl-N4*-{3-[(2-piperidin-1-ylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	439.3 (0.25)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
120		N4*-{3-[(2-morpholin-4-ylpyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	441.3 (0.25)
121		6-phenyl-N4*-{3-[(2-pyrrolidin-1-ylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	425.3 (0.25)
122		N4*-[3-({2-[2-(methoxymethyl)pyrrolidin-1-yl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	469.3 (0.21)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
123	 Chiral	[(2R)-1-(4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)pyrrolidin-2-yl]methanol	455.4 (2.34)
124		N4*-[3-({2-[(2-methoxyethyl)(methyl)amino]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	443.4 (1.71)
125		N4*-(3-{[2-(2-methylpyrrolidin-1-yl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	439.3 (2.01)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
126		N4*-[3-({2-[bis(2-methoxyethyl)amino]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	487.2 (1.03)
127		2,2'-[(4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)imino]diethanol	459.3 (0.93)
128	 Chiral	[(2S)-1-(4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)pyrrolidin-2-yl]methanol	455.4 (2.41)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
129		2,2'-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)imino]diethanol	459.4 (2.34)
130		N4*-(3-{[2-(4-methylpiperazin-1-yl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	454.3 (2.37)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
131		N4*-(4-{[2-(1-benzylhydrazino)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	476.2 (2.19)
132		N4*-(4-{[2-(benzylamino)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine trifluoroacetate	461.3 (2.10)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
133	Chiral	N4*-[4-({2-[(2R)-2-(methoxymethyl)pyrrolidin-1-yl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	469.3 (0.96)
134		N4*-{4-[(2-morpholin-4-ylpyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	441.3 (1.74)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
135		N4*-(4-{[2-(4-methylpiperazin-1-yl)pyridin-4-yl]oxy}phenyl)-6-phenylpyrimidine-2,4-diamine	454.2 (1.47)
136		N4*-[4-({2-[4-(2-methoxyethyl)piperazin-1-yl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	498.2 (1.59)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
137		N4*-[4-({2-[(2-methoxyethyl)(methyl)amino]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	443.3 (0.82)
138		N4*-{4-[(2-{[2-(dimethylamino)ethyl]amino}pyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	442.2 (0.82)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
139		6-phenyl-N4*-{4-[(2-piperidin-1-ylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	439.3 (1.82)
140		N4*-[4-({2-[bis(2-methoxyethyl)amino]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	487.2 (1.34)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
141		N4*-{4-[(2-{[3-(diethylamino)propyl]amino}pyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine	484.2 (0.86)
142		N4*-[4-({2-[(3-morpholin-4-ylpropyl)amino]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	498.2 (0.94)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
143		4-[2-amino-6-({4-[(2-morpholin-4-ylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]-2-methoxyphenol	487.3 (0.97)
144	 Chiral	N4*-[4-({2-[(3S)-3-(dimethylamino)pyrrolidin-1-yl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	468.2 (1.01)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
145	Chiral	[(2S)-1-(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)pyrrolidin-2-yl]methanol	455.3 (1.91)
146	Chiral	(3R)-1-(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)pyrrolidin-3-ol	441.3 (1.31)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
147	Chiral	N4*-[3-({2-[(2R)-2-(methoxymethyl)pyrrolidin-1-yl]pyridin-4-yl}oxy)phenyl]-6-phenylpyrimidine-2,4-diamine	469.4 (1.71)
148		4-[(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)amino]-4-oxobutanoic acid	539.4 (2.25)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
149		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-2-methoxyphenyl N-(tert-butoxycarbonyl)glycinate	627.0 (2.8)
150		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-2-methoxyphenyl glycinate hydrochloride	527.2 (2.54)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
151		N-(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)glycinamide trifluoroacetate	496.2 (2.11)
152		4-amino-N-(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)butanamide trifluoroacetate	524.1 (2.15)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
153		6-(4-amino-3-methoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	469.4 (2.67)
154		N4*-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-(2-methylphenyl)pyrimidine-2,4-diamine	399.6 (2.15)
155		6-[2-(trifluoromethyl)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	493.0 (2.96)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
156		6-(2-ethylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	452.9 (2.89)
157		6-(2-ethoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	469.0 (2.81)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
158		6-(2-fluoro-5-methylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	456.7 (2.87)
159		6-(2,3-difluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	460.0 (2.89)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
160		6-(2,5-difluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	460.7 (2.78)
161		6-[2-(trifluoromethoxy)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	508.6 (2.78)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
162		6-(2-isopropylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	466.7 (2.99)
163		N4*-(4-{[2-(aminomethyl)pyridin-4-yl]oxy}phenyl)-6-(2-ethylphenyl)pyrimidine-2,4-diamine	413.3 (1.99)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
164	 ClH	6-(4-aminophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine hydrochloride	439.3 (2.47)
165		6-(4-aminophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	439.3 (2.37)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
166		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-2-methoxyphenol	470.2 (1.46)
167		3-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenol	440.3 (1.08)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
168		6-(3-amino-4-methoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	469.3 (2.45)
169		6-(3-aminophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	439.3 (2.37)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
170	 ClH	6-(3-aminophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine hydrochloride	439.4 (2.36)
171		4-(4-{[2-amino-6-(4-hydroxy-3-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	427.3 (2.16)
172		4-(4-{[2-amino-6-(3-amino-4-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	426.3 (2.10)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
173		4-(4-{[2-amino-6-(4-amino-3-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	444.2 (2.22)
174		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-3-methylphenol	454.3 (2.48)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
175		(2-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)methanol	454.2 (2.38)
176		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-3-(trifluoromethyl)phenol	508.2 (2.62)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
177		6-(4-amino-3-fluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	457.3 (2.52)
178		tert-butyl {4-[2-amino-6-({4-[(2-cyanopyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]-2-methoxyphenyl}carbamate	526.2 (2.79)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
179		4-(4-{[2-amino-6-(4-amino-3-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile trifluoroacetate	426.2 (2.31)
180		6-(4-amino-3-methylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	453.4 (2.26)

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
181		6-(6-aminopyridin-3-yl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	440.3 (2.01)
182		6-(3-amino-4-chlorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	473.3 (2.61)
183		4-(4-{[2-amino-6-(6-aminopyridin-3-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	397.3 (1.85)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
184		2-amino-5-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}benzonitrile	464.4 (2.54)
185		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-2-methoxyphenyl sulfamate	549.1 (2.66)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
186		N-(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)sulfamide	518.1 (2.42)
187		6-biphenyl-4-yl-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	499.5 (3.29)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
188		6-(2-thienyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	429.4 (2.97)
189		6-(3,4-dimethoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	483.4 (3.03)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
190		6-(4-methoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	453.4 (3.07)
191		6-(6-methoxypyridin-3-yl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	454.4 (3.00)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
192		6-(3-fluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	441.4 (3.03)
193		6-(2-fluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	441.4 (3.03)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
194		6-(2-aminophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	438.4 (2.96)
195		6-(3-furyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	413.4 (2.94)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
196		6-(4-fluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	441.4 (3.03)
197		6-[3-(trifluoromethyl)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	491.4 (3.18)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
198		3-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}benzonitrile	448.4 (3.00)
199		6-[4-(dimethylamino)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	466.4 (3.11)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
200		6-(3-ethylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	451.4 (3.18)
201		6-(5-isopropyl-2-methoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	495.5 (3.29)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
202		6-(2-methylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	437.4 (3.07)
203		6-(4-methoxypyridin-3-yl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	454.4 (2.67)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
204		6-(2,4-difluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	459.3 (3.03)
205		(3-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)methanol	453.4 (2.92)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
206		6-(3-methoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	453.4 (3.07)
207		6-[4-(ethylthio)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	483.5 (3.18)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
208		6-(3-amino-4-methylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	452.4 (3.03)
209		6-(4-isopropylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	465.5 (3.22)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
210		(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)acetonitrile	462.4 (3.00)
211		6-(4-methylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	437.4 (3.11)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
212		6-(3-methylphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	437.4 (3.11)
213		6-pyridin-3-yl-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	424.4 (2.81)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
214		6-(3-thienyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	429.4 (3.00)
215		6-(2,5-dimethoxyphenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	483.4 (3.07)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
216		6-[3-(trifluoromethoxy)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	507.4 (3.22)
217		6-[4-(trifluoromethoxy)phenyl]-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	507.4 (3.22)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
218		6-(3,5-difluorophenyl)-N4*-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine	459.4 (3.07)
219	 ClH	4-(4-{[2-amino-6-(1H-pyrrol-2-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile hydrochloride	370.3 (2.65)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
220		4-(4-{[2-amino-6-(4-butylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	437.5 (3.47)
221		4-(4-{[2-amino-6-(2-fluorobiphenyl-4-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	475.5 (3.36)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
222		4-[4-({2-amino-6-[3-(benzyloxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	505.1 (3.29)
223		4-(4-{[2-amino-6-(6-methoxypyridin-3-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	412.4 (2.92)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
224		4-(4-{[2-amino-6-(3,4-dimethoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	441.4 (2.92)
225		4-(4-{[2-amino-6-(4-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile 4-(4-{[6-amino-2-(4-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	411.4 (2.96)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
226		4-(4-{[2-amino-6-(3-hydroxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile 4-(4-{[6-amino-2-(3-hydroxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	397.4 (3.07)
227		4-(4-{[2-amino-6-(4-hydroxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile 4-(4-{[6-amino-2-(4-hydroxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	397.4 (2.89)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
228		4-(4-{[2-amino-6-(3-furyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile 4-(4-{[6-amino-2-(3-furyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	371.4 (2.89)
229		4-(4-{[2-amino-6-(4-aminophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile 4-(4-{[6-amino-2-(4-aminophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	396.4 (2.89)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
230		4-(4-{[2-amino-6-(6-methoxypyridin-3-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	430.4 (2.74)
231		4-[4-({2-amino-6-[4-(methylthio)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	427.5 (3.00)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
232		4-[4-({2-amino-6-[4-(dimethylamino)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	424.4 (3.00)
233		4-(4-{[2-amino-6-(2,5-dimethoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	441.4 (3.00)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
234		4-(4-{[2-amino-6-(3-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	411.4 (2.92)
235		4-[4-({2-amino-6-[4-(cyanomethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	420.4 (2.89)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
236		4-{4-[(2-amino-6-pyridin-3-ylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carbonitrile	382.4 (2.78)
237		4-{4-[(2-amino-2',4'-dimethoxy-4,5'-bipyrimidin-6-yl)amino]phenoxy}pyridine-2-carbonitrile	443.4 (2.85)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
238		4-(4-{[2-amino-6-(3-aminophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	396.4 (3.00)
239		4-(4-{[2-amino-6-(2-thienyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	387.4 (3.07)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
240		4-(4-{[2-amino-6-(3-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	399.4 (3.14)
241		4-(4-{[2-amino-6-(2-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	399.4 (3.14)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
242		4-(4-{[2-amino-6-(4-chlorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	415.8 (3.14)
243		4-(4-{[2-amino-6-(2-aminophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	396.4 (3.11)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
244		4-[4-({2-amino-6-[3-(trifluoromethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	449.4 (3.29)
245		4-(4-{[2-amino-6-(5-isopropyl-2-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	453.5 (3.36)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
246		4-(4-{[2-amino-6-(2-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	395.4 (3.14)
247		4-(4-{[2-amino-6-(2,4-difluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	417.4 (3.11)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
248		4-[4-({2-amino-6-[3-(hydroxymethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	411.4 (3.03)
249		4-(4-{[2-amino-6-(4-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	395.4 (3.18)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
250		4-(4-{[2-amino-6-(3-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	395.4 (3.22)
251		4-(4-{[2-amino-6-(3-thienyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	387.4 (3.07)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
252		4-(4-{[2-amino-6-(3-amino-4-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	410.5 (3.03)
253		4-(4-{[2-amino-6-(3,4-dichlorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	450 (3.33)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
254		4-(4-{[2-amino-6-(3-aminophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	414.2 (2.85)
255		4-(4-{[2-amino-6-(2-thienyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	405.1 (3.00)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
256		4-(4-{[2-amino-6-(3-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	417.2 (2.96)
257		4-(4-{[2-amino-6-(4-chlorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	433.1 (3.07)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
258		4-[4-({2-amino-6-[3-(trifluoromethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	467.1 (3.11)
259		4-(4-{[2-amino-6-(4-butylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	455.2 (3.25)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
260		4-(4-{[2-amino-6-(2-fluorobiphenyl-4-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	493.1 (3.29)
261		4-(4-{[2-amino-6-(3-ethylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	427.2 (3.11)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
262		4-(4-{[2-amino-6-(5-isopropyl-2-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	471.2 (3.25)
263		4-(4-{[2-amino-6-(2-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	413.2 (3.00)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
264		4-(4-{[2-amino-6-(2,4-difluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	435.1 (3.00)
265		4-[4-({2-amino-6-[3-(hydroxymethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	429.2 (2.89)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
266		4-(4-{[2-amino-6-(4-isopropylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	441.2 (3.18)
267		4-(4-{[2-amino-6-(4-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	413.2 (3.07)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
268		4-(4-{[2-amino-6-(3-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	413.2 (3.03)
269		4-(4-{[2-amino-6-(3-thienyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	405.1 (2.96)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
270		4-(4-{[2-amino-6-(3-isopropylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	441.2 (3.22)
271		4-(4-{[2-amino-6-(3-amino-4-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	428.2 (2.89)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
272		4-(4-{[2-amino-6-(3,4-dichlorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	467.1 (3.18)
273		4-(4-{[2-amino-6-(3-ethoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	443.4 (2.81)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
274		4-[4-({2-amino-6-[3-(trifluoromethoxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	465.1 (3.00)
275		4-[4-({2-amino-6-[2-(trifluoromethoxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carbonitrile	465.1 (2.92)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
276		4-(4-{[2-amino-6-(3,5-difluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carbonitrile	417.2 (2.89)
277		4-[4-({2-amino-6-[3-(trifluoromethoxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	483.4 (2.89)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
278		4-[4-({2-amino-6-[2-(trifluoromethoxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	483.4 (2.81)
279		4-(4-{[2-amino-6-(3,5-difluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	435.4 (2.74)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
280		4-{3-[(2-amino-6-biphenyl-4-ylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxamide	475.5 (3.00)
281		4-[3-({2-amino-6-[3-(benzyloxy)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	505.5 (3.07)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
282		6-(4-fluorophenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	388.4 (2.54)
283		4-(3-{[2-amino-6-(3-aminophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	414.4 (2.64)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
284		4-(3-{[2-amino-6-(6-methoxypyridin-3-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	430.4 (2.70)
285		4-(3-{[2-amino-6-(3-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	417.4 (2.77)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
286		4-(3-{[2-amino-6-(2-fluorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	417.4 (2.74)
287		4-[3-({2-amino-6-[4-(dimethylamino)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	442.5 (2.85)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
288		4-(3-{[2-amino-6-(3-ethylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	427.5 (2.88)
289		4-(3-{[2-amino-6-(5-isopropyl-2-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	471.5 (3.00)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
290		4-(3-{[2-amino-6-(2-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	413.5 (2.79)
291		4-(3-{[2-amino-6-(2-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	429.5 (2.78)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
292		4-(3-{[2-amino-6-(3-methoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	429.5 (2.78)
293		4-(3-{[2-amino-6-(4-isopropylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	441.5 (2.97)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
294		4-[3-({2-amino-6-[4-(cyanomethyl)phenyl]pyrimidin-4-yl}amino)phenoxy]pyridine-2-carboxamide	438.5 (2.75)
295		4-(3-{[2-amino-6-(4-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	413.5 (2.81)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
296		4-(3-{[2-amino-6-(3-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	413.5 (2.80)
297		4-{3-[(2-amino-6-pyridin-3-ylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxamide	400.4 (2.58)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
298		4-(3-{[2-amino-6-(3-thienyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	405.5 (2.72)
299		4-(3-{[2-amino-6-(3-isopropylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	441.5 (2.95)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
300		4-(3-{[2-amino-6-(2,5-dimethoxyphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	459.5 (2.80)
301		4-(3-{[2-amino-6-(1H-pyrrol-2-yl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	388.401 (2.69)

Table 1



Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
302		4-(3-{[2-amino-6-(3-amino-4-methylphenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	428.5 (2.71)
303		4-(3-{[2-amino-6-(3,4-dichlorophenyl)pyrimidin-4-yl]amino}phenoxy)pyridine-2-carboxamide	468.3 (2.92)

Table 1





Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
304		N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-(4-phenoxyphenyl)pyrimidine-2,4-diamine	462.5 (2.71)
305		6-[3-(dimethylamino)phenyl]-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	413.5 (2.54)
306		3-[2-amino-6-({4-[(2-methylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenol	386.4 (2.48)
307		N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-[4-(trifluoromethoxy)phenyl]pyrimidine-2,4-diamine	454.4 (2.71)

Table 1





Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
308		N-{4-[2-amino-6-({4-[(2-methylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl}acetamide	427.5 (2.47)
309		N4-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-[2-(trifluoromethoxy)phenyl]pyrimidine-2,4-diamine	454.4 (2.61)
310		4-[2-amino-6-({4-[(2-methylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenol	386.4 (2.49)
311		6-(2-methylphenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	384.5 (2.54)

Table 1





Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
312		6-(6-methoxypyridin-3-yl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	401.4 (2.50)
313		6-(4-aminophenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	385.4 (2.47)
314		N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-[3-(trifluoromethyl)phenyl]pyrimidine-2,4-diamine	438.4 (2.66)
315		6-(3-fluorophenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	388.4 (2.55)

Table 1





Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
316		6-(3-methoxyphenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	400.5 (2.57)
317		6-(4-ethylphenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	398.5 (2.66)
318		6-[4-(dimethylamino)phenyl]-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	413.5 (2.63)
319		6-(3,4-dimethoxyphenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	430.5 (2.55)

Table 1





Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
320		6-(3-aminophenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	385.4 (2.41)
321		6-(3-furyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	360.4 (2.49)
322		6-(2-aminophenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	385.4 (2.49)
323		6-(3-amino-4-methylphenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	399.5 (2.48)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
324		6-(3-isopropoxyphenyl)-N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}pyrimidine-2,4-diamine	428.5 (2.68)
325		N4*-{4-[(2-methylpyridin-4-yl)oxy]phenyl}-6-[3-(trifluoromethoxy)phenyl]pyrimidine-2,4-diamine	454.4 (2.69)
326		4-[2-amino-6-({4-[(2-cyanopyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl sulfamate	

Table 1




Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
327		3-[2-amino-6-({4-[(2-methylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl sulfamate	
328		3-[2-amino-6-({4-[(2-cyanopyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl sulfamate	
329		4-[2-amino-6-({4-[(2-methylpyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl sulfamate	

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
330		3-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl sulfamate	
331		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}-3-methylphenyl sulfamate	
332		4-[2-amino-6-({4-[(2-chloropyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl sulfamate	485.3 (2.42)

Table 1

Example No.	Structure	IUPAC Name*	LCMS: $[M+H]^+$ (RT, min)
333		4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl dimethylsulfamate	
334		3-[2-amino-6-({4-[(2-chloropyridin-4-yl)oxy]phenyl}amino)pyrimidin-4-yl]phenyl sulfamate	

Cytotoxic Activity of the Invention Compounds

The following section describes an assay that can be used to characterize compounds of the invention, e.g., to test for the cytotoxic activity of compounds on cells.

Human tumor cells, e.g., HCT116 cells, are seeded in a 96-well plate at $3.0x10^3$ cells/well and grown in 100 µl of RPMI complete media (Invitrogen Corporation, Grand Island, NY) containing 10% fetal bovine serum (Hyclone, Logan, Utah) and 10 mM HEPES and at 37 °C for 16 h in an incubator with 5% CO_2. To each well, 50 µl of additional growth media containing 20 µM to 60 nM concentrations of compound with 0.2% DMSO is added. Cells are grown for another 72 h at 37 °C. 20 µl of Alamar Blue (Trek Diagnostic Systems, Inc., Cleveland, Ohio) reagent is added to each well and incubated for 4 h at 37 °C. Plates are read in a SpectraMax Gemini (Molecular Devices, CA) with 544 nm excitation and 590 nm emission wavelength. IC_{50} values are determined by linear regression analysis of log drug concentration versus percent inhibition.

Representative compounds of this invention were tested for cytotoxicity using the above-described assay procedure with the following results:

Examples 1, 2, 3, 4, 6, 7, 8, 9, 11, 13, 19, 21, 33, 34, 35, 37, 38, 39, 40, 45, 46, 48, 49, 50, 51, 52, 53, 58, 62, 63, 64, 66, 67, 69, 76, 80, 84, 88, 92, 94, 95, 96, 97, 98, 99, 113, 114, 116, 117, 118, 119, 120, 121, 123, 124, 125, 127, 129, 133, 134, 137, 139, 140, 142, 143, 145, 146, 149, 150, 151, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 173, 174, 175, 177, 179, 180, 181, 183, 185, 187, 188, 190, 191, 192, 193, 195, 196, 199, 200, 202, 204, 205, 206, 208, 209, 210, 211, 212, 213, 214, 215, 218, 219, 223, 227, 228, 229, 233, 234, 238, 240, 241, 246, 249, 250, 251, 252, 254, 259, 261, 262, 263, 264, 266, 267, 268, 269, 271, 273, 275, 278, 282, 288, 290, 292, 296, 298, 302, 305, 310, 311, 312, 313, 315, 316, 317, 318, 320, 321, 323 and 332 show an IC_{50} of less than or equal to 500 nM in the HCT116 cytotoxic activity assay:


Examples 5, 10, 12, 23, 26, 30, 32, 41, 42, 47, 60, 61, 65, 68, 70, 73, 75, 79, 82, 85, 86, 89, 90, 91, 100, 101, 105, 107, 109, 112, 115, 128, 130, 131, 132, 135, 136, 138, 141, 144, 148, 152, 172, 176, 189, 194, 197, 198, 201, 203, 207, 220, 222, 226, 230, 231, 232, 235, 239, 242, 243, 245, 247, 248, 256, 257, 258, 276, 279, 281, 283, 284, 286, 291, 294, 297, 300, 304, 306, 308, 309, 319, 322, 324, and 325 show an IC_{50} greater than 500 nM but less than or equal to 2 µM in the HCT116 cytotoxic activity assay.

Examples, 43, 44, 71, 72, 74, 77, 81, 83, 87, 93, 106, 108, 110, 111, 122, 126, 147, 178, 216, 217, 221, 224, 225, 236, 237, 244, 253, 255, 260, 265, 270, 272, 274, 277, 280, 285, 287, 289, 293, 295, 299, 301, 303, 307, and 314 show an IC_{50} greater than 2 μM in the HCT116 cytotoxic activity assay.

Other embodiments of the invention will be apparent to the skilled in the art from a consideration of this specification or practice of the invention disclosed herein. It is intended that the specification and examples be considered as exemplary only, with the true scope and spirit of the invention being indicated by the following claims.

WE CLAIM:

1. A compound having the structure



(I)

wherein

R^1 represents H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^2 represents $(C_1\text{-}C_3)$alkyl, cyclopropyl, $O(C_1\text{-}C_3)$alkyl, or NR^3R^4

wherein R^3 and R^4 are H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^{2a} represents H or halogen;


M represents CH or N;

L represents a carbonyl group, O, NR^5 , CR^6R^7 , or $(C_2\text{-}C_3)$alkylenyl which is optionally substituted up to twice by groups independently selected from halogen and OH; wherein

R^5 is H or $(C_1\text{-}C_3)$alkyl; and


R^6 and R^7 are independently H, CH_3, halogen, or OH;

J represents an aromatic or heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

wherein R^8 represents H or $(C_1\text{-}C_3)$alkyl;

G" represents a substituent selected from the group consisting of $(C_1$-$C_3)$alkyl, cyclopropyl, $O(C_1$-$C_3)$alkyl, halogen, CF_3, CN and CO_2R^9;

wherein

R^9 represents H or $(C_1$-$C_3)$alkyl; and

m represents the number of substituents G", and is 0, 1, or 2;

G represents a substituent located on ring J;

G' represents a substituent located on ring Y;

n represents the number of substituents G; and

n' represents the number of substituents G';

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,
2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G11, to a maximum total of 3 substituents on rings J and Y, and
3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12-G37;

and subject to the further provisos

4) when J is phenyl, G is other than OH or alkylthio; and when J is phenyl or pyridyl, n is 1, 2, or 3;
5) when J is phenyl, and G is G4 shown below, then R^2 is NR^3R^4;

G and G' moieties are independently selected from the group consisting of:

G1) halogen;

G2) $O(C_1$-$C_4)$alkyl which optionally is substituted up to two times by $O(C_1$-$C_2)$alkyl;

G3) OH;

G4) $(C_1$-$C_5)$alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3;

G6) $NHC(O)(C_1\text{-}C_3)$alkyl;

G7) $NHSO_2(C_1\text{-}C_3)$alkyl;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,

$NR^{12}R^{13}$ wherein


R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula



wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,


or by

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G9) $(CH_2)_a\text{-}NR^{19}R^{20}$ wherein

R^{19} and R^{20} are independently H, $(C_1\text{-}C_5)$alkyl, or $(C_3\text{-}C_6)$cycloalkyl, or may be joined to form a saturated 5-6-membered N-containing ring; and

the subscript "a" is an integer of 1-4;

G10)  wherein

Q' is O or NR^{21} ;

R^{21} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl; and

the subscript "b" is an integer of 1-3;

G11) $CH_2NR^{22}(CH_2)_cOCH_3$ wherein

R^{22} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl; and

the subscript "c" is an integer of 2-4;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G13) CN ;

G14) NO_2 ;

G15) cyclopropyl ;


G16) OR^{27}, wherein


R^{27} represents phenyl or benzyl;

G17) $S(C_1\text{-}C_3)alkyl$;

G18) $CH{=}CH\text{-}(CH_2)_{1\text{-}3}\text{-}OR^5$; wherein

R^5 represents H or $(C_1\text{-}C_3)alkyl$;

G19)  ;


G20)  ;

G21) $C(O)NR^{28}R^{29}$, wherein

R^{28} and R^{29} are independently selected from

H,

cyclopropyl, provided that both R^{28} and R^{29} are not simultaneously cyclopropyl,


, provided that this group does not constitute both R^{28} and R^{29} simultaneously,

and

$(C_1\text{-}C_3)alkyl$ which is optionally substituted up to two times by OH;

or

R^{28} and R^{29} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by OH, or by $(C_1\text{-}C_3)alkyl$ which in turn is optionally substituted up to two times by OH or $O(C_1\text{-}C_3)alkyl$;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is

H,


cyclopropyl, or

$(C_1\text{-}C_3)$alkyl which is optionally substituted once by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G23) $O\text{-}(CH_2)_d\text{-}NR^{31}R^{32}$ wherein

R^{31} and R^{32} are independently H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl, or may be joined to form a saturated 5-6-membered N-containing ring; and


the subscript "d" is an integer of 2-4;

G24)  wherein

the subscript "e" is an integer of 2-3; and

Q''' is O or NR^{33} ; and

R^{33} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

G25)  wherein

Q^{iv} is O or NR^{34} ; and

R^{34} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

G26) $C(O)NR^{35}(CH_2)_fOR^{36}$ wherein

R^{35} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^{36} is $(C_1\text{-}C_6)$alkyl optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl, and

the subscript "f" is an integer of 2-4;

G27) CO_2R^{37} wherein

R^{37} is H or $(C_1\text{-}C_3)$alkyl;

G28) phenyl, which is optionally substituted by up to 2 groups selected from halogen, $(C_1\text{-}C_3)$alkyl, OR^{38}, CN, CF_3, and $NR^{39}R^{40}$
wherein
R^{38} represents H or $(C_1\text{-}C_3)$alkyl; and
$R^{39}$and R^{40} represent H or $(C_1\text{-}C_3)$alkyl;

G29) $NR^{41}SO_2NR^{42}R^{43}$ wherein
R^{41}represents H, or $(C_1\text{-}C_4)$alkyl, and
R^{42} and R^{43} independently represent H, CH_3, or $(C_2\text{-}C_3)$alkyl which may optionally be substituted once by -OH or $NR^{44}R^{45}$, wherein
R^{44} and R^{45} independently represent H or $(C_1\text{-}C_3)$alkyl;

G30) $OC(O)\text{-}CH_2\text{-}NR^{46}R^{47}$ wherein
R^{46} and R^{47} independently represent H, $(C_1\text{-}C_3)$alkyl, or CO_2(t-butyl), provided that R^{46} and R^{47} are not both simultaneously CO_2(t-butyl);

G31) $N(R^{48})C(O)R^{49}$ wherein
R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and
R^{49} represents
$(CH_2)_{1\text{-}3}\text{-}CO_2H$,
$O(C_2\text{-}C_4)$alkyl,
$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein
R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or
$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein
R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and
R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl;

G32) $C(O)$-$(C_1$-$C_3)$alkyl;

G33) $(CH_2)_g$-$N(R^{55})$-$C(O)$-R^{56} wherein

g represents 1, 2, or 3;


R^{55} represents H or $(C_1$-$C_3)$alkyl;

R^{56} represents

$(C_1$-$C_3)$alkyl optionally substituted up to two times by OR^{57} or $NR^{58}R^{59}$, wherein

R^{57} represents H or $(C_1$-$C_3)$alkyl, and

R^{58} and R^{59} each represents H or $(C_1$-$C_3)$alkyl,

or R^{56} represents  wherein

R^{60} represents halogen, $(C_1$-$C_3)$alkyl, $O(C_1$-$C_3)$alkyl, CN, OH, CF_3, or $NR^{61}R^{62}$, wherein

R^{61} and R^{62} represent H or $(C_1$-$C_3)$alkyl; and

h represents 0, 1, or 2;


G34) $(CH_2)_i$-$N(R^{63})$-$C(O)$-$NR^{64}R^{65}$ wherein

i represents 1, 2, or 3;

R^{63} represents H or $(C_1$-$C_3)$alkyl;


R^{64} and R^{65} each represents H or $(C_1$-$C_3)$alkyl;

or

R^{64} and R^{65} may be joined to form  wherein

Q^V represents CH_2, O or NR^{66} wherein

R^{66} represents H or $(C_1$-$C_3)$alkyl;

G35)  wherein

j represents 1, 2, or 3;

R^{67}represents H or $(C_1$-$C_3)$alkyl; and

R^{68} represents H or $(C_1\text{-}C_3)$alkyl;

G36) $(CH_2)_k\text{-}N(R^{69})\text{-}SO_2\text{-}R^{70}$ wherein

k represents 1, 2, or 3;

R^{69} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{70} represents $(C_1\text{-}C_4)$alkyl, or phenyl which is optionally substituted up to perhalo by halogen or up to three times by OR^{71}, CN, CF_3, or $NR^{72}R^{73}$, wherein

R^{71} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{72} and R^{73} each represents H or $(C_1\text{-}C_3)$alkyl;

G37) $CH{=}CH\text{-}(CH_2)_{1\text{-}3}\text{-}NR^{74}R^{75}$ wherein

R^{74} and R^{75} represent H or $(C_1\text{-}C_3)$alkyl;

or a pharmaceutically acceptable salt, solvate, solvate of a salt, or stereoisomer thereof.


2. The compound of claim 1

wherein

R^1 represents H;


M represents CH;

J represents a heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,

2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G5 and G8, to a maximum total of 3 substituents on rings J and Y, and

3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12, G13, G22, G29, and G31;

and subject to the further proviso

4) when J is pyridyl, n is 1, 2, or 3;

and proviso 5 does not apply;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1-C_4)$alkyl which optionally is substituted up to two times by $O(C_1-C_2)$alkyl;

G3) OH ;

G4) (C_1-C_5)alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,

$NR^{12}R^{13}$ wherein


R^{12} and R^{13} are independently H or (C_1-C_3)alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

(C_2-C_4)alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups

independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,

or by


$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G13) CN;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is

H,

cyclopropyl, or

(C_1-C_3)alkyl which is optionally substituted once by halogen, OH, or $O(C_1-C_3)$alkyl;

G29) $NR^{41}SO_2NR^{42}R^{43}$ wherein

R^{41}represents H, or (C_1-C_4)alkyl, and

R^{42} and R^{43} independently represent H, CH_3, or (C_2-C_3)alkyl which may optionally be substituted once by -OH or $NR^{44}R^{45}$, wherein

R^{44} and R^{45} independently represent H or (C_1-C_3)alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or (C_1-C_3)alkyl; and

R^{49} represents

$(CH_2)_{1-3}-CO_2H$,

$O(C_2-C_4)$alkyl,

$(CH_2)_{1-4}-NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or (C_1-C_3)alkyl, or

$CH(R^{52})-NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1-4}-NH_2$, CH_2OH, $CH(CH_3)OH$, or (C_1-C_3)alkyl; and

R^{53} and R^{54} independently represent H or (C_1-C_3)alkyl.

3. The compound of claim 2

wherein

R^1 represents H;

R^2 represents $O(C_1-C_3)$alkyl or NR^3R^4

wherein R^3 and R^4 are H or (C_1-C_3)alkyl;

R^{2a} represents H;

L represents O or CR^6R^7 wherein

R^6 and R^7 are independently H, CH_3 , or OH;

G" represents a substituent selected from the group consisting of $O(C_1-C_3)$alkyl, halogen, and CF_3;

n and n' are independently 0 or 1, and provisos 1-3 do not apply;

G and G' moieties are independently selected from the group consisting of:

G1) Cl or F;

G2) $O(C_1-C_3)$alkyl;

G3) OH ;

G4) (C_1-C_3)alkyl, which is optionally substituted up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or (C_1-C_3)alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

(C_2-C_4)alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1-C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or (C_1-C_3)alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and


R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G13) CN ;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is H or $(C_1\text{-}C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and


R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

4. The compound of claim 1 wherein

R^1 represents H;


M represents CH;

J represents a heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,
2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G5 and G8, to a maximum total of 3 substituents on rings J and Y, and
3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12, G21, G25, G26, and G31;

and subject to the further proviso

4) when J is pyridyl, n is 1, 2, or 3;

and proviso 5 does not apply;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1\text{-}C_4)$alkyl which optionally is substituted up to two times by $O(C_1\text{-}C_2)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_5)$alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,

or by

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein


R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G21) $C(O)NR^{28}R^{29}$, wherein

R^{28} and R^{29} are independently selected from

H,

cyclopropyl, provided that both R^{28} and R^{29} are not simultaneously cyclopropyl,




, provided that this group does not constitute both R^{28} and R^{29} simultaneously,

and

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by OH;

or

R^{28} and R^{29} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by OH, or by $(C_1\text{-}C_3)$alkyl which in turn is optionally substituted up to two times by OH or $O(C_1\text{-}C_3)$alkyl;

G25)  wherein

Q^{iv} is O or NR^{34}; and

R^{34} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

G26) $C(O)NR^{35}(CH_2)_fOR^{36}$ wherein

R^{35} is H, $(C_1\text{-}C_3)$alkyl, or cyclopropyl;

R^{36} is $(C_1\text{-}C_6)$alkyl optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl, and

the subscript "f" is an integer of 2-4; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl, or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

5. The compound of claim 4

wherein

R^1 represents H;

R^2 represents $O(C_1\text{-}C_3)$alkyl or NR^3R^4

wherein R^3 and R^4 are H or $(C_1\text{-}C_3)$alkyl;

R^{2a} represents H;

L represents O or CR^6R^7, wherein

R^6 and R^7 are independently H, CH_3, or OH;

G" represents a substituent selected from the group consisting of $O(C_1\text{-}C_3)$alkyl, halogen, and CF_3;

n and n' are independently 0 or 1, and provisos 1-3 do not apply;

G and G' moieties are independently selected from the group consisting of:

G1) Cl or F;

G2) $O(C_1\text{-}C_3)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_3)$alkyl, which is optionally substituted up to three times by halogen;

G5) OCF_3 ;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and


R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G21) $C(O)NR^{28}R^{29}$, wherein
R^{28} and R^{29} are independently selected from
H
and
$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by OH;

G25)  wherein
Q^{iv} is O or NR^{34} ; and
R^{34} is H or $(C_1\text{-}C_3)$alkyl;

G26) $C(O)NR^{35}(CH_2)_fOR^{36}$ wherein
R^{35} is H or $(C_1\text{-}C_3)$alkyl;
R^{36} is $(C_1\text{-}C_6)$alkyl optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl, and
the subscript "f" is an integer of 2-4; and

G31) $N(R^{48})C(O)R^{49}$ wherein
R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and
R^{49} represents
$(CH_2)_{1\text{-}3}\text{-}CO_2H$,
$O(C_2\text{-}C_4)$alkyl,
$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein
R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl,
or
$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein
R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and
R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.


6. The compound of claim 1

wherein

R^1 represents H;


M represents CH;

J represents an aromatic or heteroaromatic ring selected from the group consisting of



;

Y represents an aromatic or heteroaromatic ring selected from the group consisting of



;

n and n' are independently 0, 1, 2, or 3, subject to the provisos that

1) ring J and ring Y each may be substituted independently up to 3 times by substituents listed below as numbers G1-G2, to a maximum total of 4 substituents on rings J and Y,
2) ring J and ring Y each may be substituted independently up to 2 times by substituents listed below as numbers G3-G5 and G8, to a maximum total of 3 substituents on rings J and Y, and
3) ring J and ring Y each may be substituted independently once by a substituent selected from those listed below as numbers G12, G22, and G31;

and subject to the further proviso

4) when J is pyridyl, n is 1, 2, or 3;

and proviso 5 does not apply;

G and G' moieties are independently selected from the group consisting of:

G1) halogen ;

G2) $O(C_1\text{-}C_4)$alkyl which optionally is substituted up to two times by $O(C_1\text{-}C_2)$alkyl;

G3) OH ;

G4) $(C_1\text{-}C_5)$alkyl, which is optionally substituted independently up to two times by groups selected from hydroxyl and cyano, or up to three times by halogen;

G5) OCF_3;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2\text{-}C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1\text{-}C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1\text{-}C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein

Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1\text{-}C_3)$alkyl,

or

R^{10} and R^{11} may be joined to form a saturated 5-6-membered N-containing ring which is optionally substituted up to two times by

OH,

$NR^{17}R^{18}$, wherein

R^{17} and R^{18} are H or $(C_1\text{-}C_3)$alkyl,


or by

$(C_1\text{-}C_3)$alkyl which is optionally substituted up to two times by halogen, OH, or $O(C_1\text{-}C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2\text{-}C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1\text{-}C_3)$alkyl;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is

H,

cyclopropyl, or

$(C_1\text{-}C_3)$alkyl which is optionally substituted once by halogen, OH, or $O(C_1\text{-}C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1\text{-}C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}\text{-}CO_2H$,

$O(C_2\text{-}C_4)$alkyl,

$(CH_2)_{1\text{-}4}\text{-}NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1\text{-}C_3)$alkyl,

or

$CH(R^{52})\text{-}NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1\text{-}4}\text{-}NH_2$, CH_2OH, $CH(CH_3)OH$, or $(C_1\text{-}C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1\text{-}C_3)$alkyl.

7. The compound of claim 6

wherein

R^1 represents H;

R^2 represents $O(C_1\text{-}C_3)$alkyl, or NR^3R^4

wherein R^3 and R^4 are H or $(C_1\text{-}C_3)$alkyl;

R^{2a} represents H;

L represents O or CR^6R^7, wherein

R^6 and R^7 are independently H, CH_3, or OH;

G" represents a substituent selected from the group consisting of $O(C_1\text{-}C_3)$alkyl, halogen, and CF_3;

n and n' are independently 0 or 1, and provisos 1-3 do not apply;

G and G' moieties are independently selected from the group consisting of:

G1) Cl or F;

G2) $O(C_1\text{-}C_3)$alkyl;

G3) OH;

G4) $(C_1\text{-}C_3)$alkyl, which is optionally substituted up to three times by halogen;

G5) OCF_3;

G8) $NR^{10}R^{11}$, wherein

R^{10} and R^{11} are independently selected from

H,

CH_3,

cyclopropyl,

benzyl,


$NR^{12}R^{13}$ wherein

R^{12} and R^{13} are independently H or $(C_1\text{-}C_3)$alkyl, provided that both R^{10} and R^{11} are not $NR^{12}R^{13}$ simultaneously,

and

$(C_2$-$C_4)$alkyl which is optionally substituted up to three times by halogen, and up to two times by substituent groups independently selected from hydroxyl, $O(C_1$-$C_3)$alkyl, and $NR^{14}R^{15}$, wherein

R^{14} and R^{15} are independently H or $(C_1$-$C_3)$alkyl, or

R^{14} and R^{15} can join to form a heterocycle of formula  wherein


Q represents CH_2, O, or NR^{16}, and

R^{16}represents H or $(C_1$-$C_3)$alkyl;

G12) $OSO_2NR^{23}R^{24}$ wherein

R^{23} and R^{24} independently represent H, CH_3, or $(C_2$-$C_4)$alkyl which may optionally be substituted once by OH or $NR^{25}R^{26}$, wherein

R^{25} and R^{26} independently represent H or $(C_1$-$C_3)$alkyl;

G22)  wherein

Q'' is O or NR^{30}, and

R^{30} is H or $(C_1$-$C_3)$alkyl; and

G31) $N(R^{48})C(O)R^{49}$ wherein

R^{48} represents H or $(C_1$-$C_3)$alkyl; and

R^{49} represents

$(CH_2)_{1\text{-}3}$-CO_2H,

$O(C_2$-$C_4)$alkyl,

$(CH_2)_{1\text{-}4}$-$NR^{50}R^{51}$ wherein

R^{50} and R^{51} independently represent H or $(C_1$-$C_3)$alkyl, or

$CH(R^{52})$-$NR^{53}R^{54}$ wherein

R^{52} represents $(CH_2)_{1-4}$-NH_2, CH_2OH, $CH(CH_3)OH$, or $(C_1$-$C_3)$alkyl; and

R^{53} and R^{54} independently represent H or $(C_1$-$C_3)$alkyl.

8. A compound selected from the group consisting of

4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}-*N*-methylpyridine-2-carboxamide;

4-{3-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carboxamide;

4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridine-2-carbonitrile;

6-phenyl-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine;

N^4-{4-[(2-chloropyridin-4-yl)oxy]phenyl}-6-phenylpyrimidine-2,4-diamine;

4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl sulfamate;

N-(4-{2-amino-6-[(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)amino]pyrimidin-4-yl}phenyl)glycinamide trifluoroacetate;

6-(4-aminophenyl)-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine;

6-(6-aminopyridin-3-yl)-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine;

6-pyridin-3-yl-N^4-(4-{[2-(trifluoromethyl)pyridin-4-yl]oxy}phenyl)pyrimidine-2,4-diamine;

N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]-4-methoxybenzenesulfonamide trifluoroacetate;

N-[(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methyl]methanesulfonamide trifluoroacetate;

and

(4-{4-[(2-amino-6-phenylpyrimidin-4-yl)amino]phenoxy}pyridin-2-yl)methanol trifluoroacetate (salt).

9. A pharmaceutical composition comprising a compound of claim 1 and a pharmaceutically acceptable carrier.

10. A method of treatment for a hyperproliferative disorder comprising administering an effective amount of a compound of claim 1 to a subject in need thereof.

11. The method of claim 10 wherein said hyperproliferative disorder is cancer.