




(86) Date de dépôt PCT/PCT Filing Date: 2007/01/31
(87) Date publication PCT/PCT Publication Date: 2007/08/09
(85) Entrée phase nationale/National Entry: 2008/07/23
(86) N° demande PCT/PCT Application No.: US 2007/002790
(87) N° publication PCT/PCT Publication No.: 2007/089904
(30) Priorité/Priority: 2006/01/31 (US60/763,782)

(51) Cl.Int./Int.Cl. *C07D 213/44* (2006.01),
A61K 31/44 (2006.01)
(71) Demandeur/Applicant:
SYNTA PHARMACEUTICALS CORP., US
(72) Inventeurs/Inventors:
JIANG, JUN, US;
XIA, ZHI-QIANG, US;
ZHANG, JUNYI, US;
BOHNERT, GARY, US;
CHEN, SHOUJUN, US;
XIE, YU, US
(74) Agent: BERESKIN & PARR

(54) Titre : COMPOSES DE PYRIDYLPHENYL DESTINES A TRAITER UNE INFLAMMATION ET UTILISATIONS LIEES
A L'IMMUNITE

(54) Title: PYRIDYLPHENYL COMPOUNDS FOR INFLAMMATION AND IMMUNE-RELATED USES



(57) Abrégé/Abstract:

The invention relates to compounds of structural formula (I): or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein Y, L, X₁, X₂, Z, R₃, R₄, and n are defined herein. These compounds are useful as immunosuppressive agents and for treating and preventing inflammatory conditions, allergic disorders, and immune disorders.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 August 2007 (09.08.2007)

PCT

(10) International Publication Number
WO 2007/089904 A3

(51) International Patent Classification:

C07D 213/44 (2006.01) A61K 31/44 (2006.01)

(21) International Application Number:

PCT/US2007/002790

(22) International Filing Date: 31 January 2007 (31.01.2007)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

60/763,782 31 January 2006 (31.01.2006) US

(71) Applicant (for all designated States except US): **SYNTA PHARMACEUTICALS CORP.** [US/US]; 45 Hartwell Ave., Lexington, MA 02421 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JIANG, Jun** [CN/US]; 211 Central Street, A202, Norwood, MA 02062 (US). **XIA, Zhi-Qiang** [CN/US]; 634 Massachusetts Avenue, Acton, MA 01720 (US). **ZHANG, Junyi** [CN/US]; 130 Bowden Street, 304, Lowell, MA 01852 (US). **BOHNERT, Gary** [US/US]; 379A Cardinal Medeiros, #4, Cambridge, MA 02141 (US). **CHEN, Shoujun** [US/US]; 19 Dunster Road, Bedford, MA 01730 (US). **XIE, Yu** [US/US]; 28 Algonquian Drive, Natick, MA 01760 (US).

(74) Agent: **HSI, Jeffrey D.**; Edwards Angell Palmer & Dodge LLP, P.o. Box 55874, Boston, MA 02205 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:

3 January 2008

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: PYRIDYLPHENYL COMPOUNDS FOR INFLAMMATION AND IMMUNE-RELATED USES



(I)

(57) Abstract: The invention relates to compounds of structural formula (I): or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein Y, L, X₁, X₂, Z, R₃, R₄, and n are defined herein. These compounds are useful as immunosuppressive agents and for treating and preventing inflammatory conditions, allergic disorders, and immune disorders.

WO 2007/089904 A3

IL-2, although useful in the immune response, can cause a variety of problems. IL-2 damages the blood-brain barrier and the endothelium of brain vessels. These effects may be the underlying causes of neuropsychiatric side effects observed under IL-2 therapy, e.g. fatigue, disorientation and depression. It
5 also alters the electrophysiological behaviour of neurons.

Due to its effects on both T and B cells, IL-2 is a major central regulator of immune responses. It plays a role in inflammatory reactions, tumour surveillance, and hematopoiesis. It also affects the production of other
10 cytokines, inducing IL-1, TNF- α and TNF- β secretion, as well as stimulating the synthesis of IFN- γ in peripheral leukocytes.

T cells that are unable to produce IL-2 become inactive (anergic). This renders them potentially inert to any antigenic stimulation they might receive in the
15 future. As a result, agents which inhibit IL-2 production can be used for immunosuppression or to treat or prevent inflammation and immune disorders. This approach has been clinically validated with immunosuppressive drugs such as cyclosporin, FK506, and RS61443. Despite this proof of concept, agents that inhibit IL-2 production remain far from ideal. Among other
20 problems, efficacy limitations and unwanted side effects (including dose-dependant nephrotoxicity and hypertension) hinder their use.

Over production of proinflammatory cytokines other than IL-2 has also been implicated in many autoimmune diseases. For example, Interleukin 5 (IL-5), a
25 cytokine that increases the production of eosinophils, is increased in asthma. Overproduction of IL-5 is associated with accumulation of eosinophils in the asthmatic bronchial mucosa, a hall mark of allergic inflammation. Thus, patients with asthma and other inflammatory disorders involving the accumulation of eosinophils would benefit from the development of new drugs
30 that inhibit the production of IL-5.

Interleukin 4 (IL-4) and interleukin 13 (IL-13) have been identified as mediators of the hypercontractility of smooth muscle found in inflammatory bowel disease

and asthma. Thus, patients with asthma and inflammatory bowel disease would benefit from the development of new drugs that inhibit IL-4 and IL-13 production.

- 5 Granulocyte macrophage-colony stimulating factor (GM-CSF) is a regulator of maturation of granulocyte and macrophage lineage population and has been implicated as a key factor in inflammatory and autoimmune diseases. Anti-GM-CSF antibody blockade has been shown to ameliorate autoimmune disease. Thus, development of new drugs that inhibit the production of
10 GM-CSF would be beneficial to patients with an inflammatory or autoimmune disease.

There is therefore a continuing need for new drugs which overcome one or more of the shortcomings of drugs currently used for immunosuppression or in
15 the treatment or prevention of inflammatory disorders, allergic disorders and autoimmune disorders. Desirable properties of new drugs include efficacy against diseases or disorders that are currently untreatable or poorly treatable, new mechanism of action, oral bioavailability and/or reduced side effects.


20

SUMMARY OF THE INVENTION

This invention meets the above-mentioned needs by providing certain pyridylphenyl derivatives that inhibit the activity of CRAC ion channels and inhibit the production of IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , and IFN γ .

- 5 These compounds are particularly useful for immunosuppression and/or to treat or prevent inflammatory conditions and immune disorders.

The invention relates to compounds of formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of $-NRCR_2-$, $-CR_2NR-$, $-C(O)-$, $-NR-C(O)-$, $-C(O)NR-$, $-OC(O)-$, $-C(O)O-$, $-C(S)-$, $-NR-C(S)-$, $-C(S)NR-$;

one of X_1 or X_2 is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$, cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from $-H$ or an alkyl;

R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$, or $-NR_6R_7$;

5 R₄ is a halo, a haloalkyl, cyano, nitro, -C(O)R₅, -C(O)OR₅, -C(O)SR₅, -C(O)NR₉R₁₀, -C(S)R₅, -C(S)OR₅, -C(S)SR₅, -C(S)NR₆R₇, -C(NR₈)R₅, -C(NR₈)OR₅, -C(NR₈)SR₅, -C(NR₈)NR₆R₇, -S(O)_pR₅, -S(O)_pNR₅, -S(O)_pOR₅, -P(O)(OR₅)₂, -OP(O)(OR₅)₂, -P(O)(R₅)₂, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

10 R₅, for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

15 R₆ and R₇, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₆ and R₇ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

20 R₈, for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

25 R₉ and R₁₀, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indoliziny, an

30

optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an
 optionally substituted isothiazolyl, an optionally substituted pyridazinyl,
 an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl,
 an optionally substituted triazinyl, an optionally substituted triazolyl, an
 5 optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an
 optionally substituted pyrazinyl, an optionally substituted quinolinyl, an
 optionally substituted isoquinolinyl, an optionally substituted indazolyl,
 an optionally substituted benzoxazolyl, an optionally substituted
 benzofuryl, an optionally substituted benzothiazolyl, an optionally
 10 substituted indolizinyl, an optionally substituted imidazopyridinyl, an
 optionally substituted tetrazolyl, an optionally substituted
 benzimidazolyl, an optionally substituted benzoxazolyl, an optionally
 substituted benzothiazolyl, an optionally substituted benzothiadiazolyl,
 an optionally substituted benzoxadiazolyl, an optionally substituted
 15 indolyl, an optionally substituted tetrahydroindolyl, an optionally
 substituted azaindolyl, an optionally substituted imidazopyridyl, an
 optionally substituted quinazolinyl, an optionally substituted purinyl, an
 optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted
 pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉
 20 and R₁₀ taken together with the nitrogen to which they are attached are
 an optionally substituted heterocyclyl or optionally substituted
 heteroaryl;
 q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 25 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.

A compound of the invention or a pharmaceutically acceptable salt, solvate,
 clathrate, or prodrug thereof is particularly useful inhibiting immune cell (e.g.,
 30 T-cells and/or B-cells) activation (e.g., activation in response to an antigen). In
 particular, a compound of the invention or a pharmaceutically acceptable salt,
 solvate, clathrate, or prodrug thereof can inhibit the production of certain
 cytokines that regulate immune cell activation. For example, a compound of

the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can inhibit the production of IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , INF- γ or combinations thereof. Moreover, a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can
5 modulate the activity of one or more ion channel involved in activation of immune cells, such as CRAC ion channels.

A compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof is particularly useful for immunosuppression or for
10 treating or preventing inflammatory conditions, allergic disorders, and immune disorders.

The invention also encompasses pharmaceutical compositions comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof; and a pharmaceutically acceptable carrier or
15 vehicle. These compositions may further comprise additional agents. These compositions are useful for immunosuppression and treating or preventing inflammatory conditions, allergic disorders and immune disorders.

20 The invention further encompasses methods for treating or preventing inflammatory conditions, allergic disorders, and immune disorders, comprising administering to a subject in need thereof an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, or a pharmaceutical composition comprising a compound of
25 the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof. These methods may also comprise administering to the subject an additional agent separately or in a combination composition with the compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

30

The invention further encompasses methods for suppressing the immune system of a subject, comprising administering to a subject in need thereof an effective amount of a compound of the invention or a pharmaceutically

acceptable salt, solvate, clathrate, or prodrug thereof, or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof. These methods may also comprise administering to the subject an additional agent separately or in
5 a combination composition with the compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

The invention further encompasses methods for inhibiting immune cell activation, including inhibiting proliferation of T cells and/or B cells, *in vivo* or *in*
10 *vitro* comprising administering to the cell an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

15 The invention further encompasses methods for inhibiting cytokine production in a cell, (e.g., IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , and/or INF- γ production) *in vivo* or *in vitro* comprising administering to a cell an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate,
20 clathrate, or prodrug thereof or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

The invention further encompasses methods for modulating ion channel
25 activity (e.g., CRAC) *in vivo* or *in vitro* comprising administering an effective amount of a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof or a pharmaceutical composition comprising a compound of the invention or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

30 All of the methods of this invention may be practice with a compound of the invention alone, or in combination with other agents, such as other

immunosuppressive agents, anti-inflammatory agents, agents for the treatment of allergic disorders or agents for the treatment of immune disorders.

DETAILED DESCRIPTION OF THE INVENTION

5

DEFINITIONS

Unless otherwise specified, the below terms used herein are defined as follows:

10 As used herein, the term an "aromatic ring" or "aryl" means a monocyclic or polycyclic-aromatic ring or ring radical comprising carbon and hydrogen atoms. Examples of suitable aryl groups include, but are not limited to, phenyl, tolyl, anthacenyl, fluorenyl, indenyl, azulenyl, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl. An aryl group can be
15 unsubstituted or substituted with one or more substituents (including without limitation alkyl (preferably, lower alkyl or alkyl substituted with one or more halo), hydroxy, alkoxy (preferably, lower alkoxy), alkylthio, cyano, halo, amino, and nitro. In certain embodiments, the aryl group is a monocyclic ring, wherein the ring comprises 6 carbon atoms.

20

As used herein, the term "alkyl" means a saturated straight chain or branched non-cyclic hydrocarbon typically having from 1 to 10 carbon atoms. Representative saturated straight chain alkyls include methyl, ethyl, n-propyl, n-butyl, n-pentyl, n-hexyl, n-heptyl, n-octyl, n-nonyl and n-decyl; while saturated
25 branched alkyls include isopropyl, sec-butyl, isobutyl, *tert*-butyl, isopentyl, 2-methylbutyl, 3-methylbutyl, 2-methylpentyl, 3-methylpentyl, 4-methylpentyl, 2-methylhexyl, 3-methylhexyl, 4-methylhexyl, 5-methylhexyl, 2,3-dimethylbutyl, 2,3-dimethylpentyl, 2,4-dimethylpentyl, 2,3-dimethylhexyl, 2,4-dimethylhexyl, 2,5-dimethylhexyl, 2,2-dimethylpentyl, 2,2-dimethylhexyl, 3,3-dimethylpentyl, 3,3-dimethylhexyl, 4,4-dimethylhexyl, 2-ethylpentyl, 3-ethylpentyl, 2-ethylhexyl,
30 3-ethylhexyl, 4-ethylhexyl, 2-methyl-2-ethylpentyl, 2-methyl-3-ethylpentyl, 2-methyl-4-ethylpentyl, 2-methyl-2-ethylhexyl, 2-methyl-3-ethylhexyl, 2-methyl-4-ethylhexyl, 2,2-diethylpentyl, 3,3-diethylhexyl, 2,2-diethylhexyl,

3,3-diethylhexyl and the like. Alkyl groups included in compounds of this invention may be optionally substituted with one or more substituents, such as amino, alkylamino, alkoxy, alkylthio, oxo, halo, acyl, nitro, hydroxyl, cyano, aryl, alkylaryl, aryloxy, arylthio, arylamino, carbocyclyl, carbocycloxy, carbocyclylthio, carbocyclylamino, heterocyclyl, heterocycloxy, heterocyclylamino, heterocyclylthio, and the like. In addition, any carbon in the alkyl segment may be substituted with oxygen (=O), sulfur (=S), or nitrogen (=NR²³, wherein R²³ is -H, an alkyl, acetyl, or aralkyl). Lower alkyls are typically preferred for the compounds of this invention.

The term alkylene refers to an alkyl group that has two points of attachment to two moieties (e.g., {-CH₂-}, {-CH₂CH₂-},



, etc., wherein the brackets indicate the points of attachment). Alkylene groups may be substituted or unsubstituted.

An aralkyl group refers to an aryl group that is attached to another moiety via an alkylene linker. Aralkyl groups can be substituted or unsubstituted.

The term "alkoxy," as used herein, refers to an alkyl group which is linked to another moiety through an oxygen atom. Alkoxy groups can be substituted or unsubstituted.

The term "alkoxyalkoxy," as used herein, refers to an alkoxy group in which the alkyl portion is substituted with another alkoxy group.

The term "alkyl sulfanyl," as used herein, refers to an alkyl group which is linked to another moiety through a divalent sulfur atom. Alkyl sulfanyl groups can be substituted or unsubstituted.

The term "alkylamino," as used herein, refers to an amino group in which one hydrogen atom attached to the nitrogen has been replaced by an alkyl group. The term "dialkylamino," as used herein, refers to an amino group in which two
5 hydrogen atoms attached to the nitrogen have been replaced by alkyl groups, in which the alkyl groups can be the same or different. Alkylamino groups and dialkylamino groups can be substituted or unsubstituted.

As used herein, the term "alkenyl" means a straight chain or branched,
10 hydrocarbon radical typically having from 2 to 10 carbon atoms and having at least one carbon-carbon double bond. Representative straight chain and branched alkenyls include vinyl, allyl, 1-butenyl, 2-butenyl, isobutylenyl, 1-pentenyl, 2-pentenyl, 3-methyl-1-butenyl, 1-methyl-2-butenyl, 2,3-dimethyl-2-butenyl, 1-hexenyl, 2-hexenyl, 3-hexenyl, 1-heptenyl,
15 2-heptenyl, 3-heptenyl, 1-octenyl, 2-octenyl, 3-octenyl, 1-nonenyl, 2-nonenyl, 3-nonenyl, 1-decenyl, 2-decenyl, 3-decenyl and the like. Alkenyl groups can be substituted or unsubstituted.

As used herein, the term "alkynyl" means a straight chain or branched,
20 hydrocarbon radical typically having from 2 to 10 carbon atoms and having at least one carbon-carbon triple bond. Representative straight chain and branched alkynyls include acetylenyl, propynyl, 1-butyne, 2-butyne, 1-pentyne, 2-pentyne, 3-methyl-1-butyne, 4-pentyne, 1-hexyne, 2-hexyne, 5-hexyne, 1-heptyne, 2-heptyne, 6-heptyne, 1-octyne, 2-octyne, 7-octyne,
25 1-nonyne, 2-nonyne, 8-nonyne, 1-decynyl, 2-decynyl, 9-decynyl and the like. Alkynyl groups can be substituted or unsubstituted.

As used herein, the term "cycloalkyl" means a saturated, mono- or polycyclic alkyl radical typically having from 3 to 10 carbon atoms. Representative
30 cycloalkyls include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl, adamantyl, decahydronaphthyl, octahydronaphthalene, bicyclo[1.1.1]pentyl, and the like. Cycloalkyl groups can be substituted or unsubstituted.

As used herein, the term "cycloalkenyl" means a cyclic non-aromatic alkenyl radical having at least one carbon-carbon double bond in the cyclic system and typically having from 5 to 10 carbon atoms. Representative cycloalkenyls include cyclopentenyl, cyclopentadienyl, cyclohexenyl, cyclohexadienyl, cycloheptenyl, cycloheptadienyl, cycloheptatrienyl, cyclooctenyl, cyclooctadienyl, cyclooctatrienyl, cyclooctatetraenyl, cyclononenyl, cyclononadienyl, cyclodecenyl, cyclodecadienyl and the like. Cycloalkenyl groups can be substituted or unsubstituted.

10

As used herein, the term "heterocycle" or "heterocyclyl" means a monocyclic or polycyclic heterocyclic ring (typically having 3- to 14-members) which is either a saturated ring or an unsaturated non-aromatic ring. A 3-membered heterocycle can contain up to 3 heteroatoms, and a 4- to 14-membered heterocycle can contain from 1 to about 8 heteroatoms. Each heteroatom is independently selected from nitrogen, which can be quaternized; oxygen; and sulfur, including sulfoxide and sulfone. The heterocycle may be attached via any heteroatom or carbon atom. Representative heterocycles include morpholinyl, thiomorpholinyl, pyrrolidinonyl, pyrrolidinyl, piperidinyl, piperazinyl, hydantoinyl, valerolactamyl, oxiranyl, oxetanyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydropyrindinyl, tetrahydropyrimidinyl, tetrahydrothiophenyl, tetrahydrothiopyranyl, and the like. A heteroatom may be substituted with a protecting group known to those of ordinary skill in the art, for example, the hydrogen on a nitrogen may be substituted with a tert-butoxycarbonyl group. Furthermore, the heterocyclyl may be optionally substituted with one or more substituents (including without limitation a halogen atom, an alkyl radical, or aryl radical). Only stable isomers of such substituted heterocyclic groups are contemplated in this definition. Heterocyclyl groups can be substituted or unsubstituted.

30

As used herein, the term "heteroaromatic" or "heteroaryl" means a monocyclic or polycyclic heteroaromatic ring (or radical thereof) comprising carbon atom ring members and one or more heteroatom ring members (such as, for

example, oxygen, sulfur or nitrogen). Typically, the heteroaromatic ring has from 5 to about 14 ring members in which at least 1 ring member is a heteroatom selected from oxygen, sulfur and nitrogen. In another embodiment, the heteroaromatic ring is a 5 or 6 membered ring and may contain from 1 to about 4 heteroatoms. In another embodiment, the heteroaromatic ring system has a 7 to 14 ring members and may contain from 1 to about 7 heteroatoms. Representative heteroaryls include pyridyl, furyl, thienyl, pyrrolyl, oxazolyl, imidazolyl, indoliziny, thiazolyl, isoxazolyl, pyrazolyl, isothiazolyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl, triazolyl, pyridinyl, thiadiazolyl, pyrazinyl, quinolyl, isoquinolyl, indazolyl, benzoxazolyl, benzofuryl, benzothiazolyl, indoliziny, imidazopyridinyl, isothiazolyl, tetrazolyl, benzimidazolyl, benzoxazolyl, benzothiazolyl, benzothiadiazolyl, benzoxadiazolyl, indolyl, tetrahydroindolyl, azaindolyl, imidazopyridyl, quinazolinyl, purinyl, pyrrolo[2,3]pyrimidyl, pyrazolo[3,4]pyrimidyl or benzo(b)thienyl and the like. These heteroaryl groups may be optionally substituted with one or more substituents

A heteroaralkyl group refers to a heteroaryl group that is attached to another moiety via an alkylene linker. Heteroaralkyl groups can be substituted or unsubstituted.

As used herein, the term "halogen" or "halo" means -F, -Cl, -Br or -I.

As used herein, the term "haloalkyl" means an alkyl group in which one or more -H is replaced with a halo group. Examples of haloalkyl groups include -CF₃, -CHF₂, -CCl₃, -CH₂CH₂Br, -CH₂CH(CH₂CH₂Br)CH₃, -CHICH₃, and the like.

As used herein, the term "haloalkoxy" means an alkoxy group in which one or more -H is replaced with a halo group. Examples of haloalkoxy groups include -OCF₃ and -OCHF₂.

The terms "bioisostere" and "bioisosteric replacement" have the same meanings as those generally recognized in the art. Bioisosteres are atoms,

ions, or molecules in which the peripheral layers of electrons can be considered substantially identical. The term bioisostere is usually used to mean a portion of an overall molecule, as opposed to the entire molecule itself. Bioisosteric replacement involves using one bioisostere to replace another with the
5 expectation of maintaining or slightly modifying the biological activity of the first bioisostere. The bioisosteres in this case are thus atoms or groups of atoms having similar size, shape and electron density. Preferred bioisosteres of esters, amides or carboxylic acids are compounds containing two sites for hydrogen bond acceptance. In one embodiment, the ester, amide or carboxylic
10 acid bioisostere is a 5-membered monocyclic heteroaryl ring, such as an optionally substituted 1H-imidazolyl, an optionally substituted oxazolyl, 1H-tetrazolyl, [1,2,4]triazolyl, or an optionally substituted [1,2,4]oxadiazolyl.

As used herein, the terms "subject", "patient" and "animal", are used
15 interchangeably and include, but are not limited to, a cow, monkey, horse, sheep, pig, chicken, turkey, quail, cat, dog, mouse, rat, rabbit, guinea pig and human. The preferred subject, patient or animal is a human.

As used herein, the term "lower" refers to a group having up to four carbon
20 atoms. For example, a "lower alkyl" refers to an alkyl radical having from 1 to 4 carbon atoms, and a "lower alkenyl" or "lower alkynyl" refers to an alkenyl or alkynyl radical having from 2 to 4 carbon atoms, respectively. A lower alkoxy or a lower alkyl sulfanyl refers to an alkoxy or a alkyl sulfanyl having from 1 to 4 carbon atoms. Lower substituents are typically preferred.

25

Where a particular substituent, such as an alkyl substituent, occurs multiple times in a given structure or moiety, the identity of the substituent is independent in each case and may be the same as or different from other occurrences of that substituent in the structure or moiety. Furthermore,
30 individual substituents in the specific embodiments and exemplary compounds of this invention are preferred in combination with other such substituents in the compounds of this invention, even if such individual substituents are not

expressly noted as being preferred or not expressly shown in combination with other substituents.

5 The compounds of the invention are defined herein by their chemical structures and/or chemical names. Where a compound is referred to by both a chemical structure and a chemical name, and the chemical structure and chemical name conflict, the chemical structure is determinative of the compound's identity.

10 Suitable substituents for an alkyl, alkoxy, alkyl sulfanyl, alkylamino, dialkylamino, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and heteroarylalkyl groups include any substituent which will form a stable compound of the invention. Examples of substituents for an alkyl, alkoxy, alkylsulfanyl, alkylamino, dialkylamino, alkylene, alkenyl, alkynyl, cycloalkyl, cycloalkenyl, heterocyclyl, aryl, aralkyl, heteroaryl, and
 15 heteroarylalkyl include an alkyl, alkoxy, alkyl sulfanyl, alkylamino, dialkylamino, an alkenyl, an alkynyl, an cycloalkyl, an cycloalkenyl, an heterocyclyl, an aryl, an heteroaryl, an aralkyl, an heteraralkyl, a haloalkyl, $-C(O)NR_{13}R_{14}$, $-NR_{15}C(O)R_{16}$, halo, $-OR_{15}$, cyano, nitro, haloalkoxy, $-C(O)R_{15}$, $-NR_{13}R_{14}$, $-SR_{15}$, $-C(O)OR_{15}$, $-OC(O)R_{15}$, $-NR_{15}C(O)NR_{13}R_{14}$,
 20 $-OC(O)NR_{13}R_{14}$, $-NR_{15}C(O)OR_{16}$, $-S(O)_pR_{15}$, or $-S(O)_pNR_{13}R_{14}$, wherein R_{13} and R_{14} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;
 25 substituted heteraralkyl; or R_{13} and R_{14} taken together with the nitrogen to which they are attached is optionally substituted heterocyclyl or optionally substituted heteroaryl; and R_{15} and R_{16} for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl,
 30 alkynyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

In addition, alkyl, cycloalkyl, alkylene, a heterocyclyl, and any saturated portion of a alkenyl, cycloalkenyl, alkynyl, aralkyl, and heteroaralkyl groups, may also be substituted with =O, =S, =N-R₁₅.

5

When a heterocyclyl, heteroaryl, or heteroaralkyl group contains a nitrogen atom, it may be substituted or unsubstituted. When a nitrogen atom in the aromatic ring of a heteroaryl group has a substituent the nitrogen may be a quaternary nitrogen.

10

Choices and combinations of substituents and variables envisioned by this invention are only those that result in the formation of stable compounds. The term "stable", as used herein, refers to compounds which possess stability sufficient to allow manufacture and which maintains the integrity of the compound for a sufficient period of time to be useful for the purposes detailed herein (e.g., therapeutic or prophylactic administration to a subject). Typically, such compounds are stable at a temperature of 40 °C or less, in the absence of excessive moisture, for at least one week. Such choices and combinations will be apparent to those of ordinary skill in the art and may be determined without undue experimentation.

20

Unless indicated otherwise, the compounds of the invention containing reactive functional groups (such as, without limitation, carboxy, hydroxy, and amino moieties) also include protected derivatives thereof. "Protected derivatives" are those compounds in which a reactive site or sites are blocked with one or more protecting groups. Suitable protecting groups for carboxy moieties include benzyl, tert-butyl, and the like. Suitable protecting groups for amino and amido groups include acetyl, tert-butoxycarbonyl, benzyloxycarbonyl, and the like. Suitable protecting groups for hydroxy include benzyl and the like. Other suitable protecting groups are well known to those of ordinary skill in the art and include those found in T. W. Greene, *Protecting Groups in Organic Synthesis*, John Wiley & Sons, Inc. 1981, the entire teachings of which are incorporated herein by reference.

25

30

As used herein, the term "compound(s) of this invention" and similar terms refers to a compound of any one of formulas (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof and
5 also include protected derivatives thereof.

As used herein and unless otherwise indicated, the term "prodrug" means a derivative of a compound that can hydrolyze, oxidize, or otherwise react under biological conditions (*in vitro* or *in vivo*) to provide a compound of this invention.
10 Prodrugs may only become active upon such reaction under biological conditions, but they may have activity in their unreacted forms. Examples of prodrugs contemplated in this invention include, but are not limited to, analogs or derivatives of compounds of any one of formulas (I) through (II), or Table 1 that comprise biohydrolyzable moieties such as biohydrolyzable amides,
15 biohydrolyzable esters, biohydrolyzable carbamates, biohydrolyzable carbonates, biohydrolyzable ureides, and biohydrolyzable phosphate analogues. Other examples of prodrugs include derivatives of compounds of any one of formulas (I) through (II), or of Table 1 that comprise -NO, -NO₂, -ONO, or -ONO₂ moieties. Prodrugs can typically be
20 prepared using well-known methods, such as those described by 1 BURGER'S MEDICINAL CHEMISTRY AND DRUG DISCOVERY (1995) 172-178, 949-982 (Manfred E. Wolff ed., 5th ed), the entire teachings of which are incorporated herein by reference.

25 As used herein and unless otherwise indicated, the terms "biohydrolyzable amide", "biohydrolyzable ester", "biohydrolyzable carbamate", "biohydrolyzable carbonate", "biohydrolyzable ureide" and "biohydrolyzable phosphate analogue" mean an amide, ester, carbamate, carbonate, ureide, or phosphate analogue, respectively, that either: 1) does not destroy the biological activity of
30 the compound and confers upon that compound advantageous properties *in vivo*, such as uptake, duration of action, or onset of action; or 2) is itself biologically inactive but is converted *in vivo* to a biologically active compound. Examples of biohydrolyzable amides include, but are not limited to, lower alkyl

amides, α -amino acid amides, alkoxyacyl amides, and alkylaminoalkylcarbonyl amides. Examples of biohydrolyzable esters include, but are not limited to, lower alkyl esters, alkoxyacyloxy esters, alkyl acylamino alkyl esters, and choline esters. Examples of biohydrolyzable carbamates include, but are not limited to, lower alkylamines, substituted ethylenediamines, aminoacids, hydroxyalkylamines, heterocyclic and heteroaromatic amines, and polyether amines.

As used herein, the term "pharmaceutically acceptable salt," is a salt formed from an acid and a basic group of one of the compounds of any one of formulas (I) through (II) or of Table 1. Illustrative salts include, but are not limited, to sulfate, citrate, acetate, oxalate, chloride, bromide, iodide, nitrate, bisulfate, phosphate, acid phosphate, isonicotinate, lactate, salicylate, acid citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucaronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate, and pamoate (*i.e.*, 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. The term "pharmaceutically acceptable salt" also refers to a salt prepared from a compound of any one of formulas (I) through (II) or Table 1 having an acidic functional group, such as a carboxylic acid functional group, and a pharmaceutically acceptable inorganic or organic base. Suitable bases include, but are not limited to, hydroxides of alkali metals such as sodium, potassium, and lithium; hydroxides of alkaline earth metal such as calcium and magnesium; hydroxides of other metals, such as aluminum and zinc; ammonia, and organic amines, such as unsubstituted or hydroxy-substituted mono-, di-, or trialkylamines; dicyclohexylamine; tributyl amine; pyridine; N-methyl,N-ethylamine; diethylamine; triethylamine; mono-, bis-, or tris-(2-hydroxy-lower alkyl amines), such as mono-, bis-, or tris-(2-hydroxyethyl)- amine, 2-hydroxy-tert-butylamine, or tris-(hydroxymethyl)methylamine, N, N,-di-lower alkyl-N-(hydroxy lower alkyl)-amines, such as N,N-dimethyl-N-(2-hydroxyethyl)- amine, or tri-(2-hydroxyethyl)amine; N-methyl-D-glucamine; and amino acids such as

arginine, lysine, and the like. The term "pharmaceutically acceptable salt" also refers to a salt prepared from a compound of any one of formulas (I) through (II) or Table 1 having a basic functional group, such as an amino functional group, and a pharmaceutically acceptable inorganic or organic acid. Suitable acids
5 include, but are not limited to, hydrogen sulfate, citric acid, acetic acid, oxalic acid, hydrochloric acid, hydrogen bromide, hydrogen iodide, nitric acid, phosphoric acid, isonicotinic acid, lactic acid, salicylic acid, tartaric acid, ascorbic acid, succinic acid, maleic acid, besylic acid, fumaric acid, gluconic acid, glucaronic acid, saccharic acid, formic acid, benzoic acid, glutamic acid,
10 methanesulfonic acid, ethanesulfonic acid, benzenesulfonic acid, and *p*-toluenesulfonic acid.

As used herein, the term "pharmaceutically acceptable solvate," is a solvate formed from the association of one or more solvent molecules to one or more
15 molecules of a compound of any one of formulas (I) through (II) or Table 1. The term solvate includes hydrates (*e.g.*, hemi-hydrate, mono-hydrate, dihydrate, trihydrate, tetrahydrate, and the like).

As used herein, the term "clathrate" means a compound of the present
20 invention or a salt thereof in the form of a crystal lattice that contains spaces (*e.g.*, channels) that have a guest molecule (*e.g.*, a solvent or water) trapped within.

As used herein, the term "asthma" means a pulmonary disease, disorder or
25 condition characterized by reversible airway obstruction, airway inflammation, and increased airway responsiveness to a variety of stimuli.

"Immunosuppression" refers to impairment of any component of the immune system resulting in decreased immune function. This impairment may be
30 measured by any conventional means including whole blood assays of lymphocyte function, detection of lymphocyte proliferation and assessment of the expression of T cell surface antigens. The antisheep red blood cell (SRBC) primary (IgM) antibody response assay (usually referred to as the plaque

assay) is one specific method. This and other methods are described in Luster, M.I., Portier, C., Pait, D.G., White, K.L., Jr., Gennings, C., Munson, A.E., and Rosenthal, G.J. (1992). "Risk Assessment in Immunotoxicology I: Sensitivity and Predictability of Immune Tests." *Fundam. Appl. Toxicol.*, 18, 200-210.

5 Measuring the immune response to a T-cell dependent immunogen is another particularly useful assay (Dean, J.H., House, R.V., and Luster, M.I. (2001). "Immunotoxicology: Effects of, and Responses to, Drugs and Chemicals." In *Principles and Methods of Toxicology: Fourth Edition* (A.W. Hayes, Ed.), pp. 1415-1450, Taylor & Francis, Philadelphia, Pennsylvania).

10

The compounds of this invention can be used to treat subjects with immune disorders. As used herein, the term "immune disorder" and like terms means a disease, disorder or condition caused by the immune system of an animal, including autoimmune disorders. Immune disorders include those diseases,
15 disorders or conditions that have an immune component and those that are substantially or entirely immune system-mediated. Autoimmune disorders are those wherein the animal's own immune system mistakenly attacks itself, thereby targeting the cells, tissues, and/or organs of the animal's own body. For example, the autoimmune reaction is directed against the nervous system
20 in multiple sclerosis and the gut in Crohn's disease. In other autoimmune disorders such as systemic lupus erythematosus (lupus), affected tissues and organs may vary among individuals with the same disease. One person with lupus may have affected skin and joints whereas another may have affected skin, kidney, and lungs. Ultimately, damage to certain tissues by the immune
25 system may be permanent, as with destruction of insulin-producing cells of the pancreas in Type 1 diabetes mellitus. Specific autoimmune disorders that may be ameliorated using the compounds and methods of this invention include without limitation, autoimmune disorders of the nervous system (e.g., multiple sclerosis, myasthenia gravis, autoimmune neuropathies such as
30 Guillain-Barré, and autoimmune uveitis), autoimmune disorders of the blood (e.g., autoimmune hemolytic anemia, pernicious anemia, and autoimmune thrombocytopenia), autoimmune disorders of the blood vessels (e.g., temporal arteritis, anti-phospholipid syndrome, vasculitides such as Wegener's

granulomatosis, and Behcet's disease), autoimmune disorders of the skin (e.g., psoriasis, dermatitis herpetiformis, pemphigus vulgaris, and vitiligo), autoimmune disorders of the gastrointestinal system (e.g., Crohn's disease, ulcerative colitis, primary biliary cirrhosis, and autoimmune hepatitis),
5 autoimmune disorders of the endocrine glands (e.g., Type 1 or immune-mediated diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, and autoimmune disorder of the adrenal gland); and autoimmune disorders of multiple organs (including connective tissue and musculoskeletal system diseases) (e.g., rheumatoid arthritis,
10 systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, spondyloarthropathies such as ankylosing spondylitis, and Sjogren's syndrome). In addition, other immune system mediated diseases, such as graft-versus-host disease and allergic disorders, are also included in the definition of immune disorders herein. Because a number of immune disorders
15 are caused by inflammation, there is some overlap between disorders that are considered immune disorders and inflammatory disorders. For the purpose of this invention, in the case of such an overlapping disorder, it may be considered either an immune disorder or an inflammatory disorder. "Treatment of an immune disorder" herein refers to administering a compound or a composition
20 of the invention to a subject, who has an immune disorder, a symptom of such a disease or a predisposition towards such a disease, with the purpose to cure, relieve, alter, affect, or prevent the autoimmune disorder, the symptom of it, or the predisposition towards it.

25 As used herein, the term "allergic disorder" means a disease, condition or disorder associated with an allergic response against normally innocuous substances. These substances may be found in the environment (such as indoor air pollutants and aeroallergens) or they may be non-environmental (such as those causing dermatological or food allergies). Allergens can enter
30 the body through a number of routes, including by inhalation, ingestion, contact with the skin or injection (including by insect sting). Many allergic disorders are linked to atopy, a predisposition to generate the allergic antibody IgE. Because IgE is able to sensitize mast cells anywhere in the body, atopic individuals often

- express disease in more than one organ. For the purpose of this invention, allergic disorders include any hypersensitivity that occurs upon re-exposure to the sensitizing allergen, which in turn causes the release of inflammatory mediators. Allergic disorders include without limitation, allergic rhinitis (e.g.,
5 hay fever), sinusitis, rhinosinusitis, chronic or recurrent otitis media, drug reactions, insect sting reactions, latex reactions, conjunctivitis, urticaria, anaphylaxis and anaphylactoid reactions, atopic dermatitis, asthma and food allergies.
- 10 The compounds of this invention can be used to prevent or to treat subjects with inflammatory disorders. As used herein, an "inflammatory disorder" means a disease, disorder or condition characterized by inflammation of body tissue or having an inflammatory component. These include local inflammatory responses and systemic inflammation. Examples of such inflammatory
15 disorders include: transplant rejection, including skin graft rejection; chronic inflammatory disorders of the joints, including arthritis, rheumatoid arthritis, osteoarthritis and bone diseases associated with increased bone resorption; inflammatory bowel diseases such as ileitis, ulcerative colitis, Barrett's syndrome, and Crohn's disease; inflammatory lung disorders such as asthma,
20 adult respiratory distress syndrome, and chronic obstructive airway disease; inflammatory disorders of the eye including corneal dystrophy, trachoma, onchocerciasis, uveitis, sympathetic ophthalmitis and endophthalmitis; chronic inflammatory disorders of the gums, including gingivitis and periodontitis; tuberculosis; leprosy; inflammatory diseases of the kidney including uremic complications, glomerulonephritis and nephrosis; inflammatory disorders of the
25 skin including sclerodermatitis, psoriasis and eczema; inflammatory diseases of the central nervous system, including chronic demyelinating diseases of the nervous system, multiple sclerosis, AIDS-related neurodegeneration and Alzheimer's disease, infectious meningitis, encephalomyelitis, Parkinson's
30 disease, Huntington's disease, amyotrophic lateral sclerosis and viral or autoimmune encephalitis; autoimmune disorders, immune-complex vasculitis, systemic lupus and erythematodes; systemic lupus erythematosus (SLE); and inflammatory diseases of the heart such as cardiomyopathy, ischemic heart

- disease hypercholesterolemia, atherosclerosis); as well as various other diseases with significant inflammatory components, including preeclampsia; chronic liver failure, brain and spinal cord trauma, cancer). There may also be a systemic inflammation of the body, exemplified by gram-positive or gram negative shock, hemorrhagic or anaphylactic shock, or shock induced by cancer chemotherapy in response to pro-inflammatory cytokines, e.g., shock associated with pro-inflammatory cytokines. Such shock can be induced, e.g., by a chemotherapeutic agent used in cancer chemotherapy. "Treatment of an inflammatory disorder" herein refers to administering a compound or a composition of the invention to a subject, who has an inflammatory disorder, a symptom of such a disorder or a predisposition towards such a disorder, with the purpose to cure, relieve, alter, affect, or prevent the inflammatory disorder, the symptom of it, or the predisposition towards it.
- 15 An "effective amount" is the quantity of compound in which a beneficial outcome is achieved when the compound is administered to a subject or alternatively, the quantity of compound that possess a desired activity in-vivo or in-vitro. In the case of inflammatory disorders and autoimmune disorders, a beneficial clinical outcome includes reduction in the extent or severity of the symptoms associated with the disease or disorder and/or an increase in the longevity and/or quality of life of the subject compared with the absence of the treatment. The precise amount of compound administered to a subject will depend on the type and severity of the disease or condition and on the characteristics of the subject, such as general health, age, sex, body weight and tolerance to drugs. It will also depend on the degree, severity and type of inflammatory disorder or autoimmune disorder or the degree of immunosuppression sought. The skilled artisan will be able to determine appropriate dosages depending on these and other factors. Effective amounts of the disclosed compounds typically range between about 1 mg/mm² per day and about 10 grams/mm² per day, and preferably between 10 mg/mm² per day and about 1 gram/mm².

The compounds of the invention may contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as double-bond isomers (*i.e.*, geometric isomers), enantiomers, or diastereomers. According to this invention, the chemical structures depicted herein, including the
5 compounds of this invention, encompass all of the corresponding compounds' enantiomers and stereoisomers, that is, both the stereomerically pure form (*e.g.*, geometrically pure, enantiomerically pure, or diastereomerically pure) and enantiomeric, diastereomeric, and geometric isomeric mixtures. In some cases, one enantiomer, diastereomer, or geometric isomer will possess
10 superior activity or an improved toxicity or kinetic profile compared to others. In those cases, such enantiomers, diastereomers, and geometric isomers of a compound of this invention are preferred.

The term "inhibit production of IL-2" and like terms means inhibiting IL-2
15 synthesis (*e.g.* by inhibiting transcription (mRNA expression), or translation (protein expression)) and/or inhibiting IL-2 secretion in a cell that has the ability to produce and/or secrete IL-2 (*e.g.*, T lymphocyte). Likewise, the term "inhibiting production of IL-4, IL-5, IL-13, GM-CSF, TNF- α or INF- γ means inhibiting the synthesis (*e.g.* by inhibiting transcription, or translation) and/or
20 inhibiting the secretion in a cell that has the ability to produce and/or secrete these cytokines.

As used herein, a composition that "substantially" comprises a compound means that the composition contains more than about 80% by weight, more
25 preferably more than about 90% by weight, even more preferably more than about 95% by weight, and most preferably more than about 97% by weight of the compound.

As used herein, a composition that is "substantially free" of a compound means
30 that the composition contains less than about 20% by weight, more preferably less than about 10% by weight, even more preferably less than about 5% by weight, and most preferably less than about 3% by weight of the compound.

As used herein, a reaction that is "substantially complete" means that the reaction contains more than about 80% by weight of the desired product, more preferably more than about 90% by weight of the desired product, even more preferably more than about 95% by weight of the desired product, and most
5 preferably more than about 97% by weight of the desired product.

As used herein, a racemic mixture means about 50% of one enantiomer and about 50% of its corresponding enantiomer relative to all chiral centers in the molecule. The invention encompasses all enantiomerically-pure,
10 enantiomerically-enriched, diastereomerically pure, diastereomerically enriched, and racemic mixtures of the compounds of any one of formulas (I) through (II) or Table 1.

Enantiomeric and diastereomeric mixtures can be resolved into their
15 component enantiomers or stereoisomers by well known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Enantiomers and diastereomers can also be obtained from diastereomerically- or enantiomerically-pure
20 intermediates, reagents, and catalysts by well known asymmetric synthetic methods.

When administered to a patient, e.g., to a non-human animal for veterinary use or for improvement of livestock, or to a human for clinical use, the compounds
25 of the invention are typically administered in isolated form or as the isolated form in a pharmaceutical composition. As used herein, "isolated" means that the compounds of the invention are separated from other components of either (a) a natural source, such as a plant or cell, preferably bacterial culture, or (b) a synthetic organic chemical reaction mixture. Preferably, via conventional
30 techniques, the compounds of the invention are purified. As used herein, "purified" means that when isolated, the isolate contains at least 95%, preferably at least 98%, of a single compound of the invention by weight of the isolate.


Only those choices and combinations of substituents that result in a stable structure are contemplated. Such choices and combinations will be apparent to those of ordinary skill in the art and may be determined without undue experimentation.

The invention can be understood more fully by reference to the following detailed description and illustrative examples, which are intended to exemplify non-limiting embodiments of the invention.

SPECIFIC EMBODIMENTS

The invention relates to compounds and pharmaceutical compositions that are particularly useful for immunosuppression or to treat or prevent inflammatory conditions, immune disorders, and allergic disorders.

One embodiment of the invention relates to compounds of formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of -NR₂CR₂-, -CR₂NR-, -C(O)-, -NR-C(O)-, -C(O)-NR-, -OC(O)-, -C(O)O-, -C(S)-, -NR-C(S)-, -C(S)-NR-;

one of X_1 or X_2 is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$, cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from $-H$ or an alkyl;

R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$, or $-NR_6R_7$;

R_4 is a halo, a haloalkyl, cyano, nitro, $-C(O)R_5$, $-C(O)OR_5$, $-C(O)SR_5$, $-C(O)NR_6R_7$, $-C(S)R_5$, $-C(S)OR_5$, $-C(S)SR_5$, $-C(S)NR_6R_7$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-C(NR_8)SR_5$, $-C(NR_8)NR_6R_7$, $-S(O)_pR_5$, $-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

R_8 , for each occurrence, is independently $-H$, a halo, an alkyl, $-OR_5$, $-NR_6R_7$, $-C(O)R_5$, $-C(O)OR_5$, or $-C(O)NR_6R_7$;

R_9 and R_{10} , for each occurrence are, independently, H, an

optionally substituted alkyl, an optionally substituted alkenyl, an
 optionally substituted alkynyl, an optionally substituted 5- to
 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an
 optionally substituted heterocyclyl, an optionally substituted aryl, an
 5 optionally substituted aralkyl, an optionally substituted heteraralkyl, or
 an optionally substituted heteroaryl selected from the group consisting
 of an optionally substituted pyridinyl, an optionally substituted thienyl, an
 optionally substituted pyrrolyl, an optionally substituted oxazolyl, an
 optionally substituted imidazolyl, an optionally substituted indolizinyl, an
 10 optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an
 optionally substituted isothiazolyl, an optionally substituted pyridazinyl,
 an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl,
 an optionally substituted triazinyl, an optionally substituted triazolyl, an
 optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an
 15 optionally substituted pyrazinyl, an optionally substituted quinolinyl, an
 optionally substituted isoquinolinyl, an optionally substituted indazolyl,
 an optionally substituted benzoxazolyl, an optionally substituted
 benzofuryl, an optionally substituted benzothiazolyl, an optionally
 substituted indolizinyl, an optionally substituted imidazopyridinyl, an
 20 optionally substituted tetrazolyl, an optionally substituted
 benzimidazolyl, an optionally substituted benzoxazolyl, an optionally
 substituted benzothiazolyl, an optionally substituted benzothiadiazolyl,
 an optionally substituted benzoxadiazolyl, an optionally substituted
 indolyl, an optionally substituted tetrahydroindolyl, an optionally
 25 substituted azaindolyl, an optionally substituted imidazopyridyl, an
 optionally substituted quinoxalinyl, an optionally substituted purinyl, an
 optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted
 pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉
 and R₁₀ taken together with the nitrogen to which they are attached are
 30 an optionally substituted heterocyclyl or optionally substituted
 heteroaryl;

q, for each occurrence, is independently, an integer from 1 to 3;


k is for each occurrence, is independently, an integer from 1 to 4;

n is zero, 1 or 2; and

p, for each occurrence, is independently 1 or 2.

Another embodiment of the invention relates to compounds of formula (II):

5



(II)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

10

Y₁ is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl, provided that Y₁ is not a substituted isoxazolyl or a substituted 1H-pyrrolyl;

L₁ is a linker selected from the group consisting of -NHCH₂-, -CH₂NH-, -NR-C(O)-, -OC(O)-, -C(O)O-, -C(S)-, -NR-C(S)-, or -C(S)-NR-;

15

one of X₁ or X₂ is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, -S(O)_p-alkyl, -C(O)NRR, -(CH₂)_kNRR, -(CH₂)_kOR, -(CH₂)_kSR, cyano, nitro, or lower haloalkoxy;

20

R, for each occurrence is independently selected from -H or an alkyl;

R₃ is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, -OR₅, -SR₅, or -NR₆R₇;

25

R₄ is a halo, a haloalkyl, cyano, nitro, -C(O)R₅, -C(O)OR₅, -C(O)SR₅, -C(O)NR₉R₁₀, -C(S)R₅, -C(S)OR₅, -C(S)SR₅, -C(S)NR₆R₇, -C(NR₈)R₅, -C(NR₈)OR₅, -C(NR₈)SR₅, -C(NR₈)NR₆R₇, -S(O)_pR₅,

$-\text{S}(\text{O})_p\text{NR}_5$, $-\text{S}(\text{O})_p\text{OR}_5$, $-\text{P}(\text{O})(\text{OR}_5)_2$, $-\text{OP}(\text{O})(\text{OR}_5)_2$, $-\text{P}(\text{O})(\text{R}_5)_2$, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

5 R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

20 R_8 , for each occurrence, is independently $-\text{H}$, a halo, an alkyl, $-\text{OR}_5$, $-\text{NR}_6\text{R}_7$, $-\text{C}(\text{O})\text{R}_5$, $-\text{C}(\text{O})\text{OR}_5$, or $-\text{C}(\text{O})\text{NR}_6\text{R}_7$;

25 R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indolizinyl, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl,

- an optionally substituted triazinyl, an optionally substituted triazolyl, an optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally substituted indolizinyl, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;
- q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.
- In one embodiment, in compounds represented by formula (I) or (II), X₁ is N. In one aspect, X₂ is CH. In one aspect, X₂ is C(CH₃).
- In one embodiment, in compounds represented by formula (I) or (II), X₂ is N. In one aspect, X₁ is CH. In one aspect, X₁ is C(CH₃).
- In one embodiment, in compounds represented by formula (I) or (II), X₁ is N; X₂ is CH; Z is methyl; and n is 1.

In one embodiment, in compounds represented by formula (I) or (II), X_2 is N; X_1 is CH; Z is methyl; and n is 1.

In one embodiment, in compounds represented by formula (I) or (II), n is 0.

5

In one embodiment, in compounds represented by formula (I) or (II), n is 1 and Z is chloro, fluoro, bromo, methyl, methoxy, cyano, nitro, methylsulfanyl, or trifluoromethyl. In one aspect, Z is methyl.

10 In one embodiment, in compounds represented by formula (I) or (II), n is 2 and each Z is independently chloro, fluoro, bromo, methyl, methoxy, cyano, nitro, methylsulfanyl, or trifluoromethyl.

15 In one embodiment, in compounds represented by formula (I) or (II), Z is chloro, fluoro, bromo, methyl, methoxy, cyano, nitro, methylsulfanyl, or trifluoromethyl. In one aspect, Z is methyl.

20 In one embodiment, in compounds represented by formula (I) or (II), R_4 is $-C(O)OR_5$, a haloalkyl, an optionally substituted oxazolyl, an optionally substituted isoxazolyl, an optionally substituted thiazolyl, an optionally substituted imidazolyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyrrolyl, an optionally substituted thiophenyl, an optionally substituted furanyl, an optionally substituted thiadiazolyl, an optionally substituted oxadiazolyl, or an optionally substituted tetrazolyl. In another aspect, R_4 is $-C(O)OR_5$, a haloalkyl, an optionally substituted oxazolyl, an optionally substituted thiazolyl, an optionally substituted imidazolyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyrrolyl, an optionally substituted thiophenyl, an optionally substituted furanyl, an optionally substituted thiadiazolyl, an optionally substituted oxadiazolyl, or an optionally substituted tetrazolyl. In one aspect, R_4 is oxazol-2-yl, oxazol-5-yl, 3-methyl-isoxazol-5-yl, isoxazol-5-yl, thiazol-2-yl, thiazol-5-yl, [1,3,4]oxadiazol-2-yl, [1,3,4]thiadiazol-2-yl, trifluoromethyl, pyridin-3-yl,

25

30

pyridin-2-yl, furan-2-yl, furan-3-yl, thiophen-2-yl, thiophen-3-yl,
 -C(O)OCH₂CH₂OCH₃, -C(O)OCH₂CH₂CH₃, -C(O)OCH₂CH₃, or -C(O)OCH₃. In
 one aspect, R₄ is oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, thiazol-5-yl,
 [1,3,4]oxadiazol-2-yl, trifluoromethyl, pyridin-3-yl, pyridin-2-yl,
 5 -C(O)OCH₂CH₂CH₃, -C(O)OCH₂CH₃, or -C(O)OCH₃. In one aspect, R₄ is
 oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, trifluoromethyl, or -C(O)OCH₃. In one
 aspect, R₄ is oxazol-2-yl. In one aspect, R₄ is oxazol-5-yl. In one aspect, R₄ is
 thiazol-2-yl. In one aspect, R₄ is thiazol-5-yl. In one aspect, R₄ is
 [1,3,4]oxadiazol-2-yl. In one aspect, R₄ is trifluoromethyl. In one aspect, R₄ is
 10 pyridin-3-yl. In one aspect, R₄ is pyridin-2-yl. In one aspect, R₄ is
 -C(O)OCH₂CH₂CH₃, -C(O)OCH₂CH₃, or -C(O)OCH₃. In one aspect, R₄ is
 -C(O)OCH₃.

15 In one embodiment, in compounds represented by formula (I) or (II), R₄ is a
 biostere of an ester, amide, or carboxylic acid.

In one embodiment, in compounds represented by formula (I) or (II), R₃ is -H,
 chloro, fluoro, bromo, methyl, or methoxy. In one aspect, R₃ is -H, chloro,
 methyl, or methoxy. In one aspect, R₃ is -H. In one aspect, R₃ is chloro. In one
 20 aspect, R₃ is fluoro. In one aspect, R₃ is bromo. In one aspect, R₃ is methyl.
 In one aspect, R₃ is methoxy.

In one embodiment, in compounds represented by formula (I), Y is an
 optionally substituted phenyl, an optionally substituted pyridinyl, an optionally
 25 substituted pyrazolyl, an optionally substituted pyridazinyl, an optionally
 substituted isothiazolyl, an optionally substituted thiazolyl, an optionally
 substituted triazolyl, an optionally substituted oxadiazolyl, an optionally
 substituted oxazolyl, an optionally substituted isoxazolyl, an optionally
 substituted thiadiazolyl, or an optionally substituted thiophenyl. In another
 30 aspect, Y is an optionally substituted phenyl, an optionally substituted pyridinyl,
 an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an
 optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally
 substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally

substituted isoxazolyl, an optionally substituted thiadiazolyl, or an optionally substituted thiophenyl. In one aspect, Y is an optionally substituted phenyl or an optionally substituted thiadiazolyl. In one aspect, Y is an optionally substituted phenyl. In one aspect, Y is a phenyl substituted with one to two substituents. In one aspect, Y is a phenyl substituted with one to two substituents, wherein the one to two substituents are each independently a lower alkyl or a halo. In one aspect, Y is 2,6-difluorophenyl or 4-methyl-1,2,3-thiadiazol-5-yl. In one aspect, Y is a difluorophenyl. In one aspect, Y is 2,6-difluorophenyl. In one aspect, Y is 4-methyl-1,2,3-thiadiazol-5-yl. In one aspect, Y is an optionally substituted pyridinyl. In one aspect, Y is an optionally substituted triazolyl. In one aspect, Y is an optionally substituted oxadiazolyl. In one aspect, Y is an optionally substituted thiazolyl. In one aspect, Y is an optionally substituted oxazolyl. In one aspect, Y is an optionally substituted isoxazolyl. In one aspect, Y is a mono-substituted pyridinyl. In one aspect, Y is a methyl pyridin-2yl or a fluoro pyridin-2yl.

In one embodiment, in compounds represented by formula (I), L is a linker selected from the group consisting of -NHCH₂-, -CH₂NH-, -C(O)-, -NH-C(O)-, -C(O)-NH-, -OC(O)-, -C(O)O-, -C(S)-, -NH-C(S)-, -C(S)-NH-. In one aspect, L is -NHCH₂-. In one aspect, L is -CH₂NH-. In one aspect, L is -C(O)-. In one aspect, L is -C(O)-NH-. In one aspect, L is -OC(O)-. In one aspect, L is -C(O)O-. In one aspect, L is -C(S)-. In one aspect, L is -NH-C(S)-. In one aspect, L is -C(S)-NH-.

In one embodiment, in compounds represented by formula (I), L is -NR-C(O)-. In one aspect, L is -NH-C(O)-.

In one embodiment, in compounds represented by formula (I), Y is an optionally substituted phenyl, an optionally substituted pyridinyl, an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally substituted

thiadiazolyl, or an optionally substituted thiophenyl; R₄ is -C(O)OR₄, a haloalkyl, an optionally substituted oxazolyl, an optionally substituted thiazolyl, an optionally substituted imidazolyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyrrolyl, an optionally substituted thiophenyl, an optionally substituted furanyl, an optionally substituted thiadiazolyl, an optionally substituted oxadiazolyl, or an optionally substituted tetrazolyl; one of X₁ or X₂ is CH; and n is 0. In one aspect, L is -NH-C(O)-. In one aspect, R₃ is -H, chloro, fluoro, bromo, methyl, or methoxy. In one aspect, R₄ is oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, thiazol-5-yl, [1,3,4]oxadiazol-2-yl, trifluoromethyl, pyridin-3-yl, pyridin-2-yl, -C(O)OCH₂CH₂CH₃, -C(O)OCH₂CH₃, or -C(O)OCH₃. In one aspect, Y is a difluorophenyl. In one aspect, Y is 4-methyl-1,2,3-thiadiazol-5-yl.

In one embodiment, in compounds represented by formula (I), Y is 2,6-difluorophenyl or 4-methyl-1,2,3-thiadiazol-5-yl; L is -NH-C(O)-; one of X₁ or X₂ is CH; R₃ is -H, chloro, methyl, or methoxy; R₄ is oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, trifluoromethyl, or -C(O)OCH₃; and n is 0.

In one embodiment, in compounds represented by formula (II), Y₁ is an optionally substituted phenyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally substituted isoxazolyl, an optionally substituted thiadiazolyl, or an optionally substituted thiophenyl. In another aspect, Y₁ is an optionally substituted phenyl, an optionally substituted pyridinyl, an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally substituted thiadiazolyl, or an optionally substituted thiophenyl. In one aspect, Y₁ is an optionally substituted phenyl or an optionally substituted thiadiazolyl. In one aspect, Y₁ is an optionally substituted phenyl. In one aspect, Y₁ is a phenyl substituted with one to two substituents.

In one aspect, Y₁ is a phenyl substituted with one to two substituents, wherein the one to two substituents are each independently a lower alkyl or a halo. In one aspect, Y₁ is 2,6-difluorophenyl or 4-methyl-1,2,3-thiadiazol-5-yl. In one aspect, Y₁ is a difluorophenyl. In one aspect, Y₁ is 2,6-difluorophenyl. In one aspect, Y₁ is 4-methyl-1,2,3-thiadiazol-5-yl. In one aspect, Y₁ is an optionally substituted pyridinyl. In one aspect, Y₁ is an optionally substituted triazolyl. In one aspect, Y₁ is an optionally substituted oxadiazolyl. In one aspect, Y₁ is an optionally substituted thiazolyl. In one aspect, Y₁ is an optionally substituted oxazolyl. In one aspect, Y₁ is a mono-substituted pyridinyl. In one aspect, Y₁ is a methyl pyridin-2-yl or a fluoro pyridin-2-yl.

In one embodiment, in compounds represented by formula (II), L₁ is a linker selected from the group consisting of -NHCH₂-, -CH₂NH-, -NH-C(O)-, -OC(O)-, -C(O)O-, -C(S)-, -NH-C(S)-, or -C(S)-NH-. In one aspect, L₁ is -NHCH₂-. In one aspect, L₁ is -CH₂NH-. In one aspect, L₁ is -OC(O)-. In one aspect, L₁ is -C(O)O-. In one aspect, L₁ is -C(S)-. In one aspect, L₁ is -NH-C(S)-. In one aspect, L₁ is -C(S)-NH-.

In one embodiment, in compounds represented by formula (II), L₁ is -NR-C(O)-. In one aspect, L₁ is -NH-C(O)-.

In one embodiment, in compounds represented by formula (II), Y₁ is an optionally substituted phenyl, an optionally substituted pyridinyl, an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally substituted thiadiazolyl, or an optionally substituted thiophenyl; R₄ is -C(O)OR₄, a haloalkyl, an optionally substituted oxazolyl, an optionally substituted thiazolyl, an optionally substituted imidazolyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyrrolyl, an optionally substituted thiophenyl, an optionally substituted furanyl, an optionally substituted thiadiazolyl, an optionally substituted oxadiazolyl, or an optionally substituted tetrazolyl; one of X₁ or X₂ is CH; and n is 0. In one aspect, L₁ is

-NH-C(O)-. In one aspect, R₃ is -H, chloro, fluoro, bromo, methyl, or methoxy.
 In one aspect, R₄ is oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, thiazol-5-yl,
 [1,3,4]oxadiazol-2-yl, trifluoromethyl, pyridin-3-yl, pyridin-2-yl,
 -C(O)OCH₂CH₂CH₃, -C(O)OCH₂CH₃, or -C(O)OCH₃. In one aspect, Y₁ is a
 5 difluorophenyl. In one aspect, Y₁ is 4-methyl-1,2,3-thiadiazol-5-yl.

In one embodiment, in compounds represented by formula (II), Y₁ is
 2,6-difluorophenyl or 4-methyl-1,2,3-thiadiazol-5-yl; L₁ is -NH-C(O)-; one of X₁
 or X₂ is CH; R₃ is -H, chloro, methyl, or methoxy; R₄ is oxazol-2-yl, oxazol-5-yl,
 10 thiazol-2-yl, trifluoromethyl, or -C(O)OCH₃; and n is 0.

In another embodiment, the invention relates to compounds selected from the group consisting of:

N-[6-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-3-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-3-methyl-isonicotinamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-3-fluoro-isonicotinamide;
 2,6-Difluoro-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester;
 3-[6-(2,6-Difluoro-benzoylamino)-pyridin-3-yl]-4-methyl-benzoic acid methyl ester;
 N-[5-(2-Chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-6-methyl-pyridin-2-yl]-2,6-difluoro-
 benzamide;
 N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-4-methyl-pyridin-2-yl]-2,6-difluoro-
 benzamide;
 3-[5-(2,6-Difluoro-benzoylamino)-pyridin-2-yl]-4-methyl-benzoic acid methyl ester;
 2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[4-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[6-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-trifluoromethyl-phenyl)-
 pyridin-2-yl]-amide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-thiazol-2-yl-
 phenyl)-pyridin-2-yl]-amide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-methyl-5-oxazol-2-yl-
 phenyl)-pyridin-2-yl]-amide;

4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide;

2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide hydrochloride;

4-methyl-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-2-yl)-1,2,3-thiadiazole-5-carboxamide;

3-(6-Benzoylamino-pyridin-3-yl)-4-chloro-benzoic acid methyl ester;

4-Chloro-3-[6-(2-methyl-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester;

4-Chloro-3-{6-[(3-methyl-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(3-fluoro-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(3-chloro-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-[1,2,3]oxadiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(5-methyl-3H-[1,2,3]triazole-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-isothiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-thiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-oxazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-isoxazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(2-methyl-2H-pyrazole-3-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-fluoro-isothiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

3-Methyl-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-isonicotinamide;

3-Methyl-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-isonicotinamide;

2,6-Difluoro-N-[5-methyl-6-(3-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide;

2,6-Difluoro-N-[4-methyl-5-(3-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;

2,6-Difluoro-N-[6-methyl-5-(3-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;

2,6-Difluoro-N-[5-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;

3-Methyl-N-[5-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-isonicotinamide;

3-Methyl-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-isonicotinamide;
 2,6-Difluoro-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-benzamide
 hydrochloride;
 N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-3-methyl-isonicotinamide;
 N-[6-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-3-methyl-isonicotinamide;
 N-[6-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-2,6-difluoro-benzamide;
 3-[5-(2,6-Difluoro-benzoylamino)-pyridin-2-yl]-4-methyl-benzoic acid ethyl ester;
 3-[6-(2,6-Difluoro-benzoylamino)-pyridin-3-yl]-4-methyl-benzoic acid ethyl ester;
 2,6-Difluoro-N-[5-(2-methyl-5-oxazol-5-yl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-oxazol-5-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[6-(5-isoxazol-5-yl-2-methyl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[5-(5-isoxazol-5-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-[2-methyl-5-(3-methyl-isoxazol-5-yl)-phenyl]-pyridin-2-yl]-
 benzamide;
 2,6-Difluoro-N-[6-[2-methyl-5-(3-methyl-isoxazol-5-yl)-phenyl]-pyridin-3-yl]-
 benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-pyridin-3-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-pyridin-2-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-[1,3,4]oxadiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-[1,3,4]thiadiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid propyl ester;
 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid 2-methoxy-ethyl
 ester;
 2,6-Difluoro-N-[5-(5-furan-2-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-thiophen-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(5-furan-3-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-thiophen-3-yl-phenyl)-pyridin-2-yl]-benzamide;
 N-(5-(2-methoxy-5-(oxazol-5-yl)phenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-
 carboxamide;
 N-(5-(5-(isoxazol-5-yl)-2-methylphenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-carb
 oxamide
 2,6-difluoro-N-(5-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl) pyridin-2-yl)benzamide;
 2,6-difluoro-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl) pyridin-2-yl) benzamide;
 N-(5-(2-chloro-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)-2,6-difluorobenzamide;
 N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide;

N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt;

2,6-Difluoro-N-(6-(2-methyl-5-(thiazol-2-yl)phenyl)pyridin-3-yl)benzamide;

2,6-Difluoro-N-(6-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-3-yl)benzamide;

N-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide;

N-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt;

2,6-Difluoro-N-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide;

2,6-Difluoro-N-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide hydrochloric acid salt; or

4-chloro-2-fluoro-N-(6-(2-methyl-5-(oxazol-2-yl)phenyl)pyridin-3-yl)benzamide;

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

5 All of the features, specific embodiments and particular substituents disclosed herein may be combined in any combination. Each feature, embodiment or substituent disclosed in this specification may be replaced by an alternative feature, embodiment or substituent serving the same, equivalent, or similar purpose. In the case of chemical compounds, specific values for variables (e.g., values shown in the exemplary compounds disclosed herein) in any chemical formula disclosed herein can be combined in any combination
10 resulting in a stable structure. Furthermore, specific values (whether preferred or not) for substituents in one type of chemical structure may be combined with values for other substituents (whether preferred or not) in the same or different type of chemical structure. Thus, unless expressly stated otherwise, each feature, embodiment or substituent disclosed is only an example of a generic
15 series of equivalent or similar features, embodiments or substituents.

In another embodiment, the invention relates to pharmaceutical compositions that comprise a compound of any one of formulas (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, as an
20 active ingredient, and a pharmaceutically acceptable carrier or vehicle. The compositions are useful for immunosuppression or to treat or prevent inflammatory conditions, allergic conditions and immune disorders.

In another embodiment, the invention relates to methods for immunosuppression or for treating or preventing inflammatory conditions, immune disorders, or allergic disorders in a patient in need thereof comprising administering an effective amount of a compound represented by any one of
5 formulas (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

In another embodiment, the invention relates to methods for immunosuppression or for treating or preventing inflammatory conditions,
10 immune disorders, or allergic disorders in a patient in need thereof comprising administering an effective amount of a pharmaceutical composition that comprises a compound represented by any one of formulas (I) through (II), or in or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

15 In another embodiment, compounds of any one of formulas (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, are particularly useful inhibiting immune cell (e.g., T-cells and/or B-cells) activation (e.g., activation in response to an antigen) and/or T cell
20 and/or B cell proliferation. Indicators of immune cell activation include secretion of IL-2 by T cells, proliferation of T cells and/or B cells, and the like. In one embodiment, a compound of any one of formulas (I) through (II) or Table 1, inhibits immune cell activation and/or T cell and/or B cell proliferation in a mammal (e.g., a human).

25 In another embodiment, compounds of any one of formula (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can inhibit the production of certain cytokines that regulate immune cell activation. For example, compounds of any one of formulas (I) through (II), or
30 Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can inhibit the production of IL-2, IL-4, IL-5, IL-13, GM-CSF, IFN- γ , TNF- α and combinations thereof. In one embodiment, a compound of any one



of formulas (I) through (II), or Table 1, inhibits cytokine production in a mammal (e.g., a human).









In another embodiment, compounds of any one of formulas (I) through (II), or
 5 Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, can modulate the activity of one or more ion channel involved in activation of immune cells, such as CRAC ion channels. In one embodiment, a compound of any one of formulas (I) through (II) or Table 1 can inhibit the influx of calcium ions into an immune cell (e.g., T cells and/or B cells) by
 10 inhibiting the action of CRAC ion channels. In general, a decrease in I_{CRAC} current upon contacting a cell with a compound is one indicator that the compound inhibitions CRAC ion channels. I_{CRAC} current can be measured, for example, using a patch clamp technique, which is described in more detail in the examples below. In one embodiment, a compound of any one of formulas
 15 (I) through (II) or Table 1 modulates an ion channel in a mammal (e.g., a human).










EXEMPLARY COMPOUNDS OF THE INVENTION

20 Exemplary compounds of the invention are depicted in Table 1 below.











Table 1








Compound No.	Structure	Chemical Name
1		N-[6-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-3-yl]-2,6-difluorobenzamide
2		N-[5-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-2,6-difluorobenzamide









3		N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-3-methyl-isonicotinamide
4		N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-3-fluoro-isonicotinamide
5		2,6-Difluoro-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide
6		4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester
7		3-[6-(2,6-Difluoro-benzoylamino)-pyridin-3-yl]-4-methyl-benzoic acid methyl ester
8		N-[5-(2-Chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide
9		N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-6-methyl-pyridin-2-yl]-2,6-difluoro-benzamide
10		N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide








11		N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-4-methyl-pyridin-2-yl]-2,6-difluoro-benzamide
12		3-[5-(2,6-Difluoro-benzoylamino)-pyridin-2-yl]-4-methyl-benzoic acid methyl ester
13		2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide
14		2,6-Difluoro-N-[4-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide
15		2,6-Difluoro-N-[6-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide
16		4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-amide
17		4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-amide
18		4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide
19		4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide








20		2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide hydrochloride
21		4-methyl-N-(5-(2-methyl-5-oxazol-5-yl)phenyl)pyridin-2-yl)-1,2,3-thiadiazole-5-carboxamide
22		3-(6-Benzoylamino-pyridin-3-yl)-4-chloro-benzoic acid methyl ester
23		4-Chloro-3-[6-(2-methyl-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester
24		4-Chloro-3-{6-[(3-methyl-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
25		4-Chloro-3-{6-[(3-fluoro-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
26		4-Chloro-3-{6-[(3-chloro-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
27		4-Chloro-3-{6-[(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
28		4-Chloro-3-{6-[(4-methyl-[1,2,3]oxadiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester







29		4-Chloro-3-{6-[(5-methyl-3H-[1,2,3]triazole-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
30		4-Chloro-3-{6-[(4-methyl-isothiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
31		4-Chloro-3-{6-[(4-methyl-thiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
32		4-Chloro-3-{6-[(4-methyl-oxazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
33		4-Chloro-3-{6-[(4-methyl-isoxazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
34		4-Chloro-3-{6-[(2-methyl-2H-pyrazole-3-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
35		4-Chloro-3-{6-[(4-fluoro-isothiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester
36		3-Methyl-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-isonicotinamide
37		3-Methyl-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-isonicotinamide
38		2,6-Difluoro-N-[5-methyl-6-(3-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide







39		2,6-Difluoro-N-[4-methyl-5-(3-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide
40		2,6-Difluoro-N-[6-methyl-5-(3-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide
41		2,6-Difluoro-N-[5-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide
42		3-Methyl-N-[5-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-isonicotinamide
43		3-Methyl-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-isonicotinamide
44		2,6-Difluoro-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-benzamide
45		2,6-Difluoro-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-benzamide hydrochloride


46		N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-3-methyl-isonicotinamide
47		N-[6-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-3-methyl-isonicotinamide
48		N-[6-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-2,6-difluorobenzamide
49		3-[5-(2,6-Difluorobenzoylamino)-pyridin-2-yl]-4-methylbenzoic acid ethyl ester
50		3-[6-(2,6-Difluorobenzoylamino)-pyridin-3-yl]-4-methylbenzoic acid ethyl ester
51		2,6-Difluoro-N-[5-(2-methyl-5-oxazol-5-yl-phenyl)-pyridin-2-yl]-benzamide
52		2,6-Difluoro-N-[6-(2-methyl-5-oxazol-5-yl-phenyl)-pyridin-3-yl]-benzamide
53		2,6-Difluoro-N-[6-(5-isoxazol-5-yl-2-methyl-phenyl)-pyridin-3-yl]-benzamide

54		2,6-Difluoro-N-[5-(5-isoxazol-5-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide
55		2,6-Difluoro-N-{5-[2-methyl-5-(3-methyl-isoxazol-5-yl)-phenyl]-pyridin-2-yl}-benzamide
56		2,6-Difluoro-N-{6-[2-methyl-5-(3-methyl-isoxazol-5-yl)-phenyl]-pyridin-3-yl}-benzamide
57		2,6-Difluoro-N-[6-(2-methyl-5-pyridin-3-yl-phenyl)-pyridin-3-yl]-benzamide
58		2,6-Difluoro-N-[6-(2-methyl-5-pyridin-2-yl-phenyl)-pyridin-3-yl]-benzamide
59		2,6-Difluoro-N-[5-(2-methyl-5-[1,3,4]oxadiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide
60		2,6-Difluoro-N-[5-(2-methyl-5-[1,3,4]thiadiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide

61		4-Chloro-3-[6-(2,6-difluorobenzoylamino)-pyridin-3-yl]-benzoic acid propyl ester
62		4-Chloro-3-[6-(2,6-difluorobenzoylamino)-pyridin-3-yl]-benzoic acid 2-methoxy-ethyl ester
63		2,6-Difluoro-N-[5-(5-furan-2-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide
64		2,6-Difluoro-N-[5-(2-methyl-5-thiophen-2-yl-phenyl)-pyridin-2-yl]-benzamide
65		2,6-Difluoro-N-[5-(5-furan-3-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide
66		2,6-Difluoro-N-[5-(2-methyl-5-thiophen-3-yl-phenyl)-pyridin-2-yl]-benzamide
67		N-(5-(2-methoxy-5-(oxazol-5-yl)phenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-carboxamide

68		N-(5-(5-(isoxazol-5-yl)-2-methylphenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-carboxamide
69		2,6-difluoro-N-(5-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)benzamide
70		2,6-difluoro-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-2-yl)benzamide
71		N-(5-(2-chloro-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)-2,6-difluorobenzamide
72		N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide
73		N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt

74		2,6-Difluoro-N-(6-(2-methyl-5-(thiazol-2-yl)phenyl)pyridin-3-yl)benzamide
75		2,6-Difluoro-N-(6-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-3-yl)benzamide
76		N-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide
77	 HCl	N-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt
78		2,6-Difluoro-N-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide
79	 HCl	2,6-Difluoro-N-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide hydrochloric acid salt

80		4-chloro-2-fluoro-N-(6-(2-methyl-5-(oxazol-2-yl)phenyl)pyridin-3-yl)benzamide
----	--	---

MECHANISM OF ACTION

Activation of T-lymphocytes in response to an antigen is dependent on calcium ion oscillations. Calcium ion oscillations in T-lymphocytes are triggered through stimulation of the T-cell antigen receptor, and involve calcium ion influx through the stored-operated Ca^{2+} -release-activated Ca^{2+} (CRAC) channel. Although the molecular structure of the CRAC ion channel has not been identified, a detailed electrophysiological profile of the channel exist. Thus, inhibition of CRAC ion channels can be measured by measuring inhibition of the I_{CRAC} current. Calcium ion oscillations in T-cells have been implicated in the activation of several transcription factors (e.g., NFAT, Oct/Oap and NFkB) which are critical for T-cell activation (Lewis, *Biochemical Society Transactions* (2003), 31:925-929, the entire teachings of which are incorporated herein by reference). Without wishing to be bound by any theory, it is believed that because the compounds of the invention inhibit the activity of CRAC ion channels, they inhibit immune cell activation.

METHODS OF TREATMENT AND PREVENTION

In accordance with the invention, an effective amount of a compound of any one of formulas (I) through (II) or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, and prodrug thereof, or a pharmaceutical composition comprising a compound of any one of formulas (I) through (II) or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, and prodrug thereof, is administered to a patient in need of immunosuppression or in need of treatment or prevention of an inflammatory condition, an immune disorder, or an allergic disorder. Such patients may be treatment naïve or may experience partial or no response to conventional therapies.

Responsiveness of a particular inflammatory condition, immune disorder, or allergic disorder in a subject can be measured directly (e.g., measuring blood levels of inflammatory cytokines (such as IL-2, IL-4, IL-5, IL-13, GM-CSF, 5 TNF- α , IFN- γ and the like) after administration of a compound of this invention), or can be inferred based on an understanding of disease etiology and progression. The compounds of any one of formulas (I) through (II), or Table 1, or pharmaceutically acceptable salts, solvates, clathrates, and prodrugs thereof can be assayed *in vitro* or *in vivo*, for the desired therapeutic or 10 prophylactic activity, prior to use in humans. For example, known animal models of inflammatory conditions, immune disorders, or allergic disorders can be used to demonstrate the safety and efficacy of compounds of this invention.

PHARMACEUTICAL COMPOSITIONS AND DOSAGE FORMS

15 Pharmaceutical compositions and dosage forms of the invention comprise one or more active ingredients in relative amounts and formulated in such a way that a given pharmaceutical composition or dosage form can be used for immunosuppression or to treat or prevent inflammatory conditions, immune disorders, and allergic disorders. Preferred pharmaceutical compositions and 20 dosage forms comprise a compound of any one of formulas (I) through (II), or Table 1, or a pharmaceutically acceptable prodrug, salt, solvate, or clathrate thereof, optionally in combination with one or more additional active agents.

Single unit dosage forms of the invention are suitable for oral, mucosal (e.g., 25 nasal, sublingual, vaginal, buccal, or rectal), parenteral (e.g., subcutaneous, intravenous, bolus injection, intramuscular, or intraarterial), or transdermal administration to a patient. Examples of dosage forms include, but are not limited to: tablets; caplets; capsules, such as soft elastic gelatin capsules; cachets; troches; lozenges; dispersions; suppositories; ointments; cataplasms 30 (poultices); pastes; powders; dressings; creams; plasters; solutions; patches; aerosols (e.g., nasal sprays or inhalers); gels; liquid dosage forms suitable for oral or mucosal administration to a patient, including suspensions (e.g., aqueous or non-aqueous liquid suspensions, oil-in-water emulsions, or a

water-in-oil liquid emulsions), solutions, and elixirs; liquid dosage forms suitable for parenteral administration to a patient; and sterile solids (e.g., crystalline or amorphous solids) that can be reconstituted to provide liquid dosage forms suitable for parenteral administration to a patient.

5

The composition, shape, and type of dosage forms of the invention will typically vary depending on their use. For example, a dosage form suitable for mucosal administration may contain a smaller amount of active ingredient(s) than an oral dosage form used to treat the same indication. This aspect of the invention will be readily apparent to those skilled in the art. See, e.g., Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing, Easton PA.

10

Typical pharmaceutical compositions and dosage forms comprise one or more excipients. Suitable excipients are well known to those skilled in the art of pharmacy, and non-limiting examples of suitable excipients are provided herein. Whether a particular excipient is suitable for incorporation into a pharmaceutical composition or dosage form depends on a variety of factors well known in the art including, but not limited to, the way in which the dosage form will be administered to a patient. For example, oral dosage forms such as tablets may contain excipients not suited for use in parenteral dosage forms.

15

20

The suitability of a particular excipient may also depend on the specific active ingredients in the dosage form. For example, the decomposition of some active ingredients can be accelerated by some excipients such as lactose, or when exposed to water. Active ingredients that comprise primary or secondary amines (e.g., N-desmethylvenlafaxine and N,N-didesmethylvenlafaxine) are particularly susceptible to such accelerated decomposition. Consequently, this invention encompasses pharmaceutical compositions and dosage forms that contain little, if any, lactose. As used herein, the term "lactose-free" means that the amount of lactose present, if any, is insufficient to substantially increase the degradation rate of an active ingredient. Lactose-free compositions of the invention can comprise excipients that are well known in the art and are listed, for example, in the U.S. Pharmacopia (USP) SP (XXI)/NF (XVI). In general,

25

30

lactose-free compositions comprise active ingredients, a binder/filler, and a lubricant in pharmaceutically compatible and pharmaceutically acceptable amounts. Preferred lactose-free dosage forms comprise active ingredients, microcrystalline cellulose, pre-gelatinized starch, and magnesium stearate.

5

This invention further encompasses anhydrous pharmaceutical compositions and dosage forms comprising active ingredients, since water can facilitate the degradation of some compounds. For example, the addition of water (*e.g.*, 5%) is widely accepted in the pharmaceutical arts as a means of simulating long-term storage in order to determine characteristics such as shelf-life or the stability of formulations over time. See, *e.g.*, Jens T. Carstensen (1995) *Drug Stability: Principles & Practice*, 2d. Ed., Marcel Dekker, NY, NY, 379-80. In effect, water and heat accelerate the decomposition of some compounds. Thus, the effect of water on a formulation can be of great significance since moisture and/or humidity are commonly encountered during manufacture, handling, packaging, storage, shipment, and use of formulations.

Anhydrous pharmaceutical compositions and dosage forms of the invention can be prepared using anhydrous or low moisture containing ingredients and low moisture or low humidity conditions. Pharmaceutical compositions and dosage forms that comprise lactose and at least one active ingredient that comprises a primary or secondary amine are preferably anhydrous if substantial contact with moisture and/or humidity during manufacturing, packaging, and/or storage is expected.

25

An anhydrous pharmaceutical composition should be prepared and stored such that its anhydrous nature is maintained. Accordingly, anhydrous compositions are preferably packaged using materials known to prevent exposure to water such that they can be included in suitable formulary kits. Examples of suitable packaging include, but are not limited to, hermetically sealed foils, plastics, unit dose containers (*e.g.*, vials), blister packs, and strip packs.

30

The invention further encompasses pharmaceutical compositions and dosage forms that comprise one or more compounds that reduce the rate by which an active ingredient will decompose. Such compounds, which are referred to herein as "stabilizer" include, but are not limited to, antioxidants such as ascorbic acid, pH buffers, or salt buffers.

Like the amounts and types of excipients, the amounts and specific types of active ingredients in a dosage form may differ depending on factors such as, but not limited to, the route by which it is to be administered to patients. However, typical dosage forms of the invention comprise a compound of any one of formulas (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof in an amount of from about 1 mg to about 1000 mg, preferably in an amount of from about 50 mg to about 500 mg, and most preferably in an amount of from about 75 mg to about 350 mg. The typical total daily dosage of a compound of any one of formulas (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof can range from about 1 mg to about 5000 mg per day, preferably in an amount from about 50 mg to about 1500 mg per day, more preferably from about 75 mg to about 1000 mg per day. It is within the skill of the art to determine the appropriate dose and dosage form for a given patient.

ORAL DOSAGE FORMS

Pharmaceutical compositions of the invention that are suitable for oral administration can be presented as discrete dosage forms, such as, but are not limited to, tablets (e.g., chewable tablets), caplets, capsules, and liquids (e.g., flavored syrups). Such dosage forms contain predetermined amounts of active ingredients, and may be prepared by methods of pharmacy well known to those skilled in the art. See *generally*, Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing, Easton PA.

Typical oral dosage forms of the invention are prepared by combining the active ingredient(s) in an admixture with at least one excipient according to conventional pharmaceutical compounding techniques. Excipients can take a

wide variety of forms depending on the form of preparation desired for administration. For example, excipients suitable for use in oral liquid or aerosol dosage forms include, but are not limited to, water, glycols, oils, alcohols, flavoring agents, preservatives, and coloring agents. Examples of excipients
5 suitable for use in solid oral dosage forms (e.g., powders, tablets, capsules, and caplets) include, but are not limited to, starches, sugars, micro-crystalline cellulose, diluents, granulating agents, lubricants, binders, and disintegrating agents.

10 Because of their ease of administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid excipients are employed. If desired, tablets can be coated by standard aqueous or nonaqueous techniques. Such dosage forms can be prepared by any of the methods of pharmacy. In general, pharmaceutical compositions and dosage
15 forms are prepared by uniformly and intimately admixing the active ingredients with liquid carriers, finely divided solid carriers, or both, and then shaping the product into the desired presentation if necessary.

For example, a tablet can be prepared by compression or molding.
20 Compressed tablets can be prepared by compressing in a suitable machine the active ingredients in a free-flowing form such as powder or granules, optionally mixed with an excipient. Molded tablets can be made by molding in a suitable machine a mixture of the powdered compound moistened with an inert liquid diluent.

25 Examples of excipients that can be used in oral dosage forms of the invention include, but are not limited to, binders, fillers, disintegrants, and lubricants. Binders suitable for use in pharmaceutical compositions and dosage forms include, but are not limited to, corn starch, potato starch, or other starches,
30 gelatin, natural and synthetic gums such as acacia, sodium alginate, alginic acid, other alginates, powdered tragacanth, guar gum, cellulose and its derivatives (e.g., ethyl cellulose, cellulose acetate, carboxymethyl cellulose calcium, sodium carboxymethyl cellulose), polyvinyl pyrrolidone, methyl

cellulose, pre-gelatinized starch, hydroxypropyl methyl cellulose, (e.g., Nos. 2208, 2906, 2910), microcrystalline cellulose, and mixtures thereof.

5 Suitable forms of microcrystalline cellulose include, but are not limited to, the materials sold as AVICEL-PH-101, AVICEL-PH-103 AVICEL RC-581, AVICEL-PH-105 (available from FMC Corporation, American Viscose Division, Avicel Sales, Marcus Hook, PA), and mixtures thereof. One specific binder is a mixture of microcrystalline cellulose and sodium carboxymethyl cellulose sold as AVICEL RC-581. Suitable anhydrous or low moisture excipients or
10 additives include AVICEL-PH-103J and Starch 1500 LM.

Examples of fillers suitable for use in the pharmaceutical compositions and dosage forms disclosed herein include, but are not limited to, talc, calcium carbonate (e.g., granules or powder), microcrystalline cellulose, powdered
15 cellulose, dextrates, kaolin, mannitol, silicic acid, sorbitol, starch, pre-gelatinized starch, and mixtures thereof. The binder or filler in pharmaceutical compositions of the invention is typically present in from about 50 to about 99 weight percent of the pharmaceutical composition or dosage form.

20

Disintegrants are used in the compositions of the invention to provide tablets that disintegrate when exposed to an aqueous environment. Tablets that contain too much disintegrant may disintegrate in storage, while those that contain too little may not disintegrate at a desired rate or under the desired
25 conditions. Thus, a sufficient amount of disintegrant that is neither too much nor too little to detrimentally alter the release of the active ingredients should be used to form solid oral dosage forms of the invention. The amount of disintegrant used varies based upon the type of formulation, and is readily discernible to those of ordinary skill in the art. Typical pharmaceutical
30 compositions comprise from about 0.5 to about 15 weight percent of disintegrant, preferably from about 1 to about 5 weight percent of disintegrant.

Disintegrants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, agar-agar, alginic acid, calcium carbonate, microcrystalline cellulose, croscarmellose sodium, crospovidone, polacrillin potassium, sodium starch glycolate, potato or tapioca starch, other starches, pre-gelatinized starch, other starches, clays, other alginates, other celluloses, gums, and mixtures thereof.

Lubricants that can be used in pharmaceutical compositions and dosage forms of the invention include, but are not limited to, calcium stearate, magnesium stearate, mineral oil, light mineral oil, glycerin, sorbitol, mannitol, polyethylene glycol, other glycols, stearic acid, sodium lauryl sulfate, talc, hydrogenated vegetable oil (e.g., peanut oil, cottonseed oil, sunflower oil, sesame oil, olive oil, corn oil, and soybean oil), zinc stearate, ethyl oleate, ethyl laureate, agar, and mixtures thereof. Additional lubricants include, for example, a syloid silica gel (AEROSIL 200, manufactured by W.R. Grace Co. of Baltimore, MD), a coagulated aerosol of synthetic silica (marketed by Degussa Co. of Plano, TX), CAB-O-SIL (a pyrogenic silicon dioxide product sold by Cabot Co. of Boston, MA), and mixtures thereof. If used at all, lubricants are typically used in an amount of less than about 1 weight percent of the pharmaceutical compositions or dosage forms into which they are incorporated.

CONTROLLED RELEASE DOSAGE FORMS

Active ingredients of the invention can be administered by controlled release means or by delivery devices that are well known to those of ordinary skill in the art. Examples include, but are not limited to, those described in U.S. Patent Nos.: 3,845,770; 3,916,899; 3,536,809; 3,598,123; and 4,008,719, 5,674,533, 5,059,595, 5,591,767, 5,120,548, 5,073,543, 5,639,476, 5,354,556, and 5,733,566, each of which is incorporated herein by reference. Such dosage forms can be used to provide slow or controlled-release of one or more active ingredients using, for example, hydropropylmethyl cellulose, other polymer matrices, gels, permeable membranes, osmotic systems, multilayer coatings, microparticles, liposomes, microspheres, or a combination thereof to provide the desired release profile in varying proportions. Suitable controlled-release

formulations known to those of ordinary skill in the art, including those described herein, can be readily selected for use with the active ingredients of the invention. The invention thus encompasses single unit dosage forms suitable for oral administration such as, but not limited to, tablets, capsules, 5 gelcaps, and caplets that are adapted for controlled-release.

All controlled-release pharmaceutical products have a common goal of improving drug therapy over that achieved by their non-controlled counterparts. Ideally, the use of an optimally designed controlled-release preparation in 10 medical treatment is characterized by a minimum of drug substance being employed to cure or control the condition in a minimum amount of time. Advantages of controlled-release formulations include extended activity of the drug, reduced dosage frequency, and increased patient compliance. In addition, controlled-release formulations can be used to affect the time of onset 15 of action or other characteristics, such as blood levels of the drug, and can thus affect the occurrence of side (e.g., adverse) effects.

Most controlled-release formulations are designed to initially release an amount of drug (active ingredient) that promptly produces the desired 20 therapeutic effect, and gradually and continually release of other amounts of drug to maintain this level of therapeutic or prophylactic effect over an extended period of time. In order to maintain this constant level of drug in the body, the drug must be released from the dosage form at a rate that will replace the amount of drug being metabolized and excreted from the body. 25 Controlled-release of an active ingredient can be stimulated by various conditions including, but not limited to, pH, temperature, enzymes, water, or other physiological conditions or compounds.

A particular extended release formulation of this invention comprises a 30 therapeutically or prophylactically effective amount of a compound of formula (I) through (II), or Table 1, or a pharmaceutically acceptable salt, solvate, hydrate, clathrate, or prodrug thereof, in spheroids which further comprise microcrystalline cellulose and, optionally, hydroxypropylmethyl-cellulose

coated with a mixture of ethyl cellulose and hydroxypropylmethylcellulose. Such extended release formulations can be prepared according to U.S. Patent No. 6,274,171, the entire teachings of which are incorporated herein by reference.

5

A specific controlled-release formulation of this invention comprises from about 6% to about 40% a compound of any one of formulas (I) through (II), or Table 1 by weight, about 50% to about 94% microcrystalline cellulose, NF, by weight, and optionally from about 0.25% to about 1% by weight of hydroxypropyl-methylcellulose, USP, wherein the spheroids are coated with a film coating composition comprised of ethyl cellulose and hydroxypropylmethylcellulose.

10

PARENTERAL DOSAGE FORMS

Parenteral dosage forms can be administered to patients by various routes including, but not limited to, subcutaneous, intravenous (including bolus injection), intramuscular, and intraarterial. Because their administration typically bypasses patients' natural defenses against contaminants, parenteral dosage forms are preferably sterile or capable of being sterilized prior to administration to a patient. Examples of parenteral dosage forms include, but are not limited to, solutions ready for injection, dry products ready to be dissolved or suspended in a pharmaceutically acceptable vehicle for injection, suspensions ready for injection, and emulsions.

20

Suitable vehicles that can be used to provide parenteral dosage forms of the invention are well known to those skilled in the art. Examples include, but are not limited to: Water for Injection USP; aqueous vehicles such as, but not limited to, Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, and Lactated Ringer's Injection; water-miscible vehicles such as, but not limited to, ethyl alcohol, polyethylene glycol, and polypropylene glycol; and non-aqueous vehicles such as, but not limited to, corn oil, cottonseed oil, peanut oil, sesame oil, ethyl oleate, isopropyl myristate, and benzyl benzoate.

30

Compounds that increase the solubility of one or more of the active ingredients disclosed herein can also be incorporated into the parenteral dosage forms of the invention.

5

TRANSDERMAL, TOPICAL, AND MUCOSAL DOSAGE FORMS

Transdermal, topical, and mucosal dosage forms of the invention include, but are not limited to, ophthalmic solutions, sprays, aerosols, creams, lotions, ointments, gels, solutions, emulsions, suspensions, or other forms known to one of skill in the art. See, e.g., Remington's Pharmaceutical Sciences (1980 & 1990) 16th and 18th eds., Mack Publishing, Easton PA and Introduction to Pharmaceutical Dosage Forms (1985) 4th ed., Lea & Febiger, Philadelphia. Dosage forms suitable for treating mucosal tissues within the oral cavity can be formulated as mouthwashes or as oral gels. Further, transdermal dosage forms include "reservoir type" or "matrix type" patches, which can be applied to the skin and worn for a specific period of time to permit the penetration of a desired amount of active ingredients.

Suitable excipients (e.g., carriers and diluents) and other materials that can be used to provide transdermal, topical, and mucosal dosage forms encompassed by this invention are well known to those skilled in the pharmaceutical arts, and depend on the particular tissue to which a given pharmaceutical composition or dosage form will be applied. With that fact in mind, typical excipients include, but are not limited to, water, acetone, ethanol, ethylene glycol, propylene glycol, butane-1,3-diol, isopropyl myristate, isopropyl palmitate, mineral oil, and mixtures thereof to form lotions, tinctures, creams, emulsions, gels or ointments, which are non-toxic and pharmaceutically acceptable. Moisturizers or humectants can also be added to pharmaceutical compositions and dosage forms if desired. Examples of such additional ingredients are well known in the art. See, e.g., Remington's Pharmaceutical Sciences (1980 & 1990) 16th and 18th eds., Mack Publishing, Easton PA.

Depending on the specific tissue to be treated, additional components may be used prior to, in conjunction with, or subsequent to treatment with active ingredients of the invention. For example, penetration enhancers can be used to assist in delivering the active ingredients to the tissue. Suitable penetration
5 enhancers include, but are not limited to: acetone; various alcohols such as ethanol, oleyl, and tetrahydrofuryl; alkyl sulfoxides such as dimethyl sulfoxide; dimethyl acetamide; dimethyl formamide; polyethylene glycol; pyrrolidones such as polyvinylpyrrolidone; Kollidon grades (Povidone, Polyvidone); urea; and various water-soluble or insoluble sugar esters such as Tween 80
10 (polysorbate 80) and Span 60 (sorbitan monostearate).

The pH of a pharmaceutical composition or dosage form, or of the tissue to which the pharmaceutical composition or dosage form is applied, may also be adjusted to improve delivery of one or more active ingredients. Similarly, the
15 polarity of a solvent carrier, its ionic strength, or tonicity can be adjusted to improve delivery. Compounds such as stearates can also be added to pharmaceutical compositions or dosage forms to advantageously alter the hydrophilicity or lipophilicity of one or more active ingredients so as to improve delivery. In this regard, stearates can serve as a lipid vehicle for the
20 formulation, as an emulsifying agent or surfactant, and as a delivery-enhancing or penetration-enhancing agent. Different salts, hydrates or solvates of the active ingredients can be used to further adjust the properties of the resulting composition.

25 COMBINATION THERAPY

The methods for immunosuppression or for treating or preventing inflammatory conditions and immune disorders in a patient in need thereof can further comprise administering to the patient being administered a compound of this invention, an effective amount of one or more other active agents. Such active
30 agents may include those used conventionally for immunosuppression or for inflammatory conditions or immune disorders. These other active agents may also be those that provide other benefits when administered in combination with the compounds of this invention. For example, other therapeutic agents

may include, without limitation, steroids, non-steroidal anti-inflammatory agents, antihistamines, analgesics, immunosuppressive agents and suitable mixtures thereof. In such combination therapy treatment, both the compounds of this invention and the other drug agent(s) are administered to a subject (e.g.,
5 humans, male or female) by conventional methods. The agents may be administered in a single dosage form or in separate dosage forms. Effective amounts of the other therapeutic agents and dosage forms are well known to those skilled in the art. It is well within the skilled artisan's purview to determine the other therapeutic agent's optimal effective-amount range.

10

In one embodiment of the invention where another therapeutic agent is administered to a subject, the effective amount of the compound of this invention is less than its effective amount when the other therapeutic agent is not administered. In another embodiment, the effective amount of the
15 conventional agent is less than its effective amount when the compound of this invention is not administered. In this way, undesired side effects associated with high doses of either agent may be minimized. Other potential advantages (including without limitation improved dosing regimens and/or reduced drug cost) will be apparent to those of skill in the art.

20

In one embodiment relating to autoimmune and inflammatory conditions, the other therapeutic agent may be a steroid or a non-steroidal anti-inflammatory agent. Particularly useful non-steroidal anti-inflammatory agents, include, but are not limited to, aspirin, ibuprofen, diclofenac, naproxen, benoxaprofen,
25 flurbiprofen, fenoprofen, flubufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pramoprofen, muprofen, trioxaprofen, suprofen, aminoprofen, tiaprofenic acid, fluprofen, bucloxic acid, indomethacin, sulindac, tolmetin, zomepirac, tiopinac, zidometacin, acetaminophen, fentiazac, clidanac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid,
30 tolfenamic acid, diflunisal, flufenisal, piroxicam, sudoxicam, isoxicam; salicylic acid derivatives, including aspirin, sodium salicylate, choline magnesium trisalicylate, salsalate, diflunisal, salicylsalicylic acid, sulfasalazine, and olsalazin; para-aminophenol derivatives including acetaminophen and

phenacetin; indole and indene acetic acids, including indomethacin, sulindac, and etodolac; heteroaryl acetic acids, including tolmetin, diclofenac, and ketorolac; anthranilic acids (fenamates), including mefenamic acid, and meclofenamic acid; enolic acids, including oxicams (piroxicam, tenoxicam),
5 and pyrazolidinediones (phenylbutazone, oxyphenthartazone); and alkanones, including nabumetone and pharmaceutically acceptable salts thereof and mixtures thereof. For a more detailed description of the NSAIDs, see *Paul A. Insel, Analgesic-Antipyretic and Antiinflammatory Agents and Drugs Employed in the Treatment of Gout, in Goodman & Gilman's The Pharmacological Basis of Therapeutics* 617-57 (Perry B. Molinoff and Raymond W. Ruddon eds., 9th
10 ed 1996) and Glen R. Hanson, *Analgesic, Antipyretic and Anti-Inflammatory Drugs in Remington: The Science and Practice of Pharmacy Vol II* 1196-1221 (A.R. Gennaro ed. 19th ed. 1995) which are hereby incorporated by reference in their entireties.

15

Of particular relevance to allergic disorders, the other therapeutic agent may be an antihistamine. Useful antihistamines include, but are not limited to, loratadine, cetirizine, fexofenadine, desloratadine, diphenhydramine, chlorpheniramine, chlorcyclizine, pyrilamine, promethazine, terfenadine,
20 doxepin, carbinoxamine, clemastine, tripeleminamine, brompheniramine, hydroxyzine, cyclizine, meclizine, cyproheptadine, phenindamine, acrivastine, azelastine, levocabastine, and mixtures thereof. For a more detailed description of antihistamines, see Goodman & Gilman's *The Pharmacological Basis of Therapeutics* (2001) 651-57, 10th ed).

25

Immunosuppressive agents include glucocorticoids, corticosteroids (such as Prednisone or Solumedrol), T cell blockers (such as cyclosporin A and FK506), purine analogs (such as azathioprine (Imuran)), pyrimidine analogs (such as cytosine arabinoside), alkylating agents (such as nitrogen mustard,
30 phenylalanine mustard, buslfan, and cyclophosphamide), folic acid antagonists (such as aminopterin and methotrexate), antibiotics (such as rapamycin, actinomycin D, mitomycin C, puramycin, and chloramphenicol), human IgG, antilymphocyte globulin (ALG), and antibodies (such as anti-CD3 (OKT3),

anti-CD4 (OKT4), anti-CD5, anti-CD7, anti-IL-2 receptor, anti-alpha/beta TCR, anti-ICAM-1, anti-CD20 (Rituxan), anti-IL-12 and antibodies to immunotoxins).

5 The foregoing and other useful combination therapies will be understood and appreciated by those of skill in the art. Potential advantages of such combination therapies include a different efficacy profile, the ability to use less of each of the individual active ingredients to minimize toxic side effects, synergistic improvements in efficacy, improved ease of administration or use and/or reduced overall expense of compound preparation or formulation.

10

OTHER EMBODIMENTS

The compounds of this invention may be used as research tools (for example, as a positive control for evaluating other potential CRAC inhibitors, or IL-2, IL-4, IL-5, IL-13, GM-CSF, TNF- α , and/or INF- γ inhibitors). These and other uses
15 and embodiments of the compounds and compositions of this invention will be apparent to those of ordinary skill in the art.

The invention is further defined by reference to the following examples describing in detail the preparation of compounds of the invention. It will be
20 apparent to those skilled in the art that many modifications, both to materials and methods, may be practiced without departing from the purpose and interest of this invention. The following examples are set forth to assist in understanding the invention and should not be construed as specifically limiting the invention described and claimed herein. Such variations of the invention,
25 including the substitution of all equivalents now known or later developed, which would be within the purview of those skilled in the art, and changes in formulation or minor changes in experimental design, are to be considered to fall within the scope of the invention incorporated herein.

EXAMPLES

EXPERIMENTAL RATIONALE

- 5 Without wishing to be bound by theory, it is believed that the compounds of this invention inhibit CRAC ion channels, thereby inhibiting production of IL-2 and other key cytokines involved with inflammatory and immune responses. The examples that follow demonstrate these properties.

10 MATERIALS AND GENERAL METHODS


- Reagents and solvents used below can be obtained from commercial sources such as Aldrich Chemical Co. (Milwaukee, Wisconsin, USA). ¹H-NMR and ¹³C-NMR spectra were recorded on a Varian 300MHz NMR spectrometer. Significant peaks are tabulated in the order: δ (ppm); chemical shift, multiplicity
15 (s, singlet; d, doublet; t, triplet; q, quartet; m, multiplet; br s, broad singlet), coupling constant(s) in Hertz (Hz) and number of protons.

- Patch clamp experiments were performed in the tight-seal whole-cell configuration at 21-25°C. High resolution current recordings were acquired by
20 a computer-based patch clamp amplifier system (EPC-9, HEKA, Lambrecht, Germany). Patch pipettes had resistances between 2-4 M Ω after filling with the standard intracellular solution. Immediately following establishment of the whole-cell configuration, voltage ramps of 50-200 ms duration spanning the voltage range of -100 to +100 mV were delivered at a rate of 0.5 Hz over a
25 period of 300-400 seconds. All voltages were corrected for a liquid junction potential of 10 mV between external and internal solutions when using glutamate as the intracellular anion. Currents were filtered at 2.9 kHz and digitized at 10 μ s intervals. Capacitive currents and series resistance were determined and corrected before each voltage ramp using the automatic
30 capacitance compensation of the EPC-9. The low resolution temporal development of membrane currents was assessed by extracting the current amplitude at -80 mV or +80 mV from individual ramp current records.

Compounds of the invention can also be prepared as in U.S. Application No. 10/897,681, filed July 22, 2004 and U.S. Application No. --- entitled "Compounds for Inflammation and Immune-Related Uses," by Lijun Sun, *et al.*, filed on January 6, 2006, the entire teachings of which are incorporated herein by reference.

EXAMPLE 1: SYNTHESIS OF REPRESENTATIVE EXEMPLARY COMPOUNDS OF THIS INVENTION

10 Compound 1: N-[6-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-3-yl]-2,6-difluoro-benzamide



A mixture of 2-chloro-5-trifluoromethyl-phenylboronic acid (**a**, 1 mmol), 6-bromo-3-aminopyridine (**b**, 1 mmol), palladium catalyst (0.1 mmol), sodium bicarbonate (1 mmol) in a mixture of toluene (5 mL), water (1 mL), ethanol (1 mL) was heated at 100°C for 24 h. The mixture was taken up with EtOAc (100 mL), washed with water (2x100 mL) and dried (Na_2SO_4). The oil obtained on concentration was purified by flash chromatography to give **c** as a yellowish oil (0.20 mg).

The above oil was treated with 2,6-difluorobenzoyl chloride (0.6 mmol) in DCM (5 mL) and pyridine (0.2 mL) at rt for 2 h. Removal of solvents and purification of the residue by chromatography gave product **d** as yellow solid (0.15 g). $^1\text{H-NMR}$ (CDCl_3) δ 8.8 (d, 1H, $J = 2$), 8.5 (dd, 1H, $J_1 = 9$, $J_2 = 2$), 7.91 (s, 2H), 7.8 (d, 1H, $J = 8$), 7.6 (m, 2H), 7.5 (m, 1H), 7.0 (t, 2H, $J = 8$) ppm; ESMS calcd for $\text{C}_{19}\text{H}_{10}\text{ClF}_5\text{N}_2\text{O}$: 412.0; found: 413.0 ($\text{M} + \text{H}^+$).

Compound 2 through Compound 22 were synthesized in a similar manner:

Compound 2: N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-
2,6-difluoro-benzamide

5

$^1\text{H-NMR}$ (CDCl_3) δ 9.2 (m, 1H), 8.5 (m, 1H), 8.18 (s, 1H), 7.9 (m, 1H), 7.6 (m, 3H), 7.4 (m, 1H), 7.0 (t, 2H, $J = 8$) ppm; ESMS calcd for $\text{C}_{19}\text{H}_{10}\text{ClF}_5\text{N}_2\text{O}$: 412.0; found: 413.0 ($\text{M} + \text{H}^+$).

10 Compound 3: N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-
3-methyl-isonicotinamide

15 $^1\text{H-NMR}$ (CDCl_3) δ 9.20 (s, 1H), 8.6 (m, 2H), 8.5 (d, 1H, $J = 9$), 8.11 (s, 1H), 7.9 (d, 1H, $J = 9$), 7.6 (m, 3H), 7.4 (d, 1H, $J = 5$), 2.52 (s, 3H) ppm; ESMS calcd for $\text{C}_{19}\text{H}_{13}\text{ClF}_3\text{N}_3\text{O}$: 391.1; found: 392.0 ($\text{M} + \text{H}^+$).

Compound 4: N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-
3-fluoro-isonicotinamide

20

$^1\text{H-NMR}$ (CDCl_3) δ 9.2 (br, 1H), 8.7 (m, 2H), 8.4 (m, 2H), 8.0 (t, 1H, $J = 6$), 7.9 (dd, 1H, $J_1 = 9$, $J_2 = 2$), 7.6 (m, 3H) ppm; ESMS calcd for $\text{C}_{18}\text{H}_{10}\text{ClF}_4\text{N}_3\text{O}$: 395.0; found: 396.0 ($\text{M} + \text{H}^+$).

25 Compound 5: 2,6-Difluoro-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-
pyridin-2-yl]-benzamide

30 $^1\text{H-NMR}$ (CDCl_3) δ 8.6 (br, 1H), 8.4 (d, 1H, $J = 8$), 8.26 (s, 1H), 8.0 (d, 1H, $J = 8$), 7.91 (s, 1H), 7.8 (d, 1H, $J = 8$), 7.71 (s, 1H), 7.4 (m, 2H), 7.3 (m, 1H), 7.0 (m, 2H), 2.33 (s, 3H) ppm; ESMS calcd for $\text{C}_{20}\text{H}_{15}\text{F}_2\text{N}_3\text{O}_2$: 391.1; found: 392.1 ($\text{M} + \text{H}^+$).

Compound 6: 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester

¹H-NMR (CDCl₃) δ 8.7 (br, 1H), 8.5 (d, 1H, J = 8), 8.3 (m, 1H), 8.0 (m, 2H), 7.9 (d, 1H, J = 8), 7.6 (d, 1H, J = 9), 7.4 (m, 1H), 7.0 (t, 2H, J = 8), 3.94 (s, 3H) ppm; ESMS calcd for C₂₀H₁₃ClF₂N₂O₃: 402.1; found: 403.0 (M + H⁺).

Compound 7: 3-[6-(2,6-Difluoro-benzoylamino)-pyridin-3-yl]-4-methyl-benzoic acid methyl ester

10

¹H-NMR (CDCl₃) δ 9.5 (br, 1H), 8.4 (d, 1H, J = 8), 8.0 (m, 2H), 7.9 (m, 2H), 7.4 (m, 2H), 7.0 (t, 2H, J = 8), 3.94 (s, 3H), 2.30 (s, 3H) ppm; ESMS calcd for C₂₁H₁₆F₂N₂O₃: 382.1; found: 383.1 (M + H⁺).

15 Compound 8: N-[5-(2-Chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide

¹H-NMR (CDCl₃) δ 9.8 (br, 1H), 8.5 (m, 1H), 8.0 (m, 4H), 7.8 (m, 1H), 7.6 (m, 1H), 7.4 (m, 2H), 7.0 (m, 2H) ppm; ESMS calcd for C₂₁H₁₂ClF₂N₃O₂: 411.1; found: 412.0 (M + H⁺).

20

Compound 9: N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-6-methyl-pyridin-2-yl]-2,6-difluoro-benzamide

25 ¹H-NMR (CDCl₃) δ 9.1 (br, 1H), 8.3 (d, 1H, J = 8), 7.3-7.6 (m, 5H), 7.0 (t, 2H, J = 8), 2.16 (s, 3H) ppm; ESMS calcd for C₂₀H₁₂ClF₅N₂O: 426.1; found: 427.0 (M + H⁺).

Compound 10: N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide

30

¹H-NMR (CDCl₃) δ 9.6 (br, 1H), 8.5 (d, 1H, J = 8), 8.07 (s, 1H), 7.9 (m, 4H), 7.6 (d, 1H, J = 9), 7.4 (m, 2H), 7.0 (t, 2H, J = 8) ppm; ESMS calcd for C₂₀H₁₂ClF₂N₃OS: 427.0; found: 428.0 (M + H⁺).

5 Compound 11: N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-4-methyl-pyridin-2-yl]-2,6-difluoro-benzamide

10 ¹H-NMR (CDCl₃) δ 10.6 (br, 1H), 8.40 (s, 1H), 7.6 (m, 2H), 7.4 (m, 3H), 6.9 (m, 2H), 2.21 (s, 3H) ppm; ESMS calcd for C₂₀H₁₂ClF₅N₂O: 426.1; found: 427.0 (M + H⁺).

Compound 12: 3-[5-(2,6-Difluoro-benzoylamino)-pyridin-2-yl]-4-methyl-benzoic acid methyl ester

15 ¹H-NMR (CDCl₃) δ 8.72 (s, 1H), 8.4 (d, 1H, J = 8), 8.09 (s, 1H), 8.0 (d, 1H, J = 8), 7.8 (br, 1H), 7.5 (m, 2H), 7.4 (d, 1H, J = 8), 7.0 (t, 2H, J = 8), 3.91 (s, 3H), 2.45 (s, 3H) ppm; ESMS calcd for C₂₁H₁₆F₂N₂O₃: 382.1; found: 383.1 (M + H⁺).

20 Compound 13: 2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide

25 ¹H-NMR (CDCl₃) δ 8.72 (s, 1H), 8.4 (d, 1H, J = 8), 8.32 (s, 1H), 8.04 (s, 1H), 8.0 (d, 1H, J = 8), 7.64 (s, 1H), 7.4 (m, 3H), 7.20 (s, 1H), 7.0 (t, 2H, J = 8), 2.38 (s, 3H) ppm; ESMS calcd for C₂₂H₁₅F₂N₃O₂: 391.1; found: 392.1 (M + H⁺).

Compound 14:

2,6-Difluoro-N-[4-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide

30 ¹H-NMR (CDCl₃) δ 9.4 (br, 1H), 8.37 (s, 1H), 7.80 (s, 1H), 7.3-7.7 (m, 5H), 7.0 (t, 2H, J = 8), 2.34 (s, 3H) ppm; ESMS calcd for C₂₀H₁₃F₅N₂O: 392.1; found: 393.0 (M + H⁺).

Compound 15:

2,6-Difluoro-N-[6-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide

- 5 $^1\text{H-NMR}$ (CDCl_3) δ 8.9 (br, 1H), 8.3 (d, 1H, $J = 8$), 7.3-7.7 (m, 6H), 7.0 (t, 2H, $J = 8$), 2.32 (s, 3H) ppm; ESMS calcd for $\text{C}_{20}\text{H}_{13}\text{F}_5\text{N}_2\text{O}$: 392.1; found: 393.0 ($\text{M} + \text{H}^+$).

Compound 16: 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid

- 10 [5-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-amide

$^1\text{H-NMR}$ (CDCl_3) δ 9.1 (br, 1H), 8.4 (d, 1H, $J = 9$), 8.29 (s, 1H), 8.0 (d, 1H, $J = 8$), 7.6 (m, 3H), 3.02 (s, 3H) ppm; ESMS calcd for $\text{C}_{16}\text{H}_{10}\text{ClF}_3\text{N}_4\text{OS}$: 398.0; found: 399.0 ($\text{M} + \text{H}^+$).

15

Compound 17: 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid

[5-(2-chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-amide

- 20 $^1\text{H-NMR}$ (CDCl_3) δ 8.4 (m, 3H), 7.9 (m, 4H), 7.6 (d, 1H, $J = 8$), 7.4 (d, 1H, $J = 5$), 3.05 (s, 3H) ppm; ESMS calcd for $\text{C}_{18}\text{H}_{12}\text{ClN}_5\text{OS}_2$: 413.0; found: 414.0 ($\text{M} + \text{H}^+$).

Compound 18: 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid

[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide

25

$^1\text{H-NMR}$ (CDCl_3) δ 9.4 (br, 1H), 8.4 (d, 1H, $J = 8$), 8.31 (s, 1H), 7.9 (m, 3H), 7.72 (s, 1H), 7.4 (d, 1H, $J = 8$), 7.24 (s, 1H), 3.03 (s, 3H), 2.34 (s, 3H) ppm; ESMS calcd for $\text{C}_{19}\text{H}_{15}\text{N}_5\text{O}_2\text{S}$: 377.1; found: 378.1 ($\text{M} + \text{H}^+$).

- 30 Compound 19: 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid

[5-(2-chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide

¹H-NMR (CDCl₃) δ 8.6 (d, 1H, J = 9), 8.42 (s, 1H), 8.1 (m, 3H), 7.77 (s, 1H), 7.6 (s, 1H, J = 8), 7.30 (s, 1H), 3.03 (s, 3H) ppm; ESMS calcd for C₁₈H₁₂ClN₅O₂S: 397.0; found: 398.1 (M + H⁺).

5 **Compound 20:** 2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide hydrochloride


¹H-NMR (DMSO-d₆) δ 11.6 (br, 1H), 9.12 (s, 1H), 8.4 (d, 1H, J = 8), 8.12 (s, 1H), 8.0 (m, 2H), 7.9 (d, 1H, J = 8), 7.6 (m, 1H), 7.5 (d, 1H, J = 8), 7.41 (s, 1H), 7.3 (t, 2H, J = 8), 2.41 (s, 3H) ppm; ESMS calcd for C₂₂H₁₆ClF₂N₃O₂: 427.1; found: 398.1 (M + H⁺).

Compound 21: 4-methyl-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-2-yl)-1,2,3-thiadiazole-5-carboxamide

15

¹H NMR (300 MHz, (CD₃)₂SO) 11.55 (s, 1 H), 8.45-8.43 (m, 2 H), 8.22-8.19 (m, 1 H), 7.98-7.94 (m, 1 H), 7.71-7.63 (m, 3 H), 7.46-7.43 (m, 1 H), 2.82 (s, 3 H), 2.28 (s, 3 H); ESMS calcd (C₁₉H₁₅N₅O₂S): 377.1; found: 378.2 (M+H).

20 **Compound 68:** N-(5-(5-(isoxazol-5-yl)-2-methylphenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-carboxamide



Compound 68

25


General Procedure for Suzuki cross coupling: To a solution of 5-(3-bromo-4-methylphenyl)isoxazole (200 mg, 0.84 mmol), dichlorobis(triphenylphosphine)palladium (II) (Pd(PPh₃)₂Cl₂, 60 mg, 0.09 mmol), and

6-aminopyridin-3-ylboronic acid pinacol ester (185 mg, 0.84 mmol) in toluene (10 mL) was added Na_2CO_3 (2 N, 1.0 mL) and ethanol (1.0 mL). The stirred mixture was heated up to 80 °C for 16 hr. The solution was cooled to room temperature and diluted with H_2O (10 mL) and EtOAc (10 mL). The organic phase was dried over Na_2SO_4 , concentrated, and chromatographed to give the pure product (120 mg, 57%).

General Procedure A for amide bond formation: To a solution of 5-(5-(isoxazol-5-yl)-2-methylphenyl)pyridin-2-amine (60 mg, 0.24 mmol) in DCM (4 mL) was added EDC (85 mg, 0.45 mmol) and 4-methyl-1,2,3-thiadiazole-5-carboxylic acid (70 mg, 0.48 mmol). The solution was stirred at room temperature for 6 hr before it was concentrated and chromatographed to give the pure product (50 mg, 55%).

^1H NMR (300 MHz, CDCl_3) δ 8.58 (d, J = 9 Hz, 1 H), 8.32-8.28 (m, 2 H), 7.99-7.95 (m, 1 H), 7.78-7.74 (m, 1 H), 7.68 (d, J = 1.8 Hz, 1 H), 7.45-7.43 (m, 1 H), 6.54 (d, J = 1.8 Hz, 1 H), 2.35 (d, 3 H);). ESMS calcd ($\text{C}_{19}\text{H}_{15}\text{N}_5\text{O}_2\text{S}$): 377.1; found: 378.1 ($\text{M}+\text{H}$).

Compound 69: 2,6-difluoro-N-(5-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)benzamide



General Procedure B for amide bond formation: To the solution of 5-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-amine (100 mg, 0.4 mmol) in 3 DCM (3 mL) was added 2,6-difluorobenzoyl chloride (0.1 mL, 0.8 mmol). The reaction solution was stirred at room temperature for 60 min before it was concentrated and chromatographed to afford the mixture of mono amide and

di-amide products. The mixture was dissolved in 5 mL MeOH and heated at 50 °C for 25 min before it was concentrated and extracted between ethyl acetate and H₂O. The organic phase was dried and concentrated to afford the pure final product (117 mg, 75%).

5

¹H NMR (300 MHz, CDCl₃) δ 9.16 (s, 1 H), 8.51-8.48 (m, 2 H), 8.14-8.12 (m, 1 H), 8.01 (dd, *J* = 8.1 and 2.1 Hz, 1 H), 7.91 (d, *J* = 1.8 Hz, 1 H), 7.79 (dd, *J* = 8.7 and 2.4 Hz, 1 H), 7.45-7.42 (m, 2 H), 7.05-6.98 (m, 2 H), 2.35 (s, 3 H); ESMS calcd (C₂₁H₁₄F₂N₄O₂): 392.1; found: 393.2 (M+H).

10

Compound 70: 2,6-difluoro-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-2-yl)benzamide

¹H NMR (300 MHz, CDCl₃) δ 8.81 (s, 1 H), 8.47 (d, *J* = 8.4 Hz, 1 H), 8.21-8.20 (m, 1 H), 7.92 (s, 1 H), 7.78 (dd, *J* = 8.4 and 2.1 Hz, 1 H), 7.59 (dd, *J* = 7.8 and 1.8 Hz, 1 H), 7.50-7.35 (m, 4 H), 7.05-6.99 (m, 2 H), 2.30 (s, 3 H); ESMS calcd (C₂₂H₁₅F₂N₃O₂): 391.1; found: 392.2 (M+H).

15

Compound 71: N-(5-(2-chloro-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)-2,6-difluorobenzamide

20

¹H NMR (300 MHz, CDCl₃) δ 9.07 (s, 1 H), 8.52-8.49 (m, 2 H), 8.26 (d, *J* = 1.8 Hz, 1 H), 8.07-8.03 (m, 2 H), 7.93 (dd, *J* = 8.7 and 2.4 Hz, 1 H), 7.69-7.66 (m, 1 H), 7.48-7.42 (m, 1 H), 7.05-6.98 (m, 2 H); ESMS calcd (C₂₀H₁₁ClF₂N₄O₂): 412.1; found: 413.2 (M+H).

25

Compound 72: N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide

¹H NMR (300 MHz, CDCl₃) δ 8.76 (d, *J* = 2.7 Hz, 1 H), 8.42 (dd, *J* = 8.7 and 2.7 Hz, 1 H), 8.34 (s, 1 H), 8.14 (d, *J* = 2.4 Hz, 1 H), 7.92 (dd, *J* = 8.6 and 2.3 Hz,

30

1 H), 7.85 (d, $J = 3.3$ Hz, 1 H), 7.73 (d, $J = 8.7$ Hz, 1 H), 7.55 (d, $J = 8.4$ Hz, 1 H), 7.45-7.35 (m, 2 H), 7.02-6.95 (m, 2 H); ESMS calcd ($C_{21}H_{12}ClF_2N_3OS$): 427.0; found: 428.1 (M+H).

- 5 Compound 73: *N*-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt

^1H NMR (300 MHz, CD_3OD) δ 9.62 (d, $J = 2.1$ Hz, 1 H), 8.72-8.67 (m, 1 H), 8.36-8.23 (m, 3 H), 8.00 (d, $J = 3.3$ Hz, 1 H), 7.86 (d, $J = 8.1$ Hz, 1 H), 7.79 (d, $J = 3.3$ Hz, 1 H), 7.66-7.61 (m, 1 H), 7.19 (t, $J = 8.1$ Hz, 2 H); ESMS calcd ($C_{21}H_{12}ClF_2N_3OS$): 427.0; found: 428.1 (M+H).

Compound 74: 2,6-Difluoro-*N*-(6-(2-methyl-5-(thiazol-2-yl)phenyl)pyridin-3-yl)benzamide

15

^1H NMR (300 MHz, $CDCl_3$) δ 8.74 (d, $J = 2.4$ Hz, 1 H), 8.43 (dd, $J = 8.6$ and 2.6 Hz, 1 H), 8.02-7.98 (m, 2 H), 7.89 (dd, $J = 5.0$ and 2.1 Hz, 1 H), 7.84 (d, $J = 3.3$ Hz, 1 H), 7.53-7.30 (m, 4 H), 7.03 (t, $J = 8.3$ Hz, 2 H); ESMS calcd ($C_{22}H_{15}F_2N_3OS$): 407.1; found: 408.2 (M+H).

20

Compound 75: 2,6-Difluoro-*N*-(6-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-3-yl)benzamide

^1H NMR (300 MHz, $CDCl_3$) δ 8.76 (d, $J = 2.4$ Hz, 1 H), 8.45-8.41 (m, 2 H), 8.29 (s, 1 H), 8.10 (d, $J = 1.8$ Hz, 1 H), 8.00 (dd, $J = 8.0$ and 2.0 Hz, 1 H), 7.51-7.42 (m, 3 H), 7.01 (t, $J = 8.1$ Hz, 2 H), 2.45 (s, 3 H); ESMS calcd ($C_{21}H_{14}F_2N_4O_2$): 392.1; found: 393.2 (M+H).

Compound 76: *N*-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide

30

^1H NMR (300 MHz, CD_3OD) δ 8.96-8.94 (m, 1 H), 8.37 (dd, $J = 8.6$ and 2.6 Hz, 1 H), 8.22 (d, $J = 2.1$ Hz, 1 H), 8.08-8.04 (m, 1 H), 8.02 (d, $J = 0.9$ Hz, 1 H),

7.80-7.76 (m, 1 H), 7.70 (d, $J = 8.4$ Hz, 1 H), 7.60-7.54 (m, 1 H), 7.33 (d, $J = 0.9$ Hz, 1 H), 7.17-7.12 (m, 2 H); ESMS calcd ($C_{21}H_{12}ClF_2N_3O_2$): 411.1; found: 412.2 (M+H).

- 5 Compound 77: *N*-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt

^1H NMR (300 MHz, CD_3OD) δ 9.63-9.61 (m, 1 H), 8.74-8.68 (m, 1 H), 8.39-8.27 (m, 3 H), 8.10 (d, $J = 0.6$ Hz, 1 H), 7.89 (d, $J = 8.7$ Hz, 1 H), 8.67-8.59 (m, 1 H),
10 7.40 (d, $J = 0.6$ Hz, 1 H), 7.19 (t, $J = 8.1$ Hz, 2 H); ESMS calcd ($C_{21}H_{12}ClF_2N_3O_2$): 411.1; found: 412.2 (M+H).

Compound 78: 2,6-Difluoro-*N*-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide

15

^1H NMR (300 MHz, CD_3OD) δ 8.91-8.90 (m, 1 H), 8.35 (dd, $J = 8.6$ and 2.4 Hz, 1 H), 8.24 (s, 1 H), 7.75-7.67 (m, 2 H), 7.61-7.52 (m, 3 H), 7.42 (d, $J = 8.1$ Hz, 1 H), 7.14 (t, $J = 8.1$ Hz, 2 H), 2.36 (s, 3 H); ESMS calcd ($C_{22}H_{15}F_2N_3O_2$): 391.1; found: 392.3 (M+H).

20

Compound 79: 2,6-Difluoro-*N*-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide hydrochloric acid salt

^1H NMR (300 MHz, CD_3OD) δ 9.60 (s, 1 H), 8.69 (d, $J = 7.8$ Hz, 1 H), 8.57 (s, 1 H), 8.23 (d, $J = 8.4$ Hz, 1 H), 7.98-7.92 (m, 2 H), 7.76 (s, 1 H), 7.66-7.60 (m, 2 H), 7.19 (t, $J = 8.1$ Hz, 2 H), 2.43 (s, 3 H); ESMS calcd ($C_{22}H_{15}F_2N_3O_2$): 391.1; found: 392.3 (M+H).

25

Compound 80: 4-chloro-2-fluoro-*N*-(6-(2-methyl-5-(oxazol-2-yl)phenyl)pyridin-3-yl)benzamide

30

$^1\text{H-NMR}$ (CDCl_3) δ (ppm) 8.8 (d, 1H, $J=2$), 8.5 (br, 1H), 8.4 (dd, 1H, $J_1=8$, $J_2=2$), 7.9-8.2 (m, 3H), 7.70 (s, 1H), 7.2-7.5 (m, 5H), 2.42 (s, 3H); ESMS clcd for $\text{C}_{22}\text{H}_{15}\text{ClFN}_3\text{O}_2$: 407.1; Found: 408.0 ($\text{M}+\text{H}$) $^+$.

5


Compounds of the invention in which L is $-\text{NHC}(\text{S})-$ or $-\text{C}(\text{S})\text{NH}-$ can be prepared by treating compounds having an amide linker with Lawesson's reagent.

- 10 Compounds having $-\text{CH}_2\text{NH}-$ or $-\text{NHCH}_2-$ linkers can be prepared by contacting compounds having $-\text{NHC}(\text{S})-$ or $-\text{C}(\text{S})\text{NH}-$ linkers with Raney Ni. Alternatively, compounds of the invention having a $-\text{CH}_2\text{NH}-$ or $-\text{NHCH}_2-$ linker can be prepared by reducing a compound having a $-\text{C}(\text{O})\text{NH}-$ or $-\text{NH}\text{C}(\text{O})-$ linker, respectively, with, for example, sodium borohydride (see U.S. Patent Application No. 10/897,681, filed on July 22, 2004, the entire teachings of which are incorporated herein by reference).
- 15

Compounds of the invention having $-\text{C}(\text{O})-$ linkers can be prepared by a Friedel-Craft acylation reaction, for example, by reacting a pyridinyl derivative (XXIII) with an acid chloride (XXIV) in the presence of AlCl_3 to form an intermediate which can then be reacted with an [1,3,2]dioxaborolan-2-yl-aryl or -heteroaryl (XXV) in the presence of a palladium catalyst and a base to form a compound of the invention having a carbonyl linker (XXVI) (see Scheme IV).

20

Scheme IV




Compounds of the invention that have $-C(S)-$ can be prepared from compounds that have carbonyl linkers by treating them with Lawesson's reagent or P_2S_5 in pyridine.

5

Compounds of the invention having $-CH_2-NH-$ or $-NH-CH_2-$ linkers can be prepared by contacting compounds having $-NHC(S)-$ or $-C(S)NH-$ linkers with Raney Ni. Alternatively, compounds of the invention having a $-CH_2-NH-$ or $-NH-CH_2-$ linker can be prepared by reducing a compound having a $-C(O)-NH-$ or $-NH-C(O)-$ linker, respectively, with, for example, sodium borohydride. Alternatively, compounds that have $-NHCH_2-$ linkers can be prepared by reacting aldehyde (f) with amine (XX) followed by reduction of the shift base with sodium borohydride as shown in Scheme VIa (see U.S. Patent Application No. 10/897,681, filed on July 22, 2004, the entire teachings of which are incorporated herein by reference).

15

Scheme VIa



20

EXAMPLE 2: INHIBITION OF IL-2 PRODUCTION

Jurkat cells were placed in a 96 well plate (0.5 million cells per well in 1% FBS medium) then a test compound of this invention was added at different concentrations. After 10 minutes, the cells were activated with PHA (final concentration 2.5 $\mu g/mL$) and incubated for 20 hours at 37°C under CO_2 . The final volume was 200 μL . Following incubation, the cells were centrifuged and the supernatants collected and stored at -70°C prior to assaying for IL-2 production. A commercial ELISA kit (IL-2 Eli-pair, Diaclone Research, Besancon, France) was used to detect production of IL-2, from which dose response curves were obtained. The IC_{50} value was calculated as the

25

30

concentration at which 50% of maximum IL-2 production after stimulation was inhibited versus a non-stimulation control.

Compound #	IC ₅₀ (nM)
6	6
7	4
8	3
10	2
13	14
14	16
15	98
16	15
17	2
18	12
19	7
20	12
21	31
67	7
68	38
69	73
70	9
71	17
72	17
73	17
74	25
75	98
76	9
77	11
78	10
79	10
80	180

Inhibition of other cytokines, such as IL-4, IL-5, IL-13, GM-CSF, TNF- α , and INF- γ , can be tested in a similar manner using a commercially available ELISA kit for each cytokine.

5 EXAMPLE 3: PATCH CLAMP STUDIES OF INHIBITION OF I_{CRAC} CURRENT IN RBL CELLS, JURKAT CELLS, AND PRIMARY T CELLS

In general, a whole cell patch clamp method is used to examine the effects of a compound of the invention on a channel that mediates I_{crac} . In such experiments, a baseline measurement is established for a patched cell. Then
10 a compound to be tested is perfused (or puffed) to cells in the external solution and the effect of the compound on I_{crac} is measured. A compound that modulates I_{crac} (e.g., inhibits) is a compound that is useful in the invention for modulating CRAC ion channel activity.

15 **1) RBL cells**

Cells

Rat basophilic leukemia cells (RBL-2H3) were grown in DMEM media
20 supplemented with 10% fetal bovine serum in an atmosphere of 95% air/5% CO₂. Cells were seeded on glass coverslips 1-3 days before use.

Recording Conditions

Membrane currents of individual cells were recorded using the whole-cell configuration of the patch clamp technique with an EPC10 (HEKA Elektronik,
25 Lambrecht, Germany). Electrodes (2-5 M Ω in resistance) were fashioned from borosilicate glass capillary tubes (Sutter Instruments, Novato, Ca). The recordings were done at room temperature.

Intracellular pipette solution

The intracellular pipette solution contained Cs-Glutamate 120mM; CsCl
30 20mM; CsBAPTA 10mM; CsHEPES 10mM; NaCl 8mM; MgCl₂ 1mM; IP3 0.02mM; pH=7.4 adjusted with CsOH. The solution was kept on ice and shielded from light before the experiment was preformed.

Extracellular solution

The extracellular solution contained NaCl 138mM; NaHEPES, 10mM; CsCl 10mM; CaCl₂ 10mM; Glucose 5.5mM; KCl 5.4mM; KH₂PO₄ 0.4mM; Na₂HPO₄·H₂O 0.3mM at pH=7.4 adjusted with NaOH.

5 Compound treatment

Each compound was diluted from a 10 mM stock in series using DMSO. The final DMSO concentration was always kept at 0.1 %.

Experimental procedure

10 I_{CRAC} currents were monitored every 2 seconds using a 50 msec protocol, where the voltage was ramped from -100 mV to +100 mV. The membrane potential was held at 0 mV between the test ramps. In a typical experiment, the peak inward currents would develop within 50-100 seconds. Once the I_{CRAC} currents were stabilized, the cells were perfused with a test compound in the extracellular solution. At the end of an experiment, the remaining I_{CRAC} currents
15 were then challenged with a control compound (SKF96365, 10 μ M) to ensure that the current could still be inhibited.

Data analysis

The I_{CRAC} current level was determined by measuring the inward current amplitude at -80 mV of the voltage ramp in an off-line analysis using MATLAB.
20 The I_{CRAC} current inhibition for each concentration was calculated using peak amplitude in the beginning of the experiment from the same cell. The IC₅₀ value and Hill coefficient for each compound was estimated by fitting all the individual data points to a single Hill equation.

25 Results

The table below shows the concentration of compounds of the invention which inhibits 50 % of the I_{CRAC} current in RBL cells.

Compound Number	IC ₅₀
8	80 nM

17	40 nM
18	140 nM
20	150 nM
SKF96365	4 μ M

2) Jurkat cells

Cells

Jurkat T cells are grown on glass coverslips, transferred to the recording
 5 chamber and kept in a standard modified Ringer's solution of the following
 composition: NaCl 145mM, KCl 2.8mM, CsCl 10mM, CaCl₂ 10mM, MgCl₂
 2mM, glucose 10mM, HEPES-NaOH 10mM, pH 7.2.

Extracellular Solution

The external solution contains 10 mM CaNaR, 11.5 mM glucose and a test
 10 compound at various concentrations.

Intracellular Pipette Solution

The standard intracellular pipette solution contains: Cs-glutamate 145 mM,
 NaCl 8 mM, MgCl₂ 1 mM, ATP 0.5 mM, GTP 0.3 mM, pH 7.2 adjusted with
 CsOH. The solution is supplemented with a mixture of 10 mM Cs-BAPTA and
 15 4.3-5.3 mM CaCl₂ to buffer [Ca²⁺]_i to resting levels of 100-150 nM.

Patch-clamp recordings

Patch-clamp experiments are performed in the tight-seal whole-cell
 configuration at 21-25 °C. High-resolution current recordings are acquired by
 a computer-based patch-clamp amplifier system (EPC-9, HEKA, Lambrecht,
 20 Germany). Sylgard®-coated patch pipettes have resistances between 2-4 M Ω
 after filling with the standard intracellular solution. Immediately following
 establishment of the whole-cell configuration, voltage ramps of 50 ms duration
 spanning the voltage range of -100 to +100 mV are delivered from a holding
 potential of 0 mV at a rate of 0.5 Hz over a period of 300 to 400 seconds. All
 25 voltages are corrected for a liquid junction potential of 10 mV between external

and internal solutions. Currents are filtered at 2.3 kHz and digitized at 100 μ s intervals. Capacitive currents and series resistance are determined and corrected before each voltage ramp using the automatic capacitance compensation of the EPC-9.

5 **Data analysis**

The very first ramps before activation of I_{CRAC} (usually 1 to 3) are digitally filtered at 2 kHz, pooled and used for leak-subtraction of all subsequent current records. The low-resolution temporal development of inward currents is extracted from the leak-corrected individual ramp current records by measuring
10 the current amplitude at -80 mV or a voltage of choice.

3) Primary T Cells

Preparation of Primary T Cells

15 Primary T cells are obtained from human whole blood samples by adding 100 μ L of RosetteSep® human T cell enrichment cocktail to 2 mL of whole blood. The mixture is incubated for 20 minutes at room temperature, then diluted with an equal volume of PBS containing 2% FBS. The mixture is layered on top of RosetteSep® DM-L density medium and then centrifuged for
20 20 minutes at 1200 g at room temperature. The enriched T cells are recovered from the plasma/density medium interface, then washed with PBS containing 2% FBS twice, and used in patch clamp experiments following the procedure described for RBL cells.

25 **EXAMPLE 4: INHIBITION OF MULTIPLE CYTOKINES IN PRIMARY HUMAN PBMCs**

Peripheral blood mononuclear cells (PBMCs) are stimulated with phytohemagglutinin (PHA) in the presence of varying concentrations of compounds of the invention or cyclosporine A (CsA), a known inhibitor of
30 cytokine production. Cytokine production is measured using commercially available human ELISA assay kits (from Cell Science, Inc.) following the manufacturers instructions.

The compounds of the invention are expected to be potent inhibitors of IL-2, IL-4, IL-5, IL-13, GM-CSF, INF- γ and TNF- α in primary human PBM cells. In addition, compounds of the invention are not expected to inhibit the anti-inflammatory cytokine, IL-10.

5

EXAMPLE 5: INHIBITION OF DEGRANULATION IN RBL CELLS

Procedure:

The day before the assay is performed, RBL cells, that have been grown to confluence in a 96 well plate, are incubated at 37°C for at least 2 hours. The medium is replaced in each well with 100 μ L of fresh medium containing 2 μ Lg/mL of anti-DNP IgE.

On the following day, the cells are washed once with PRS (2.6 mM glucose and 0.1% BSA) and 160 μ L of PRS is added to each well. A test compound is added to a well in a 20 μ L solution at 10X of the desired concentration and incubated for 20 to 40 minutes at 37°C. 20 μ L of 10X mouse anti-IgE (10 μ L/mL) is added. Maximum degranulation occurs between 15 to 40 minutes after addition of anti-IgE.

20

Compounds of the invention are expected to inhibit degranulation.

EXAMPLE 6: INHIBITION OF CHEMOTAXIS IN T CELLS

T-cell isolation:

Twenty ml aliquots of heparinized whole blood (2 pig, 1 human) are subjected to density gradient centrifugation on Ficoll Hypaque. The buffy coat layers representing peripheral blood mononuclear cells (PBMCs) containing lymphocytes and monocytes are washed once, resuspended in 12 ml of incomplete RPMI 1640 and then placed in gelatin-coated T75 culture flasks for 1 hr at 37°C. The non-adherent cells, representing peripheral blood lymphocytes (PBLs) depleted of monocytes, are resuspended in complete RPMI media and placed in loosely packed activated nylon wool columns that

30

have been equilibrated with warm media. After 1 hr at 37°C, the non-adherent T cell populations are eluted by washing of the columns with additional media. The T cell preparations are centrifuged, resuspended in 5 ml of incomplete RPMI, and counted using a hemocytometer.

5

Cell migration assay:

Aliquots of each T cell preparation are labeled with Calciën AM (TefLabs) and suspended at a concentration of 2.4×10^6 /ml in HEPES-buffered Hank's Balanced Salt Solution containing 1.83 mM CaCl_2 and 0.8 mM MgCl_2 , pH 7.4 (HHBSS). An equal volume of HHBSS containing 0, 20 nM, 200 nM or 2000 nM of compound 1 or 20 nM EDTA is then added and the cells incubated for 30 min at 37 °C. Fifty μl aliquots of the cell suspensions (60,000 cells) are placed on the membrane (pore size 5 μm) of a Neuroprobe ChemoTx 96 well chemotaxis unit that have been affixed over wells containing 10 ng/ml MIP-1 α in HHBSS. The T cells are allowed to migrate for 2 hr at 37 °C, after which the apical surface of the membrane is wiped clean of cells. The chemotaxis units are then placed in a CytoFlour 4000 (PerSeptive BioSystems) and the fluorescence of each well measured (excitation and emission wavelengths of 450 and 530 nm, respectively). The number of migrating cells in each well is determined from a standard curve generated from measuring the fluorescence of serial two-fold dilutions of the labeled cells placed in the lower wells of the chemotaxis unit prior to affixing the membrane.

25

Compounds of the invention are expected to inhibit chemotactic response of T cells.


30

All publications, patent applications, patents, and other documents cited herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting in any way.

From the foregoing description, it will be apparent that variations and modifications may be made to the invention described herein. Such embodiments are also within the scope of the following claims.

WE CLAIM:

1. A compound represented by structural formula (II):



or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

Y_1 is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl, provided that Y_1 is not a substituted isoxazolyl or a substituted 1H-pyrrolyl;

L_1 is a linker selected from the group consisting of $-NHCH_2-$, $-CH_2NH-$, $-NR-C(O)-$, $-OC(O)-$, $-C(O)O-$, $-C(S)-$, $-NR-C(S)-$, or $-C(S)-NR-$;

one of X_1 or X_2 is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sufanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$, cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from $-H$ or an alkyl;

R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$, or $-NR_6R_7$;

R_4 is a halo, a haloalkyl, cyano, nitro, $-C(O)R_5$, $-C(O)OR_5$, $-C(O)SR_5$, $-C(O)NR_9R_{10}$, $-C(S)R_5$, $-C(S)OR_5$, $-C(S)SR_5$, $-C(S)NR_6R_7$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-C(NR_8)SR_5$, $-C(NR_8)NR_6R_7$, $-S(O)_pR_5$, $-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an

optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

5 R₅, for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 R₆ and R₇, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₆ and R₇ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

20 R₈, for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

25 R₉ and R₁₀, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indoliziny, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an

- optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally substituted indolizinyl, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;
- q, for each occurrence, is independently, an integer from 1 to 3;
- k is for each occurrence, is independently, an integer from 1 to 4;
- n is zero, 1 or 2; and
- p, for each occurrence, is independently 1 or 2.
2. The compound of Claim 1, wherein Y₁ is an optionally substituted phenyl, an optionally substituted pyridinyl, an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally substituted thiadiazolyl, or an optionally substituted thiophenyl.
3. The compound of Claim 2, wherein Y₁ is an optionally substituted phenyl or an optionally substituted thiadiazolyl.

4. The compound of Claim 3, wherein Y_1 is an optionally substituted phenyl.
5. The compound of Claim 4, wherein Y_1 is substituted with one to two substituents.
6. The compound of Claim 5, wherein the one to two substituents are each independently a lower alkyl or a halo.
7. The compound of Claim 6, wherein Y_1 is a difluorophenyl.
8. The compound of Claim 3, wherein Y_1 is 4-methyl-1,2,3-thiadiazol-5-yl.
9. The compound of Claim 1, wherein L_1 is $-NR-C(O)-$.
10. The compound of Claim 1, wherein R is $-H$.
11. The compound of Claim 1, wherein R_4 is $-C(O)OR_5$, a haloalkyl, an optionally substituted oxazolyl, an optionally substituted thiazolyl, an optionally substituted imidazolyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyrrolyl, an optionally substituted thiophenyl, an optionally substituted furanyl, an optionally substituted thiadiazolyl, an optionally substituted oxadiazolyl, or an optionally substituted tetrazolyl.
12. The compound of Claim 11, wherein R_4 is oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, thiazol-5-yl, [1,3,4]oxadiazol-2-yl, trifluoromethyl, pyridin-3-yl, pyridin-2-yl, $-C(O)OCH_2CH_2CH_3$, $-C(O)OCH_2CH_3$, or $-C(O)OCH_3$.
13. The compound of Claim 1, wherein R_3 is $-H$, chloro, fluoro, bromo, methyl, or methoxy.

14. The compound of Claim 1, wherein n is 0.
15. The compound of Claim 1, wherein n is 1 and Z is methyl.
- 5 16. The compound of Claim 1, wherein X₁ is N.
17. The compound of Claim 16, wherein X₂ is CH.
18. The compound of Claim 1, wherein X₂ is N.
- 10 19. The compound of Claim 18, wherein X₁ is CH.
20. The compound of Claim 1, wherein:
 - 15 Y₁ is an optionally substituted phenyl, an optionally substituted pyridinyl, an optionally substituted pyridazinyl, an optionally substituted isothiazolyl, an optionally substituted thiazolyl, an optionally substituted triazolyl, an optionally substituted oxadiazolyl, an optionally substituted oxazolyl, an optionally substituted thiadiazolyl, or an optionally substituted thiophenyl;
 - 20 R₄ is -C(O)OR₅, a haloalkyl, an optionally substituted oxazolyl, an optionally substituted thiazolyl, an optionally substituted imidazolyl, an optionally substituted pyridinyl, an optionally substituted pyrazolyl, an optionally substituted pyrrolyl, an optionally substituted thiophenyl, an optionally substituted furanyl, an optionally substituted thiadiazolyl, an
 - 25 optionally substituted oxadiazolyl, or an optionally substituted tetrazolyl;
 - one of X₁ or X₂ is CH; and
 - n is 0.
21. The compound of Claim 20, wherein L₁ is -NH-C(O)-.
- 30 22. The compound of Claim 21, wherein R₃ is -H, chloro, fluoro, bromo, methyl, or methoxy.

23. The compound of Claim 22, wherein, R₄ is oxazol-2-yl, oxazol-5-yl, thiazol-2-yl, thiazol-5-yl, [1,3,4]oxadiazol-2-yl, trifluoromethyl, pyridin-3-yl, pyridin-2-yl, -C(O)OCH₂CH₂CH₃, -C(O)OCH₂CH₃, or -C(O)OCH₃.
- 5 24. The compound of Claim 23, wherein Y₁ is a difluorophenyl.
25. The compound of Claim 23, wherein Y₁ is 4-methyl-1,2,3-thiadiazol-5-yl.
- 10 26. The compound of Claim 1, wherein the compound is selected from the group consisting of:
- N-[6-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-3-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-3-methyl-isonicotinamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-3-fluoro-isonicotinamide;
 2,6-Difluoro-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester;
 3-[6-(2,6-Difluoro-benzoylamino)-pyridin-3-yl]-4-methyl-benzoic acid methyl ester;
 N-[5-(2-Chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-6-methyl-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-2,6-difluoro-benzamide;
 N-[5-(2-Chloro-5-trifluoromethyl-phenyl)-4-methyl-pyridin-2-yl]-2,6-difluoro-benzamide;
 3-[5-(2,6-Difluoro-benzoylamino)-pyridin-2-yl]-4-methyl-benzoic acid methyl ester;
 2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[4-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[6-methyl-5-(3-trifluoromethyl-phenyl)-pyridin-2-yl]-benzamide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-trifluoromethyl-phenyl)-pyridin-2-yl]-amide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-amide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide;
 4-Methyl-[1,2,3]thiadiazole-5-carboxylic acid [5-(2-chloro-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-amide;

2,6-Difluoro-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide hydrochloride;

4-methyl-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-2-yl)-1,2,3-thiadiazole-5-carboxamide;

3-(6-Benzoylamino-pyridin-3-yl)-4-chloro-benzoic acid methyl ester;

4-Chloro-3-[6-(2-methyl-benzoylamino)-pyridin-3-yl]-benzoic acid methyl ester;

4-Chloro-3-{6-[(3-methyl-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(3-fluoro-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(3-chloro-pyridine-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-[1,2,3]thiadiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-[1,2,3]oxadiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(5-methyl-3H-[1,2,3]triazole-4-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-isothiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-thiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-methyl-oxazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(2-methyl-2H-pyrazole-3-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

4-Chloro-3-{6-[(4-fluoro-isothiazole-5-carbonyl)-amino]-pyridin-3-yl}-benzoic acid methyl ester;

3-Methyl-N-[5-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-2-yl]-isonicotinamide;

3-Methyl-N-[6-(2-methyl-5-oxazol-2-yl-phenyl)-pyridin-3-yl]-isonicotinamide;

2,6-Difluoro-N-[5-methyl-6-(3-oxazol-2-yl-phenyl)-pyridin-3-yl]-benzamide;

2,6-Difluoro-N-[4-methyl-5-(3-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;

2,6-Difluoro-N-[6-methyl-5-(3-oxazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;

2,6-Difluoro-N-[5-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;

3-Methyl-N-[5-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-isonicotinamide;

3-Methyl-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-isonicotinamide;

2,6-Difluoro-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-benzamide;

2,6-Difluoro-N-[6-(2-methyl-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-benzamide hydrochloride;

N-[5-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-2-yl]-3-methyl-isonicotinamide;
 N-[6-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-3-methyl-isonicotinamide;
 N-[6-(2-Chloro-5-thiazol-2-yl-phenyl)-pyridin-3-yl]-2,6-difluoro-benzamide;
 3-[5-(2,6-Difluoro-benzoylamino)-pyridin-2-yl]-4-methyl-benzoic acid ethyl ester;
 3-[6-(2,6-Difluoro-benzoylamino)-pyridin-3-yl]-4-methyl-benzoic acid ethyl ester;
 2,6-Difluoro-N-[5-(2-methyl-5-oxazol-5-yl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-oxazol-5-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[6-(5-isoxazol-5-yl-2-methyl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[5-(5-isoxazol-5-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-[2-methyl-5-(3-methyl-isoxazol-5-yl)-phenyl]-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[6-[2-methyl-5-(3-methyl-isoxazol-5-yl)-phenyl]-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-pyridin-3-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[6-(2-methyl-5-pyridin-2-yl-phenyl)-pyridin-3-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-[1,3,4]oxadiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-[1,3,4]thiadiazol-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid propyl ester;
 4-Chloro-3-[6-(2,6-difluoro-benzoylamino)-pyridin-3-yl]-benzoic acid 2-methoxy-ethyl ester;
 2,6-Difluoro-N-[5-(5-furan-2-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-thiophen-2-yl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(5-furan-3-yl-2-methyl-phenyl)-pyridin-2-yl]-benzamide;
 2,6-Difluoro-N-[5-(2-methyl-5-thiophen-3-yl-phenyl)-pyridin-2-yl]-benzamide;
 N-(5-(2-methoxy-5-(oxazol-5-yl)phenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-carboxamide;
 N-(5-(5-(isoxazol-5-yl)-2-methylphenyl)pyridin-2-yl)-4-methyl-1,2,3-thiadiazole-5-carboxamide;
 2,6-difluoro-N-(5-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)benzamide;
 2,6-difluoro-N-(5-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-2-yl)benzamide;
 N-(5-(2-chloro-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-2-yl)-2,6-difluorobenzamide;
 N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide;
 N-(6-(2-chloro-5-(thiazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt;
 2,6-Difluoro-N-(6-(2-methyl-5-(thiazol-2-yl)phenyl)pyridin-3-yl)benzamide;
 2,6-Difluoro-N-(6-(2-methyl-5-(1,3,4-oxadiazol-2-yl)phenyl)pyridin-3-yl)benzamide;
 N-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide;

N-(6-(2-chloro-5-(oxazol-2-yl)phenyl)pyridin-3-yl)-2,6-difluorobenzamide hydrochloric acid salt;

2,6-Difluoro-N-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide;

2,6-Difluoro-N-(6-(2-methyl-5-(oxazol-5-yl)phenyl)pyridin-3-yl)benzamide hydrochloric acid salt; or

4-chloro-2-fluoro-N-(6-(2-methyl-5-(oxazol-2-yl)phenyl)pyridin-3-yl)benzamide;

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof.

27. A pharmaceutical composition, comprising a pharmaceutically acceptable carrier and a compound of any one of Claims 1 through 26.

28. The pharmaceutical composition of Claim 27, further comprising one or more additional therapeutic agents.

29. The pharmaceutical composition according to Claim 28, wherein the additional therapeutic agent is selected from the group consisting of immunosuppressive agents, anti-inflammatory agents and suitable mixtures thereof.

30. The pharmaceutical composition of Claim 29, wherein the additional therapeutic agent is selected from the group consisting of steroids, non-steroidal anti-inflammatory agents, antihistamines, analgesics, and suitable mixtures thereof.

31. A method of inhibiting immune cell activation comprising administering to the cell a compound of any one of Claims 1 through 26.


32. The method of Claim 31, wherein immune cell activation is inhibited in a subject by administering the compound to the subject.

33. The method of Claim 32, wherein the subject is human.

34. A method of inhibiting cytokine production in a cell, comprising administering to the cell a compound of any one of Claims 1 through 26.
- 5 35. The method of Claim 34, wherein cytokine production is inhibited in a subject by administering the compound to the subject.
36. The method of Claim 35, wherein the subject is human.
- 10 37. The method of Claim 35, wherein the cytokine is selected from the group consisting of IL-2, IL-4, IL-5, IL-13, GM-CSF, IFN- γ , TNF- α , and combinations thereof.
38. The method of Claim 37, wherein the cytokine is IL-2.
- 15 39. A method of modulating an ion channel in a cell, wherein the ion channel is involved in immune cell activation, comprising administering to the cell a compound of any one of Claims 1 through 26.
- 20 40. The method of Claim 39, wherein the ion channel is in a subject and it is modulated by administering the compound to the subject.
41. The method of Claim 40, wherein the subject is human.
- 25 42. The method of Claim 40, wherein the ion channel is a Ca^{2+} -release-activated Ca^{2+} channel (CRAC).
- 30 43. A method of inhibiting T-cell and/or B-cell proliferation in response to an antigen, comprising administering to the cell a compound of any one of Claims 1 through 26.
44. The method of Claim 43, wherein T-cell and/or B-cell proliferation is inhibited in a subject by administering the compound to the subject.

45. The method of Claim 44, wherein the subject is human.
46. A method for treating or preventing an immune disorder in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 26.
47. The method of Claim 46, wherein the subject is human.
48. The method of Claim 46, wherein the disorder is selected from the group consisting of multiple sclerosis, myasthenia gravis, Guillain-Barré, autoimmune uveitis, autoimmune hemolytic anemia, pernicious anemia, autoimmune thrombocytopenia, temporal arteritis, anti-phospholipid syndrome, vasculitides such as Wegener's granulomatosis, Behcet's disease, psoriasis, dermatitis herpetiformis, pemphigus vulgaris, vitiligo, Crohn's disease, ulcerative colitis, primary biliary cirrhosis, autoimmune hepatitis, Type 1 or immune-mediated diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, autoimmune disorder of the adrenal gland, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, ankylosing spondylitis, and Sjogren's syndrome.
49. A method for treating or preventing an inflammatory condition in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 26.
50. The method of Claim 49, wherein the subject is human.
51. The method according to Claim 49, wherein the disorder is selected from transplant rejection, skin graft rejection, arthritis, rheumatoid arthritis, osteoarthritis and bone diseases associated with increased bone resorption; inflammatory bowel disease, ileitis, ulcerative colitis, Barrett's syndrome, Crohn's disease; asthma, adult respiratory distress syndrome, chronic obstructive airway disease; corneal dystrophy,

- trachoma, onchocerciasis, uveitis, sympathetic ophthalmitis, endophthalmitis; gingivitis, periodontitis; tuberculosis; leprosy; uremic complications, glomerulonephritis, nephrosis; sclerodermatitis, psoriasis, eczema; chronic demyelinating diseases of the nervous system, multiple sclerosis, AIDS-related neurodegeneration, Alzheimer's disease, infectious meningitis, encephalomyelitis, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis viral or autoimmune encephalitis; autoimmune disorders, immune-complex vasculitis, systemic lupus and erythematoses; systemic lupus erythematosus (SLE); cardiomyopathy, ischemic heart disease hypercholesterolemia, atherosclerosis, preeclampsia; chronic liver failure, brain and spinal cord trauma, and cancer.
52. A method for suppressing the immune system of a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 26.
53. The method of Claim 52, wherein the subject is human.
54. A method for treating or preventing an allergic disorder in a subject in need thereof, comprising administering to the subject an effective amount of a compound of any one of Claims 1 through 26.
55. The method of Claim 54, wherein the subject is human.
56. The method of Claim 54, wherein the disorder is allergic rhinitis, sinusitis, rhinosinusitis, chronic otitis media, recurrent otitis media, drug reactions, insect sting reactions, latex reactions, conjunctivitis, urticaria, anaphylaxis reactions, anaphylactoid reactions, atopic dermatitis, asthma, or food allergies.
57. A method of inhibiting immune cell activation comprising administering to the cell a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or
 5 prodrug thereof, wherein:

Y is a monocyclic optionally substituted aryl or a monocyclic
 optionally substituted heteroaryl;

L is a linker selected from the group consisting of $-NRCR_2-$,
 $-CR_2NR-$, $-C(O)-$, $-NR-C(O)-$, $-C(O)-NR-$, $-OC(O)-$, $-C(O)O-$, $-C(S)-$,
 10 $-NR-C(S)-$, $-C(S)-NR-$;

one of X_1 or X_2 is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a
 lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl
 sufanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$,
 15 cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from $-H$ or an
 alkyl;

R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$,
 or $-NR_6R_7$;

R_4 is a halo, a haloalkyl, cyano, nitro, $-C(O)R_5$, $-C(O)OR_5$,
 $-C(O)SR_5$, $-C(O)NR_9R_{10}$, $-C(S)R_5$, $-C(S)OR_5$, $-C(S)SR_5$, $-C(S)NR_6R_7$,
 $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-C(NR_8)SR_5$, $-C(NR_8)NR_6R_7$, $-S(O)_pR_5$,
 $-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an
 20 optionally substituted heterocyclyl, or an optionally substituted
 25 heteroaryl;

R_5 , for each occurrence, is independently, H, an optionally
 substituted alkyl, an optionally substituted alkenyl, an optionally


substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

5 R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

15 R_8 , for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indoliziny, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted

- benzofuryl, an optionally substituted benzothiazolyl, an optionally substituted indoliziny, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;
- q, for each occurrence, is independently, an integer from 1 to 3;
k is for each occurrence, is independently, an integer from 1 to 4;
n is zero, 1 or 2; and
p, for each occurrence, is independently 1 or 2.
58. The method of Claim 57, wherein immune cell activation is inhibited in a subject by administering the compound to the subject.
59. The method of Claim 58, wherein the subject is human.
60. A method of inhibiting cytokine production in a cell, comprising administering to the cell a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

5 Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of $-NRCR_2-$, $-CR_2NR-$, $-C(O)-$, $-NR-C(O)-$, $-C(O)NR-$, $-OC(O)-$, $-C(O)O-$, $-C(S)-$, $-NR-C(S)-$, $-C(S)NR-$;

10 one of X_1 or X_2 is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$, cyano, nitro, or lower haloalkoxy;

15 R, for each occurrence is independently selected from $-H$ or an alkyl;

R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$, or $-NR_6R_7$;

20 R_4 is a halo, a haloalkyl, cyano, nitro, $-C(O)R_5$, $-C(O)OR_5$, $-C(O)SR_5$, $-C(O)NR_9R_{10}$, $-C(S)R_5$, $-C(S)OR_5$, $-C(S)SR_5$, $-C(S)NR_6R_7$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-C(NR_8)SR_5$, $-C(NR_8)NR_6R_7$, $-S(O)_pR_5$, $-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

25 R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally

substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

5 R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;


R_8 , for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

15 R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indolizinyl, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally

5 substituted indolizinyl, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

15 q, for each occurrence, is independently, an integer from 1 to 3;
k is for each occurrence, is independently, an integer from 1 to 4;
n is zero, 1 or 2; and
p, for each occurrence, is independently 1 or 2.

- 20 61. The method of Claim 60, wherein cytokine production is inhibited in a subject by administering the compound to the subject.
62. The method of Claim 61, wherein the subject is human.
- 25 63. The method of Claim 61, wherein the cytokine is selected from the group consisting of IL-2, IL-4, IL-5, IL-13, GM-CSF, IFN- γ , TNF- α , and combinations thereof.
64. The method of Claim 63, wherein the cytokine is IL-2.
- 30 65. A method of modulating an ion channel in a cell, wherein the ion channel is involved in immune cell activation, comprising administering to the cell a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

5 Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of -NR₁CR₂-, -CR₂NR₁-, -C(O)-, -NR₁-C(O)-, -C(O)-NR₁-, -OC(O)-, -C(O)O-, -C(S)-, -NR₁-C(S)-, -C(S)-NR₁-;

10 one of X₁ or X₂ is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, -S(O)_p-alkyl, -C(O)NRR, -(CH₂)_kNRR, -(CH₂)_kOR, -(CH₂)_kSR, cyano, nitro, or lower haloalkoxy;

15 R, for each occurrence is independently selected from -H or an alkyl;

R₃ is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, -OR₅, -SR₅, or -NR₆R₇;

20 R₄ is a halo, a haloalkyl, cyano, nitro, -C(O)R₅, -C(O)OR₅, -C(O)SR₅, -C(O)NR₉R₁₀, -C(S)R₅, -C(S)OR₅, -C(S)SR₅, -C(S)NR₆R₇, -C(NR₈)R₅, -C(NR₈)OR₅, -C(NR₈)SR₅, -C(NR₈)NR₆R₇, -S(O)_pR₅, -S(O)_pNR₅, -S(O)_pOR₅, -P(O)(OR₅)₂, -OP(O)(OR₅)₂, -P(O)(R₅)₂, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

25 R₅, for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally

substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

5 R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

R_8 , for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

15 R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indoliziny, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally

substituted indoliziny, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;


q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.

66. The method of Claim 65, wherein the ion channel is in a subject and it is modulated by administering the compound to the subject.

67. The method of Claim 66, wherein the subject is human.

68. The method of Claim 66, wherein the ion channel is a Ca²⁺-release-activated Ca²⁺ channel (CRAC).

69. A method of inhibiting T-cell and/or B-cell proliferation in response to an antigen, comprising administering to the cell a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

5 Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of -NR₁CR₂-, -CR₂NR₁-, -C(O)-, -NR-C(O)-, -C(O)-NR-, -OC(O)-, -C(O)O-, -C(S)-, -NR-C(S)-, -C(S)-NR-;

10 one of X₁ or X₂ is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, -S(O)_p-alkyl, -C(O)NRR, -(CH₂)_kNRR, -(CH₂)_kOR, -(CH₂)_kSR, cyano, nitro, or lower haloalkoxy;

15 R, for each occurrence is independently selected from -H or an alkyl;

R₃ is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, -OR₅, -SR₅, or -NR₆R₇;

20 R₄ is a halo, a haloalkyl, cyano, nitro, -C(O)R₅, -C(O)OR₅, -C(O)SR₅, -C(O)NR₉R₁₀, -C(S)R₅, -C(S)OR₅, -C(S)SR₅, -C(S)NR₆R₇, -C(NR₈)R₅, -C(NR₈)OR₅, -C(NR₈)SR₅, -C(NR₈)NR₆R₇, -S(O)_pR₅, -S(O)_pNR₅, -S(O)_pOR₅, -P(O)(OR₅)₂, -OP(O)(OR₅)₂, -P(O)(R₅)₂, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

25 R₅, for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally

substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

5 R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

R_8 , for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

15 R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indolizinyl, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally

20


25

30

substituted indoliziny, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.

70. The method of Claim 69, wherein T-cell and/or B-cell proliferation is inhibited in a subject by administering the compound to the subject.
71. The method of Claim 70, wherein the subject is human.
72. A method for treating or preventing an immune disorder in a subject in need thereof, comprising administering to the subject an effective amount of a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

5 Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of $-NRCR_2-$, $-CR_2NR-$, $-C(O)-$, $-NR-C(O)-$, $-C(O)-NR-$, $-OC(O)-$, $-C(O)O-$, $-C(S)-$, $-NR-C(S)-$, $-C(S)-NR-$;

10 one of X_1 or X_2 is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$, cyano, nitro, or lower haloalkoxy;

15 R, for each occurrence is independently selected from $-H$ or an alkyl;

R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$, or $-NR_6R_7$;

20 R_4 is a halo, a haloalkyl, cyano, nitro, $-C(O)R_5$, $-C(O)OR_5$, $-C(O)SR_5$, $-C(O)NR_9R_{10}$, $-C(S)R_5$, $-C(S)OR_5$, $-C(S)SR_5$, $-C(S)NR_6R_7$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-C(NR_8)SR_5$, $-C(NR_8)NR_6R_7$, $-S(O)_pR_5$, $-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

25 R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally

substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

5 R₆ and R₇, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₆ and R₇ taken together with the nitrogen to which they
10 are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

 R₈, for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;


15 R₉ and R₁₀, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an
20 optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indoliziny, an
25 optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an
30 optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally

5 substituted indoliziny, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉ and R₁₀ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

15 q, for each occurrence, is independently, an integer from 1 to 3;
k is for each occurrence, is independently, an integer from 1 to 4;
n is zero, 1 or 2; and
p, for each occurrence, is independently 1 or 2.

- 20 73. The method of Claim 72, wherein the subject is human.
74. The method of Claim 72, wherein the disorder is selected from the group consisting of multiple sclerosis, myasthenia gravis, Guillain-Barré, autoimmune uveitis, autoimmune hemolytic anemia, pernicious anemia, autoimmune thrombocytopenia, temporal arteritis, anti-phospholipid syndrome, vasculitides such as Wegener's granulomatosis, Behcet's disease, psoriasis, dermatitis herpetiformis, pemphigus vulgaris, vitiligo, Crohn's disease, ulcerative colitis, primary biliary cirrhosis, autoimmune hepatitis, Type 1 or immune-mediated diabetes mellitus, Grave's disease, Hashimoto's thyroiditis, autoimmune oophoritis and orchitis, autoimmune disorder of the adrenal gland, rheumatoid arthritis, systemic lupus erythematosus, scleroderma, polymyositis, dermatomyositis, ankylosing spondylitis, and Sjogren's syndrome.
- 25
30

75. A method for treating or preventing an inflammatory condition in a subject in need thereof, comprising administering to the subject an effective amount of a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of -NR₁CR₂-, -CR₂NR₁-, -C(O)-, -NR₁-C(O)-, -C(O)-NR₁-, -OC(O)-, -C(O)O-, -C(S)-, -NR₁-C(S)-, -C(S)-NR₁-;

one of X₁ or X₂ is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, -S(O)_p-alkyl, -C(O)NRR, -(CH₂)_kNRR, -(CH₂)_kOR, -(CH₂)_kSR, cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from -H or an alkyl;

R₃ is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, -OR₅, -SR₅, or -NR₆R₇;

R₄ is a halo, a haloalkyl, cyano, nitro, -C(O)R₅, -C(O)OR₅, -C(O)SR₅, -C(O)NR₉R₁₀, -C(S)R₅, -C(S)OR₅, -C(S)SR₅, -C(S)NR₆R₇, -C(NR₈)R₅, -C(NR₈)OR₅, -C(NR₈)SR₅, -C(NR₈)NR₆R₇, -S(O)_pR₅, -S(O)_pNR₅, -S(O)_pOR₅, -P(O)(OR₅)₂, -OP(O)(OR₅)₂, -P(O)(R₅)₂, an

optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

5 R₅, for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 R₆ and R₇, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R₆ and R₇ taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;


20 R₈, for each occurrence, is independently -H, a halo, an alkyl, -OR₅, -NR₆R₇, -C(O)R₅, -C(O)OR₅, or -C(O)NR₆R₇;

25 R₉ and R₁₀, for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indolizinyl, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl, an optionally substituted triazinyl, an optionally substituted triazolyl, an

- optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an optionally substituted pyrazinyl, an optionally substituted quinolinyl, an optionally substituted isoquinolinyl, an optionally substituted indazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzofuryl, an optionally substituted benzothiazolyl, an optionally substituted indolizinyl, an optionally substituted imidazopyridinyl, an optionally substituted tetrazolyl, an optionally substituted benzimidazolyl, an optionally substituted benzoxazolyl, an optionally substituted benzothiazolyl, an optionally substituted benzothiadiazolyl, an optionally substituted benzoxadiazolyl, an optionally substituted indolyl, an optionally substituted tetrahydroindolyl, an optionally substituted azaindolyl, an optionally substituted imidazopyridyl, an optionally substituted quinazolinyl, an optionally substituted purinyl, an optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R_9 and R_{10} taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;
- q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.
76. The method of Claim 75, wherein the subject is human.
77. The method according to Claim 75, wherein the disorder is selected from transplant rejection, skin graft rejection, arthritis, rheumatoid arthritis, osteoarthritis and bone diseases associated with increased bone resorption; inflammatory bowel disease, ileitis, ulcerative colitis, Barrett's syndrome, Crohn's disease; asthma, adult respiratory distress syndrome, chronic obstructive airway disease; corneal dystrophy, trachoma, onchocerciasis, uveitis, sympathetic ophthalmitis, endophthalmitis; gingivitis, periodontitis; tuberculosis; leprosy; uremic

complications, glomerulonephritis, nephrosis; sclerodermatitis, psoriasis, eczema; chronic demyelinating diseases of the nervous system, multiple sclerosis, AIDS-related neurodegeneration, Alzheimer's disease, infectious meningitis, encephalomyelitis, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis viral or autoimmune encephalitis; autoimmune disorders, immune-complex vasculitis, systemic lupus and erythematoses; systemic lupus erythematosus (SLE); cardiomyopathy, ischemic heart disease hypercholesterolemia, atherosclerosis, preeclampsia; chronic liver failure, brain and spinal cord trauma, and cancer.

78. A method for suppressing the immune system of a subject in need thereof, comprising administering to the subject an effective amount of a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of -NR₂CR₂-, -CR₂NR-, -C(O)-, -NR-C(O)-, -C(O)-NR-, -OC(O)-, -C(O)O-, -C(S)-, -NR-C(S)-, -C(S)-NR-;

one of X₁ or X₂ is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl

sufanyl, $-S(O)_p$ -alkyl, $-C(O)NRR$, $-(CH_2)_kNRR$, $-(CH_2)_kOR$, $-(CH_2)_kSR$, cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from $-H$ or an alkyl;

5 R_3 is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, $-OR_5$, $-SR_5$, or $-NR_6R_7$;

R_4 is a halo, a haloalkyl, cyano, nitro, $-C(O)R_5$, $-C(O)OR_5$, $-C(O)SR_5$, $-C(O)NR_9R_{10}$, $-C(S)R_5$, $-C(S)OR_5$, $-C(S)SR_5$, $-C(S)NR_6R_7$, $-C(NR_8)R_5$, $-C(NR_8)OR_5$, $-C(NR_8)SR_5$, $-C(NR_8)NR_6R_7$, $-S(O)_pR_5$, $-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

10

R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

15

R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

20

25


R_8 , for each occurrence, is independently $-H$, a halo, an alkyl, $-OR_5$, $-NR_6R_7$, $-C(O)R_5$, $-C(O)OR_5$, or $-C(O)NR_6R_7$;

30 R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an

optionally substituted heterocyclyl, an optionally substituted aryl, an
 optionally substituted aralkyl, an optionally substituted heteraralkyl, or
 an optionally substituted heteroaryl selected from the group consisting
 of an optionally substituted pyridinyl, an optionally substituted thienyl, an
 5 optionally substituted pyrrolyl, an optionally substituted oxazolyl, an
 optionally substituted imidazolyl, an optionally substituted indoliziny, an
 optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an
 optionally substituted isothiazolyl, an optionally substituted pyridazinyl,
 an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl,
 10 an optionally substituted triazinyl, an optionally substituted triazolyl, an
 optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an
 optionally substituted pyrazinyl, an optionally substituted quinolinyl, an
 optionally substituted isoquinolinyl, an optionally substituted indazolyl,
 an optionally substituted benzoxazolyl, an optionally substituted
 15 benzofuryl, an optionally substituted benzothiazolyl, an optionally
 substituted indoliziny, an optionally substituted imidazopyridinyl, an
 optionally substituted tetrazolyl, an optionally substituted
 benzimidazolyl, an optionally substituted benzoxazolyl, an optionally
 substituted benzothiazolyl, an optionally substituted benzothiadiazolyl,
 20 an optionally substituted benzoxadiazolyl, an optionally substituted
 indolyl, an optionally substituted tetrahydroindolyl, an optionally
 substituted azaindolyl, an optionally substituted imidazopyridyl, an
 optionally substituted quinazolinyl, an optionally substituted purinyl, an
 optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted
 25 pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉
 and R₁₀ taken together with the nitrogen to which they are attached are
 an optionally substituted heterocyclyl or optionally substituted
 heteroaryl;

30 q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.

79. The method of Claim 78, wherein the subject is human.
80. A method for treating or preventing an allergic disorder in a subject in need thereof, comprising administering to the subject an effective amount of a compound represented by structural formula (I):



(I)

or a pharmaceutically acceptable salt, solvate, clathrate, or prodrug thereof, wherein:

Y is a monocyclic optionally substituted aryl or a monocyclic optionally substituted heteroaryl;

L is a linker selected from the group consisting of -NRCR₂-, -CR₂NR-, -C(O)-, -NR-C(O)-, -C(O)-NR-, -OC(O)-, -C(O)O-, -C(S)-, -NR-C(S)-, -C(S)-NR-;

one of X₁ or X₂ is CH or CZ and the other is N;

each Z is independently selected from the group consisting of a lower alkyl, a lower haloalkyl, a halo, a lower alkoxy, a lower alkyl sulfanyl, -S(O)_p-alkyl, -C(O)NRR, -(CH₂)_kNRR, -(CH₂)_kOR, -(CH₂)_kSR, cyano, nitro, or lower haloalkoxy;

R, for each occurrence is independently selected from -H or an alkyl;

R₃ is H, an alkyl, a haloalkyl, a halo, a haloalkoxy, -OR₅, -SR₅, or -NR₆R₇;

R₄ is a halo, a haloalkyl, cyano, nitro, -C(O)R₅, -C(O)OR₅, -C(O)SR₅, -C(O)NR₉R₁₀, -C(S)R₅, -C(S)OR₅, -C(S)SR₅, -C(S)NR₆R₇, -C(NR₈)R₅, -C(NR₈)OR₅, -C(NR₈)SR₅, -C(NR₈)NR₆R₇, -S(O)_pR₅,

$-S(O)_pNR_5$, $-S(O)_pOR_5$, $-P(O)(OR_5)_2$, $-OP(O)(OR_5)_2$, $-P(O)(R_5)_2$, an optionally substituted heterocyclyl, or an optionally substituted heteroaryl;

5 R_5 , for each occurrence, is independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl;

10 R_6 and R_7 , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted heteroaryl, an optionally substituted aralkyl, or an optionally substituted heteraralkyl; or R_6 and R_7 taken together with the nitrogen to which they are attached are an optionally substituted heterocyclyl or optionally substituted heteroaryl;

20 R_8 , for each occurrence, is independently $-H$, a halo, an alkyl, $-OR_5$, $-NR_6R_7$, $-C(O)R_5$, $-C(O)OR_5$, or $-C(O)NR_6R_7$;


25 R_9 and R_{10} , for each occurrence are, independently, H, an optionally substituted alkyl, an optionally substituted alkenyl, an optionally substituted alkynyl, an optionally substituted 5- to 14-membered cycloalkyl, an optionally substituted cycloalkenyl, an optionally substituted heterocyclyl, an optionally substituted aryl, an optionally substituted aralkyl, an optionally substituted heteraralkyl, or an optionally substituted heteroaryl selected from the group consisting of an optionally substituted pyridinyl, an optionally substituted thienyl, an optionally substituted pyrrolyl, an optionally substituted oxazolyl, an optionally substituted imidazolyl, an optionally substituted indoliziny, an optionally substituted isoxazolyl, an optionally substituted pyrazolyl, an optionally substituted isothiazolyl, an optionally substituted pyridazinyl, an optionally substituted pyrimidinyl, an optionally substituted pyrazinyl,

an optionally substituted triazinyl, an optionally substituted triazolyl, an
 optionally substituted pyridinyl, an optionally substituted thiadiazolyl, an
 optionally substituted pyrazinyl, an optionally substituted quinolinyl, an
 optionally substituted isoquinolinyl, an optionally substituted indazolyl,
 5 an optionally substituted benzoxazolyl, an optionally substituted
 benzofuryl, an optionally substituted benzothiazolyl, an optionally
 substituted indolizinyl, an optionally substituted imidazopyridinyl, an
 optionally substituted tetrazolyl, an optionally substituted
 benzimidazolyl, an optionally substituted benzoxazolyl, an optionally
 10 substituted benzothiazolyl, an optionally substituted benzothiadiazolyl,
 an optionally substituted benzoxadiazolyl, an optionally substituted
 indolyl, an optionally substituted tetrahydroindolyl, an optionally
 substituted azaindolyl, an optionally substituted imidazopyridyl, an
 optionally substituted quinazolinyl, an optionally substituted purinyl, an
 15 optionally substituted pyrrolo[2,3]pyrimidyl, an optionally substituted
 pyrazolo[3,4]pyrimidyl or an optionally substituted benzo(b)thienyl; or R₉
 and R₁₀ taken together with the nitrogen to which they are attached are
 an optionally substituted heterocyclyl or optionally substituted
 heteroaryl;

20 q, for each occurrence, is independently, an integer from 1 to 3;
 k is for each occurrence, is independently, an integer from 1 to 4;
 n is zero, 1 or 2; and
 p, for each occurrence, is independently 1 or 2.

25 81. The method of Claim 80, wherein the subject is human.

82. The method of Claim 80, wherein the disorder is allergic rhinitis,
 sinusitis, rhinosinusitis, chronic otitis media, recurrent otitis media, drug
 reactions, insect sting reactions, latex reactions, conjunctivitis, urticaria,
 30 anaphylaxis reactions, anaphylactoid reactions, atopic dermatitis,
 asthma, or food allergies.



(I)