



(86) Date de dépôt PCT/PCT Filing Date: 2001/03/15
(87) Date publication PCT/PCT Publication Date: 2001/09/20
(45) Date de délivrance/Issue Date: 2012/08/07
(85) Entrée phase nationale/National Entry: 2002/09/09
(86) N° demande PCT/PCT Application No.: KR 2001/000408
(87) N° publication PCT/PCT Publication No.: 2001/068802
(30) Priorités/Priorities: 2000/03/15 (KR2000/13133);
2001/03/13 (KR2001/12987)


(51) Cl.Int./Int.Cl. *C12N 15/24* (2006.01),
A61K 38/20 (2006.01), *A61K 39/39* (2006.01),
C07K 14/54 (2006.01), *C12N 15/00* (2006.01),
C12N 15/63 (2006.01), *C12N 15/85* (2006.01)

(72) Inventeurs/Inventors:
SUNG, YOUNG CHUL, KR;
LEE, SUNG HEE, KR;
HA, SANG JUN, KR;
SONG, MAN KI, KR;
CHANG, JUN, KR

(73) Propriétaire/Owner:
GENEXINE CO., LTD., KR

(74) Agent: SMART & BIGGAR

(54) Titre : GENES DE LA SOUS-UNITE IL-12P40 SOUMIS A MUTATION AFIN D'AMELIORER L'ACTIVITE D'IL-12, ET
UTILISATION DESDITS GENES DANS UN ADJUVANT DE VACCIN GENETIQUE
(54) Title: GENES OF IL-12P40 SUBUNIT MUTATED FOR IMPROVING THE ACTIVITY OF IL-12 AND USE THEREOF
FOR DNA VACCINE ADJUVANT



(57) Abrégé/Abstract:

The present invention relates to the IL-12p40 subunit mutant gene which can produce IL-12 (interleukin 12) of human and mouse origin with high activity, the expression vector including above mutant gene and the use of them to DNA vaccine adjuvant.

(57) Abrégé(suite)/Abstract(continued):

Particularly, it relates to IL-12p40 mutant gene which inhibits the secretion of IL-12p40 but normally secretes active IL-12p70 by making mutation at Asn-222 (human) or Asn-220 (mouse) amino acid of IL-12p40, which acts competitive inhibitor of active form of IL-12, IL-12p70. Therefore, the IL-12p40 mutant gene of the present invention can be useful for DNA vaccination and gene therapy to various diseases, for example, AIDS, hepatitis C or hepatitis B, cancer, influenza, tuberculosis and malaria, which essentially require cellular immune responses for thier therapy.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
20 September 2001 (20.09.2001)

PCT

(10) International Publication Number
WO 01/68802 A3(51) International Patent Classification⁷: **C12N 15/00,**
15/24[KR/KR]; 47-5, Sanggal-Ri, Kiheung-Up, Yongin-si
449-900, Kyunggi-do (KR).

(21) International Application Number: PCT/KR01/00408

(72) Inventors; and

(22) International Filing Date: 15 March 2001 (15.03.2001)

(75) Inventors/Applicants (for US only): **SUNG, Young, Chul**


(25) Filing Language: Korean

[KR/KR]; #101-601 Hyundai-leedong Apt., Daezam-dong,
Nam-ku, Pohang-si 790-310, Kyungsangbuk-do (KR).

(26) Publication Language: English

LEE, Sung, Hee [KR/KR]; #Na-101, Songee Man-
sion 148-148, Bongchun 6 dong, Kwanak-ku, Seoul(30) Priority Data:
2000/13133 15 March 2000 (15.03.2000) KR
2001/12987 13 March 2001 (13.03.2001) KR151-056 (KR). **HA, Sang, Jun** [KR/KR]; #236-702
Green Apt., Jigok-dong, Nam-ku, Pohang-si 790-390,(71) Applicants (for all designated States except US):
POHANG UNIVERSITY OF SCIENCE AND TECH-
NOLOGY [KR/KR]; San 31, Hyoja-dong, Pohang
790-784, Kyung-buk (KR). **PROGEN CO., LTD.**Kyungsangbuk-do (KR). **SONG, Man, Ki** [KR/KR]; Dae-
hakwon Apt., Jikok-dong, Nam-ku, Pohang-si 790-390,Kyungsangbuk-do (KR). **CHANG, Jun** [KR/KR]; Po-
hang University of Science and Technology, Dept. ofLife science, San 31, Hyoja-dong, Pohang-si 790-330,
Kyungsangbuk-do (KR).(74) Agent: **LEE, Won-Hee**; 8th Floor, Sung-ji Heights II,
642-16 Yoksam-dong, Kangnam-ku, Seoul 135-080 (KR).

[Continued on next page]

(54) Title: GENES OF IL-12P40 SUBUNIT MUTATED FOR IMPROVING THE ACTIVITY OF IL-12 AND USE THEREOF
FOR DNA VACCINE ADJUVANT

(57) Abstract: The present invention relates to the IL-12p40 subunit mutant gene which can produce IL-12 (interleukin 12) of human and mouse origin with high activity. the expression vector including above mutant gene and the use of them to DNA vaccine adjuvant. Particularly, it relates to IL-12p40 mutant gene which inhibits the secretion of IL-12p40 but normally secretes active IL-12p70 by making mutation at Asn-222 (human) or Asn-220 (mouse) amino acid of IL-12p40, which acts competitive inhibitor of active form of IL-12, IL-12p70. Therefore, the IL-12p40 mutant gene of the present invention can be useful for DNA vaccination and gene therapy to various diseases, for example, AIDS, hepatitis C or hepatitis B, cancer, influenza, tuberculosis and malaria, which essentially require cellular immune responses for thier therapy.

WO 01/68802 A3



(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

– with international search report

(88) Date of publication of the international search report:

21 February 2002

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

**Genes of IL-12p40 subunit mutated for improving the activity of
IL-12 and use thereof for DNA vaccine adjuvant**

FIELD OF THE INVENTION

5 The present invention relates to mutant IL-12p40
subunit genes producing highly active human and mouse
Interleukin 12(IL-12), expression vector which contains
those mutant genes and methods to use thereof for DNA
vaccine adjuvant. Particularly, this invention relates
10 to mutant IL-12p40 subunit genes which can secrete
immunoactive IL-12p70 normally but reduce secretion of
IL-12p40 by inducing mutation for amino acids Asn-
222(mouse) or Asn-220(human) of IL-12p40 working as a
competitive inhibitor to active IL-12.

15

BACKGROUND ART OF THE INVENTION

IL-12 is secreted by antigen presenting cells
(APC) such as macrophages, monocytes and B cells after
getting appropriate stimulation, and works as a
20 regulator for many kinds of immune response *in vivo*.
Particularly, IL-12 has a broad range of activities
including proliferation of activated type 1 helper T
(Th1) and natural killer(NK) cells, regulation of the
production of many cytokines, induction of type 1 T
25 helper cell immune responses, differentiation of CD8+ T

WO 01/68802

PCT/KR01/00408

cells, stimulation of hematopoietic stem cells(Hsieh, C. S. et al., *Science*, 260:547-549, 1993), and especially, regulation of immune response by promoting lytic activity of cytotoxic T lymphocytes(CTL) and NK
5 cells(Robertson, M. J. and J. Ritz., *Oncologist*, 1:88-97, 1999; Trinchieri, G., *Annu. Rev. Immunol.*, 13:251-276, 1995). According to recent reports, peripheral blood mononuclear cells (PBMC) from human immunodeficiency virus (HIV)-infected patients produced
10 five-fold less biologically active IL-12(Chehimi, J. et al., *J. Exp. Med.*, 179:1361-1366, 1994), and mycobacterial and salmonella infections were found to lack of IL-12 receptor expression(de Jong, R. et al., *Science*, 280:1435-1438). Due to these roles of IL-12,
15 it could induce strong immune response *in vivo* against virus, bacteria and various cancers. Therefore, the present inventors can expect development of many treatment agents using the present invention.

On the basis of the theory that IL-12 is related
20 to the growth of memory Th1 and memory CTL cells(Stobie, L. et al., *Proc. Natl. Acad. Sci. USA*, 97:8427-8432, 2000; Mortarini, R. et al., *Cancer Res.*, 60:3559-3568, 2000; Mbawuike, I. N. et al., *J. Infect. Dis.*, 180:1477-1486, 1999), IL-12 is believed to be used as a
25 effective vaccine adjuvant or a treatment agent against various diseases requiring cellular immune response. Considering metastasis or recurrence, which is the most

WO 01/68802

PCT/KR01/00408

difficult part of the cancer treatment, the inducing of memory immune response is especially necessary. However, as of today the exact mechanism of IL-12, which is connected to those effects, has not been cleared. But, based on current available data, these are some clues to the mechanism by which IL-12 sustains CD4+ Th1 cells. In the course of Th1 differentiation, production of IFN- γ is enhanced and IL-2 is diminished. Because IL-2 is a potent growth factor and IFN- γ is antiproliferative, IL-12 could function to sustain cell growth and viability or to prevent apoptosis of CD4+ IFN- γ T cells (Fuss, I. J. et al., *Gastroenterology*, 117:1078-1088, 1999; Marth, T. et al., *J. Immunol.*, 162:7233-7240, 1999). Also, IFN- γ which is increased by IL-12 augments expression of IL-15 (Zhang, X. et al., *Immunity*, 8:591-599, 1998), which is related to memory CD8+ T cell. Based on those reports, IL-12 is very useful in vaccine immunization since IL-12 is related not only to the early immune response but also to the memory immune response.

The biologically functional form of IL-12 is a 70 kDa heterodimer, IL-12p70, which consists of disulfide-bonded p40 and p35 subunits. The p40 subunit shares amino acid sequence homology with the IL-6 receptor, thus it belongs to the cytokine receptor superfamily, whereas p35 subunit has a distant but a significant relationship to the IL-6/granulocyte colony stimulating

WO 01/68802

PCT/KR01/00408

factor cytokine family(Gearing, D. P., and Cosman, D.,
Cell, 66:9-10, 1991).

The IL-12p40 is secreted as both a monomer and a
homodimer in large excess over IL-12p70 both *in*
5 *vitro*(D'Andrea et al., *J. Exp. Med.*, 176:1387-1398,
1992; Podlasky, F. J. et al., *Arch. Biochem. Biophys.*,
294:230-237, 1992) and *in vivo*(Mattner, F. et al., *Eur.*
J. Immunol., 23:2202-2208, 1993; Heinzl, F. P. et al.,
Infect. Immun., 62:4244-4249, 1994). It has been
10 demonstrated that IL-12p40 strongly antagonizes IL-
12p70-mediated responses by binding competitively to
IL-12 receptor *in vitro*(Gillesen, S. et al., *Eur. J.*
Immunol., 25:200-206, 1995; Ling, P. et al., *J.*
Immunol., 154:116-127, 1995), suggesting the critical
15 role of IL-12p40 as a natural antagonist of IL-12p70.
This is supported by several observations including a
reduction in Th1 responses in IL-12p40 transgenic
mice(Yoshimoto, T. et al., *J. Immunol.*, 160:588-594,
1998), the prevention of allogeneic rejection in
20 transplanted myoblasts engineered to produce IL-
12p40(Kato, K. et al., *Proc. Natl. Acad. Sci., USA*,
93:9085-9089, 1996), and the inhibition of tumor
suppression activity of IL-12p70 by adenovirus
expressing IL-12p40(Chen, L. et al., *J. Immunol.*,
25 159:351-359, 1997). In contrast, there is a report
describing a positive role of IL-12p40 in IL-12p70-
mediated responses resulting in alloantigen-specific

WO 01/68802

PCT/KR01/00408

Th1 development under certain condition(Piccotti, J. R.
et al., *J. Immunol.*, 157:1951-1957, 1996). Furthermore,
a novel function of IL-12p40 as a chemotactic molecule
for macrophage was known by this inventors(Ha, S. J. et
5 al., *J. Immunol.*, 163:2902-2908, 1999).

Various *in vivo* systems have been established in
order to prove the anti-cancer effect of IL-12p70.
However, there is a significant side-effect which could
result in death if human recombinant IL-12p70 protein
10 is directly injected into a cancer patient(Robertson, M.
J. and Ritz, J., *Oncologist*, 1:88-97, 1996; Cohen, J.,
Science, 270:908, 1995). To prevent this side-effect
and to be effective in economy, lots of studies have
been conducted on gene therapy by using IL-12 gene,
15 which now proves that IL-12 gene treatment is very
effective and has no harm(Rakhmilevich, A. L. et al.,
Proc. Natl. Acad. Sci. USA, 93:6291-6296, 1996; Tahara,
H. et al., *J. Immunol.*, 154:6466-6474, 1995; Lotze, M.
T. et al., *Ann. N. Y. Acad. Sci.*, 795:440-454, 1996).

20 IL-12p70, a biologically active form of IL-12, is
essential(Gulber, U. et al., *Proc. Natl. Acad. Sci. USA*,
88:4143-4147) for the gene-therapy using IL-12 and,
cDNAs of p35 and p40 should be expressed in a cell at
the same time. Many methods have been used to
25 simultaneously express p35 and p40 subunit together in
one cell. One of them is to use expression cassette
through which p35 and p40 are placed in consecutive row

WO 01/68802

PCT/KR01/00408

and expressed each by each(Rakhmievich, A. L. *et al.*,
Proc. Natl. Acad. Sci. USA, 93:6291-6296, 1996).
Another one is to use internal ribosomal entry
site(IRES) of encephalomyocarditis virus(EMCV) for
5 simultaneous expression of both genes. These methods
were successful in expressing IL-12p70 in one cell, but
as mentioned above, not successful in removing the
possibility that continuously expressed excessive IL-
12p40 inhibits the biological action of IL-12p70.

10 To overcome this intrinsic defect, the genetic
linkage of p35 subunit followed by p40 subunit was
recently proposed by means of a DNA sequence encoding a
protein linker commonly used in antibody
engineering(Lieschke, G. J. *et al.*, *Nat. Biotechnol.*,
15 15:35-40, 1997; Lode, H. N. *et al.*, *Proc. Natl. Acad.*
Sci. USA, 95:2475-2480, 1998; Lee, Y. L. *et al.*, *Human*
Gene Ther., 9:457-465, 1998). Antagonistic effect of
excessive IL-12p40 against biological activity of IL-
12p70 could be surmounted in the above method. However,
20 it is still not good enough having a problem that the
activity of IL-12p70 was 5-100 times more decreased
since it's structure may be changed when p35 and p40
are linked together. For the effective treatment
against cancer or other diseases with IL-12 gene, it is
25 required to induce IL-12p70 still having the same
activity and prevent the secretion of IL-12p40.

Glycosylation has been known to contribute protein

WO 01/68802

PCT/KR01/00408

folding, secretion, conformation, stability and biological activity. The p35 and p40 subunits of human IL-12 express 219 and 328 amino acids that contain 56 and 22 amino acids of hydrophobic signal sequences, respectively. The analysis of human IL-12 amino acid sequence reveals three and four putative N-glycosylation sites within p35 and p40 subunits, respectively(Podlasky, F. J. et al., *Arch. Biochem. Biophys.*, 294:230-237, 1992). Stern et al. reported that after treatment of human IL-12 with trifluoromethanesulfonic acid or glycosidase F, IL-12p35 and IL-12p40 were reduced in molecular weight(Podlasky, F. J. et al., *Arch. Biochem. Biophys.*, 294:230-237, 1992). This suggests that p35 and p40 subunits of human IL-12 are composed of carbohydrates. In the same report, it was also demonstrated that the digestion of IL-12p35 with neuramidase followed by endo- α -N-acetylgalactosaminidase reduced its molecular weight, whereas IL-12p40 was unaffected by such treatment. These experiments indicate that the glycosylation of IL-12p35 is contributed by O-linked oligosaccharides and that IL-12p40 has no O-linked carbohydrates. And, in the analysis of N-glycosylation at Asn-135 and Asn-222 amino acid residues, it was revealed that Asn-222 is the N-glycosylation site. However the exact N-glycosylation sites of human and mouse IL-12 and its effects on the synthesis, secretion and biological

WO 01/68802

PCT/KR01/00408

activity of IL-12 have not been defined.

On the other hand, the study using IL-12 gene has been accelerated since cell-mediated immune response is required rather than humoral immune response for the prevention and treatment of many viral or bacterial diseases along with suppression of cancer formation. Hepatitis C is the representative virus-mediated disease and once infected with HCV, more than 50% patients become chronic and finally lead to liver cirrhosis or liver cancer(Alter, H.J. et al., *N. Engl. J. Med.*, 321:1494-1500, 1989). As of today, only α -interferon is known as a treatment agent for hepatitis C, but the effect is not good enough(10-30%)(Weiland, E. et al. *J. Virol.*, 66:3677-3682, 1992). So, more effective vaccine or treatment agents against HCV are urgently required. According to the medical test reports including both human and chimpanzee, HCV is related to specific humoral immune response and cell-mediated immune response as well(Prince, A. M. et al., *J. Infect. Dis.*, 165:438-443, 1992), and E1 and E2, structural protein of HCV is reported as a major antigen to induce protective immunity(Choo, Q. L. et al., *Proc. Natl. Acad. Sci. USA*, 91:1294-1298, 1994). Once more, cell-mediated immune response including CTL is preferably needed to remove HCV compared to humoral immune response(Cooper, S. et al., *Immunity*, 10:439-449, 1999; Rinaldo, C. et al., *J. Virol.*, 69:5838-5842,

WO 01/68802

PCT/KR01/00408

1995).

DNA immunization is the most recent method to induce cell-mediated immune response. DNA immunization is differentiated with the existing one using dead or
5 detoxicated pathogen or certain parts of pathogen in the matter of inserting DNA coding a specific component of pathogen directly to the human body. DNA immunization is also known to induce strong immune response against various infectious virus such as
10 influenza, hepatitis B and human immunodeficiency virus (Ulmer, J. B. et al., *Science*, 259:1745-1749, 1993; Michel, M. L. et al., *Proc. Natl. Acad. Sci. USA*, 92:5307-5311, 1995; Irwin, M. J. et al., *J. Virol.*, 68:5306-5044, 1994). In addition, DNA immunization is
15 also reported to induce special immune response against capsid and E2 protein of HCV (Major, M. E. et al., *J. Virol.*, 69:5798-5805, 1995; Tedeschi, V. et al., *Hepatology*, 25:459-462, 1997).

However, DNA immunization is limited in use
20 because expression frequency of an antigen is low *in vivo*. Some kind of costimulatory molecule genes which are necessary for the activation of immune cells were used to increase the effect of DNA immunization (Geissler, M. et al., *J. Immunol.*,
25 159:5107-5113, 1997; Iwasaki, A. et al., *J. Immunol.*, 158:4591-4601, 1997; Lee, S. W. et al., *J. Virol.*, 72:8430-8436, 1997). IL-12 gene was also used to induce

WO 01/68802

PCT/KR01/00408

immune response against HCV effectively(Lasartte, J. J.
et al., *J. Immunol.*, 162:270-277, 1999). Those, however,
has given unsatisfactory results so far especially in
the DNA immunization with human and primates(Boyer, J.
5 *et al.*, *Keystone Symposium on DNA Vaccines April*, 12-17,
1998).

The present inventors worked hard to produce genes
which can express active IL-12p70 through the control
of glycosylation and can minimize the secretion of IL-
10 12p40 decreasing immune activity by action of IL-12. As
a result, the mutant gene was obtained through the
mutation of Asn-222, glycosylation site of human IL-
12p40 subunit and Asn-220, glycosylation site of mouse
IL-12p40 subunit. The obtained mutant genes which
15 increase the expression of active IL-12p70 and decrease
the secretion of IL-12p40 was used in small animal
model, mice, along with HCV E2 gene for DNA
immunization. Through this trial, the best cell-
mediated immune response was induced and even this
20 immune response was continued long period. Thus, it is
certain that the mutant gene of the present invention
is very useful as an adjuvant for DNA vaccine.

SUMMARY OF THE INVENTION

25 It is an object of this invention to provide
mutant IL-12p40 gene which activates the basic role of

WO 01/68802

PCT/KR01/00408

IL-12 such as activation of CTL cells or enhancement of
immune response through Th1 cells by mutating the
glycosylation site which is essential in secretion of
IL-12p40 in human or mouse, to decrease the secretion
5 of IL-12p40 which inhibits the activation of IL-12p70
by binding the receptor of IL-12p70 competitively.

It is a further object of this invention to
provide adjuvant which keeps and increases antigen-
specific immune response by DNA immunization using
10 expression vector containing the mutant IL-12p40 gene
and DNA vaccine together.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the sequence homology of amino acids
15 between human and mouse IL-12p40.

FIG. 2 shows p40 and p35 subunits of wild type
human IL-12, and a amino acid changes at putative N-
glycosylation sites of p40 and p35 subunits on the
20 present invention.

FIG. 3a shows a result of western blot analysis of
human IL-12p35 and its derivatives with mutations at
putative N-glycosylation sites in cell lysate of the
25 present invention.

WO 01/68802

PCT/KR01/00408

FIG. 3b shows a result of western blot analysis of human IL-12p40 and its derivatives with mutations at putative N-glycosylation sites in cell lysate of the present invention.

5

FIG. 3c shows a result of western blot analysis of human IL-12p40 and its derivatives with mutations at putative N-glycosylation sites in cell supernatant of the present invention.

10

FIG. 4 shows a expression vector of the present invention which is inserted with envelope glycoprotein 2(E2) gene of hepatitis C virus(HCV) and wild-type or mutant mouse IL-12 gene.

15

FIG. 5a shows a result measuring titer of total IgG antibody to HCV E2 in mouse serum which is immunized with DNA vector of the present invention.

GI; Mouse which is immunized with 200 μ g of pTV2.

20

GII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

GIII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

25 GIV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG. 5b shows a result measuring titer of IgG1

WO 01/68802

PCT/KR01/00408

antibody to HCV E2 in mouse serum which is immunized with DNA vector of the present invention.

G I; Mouse which is immunized with 200 μ g of pTV2.

G II; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

G III; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

G IV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

10

FIG. 5c shows a result measuring titer of IgG2a antibody to HCV E2 in mouse serum which is immunized with DNA vector of the present invention.

G I; Mouse which is immunized with 200 μ g of pTV2.

G II; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

G III; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

G IV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

20

FIG. 5d shows a result measuring ratio of titer of IgG2a and IgG1 antibody(IgG2a/IgG1) to HCV E2 with mouse serum which is immunized with DNA vector of the present invention.

25

G I; Mouse which is immunized with 200 μ g of pTV2.

G II; Mouse which is immunized with 100 μ g of pTV2-

WO 01/68802

PCT/KR01/00408

HCV-gDsE2t and 100 μ g of pTV2.

GIII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

5 GIV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG. 6a shows a result measuring IFN- γ production level from mouse splenocytes obtained 3 weeks after immunized with DNA vector of the present invention.

10 GI; Mouse which is immunized with 200 μ g of pTV2.

GII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

GIII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

15 GIV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG. 6b shows a result measuring IFN- γ production level from mouse splenocytes obtained 6 weeks after immunized with DNA vector of the present invention.

20 GI; Mouse which is immunized with 200 μ g of pTV2.

GII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

25 GIII; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

GIV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

WO 01/68802

PCT/KR01/00408

FIG. 6c shows a result measuring IFN- γ production level from mouse splenocytes obtained 10 weeks after immunized with DNA vector of the present invention.

5 G I ; Mouse which is immunized with 200 μ g of pTV2.

 G II ; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

 G III ; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

10 G IV ; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG. 7 shows a result measuring specific CTL activity to HCV E2 using engineered CT26 tumor cells expressing hghE2t after immunization with DNA vector of the present invention.

15 G I ; Mouse which is immunized with 200 μ g of pTV2.

 G II ; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

20 G III ; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

 G IV ; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

25 a ; right after immunization, b ; 3 weeks after immunization,

 c ; 6 weeks after immunization, d ; 10 weeks after immunization,

WO 01/68802

PCT/KR01/00408

e; 14 weeks after immunization, f; 14 weeks after immunization, control(CT26-neo cells)

FIG. 8a shows a result measuring levels of total
5 IgG, IgG1 and IgG2a to HCV E2 in mouse serum 2 weeks after challenge with engineered CT26 tumor cells expressing hghE2t into mice immunized with DNA vector of the present invention.

G I; Mouse which is immunized with 200 μ g of pTV2.

10 G II; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

G III; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

15 G IV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG. 8b shows a result measuring ratio of titers of IgG2a and IgG1(IgG2a/IgG1) to HCV E2 in mouse serum
2 weeks after challenge with engineered CT26 tumor
20 cells expressing hghE2t into mice immunized with DNA vector of the present invention.

G I; Mouse which is immunized with 200 μ g of pTV2.

G II; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

25 G III; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

G IV; Mouse which is immunized with 100 μ g of pTV2-

WO 01/68802

PCT/KR01/00408

HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG.8c shows a result measuring change of the IFN- γ production level of tumor cells expressing hghE2t into mice immunized with DNA vector of the present invention.

G I; Mouse which is immunized with 200 μ g of pTV2.

G II; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

10 G III; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

G IV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

15 FIG. 8d shows a result measuring survival rate of mice challenged with engineered CT26 tumor cells expressing hghE2t after immunization with DNA vector of the present invention.

G I; Mouse which is immunized with 200 μ g of pTV2.

20 G II; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2.

G III; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12wt.

25 G IV; Mouse which is immunized with 100 μ g of pTV2-HCV-gDsE2t and 100 μ g of pTV2-mIL-12mut.

FIG. 9 shows a schematic view of biological

WO 01/68802

PCT/KR01/00408

function of IL-12 *in vivo*.

▲; p35,

○; p40,

○○; (p40)₂,

▲○; IL-12(p70)

5

DETAILED DESCRIPTION OF THE INVENTION

To achieve those objects, the present invention provides human or mouse mutant IL-12p40 subunit genes replaced Asn-222(human) or Asn-220(mouse) which is essential for the secretion of IL-12p40 with other amino acids.

The present invention provides gene construct and expression vector containing IRES sequence for co-expression of human or mouse genes of mutated IL-12p40 subunit and subunit itself.

And also, the present invention provides DNA vaccine vector containing and expressing HCV E2 gene, gene construct and expression vector containing p40 subunit gene which has mutated Asn-222(human) or Asn-220(mouse) sites.

Finally, the present invention provides a method for gene therapy or DNA vaccine immunization using this gene construct as an adjuvant for the enhancement of immune response.

Further features of the present invention will appear hereinafter.

Once again, the present invention is to provide

WO 01/68802

PCT/KR01/00408

human or mouse mutant IL-12p40 genes in which Asn-222(human) or Asn-220(mouse) playing a very important role in secretion of human IL-12p40 having SEQ. NO. 1 or mouse IL-12p40 having SEQ. NO. 2 are replaced with
5 other amino acids.

Comparing with the amino acid sequence of human and mouse IL-12p40, Asn-220 of mouse IL-12p40 is located in the very similar context to Asn-222 of human IL-12p40 (FIG. 1).

10 More specifically, codon AAC designating Asn-222 of human IL-12p40 subunit can be changed with CUC, CAG or AUA etc, and amino acids coming under each of those are Leu, Gln and Ile. The present invention gives test examples demonstrating that AAC was changed into CTC or
15 CAG on cDNA, and thus codon AAC was exchanged with CUC or CAG. Finally, the present invention provides mutant gene in which Asn-222 amino acid was replaced with Leu-222(hp40-N222L) or Gln-222(hp40-N222Q).

Codon ACC designating Asn-220 of mouse IL-12p40
20 subunit which is compared with Asn-222 of human IL-12p40 subunit can be changed with CUC, CAG or AUA, and amino acids coming under each of those are Leu, Gln and Ile. The present invention provides mutant gene in which Asn-220 amino acid was replaced with Leu-
25 220(mp40-N220L) by changing AAC with CTC on cDNA.

The mutant genes, hp40-N222L, hp40-N222Q and mp40-N220L of the present invention are coding amino acid

WO 01/68802

PCT/KR01/00408

sequences noted as SEQ. NO. 3, SEQ. NO. 4 and SEQ. NO. 5 respectively.

The present invention provides gene construct containing IRES for simultaneous expression of subunit
5 itself and human or mouse mutant IL-12p40 subunit gene, and expression vector including this gene construct.

First of all, the present invention provides hp40-N222L/IRES/hp35, hp35/IRES/hp40-N222L and mp35/IRES/mp40-N220L genes containing genes explained
10 above. hp40-N222L/IRES/hp35 gene and hp35/IRES/hp40-N222L gene include gene encoding p35 subunit along with human p40 subunit having Leu-222 instead of Asn-222, while mp35/IRES/mp40-N220L gene includes gene encoding p35 subunit along with mouse p40 subunit whose Asn-220
15 sites were replaced with Leu-220. Both genes have IRES sequence for simultaneous expression of subunits. IRES from EMCV(*encephalomyocarditis*) is preferable for genes such as hp40-N222L/IRES/hp35, hp35/IRES/hp40-N222L and mp35/IRES/mp40-N220L, but not limited to thereof. IRES
20 sequence is important in co-expression of genes coding p40 subunit and p35 subunit.

The present invention provides expression vectors, for example, pGX0-hp40-N222L/IRES/hp35 containing hp40-N222L/IRES/hp35, pGX0-hp35/IRES/hp40-N222L containing
25 hp35/IRES/hp40-N222L and pTV2-mp35/IRES/mp40-N220L containing mp35/IRES/mp40-N220L as well.

The present invention shows an example in which

pGX0 plasmid, a DNA vaccine vector becomes possible to be used in medical experiment by inserting kanamycin resistant gene instead of ampicillin resistant gene of pTV2 DNA vaccine vector(Song, M. K. et al., *J. Virol.*, 74:2920-2925, 2000), and hp40-N222L/IRES/hp35 gene and hp35/IRES/hp40-N222L gene were inserted in the part of that pGX0 plasmid which is ready for receiving foreign genes. And mp35/IRES/mp40-N220 gene was inserted into the foreign gene receiving part of pTV2 DNA vaccine vector. Besides the above plasmids used for making expression vectors, there are various vectors including for prokaryotes or eukaryotes, which makes possible to use a different vector for a different purpose. It is also possible to change the size and nucleotide sequence of gene which is to be inserted into the foreign gene receiving part of expression vector.

The expression vectors, pGX0-hp35/IRES/hp40-N222L and pGX0-hp40-N222L/IRES/hp35 of the present invention were deposited at Gene Bank of Korea Research Institute of Bioscience and Biotechnology on February, 26, 2001(Accession No: KCTC 0969BP and KCTC 0970BP). And the pTV2-mp35/IRES/mp40-N220L was deposited on February, 29, 2000(Accession NO: KCTC 0745BP).

Since the co-expression of p35 and p40 is essential to obtain a biologically active IL-12p70, bicistronic expression vectors, pGX0-hp40/IRES/hp35 and

WO 01/68802

PCT/KR01/00408

pGX0-hp35/IRES/hp40, were generated by using IRES of EMCV besides pCIN-hp35 or pCIN-hp40 expressing hp35 or hp40 respectively. To understand the influence of N-glycosylation on the synthesis, secretion and specific
5 activity of human IL-12, Asn amino acid residue in putative N-glycosylation site existing on p35 and p40 subunit was replaced with other amino acid residues by site-directed mutagenesis(FIG. 2). Through the immunoblot of wild-type protein on mutant protein(FIG.
10 3a, 3b and 3c), Asn-127, -141 of hp35 subunit and Asn-222, -303 of hp40 subunit were sure to be used for N-glycosylation.

To detect the effect of N-glycosylation of hp35 or hp40 on synthesis of heterodimerization and secretion
15 of IL-12p70, cells were transfected with hIL-12 expression vector containing each wild-type gene or mutant gene, and then analyzed by ELISA using culture supernatant and cell lysate obtained therefrom.

As a result, it has been clarified that Asn-127 of
20 hp35 causes decrease of heterodimerization and secretion of IL-12p70. However, compared with wild-type hp35, Asn-141 of hp35 seems relatively not to affect much to the heterodimerization and secretion of IL-12p70 and IL-12p40(TABLE 1).

25 In addition, the mutation of Asn-135 or Asn-222 of hp40 subunit significantly decreases the secretion of IL-12p40, while does not affect the secretion of IL-

WO 01/68802

PCT/KR01/00408

12p70. Especially, the secretion of hp40 dropped to the degree of 8% relatively to the wild-type hp40 by Asn-222. Although Asn-135 of hp40 is not a N-glycosylation site, it still causes decrease of the secretion of hp40, which means Asn of Asn-135 itself plays an important role in the secretion of hp40. Even though amino acid of Asn-222 was replaced with Gln or Leu, the secretion of hp40 was still decreased, suggesting that was because of the loss of N-glycosylation at Asn-222 (TABLE 1). Thus, N-glycosylation at Asn-222 is required not for the secretion of heterodimer hIL-12, hIL-12p70 but for the secretion of hIL-12p40 only.

Considering all those results together, the present inventors verified that N-glycosylation at Asn-222 is necessary for the secretion of hIL-12p40, but not required for the secretion and heterodimerization of hIL-12p70, in the meantime N-glycosylation at Asn-127 of hp35 plays an important role in the secretion and heterodimerization of hIL-12p70.

20

To investigate the effect on biological activity of N-glycosylation of hIL-12, IFN- γ induction ability of wild-type hIL-12 and its derivatives containing putative N-glycosylation site mutation were analyzed by ELISA.

25

As a result, in the matter of IFN- γ induction ability, IFN- γ induction ability in culture

WO 01/68802

PCT/KR01/00408

supernatant obtained by co-transfection with wild-type hp35 and hp40 mutants in which Asn-135 and/or Asn-222 were mutated was increased in comparison with that of wild-type or other hp40 mutants (TABLE 1). And, when
5 hp40 was supplied in the culture supernatant, the increased IFN- γ induction ability had come down to the similar level with that of wild-type (TABLE 1). Thus, it is suggested that these results are attributed to the relatively low level of hIL-12p40, known as an
10 antagonist against hIL-12p70, in culture supernatant of mutants containing hp40-N135Q and/or hp40-N222Q, and the activity of hp70 produced from mutation of Asn-135 and/or Asn-222 does not increased by the mutation itself.

15

Meanwhile, to examine the effect of glycosylation of hp40 at Asn-222 on the decrease of secretion of hIL-12p40, cells were co-transfected with expression vector containing certain amount of mutant gene and various
20 amount of wild-type hp35 DNA, then cultured. Finally, the induced amount of IFN- γ was measured by ELISA.

Generally, p35 subunit is not secreted solely but secreted in the form of IL-12p70 binding to p40, while p40 subunit is secreted in the form of monomer or
25 homodimer, which suggests that not p35 but p40 subunit is a major factor inducing secretion of IL-12p70. Corresponding to this, the amount of hIL-12p70

WO 01/68802

PCT/KR01/00408

secretion was increased in proportion to the amount of transfected hp35 DNA in both wild-type hp40 and hp40-N222Q mutant (TABLE 1). That means once hp40 having secretion defect is bound with hp35 subunit, it can be secreted in the form of IL-12p70, and thus, hp35 subunit seems to perform its another function in secretion of hIL-12p70. According to a recent report, the conformational change of hp40 is due to the binding with hp35 (Yoon, C. et al., *EMBO J.*, 19:3530-3534, 2000), which also suggests that hp35 subunit can contribute to the secretion of IL-12p70. Conclusively, hp40 including glycosylation at Asn-222 itself has a defect in secretion but once it is bound with hp35, hp40 may be conformationally changed in its form and its conformational change could expose or generate the covered or new secretion signal, respectively, and then induce the secretion of heterodimer.

To decrease the secretion of p40 by inducing co-expression of p35 and p40 in a cell, and to decrease the amount of p40 secretion by inducing formation of p40 in which p40 gene is located behind IRES because the degree of gene expression using IRES is rather lower than using cytomegalovirus (CMV) promoter, hp40/IRES/hp35 and hp35/IRES/hp40 vectors were constructed. And also, hp40-N222L/IRES/hp35 and hp40/IRES/hp40-N222L in which hp40 gene had been

WO 01/68802

PCT/KR01/00408

replaced with hp40-N222L gene in each plasmid were generated (TABLE 1). Compared hp40/IRES/hp35 with hp40-N222L/IRES/hp35, hp40 was less secreted to the degree of 5% in the later. On the other hand, when hp35 and hp40-N222L were electroporated, hp40 was secreted to the degree of 8%. When hp35 and hp40-N222L were electroporated, it is possible for the two plasmids not to be transfected both of them into a cell. Besides, as hp40-N222L gene is expressed alone in a cell, the small amount of hp40 is secreted without the secretion of hp70. Therefore hp40 is supposed to be secreted more in hp40-N222L plus hp35 than in hp40-N222L/IRES/hp35 in which hp35 and hp40-N222L genes are expressed at the same time. Comparing hp40/IRES/hp35 with hp35/IRES/hp40, the secretion and expression of hp70 were not much different, but the secretion and expression of hp40 were significantly decreased in hp35/IRES/hp40 since the expression of hp40 was depressed by locating hp40 gene behind IRES. In the case of hp35/IRES/hp40-N222L made by inserting hp40-N222L gene instead of hp40 gene, the secretion level of hp40 dropped to 0.3%. Thus, it is confirmed through the present experiment that hp35/IRES/hp40-N222L is the one that can keep the secretion level of hp70 and at the same time minimize the secretion of hp40.

In addition, the present invention provides DNA

WO 01/68802

PCT/KR01/00408

vaccine vector, containing and expressing E2 gene of HCV-1b, and gene construct having p40 subunit gene which is mutated at Asn-222(human) or Asn-220(mouse).

5 To inspect the possibility for the use of hIL-12 mutant gene of the present invention in gene therapy as a DNA vaccine, the sequence of mouse IL-12p40(mp40) gene homologous to Asn-222 of hp40 gene was searched. Asn-220 of mp40 was located in the very similar context to that of Asn-222 of hp40, but this amino acid was not
10 known to be N-glycosylated until now(FIG. 1). Therefore, the present inventors generated pCIN-mp40-N220L vector containing the mutant mp40 gene, mp40-N220L, in which amino acid of Asn-220 was replaced with Leu by site-directed mutagenesis.

15 To generate a vector encoding mouse p35 and p40 subunits and being used for the DNA immunization, pTV2-mp35/IRES/mp40 vector was constructed by inserting mp35/IRES/mp40 fragment into pTV2 vector, an eukaryotic expression vector used as a DNA vaccine vector in small
20 animals(Lee et al., *J. Virol.*, 72:8430-8436, 1998; Cho et al., *Vaccine*, 17:1136-1144, 1999). And, the pTV2-mp35/IRES/mp40-N220L vector, containing Asn-220 mutant gene of mouse IL-12p40 and expressing p35, was constructed based on the observation that
25 hp35/IRES/hp40-N222L gene can sustain the secretion of hp70 while can minimize the secretion of hp40.

The pTV2-mp35/IRES/mp40-N220L vector of the

WO 01/68802

PCT/KR01/00408

present invention was deposited at Gene Bank of Korea Research Institute of Bioscience and Biotechnology on February, 29, 2000 (Accession No: KCTC 0745BP).

5 pTV2-HCV-E2 DNA vaccine vector which can express HCV-E2 protein in eukaryotic cells was constructed (Song M. K., et al., *J. Virol.*, 74:2920-2925, 2000). As seen in FIG. 4, pTV2-HCV-E2 DNA vaccine vector is consist of simian virus 40 replication origin (SV40 ori),
10 cytomegalovirus (CMV) promoter, tripartite leader sequence (TPL) of adenovirus, multiple cloning sequence (MCS), SV40 polyadenylation sequence (poly A) and ampicillin resistance gene (AmpR). And also, HCV-E2 gene is cloned in MCS of this vector. Carboxyl-
15 terminal (C-terminal) containing hydrophobic amino acid residue of E2 gene used in the present invention was removed to facilitate protein secretion. And to help protein expression and cell secretion, aminocyl-
terminal (N-terminal) and signal sequence (S) of
20 herpesvirus (HSV) glycoprotein D (gD) were linked.

 To analyze the effect of Asn-220 mutation of mouse IL-12p40 on the secretion of IL-12p40 or IL-12p70, the secretion level of mouse IL-12 was analyzed by ELISA with both culture supernatants and lysates of cells
25 transfected with above vectors. As a result, mp40-N220L mutants showed similar characteristics to Asn-222 mutants of hp40 in the point of the secretion of IL-

WO 01/68802

PCT/KR01/00408

12p40 or IL-12p70 and its biological activity(TABLE 1).

Finally, the present invention provides a method for gene therapy or DNA vaccine immunization using this gene construct containing mutant p40 subunit gene whose
5 Asn-222(human) or Asn-220(mouse) is mutated as an immune enhancer.

And this method can be useful for prevention and treatment against various diseases, for example, AIDS, hepatitis C or hepatitis B, cancer, influenza,
10 tuberculosis and malaria, which essentially require cellular immune responses for their therapy.

In a previous report, DNA vaccination of plasmid encoding HCV E2 antigen(pTV2-gDsE2t) was sufficient to induce the antigen-specific and cell-mediated immune
15 responses after 3 weeks post-immunization. To determine whether the mutant mIL-12 gene can affect the efficient antigen-specific immune response *in vivo* compared with wild-type mIL-12 gene, and to inspect whether the effect can be maintained for a long time, mice were
20 immunized and boosted with the DNA vaccine vector of the present invention and production of HCV E2 specific antigen was analyzed by ELISA.

As a result, HCV E2 DNA vaccine induced systemic HCV E2-specific total IgG, IgG1 and IgG2a levels
25 significantly higher than negative control values, and coinjection with mIL-12mut gene or mIL-12wt gene showed the similar total IgG level compared to HCV E2 DNA

WO 01/68802

PCT/KR01/00408

vaccine only. However, the level of IgG1 was similar among the groups injected with HCV E2 DNA. In contrast, the level of IgG2a against HCV E2 slightly increased in mIL-12wt group and significantly increased in mIL-12mut group compared with HCV E2 DNA immunized only group. In addition, the ratio of IgG2a/IgG1, which is generally accepted as an indirect indicator of Th1 immunity, was the highest in the mIL-12mut group, which showed the similar pattern of IgG2 level (FIG. 5a, 5b, 5c and 5d). These data represent that mIL-12mut gene significantly affected the shift of IgG subclasses from IgG1 to IgG2a in humoral immune response compared to mIL-12wt or HCV E2 only group, suggesting that IL-12mut may induce Th1 type immune response. And these effects maintained for 0, 3, 6, 10 weeks after booster immunization.

To investigate the effect of mIL-12mut gene on Th1 immune response, which is one of the parameter used to evaluate the potency of cell-mediated immune response, IFN- γ expression of splenocytes was analyzed.

As a result, HCV E2 DNA immunized group without cytokine gene showed the increased level of IFN- γ in proportion to the concentration of hghE2t protein, whereas mock plasmid immunized group did not. Expectedly, the level of IFN- γ induction in mIL-12wt group was more enhanced than in HCV E2 only group and mIL-12mut group showed up to 2-3 times higher IFN- γ production than mIL-12wt group (FIG. 6a, 6b and 6c),

WO 01/68802

PCT/KR01/00408

suggesting that mIL-12p70 increases the antigen-specific Th1 immune response and mIL-12p40 inhibits the induction of Th1 immune response by IL-12p70 *in vivo*.

5 As noted above, mIL-12mut gene contributed to the long-term Th1 immune response in HCV E2 DNA immunization. To determine whether the long-term Th1 immune response induction by the expression of mIL-12mut gene correlates with CTL immunity, the major
10 cell-mediated immune response, and if so, to find out whether mIL-12mut gene can affect the maintenance of CTL activity in DNA immunization model, the present inventors performed the CTL assay with splenocytes of DNA immunized mice at various weeks after booster
15 immunization.

As a result, two weeks after boost, all groups, except for mock plasmid immunized group, showed very strong antigen-specific CTL activity. However, there was little significant difference among HCV E2 only,
20 mIL-12wt and mIL-12mut groups. In mIL-12wt coimmunized group, CTL response was more increased than in HCV E2 only group in overall period, indicating that mIL-12 gene played a role in the enhanced CTL generation. Interestingly, the difference in CTL activity between
25 mIL-12mut group and the other two groups, HCV E2 only and mIL-12wt group, was more and more extensive as the time after booster immunization is longer. Especially,

WO 01/68802

PCT/KR01/00408

at 10 weeks, CTL response was very low in HCV E2 only and mIL-12wt groups, suggesting that the frequency of antigen-specific CTL significantly decreased after long period, whereas mIL-12mut group showed the CTL activity
5 5 to 10 times higher than the other two groups, sustaining the antigen-specific CTL response(FIG. 7). As a control, when CT26-neo cell was used as the target cell, no lysis was observed in all groups, suggesting that CTL activity observed in this experiment is HCV
10 E2-specific.

And, the present inventors measured the frequency of HCV E2 specific CD8+ cells *in vivo* to confirm the effect of induction and maintenance of antigen-specific
15 CD8+ T cell stimulated by mIL-12. Two types of immuno assay methods were used. First one is counting the number of antigen-specific IFN- γ producing cells. To investigate whether the enhancement of CTL activity is originated from the secretion of antigen-specific CD8+
20 T cell, CD8+ cells were isolated, and stained with PE conjugated anti-mouse IFN- γ antibody or control PE-conjugated isotype-matched antibody. Stained cells were analyzed by FACSCalibur flow cytometry(Becton Dickinson), and then induction of IFN- γ was observed.
25 As a result, mice coimmunized with mIL-12mut gene had a 3 to 7-fold enhancement in the frequency of CD8+ IFN- γ producing cells compared with HCV E2 only and mIL-12wt

WO 01/68802

PCT/KR01/00408

groups at 0, 3, 6, 10 or 14 weeks after booster immunization, which showed correlation with the result in CTL response. By contrast, in isotype-matched control experiment, there was no difference among all groups. Similarly to the result of CTL assay, the difference of IFN- γ producing CD8+ T cell frequency among immunization groups was not extensive at 2-3 weeks after booster immunization. These data demonstrate that the expression of mIL-12mut gene sustained the frequency of CD8+ IFN- γ producing T cells after DNA immunization for a long time.

To investigate the frequency of antigen-specific CD8+ T cells, the frequency of HCV E2 specific CD8+ T cells was measured by the other assay method, limiting dilution assay(LDA). Splenocytes of booster immunized mice were diluted with various concentration, and cultured with CT-26 cells expressing E2. And then, CTL activity of antigen-specific stimulated CD8+ T cell was measured. The result of limiting dilution assay was similar with that of intracellular staining assay. Namely, the highest frequency of antigen-specific CD8+ T cell was observed in mIL-12mut group even in early stage of immunization, and this frequency had been remained over 14 weeks after booster immunization(TABLE 2).

The frequency of HCV E2 specific CD8+ T cells without booster immunization was also observed

WO 01/68802

PCT/KR01/00408

according to the time. Overall frequency of antigen-specific CD8+ T cell was slightly decreased, but the highest frequency of antigen-specific CD8+ T cell was observed in mIL-12mut group(TABLE 2).

5 In this regard, it might be suggested that the role of IL-12p70 itself in cell-mediated immune response *in vivo* is to induce the activation of Th1 and CTL from the very beginning and to maintain it for a long time, whereas IL-12p40 inhibits the IL-12p70 as an
10 antagonist *in vivo*.

To confirm the Th1 and CTL immunity induced by mIL-12mut gene *in vivo*, and to examine that the correlation of protective immunity with Th1 and CTL
15 immune response, CT26-hghE2t tumor cells expressing hghE2t were injected in the groups of the immunized mice at 12 weeks after the booster immunization. 2 weeks after injection, relative levels of antigen-specific IgG, IgG1, IgG2 and the ratio of IgG2a/IgG1
20 were determined(FIG. 8a and FIG. 8b). As a result, highest level of IgG2a/IgG1 ratio was observed in mIL-12mut group, suggesting that Th1 immune response can be induced by mIL-12mut even in tumor injection.

Tumor size has been measured for 30 days.
25 Particularly, the mean local tumor growth was determined by measuring the volume and diameter of tumors with calipers every three days. Also, the

WO 01/68802

PCT/KR01/00408

survival rates of these mice were determined by observing for about 70 days. The group of mice immunized with mIL-12mut induced strong Th1 immune response. The group of mIL-12wt immunization displayed the delayed tumor growth in contrast to the group of only pTV2-gDsE2t immunization(FIG. 8c), whereas the group of mIL-12mut immunization showed the significantly delayed tumor growth. In control group, most of mice had tumor and died within 50 days, but 90% of mice in the group of mIL-12mut could survive after 70 days(FIG. 8d). Thus, these data suggest that HCV E2-specific Th1 and CTL responses induced by mIL-12mut gene of the present invention confers *in vivo* protection against the challenge of modified tumor cells expressing specific antigen.

The present inventors verified that human or mouse mutant IL-12p40 subunit containing mutant Asn-222(human) or Asn-220(mouse) which is essential for the secretion of IL-12p40 is very useful for DNA vaccine immunization and gene therapy as an adjuvant. Thus, the present invention provides an adjuvant containing human or mouse mutant IL-12p40 gene as an effective component for DNA immunization and gene therapy which can induce antigen-specific immune response for prevention and treatment against various diseases.

This adjuvant can induce immune response, if used

WO 01/68802

PCT/KR01/00408

with DNA vaccine, by increasing the secretion of IFN- γ from CD8+ or T helper cells and the hydrolysis activity of cytotoxic T lymphocytes(CTL).

5

EXAMPLES

Practical and presently preferred embodiments of the present invention are illustrative as shown in the following Examples.

10

However, it will be appreciated that those skilled in the art, on consideration of this disclosure, may make modifications and improvements within the spirit and scope of the present invention.

Example 1: Construction of human IL-12 expression

15

vectors

<1-1> Construction of human IL-12 expression vectors

20

The cDNAs of human p35(820bp) and p40(1050bp) subunits were cloned and amplified using reverse transcriptase-polymerase chain reaction(RT-PCR, PCR System 2400, Perkin Elmer). The each amplified cDNA was subcloned into Sma I region of the universal vector, pSK(Stratagene, La Jolla, California) and then pSK-hp35 and pSK-hp40 were constructed.

25

To generate a bicistronic vector encoding hp35 and hp40 genes, pSK-internal ribosomal entry site(IRES)

WO 01/68802

PCT/KR01/00408

vector was constructed. IRES gene of encephalomyocarditis virus (EMCV) obtained by RT-PCR was subcloned between Sma I and Pst I sites of pSK plasmid. pSK-IRES vector was cut by EcoRV, and added end of p40 DNA fragment obtained by treating pSK-hp40 with Xba I and BamH I was added to thereof using T4 DNA polymerase, then pSK-hp40/IRES was produced. And then pSK-hp40/IRES/hp35 plasmid in which p40, IRES, p35 genes are arranged in order, was constructed by the insertion of p35 DNA fragment from pSK-hp35 treated with Nco I and Sac I, to pSK-hp40/IRES.

pSK-hp35/IRES/hp40 was also constructed. pSK-IRES vector was cut by EcoRV, and hp35 DNA fragment obtained from pSK-hp40/IRES/hp35 treated with Nco I and Not I was added to thereof. pSK-hp35/IRES/hp40 plasmid was completed by adding hp40 fragment obtained by cutting pSK-hp40 with Nco I and BamH I. hp40/IRES/hp35 and hp35/IRES/hp40 genes were cloned into Spe I/Not I and Xho I/Not I sites of pGX0 vector to create pGX0-hp40/IRES/hp35 and pGX0-hp35/IRES/hp40 expression vectors which can express active IL-12p70 in mammalian cells. pGX0 vector was constructed by inserting anticanamycin gene into pTV2 vector (Song, M. K. et al., *J. Virol.*, 74:2920-2925, 2000).

25

<1-2> Construction of p40 subunit expression vector

WO 01/68802

PCT/KR01/00408

To create p40 subunit expression vector, pCIN-hp40/IRES/hp35 vector was cut by SacII and Not I (to eliminate gene encoding p35 subunit), and self-ligated with T4 DNA polymerase, and then named as pCIN-hp40.

5

<1-3> Construction of p35 subunit expression vector

To create p35 subunit expression vector, p35 DNA fragment was obtained from pCIN-hp40/IRES/hp35 vector by cutting and ligation with Nco I and T4 DNA polymerase. This p35 DNA fragment was inserted into XhoI and Not I restriction enzyme sites of pCI-neo vector, and then named as pCIN-hp35.

10

Example 2: Construction of mouse IL-12 expression vectors

15

<2-1> Construction of human IL-12 expression vectors

20

To generate a bicistronic vector encoding mouse p35 and p40 genes, pSK-IRES/mp40 vector was constructed by inserting p40 DNA fragment obtained from mouse IL-12p40 PCR product (Schoenhaent, D. S. et al., *J. Immunol.*, 148:3433-3440, 1999) treated with NcoI and BamH I into cleaved pSK-IRES vector containing IRES of EMCV. And pSK-mp35/IRES/mp40 plasmid was constructed by inserting mouse p35 DNA fragment into pSK-IRES/mp40 using BamH I and T4 DNA polymerase. Finally, pCIN-mp35/IRES/mp40 expression vector which can express

25

WO 01/68802

PCT/KR01/00408

active IL-12p70 in mammalian cells was generated by inserting mp35/IRES/mp40 gene into XhoI and NotI sites of pCI-neo vector (Promega).

5 <2-2> Construction of p40 subunit expression vector

To create wild type mouse p40 subunit expression vector, p40 DNA fragment was obtained from pSK-mp35/IRES/mp40 vector treated with NcoI and SacI. pGEX-KG-mp40 was created by inserting this p40 DNA fragment
10 into pGEX-KG vector (Clontech) treated with same restriction enzymes. pGEX-KG-mp40 was treated with EcoRI and NotI, and inserted into EcoRI and NotI sites of pCI-neo vector. So, pCIN-mp40 expression vector was generated.

15

<2-3> Construction of p35 subunit expression vector

To create wild type mouse p35 subunit expression vector, p35 DNA fragment was obtained from pSK-mp35/IRES/mp40 vector treated with XhoI and EcoRI.
20 pCIN-mp35 expression vector was constructed by inserting this DNA fragment into XhoI and EcoRI sites of pCI-neo vector treated with same restriction enzymes.

25 Example 3: Construction of IL-12p40 and IL-12p70 which have partially mutated glycosylation

Seven Asn codons which are expected to be used as

WO 01/68802

PCT/KR01/00408

N-glycosylation sites of hp35 and hp40 subunits were replaced with unrelated codons by site-directed mutagenesis.

For the construction of glutamine mutant genes for putative N-glycosylation sites of hp40 and hp35, amino acid substitutions were performed by using PCR according to the method of Haraguchi *et al.* (Haraguchi, *et al.*, *J. Immunol.*, 163:2092-2098, 1999). The primers for mutagenesis were used by synthesis of nucleotides, such as T7 represented by the SEQ. ID NO.6, T3 represented by the SEQ. ID NO.7, hp40-N125Q(S) represented by the SEQ. ID NO.8, hp40-N125Q(AS) represented by the SEQ. ID NO.9, hp40-N135Q(S) represented by the SEQ. ID NO.10, hp40-N135Q(AS) represented by the SEQ. ID NO.11, hp40-N222Q(S) represented by the SEQ. ID NO.12, hp40-N222Q(AS) represented by the SEQ. ID NO.13, hp40-N303Q(S) represented by the SEQ. ID NO.14, hp40-N303Q(AS) represented by the SEQ. ID NO.15, hp40-N127Q(S) represented by the SEQ. ID NO.16, hp40-N127Q(AS) represented by the SEQ. ID NO.17, hp40-N141Q(S) represented by the SEQ. ID NO.18, hp40-N141Q(AS) represented by the SEQ. ID NO.19, hp40-N251Q(S) represented by the SEQ. ID NO.20 and hp40-N251Q(AS) represented by the SEQ. ID NO.21. (S) and (AS) mean sense and antisense primers, respectively.

To construct single glutamine mutant genes of hp40

WO 01/68802

PCT/KR01/00408

and hp35, T7 primer and each sense primer were used for PCR using pCIN-mp40 produced from example <2-2> or pCINmp35 produced from example <2-3> as a template. Similarly, T3 primer and each antisense primer were used for PCR. Two PCR fragments sharing a common site containing the mutational point were generated. The second PCR was performed with a mixture of these products as templates and the flanking primers, resulting in the generation of a fusion product. This product was inserted into pCI-neo plasmid. Similarly, double and triple glutamine mutant genes were constructed by using single or double glutamine mutant genes as PCR templates. Mutant genes were verified by DNA sequencing.

For the construction of mouse IL-12p70 gene containing N-glycosylation defect at Asn-220 of mp40, the region of mp40 gene in pCIN-mp35/IRES/mp40 plasmid produced from example <2-1> was replaced with mp40-N222L. In the mutagenesis experiment to construct pCIN-mp40-N222L, mp40-N220L(S) represented by the SEQ. ID NO. 22 and mp40-N220L(AS) represented by the SEQ. ID NO. 23 containing SacI site were used for PCR as primers. Amplified mutant genes were verified by treatment of restriction enzyme to specific recognition site produced after mutagenesis and DNA sequence analysis.

FIG. 2 shows amino acid composition of normal and mutant genes of human IL-12p40 or IL-12p35 subunit of

WO 01/68802

PCT/KR01/00408

the present invention. Putative N-glycosylation sites are indicated by Y shape with the number of amino acid, and amino acid substituted in each mutant gene is demonstrated in a square.

5

Example 4: The effect of N-glycosylation at Asn222 of human IL-12p40

COS-7(ATCC) cells were cultured in Dulbecco's modified Eagle's medium(DMEM, GIBCO-BRL) containing 10% heat inactivated fetal bovine serum. Transfection into COS-7 cells was carried out by electroporation. The suspension of approximately 5×10^6 cells in culture medium was pulsed at 250 V, 960 μ F in the presence of 20 μ g of specimen DNA in addition to 2 μ g of pNEB-SEAP(pNEB-secreted alkaline phosphatase, New England Biolabs), which expresses secreted alkaline phosphatase and serves as an internal control(electroporator and 0.4 electroporation cuvettes by Bio-Lad).

At 24 hours post-transfection, the medium was replaced with 1.5 ml of serum-free CHO-SFMII medium(GIBCO-BRL). 1.5 μ g/ml of tunicamycin was added in some experiments. After incubation for another 24 hours, supernatants and cell pellets were harvested by centrifugation. Supernatants were used for the SEAP assay and cell pellets were resuspended in 200 μ l of lysis solution(Promega). The levels of IL-12p70 and IL-

WO 01/68802

PCT/KR01/00408

12p40 in both supernatants and cell lysates were measured by ELISA(R&D system). For immunoblotting, 10% or 12% sodium dodecyl-sulfate-polyacrylamide electrophoresis(SDS-PAGE) was conducted with
5 supernatants and cell lysates. The proteins separated from the above experiment was electrotransferred to nylon membrane(Amersham). The proteins absorbed to the membrane were obtained by using biotin-labeled human IL-12 antibody(Amersham), horseradish peroxidase(HRP)-
10 labeled streptavidin(PharMingen) and ECL kit(Amersham). The results are shown in Table 1. The levels of IL-12p70 or IL-12p40 expression were measured by ELISA and presented relatively to the level of wild type, which is arbitrarily set at 100%.

15

<Table 1> The changes of expression level, secretion of IL-12p40 and IL-12p70, and IFN- γ induction capacity.

20

25

WO 01/68802

PCT/KR01/00408

Consturcts ^a	Cell Lysates (%) ^b		Supernatants (%) ^b		IFN- γ induction (%) ^c
	IL-12p40 _d	IL-12p70 _e	IL-12p40	IL-12p70	
mock ^f	< 1 ^g	< 1	< 1	< 1	5.2 $\pm$ 2.3
hp35 + hp40	100 $\pm$ 8.3	100 $\pm$ 12	100 $\pm$ 13	100 $\pm$ 7.4	100 $\pm$ 11
5					
hp40	102 $\pm$ 11	< 1	91 $\pm$ 16	< 1	8.3 $\pm$ 2.1
hp35-N127Q + hp40	99 $\pm$ 4.3	72 $\pm$ 7.7	98 $\pm$ 16	55 $\pm$ 14	98 $\pm$ 19
hp35-N141Q + hp40	102 $\pm$ 18	90 $\pm$ 12	87 $\pm$ 11	95 $\pm$ 8.1	91 $\pm$ 21
hp35-N251Q + hp40	97 $\pm$ 10	88 $\pm$ 8.4	88 $\pm$ 12	89 $\pm$ 8.9	102 $\pm$ 18
hp35-N127,141Q + hp40	88 $\pm$ 6.7	71 $\pm$ 4.6	103 $\pm$ 13	52 $\pm$ 11	90 $\pm$ 7.9
10					
hp35	< 1	< 1	< 1	< 1	3.4 $\pm$ 1.4
hp40-N125Q + hp35	89 $\pm$ 11	97 $\pm$ 17	108 $\pm$ 17	101 $\pm$ 11	108 $\pm$ 21
hp40-N135Q + hp35	92 $\pm$ 13	99 $\pm$ 4.4	29 $\pm$ 4.7	102 $\pm$ 14	121 $\pm$ 9.1
hp40-N135Q + hp35 + rhp40 ^h					97 $\pm$ 14
hp40-N222L + hp35	90 $\pm$ 16	89 $\pm$ 21	8 $\pm$ 6.2	93 $\pm$ 12	142 $\pm$ 24
hp40-N222Q + hp35 (10 : 10) ⁱ	88 $\pm$ 17	90 $\pm$ 14	9 $\pm$ 4.3	94 $\pm$ 7.5	146 $\pm$ 12
hp40-N222Q + hp35 (10 : 0)	92 $\pm$ 6.7	< 1	10 $\pm$ 5.3	< 1	5.3 $\pm$ 3.1
hp40-N222Q + hp35 (10 : 2)	87 $\pm$ 11	39 $\pm$ 7.0	11 $\pm$ 2.7	48 $\pm$ 16	125 $\pm$ 14
hp40-N222Q + hp35 (10 : 4)	84 $\pm$ 14	62 $\pm$ 11	9 $\pm$ 11	72 $\pm$ 9.2	139 $\pm$ 24
hp40-N222Q + hp35 (2 : 10)	41 $\pm$ 4.2	92 $\pm$ 11	2.4 $\pm$ 1.0	91 $\pm$ 7.4	166 $\pm$ 19
hp40-N222Q + hp35 (4 : 10)	65 $\pm$ 7.8	99 $\pm$ 17	7.2 $\pm$ 0.8	97 $\pm$ 11	151 $\pm$ 23
hp40-N222Q + hp35 + rhp40 ^h					103 $\pm$ 18
hp40-N303Q + hp35	108 $\pm$ 12	102 $\pm$ 22	104 $\pm$ 11	121 $\pm$ 19	97 $\pm$ 7.8
hp40-N135,222Q + hp35	86 $\pm$ 11	88 $\pm$ 6.9	4.0 $\pm$ 2.1	81 $\pm$ 8.4	147 $\pm$ 15
hp40-N135,303Q + hp35	89 $\pm$ 7.1	97 $\pm$ 11	23 $\pm$ 4.5	82 $\pm$ 11	103 $\pm$ 8.3
hp40-N222,303Q + hp35	93 $\pm$ 7.9	102 $\pm$ 24	19 $\pm$ 6.7	85 $\pm$ 4.8	125 $\pm$ 12
hp40-N135,222,303Q + hp35	85 $\pm$ 7.4	65 $\pm$ 4.3	2.2 $\pm$ 1.1	65 $\pm$ 7.4	148 $\pm$ 28
hp35-N127,141Q + hp40-N222,303Q	89 $\pm$ 11	65 $\pm$ 12	17 $\pm$ 8.8	51 $\pm$ 6.5	120 $\pm$ 13
15					
mock ^j	< 1	< 1	< 1	< 1	5.4 $\pm$ 3.2
hp40/IRES/hp35	100 $\pm$ 11	100 $\pm$ 21	100 $\pm$ 14	100 $\pm$ 17	100 $\pm$ 16
hp40-N222L/IRES/hp35	88 $\pm$ 13	87 $\pm$ 12	5.3 $\pm$ 3.2	95 $\pm$ 15	134 $\pm$ 17
hp35/IRES/hp40	6.2 $\pm$ 2.3	99 $\pm$ 14	4.5 $\pm$ 1.9	101 $\pm$ 13	138 $\pm$ 21
mp35/IRES/hp40-N222L	5.8 $\pm$ 3.1	97 $\pm$ 17	0.3 $\pm$ 0.2	97 $\pm$ 16	169 $\pm$ 33
mock ^k	< 1	< 1	< 1	< 1	5.2 $\pm$ 2.4
mp35/IRES/mp40 (mIL-12wt)	100 $\pm$ 17	100 $\pm$ 11	100 $\pm$ 22	100 $\pm$ 12	100 $\pm$ 22
mp35/IRES/mp40-N220L (mIL-12mut)	94 $\pm$ 22	96 $\pm$ 7.9	2.2 $\pm$ 0.5	98 $\pm$ 18	148 $\pm$ 19

a; Total 20 μ g of each DNA construct was used for transfection into COS-7 cells by electroporation. For simultaneous transfection of two DNA constructs, 10 μ g of each DNA construct was used.

b; The levels of human and mouse IL-12p70 or IL-12p40 expression were measured by ELISA and presented relative to the wild-type, which is arbitrarily set at 100%.

c; The equal amount of p70 in each mutant supernatant

WO 01/68802

PCT/KR01/00408

was used in the IFN- γ induction assay. The levels of induced IFN- γ were measured by ELISA and presented relative to the level of wild type, which is arbitrarily set at 100%.

5 d; IL-12p40 indicates the monomeric and homodimeric p40 form of IL-12p40, but not the p40 part of IL-12p70.

e; IL-12p70 indicates the heterodimer which consists of IL-12p35 and IL-12p40.

10 f; The plasmid backbone used for independent expression of hIL-12p40, hIL-12p35, and their derivatives is pCIN-neo and is described as mock.

g; <1 indicates values below the detectable range of ELISA assay.

15 h; When the equal amount of p70 in each indicated mutant (hp40-N135Q or hp40-N222Q) and wild-type supernatant is used in IFN- γ induction assay, the insufficient hp40 in each mutant supernatant compared to wild-type supernatant was reconstituted with hp40 supernatant obtained after transfection with hp40 only.

20 i; Numbers in () mean each amount (μ g) of cotransfected plasmid, pCIN-hp40-N222Q and pCIN-hp35. Total 20 μ g of DNA was cotransfected and insufficient amount of DNA was supplemented with pCI-neo plasmid.

25 j; The plasmid backbone used for co-expression of hIL-12p40 and hIL-12p35 or their derivatives is pGX0 and is described as mock.

k; The plasmid backbone used for co-expression of mLIL-

WO 01/68802

PCT/KR01/00408

12p40 and mIL-12p35 or their derivatives is pTV2 and is described as mock.

As seen in TABLE 1, mutation of Asn-141 of p35 and
5 Asn-303 of p40 had no effect on IL-12p70 or IL-12p40 secretion. But, the secretion of IL-12p40 in Asn-135 and Asn-222 mutants was decreased. Especially, Asn-222 mutant showed same result when its amino acid was replaced with leucine(Leu) or glutamine(Gln),
10 indicating the loss of N-glycosylation at Asn-222 is very important.

FIG. 3a shows immunoblotting results of cell lysates after that wild type and mutant human IL-12p35 genes were transfected into COS-7 cells. Column 1 and 2
15 show results from cell lysates transfected with pCI-neo and pCIN-hp35 respectively. Column 3, 4, 5 and 6 show results from cell lysates transfected with expression vectors containing mutant gene such as pCIN-hp35-N127Q, pCIN-hp35-N141Q, pCIN-hp35-N251Q and pCIN-hp35-
20 N127,251Q respectively. About 33.2 kDa band in column 2 and 5 is the protein of N-glycosylated p35 subunit. When N-glycosylation was inhibited by tunicamycin, as shown in column 3 and 4, 28 kDa band was formed. Therefore, Asn-127 and Asn-141 amino acids are the N-
25 glycosylation site.

FIG. 3b shows immunoblotting results of cell lysates obtained from COS-7 cells transfected with wild

WO 01/68802

PCT/KR01/00408

type and mutant human IL-12p40 genes. Column 1 and 2 show results from cell lysates transfected with pCI-neo and pCIN-hp40 respectively. Column 3, 4, 5 and 6 show results from cell lysates transfected with expression
5 vectors containing mutant gene such as pCIN-hp40-N125Q, pCIN-hp40-N135Q, pCIN-hp40-N222Q and pCIN-hp40-N303Q respectively. Again, column 7, 8, 9 and 10 show results of cell lysates transfected with double or triple mutant genes. A result of cell lysates transfected with
10 pCIN-hp40 and tunicamycin, an antagonistic agent against N-glycosylation, is shown in column 11.

In many reports, 3-4 bands having 36-45 kDa are known in immunoblot assay of IL-12p40. Similarly, 36, 37.5, 40 and 43.1 kDa bands are shown in immunoblot
15 assay of cell lysates transfected with wild type IL-12p40 and IL-12p35. In cell lysates treated with tunicamycin, only 36 kDa band was remained while the other three bands were disappeared(Column 11). So, this band is p40 subunit which is not N-glycosylated. Bands
20 obtained from Asn-125 or Asn-135 mutation, were almost same as those of wild type(Column 3, 4). Bands were shifted in Asn-222 and Asn-303(Column 5, 6), indicating amino acids of 135, 222 and 303-Asn are the N-glycosylation site.

25 FIG. 3c shows immunoblotting results of supernatants obtained from COS-7 cell culture transfected with wild type and mutant human IL-12p40

WO 01/68802

PCT/KR01/00408

genes. Explanations for each column are same as FIG. 3b. Similar to the result of cell lysates, 3-4 bands having 36-45 kDa were made. Bands obtained from Asn-125 or Asn-135 mutation, were almost same as those of wild-type(Column 3, 4). Bands were shifted in Asn-222 and Asn-303(Column 5, 6). Especially, Asn-135 and Asn-222 mutants were not secreted to the cell culture medium, contrary to the results of cell lysates. This result is consistent with the quantification result obtained by ELISA.

Expression vectors, pGX0-hp35/IRES/hp40-N222L and pGX0-hp40-N222L/IRES/hp35, containing mutant gene of the present invention in which Asn-222 is replaced with Leu-222, were deposited at Gene Bank of Korea Research Institute of Bioscience and Biotechnology on February, 29, 2001(Accession No: KCTC 0969BP and KCTC 0970BP).

<4-1> The effect of N-glycosylation of hp35 on the synthesis, heterodimerization, and secretion of hIL-12p70.

The effect of N-glycosylation of hp35 on the synthesis, heterodimerization, and secretion of hIL-12p70 was analyzed by ELISA with both culture supernatants and lysates of cells transfected with hIL-12 expression vectors including wild type hp35 gene or its N-glycosylation mutant genes.

As seen in TABLE 1, the removal of potential N-

WO 01/68802

PCT/KR01/00408

glycosylation residues of hp35, except for Asn-127, did not significantly affect the synthesis, heterodimerization and secretion of hIL-12p70. However, deglycosylation of Asn-127 was shown to decrease the heterodimerization and secretion of hIL-12p70 to some degree, since the expression levels of hp35-N127Q and hp35-N127,141Q on western blot, in which transfection efficiency was normalized, were similar to those of other mutants. Accordingly, these results indicate that the N-glycosylation at Asn-127 of hp35 is important for the heterodimerization and secretion of hIL-12p70, but not at Asn-141.

<4-2> The effect of N-glycosylation of hp40 on the synthesis, heterodimerization, and secretion of hIL-12p70

To determine the effect of N-glycosylation of hp40 on the synthesis, heterodimerization, and secretion of hIL-12p70, the levels of hIL-12p70 and hIL-12p40 existing in both culture supernatants and lysates of cells transfected with hIL-12 expression vectors including wild type hp40 gene or its N-glycosylation mutant genes were analyzed by ELISA.

Mutation of Asn-135 or Asn-222 had little effect on hIL-12p70 secretion, as seen in TABLE 1, because the extracellular level of hIL-12p70 of these mutants was similar to that of wild type hp40. Interestingly, the

WO 01/68802

PCT/KR01/00408

secretion of hIL-12p40 in Asn-135 and Asn-222 mutants was significantly decreased, specially, Asn-222 mutant showed the secretion level less than about 9% compared with that of wild type hp40, indicating that N-glycosylation at Asn-222 is required for the secretion of hIL-12p40 alone, but not heterodimeric hIL-12, hIL-12p70. Also, double and triple mutants containing Asn-135 and/or Asn-222 showed the low level of hIL-12p40, but not that of hIL-12p70. In contrast, the other mutants appeared to produce an equal amount of hIL-12p40 and hIL-12p70 in cell lysates compared with wild type hp40. The mutations of Asn-125 and Asn-303 did not cause significant difference in the expression, heterodimerization and secretion of hIL-12p40 and hIL-12p70.

As a whole, these data indicate that N-glycosylation of hp40 at Asn-222 is critical for the secretion of IL-12p40 but not required for the heterodimerization and secretion of IL-12p70, whereas N-glycosylation of hp35 at Asn-127 is important for the heterodimerization and secretion of IL-12p70. It is also consistent with the report that N-glycosylation of hp35 appears to be a key requirement for the secretion of IL-12p70 (Carra, G., et al., *J. Immunol.*, 164:4752-4761, 2000).

<4-3> The effect of N-glycosylation of hIL-12 on its

WO 01/68802

PCT/KR01/00408

biological activity

To investigate the role of N-glycosylation of hIL-12 on its biological activity, the IFN- γ induction ability of wild type hIL-12 and its derivatives with
5 mutation at putative N-glycosylation sites was analyzed. The culture supernatants containing equal amount of wild-type hIL-12p70 and its mutants, 100 ng/ml each, were incubated with human PBLs. The level of induced IFN- γ in each culture supernatant was determined by
10 ELISA. As seen in TABLE 1, there was little difference between wild-type and all of its derivatives in terms of IFN- γ induction. However, hp40 derivatives mutated at Asn-135 and/or Asn-222 showed some increased IFN- γ induction compared with wild-type or the other hp40
15 mutants. It is probably due to the fact that the level of hIL-12p40, which was known to be an antagonist of hIL-12p70, in culture supernatants of mutants containing hp40-N135Q and/or hp40-N222Q was relatively very lower than that of the other mutants.

20

Example 5: Role of IL-12p35 in secretion of IL-12p70
containing mutated glycosylation

To investigate how deglycosylation at Asn-222 of hp40 decreases the secretion of hIL-12p40, but not that
25 of hIL-12p70, present inventors co-transfected the restricted amount of hp40-N222Q-expressing plasmid with

WO 01/68802

PCT/KR01/00408

the various amount of wild-type hp35 DNA.

Human peripheral blood mononuclear cells(PBMC) were isolated by Ficoll-Hypaque(Sigma) density gradient centrifugation from fresh blood and resuspended in
5 RPMI-1640 medium(GIBCO-BRL) supplemented with 10% heat inactivated FBS and penicillin/streptomycin(GIBCO-BRL). For the human IFN- γ induction assay, human PBM cells(4×10^5) were incubated with culture supernatants containing 100 ng/ml of human IL-12p70 or its mutant
10 derivatives for 16 hours.

For the mouse IFN- γ detection, spleens of 6 to 8 weeks old female BALB/c mice were obtained and 1×10^5 splenocytes were incubated with culture supernatants containing 100 ng/ml of mouse IL-12p70 or its mutant
15 derivatives for 24 hours. The amounts of induced human and mouse IFN- γ were estimated by human and mouse IFN- γ ELISA kit(R&D systems), respectively. The results are given in TABLE 1. The numbers in TABLE 1 is the amount of IFN- γ measured by ELISA and presented
20 relative to the level of wild type, which is arbitrarily set at 100%.

Generally, it was known that p35 subunit is not secreted alone and is secreted as a form of IL-12p70 in association with p40 subunit, whereas p40 subunit is
25 secreted in the form of either monomer or homodimer, suggesting that p40 subunit is the major factor in the secretion of IL-12p70. As shown in TABLE 1, the

WO 01/68802

PCT/KR01/00408

secretion level of hIL-12p70 increased in proportion to the amount of transfected hp35 DNA in the case of both wild-type hp40 and hp40-N222Q. This result indicates that hp40 subunit with secretion defect is secreted in the form of IL-12p70 if it is associated with hp35 subunit, suggesting that hp35 subunit also gives another help to the secretion of hIL-12p70. Recently, the report that the conformational change in hp40 subunit happens upon the binding of hp35 suggests the possibility for the contribution of hp35 subunit to IL-12p70 secretion(Yoon, C. et al., *EMBO J.*, 19:3530-3534, 2000). On the basis of this report and the data of the present invention, it is verified that hp40 containing deglycosylation at Asn-222 is defect in the secretion of itself but can be secreted in the association with hp35 due to the conformational change of the hp40 region and subsequent exposure or generation of covered or new secretion signal, respectively.

20 Example 6: Construction of HCV-E2 DNA vaccine and
expression vector containing mouse IL-12p40 Asn-220
mutant gene

<6-1> Construction of pCIN-mp40-N220L

To inspect the possibility for the use of hIL-12 mutant gene of the present invention in gene therapy as a DNA vaccine, the sequence of mouse IL-12p40(mp40)

WO 01/68802

PCT/KR01/00408

gene homologous to Asn-222 of hp40 gene was searched. Asn-220 of mp40 was located in the very similar context to that of Asn-222 of hp40, but this amino acid was not known to be N-glycosylated until now. Therefore, the
5 inventors generated the mutant mp40 gene, mp40-N220L, in which Asn at 220 in amino acid sequence was replaced with Leu by site-directed mutagenesis. For mutagenesis, SEQ. NO. 22 and SEQ. NO. 23 were used as primers. For easy identification of mutant gene, SacI restriction
10 site was generated. Amplified mutant genes were verified by the treatment of restriction enzyme to specific recognition site produced after mutagenesis and DNA sequence analysis. Eventually, the pCIN-mp40-N220L vector containing mouse IL-12p40 mutant gene
15 which can be expressed in animal cells was constructed.

<6-2> Construction of pTV2-mp35/IRES/mp40-N220L vector

To generate a vector encoding mouse p35 and p40 subunits and being used for the DNA immunization, pTV2
20 vector, an eukaryotic expression vector used as a DNA vaccine vector in small animals (Lee, et al., *J. Virol.*, 72:8430-8436, 1998; Cho, et al., *Vaccine*, 17:1136-1144, 1999), was treated with Asp718 and NotI. pTV2-mp35/IRES/mp40 vector was constructed by inserting
25 mp35/IRES/mp40 fragment obtained from pSK-mp35/IRES/mp40 treated with restriction enzymes into therein. And, pSK-mp35/IRES/mp40-N220L vector

WO 01/68802

PCT/KR01/00408

containing Asn-220 mutant gene, which can express p35, was generated by inserting mp40-N220L fragment into pSK-mp35/IRES/mp40 treated with NcoI and NotI. To delete mp40 fragment, pTV2-mp35/IRES/mp40 vector was treated with EcoRV and NotI, followed by added mp40-N220L. As a result, the pTV2-mp35/IRES/mp40-N220L was constructed.

The pTV2-mp35/IRES/mp40-N220L vector of the present invention was deposited at Gene Bank of Korea Research Institute of Bioscience and Biotechnology on February, 29, 2000 (Accession No: KCTC 0745BP).

<6-3> Construction of pTV2-HCV-E2 vector

pTV2-HCV-E2 DNA vaccine vector which can express HCV-E2 protein in eukaryotic cells was constructed (Song M. K., et al., *J. Virol.*, 74:2920-2925, 2000). As seen in FIG. 4, pTV2-HCV-E2 DNA vaccine vector is consist of simian virus 40 replication origin (SV40 ori), cytomegalovirus (CMV) promoter, tripartite leader sequence (TPL) of adenovirus, multiple cloning sequence (MCS), SV40 polyadenylation sequence (poly A) and ampicillin resistance gene (AmpR). And also, HCV-E2 gene is cloned in MCS of this vector. Carboxyl-terminal (C-terminal) containing hydrophobic amino acid residue of E2 gene used in the present invention was removed to facilitate protein secretion. And to help protein expression and cell secretion, aminocyl-

WO 01/68802

PCT/KR01/00408

terminal (N-terminal) and signal sequence (S) of herpesvirus (HSV) glycoprotein D (gD) were linked.

<6-4> Secretion of IL-12p40 and IL-12p70 according to
5 the mutation of mouse IL-12p40 Asn-220

Vectors listed in TABLE 1 were transfected into COS-7 cells using the same method in <Example 4>. The culture supernatants and cell lysates were analyzed through ELISA (PharMingen). Expression levels of wild-type IL-12p70 and 40 were shown in TABLE 1. <1 in pCI-
10 neo means that proteins were not detected by ELISA.

As seen in TABLE 1, mp40-N220L mutants showed the similar characteristics to Asn-222 mutant of hp40 in the point of the secretion of IL-12p40 and IL-12p70 and
15 its biological activity.

To apply the mp40-N222L mutant gene into the DNA vaccine model and to compare it with wild-type mp40 gene in a view of the induction of Th1 and CTL immune responses, this inventors inserted mp35/IRES/mp40 (mIL-
20 12wt) or mp35/IRES/mp40-N220L (mIL-12mut) genes into the pTV2 DNA vaccine vector and observed their characteristics *in vitro*. In result, as seen in TABLE 1, mIL-12mut (mp40-N220L mutant) gene in pTV2 vector also showed the similar characteristics to Asn-222 mutant of
25 hp40 in the point of the secretion of IL-12p40 and IL-12p70 and its biological activity.

Example 7: Effect of mutant IL-12 on antigen-specific humoral immune response

In a previous report, DNA vaccination of plasmid encoding HCV-E2 antigen(pTV2-gDsE2t) was sufficient to induce the antigen-specific humoral and cell-mediated immune responses after 3 weeks post-immunization(Song, M. K. et al., *J. Virol.*, 74:2920-2925, 2000). To determine whether the mIL-12mut gene also can affect the efficient antigen-specific immune response *in vivo* compared with mIL-12wt gene, mice were initially immunized and boosted with pTV2-mIL-12mut or pTV2-mIL-12wt, together with pTV2-gDsE2 plasmid 4 weeks later.

To inspect the induction of immune response by DNA vaccine with expression vector containing mutant genes of the present invention, 6 to 8-weeks old BALB/c mice were immunized with various pTV2 or pTV2-gDsE2t DNAs with mutant mIL-12 or wild-type mIL-12 DNAs. Particularly, the anterior tibialis muscles of each mouse were injected with a total 200 μ g of DNA, formulated in a final volume of 100 μ l of phosphate-buffered saline and boosted on time with an identical dose of DNA at 4 week interval. Humoral immune responses were monitored 3 weeks after booster immunization by ELISA. In particular, 100 ng of hghE2t protein, which is a fusion protein of human growth hormone(hgh) and C-terminal truncated HCV E2(HCV E2t),

WO 01/68802

PCT/KR01/00408

was coated to each well of 96-well plate (Dynex Technologies). Sera from immunized mice were diluted to 1:100 for reaction with hghE2t protein and introduced to the determination of relative levels of HCV E2t-specific IgG, and its subclasses, such as IgG1 and IgG2a.

As shown in FIG 5a, 5b, 5c and 5d, HCV E2 DNA vaccine induced systemic HCV E2-specific total IgG, IgG1 and IgG2a levels significantly higher than negative control values, and co-injection with mIL-12mut gene or mIL-12wt gene showed the similar total IgG level compared to HCV E2 DNA vaccine only. And, the level of IgG1 was similar among the groups injected with HCV E2 DNA. In contrast, the level of IgG2a against HCV E2 slightly increased in mIL-12wt group and significantly increased in mIL-12mut group compared with HCV E2 DNA immunized only group. In addition, the ratio of IgG2a/IgG1, which is generally accepted as an indirect indicator of Th1 immunity, was the highest in the mIL-12mut group, which showed the similar pattern of IgG2 level. These data represent that mIL-12mut gene significantly affected the shift of IgG subclasses from IgG1 to IgG2a in humoral immune response compared with mIL-12wt or HCV E2 only group, suggesting that IL-12mut may induce Th1 type immune response.

Example 8: Cell-mediated immune response of immunized

WO 01/68802

PCT/KR01/00408

mice

<8-1> Induction of antigen-specific Th1 immune response by mutant IL-12

To investigate the effect of mIL-12mut gene on Th1 immune response, which is one of the parameter used to evaluate the potency of cell mediated immunity, IFN- γ expression of splenocytes was measured. 1×10^5 splenocytes obtained at 8 weeks after the booster immunization were added to wells of U-bottomed 96-well plate. Then, 1 or 5 μ g of hgh-E2t protein purified from CHO cells was added to each well and the cells were incubated at 37°C, in 5% CO₂ incubator for 3 days. And then, cell supernatants were secured and used for the detection of IFN- γ levels using ELISA kit(R&D systems). It was known that IFN- γ induced after stimulation with specific antigen is produced from antigen-specific CD4+ T cell and it is the indicator of Th1 immune response.

As shown in FIG. 6, HCV E2 DNA immunized group without cytokine gene showed the increased level of IFN- γ in proportion to the concentration of hghE2t protein, whereas mock plasmid immunized group did not. Expectedly, the level of IFN- γ induction in mIL-12wt group was more enhanced than in HCV E2 only group, and mIL-12mut group showed up to 2-3 times higher production of IFN- γ than mIL-12wt group, suggesting that mIL-12p70 increases the antigen-specific Th1

WO 01/68802

PCT/KR01/00408

immune response and mIL-12p40 inhibits the induction of Th1 immune response by IL-12p70 *in vivo*.

<8-2> Long-term enhancement of antigen-specific CD8+ T cell function by mutant IL-12

5 As noted above, mIL-12mut gene contributed to the long-term Th1 immune response in HCV E2 DNA immunization. To determine whether the long-term Th1 immune response induced by the expression of mIL-12mut gene correlates with CTL immunity and the major cell-mediated immune response, and therefore mIL-12mut gene can affect the maintenance of CTL activity in DNA immunization model, the present inventors performed the CTL assay with splenocytes of DNA immunized mice at various weeks after booster immunization.

10 Splenocytes(2×10^7) were restimulated *in vitro* at 37°C with mitomycin C-treated($25 \mu\text{g/ml}$) CT26-hghE2t cells(1×10^6), which express truncated form of HCV envelope protein 2(E2t). After 5 days *in vitro* culture, effector cells were tested in a conventional

15 cytotoxicity assay against different target cell, such as CT26-hghE2t or CT26-neo. Various numbers of effector cells were plated in triplicate to achieve the desired E/T ratio. ^{51}Cr -labeled target cells(5×10^3) were added to each well of U-bottomed 96-well plate, and after 6

20 hours incubation at 37°C, the supernatant was harvested and counted with a γ -counter(Wallac, Turku, Finland). The percentage of specific lysis was calculated as

following mathematical formula 1. Minimum lysis was obtained by incubating the target cells with the culture medium alone. Maximum lysis was obtained by exposing the target cells to 1% Nonidet-P40TM.

5

<Mathematical formula 1>

Percentage of specific lysis = (Experimental lysis-Minimum lysis)x100/(Maximum lysis-Minimum lysis)

10

As a result, as shown in FIG. 7, two weeks after boost, all groups except for mock plasmid immunized group showed very strong antigen-specific CTL activity. However, there was little difference among HCV E2 only, mIL-12wt and mIL-12mut groups. CTL response was induced more in mIL-12wt coimmunized group than in HCV E2 only group in overall period, indicating that mIL-12 gene played a role in the enhanced CTL generation. Interestingly, the difference in CTL activity between mIL-12mut group and the other two groups, HCV E2 only and mIL-12wt group, became bigger and bigger as the time after booster immunization was longer. Especially, at 10 weeks, CTL response was very low in HCV E2 only and mIL-12wt groups, suggesting that the frequency of antigen-specific CTL decreased significantly after long period, whereas mIL-12mut group showed 5 to 10 times higher CTL activity than the other two groups sustaining the antigen-specific CTL response. As a

25

control, when CT26-neo cell was used as the target cell, no lysis was observed in all groups, suggesting that CTL activity observed in this experiment is HCV E2-specific.

5

<8-3> FACSCaliburTM flow cytometry analysis for IFN- γ production of CD8⁺ cells of immunized mice

To investigate whether the long-term enhancement of CTL activity is originated from the frequency of antigen-specific CD8⁺ T cell, and for determination of CD8⁺ antigen-specific T lymphocyte frequency, following experiment was performed. Splenocytes(2×10^7) obtained at the indicated week after the booster immunization were stimulated with CT26-hghE2t cells(1×10^6) in the presence of 10 U/ml recombinant IL-12(PharMingen) for 40 hours, and then 4 μ l of GolgiStoptm(PharMingen) was added and the cells were incubated for another 8 hours at 37°C. For the direct purification of CD8⁺ T cells, the stimulated splenocytes were incubated with anti-CD8 microbeads(Miltenyi Biotech, Inc) and then passed through a column of miniMACSTM system (Miltenyi Biotech, Inc), and the remaining CD8⁺ T cells were isolated. To block nonspecific staining, cells were preincubated with Fc BlockTM(PharMingen) and stained with FITC-conjugated anti-mouse CD8. After the incubation, the cell suspensions were fixed and permeablized with Cytofix/CytopermTM(PharMingen) before adding PE-

25

20

15

10

conjugated anti-mouse IFN- γ mAb or control PE-conjugated isotype-matched mAb. Stained cells were analyzed by FACSCaliburTM flow cytometry (Becton Dickinson), and then induction of IFN- γ was observed.

5

<Table 2> Frequency kinetics of HCV-E2-specific precursor CD8+ T cells

No. of immunization	Weeks after final immunization	Group	Frequency of HCV-E2-specific CD8 ⁺ T cells							
			Intracellular IFN- γ staining assay (%)				Limiting dilution assay (No.)			
			I	II	III	IV	I	II	III	IV
10	0		0.08	0.34	0.42	0.58	7.1	36.1	39.0	47.1
	3		0.05	0.34	0.48	0.72	8.4	39.0	48.8	66.7
	6		0.06	0.29	0.39	0.73	6.9	34.8	46.8	64.9
	10		0.07	0.21	0.20	0.68	7.2	18.2	19.8	64.3
	14		0.07	0.09	0.09	0.57	8.1	9.6	8.8	45.7
15	4		0.06	0.33	0.42	0.55	7.5	33.2	41.0	48.7
	8		0.08	0.26	0.27	0.51	8.4	11.3	12.5	48.2
	14		0.06	0.10	0.07	0.42	7.1	6.8	8.1	44.7

a; Six- to 8-week-old female BALB/c mice were immunized with various plasmids of the present invention at 4-week interval.

20 b; The expression of each group is as follows. Group I; pTV2, Group II; pTV2-HCV-E2t+pTV2, Group III; pTV2-HCV-E2t+pTV2-mIL-12wt, Group IV; pTV2-HCV-E2t+pTV2-mIL-12mut

25 c; 2×10^7 of splenocytes obtained from DNA-immunized mice were restimulated with 1×10^6 of mitomycin C-treated CT26-hghE2t cells in vitro. After 48 hours in culture, CD8+ T cells were isolated by using MACSing.

WO 01/68802

PCT/KR01/00408

After fixation and permeabilization, the cells were stained with anti-CD8 and anti-IFN- γ antibody. Live CD8+ T cells were gated by plotting of cells with FSC and CD8 and then, the percentage of IFN- γ -producing CD8+ T cells among live CD8+ T cells was calculated by plotting of live CD8+ T cells with CD8 and IFN- γ . Data are represented as the average value obtained with 2 mice per group in two independent experiments.

d; Splenocytes obtained from DNA-immunized mice were diluted, mixed with mitomycin C-treated CT26-hghE2t cells, and incubated for 5 days. The specificity of resulting CTL was determined by specific lysis of ^{51}Cr -labeled CT26-hghE2t cells. Wells were scored as positive for CTL recognition if the level of specific lysis exceeds the mean lysis value plus 3SD obtained from naive mice. The frequency of precursor CTL per 1×10^7 spleen cells was calculated by regression analysis of the number of negative wells at each dilution of responder cells. Data are represented as the average value obtained with 2 mice per group in two independent experiments.

As shown in TABLE 2, mice coimmunized with mIL-12mut gene had a 3 to 7-fold enhancement in the frequency of CD8+ IFN- γ producing cells compared with HCV E2 only and mIL-12wt groups at 0, 3, 6, 10 or 14 weeks after booster immunization, correlating with the

WO 01/68802

PCT/KR01/00408

result in CTL response. By contrast, in isotype-matched control experiment, there was no difference among all groups. Like the result of CTL assay, the frequency of IFN- γ producing CD8+ T cells among immunization groups was not much different at 2-3 weeks after booster immunization. These data demonstrate that the expression of mIL-12mut gene sustained the frequency of CD8+ IFN- γ producing T cells after DNA immunization for long period. In this regard, it might be suggested that *in vivo* role of IL-12p70 itself in cell-mediated immune response is the long-term maintenance of Th1 and CTL, whereas IL-12p40 inhibits the IL-12p70 as an antagonist *in vivo*.

<8-4> The frequency of antigen-specific CD8+ T cells of splenocytes of immunized mice

To investigate the frequency of other antigen-specific CD8+ T cells, limiting dilution assay(LDA) was conducted. In this experiment, the frequency of antigen-specific CD8+ T cells was measured by using lysis ability of HCV E2 specific CD8+ T cells(Kuzushima, K. *et al.*, Blood, 94:3094-3100, 1999). Splenocytes of booster immunized mice were diluted with various concentration, and placed into wells of U-bottomed 96-well plate to 20 wells/dilution. CT-26 cells expressing E2, were treated with mytomycin C(500 $\mu\text{g}/\text{ml}$) to block cell division, and activated with diluted splenocytes

WO 01/68802

PCT/KR01/00408

for 5 days. ^{51}Cr -labeled target cells (5×10^3) were added to each well of the above well plate, and after 6 hours at 37°C , the supernatant was harvested and counted with a γ -counter (Wallac, Turku, Finland). CTL frequency was
5 calculated and regarded as positive when the level of specific lysis was higher than that of mean lysis value + 3 x standard deviation (Kuzushima, K. et al., *Blood*, 94:3094-3100, 1999).

The result of limiting dilution assay was similar
10 to the result of intracellular staining assay. Namely, the highest frequency of antigen-specific CD8+ T cell was observed in mIL-12 group even in early stage of immunization, and this frequency had been remained over 14 weeks after booster immunization (TABLE 2).

15

<8-5> Enhancement of protective immune response by mutant IL-12

To confirm the Th1 and CTL immunity induced by mIL-12mut gene *in vivo*, and to examine that the
20 correlation of protective immunity with Th1 and CTL immunity, CT26-hgHE2t tumor cells expressing hgHE2t were injected in the groups of the immunized mice at 12 weeks after the booster immunization. 2 weeks after, relative levels of antigen-specific IgG, IgG1, IgG2 and
25 the ratio of IgG2a/IgG1 were determined, and the size of tumor was measured for 30 days. Particularly, the mean local tumor growth was determined by measuring the

WO 01/68802

PCT/KR01/00408

volume and diameter of tumors with calipers every three days. Also, the survival rates of these mice were determined by observing for about 70 days.

As shown in FIG. 8a, 8b and 8c, the group of mice immunized with mIL-12mut induced strong Th1 immune response. The group of mIL-12wt immunization displayed the delayed tumor growth in contrast to the group of only pTV2-gDsE2t immunization, whereas the group of mIL-12mut immunization showed the significantly delayed tumor growth. In control group, most of mice had tumor and died within 50 days, but 90% of mice in the group of mIL-12mut could survive after 70 days. Thus, These data suggest that HCV E2-specific Th1 and CTL responses induced by mIL-12mut gene confers *in vivo* protection against the challenge of modified tumor cells expressing specific antigen. Although it is not easy to evaluate relative effects of Th1 and CTL responses on tumor protection *in vivo*, it is likely that E2-specific CD8+ CTLs and Th1 cells could directly kill the CT26-hghE2t cells and help the CTLs, respectively as shown in the *in vitro* Th1 and CTL assay. Also, IgG2a antibodies, which strongly binds to Fc γ R on macrophages and natural killer cells, might mediate antibody-dependent cell-mediated cytotoxicity.

25

INDUSTRIAL APPLICABILITY

As described hereinbefore, the present invention relates to the IL-12p40 subunit mutant gene which can produce interleukin 12(IL-12) of human and mouse origin with high activity, and the expression vector including above mutant gene as well as the use of them to DNA vaccine adjuvant. Particularly, it relates to IL-12p40 mutant gene which inhibits the secretion of IL-12p40 but normally secretes active IL-12p70 by making mutation at Asn-222(human) or Asn-220(mouse) amino acid of IL-12p40, which acts as a competitive inhibitor of active form of IL-12, IL-12p70. IL-12p40 mutant gene of the present invention can induce optimal cell-mediated immune response at early stage and for a long time if immunized with DNA vaccine. Therefore, the IL-12p40 mutant gene of the present invention can be useful for DNA vaccination and gene therapy to various diseases, for example, AIDS, hepatitis C or hepatitis B, cancer, influenza, tuberculosis and malaria, which essentially require cellular immune responses for their therapy. Those skilled in the art will appreciate that the conceptions and specific embodiments disclosed in the foregoing description may be readily utilized as a basis for modifying or designing other embodiments

WO 01/68802

PCT/KR01/00408

for carrying out the same purposes of the present
invention. Those skilled in the art will also
appreciate that such equivalent embodiments do not
depart from the spirit and scope of the invention as
5 set forth in the appended claims.

10

15

issued pursuant to Rule 7.1

WO 01/68802

PCT/KR01/00408


BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT
OF MICROORGANISMS FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued pursuant to Rule 7.1

TO : SUNG, Young Chul
Department of Life Science, POSTECH,
#San31, Hyoja-dong, Nam-ku, Pohang-si, Kyungbuk 790-784,
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: <i>Escherichia coli</i> DH5@/pGX0-hp40-N222L/IRES/hp35	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCTC 0970BP
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: [x] a scientific description [] a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on February 26 2001 .	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Korean Collection for Type Cultures Address: Korea Research Institute of Bioscience and Biotechnology (KRIBB) #52, Oun-dong, Yusong-ku, Taejon 305-333, Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official(s):  BAE, Kyung Sook, Director Date: March 02 2001

WO 01/68802

PCT/KR01/00408


BUDAPEST TREATY ON THE INTERNATIONAL RECOGNITION OF THE DEPOSIT
OF MICROORGANISMS FOR THE PURPOSE OF PATENT PROCEDURE

INTERNATIONAL FORM

RECEIPT IN THE CASE OF AN ORIGINAL DEPOSIT

issued pursuant to Rule 7.1

TO : SUNG, Young Chul
Department of Life Science, POSTECH,
#San31, Hyoja-dong, Nam-ku, Pohang, Kyung-buk 790-784,
Republic of Korea

I. IDENTIFICATION OF THE MICROORGANISM	
Identification reference given by the DEPOSITOR: <i>Escherichia coli</i> DH5 α /pTV2-mp35/TRES/np40-N220L	Accession number given by the INTERNATIONAL DEPOSITARY AUTHORITY: KCTC 0745BP
II. SCIENTIFIC DESCRIPTION AND/OR PROPOSED TAXONOMIC DESIGNATION	
The microorganism identified under I above was accompanied by: ☒ a scientific description ☐ a proposed taxonomic designation (Mark with a cross where applicable)	
III. RECEIPT AND ACCEPTANCE	
This International Depositary Authority accepts the microorganism identified under I above, which was received by it on February 29 2000 .	
IV. RECEIPT OF REQUEST FOR CONVERSION	
The microorganism identified under I above was received by this International Depositary Authority on _____ and a request to convert the original deposit to a deposit under the Budapest Treaty was received by it on _____	
V. INTERNATIONAL DEPOSITARY AUTHORITY	
Name: Korean Collection for Type Cultures Address: Korea Research Institute of Bioscience and Biotechnology (KRIBB) #52, Oun-dong, Yusong-ku, Taejeon 305-333, Republic of Korea	Signature(s) of person(s) having the power to represent the International Depositary Authority of authorized official(s):  BAE, Kyung Soek, Director Date: March 06 2000

WO 01/68802

PCT/KR01/00408

SEQUENCE LISTING

<110> Pohang University of Science and Technology et al.

<120> Genes of IL-12p40 subunit mutated for improving the activity of IL-12 and use thereof for DNA vaccine adjuvant

<130> 1p-02-14

<150> KR 2000-13133

<151> 2000-03-15

<160> 23

<170> KopatentIn 1.55

<210> 1

<211> 1007

<212> DNA

<213> Homo sapiens

<400> 1

agcaagatgt gtcaccagca gttggtcate tcttggtttt cccctggtttt tctggcatct 60

cccctcgtgg ccatatggga actgaagaaa gatgtttatg tcgtagaatt ggattggtat 120

ccggatgccc ctggagaaat ggtggtcctc acctgtgaca cccctgaaga agatggtatc 180

acctggacct tggaccagag cagtgaggtc ttaggctctg gcaaaaccct gaccatccaa 240

gtcaaagagt ttggagatgc tggccagtac acctgtcaca aaggaggcga ggttctaagc 300

cattcgctcc tgctgcttca caaaaaggaa gatggaattt ggtccactga tattttaag 360

gaccagaaag aacccaaaaa taagaccttt ctaagatgcg aggccaagaa ttattctgga 420

WO 01/68802

PCT/KR01/00408

cgtttcacct gctgggtggct gacgacaatc agtactgatt tgacattcag tgtcaaaagc 480
 agcagaggct cttctgaccc ccaaggggtg acgtgcggag ctgctacact ctctgcagag 540
 agagtcagag gggacaacaa ggagtatgag tactcagtgg agtgccagga ggacagtgcc 600
 tgcccagctg ctgaggagag tctgcccatt gaggtcatgg tggatgccgt tcacaagctc 660
 aagtatgaaa actacaccag cagcttcttc atcagggaca tcacaaacc tgaccacccc 720
 aagaactgc agctgaagcc attaaagaat tctcggcagg tggaggtcag ctgggagtag 780
 cctgacacct ggagtactcc acattcctac ttctccctga cattctgcgt tcaggtccag 840
 ggcaagagca agagagaaaa gaaagataga gtcttcacgg acaagacctc agccacggtc 900
 atctgccgca aaaatgccag cattagcgtg cgggcccagg accgtacta tagctcatct 960
 tggagcgaat gggcatctgt gccctgcagt taggttctga tccagga 1007

<210> 2
 <211> 900
 <212> DNA
 <213> mouse

<400> 2
 cccaaaagag tgaccttga atcatctgta cgcttgccata tgtctagctc agttcatgct 60
 gctatcaatc catgagtaag gacctataag cataagagac gccctcaaaa cactatgact 120
 ttattagtt attcacctcc ccagagctgt ccctggatac agacaacata ggtatgaggt 180
 agggggtagc tggagccaaa caggaggtaa taccttctga atttagatgc taacaagaaa 240

WO 01/68802

PCT/KR01/00408

acatggggaa aggtggccca gatacactag gccctttatt ctttgggcct gtaacaccta 300
 cttatttgat tgtggcatga accatgaact cggttgggg caagtccttc cttttctgc 360
 agtctgtgga atcgggagag gttagccatt gccgcctcta ttcacctag gcatgatgta 420
 aacagaaatt agtatctctg cctccttctt tttccacac cccgaagtca tttcctctta 480
 acctgggatt tcgacgtcta tattccctct gtatgataga tgcactcagg gaggcaaggg 540
 ggggagggag gaacttctta aaattccccc agaatgtttt gacactagtt ttcagtgttg 600
 caattgagac tagtcagttt ctactttggg tttccatcag aaagttctgt aggagtagag 660
 tatataagca ccaggagcag ccaaggcagc agaaggaaca gtgggtgtcc aggcacatca 720
 gaccaggcag ctgcagcaa agcaaggtaa gttctctcct cttccctgtc gctaactccc 780
 tgcacttaga ggctgtccag attcagactc caggggacag gctacccttg aaccaggcag 840
 cgtgggagtg gggtaagtgg attctgggag catctcgat ggctttcccc gctggtggaa 900
 900

<210> 3

<211> 328

<212> PRT

<213> human IL-12p40-N222L

<400> 3

Met Cys His Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Phe Leu

1

5

10

15

Ala Ser Pro Leu Val Ala Ile Trp Glu Leu Lys Lys Asp Val Tyr Val

WO 01/68802

PCT/KR01/00408

	20	25	30
Val Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu Met Val Val Leu			
	35	40	45
Thr Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln			
	50	55	60
Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys			
	65	70	75
			80
Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val			
	85	90	95
Leu Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp			
	100	105	110
Ser Thr Asp Ile Leu Lys Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe			
	115	120	125
Leu Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp			
	130	135	140
Leu Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser Arg			
	145	150	155
			160
Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser			
	165	170	175
Ala Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu			
	180	185	190
Cys Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile			
	195	200	205

WO 01/68802

PCT/KR01/00408

Glu Val Met Val Asp Ala Val His Lys Leu Lys Tyr Glu Leu Tyr Thr
 210 215 220

Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn
 225 230 235 240

Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp
 245 250 255

Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr
 260 265 270

Phe Cys Val Gln Val Gln Gly Lys Ser Lys Arg Glu Lys Lys Asp Arg
 275 280 285

Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys Asn Ala
 290 295 300

Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser
 305 310 315 320

Glu Trp Ala Ser Val Pro Cys Ser
 325

<210> 4

<211> 328

<212> PRT

<213> human IL-12p40-N222Q

<400> 4

Met Cys His Gln Gln Leu Val Ile Ser Trp Phe Ser Leu Val Phe Leu
 1 5 10 15

Ala Ser Pro Leu Val Ala Ile Trp Glu Leu Lys Lys Asp Val Tyr Val
 20 25 30

WO 01/68802

PCT/KR01/00408

Val Glu Leu Asp Trp Tyr Pro Asp Ala Pro Gly Glu Met Val Val Leu
 35 40 45

Thr Cys Asp Thr Pro Glu Glu Asp Gly Ile Thr Trp Thr Leu Asp Gln
 50 55 60

Ser Ser Glu Val Leu Gly Ser Gly Lys Thr Leu Thr Ile Gln Val Lys
 65 70 75 80

Glu Phe Gly Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Val
 85 90 95

Leu Ser His Ser Leu Leu Leu Leu His Lys Lys Glu Asp Gly Ile Trp
 100 105 110

Ser Thr Asp Ile Leu Lys Asp Gln Lys Glu Pro Lys Asn Lys Thr Phe
 115 120 125

Leu Arg Cys Glu Ala Lys Asn Tyr Ser Gly Arg Phe Thr Cys Trp Trp
 130 135 140

Leu Thr Thr Ile Ser Thr Asp Leu Thr Phe Ser Val Lys Ser Ser Arg
 145 150 155 160

Gly Ser Ser Asp Pro Gln Gly Val Thr Cys Gly Ala Ala Thr Leu Ser
 165 170 175

Ala Glu Arg Val Arg Gly Asp Asn Lys Glu Tyr Glu Tyr Ser Val Glu
 180 185 190

Cys Gln Glu Asp Ser Ala Cys Pro Ala Ala Glu Glu Ser Leu Pro Ile
 195 200 205

Glu Val Met Val Asp Ala Val His Lys Leu Lys Tyr Glu Gln Tyr Thr

WO 01/68802

PCT/KR01/00408

210 215 220
 Ser Ser Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn
 225 230 235 240
 Leu Gln Leu Lys Pro Leu Lys Asn Ser Arg Gln Val Glu Val Ser Trp
 245 250 255
 Glu Tyr Pro Asp Thr Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Thr
 260 265 270
 Phe Cys Val Gln Val Gln Gly Lys Ser Lys Arg Glu Lys Lys Asp Arg
 275 280 285
 Val Phe Thr Asp Lys Thr Ser Ala Thr Val Ile Cys Arg Lys Asn Ala
 290 295 300
 Ser Ile Ser Val Arg Ala Gln Asp Arg Tyr Tyr Ser Ser Ser Trp Ser
 305 310 315 320
 Glu Trp Ala Ser Val Pro Cys Ser
 325

<210> 5
 <211> 335
 <212> PRT
 <213> mouse IL-12p40-N220L

<400> 5
 Met Cys Pro Gln Lys Leu Thr Ile Ser Trp Phe Ala Ile Val Leu Leu
 1 5 10 15
 Val Ser Pro Leu Met Ala Met Trp Glu Leu Glu Lys Asp Val Tyr Val
 20 25 30

WO 01/68802

PCT/KR01/00408

Val Glu Val Asp Trp Thr Pro Asp Ala Pro Gly Glu Thr Val Asn Leu
 35 40 45

Thr Cys Asp Thr Pro Glu Glu Asp Asp Ile Thr Trp Thr Ser Asp Gln
 50 55 60

Arg His Gly Val Ile Gly Ser Gly Lys Thr Leu Thr Ile Thr Val Lys
 65 70 75 80

Glu Phe Leu Asp Ala Gly Gln Tyr Thr Cys His Lys Gly Gly Glu Thr
 85 90 95

Leu Ser His Ser His Leu Leu Leu His Lys Lys Glu Asn Gly Ile Trp
 100 105 110

Ser Thr Glu Ile Leu Lys Asn Phe Lys Asn Lys Thr Phe Leu Lys Cys
 115 120 125

Glu Ala Pro Asn Tyr Ser Gly Arg Phe Thr Cys Ser Trp Leu Val Gln
 130 135 140

Arg Asn Met Asp Leu Lys Phe Asn Ile Lys Ser Ser Ser Ser Pro
 145 150 155 160

Asp Ser Arg Ala Val Thr Cys Gly Met Ala Ser Leu Ser Ala Glu Lys
 165 170 175

Val Thr Leu Asp Gln Arg Asp Tyr Glu Lys Tyr Ser Val Ser Cys Gln
 180 185 190

Glu Asp Val Thr Cys Pro Thr Ala Glu Glu Thr Leu Pro Ile Glu Leu
 195 200 205

Ala Leu Glu Ala Arg Gln Gln Asn Lys Tyr Glu Leu Tyr Ser Thr Ser

WO 01/68802

PCT/KR01/00408

210	215	220	
Phe Phe Ile Arg Asp Ile Ile Lys Pro Asp Pro Pro Lys Asn Leu Gln			
225	230	235	240
Met Lys Pro Leu Lys Asn Ser Gln Val Glu Val Ser Trp Glu Tyr Pro			
	245	250	255
Asp Ser Trp Ser Thr Pro His Ser Tyr Phe Ser Leu Lys Phe Phe Val			
	260	265	270
Arg Ile Gln Arg Lys Lys Glu Lys Met Lys Glu Thr Glu Glu Gly Cys			
	275	280	285
Asn Gln Lys Gly Ala Phe Leu Val Glu Lys Thr Ser Thr Glu Val Gln			
290	295	300	
Cys Lys Gly Gly Asn Val Cys Val Gln Ala Gln Asp Arg Tyr Tyr Asn			
305	310	315	320
Ser Ser Cys Ser Lys Trp Ala Cys Val Pro Cys Arg Val Arg Ser			
	325	330	335

<210> 6
 <211> 24
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> T7 primer

<400> 6
 gtacttaata cgactcacta tagg

24

WO 01/68802

PCT/KR01/00408

<210> 7
 <211> 24
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> T3 primer

<400> 7
 gaagcattaa ccctcactaa aggg

24

<210> 8
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> hp40-N125Q sense primer

<400> 8
 cccaaacaga aaacgtttct a

21

<210> 9
 <211> 21
 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> hp40-N125Q antisense primer

<400> 9
 cgttttctgt ttgggttctt t

21

WO 01/68802

PCT/KR01/00408

<210> 10

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N135Q sense primer

<400> 10

gccaagcagt attctggacg t

21

<210> 11

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> ho40-N135Q antisense primer

<400> 11

agaatactgc ttggcctcgc a

21

<210> 12

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N222Q sense primer

<400> 12

tatgaacagt acaccagcag c

21

WO 01/68802

PCT/KR01/00408

<210> 13

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N222Q antisense primer

<400> 13

ggtgtactgt tcatacttga g

21

<210> 14

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N303Q sense primer

<400> 14

cgcaaacagg ccagcattag c

21

<210> 15

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N303Q antisense primer

<400> 15

WO 01/68802

PCT/KR01/00408

gctggcctgt ttgcggcaga t

21

<210> 16

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N127Q sense primer

<400> 16

accaagcagg agagttgcct a

21

<210> 17

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N127Q antisense primer

<400> 17

actctcctgc ttggtaatt c

21

<210> 18

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N141Q sense primer

WO 01/68802

PCT/KR01/00408

<400> 18

ataactcagg ggagttgcct g

21

<210> 19

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N141Q antisense primer

<400> 19

actcccctga gttatgaaag a

21

<210> 20

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N251Q sense primer

<400> 20

tatctgcagg cttcctaaaa a

21

<210> 21

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> hp40-N251Q antisense primer

WO 01/68802

PCT/KR01/00408

<400> 21

ggaagcctgc agatagctca t

21

<210> 22

<211> 21

<212> DNA

<213> Artificial Sequence

<220>

<223> mp40-N220L sense primer

<400> 22

tatgagctct acagcaccag c

21

<210> 23

<211> 24

<212> DNA

<213> Artificial Sequence

<220>

<223> mp40-N220L antisense primer

<400> 23

gctgtagagc tcatattttt actg

24

What is claimed is:

1. An isolated nucleic acid comprising a mutated human IL-12p40 encoding sequence, wherein the mutated human IL-12p40 encoding sequence comprises the nucleic acid sequence of SEQ ID NO: 1 with the exception that the codon for the amino acid 222 of the encoded IL-12p40 sequence encodes an amino acid with uncharged polar side chain or an amino acid with nonpolar side chain.
2. An isolated nucleic acid comprising a mutated mouse IL-12p40 encoding sequence, wherein the mutated mouse IL-12p40 encoding sequence comprises the nucleic acid sequence of SEQ ID NO: 2 with the exception that the codon for the amino acid 220 of the encoded IL-12p40 sequence encodes an amino acid with uncharged polar side chain or an amino acid with nonpolar side chain.
3. The isolated nucleic acid according to claim 1, wherein the codon for the amino acid 222 encodes leucine, glutamine or isoleucine.
4. The isolated nucleic acid according to claim 2, wherein the codon for the amino acid 220 encodes leucine or isoleucine.
5. The isolated nucleic acid according to claim 1 or 3, further comprising a human IL-12p35 encoding sequence, and an internal ribosome entry site (IRES) operably linked between the mutated human IL-12p40 encoding sequence and the human IL-12p35 encoding sequence.
6. The isolated nucleic acid according to claim 2 or 4, further comprising a mouse IL-12p35 encoding sequence, and an internal ribosome entry site (IRES) operably linked between the mutated mouse IL-12p40 encoding sequence and the mouse IL-12p35 encoding sequence.
7. An expression vector comprising the isolated nucleic acid of claim 5.


8. The expression vector according to claim 7, which is pGXO-hp35/IRES/hp40-n222L deposited under the accession No. KCTC 0969BP, wherein the isolate nucleic acid comprises the mutated human IL-12p40 encoding sequence whose codon for the amino acid 222 encodes leucine.
9. The expression vector according to claim 7, which is pGXO-hp40-N222L/IRES/hp35 deposited under the accession No. KCTC 0970BP wherein the isolate nucleic acid comprises the mutated human IL-12p40 encoding sequence whose codon for the amino acid 222 encodes leucine.
10. An expression vector comprising the isolated nucleic acid of claim 6.
11. The expression vector according to claim 10, which is pTV2-mp35/IRES/mp40-n220L deposited under the accession No. KCTC 07450BP wherein the isolate nucleic acid comprises the mutated mouse IL-12p40 encoding sequence whose codon for the amino acid 220 encodes leucine.
12. A therapeutic or prophylactic adjuvant for DNA immunization or gene therapy, which comprises an expression vector comprising the isolated nucleic acid of claim 1 or 2 as an active ingredient.
13. The adjuvant according to claim 12, for use to induce an immune response by increasing the hydrolysis activity of cytotoxic T lymphocytes, wherein the expression vector is for use by injection as a DNA vaccine.
14. The adjuvant according to claim 12, for use to induce an immune response by increasing the secretion of IFN- γ from T helper cells, wherein the expression vector is for use by injection as a DNA vaccine.
15. The adjuvant according to claim 12, for use to induce an immune response by increasing the secretion of IFN- γ from CD8 $^{+}$ cells, wherein the expression vector is for use by injection as a DNA vaccine.

16. The adjuvant according to claim 12, for use against a disease that is: a cancer, AIDS, hepatitis C, hepatitis B, influenza, tuberculosis or malaria.

FIG. 1

2/11

FIG. 2



3/11

FIG. 3a





FIG. 3b



4/11

FIG. 3c





FIG. 4



6/11

FIG. 5a





FIG. 5b



7/11

FIG. 5c






FIG. 5d



8/11

FIG. 6




PCT/KR01/00408

IPEA/KR 21.09.2001


9/11

FIG. 7




10/11

FIG. 8



11/11

FIG. 9



Viruses, Bacteria
Fungi, Protozoa

- ▲ : (p35)
- : (p40)
- ⊖ : (p40)₂
- ▲○ : IL-12 (p70)

