



Office de la Propriété
Intellectuelle
du Canada

Un organisme
d'Industrie Canada

Canadian
Intellectual Property
Office

An agency of
Industry Canada

CA 2623197 A1 2007/03/29

(21) **2 623 197**

(12) **DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION**

(13) **A1**

(86) Date de dépôt PCT/PCT Filing Date: 2006/09/22
(87) Date publication PCT/PCT Publication Date: 2007/03/29
(85) Entrée phase nationale/National Entry: 2008/03/19
(86) N° demande PCT/PCT Application No.: US 2006/037302
(87) N° publication PCT/PCT Publication No.: 2007/035930
(30) Priorité/Priority: 2005/09/22 (US60/719,952)


(51) Cl.Int./Int.Cl. *G01N 33/53* (2006.01)
(71) Demandeur/Applicant:
PROSCI INCORPORATED, US
(72) Inventeurs/Inventors:
GENG, YU, US;
LUAllen, ROBERT JAMES, US
(74) Agent: BORDEN LADNER GERVAIS LLP

(54) Titre : POLYPEPTIDES GLYCOSYLES PRODUITS DANS DES MUTANTS DE LEVURE ET METHODES
D'UTILISATION

(54) Title: GLYCOSYLATED POLYPEPTIDES PRODUCED IN YEAST MUTANTS AND METHODS OF USE THEREOF

(57) **Abrégé/Abstract:**

The invention provides various high mannose glycosylated polypeptides that are useful in a vaccine formulations. The invention also provides methods for making such glycosylated polypeptides and its uses in eliciting HIV-neutralizing antibodies.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
29 March 2007 (29.03.2007)

(10) International Publication Number
WO 2007/035930 A3

(51) International Patent Classification:
G01N 33/53 (2006.01)

(21) International Application Number:
PCT/US2006/037302

(22) International Filing Date:
22 September 2006 (22.09.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/719,952 22 September 2005 (22.09.2005) US

(71) Applicant (for all designated States except US): **PROSCI INCORPORATED** [US/US]; 12170 Flint Place, Poway, CA 90264 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GENG, Yu** [US/US]; 11614 Cohansey Road, San Diego, California 92131 (US). **LUAllen, Robert, James** [US/US]; 3630 Georgia Street, #6, San Diego, CA 92103 (US).

(74) Agents: **ZHOU, Jie** et al.; Morrison & Foerster LLP, 755 Page Mill Road, Palo Alto, CA 94304-1018 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(88) Date of publication of the international search report:
29 November 2007

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: GLYCOSYLATED POLYPEPTIDES PRODUCED IN YEAST MUTANTS AND METHODS OF USE THEREOF

(57) Abstract: The invention provides various high mannose glycosylated polypeptides that are useful in a vaccine formulations. The invention also provides methods for making such glycosylated polypeptides and its uses in eliciting HIV-neutralizing antibodies.



WO 2007/035930 A3

GLYCOSYLATED POLYPEPTIDES PRODUCED IN YEAST MUTANTS AND METHODS OF USE THEREOF

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the priority benefit of provisional application U.S. Serial No. 60/719,952, filed September 22, 2005, the contents of which are incorporated by reference in their entirety.

TECHNICAL FIELD

[0002] The present invention relates to the fields of immunology and medicine. More particularly, the invention relates to methods of producing glycosylated antigens useful in eliciting effective immune responses and vaccination strategies using such antigens.

STATEMENT REGARDING FEDERALLY SPONSORED RESEARCH OR DEVELOPMENT

[0003] This invention was made with U.S. Government support under National Institutes of Health grant numbers AI51903 and AI58724. The U.S. Government may have certain rights in this invention.

BACKGROUND OF THE INVENTION

[0004] Complex carbohydrates expressed by some pathogens facilitate the evasion of host immune responses and contribute to the destructive sequelae following infection with organisms such as *Streptococcus pneumoniae*, *Neisseria meningitidis*, *Haemophilus influenza*, *Salmonella typhi*, and human immunodeficiency virus ("HIV"). In particular, carbohydrate rich regions of glycoproteins are poorly immunogenic. Factors contributing to this lack of immunogenicity include dilution of single antigenic response due to carbohydrate microheterogeneity and steric interference with highly immunogenic protein epitopes. See, e.g., Rudd, *et al.*, *Crit. Rev. Biochem. Mol. Biol.* 32:1-100 (1997); Woods, *et al.*, *Nature Struct. Biol.* 1:499-501 (1994). Additionally, viruses frequently fool the host immune system by using the cellular glycosylation machinery to express endogenous glycans for which the host is already tolerized.

[0005] The role of glycoproteins in HIV pathogenicity, particularly that of gp120, exemplifies the difficulty and importance of complex carbohydrates in eliciting protective host immune responses. HIV, like all viruses, requires the transcriptional and translation machinery of a cell in order to successfully propagate itself in the host. HIV accomplishes this by entering a cell through interactions with specific cellular receptors. Specifically, the HIV virion enters a cell, usually a T lymphocyte, through the interaction of its viral envelope protein ("Env") with the cellular CD4 receptor and the CCR5 cellular coreceptor during early infection, and through the CD4 receptor-CXCR4 coreceptor complex later in infection. *See, e.g., Pöhlmann, et al., J. Virol. 75:4664-72 (2001); Teunis, et al., Cell 100:587-97 (2000).* The Env protein is a glycoprotein known as gp120. On the surface of the HIV virion, three molecules of gp120 are noncovalently linked to another cell surface protein, gp41. *See, e.g., Zolla-Pazner, Nature Rev. Immunol. 4:199-210 (2004).* gp41 is a transmembrane glycoprotein found as a homotrimeric complex in the viral envelope. This gp120-gp41 complex forms hetero-oligomeric spikes on the HIV virion that first binds the CD4/coreceptor complex and then subsequently undergoes a conformational change resulting in the exposure of the viral fusion peptide that mediates the entry of the virus into the cell.

[0006] Recent evidence suggests that a second interaction distinct from the CD4/coreceptor interaction is also critical to HIV transmission and infection, particularly at the earliest stages. Dendritic cells (DCs), a highly specialized antigen presenting cell, is the first cell targeted by HIV upon infection. *See Geijtenbeek, et al., Cell 100:587-97 (2000).* DCs at the mucosa capture, internalize, and transport HIV from the mucosal surfaces to remote lymph nodes via an interaction between gp120 and DC-SIGN, a cell surface molecule on DCs. The delivery of intact virus by the DCs then results in the infection of CD4⁺ T-lymphocytes. *See Bashirova, et al., J. Exp. Med. 193:671-78 (2001).* In other words, DC-SIGN acts *in trans* to mediate efficient infection of CD4⁺ cells by HIV. *See Pöhlmann, et al., J. Virol. 75:4664-72 (2001).* Later studies showed that DC-SIGN also acts *in cis* to promote efficient viral infection. *See, e.g., Lee, et al., J. Virol. 75:12028-38 (2001).*

[0007] Given the difficulty in eliminating infected cells, an immune response is likely to be most effective if the elicited response impedes the entry of the virus into the cell. To date, attempts to elicit such protective responses have been hampered by the poor immunogenicity of gp120. gp120 contains extensive glycosylation and highly variable loops interspersed with more conserved, functionally constrained regions acting as physical

shields for critical gp120 epitopes, *i.e.*, those epitopes that interact with the receptor/coreceptor complex, from antibodies that can block viral entry or neutralize the virus. *See, e.g.*, Garber, *et al.*, *Lancet Infect. Dis.* 4:397-413 (2004). However, naturally elicited neutralizing antibodies have been identified, confirming the potential for effective neutralizing responses. *See, e.g.*, Burton, *et al.*, *Nature Immunol.* 5:233-36 (2004).

[0008] Development of HIV/AIDS vaccine to induce neutralizing antibodies against a broad spectrum of HIV-1 primary isolates is still a highly challenging endeavor, 25 years after the discovery of AIDS. The challenge for developing effective vaccines lies in the identification of appropriate antigenic epitopes that can be presented immunogenically such that neutralizing antibodies are elicited in the host. The challenge in developing successful HIV vaccines is further complicated by an expansive diversity in primary HIV isolates. *See, e.g.*, Gaschen, *et al.*, *Science* 296:2354-60 (2002). To date, traditional approaches to vaccine design have not proven successful in eliciting neutralizing antibody responses. Among the approximately 30 clinical trials of HIV vaccines, none are able to induce broadly neutralizing antibodies. One of the major challenges is the lack of an appropriate design of an antigen with neutralizing epitopes that are exposed on the surface of the antigen and highly conserved in most or all subtypes of HIV-1. Pantophlet R *et al.*, *Annu Rev Immunol.* 24:739-69, 2006. Up to date only four monoclonal antibodies (MAbs) with broad and potent neutralizing activity were isolated from HIV-1 infected humans. Douek DC *et al.*, *Cell* 124:677-81 (2006). None of them can be duplicated in all tested species of animals. Among the four MAbs, one targets a conformational epitope on the HIV-1 env gp120, two recognize gp41, and one, 2G12, binds to the high mannose-type carbohydrates on gp120.

[0009] The earliest target cell for HIV infection is the dendritic cell (DC), and therefore the most potent vaccine is one that disrupts the ability of HIV to target DCs in a host. The high mannose oligosaccharides of gp120 provide epitopes essential for HIV-DC interaction, and thus provide suitable vaccine targets. High mannose oligosaccharides mediate the interaction between DC-SIGN and gp120. *See* Geijtenbeek, *et al.*, *Cell* 100:587-97 (2000). Yet, the naturally occurring gp120 only expresses about 20% Man₈GlcNAc₂ (Man₈) and 10% Man₉GlcNAc₂ (Man₉). *See* Scalan, *et al.*, *J. Virol.* 76:7306-21 (2002). Cyanovirin-N (CV-N), a cyanobacterial protein, binds to high mannose oligosaccharides of gp120, specifically recognizing the Man α 1,2-Man structures on Man₉GlcNAc₂ (Man₉) and the D1D3 isomer of Man₈GlcNAc₂ (Man₈), and through this interaction acts as a potent microbicide against HIV. *See* Bewley, *et al.*, *J. Am. Chem. Soc.* 123:3982-902 (2001);

Sandstrom, *et al.*, *Biochem.* 43:13926-31 (2004). Furthermore, one of the naturally occurring, neutralizing antibodies specifically recognizes a cluster of Man α 1,2-Man high mannose oligosaccharides of gp120. See Scanlan, *et al.*, *J. Virol.* 76:7306-21 (2002). This antibody, known as the 2G12 antibody, potently neutralizes a broad range of HIV-1 primary isolates by inhibiting the HIV virion interaction with DCs and CD4⁺ T cells. See, e.g., Trkola, *et al.*, *J. Virol.* 70:1100-08 (1996); Sanders, *et al.*, *J. Virol.* 76:7293-305 (2002).

High mannose type glycans as a target for HIV-1 vaccine

[0010] Development of a carbohydrate-based HIV vaccine is considered to be one of the novel approaches for a prophylactic vaccine. Wang, *Curr Opin Drug Discov Devel.* 9(2):194-206, 2006. Several lines of evidence have demonstrated that the terminal Man α 1,2-Man structures (α 1,2-linked mannose) found on the D1 and D3 arm of Man₈NAcGlc₂ present novel targets on the gp120 glycoprotein, with the possibility of inducing potent, neutralizing antibodies against HIV-1 from different strains and subtypes.

[0011] High-mannose glycans on gp120 are recognized by the broadly neutralizing MAb 2G12. Among the hundreds of MAbs against gp120 that have been generated in rodents and isolated from HIV-1 infected humans, 2G12 is the only one that recognizes virus carbohydrates and potently neutralizes a broad range of HIV-1 primary isolates. It does so by inhibiting the interactions of HIV-1 with DCs and CD4⁺ T cells. Trkola *et al.*, *J Virol.* 70(2):1100-8, 1996; Scanlan *et al.*, *J Virol.* 76(14):7306-21, 2002; Sanders *et al.*, *J Virol.* 76(14):7293-305, 2002;. The binding site of 2G12 has been identified as high mannose-type glycans on the HIV-1 env gp120 glycoprotein. More specifically, the 2G12 MAb binds to a cluster of terminal α 1,2-linked mannose residues from at least three high-mannose glycans (Scanlan *et al.*, *J Virol.* 76(14):7306-21, 2002); it does not recognize other carbohydrates or mannose residues with different terminal linkages, α 1,3-linked or α 1,6-linked mannose.

[0012] The Man α 1,2-Man structures on high-mannose glycans are also the binding sites for a potent HIV-1 inhibitor. The cyanobacterial protein termed Cyanovirin-N (CV-N) can inactivate a diverse array of laboratory strains and primary isolates of HIV-1, HIV-2 and SIV. Boyd *et al.*, *Antimicrob Agents Chemother.* 41(7):1521-30, 1997. The protein can also block HIV-1 gp120 interaction with CD4 and coreceptors, prevent virus-to-cell fusion, and stop infection of cells. Esser *et al.*, *J Virol.* 73(5):4360-71, 1999. Dey *et al.*, *J Virol.* 74(10):4562-9 2000. These potent properties of CV-N are attributed to its ability to bind with extremely high affinity to the high-mannose oligosaccharides on gp120. Specifically,

the inhibitor recognizes the $\text{Man}\alpha 1,2\text{-Man}$ structures on $\text{Man}_9\text{GlcNAc}_2$ (Man9) and the D1D3 isomer of $\text{Man}_8\text{GlcNAc}_2$ (Man8), but not other forms of high mannoses, including Man7, Man6, and Man5. Bewley et al., *J Am Chem Soc.* 123(17):3892-902.2001; Sandstrom et al., *Biochemistry.* 43(44):13926-13931 2004. This ability of CV-N to inhibit HIV infection presents evidence of the potency of such molecules that are able to bind to these terminal glycan structures.

[0013] Dendritic cells have been shown to enhance infection through the interaction of DC-SIGN with the high-mannose glycans on gp120. Recently, DCs were found to be the first cell type targeted by HIV in the body. DCs at the mucosa are found to capture, internalize and transport HIV to remote lymph nodes where they deliver the intact virus to CD4^+ T-lymphocytes. Geijtenbeek et al., *Cell.* 100:587-97, 2000. It was found that all tested strains of HIV-1, HIV-2, SIV and SHIV bind to DCs, with DC-SIGN playing an important role in this process. Pohlmann, et al., *J Virol.* 75(10):4664-72, 2001. The interaction between HIV and DC-SIGN is mediated by the high-mannose glycans on gp120. Geijtenbeek et al., *Cell.* 100:587-97, 2000. In fact, synthetic high mannose oligosaccharides are able to bind DC-SIGN and prevent subsequent HIV interactions. Feinberg et al., *Science.* 294(5549):2163-6, 2001.

[0014] The gp120 protein of HIV-1 is heavily glycosylated and contains an average of 25 N-linked glycosylation sites. Approximately half of them are occupied by high mannose-type or hybrid-type glycans (Leonard et al., *J Biol Chem.* 265(18):10373-82, 1990), with the high mannose glycans interacting with 2G12, CV-N, and DC-SIGN through different binding sites. MAb 2G12 binds a cluster of D1 arms from at least three Man9 or Man8 residues, CV-N binds with a high affinity to the D1 and D3 arms from a single Man9 and Man8 residues, and DC-SIGN binds several mannoses residues through its tetramer.

[0015] Altogether, these results indicate the strong possibility of inhibiting an early stage of HIV infection with highly specific neutralization antibodies against the $\text{Man}\alpha 1,2\text{-Man}$ structures found on high mannose glycans. The major challenge is to develop an antigen containing strictly high mannoses with terminal $\alpha 1,2$ -linked mannose structures, and eliciting an immunogenic response to this epitope that can cross-react to gp120.

[0016] Two approaches have taken to construct homogenous HIV-1 glycopeptides for establishing glycopeptide-based HIV vaccines. Wang, *Curr. Opin. Drug Discov. Devel.* 9(2):194-206, 2006. These approaches are total chemical synthesis of HIV-1 gp120 glycopeptides carrying either a hybrid-type or a high-mannose-type N-glycan, and

chemoenzymatic approach to construct various HIV-1 glycopeptides. See Mandal et al., *Angew. Chem. Int. Ed.* 43:2557-2561, 2004; and Geng et al., *Angew. Chem. Int. Ed.* 43:2562-2565, 2004; Singh et al., *Bioorg. Med. Chem. Lett.* 13:327-330, 2003; Wang et al., *ChemBioChem.* 6:1068-1074 (2005); Zeng et al., *J. Am. Chem. Soc.* 127:9692-9693, 2005; Li et al., *J. Org. Chem.* 70:9990-9996, 2005. However, these synthetic glycopeptides need to be further evaluated for their immunogenicity in animal models.

SUMMARY OF THE INVENTION

[0017] An effective HIV vaccine capable of eliciting neutralizing antibodies to a broad spectrum of viral isolates is the best hope for controlling the HIV pandemic. Unfortunately, major obstacles continue to hinder the development of such a vaccine, including poor immunogenicity of the Env glycoprotein, viral antigen diversity and immune escape. In response to these challenges, the compositions provided herein offer a distinct recombinant antigen equipped with multiple complex carbohydrate epitopes critical for HIV entry into host cells. The increased number of carbohydrate epitopes maximizes the immunogenicity of these epitopes and its relative potency in a vaccine composition to elicit neutralizing antibodies.

[0018] For example, the gp120 provided herein expresses essentially uniform mannose oligosaccharides with a terminal α 1-2 glycan, *e.g.*, mannose-9-*N*-acetylglucosamine-2 (“Man₉GlcNAc₂”) or Man₈GlcNAc. The methods of making the high mannose gp120 compositions and using the resulting compositions to elicit effective anti-HIV neutralizing antibody responses are also described herein.

[0019] Therefore, provided herein is an essentially uniformly glycosylated recombinant protein, wherein the terminal glycan is a terminal α 1-2 glycan structure. The recombinant protein can be obtained by the transformation of *Saccharomyces cerevisiae* variants with a nucleotide sequence that encodes the protein, fermentation of the transformed cells, and isolation of the protein from the cells or culture supernatant. In some embodiments, the protein is a virulence factor. The protein can be a viral protein. In a specific embodiment, the viral protein is a surface protein. In one embodiment, the protein is gp120 of the human immunodeficiency virus. The terminal glycan can be mannose. In some embodiments, the glycan is oligomannose. In some embodiments, the terminal α 1-2 structure is on Man₉GlcNAc₂ or Man₈GlcNAc₂, or a combination.

[0020] Further provided herein is a method of preparing a uniformly glycosylated recombinant protein, wherein the terminal glycan is a terminal α 1-2 glycan structure, comprising: a) providing a vector comprising a nucleotide sequence encoding the protein; b) transforming a cell with a defect in protein glycosylation; c) fermenting the transformed cells; and d) purifying the secreted recombinant protein from the cell supernatant. In some embodiments, the nucleotide sequence encodes a protein that is glycosylated in its native state. The nucleotide sequence can encode a protein that is a virulence factor. In a specific embodiment, the nucleotide sequence encodes the gp120 protein. In some embodiments, the cell expresses the gp120 protein with largely $\text{Man}_9\text{GlcNAc}_2$ or $\text{Man}_8\text{GlcNAc}_2$. The cell of the instant method can be a yeast cell such as *S. cerevisiae mns1* Δ or *S. cerevisiae pmr1* Δ . The vector can be YEpl and, in some embodiments, may comprise the GAL1 promoter. In some embodiments, the glycoprotein is purified using a His tag.

[0021] Also provided herein is a method of preparing an antibody which specifically binds a protein comprising a terminal α 1-2 glycan structure, the method comprising immunizing an animal with an effective amount of a composition comprising a recombinant protein made by the method or the protein as disclosed herein. The composition can further comprise an adjuvant, a carrier, or both.

[0022] Provided herein is the isolated antibody generated by the method disclosed herein as well as the hybridoma that produces the isolated antibody.

[0023] Also provided herein is a composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising an isolated glycosylated polypeptide comprising at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the glycosylated polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are $\text{Man}_9\text{GlcNAc}_2$, $\text{Man}_8\text{GlcNAc}_2$, or a combination thereof.

[0024] Further provided herein is a composition for eliciting antibodies specific for a protein having an epitope recognized by antibody 2G12 in a subject, comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted *pmr1* gene of *Saccharomyces cerevisiae* or disrupted homolog gene of *pmr1* gene of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12, and a pharmaceutically acceptable excipient.

[0025] Further provided herein is a composition for eliciting antibodies specific for a protein having an epitope recognized by antibody 2G12 in a subject, comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted *pmr1* and *mn1* genes of *Saccharomyces cerevisiae* or disrupted homolog genes of *pmr1* and *mn1* genes of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12, and a pharmaceutically acceptable excipient.

[0026] Further provided herein is a composition for eliciting antibodies specific for a protein having an epitope recognized by antibody 2G12 in a subject, comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted *och1* and *mn1* genes, or *och1*, *mn1*, and *mn4* genes of *Saccharomyces cerevisiae* or disrupted homolog genes of *och1* and *mn1* genes, or *och1*, *mn1*, and *mn4* genes of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12, and a pharmaceutically acceptable excipient.

[0027] Further provided herein is a method of preparing a composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, said method comprising: a) fermenting a mutant fungus, wherein the mutant fungus is mutated to produce a glycosylated polypeptide, wherein the glycosylated polypeptide comprises at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the glycosylated polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are $\text{Man}_9\text{GlcNAc}_2$, $\text{Man}_8\text{GlcNAc}_2$, or a combination thereof; and b) purifying the glycosylated polypeptide from the mutant fungus cell lysate or supernatant. In some embodiments, the method further comprises a step of formulating the glycosylated polypeptide with a pharmaceutically acceptable excipient. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae* having disrupted *och1*, *mn1*, and *mn4* genes, *Saccharomyces cerevisiae* having disrupted *och1* and *mn1* genes, *Saccharomyces cerevisiae* having disrupted *pmr1* and *mn1* genes, or *Pichia pastoris* or *Candida albicans* having a disrupted *och1* gene.

[0028] Further provided herein is a method of preparing a composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, said method comprising: a) fermenting a mutant fungus, wherein the mutant fungus has disrupted homolog gene of *pmr1* gene of *Saccharomyces cerevisiae*; and b) isolating a glycosylated polypeptide recognized by antibody 2G12 from the mutant fungus cell. The

method may further comprises a step of formulating the isolated glycosylated polypeptide with a pharmaceutically acceptably excipient. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae* or *Pichia pastoris*, wherein the *pmr1* gene is disrupted. In some embodiments, the mutant fungus has been transformed with a vector comprising a nucleotide sequence encoding the glycosylated polypeptide before step a). In some embodiments, the glycosylated polypeptide is HIV gp120 or a fragment thereof.

[0029] Further provided herein is a method of preparing a composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, said method comprising: a) fermenting a mutant fungus, wherein the mutant fungus has disrupted homolog genes of *pmr1* and *mn1* genes of *Saccharomyces cerevisiae*; and b) isolating a glycosylated polypeptide recognized by antibody 2G12 from the mutant fungus cell. The method may further comprises a step of formulating the isolated glycosylated polypeptide with a pharmaceutically acceptably excipient. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae*, wherein the *pmr1* and *mn1* genes are disrupted. In some embodiments, the mutant fungus has been transformed with a vector comprising a nucleotide sequence encoding the glycosylated polypeptide before step a). In some embodiments, the glycosylated polypeptide is HIV gp120 or a fragment thereof.

[0030] Further provided herein is a method of preparing a composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, said method comprising: a) fermenting a mutant fungus, wherein the mutant fungus has disrupted homolog genes of *och1* and *mn1* genes, or *och1*, *mn1*, and *mn4* genes of *Saccharomyces cerevisiae*; and b) isolating a glycosylated polypeptide recognized by antibody 2G12 from the mutant fungus cell. The method may further comprises a step of formulating the glycosylated polypeptide with a pharmaceutically acceptably excipient. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae* having disrupted *och1*, *mn1*, and *mn4* genes, or *Saccharomyces cerevisiae* having disrupted *och1* and *mn1* genes. In some embodiments, the mutant fungus has been transformed with a vector comprising a nucleotide sequence encoding the glycosylated polypeptide before step a). In some embodiments, the glycosylated polypeptide is HIV gp120 or a fragment thereof.

[0031] Further provided herein is a mutant fungus cell comprising a polynucleotide comprising a nucleotide sequence encoding gp120 or a fragment thereof comprising at least two N-glycosylation site, wherein the mutant yeast cell is capable of producing the gp120 or the fragment comprising at least two N-linked high mannose oligosaccharides that are

recognized by antibody 2G12, wherein the high-mannose oligosaccharides are $\text{Man}_9\text{GlcNAc}_2$, $\text{Man}_8\text{GlcNAc}_2$, or a combination thereof. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae* having disrupted och1, mnn1, and mnn4 genes, *Saccharomyces cerevisiae* having disrupted och1 and mnn1 genes, or *Pichia pastoris* or *Candida albicans* having a disrupted och1 gene. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae* having disrupted pmr1 and mnn1 genes. In some embodiments, the mutant fungus is *Saccharomyces cerevisiae* or *Pichia pastoris* having disrupted pmr1 gene.

[0032] Further provided herein is a composition comprising a whole cell of a mutant yeast and a pharmaceutically acceptable excipient, wherein the mutant yeast has disrupted pmr1 and mnn1 genes, och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*, or disrupted homolog genes of pmr1 and mnn1 genes, och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*.

[0033] Further provided herein is a method of generating antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising administering to the subject an effective amount of any of the compositions described herein.

[0034] Further provided herein is a method of inducing neutralizing antibodies against a pathogen (such as HIV) in a subject, comprising administering to the subject an effective amount of any of the compositions described herein.

[0035] Further provided herein is a method of inducing neutralizing antibodies against a pathogen in a subject in need thereof, comprising administering an effective amount of a pharmaceutical composition comprising the protein provided herein or a protein prepared by the method provided herein, and a suitable excipient. In some embodiments, the method further comprises administering an adjuvant. In a specific embodiment, the pathogen is HIV.

[0036] Further provided herein is a method of preventing or treating pathogen-induced disease in a subject, comprising administering to the subject an effective amount of any of the compositions described herein.

[0037] Further provided herein is a method of preventing or treating pathogen-induced disease in a subject in need thereof, comprising administering an effective amount of a pharmaceutical composition comprising the protein provided herein or a protein prepared by the method provided herein, and a suitable excipient. In some embodiments, the method

further comprises administering an adjuvant. In a specific embodiment, the pathogen is HIV. In one embodiment, the protein is gp120. In some embodiments, the pharmaceutical composition further comprises multiple gp120 proteins from different HIV strains.

[0038] Further provided is a method for identifying an antibody that specifically binds to N-linked high mannose oligosaccharide, comprising detecting binding of an antibody to a glycosylated polypeptide comprising at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the glycosylated polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.

[0039] Further provided is a method for identifying an antibody that specifically binds to N-linked high mannose oligosaccharide, comprising detecting binding of an antibody to a glycoprotein produced in a mutant fungus having disrupted pmr1 gene of *Saccharomyces cerevisiae* or disrupted homolog gene of pmr1 gene of *Saccharomyces cerevisiae*, wherein the glycoprotein is recognized by antibody 2G12.

[0040] Further provided is a method for identifying an antibody that specifically binds to N-linked high mannose oligosaccharide, comprising detecting binding of an antibody to a glycosylated polypeptide produced in a mutant fungus having disrupted pmr1 and mnn1 genes of *Saccharomyces cerevisiae* or disrupted homolog genes of pmr1 and mnn1 genes of *Saccharomyces cerevisiae*, wherein the glycoprotein is recognized by antibody 2G12.

[0041] Further provided is a method for identifying an antibody that specifically binds to N-linked high mannose oligosaccharide, comprising detecting binding of an antibody to a glycosylated polypeptide produced in a mutant fungus having disrupted och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae* or disrupted homolog genes of och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*, wherein the glycoprotein is recognized by antibody 2G12.

[0042] In some embodiments, the antibody is in a serum of a subject infected by HIV. In some embodiments, the glycosylated polypeptide is in a mutant yeast cell lysate.

[0043] Further provided is a method of identifying a neutralizing monoclonal antibody for HIV transmission, comprising: a) contacting a first cell, wherein the first cell expresses dendritic cell-specific C-type lectin (DC-SIGN), with a second cell expressing CD4+ and CCR5+, b) coculturing the first and second cells with infectious viral particles; c) contacting coculture with a candidate antibody prepared by the method of claim 20; and d) determining

relative infectivity, whereby the neutralizing antibody is one that reduces or eliminates relative infectivity of the second cell by infectious viral particles.

BRIEF DESCRIPTION OF THE DRAWINGS

[0044] FIGURE 1 illustrates the structures and processing of high mannoses in mammalian cells. The left box shows the processes of glycan from precursor to Man8 in the ER. Then, proteins with mainly Man8 are transported to Golgi for further processing (top-right box, an example in mammalian cells). When protein transport from the ER to Golgi is blocked, such as by a mutation of *Pmr1* gene in yeast, these glycans will not be further processed resulting in a glycoprotein that contains only Man8 and Man9 forms of glycans (bottom-right).

[0045] FIGURE 2 shows the expression of HIV-1 gp120 glycoproteins in yeast. **A.** Expression of gp120 proteins from 3 HIV-1 strains in $\Delta pmr1$ was induced by galactose for the indicated days. The proteins in the culture supernatants were detected with 2G12. **B.** Purification of HIV_{YU2} gp120. Lane 1 shows the concentrated yeast culture media, lane 2 after dialysis and centrifugation, lanes 3 and 4 after GNL and gel filtration at 200 and 400 ng respectively, and lanes 5 and 6 HIV_{YU2} gp120 produced in 293T cells at 200 and 400 ng.

[0046] FIGURE 3 shows EndoH digestion assay of gp120 protein produced in yeast. The same lot of purified gp120 proteins used for MS/MS analysis was digested with EndoH. Then the non-digested and digested proteins were separated by SDS-PAGE followed by Coomassie blue staining. Lane 2 is the non-digested at 200 ng, lane 3 digested at 400 ng, and lane 4 is a non-glycosylated gp120 from HIV_{SF2} expressed in yeast.

[0047] FIGURE 4 shows the antibodies against mannoses. (A) Two rabbits were immunized with 200 μ g of zymosan A containing CFA at week 0, and boosted with 100 μ g of antigen containing incomplete Freund's adjuvant ("IFA") at weeks 2 and 4 and thereafter every 4 weeks. Immune sera were collected at week 5 and 1 week after each booster. Antibodies were tested in duplicate using ELISA in microwell plates coated with zymosan and yeast mannan. The mean antibody titers from two rabbits were presented. (B) Antibodies against mannoses are reactive to some of tested gp120 glycoproteins. Microwell plates were coated with 5 μ g/ml of gp120 proteins. (C) This panel shows the comparison of the antigenicity of gp120 proteins from YU2 strain in mammalian cells and in yeast cells tested by anti-mannan antibody.

[0048] **FIGURE 5** shows the analysis of BSA-Man₉ conjugate. The BSA-Man₉ conjugate was analyzed with 10% SDS-PAGE gel followed by Coomassie blue staining. Lane 1 is non-conjugated BSA, lane 2 BSA-SMCC, and lane 3 BSA-SMCC-Man₉.

[0049] **FIGURE 6A** shows the expression and purification of ECD recombinant protein of DC-SIGN. Expression and purification of DC-SIGN ECD recombinant protein. The ECD protein was expressed in *E. coli* at 25°C in the absence (lane 2) and presence of IPTG (lane 3) for 4 h. Inclusion body was isolated (lane 4) and refolded proteins from insoluble fraction was purified using Ni-NTA columns (lanes 5). Proteins were separated onto 12% SDS-PAGE gel and stained with Coomassie blue. 6B shows the ELISA using DC-SIGN and anti-gp120 polyclonal antibody coated microwell plates. HIV-1 gp120 of YU2 strain was used to bind to the coating proteins.

[0050] **FIGURE 7** shows the development of DC-SIGN MAbs. The purified DC-SIGN ECD protein was used to immunize Balb/c mice. The splenocytes from a mouse with good antibody response were used for cell fusion. After ELISA screening of the culture supernatants, the cells from positive wells were expanded and subcloned. The supernatants from wells with single clones were screened using Western blots. Each separate lane represents an individual clone.

[0051] **FIGURE 8** shows a neutralization assay. Neutralizing activities of two polyclonal (P) antibodies were tested using a LuSIV cell line infected by pseudovirus bearing pHXB2 gp160. Three well-known neutralizing MAbs (M) were used as positive controls. Four doses (50, 25, 12.5 and 6.25 µg/ml) of each antibody were examined.

[0052] **FIGURE 9** shows structure and processing of N-linked oligosaccharides. Figure 9A shows the processes of glycan from Man₉GlcNAc₂ to Man₈GlcNAc₂ by MNS1 in the ER. Then, proteins with mainly Man₈GlcNAc₂ form of high mannose core are transported to Golgi for further processing. Figure 9B lower box shows the superglycosylation, which is the major (thick arrow) pathway. Figure 9B right upper box shows the addition of α1,3-Man on the core, which is the minor pathway (thin arrow). When the genes of Och1, Mnn1 and Mnn4 are all deleted, no sugars can be added on the Man₈GlcNAc₂ core resulting in an almost homogenous Man₈GlcNAc₂ type of glycans (Figure 9C).

[0053] **FIGURE 10** shows screening 2G12 cross-reactive protein in yeast mutant strains using Western Blot. A. Yeast culture media from various mutant strains were screened with HIV-1 MAb 2G12 (Polymun Scientific Inc. Forschung GmbH).. Secreted proteins in the culture media were precipitated with ethetone and loaded onto a 4-20% gradient SDS-PAGE

gel. The blot was probed with 2G12 followed by incubation with horseradish peroxidase (HRP) labeled anti-human IgG. Signals were developed with ECL and exposed to X-ray film. B. Cell lysates were screened from the same set of yeast mutant strains with control of HIV-1 gp120 glycoprotein.

[0054] FIGURE 11 shows induction of 2G12 cross-reactive protein in $\Delta pmr1$ yeast cells. A. gp120 transformed $\Delta pmr1$ yeast cells were cultured at the indicated time points. Secreted proteins in the culture media were precipitated with ethetone and loaded onto a 4-20% gradient SDS-PAGE gel. The blot was probed with 2G12 followed by incubation with HRP-anti-human IgG. Signals were developed with ECL and exposed to X-ray film. B. Proteins in the culture media from non-transformed (lanes 1-3) and gp120 transformed (lanes 4-6) $\Delta pmr1$ cells were detected with 2G12 as described in A.

[0055] FIGURE 12 shows purification and identification of the 2G12 reactive glycoprotein. The approximately 100 kDa protein (designated Yp100) was partially purified using GNL lectin affinity chromatography. The culture media were incubated with agarose bound GNL for 16 hrs. The samples of starting material (lane 1), flow through (lane 2), washes (lanes 3-6) and elutes (lanes 7-11) were loaded onto 4-20% SDS-PAGE gel. The proteins on transferred membrane were probed with MAb 2G12.

[0056] FIGURE 13 shows identification of the 2G12 reactive glycoprotein. The proteins in the 100 kDa band were subjected to in-gel digestion. Digested peptides were extracted and analyzed using Nano-LC-MS/MS. Analysis was done on a Micromass Q-ToF hybrid quadrupole/time-of-flight mass spectrometer with a nanoelectrospray source. Raw files were processed with ProteinLynx software (Waters) and submitted to a MASCOT search. Totally 6 peptides are identified in the yeast protein PST1 (SEQ ID NO:1) in *Saccharomyces cerevisiae* with Accession NP_010340 and 45,749 Da of mass. The bold letters at the N-terminus is the identified signaling peptide. The Bold letters at the C-terminus is the identified GPI anchor signal. Single underlines indicate the identified peptides by mass spectrometry. Highlighted letters are the potential N-linked glycosylation sites.

[0057] FIGURE 14 shows confirmation of the 2G12 cross-reactive protein by 2D gel. A. Partially purified Yp100 protein was separated by a 2D gel and subjected to Commassie blue staining (A), western blots with 2G12 MAb (B) and anti-PST1 polyclonal antibody raised against a synthetic peptide in rabbits. Although the partially purified Yp100 proteins

have contamination by other glycoproteins from yeast but none of them has similar PI as PST1 does.

[0058] FIGURE 15 shows glycan analysis of 2G12 cross-reactive protein PST1. A. Partially purified Yp100 protein was digested with Endo H and the non-digested (lanes 2, 4 and 6) and digested (lanes 3, 5 and 7) proteins were separated by SDS-PAGE gel and subjected to Commassie blue staining (lanes 1-3), Western blots with anti-PST1 polyclonal antibody (lanes 4 and 5), and with MAb 2G12 (lanes 6 and 7). This is followed by incubation with HRP labeled secondary antibody, ECL, and exposed to X-ray films.

[0059] FIGURE 16 shows cross-reactivity of PST1 to 2G12 in *pmr1* and *mnn1* double mutant. Cell lysates from 5 different double mutant stains were prepared and separated on 4-20% gradient SDS-PAGE. The blot was probed with MAb 2G12 followed by incubation with HRP labeled secondary antibody, ECL, and exposed to X-ray films.

[0060] FIGURE 17 shows up-regulation and migration shifts of PST1 in double mutant. A. PST1 in yeast cell lysates. Cell lysates from different stains were separated on 4-20% gradient SDS-PAGE. The levels of PST1 expression in wild type and mutants were analyzed using Western blot with anti-PST1 polyclonal antibody. Lane 1, wild type; lane 2, $\Delta mnn1$; lane 3, $\Delta och1$; lane 4, $\Delta pmr1$; lane 5, $\Delta pmr1$ and $\Delta mnn1$ double mutant. Upper panel shows PST1 was only detected in the lysates of *pmr1* and *mnn1* double mutant. Lower panel shows another glycosylated protein GP38 that is detectable in all 5 strains. B. PST1 in yeast cell culture media. Yeast cells from different strains were cultured in rich media for 24 h and the proteins in media were precipitated with acetone. The levels of protein secretion were detected using Western blot with anti-PST1 antibody (upper panel) and anti-GP38 (lower panel).

[0061] FIGURE 18 shows analysis of $\alpha 1,3$ -Man and $\alpha 1,2$ -Man on PST1 in double mutant. Yeast cell culture media from *pmr1* single mutant (lane 1) and *pmr1* + *mnn1* double mutant (lane 2) were detected for terminal $\alpha 1,3$ -Man and $\alpha 1,2$ -Man by Western blot using anti- $\alpha 1,3$ -linked mannose polyclonal antibody and 2G12. *Och1* single mutant and purified PST1 were used as controls. Proteins in media were precipitated with acetone and loaded onto a 4-20% SDS-PAGE gradient gel. After blotting to nitrocellulose, the membranes were probed with 2G12 MAb, and rabbit anti- $\alpha 1,3$ Man antibody or anti-PST1 antibody as indicated, followed by goat anti-human IgG-HRP and goat anti-rabbit IgG-HRP, respectively.

[0062] FIGURE 19 shows confirmation of the 2G12 cross-reactive protein in double mutant by immunoblot. Culture supernatant from double mutant was centrifuged and the precipitate was discarded. The supernatant (lane 1) was pre-cleared with agarose (lane 2). Then the sample was incubated with MAbs 2G12 for 16 h at 4°C. The immunocomplex was incubated with protein A-agarose for 1 h at room temperature. The sample was centrifuged briefly and the supernatant was used as flow through (lane 3). The protein A-agarose bound immunocomplex was washed twice (lanes 4 and 5). The bound proteins were eluted with SDS sample buffer (lane 6). Then samples were separated on 4-20% gradient gel, and probed with anti-PST1 polyclonal antibody.

[0063] FIGURE 20 shows PST1 homolog gene among 18 species. The protein sequence of PST1 (NP_010340) and ECM33 (NP_009634) were used to search homolog genes from HomoloGene database (release 50.1) of National Center for Biotechnology Information (NCBI). HomoloGene is a system for automated detection of homologs among the annotated genes of several completely sequenced eukaryotic genomes. Currently HomoloGene database contains 165,820 HomoloGene groups from 18 species. The HomoloGene of interest is obtained by entering protein Reference Sequence (RefSeq) number at NCBI home page (<http://www.ncbi.nlm.nih.gov/>). Multiple alignment of the HomoloGene was performed by clustalw program. PST1 and 3 homolog genes (Proteins of *S. pombe* and *E. gossypii*) were obtained from HomoloGene database searching. Protein from *Candida glabrata* was obtained using blastp searching against NCBI nr sequence database).

[0064] FIGURE 21 shows analysis of double and triple mutants using PCR. Screening and verification of the genotype for triple mutant haploids. **Panel A** shows a 1.5% agarose gel with the PCR screening of putative triple mutants for the $\Delta mnn4$ genotype (top panel) using the MNN4A and KanB primers (expected size was 609bp), and the $\Delta och1$ genotype (bottom panel) using the OCH1A and KanB primers (697bp). Lane 1 contains haploid clone 2-2. Lane 2 contains haploid clone 2-3. Lane 3 contains haploid clone 2-4. Lane 4 contains haploid clone 2-5. **Panel B** shows a 1.5% agarose gel with PCR verification of the $\Delta mnn1$, $\Delta mnn4$ and $\Delta och1$ genotype for the triple mutant haploid clones 2-5 and 2-6 (top and bottom panels, respectively). Each lane contains a different pair of PCR primers, as indicated. The combination of an internal ORF-specific primer (MNN1B, MNN4B, OCH1B, MNN1C, MNN4C and OCH1C) and an upstream or downstream primer (MNN1A, MNN4A, OCH1A, MNN1D, MNN4D and OCH1D) result in no PCR product if the ORF is deleted.

Lane 1 is MNN1A + MNN1B. Lane 2 is MNN4A + MNN4B. Lane 3 is OCH1A + OCH1B. Lane 4 is MNN1C + MNN1D. Lane 5 is MNN4C + MNN4D. Lane 6 is OCH1C + OCH1D. By contrast, the combination of an upstream or downstream primer and an internal KanMX specific primer (KanB and KanC) result in a PCR product if a deletion is present. Lane 7 is MNN1A + KanB (expected size of 591bp). Lane 8 is MNN1D + KanC (944bp). Lane 9 is MNN4D + KanC (931bp). Lane 10 is OCH1D + KanC (854bp).

[0065] **FIGURE 22** shows whole-Cell ELISA. ELISA was used to test the specificity of 2G12 and α 1,3-linked mannose antibodies against whole-cell yeast. INVSc1 (diploid) and Δ mn1 Δ mn4 Δ och1-DIP yeast were diluted to 5.0×10^7 cells/ml in PBS and used to coat ELISA plates. SF162 gp120 and wild-type yeast mannan were diluted to 5 μ g/ml in 50mM carbonate buffer. 2G12 and anti- α 1,3-mannose were used at 10 μ g/ml for capture with a three-fold serial dilution in blocking buffer. **Panel A** shows that the anti- α 1,3 antibody has a high affinity towards the wild-type cells and mannan, with the triple mutant showing low affinity and SF162 showing no binding. By contrast, **Panel B** shows that the triple mutant cells and SF162 gp120 respectively have moderate and high affinities towards the 2G12 antibody, which is specific to terminal α 1,2-linked mannose residues. Both the wild-type cells and mannan show no binding to 2G12. This experiment was done in duplicate, and the above OD₄₅₀ reading represent the average.

[0066] **FIGURE 23** shows loss of α 1,3-Man reactivity in triple mutant cells. Immunofluorescence of triple mutant and wild-type yeast using anti- α 1,3-linked mannose immune sera and anti-Zymosan antibody. Log-phase Δ mn1 Δ mn4 Δ och1-DIP cells (Panel A, C) and INVScI cells (Panel B, D) were fixed with 4% paraformaldehyde and transferred to polylysine-coated slides. The cells were incubated with 0.5% SDS and stained with α 1,3-linked mannose antisera at a 1:500 dilution (Panel A, B) and purified anti-Zymosan at 10 μ g/ml (Panel C, D), followed by Alexa Fluor 568-conjugated goat anti-rabbit IgG.

[0067] **FIGURE 24** shows exposure of terminal α 1,2-Man in triple mutant cells. Immunofluorescence of Triple Mutant and wild-type yeast using 2G12 and anti-Zymosan. Log-phase Δ mn1 Δ mn4 Δ och1-DIP cells (Panel A, B and C) and INVScI cells (Panel D, E and F) were fixed with 4% paraformaldehyde and transferred to polylysine-coated slides. The cells were incubated with 0.5% SDS and stained with 2G12 at 100 μ g/ml (Panel A, D) and rabbit anti-Zymosan at 10 μ g/ml (Panel B, E), followed Alexa Fluor 568-conjugated goat anti-human IgG and Alexa Fluor 488-conjugated goat anti-rabbit IgG, respectively. Panel C and F show the co-localization of the antibodies.

[0068] **FIGURE 25** shows analyses of α 1,2-Man and α 1,3-Man on the glycoproteins of yeast mutants. Western blot using 2G12 and α 1,3-linked mannose antisera on all yeast glycosylation mutants. Log phase yeast cells were lysed in RIPA buffer and loaded onto a 4-20% SDS-PAGE gradient gel at 2.4 μ g/lane. After blotting to nitrocellulose, the membranes are probed with 2G12 MAb or α 1,3Man antisera, followed by goat anti-human IgG-HRP and goat anti-rabbit IgG-HRP, respectively. Lane 1 contains INVSc1 wild-type lysate. Lane 2 contains Δ mnn1 lysate. Lane 3 is Δ och1 lysate. Lane 4 contains Δ mnn4 lysate. Lane 5 contains Δ mnn1 Δ mnn4 lysate. Lane 6 contains Δ mnn1 Δ och1 lysate. Lane 7 contains Δ mnn1 Δ mnn4 Δ och1 lysate. There are at least four proteins that bind strongly to 2G12 found in the cell lysate of Δ mnn1 Δ och1 and Δ mnn1 Δ och1 Δ mnn4 (Left Panel). By contrast, only WT yeast and mutant without the Δ mnn1 genotype show binding to antisera specific to α 1,3-linked mannose (Right Panel).

[0069] **FIGURE 26** shows 2G12 reactivity in all isolated triple mutants. Western blot using 2G12 on six yeast triple mutant clones. Log phase yeast triple mutant clones were crudely lysed in SDS loading buffer and loaded onto a 4-20% SDS-PAGE gradient gel. After blotting to nitrocellulose, the membrane was probed with 2G12 at 1 μ g/ml followed by goat anti-human IgG-HRP. Lane 1 contains the haploid clone 1-2. Lane 2 contains the haploid clone 1-3. Lane 3 contains the haploid clone 2-2. Lane 4 contains the haploid clone 2-4. Lane 5 contains the haploid clone 2-5. Lane 6 contains the haploid clone 2-6. There are at least four proteins that bind strongly to 2G12 found in the crude cell lysate of each triple mutant haploid clone.

[0070] **FIGURE 27** shows homogenous of Man₈ form of high mannose in yeast triple mutant cells. MALDI-TOF Mass Spectrometry of Triple Mutant Yeast Lysate. Log-phase Δ mnn1 Δ mnn4 Δ och1-DIP cells were homogenized, with lipid extraction, and the resulting lysate was digested with PNGase F. N-glycan profiling was conducted by Maldi-TOF Mass Spectrometry after permethylation. The major peak, which represents over 90% of the total glycans is Man₈GlcNAc₂, while there are two minor peak representative of Man₅GlcNAc₂ and Man₉GlcNAc₂.

[0071] **FIGURE 28** shows partial purification and separation of 2G12-reactive proteins from the cell lysate of triple mutant yeast. In **Panel A**, the subcellular localization of 2G12-reactive proteins was estimated using differential centrifugation. Log-phase Δ mnn1 Δ mnn4 Δ och1 cells were lysed in sucrose lysis using glass beads. After removal of unlysed cells and large aggregates, the proteins were separated into two fractions, the

22,000g supernatant (Lane 1) and the 22,000g pellet (Lane 2). From this fraction, the pellet was resolubilized in 1.0% Triton X-100 and separated into two additional fractions, the Triton-soluble pellet (Lane 3) and the Triton-insoluble pellet (Lane 4). All lanes represent a Western blot using 2G12 at 1 μ g/ml. In **Panel B**, the 2G12-reactive proteins were partially purified using ConA-agarose. The Triton-soluble fraction (Lane 1) from Δ mnn1 Δ mnn4 Δ och1 cells were incubated with ConA-agarose beads and the flowthrough was collected (Lane 2). After washing with binding buffer (Lane 3), proteins were eluted with either 0.5M methyl manno-pyranoside (Lane 4, 5), or 2mM EDTA + 1.0% SDS (Lane 6). All lanes represent a Western blot using 2G12 at 1 μ g/ml. In **Panel C**, the 2G12-reactive proteins were separated using 2D electrophoresis. The elute fractions from the ConA purification (Panel B, Lane 4-6) were run onto Invitrogen IPG ZOOM 4-7 IPG strips followed by size separation on the IPG ZOOM system using 4-12%. One gel was stained by Western blotting using 2G12 at 1 μ g/ml. Two large spots that showed a positive signal by Western blot were excised from the Sypro Ruby gel, digested with Trypsin, and analyzed by nano LC/MS/MS on a Micromass Q-ToF 2 for peptide identification. Three glycoproteins (ECM33, Gas1, and Gas5) in the upper spot and a single glycoprotein (YJL171c) in the lower spot were identified using MS/MS.

[0072] **FIGURE 29** shows expression and partial purification of 2G12-reactive glycoproteins in the culture media of triple mutant yeast. In **Panel A**, 100 μ l of culture supernatant from log phase INVSc1 and Δ mnn1 Δ mnn4 Δ och1 yeast were subjected to acetone precipitation by incubating with two-times the volume of ice-cold acetone for 30min at -80°C. The resulting pellets were resuspended in 10 μ l of SDS sample buffer and loaded onto 4-20% SDS-PAGE gradient gels. After transfer to nitrocellulose, proteins were probed with 2G12 at 2 μ g/ml. The triple mutant yeast has at least four proteins with strong reactivity to the 2G12 MAbs (Lane 2), while the wild-type yeast supernatant shows no signal (Lane 1). In **Panel B**, the 2G12-reactive proteins were partially purified using ConA-agarose. The culture supernatant (Lane 1) from log-phase Δ mnn1 Δ mnn4 Δ och1 cells was directly incubated with ConA-agarose beads. The flowthrough (Lane 2) was collected and the beads were washed with binding buffer (Lane 3). The bound 2G12-reactive proteins (lane 4) were eluted by boiling the ConA beads with SDS-PAGE loading buffer. The proteins were detected by Western blot using 2G12 at 1 μ g/ml. Two glycoproteins (ECM33 and GP38) were identified using MS/MS from an aliquot of 2G12 precipitates shown on lane 4.

[0073] **FIGURE 30** shows immunoprecipitation of 2G12-reactive proteins. The cell lysate and culture media of triple mutant cells were immunoprecipitated with 2G12 and blotted with antibodies against the identified 2G12-reactive proteins. Cell lysate (Panel A) and culture media (Panel B) from $\Delta mnn1\Delta mnn4\Delta och1$ cells (Lane 1) were pre-cleared by incubation with Protein A-Sepharose 4B. 2G12 was added to the samples, incubated for 1 hour, followed by Protein A-Sepharose. The unbound proteins were removed as the supernatant (Lane 2) and washed with PBS (Lane 3). Bound proteins were eluted by boiling with 40 μ l of 1xSDS sample buffer (Lane 4). All samples were loaded onto 4-20% gradient SDS-PAGE gels and blotted for Western analysis using anti-ECM33, anti-gp38, anti-YJL171C, anti-Gas1, and anti-PST1.

[0074] **FIGURE 31** shows migration shift of glycoproteins in the mutant of different genes. Yeast cells were grown to log phase and the cells and culture media were collected. Cells were lysed in RIPA buffer and loaded onto a 4-20% SDS-PAGE gradient gel at 2.4 μ g/lane. The culture media was precipitated with acetone and proteins were also loaded onto a 4-20% SDS-PAGE gradient gel. Lane 1 is WT. Lane 2 is $\Delta mnn1$. Lane 3 is $\Delta och1$. Lane 4 is $\Delta mnn4$. Lane 5 is $\Delta mnn1mnn4$. Lane 6 is $\Delta mnn1\Delta och1$. Lane 7 is $\Delta mnn1\Delta mnn4\Delta och1$. After blotting to nitrocellulose, the membranes are probed with anti-ECM33 (upper panels) or anti-GP38 (lower panels), followed by goat anti-rabbit IgG-HRP.

[0075] **FIGURE 32** shows the analysis of antibodies against α 1,2-linked mannoses raised in rabbits immunized with triple mutant yeast cells. Rabbits were immunized with whole cells from wild type (WT) and triple mutant (TM) of *S cerevisiae*. Microwell plates were coated with 5 μ g/ml of HIV-1 gp120 from JRFL strain produced in 293T cells. The immune sera were diluted to 1:500 and the antibodies against α 1,2-Man were detected with 2G12.

[0076] **FIGURE 33** shows analyses of potential N-linked and O-linked glycosylation sites of PST1. **Panel A** shows the amino acid sequence of precursor (SEQ ID NO:2). Bold letters at the N-terminus is the identified signaling peptide. The Bold letters at the C-terminus is the identified GPI anchor signal. Single underlines indicate the identified peptides by mass spectrometry, while double underlines indicate the peptides were identified twice in the same or different mass spectrometry analyses. Highlighted letters are the potential N-linked glycosylation sites. **Panel B** shows the analysis of O-linked glycosylation sites using software NetOGlyc 3.1 developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark in a public Website

<http://www.cbs.dtu.dk/services/netoglyc>. Julenius K., A. Mølgaard, R. Gupta and S. Brunak. Prediction, conservation analysis and structural characterization of mammalian mucin-type O-glycosylation sites. *Glycobiology*, 15:153-164, 2005.

[0077] **FIGURE 34** shows analyses of potential N-linked and O-linked glycosylation sites of ECM33. **Panel A** shows the amino acid sequence of precursor (SEQ ID NO:3). Bold letters at the N-terminus is the identified signaling peptide. The Bold letters at the C-terminus is the identified GPI anchor signal. Single underlines indicate the identified peptides by mass spectrometry, while double underlines indicate the peptides were identified twice in the same or different mass spectrometry analyses. Highlighted letters are the potential N-linked glycosylation sites. **Panel B** shows the analysis of O-linked glycosylation sites using software NetOGlyc 3.1 developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark in a public Website <http://www.cbs.dtu.dk/services/netoglyc>.

[0078] **FIGURE 35** shows analyses of potential N-linked and O-linked glycosylation sites of GP38. **Panel A** shows the amino acid sequence of precursor (SEQ ID NO:4). Bold letters at the N-terminus is the identified signaling peptide. Single underlines indicate the identified peptides by mass spectrometry, while double underlines indicate the peptides were identified twice in the same or different mass spectrometry analyses. Highlighted letters are the potential N-linked glycosylation sites. **Panel B** shows the analysis of O-linked glycosylation sites using software NetOGlyc 3.1 developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark in a public Website <http://www.cbs.dtu.dk/services/netoglyc>.

[0079] **FIGURE 36** shows analyses of potential N-linked and O-linked glycosylation sites of YJL171c. **Panel A** shows the amino acid sequence of precursor (SEQ ID NO:5). Bold letters at the N-terminus is the identified signaling peptide. The Bold letters at the C-terminus is the identified GPI anchor signal. Single underlines indicate the identified peptides by mass spectrometry, while double underlines indicate the peptides were identified twice in the same or different mass spectrometry analyses. Highlighted letters are the potential N-linked glycosylation sites. **Panel B** shows the analysis of O-linked glycosylation sites using software NetOGlyc 3.1 developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark in a public Website <http://www.cbs.dtu.dk/services/netoglyc>.

[0080] **FIGURE 37** shows analyses of potential N-linked and O-linked glycosylation sites of Gas1. **Panel A** shows the amino acid sequence of precursor (SEQ ID NO:6). Bold letters at the N-terminus is the identified signaling peptide. Bold letters at the C-terminus is the identified GPI anchor signal. Single underlines indicate the identified peptides by mass spectrometry, while double underlines indicate the peptides were identified twice in the same or different mass spectrometry analyses. Highlighted letters are the potential N-linked glycosylation sites. **Panel B** shows the analysis of O-linked glycosylation sites using software NetOGlyc 3.1 developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark in a public Website <http://www.cbs.dtu.dk/services/netoglyc>.

[0081] **FIGURE 38** shows analyses of potential N-linked and O-linked glycosylation sites of Gas5. **Panel A** shows the amino acid sequence of precursor (SEQ ID NO:7). Bold letters at the N-terminus is the identified signaling peptide. Bold letters at the C-terminus is the identified GPI anchor signal. Single underlines indicate the identified peptides by mass spectrometry, while double underlines indicate the peptides were identified twice in the same or different mass spectrometry analyses. Highlighted letters are the potential N-linked glycosylation sites. **Panel B** shows the analysis of O-linked glycosylation sites using software NetOGlyc 3.1 developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark in a public Website <http://www.cbs.dtu.dk/services/netoglyc>.

[0082] **FIGURE 39** shows cloning and expression of gp120 glycoproteins in triple mutant yeast. **Panel A** shows colony PCR results on yeast transformants run onto a 1.0% agarose gel. Four separate yeast clones of $\Delta mnn1 \Delta mnn4 \Delta och1$, transformed with YU2-gp120 or JRFL-gp120, were picked from -ura/glucose plates and incubated with PCR master mix. The PCR mixture contained the primers MF α 1-Kpn-5 and JRFL-Xba-3 for pJRFL-gp120 verification and MF α 1-Kpn-5 and YU2-Xba-3 for the pYU2-gp120. The expected PCR product sizes were 1724kB and 1697kB for JRFL and YU2 transformants, respectively. **Panel B** shows the detection of gp120 using anti-gp120-IIIB (Virostat Inc.). One $\Delta mnn1 \Delta mnn4 \Delta och1$ transformant of each plasmid (JRFL-gp120 and YU2-gp120) was induced for gp120 expression by growth in -ura/galactose media. A sample of the culture media was collected at four different time points: 0, 3, 6, and 9 days. Proteins were precipitated with acetone, run onto a 4-20% SDS-PAGE gel, and detected using anti-gp120-IIIB. **Panel C** shows the detection of gp120 using anti-gp120-YU2. The same

$\Delta mnn1 \Delta mnn4 \Delta och1$ transformants were induced for the indicated days. Proteins were precipitated with acetone, run onto a 4-20% SDS-PAGE gel, and detected using anti-gp120-YU2. The resulting proteins detected by both anti-gp120 show a clear band at ~110kDa, with optimal expression at 6 days to 9 days. .

[0083] **FIGURE 40** shows detection of expressed gp120 with 2G12. One $\Delta mnn1 \Delta mnn4 \Delta och1$ transformant of each plasmid (JRFL-gp120 and YU2-gp120) was induced for gp120 expression by growth in -ura/galactose media, and one clone of non-transformed $\Delta mnn1 \Delta mnn4 \Delta och1$ was grown in SC/galactose. A sample of the culture media was collected at 3, 6, and 9 days, as indicated. Proteins were precipitated with acetone, run onto a 4-20% SDS-PAGE gel, and detected using 2G12. These gp120 glycoproteins have a predicted molecular weight of 96.4 and 99.6 kDa for JR-FL and YU-2 gp120, respectively.

MODES OF CARRYING OUT THE INVENTION

[0084] The ability of CV-N and the 2G12 to inhibit HIV infection suggests that the elicitation of a neutralizing antibody response to Man9, Man8, or both would provide effective immunity against HIV transmission. However, to date there is no reproducible manner in which to provide the gp120 antigen with a specific glycosylation profile. Thus, the recombinant high mannose gp120 composition provided herein provides a potent inducer of protective immunity against HIV infection.

[0085] For clarity of disclosure, and not by way of limitation, the detailed description of the invention is divided into the subsections that follow.

A. Definitions

[0086] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of ordinary skill in the art to which this invention belongs. All patents, published patent applications and other publications and sequences from GenBank and other databases referred to herein are incorporated by reference in their entirety. If a definition set forth in this section is contrary to or otherwise inconsistent with a definition set forth in patents, published patent applications and other publications and sequences from GenBank and other data bases that are herein incorporated by reference, the definition set forth in this section prevails over the definition that is incorporated herein by reference.

[0087] Other features and advantages of the invention will be apparent from the detailed description, and from the claims.

[0088] As used herein, "a" or "an" means "at least one" or "one or more."

[0089] As used herein, the term "adjuvant" refers to a substance which, when added to an immunogenic agent, nonspecifically enhances or potentiates an immune response to the agent in the recipient host upon exposure to the mixture. Adjuvants can include, for example, oil-in-water emulsions, water-in-oil emulsions, alum (*e.g.*, aluminum hydroxide/phosphate), liposomes and microparticles. Exemplary adjuvants include, but are not limited to squalene mixtures (SAF-I), muramyl peptide, saponin derivatives, mycobacterium cell wall preparations, monophosphoryl lipid A, mycolic acid derivatives, nonionic block copolymer surfactants, Quil A, cholera toxin B subunit, polyphosphazene and derivatives, immunostimulating complexes (ISCOMs), and cytokines. *See, e.g.*, Vogel, *et al.*, *Immunological Adjuvants* IN VACCINES 69-79 (Plotkin, *et al.*, eds., 4th Ed. Saunders (2004)). For veterinary use and for production of antibodies in animals, mitogenic components of Freund's adjuvant (both complete and incomplete) also can be used.

[0090] As used herein, "AIDS" refers to the symptomatic phase of HIV infection, and includes both Acquired Immune Deficiency Syndrome (commonly known as AIDS) and AIDS-Related Complex ("ARC"). *See, e.g.*, Kilby, *et al.*, *Natural History of HIV-1 Disease* IN TEXTBOOK OF AIDS MEDICINE 49-58 (Merrigan, *et al.*, eds., Williams & Wilkins 2nd Ed. 1999). The immunological and clinical manifestations of AIDS are well known in the art and include, for example, opportunistic infections and cancers resulting from immune deficiency.

[0091] As used herein, the term "antibody" refers to any form of a peptide or polypeptide derived from, modeled after or substantially encoded by, an immunoglobulin gene or immunoglobulin genes, or fragments thereof, capable of specifically binding an antigen or epitope. *See, e.g.*, FUNDAMENTAL IMMUNOLOGY (W. E. Paul, ed., 5th Ed., Lippincott, Williams & Wilkins (2003)); CURRENT PROTOCOLS IN IMMUNOLOGY (Coligan, *et al.*, eds., John Wiley & Sons, most recent edition); ANTIBODY ENGINEERING: A PRACTICAL APPROACH (J. McCafferty, *et al.*, eds., Oxford University Press 1996). Examples of antibody fragments are those that retain antigen-binding and include Fab, Fab', F(ab')₂, Fd, and Fv fragments; diabodies; linear antibodies; single-chain antibody molecules, *e.g.*, scFv and V_{HH}; minibodies; nanobodies; and multispecific antibodies formed from antibody fragments. Typically, a binding fragment or derivative retains at least 50% of its binding activity. Preferably, a binding fragment or derivative retains at least 60%, 70%, 80%, 90%, 95%, 99% or 100% of its biological activity. The antibody or antigen-binding fragment

thereof can include conservative amino acid substitutions that do not substantially alter its binding and/or biologic activity. Thus, the term "antibody" is used in the broadest sense and specifically covers monoclonal (including full length monoclonal antibodies), polyclonal, multispecific (*e.g.*, bispecific), heteroconjugate, chimeric, humanized, human, murine, and synthetic antibodies as well as antibody fragments that specifically bind the desired antigen and exhibit the desired binding and/or biological activity.

[0092] The term "antigen" refers to any molecule that is specifically recognized and bound by an antibody. An antigen which elicits an immune response in an organism, as evidenced by production of specific antibodies within the organism is termed an "immunogen." The specific portion of the antigen or immunogen which is bound by the antibody is termed the "binding epitope" or "epitope." An antibody is specific for a particular antigen when it selectively binds the antigen, thereby distinguishing it from other antigens. Preferably, the antibody lacks significant binding to unrelated antigens.

[0093] The term "expression cassette" as used herein refers to a nucleotide sequence which is capable of affecting expression of a structural gene (*i.e.*, a protein coding sequence, such as the antibody provided herein) in a host compatible with such sequences. Expression cassettes include at least a promoter operably linked with the polypeptide coding sequence; and, optionally, with other sequences, *e.g.*, transcription termination signals. As used herein, the term "promoter" includes all sequences capable of driving transcription of a coding sequence in a cell, *e.g.*, a mammalian or plant cell. Thus, promoters used in the constructs of the invention include cis-acting transcriptional control elements and regulatory sequences that are involved in regulating or modulating the timing and/or rate of transcription of a gene. For example, a promoter can be a cis-acting transcriptional control element, including an enhancer, a promoter, a transcription terminator, an origin of replication, a chromosomal integration sequence, 5' and 3' untranslated regions, or an intronic sequence, which are involved in transcriptional regulation. Promoters can be constitutive or inducible. Additional factors necessary or helpful in effecting expression also may be used, *e.g.*, enhancers. A promoter is operably linked to a coding sequence, such as a nucleic acid of the invention, if it stimulates or modulates the transcription of the coding sequence in an appropriate host cell or other expression system. Generally, promoter transcriptional regulatory sequences that are operably linked to a transcribed sequence are physically contiguous to the transcribed sequence, *i.e.*, they are cis-acting. However, some transcriptional regulatory sequences, *e.g.*, enhancers, need not be physically contiguous or

located in close proximity to the coding sequences whose transcription they enhance. Expression cassettes also include plasmids, expression vectors, recombinant viruses, any form of recombinant "naked DNA" vector, and the like.

[0094] The term "isolated" includes a material removed from its original environment, *e.g.*, the natural environment if it is naturally occurring. For example, an isolated polynucleotide or polypeptide is one that is separated from some or all of the coexisting materials in the natural system. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides could be part of a composition, and still be isolated in that such vector or composition is not part of its natural environment.

[0095] The phrases "nucleic acid" or "nucleic acid sequence" includes oligonucleotide, nucleotide, polynucleotide, or to a fragment of any of these, to DNA or RNA (*e.g.*, mRNA, rRNA, tRNA) of genomic or synthetic origin which may be single-stranded or double-stranded which encodes an antibody of the present invention, or a biologically active fragment thereof. The term encompasses nucleic acids, *i.e.*, oligonucleotides, containing known analogues of natural nucleotides. The term also encompasses nucleic-acid-like structures with synthetic backbones. *See e.g.*, Mata, *Toxicol. Appl. Pharmacol.* 144:189-97 (1997); Strauss-Soukup, *Biochemistry* 36:8692-98 (1997); and Samstag, *Antisense Nucleic Acid Drug Dev* 6:153-56 (1996).

[0096] "Recombinant" polypeptides or proteins refer to polypeptides or proteins produced by recombinant DNA techniques; *e.g.*, produced from cells transformed by an exogenous DNA construct encoding the desired polypeptide or protein. *See, e.g.*, CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (F.M. Ausubel, *et al.*, eds., John Wiley & Sons, most current edition).

[0097] The terms "polypeptide", "oligopeptide", "peptide" and "protein" are used interchangeably herein to refer to polymers of amino acids of any length. The polymer may be linear or branched, it may comprise modified amino acids, and it may be interrupted by non-amino acids. The terms also encompass an amino acid polymer that has been modified naturally or by intervention; for example, disulfide bond formation, glycosylation, lipidation, acetylation, phosphorylation, or any other manipulation or modification, such as conjugation with a labeling component. Also included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), as well as other modifications known in the art. It is

understood that, because the polypeptides of this invention are based upon an antibody, the polypeptides can occur as single chains or associated chains.

[0098] The term "homolog" is used to refer a gene or a product of a gene from one species that has a common origin and functions the same as a gene from another species. For example, a fungus homolog gene of *pmr1* gene of *S. cerevisiae* refers to a fungus gene that has a common origin and encodes a protein that has the same function as the *pmr1* gene of *S. cerevisiae*.

[0099] A "vector" comprises a nucleic acid which can infect, transfect, transiently or permanently transduce a cell. It will be recognized that a vector can be a naked nucleic acid, or a nucleic acid complexed with protein or lipid. The vector optionally comprises viral or bacterial nucleic acids and/or proteins, and/or membranes (*e.g.*, a cell membrane, a viral lipid envelope, etc.).

[0100] As used herein, the term "synergize" refers to the ability of one agent to increase the anti-pathogenic or neutralizing effect of a second agent. Synergistic activity, thus, includes but is not limited to an increased biological effect (*e.g.*, more potent or longer lasting) using the two agents together that is not observed when the agents are used separately, a more effective biological effect, *e.g.*, elimination of multiple types of toxicity not achievable with the administration of a single agent, or a reduction in the amount of agents necessary for administration to achieve the biological effect observed with a single agent.

[0101] The term "pathogen" refers to any organism that induces or elicits a undesired symptom or disease state. A pathogen may be a bacteria, virus, or fungus.

[0102] As employed herein, the term "subject" embraces human as well as other animal species, such as, for example, canine, feline, bovine, porcine, rodent, and the like. It will be understood by the skilled practitioner that the subject having a pathogen or disease targeted by the antibody of the invention.

[0103] As used herein, the term "ameliorating, treating or preventing" include a postponement of one or more symptoms associated with the infection or other disorder, a reduction in the severity of such symptoms that will or are expected to develop, or a complete elimination of such symptoms. These terms further include ameliorating existing pathogen-related symptoms, reducing duration of disease, preventing additional symptoms, ameliorating or preventing serious sequelae, preventing or reversing mortality, and reducing or preventing pathogen transmission. Thus, the terms denote that a beneficial

result has been conferred on a subject with a pathogen, or with the potential of exposure to such a pathogen. In particular, ameliorating, treating or preventing include, *e.g.*, preventing initial infection of an individual exposed to HIV; reducing viral burden in an individual infected with HIV; prolonging the asymptomatic phase of HIV infection; increasing overall health or quality of life in an individual with AIDS; and prolonging life expectancy of an individual with AIDS. A clinician can compare the effect of immunization with the patient's condition prior to treatment, or with the expected condition of an untreated patient, to determine whether the treatment is effective in inhibiting AIDS.

[0104] As used herein, an "effective amount" is an amount sufficient to inhibit or prevent, partially or totally, transmission of the pathogen, tissue damage or other symptoms associated with the action of the virulence factor within or on the body of the subject or to prevent or reduce the further progression of such symptoms. When applied to an individual active ingredient administered alone, an effective dose refers to the dose required for that ingredient alone. When applied to a combination of active agents, an effective dose refers to the combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously. The term "therapeutically effective amount" and "effective amount" are employed interchangeably.

[0105] As used herein, the term "bioactive agent" refers to any synthetic or naturally occurring compound that enhances or mediates a desired biological effect to reduce or eliminate the infectivity or pathogenicity of a particular organism. Bioactive agents include, for example, a pharmaceutical agent, such as a chemotherapeutic drug, a microbicidal drug, a antiviral drug, a toxin, a cytokine, a ligand, an antibody, or some combination thereof.

[0106] As used herein, "prevents the productive interaction" means that the amount of interaction is reduced as compared to the amount that would occur without the antibody or biologically active fragment thereof. The interactions may be reduced or prevented by any means including, but not limited to masking or altering interactive regions on the virulence factor, and altering the expression, aggregation, conformation, or association state of the virulence factor.

[0107] As used herein, the term "treat" refers to the application or administration of a therapeutic agent to a subject, or application or administration of a therapeutic agent to an isolated tissue or cell line from a subject, who has a disease or disorder, a symptom of disease or disorder or a predisposition toward a disease or disorder, with the purpose of

curing, healing, alleviating, relieving, altering, remedying, ameliorating, improving or affecting the disease or disorder.

B. Compositions

[0108] Provided herein is a uniformly glycosylated recombinant protein, wherein the terminal sugar is a terminal α 1-2-Mannose structure. The recombinant protein can be obtained by the transformation of *Saccharomyces cerevisiae* variant with a nucleotide sequence that encodes the protein, fermentation of the transformed cells, and isolation of the protein from the cells or culture supernatant. In some embodiments, the protein is a virulence factor. The protein can be a viral protein. In a specific embodiment, the viral protein is a surface protein. In one embodiment, the protein is gp120 of the human immunodeficiency virus. In some embodiments, the glycan is oligomannose. In one embodiment, the terminal α 1-2-Mannose structures are on Man₉GlcNAc₂ or Man₈GlcNAc₂.

[0109] Any suitable virulence factor may be employed in the composition disclosed herein. A virulence factor is any glycoprotein that mediates or participates in the ability of a microorganism to cause disease in a host under suitable conditions. For example, a virulence factor can be an adherence factor, a coat protein, an invasion factor, a capsule, an exotoxin, or an endotoxin. Exemplary organisms expressing glycoprotein virulence factors include, but are not limited to lentiviruses (e.g., HIV), ebola virus, gram negative bacteria (e.g., *Campylobacter jejuni*, *Campylobacter coli*, *Neisseria meningitidis*, *Helicobacter pylori*, *Escherichia coli*, *Haemophilus influenzae*, *Pseudomonas aeruginosa*, *Mycobacteria tuberculosis*, *Mycobacteria bovis*, *Borrelia burgdorferi*, *Chlamydia* spp., and *Streptococcus parasanguis*).

[0110] The glycosylation of the protein virulence factor can be N-linked or O-linked. A glycan is a polysaccharide consisting of more than about five monosaccharide residues joined to one another by glycosidic linkages. Representative glycans include mannose. Essentially uniform glycosylation of the protein virulence factor refers to greater than 70%, 80%, 90% or 100% glycosylation with a single glycoform.

[0111] In one aspect, the composition provided herein is a gp120 essentially uniformly glycosylated with Man₉GlcNAc₂, Man₈GlcNAc₂, or some combination thereof, having a terminal α 1-2-Man structure. As used herein, the term “essentially uniformly” refers to greater than 70% of the glycans on gp120 are Man₉GlcNAc₂, Man₈GlcNAc₂, or some

combination thereof. In a specific embodiment, the glycosylated gp120 composition provided herein has greater than 90% Man₉GlcNAc₂ or Man₈GlcNAc₂, preferably 100% Man₉GlcNAc₂ or Man₈GlcNAc₂. The term "high mannose" refers to a glycan of 5 to 9 mannose residues without other kinds of terminal sugars. See, e.g., Figure 1. Mammalian cell surface and serum glycoproteins rarely contain terminal mannose residues. See, e.g., Weis, *et al.*, *Immunol Rev.* 163:19-34 (1998). Mammalian glycoproteins contain mostly complex-type oligosaccharides with few Man₅ or Man₆, while HIV-1 gp120 proteins contain a larger number of Man₇₋₉, indicating that the high mannoses on HIV-1 virus and human glycoproteins are significantly different in quantity and quality. See, e.g., Scanlan, *et al.*, *J Virol.* 76:7306-21 (2002); Cutalo, *et al.*, *J Am Soc Mass Spectrom.* 15:1545-55 (2000).

[0112] Also provided herein is a composition comprising an isolated glycosylated polypeptide comprising at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the glycosylated polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.

[0113] Further provided herein is a composition comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted pmr1 gene of *Saccharomyces cerevisiae* or disrupted homolog gene of pmr1 gene of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12.

[0114] Further provided herein is a composition comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted pmr1 and mnn1 genes of *Saccharomyces cerevisiae* or disrupted homolog genes of pmr1 and mnn1 genes of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12.

[0115] Further provided herein is a composition comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted any combination of och1, mnn1, and/or mnn4 genes of *Saccharomyces cerevisiae* or disrupted any combination of the homolog genes of och1, mnn1, and/or mnn4 genes of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12.

[0116] The glycosylated polypeptide may comprise at least three, at least four, at least five, at least six, at least seven, at least eight, at least nine, or at least ten N-linked high-mannose oligosaccharides. The glycosylated polypeptide may be a fungus (such as a yeast)

or a non-fungus glycoprotein or a fragment thereof. For example, the glycosylated polypeptide is PST1, ECM33, Gas1, Gas5, GP38, or YJL171c of *S. cerevisiae*, or a homolog of any of these proteins. The glycosylated polypeptide may be a virulence factor. In some embodiments, the glycosylated polypeptide is a viral protein or a fragment thereof. In some embodiments, the glycosylated polypeptide is HIV gp120 or a fragment thereof.

[0117] In some embodiments, at least 50%, at least 60%, at least 70%, at least 80%, at least 90 percent, or at least 95% of the N-linked glycans on the glycopeptide are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.

[0118] The compositions of the present invention exclude any of the glycosylated polypeptides described in Wang, *Curr. Opin. Drug Discov. Devel.* 9(2):194-206, 2006; Mandal et al., *Angew. Chem. Int. Ed.* 43:2557-2561, 2004; Geng et al., *Angew. Chem. Int. Ed.* 43:2562-2565, 2004; Singh et al., *Bioorg. Med. Chem. Lett.* 13:327-330, 2003; Wang et al., *ChemBioChem.* 6:1068-1074 (2005); Zeng et al., *J. Am. Chem. Soc.* 127:9692-9693, 2005; and Li et al., *J. Org. Chem.* 70:9990-9996, 2005.

[0119] Further provided herein is a composition comprising a whole cell of a mutant yeast and a pharmaceutically acceptable excipient, wherein the mutant yeast has disrupted pmr1 and mnn1 genes, och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*, or disrupted homolog genes of pmr1 and mnn1 genes, och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*. The yeast whole cell may be alive or killed. The composition may be used for therapeutic or preventive vaccine (e.g., vaccine for HIV).

[0120] The composition described here comprises a pharmaceutically acceptable excipient. Any pharmaceutically acceptable carriers, excipients, or stabilizers (*Remington: The Science and Practice of Pharmacy* 20th Ed. (2000) Lippincott Williams and Wilkins, Ed. K. E. Hoover.), in the form of lyophilized formulations or aqueous solutions may be used. Acceptable carriers, excipients, or stabilizers are nontoxic to recipients at the dosages and concentrations, and may comprise buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone;

amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrans; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or sorbitol; salt-forming counter-ions such as sodium; metal complexes (e.g. Zn-protein complexes); and/or non-ionic surfactants such as TWEEN™, PLURONICS™ or polyethylene glycol (PEG). In some embodiments, the pharmaceutically acceptable excipient is an adjuvant. The compositions comprising adjuvant may be used vaccines.

[0121] Further provided herein is a method of preparing a uniformly glycosylated recombinant protein, wherein the terminal glycan is a terminal α 1-2-Man structure, comprising: a) providing a vector comprising a nucleotide sequence encoding the protein; b) transforming a cell with a defect in protein glycosylation; c) fermenting the transformed cells; and d) purifying the secreted recombinant protein from the cell supernatant. In some embodiments, the nucleotide sequence encodes a protein that is glycosylated in its native state. The nucleotide sequence can encode a protein that is a virulence factor. In some embodiments, the mutated cell expresses glycoproteins with largely Man₉GlcNAc₂ or Man₈GlcNAc₂. In a specific embodiment, the nucleotide sequence encodes the gp120 protein. In some embodiments, the cell expresses the gp120 protein with largely Man₉GlcNAc₂ or Man₈GlcNAc₂. The cell of the instant method can be a yeast cell such as *S. cerevisiae mns1* Δ or *S. cerevisiae pmr1* Δ . The vector can be YEpl and, in some embodiments, may comprise the GAL1 promoter. In some embodiments, the glycoprotein is purified using a His tag.

[0122] Further provided herein is a method for preparing any of the compositions described herein by fermenting a mutant fungus, wherein the mutant fungus is mutated to produce the glycosylated polypeptide; isolating or purifying the glycosylated polypeptide; and combining the polypeptide with a pharmaceutically acceptable excipient to form the composition. The compositions comprising the whole yeast cells may be prepared by fermenting a mutant yeast; preparing whole yeast cell with a pharmaceutically acceptable excipient to form the composition.

[0123] Any suitable cell may be employed. Preferably, the cell has a defect that permits truncates or otherwise modulates the typical intracellular processing of glycans. In some embodiments, the cell is a fungus cell (e.g., a yeast or a mold cell). In a specific embodiment, the cell is a yeast cell. See, e.g., U.S. Patent Nos. 5,919,651; 5,705,616. The outer chain glycosylation of secreted yeast proteins is a high (or long) mannose type

oligosaccharide chain. Thus, yeast production provides heterologous proteins with this yeast-specific outer chain glycosylation of the high mannose type, which is sometimes denoted “hyperglycosylation.” Briefly, the O-glycosidic carbohydrate structures of yeast proteins consist of an unbranched mannose chain of 1-5 mannose residues. O-glycosylation begins in the ER (transfer of the first mannose residue) and is completed in the Golgi apparatus. N-glycosylation begins with a core unit of N-acetylglucosamine, mannose and glucose being built up on a lipid carrier intermediate that is subsequently transferred to asparagine (Asn) residues of proteins in the endoplasmic reticulum. The protein-bound core unit then is cleavage at specific glucose and mannose residues followed by elongation of the polysaccharide in the Golgi apparatus to result in “outer chain” glycosylation.

[0124] A number of yeast strains can be employed to produce the composition provided herein. The cells of *Saccharomyces cerevisiae* are typically used, but those of *Pichia pastoris* and *Schizosaccharomyces pombe* are also commercially available for production of heterologous proteins. See, e.g., Tuite, *et al.*, *Expressing Cloned Genes in the Yeasts Saccharomyces cerevisiae and Pichia pastoris* IN PROTEIN EXPRESSION: A PRACTICAL APPROACH 61-100 (Higgins, *et al.*, eds., Oxford University Press 1999); Hitzeman, *et al.*, *Methods Enzymol.* 185:421-440 (1990); Barr, *et al.*, *Vaccine* 5:90-101 (1987); Liu, *et al.*, *Clin. Diagn. Lab. Immunol.* 5:592-94 (1998). Yeast strains with defects in Mns1p or Pmr1p are useful. Mns1p is an α 1,2 mannosidase that localizes in the ER where it trims Man9 to a single isomer of Man8 by removing the central arm of α 1,2-linked mannose. Therefore, the *S. cerevisiae* mns1p mutant strain *Mns1* Δ produces proteins glycosylated only with Man9. The protein Pmr1p is a calcium-dependent ATPase important for the transport of Ca^{++} from the ER to the Golgi. See Herscovics, *Biochim. Biophys. Acta* 1426:275-85 (1999); Camirand, *et al.*, *J. Biol. Chem.* 266:15120-27 (1991). The *S. cerevisiae* pmr1p mutant strain *pmr1* Δ secretes glycoproteins consisting solely of Man8, trimmed from Man9 by mannosidase I in the ER. See, e.g., Harmsen, *et al.*, *Gene* 125:115-23 (1993). However, any yeast strain with a suitable defect in N-glycosylation may be employed to produce the composition provided herein. See, e.g., U.S. Patent Nos: 5,705,616, 5,798,226.

[0125] Yeast strains with disrupted pmr1 gene, disrupted pmr1 and mnn1 genes, disrupted och1 genes, disrupted och1 genes in combination with one or more of mnn1, mns1, and mnn4 genes, disrupted och1 and mnn1 genes, or disrupted och1, mnn1, and

mn4 genes of *Saccharomyces cerevisiae*, *Candida albicans* and *Pichia pastoris*, or other yeast strains with disrupted homolog gene(s) may also be used to produce the glycosylated polypeptide or the whole yeast cell compositions.

[0126] Methods of generating mutant fungus cells are known in the art. The Examples of the invention provide details of generating several mutant yeast cells. See, e.g., Chiba et al., J. Biol. Chem. 273:26298-26304, 1998; Rudolph et al., Cell 58:133-145, 1989; Nakanishi-Shindo et al., J. Biol. Chem. 268:26338-26345, 1993.

[0127] Any suitable vector may be employed to transfect the mutant cells to produce recombinant glycosylated polypeptides. Yeast cells typically employ plasmid-based vectors. The vectors can be autonomous, multicopy plasmids (e.g., YEp or YRp), autonomous, single copy plasmids (e.g., YCp), or integrative (usually single copy) plasmids (e.g., YIp). See, e.g., *Yeast Vectors* (Unit 13.4) IN CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (Ausebel, et al., eds. John Wiley & Sons, most recent edition). In one embodiment, the vector is YEpL or pYES2-CT. Any promoter known in the art to result in heterologous protein production in the transformed cell may be employed. For yeast cells, natural and engineered promoters are available. Such promoters include, but are not limited to *PGK*, *GAP*, *TEF1*, *GAL1*, *ADH2*, *PHO5*, *CUP1*, *MF α 1*, *TRP1*, *PAL*, *GAP/GAL*, *GAP/ADH2*, *CYC1/GRE*, and *PGK/ARE*. In one embodiment, the promoter is *GAL1*. See, e.g., US Patent No. 5,139,936. The vector may also include additional sequences such as an origin of replication and various tag sequences useful for purification of the heterologous virulence factor, e.g., gp120.

[0128] Cells are transformed with the virulence factor-encoding vector and fermented using well known methods in the art. For yeast, cells may be transformed using lithium acetate, spheroplast transformation, or electroporation followed by standard fermentation. See, e.g., *Yeast* (Unit 13) IN CURRENT PROTOCOLS IN MOLECULAR BIOLOGY (Ausebel, et al., eds. John Wiley & Sons, most recent edition).

[0129] Exemplary virulence factors include, but are not limited to gp120, ebola virus envelope protein, or other glycosylated viral envelope proteins or viral proteins. See, e.g., U.S. Patent No. 6,630,144; 6,713,069.

[0130] In one aspect, any suitable gp120 nucleotide sequence can be employed in the composition provided herein. The gp120 molecule consists of a polypeptide core of 60 kD extensively modified by N-linked glycosylation thereby increasing its apparent molecular weight to 120 kD. See, e.g., Zolla-Pazner, et al., *Nature Rev. Immunol.* 4:199-209 (2004).

The amino acid sequence of gp120 contains five relatively conserved domains interspersed with five hypervariable domains. The positions of the 18 cysteine residues in the gp120 primary sequence, and the positions of 13 of the approximately 24 N-linked glycosylation sites in the gp120 sequence are common to all gp120 sequences. *See* Modrow, *et al.*, *J. Virol.* 61:570-78 (1987). The hypervariable domains contain extensive amino acid substitutions, insertions and deletions. Sequence variations in these domains result in up to 30% overall sequence variability between gp120 molecules from the various viral isolates. *See, e.g.*, Gaschen, *et al.*, *Science* 296:2354-60 (2002). Nonetheless, gp120 sequences maintain the virus's ability to bind to the viral receptor CD4 and DC-SIGN. Exemplary sequences include, but are not limited to those disclosed in Ratner, *et al.*, *Nature* 313:227 (1985); Muesing, *et al.*, *Nature* 313:450-58 (1985); McCutchan, *et al.*, *AIDS Res. Human Retroviruses* 8:1887-95 (1992); Gurgo, *et al.*, *Virol.* 164: 531-36 (1988); and the HIV SEQUENCE COMPENDIUM 1987-PRESENT (Myers, *et al.*, eds., Theoretical Biology and Biophysics Group, Los Alamos National Laboratory, Los Alamos, NM). Sequences from newly isolated strains also can be employed. *See, e.g.*, Ou, *et al.*, *Science* 256:1165-71 (1992); Zhang, *et al.*, *AIDS* 5:675-81 (1991); and Wolinsky, *Science* 255:1134-37 (1992).

[0131] Any suitable purification and glycosylation analysis methods may be employed with the glycosylated polypeptide (such as uniformly glycosylated) provided herein. For example, NP-HPLC of fluorescence labeled N-glycans from recombinant gp120 employs exoglycosidase digestion with HPLC to identify the various forms of glycans. *See, e.g.*, Scalan, *et al.*, *J. Virol.* 76:7306-21 (2002). Alternatively, the expression levels may be determined using flow cytometric analysis, ELISA, or Western blot analysis using glycan specific antibodies like, *e.g.*, the 2G12 antibody. Other exemplary methods include lectin-based assays, rocket affinoelectrophoresis, two-dimensional electrophoresis, fluorophore-assisted carbohydrate electrophoresis (FACE), capillary electrophoresis, nuclear magnetic resonance, mass spectrometry, gas-liquid chromatography with mass spectrometry (GLC/MS), fast atom bombardment (FAB), and electrospray ionization (ESI). *See, e.g.*, Brooks, *et al.*, FUNCTIONAL & MOLECULAR GLYCOBIOLOGY (BIOS Scientific Publishers Ltd 2002).

[0132] The glycosylated polypeptides (such as uniformly glycosylated protein) provided herein can also be used in assays for detecting the presence of neutralizing antibodies to the pathogen, *e.g.*, HIV. In such assays, the protein will normally be labeled with one of a variety of labels which find use in assays. These labels have been extensively

reported in the patent and technical literature, and include radionuclides, fluorescers, enzymes, enzyme substrates, particles, small molecules, and the like.

[0133] The wild type protein serotypes of the virulence factor need not be employed since one or more amino acids may be added, deleted or substituted so long as the relevant binding property, glycosylation sites, and other relevant immunological properties are retained. Thus, at least 90 %, usually at least 95%, more usually at least 99 % of the amino acids will be the correct amino acids and in the correct sequence. Usually, any changes will be at the N-terminus where from 0 to 5 amino acids may differ.

[0134] In one embodiment, the uniformly glycosylated gp120 provided herein displays conformational epitopes which elicit highly neutralizing antibodies capable of neutralizing primary HIV-1 isolates. In some embodiments, the preferred proteins are those which can elicit antibodies which neutralize primary isolates in two or more different clades (*e.g.*, two or more of clades A, B, C, D, and E). Neutralization is determined as disclosed herein.

[0135] The invention provides kits comprising any of compositions described herein, *e.g.*, uniformly glycosylated proteins. The kits also can contain instructional material teaching the methodologies and industrial uses of the invention, as described herein.

[0136] Specifically provided herein is a compartment kit comprising one or more containers, wherein a first container comprises one or more antibodies engineered by the present methods, and one or more other containers comprising one or more of the following: wash reagents, reagents necessary for administration of the composition or capable of detecting presence of antibody specific for the composition. The containers can be glass, plastic, or strips of plastic or paper. Types of detection agents include labeled secondary antibodies, other labeled secondary binding agents, or in the alternative, if the primary antibody is labeled, the enzymatic, or antibody binding reagents that are capable of reacting with the labeled antibody. Ancillary materials to assist in or to enable performing such a method may be included within a kit provided herein.

C. Vaccines using glycosylated polypeptides or whole mutant yeast cells

[0137] Provided herein is a method of inducing neutralizing antibodies against a pathogen (such as HIV) in a subject, comprising administering to the subject an effective amount of any of the compositions described herein.

[0138] Provided herein is a method of inducing neutralizing antibodies against a pathogen in a subject in need thereof, comprising administering an effective amount of a

pharmaceutical composition comprising a uniformly glycosylated recombinant protein, wherein the terminal glycan is a terminal α 1-2 glycan structure or a protein prepared by the method of preparing a uniformly glycosylated recombinant protein, wherein the glycan is oligomannose with a terminal α 1-2 glycan structure, comprising: a) providing a vector comprising a nucleotide sequence encoding the protein; b) transforming a cell with a defect in protein glycosylation; c) fermenting the transformed cells; and d) purifying the secreted recombinant protein from the cell supernatant as disclosed herein, and a suitable excipient. The method can further comprise administering an adjuvant. In one embodiment, the pathogen is HIV. In a specific embodiment, the uniformly glycosylated recombinant protein is Man₉GlcNAc₂ gp120 or Man₈GlcNAc₂ gp120.

[0139] Also provided herein is a method of preventing or treating pathogen-induced (such as HIV-induced) disease in a subject, comprising administering to the subject an effective amount of any of the compositions described herein.

[0140] Also provided herein is a method of preventing or treating pathogen-induced disease in a subject in need thereof, comprising administering an effective amount of a pharmaceutical composition comprising a uniformly glycosylated recombinant protein, wherein the terminal glycan is a terminal α 1-2 glycan structure or protein prepared by the method of preparing a uniformly glycosylated recombinant protein, wherein the glycan is oligomannose with a terminal α 1-2 glycan structure, comprising: a) providing a vector comprising a nucleotide sequence encoding the protein; b) transforming a cell with a defect in protein glycosylation; c) fermenting the transformed cells; and d) purifying the secreted recombinant protein from the cell supernatant as disclosed herein, and a suitable excipient. The method can further comprise administering an adjuvant. In one embodiment, the pathogen is HIV. In a specific embodiment, the uniformly glycosylated recombinant protein is Man₉GlcNAc₂ gp120 or Man₈GlcNAc₂ gp120.

[0141] In some embodiments, the pharmaceutical compositions provided herein further comprises multiple gp120 proteins from different HIV strains.

[0142] The vaccine compositions of the invention may be administered by any route clinically indicated, such as by application to the surface of mucosal membranes including, but not limited to intranasal, oral, ocular, gastrointestinal, rectal, vaginal, or genito-urinary routes. Alternatively, parenteral modes of administration such as intravenous, subcutaneous, intraperitoneal, or intramuscular may also be used. In some embodiments, the protein is administered transdermally or transmucosally.

[0143] Typically, vaccine protection against HIV infection induces a mucosal immune response, a systemic immune responses, or both. Useful routes for administration can include parenterally or non-parenterally routes, including but not limited to subcutaneous, intracutaneous, intravenously, intramuscular, peroral, mucosal or intranasal administration.

[0144] Vaccines according to the present invention may be formulated according to any suitable method known in the art, and may contain any suitable and compatible diluent, carrier, preservative, pharmaceutical excipient or other ingredient. *See, e.g.*, REMINGTON: THE SCIENCE AND PRACTICE OF PHARMACY 20th Ed. (Gennaro et al. (eds.), University of Sciences 2000). Suitable pharmaceutical carriers for administration intravenously, intraperitoneally, intramuscularly, transmucosally (for example intranasally, rectally, vaginally, buccally), transdermally, subcutaneously or any other route of administration are well known in the art and are contemplated for use with the invention. *See, e.g.*, Robinson, *et al.* (eds.), VACCINE PROTOCOLS 2nd. Ed. (Humana Press 2003); Powell, *et al.* (eds.), VACCINE DESIGN: THE SUBUNIT AND ADJUVANT APPROACH (Plenum Press 1995); Plotkin, *et al.* (eds.) VACCINES 4th Ed. (Saunders 2004); U.S. Pat. No. 4,235,877; 4,372,945; and 4,474,757. Suitable formulations, which may include an adjuvant, can be determined empirically.

[0145] The amounts of vaccine administered depend on the particular vaccine antigen and any adjuvant employed, the mode and frequency of administration, the age and weight of the recipient, and the desired effect (*e.g.*, prophylaxis and/or treatment), as determined by one skilled in the art. The amount of protein in a vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccinees. Such amount will vary depending upon which specific immunogen is employed. Generally, it is expected that each dose will comprise between 1 μ g and 100 mg of protein, sometimes between 2-200 μ g, and other times 4-40 μ g. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of antibody titers and other responses in subjects.

[0146] Administration can repeated as is determined to be necessary by one skilled in the art. In many instances it will be desirable to have multiple administrations of the vaccine, usually not exceeding six vaccinations, more usually not exceeding four vaccinations, and preferably one or more, usually about three vaccinations. The vaccinations will normally be at from 2 to 12 week intervals, more usually from 3 to 5 week intervals, with optional periodic boosters at intervals of 1 to 5 years. The course of

the immunization may be followed by assays for antibodies for pathogen, *e.g.*, HIV gp120 antibodies.

[0147] The active vaccines of the invention can be used alone or in combination with bioactive agents including but not limited to other relevant virulence factors, attenuated or killed forms of the virus, or adjuvants. Such additional agents can be administered simultaneously or sequentially through the same or a direct route of administration. In some embodiments, the administered proteins originate from different serotypes of the same or related pathogens. In one embodiment, the bioactive agent synergizes with the glycosylated protein provided herein to stimulate an effective immune response against the pathogen.

[0148] In some embodiments, glycosylated protein (such as uniformly glycosylated recombinant protein) provided herein is administered with an adjuvant. In the formulation of vaccines for use in the invention it is preferred that the adjuvant composition induces a humoral or Th2 response. Such adjuvants can include those that induce IL-4, IL-6, and/or IL-10 production as well as other means of enhancing the immunogenicity of T-independent antigens. *See, e.g.*, Mond et al., Annual Reviews 13:655-92 (1995); Lesinski et al., J. Microbiolog. Methods 47:135-49 (2001). However it will be understood that other responses, including cellular responses, are not excluded, particularly if other virulence factors are co-administered in the vaccine regimen. Therefore, well known adjuvants such as CpG oligodeoxynucleotides, tetanus toxoid, saponin, lipopeptide, and IL-12 are also contemplated. In a specific embodiment, the glycosylated protein provided herein is conjugated to stimulatory epitopes of tetanus toxin. *See, e.g.*, Kumat et al., J. Immunol. 148:1499-505 (1992).

[0149] The proteins provided herein may be delivered in a formulation with an adjuvant or separately from the adjuvant, either simultaneously or sequentially. Suitable adjuvants include an aluminium salt such as aluminium hydroxide gel (alum) or aluminium phosphate, but may also be a salt of calcium, iron or zinc, or may be an insoluble suspension of acylated tyrosine, or acylated sugars, cationically or anionically derivatized polysaccharides, or polyphosphazenes. *See, e.g.*, Robinson, *et al.* (eds.), VACCINE PROTOCOLS 2nd. Ed. (Humana Press 2003); Powell, *et al.* (eds.), VACCINE DESIGN: THE SUBUNIT AND ADJUVANT APPROACH (Plenum Press 1995); Plotkin, *et al.* (eds.) VACCINES 4th Ed. (Saunders 2004); U.S. Patent Nos: 6,797,276; 6,596,278; 6,491,919; 4,894,229; 5,814,321; 5,679,355; and 4,803,070. In some embodiments, the adjuvant is incomplete

Freund's adjuvant (IFA) or alum. The amount of the adjuvant employed can vary widely depending upon the nature of the adjuvant, generally ranging from 0.1 to 100 times the weight of the immunogen, more usually from about 1 to 10 times.

[0150] In some embodiments, the whole mutant yeast cells described herein (e.g., alive or killed) may be used in preventive or therapeutic vaccine. For example, the whole mutant yeast cells are formulated in a capsule, a pill, a gel, or a cream for oral, vaginal, or rectal administration.

[0151] The neutralizing capacity of the induced antibodies can be assessed using routine methods. Thus, it is contemplated that antibodies that neutralize HIV may inhibit one or more of the following: viral attachment, coreceptor engagement, membrane fusion, or some productive interaction. *See, e.g.*, US Patent No. 6,391,567; Lin, *et al.*, *Virology* 77:1337 (2003); Lee, *et al.*, *J. Virol.* 75:12028 (2001); Geijtenbeek, *et al.*, *Cell* 100:587 (2000); Brewley, *et al.*, *J. Am. Chem. Soc.* 123:3892-3902 (2001); US Patent Application Nos. 20030096221; 20050064390; 20040259785; 20040228869; and 20040096823. Cell fusion encompasses the joining or union of the lipid bilayer membranes found on mammalian cells or viruses such as HIV-1. This process is distinguishable from the attachment of HIV-1 to a target cell. In one embodiment, attachment is mediated by the binding of the HIV-1 exterior glycoprotein to the human CD4 receptor. The interaction can also include coreceptors including but not limited to CCR5, CXCR4, CCR2, CCR3, CCR8, STRL33, GPR-15, CX3CR1 and APJ. *See, e.g.*, Opperman *et al.*, *Cell Signal.* 16:1201-10 (2004); Philpott *et al.*, *Curr. HIV Res.* 1:217-27 (2003). In a particular embodiment, a neutralizing antibody is one that prevents or inhibits the binding of the glycosylated gp120 to DC-SIGN. *See, e.g.*, US Patent No. 6,391,567; Lin *et al.*, *J. Virol.* 77:1337-46 (2003); Mitchell *et al.*, *J. Biol. Chem.* 276:31:28939-45 (2001).

[0152] The neutralizing antibody can be any isotype. In some embodiments, the antibody is an IgG antibody. More particularly, the antibody can be an IgG₁, IgG₂, IgG₃, or IgG₄ antibody.

D. Methods of making and identifying antibodies specific for essentially uniformly glycosylated compositions

[0153] Also provided herein is a method of preparing an antibody which specifically binds a protein comprising a terminal α 1-2 glycan structure, the method comprising immunizing an animal with an effective amount of a composition described herein (e.g., a

composition comprising a recombinant protein made by the method or the protein as disclosed herein). The composition can further comprise an adjuvant, a carrier, or both. Provided herein is the isolated antibody generated by the method disclosed herein as well as the hybridoma that produces the isolated antibody. In one embodiment, the antibody is specific for $\text{Man}_9\text{GlcNAc}_2$ gp120 or $\text{Man}_8\text{GlcNAc}_2$ gp120 and preferably has no substantial cross-reactivity with human glycoproteins.

[0154] Any suitable method of generating an antibody specific for the glycosylated proteins provided herein can be employed. Methods of immunization, producing and isolating antibodies (polyclonal and monoclonal) are known to those of skill in the art and described in the scientific and patent literature. *See, e.g.*, Coligan, CURRENT PROTOCOLS IN IMMUNOLOGY (John Wiley 2005); Stites (eds.), BASIC AND CLINICAL IMMUNOLOGY (9th ed.) (Lange Medical Publications 1998); Goding, MONOCLONAL ANTIBODIES: PRINCIPLE AND PRACTICE (3RD ED. (Academic Press 1996); Harlow, ANTIBODIES: A LABORATORY MANUAL, (Cold Spring Harbor 1998).

[0155] Antibodies provided herein can be isolated from natural sources, be synthetic, or be recombinantly generated polypeptides using methods known in the art. The antibodies can be recombinantly expressed *in vitro* or *in vivo*. *See e.g.*, Caruthers (1980) Nucleic Acids Res. Symp. Ser. 215-223; Horn (1980) Nucleic Acids Res. Symp. Ser. 225-232; Banga, A. K., THERAPEUTIC PEPTIDES AND PROTEINS, FORMULATION, PROCESSING AND DELIVERY SYSTEMS (1995) Technomic Publishing Co., Lancaster, Pa. For example, antibody synthesis can be performed using various solid-phase techniques (*see e.g.*, Roberge (1995) *Science* 269:202; Merrifield (1997) *Methods Enzymol.* 289:3-13) and automated synthesis may be achieved, *e.g.*, using the ABI 431A Peptide Synthesizer (Perkin Elmer) in accordance with the instructions provided by the manufacturer. Exemplary descriptions of recombinant means of antibody generation and production include Delves, ANTIBODY PRODUCTION: ESSENTIAL TECHNIQUES (Wiley, 1997); Shephard, *et al.*, MONOCLONAL ANTIBODIES (Oxford University Press, 2000); Goding, MONOCLONAL ANTIBODIES: PRINCIPLES AND PRACTICE (Academic Press, 1993); CURRENT PROTOCOLS IN IMMUNOLOGY (John Wiley & Sons, most recent edition). Antibodies also can be generated *in vitro*, *e.g.*, using recombinant antibody binding site expressing phage display libraries, in addition to the traditional *in vivo* methods using animals including but not limited to genetically engineered animals. *See, e.g.*, Hoogenboom *Trends Biotechnol.* 15:62-70 (1997); Katz (1997) *Annu. Rev. Biophys. Biomol. Struct.* 26:27-45 (1997);

Fishwild, *et al.*, *Nat. Biotech.* 14:845 (1996); Mendez, *et al.*, *Nat. Genet.* 15:146 (1997); US Patent No: 6,632,976.

[0156] Once produced, polyclonal or monoclonal antibodies are tested for specific recognition of the glycosylated protein by Western blot or immunoprecipitation analysis by standard methods, *e.g.*, as described in Ausubel, *et al.*, *supra*. In some embodiments, the antibodies can be humanized. U.S. Pat. No. 5,585,089; U.S. Pat. No. 5,693,761; and WO 90/07861.

[0157] The antigen with essentially uniform glycosylation is produced using the methods provided herein. The eliciting antigen may be a single epitope, multiple epitopes, or the entire protein alone or in combination with one or more immunogenicity enhancing agents known in the art. The eliciting antigen may be an isolated full-length protein, a cell surface protein (*e.g.*, immunizing with cells transfected with at least a portion of the antigen), or a soluble protein (*e.g.*, immunizing with only the extracellular domain portion of the protein).

[0158] The antibody of the claimed invention can be an IgG, IgM, IgD, IgE, or IgA antibody. In some embodiments, the antibody is an IgG antibody. More particularly, the antibody can be an IgG₁, IgG₂, IgG₃, or IgG₄ antibody. Any suitable source can be used for the antibody. For example, the antibody can be a human, murine, rat, rabbit, bovine, camel, llama, dromedary, or simian antibody. Once a suitable antigen binding region is identified, *e.g.*, the CDRs from an antibody which is neutralizing *in vitro*, *in vivo* or both, the antibody can be modified in any suitable manner to enhance its therapeutic efficacy. The modifications in the antibody of the claimed invention comprise at least one mutation in the amino acid sequence of the antibody. For example, one or more mutations can be introduced by modifications, additions or deletions to a nucleic acid encoding the antibody to alter one or more properties of the antibody. The antibody of the present invention can be modified in any suitable manner to confer or enhance a desirable effector function or physical characteristic. *See, e.g.*, U.S. Patent No. 6,737,056; and US 2004/0132101. Thus, a nucleic acid encoding the antibody modified by the method of the invention can be altered by any suitable means.

[0159] Therefore, the antibody can be a humanized antibody, a chimeric antibody, a bispecific antibody, a fusion protein, or a biologically active fragment thereof. In some embodiments, the antibody (or biologically active fragment thereof) is a fusion protein. The fusion protein can encompass additional peptide sequence that simplifies purification

or production. Fusion proteins also may include domains and/or whole polypeptides that are biologically active in a manner that complements the activity of the antibody. For example, the antibody can be fused to a cytokine, ligand, adhesion molecule, peptide, receptors, enzymes, therapeutic proteins, dyes, small organic molecules, or any biologically active portion thereof.

[0160] Any suitable means can be used to determine the binding affinity of the antibody of the present invention. In one example, the affinity is determined by surface plasmon resonance (Biacore). The antibody of the present invention can also be modified further to increase binding affinity using methods known in the art. *See, e.g.*, U.S. Patent No. 6,350,861.

[0161] The antibody provided herein can be one that is suitable for immunoassays. Particular assays include ELISA assays, sandwich assays, radioimmunoassays, and Western Blots. *See e.g.*, ANTIBODY ENGINEERING: A PRACTICAL APPROACH (Oxford University Press, 1996). These monoclonal antibodies will usually bind with at least a K_d of about 1 μ M, more usually at least about 300 nM, typically at least about 30 nM, preferably at least about 10 nM, more preferably at least about 3 nM or better, usually determined by ELISA

[0162] Any suitable method may be employed to determine the biological activity of the antibody in the presence of the virulence factor. In one embodiment, an antibody has anti-virulence factor activity if the antibody reduces the pathogenicity of the organism and/or the toxicity of the virulence factor by at least 20%, at least 50%, 60%, 70%, 80%, 90%, or 100%. In another embodiment, the antibody of the present methods has anti-virulence activity if the antibody reduces the pathogenicity of the organism and/or the toxicity of the virulence factor by at least 20%, at least 50%, 60%, 70%, 80%, 90%, or 100% in presence of one or more other anti-virulence factors.

[0163] The antibody of the present invention can specifically bind any suitable glycosylated virulence factor. The virulence factor can be an adherence factor, a coat protein, an invasion factor, a capsule, an exotoxin, or an endotoxin. The anti-pathogenic effect of the antibody can result from the antibody binding a virulence factor, clearance of the factor, inactivation of the factor, and the like.

[0164] In another aspect, the present invention provides an isolated or recombinant nucleic acid comprising a sequence encoding the antibody disclosed herewith, a vector comprising the encoding nucleic acid, and a cell comprising the encoding nucleic acid or

the vector comprising the encoding nucleic acid. Typically, the vector comprises the antibody-encoding nucleic acid operably linked to a promoter suitable for expression in the designated host cell. Host cells for expressing the nucleic acids, expression cassettes and vectors of the invention include bacteria, yeast, fungi, plant cells, insect cells and mammalian cells. Thus, the invention provides methods for optimizing codon usage in all of these cells, codon-altered nucleic acids and polypeptides made by the codon-altered nucleic acids. Exemplary host cells include gram negative bacteria, such as *Escherichia coli* and *Pseudomonas fluorescens*; gram positive bacteria, such as *Streptomyces diversa*, *Lactobacillus gasseri*, *Lactococcus lactis*, *Lactococcus cremoris*, and *Bacillus subtilis*. Exemplary host cells also include eukaryotic organisms, e.g., various yeast, such as *Saccharomyces* spp., including *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, *Pichia pastoris*, and *Kluyveromyces lactis*, *Hansenula polymorpha*, *Aspergillus niger*, and mammalian cells and cell lines, and insect cells and cell lines. Thus, the invention also includes antibodies and their encoding nucleic acids optimized for expression in these organisms and species. See, e.g., U.S. Pat. No. 5,795,737; Baca (2000) *Int. J. Parasitol.* 30:113-118; Hale (1998) *Protein Expr. Purif.* 12:185-188; Narum (2001) *Infect. Immun.* 69:7250-7253; Outchkourov (2002) *Protein Expr. Purif.* 24:18-24; Feng (2000) *Biochemistry* 39:15399-15409; and Humphreys (2000) *Protein Expr. Purif.* 20:252-264.

[0165] In one aspect, the present invention provides a pharmaceutical composition comprising the antibody of the present invention, and a suitable excipient which is administered as a therapeutic agent. In one aspect, the present invention provides a pharmaceutical composition comprising the engineered antibody of the present invention, and a suitable excipient. The present invention provides a pharmaceutical composition comprising at least one monoclonal antibody as set forth above, and a suitable excipient. Formulations and excipients useful in the pharmaceutical compositions are those well known in the art. An antibody useful in the present methods (from whatever source derived, including without limitation from recombinant sources) may be administered to a subject in need, by itself, or in pharmaceutical compositions where it is mixed with suitable carriers or excipient(s) at doses to treat or ameliorate a variety of disorders. See e.g., REMINGTON: THE SCIENCE AND PRACTICE OF PHARMACY (latest edition). Such a composition may also contain (in addition to protein and a carrier) diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The term “pharmaceutically acceptable” means a non-toxic material that does not interfere with the

effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. The pharmaceutical composition of the invention may also contain other anti-pathogen such cytokines or antimicrobial agents as is desirable.

[0166] When the antibody of the present invention is co-administered with one or more biologically active agents, the antibody provided herein may be administered either simultaneously with the biologically active agent(s), or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering protein of the present invention in combination with the biologically active agent(s).

[0167] Toxicity and therapeutic efficacy of such antibodies can be determined using routine methods. The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. *See, e.g., Fingl., et al., THE PHARMACOLOGICAL BASIS OF THERAPEUTICS 1* (latest edition).

[0168] Further provided herein is a method of inducing neutralizing antibodies against a pathogen in a subject in need thereof, comprising administering an effective amount of a pharmaceutical composition comprising the protein provided herein or a protein prepared by the method provided herein, and a suitable excipient. In some embodiments, the method further comprises administering an adjuvant. In a specific embodiment, the pathogen is HIV.

[0169] Yet further provided herein is a method of preventing or treating pathogen-induced disease in a subject in need thereof, comprising administering an effective amount of a pharmaceutical composition comprising the protein provided herein or a protein prepared by the method provided herein, and a suitable excipient. In some embodiments, the method further comprises administering an adjuvant. In a specific embodiment, the pathogen is HIV. In one embodiment, the protein is gp120. In some embodiments, the pharmaceutical composition further comprises multiple gp120 proteins from different HIV strains.

[0170] Further provided is a method of identifying a neutralizing monoclonal antibody for HIV transmission, comprising: a) contacting a first cell, wherein the first cell expresses dendritic cell-specific C-type lectin (DC-SIGN), with a second cell expressing CD4+ and CCR5+, b) coculturing the first and second cells with infectious viral particles; c) contacting coculture with a candidate antibody prepared by the method of claim 20; and d)

determining relative infectivity, whereby the neutralizing antibody is one that reduces or eliminates relative infectivity of the second cell by infectious viral particles. See, *e.g.*, US 6,391,567; Bashirova, *et al.*, J. Exp. Med. 193:671 (2001); Lee, *et al.*, J. Virol. 75:12028 (2001).

[0171] Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention, unless specified.

EXAMPLES

Example 1

[0172] **Selection of an expression system.** A strain of *S. cerevisiae* deficient for Pmr1p protein, a calcium-dependent ATPase important for the transport of Ca^{++} from the ER to the Golgi was selected. (55,56). The *pmr1* Δ strain is a so-called super-secretor of heterologous proteins, secreting nearly 100% of heterologous proteins. (57-58). This strain secretes glycoproteins with only $\text{Man}_8\text{GlcNAc}_2$ trimmed from $\text{Man}_9\text{GlcNAc}_2$ by mannosidase I in the ER as reported for several proteins (58-60).

[0173] **Cloning of HIV-1 gp120 genes to express secreted glycoproteins.** The signal sequence of the *S. cerevisiae* α mating factor (MF α) was cloned into an expression vector pYES2/CT (Invitrogen). The coding sequence for the MF α was amplified from total yeast DNA by PCR. The PCR primers contained KpnI and BamHI sites. A 250 bp fragment was amplified, purified and ligated into the pGEM-T (Promega). After transformation and miniprep identification, a positive clone was identified. The pGEM-T-MF α and pYES2/CT plasmids were then digested with BamHI and KpnI, simultaneously. After purification of the MF α band and pYES2 band, the plasmids were ligated, transformed into NovaBlue competent cells and seeded onto LB-ampicillin plates. A positive clone containing MF α was identified and used to clone gp120.

[0174] The sequences encoding HIV-1 gp120 proteins from strains of JR-FL, JR-CSF and YU2 were amplified by PCR using their corresponding plasmids (AIDS Research and Reference Reagent Program, ARRRP) as the templates. The PCR amplified gp120 DNA fragments were cloned into pYES2/CT-MF α plasmid containing EcoRI and XbaI sites. Then the three pYES2/CT-MF α -gp120 plasmids were transformed into NovaBlue competent cells, and positive clones were identified by digestion analysis of minipreps.

These plasmids were used for transformation into a diploid yeast strain *pmr1*Δ (Invitrogen) using a quick Lithium acetate transformation protocol. The pYES2/CT vectors contained the inducible *GALI* promoter, permitting high expression levels of the recombinant protein following induction with galactose. To produce and secrete gp120 into culture media, transformed yeast cells were grown overnight at 30°C in –Ura/raffinose media. After centrifugation, the cells were resuspended with –Ura/galactose media for induction of gp120 expression.

[0175] Expression, purification, and analysis of uniformly glycosylated HIV-1 Env proteins. The *pmr1*Δ strain is temperature sensitive and grows slowly at permissive temperatures. Therefore, gp120 expression was examined at different temperatures and time points in order to determine the conditions under which these yeast cells can express the highest level of proteins and secrete them most efficiently into culture media. The *pmr1*Δ yeast cells secreted the greatest amount of gp120 proteins at 25°C for 7 to 14 days (Figure 2A). gp120 proteins were detected after 1-day of culture and continued to accumulate in the culture media for at least 14 days, indicating that the yeast cells continued production and secretion of the heterologous proteins. It also indicated that the glycosylated gp120 proteins were stable at room temperature. If the gp120 glycoproteins produced in the *pmr1*Δ strain contained only Man₈GlcNAc₂ oligosaccharides, the proteins should have predicted molecular masses of 99603 Da, 96385 Da, and 99621 Da for JR-CSF, JR-FL, and YU2, respectively. Western blots showed that the proteins migrated at these predicted sizes in SDS-PAGE. The blot was probed with MAb 2G12 which readily detected the gp120 proteins produced in the *pmr1*Δ strain (Figure 2A).

[0176] Purification procedures were simple because yeast culture media was protein poor. A combination of mannose binding lectin affinity chromatography and gel filtration was sufficient to obtain purities greater than 95%. In brief, the culture supernatants at 14 days were concentrated and dialyzed against dialysis buffer (1 mM CaCl₂ and 1 mM MnCl₂) for 16 h at 4°C. After centrifugation at 40,000 g for 30 min at 4°C, the precipitate containing proteins of non-interest was removed. Agarose–Con-A column (Vector Laboratories, CA) was then used to purify these gp120 proteins. Supernatant containing gp120 proteins was loaded onto pre-equilibrated Con-A-agarose columns and incubated at 4°C for 16 h. The column was then washed with dialysis buffer and the bound proteins were eluted with elution buffer (500 mM α-methyl mannoside, 250 mM NaCl, 20 mM TrisCl, 1 mM CaCl₂ and 1 mM MnCl₂, pH6). The purified materials contained mainly a

100 kDa and a 160 kDa protein. Gel filtration was then used to remove the 160 kDa protein. The prepacked column of Sephacryl S-100 HR (Amersham Pharmacia Biotech) was washed with washing buffer (50 mM imidazole, 100 mM NaCl, 10 mM sodium phosphate, pH 8.0). The bound proteins were eluted with elution buffer (500 mM imidazole, 100 mM NaCl, 10 mM sodium phosphate, pH 8.0) and 1 ml fractions were collected. The fractions were analyzed by SDS-PAGE followed by Coomassie blue staining (Figure 2B).

[0177] Following purification, the carbohydrate types will be analyzed using PGNase F digestion and mass spectrometry. Typically, greater than 90% of the Man will be Man₈ or Man₉.

[0178] To estimate the quantity of gp120 and contaminants, the purified proteins were digested with an enzyme endoglycosidase H (Endo H) that is known to cleave high mannose and hybrid types of carbohydrates. The results from Figure 3 show that the predominant band was a 55-60 kDa protein, correlating to the size of gp120 polypeptide. Based on the data from Figure 3, the estimated gp120 purity was greater than 90%. In later experiments, this band was further identified to be mostly PST1, a yeast protein. See figure 15.

[0179] Antibodies specific for glycosylated gp120 proteins. In order to test whether mannose-specific antibodies can be induced and react with HIV-1 Env proteins, rabbits were immunized with a yeast cell wall extract, zymosan A, which contains β -glucan and α -mannan. These mannose-rich structures contained mostly terminal α 1,3 and α 1,6 linkages, with a few α 1,2 structures such as those commonly found in gp120. Two rabbits were immunized with 200 μ g of zymosan A containing CFA at week 0, and boosted with 100 μ g of antigen containing IFA at weeks 2 and 4 and thereafter every 4 weeks. Immune sera were collected at week 5 and 1 week after each booster. The antigen elicited a good antibody response as determined by ELISA analysis, using either zymosan A or mannan captured to the plates (Figure 4A). Notably, the immune sera also recognized some of the tested gp120 glycoproteins (Figure 4B). The anti-mannan sera showed a broad, though relatively low level of reactivity to clades E (ChenMai and 93TH975), C (CN54 and 96ZM651) and one of clade B (LAV). In contrast, 2G12 MAb mainly reacted with clade B (ADA, JR-FL, BAL, LAV) strains, but had no response to the clade C (CN54 and 96ZM651) or little response to the clade E strains tested (93TH975 and ChenMai). The high titer of antibody to the clade E Chen strain was further confirmed by Western blot and

repeated ELISA. These results suggest that antibodies against mannose can be induced in animals, and that reactivity to HIV gp120 can be obtained. These findings provide a strong rationale for moving forward with our novel, mannose-rich gp120 immunogens.

[0180] The gp120 proteins from strains YU2, JR-FL and JR-CSF expressed in *pmr1*Δ yeast cells contained 22-24 Man8 structures while those expressed in mammalian cells from the same HIV strains contained about half Man5-9 and half complex carbohydrate structures. To determine if gp120 proteins expressed in yeast were recognized more efficiently by antibodies against mannan and as well as 2G12, we used the purified gp120 protein from YU2 strain from yeast to compare with YU2 expressed in mammalian cells. The results from this ELISA analysis (Figure 4C) show that gp120 protein from yeast has much higher antigenicity to anti-mannan antibody than that from mammalian cells. This figure also showed that the anti-mannan antibody recognized the Man9-BSA conjugate but not BSA alone.

[0181] Isolation of Man9 and conjugation of Man9 to BSA. In order to use Man₉GlcNAc₂ as a screening tool for anti-α1,2-Man structure antibodies, Man9 was isolated from a plant protein and conjugated to BSA. Specifically, the Man9 glycan was isolated from soybean agglutinin (SBA, EY labs) by the Glycobiology Core Facility at the University of California San Diego (UCSD). Briefly, the N-glycans on purified SBA were released using PNGase F and purified using TCA precipitation to remove the proteins. The supernatant was passed over a Sep-Pak C18 cartridge to remove residual detergent, and the glycans were purified on a porous graphitized carbon cartridge. Monosaccharide analysis indicated that the yield was 0.2 mg from 25 mg of protein. The glycan was analyzed by HPAEC-PAD and compared with reduced Man₉GlcNAc₂ standard and reduced RNase B glycans. An estimated 90% of the glycan was in fact Man₉GlcNAc₂ and the ratio of Man₉GlcNAc₂ to Man₈GlcNAc₂ was 9:1. An aliquot was subjected to MALDI-TOF mass spectrometric analysis. The major species was Hex₉HexNAc₂ although smaller glycans were also observed in lower amounts.

[0182] Man₉GlcNAc₂ was then conjugated to BSA using sulfo-SMCC (Pierce Biochemicals) method. Briefly, Man₉GlcNAc₂ was reacted with 2-aminoethanethiol-HCl at 85°C for 30 min. The reaction mixture was dialyzed against water to remove the reaction reagent and salt, and then vacuum dried. The Man9 derivative was then dissolved in 100 μl SMCC reaction buffer (0.1 M sodium phosphate, 50 mM EDTA, pH 7.4). BSA was activated with sulfo-SMCC in the reaction buffer for 2 h. The excessive SMCC was

removed by passing the reaction mixture through Sepharose G25 column. Fractions containing BSA-SMCC were combined and concentrated. 100 µg BSA-SMCC was reacted with 50 µg of derived Man9 for 30 min. The reaction was stopped by addition of 0.5 M cysteine, and dialyzed against phosphate buffer saline. The solution was vacuum dried and the BSA-Man9 conjugate was dissolved with 0.1 ml of water to have the concentration of BSA at 1 mg/ml. Figure 5 shows the migration shift of the conjugate of BSA-Man9 compared to BSA-SMCC and non-conjugated BSA.

[0183] Preparation of DC-SIGN antibodies and recombinant proteins. Antibodies against DC-SIGN were used to monitor expression of the recombinant DC-SIGN protein. Two antibodies were produced in rabbits using synthetic peptides. The immune sera were purified by immunoaffinity chromatography. Both peptide antibodies recognized a 44 kDa band in the lysates from human mucosal tissues. This band was blocked by the antigenic peptides indicating that these two antibodies are specific.

[0184] Cloning and expression of DC-SIGN. In order to perform gp120-DC-SIGN binding assays and test the neutralizing activity of elicited antibodies, the extracellular domain of DC-SIGN was cloned and expressed. The cDNA fragment of DC-SIGN was prepared by PCR amplification and ligated into the pET100 vector (Invitrogen). BL21(DE3) bacteria were used to express the recombinant protein in the presence of 0.5 M IPTG. Bacterial cells were harvested after 4 h of IPTG induction. The bacterial lysates were separated into soluble and insoluble fractions and analyzed by SDS-PAGE. The recombinant ECD protein was found in the insoluble fraction and purified using Ni-NTA columns (Figure 6A).

[0185] Refolding and purification of DC-SIGN from bacteria inclusion bodies. Because the insoluble proteins in inclusion bodies were incorrectly folded. The denatured proteins were refolded to obtain biologically functional proteins, and the refolded proteins were purified with Ni-NTA and D-mannose affinity chromatography. In brief, bacterial cells were lysed with lysis buffer (50 mM Tris-Cl, 150 mM NaCl, 5 mM 2ME, pH 7.45) containing 1 mg/ml lysozyme and 5 µg/ml DNase I to release inclusion bodies and soluble proteins. The inclusion bodies were precipitated by centrifugation and thoroughly washed to remove soluble proteins. The inclusion body was lysed in refolding buffer (20 mM Tris, 10 mM 2ME, 10 mM DTT, 1 mM glycine, 1 mM GSH, 0.1 mM GSSG, 1 mM EDTA, pH 10.5) containing 8 M urea. The urea concentration was diluted with refolding buffer and the solution was adjusted with 1 N HCl sequentially to pH 9.0, 8.8, 8.5 and 8.0. The

refolding solution was centrifuged at 40,000 g 30 min to remove the insoluble materials. The refolded proteins were then purified with Ni-NTA agarose column and eluted with lysis buffer containing 500 mM imidazole. The quantity and purity of purified ECD protein was analyzed with SDS-PAGE followed by Coomassie blue staining. Figure 6A shows that the refolded protein had purity greater than 90% after Ni-NTA purification. The elute from Ni-NTA column containing ECD protein was further purified using a D-Mannose-agarose column pre-equilibrated with loading buffer in order to select proteins competent to bind ligand. The column was washed with loading buffer and the bound proteins were eluted with 50 mM α -D-mannose. Fractions were analyzed by SDS-PAGE followed by Coomassie blue staining. The binding assay in Figure 6B shows this recombinant protein is functional since it binds to gp120 with a similar affinity to a polyclonal antibody.

[0186] Binding assay of DC-SIGN-gp120 interaction. To test the DC-SIGN activity, a binding assay of DC-SIGN-gp120 is performed. Different dilutions of immune sera with different concentrations of 2G12 MAb as a control are incubated with 200 ng of gp120 proteins to determine IC_{50} values. gp120 proteins from YU2 and JR-FL strains produced in mammalian cells and yeast cells are used in parallel to compare. Then the wells are incubated with anti-gp120 polyclonal antibody (US Biological) followed by goat anti-rabbit IgG-HRP conjugates. Each condition is in duplicate. The quantity of gp120 bound to DC-SIGN is determined using a standard curve created based upon the gp120 concentration and OD value. The percent inhibition at a given antibody concentration is expressed as $[(gp120 \text{ quantity in the absence of antibody} - gp120 \text{ quantity in the presence of a given antibody concentration}) / gp120 \text{ quantity in the absence of antibody}] \times 100$. The inhibition potency, *i.e.*, 50% (IC_{50}) and 90% (IC_{90}) inhibitory doses, are determined. To test the broadly neutralizing reactivity of each antibody, recombinant gp120 glycoproteins from clades A, B, and C will be used.

[0187] The purified ECD protein at 500 ng/well is used to coat microwell plates for 16 h. The wells are blocked with TBS-Tween containing 0.1% BSA. The gp120 protein from YU2 strain produced in mammalian cells is used to bind to the immobilized proteins. Then the wells are incubated with anti-gp120-MN polyclonal antibody followed by goat anti-rabbit IgG-HRP conjugates. After color development with substrate, optical density (OD_{450}) is read with a microwell reader. Each condition is set in duplicate.

[0188] Besides the two polyclonal antibodies, MAbs against the purified ECD protein have also been developed. After first screening of culture supernatants using ELISA and expansion of the positive clones, hybridomas were further screened using Western blot (Figure 7). These MAbs showed high specificity since a longer exposure of film (30 min) did not show other detectable bands.

[0189] **Recombinant gp120 proteins and neutralization assay.** gp120 proteins from different clades will be used to perform the primary screening of the immune sera and the hybridoma supernatants. These proteins are ideally expressed in mammalian cells to be fully glycosylated. Many of these proteins are available from commercial sources. Less than 1 mg of each protein will be needed for screening. The HIV neutralization assay are typically performed as described in Binley et al., *J. Virol.* 78:13232-52 (2004), Li et al., *J. Virol.* 79:10108-25 (2005); Mascola et al., *J. Virol.* 79:10103-7 (2005). Briefly, the retrovirus plasmid pHIV-gpt and gp160 env pHXB2 to produce a high titer pseudovirus after co-transfection of Cos-7 cell. This pseudovirus infects luciferase indicator cell lines LuSIV very well as measured by luciferase activity using Bright-Glo Luciferase assay kits (Promega) and a recently purchased Veritas Luminometer (Tuner Biosystem). This system was used to test anti-HIV-1 polyclonal antibodies recently produced in our lab (Figure 8). The two polyclonal antibodies showed dose dependent neutralizing activity. Trim5 α polyclonal antibody was used as a negative control and human MAbs (2F5, 4E10, and 2G12) as positive controls.

[0190] **Preparation of gp120 conjugated to immunostimulant.** To enhance the immunogenicity of the uniformly glycosylated gp120, multiple antigenic peptide (MAP) containing universal T cell epitopes from tetanus toxin will be synthesized, conjugates of gp120-tetanus toxoid and gp120-MAP will be prepared using different coupling procedures, and will be analyzed for the coupling efficiency and the antigenicity. Briefly, a multiple antigenic peptide (MAP) with 4 branches of the p2 sequence (p24) will be synthesized by a well-established peptide company (AnaSpec, Inc.). The MAP peptide will be purified greater than 90% purity by HPLC and analyzed by mass spectroscopy. The MAP peptide will be covalently coupled to the gp120 protein by using the cysteine residue that has been added to the core matrix. The cysteine sulfhydryl group will be cross-linked to free amino groups on the gp120 proteins using the SMCC method, and thus should not modify the carbohydrate structures that we hope to develop antibodies against. In brief, the gp120 proteins at 10 mg/ml will be reacted with Sulfo-SMCC (Pierce) solution at 10 mg/ml

in coupling buffer (1 x PBS, 0.1 M EDTA, pH7.2) for 2 h. The reaction solution will be passed through a Sephadex G-25 column to remove excess SMCC. Fractions containing protein will be collected. The synthetic MAP peptide will be dissolved in coupling buffer at 10 mg/ml and mixed with activated antigen at a 30:1 molar ratio, which allows most of the free amine groups on the gp120 proteins to be coupled. The cross-linking reaction proceeds for 2 h. All reactions will be performed at room temperature.

[0191] A gp120-tetanus toxoid conjugate will be prepared using a similar approach. The tetanus toxoid protein (TT, Statens Serum Institute, Denmark) will be coupled to gp120 by crosslinking through their free amino groups. Briefly, fresh glutaraldehyde in 100-fold molar excess, to prevent self-conjugates, is reacted to TT in 50 mM borate buffer, pH 8.0. Free glutaraldehyde is removed by passing through a Sephadex G-25 column. TT-glutaraldehyde is then coupled to gp120. Free aldehyde groups are blocked by the addition of 1 M glycine. The reaction mixture is re-passed through the Sephadex G-25 column. The fractions containing TT-gp120 conjugate are collected and dialyzed against PBS.

[0192] The coupling efficiency is analyzed using gel shift assays. Briefly, the conjugates are analyzed with 4-20% gradient SDS-PAGE followed by Coomassie blue staining, and Western blot using antibodies against gp120. The conjugates migrate slower than the non conjugated protein. The size changes of the conjugates depend on the numbers of MAP peptide and protein molecules that have been coupled to the gp120 protein. The antigenicity of the conjugates will be tested by comparison to non-conjugated gp120 protein using ELISA with the polyclonal antibody against mannan and 2G12 MAbs.

[0193] **Screening the immunogenicity of the gp120 proteins and their conjugates *in vivo*.** Six groups of 6 BALB/c mice at 5 to 6 weeks old will be immunized (Table 1). Antigens are injected subcutaneously at the doses indicated. The mice are boosted with the same dose of antigen 14 days after the primary immunization and every three weeks thereafter. They are bled 1 week after each booster injection. Thus, the mice are immunized at week 0, 2, 5, 8 and 11 and immune sera are collected at week 0, 3, 6, 9, and 12. Both YU2 and JR-FL can be employed in order to have more Man α 1,2-Man epitopes and develop more numbers of broadly neutralizing MAbs.

Table 1.
Groups of immunization

Gro up	gp120/ conjugat e	Adjuvant	Significance
1.	10 µg	None	No adjuvant control
2.	10 µg	CFA/IFA	Co-administration of gp120 with CFA/IFA adjuvant as control
3.	20 µg	gp120-TT	Compare the effect of TT with CFA/IFA
4.	50 µg	gp120-TT	Compare the dose effect of the conjugate
5.	20 µg	gp120-MAP	Compare the effect of TT-MAP with CFA/IFA
6.	50 µg	gp120-MAP	Compare the dose effect of the conjugate

[0194] Antibody titers will be tested using ELISA with gp120 antigens coated microwell plates. The ELISA procedures are briefly described. Costar microwell plates are coated with 300 ng of antigen/well in 100 µl of 50 mM carbonate (pH9.5) for 24 h. The wells are blocked with 200 µl of blocking buffer (1% BSA, 0.02% thimerosal in PBS) for 2 h. After washing, 100 µl of serially diluted immune sera is incubated with antigen-coated wells for 2 h. After 5 washes with PBS-Tween, the wells are incubated with 100 µl of goat anti-mouse IgG-HRP conjugates (Sigma) at 1:5000 in blocking buffer for 2 h. After washing, the wells are incubated with 100 µl of HRP substrate TMB solution (BD Bioscience) for 10 min and the reaction is stopped by addition of 100 µl of 1 N HCl. The optical density is read at A₄₅₀ by microwell reader. All reactions are performed at room temperature.

[0195] Screen immune sera for cross-reactive antibodies to gp120 proteins from different clades. Up to 12 clades of HIV-1 isolates globally have been reported and clades A (27%), B (12%) and C (42%) comprise more than 80% of all HIV-1 infections. The Man9 and Man8 forms of high mannose oligosaccharides are highly conserved in all HIV-1 strains and isolates. Thus, antibodies against the Man α 1,2-Man structures could recognize an epitope from a single Man8 or Man9, or a conformational epitope formed by 2 or 3 of these glycans. Therefore, the antibodies against Man α 1,2-Man epitopes could recognize most HIV-1 strains and isolates.

[0196] To develop MAbs with broadly neutralizing activity, we will screen the immunized mice and hybridoma supernatant with gp120 glycoproteins from strains of clade A (e.g., Q2317, B2539, and Q1769), B (e.g., ADA, YU2, JR-FL, 89.6, BaL, SF162) and C (e.g., DU123, DU422, DU179, and 98IN012). In addition, antibodies against high mannose Man8 structures can be tested. Man8 will be isolated from gp120 protein produced in yeast, and conjugated to BSA using sulfo-SMCC method disclosed herein.

BSA-Man8 will be used to coat the microwell plate and to screen immune sera for antibodies against Man α 1,2-Man epitopes.

[0197] Mice with a serum titer $\geq 1:10,000$ to the immunogen, and $\geq 1:1,000$ to gp120 proteins from different clades and BSA-Man8 will be selected for creation of hybridomas. Briefly, the mice are boosted with same dose of antigen three days before fusion, after which they are sacrificed and the spleens are aseptically harvested. Cell suspensions are prepared by injection of 20 ml of sterile complete serum-free DMEM (CSFD) into a spleen. Concurrently, cultures of myeloma cells SP2/0-Ag14 (ATCC) are harvested. The splenocytes and myeloma cells are washed three times with DMEM and resuspended separately in 10 ml of CSFD. After assessing the cell viability, they are mixed at 1:1 ratio, supplemented with 30 ml DMEM, and pelleted by centrifugation. The cell pellet is resuspended by slowly adding 1 ml of 50% polyethylene glycol (PEG 400, Roche) with stirring after each drop. Sterile CSFD is added to the mixture drop-by drop with stirring to bring the final volume to 10 ml. The fused cells are centrifuged and resuspended with complete DMEM-20/HEPES/pyruvate/HAT. The cell suspension is adjusted to 2.5×10^6 total cells/ml. The fused cells are seeded in 96-well plate and incubated in a humidified 37°C, 5% CO₂ incubator for 7 days. Then, the media is replaced with complete DMEM-20/HEPES/pyruvate/HT. On day 12 to 15 the hybridomas are ready for screening as witnessed by visible hybridoma colonies. Typically, around 1×10^8 cells per spleen are seeded into ten 96-well plates, and obtain $\geq 80\%$ efficiency in fusion.

[0198] The supernatants in the wells with visible colonies are screened using ELISA with immunogen. The procedure is the same as screening immune sera except the antigen-coated wells are probed with 100 μ l of non-diluted hybridoma supernatant. The candidate hybridoma cells in positive wells are transferred to 24 wells with DMEM-20/HEPES/pyruvate and cloning/expansion medium, and expanded 3 to 5 days. The supernatants are screened again with gp120s from different clades and BSA-Man8, as above. Cells in the wells that show positive to all or most tested gp120 proteins, and BSA-Man8, are subjected to cloning by plating 1 and 3 cells per well. Cells are cultured for 7 to 10 days and candidate clones are screened again using the same panel of antigens as above. Subcloning and screening are repeated until stable hybridoma cell lines of interest are established.

[0199] To test the neutralizing activity of these MAbs against a large number of HIV-1 strains and primary isolates, MAbs will be produced in mice and purified with Protein A affinity chromatography.

[0200] Identify MAbs that can potently neutralize a broad range of HIV-1 stains and primary isolates. Culture supernatant and MAbs the HIV neutralization assay will be tested for the ability to neutralize homologous HIV strains. Thus, the hybridoma culture supernatant from animals immunized with YU2 or JR-FL gp120 produced in yeast will be tested for the ability to neutralize HIV pseudotypes bearing the YU2 or JR-FL Env protein as well as a luciferase reporter gene. Supernatants and MAbs will be tested for the ability to neutralize infection of p24-normalized virus stocks on U87 cells bearing CD4 and CCR5 (or CXCR4 when X4 virus strains are used). The virus pseudotypes are first incubated with non-diluted supernatant or serial dilutions of MAbs for one hour prior to infection. The amount of luciferase produced in infected cells is measured 48 to 72 hours later. Neutralizing activity is displayed as the percent inhibition of viral replication (luciferase activity) at each antibody dilution compared with an antibody-negative control: % inhibition $\{1 [\text{luciferase Ab luciferase Ab}] \}$ 100. Titers are calculated as the reciprocal of the MAb dilution conferring 50% inhibition (IC_{50}). Supernatants and MAbs that neutralize homologous virus will then be tested for the ability to neutralize a variety of primary HIV-1 strains.

Materials and Methods Used in Examples 2-4

Generation of Double and Triple Yeast Mutants:

[0201] Yeast strains and media. Four yeast deletion strains used in this study were acquired from Open BioSystems: MAT α , his3 Δ 1, leu2 Δ 0, lys2 Δ 0, ura3 Δ 0, mnn1:kanMX4 (10322); MATa, his3 Δ 1, leu2 Δ 0, met15 Δ 0, ura3 Δ 0, mnn4:kanMX4 (7034); MATa, his3 Δ 1, leu2 Δ 0, met15 Δ 0, ura3 Δ 0, och1:kanMX4 (4406); and MATa his3 Δ 1 leu2 Δ 0 met15 Δ 0 ura3 Δ 0 pmr1:kanMX4 (14534). Two parental yeast strains were acquired from ATCC for verification and screening of mating type: MATa, trp1 Δ 63 (Designation: BY4710) and MAT α , trp1 Δ 63 (Designation: BY4711). The wild-type, diploid yeast strain, INVSc1, was obtained from Invitrogen. Knockout of the indicated genes was verified by PCR using synthesized primers specific to the MNN1, OCH1, MNN4 and PMR1 genes (see sequences in the Yeast Deletion Project at

http://www.sequence.stanford.edu/group/yeast_deletion_project), and to the KanMX4 cassette (KanB, 5'-CTGCAGCGAGGAGCCGTAAT-3' (SEQ ID NO:8) and KanC, 5'-TGATTTTGATGACGAGCGTAAT-3' (SEQ ID NO:9)). Colony PCR was conducted as described (Giaever, 2002) using JumpStart Taq DNA Polymerase (Sigma). All yeast cultures were grown in YPD (1% Bacto-yeast extract, 2% Bacto-peptone, 2% glucose) or YPD + 0.3M KCl, supplemented with 200µg/ml G418 (Invitrogen) as indicated. Selection of diploids and haploids was conducted using CSM-met-lys plates (.67% yeast nitrogen base without amino acids, 0.07% CSM-met-lys dropout mix, 2% glucose, 2% Bacto-agar), supplemented with 20µg/ml L-methionine (CSM-lys) or 50µg/ml L-lysine-HCl (CSM-met). CSM-met-lys Dropout Mix was obtained from QBioGene. For yeast mating type tests, strains were selected on SD plates (0.67% yeast nitrogen base without amino acids, 2% glucose, 2% Bacto-agar).

[0202] Yeast mating. Yeast strain 10322 ($\Delta mnn1$) was mated with strains 7034 ($\Delta mnn4$), 4406 ($\Delta och1$), and 14534 ($\Delta pmr1$) to form the diploid strains DIP- $mnn1mnn4$ (MATa/ α ; $his3\Delta1/his3\Delta1$; $leu2\Delta0/leu2\Delta0$; $LYS2/lys2\Delta0$; $met15\Delta0/MET15$; $ura3\Delta0/ura3\Delta0$; $mnn4:kanMX4/MNN4$; $MNN1/mnn1:kanMX4$), DIP- $mnn1och1$ (MATa/ α ; $his3\Delta1/his3\Delta1$; $leu2\Delta0/leu2\Delta0$; $met15\Delta0/MET15$; $LYS2/lys2\Delta0$; $ura3\Delta0/ura3\Delta0$; $och1:kanMX4/OCH1$; $MNN1/mnn1:kanMX4$), and DIP- $mnn1pmr1$ (MATa/ α ; $his3\Delta1/his3\Delta1$; $leu2\Delta0/leu2\Delta0$; $LYS2/lys2\Delta0$; $met15\Delta0/MET15$; $ura3\Delta0/ura3\Delta0$; $pmr1:kanMX4/PMR1$; $MNN1/mnn1:kanMX4$). The diploids were selected on CSM-met-lys and sporulated as described (Giaever, 2002). The ascus of the sporulated diploids were digested with Zymolyase 100T (US Biological) and random spore analysis was conducted as described in Current Protocols in Molecular Biology. Spores from DIP- $mnn1mnn4$ and DIP- $mnn1pmr1$ were plated on YPD, while spores from DIP- $mnn1och1$ were plated on YPD + 0.3M KCl. The resulting haploid spores were selected for mating-type by crossing with parental strains (BY4710 and BY4711) on SD, and auxotrophic requirements by replica plating onto CSM-met and CSM-lys. The $\Delta och1$ mutation was screened by slow growth on YPD at 37°C, while all ORF knockouts were screened and verified using colony PCR. The final double mutants were $\Delta mnn1\Delta mnn4$ -9 (MATa, $his3\Delta1$, $leu2\Delta0$, $met15\Delta0$, $ura3\Delta0$, $mnn1:kanMX4$, $mnn4:kanMX4$), $\Delta mnn1\Delta och1$ -3 (MATa, $his3\Delta1$, $leu2\Delta0$, $lys2\Delta0$, $ura3\Delta0$, $mnn1:kanMX4$, $och1:kanMX4$), and $\Delta mnn1\Delta pmr1$ -5 (MATa, $his3\Delta1$, $leu2\Delta0$, $lys2\Delta0$, $ura3\Delta0$, $mnn1:kanMX4$, $pmr1:kanMX4$).

[0203] The haploid double mutants, $\Delta mnn1\Delta mnn4-9$ and $\Delta mnn1\Delta och1-3$, were mated to form the diploid strain DIP- $mnn1mnn4och1$ (MATa/ α ; his3 $\Delta 1$ /his3 $\Delta 1$; leu2 $\Delta 0$ /leu2 $\Delta 0$; met15 $\Delta 0$ /MET15; LYS2/lys2 $\Delta 0$; ura3 $\Delta 0$ /ura3 $\Delta 0$; $mnn1:kanMX4/mnn1:kanMX4$; $mnn4:kanMX4/MNN4$; OCH1/ $och1:kanMX4$), followed by sporulation, digestion, random spore analysis, and selection as before. The final, triple mutant haploids were: $\Delta mnn1\Delta mnn4\Delta och1-1-2$, $\Delta mnn1\Delta mnn4\Delta och1-1-3$, and $\Delta mnn1\Delta mnn4\Delta och1-2-5$ (MATa, his3 $\Delta 1$, leu2 $\Delta 0$, met15 $\Delta 0$, ura3 $\Delta 0$, $mnn1:kanMX4$, $mnn4:kanMX4$, $och1:kanMX4$); $\Delta mnn1\Delta mnn4\Delta och1-2-4$ and $\Delta mnn1\Delta mnn4\Delta och1-2-6$ (MAT α , his3 $\Delta 1$, leu2 $\Delta 0$, lys2 $\Delta 0$, ura3 $\Delta 0$, $mnn1:kanMX4$, $mnn4:kanMX4$, $och1:kanMX4$); and $\Delta mnn1\Delta mnn4\Delta och1-2-2$ (MATa, his3 $\Delta 1$, leu2 $\Delta 0$, lys2 $\Delta 0$, ura3 $\Delta 0$, $mnn1:kanMX4$, $mnn4:kanMX4$, $och1:kanMX4$). All clones were verified for KanMX knockout of each ORF by PCR at the 3' and 5' ends (Giaevet, 2002).

[0204] The two haploid triple mutants, $\Delta mnn1\Delta mnn4\Delta och1-2-4$ and $\Delta mnn1\Delta mnn4\Delta och1-2-5$ were mated to form the diploid triple mutant, $\Delta mnn1\Delta mnn4\Delta och1-DIP$ (MATa/ α ; his3 $\Delta 1$ /his3 $\Delta 1$; leu2 $\Delta 0$ /leu2 $\Delta 0$; met15 $\Delta 0$ /MET15; LYS2/lys2 $\Delta 0$; ura3 $\Delta 0$ /ura3 $\Delta 0$; $mnn1:kanMX4/mnn1:kanMX4$; $mnn4:kanMX4/mnn4:kanMX4$; $och1:kanMX4/och1:kanMX4$).

Preparation of Immune Sera:

[0205] *Antibody against yeast zymosan.* Rabbits were immunized using a yeast cell wall extract, zymosan A, which is composed of β -glucan and α -mannan of wild-type yeast. Two rabbits were immunized with 200 μ g of zymosan A containing CFA at week 0, and boosted with 100 μ g of antigen containing IFA at weeks 2 and 4, and thereafter every 4 weeks. Immune sera were collected at week 5, and 1 week after each booster. The resulting sera were pooled and purified with Protein A-agarose.

[0206] *Antibodies against yeast glycoproteins.* Synthetic peptides were used to generate immune sera in rabbits against the yeast proteins of interest. An accelerated immunization schedule was used, with an initial immunization of 200 μ g at the outset and 100 μ g immunizations at week 2, 4, 6 and 10, and test bleeds taken at week 5, 7, 8, and 11. The sequences of the peptides used are listed: CDSLESITDSLNLQSLT (SEQ ID NO:10) for PST1, CDSIKKITGDLNMQE (SEQ ID NO:11) for ECM33, CAVNGVTSKSALESIFP (SEQ ID NO:12) for Gp38, CTPKEQLSFVMNLYYEKSGGSKSD (SEQ ID NO:13) and

CPATGKYWSAATELPPTPNG (SEQ ID NO:14) for Gas1, CPAMPSAASAYFTSGAGSPMGTGIATQQS (SEQ ID NO:15) and CEIKGNAFFNSESGERFYIRGVQPGGS (SEQ ID NO:16) for Gas5, and CSGPQSYQKLDFTNVGFTGS (SEQ ID NO:17) and CEVGDRVWFSGKNAPLADY (SEQ ID NO:18) for YJL171c. Antigen specific antibodies were purified by immunoaffinity chromatography using the antigenic peptide coupled SulfoLink Gel (Pierce).

[0207] *Antibody specific to α 1,3-linked mannose.* Immune sera specific to α 1,3-linked mannose residues was prepared as previously described, with slight variations (Raschke et al., *J. Biol. Chem.* 248(13):4660-6, 1973 and Ballou, *J Biol Chem.* 245:1197-1203, 1970). Briefly, log phase INVSc1 cells suspended in 0.9% NaCl at 3.0×10^7 cells/ml were heat-killed at 70°C for 90 min. Rabbits were injected with this suspension in the marginal ear vein three times a week at 0.25ml, 0.5ml, 0.75ml and 1.0ml for week 1, week 2, week 3, and week 4, respectively. Three days after the last injection, the rabbits were bled. The resulting sera were then adsorbed to Δ mn1 yeast cells. Each 5 ml of sera was incubated with 5 g (wet weight) of heat-killed Δ mn1 yeast in 50 ml of 0.9% NaCl. Sodium azide was added to 0.2% and the suspension was incubated overnight at room temperature with end-over-end mixing. The cells were removed by centrifuging at 6,000g for 20 min and the adsorption was repeated. The final sera were dialyzed against 0.9% NaCl and sterilized with a 0.2 micron filter.

Determination of Carbohydrate Types on Yeast Whole Cells.

[0208] *Materials.* 2G12 monoclonal antibody was purchased from Polymun Scientific Inc. (Forschung GmbH). Protein A–Sepharose 4B conjugate was purchased from Invitrogen (Carlsbad, CA). Goat anti-Rabbit IgG HRP was purchased from Jackson ImmunoResearch (West Grove, PA).

[0209] *Immunofluorescence.* Log-phase Δ mn1 Δ mn4 Δ och1-DIP and INVScI cells grown at 30°C in YPD + 0.3 M KCl were fixed with 1 ml fresh 4% paraformaldehyde for 1 h and pelleted by centrifugation. The cells were washed once with PBS and transferred to polylysine-coated slides. After air-drying, the cells were incubated with 0.5% SDS at room temperature (RT) for 15 min. For single staining, rabbit anti-Zymosan antibody was diluted to 10 μ g/ml and α 1,3-linked mannose anti-serum was used at a 1:500 dilution. These antibodies were incubated with the cells for 1 h at RT. The slides were

washed and incubated for 1 h at RT with the secondary antibodies, Alexa Fluor 568-conjugated goat anti-rabbit IgG (Molecular Probe, USA). Cells were washed with PBS and mounted onto slides. For co-localized staining, the anti-Zymosan and 2G12 antibodies were diluted to 10 µg/ml and 20 µg/ml, respectively, and incubated with the cells for 1 h. The slides were washed, probed, and mounted onto slides as above, using Alexa Fluor 568-conjugated goat anti-human IgG and Alexa Fluor 488-conjugated goat anti-rabbit IgG as secondary antibodies. Images were obtained using a Zeiss Axioskope fluorescence microscope (McBain, USA).

[0210] Whole-cell ELISA. Enzyme-linked immunosorbent assays (ELISA) were performed on heat-killed, whole yeast cells. Log-phase $\Delta mnn1 \Delta mnn4 \Delta och1$ -DIP and INVScI cells were heat-killed at 70°C for 1 h in 0.9% NaCl and diluted to 5.0×10^7 cells/ml in PBS for coating ELISA plates. HIV-1 gp120 from SF162 strain (ARRRP) and wild-type yeast mannan (Sigma) were diluted to 5 µg/ml in 50 mM carbonate buffer, pH 9.5. One-hundred microliters of the antigens were incubated in each well overnight at RT. Wells were blocked with ELISA blocking buffer (PBS, 0.1% BSA, 0.02% thimerosal) and incubated for 2 hrs at RT in a humid chamber. After two dH₂O rinses, 2G12 and anti- $\alpha 1,3$ -mannose antibody were added at 10 µg/ml with a three-fold serial dilution in blocking buffer and incubated overnight at room temperature. Wells were washed with ELISA wash buffer (PBS, 0.05% Tween 20, 0.02% thimerosal) and incubated at 37°C for 1 h with a 1:10,000 dilution of goat anti-human IgG-HRP or goat anti-rabbit IgG-HRP (Jackson ImmunoResearch) for 2G12 or anti- $\alpha 1,3$ -mannose, respectively. Wells were washed as before, incubated with TMB for color development, and stopped with HCl. HRP activity was read by absorbance at 450nm using an EMax Microplate Reader (Molecular Devices).

[0211] Glycan profiling. Log-phase $\Delta mnn1 \Delta mnn4 \Delta och1$ -DIP cells grown at 30°C were diluted to $\sim 3.0 \times 10^7$ cells/ml in 0.9% NaCl. Cells were heat-killed at 70°C for 1 h. N-linked glycans from cell were released by PNGase F digestion. Released N-linked glycans were permethylated and analyzed by Maldi-TOF Mass Spectrometry (MS) by Glycotechnology Core Resource at UCSD.

Identification and Confirmation of 2G12 Cross-Reactive Glycoproteins

[0212] Western blot. Yeast cells from strains of INVScI, $\Delta mnn1$, $\Delta mnn4$, $\Delta och1$, $\Delta mnn1 \Delta mnn4$, $\Delta mnn1 \Delta och1$ and $\Delta mnn1 \Delta mnn4 \Delta och1$ were grown at 30°C in YPD + 0.3 M KCl to an OD₆₀₀ of 3.0. Yeast cells were centrifuged at 14,000g for 2 min. The culture

supernatant was subjected to acetone precipitation, by incubating with two-volumes of ice-cold acetone for 30 min at -80°C. The samples were centrifuged at 14,000g for 15 min and the supernatant was removed. After air-drying for 15 min, the pellet was boiled for 5 min with 2x Laemmli SDS-sample buffer at 1/10 the original sample volume.

[0213] The yeast cell lysate was prepared by lysing in RIPA buffer. Briefly, each 1 ml of cell pellet is resuspended in 100 µl ice-cold RIPA buffer (150 mM NaCl, 10 mM Tris pH7.4, 1 mM EDTA pH 8.0, 1% Triton X-100, 1% DOC, 0.1% SDS, 1 mM PMSF, 1 µg/ml Aprotinin, and 1 µg/ml Leupeptin), and cell walls were broken by vortexing with acid-washed glass beads (Sigma) for 10 min. The resulting lysate was centrifuged at 13,000g for 30 min to remove any debris and the protein concentration of each was diluted to 0.3 mg/ml in dilution buffer (50 mM Tris pH7.4, 1 mM EDTA pH8.0, 1 µg/ml Aprotinin, 1 µg/ml Leupeptin, and 0.1% SDS). Laemmli SDS-buffer is added to 1x and the samples containing 0.24 mg/ml protein were boiled for 5 min.

[0214] The supernatant and cell lysate fraction were subjected to SDS-PAGE separation on 4-20% gradient gels and transferred to nitrocellulose membranes at 250 mA for 2 hrs. Western blotting was conducted using the primary antibodies including MAb 2G12 and polyclonal antibodies raised in rabbits as described above followed by the secondary antibodies, goat anti-human IgG-HRP and goat anti-rabbit IgG-HRP (Jackson ImmunoResearch), at the indicated dilutions. Chemilluminiscent detection was employed using ECL (GE Biosciences).

[0215] *Partial purification of 2G12 cross-reactive proteins.* The subcellular localization of endogenous yeast proteins was estimated using differential centrifugation as described in Current Protocols in Cell Biology. Log-phase cells from INVSc1 and $\Delta mnn1 \Delta mnn4 \Delta och1$ -DIP strains were grown as described above to an OD₆₀₀ of 3.0. Each 1 ml of cell pellet was resuspended in 100 µl sucrose lysis buffer (0.4 M sucrose, 2 mM EDTA, 25 mM imidazole, pH7.0, 1 mM PMSF, 1 µg/ml Aprotinin, and 1 µg/ml Leupeptin). An equal volume of acid-washed glass beads (Sigma) were added and cell walls were broken with 15 min of high-speed vortexing alternated with ice-bath incubation. The lysates were centrifuged at 500g for 5 min to pellet the unlysed cells and large aggregates, and the supernatant was centrifuged at 22,000g for 30 min. The resulting supernatant that contains cytoplasmic, Golgi complex and endosomal proteins was saved for analysis. The pellet, containing mostly vacuolar, nuclear, ER and plasma membrane proteins, was resuspended in a similar volume of PBS + 1.0% Triton to solubilize the

membrane proteins. The fraction was centrifuged as above to segregate the soluble and insoluble membrane fractions.

[0216] *Lectin affinity chromatography.* For Concanavalin A (ConA) lectin affinity chromatography, 9.5 ml of Triton-soluble membrane fraction was incubated overnight at 4°C with 1 ml of ConA-agarose (Vector Labs) in 5 ml of Triton wash buffer (20 mM Tris, pH7.0, 50 mM NaCl, 10 mM CaCl₂, and 1% Triton X-100). The column was washed extensively with Triton wash buffer. To elute ConA-bound proteins, 5 ml of Triton wash buffer including 0.5 M methyl manno-pyranoside was added to the column followed by 5 ml of Triton wash buffer including 0.2 mM EDTA and 1.0% SDS.

[0217] For purification from the culture supernatant, 300 ml of $\Delta mnn1 \Delta mnn4 \Delta och1-2$ -5 yeast supernatant with 10 mM CaCl₂ was loaded on a 5 ml ConA-agarose column (Vector Labs) and washed extensively with ConA wash buffer (20 mM Tris, pH7.0, 50 mM NaCl, 10 mM CaCl₂), followed by ConA wash buffer including 0.5M methyl manno-pyranoside. To elute ConA-bound proteins, 5 ml of 2x SDS loading buffer was added to the column and boiled for 5 min.

[0218] *Two-dimensional gel electrophoresis and mass spectrometry.* ConA-agarose purified protein fractions were precipitated with TCA and resolubilized in 200 μ l Urea/Thiourea Rehydration Solution (7 M urea, 2 M thiourea, 2% CHAPS, 60 mM DTT; 1% Ampholytes pH 3-10; 0.01% Bromophenol blue). IPG ZOOM 4-7 IPG strips (Invitrogen) were rehydrated overnight with 120 μ l of sample and 20 μ l of rehydration buffer (7 M urea, 2 M thiourea, 2% CHAPS, 60 mM DTT, 1% Ampholytes pH 3-10, 0.01% Bromophenol blue) for the sypro ruby gel, and 80 μ l of sample and 60 μ l of rehydration buffer for the Western blot. The strips were loaded on the ZOOM IPG Runner (Invitrogen) and focused as follows: 200V for 30 min, 450V for 25 min, 750V for 25 min, and 2000V for 60 min.

[0219] For the second dimension, the IPG strips were reduced twice in LDS-equilibration buffer containing DTT for 15 min on an orbital shaker. Then, the strips were alkylated in LDS-equilibration buffer containing iodoacetamide for 15 min and electrophoresed on the IPG ZOOM system using 4-12% gradient acrylamide gels with an MES running buffer. The gels were electrophoresed until the dye front migrated to approximately 0.3 cm from the bottom of the gel.

[0220] For Sypro Ruby staining, the gel was fixed in 40% methanol, 10% acetic acid for one hour and stained overnight with SYPRO Ruby fluorescent stain. The gel was

destained for 120 min in 10% methanol, 6% acetic acid with one change of destaining solution and imaged on the Molecular Imager PharosFX laser imager with the following parameters: and excitation wavelength of 532 nm, an emission wavelength of 605 nm, and a resolution of 100 μ m.

[0221] For the Western blot, the gel was soaked in transfer buffer (Invitrogen) immediately following the second dimension run. Proteins were transferred to nitrocellulose and probed with 2G12 at 1 μ g/ml.

[0222] The proteins on 2D blot detected by 2G12 were matched to the spots stained with Sypro Ruby. The matching spots were sliced and subjected to in gel digestion with trypsin. The digested peptides were analyzed by Nano-LC MS/MS. The above experiments were performed by Proteomic Research Services Incorporated.

[0223] **Immunoprecipitation.** Culture supernatant and cell lysate were prepared from log-phase Δ mnn1 Δ mnn4 Δ och1-2-5 as described above. These samples were pre-cleared by incubation with protein A-Sepharose 4B conjugate at RT for 2 hrs. For antigen-antibody complex formation, 10 μ g of 2G12 was added to either cell lysate or media sample, and incubated on a shaker at RT for 1 h. After incubation, 20 μ l of protein A-Sepharose 4B conjugate was added to antigen-antibody complex, and incubated again at RT for 1 h on a shaker. The samples were spun 10,000g for 5 min and the supernatant was aspirated. The beads were washed twice with 800 μ l of PBS and the bound proteins were eluted by boiling with 40 μ l of 1x SDS sample buffer. All samples were loaded onto 4-20% gradient SDS-PAGE gels and blotted for Western analysis using antibodies against the identified yeast glycoproteins including ECM33, PST1, GP38, YJL171C, Gas1, and Gas5.

Bioinformatic Analyses of N-linked and O-linked glycosylation sites

[0224] **Analysis of N-linked glycosylation sites.** N-linked glycosylation sites in the yeast *S. cerevisiae* genome and on the glycoproteins identified in the study were analyzed using software that we developed recently.

[0225] **Analysis of O-linked glycosylation sites.** O-linked glycosylation sites on some proteins in the *S. cerevisiae* genome and on the glycoproteins identified in the study were analyzed using software NetOGlyc 3.1 that was developed by Center for Biological Sequence Analysis (CBS) at Technical University of Denmark, which is available at <http://www.cbs.dtu.dk/services/netoglyc>.

Cloning, Expression, characterization of gp120 produced in yeast triple mutant.

[0226] **Reagents.** The pYES2/CT vector was obtained from Invitrogen and the pJR-FLsyngp120 and pYU-2 plasmids were obtained from the AIDS Research and Reference Program (ARRRP). Yeast Nitrogen Base Without Amino Acids and -Ura Dropout Mix were obtained from U.S. Biological. A rabbit polyclonal antibody against gp120 from HIV-1_{IIIB} (anti-gp120-IIIB) was obtained from Virostat Inc. Another rabbit polyclonal antibody against HIV-1_{YU-2} (anti-gp120-YU2) was produced in house using a synthesized peptide, CEQMHEDIISLWDQSLK (SEQ ID NO:19), and affinity purified using the peptide coupled to Sulfolink Gel (Pierce).

[0227] **PCR cloning of the α -mating factor and gp120.** The signal sequence of the *S. cerevisiae* alpha-mating factor (MF α) was added upstream of the multiple cloning site (MCS) of the pYES2/CT vector. The MF α was cloned from total yeast DNA (extracted from INVSc1) by PCR using the primers MF α 1-Kpn-5 (5'-ATGGTACCAAAGAATGAGATTCCTTCAATT-3' (SEQ ID NO:20)) and MF α 1-Bam-3 (5'-ATGGATCCAGCTTCAGCCTCTCTTTTATC-3' (SEQ ID NO:21)), with KpnI and BamHI sites added, respectively. This fragment contains the Kex2 cleavage site to remove the signal sequence from gp120. PCR was conducted using T4 DNA Polymerase (Sigma) according the manufacturer's protocol. PCR products were purified, digested with BamHI and KpnI, and ligated into a similarly digested pYES2/CT plasmid using standard molecular biology techniques. The resulting plasmid was pYES2/CT- α .

[0228] The coding sequence for JR-FL was amplified from pJR-FLsyngp120 by PCR using the primers JRFL-Eco-5 (5'-TGGAATTCTGTGGGTGACTGTATACTAT (SEQ ID NO:22)) and JRFL-Xba-3 (5'-TATCTAGACCCCACAGCGCGCTTCTCCCT-3' (SEQ ID NO:23)), with EcoRI and XbaI sites added, respectively. The coding sequence for YU-2 was amplified from pYU-2 using the primers YU2-Eco-5 (5'-TGGAATTCTGTTGTGGGTCACAGTCTATTAT-3' (SEQ ID NO:24)) and YU2-Xba-3 (5'-ATTCTAGATTCTCTTTGCACCACTCTTCT-3' (SEQ ID NO:25)), with EcoRI and XbaI sites added, respectively. The gp120 PCR products were cloned into the pYES2/CT- α using digestion with EcoRI and XbaI, as above. The final plasmids were pYES2/CT- α -JRFL (pJRFL-gp120) and pYES2/CT- α -YU2 (pYU2-gp120).

[0229] **Lithium acetate transformation of $\Delta mnn1\Delta mnn4\Delta och1-2-5$ yeast.** The plasmids pJRFL-gp120 and pYU2-gp120 were used to transform $\Delta mnn1\Delta mnn4\Delta och1-2-5$

yeast cells. Briefly, 1.5 ml of an overnight culture of yeast grown in YPD + 0.3 M KCl was pelleted. All but ~ 50 μ l of the supernatant was removed, and the cells were resuspended. Then, 2 μ l of 10 mg/ml single stranded salmon sperm DNA (Sigma) was added, followed by 1 μ g of pJRFL-gp120 or pYU2-gp120. After vortexing, 500 μ l of PLATE mixture (40% PEG 4000, 100 mM LiAc, 10 mM Tris, pH 7.5, 1 mM EDTA) and 20 μ l of 1M DTT were added, and the mixtures were incubated for 4 hrs at RT. The cells were heat shocked for 10 min at 42°C, pelleted, and resuspended in 100 μ l of sterile dH₂O. Cells were plated onto –ura/glucose (2% glucose, 0.67% yeast nitrogen base without amino acids, 0.2% -Ura Dropout Mix, 2% Bacto-Agar). Plates were incubated at 30°C until colonies appeared. PCR was used directly on the colonies to verify the presence of each plasmid using the primers MF α 1-Kpn-5 and JRFL-Xba-3 for pJRFL-gp120 transformant, and MF α 1-Kpn-5 and YU2-Xba-3 for pYU2-gp120.

[0230] *Expression of gp120 proteins in $\Delta mnn1\Delta mnn4\Delta och1$ yeast.* Triple mutant yeast cells transformed with pJRFL-gp120 and pYU2-gp120 were grown to log phase in –ura/glucose media (2% glucose, 0.67% Yeast Nitrogen Base without amino acids, 0.2% -Ura Dropout Mix). The cells were centrifuged and washed twice with –ura/galactose media (2% galactose, 0.67% Yeast Nitrogen Base without amino acids, 0.2% -Ura Dropout Mix), and resuspended in –ura/galactose at an OD₆₀₀=0.4. The cells were grown at 30°C and samples of the media were taken at different time points to test for protein induction.

Purification of Yeast Expressed Glycoproteins

[0231] *Ammonium Sulfate Precipitation.* PST1 protein in the yeast culture supernatant from $\Delta mnn1\Delta pmr1$ and gp120 expressed in transformed $\Delta mnn1\Delta mnn4\Delta och1$, were collected and centrifuged (Beckman Coulter, Allegra™ 64R, rotor F0650) at 3000g, 4°C for 10 min to remove any insoluble particles. The supernatant was transferred to a clean beaker, 10 mM EDTA was added and the pH was adjusted to pH 7.4 using 1M Tris Base. Ammonium sulfate was slowly added while stirring until a final concentration of 55% saturation at 4°C was reached. The solution was stirred for additional 30 min and centrifuged at 15,000 x g at 4 °C for 30 min. The pellet was resuspended in 2 ml PBS and the insoluble material was removed by centrifugation at 15,000 x g at 4°C for 1 h. This material was designated AS55 precipitate. Ammonium sulfate was slowly added to the 55% AS supernatant while stirring until a final concentration of 90% saturation at 4°C was reached. The pellet was resuspended in 2 ml PBS and the insoluble material was removed

by centrifugation at 15,000 x g at 4°C for 1 h. This material was designated AS90 precipitate. The AS55 and AS90 precipitates were desalted into 5 mM Sodium Phosphate, pH 7.5 using 5 ml HiTrap desalting column (GE Life Sciences) and protein concentration was adjusted to 1 mg/ml.

[0232] *Immunoaffinity purification of yeast expressed gp120.* Anti-gp120-IIIB was conjugated to cyanogen bromide-activated Sepharose 4B (Sigma) according to the manufacturer's protocol at 5mg antibody per 1ml gel. After extensive washing of the columns with PBS, the AS55 sample from $\Delta mnn1 \Delta mnn4 \Delta och1$ cells was incubated with the immunoaffinity column for 2 hours at room temperature. The flowthrough was collected and the columns were washed again with PBS. Bound proteins were eluted with 100mM glycine, pH2.5 in 1ml fractions containing 50 μ l of 1M Tris, pH9.5. These fractions were tested for gp120 by Western blot using 2G12 and anti-gp120-YU2. The gp120 containing fractions were pooled and designated AS55-AP.

[0233] *Cation-Exchange Chromatography.* Separate chromatography was performed for different samples, with AS55-AP used for gp120 purification and AS90 used for PST1 purification. A 5 ml HiTrap SP FF column (GE Life Sciences) was equilibrated with 5 mM Sodium Phosphate, pH 7.5. The samples were loaded at 1 ml/min onto the column and washed with 15 ml of 5 mM sodium phosphate, pH 7.5. Proteins were eluted with an 80-ml gradient of 0-500 mM NaCl in 5 mM sodium phosphate pH 7.5 at a flow rate of 1 ml/min. Fractions of 2 ml were collected throughout, and 3 μ l samples were tested for the presence of desired glycoprotein employing a dot blot assay. Fractions exhibiting a significant amount protein were pooled and designated GP-IEC-55 and GP-IEC-90. These fractions were further purified using ConA purification of glycoproteins as described earlier. The proteins were eluted into 1M methyl mannopyranoside with a 60 min. incubation.

[0234] *Gel Filtration Chromatography of Glycoproteins.* As a final step, the two GP-IEC-55 and GP-IEC-90 fractions were desalted into the loading buffer (50 mM Tris, 150 mM NaCl, 5 mM 2-bME, pH 7.5) and concentrated by Amicon® Centricon® 30 (Millipore) to a final concentration of ~1 mg/ml. A 150 ml Sephacryl S-HR-200 (GE Life Sciences) was equilibrated with the same buffer and the GP-IEC pools were loaded onto the column. Proteins were eluted with the loading buffer at a flow rate of 0.3 ml/min, 2 ml fractions were collected and 3 μ l samples of each fraction were tested for the presence of a glycoprotein employing the dot blot assay. Fractions exhibiting significant amount of the glycoprotein were pooled and collectively designated GP-SHR200-55 and GP-SHR200-90.

The following standards were used and co-eluted from the filtration column: IgG (MW150kD), AP (MW 140kD), Egg white avidin (MW68) and Streptavidin (MW 53).

[0235] Glycan profiling. The GP-SHR200-55 and GP-SHR200-90 fractions, containing purified gp120 PST1, respectively, were digested with PNGase F to release all N-linked glycans. These glycans were permethylated and analyzed by Maldi-TOF Mass Spectrometry (MS) by Glycotechnology Core Research Facility at UCSD.

Immunization and preparation of antibodies specific to α 1,2-linked terminal mannose

[0236] Raising antibody against α 1,2-linked mannose using yeast whole cell. Immune sera specific to α 1,2-linked mannose residues was prepared using the triple mutant (Raschke et al., *J. Biol. Chem.* 248(13):4660-6, 1973; and Ballou, *J Biol Chem.* 245:1197-1203, 1970). Briefly, log phase Δ mn1 Δ mn4 Δ och1-DIP suspended in 0.9% NaCl at 3.0×10^7 cells/ml were heat-killed at 70°C for 90 min. Rabbits were injected with this suspension in the marginal ear vein three times a week at 0.25ml, 0.5ml, 0.75ml and 1.0ml for week 1, week 2, week 3, and week 4, respectively. Three days after the last injection from week 2, 3, and 4, the rabbits were bled. After four weeks of rest, the rabbits were injected with the same suspension in the marginal ear vein three times a week at 0.5ml, 0.75ml and 1.0ml for week 9, week 10, and week 11. A fourth, final bleed was collected 3 days after the final injection. The resulting sera were tested for gp120 binding by ELISA and Western blot.

[0237] Raising antibody against α 1,2-linked mannose using yeast cell extract. Immune sera specific to α 1,2-linked mannose residues was prepared using the membrane fractions of the triple mutant. Log-phase cells from Δ mn1 Δ mn4 Δ och1-DIP strains were lysed in sucrose lysis buffer (0.4 M sucrose, 2 mM EDTA, 25 mM imidazole, pH7.0, 1 mM PMSF, 1 μ g/ml Aprotinin, and 1 μ g/ml Leupeptin) using acid-washed glass beads (Sigma). Lysates were centrifuged at 500g for 5 min to pellet the unlysed cells and large aggregates, and the supernatant was centrifuged at 22,000g for 30 min. The resulting pellet, containing mostly vacuolar, nuclear, ER and plasma membrane proteins, was resuspended in a similar volume of PBS + 1.0% Triton to solubilize the membrane proteins. The fraction was centrifuged as above to segregate the Triton-soluble and -insoluble membrane fractions. Four groups of two rabbits were immunized with these fractions, one group with the Triton-soluble fraction of wild-type (INVSc1), two groups with the Triton-soluble fraction of the triple mutant, and one group with the Triton-insoluble fraction of the triple mutant.

An accelerated immunization schedule was used, with an initial immunization of 200 μ g at the outset and 100 μ g immunizations at week 2, 4, 6 and 10, and test bleeds taken at week 5, 7, 8, and 11. The resulting sera were tested for gp120 binding by ELISA and Western blot.

[0238] *Adsorption of antisera to heat-killed, whole cell yeast.* In order to enrich for antibodies specific to α 1,2 linked mannose residues, any sera that showed gp120 binding were then adsorbed to Δ och1 and INVSc1 (WT) yeast cells. Each 1 ml of sera was incubated with 1.0-2.0 g (wet weight) of heat-killed Δ och1 and/or WT yeast in 5 ml of 0.9% NaCl. Sodium azide was added to 0.2% and the suspension was incubated overnight at room temperature with end-over-end mixing. The cells were removed by centrifuging at 6,000g for 20 min and the adsorption was repeated. The final sera were dialyzed against 0.9% NaCl and sterilized with a 0.2 micron filter. Then, the antibodies were purified on Protein A-agarose (Invitrogen) and tested again for gp120 binding by ELISA and Western blot.

[0239] *Affinity purification of antibodies against α 1,2-linked mannose residues.* In order to further purify antibodies specific to α 1,2 linked mannose residues, the sera from any adsorbed antibodies that showed gp120 binding by ELISA were purified using ConA-eluted glycoproteins from the triple mutant. Briefly, the culture supernatant and cell lysate from Δ mn1 Δ mn4 Δ och1-2-5 cells were purified using ConA lectin affinity chromatography as described earlier. The proteins from each fraction (supernatant and lysate) were separately conjugated to cyanogen bromide-activated Sepharose 4B (Sigma) according to the manufacturer's protocol at 10mg total protein per 1ml gel. After extensive washing of the columns with PBS, 10-20 ml of selected sera were filter sterilized and incubated with the media for 2 hours per 5ml at room temperature. The flowthrough was collected and the columns were washed again with PBS. Bound antibodies were eluted with 100mM glycine, pH2.5 in 1ml fractions containing 50 μ l of 1M Tris, pH9.5. After dialysis against PBS, these antibodies were tested for gp120 binding by ELISA and Western blot.

Analysis of Immune Sera for gp120 Reactivity and Pseudovirus Neutralization

[0240] *ELISA against gp120.* Enzyme-linked immunosorbent assay (ELISA) was performed to test for gp120 specificity of anti- α 1,2-linked mannose sera. Different HIV-1 gp120 proteins (from various sources, including Dr. Doms' laboratory, ARRRP and commercial companies) were diluted to 3-5 μ g/ml in 50 mM carbonate buffer, pH 9.5. One-hundred microliters of the antigens were incubated in each well overnight at RT.

Wells were blocked with ELISA blocking buffer (PBS, 0.1% BSA, 0.02% thimerosal) and incubated for 2 hrs at RT in a humid chamber. After two dH₂O rinses, 2G12 and anti-sera against $\Delta mnn1 \Delta mnn4 \Delta och1$ -DIP were added at varying dilutions in blocking buffer and incubated overnight at room temperature. Wells were washed with ELISA wash buffer (PBS, 0.05% Tween 20, 0.02% thimerosal) and incubated at 37°C for 1 h with a 1:10,000 dilution of goat anti-human IgG-HRP or goat anti-rabbit IgG-HRP (Jackson ImmunoResearch). Wells were washed as before, incubated with TMB for color development, and stopped with HCl. HRP activity was read by absorbance at 450nm using an EMax Microplate Reader (Molecular Devices). SF162 gp120 expressed in CHO cells was obtained through the NIH AIDS Research and Reference Reagent Program (ARRRP). ADA, JR-FL, and YU2 gp120 expressed in 293T cells were obtained from Dr. Doms at University of Philadelphia.

[0241] In a similar method used above, the binding affinity of gp120 proteins expressed in mammalian cells will be compared to the triple mutant yeast expressed by quantitative ELISA. An equal quantity of gp120-yeast (expressed in $\Delta mnn1 \Delta mnn4 \Delta och1$) and gp120-293 (expressed in 293T cells), from different strains, will be used to coat microwell plates. Then, 2G12, anti-gp120-IIIB, and anti-gp120-YU2 will be used to probe. By capturing equivalent amounts of protein and serially diluting the binding antibodies, the binding affinities 2G12 toward each gp120 will be calculated and compared.

[0242] *Neutralization assay.* The 293T/17 cells were purchased from the American Type Culture Collection (Manassas, VA) and maintained in Dulbecco's modified Eagle's medium (DMEM) with 10% fetal bovine serum. The plasmid pNL4-3.luc.R-E- was purchased from Dr. Landau's laboratory. The various HIV-1 env gp160 plasmid used to produce pseudoviruses were obtained from NIH AIDS Research Reference Reagent Program (NIH, ARRRP). The U87.CD4.CCR5 and U87.CD4.CXCR4 cells line expressing CCR5 and CD4 used for pseudovirus infectivity were a gift of Dr. Doms at University of Philadelphia.

[0243] For pseudoviruses, a 10 cm dish of 70–80% confluent 293T/17 cells were cotransfected with gp160 env-expressing plasmids and with the complementing viral genome-reporter gene vector, pNL4-3.Luc.E⁻R⁻ using calcium phosphate method according to manufacturer's instruction (Invitrogen). The culture supernatant containing recombinant viruses pseudotyped with Env proteins were harvested 48 h posttransfection, centrifuged, filtrated through 0.45 μ m filters, and frozen at -80°C. To measure the

pseudovirus infectivity, aliquots of harvested supernatant were used to infect 10,000 cells per 96 well plates of CCR5 (U87.CD4.CCR5) or CXCR4 (U87.CD4.CXCR4) expressing cell lines. The luciferase activity from a single round infected cells was measured after 72 h of cell culture using the Bright-Glo Luciferase Assay System reagent (Promega, Madison, WI) and Veritas microplate luminometer (Turner Biosystems) equipment according to manufacturer. Mean relative light units (RLU) for triplicate wells were determined and pseudovirus stock $\geq 2 \times 10^5$ rlu/ml was used for subsequent neutralization assay.

[0244] For the neutralization assay, animal serum or purified antibodies were assayed for neutralizing antibody (NAb) activity against HIV-1 virus using a single-round pseudotype reporter assay. Briefly, CCR5 (U87.CD4.CCR5) or CXCR4 (U87.CD4.CXCR4) express cell lines cells were plated at 10,000 cell/well on 96 flat bottom plate and cultured overnight. For testing with sera, samples were heat-inactivated for 45 minutes at 56°C before use. The next day and on a separated 96 well flat bottom plate, 25 μ l of appropriately diluted pseudovirus suspensions were mixed in triplicate with 25 μ l of serially diluted antibody sample (this brings the final antibody concentration to 200 μ g/ml or 50 μ g/ml for purified antibody or to 1:5 dilution for plasma sample) and incubated for 1 h at 37°C. After incubation of antibody/virus mixture, the media was removed from pre-incubated plate cells; the antibody mixture was added to cells and incubated for four hours at 37°C in CO₂ incubator. After incubation, the mixture was then removed from the cells, and replaced by fresh medium. The cells were harvested 72h postinfection by adding equal volume (usually 50-100 μ l) of Bright-Glo Luciferase Assay System reagent (Promega, Madison, WI) to culture cells and luciferase activity (in RLU) was measured by using a Veritas microplate luminometer (Turner Biosystems) as prescribed by manufacturer. The percentage of inhibition was determined for each sample by averaging the luciferase activity measurement on tested sample wells (cell/antibody/virus), virus control (cell/virus), and cell control (cell only) wells using the following formula: $[(\text{virus control wells} - \text{test sample well}) / (\text{virus control well} - \text{cell control}) \times 100]$.

Example 2. A yeast glycoprotein PST1 cross-reacts with 2G12 in a *S. cerevisiae* mutant strain with pmr1 gene disrupted or both pmr 1 and mnn1 genes disrupted

[0245] Screening 2G12 cross-reactive protein in yeast mutant strains. The glycosylation pathway of *S. cerevisiae* has been extensively studied. The deletion of nearly

all ORFs in the *S. cerevisiae* genome (Winzeler et al., *Science* 285:901-906, 1999 and Giaever et al., *Nature*, 418:387-391, 2002), and the availability of each mutant strain, have provided an excellent opportunity to examine the changes in glycan profile in the yeast. Different mutants were first screened using 2G12 for possible cross-reactive glycoproteins in any of the mutants. As shown in Figure 10, cell lysates and culture media from five single mutants with deletion of enzymes that are involved in the process of carbohydrate in ER and Golgi including Mns1, Och1, Mnn9, Mnn1, and Pmr1 were tested. The results show that two faint bands at approximately 100 and 120 kDa were detected in the cell lysate of the $\Delta pmr1$ mutant, although the signals are weaker than that from the HIV-1 gp120 glycoprotein (293T-expressed) used as a positive control.

[0246] The PMR1 gene encodes a high affinity Ca^{2+}/Mn^{2+} P-type ATPase that is required for Ca^{2+} and Mn^{2+} transport into Golgi and involved in Ca^{2+} dependent protein sorting and processing (Rudolph et al., *Cell*. 58:133-145, 1989). Deletion of PMR1 gene is thought to cause the secretion of some glycoproteins directly from the ER, thereby bypassing the Golgi apparatus and all glycan processing therein (Rudolph et al., *Cell*. 58:133-145, 1989). This would result in the lack of a polymannose outer chain and terminal α -1,3-linked caps on N-linked glycans (Verostek et al., *Glycobiology* 5(7):671-81, 1995). HIV-1 gp120 gene, from the YU-2 strain, containing the *S. cerevisiae* alpha-mating factor signal sequence was used to transform the $\Delta pmr1$ yeast. As shown in the Figure 11A, an approximately 100 kDa protein in the media after 24 h culture was recognized by 2G12 in Western blots (herein dubbed YP100). In a parallel growth condition, the 2G12 reactive YP100 was shown in the media from transformed cells, but was not detectable in the media from non-transformed *pmr1* mutant cells after 3 to 9 days culture (Figure 11B). Several bands in the range of 90 to 160 kDa were shown in the non-transformed and transformed cells in this experiment but not shown in others. These results suggest that the gp120 plasmid transformation cause either an increase in expression or secretion of this protein.

[0247] The 2G12 reactive protein (YP100) in the gp120 transformed *pmr1* mutant was initially thought to be the gp120 glycoprotein due to the following evidence: a 2G12 detectable band was induced by the transformation of gp120 plasmid (Figure 2A, 11A), the band was detected in the gp120 transformed yeast cells but not in non-transformed cells (Figure 11B), it showed an expected size at 96 to 100 kDa of glycoprotein (Figure 2A and 11), it showed an expected Endo H digested protein size at ~60 kDa, a similar molecular weight to the gp120 without glycosylation (Figure 3). Further studies presented doubt that

the induced YP100 protein was indeed gp120 since inconsistent results were observed using antibodies against gp120 (data not shown). As shown in Figure 10, two faint bands at ~100 kDa and 120 kDa in the Δ pmr1 yeast cell lysate were recognized by 2G12. These results indicate that the 2G12 reactive 100 kDa band has the possibility to be either expressed exogenous HIV-1 gp120 or an unknown endogenous yeast protein.

[0248] Identification of a 2G12 cross-reactive protein in Pmr1 mutant. To determine the protein recognized by 2G12 was gp120 or a yeast glycoprotein, Yp100 was partially purified using lectin affinity chromatography (Figures 2B and 12) and gel filtration. Proteins in the purified material were identified using Nano-LC-MS/MS. Six peptides were identified to belong to an unknown yeast protein, PST1 (Fig. 13), although some other peptides matched gp120 and yeast proteins. The PST1 protein contains a signaling peptide and a GPI anchor at its carboxy terminus indicating that it is a cell wall or membrane protein and can be secreted. There are 15 potential N-linked glycosylation sites in the 400 amino acids of mature protein, which accounts for a 3.8% density, indicating that PST1 is heavily glycosylated.

[0249] To confirm that Yp100 is indeed PST1, a polyclonal antibody was raised in rabbits against a synthetic peptide and purified using antigen immunoaffinity chromatography. The partially purified Yp100 protein was separated by 2D followed by Coomassie blue staining and Western blot analyses. Figure 14A shows a predominant spot at molecular mass of 100 kDa and isoelectric point (PI) of 9-10. The same spot was identically recognized by 2G12 (Fig. 14B) and anti-PST1 polyclonal antibody (Fig. 14C). PST1 protein has a calculated PI of 9.25. All these results are consistent and suggest that the 2G12 cross-reactive glycoprotein is very likely the yeast glycoprotein PST1. These results present the first evidence of an endogenous yeast protein showing a binding affinity to the HIV-specific neutralizing antibody, 2G12.

[0250] To further confirm the 2G12 specificity of PST1, YP100 was digested with an enzyme, endo-L-N-acetylglucosaminidase H (Endo H), which specifically cleaves the high mannose type of glycans. As shown in Figure 15 by Commassie blue and anti-PST1 antibody staining, after digestion, the migration of PST1 protein changed from ~ 100 kDa to ~ 60 kDa (compared lanes 2 and 4 with lanes 3 and 5). While the PST1 protein was still recognizable by anti-PST1 after Endo H digestion, it was no longer recognized by 2G12 (lane 7), which suggest that the N-glycans on the PST1 protein are the epitopes being recognized by HIV-1 neutralizing MAb 2G12. The molecular weight (~60kDa) of PST1

after Endo H digestion in SDS-PAGE is higher than the calculated molecular mass (41 kDa) of amino acids from mature protein. This could be due to the O-linked glycosylation since PST contains potential O-linked glycosylation sites (Figure 33). These results further confirmed that the 2G12-reactive protein shown in figures 2 and 11 is actually a yeast glycoprotein PST1 with unknown function. This protein is induced by the transformation, of exogenous genes, at least, HIV-1 gp120 Env.

[0251] Generation of a double mutant and characterization of PST1 in the double mutant. The binding site of 2G12 on gp120 was identified as a conformation of terminal α 1,2-linked mannose residues found on a cluster of at least three N-linked glycans. The Δ pmr1 mutation in yeast is thought to bypass the Golgi, thus leaving only Man8 glycans on yeast proteins. However other evidence suggests that this mutation may only bypass the early Golgi compartment, or that the resulting change in Ca^{++} affects the efficiency of the enzymes found in the Golgi. With this in mind, one of the most debilitating glycosylation enzymes for 2G12 binding in the Golgi would be Mnn1p, which is responsible for all terminal α 1,3-linked mannose capping of α 1,2 residues (Nakajima et al, Proc Nat Acad Sci USA 72:3912-3916, 1975). In order to prevent this capping of PST1, a yeast double mutant strain, Δ mnn1 Δ pmr1, was created. Yeast deletion strains with ORF knockouts in the *MNN1* and *PMR1* genes were mated and sporulated to produce a Δ mnn1 Δ pmr1 haploid. Colony PCR was conducted for selection and verification of the Δ mnn1 and Δ pmr1 phenotypes (data not shown). Four unique primers per gene deletion confirmed that each ORF was completely deleted and replaced with the KanMX4 module.

[0252] Five double mutant strains (Δ mnn1 Δ pmr1) were screened using Western blot with 2G12. As expected, the ~ 100 kDa band seen in the previous experiments (see Figure 10) was detected by 2G12 in all five mutant strains (Fig. 16). In fact, there appears to be an enhancement of signal, which is likely due to the loss of terminal α 1,3-Man caps and the exposure of more α 1,2-Man on that protein. This hypothesis was confirmed by the following experimental study shown in Figures 17 and 18.

[0253] To analyze the effect of deletions of the genes involved in glycosylation pathway, size changes in PST1 protein in this double mutant, along with other yeast single mutants, were examined. Cell lysates (Fig. 17A, upper) and culture media (Fig. 17B, upper) from different strains were analyzed using Western blots with anti-PST1 polyclonal antibody. PST1 was only detected in the cell lysate of the Δ mnn1 Δ pmr1 double mutant. The culture media shows the presence of a hyperglycosylated PST1 at > 200kDa in the

wild type strain. PST1 in the Och1 mutant shows a significant migration shift to approximately 105 kDa, which migrated at similar size to that in pmr1 single mutant. These results suggest that the outer chain of high mannose on PST1 in the pmr1 mutant was not added, like in the och1 mutant. PST1 in the double mutant migrated even faster than it did in the pmr1 or och1, indicating a loss of terminal α 1,3-Man on the Man₈GlcNAc₂ core in double mutant.

[0254] In contrast to PST1, another glycoprotein, GP38, was present in the cell lysate from all tested strains (Fig. 17A, lower). The size of this protein varied dramatically as the strains were mutated. Wild type and $\Delta mnn1$ showed a large, hyperglycosylated size of greater than 200 kDa. A band at ~ 90 kDa was shown in $\Delta och1$ while a band at ~ 110 kDa shown in $\Delta pmr1$ suggesting a complete loss and truncation of hyperglycosylation in these two mutants, respectively. GP38 in pmr1mnn1 double mutant showed a slightly smaller size than that in pmr1 single mutant, indicating the loss of the α 1,3-linked mannose caps, which is consistent with PST1 seen in these two mutants. In addition, GP38 was detectable in media neither from wild type nor from mnn1 mutant (Fig. 17B, lower). Interestingly, it was strongly induced by mutation of pmr1 or och1. This further indicates that Pmr1p, similar to Och1p, plays a critical role in glycosylation, at least in some proteins. However, while the $\Delta och1$ mutation almost guarantees the loss of hyperglycosylation, due to the deletion of the gene responsible for outer chain initiation, $\Delta pmr1$ appears to affect the efficiency of such chain elongation. This results in some proteins, like PST1, having a complete loss of hyperglycosylation, and other proteins, like Gp38, having reduced hyperglycosylation. The size changes of GP38 in pmr1 single mutant and pmr1mnn1 double mutant showed the same pattern as PST1 in these two mutants indicating that the deletion of α 1,3-Man that is added in the Golgi.

[0255] To confirm the loss of mnn1p function, the presence of terminal α 1,3-Man and α 1,2-Man were analyzed by Western blot with antibodies specific to each structure. Figure 18 lower panel shows that the levels of PST1 expression in pmr1 single mutant and pmr1mnn1 double mutant are similar. However, a band was detected by the anti- α 1,3-Man specific polyclonal antibody in pmr1 single mutant while this activity was completely lost in the double mutant (Fig 18, upper panel). In contrast, anti- α 1,2-Man MAb 2G12 showed the opposite results. The 2G12 signal in the double mutant was much stronger than that in single mutant (Fig. 18, middle panel). In the och1 mutant, anti- α 1,3-Man antibody, but not anti- α 1,2-Man (2G12), detected a strong smear ranging from 70 to over 150 kDa indicating

multiple glycoproteins with α 1,3-Man structure are detected by anti- α 1,3-Man antibody. The partially purified Yp100 showed reactivity to PST1 antibody and 2G12, but not to anti- α 1,3-Man antibody.

[0256] 2G12 MAb cross-reacts with both native and denatured PST1 glycoprotein.

To test whether 2G12 recognizes the glycans on the native form of PST1, immunoblot was performed. Proteins in culture media from double mutant were precipitated with 2G12 followed by incubation with anti-human IgG-agarose. The immune complex was eluted using SDS-sample buffer and separated on 4-20% SDS-PAGE. Then the blot was probed with anti-PST1 polyclonal antibody. As shown in Figure 19, strong signals were detected by anti-PST1 in the starting material (lane 1), flow through of preclear beads (lane 2), and elute (lane 6), but not detected, or weakly detected, in the supernatant and washes (lanes 3-5). These results indicate that 2G12 recognizes the native PST1 protein and has high affinity to the carbohydrate epitopes since most MAbs do not precipitate proteins.

Background information of PST1 gene and protein:

[0257] Pst1p (*protoplast secreted protein 1*), belongs to the SPS2 family. It is a protein attached to the membrane by glycosylphosphatidyl inositol (GPI) anchor, and secreted by regenerating protoplasts. Pst1p is thought to be important for cell wall integrity due to its upregulation upon activation of the cell integrity pathway, mediated by Rlm1p, and its upregulation by cell wall damage as mediated by Fks1p.

[0258] Gene and Family: Its precursor has 444 amino acids, with a predicted signaling peptide of 19 amino acids and a C-terminal domain of 25 amino acids for GPI anchor attachment. The molecular mass and isoelectric point of precursor are 45,776 Da and 9.91, respectively. It has a large number of N-linked glycosylation sites and rich in serine and threonine, which are likely to be heavily *O*-glycosylated. Its mature protein has 400 amino acids with predicted molecular mass of 41,260 Da and isoelectric point of 9.25. PST1 contains 15 potential N-linked glycosylation sites, which count 3.8% of the 400 amino acids. There are a total of 118 of serine and threonine residues, which count 29.5% of the 400 residues. Some of these serine and threonine are *O*-linked glycosylation sites (Figure 33).

[0259] PST1 is most similar to Ecm33, showing a 58% identity and 79% similarity over the whole protein. The homologs of PST1 and Ecm33 were found in different species of yeasts and fungi but not in any other species including mammals, plants, insects, worms,

and birds (Figure 20). They have similar or overlapping activities, but they are not functionally redundant (Pardo M, *Microbiology*, 150:4157-4170, 2004).

[0260] Function: Molecular function of PST1 is unknown. It is involved in the cell wall organization and biogenesis. Both PST1 and Ecm33 are GPI-anchored cell wall proteins and localize to the cell surface.

[0261] Expression, induction, and secretion: PST1 (YDR055W) was initially found in the culture media of *S cerevisiae* that was incubated in regenerating conditions with glucanase treatment to obtain protoplasts (Padro et al., *Yeast*, 15:459-472, 1999). Disruption of *FKS1*, one of subunits of glucan synthases, up-regulated the expression levels of YDR055W RNA and protein (Terashima et al., *Mol Gen Genet*, 264:64-74, 2000). The expression levels of its mRNA were found to be up-regulated by activation of MAP kinase, Mpk1/Slt2, or elevated temperature which cause environmental stresses that challenge the integrity of yeast cell wall (Jung et al., *Mol. Microbiol.* 34(5):1049-57, 1999).

[0262] Taken together, a yeast glycoprotein was found to cross-react with HIV-1 broadly neutralizing MAb 2G12, which binds an epitope composed of Man α 1,2-Man on high mannose type of glycan. This heavily glycosylated protein was identified to be PST1, a cell wall or membrane protein attached through GPI anchor. It is a secreted protein and up-regulated after transformation or deletion of the genes that are involved in the glycosylation pathway.

Example 3. Construction of triple mutants and identification 2G12 cross-reactive glycoproteins in Och1/Mnn1/Mnn4 triple mutant.

I. Construction and Characterization of Yeast Triple Mutant:

Yeast mating and verification of the $\Delta mnn1\Delta mnn4\Delta och1$ triple mutant.

[0263] In order to explore the possibility of using yeast to produce glycoproteins similar to those found on gp120, the glycosylation pathway had to be mutated. Three proteins are vital for creating N-linked glycans that are specific to *S. cerevisiae*: OCH1p initiates the first α 1,6 mannose residue necessary for hypermannosylation (Lehle, et al., *FEBS Lett.* , 370(1-2):41-5, 1995), MNN1p is responsible for all terminal α 1,3-linked mannose capping of α 1,2 residues (Nakajima et al, *Proc Nat Acad Sci USA*, 72:3912-3916, 1975), and MNN4p is a positive regulator of mannosylphosphorylation (Jigami, et al., *Biochim Biophys Acta*, 1426(2):335-45, 1999). Yeast deletion strains with ORF knockouts

of these genes were mated and sporulated to produce $\Delta mnn1\Delta mnn4$ and $\Delta mnn1\Delta och1$ haploids. Due to the temperature sensitivity exhibited by the $\Delta och1$ phenotype (Lee, et al., *Proc Natl Acad Sci*, 97(22):12679-84, 1999), selection of these clones was determined by slow growth on YPD at 37°C. Colony PCR was conducted for selection and verification of the $\Delta mnn1$ and $\Delta mnn4$ phenotypes (Data not shown).

[0264] The resulting double mutant haploids were mated and sporulated to produce the haploid, triple mutants, $\Delta mnn1\Delta mnn4\Delta och1$. Screening of $\Delta mnn4$ and $\Delta och1$ phenotype was conducted as before, using slow growth on YPD at 37°C for $\Delta och1$, and PCR for $\Delta mnn4$ and $\Delta och1$. Figure 23A shows a sample of the colony PCR screening for four clones, with three clones showing the presence of both the $\Delta mnn4$ and $\Delta och1$ phenotype.

[0265] Six putative triple mutants were tested by PCR to verify the $\Delta mnn1\Delta mnn4\Delta och1$ genotype. Figure 21B shows the results for two of these clones, $\Delta mnn1\Delta mnn4\Delta och1$ -2-5 and $\Delta mnn1\Delta mnn4\Delta och1$ -2-6. Four unique ORF-specific primers per gene deletion confirmed that each gene was completely deleted. The combination of an internal ORF-specific primer (MNN1B, MNN4B, OCH1B, MNN1C, MNN4C and OCH1C) and an upstream or down stream primer (MNN1A, MNN4A, OCH1A, MNN1D, MNN4D and OCH1D) results in a PCR product if the ORF is present (i.e. MNN1A + MNN1B, MNN1C + MNN1D, etc.). As seen in the figure, both clones showed no PCR product for both primer pairs against MNN1, MNN4 and OCH1, verifying the deletion genotype. Each ORF specific primer pair was also tested against WT as a positive control (results not shown).

[0266] By contrast, the combination of an upstream or downstream primer and an internal KanMX specific primer (KanB and KanC) results in a PCR product if the ORF has been replaced with the KanMX module. As seen in Figure 23B, both of these clones show a PRC product at the expected size when tested with both primer pairs per gene deletion.

[0267] Two of the haploid triple mutants, $\Delta mnn1\Delta mnn4\Delta och1$ -2-6 and $\Delta mnn1\Delta mnn4\Delta och1$ -2-5, were mated to form the diploid triple mutant, $\Delta mnn1\Delta mnn4\Delta och1$ -DIP. This strain would be used for any future comparison studies to the wild-type diploid strain, INVSc1, including ELISA, immunofluorescence, and immunization.

Loss of α 1,3-Mannose and increase of 2G12 cross-reactivity in the triple mutant by ELISA.

[0268] The $\Delta mnn1\Delta mnn4\Delta och1$ mutant yeast strain is expected to produce glycoproteins with a majority of $Man_8GlcNAc_2$ glycans. In order to indirectly verify the presence of these oligosaccharides, the triple mutant was tested for cross-reactivity to two mannose-specific antibodies, 2G12 and anti- $\alpha 1,3$ -linked mannose (anti- $\alpha 1,3$). Immune sera specific to $\alpha 1,3$ -linked mannose residues was prepared by immunizing rabbits with INVSc1 whole-cells and adsorbing the sera to $\Delta mnn1$ cells. Raschke et al., *J Biol Chem.* 248(13):4660-6, 1973; and Ballou, *J Biol Chem*, 245:1197-1203, 1970). The resulting sera are expected to be enriched for antibodies specific to the terminal $\alpha 1,3$ -linked mannose residues used to cap $\alpha 1,2$ -linked residues by Mnn1p. By contrast, 2G12 is known to be specific to the terminal $\alpha 1,2$ -linked mannose residues found on the core N-linked glycan, albeit a cluster of such residues (Scanlan et al., *J Virol*, 76:7306-21, 2002).

[0269] Intact, whole yeast cells for WT (INVSc1) and $\Delta mnn1\Delta mnn4\Delta och1$ -DIP cells were used to coat ELISA plates, and 2G12 and $\alpha 1,3$ specific Ab were used to probe. As seen in Figure 22A, the anti- $\alpha 1,3$ antibody showed a high affinity towards the wild-type cells, with the triple mutant showing low affinity. SF162 and wild-type yeast mannan were included as negative and positive controls, respectively. These results show the loss of terminal $\alpha 1,3$ -linked mannose residues in the triple mutant. Any residual affinity of the anti- $\alpha 1,3$ towards the triple mutant can be attributed to the fact that the adsorption process is not completely efficient; other yeast-specific antibodies that are still present can bind to the epitopes exposed on the triple mutant (i.e. cell wall proteins and β -glucan).

[0270] Figure 22B shows that 2G12 was able to bind to the triple mutant whole cells with no cross-reaction in binding to the wild-type. This affinity was seen between 10 μ g/ml to $\sim$ 1 μ g/ml. SF162 and wild-type yeast mannan were included as positive and negative controls, respectively. These results not only confirm the loss of terminal $\alpha 1,3$ -linked mannose residues in the triple mutant, but also show the resulting exposure of the $\alpha 1,2$ -linked mannoses found on the core Man_8 glycan. More importantly, the ability of the $\Delta mnn1\Delta mnn4\Delta och1$ yeast to bind to 2G12 implies the presence of 2G12-like epitopes on one or more proteins found on the yeast cell wall.

Loss of $\alpha 1,3$ -Mannose and increase of 2G12 cross-reactivity in the triple mutant by Immunofluorescence.

[0271] Terminal $\alpha 1,3$ mannose residues in yeast not only cap $\alpha 1,2$ -linked mannose residues, but are also highly immunogenic, making them divergent from the glycans found

on HIV gp120. In order to further verify the loss of α 1,3-linked mannose residues in the Δ mnn1 Δ mnn4 Δ och1 strain, we conducted immunofluorescence using anti- α 1,3-Man on intact, whole cells. As shown in Figure 23 (Panel A and C), the wild-type cell wall shows a strong reaction to the antisera specific to α 1,3-linked mannose residues, while the diploid triple mutant shows no signal at the same dilution. As a control, antibodies specific to yeast Zymosan, which contains common components of the yeast cell wall, α -mannan and β -glucan, were found to give a similar signal to both the INVSc1 and Δ mnn1 Δ mnn4 Δ och1 whole-cells by immunofluorescence (Figure 23, Panel B and D). These results indicate the loss of the α 1,3-mannosyltransferase activity specific to the MNN1 protein. Such a loss would almost strictly leave α 1,2-linked terminal mannose residues on all glycoproteins found in the cell wall of the triple mutant.

[0272] To confirm this exposure of the α 1,2-mannose residues on the triple mutant, 2G12 was then tested for co-localization with anti-Zymosan by immunofluorescence. Figure 24 (Panel A, B, C) shows that 2G12 binds strongly to Δ mnn1 Δ mnn4 Δ och1-DIP cells and is co-localized with anti-Zymosan on the cell wall. The wild-type yeast only shows binding to anti-Zymosan, with no signal seen for 2G12 (Panel D, E, F). These data, similar to the whole-cell ELISA results, show that the glycoproteins found on the triple mutant contain exposed α 1,2-linked mannose structures with the ability to bind 2G12 under native conditions. It is interesting to note that the sizes of the triple mutant yeast cells are noticeably larger than that of wild-type cells.

Loss of α 1,3-Mannose and exposure of α 1,2-Mannose on glycoproteins in the triple mutant by Western blot.

[0273] These 2G12-reactive proteins appear in the cell wall of Δ mnn1 Δ mnn4 Δ och1 yeast, but are absent in the wild-type. However, this does not directly implicate the exposed α 1,2-linked mannose residues as the primary reason for affinity towards 2G12. Using Western blots, the presence of 2G12-reactive proteins among a panel of yeast glycosylation mutants was tested. As seen in Figure 25 left panel, only the lysate from the double mutant Δ mnn1 Δ och1 and the triple mutant Δ mnn1 Δ mnn4 Δ och1 showed specific protein bands that are detected by 2G12. In fact, both of these glycosylation mutants appear to have the same number of proteins (~4-5) that are reactive to 2G12. This, along with the lack of signal among the other yeast mutants, implies that the presence of core Man8 in Δ mnn1 Δ mnn4 Δ och1, and Δ mnn1 Δ och1 mutants plays an important role in the specificity

of 2G12 toward these proteins. In addition, the lack of signal seen on the $\Delta mnn1$ and $\Delta mnn1 \Delta mnn4$ mutant, which should have hyperglycosylated N-linked glycans with numerous exposed $\alpha 1,2$ -linked mannose residues, further implicates the sole role of Man₈ glycans in 2G12 binding.

[0274] This panel of yeast glycosylation mutants was also probed with anti- $\alpha 1,3$ to verify the presence or loss of $\alpha 1,3$ -linked mannose caps. Figure 25 right panel shows that the anti- $\alpha 1,3$ -Man serum does not detect proteins on any mutant with the $\Delta mnn1$ phenotype, including $\Delta mnn1 \Delta och1$ and the triple mutant. This further implies the exposure of $\alpha 1,2$ epitopes on the triple mutant and the $\Delta mnn1 \Delta och1$ strain, along with the $\Delta mnn1$ and $\Delta mnn1 \Delta mnn4$ strains, which 2G12 does not recognize. A second, identical Western blot was mock incubated (without a primary antibody) and showed no significant signal in any of the strains, thereby ruling out interaction with the goat anti-human secondary antibody (results not shown).

[0275] In order to verify that the reactivity of the triple mutant proteins towards 2G12 is due to the genotype of the yeast strain, and that no other protein with complementary function are expressed, different triple mutant clones were analyzed. A total of six $\Delta mnn1 \Delta mnn4 \Delta och1$ yeast strains were obtained from the $\Delta mnn1 \Delta mnn4 \times \Delta mnn1 \Delta och1$ cross. Each was picked from a separate haploid colony growing on YPD + KCl plates and was verified to contain the three deletions (see Figure 21 for clones 2-5 and 2-6). In Figure 26, all six clones show 2G12 reactivity to the same proteins by Western blot. This indicates that the reactivity of these proteins to 2G12 is most likely due to the genotype of the triple mutant resulting in mutated glycans.

Verification of the glycans on the triple mutant.

[0276] As a final confirmation of the oligosaccharides present on the $\Delta mnn1 \Delta mnn4 \Delta och1$ glycoproteins, MALDI-TOF and NP-HPLC profiling were conducted on N-linked glycans extracted from whole yeast cell extract. In Figure 26, the MALDI-TOF results show that Man₈GlcNAc₂ is predominant glycan in the triple mutant extract, with a very minor amount of Man₉GlcNAc₂ and Man₅GlcNAc₂. The level of Man₈ glycans is 20 times more than that of either Man₉ or Man₅, such that Man₈GlcNAc₂ represents over 90% of the total glycans in the triple mutant. The NP-HPLC results appear consistent with the MALDI-TOF, such that Man₈ is the predominant N-linked glycan present in the $\Delta mnn1 \Delta mnn4 \Delta och1$ cell extract.

[0277] Taken together, these results verify that the types of glycans present in the $\Delta mnn1 \Delta mnn4 \Delta och1$ cells, are predominantly $Man_8GlcNAc_2$, similar to those found on core N-linked glycans. This indicates not only the loss of the highly immunogenic terminal $\alpha 1,3$ -linked mannose residues, but also the loss of the hypermannosylated outer chain. The resulting exposure of terminal $\alpha 1,2$ -linked mannose residues is most likely the reason behind the affinity of 2G12 for specific cell wall proteins found in this mutant.

II. Identification and Verification of 2G12 Cross-Reactive Glycoproteins in the Yeast Triple Mutant

2D separation of 2G12-reactive proteins from yeast cells.

[0278] These proteins in $\Delta mnn1 \Delta mnn2 \Delta och1$ yeast that show binding to 2G12 by ELISA, immunofluorescence and Western blot are promising antigens for the production of 2G12-like antibodies. They would represent the first such antigens produced in yeast with the ability to mimic the 2G12 epitope, which consists of a conformation of $\alpha 1,2$ -linked mannose residues found on a cluster of N-linked glycans. Thus, this study was continued with the partial purification and identification of these proteins. Due to the predominance of glycoproteins in the cell wall and plasma membrane, differential centrifugation was used to largely separate the membrane-associated proteins from the cytoplasmic proteins. Log-phase $\Delta mnn1 \Delta mnn4 \Delta och1$ -DIP cells were lysed and subjected to centrifugation to remove cytoplasmic, Golgi complex and endosomal proteins. The pellet, containing mostly vacuolar, nuclear, ER and plasma membrane proteins, was resolubilized in 1% Triton X-100 to separate out the highly insoluble proteins. As seen in Figure 28A, the 2G12-binding proteins were found mainly in the membrane fraction, and a small majority of these proteins were found to be soluble in Triton. These results are consistent with the whole-cell ELISA and immunofluorescence, in that these proteins appear to be mainly cell wall- or plasma membrane-associated.

[0279] In addition to being membrane-associated, previous results also implied that these 2G12-reactive proteins are highly glycosylated. Therefore, Concanavalin A (ConA) lectin affinity chromatography was used to purify the 2G12-reactive proteins found in the Triton-soluble fraction. Figure 28B shows that these proteins have a high affinity for ConA, such that the majority of proteins cannot be fully eluted by competitive binding (using 0.5M methyl manno-pyranoside), requiring harsher conditions to elute (1% SDS).

[0280] The resulting proteins from the ConA eluate would represent the highly glycosylated membrane proteins found in the $\Delta mnn1\Delta mnn4\Delta och1$ cells, including those that bind to 2G12. In order to efficiently separate these proteins for identification, 2-D gel electrophoresis was used on this fraction. Two separate aliquots were run onto pH4.0-7.0 IPG strips followed by SDS-PAGE separation on 4-12% gradient gels. One gel was stained by Sypro Ruby while the other gel was transferred to nitrocellulose for a Western blot using 2G12. Figure 28C shows that two large spots were identified by Western blot, which corresponded to two spots visualized on the Sypro Ruby stained gel (not shown). These spots were excised for Trypsin digestion and nano LC/MS/MS identification. The higher 2D spot (at ~110 kDa) identified peptides from three yeast glycoproteins, Gas5 (5 peptides), ECM33 (4 peptides), and Gas1 (2 peptides). The lower 2D spot (at ~95 kDa) identified 7 peptides from a single yeast glycoprotein, YJL171c. These proteins represent all or most of the higher molecular weight proteins with 2G12-reactivity. There may be more such proteins at this size range that were not identified, but there is definitely an ~35kDa protein (see Figures 25 and 28) that was not identified in this screen. The identity of this protein is not known, but is currently under investigation.

1-D separation of 2G12-reactive proteins from the culture supernatant.

[0281] In addition to the cell lysate, the culture supernatant of the yeast triple mutant was found to contain glycoproteins with 2G12-reactivity. The culture media from INVSc1 and $\Delta mnn1\Delta mnn4\Delta och1$ cells were tested for 2G12-reactivity by Western blot. Figure 29A shows the expression of at least four bands with strong reactivity to the 2G12 MAbs in the culture media of the triple mutant, while the wild-type yeast supernatant shows no signal.

[0282] These 2G12-reactive proteins in the media may represent secreted glycoproteins, or cell wall proteins released into the media, either naturally or due to changes in cell wall structure of the triple mutant. In fact, the sizes of these four proteins appear similar to the 2G12-reactive proteins found in the cell lysate (see Figure 25A & 28A), with a predominance of signal seen between 90 to 115kDa and one band at ~37kDa. This suggests that these could be the same proteins found in the cell lysate. Regardless, the fact that these proteins may represent other antigens that mimic the 2G12 epitope warrants their identification. The supernatant from $\Delta mnn1\Delta mnn4\Delta och1$ yeast was applied directly to a ConA column for partial purification of the glycoproteins. The proteins in the culture supernatant showed a very high affinity to the ConA column and could not be eluted thought competitive binding using methyl manno-pyranoside. Therefore, the 2G12-reactive

proteins were eluted by boiling the ConA beads with SDS-PAGE loading buffer (Figure 29B). This SDS elute was separated using a large-format (10cm x16cm) 10% SDS-PAGE gel. A Coomassie blue stain of the gel showed five separate protein bands, each of which was excised from the gel (data not shown). These bands were tested for 2G12 reactivity by re-loading onto an SDS-PAGE minigel and probing with 2G12 by Western blot. Two protein bands (at 110kDa and 95kDa) were found to bind 2G12 (results not shown) and excised. These proteins were digested with Trypsin, and analyzed by nano LC/MS/MS on a Micromass Q-Tof 2 for peptide identification. The 110kDa band identified only one protein in the gel slice, with 4 peptides matching ECM33, the same protein identified by 2D at 110kDa in the membrane fraction. The 95kDa band identified only one protein in the gel slice, with 4 peptides matching the yeast glycoprotein, Gp38.

[0283] These two proteins probably represent only half of the 2G12-reactive proteins in the culture media (see Figure 29A); in fact there are four bands that appear to be 2G12-reactive in the culture supernatant. These two unidentified proteins may be the same as those found in the cell lysate, or, like GP38, may represent unknown proteins with the ability to bind 2G12. These unidentified proteins are still under investigation.

Verification of 2G12-reactive proteins by immunoprecipitation.

[0284] The identification of ECM33, Gas1, Gas5, GP38 and YJL171c as 2G12-reactive yeast glycoproteins was conducted indirectly; LC/MS/MS was used to identify bands in gel electrophoresis that bound to 2G12 by Western blotting. In order to confirm the affinity of these proteins to 2G12, the cell lysate and culture media of $\Delta mnn1\Delta mnn4\Delta och1$ cells were immunoprecipitated with 2G12 and probed the resulting Western blots with antibodies against these proteins. Figure 30A shows the identification of 4 proteins, Gas1, ECM33, YJL171c, and GP38, that were precipitated from the culture supernatant by 2G12. Gas1, ECM33 and YJL171c were identified in the cell lysate of the triple mutant by 2D separation, and show a similar size in both experiments, ~115kDa, ~110kDa, and ~95kDa, respectively. GP38 was originally identified in the culture supernatant of the triple mutant, but here shows that can be also be precipitated from the cell lysate by 2G12. Gas5 was also precipitated with 2G12 with relatively low affinity (data not shown), which could be due to the less number and lower density of N-linked glycans than those from other identified proteins (Figure 38).

[0285] Figure 30B shows the identification of 3 proteins, PST1, ECM33, and GP38, with immunoblots. These proteins were precipitated from the culture supernatant by 2G12.

ECM33 and GP38 were also identified in the culture media of the triple mutant by 1D separation, and show a similar size in both experiments, ~110kDa and ~95kDa, respectively. Interestingly, PST1 was originally identified as a 2G12-binding protein in the culture supernatant of Δ pmr1 yeast (Example 2). It can be seen here that the expression level of PST1 is very low in the supernatant of the triple mutant (Figure 30B, Lane1) when compared to ECM33 and GP38, which may explain why this protein wasn't originally identified in this mutant.

[0286] These results confirm the ability of ECM33, Gas1, Gas5, GP38, PST1 and YJL171c to bind 2G12 under the native conditions of this immunoprecipitation experiment. When analyzed with our previous experiments, it appears that these yeast glycoproteins have the ability to mimic the 2G12 epitope found on the HIV-1 gp120 glycoproteins, thereby causing binding to this antibody by Western blotting, immunoprecipitation, immunofluorescence and ELISA.

Glycoprotein size shifts for two 2G12-reactive yeast proteins.

[0287] The discovery and identification of six yeast glycoproteins with the potential to mimic the 2G12 epitope warrants further investigation into the glycan structures on these proteins. Two of these proteins, ECM33 and GP38, were analyzed against a panel of yeast glycosylation mutants by Western blot. ECM33 is a glycosyl phosphatidyl inositol (GPI)-anchored protein important for cell wall stability (Pardo, *Microbiology* 150:4157-70, 2004), and gp38 is a cell wall-related secretory protein (Destruelle, *Mol. Cell. Biol.* 14(4):2740-54, 1994). Figure 31 (left panels) shows the results. In the cell lysate, both of these proteins are present in all the strains, albeit at different levels. In addition, these proteins are hypermannosylated in any strain without the Δ och1 genotype. In fact, the Δ och1 genotype causes a drastic shift in size of greater than 80kDa for both protein, with GP38 showing a more significant change. A closer analysis of these strains shows that in both proteins there is a noticeable size shift between the Δ och1 and Δ och1 Δ mn1 strain, indicating a loss of terminal α 1,3-linked manose residues. The final protein sizes in the Δ mn1 Δ mn4 Δ och1 mutant appear to be the same size as the proteins precipitated by 2G12; ~110kDa for ECM33 and ~95kDa for GP38 (see Figure 31). Any size shift caused by the loss of phosphomannose in Δ mn4 mutants is not seen on the SDS-PAGE gels, suggesting that the change is too slight.

[0288] In the culture supernatant (Figure 31, right panels), there is little to no GP38 or ECM33 present from any strains with the OCH1 gene. It appears that the weak cell walls of

Δ och1 containing yeast mutants results in the release of these proteins into the media. However, a similar size shift can be seen in ECM33 by comparing the ~200kDa band found in the Δ mnn1 Δ mnn4 mutant and the 110-115kDa band found in Δ och1 containing mutants. Again, this shows the lack of outer chain hypermannosylation. Also, a similar size shift from the Δ mnn1 mutation can be seen between Δ och1 and Δ och1 Δ mnn1 mutants or triple mutants.

[0289] Taken together, these results help us visualize the loss of hypermannosylation and α 1,3-linked mannose caps as the yeast strains are mutated. By comparing these results with Figure 25, 2G12 reactivity is not present unless both of these changes in glycan structure are present. Thus, the ability of ECM33 and GP38 to bind 2G12, along with Gas1, Gas5, YJL171c, and PST1, is dependent upon the presence of core Man8 oligosaccharides preserved in the yeast triple mutant and Δ och1 Δ mnn1 double mutants.

Production of α 1,2-mannose specific anti-sera in rabbits that cross-reacts with gp120 by ELISA.

[0290] We identified at least six yeast glycoproteins, PST1, ECM33, Gas1, Gas5, GP38 and YJL171c, from Δ mnn1 Δ mnn4 Δ och1 cells that showed cross-reactivity to the HIV MAb 2G12. In addition, the cell wall of the yeast triple mutant was able to cross-react with 2G12 by whole cell ELISA and immunofluorescence. Therefore, in order to attempt to induce antibodies similar to 2G12 - such that they are specific to α 1,2-linked terminal mannose sugars - whole cell, heat-killed triple mutant cells were used to immunize rabbits. Serum samples from 0, 4, 5, and 12 weeks were tested for HIV Env-specific total IgG antibodies. As shown in Figure 32, the levels of antibodies specific to gp120 show a significant increase from week 0 to week 5, and a subsequent decrease at week 12. There was a significant increase in the gp120 binding in the sera from week 5 when compared to week 0 ($P < 0.001$), while the binding at week 12 showed a significant increase too ($P < 0.05$) although the titer is lower.

[0291] For comparison, rabbits were immunized in parallel with whole cell, heat-killed WT cells. The resulting sera showed that there is no significant increase in the levels of gp120 specific antibodies from week 0 to week 12 (see Figure 32). By comparing the parallel bleeds from WT and triple mutant immunized rabbits, we again see a significant increase in gp120-specific IgG in the triple mutant, particularly at week 5 ($P < 0.001$), when compared to WT. All of these p values were calculated by the Student's two-tailed *t* test. Altogether, these results may present the first instance of the production of antibodies in

animals that are able to cross-react with the α 1,2-linked terminal mannose residues on gp120.

III. Bioinformatic Analyses of N-linked and O-linked glycosylation sites in the Glycoproteins Recognized by 2G12 in the Yeast Triple Mutant.

[0292] Figures 33-38 show the analyses of potential N-linked and O-linked glycosylation sites of PST1, ECM33, GP38, YJL171c, Gas1, and Gas5. Table 2 below shows a summary of identified 2G12-reactive glycoproteins in yeast mutants. There are several noticeable features of the identified 2G12 cross-reactive glycoproteins. All have a high number and density of N-linked and/or O-linked glycosylation sites. Among them, PST1 and ECM33 are members in the same family and have high number and density of both N-linked and O-linked glycosylation sites. GP38 and YJL171c have only high number and density of N-linked glycosylation sites, and Gas1 and Gas5 in the same family have high number and density of O-linked glycosylation sites and lower percentage (approximately 50% versus 60% of the other four) of molecular mass from glycans.

Table 2. Summary of identified 2G12-reactive glycoproteins in yeast mutants.

Characterization of 2G12 Cross-Reactive Glycoproteins in Yeast Mutants											
ID	Name	Precursor	Mature Protein			N-sites	O-sites	Molecular weight (kDa)			GPI
		aa No.	aa No.	MW	PI	No. (%)	No. (%)	PAGE	MW	Glycan (%)	
P38616	GP38	354	315	33,124	4.67	15 (4.8)	4 (1.3)	90	33	57 (63)	No
P46992	YJL171c	396	349	37,723	4.69	11 (3.2)	0 (0)	90	38	52 (58)	Yes
Q12355	PST1	444	399	41,260	9.25	15 (3.8)	14 (3.5)	100	41	59 (59)	Yes
P38248	ECM33	469	386	39,628	4.68	13 (3.4)	21 (5.4)	100	40	60 (60)	Yes
P22146	Gas1	559	506	54,325	4.37	10 (2.0)	26 (5.1)	105	54	51 (49)	Yes
Q08193	Gas5	484	440	47,298	4.43	6 (1.4)	35 (8)	100	47	53 (53)	Yes

[0293] Table 3 below shows the homologs of 2G12 cross-reactive glycoproteins among 18 genomes. The protein sequences of the six 2G12 cross-reactive glycoproteins were used to search homolog genes from HomoloGene database (release 50.1) of National Center for Biotechnology Information (NCBI). HomoloGene is a system for automated detection of homologs among the annotated genes of several completely sequenced eukaryotic genomes. Currently HomoloGene database contains 165,820 HomoloGene groups from 18 species. The HomoloGene of interest is obtained by entering protein Reference Sequence

(RefSeq) number at NCBI home page. Multiple alignment of the HomoloGene was performed by clustalw program. The homologs of each gene were obtained from HomoloGene database searching and listed in the table. Homologs in *Candida glabrata* of the six genes were found in NCBI database but the genome of *Candida glabrata* is not included in the HomoloGene database.

Table 3. Homologs of 2G12 cross-reactive glycoproteins among 18 genomes.

Homologs of 2G12 Cross-Reactive Glycoproteins among 18 Genomes								
	Genome	Species	PST1	ECM33	Gas1	Gas5	GP38	YJL171c
Mammalian	Homo sapiens	Human	-	-	-	-	-	-
	Pan troglodytes	Chimpanzee	-	-	-	-	-	-
	Canis familiaris	Dog	-	-	-	-	-	-
	Mus musculus	House mouse	-	-	-	-	-	-
	Rattus norvegicus	Rat	-	-	-	-	-	-
Bird	Gallus gallus	Chicken	-	-	-	-	-	-
Insect	Drosophila melanogaster	Fly	-	-	-	-	-	-
	Anopheles gambiae	Mosquito	-	-	-	-	-	-
Worm	Caenorhabditis elegans	Small soil nematode	-	-	-	-	-	-
Fungus	Saccharomyces cerevisiae	Yeast	+	+	+	+	+	+
	Schizosaccharomyces pombe	Yeast	+	+	+	+	-	-
	Kluyveromyces lactis	Yeast	-	+	+	+	-	+
	Eremothecium gossypii	Cotton pathogen	+	-	-	+	-	-
	Magnaporthe grisea	Rice blast fungus	-	+	+	-	-	-
	Neurospora crassa	Red bread mold	-	+	+	+	-	-
	Arabidopsis thaliana	Small flowering plant	-	-	-	-	-	-
Plant	Oryza sativa	Japanese rice	-	-	-	-	-	-
Parasite	Plasmodium falciparum	malaria parasite	-	-	-	-	-	-

Example 4. Production and Characterization of HIV-1 gp120 Glycoproteins with Homogenous Man8 type Glycans in Och1/Mnn1/Mnn4 Triple Mutant.

[0294] Unlike in the mammalian expression system, the carbohydrates on glycoproteins produced in *S. cerevisiae* contain only mannose residues built upon the core GlcNAc₂ sugars. Full-length gp120 proteins have been successfully expressed in *S. cerevisiae*, although they appear to be hyperglycosylated. In order to create gp120 proteins in yeast with strictly Man₈GlcNAc₂, the glycosylation pathway had to be mutated. Three proteins are responsible for creating N-linked glycans that are specific to *S. cerevisiae*: OCH1p initiates the first α 1,6 mannose residue necessary for hypermannosylation (Lehle, et al., *FEBS Lett.*, 370(1-2):41-5, 1995), MNN1p is responsible for all terminal α 1,3-linked mannose capping of α 1,2 residues (Nakajima et al., *Proc Nat Acad Sci USA*, 3912-3916,

1975), and MNN4p is a positive regulator of mannosylphosphorylation (Jigami, et al., *Biochim Biophys Acta*, 1426(2):335-45, 1999). By deleting these three genes in *S. cerevisiae*, a strain for the production of gp120 with strictly Man8 oligosaccharides was created.

[0295] In order for gp120 to be efficiently glycosylated in the ER and secreted into the culture media, we first cloned the signal sequence of the *S. cerevisiae* alpha-mating factor (MF α) into the pYES2/CT vector (Invitrogen), which contains the GAL1 promoter for high-level expression of recombinant proteins following galactose induction. Then, HIV-1 gp120 genes from strains of JR-FL and YU2 (ARRRP) were PCR cloned into the pYES2/CT- α plasmid. The resulting gp120 expression plasmids were transformed into the haploid yeast strain $\Delta mnn1 \Delta mnn4 \Delta och1$, using a quick lithium acetate transformation protocol. As seen in Panel A of Figure 40, four separate clones of $\Delta mnn1 \Delta mnn4 \Delta och1$ -2-5, transformed with pYU2-gp120 or pJRFL-gp120, were verified to contain their respective plasmids by colony PCR. JRFL-gp120 transformants show an expected PCR product size of 724kB while YU2-gp120 transformants show a product at 1697kB.

[0296] These yeast transformants containing the pYU2-gp120 and pJRFL-gp120 were induced for protein expression by transferring the cells to galactose containing media. Western blotting using anti-gp120-IIIB (Virostat Inc.) and anti-gp120-YU2 were used to detect the induction and secretion of gp120 glycoproteins in the media. The $\Delta mnn1 \Delta mnn4 \Delta och1$ yeast transformants were found to secrete the greatest amount of gp120 by growth at 30°C for 6 to 9 days (Figure 39, Panel B and C). As seen in the Western blot, these gp120 proteins migrate to their predicted molecular mass of 96.4 and 99.6 kDa for JR-FL and YU-2 gp120, respectively, assuming they contain mainly Man8 oligosaccharides.

[0297] In addition to being reactive to anti-gp120 antibodies, these expressed gp120 proteins also show reactivity to 2G12 (Figure 40). Two $\Delta mnn1 \Delta mnn4 \Delta och1$ transformants (with pYU2-gp120 and pJR-FL-gp120) and a non-transformed strain were grown in parallel, with uracil added to the media of the non-transformant. Due to the presence of other 2G12-reactive proteins in the culture supernatant of $\Delta mnn1 \Delta mnn4 \Delta och1$ cells, one of which has a molecular weight of ~110kDa, the induction of gp120 can only be seen indirectly when testing with 2G12. The non-transformed yeast shows a steady basal expression level from 3-9 days, strongly suggesting that the basal level protein at ~110kDa (ECM33) does not increase in this time period. By contrast, both of the gp120 transformed

yeast show a marked increase in the expression of the 110kDa protein from 3-9 days, suggesting the induction of gp120 glycoproteins that are recognized by the 2G12 antibody.

LITERATURE CITED

1. UN AIDS Report on Global AIDS Epidemic. 2004.
2. McMichael AJ, Hanke T. HIV vaccines 1983-2003. *Nat Med*. 2003;9(7):874-80.
3. Garber DA, Silvestri G, Feinberg MB. Prospects for an AIDS vaccine: three big questions, no easy answers. *Lancet Infect Dis*. 2004;4(7):397-413.
4. <http://www.niaid.nih.gov/daids/vaccine>
5. <http://www.iavi.org>
6. Gaschen B, Taylor J, Yusim K, Foley B, Gao F, Lang D, Novitsky V, Haynes B, Hahn BH, Bhattacharya T, Korber B. Diversity considerations in HIV-1 vaccine selection. *Science*. 2002;296(5577):2354-60.
7. Burton DR, Desrosiers RC, Doms RW, Koff WC, Kwong PD, Moore JP, Nabel GJ, Sodroski J, Wilson IA, Wyatt RT. HIV vaccine design and the neutralizing antibody problem. *Nat Immunol*. 2004;5(3):233-6.
8. Zolla-Pazner S. Identifying epitopes of HIV-1 that induce protective antibodies. *Nat Rev Immunol*. 2004;4(3):199-210.
9. Geijtenbeek TB, Kwon DS, Torensma R, van Vliet SJ, van Duijnhoven GC, Middel J, Cornelissen IL, Nottet HS, KewalRamani VN, Littman DR, Figdor CG, van Kooyk Y. DC-SIGN, a dendritic cell-specific HIV-1-binding protein that enhances trans-infection of T cells. *Cell*. 2000;100:587-97.
10. Pohlmann S, Baribaud F, Lee B, Leslie GJ, Sanchez MD, Hiebenthal-Millow K, Munch J, Kirchhoff F, Doms RW. DC-SIGN interactions with human immunodeficiency virus type 1 and 2 and simian immunodeficiency virus. *J Virol*. 2001;75(10):4664-72.
11. Pohlmann S, Soilleux EJ, Baribaud F, Leslie GJ, Morris LS, Trowsdale J, Lee B, Coleman N, Doms RW. DC-SIGNR, a DC-SIGN homologue expressed in endothelial cells, binds to human and simian immunodeficiency viruses and activates infection in trans. *Proc Natl Acad Sci USA*. 2001;98(5):2670-2675.
12. Curtis BM, Scharnowske S, Watson AJ. Sequence and expression of a membrane-associated C-type lectin that exhibits CD4-independent binding of human immunodeficiency virus envelope glycoprotein gp120. *Proc Natl Acad Sci USA*. 1992;89:8356-60.
13. Bashirova AA, Geijtenbeek TB, van Duijnhoven GC, van Vliet SJ, Eilering JB, Martin MP, Wu L, Martin TD, Viebig N, Knolle PA, KewalRamani VN, van Kooyk Y,

- Carrington M. A dendritic cell-specific intercellular adhesion molecule 3-grabbing nonintegrin (DC-SIGN)-related protein is highly expressed on human liver sinusoidal endothelial cells and promotes HIV-1 infection. *J Exp Med*. 2001;193(6):671-8.
14. Mitchell DA, Fadden AJ, Drickamer K. A novel mechanism of carbohydrate recognition by the C-type lectins DC-SIGN and DC-SIGNR. Subunit organization and binding to multivalent ligands. *J Biol Chem*. 2001;276(31):28939-45.
 15. Feinberg H, Mitchell DA, Drickamer K, Weis WI. Structural basis for selective recognition of oligosaccharides by DC-SIGN and DC-SIGNR. *Science*. 2001;294(5549):2163-6.
 16. Pohlmann S, Doms RW. Evaluation of current approaches to inhibit HIV entry. *Curr Drug Targets Infect Disord*. 2002;2(1):9-16.
 17. Trkola A, Purtscher M, Muster T, Ballaun C, Buchacher A, Sullivan N, Srinivasan K, Sodroski J, Moore JP, Katinger H. Human monoclonal antibody 2G12 defines a distinctive neutralization epitope on the gp120 glycoprotein of human immunodeficiency virus type 1. *J Virol*. 1996;70(2):1100-8.
 18. Scanlan CN, Pantophlet R, Wormald MR, Ollmann Saphire E, Stanfield R, Wilson IA, Katinger H, Dwek RA, Rudd PM, Burton DR. The broadly neutralizing anti-human immunodeficiency virus type 1 antibody 2G12 recognizes a cluster of alpha1-->2 mannose residues on the outer face of gp120. *J Virol*. 2002;76(14):7306-21.
 19. Sanders RW, Venturi M, Schiffner L, Kalyanaraman R, Katinger H, Lloyd KO, Kwong PD, Moore JP. The mannose-dependent epitope for neutralizing antibody 2G12 on human immunodeficiency virus type 1 glycoprotein gp120. *J Virol*. 2002;76(14):7293-305.
 20. Shattock RJ, Moore JP. Inhibiting sexual transmission of HIV-1 infection. *Nat Rev Microbiol*. 2003;1(1):25-34.
 21. Boyd MR, Gustafson KR, McMahon JB, Shoemaker RH, O'Keefe BR, Mori T, Gulakowski RJ, Wu L, Rivera MI, Laurencot CM, Currens MJ, Cardellina JH 2nd, Buckheit RW Jr, Nara PL, Pannell LK, Sowder RC 2nd, Henderson LE. Discovery of cyanovirin-N, a novel human immunodeficiency virus-inactivating protein that binds viral surface envelope glycoprotein gp120: potential applications to microbicide development. *Antimicrob Agents Chemother*. 1997 Jul;41(7):1521-30.
 22. Esser MT, Mori T, Mondor I, Sattentau QJ, Dey B, Berger EA, Boyd MR, Lifson JD. Cyanovirin-N binds to gp120 to interfere with CD4-dependent human

- immunodeficiency virus type 1 virion binding, fusion, and infectivity but does not affect the CD4 binding site on gp120 or soluble CD4-induced conformational changes in gp120. *J Virol.* 1999;73(5):4360-71.
23. Dey B, Lerner DL, Lusso P, Boyd MR, Elder JH, Berger EA. Multiple antiviral activities of cyanovirin-N: blocking of human immunodeficiency virus type 1 gp120 interaction with CD4 and coreceptor and inhibition of diverse enveloped viruses. *J Virol.* 2000;74(10):4562-9.
24. Tsai CC, Emau P, Jiang Y, Agy MB, Shattock RJ, Schmidt A, Morton WR, Gustafson KR, Boyd MR. Cyanovirin-N inhibits AIDS virus infections in vaginal transmission models. *AIDS Res Hum Retroviruses.* 2004;20(1):11-8.
25. Tsai CC, Emau P, Jiang Y, Tian B, Morton WR, Gustafson KR, Boyd MR. Cyanovirin-N gel as a topical microbicide prevents rectal transmission of SHIV89.6P in macaques. *AIDS Res Hum Retroviruses.* 2003;19(7):535-41.
26. Bewley CA, Otero-Quintero S. The potent anti-HIV protein cyanovirin-N contains two novel carbohydrate binding sites that selectively bind to Man(8) D1D3 and Man(9) with nanomolar affinity: implications for binding to the HIV envelope protein gp120. *J Am Chem Soc.* 2001;123(17):3892-902.
27. Sandstrom C, Berteau O, Gemma E, Oscarson S, Kenne L, Gronenborn AM. Atomic Mapping of the Interactions between the Antiviral Agent Cyanovirin-N and Oligomannosides by Saturation-Transfer Difference NMR. *Biochemistry.* 2004;43(44):13926-13931.
28. Li H, Wang LX. Design and synthesis of a template-assembled oligomannose cluster as an epitope mimic for human HIV-neutralizing antibody 2G12. *Org Biomol Chem.* 2004;2(4):483-8.
29. Wang LX, Ni J, Singh S, Li H. Binding of high-mannose-type oligosaccharides and synthetic oligomannose clusters to human antibody 2G12: implications for HIV-1 vaccine design. *Chem Biol.* 2004;11(1):127-34.
30. Lee HK, Scanlan CN, Huang CY, Chang AY, Calarese DA, Dwek RA, Rudd PM, Burton DR, Wilson IA, Wong CH. Reactivity-based one-pot synthesis of oligomannoses: defining antigens recognized by 2G12, a broadly neutralizing anti-HIV-1 antibody. *Angew Chem Int Ed Engl.* 2004;43(8):1000-3.
31. Calarese DA, Scanlan CN, Zwick MB, Deechongkit S, Mimura Y, Kunert R, Zhu P, Wormald MR, Stanfield RL, Roux KH, Kelly JW, Rudd PM, Dwek RA, Katinger H,

- Burton DR, Wilson IA. Antibody domain exchange is an immunological solution to carbohydrate cluster recognition. *Science*. 2003;300(5628):2065-71.
32. Roux KH, Zhu P, Seavy M, Katinger H, Kunert R, Seamon V. Electron microscopic and immunochemical analysis of the broadly neutralizing HIV-1-specific, anti-carbohydrate antibody, 2G12. *Mol Immunol*. 2004;41(10):1001-11.
 33. Veazey RS, Shattock RJ, Pope M, Kirijan JC, Jones J, Hu Q, Ketas T, Marx PA, Klasse PJ, Burton DR, Moore JP. Prevention of virus transmission to macaque monkeys by a vaginally applied monoclonal antibody to HIV-1 gp120. *Nat Med*. 2003;9(3):343-6.
 34. Dacheux L, Moreau A, Ataman-Onal Y, Biron F, Verrier B, Barin F. Evolutionary Dynamics of the Glycan Shield of the Human Immunodeficiency Virus Envelope during Natural Infection and Implications for Exposure of the 2G12 Epitope. *J Virol*. 2004;78(22):12625-37.
 35. Binley JM, Wrin T, Korber B, Zwick MB, Wang M, Chappey C, Stiegler G, Kunert R, Zolla-Pazner S, Katinger H, Petropoulos CJ, Burton DR. Comprehensive cross-clade neutralization analysis of a panel of anti-human immunodeficiency virus type 1 monoclonal antibodies. *J Virol*. 2004;78(23):13232-52.
 36. Hindsgaul O. 1999. Protein-glycan interactions. In *Essentials of Glycobiology* (ed Varki A, *et al*), pp. 41-56, Cold Spring Harbor Laboratory Press, New York.
 37. Winzeler EA, Shoemaker DD, Astromoff A, *et al*. Functional characterization of the *S. cerevisiae* genome by gene deletion and parallel analysis. *Science*. 1999;285(5429):901-6.
 38. Muller WE, Schroder HC, Reuter P, Maidhof A, Uhlenbruck G, Winkler I. Polyclonal antibodies to mannan from yeast also recognize the carbohydrate structure of gp120 of the AIDS virus: an approach to raise neutralizing antibodies to HIV-1 infection in vitro. *AIDS*. 1990;4(2):159-62.
 39. Ozinsky A, Underhill DM, Fontenot JD, Hajjar AM, Smith KD, Wilson CB, Schroeder L, Aderem A. The repertoire for pattern recognition of pathogens by the innate immune system is defined by cooperation between toll-like receptors. *Proc Natl Acad Sci U S A*. 2000;97(25):13766-71.
 40. Sato M, Sano H, Iwaki D, Kudo K, Konishi M, Takahashi H, Takahashi T, Imaizumi H, Asai Y, Kuroki Y. Direct binding of Toll-like receptor 2 to zymosan, and zymosan-induced NF-kappa B activation and TNF-alpha secretion are down-regulated by lung collectin surfactant protein A. *J Immunol*. 2003;171(1):417-25.

41. Mond JJ, Lees A, Snapper CM. T cell-independent antigens type 2. *Annu Rev Immunol.* 1995;13:655-92.
42. Berland R, Wortis HH. Origins and functions of B-1 cells with notes on the role of CD5. *Annu Rev Immunol.* 2002;20:253-300.
43. Lesinski GB, Westerink MA. Novel vaccine strategies to T-independent antigens. *J Microbiol Methods.* 2001;47(2):135-49.
44. Lockhart S. Conjugate vaccines. *Expert Rev Vaccines.* 2003;2(5):633-48.
45. Ada G, Isaacs D. Carbohydrate-protein conjugate vaccines. *Clin Microbiol Infect.* 2003;9(2):79-85.
46. Weis WI, Taylor ME, Drickamer K. The C-type lectin superfamily in the immune system. *Immunol Rev.* 1998;163:19-34.
47. Cutalo JM, Deterding LJ, Tomer KB. Characterization of glycopeptides from HIV-I(SF2) gp120 by liquid chromatography mass spectrometry. *J Am Soc Mass Spectrom.* 2000;15(11):1545-55.
48. Ji X, Gewurz H, Spear GT. Mannose binding lectin (MBL) and HIV. *Mol Immunol.* 2005;42(2):145-52.
49. Botos I, Wlodawer A. Proteins that bind high-mannose sugars of the HIV envelope. *Prog Biophys Mol Biol.* 2005;88(2):233-82.
50. Funatsu O, Sato T, Kotovuori P, Gahmberg CG, Ikekita M, Furukawa K. Structural study of N-linked oligosaccharides of human intercellular adhesion molecule-3 (CD50). *Eur J Biochem.* 2001;268(4):1020-9.
51. Herscovics A. Importance of glycosidases in mammalian glycoprotein biosynthesis. *Biochim Biophys Acta.* 1999 Dec 6;1473(1):96-107.
52. Hitzeman, R.A., Chen, C.Y., Dowbenko, D.J., Renz, M.E., Lui, C., Pai, R., Simpson, N.J., Kohr, W.J., Singh, A., Chisolm, V., Hamilton, R. and Chang, C.N. Use of heterologous and homologous signal sequences for secretion of heterologous proteins from yeast. *Methods Enzymol.* 1990;185:421-440.
53. Barr, P.J., Steimer, K.S., Sabin, E.A., Parkes, D., George-Nascimento, C., Stephans, J.C., Powers, M.A., Gyenes, A., Van Nest, G.A., Miller, E.T., Higgins, K.W. and Luciw, P.A. Antigenicity and immunogenicity of domains of the human immunodeficiency virus (HIV) envelope polypeptide expressed in the yeast *Saccharomyces cerevisiae*. *Vaccine* 1987;5:90-101.
54. Liu, W.F., Gao, D. and Wang, Z.N. Expression of the extracellular domain of the

- human immunodeficiency virus type 1 envelope protein and its fusion with β -galactosidase in *Saccharomyces cerevisiae*. *Clin Diagn lab Immunol*. 1998;5:592-594.
55. Herscovics A. Processing glycosidases of *Saccharomyces cerevisiae*. *Biochim Biophys Acta*. 1999;1426(2):275-85.
56. Camirand A, Heysen A, Grondin B, Herscovics A. Glycoprotein biosynthesis in *Saccharomyces cerevisiae*. Isolation and characterization of the gene encoding a specific processing alpha-mannosidase. *J Biol Chem*. 1991;266(23):15120-7.
57. Smith, R.A., Duncan, M.J. and Moir, D.T. Heterologous protein secretion from yeast. *Science* 1985;229:1219-1224.
58. Rudolph, H.K., Antebi, A., Fink, G.R., Buckley, C.M., Dorman, T.E., LeVitre, J., Davidow, S., Mao, J., Moir, D.T. The yeast secretory pathway is perturbed by mutations in *PMR1*, a member of a Ca^{2+} ATPase family. *Cell*. 1989; 58:133-145.
59. Livi, G.P., Ferrara, A., Roskin, R., Simon, P.L. and Young P.R. Secretion of *N*-glycosylated human recombinant interleukin-1 α in *Saccharomyces cerevisiae*. *Gene* 1990; 88:297-301.
60. Harmsen, M.M., Langedijk, A.C., van Tuinen, E., Geerse, R.H., Raué, H.A. and Maat, J. Effect of a *pmr1* disruption and different signal sequences on the intracellular processing and secretion of *Cyamopsis tetragonoloba* α -galactosidase by *Saccharomyces cerevisiae*. *Gene* 1993; 125:115-123.

Claims

1. A composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising an isolated glycosylated polypeptide comprising at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.
2. The composition of claim 1, wherein the polypeptide comprises at least three N-linked high-mannose oligosaccharides.
3. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of gp120 or a fragment thereof.
4. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of a fungus glycoprotein or a fragment thereof.
5. The composition of claim 1, wherein the polypeptide comprises the amino acid sequence of a yeast glycoprotein or a fragment thereof.
6. The composition of claim 5, wherein the yeast is *Saccharomyces cerevisiae*.
7. The composition of claim 6, wherein the yeast glycoprotein is selected from the group consisting of PST1, ECM33, Gas1, Gas5, GP38, and YJL171c.
8. The composition of claim 5, wherein the yeast is *Pichia pastoris*.
9. The composition of claim 5, wherein the yeast is *Candida albicans*.

10. The composition of claim 4, wherein the polypeptide comprises the amino acid sequence of a fungus glycoprotein which is a homolog to PST1, ECM33, Gas1, Gas5, GP38, or YJL171c of *Saccharomyces cerevisiae*.

11. The composition of claim 1, wherein at least 90% of the N-linked glycans on the polypeptide are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.

12. The composition of claim 1, wherein at least 90% of the N-linked glycans on the polypeptide are Man₈GlcNAc₂.

13. The composition of claim 1, wherein the pharmaceutically acceptable excipient is an adjuvant.

14. A composition for eliciting antibodies specific for a protein having an epitope recognized by antibody 2G12 in a subject, comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted pmr1 gene of *Saccharomyces cerevisiae* or disrupted homolog gene of pmr1 gene of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12, and a pharmaceutically acceptable excipient.

15. The composition of claim 14, wherein the mutant fungus is a mutant yeast.

16. The composition of claim 15, wherein the mutant yeast is *Saccharomyces cerevisiae* having disrupted pmr1 gene.

17. The composition of claim 16, wherein the glycosylated polypeptide comprises the amino acid sequence of PST1 or a fragment thereof.

18. A composition for eliciting antibodies specific for a protein having an epitope recognized by antibody 2G12 in a subject, comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted pmr1 and mnn1 genes of *Saccharomyces cerevisiae* or disrupted homolog genes of pmr1 and mnn1 genes of *Saccharomyces*

cerevisiae, wherein the glycosylated polypeptide is recognized by antibody 2G12, and a pharmaceutically acceptable excipient.

19. The composition of claim 18, wherein the mutant fungus is a mutant yeast.

20. The composition of claim 19, wherein the mutant yeast is *Saccharomyces cerevisiae* having disrupted *pmr1* and *mn1* genes.

21. The composition of claim 20, wherein the glycosylated polypeptide comprises the amino acid sequence of PST1 or a fragment thereof.

22. A composition for eliciting antibodies specific for a protein having an epitope recognized by antibody 2G12 in a subject, comprising a glycosylated polypeptide isolated from a mutant fungus having disrupted *och1* and *mn1* genes, or *och1*, *mn1*, and *mn4* genes of *Saccharomyces cerevisiae* or disrupted homolog genes of *och1* and *mn1*, or *och1*, *mn1*, and *mn4* genes of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12, and a pharmaceutically acceptable excipient.

23. A method of preparing a composition for eliciting antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, said method comprising:

- a) fermenting a mutant fungus, wherein the mutant fungus is mutated to produce a glycosylated polypeptide, wherein the glycosylated polypeptide comprises at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the glycosylated polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are $\text{Man}_9\text{GlcNAc}_2$, $\text{Man}_8\text{GlcNAc}_2$, or a combination thereof;
- b) purifying the glycosylated polypeptide from the mutant fungus cell lysate or supernatant; and
- c) formulating the glycosylated polypeptide with a pharmaceutically acceptable excipient.

24. The method of claim 23, wherein the mutant fungus is a mutant yeast.
25. The method of claim 24, wherein the mutant yeast is *Saccharomyces cerevisiae* having disrupted och1, mnn1, and mnn4 genes.
26. The method of claim 24, wherein the mutant yeast is *Saccharomyces cerevisiae* having disrupted pmr1 and mnn1 genes.
27. The method of claim 24, wherein the mutant yeast is *Saccharomyces cerevisiae* having disrupted och1 and mnn1 genes.
28. The method of claim 24, wherein the mutant yeast is *Pichia pastoris* having disrupted och1 gene.
29. The method of claim 24, wherein the mutant yeast is *Candida albicans* having disrupted och1 gene.
30. The method of claim 23, wherein the glycosylated polypeptide is a yeast glycoprotein or fragment thereof.
31. The method of claim 30, wherein the yeast glycoprotein is selected from the group consisting of PST1, ECM33, Gas1, Gas5, GP38, and YJL171c of *Saccharomyces cerevisiae*.
32. The method of claim 30, wherein the yeast glycoprotein is a homolog of PST1, ECM33, Gas1, Gas5, GP38, or YJL171c of *Saccharomyces cerevisiae*.
33. The method of claim 23, wherein the mutant fungus has been transformed with a vector comprising a nucleotide sequence encoding the glycosylated polypeptide before step a).
34. The method of claim 33, wherein the nucleotide sequence encodes gp120 or a fragment thereof.

35. A mutant fungus cell comprising a polynucleotide comprising a nucleotide sequence encoding gp120 or a fragment thereof comprising at least two N-glycosylation site, wherein the mutant yeast cell is capable of producing the gp120 or the fragment comprising at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, wherein the high-mannose oligosaccharides are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.

36. The mutant fungus cell of claim 35, wherein the mutant fungus is *Saccharomyces cerevisiae* having disrupted och1, mnn1, and mnn4 genes.

37. The mutant fungus cell of claim 35, wherein the mutant fungus is *Saccharomyces cerevisiae* having disrupted pmr1 and mnn1 genes.

38. The mutant fungus cell of claim 35, wherein the mutant fungus is *Saccharomyces cerevisiae* having disrupted och1 and mnn1 genes.

39. The mutant fungus cell of claim 35, wherein the mutant fungus is *Pichia pastoris* having disrupted och1.

40. A method of generating antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising administering to the subject an effective amount of the composition of claim 1.

41. A method of generating antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising administering to the subject an effective amount of the composition of claim 14.

42. A method of generating antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising administering to the subject an effective amount of the composition of claim 18.

43. A method of generating antibodies specific for a protein comprising an epitope recognized by antibody 2G12 in a subject, comprising administering to the subject an effective amount of the composition of claim 22.

44. A method of inducing neutralizing antibodies against a pathogen in a subject, comprising administering to the subject an effective amount of the composition of claim 1.

45. The method of claim 44, wherein the pathogen is HIV.

46. A method of inducing neutralizing antibodies against a pathogen in a subject, comprising administering to the subject an effective amount of the composition of claim 14.

47. The method of claim 46, wherein the pathogen is HIV.

48. A method of inducing neutralizing antibodies against a pathogen in a subject, comprising administering to the subject an effective amount of the composition of claim 18.

49. The method of claim 48, wherein the pathogen is HIV.

50. A method of inducing neutralizing antibodies against a pathogen in a subject, comprising administering to the subject an effective amount of the composition of claim 22.

51. The method of claim 50, wherein the pathogen is HIV.

52. A method of preventing or treating pathogen-induced disease in a subject, comprising administering to the subject an effective amount of the composition of claim 1.

53. The method of claim 52, wherein the pathogen is HIV.

54. A method of preventing or treating pathogen-induced disease in a subject, comprising administering to the subject an effective amount of the composition of claim 14.

55. The method of claim 54, wherein the pathogen is HIV.

56. A method of preventing or treating pathogen-induced disease in a subject, comprising administering to the subject an effective amount of the composition of claim 18.

57. The method of claim 56, wherein the pathogen is HIV.

58. A method of preventing or treating pathogen-induced disease in a subject, comprising administering to the subject an effective amount of the composition of claim 22.

59. The method of claim 58, wherein the pathogen is HIV.

60. A method for identifying an antibody that specifically binds to N-linked high mannose oligosaccharide, comprising detecting binding of an antibody to a glycosylated polypeptide comprising at least two N-linked high mannose oligosaccharides that are recognized by antibody 2G12, and a pharmaceutically acceptable excipient, wherein greater than 50% of the N-linked glycans on the glycosylated polypeptide are the high-mannose oligosaccharides, and wherein the high-mannose oligosaccharides are Man₉GlcNAc₂, Man₈GlcNAc₂, or a combination thereof.

61. The method of claim 60, wherein the antibody is in a serum of a subject infected by HIV.

62. The method of claim 60, wherein the glycosylated polypeptide is in a mutant yeast cell lysate.

63. A method for identifying an antibody that specifically binds to N-linked high mannose oligosaccharide, comprising detecting binding of an antibody to a glycosylated polypeptide produced in a mutant fungus having disrupted och1 and mnn1, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae* or disrupted homolog genes of och1 and mnn1, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*, wherein the glycosylated polypeptide is recognized by antibody 2G12.

64. The method of claim 63, wherein the antibody is in a serum of a subject infected by HIV.

65. A composition comprising a whole cell of a mutant yeast, and a pharmaceutically acceptable excipient, wherein the mutant yeast has disrupted pmr1 and mnn1 genes, och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*, or disrupted homolog genes of pmr1 and mnn1 genes, och1 and mnn1 genes, or och1, mnn1, and mnn4 genes of *Saccharomyces cerevisiae*.


**Figure 1**



FIGURE 2

FIGURE 3

4/30




A**B****C****Figure 4**

FIGURE 5






FIGURE 6**A****B**

FIGURE 7


**Figure 8**



Figure 9

Figure 10




Figure 11



Figure 12

MQLHSLIAST**ALLITSALAATSSSSSIPSSCTISSHATATAQSDLDKYSRCDTLVGNL**TIGGGLKT**GALANVKEI
 NGSLTIFNATNLTSFAADSLESITDSLNLQSLTILTSASFGSLQSVDSIKLITLPAISSFTSNIKSANNIYISDT
 SLOSVDGFSALKKVVNFNVNNK**LTSIK**SPVETVSDSLQFSFNGNQTKITFDDLWANNISLTDVHSVSVFANLQ
 KINSSLGFINNSISSLNFTKL**NTIGOTFSIVS**NDYLKNLSFSNLSTIGGALVVANNTGLQKIGGLDNLT**TIGGTL**
 EVVGNFTSLNLD**SLKSVKGGADVESKSSNFSCN**ALKALQKKGGIKGESFVCKNGASSTSVKLSSTSKSQSSQTTA
 KVS**KSSSKAE**EKKFTSGDIKAAASASSVSSSGASSSSSKSSKGNA**AIMAPIGQTTPLVGLLTA**IMSIM**

Figure 13




Figure 14




Figure 15




Figure 16

14/30




Figure 17



Figure 18




Figure 19

PST1 HomoloGene among 18 Species		
Mammalian	Homo sapiens	Human
	Pan troglodytes	Chimpanzee
	Canis familiaris	Dog
	Mus musculus	House mouse
	Rattus norvegicus	Rat
Bird	Gallus gallus	Chicken
Insect	Drosophila melanogaster	Fly
	Anopheles gambiae	Mosquito
Worm	Caenorhabditis elegans	Small soil nematode
Fungus	Saccharomyces cerevisiae	Yeast
	Schizosaccharomyces pombe	Yeast
	Kluyveromyces lactis	Yeast
	Eremothecium gossypii	Cotton pathogen
	Magnaporthe grisea	Rice blast fungus
Plant	Neurospora crassa	Red bread mold
	Arabidopsis thaliana	Small flowering plant
	Oryza sativa	Japanese rice
Parasite	Plasmodium falciparum	malaria parasite

Figure 20






Figure 21

17/30

A**B****Figure 22**




Figure 23




Figure 24




Figure 25




Figure 26




Figure 27




Figure 28




Figure 29




Figure 30





Figure 31

**Figure 32**


PST1**A.**

MQLHSLIA**STALLITS**SALAATSSSSSIPSSCTISSHATATAQSDLDKYSRCDTLVG**NLTIGGGLKT**GALANVKEI
 NGSLTIFNATNLTSFAADSLESITDSLNLQSLTILTSASFGSLQSVDSIKLITLPAISSFTSNIKSANNIYISDT
 SLOVDGFSALKKVNVFNVNNNKKLTSIKSPVETVSDSLQFSFNGNQTKITFDDLWANNISLTDVHVSFANLQ
 KINSSLGFINNSISSLNFTKLNTIGOTFSIVSNDYLKNLSFSNLSTIGGALVVANNTGLQKIGGLDNLTTIGGTL
 EVVG**NFTSLNLD**SLKSVKGGADVESKSSNFSCNALKALQKKGGIKGESFVCKNGASSTSVKLSSTSKSQSSQTTA
 KVS**KSSSKAEKKFTSG**DIKAAASASSVSSSGASSSSSSKSSKGNA**AIMAPIGQTTPLVGLLTAIIM**SIM

B.

.....T.....SS.T.....

T.....SS.T.....
T.....S....SSS...S.....

**Figure 33**

ECM33:

A.

MQFKNALTATAILSASALAAANSTTSIPSSCSIGTSATATAQADLDKISGCSTIVGNLTITGDLGSAALASIQEI
DGLTIFNSSSLSSFSADSIKKITGDLNMQELIILTSASFGSLQEVDSINMVTLPALSTFSTDLOANNIIVSDT
TLESVEGFSTLKKVNVFNINNNRYLNSFQSSLESVSDSLQFSSNGDNTTLAFDNLVWANNITLRDVNSISFGSLQ
TVNASLGFINNLTPLSLNLTQLSKVGQSLIVSNDELSKAAFSNLTTVGGGFIIANNNTOLKVIDGENKVQTVGGAI
EVTGNFSTLDLSSLKSVRGGANFDSSSSNFSCNALKKLQSNGLQGDSEFVCKNGATSTSVKLSSTSTESSKSSAT
SSASSSGDASNAQANVSASASSSSSSSSKSKSGAAPELVPATSFMGVVAAGVAVYKIKATICVSIITLISSLMIS
LPFLFYETVGSSSLNFICR

B.

.....TT.....T..T.....
.....
.....
.....
.....SSTST.SS.SS.T
SS.SSS...S.....S.....




Figure 34

GP38

A.

MKFQVVL~~S~~ALLACSSAVVA~~S~~PIENL~~F~~KYRAVKASH~~S~~KNIN~~S~~TLP~~A~~WN~~G~~SN~~S~~SNVTYANGTNSTNTTTAESSQLQ
 IIVTGGQVPITN~~S~~SLTHTNYTRLFN~~S~~SSALNITELYNVARVVNETIQDKSSAGAVV~~V~~ANAKSLEAVSFFF~~S~~IIFD
 TEKPIVVTE~~D~~SAYAI~~P~~VANNKNAT~~K~~RGVLSVTSDKLVYSGVFTPTACSYGAGLPVAIVDDQDEVKWF~~F~~DASKPT
 LISSDSIIRKEYSN~~F~~TTPYGLLENGVPIVPIVYDGGYSSSLIDSLSSAVQGLVVVSSGSTNST~~S~~STIESTEIPVV
 YAQANTPLN~~F~~IDNKDVPKNAV~~G~~AGYLSPIKAQILL~~S~~IAAVNGVT~~S~~KSALESIFP

B.

.....

T..T..T...T.....




Figure 35

YJL171c

A.

MLQSIVLSVCMFMLHTVAASGPQSYQKLDFTNVGFTGSYVDNKFKDITNNESCTCEVGDRVWFSGKNAPLADYL
SVHERGPLKLKQFAFYTSPGFTVNNSRSSSDNRLAYYESSSKTADNVTFLNHGGEASPCLGNALSYASSNGTGS
ASEATVLADGTLISSDQEYIIYSNVSCPKSGYDKGCGVRSGIPAYYGYGGTTKMFLFEFEMPTETEKNSSSIGY
YDLPAIWLLLNDHIARTSQYPTNANCSCWASGCGEYDIFEAMNGTEKNHLYSTFHTFOGIEDLGTGIOSYGYITRN
TTGTMKGGVVFDSSGNVVSFISDATPFNGTVSADTVNDLLAIPENETYSSQLMSISATAPSTTSLSNGVRLTNM
QNGVWYYILAIFTATTQVVLI

B.

.....
.....
.....
.....
.....
.....

NetOGlyc 3.1: predicted O-glycosylation sites in Sequence





Figure 36

Gas1**A.**

MLFKSLSKLATAAAFFAGVATADDVPAIEVVGNKFFYSNNGSQFYIRGVAYQADTANETSGSTVNDPLANYESCS
RDIPYLKKLNTNVLIRVYAINTTLDHSECMKALNDADIYVIADLAAPATSINRDDPTWTVDLFNSYKTVVDTFANY
TNVLGFFAGNEVTNNYTNTDASAFVKAIRDVROYSIDKNYRKIPVGYSSNDDTRVKMADYFACGDDDDVKADF
YGINMYEWCGKSDFKTSYADRTAEFKNLSIPVFFSEYGCNEVTPRLFTEVEALYGSNMTDVWSGGIVYMYFEET
NKYGLVSIIDGNDVKTLDDEFNNYSSEINKISPTSAANTKSYSATTSDVACPATGKYWSAATELPPTPNGGLCSCMNA
ANSCVVSDDVDSDDYETLFNWCNEVDCSGISANGTAGKYGAYSFCTPKEQLSFVMNLYYEKSGGSKSDCSFSGS
ATLQTATTQASCSSALKEIGSMGTNSASGSVDLGSGTESSTASSNASGSSSSKSNSSGSSGSSSSSSSSSSSSSSSSSSSS
KKNAATNVKANLAQVVFTSIISLSIAAGVGFALV

B.

.....
.....
.....
.....
.....T.....T.....
.....T...T.SS..S.SSS.S.S.SS.SSSSSSSSS.SSS...
.....

**Figure 37**

MLRLSLTSAFVLSAGLAQAASSN̄S̄STPSIEIKGNAFFNSESGERFYIRGVDYQPGGSSN̄LTDPLADASVCDRDV
PVLKDLGINTVRVYTVDNSQDHSCHMKLLQENGIYLILDVNTPTSAISRYDPACSYNADYLQNVFATIDTFADYD
NVLGFFAGNEVINSVNTTNTATYVKAVVRDMKKYIKARKYROI PVGYSAADIVANRQLAAEYFNCGDEADARIDM
FGVNDYSWCGESSFVVS̄GYSTKMKLYQDYSVPVFLSEFGCNQVKSSRPFTEIEAIYSTQMSSVFSGGLVYEYSN̄Ē
TN̄NYGLVQIDGDKVTKLTD̄FENLKNEYSKVSNPEGNNGYSTSN̄NYSTCPDYEKGVWEANN̄TL̄PAMPSAASAYFTS
GAGSPMGTGIATQQSCDAKDDDDDEEDDDTSSSSSSSSSSSSSSASSSESSSSSTSKASSSSPSASETSLKSAASA
TSSSQSSSKSKGAAGIIEIPLIFRALAELYNLVL

TSSSSSSSSSSSSSSSS.SSSS.SSSSTS..SSSS.S.S.TS...S..S.

C.




Figure 38




Figure 39




Figure 40