

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
27 February 2003 (27.02.2003)

PCT

(10) International Publication Number
WO 03/016860 A2

- (51) International Patent Classification⁷: **G01N**
- (21) International Application Number: PCT/US02/25761
- (22) International Filing Date: 14 August 2002 (14.08.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
60/312,322 14 August 2001 (14.08.2001) US
- (71) Applicant (for all designated States except US): **WASHINGTON UNIVERSITY IN ST. LOUIS** [US/US]; One Brookings Drive, St. Louis, MO 63130 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **ELSON, Elliot** [US/US]; 6322 McPherson, University City, MO 63130 (US). **MCCONNAUGHEY, William, B.** [US/US]; 3814 Utah Street, St Louis, MO 63116 (US). **WAKATSUKI,**

Tetsuro [JP/US]; 546 Warder Avenue, University City, MO 63130 (US).


(74) Agents: **BEULICK, John, S.** et al.; Armstrong Teasdale LLP, Suite 2600, One Metropolitan Square, St. Louis, MO 63102 (US).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

[Continued on next page]

(54) Title: SYSTEMS AND METHODS FOR SCREENING PHARMACEUTICAL CHEMICALS



(57) Abstract: A method for obtaining a response of a tissue model system to an activator includes contacting a bio-artificial tissue model system with an activator and measuring cellular mechanical response thereto of at least one of contractile force and tissue stiffness. A method for obtaining a response of a tissue model system to an activator includes contacting a bio-artificial tissue model system with an activator and measuring cellular mechanical response thereto of at least one of contractile force and hysteresis.



Published:

— without international search report and to be republished
upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

SYSTEMS AND METHODS FOR SCREENING PHARMACEUTICAL CHEMICALS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. provisional patent application No. 60/312,322 filed August 14, 2001, which is incorporated herein in its entirety by reference.

BACKGROUND OF THE INVENTION

[0002] This invention relates to systems and methods for screening pharmaceutical chemicals. More in particular, this invention relates to systems and methods for screening pharmaceutical chemicals in tissue models.

[0003] The number of pharmaceutical chemicals that must be tested for efficacy has increased enormously, mainly due to the development of large libraries of chemicals obtained using parallel and combinatorial chemical synthesis methods. Correspondingly, the number of identified therapeutic targets such as receptor and intracellular regulatory proteins has been greatly increased by the application of functional genomics. Therefore, there is a great need for rapid and quantitative methods with which to screen pharmaceutical chemical(s) for their ability to elicit specific cellular responses and to identify leading pharmaceutical candidates.

[0004] Initial screening of large libraries of chemicals is carried out by testing for specific binding to target molecules using assay methods that operate in solution. High throughput methods have been developed based on scintillation proximity assay or fluorescence detection techniques (Sundberg, 2000). These methods, while readily adapted to screen thousands of compounds per day, provide information only about the strength and specificity of chemical interaction, not about cell response. Hence, chemicals that are initially selected based on their ability to bind to a target in solution must be rescreened to assess their ability to elicit a desired cellular response. These secondary and tertiary levels of screening add increased expense and time to the process of detecting promising or lead pharmaceutical chemicals.

[0005] Stimulation of receptors and activation of ion channels have been assessed using fluorescence methods to detect changes in, e.g., calcium ion concentration and, membrane potential and pH (Sundberg, 2000). These changes in ion concentration and transport often occur relatively early in the process of signal transduction and lead to more specific end responses such as the activation of specific enzymes. Hence, measurement of these responses does not necessarily provide information about the ultimate cellular responses that are activated or inhibited by a test pharmaceutical compound.

BRIEF SUMMARY OF THE INVENTION

[0006] In one embodiment, a method is provided for obtaining a response of a multi-cell tissue model system to an agent which comprises contacting a tissue model with the agent and determining the cellular mechanical response to that contact of at least one of contractile force and tissue stiffness.

[0007] In another embodiment, a system is provided for obtaining a response of a tissue model system to an agent which comprises constructing a tissue model, providing an agent, and measuring cellular mechanical response to contact of the agent to the system of at least one of contractile force and hysteresis.

[0008] In a further embodiment, a system is provided for obtaining a mechanical response profile based on mechanical measurements of the response of reconstituted muscle and nonmuscle tissue models to an agent, which comprises constructing a tissue model system having cells reconstituted in collagen and contacting the reconstituted cells with an agent. The mechanical response comprises at least one of contractile force and tissue stiffness.

[0009] In a further embodiment, a method is provided for screening pharmaceuticals, which comprises contacting a tissue model comprising reconstituted cells in collagen with an effective amount of a pharmaceutical chemical and measuring cellular response to the chemical in terms of at least one of contractile force or tissue stiffness.

[0010] In a further embodiment, a method is provided for managing a library of one or more pharmaceuticals or pharmaceutical chemicals which comprises obtaining a profile of a mechanical response to the contact of an agent with a tissue model, storing that profile in a database, storing at least one additional profile of another pharmaceutical in the database, setting up a means for comparing more than one profile with another profile, comparing the profile of a first pharmaceutical with a profile of a second pharmaceutical based on a pre-established or ordered standard/hierarchy of comparison and ranking the pharmaceuticals in an order of activity with respect to mechanical effect on a tissue model.

[0011] In a further embodiment, a method is provided for obtaining a multi-parameter mechanical response profile for a tissue model contacted with a pharmaceutical measuring the cellular response thereof.

[0012] In a further embodiment, a tissue model is provided, the tissue model comprises tissue assembled in the form of a ring mounted on a system comprising an isometric force transducer electrically coupled to a computer. The tissue model spans the isometric force transducer and a computer-controlled stepping motor provides an application of stretching and strain to the tissue.

[0013] In a further embodiment, a tissue model is provided wherein said tissue model comprises a membrane of reconstituted tissue supported by a frame. The mechanical properties of the membrane of reconstituted tissue are determined from resistance to stretching as the tissue is moved against a probe attached to an isometric force transducer.

[0014] In a further embodiment, a method is provided for preparing a tissue model, said method comprises placing self assembling tissue in a shape on a support, and subjecting said shaped tissue to an application of stretch by application of indentation.

[0015] In a further embodiment, a method is provided for establishing a mechanical response profile of a pharmaceutical which comprises contacting or relaxing a tissue model with a pharmaceutical and determining the

mechanical response(s) of the tissue model in terms of at least one of contractile force and stiffness.

[0016] In another aspect, a method of culturing a cardiac tissue is provided using a medium(s) conditioned by a fibroblast(s).

[0017] In another aspect, this invention comprises a method to identify the effect of a drug upon a tissue which comprises treating a tissue using this invention and measuring the effect so resulting, and thereafter comparing such measured effect with a standard drug effect on the same or similar tissue.

[0018] In another aspect, this invention comprises a method to data mine a library of pharmaceutical moieties for activity in tissue, which comprises treating a tissue using this invention and measuring the effect so resulting. Optionally, the measured effect may be compared with a correspondingly measured effect of a standard drug on the same or similar tissue.

[0019] In another aspect, this invention comprises a method of optimizing tissue culture conditions for constructing implantable artificial cardiac tissue, with additional different factors influencing cardiac tissue development including growth factors, and matrix proteins and hormones to a tissue model of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] Figure 1 is a schematic of an illustrative method for preparing and measuring fibroblast populated matrices (FPMs) (including live tissue models). Figure 1 illustrates an exemplary method of preparation and measurement overall.

[0021] Figure 2 illustrates a force response to stretch from a tissue model and illustrates a profile obtained from a sequence of four stretch cycles applied to one FPM plotted over time. Profile a of Figure 2 illustrates a substantially larger peak force for a first stretch than for subsequent cycles. Profile b of Figure 2 illustrates the excellent reproducibility of the measurements by showing force responses of measurements carried out on two FPM's.

[0022] Figure 3 is a graph, which displays tissue stiffness as a function of cell generated force. Figure 3 a shows the change in force and dynamic stiffness in response to activation by fetal bovine serum. Figure 3 b shows the change in force and dynamic stiffness resulting from disruption of the actin cytoskeleton by 2 μ M cytochalasin D (CD) Figure 3 c is a summary of the data from both 3a and 3b to show the overall (linear) dependence of dynamic stiffness on force.

[0023] Figure 4 displays a dependence of contractile force and dynamic stiffness on CD and Latrunculin B (LA-B, Calbiochem-Novabiochem Corporation, San Diego, CA) concentrations. Figures 4a and 4b show contractile force and stiffness, respectively, generated by the tissue models after treatment with various concentrations of CD. Figure 4c and 4d show contractile force and stiffness, respectively, generated by the tissue models after treatment with various concentrations of LA-B.

[0024] Figure 5 shows the degrees of cardiac myocytes spreading in various tissue culture conditions. Figure 5a shows well spread cardiac myocytes in a cardiac tissue model made in a conditioned medium prepared with cardiac fibroblasts. Cardiac myocytes cultured in Dulbecco's modified Eagle's medium (DMEM) with 10 % fetal bovine serum, shown in Figure 5b, do not spread well. Cardiac myocytes cultured on tissue culture dishes with and without conditioned medium are shown in Figures 5c and 5d, respectively. The supplemental table in the Figure 5 shows spontaneous contractility of cardiac tissue models made with various culture conditions.

[0025] Figure 6 is a stainless steel spacer used to culture model tissues after removal from the molds shown in Figures 1c and 1e.

[0026] Figure 7 shows a time sequence of images (1 through 32 with a 200-msec. interval) capturing spontaneous contraction of cardiac tissue models made using cells from embryonic mouse hearts.

[0027] Figure 8 shows photo images of regular and small size tissue molds and the tissue models. Figure 8a shows disassembled small and regular size

molds schematically shown in Figure 1e. Figure 8b shows assembled small and regular size molds schematically shown in Figure 1c. Figure 8c shows photo images of small size tissue models made using small size mold held by a spacer. Figure 8d shows photo image of regular size tissue models held by the spacer schematically shown in Figure 6.

[0028] Figure 9 is a comparison of force traces observed using small and regular size cardiac tissue models made using these molds. The change of force with time shown in Figure 9a is observed using a cardiac tissue model made with the regular size mold. The change of force with time in response to stretch and treatment with Y27632, an inhibitor of the activation of myosin activation by Rho kinase, is shown in Figure 9b. The insert of Figure 9b shows an enlarged plot of force and time during the sample stretch spontaneously contracting.

[0029] Figure 10 is a comparison of data variance for measurements of tissue stiffness using the reconstituted tissue ring method (left side) and a single cell indentation method (right side).

[0030] Figure 11a shows a high throughput system illustrating use of triangular and rectangular (alternative shape) frames shown in Figure 11b, made of stainless steel wire about one millimeter in diameter, which provide scaffold supports on which reconstituted tissues form to provide samples from an indentation method of measuring tissue contractile force and stiffness.

[0031] Figure 12 shows that after the peak contractile force reaches its steady level the tissue model is activated with 20% Fetal Bovine Serum (FBS) (arrow a in Figure 12(a) to cause an increase in force. This amount of FBS activates fibroblast non-muscle myosin. Figure 12(b) shows a plot of force verses indentation depth for the same data as for Figure 12(a) including also the effect of adding 40 nM CD and 2 μ M CD.

[0032] Figures 13(a), 13(b) and 13(c) show changes in the peak tissue contractile force and area (arbitrary unit) of hysteresis as relates to Figure 12.

[0033] Figure 14 illustrates a series of force responses to indentations on reconstituted tissue membranes. A small membrane in a well of a 96-well plate was repeatedly indented (each peak represents one indentation.) Force responses to the indentations were recorded over 70 minutes. Peaks of indentation increased due to activation of myosin by addition of 10% FBS, whereas cytochalasin D (CD) reduced peaks in a dose dependent manner by disrupting the actin cytoskeleton. (Force scale was not offset to zero.)

[0034] As used herein, the term “agonist” includes a chemical substance that activates a cellular response.

[0035] Figure 15 shows the resulting forces (dynes) of a series of tests wherein several agonists are added to a tissue model of a ring system with smooth muscle cells. Panel a, b, c, d, e, f and g are profiles of force response by treatments of thrombin, vasopressin, LPA, bradykinin, endothelin, norepinephrine and phenylephrine, respectively.

[0036] Figure 16 shows profiles of contractile responses wherein a series of agonists were applied to a tissue model of a ring system made from chicken (panel a-d) and rat (panel e-h) cardiac fibroblasts, respectively.

[0037] Figure 17 shows a reduction in contractile force by inhibition of myosin contractility initially stimulated by 20% FBS. Figure 17 shows the inhibition by different butane dione monoxime (“BDM”) concentrations of a contractile response previously stimulated by 20% FBS. Panel a, b, c and d are treated with 2, 4, 20, and 40 μ m BDM, respectively.

[0038] Figure 18 shows a profile of the increase in contractile force during tissue development in a ring system treated with no chemicals (a), herbimycin A (b), Nacodazole (C), and cytochalasin (D). The ring was mounted on the force measuring system about one hour after gelation of the collagen and prior to significant remodeling and compression of the matrix by the cells.

[0039] Figure 19a shows the dependence of force on strain for a tissue model ring measured after stimulation by FBS (labeled “Total”). The ring was then treated with 2 μ M CD and measured to yield the curve labeled “passive”. The difference between total and passive is labeled “active.” Figure 19b shows dynamic stiffness derived from the same measurements.

[0040] Figure 20 shows that cells that lack the appropriate collagen-binding integrins can neither develop force during the period of tissue remodeling (Figure 20 a) nor in response to calf serum (Figure 20 b). These defects are corrected by restoring the missing subunit of the collagen binding integrin (α_2).

[0041] Figure 21 is a comparison of the change in contractile force after treatment with various concentrations of Y27632, observed using tissue model and articular strips.

[0042] Figure 22 shows a zymograph of MMPs secreted into the tissue culture medium by cells cultured in various conditions.

[0043] Figure 23 compares force responses to stretch of tissue models cultured with and without MMP inhibitor GM6001.

[0044] Figure 24 illustrates the parameters by which the change in force recorded over time in response to various treatments can be represented.

DETAILED DESCRIPTION OF THE INVENTION

[0045] There are an estimated 20-30 trillion cells in the human body apportioned among tissues with distinct characteristics and functions. These cells include muscle and non-muscle cells. Muscle cells develop contractile force. Muscle cells respond to nerve signals, which send out messengers such as calcium ion and cyclic AMP that regulate the by generation of contractile force. Non-muscle cells, e.g., fibroblasts and endothelial cells, respond to activators such as a polypeptide growth factor or hormones by generating contractile force.

[0046] As used herein the term “treated or contacted with” include exposed to, contacted with and placed in contact with. As used herein, the term “tissue model” includes reconstituted cells and tissues from living cells and extracellular matrix material. As used herein, the term “isometric force” includes the force change without substantial change in the tissue length or tissue physical dimensions. As used herein the term “extracellular matrix (ECM)” includes fibrin, fibronectin, laminin and similar constituents/components and synthetic materials such as polylactic acid and polyglycolic acid.

[0047] Tissue models such as the bio-artificial tissue models reconstituted from cells and extracellular matrix (ECM), simulate natural tissues. Such tissue models provide a polydisperse or monodisperse population of living cells in a uniform or substantially uniform collagenous matrix.

[0048] Cytoskeletal and matrix proteins control the forces exerted by a tissue model and the stiffness of a tissue model. Cells regulate cytoskeletal structure and remodel ECM to produce mechanical changes during tissue development and wound healing.

[0049] Contractile forces result from activation of non-muscle myosin. The increase of contractile force over time after activation can be measured in tissues reconstituted from muscle and non-muscle cells. Such cell responses can lead to reorganization of the cytoskeleton within the cells or of the extracellular matrix (ECM) in which the cells are embedded. The development of contractile forces and the mechanical effects of cytoskeletal and matrix reorganization provide indicators of cellular response to a candidate pharmaceutical(s). Changes of contractile forces and stiffness result from activation or deactivation of cellular myosin, from other cytoskeletal perturbations, or from perturbations of the extracellular matrix within which the cells are embedded.

[0050] Use of reconstituted tissue permits assembly of tissue models, which are tested using specifically isolated cell types or in combinations of cell types. Hence, responses of these reconstituted tissues to a candidate pharmaceutical provide mechanical measurements of these cell types without complicating contributions of

other cell types normally present in natural tissues. An association is established between a specific cell type and a profile of mechanical measurements of changes of force and stiffness that result when the cells are contacted by an agent.

[0051] Contractile force and tissue stiffness change due to contraction and relaxation of the cells within reconstituted tissue upon effective contact with an agent. As used herein the term “agent” includes one or more candidate pharmaceuticals, which may or may not have any pharmaceutical activity or cellular response. Agents also include activators, antagonists and the like. As used herein, the term “effective contact” means placing an effective amount of an activator in contact with the tissue model, for example, as by adding an agent to a tissue model. As used herein the term “pharmaceutical” means having to do with medicine, which affects an animal or a person.

[0052] Cells utilized are obtained from a variety of tissue sources so that the tissue models used as tissue systems provide tissue-specific information of responses to candidate pharmaceutical(s).

[0053] As used herein, the term “tissue stiffness” is the force required to stretch tissue by a defined amount. In other words, tissue stiffness is the ratio of force applied to the tissue to the extent of stretch of the tissue. The force required to stretch an elastic object increases as the extent of stretching of the object (the “strain”) increases. Biological tissues, however, are viscoelastic. That is, viscous forces that depend on the rate of stretching also contribute to resistance to stretch. Measurements of the contribution of viscous forces may be determined from the dependence of the stiffness on the rate of stretching. For linearly elastic materials the force increases linearly with strain, i.e. stiffness is constant, independent of strain. Biological tissues and reconstituted tissue models are mechanically nonlinear and tissue stiffness increases with strain. Over a range of forces the stiffness varies linearly with the force either generated within the tissue or externally applied to it. This linear variation indicates that the force is an exponential function of strain.

[0054] One method of measuring tissue stiffness includes subjecting the tissue in a tissue model system to a relatively large steady stretch and observing

the change in force as the strain increases. Such measurements of tissue stiffness enable measurements and determinations of parameters including hysteresis (area), phase lag and dynamic stiffness.

[0055] Once a preset strain is achieved during the measurement of tissue stiffness, the extent of stretching (strain) can then be reduced at the same rate in a return to the tissue's initial strain and stiffness values. A plot of force as a function of strain as strain decreases (unloading curve) is always at lower force levels than during the increase of strain (loading curve). The area enclosed by the aforementioned two curves is a hysteresis area, which is an indicator of tissue viscosity. The hysteresis area measures a loss of energy in the tissue during the respective loading-unloading cycle.

[0056] Another method of measuring tissue stiffness uses oscillatory stretch, i.e., periodic increases and decreases of strain at a defined frequency (e.g. sinusoidal) selected by an experimenter. The force increases and decreases correspondingly at the same frequency, but possibly with a shift in phase, i.e. a phase lag. The phase shift or phase lag is another indicator of the viscosity of the tissue models. The type of tissue stiffness measured by oscillatory stretch is "dynamic stiffness". Dynamic stiffness depends on the magnitude of the stretch (because tissue models are nonlinear) and on the frequency of oscillation because the tissues are effectively viscous.

[0057] In illustrative embodiments of this invention mechanical measurements are carried out on reconstituted tissue models. The invention provides a method of quantitatively characterizing mechanical properties of connective tissue models, such as fibroblast-populated matrices (FPM's), via uniaxial stretch measurements in response to contact with one or more activators.

[0058] The connective tissue models, which are composites of selected living cells and ECM (i.e. a bio-artificial system), usually Type I collagen, respond by stiffening due to contact with activators which activate contractile forces. In one embodiment illustrative of a tissue model system of this invention, the tissue is assembled in the form of a ring that is mounted on a system in which the tissue spans

an isometric force transducer and a computer-controlled stepping motor for measurements of contractile force and strain.

[0059] In another embodiment of this invention, especially suitable for high throughput screening, tissue stiffening is measured by an indentation method using a multi-well plate system. In the indentation method, contractile force is measured (as peak force) along with tissue stiffness, which is registered as a resistance to indentation of the tissue model by a probe contacting the tissue model. The probe is attached to a force transducer. Many tissue composites can be rapidly tested (high throughput) for their mechanical responses to reagents in this embodiment. The system is suitably adapted in size and design to use small amounts of tissue model and reagents.

[0060] Candidate pharmaceuticals that inhibit a contractile response can be screened for their ability to prevent a stiffening response evoked by a well characterized activating agent. Moreover, candidate pharmaceuticals that cause cells to remodel the collagen, thereby stiffening or softening the collagen, can be tested using the inventive methods disclosed herein.

[0061] This invention provides a system and method for characterizing and profiling the mechanical response(s) of reconstituted tissue models in contact with one or more agents. The invention allows rapid and quantitative screening of many potential activators or inhibitors of cell contraction, cytoskeletal change, cell-matrix interactions, and matrix remodeling. The methods herein provide a quantitative readout of changes in tissue stiffness, which can be calibrated to supply corresponding quantitative data on the extent of activation or inhibition of myosinmodulation of cytoskeleton components or of interaction of cell-matrix and properties of the matrix itself.

[0062] The systems and methods described and claimed herein are not limited to the specific embodiments described herein. In addition, components of each system and each method can be practiced independently and separately from other components and methods described herein. Each component and method can be used in combination with other components and other methods.

[0063] Collagens useful include collagen Classes 1-4 which include all Types I-XIII and combinations thereof.

[0064] Agents useful as activators include Fetal Bovine Serum (FBS), lysophosphatidic Acid (LPA); thrombin, growth factors including epidermal growth factor (EGF), platelet derived growth factor (PDGF), angiotensin-II, endothelin-1 and vasopressin and combinations thereof.

[0065] Inhibitors include those inhibitors which act on cell surface receptors including a receptor antagonist for angiotensin II receptor and also inhibitors that act within the cell. Inhibitors useful herein include those which inhibit the process of signal transduction pathways including genistein, herbimycin and agents which act on the cytoskeleton. These include cytochalasin D, latrunculin B, paclitoxol, nocodazole, calyculin A and butane-dione-monoxime (BDM) and combinations thereof.

[0066] The amount of agent(s) provided to the reconstituted cell is an effective amount which is generally from an amount in nanomolar quantity to an amount of about 100 millimolar. An effective amount is that amount which is sufficient to elicit a response from or by a tissue model.

[0067] There are several mechanical parameters which may be determined using embodiments of this invention.

Ring type system:

Baseline force	Force measured at 0 strain (no stretch).
Dynamic stiffness	Amplitude of force response to the sinusoidal stretch divided by amplitude of applied oscillatory strain. The measurements are taken at various strain levels. Dynamic stiffness can also be obtained during a ramp stretch.
Phase angle	Phase angle indicates time dependent viscosity of a sample. It is obtained by the angle of phase delayed between force response and sinusoidal driving function. The measurements are

normally taken at 0 strain but can be measured at various strains as long as the mean force level reaches a steady state for a short period of time.

*Storage modulus, G' In-phase component stress (force divided by cross sectional area of sample) in response to a sinusoidal stretch divided by the strain.

*Loss modulus, G'' Out-of-phase component stress in response to a sinusoidal stretch divided by the strain.

Peak force Peak force in response to a ramp stretch.

Hysteresis curve area Hysteresis curve is a plot of force response to a ramp stretch plotted against strain. Force response during the sample elongation is always higher than that during the sample shortening. The area enclosed by the two lines is related to the viscosity of the sample.

Tissue Indentation procedure:

Peak force Peak force in response to tissue indentation.

Hysteresis curve, area Hysteresis curve is a plot of force response to a tissue indentation. Force response during the indentation (loading) is always higher than that during the retraction (unloading). The area enclosed by the two curves is related to the viscosity of the sample.

* G' and G'' can be calculated from dynamic stiffness and phase angle by knowing the cross sectional area of samples. G' and G'' are parameters indicating the mechanical properties of samples independent of their sizes and shapes.

[0068] More details are set forth below regarding Figures 1-15. Although specific exemplary embodiments of methods and systems for using tissue models are described herein, the methods and systems are not limited to such specific exemplary embodiments.

[0069] More in particular, Figure 1 is an illustrative schematic of a ring method for preparing and measuring FPM's (illustratively shaped as a ring). In an embodiment, CEF's (2) and monomeric collagen (4) are mixed in DMEM (6) at pH 7 to form a suspension (8). This suspension (8) is poured into casting wells (10) having a mandrel (12) and polymerized at 37°C. The casting wells (10) are incubated for a day or more during which time the cells compress and remodel the polymerized collagen matrix. After incubation the mandrel (12) is removed from the casting well (10) and the FPM-ring (14) is removed gently from the mandrel (12). The FPM-ring (14) is connected to the force measuring apparatus (an isometric force transducer) (16) and a stepper motor (18) that controls and sets the tissue strain.

[0070] Figure 2 depicts how measured a force responsive to stretch of a FPM. Typical force responses from a sequence of four stretch cycles applied to one FPM were traced in time. The first stretch cycle (as shown in panel a) produces a substantially larger peak force than subsequent cycles. A much smaller (~ 6%) reduction in peak force is seen in each of the subsequent cycles (panel a). Panel b of Figure 2 illustrates the reproducibility of the measurement by showing stretch sequences from two FPM's. (solid....) (solid line and gray broken line). The effects of treatment by both 20% calf serum (increase in contractility) and 2 uM Cytochalasin D (abolition of active contractility of the tissue) are almost identical. The cycle time was set to 30 minutes instead of the previously used 60 minutes cycle time for this test to accommodate additional test manipulations. (The force-strain and stiffness-strain curves from the 30 and the 60 minute cycle times, however, are almost identical.)

[0071] Figure 3 shows data of measured stiffness as compared to cell-generated force. In this test, the tissue model was stretched sinusoidally by 0.3% at 1 Herz. The stiffness was measured as the peak-to-peak change in the force (approximated by the breadth of the trace at this time scale). Each stiffness point represents the average stiffness for 5 seconds.

[0072] Figure 3, graph (A) shows that stiffness changes with a magnitude and time course similar to force during FBS (5%) stimulation. Figure 3,

graph (B) shows actin filament disruption by Cytochalasin D (2 μ M). As in Figure 3 (A) stiffness changes correspond to force changes.

[0073] Figure 3, graph (C) shows stiffness plotted against force for measurement data shown in Figure 3 parts (A) and (B). The force-stiffness relation for the rise in force after FBS (closed circles) is very similar to the force-stiffness relationship for the decline in force after addition of Cytochalasin D (open triangles). Each circle point and triangle point of this Figure represents both force and stiffness. Stiffness increases approximately linearly with force.

[0074] Figure 4 illustrates the dependence of force and dynamic stiffness on CD and LA-B concentrations. The force and dynamic stiffness were significantly diminished at a CD concentration of 2 nM and continued to fall as the CD concentration was increased (A, B). In contrast, for LA-B the force and dynamic stiffness began to diminish only when the concentration reached 40 nM, and reached their minimum values at a concentration of $\sim$ 600 nM (C, D). This Figure illustrates the sensitivity of the measurement method, which can detect force and stiffness changes at low concentrations of CD that cause no changes in the actin cytoskeleton which can be detected by light microscopy.

[0075] Figure 5 illustrates a comparison of data variance in tissue model stiffness measured using reconstituted tissues with a ring method (left bar graph) and in cell stiffness measured by indentation of individual cells (right bar graph). Reconstituted tissue can provide statistically significant data with less data points as compared to single cell measurements. Fewer measurements are required to obtain statistically significant data using tissue models than using single cells. This is because, even in a cultured population of cells of a single type, there is substantial cell-to-cell variation for many experimental parameters. Hence, measurements on many individual cells must be averaged to characterize the population. Therefore high throughput screening devices that measure the biological activity of chemical compounds on single cells require many measurements to obtain statistically significant data. Since the tissue models tested contained at least 10,000 cells, each measurement represents an average of many cells in the tissue. Therefore, the

efficiency of high throughput screening on tissue models is improved over that on single cells.

[0076] Figure 11 illustrates an exemplary high throughput screening system utilizing an indentation method for measuring the mechanical response of this tissue system.

[0077] A triangular frame made of stainless steel wire one millimeter in diameter provides a scaffold on which reconstituted tissue forms. In this illustration, the wells are slightly tapered toward the bottom and the frame is securely positioned 1 mm above the bottom of the well (The non-polymerized solution of collagen containing cells and appropriate cell culture media as described above was poured into the wells filling them to a level 3 mm above the bottom (Figure 11 a). The 96-well plate was incubated at 37°C with 5% CO₂. During incubation, the cells compress the collagen matrix by squeezing out liquid thereby reducing the total volume by about ten fold.

[0078] Without the wire frame, the reconstituted tissue contracted into a small sphere floating in the tissue culture medium. The collagen matrix can be compressed into different shapes using different frame shapes such as a circle or rectangle shapes of a support such as a wire frame. The support can be one of metal, nonmetal, and plastic. In an embodiment, system is a self assembly cell system in which cells form a tissue model conforming to the shape of a frame or support.

[0079] On a triangular wire frame the cells form a membrane spanning among the three edges, which is illustrated in Figure 11a. Other wire frame shapes, such as one shown in Figure 11b, produced tissue strips with different widths and shapes. A porous support material such as a Velcro fastener was not utilized to facilitate tissue adhesion even to the non-porous stainless steel surfaces of the wire. The collagen was compressed to a greater extent at the outer portion of the membrane or strip. Therefore, this outer portion of the membrane can withstand the stress produced by the cells and prevent the stress ripping the membrane off from the wire frame.

[0080] Figure 12 shows data taken using the system exemplified in Figure 11 above. The data shows that after the peak force reaches its steady level the sample is stimulated with 20% Fetal Bovine Serum (FBS) (see arrow a in Figure 12a). This amount of FBS activates fibroblast non-muscle myosin producing a contractile force that stiffens the reconstituted tissues. At about 10 minutes after the FBS addition, there is about a 25% increase in the peak force of subsequent indentations (Figure 12a). Nearly 15 minutes after the addition to the medium of 40 nM CD (see arrow b of this figure), the peak force from subsequent indentations has decreased about 40% from its initial level (Figure 12). A further reduction of peak force was recorded 20 minutes after addition of 2 μ M CD. Figure 12(b) is a plot of the same data as the data plot shown in Figure 12(a), but Figure 12(b) shows force versus indentation depth for the same conditions as for Figure 12(a).

[0081] Figure 13 is a comparison of percent changes in the peak force and area of hysteresis during the test which generated the data of Figure 12 shown above. Figure 13(a) shows the peak force (dynes) measured prior to FBS treatment (no FBS) after 20% FBS, after 40 nMCD and after 2 μ MCD. Figure 13(c) shows the corresponding changes in peak force and hysteresis relative to the values obtained prior to FBS treatment.

[0082] The area of hysteresis changes to a greater extent upon stimulation and CD addition. Therefore, the area of hysteresis is a more sensitive parameter than the peak force for monitoring the changes in mechanical properties of the tissue model.

[0083] Figure 14 shows measurements of the mechanical properties of a miniaturized reconstituted tissue using a 96-well plate. The system can be used for high throughput screening since it allows rapid and parallel measurements using the multi-well plate.

[0084] Figure 14 shows the force trace of mechanical measurements conducted using 96-well plates. A small membrane of reconstituted tissue is supported by a stainless steel wire frame. A vertical bar attached to an isometric force transducer indents the membrane and measures the force resisting the indentation.

The addition of 10% FBS increases the force peak's height by 30%. The addition of CD reduced the height of the peaks by 60%.

[0085] By automating the procedures using the 96-well plate system or going to an even higher degree of parallelism, the basic concept can be expanded to high throughput applications using the invention. Libraries of compounds can be screened and managed based on their biological activities using tissue models.

[0086] Typically, contractile force increases rapidly over a period of a few minutes following contact of an agent with a tissue model system. The contractile force reaches a maximum value and after reaching that maximum value the contractile force may be sustained over a period of an hour or more or the contractile force may decline at a rate that is specific to the activator and a cell type. The reason for the relaxation of the force is not critical but this produces further information and data for characterization of force and stiffness response.

[0087] During activation, different pathways, involving specific sets of enzymes and co-activators, which can vary among different activating agents, carry the signal received at a cell surface receptor into the cell to activate contractile force. Hence, the development and maintenance of contractile force provides an indicator (or profile) of specific cellular responses to contact with exogenous activators and to the operation of transduction pathways from cell surface receptors to the activation of myosin. The profile for non-muscle cells includes the maximum contractile force value and the maintenance of the contractile force over time.

[0088] The invention is further described in the following examples which is not intended to limit or restrict the invention in any way.

EXAMPLES

[0089] PREPARATION OF TISSUE MODELS

[0090] Bio-artificial living tissue models were prepared which were reconstituted from cells and extracellular matrix (ECM). These models simulate natural tissues.

[0091] Chicken embryo fibroblasts (CEF's) isolated from 11-day chicken embryos (Spafas Inc., Preston, Connecticut) were maintained in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal calf serum (FCS), penicillin at 50 units/ml, and streptomycin at 50 microgram/milliliter. The CEF's used to make tissue models were passaged once or twice from primary cultures. Monomeric collagen solubilized in 0.02 M acetic acid (Upstate Biotechnology Inc., Lake Placid, NY) was neutralized at 4°C with 0.1 N NaOH and mixed with concentrated DMEM stock to yield a final normal DMEM concentration. The fibroblasts suspended by trypsin (Fig. 1a) were mixed with the collagen solution (Fig. 1b), the cell suspension was poured into Teflon casting wells (Fig. 1c), and the wells were incubated at 37°C with 5% CO₂ (Fig. 1d). The collagen polymerized within 15–30 min and fibroblasts were captured within the hydrated collagen gel. The collagen gel formed a ring (3 mm thick, 3 cm diameter) between the inner wall of the cylindrical well and the central mandrel. While in culture the cells compressed this ring, reducing its volume about 10-fold (thickness 200–300 µm). The ring could then be removed from the mandrel (Fig. 1e and Fig. 1f) and mounted on the measuring instrument (Fig. 1g) as described below.

[0092] The types of cells that can be used to form reconstituted tissues are not limited to CEF. Cell types that have been used to form suitable tissues for mechanical measurements include chicken embryo fibroblasts, chicken embryo cardiac fibroblasts, chicken embryo cardiac myocytes, rat cardiac fibroblasts, mouse myotubes, mouse skeletal muscle C₂/C₁₂ cell line, normal mouse mammary gland (NMuMG) cell line and its mutant lines lacking α_1 and α_2 integrins, mouse fibroblasts and their mutant cell lines, REF52 fibroblasts, A7R5 smooth muscle cells, CCL39 fibroblasts and NR6 fibroblasts and combinations thereof. The cells can be isolated from chicken embryos or obtained from American type culture collection (Manassas, VA).

[0093] MEASUREMENTS ON AND MECHANICAL ASSEMBLY OF A RING SYSTEM

[0094] Mechanical measurements were carried out using a ring system. After two days of incubation (serum starvation for the final 12–16 hours), the mandrel was removed from the casting well and the tissue model-ring was removed gently from the mandrel. As shown in Figure 1 the tissue model-ring was looped over the triangular hook connected to an isometric force transducer (Model 52-9545, Harvard Apparatus, South Natick, MA) by a fine, flexible gold chain. The ring was also looped over a horizontal bar connected to a sliding element moveable vertically and linearly by a stepper motor (P/N 1-19-3400 24V DC 1.8° step size, Haward Industry, St. Louis, MO) controlled by a micro-stepping driver (IM483 Intelligent Motion Systems) to measure stress and dynamic modulus of the sample. The micro-stepping driver was controlled by a personal computer with software which enabled the stepper motor to achieve smooth motion. An analog-to-digital signal converter (CIO-DAS1602/16, Computer Boards, Inc., Mansfield, MA) attached to the computer translated the voltage signal from the isometric force transducer to a digital signal for recording on a data recorder. The stepper motor controlled the stretching of the tissue. Force exerted by or on the tissue was transmitted to the force transducer by the gold chain.

[0095] The tissue sample was submerged in 50 ml Hepes-buffered DMEM (pH 7.4) in a thermo-regulated organ bath (Harvard Apparatus, South Natick, MA) maintained at 37°C. The two horizontal bars over which the ring was looped were initially set to hold the ring at its original contour length (corresponding to the circumference of the mandrel).

[0096] TESTING OF TISSUE MODEL SYSTEM

[0097] Typically, for stiffness measurements, the tissue model is subjected to a sequence of stretch cycles. In each cycle the tissue model is slowly stretched from 0 to 20% over 30 minutes and the tissue model is then returned at the same rate to its original length by actuation of the stepper motor. The resulting increase in contractile force during the first stretch is substantially larger than the contractile force in subsequent stretches (Fig. 2a). For each stretch after the first stretch there is a further small decrease in maximum contractile force, but the decrease

becomes negligible after sufficient pre-stretching. The tissue models produced contractile force in response to the activation of fibroblasts by serum. This contractile force was abolished by disrupting the actin cytoskeleton with cytochalasin D (2 μ M) (Fig. 2b).

[0098] The amplitude of the force response divided by the stretching amplitude corresponded to the dynamic stiffness of the sample subjected to a preselected sinusoidal length change. The dynamic stiffness can be measured at various frequencies and amplitudes. The dynamic stiffness and tension of the FPM were measured at various strain levels by elongating and shortening the sample at a constant rate (10 micrometers/minute) up to 20% strain with superimposed sinusoidal length change (typically, 20 μ m amplitude; 0.5% strain, 0.5 Hz frequency) (Fig. 1g). The apparatus is set to change the tissue length with prescribed rate and amplitude.

[0099] This determination may be accomplished by separating the cell and matrix contributions within the generated mechanical response profile since the cells, the matrix, and the interactions between the cells and matrix all contribute to the tissue stiffness of a reconstituted tissue model.

[00100] As shown in Figure 3, dynamic stiffness correlates almost linearly to the isometric force (Fig. 3c) while their changes were measured at the original tissue length in response to the FBS stimulation and CD addition (Fig. 3a, b). Therefore, both the isometric force and the dynamic stiffness are good indicators of the mechanical properties of reconstituted tissues. The application of this method to determine the dose-dependent decrease in the force and stiffness of tissue model caused by the disruption of actin cytoskeleton with two different agents is described below. Data from the examples provide characterization of mechanical properties of connective tissues models, such as fibroblast populated matrices (FPM's) via uniaxial stretch measurements. The tissue model(s) resemble natural tissues in their exponential dependence of stress on strain and linear dependence of stiffness on force at a given strain.

[00101] DETERMINATION OF CONTRIBUTIONS OF CELLS AND MATRIX TO MECHANICAL PROPERTIES OF TISSUE MODEL USING A RING SYSTEM

[00102] Determination of cellular contributions and matrix contributions to the mechanical properties of reconstituted tissue models was carried out as it is highly desirable to obtain a more specific profile of the reconstituted cellular response to an activator. Activating cellular contractile forces by calf serum and disrupting F-actin by CD provide active and passive components which respectively emphasize cellular and matrix mechanical contributions. In Figure 4,19, the force versus strain (a) and dynamic stiffness vs. strain (b) for a pre-stretched tissue model activated by 20% CS is shown by solid lines labeled "Total". The force and stiffness curves obtained after treatment by 2 μ M CD are shown by broken lines labeled "Passive". The difference between the Total and the Passive curves, labeled "Active", is shown by dotted lines. All the curves exhibit hysteresis. The "Active" curve increases approximately linearly with strain; the "Passive" curves increases approximately exponentially. The dynamic stiffness was measured at 0.5 Hz, (b). Note that the dynamic stiffness and the force vary similarly with strain. These measurements were carried out after a single pre-stretch cycle to avoid the large difference between the first and second stretches (see Figure 2). It has been determined that the "active" and "passive" curves report primarily on the cellular and matrix portions of the tissue model.

[00103] The contributions of the cells and matrix are separately (and adequately) assessed. The cellular contributions are eliminated by reducing the stiffness of the cells and effectively disconnecting the cells from the matrix by addition of an effective amount of an actin filament disruptor. This is accomplished by adding CD as a disruptor (for example). CD disrupts actin filament organization within cells as described in detail below. It is therefore possible to determine the effects of activators and inhibitors specifically on cells or on the matrix. This ability to examine separately these two mechanical systems provides further specificity to this method.

[00104] CD and Latrunculin-B (LA-B) are two agents that disrupt the actin filament cytoskeleton and therefore both prevent the development of contractile force and weaken cellular mechanical properties. The response of a tissue model to CD and LA-B including dose-dependent mechanical properties were determined.

[00105] As used herein the term “isometric force” includes the force change without substantial change in the tissue length or tissue physical dimensions.

[00106] The isometric force and the dynamic stiffness of two day-old tissue models, which have been serum starved for 16 hours prior to the experiment, were measured over a range of CD concentrations. The measurements were carried out serially on each tissue model, beginning with the lowest concentration of CD. For each tissue model, CD was added and force and stiffness were measured and then the CD concentration was increased for the next measurement (Fig. 4 A, B). The CD was dissolved in DMSO. The total amount of DMSO added to the organ baths containing the tissue model was less than 0.1% of the total volume of the DMEM. This quantity of DMSO had no significant effect on the force and stiffness of FPM's. The data shown were averaged over quadruplicated samples, and the same experiment was repeated at least twice. The force and dynamic stiffness were significantly decreased at a CD concentration as low as 2 nM. At this concentration, no effect was observed by confocal microscopy on the actin cytoskeletons stained with Rhodamine phalloidin in mono-layer cell cultures. Both force and dynamic stiffness continued to decrease as the CD concentration increased up to 2 μ M. At this concentration, the force was reduced almost to zero and the stiffness had nearly reached its minimum value. For CD concentrations higher than 2 μ M, the stiffness did not significantly diminish further (data not shown). The concentration of CD needed to reduce the force and dynamic stiffness by 50% was approximately 0.25 μ M.

[00107] Measurements of the effects of LA-B on the mechanics of tissue models demonstrated differences between the mechanism of action of LA-B and the mode of action of CD on the actin cytoskeleton. Incremental additions of LA-B and mechanical measurements on tissue models were carried out serially as in the studies of CD. The data were averaged for at least three samples, and the same test

was repeated at least twice. The concentration of LA-B needed to produce a significant effect on the tension and the stiffness of FPM's was much higher than that required of CD (Fig. 4D, E).

[00108] Force and stiffness had a sigmoidal dependence on LA-B concentration. The estimated half maximum concentrations for reducing the force and stiffness were 53 nM and 68 nM, respectively. The LA-B-dependent decrease of tension and stiffness was confined to a single decade of LA-B concentration, whereas the response to CD ranged over almost three decades. This strongly suggests that CD and LA-B operate by different mechanisms to disrupt the actin cytoskeleton. Small differences in the values of force at the high concentration limits of CD and LA-B (Fig. 4) were observed.

[00109] For a viscoelastic system the dynamic stiffness depends on both the elastic and viscous resistance to stretching. The viscous contribution can be measured by the phase angle, δ , between force and strain.

[00110] In these tests the change of phase angle, δ , due to disruption of the actin cytoskeleton was small (data not shown). Hence, in this system CD and LA-B had a relatively minor effect on the viscosity of the tissues. Therefore, it is reasonable to suppose that the viscous contribution of the cells to tissue model force and stiffness was also minor. These measurements illustrate how force and stiffness measurements rapidly and sensitively indicate the effects of these "inhibitors" via their effects on the actin filament system.

[00111] In an embodiment of this invention using an indentation (indention) procedure in a multi-well plate system, mechanical response parameters are determined that are averaged over at least 100,000 cells. Measurements on reconstituted tissue models obtained using this system were compared with those obtained by indentation of individual cells. (Pertersen et al. 1982, Zahalak et al. 1990). For example, stiffness of the reconstituted tissues was measured with good statistical significance with ten-fold fewer measurement points using the ring system than are required for cell indentation measurements on individual cells (Figure 10).

[00112] Many signal transduction pathways contribute to the mechanical properties of reconstituted tissues by regulating contractile force, the organization of the cytoskeleton, and the integrity of the extracellular matrix. Therefore, a wide range of intracellular and extracellular target molecules can be assayed by determining their effects on the mechanical properties of reconstituted tissues using this invention.

[00113] The cells in the reconstituted tissue models of this invention are in an environment that resembles their condition in natural tissues and organs. Therefore, results of the assays using this method yield results similar to those obtained using animal models. Some of the animal testing can be replaced by using tissue models. For example, some tests of agents acting on skin can be conducted using artificial living tissues.

[00114] In an embodiment, the aforescribed methods can also be used to detect the effects of toxic materials on the mechanical properties of reconstituted tissues. For example, the inventors discovered that a 10 % solution of ethanol reduces force and stiffness of tissue models significantly. The inventors observed that infection of cells in tissue models by viruses can diminish force and stiffness. Therefore, the aforescribed method can be utilized to determine the toxicological effects of substances and biological materials.

[00115] CARDIAC TISSUE MODELS

[00116] The tissue models made using cardiac myocytes isolated from chicken embryos become contractile spontaneously. Chicken embryo extracts (CEE, Life Technologies, Rockville, MD) have been known to be required to form spontaneously contracting artificial cardiac tissues (FASEB J 1997 Jul;11(8):683-94, FASEB J 2000 Apr;14(5):669-79). Media conditioned by cardiac fibroblasts can replace the CEE. The conditioned medium (CM) was produced by incubating a confluent monolayer of cardiac fibroblasts in 100 mm dish preincubated with DMEM supplemented with 10% FBS for 2 days. The medium is changed to DMEM containing no serum and is incubated for 24 to 48 hours to make the CM. The cardiac tissue models cultured with the CM supplemented with 10% FBS start spontaneously

contracting in 4 to 5 days without adding any CEE. The cardiac myocytes grown in the tissue models spread much better (Figure 5a) than those cultured in DMEM containing only 10 % FBS (Figure 5b). A similar difference is observed for cells growing on tissue culture dishes. The cardiac myocytes cultured in the CM (Figure 5c) spread much better and cover larger areas than those cultured in DMEM supplemented with only 10% FBS (Figure 5d). This suggests that factors secreted by the fibroblasts promote cardiac myocytes spreading and spontaneous contraction. Conditioning of medium by which cardiac myocytes are induced to spread and contract spontaneously is also accomplished by coculturing cardiac fibroblasts as a monolayer at the bottom of the tissue culture dishes incubating cardiac tissue models. The conditioning of medium promoting spontaneous contraction of the cardiac tissue models is also accomplished by coculturing connective tissue models containing cardiac fibroblasts. The fibroblasts can also be mixed with myocytes to form cardiac tissue models. This induces strong contraction in 4 to 5 days yet it also stops spontaneous contraction as early as 7 to 8 days of the culture. See the description of the various conditions in supplemental table in Figure 5.

[00117] CARDIAC TISSUE WITH MOUSE CELLS

[00118] Embryonic hearts are removed from stage E17-E19 mouse embryos. Cardiac myocytes are isolated by several collagenase (167 μ g/ml) digestions after 15 min. trypsin (.25%) treatment at 37°C. Isolated cells from the hearts are cultured on tissue culture grade plastic dishes for 1 hour to remove nonmuscle cells by the rate of adhesion to the tissue culture grade dishes (nonmuscle cells adhere much faster than the muscle cells). Non adherent cells are removed with medium and sedimented with low speed centrifugation for 15 min. For 1ml of artificial tissue, 1 million myocytes were mixed with 0.75mg of rat tail collagen kept in acetic acid (0.02 N), which is neutralized by adding an appropriate amount of NaOH (0.1N), and 0.25 mg of fibrinogen kept in phosphate buffered saline. A higher concentration (more than 1X) of tissue culture medium (DMEM) is added to the sample solution to maintain a normal final concentration of the medium. One μ l of thrombin (1 unit/ml) was also added to the solution to initiate fibrin formation. The fibrin converted from fibrinogen polymerizes with collagen to form a stronger gel, which facilitates handling

the sample. The 0.5 ml of sample solution was poured into a mold described previously and incubated in the tissue culture incubator (5% CO₂ at 37°C) for 30 min. The gel is removed from the mold and cultured with the spacer -- two stainless steel bars (~1 mm in diameter) separated by a stainless steel block (Figure 6). After incubation of 5–7 days embryonic myocytes spread into the collagen/fibrin gels and spontaneously twitch. The cells make contact with one another and start twitching synchronously.

[00119] A sequence of images captured the mouse embryo cardiac tissues was shown in figure 7. A dark feature within the image is enclosed by a rectangular frame. The frame in each image is stationary. Displacement of the feature away from the frame periodically indicated by white arrows at image 10 and 18 indicates twitching of the sample by the cardiac myocytes (Figure 7).

[00120] In vitro tissue models made using mouse cells and mimicking the mechanical properties of mouse cardiac tissues are useful test system for evaluating the effects of pharmaceutical candidates. Well established methods of for knocking out specific genes in mice has created an enormous number of mutant mice lacking specific molecules by which cardiac functions are regulated. For instance, studies using a mouse lacking type 1 NO synthase (Circulation 2002 Jun 25;105(25):3011-6), Connexin43 (Development 2002 Apr;129(8):2031-42), and familial hypertrophic cardiomyopathy linked to myosin binding protein-C (Circ Res 2002 Mar 22;90(5):594-601) are just few examples published in 2002. Model cardiac tissue can be made using cells from a knockout mouse to study the role of a specific protein molecule in cardiac development and function. Many knockout mice do not survive after birth or even beyond a defined embryonic stage. Therefore, functional studies of the heart of these knockout mice are limited. Since cells isolated from embryonic or neonatal mice can be used to make functional tissues, the system can be used to study functions of molecules, which are inaccessible using intact tissue or whole animals. The study of knockout mice is useful for studying the efficacy of gene therapies in vitro.

[00121] MINIATURIZATION OF TISSUE MODELS

[00122] The samples are miniaturized by using a smaller mold (right in Figures 8 a, b) instead of the one used ordinarily (left in Figures 8 a, b). The sample sizes are determined by the diameters of the mandrels and inner diameters of the wells. The molds before (A) and after (B) assembling the parts are shown in the Figure 8 . The mandrel diameters of small and regular size molds are 3/16 and 3/8 inches, respectively. The inner diameters of the small and regular size wells are 9/32 and 17/32 inches, respectively. By using the smaller mold, the size of tissues are decreased 5-fold in volume (1ml to 0.2 ml) (* indicates the tissue samples in Figures 8 c, d). The miniaturization of the tissue model allows us to do tests similar to those described in Figures 2, 3, 4, 15, 16, 17, 18, 19, 20, 21, 23 using regular size samples but in much smaller organ baths using less medium consuming a fraction of the chemicals used in the tests. At least a 5-fold reduction, which is equal to the reduction of sample volume, can be achieved.

[00123] To demonstrate and prove this concept, the mechanical responses of the cardiac tissue models to the treatment of serum and Y27632 are observed (Figure 9). The tissue models are made using regular size mold with chicken cardiac cells (9a). They typically look like the tissue models (indicated by *) shown in Figure 8 d. Figure 9 b also shows the similar trace of force obtained with miniaturized tissues, which look like the ones in Figure 8 c. Both sizes of tissues increase force in response to 20% (v/v) calf serum and forces are decreased by Y27632 treatments. Differences in magnitudes of changes in response to these chemicals using different sizes of tissue models are not greater than 2 fold. This demonstrates that the miniaturized system hasve a large enough range of signal to detect changes induced by pharmaceutical candidates. The miniaturized system also shows force-response curve to mechanical stretches similar to that of the normal-sized tissues. Miniaturized samples made using chick cardiac cells show twitch forces. Therefore this system can be used to test the effects of pharmaceutical candidates treating cardiac dysfunctions (Insert in Figure 9 b)

[00124] TISSUE MODEL MEASUREMENTS IN MULTI-CELL INDENTATION SYSTEM (USING A MULTI-WELL PLATE)

[00125] Fabrication and measurement of miniaturized reconstituted tissue specimens were carried out using membranes and strips of reconstituted tissues in wells of a 96-well plate system.

[00126] MECHANICAL ASSEMBLY OF A MULTI-WELL PLATE SYSTEM

[00127] A triangular frame made of stainless steel wire 1 mm in diameter was employed as a scaffold on which the reconstituted tissue formed. The wells are slightly tapered toward the bottom and the frame is securely positioned 1 mm above the bottom of the well (Figure 11a). The non-polymerized solution of collagen containing cells and appropriate cell culture media as described above was poured into the wells filling the wells to a level 3 mm above the bottom (Figure 11 b). The 96-well plate was incubated at 37°C with 5% CO₂. During the incubation, cells compressed collagen matrices by squeezing liquid out from the porous collagen matrix. Without the wire frame, the reconstituted tissue contracted into a small sphere floating in the tissue culture medium. It was discovered that by utilizing different shapes of wire frames the collagen matrix was compressed into shapes corresponding to shapes of the frames. Illustratively, a triangular wire frame made a membrane spanning among the three edges is shown in Figure 11a. Other wire frame shapes, such as one shown in Figure 11b, produced tissue strips with different width. A porous support material such as Velcro fastener was not required to facilitate tissue adhesion even to the non-porous stainless steel surfaces of a wire. The collagen was compressed to a greater extent at the outer portion of the membrane or strip. Therefore, this outer portion of the membrane can withstand stress produced by the cells and prevented it from ripping the membrane off the wire frame.

[00128] MEASUREMENT OF FORCE, TISSUE STIFFNESS AND HYSTERESIS OF MEMBRANES OR STRIPS USING AN INDENTATION METHOD (IN A WELL PLATE SYSTEM)

[00129] To assess the response of the tissue model samples to test agents, the resistance of the tissue samples to stretch is measured by a probe coupled to a force transducer. The stiffness of the tissue is related to the force required to move

the probe by a specified amount once the probe contacts the tissue. The probe, consisting of a vertical glass tube whose tip has been smoothed by fire polishing, is attached to an isometric force transducer (described above). The diameter of the probe is about 2 to 3 mm and the shape of its tip can be flat or hemispherical. The probe is firmly attached to a beam and the beam attached to the force transducer by glue or wax (Figure 11a). The force transducer is attached to a stationary frame. The 96-well plates are placed on a stage that moves vertically. The stage constructed for demonstration of the method uses a micrometer driven by a computer-controlled stepper motor to raise or lower the stage at prescribed incremental rates. In this implementation the range of stage motion is 0 mm to 15 mm at a maximum velocity of 500 $\mu\text{m}/\text{sec}$. The same apparatus has been used for large-scale tissue mechanical measurements. A detailed description is provided below in the following section on force measurements.

[00130] SAMPLE PREPARATION OF TISSUE SYSTEM FOR USING THE INDENTATION METHOD (IN A WELL PLATE SYSTEM)

[00131] During the tissue formation and continued culturing of tissue models, the samples were kept in a 5% CO_2 incubator at 37 °C with bicarbonate buffered DMEM supplemented with 10% FBS, penicillin, and streptomycin. For force measurements the medium was changed to 150 μl of HEPES-buffered DMEM without serum. The temperature of the medium was kept at 37 °C by placing the wells on a heated plate connected to a temperature-controlled circulation bath.

[00132] MEASUREMENTS USING TISSUE INDENTATION METHOD (IN A WELL PLATE SYSTEM)

[00133] The stage is raised until the probe tip touches the membrane or strip of reconstituted tissue. Contact of the tip with the sample is detected by a sudden increase in the force registered by the isometric force transducer. The stage is then lowered by 5 μm , i.e. tip is withdrawn from contact with the sample. Then the stage is moved vertically according to a saw-tooth wave-form at 3.3 $\mu\text{m}/\text{sec}$ with 100 μm amplitude. During this trajectory the tip comes in contact with the tissue

specimen and stretches it as the force is continually recorded by a data recorder. The velocity can be varied to optimize the sensitivity of measurements and to measure the viscosity of the sample.

[00134] After the peak force reaches its steady level the sample is stimulated with 20% Fetal Bovine Serum (FBS) (arrow a in Figure 12a). This amount of FBS activates fibroblast non-muscle myosin producing a contractile force that stiffens the reconstituted tissues. About 10 min. after the FBS addition, there is a ~25% increase in the peak force of subsequent indentations (Figure 12a). Nearly 15 min. after the addition to the medium of 40 nM CD (arrow b), the peak force from subsequent indentations has decreased ~40% from its initial level (Figure 12). A further reduction of peak force was recorded 20 min. after addition of 2 μ M CD.

[00135] A plot of force versus indentation depth during the membrane stretching follows a different path from that seen as the stress is relaxed by retraction of the probe from the sample, establishing a hysteresis area between the two curves. Upon additions of FBS or CD, changes in the hysteresis areas are compared in Figure 12(b). Percent changes in the peak force and area of hysteresis during the experiment shown above are compared in Figure 13. The area of hysteresis changes to a greater extent upon stiffening due to myosin activation and CD addition. Therefore, the area of hysteresis is a more sensitive parameter than the peak force for monitoring the changes in mechanical properties of the sample.

[00136] TEST USING PHYSIOLOGICAL ACTIVATORS TO ACTIVATE FORCE RESPONSE WITHOUT STRETCHING

[00137] Ring samples made using a rat embryo fibroblast derived cell line (REF52) were prepared as described previously herein. The length of the tissue was maintained at a constant level during the test. Each of the drugs indicated in Figures 10-15 was added to a single tissue ring. Thrombin, vasopressin, lysophosphatidic acid (LPA), bradykinin, and endothelin contacted the ring to produce different forces that develop in different response times (Figures 15). Norepinephrine and phenylephrine relaxed the contractility of the ring, and therefore norepinephrine and phenylephrine reduced the initial base line force of tissue indicated at the

beginning of each test (Figure 15) (reagents used are obtain from Sigma, St. Louis, MO).

[00138] Profiles of contractile response to different agonists applied to ring samples were made using chicken embryo cardiac fibroblasts (CECFs) and rat adult cardiac fibroblasts (RACFs) and are shown in Figure 16(a), 16(b), 16(c), 16(d) and Figure 16(e), 16(f), 16(g), 16(h), respectively.

[00139] RACFs responded to vasopressin (Figure 1 f) and angiotensin II (Figure 16g) whereas CECFs did not response to vasopressin (Figure 16 a) and angiotensin II (Figure 16c). CECFs did not respond to any dose of EGF stimulation (Figure 16d). Endothelin stimulation contracted rings made using both CECFs (Figure 16b) and RACFs (Figure 16h). Time dependent force profiles after addition of FBS (20% v/v) were different depending on the pretreatment by different agonists.

[00140] For example, FBS addition resulted in a double peak of force after EGF pretreatment (Figure 16d). The force level is sustained after FBS addition to a system previously treated with angiotensin II. Yet the force level started to decrease from its FBS stimulated peak level after treatment with vasopressin and endothelin (Figure 16a, b).

[00141] This invention is useful for managing the profiles not only of single compounds but also of combinations of multiple compounds applied at the same time or at different times. Managing profiles of combinations of multiple pharmaceuticals is useful both for screening purposes and also to reveal unexpected consequences caused by combinations of multiple pharmaceuticals.

[00142] The myosin ATPase inhibitor BDM reduces in a dose dependent manner (Figure 17) the fully activated contractile force produced by prior addition of 20% FBS.

[00143] The contractile force developed by the ring sample during matrix remodeling and compression can be measured by connecting the ring to the force measuring apparatus as described above within about 1-2 hours after collagen

gelation. The cells begin to exert a force on the matrix after this time. Then the increase of force during tissue development (remodeling and compression) is observed over time.

[00144] A profile of force measured during the ring tissue development is shown in Figure 18(a). Maximum force is typically reached within 15 to 20 hours. This process was also disrupted by several inhibitors as illustrated in Figure 18(b), (c), (d). The tTyrosine kinase inhibitor, herbimycin A (Figure 18 b) reduces the maximum force as does the microtubule disrupting reagent nocodazole (Figure 18 c). Cytochalasin D (2 μ M) completely abolished the development of force from the beginning of the test.

[00145] This invention has broad utility including using this invention for high-throughput pharmaceutical drug screening and treatment testing. For example, reconstituted tissues in the form of membranes or strips could be mass-produced to supply 96 uniform samples, one in each well of a 96-well plate. Simultaneous force measurements could also be achieved by using 4 or more force transducers (Figure 11). The 96-well plate is placed on an x-y stage that positions the samples at the correct locations for the indentation measurements. Probes attached to force transducers move vertically to indent the samples (Figure 11a). A single force transducer can be used to indent several samples within a short period of time by repositioning the 96-well plate using the x-y stage. The addition of chemical compounds and small peptides to the sample and movement of the stage and force transducers can all be automated and controlled by a personal computer. The computer also stores and analyzes the data in a database.

[00146] Genes and proteins can be delivered to the cells as agents within the reconstituted tissues using various techniques available herein.

[00147] The number of wells per plate may vary in accordance with the size of the system being utilized in an indentation (indention) setup. Typically the number of wells per plate is from about 2 to about 1000 or more preferably from about 50 to about 500 wells per plate.

[00148] Applications of this method in addition to drug screening include testing procedures for delivering genes to cells as agents. The reconstituted tissue samples can be made using cells lacking genes that contribute to cell and tissue mechanical properties. Hence, the recovery of normal tissue mechanical characteristics provides an assay for the effective delivery of the gene.

[00149] The ring tissue model system can detect the effect of a genetic deletion on the ability of cells to generate force both while compressing and remodeling the collagen matrix and while responding to calf serum. Cells interact with and adhere to diverse ECM constituents through specific heterodimeric receptors called integrins on their surface membranes. Each integrin is composed of one α and one β subunit. The integrins that mediate binding to collagen are $\alpha_1\beta_1$ and $\alpha_1\beta_2$. NMuMG, is an immortalized but nonmalignant mouse mammary epithelial cell line that does not express α_1 and α_2 integrin. Hence, these cells interact weakly if at all with collagen matrices and therefore cannot transmit force to the ECM to generate and to maintain tissue stiffness. Figure 20 demonstrates that cells without α_1 and α_2 integrin could not develop baseline force during tissue development nor could it respond to calf serum (CS) by increasing force (Figure 20 a, b). Also shown is the rescue of these two functions by re-expressing normal α_2 . The cells containing the α_2 gene normally adhered to and compressed the matrix and responded to CS (Figure 20 a, b). This example illustrates the useful application of the invention to generate and to manage the mechanical response profiles that report on gene expression. Hence, the invention is useful to manage the mechanical response profiles as indicators of genotype and of the results of gene therapy, such as the efficiency of different methods of gene delivery.

[00150] Reconstituted tissue models can be used to assess quantitatively and rapidly the effects of many different classes of potential pharmaceuticals, toxins, and pathogens as agent(s) on the mechanical properties of cells and matrix. These mechanical properties provide general indicators of the overall organization of cellular mechanical systems, especially the cytoskeleton, of the operation of signal transduction pathways, and of the organization imposed on the

matrix by the cells during tissue development. Therefore they provide potential applications in a wide range of disease areas.

[00151] Example of potential pharmaceutical candidate testing

[00152] We demonstrated dose dependent stress relaxation of the tension in reconstituted tissues using Y27632, which has promise for the treatment of hypertension.

[00153] The connective tissue models made using NIH 3T3 cells were treated with different amounts of the candidate pharmaceutical, Y27632. Rho kinase specific inhibitor, Y27632 has been tested for reducing tension of smooth muscle strips such as rabbit aortic rings (Nature 239 (1997) 990-993). It is one of the promising drug candidates treating hypertension in the future. Depending on the agonists used to stimulate the tissue contraction, the degree of reduction in tension is different (Figure 21). Y27632 reduces the tension of connective tissue models in a similar dose-dependent fashion (Figure 21). This demonstrates the utility of technology in screening the pharmaceutical candidates.

[00154] The collagens in tissues are degraded and recycled by the family of enzymes called matrix metalloproteases (MMPs), which are secreted as latent proenzymes. The enzymes become active through proteolytic cleavage of their amino-terminal domain, and their activities depend on the presence of Zn^{++} and Ca^{++} . MMP-2 activity is known to play a role in tumor cell invasion. The presence of MMPs both in pro- and active-forms is detected by a technique commonly known as zymography. Chicken cardiac fibroblasts are cultured for 2 days with DMEMs supplemented with none or 0.5% fetal bovine serum. The MMP-2 and 9 are secreted by the cells into the medium and their presence is detected by the zymograph. In Figure 22 the control lanes 11 through 15 are loaded with purified proenzymes of MMP-2 and 9. This shows that at least 0.5 ng of enzyme (lane 11) can be detected using the assay (Figure 22). The smaller active enzyme runs ahead of pro-enzymes and a band of active enzyme appears at lower positioned of than the band of inactive MMP-2. Medium conditioned by the cells growing on tissue culture dishes with no coating, or with fibronectin, or collagen coating do not show any bands of active

enzymes. The medium conditioned by the cells grown in 3 D collagen matrix, i.e., tissue model, shows a band of active MMP-2 (lanes 9 and 10). This indicates that the cells on 2D substrata secrete inactive enzymes into the medium but they are not activated. The enzymes never get activated unless the cells are grown in 3 dimensional matrices. Therefore, the investigation of MMP activities affecting extracellular matrix degradation such as during tumor invasion requires a model system such as the tissue models in which the cells are growing in 3 dimensional matrices. This is especially important for discovering inhibitors of MMPs.

[00155] The effects of the general inhibitor of MMPs, GM6001 (Biomol Research Laboratories Inc. Plymouth Meeting, PA) (N-[(2R)-2-(Hydroxamidocarbonylmethyl)-4-methylpentanoyl]-L-tryptophan methylamide) on the mechanical properties of artificial tissues have been investigated. The tissue models made using cardiac fibroblasts are incubated with 50 μ M GM6001 for 6 days. The tissue culture medium is replenished every other day with fresh medium containing GM6001. The tissue model rings are subjected to mechanical tests as described above. The GM6001 treated samples (dash lines in Figure 23) show significant reduction in mechanical properties of the samples compared to those of controls (solid lines). Although the K_i values of GM6001 for different MMPs are around nmol/l, the effects of GM6001 in animals can be observed only with a high concentration of the inhibitor (sub mmol/l) (Circ Res 1996 Jan;78(1):38-43). This suggests that the K_i value measured in a purified system does not directly correlate to the inhibitory effect of drug in living test subjects such as animals or tissue models. MMP inhibitors affecting the mechanical properties of various tissue models can be discovered efficiently using the high throughput system.

[00156] This invention has many additional utilities. For guiding treatment cardiac diseases, reconstituted tissue models assembled from cardiac myocytes and/or cardiac fibroblasts can be used to test the effects of candidate pharmaceuticals on cell and matrix remodeling processes that result from pressure overload and on tissue restructuring that occurs in response to trauma or infarction. This invention can be employed to guide treatment of dental connective tissue diseases-, cancer metastasis- (contraction, traction force in cell motility), diabetes

(stiffening of connective tissues and skin by collagen cross linking), pulmonary diseases such as emphysema, chronic inflammation (elastase secreted from neutrophils), muscular dystrophy and aging skin.

[00157] This invention also comprises a method of managing a library of pharmaceuticals. This method comprises obtaining a profile of mechanical response to the contact of an activator with a tissue model system wherein a tissue model has been contacted with the pharmaceutical, storing that profile in a database, storing at least one additional profile of another pharmaceutical in a tissue model system in that database, setting up a means for comparing with another profile, one profile, comparing the profile of the first pharmaceutical with the profile of a second pharmaceutical based on a pre-established or ordered standard of comparison. The pharmaceuticals are ranked in an order of activity with respect to mechanical effect on the tissue model system based on their respective profile.

[00158] Possible Parameters To Analyze Curves

[00159] The force response initiated by an agonist addition plotted over time can be presented using several parameters to describe the shape of the curve. The F_0 and $\Delta F_{\max}$ represent initial the force level before the agonist addition and the maximum change in force respectively. The $\Delta T_{\max}$ represent the time to complete the change in force level. The time derivative of the force curve represents the velocity of the change of force with time. The maximum value of dF/dt , $V_{\max}$, represents a maximum slope in the force curve. These parameters are useful to make a profile of pharmaceutical candidates in terms of the changes the tension level of the artificial tissues (Figure 24)

[00160] In an embodiment, the profile of a candidate pharmaceutical is compared to the profile of a known pharmaceutical and a ranking or rating is made based on that comparison to provide information and guidance as to whether the candidate pharmaceutical would likely be an effective pharmaceutical.

[00161] Further, if desired, a test is carried out in accordance with this invention, wherein the candidate pharmaceutical is evaluated against a particular

cell system which is known to be involved in a particular disease. In this embodiment, the cellular response thereto using the system of this invention is employed to provide an indication of the probable activity of the candidate pharmaceutical to a particular cell type. In a further embodiment, the pharmaceutical is evaluated in a tissue model system and compared to the evaluation of a pharmaceutical known to be useful in treating a particular disease.

[00162] In a further embodiment, the profile of the known pharmaceutical is one which is known to provide effective treatment against a known disease. In this embodiment a comparison is thus made wherein the pharmaceutical is evaluated against cells which are known to be involved in a particular disease. This embodiment is useful in determining whether a pharmaceutical is potentially useful in the treatment of heart disease or hypertension or aging for example.

[00163] Hypertension is caused by elevated contractility and stiffness of blood vessels. Drugs for hypertension can be identified using this invention. These drugs can be used to lower the blood pressure of the animal or tension and stiffness of isolated blood vessels from the animal. This invention uses artificial tissues mimicking blood vessels using tissue cultured cells and extracellular matrices, the method herein can replace animal models or explanted tissues.

[00164] Artificial tissues are made to mimic specific biological functions or different types of organs and tissues including skin, muscle, heart, and blood vessels or to mimic more complex tissues by co-culturing different cell types in a single artificial tissue. Since the mechanical properties of the tissues are correlated to their structural integrity by using this invention, they are important parameters to indicate biological functions of tissues and organs.

[00165] In addition, this invention can be used to screen compounds at high speed based on their biological activities affecting the mechanical properties of the artificial tissues. For example, artificial tissues are made in small sizes to fit in one of the wells (4 mm in diameter and 6 mm in height) in a 96-well plate. The significantly miniaturized sample preparation reduces the amount of compounds used in each testing at least by 90% compared to known methods, such as using aortic

rings. Since animal tissues are surgically isolated from animals, their sizes and responses to the drugs are not necessarily reproducible.

[00166] Toxicity of pharmaceutical compounds can be determined using the method of this invention. For example a different dose of ethanol causes a decrease in the level of baseline force maintained by the viable cells.

[00167] In another aspect, the biochemical properties of cells and matrices comprising model tissue are measured optically using, for example, fluorescence markers.

[00168] Using the invention, one can create a new library of profiles of pharmaceuticals based on their effect on the mechanical properties of live artificial tissues. The number of active compounds in the library created by the instant invention using the indentation system is most likely to be less than that in the library created by a prior art test tube based screening system. Compounds selected in chemical screening procedures may include many that elicit no physiological response.

[00169] The chemical compounds screened by the artificial tissues based system will have a higher chance of having similar effects on the mechanical properties of real tissues and organs in animals and humans. Therefore, the use of animal models for an optimization stage of drug screening can be significantly reduced using the artificial tissue based high throughput screening. The invention can replace animal testing by the invention providing a physiological response system assembled from cultured cells.

[00170] In an aspect, the results of screening are employed to identify and advance one or more candidate pharmaceuticals or drugs to an advanced stage of further stage of testing or evaluation, including possibly commercialization. In another aspect, the results of screening are employed to terminate or alter further testing or screening on a pharmaceutical or drug candidate. In another aspect, the method and apparatus herein are used to evaluate and validate the target or locus for or of the drug or pharmaceutical candidate.

[00171] While the invention has been described in terms of various specific embodiments, those skilled in the art will recognize that the invention can be practiced with modification within the spirit and scope of the claims.

WHAT IS CLAIMED IS:

1. A method for screening pharmaceutical chemicals and for obtaining a response of a tissue model system to an agent comprises contacting a tissue model system with the agent and determining a parameter of at least one of contractile force and tissue stiffness.
2. A method in accordance with Claim 1 wherein the parameter comprises contractile force and tissue stiffness.
3. A method in accordance with Claim 1 wherein the tissue stiffness includes at least one of hysteresis, phase lag and dynamic stiffness.
4. A method in accordance with Claim 1 wherein the tissue model system is a ring system.
5. A method in accordance with Claim 1 wherein the activator is a pharmaceutical candidate or a known pharmaceutical.
6. A method in accordance with Claim 1 wherein the tissue model system comprises cells in collagen.
7. A method in accordance with Claim 6 wherein the cells are muscle cells.
8. A method in accordance with Claim 7 wherein the cells are non-muscle cells.
9. A method in accordance with Claim 7 wherein the tissue model system is a bio-artificial system.
10. A method in accordance with Claim 9 wherein the system is a living system.
11. A method in accordance with Claim 8 wherein cells are endothelial cells.

12. A method in accordance with Claim 7 wherein the cell is selected from at least one of cardiac cells.

13. A method in accordance with Claim 7 wherein the tissue model system comprises cells of extracellular matrix.

14. A method for obtaining a mechanical response of a tissue model system to contact with an agent, said method comprising constructing a tissue model system, providing an agent in contact therewith, and measuring cellular mechanical properties in response to contact of the agent with the system of at least one parameter of contractile force and hysteresis.

15. A method in accordance with Claim 14 wherein the cellular mechanical response thereof includes contractile force and hysteresis.

16. A method in accordance with Claim 15 wherein the mechanical response is measured using a tissue indentation system.

17. A method in accordance with Claim 16 wherein the tissue indentation system comprises an indentation probe coupled to a force transducer and a computer.

18. A method in accordance with Claim 17 wherein the tissue model system is a bio-artificial system.

19. A method in accordance with Claim 18 wherein the bio-artificial system is a living system.

20. A method in accordance with Claim 19 wherein the tissue model system comprises cells in collagen.

21. A method in accordance with Claim 20 wherein the cells comprise muscle and non-muscle cells.

22. A system for obtaining a mechanical response profile based on reconstituted model muscle and non-muscle tissue mechanical measurements of the

response of a tissue model to contact with an agent, which comprises constructing a reconstituted tissue model comprising reconstituted cells in collagen, contacting said reconstituted cell matrix with an agent and measuring cellular mechanical response to contact with said agent in terms of at least one of contractile force and tissue stiffness.

23. A system in accordance with Claim 22 wherein the cellular response comprises at least one of force development and tissue stiffness.

24. A system in accordance with Claim 23 wherein the tissue stiffness includes at least one of hysteresis, phase lag and dynamic stiffness.

25. A system in accordance with Claim 24 wherein the cellular response comprises force development, tissue stiffness and hysteresis.

26. A system in accordance with Claim 23 wherein the cellular response is determined using a tissue indentation method.

27. A system in accordance with Claim 23 wherein the mechanical response is measured using a tissue indentation system.

28. A system in accordance with Claim 27 wherein the tissue indentation system comprises an indentation probe, a force transducer and a computer.

29. A system in accordance with Claim 22 wherein the system is a bio-artificial system.

30. A system in accordance with Claim 23 wherein the system is a living system.

31. A method for screening pharmaceuticals comprising contacting a tissue model comprising cells in collagen with an effective amount of a candidate pharmaceutical and measuring cellular response to the candidate pharmaceutical in terms of at least one of contractile force and tissue stiffness.

32. A method in accordance with Claim 31 further comprising measuring cellular response to the pharmaceutical in terms of contractile force and tissue stiffness.

33. A method in accordance with Claim 32 wherein the tissue stiffness includes at least one of hysteresis, phase lag and dynamic stiffness.

34. A method in accordance with Claim 33 further comprising measuring cellular response to contact with said pharmaceutical by determining phase lag and dynamic stiffness.

35. A tissue model for measuring response data of a matrix comprising reconstituted cells in collagen to an activator exposure comprising:

- (a) cells
- (b) collagen; and
- (c) ECM.

36. A tissue model in accordance with Claim 35 wherein the cells comprise muscle and non-muscle cells.

37. A method of managing a library of one or more pharmaceuticals comprising obtaining a profile of a mechanical response to contact of an agent pharmaceutical with a tissue model system wherein a tissue model has been contacted with the pharmaceutical, storing that profile in a database, storing at least one additional profile of another pharmaceutical in a tissue model system in the database and comparing one profile with another profile.

38. A method in accordance with Claim 37 further comprising comparing the profile of the first pharmaceutical with the profile of a second pharmaceutical based on a pre-established or ordered standard of comparison.

39. A method in accordance with Claim 38 further comprising ranking the pharmaceuticals in an order of pharmaceutical activity based upon the

corresponding profiles of each pharmaceutical with respect to a mechanical effect on a tissue model system.

40. A method in accordance with Claim 37 wherein the tissue model system is a ring system or a tissue indentation system.

41. A method in accordance with Claim 39 further comprising advancing a pharmaceutical to a secondary level of testing based upon the ranking.

42. A method in accordance with Claim 37 wherein said pharmaceutical is a group or a class of chemicals.

43. A method in accordance with Claim 37, wherein said comparison of profile is based on contracted force or tissue stiffness.

44. A method in accordance with Claim 37 wherein the cellular matrix system uses a tissue indentation system.

45. A method in accordance with Claim 37 wherein comparing comprises establishing baseline mechanical response data comprising contractile force and stiffness.

46. A method in accordance with that of Claim 37 wherein the system is a bio-artificial system.

47. A method in accordance with Claim 37 wherein the system is a living system.

48. A method in accordance with Claim 37 wherein the cells are muscle cells.

49. A method in accordance with Claim 37 wherein the cells are non-muscle cells.

50. A method of obtaining a mechanical response profile for a tissue model comprising living cells contacted with a pharmaceutical, said method

comprising contacting a tissue model with the pharmaceutical and measuring the cellular response thereof.

51. A method in accordance with Claim 50 wherein the tissue model system is a ring system.

52. A method in accordance with Claim 50 wherein the tissue model system uses a tissue indentation system.

53. A method in accordance with Claim 50 wherein the profile is obtained from a ring system.

54. A method in accordance with Claim 50 wherein the profile is obtained using an indentation system.

55. A method in accordance with Claim 50 wherein the profile of a candidate pharmaceutical is compared to the profile of a known pharmaceutical and a ranking is created based on that comparison.

56. A method in accordance with Claim 50 wherein the profile is obtained using a tissue indentation system.

57. A method in accordance with Claim 56, wherein the candidate pharmaceutical is evaluated against a cell system known to be involved in a disease, said method comprising obtaining a profile of a candidate pharmaceutical to provide an indication of the probable activity of the candidate pharmaceutical to a particular cell and comparing the profile of the candidate pharmaceutical to a profile of a known active pharmaceutical.

58. A method in accordance with Claim 57 wherein a comparison is made of the profile of the known pharmaceutical known to provide effective treatment against a known disease to the profile of a candidate pharmaceutical whose efficiency to a disease is desired to be known.

59. A method in accordance with Claim 50, wherein a comparison is made of the profile of a pharmaceutical known to be effective against cells which are known to be involved in a particular disease.

60. A tissue model system, comprising tissue assembled in the form of a ring mounted on a system comprising an isometric force transducer electrically coupled to a computer –controlled stepping motor wherein the tissue spans the isometric force transducer and the computer-controlled stepping motor provides an application of force and strain to the tissue.

61. A system in accordance with Claim 60 wherein the application of force is linear.

62. A system in accordance with Claim 60 wherein the application of force is a ramp force.

63. A system in accordance with Claim 60 wherein the force is applied sinusoidally.

64. A system in accordance with Claim 60 wherein the force is applied to achieve a 20% stretch of tissue.

65. A method in accordance with Claim 60 wherein tissue stiffness is measured by subjecting the tissue in the tissue model system to a stretch and observing the change in force as the strain increases.

66. A system in accordance with Claim 60 wherein the tissue is muscle tissue.

67. A tissue model system, comprising a membrane of reconstituted tissue provided with a support and having a bar attached to an isometric force transducer coupled to a probe.

68. A system in accordance with Claim 67 further comprising a computer, said computer directing the probe to indent the membrane and measure the force resisting the indentation.

69. A system in accordance with Claim 67 wherein a contractile force is measured as peak force along with tissue stiffness which is registered as a resistance to indention of the tissue model by the probe contacting the tissue.

70. A system in accordance with Claim 67 wherein tissue stiffness is measured by subjecting the tissue in the system to a steady stretch and observing the change in force as the strain increases.

71. A method of preparing a tissue model system, said method comprises growing tissue in the form of a ring around a mandrel, removing the ring shaped tissue from the mandrel and thereafter subjecting said ring shaped tissue to an application of stretch force by application of a standard force.

72. A method in accordance with Claim 71, wherein the force is applied in a constant manner.

73. A method in accordance with Claim 71 wherein the tissue is a living tissue.

74. A method in accordance with Claim 71 wherein the tissue is muscle cells.

75. A method in accordance with Claim 71 wherein the cells are cardiac cells.

76. A high throughput screening system comprising at least one unit comprising a tissue model system comprising cells and ECM arranged in a membrane adapted to receive a probe, a probe adapted for receiving instructions as to advancing or retracting to/from said membrane, a transducer coupled to said probe having electrical transmission capability to an electronic recording system wherein advancement instructions are provided to said probe.

77. A system in accordance with Claim 76 wherein a multi-well plate system is engaged to house the membranes.

78. A system in accordance with Claim 76 wherein the number of units is from about 2 to about 10,000 units.

79. A system in accordance with Claim 76 wherein the number of units is from about 2 to about 96.

80. A system in accordance with Claim 76 wherein the cells are muscle cells.

81. A system in accordance with Claim 76 wherein the cells are non-muscle cells.

82. A system in accordance with Claim 76 wherein the muscle cells are cardiac cells.

83. A diagnostic kit comprising a tissue model system and cell activator, said activator contacts the tissue model system and a response is measured therefrom.

84. A kit in accordance with Claim 83 wherein said tissue model system is a living system and said kit is portable.

85. A kit in accordance with Claim 83 wherein the response determined is contractile force or tissue stiffness.

86. A kit in accordance with Claim 83 wherein the response determined is hysteresis.

87. A tissue indentation system comprising a tissue, one end of said tissue is connected to a fixed position and another other end of said tissue is connected to a transmitter probe which is connected to a computer, wherein the probe is advanced and contacted with a tissue sample to indent the tissue.

88. A system in accordance with Claim 87 wherein the tissue is living tissue.

89. A system in accordance with Claim 87 wherein the tissue is muscle tissue.

90. A system in accordance with Claim 87 wherein the tissue is muscle cells.

91. A method for establishing a mechanical response profile of a pharmaceutical compound comprising contacting the pharmaceutical with a tissue model and determining the mechanical response(s) of the tissue model comprising a bio-artificial system of living cells utilizing at least one of contractile force and stiffness.

92. A method in accordance with Claim 87 further comprising associating said profile with a cell present in the tissue model.

93. A method in accordance with Claim 87 wherein a profile is obtained using a ring system.

94. A method in accordance with Claim 87 wherein a profile is obtained using a tissue indentation system.

95. A method in accordance with Claim 87 wherein a profile is determined which reflects the contribution of cells and cytoskeleton to the overall profile.

96. A method of culturing a cardiac tissue model, said method comprising using a medium conditioned by fibroblasts.

97. A method in accordance with Claim 96 wherein fibroblasts are cardiac fibroblasts.

98. A method in accordance with Claim 97 wherein said medium is DMEM with 1% FBS.

99. A method in accordance with Claim 98 wherein a condition of said method is selected from coculture with feeding fibroblast monolayer, coculture with artificial tissue with fibroblasts and mixing fibroblasts in the tissue.

100. A method in accordance with Claim 12 wherein said cardiac cells are selected from at least one of chicken and mouse.

101. A method in accordance with Claim 100 wherein said cardiac cell is a chicken cardiac cell.

102. A method in accordance with Claim 100 wherein said cardiac cell is a mouse cardiac cell.

103. A method and apparatus to identify the effect of a drug upon a tissue which comprises treating a tissue using this invention and measuring the effect so resulting and thereafter comparing such measured effect with a standard drug effect on the same or similar tissue.

104. A method and apparatus to data mine a library of pharmaceutical moieties for activity in tissue, which comprises treating a tissue using this invention and measuring the effect so resulting. Optionally the measured effect may be compared with a correspondingly measured effect of a standard drug on the same or similar tissue.

105. A method of optimizing tissue culture conditions for constructing implantable artificial cardiac tissue, with additional different factors influencing cardiac tissue development including growth factors, and matrix proteins and hormones to a tissue model of the invention.

REFERENCES:

1. Petersen, N. O., W. B. McConnaughey, and E. L. Elson. 1982. Dependence of locally measured cellular deformability on position on the cell, temperature, and cytochalasin B. *Proceedings of the National Academy of Sciences of the United States of America*. 79:5327-31.

2. Sundberg, S. A. 2000. High-throughput and ultra-high-throughput screening: solution- and cell-based approaches. *Curr Opin Biotechnol*. 11:47-53.

3. Zahalak, G. I., W. B. McConnaughey, and E. L. Elson. 1990. Determination of cellular mechanical properties by cell poking, with an application to leukocytes. *Journal of Biomechanical Engineering*. 112:283-94.




Figure 1




Figure 2




Figure 3





Figure 4



Conditions	Contractility
DMEM supplemented with 10% FBS	No contraction
Cardiac Tissue Medium	Contraction after 7 to 8 days
Fibroblasts Conditioning Medium	Contraction after 4 to 5 days
Coculture with Feeding Fibroblast Layer	Strong Contraction after 4 to 5 days
Coculture with Artificial Tissue with Fibroblasts	Strong Contraction after 4 to 5 days
Mixing Fibroblasts in the Tissue	Contraction after 4 to 5 days but reduction in contraction in 7 to 8 days

Figure 5




Figure 6




Figure 7




Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16




Figure 17




Figure 18




Figure 19




Figure 20




Figure 21




Figure 22




Figure 23




Figure 24