

FORM 1

617023

SPRUSON & FERGUSON

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

APPLICATION FOR A STANDARD PATENT

Ciba-Geigy AG, incorporated in Switzerland, of Klybeckstrasse 141, 4002 Basle, SWITZERLAND, hereby apply for the grant of a standard patent for an invention entitled:

Process for the Preparation of Pyrimidine Derivatives
which is described in the accompanying complete specification.

Details of basic application(s):-

<u>Basic Applic. No:</u>	<u>Country:</u>	<u>Application Date:</u>
288,751	US	22 December 1988

The address for service is:-

Spruson & Ferguson
Patent Attorneys
Level 33 St Martins Tower
31 Market Street
Sydney New South Wales Australia

DATED this FOURTEENTH day of DECEMBER 1989

Ciba-Geigy AG

By:



Registered Patent Attorney

TO: THE COMMISSIONER OF PATENTS
OUR REF: 110674
S&F CODE: 52760

5845/4

COMMONWEALTH OF AUSTRALIA

Patents Act 1952 - 1969

DECLARATION IN SUPPORT OF A CONVENTION APPLICATION FOR A PATENT

In support of the Convention Application made by CIBA-GEIGY AG for a patent for an invention entitled:

Process for the preparation of pyrimidine derivatives

I, Werner Waldegg of CIBA-GEIGY AG, Klybeckstrasse 141, 4002 Basle, Switzerland do solemnly and sincerely declare as follows:

1. I am authorised by the applicant for the patent to make this declaration on its behalf.
2. The basic application(=) as defined by Section 141 on the Act was (were) made in USA
on December 22, 1988

by Gottfried Seifert and Robert Hässig


3. Gottfried Seifert, Mühlemattweg 20, 4312 Magden, Switzerland
and
Robert Hässig, Gänsackerring 3, 5264 Gipf-Oberfrick, Switzerland

is (are) the actual inventor(s) of the invention and the facts upon which the applicant is entitled to make the application are as follows: The said applicant is the assignee of the actual inventor(s).

4. The basic application(=) referred to in paragraph 2 of this Declaration was (were) the first application(=) made in a Convention country in respect of the invention the subject of the application.

DECLARED at Basle, Switzerland on
October 12, 1989

CIBA-GEIGY AG


Werner Waldegg
Single Signature, by special power


To: The Commissioner of Patents

(12) PATENT ABRIDGMENT (11) Document No. AU-B-47082/89
(19) AUSTRALIAN PATENT OFFICE (10) Acceptance No. 617023

- (54) Title
PROCESS FOR THE PREPARATION OF PYRIMIDINE DERIVATIVES
- (51)⁴ International Patent Classification(s)
C07D 239/60
- (21) Application No. : 47082/89 (22) Application Date : 21.12.89
- (30) Priority Data
- (31) Number (32) Date (33) Country
288751 22.12.88 US UNITED STATES OF AMERICA
- (43) Publication Date : 28.06.90
- (44) Publication Date of Accepted Application : 14.11.91
- (71) Applicant(s)
CIBA-GEIGY AG
- (72) Inventor(s)
GOTTFRIED SEIFERT; ROBERT HASSIG
- (74) Attorney or Agent
SPRUSON & FERGUSON , GPO Box 3898, SYDNEY NSW 2001
- (56) Prior Art Documents
AU 65703/90 C07D 239/60
- (57) Compounds used in preparation of sulfonyl urea herbicides.

CLAIM

1. A process for the preparation of a 4,6-bis(difluoromethoxy)pyrimidine of formula I



wherein R is C₁-C₄alkyl or unsubstituted or substituted phenyl or benzyl, which process comprises reacting a 4,6-dihydroxypyrimidine dialkali metal salt of formula II



wherein R is as defined for formula I and Me is an alkali metal, with chlorodifluoromethane in a solvent selected from the group consisting of ketones and alkyl cyanides, in the presence of 0.05 to 1.1 mol of water per mol of dialkali metal salt of formula II.

S & F Ref: 110674

FORM 10

COMMONWEALTH OF AUSTRALIA

PATENTS ACT 1952

COMPLETE SPECIFICATION

6 1 7 0 2 3

(ORIGINAL)

FOR OFFICE USE:

Class Int Class

Complete Specification Lodged:
Accepted:
Published:

Priority:

Related Art:

Name and Address
of Applicant:

Ciba-Geigy AG
Klybeckstrasse 141
4002 Basle
SWITZERLAND

Address for Service:

Spruson & Ferguson, Patent Attorneys
Level 33 St Martins Tower, 31 Market Street
Sydney, New South Wales, 2000, Australia

Complete Specification for the invention entitled:

Process for the Preparation of Pyrimidine Derivatives


The following statement is a full description of this invention, including the best method of performing it known to me/us

5-17348/+

Process for the preparation of pyrimidine derivatives

Abstract of the Disclosure

4,6-Bis(difluoromethoxy)pyrimidines of formula I



wherein R is C₁-C₄alkyl or unsubstituted or substituted phenyl or benzyl, are prepared by reacting a 4,6-dihoxypyrimidine dialkali metal salt of formula II




wherein R is as defined for formula I and Me is an alkali metal, with chlorodifluoromethane in a solvent selected from the group consisting of ketones and alkyl cyanides, in the presence of 0.05 to 1.1 mol of water per mol of dialkali metal salt of formula II.

5-17348/+

Process for the preparation of pyrimidine derivatives

The present invention relates to a process for the preparation of 4,6-bis(difluoromethoxy)pyrimidines of formula I



wherein R is C₁-C₄alkyl or unsubstituted or substituted phenyl or benzyl.

By alkyl is meant straight chain or branched alkyl, for example methyl, ethyl, n-propyl, isopropyl, n-butyl or the butyl isomers.

Suitable substituents of phenyl or benzyl are, for example, alkyl, halogen, nitro or alkoxy.

The 4,6-bis(difluoromethoxy)pyrimidines of formula I are valuable intermediates. They can be converted, for example by oxidation, into the corresponding sulfones which, when further reacted with ammonia or a primary amine, yield the corresponding 2-amino-4,6-bis(difluoromethoxy)pyrimidines which, in turn, when further reacted with a suitable phenylsulfonyl isocyanate or N-(phenylsulfonyl)carbamate, yield herbicidally active sulfonylureas. Such herbicidally active sulfonylureas are disclosed, for example, in published European patent applications A-0 072 347, A-0 084 020 and A-0 094 790.

It is known to prepare 4,6-bis(difluoromethoxy)pyrimidine by reacting 4,6-dihydroxypyrimidine with chlorodifluoromethane in dioxane in the presence of aqueous sodium hydroxide. This process affords 4,6-bis(difluoromethoxy)-2-methylthiopyrimidine in a yield of only 25 % of theory

(q.v. US patent specification 4 542 216, Example 5). In contrast, the conversion into the 2-methylsulfonyl-4,6-bis(difluoromethoxy)pyrimidine and the reaction thereof to 2-amino-4,6-bis(difluoromethoxy)pyrimidine give very good to quantitative yields.

It is therefore the object of the present invention to provide a process for the preparation of 4,6-bis(difluoromethoxy)pyrimidines of formula I in which these compounds can be prepared in good yield.

Accordingly, the present invention postulates preparing the 4,6-bis(difluoromethoxy)pyrimidines of formula I by reacting a 4,6-dihydroxy-pyrimidine dialkali metal salt of formula II



wherein R is as defined for formula I and Me is an alkali metal, with chlorodifluoromethane in a solvent selected from the group consisting of ketones and alkyl cyanides, in the presence of 0.05 to 1.1 mol of water per mol of dialkali metal salt of formula II.

Suitable solvents selected from the group of the ketones are, for example, acetone, methyl ethyl ketone, diethyl ketone, methyl isopropyl ketone or methyl isobutyl ketone. Suitable alkali metal cyanide solvents are, for example, acetonitrile or propionitrile. Preferred solvents are acetonitrile, propionitrile, acetone and methyl ethyl ketone. A particularly preferred solvent is acetonitrile. The above mentioned solvents are conveniently used in an amount of 400 to 3000 ml, preferably 500 to 1000 ml per mol of dialkali metal salt of formula II.

Starting materials and final products of the process of this invention are known. The dialkali metal salts of formula II can be prepared in a manner known per se from the corresponding 4,6-dihydroxypyrimidines, for example by reaction with alkali metal hydroxides or alkali metal alcoholates.

The reaction of the dialkali metal salt of formula II with chlorodifluoromethane can be carried out with particular advantage in the presence of 0.13 to 0.6 mol of water per mol of dialkali metal salt of formula II.

The process of the invention can be carried out in a wide temperature range. Particularly suitable reaction temperatures are in the range from +20° to +100°C. It is preferred to carry out the reaction in the temperature range from +40° to +60°C.

The reaction of the dialkali metal salt of formula II with chlorodifluoromethane can also be conveniently carried out in the presence of a phase transfer catalyst. The phase transfer catalyst can be used in an amount of 0.01 to 0.25 mol per mol of dialkali metal salt of formula II. It is preferred to use 0.05 to 0.15 mol of phase transfer catalyst per mol of dialkali metal salt of formula II.

Suitable phase transfer catalysts are typically quaternary ammonium salts and crown ethers. Preferred phase transfer catalysts are 18-crown-6, benzyltrimethylammonium chloride, tetrabutylammonium chloride, tetramethylammonium methyl sulfate and tetramethylammonium chloride. Tetramethylammonium chloride is especially preferred.

The reaction of the dialkali metal salt of formula II with chlorodifluoromethane can be carried out under normal pressure or under elevated pressure. The reaction is preferably carried out under elevated pressure. Suitable pressures are in the range from 1 to 100 bar. A preferred pressure range in which the reaction may be carried out is from 1 to 20 bar.

An equimolar amount or an excess of chlorodifluoromethane may be used.

It is convenient to use an amount of 1,5 to 10 mol of chlorodifluoromethane per mol of dialkali metal salt of formula II. An amount of 4 to 6 mol of chlorodifluoromethane per mol of dialkali metal salt of formula II is preferred.

A preferred embodiment of the process of this invention comprises reacting a disodium salt of formula II with chlorodifluoromethane in the presence of 0.13 to 0.6 mol of water and 0.05 to 0.15 mol of tetramethylammonium chloride per mol of disodium salt of formula II, in 500 to 1000 ml of acetonitrile per mol of disodium salt of formula II, in the temperature range from +40° to +60°C and under a pressure of 1 to 20 bar.

The process of this invention makes it possible to prepare the 4,6-bis-(difluoromethoxy)pyrimidines of formula I, starting from the 4,6-dihydroxypyrimidine alkali metal salts of formula II, in yields of up to 68 % of theory, whereas, as mentioned at the outset, a yield of only 25 % of theory is obtainable by the known prior art process.

A further advantage of the process of the invention is that the chlorodifluoromethane, which is used in excess, can be reused for a fresh batch, whereas in the known processes it is lost through hydrolysis.

The following Examples illustrate the process of the invention in more detail.

Example 1: Preparation of 4,6-bis(difluoromethoxy)-2-methylthiopyrimidine

In a stirred autoclave, 101 g of anhydrous 4,6-dihydroxy-2-methylthiopyrimidine disodium salt and 8 g of trimethylammonium chloride are mixed with 500 ml of acetonitrile and 2.5 g of water. After closing the autoclave and heating to +50°C, 215 g of chlorodifluoromethane are added over 15 minutes from a steel feed vessel, whereupon the pressure in the autoclave rises to 2 bar.

After a reaction time of 4 hours the autoclave is ventilated. The reaction mixture is then filtered and the filter residue is washed with acetonitrile. The solvent is removed by distillation under vacuum at +80°C and the product melt is washed with 200 ml of hot water, affording 98 g (68 % of theory) of 4,6-bis(difluoromethoxy)-2-methylthiopyrimidine in 90 % purity.

Example 2: The procedure of Example 1 is repeated, using acetone instead of acetonitrile, to give 78 g of 4,6-bis(difluoromethoxy)-2-methylthiopyrimidine (54 % of theory) in 90 % purity.


Example 3: The procedure of Example 1 is repeated, replacing 4,6-dihydroxy-2-methylthiopyrimidine disodium salt by the corresponding 4,6-dihydroxy-2-methylthiopyrimidine dipotassium salt, to give 54 g (38 % of theory) of 4,6-bis(difluoromethoxy)-2-methylthiopyrimidine in 90 % purity.

Example 4: The procedure of Example 1 is repeated, replacing tetramethylammonium chloride by benzyltrimethylammonium chloride, tetrabutylammonium chloride or tetramethylammonium methyl sulfate, to give the same yield of product as in Example 1.

~~What is claimed is:~~

The claims defining the invention are as follows:

1. A process for the preparation of a 4,6-bis(difluoromethoxy)pyrimidine of formula I



wherein R is C₁-C₄alkyl or unsubstituted or substituted phenyl or benzyl, which process comprises reacting a 4,6-dihydroxypyrimidine dialkali metal salt of formula II



wherein R is as defined for formula I and Me is an alkali metal, with chlorodifluoromethane in a solvent selected from the group consisting of ketones and alkyl cyanides, in the presence of 0.05 to 1.1 mol of water per mol of dialkali metal salt of formula II.

2. A process according to claim 1, wherein the solvent is selected from the group consisting of acetone, methyl ethyl ketone, diethyl ketone, methyl isopropyl ketone, methyl isobutyl ketone, acetonitrile and propionitrile.
3. A process according to claim 1, wherein the reaction of the dialkali metal salt of formula II with chlorodifluoromethane is carried out in the presence of 0.13 to 0.6 mol of water per mol of dialkali metal salt of formula II.
4. A process according to claim 1, wherein the solvent is selected from the group consisting of acetonitrile, acetone and methyl ethyl ketone.
5. A process according to claim 1, wherein the solvent is acetonitrile.

6. A process according to claim 1, wherein 400 to 3000 ml per mol of solvent is used per mol of compound of formula II.
7. A process according to claim 1, wherein 500 to 1000 ml per mol of solvent is used per mol of compound of formula II.
8. A process according to claim 1, wherein Me is sodium or potassium.
9. A process according to claim 1, wherein the reaction of the dialkali metal salt of formula II with chlorodifluoromethane is carried out in the presence of a phase transfer catalyst.
10. A process according to claim 9, wherein the phase transfer catalyst is a quaternary ammonium salt or a crown ether.
11. A process according to claim 9, wherein the phase transfer catalyst is selected from the group consisting of 18-crown-6, benzyltrimethylammonium chloride, tetrabutylammonium chloride, tetramethylammonium methyl sulfate and tetramethylammonium chloride.
12. A process according to claim 9, wherein 0.01 to 0.25 mol of phase transfer catalyst is used per mol of dialkali metal salt of formula II.
13. A process according to claim 9, wherein 0.05 to 0.15 mol of phase transfer catalyst is used per mol of dialkali metal salt of formula II.
14. A process according to claim 9, wherein the phase transfer catalyst is tetramethylammonium chloride.
15. A process according to claim 1, wherein the dialkali metal salt of formula II is reacted with chlorodifluoromethane in the temperature range from +20° to +100°C.
16. A process according to claim 1, wherein the dialkali metal salt of formula II is reacted with chlorodifluoromethane in the temperature range from +40° to +60°C.

17. A process according to claim 1, wherein the dialkali metal salt of formula II is reacted with chlorodifluoromethane under a pressure of 1 to 100 bar.
18. A process according to claim 1, wherein the dialkali metal salt of formula II is reacted with chlorodifluoromethane under a pressure of 1 to 20 bar.
19. A process according to claim 1, wherein Me is sodium.
20. A process according to claim 1, wherein 1,5 to 10 mol of chlorodifluoromethane is used per mol of dialkali metal salt of formula II.
21. A process according to claim 1, wherein 4 to 6 mol of chlorodifluoromethane is used per mol of dialkali metal salt of formula II.
22. A process according to claim 1, which comprises reacting a disodium salt of formula II with chlorodifluoromethane in the presence of 0.13 to 0.6 mol of water and 0.05 to 0.15 mol of tetramethylammonium chloride per mol of disodium salt of formula II, in 500 to 1000 ml of acetonitrile per mol of disodium salt of formula II, in the temperature range from +40° to +60°C and under a pressure of 1 to 20 bar.
23. A process for the preparation of a 4,6-bis(difluoromethoxy)pyrimidine substantially as hereinbefore described with reference to any one of the Examples.
24. The product of the process of any one of claims 1 to 23.

DATED this FOURTEENTH day of DECEMBER 1989

Ciba-Geigy AG

Patent Attorneys for the Applicant
SPRUSON & FERGUSON