

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(12) Oversættelse af
europæisk patentskrift

(10) **DK/EP 2432881 T3**

-
- (51) Int.Cl.: **C 12 N 15/113 (2010.01)** **A 61 K 31/713 (2006.01)** **C 12 N 5/074 (2010.01)**
C 12 N 15/63 (2006.01)
- (45) Oversættelsen bekendtgjort den: **2018-02-26**
- (80) Dato for Den Europæiske Patentmyndigheds bekendtgørelse om meddelelse af patentet: **2017-11-15**
- (86) Europæisk ansøgning nr.: **10778258.3**
- (86) Europæisk indleveringsdag: **2010-05-18**
- (87) Den europæiske ansøgnings publiceringsdag: **2012-03-28**
- (86) International ansøgning nr.: **US2010035264**
- (87) Internationalt publikationsnr.: **WO2010135329**
- (30) Prioritet: **2009-05-18 US 179056 P** **2009-08-14 US 233996 P**
2009-12-16 US 286852 P
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV MC MK MT NL NO PL PT RO SE SI SK SM TR**
- (73) Patenthaver: **CuRNA, Inc., 4400 Biscayne Boulevard, Miami, FL 33137, USA**
- (72) Opfinder: **COLLARD, Joseph, 1004 Brooks Lane, Delray Beach, Florida 33483, USA**
KHORKOVA SHERMAN, Olga, 18288 Se, Heritage Drive, Tequesta, Florida 33469, USA
- (74) Fuldmægtig i Danmark: **Orsnes Patent ApS, Forskerparken 10, 5230 Odense M, Danmark**
- (54) Benævnelse: **BEHANDLING AF REPROGRAMMERINGSFAKTORBESLÆGTEDE SYGDOMME VED HÆMNING AF NATURLIGE ANTISENSE-TRANSSKRIFT TIL EN OMPROGRAMMERINGSFAKTOR**
- (56) Fremdragne publikationer:
WO-A1-93/03770
WO-A1-2009/032194
US-A1- 2007 032 447
US-A1- 2007 248 590
US-A1- 2009 068 742
**NA XU ET AL.: "MicroRNA-145 regulates OCT4, SOX2, and KLF4 and represses pluripotency in human Embryonic Stem Cells", CELL, vol. 137, no. 4, 15 May 2009 (2009-05-15), pages 647-658, XP055008854, ISSN: 0092-8674, DOI: 10.1016/j.cell.2009.02.038 -& NA XU ET AL.: "Supplemental Data: MicroRNA-145 regulates OCT4, SOX2, and KLF4 and represses pluripotency in human Embryonic Stem Cells", Cell, vol. 137 15 May 2009 (2009-05-15), XP002712053, Retrieved from the Internet: URL: <http://www.sciencedirect.com/science/MiamiMultiMediaURL/1-s2.0-S0092867409002529/1-s2.0-S0092867409002529-mmc1.pdf/272196/FULL/S0092867409002529/2da421a9b77bf73ee1d9bce734f91d04/mmc1.pdf> [retrieved on 2013-08-29]
MARK RICHARDS ET AL.: "Reverse Serial Analysis of Gene Expression (SAGE) characterization of orphan SAGE tags from human Embryonic Stem Cells identifies the presence of novel transcripts and antisense transcription of key pluripotency genes", STEM CELLS, vol. 24, no. 5, 1 May 2006 (2006-05-01), pages 1162-**

Fortsættes ...

1173, XP055077025, ISSN: 1066-5099, DOI: 10.1634/stemcells.2005-0304

KAZUTOSHI TAKAHASHI ET AL.: "Induction of Pluripotent Stem Cells from mouse embryonic and adult fibroblast cultures by defined factors", CELL, vol. 126, no. 4, 25 August 2006 (2006-08-25), pages 663-676, XP008157317, CELL PRESS, US ISSN: 0092-8674, DOI: 10.1016/J.CELL.2006.07.024 -& KAZUTOSHI TAKAHASHI ET AL.: "Supplemental Data: Induction of Pluripotent Stem Cells from mouse embryonic and adult fibroblast cultures by defined factors", CELL, vol. 126 25 August 2006 (2006-08-25), XP002712054, Retrieved from the Internet: URL: <http://www.sciencedirect.com/science/MiniMultiMediaURL/1-s2.0-S0092867406009767/1-s2.0-S0092867406009767-mmc1.pdf/272196/FULL/S0092867406009767/97d3c3c1f5491871407112b47f4fea6a/mmc1.pdf> [retrieved on 2013-08-30]

KAZUTOSHI TAKAHASHI ET AL.: "Induction of Pluripotent Stem Cells from adult human fibroblasts by defined factors", CELL, CELL PRESS, US, vol. 131, no. 5, 30 November 2007 (2007-11-30), pages 861-872, XP008153593, ISSN: 0092-8674, DOI: 10.1016/J.CELL.2007.11.019

M. SACHDEVA ET AL.: "p53 represses c-Myc through induction of the tumor suppressor miR-145", PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 106, no. 9, 3 March 2009 (2009-03-03), pages 3207-3212, XP055077192, ISSN: 0027-8424, DOI: 10.1073/pnas.0808042106

NIMET MAHERALI ET AL.: "Directly reprogrammed fibroblasts show global epigenetic remodeling and widespread tissue contribution", CELL STEM CELL, vol. 1, no. ISS.1, 2007, pages 55-70, XP009091847,

STORMY J. CHAMBERLAIN ET AL.: "Induced pluripotent stem (iPS) cells as in vitro models of human neurogenetic disorders", NEUROGENETICS, vol. 9, no. 4, 2008, pages 227-235, XP019635665,

DATABASE GENBANK 21 December 2008 'Homo sapiens SRY (sex determining region Y)-box 2 (SOX2) on chromosome 3', XP008164983 Database accession no. NG-009080

NIMET MAHERALI ET AL.: 'Directly Reprogrammed Fibroblasts Show Global Epigenetic Remodeling and Widespread Tissue Contribution.' CELL STEM CELL. vol. 1, no. ISS.1, 2007, pages 55 - 70, XP009091847

STORMY J. CHAMBERLAIN ET AL.: 'Induced pluripotent stem (iPS) cells as in vitro models of human neurogenetic disorders.' NEUROGENETICS. vol. 9, no. 4, 2008, pages 227 - 235, XP019635665

DESCRIPTION

FIELD OF THE DISCLOSURE

[0001] The present application claims the priority of U.S. provisional patent application 61/179,056 filed May 18, 2009, No. 61/233,996 filed August 14, 2009 and U.S. provisional patent application No. 61/286,852 filed Dec 16, 2009.

[0002] Instances of the disclosure comprise oligonucleotides modulating expression and/or function of a Reprogramming factor and associated molecules.

BACKGROUND

[0003] DNA-RNA and RNA-RNA hybridization are important to many aspects of nucleic acid function including DNA replication, transcription, and translation. Hybridization is also central to a variety of technologies that either detect a particular nucleic acid or alter its expression. Antisense nucleotides, for example, disrupt gene expression by hybridizing to target RNA, thereby interfering with RNA splicing, transcription, translation, and replication. Antisense DNA has the added feature that DNA-RNA hybrids serve as a substrate for digestion by ribonuclease H, an activity that is present in most cell types. Antisense molecules can be delivered into cells, as is the case for oligodeoxynucleotides (ODNs), or they can be expressed from endogenous genes as RNA molecules. The FDA recently approved an antisense drug, VITRAVENETM (for treatment of cytomegalovirus retinitis), reflecting that antisense has therapeutic utility. Xu et al. (Cell (2009), 137(4):647-658) describes that MicroRNA-145 regulates OCT4, SOX2 and KLF4 and represses pluripotency in human embryonic stem cells. Richards et al. (Stem Cells (2006), 24(5):1162-1173) describes that reverse serial analysis of gene expression (SAGE) characterization of orphan SAGE tags from human embryonic stem cells identifies the presence of novel transcripts and antisense transcription of key pluripotency genes. Takahashi et al. (Cell (2006), 126(4):663-676) describes the induction of pluripotent stem cells from mouse embryonic and adult fibroblast cultures by defined factors. Takahashi et al. (Cell (2007), 131(5):861-872) describes the induction of pluripotent stem cells from adult human fibroblasts by defined factors. Sachdeva et al. (PNAS (2009), 106(9):3207-3212) describes that p53 represses c-Myc through induction of the tumor suppressor miR-145. US2009/068742 describes a nuclear reprogramming factor. Maherali et al. (Cell Stem Cell (2007), 1 (ISS. 1):55-7) describes that directly reprogrammed fibroblasts show global epigenetic remodelling and widespread tissues contribution. Chamberlain et al. (Neurogenetics (2008), 9(4):227-235) describes induced pluripotent stem cells as in vitro models of human neurogenetic disorders. Lavorgna et al. (Trends in Biochem. Sci. (2004), (29(2):88-94) related to the search for antisense. Lehner et al. (Trends in Genetics (2002), 18(2):63-65) describes antisense transcripts in the human genome.

SUMMARY

[0004] The invention is defined by the claims. Those aspects/instances of the present disclosure

which constitute the invention are defined by the claims.

[0005] In one instance, the disclosure provides methods for inhibiting the action of a natural antisense transcript by using antisense oligonucleotide(s) targeted to any region of the natural antisense transcript resulting in up-regulation of the corresponding sense gene. It is also contemplated herein that inhibition of the natural antisense transcript can be achieved by siRNA, ribozymes and small molecules, which are considered to be within the scope of the present disclosure.

[0006] One instance provides a method of modulating function and/or expression of a Reprogramming factor polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to a reverse complement of a polynucleotide comprising 5 to 30 consecutive nucleotides within nucleotides 1 to 993 of SEQ ID NO: 4, 1 to 493 of SEQ ID NO: 5, and 1 to 418 of SEQ ID NO: 6 thereby modulating function and/or expression of the Reprogramming factor polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0007] In another instance, an oligonucleotide targets a natural antisense sequence of a Reprogramming factor polynucleotide, for example, nucleotides set forth in SEQ ID NO: 4 to 6, and any variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Examples of antisense oligonucleotides are set forth as SEQ ID NOs: 7 to 17.

[0008] Another instance provides a method of modulating function and/or expression of a Reprogramming factor polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to a reverse complement of the an antisense of the Reprogramming factor polynucleotide; thereby modulating function and/or expression of the Reprogramming factor polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0009] Another instance provides a method of modulating function and/or expression of a Reprogramming factor polynucleotide in patient cells or tissues *in vivo* or *in vitro* comprising contacting said cells or tissues with an antisense oligonucleotide 5 to 30 nucleotides in length wherein said oligonucleotide has at least 50% sequence identity to an antisense oligonucleotide to a Reprogramming factor antisense polynucleotide; thereby modulating function and/or expression of the Reprogramming factor polynucleotide in patient cells or tissues *in vivo* or *in vitro*.

[0010] In one instance, a composition comprises one or more antisense oligonucleotides which bind to sense and/or antisense Reprogramming factor polynucleotides.

[0011] In another instance, the oligonucleotides comprise one or more modified or substituted nucleotides.

[0012] In another instance, the oligonucleotides comprise one or more modified bonds.

[0013] In yet another instance, the modified nucleotides comprise modified bases comprising phosphorothioate, methylphosphonate, peptide nucleic acids, 2'-O-methyl, fluoro- or carbon, methylene or other locked nucleic acid (LNA) molecules. Preferably, the modified nucleotides are

locked nucleic acid molecules, including α -L-LNA.

[0014] In another instance, the oligonucleotides are administered to a patient subcutaneously, intramuscularly, intravenously or intraperitoneally.

[0015] In another instance, the oligonucleotides are administered in a pharmaceutical composition. A treatment regimen comprises administering the antisense compounds at least once to patient; however, this treatment can be modified to include multiple doses over a period of time. The treatment can be combined with one or more other types of therapies.

[0016] In another instance, the oligonucleotides are encapsulated in a liposome or attached to a carrier molecule (e.g. cholesterol, TAT peptide).

[0017] Other aspects are described *infra*.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1:

[0018]

Figure 1A is a graph of real time PCR results showing the fold change + standard deviation in SOX2 mRNA after treatment of HepG2 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Real time PCR results show that the levels of SOX2 mRNA in HepG2 cells are significantly increased 48 h after treatment with two of the siRNAs designed to SOX2 antisense AI885646. Bars denoted as CUR-0404 and CUR-0406 correspond to samples treated with SEQ ID NOS: 7 and 8 respectively.

Figure 1B is a graph of real time PCR results showing the fold change + standard deviation in KLF4 mRNA after treatment of HepG2 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Real time PCR results show that the levels of KLF4 mRNA in HepG2 cells are significantly increased 48 h after treatment with two of the oligos designed to KLF4 antisense. Bars denoted as CUR-0933, CUR-0931, CUR-0930 and CUR-0932 correspond to samples treated with SEQ ID NOS: 9 to 12 respectively.

Figure 1C is a graph of real time PCR results showing the fold change + standard deviation in POU5F1 mRNA after treatment of HepG2 cells with phosphorothioate oligonucleotides introduced using Lipofectamine 2000, as compared to control. Bars denoted as CUR-1139 to CUR-1143 correspond to samples treated with SEQ ID NOS 13 to 17 respectively.

DETAILED DESCRIPTION

[0019] Several aspects of the disclosure are described below with reference to example applications for illustration. It should be understood that numerous specific details, relationships, and methods are set forth to provide a full understanding of the disclosure. One having ordinary skill in the relevant art, however, will readily recognize that the disclosure can be practiced without one or more of the specific details or with other methods. The present disclosure is not limited by the ordering of acts or events, as some acts may occur in different orders and/or concurrently with other acts or events. Furthermore, not all illustrated acts or events are required to implement a methodology in accordance with the present disclosure.

[0020] All genes, gene names, and gene products disclosed herein are intended to correspond to homologs from any species for which the compositions and methods disclosed herein are applicable. Thus, the terms include, but are not limited to genes and gene products from humans and mice. It is understood that when a gene or gene product from a particular species is disclosed, this disclosure is intended to be exemplary only, and is not to be interpreted as a limitation unless the context in which it appears clearly indicates. Thus, for example, for the genes disclosed herein, which in some instances relate to mammalian nucleic acid and amino acid sequences are intended to encompass homologous and/or orthologous genes and gene products from other animals including, but not limited to other mammals, fish, amphibians, reptiles, and birds. In preferred instances, the genes or nucleic acid sequences are human.

Definitions

[0021] The terminology used herein is for the purpose of describing particular instances only and is not intended to be limiting of the disclosure. As used herein, the singular forms "a", "an" and "the" are intended to include the plural forms as well, unless the context clearly indicates otherwise. Furthermore, to the extent that the terms "including", "includes", "having", "has", "with", or variants thereof are used in either the detailed description and/or the claims, such terms are intended to be inclusive in a manner similar to the term "comprising."

[0022] The term "about" or "approximately" means within an acceptable error range for the particular value as determined by one of ordinary skill in the art, which will depend in part on how the value is measured or determined, i.e., the limitations of the measurement system. For example, "about" can mean within 1 or more than 1 standard deviation, per the practice in the art. Alternatively, "about" can mean a range of up to 20%, preferably up to 10%, more preferably up to 5%, and more preferably still up to 1% of a given value. Alternatively, particularly with respect to biological systems or processes, the term can mean within an order of magnitude, preferably within 5-fold, and more preferably within 2-fold, of a value. Where particular values are described in the application and claims, unless otherwise stated the term "about" meaning within an acceptable error range for the particular value should be assumed.

[0023] As used herein, the term "mRNA" means the presently known mRNA transcript(s) of a targeted gene, and any further transcripts which may be elucidated.

[0024] By "antisense oligonucleotides" or "antisense compound" is meant an RNA or DNA molecule that binds to another RNA or DNA (target RNA, DNA). For example, if it is an RNA oligonucleotide it

binds to another RNA target by means of RNA-RNA interactions and alters the activity of the target RNA (Eguchi et al., (1991) Ann. Rev. Biochem. 60, 631-652). An antisense oligonucleotide can upregulate or downregulate expression and/or function of a particular polynucleotide. The definition is meant to include any foreign RNA or DNA molecule which is useful from a therapeutic, diagnostic, or other viewpoint. Such molecules include, for example, antisense RNA or DNA molecules, interference RNA (RNAi), micro RNA, decoy RNA molecules, siRNA, enzymatic RNA, therapeutic editing RNA and agonist and antagonist RNA, antisense oligomeric compounds, antisense oligonucleotides, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds that hybridize to at least a portion of the target nucleic acid. As such, these compounds may be introduced in the form of single-stranded, double-stranded, partially single-stranded, or circular oligomeric compounds.

[0025] In the context of this disclosure, the term "oligonucleotide" refers to an oligomer or polymer of ribonucleic acid (RNA) or deoxyribonucleic acid (DNA) or mimetics thereof. The term "oligonucleotide", also includes linear or circular oligomers of natural and/or modified monomers or linkages, including deoxyribonucleosides, ribonucleosides, substituted and alpha-anomeric forms thereof, peptide nucleic acids (PNA), locked nucleic acids (LNA), phosphorothioate, methylphosphonate, and the like. Oligonucleotides are capable of specifically binding to a target polynucleotide by way of a regular pattern of monomer-to-monomer interactions, such as Watson-Crick type of base pairing, Hoögsteeen or reverse Hoögsteeen types of base pairing, or the like.

[0026] The oligonucleotide may be "chimeric", that is, composed of different regions. In the context of this disclosure "chimeric" compounds are oligonucleotides, which contain two or more chemical regions, for example, DNA region(s), RNA region(s), PNA region(s) etc. Each chemical region is made up of at least one monomer unit, i.e., a nucleotide in the case of an oligonucleotides compound. These oligonucleotides typically comprise at least one region wherein the oligonucleotide is modified in order to exhibit one or more desired properties. The desired properties of the oligonucleotide include, but are not limited, for example, to increased resistance to nuclease degradation, increased cellular uptake, and/or increased binding affinity for the target nucleic acid. Different regions of the oligonucleotide may therefore have different properties. The chimeric oligonucleotides of the present disclosure can be formed as mixed structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotide analogs as described above.

[0027] The oligonucleotide can be composed of regions that can be linked in "register" that is, when the monomers are linked consecutively, as in native DNA, or linked via spacers. The spacers are intended to constitute a covalent "bridge" between the regions and have in preferred cases a length not exceeding about 100 carbon atoms. The spacers may carry different functionalities, for example, having positive or negative charge, carry special nucleic acid binding properties (intercalators, groove binders, toxins, fluorophors etc.), being lipophilic, inducing special secondary structures like, for example, alanine containing peptides that induce alpha-helices.

[0028] As used herein "Reprogramming factors" and "Reprogramming factor" are inclusive of all family members, mutants, alleles, fragments, species, coding and noncoding sequences, sense and antisense polynucleotide strands, etc.

[0029] As used herein, the words SOX2, Sox-2, ANOP3, MCOPS3, MGC2413 and Transcription

factor SOX-2 are used interchangeably in the present application.

[0030] As used herein, the words KLF4, Kruppel-like factor 4 (gut), Epithelial zinc finger protein EZF, EZF, GKLf, Gut-enriched krueppel-like factor, Krueppel-like factor 4 are used interchangeably in the present application.

[0031] As used herein, the words POU5F1, Oct-3, Oct-3/4, Oct-4, Octamer-binding transcription factor 3, Octamer-binding transcription factor 4, OTF3, OTF4, 'POU domain class 5, transcription factor 1', POU5F1, POU class 5 homeobox 1, DADB-104B20.2 and MGC22487 are considered the same in the literature and are used interchangeably in the present application.

[0032] As used herein, the term "oligonucleotide specific for" or "oligonucleotide which targets" refers to an oligonucleotide having a sequence (i) capable of forming a stable complex with a portion of the targeted gene, or (ii) capable of forming a stable duplex with a portion of a mRNA transcript of the targeted gene. Stability of the complexes and duplexes can be determined by theoretical calculations and/or in vitro assays. Exemplary assays for determining stability of hybridization complexes and duplexes are described in the Examples below.

[0033] As used herein, the term "target nucleic acid" encompasses DNA, RNA (comprising premRNA and mRNA) transcribed from such DNA, and also cDNA derived from such RNA, coding, noncoding sequences, sense or antisense polynucleotides. The specific hybridization of an oligomeric compound with its target nucleic acid interferes with the normal function of the nucleic acid. This modulation of function of a target nucleic acid by compounds, which specifically hybridize to it, is generally referred to as "antisense". The functions of DNA to be interfered include, for example, replication and transcription. The functions of RNA to be interfered, include all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The overall effect of such interference with target nucleic acid function is modulation of the expression of an encoded product or oligonucleotides.

[0034] RNA interference "RNAi" is mediated by double stranded RNA (dsRNA) molecules that have sequence-specific homology to their "target" nucleic acid sequences (Caplen, N. J., et al. (2001) Proc. Natl. Acad. Sci. USA 98:9742-9747). In certain instances of the present disclosure, the mediators are 5-25 nucleotide "small interfering" RNA duplexes (siRNAs). The siRNAs are derived from the processing of dsRNA by an RNase enzyme known as Dicer (Bernstein, E., et al. (2001) Nature 409:363-366). siRNA duplex products are recruited into a multi-protein siRNA complex termed RISC (RNA Induced Silencing Complex). Without wishing to be bound by any particular theory, a RISC is then believed to be guided to a target nucleic acid (suitably mRNA), where the siRNA duplex interacts in a sequence-specific way to mediate cleavage in a catalytic fashion (Bernstein, E., et al. (2001) Nature 409:363-366; Boutla, A., et al. (2001) Curr. Biol. 11:1776-1780). Small interfering RNAs that can be used in accordance with the present disclosure can be synthesized and used according to procedures that are well known in the art and that will be familiar to the ordinarily skilled artisan. Small interfering RNAs for use in the methods of the present disclosure suitably comprise between about 1 to about 50 nucleotides (nt). In examples of non limiting instances, siRNAs can comprise about 5 to about 40 nt, about 5 to about 30 nt, about 10 to about 30 nt, about 15 to about 25 nt, or about 20-25 nucleotides.

[0035] Selection of appropriate oligonucleotides is facilitated by using computer programs that automatically align nucleic acid sequences and indicate regions of identity or homology. Such programs are used to compare nucleic acid sequences obtained, for example, by searching databases such as GenBank or by sequencing PCR products. Comparison of nucleic acid sequences from a range of species allows the selection of nucleic acid sequences that display an appropriate degree of identity between species. In the case of genes that have not been sequenced, Southern blots are performed to allow a determination of the degree of identity between genes in target species and other species. By performing Southern blots at varying degrees of stringency, as is well known in the art, it is possible to obtain an approximate measure of identity. These procedures allow the selection of oligonucleotides that exhibit a high degree of complementarity to target nucleic acid sequences in a subject to be controlled and a lower degree of complementarity to corresponding nucleic acid sequences in other species. One skilled in the art will realize that there is considerable latitude in selecting appropriate regions of genes for use in the present disclosure.

[0036] By "enzymatic RNA" is meant an RNA molecule with enzymatic activity (Cech, (1988) J. American. Med. Assoc. 260, 3030-3035). Enzymatic nucleic acids (ribozymes) act by first binding to a target RNA. Such binding occurs through the target binding portion of an enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through base pairing, and once bound to the correct site, acts enzymatically to cut the target RNA.

[0037] By "decoy RNA" is meant an RNA molecule that mimics the natural binding domain for a ligand. The decoy RNA therefore competes with natural binding target for the binding of a specific ligand. For example, it has been shown that over-expression of HIV trans-activation response (TAR) RNA can act as a "decoy" and efficiently binds HIV tat protein, thereby preventing it from binding to TAR sequences encoded in the HIV RNA (Sullenger et al. (1990) Cell, 63, 601- 608). This is meant to be a specific example. Those in the art will recognize that this is but one example, and other instances can be readily generated using techniques generally known in the art.

[0038] As used herein, the term "monomers" typically indicates monomers linked by phosphodiester bonds or analogs thereof to form oligonucleotides ranging in size from a few monomeric units, e.g., from about 3-4, to about several hundreds of monomeric units. Analogs of phosphodiester linkages include: phosphorothioate, phosphorodithioate, methylphosphonates, phosphoroselenoate, phosphoramidate, and the like, as more fully described below.

[0039] The term "nucleotide" covers naturally occurring nucleotides as well as nonnaturally occurring nucleotides. It should be clear to the person skilled in the art that various nucleotides which previously have been considered "non-naturally occurring" have subsequently been found in nature. Thus, "nucleotides" includes not only the known purine and pyrimidine heterocycles-containing molecules, but also heterocyclic analogues and tautomers thereof. Illustrative examples of other types of nucleotides are molecules containing adenine, guanine, thymine, cytosine, uracil, purine, xanthine, diaminopurine, 8-oxo- N6-methyladenine, 7-deazaxanthine, 7-deazaguanine, N4,N4-ethanocytosin, N6,N6-ethano-2,6- diaminopurine, 5-methylcytosine, 5-(C3-C6)-alkynylcytosine, 5-fluorouracil, 5-bromouracil, pseudoisocytosine, 2-hydroxy-5-methyl-4-triazolopyridin, isocytosine, isoguanin, inosine and the "non-naturally occurring" nucleotides described in Benner et al., U.S. Pat No. 5,432,272. The term "nucleotide" is intended to cover every and all of these examples as well as

analogues and tautomers thereof. Especially interesting nucleotides are those containing adenine, guanine, thymine, cytosine, and uracil, which are considered as the naturally occurring nucleotides in relation to therapeutic and diagnostic application in humans. Nucleotides include the natural 2'-deoxy and 2'-hydroxyl sugars, e.g., as described in Komberg and Baker, *DNA Replication*, 2nd Ed. (Freeman, San Francisco, 1992) as well as their analogs.

[0040] "Analog" in reference to nucleotides includes synthetic nucleotides having modified base moieties and/or modified sugar moieties (see e.g., described generally by Scheit, *Nucleotide Analogs*, John Wiley, New York, 1980; Freier & Altmann, (1997) *Nucl. Acid. Res.*, 25(22), 4429-4443, Toulmé, J.J., (2001) *Nature Biotechnology* 19:17-18; Manoharan M., (1999) *Biochimica et Biophysica Acta* 1489:117-139; Freier S. M., (1997) *Nucleic Acid Research*, 25:4429-4443, Uhlman, E., (2000) *Drug Discovery & Development*, 3: 203-213, Herdewin P., (2000) *Antisense & Nucleic Acid Drug Dev.*, 10:297-310; 2'-O, 3'-C-linked [3.2.0] bicycloarabinonucleosides (see e.g. N.K Christensen., et al, (1998) *J. Am. Chem. Soc.*, 120: 5458-5463; Prakash TP, Bhat B. (2007) *Curr Top Med Chem.* 7(7):641-9; Cho EJ, et al. (2009) *Annual Review of Analytical Chemistry*, 2, 241-264). Such analogs include synthetic nucleotides designed to enhance binding properties, e.g., duplex or triplex stability, specificity, or the like.

[0041] As used herein, "hybridization" means the pairing of substantially complementary strands of oligomeric compounds. One mechanism of pairing involves hydrogen bonding, which may be Watson-Crick, Hoogsteen or reversed Hoogsteen hydrogen bonding, between complementary nucleoside or nucleotide bases (nucleotides) of the strands of oligomeric compounds. For example, adenine and thymine are complementary nucleotides which pair through the formation of hydrogen bonds. Hybridization can occur under varying circumstances.

[0042] An antisense compound is "specifically hybridizable" when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a modulation of function and/or activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays.

[0043] As used herein, the phrase "stringent hybridization conditions" or "stringent conditions" refers to conditions under which a compound of the disclosure will hybridize to its target sequence, but to a minimal number of other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances and in the context of this disclosure, "stringent conditions" under which oligomeric compounds hybridize to a target sequence are determined by the nature and composition of the oligomeric compounds and the assays in which they are being investigated. In general, stringent hybridization conditions comprise low concentrations (<0.15M) of salts with inorganic cations such as Na⁺⁺ or K⁺⁺ (i.e., low ionic strength), temperature higher than 20°C - 25°C below the T_m of the oligomeric compound:target sequence complex, and the presence of denaturants such as formamide, dimethylformamide, dimethyl sulfoxide, or the detergent sodium dodecyl sulfate (SDS). For example, the hybridization rate decreases 1.1% for each 1% formamide. An example of a high stringency hybridization condition is 0.1X sodium chloride-sodium citrate buffer (SSC)/0.1% (w/v) SDS at 60°C for 30 minutes.

[0044] "Complementary," as used herein, refers to the capacity for precise pairing between two nucleotides on one or two oligomeric strands. For example, if a nucleobase at a certain position of an antisense compound is capable of hydrogen bonding with a nucleobase at a certain position of a target nucleic acid, said target nucleic acid being a DNA, RNA, or oligonucleotide molecule, then the position of hydrogen bonding between the oligonucleotide and the target nucleic acid is considered to be a complementary position. The oligomeric compound and the further DNA, RNA, or oligonucleotide molecule are complementary to each other when a sufficient number of complementary positions in each molecule are occupied by nucleotides which can hydrogen bond with each other. Thus, "specifically hybridizable" and "complementary" are terms which are used to indicate a sufficient degree of precise pairing or complementarity over a sufficient number of nucleotides such that stable and specific binding occurs between the oligomeric compound and a target nucleic acid.

[0045] It is understood in the art that the sequence of an oligomeric compound need not be 100% complementary to that of its target nucleic acid to be specifically hybridizable. Moreover, an oligonucleotide may hybridize over one or more segments such that intervening or adjacent segments are not involved in the hybridization event (e.g., a loop structure, mismatch or hairpin structure). The oligomeric compounds of the present disclosure comprise at least about 70%, or at least about 75%, or at least about 80%, or at least about 85%, or at least about 90%, or at least about 95%, or at least about 99% sequence complementarity to a target region within the target nucleic acid sequence to which they are targeted. For example, an antisense compound in which 18 of 20 nucleotides of the antisense compound are complementary to a target region, and would therefore specifically hybridize, would represent 90 percent complementarity. In this example, the remaining noncomplementary nucleotides may be clustered or interspersed with complementary nucleotides and need not be contiguous to each other or to complementary nucleotides. As such, an antisense compound which is 18 nucleotides in length having 4 (four) noncomplementary nucleotides which are flanked by two regions of complete complementarity with the target nucleic acid would have 77.8% overall complementarity with the target nucleic acid and would thus fall within the scope of the present disclosure. Percent complementarity of an antisense compound with a region of a target nucleic acid can be determined routinely using BLAST programs (basic local alignment search tools) and PowerBLAST programs known in the art (Altschul et al., (1990) J. Mol. Biol., 215, 403-410; Zhang and Madden, (1997) Genome Res., 7, 649-656). Percent homology, sequence identity or complementarity, can be determined by, for example, the Gap program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, Madison Wis.), using default settings, which uses the algorithm of Smith and Waterman (Adv. Appl. Math., (1981) 2, 482-489).

[0046] As used herein, the term "Thermal Melting Point (T_m)" refers to the temperature, under defined ionic strength, pH, and nucleic acid concentration, at which 50% of the oligonucleotides complementary to the target sequence hybridize to the target sequence at equilibrium. Typically, stringent conditions will be those in which the salt concentration is at least about 0.01 to 1.0 M Na ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30°C for short oligonucleotides (e.g., 10 to 50 nucleotide). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide.

[0047] As used herein, "modulation" means either an increase (stimulation) or a decrease (inhibition) in the expression of a gene.

[0048] The term "variant," when used in the context of a polynucleotide sequence, may encompass a polynucleotide sequence related to a wild type gene. This definition may also include, for example, "allelic," "splice," "species," or "polymorphic" variants. A splice variant may have significant identity to a reference molecule, but will generally have a greater or lesser number of polynucleotides due to alternate splicing of exons during mRNA processing. The corresponding polypeptide may possess additional functional domains or an absence of domains. Species variants are polynucleotide sequences that vary from one species to another. Of particular utility in the disclosure are variants of wild type gene products. Variants may result from at least one mutation in the nucleic acid sequence and may result in altered mRNAs or in polypeptides whose structure or function may or may not be altered. Any given natural or recombinant gene may have none, one, or many allelic forms. Common mutational changes that give rise to variants are generally ascribed to natural deletions, additions, or substitutions of nucleotides. Each of these types of changes may occur alone, or in combination with the others, one or more times in a given sequence.

[0049] The resulting polypeptides generally will have significant amino acid identity relative to each other. A polymorphic variant is a variation in the polynucleotide sequence of a particular gene between individuals of a given species. Polymorphic variants also may encompass "single nucleotide polymorphisms" (SNPs,) or single base mutations in which the polynucleotide sequence varies by one base. The presence of SNPs may be indicative of, for example, a certain population with a propensity for a disease state, that is susceptibility versus resistance.

[0050] Derivative polynucleotides include nucleic acids subjected to chemical modification, for example, replacement of hydrogen by an alkyl, acyl, or amino group. Derivatives, e.g., derivative oligonucleotides, may comprise non-naturally-occurring portions, such as altered sugar moieties or inter-sugar linkages. Exemplary among these are phosphorothioate and other sulfur containing species which are known in the art. Derivative nucleic acids may also contain labels, including radionucleotides, enzymes, fluorescent agents, chemiluminescent agents, chromogenic agents, substrates, cofactors, inhibitors, magnetic particles, and the like.

[0051] A "derivative" polypeptide or peptide is one that is modified, for example, by glycosylation, pegylation, phosphorylation, sulfation, reduction/alkylation, acylation, chemical coupling, or mild formalin treatment. A derivative may also be modified to contain a detectable label, either directly or indirectly, including, but not limited to, a radioisotope, fluorescent, and enzyme label.

[0052] As used herein, the term "animal" or "patient" is meant to include, for example, humans, sheep, elks, deer, mule deer, minks, mammals, monkeys, horses, cattle, pigs, goats, dogs, cats, rats, mice, birds, chicken, reptiles, fish, insects and arachnids.

[0053] "Mammal" covers warm blooded mammals that are typically under medical care (e.g., humans and domesticated animals). Examples include feline, canine, equine, bovine, and human, as well as just human.

[0054] "Treating" or "treatment" covers the treatment of a disease-state in a mammal, and includes: (a) preventing the disease-state from occurring in a mammal, in particular, when such mammal is predisposed to the disease-state but has not yet been diagnosed as having it; (b) inhibiting the disease-state, e.g., arresting its development; and/or (c) relieving the disease-state, e.g., causing

regression of the disease state until a desired endpoint is reached. Treating also includes the amelioration of a symptom of a disease (e.g., lessen the pain or discomfort), wherein such amelioration may or may not be directly affecting the disease (e.g., cause, transmission, expression, etc.).

[0055] As used herein, "cancer" refers to all types of cancer or neoplasm or malignant tumors found in mammals, including, but not limited to: leukemias, lymphomas, melanomas, carcinomas and sarcomas. The cancer manifests itself as a "tumor" or tissue comprising malignant cells of the cancer. Examples of tumors include sarcomas and carcinomas such as, but not limited to: fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovioma, mesothelioma, Ewing's tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, choriocarcinoma, seminoma, embryonal carcinoma, Wilms' tumor, cervical cancer, testicular tumor, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, and retinoblastoma. Additional cancers which can be treated by the disclosed composition according to the disclosure include but not limited to, for example, Hodgkin's Disease, Non-Hodgkin's Lymphoma, multiple myeloma, neuroblastoma, breast cancer, ovarian cancer, lung cancer, rhabdomyosarcoma, primary thrombocytosis, primary macroglobulinemia, small-cell lung tumors, primary brain tumors, stomach cancer, colon cancer, malignant pancreatic insulanoma, malignant carcinoid, urinary bladder cancer, premalignant skin lesions, testicular cancer, lymphomas, thyroid cancer, neuroblastoma, esophageal cancer, genitourinary tract cancer, malignant hypercalcemia, cervical cancer, endometrial cancer, adrenal cortical cancer, and prostate cancer.

[0056] "Neurological disease or disorder" refers to any disease or disorder of the nervous system and/or visual system. "Neurological disease or disorder" include disease or disorders that involve the central nervous system (brain, brainstem and cerebellum), the peripheral nervous system (including cranial nerves), and the autonomic nervous system (parts of which are located in both central and peripheral nervous system). Examples of neurological disorders include but are not limited to, headache, stupor and coma, dementia, seizure, sleep disorders, trauma, infections, neoplasms, neuroophthalmology, movement disorders, demyelinating diseases, spinal cord disorders, and disorders of peripheral nerves, muscle and neuromuscular junctions. Addiction and mental illness, include, but are not limited to, bipolar disorder and schizophrenia, are also included in the definition of neurological disorder. The following is a list of several neurological disorders, symptoms, signs and syndromes that can be treated using compositions and methods according to the present disclosure: acquired epileptiform aphasia; acute disseminated encephalomyelitis; adrenoleukodystrophy; age-related macular degeneration; agenesis of the corpus callosum; agnosia; Aicardi syndrome; Alexander disease; Alpers' disease; alternating hemiplegia; Vascular dementia; amyotrophic lateral sclerosis; anencephaly; Angelman syndrome; angiomatosis; anoxia; aphasia; apraxia; arachnoid cysts; arachnoiditis; Anronl-Chiari malformation; arteriovenous malformation; Asperger syndrome; ataxia telegiectasia; attention deficit hyperactivity disorder; autism; autonomic dysfunction; back pain; Batten disease; Behcet's disease; Bell's palsy; benign

essential blepharospasm; benign focal; amyotrophy; benign intracranial hypertension; Binswanger's disease; blepharospasm; Bloch Sulzberger syndrome; brachial plexus injury; brain abscess; brain injury; brain tumors (including glioblastoma multiforme); spinal tumor; Brown-Sequard syndrome; Canavan disease; carpal tunnel syndrome; causalgia; central pain syndrome; central pontine myelinolysis; cephalic disorder; cerebral aneurysm; cerebral arteriosclerosis; cerebral atrophy; cerebral gigantism; cerebral palsy; Charcot-Marie-Tooth disease; chemotherapy-induced neuropathy and neuropathic pain; Chiari malformation; chorea; chronic inflammatory demyelinating polyneuropathy; chronic pain; chronic regional pain syndrome; Coffin Lowry syndrome; coma, including persistent vegetative state; congenital facial diplegia; corticobasal degeneration; cranial arteritis; craniosynostosis; Creutzfeldt-Jakob disease; cumulative trauma disorders; Cushing's syndrome; cytomegalic inclusion body disease; cytomegalovirus infection; dancing eyes-dancing feet syndrome; DandyWalker syndrome; Dawson disease; De Morsier's syndrome; Dejerine-Klumke palsy; dementia; dermatomyositis; diabetic neuropathy; diffuse sclerosis; dysautonomia; dysgraphia; dyslexia; dystonias; early infantile epileptic encephalopathy; empty sella syndrome; encephalitis; encephaloceles; encephalotrigeminal angiomas; epilepsy; Erb's palsy; essential tremor; Fabry's disease; Fahr's syndrome; fainting; familial spastic paralysis; febrile seizures; Fisher syndrome; Friedreich's ataxia; fronto-temporal dementia and other "tauopathies"; Gaucher's disease; Gerstmann's syndrome; giant cell arteritis; giant cell inclusion disease; globoid cell leukodystrophy; Guillain-Barre syndrome; HTLV-1-associated myelopathy; Hallervorden-Spatz disease; head injury; headache; hemifacial spasm; hereditary spastic paraplegia; heredopathia atactica et polyneuritiformis; herpes zoster oticus; herpes zoster; Hirayama syndrome; HIV-associated dementia and neuropathy (also neurological manifestations of AIDS); holoprosencephaly; Huntington's disease and other polyglutamine repeat diseases; hydranencephaly; hydrocephalus; hypercortisolism; hypoxia; immune-mediated encephalomyelitis; inclusion body myositis; incontinentia pigmenti; infantile phytanic acid storage disease; infantile reflux disease; infantile spasms; inflammatory myopathy; intracranial cyst; intracranial hypertension; Joubert syndrome; Keams-Sayre syndrome; Kennedy disease; Kinsbourne syndrome; Klippel Feil syndrome; Krabbe disease; Kugelberg-Welander disease; kuru; Lafora disease; Lambert-Eaton myasthenic syndrome; Landau-Kleffner syndrome; lateral medullary (Wallenberg) syndrome; learning disabilities; Leigh's disease; Lennox-Gustaut syndrome; Lesch-Nyhan syndrome; leukodystrophy; Lewy body dementia; Lissencephaly; locked-in syndrome; Lou Gehrig's disease (i.e., motor neuron disease or amyotrophic lateral sclerosis); lumbar disc disease; Lyme disease-neurological sequelae; Machado-Joseph disease; macrencephaly; megalencephaly; Melkersson-Rosenthal syndrome; Menieres disease; meningitis; Menkes disease; metachromatic leukodystrophy; microcephaly; migraine; Miller Fisher syndrome; mini-strokes; mitochondrial myopathies; Mobius syndrome; monomelic amyotrophy; motor neuron disease; Moyamoya disease; mucopolysaccharidoses; multi-infarct dementia; multifocal motor neuropathy; multiple sclerosis and other demyelinating disorders; multiple system atrophy with postural hypotension; p muscular dystrophy; myasthenia gravis; myelinoclastic diffuse sclerosis; myoclonic encephalopathy of infants; myoclonus; myopathy; myotonia congenita; narcolepsy; neurofibromatosis; neuroleptic malignant syndrome; neurological manifestations of AIDS; neurological sequelae of lupus; neuromyotonia; neuronal ceroid lipofuscinosis; neuronal migration disorders; Niemann-Pick disease; O'Sullivan-McLeod syndrome; occipital neuralgia; occult spinal dysraphism sequence; Ohtahara syndrome; olivopontocerebellar atrophy; opsoclonus myoclonus; optic neuritis; orthostatic hypotension; overuse syndrome; paresthesia; Neurodegenerative disease or disorder (Parkinson's disease, Huntington's disease, Alzheimer's disease, amyotrophic lateral sclerosis (ALS), dementia, multiple sclerosis and other diseases and disorders associated with neuronal cell death); paramyotonia congenita; paraneoplastic diseases; paroxysmal attacks; Parry

Romberg syndrome; Pelizaeus-Merzbacher disease; periodic paralyses; peripheral neuropathy; painful neuropathy and neuropathic pain; persistent vegetative state; pervasive developmental disorders; photic sneeze reflex; phytanic acid storage disease; Pick's disease; pinched nerve; pituitary tumors; polymyositis; porencephaly; post-polio syndrome; postherpetic neuralgia; postinfectious encephalomyelitis; postural hypotension; Prader- Willi syndrome; primary lateral sclerosis; prion diseases; progressive hemifacial atrophy; progressive multifocalleukoencephalopathy; progressive sclerosing poliodystrophy; progressive supranuclear palsy; pseudotumor cerebri; Ramsay-Hunt syndrome (types I and II); Rasmussen's encephalitis; reflex sympathetic dystrophy syndrome; Refsum disease; repetitive motion disorders; repetitive stress injuries; restless legs syndrome; retrovirus-associated myelopathy; Rett syndrome; Reye's syndrome; Saint Vitus dance; Sandhoff disease; Schilder's disease; schizencephaly; septo-optic dysplasia; shaken baby syndrome; shingles; Shy-Drager syndrome; Sjogren's syndrome; sleep apnea; Soto's syndrome; spasticity; spina bifida; spinal cord injury; spinal cord tumors; spinal muscular atrophy; Stiff-Person syndrome; stroke; Sturge-Weber syndrome; subacute sclerosing panencephalitis; subcortical arteriosclerotic encephalopathy; Sydenham chorea; syncope; syringomyelia; tardive dyskinesia; Tay-Sachs disease; temporal arteritis; tethered spinal cord syndrome; Thomsen disease; thoracic outlet syndrome; Tic Douloureux; Todd's paralysis; Tourette syndrome; transient ischemic attack; transmissible spongiform encephalopathies; transverse myelitis; traumatic brain injury; tremor; trigeminal neuralgia; tropical spastic paraparesis; tuberous sclerosis; vascular dementia (multi-infarct dementia); vasculitis including temporal arteritis; Von Hippel-Lindau disease; Wallenberg's syndrome; Werdnig-Hoffman disease; West syndrome; whiplash; Williams syndrome; Wildon's disease; and Zellweger syndrome.

[0057] An "Inflammation" refers to systemic inflammatory conditions and conditions associated locally with migration and attraction of monocytes, leukocytes and/or neutrophils. Examples of inflammation include, but are not limited to, Inflammation resulting from infection with pathogenic organisms (including gram-positive bacteria, gram-negative bacteria, viruses, fungi, and parasites such as protozoa and helminths), transplant rejection (including rejection of solid organs such as kidney, liver, heart, lung or cornea, as well as rejection of bone marrow transplants including graft-versus-host disease (GVHD)), or from localized chronic or acute autoimmune or allergic reactions. Autoimmune diseases include acute glomerulonephritis; rheumatoid or reactive arthritis; chronic glomerulonephritis; inflammatory bowel diseases such as Crohn's disease, ulcerative colitis and necrotizing enterocolitis; granulocyte transfusion associated syndromes; inflammatory dermatoses such as contact dermatitis, atopic dermatitis, psoriasis; systemic lupus erythematosus (SLE), autoimmune thyroiditis, multiple sclerosis, and some forms of diabetes, or any other autoimmune state where attack by the subject's own immune system results in pathologic tissue destruction. Allergic reactions include allergic asthma, chronic bronchitis, acute and delayed hypersensitivity. Systemic inflammatory disease states include inflammation associated with trauma, burns, reperfusion following ischemic events (e.g. thrombotic events in heart, brain, intestines or peripheral vasculature, including myocardial infarction and stroke), sepsis, ARDS or multiple organ dysfunction syndrome. Inflammatory cell recruitment also occurs in atherosclerotic plaques. Inflammation includes, but is not limited to, Non-Hodgkin's lymphoma, Wegener's granulomatosis, Hashimoto's thyroiditis, hepatocellular carcinoma, thymus atrophy, chronic pancreatitis, rheumatoid arthritis, reactive lymphoid hyperplasia, osteoarthritis, ulcerative colitis, papillary carcinoma, Crohn's disease, ulcerative colitis, acute cholecystitis, chronic cholecystitis, cirrhosis, chronic sialadenitis, peritonitis, acute pancreatitis, chronic pancreatitis, chronic Gastritis, adenomyosis, endometriosis, acute cervicitis, chronic cervicitis, lymphoid hyperplasia, multiple sclerosis, hypertrophy secondary to

idiopathic thrombocytopenic purpura, primary IgA nephropathy, systemic lupus erythematosus, psoriasis, pulmonary emphysema, chronic pyelonephritis, and chronic cystitis.

[0058] A cardiovascular disease or disorder includes those disorders that can either cause ischemia or are caused by reperfusion of the heart. Examples include, but are not limited to, atherosclerosis, coronary artery disease, granulomatous myocarditis, chronic myocarditis (non-granulomatous), primary hypertrophic cardiomyopathy, peripheral artery disease (PAD), stroke, angina pectoris, myocardial infarction, cardiovascular tissue damage caused by cardiac arrest, cardiovascular tissue damage caused by cardiac bypass, cardiogenic shock, and related conditions that would be known by those of ordinary skill in the art or which involve dysfunction of or tissue damage to the heart or vasculature, especially, but not limited to, tissue damage related to a Reprogramming factor activation. CVS diseases include, but are not limited to, atherosclerosis, granulomatous myocarditis, myocardial infarction, myocardial fibrosis secondary to valvular heart disease, myocardial fibrosis without infarction, primary hypertrophic cardiomyopathy, and chronic myocarditis (non-granulomatous).

[0059] A "Metabolic disease or disorder" refers to a wide range of diseases and disorders of the endocrine system including, for example, insulin resistance, diabetes, obesity, impaired glucose tolerance, high blood cholesterol, hyperglycemia, hyperinsulinemia, dyslipidemia and hyperlipidemia.

Polynucleotide and Oligonucleotide Compositions and Molecules

Targets

[0060] In one instance, the targets comprise nucleic acid sequences of a Reprogramming factor, including without limitation sense and/or antisense noncoding and/or coding sequences associated with a Reprogramming factor.

[0061] In one instance, the targets comprise nucleic acid sequences of SOX2, including without limitation sense and/or antisense noncoding and/or coding sequences associated with SOX2 gene.

[0062] In one instance, the targets comprise nucleic acid sequences of KLF4, including without limitation sense and/or antisense noncoding and/or coding sequences associated with KLF4 gene.

[0063] In one instance, the targets comprise nucleic acid sequences of POU5F1, including without limitation sense and/or antisense noncoding and/or coding sequences associated with POU5F1 gene.

[0064] SRY (sex determining region Y)-box 2, also known as SOX2, is a transcription factor that is essential to maintain self-renewal of undifferentiated embryonic stem cells. This intronless gene encodes a member of the SRY-related HMG-box (SOX) family of transcription factor involved in the regulation of embryonic development and in the determination of cell fate. The encoded protein may act as a transcriptional activator after forming a protein complex with other proteins. Mutations in this gene have been associated with bilateral anophthalmia, a severe form of structural eye

malformation. This gene lies within an intron of another gene called SOX2 overlapping transcript (SOX20T). The ectopic expression of SOX2 may be also related to abnormal differentiation of colorectal cancer cells. Sox2 is one of the key transcription factor required in induced pluripotent stem cells.

[0065] A stem cell is a relatively undifferentiated cell that can be induced to proliferate and that can produce progeny that subsequently differentiate into one or more mature cell types, while also retaining one or more cells with parental developmental potential. In many biological instances, stem cells are also multipotent because they can produce progeny of more than one distinct cell type. Self-renewal is the other classical part of the stem cell definition, and it is essential as used in this document. In theory, self-renewal can occur by either of two major mechanisms. Stem cells may divide asymmetrically, with one daughter retaining the stem state and the other daughter expressing some distinct other specific function and phenotype. Alternatively, some of the stem cells in a population can divide symmetrically into two stems, thus maintaining some stem cells in the population as a whole, while other cells in the population give rise to differentiated progeny only. Formally, it is possible that cells that begin as stem cells might proceed toward a differentiated phenotype, but then "reverse" and re-express the stem cell phenotype.

[0066] Progenitor cells have a cellular phenotype that is more primitive (i.e., is at an earlier step along a developmental pathway or progression than is a fully differentiated cell). Often, progenitor cells also have significant or very high proliferative potential. Progenitor cells may give rise to multiple distinct differentiated cell types or to a single differentiated cell type, depending on the developmental pathway and on the environment in which the cells develop and differentiate. Like stem cells, it is possible that cells that begin as progenitor cells might proceed toward a differentiated phenotype, but then "reverse" and re-express the progenitor cell phenotype.

[0067] In a preferred instance, one or more antisense oligonucleotides bind to a sense and/or antisense polynucleotide of one or more members of the sox family. Examples of the SOX (SRY-related HMG box) family include, for example, Sox-1, Sox-2, Sox-3, Sox-4, Sox6, Sox7, Sox8, Sox9, Sox1~Sox11, Sox-13, Sox14, Sox15, Sox18, Sox20, Sox21, Sox30, Sox32 or the factor Sox-11-D of *Xenopus laevis*, or a functional fragment thereof, and a preferred example includes Sox2.

[0068] Knippel-like factor 4 (KLF4) is a zinc-finger-containing transcription factor, the expression of which is enriched in the postmitotic cells of the intestinal epithelium. KLF4 is a target gene of the tumor suppressor adenomatous polyposis coli (APC). Knippel-like factor 4 (KLF4) is expressed in a wide variety of tissues in humans, including the intestine and the skin, and is important in many different physiologic processes, including development, differentiation, and maintenance of normal tissue homeostasis. KLF4 is a bi-functional transcription factor that can either activate or repress transcription, depending on the target gene, and utilizing different mechanisms. In addition, KLF4 can function as an oncogene or a tumor suppressor depending on the type of cancer involved. In concert with three other transcription factors, KLF4 can reprogram differentiated fibroblasts into a state resembling embryonic stem cells in every possible manner tested so far.

[0069] Recently, it was found that overexpression of KLF4, in combination with three other transcription factors could transform mouse fibroblasts into a state resembling embryonic stem cells (ES cells).

[0070] In preferred instances, , a method of treating diseases or disorders associated with low levels of Knippel-like factor 4 (KLF4) comprise administering to a patient an antisense oligonucleotide which increases Knippel-like factor 4 (KLF4) expression and/or function.

[0071] In accordance with instances of the disclosure, the target nucleic acid molecule is directed to Knippel-like factor 4 (KLF4) and extends to any of the isoforms, receptors, families and the like of Knippel-like factor 4 (KLF4). Synonyms of KLF4 include: Endothelial KruppelLike Zinc Finger Protein (EZF); Gut-Enriched Kruppel-Like Factor (GKLF).

[0072] In preferred instances, the disclosure is contemplated all aspects associated with the molecules described herein and encompasses all peptides, polypeptides, derivatives, variants of the polynucleotide and oligonucleotide sequences of KLF4.

[0073] In another preferred instance, the disclosure comprises antibodies and aptamers which may be generated to KLF4 molecules.

[0074] Octamer binding transcription factor 4 (hereinafter "POU5F1") belongs to the POU (Pit-Oct-Unc) family of transcription factors that bind the octamer motif, a transcription regulatory element found in the promoter and enhancer regions of many genes (Ryan AK, et al. (1997) *Genes Dev*; 11: 1207-1225; Ruvkun G, et al. (1991) *Cell* 64: 475-478). The DNA-binding activity of the POU family members is mediated by the so-called POU domain, a bipartite DNA-binding domain consisting of a 75-amino acid POU-specific domain, a linker region, and a 60-amino-acid POU homeodomain (Herr W, et al., (1995) *Genes Dev* 9: 1679-1693; ExpASy Proteomics Server. Available at <http://au.expasy.org/uniprot/Q01860> Accessed June 7, 2006; Scholer HR, et al. (1990) *Nature* 344:435-9.)

[0075] POU5F1 has been shown to be critical in the induction and maintenance of the pluripotent stem cell state. Downregulation of POU5F1 expression in embryonic stem cells (ESCs) causes them to differentiate and lose their pluripotency. POU5F1 induces and sustains pluripotency in cooperation with other transcription factors including SOX2 and NANOG, and also plays a crucial role in the early mammalian development. POU5F1 is expressed at low levels in some adult stem cell populations. (Boiani M, (2005) et al. *Nat Rev Mol Cell Biol*. 6:872-84.; Boyer LA, et al. (2005) *Cell*.; 122:947-56.)

[0076] The human POU5F1 gene consists of five exons and is located on chromosome 6 in the region of the major histocompatibility complex. The gene encodes two isoforms, A (POU5F1_iA, also known as OCT-3A) and B (POU5F1_iB, also known as OCT-3B) (Takeda J, et al. (1992) *Nucleic Acids Res* 20: 4613-4620).

[0077] In some instances, the POU5F1 antisense oligonucleotides promote stem cell proliferation.

[0078] Induced pluripotent stem (iPS) cells can be obtained from mouse fibroblasts by retrovirus-mediated expression of 4 transcription factors, Oct3/4, Sox2, c-Myc and Klf4 and subsequent selection for Fbx15 expression Selection for NANOG expression instead of Fbx15 produces iPS cells with increased embryonic stem cell-like gene expression and DNA methylation patterns Expression of the transcription factors Oct4, Sox2, c-Myc, and Klf4 reprograms a somatic genome of fibroblasts to an embryonic pluripotent state. Yu et al. OCT4, SOX2, NANOG, LIN28 are sufficient to reprogram human somatic cells to pluripotent stem cells. The induced pluripotent human stem cells have

normal karyotypes, telomerase activity, cell surface markers and typical gene expression of the human ES cells, and can differentiate into derivatives of all 3 primary germ layers.

[0079] In some instances, antisense oligonucleotides are used to prevent or treat diseases or disorders associated with Reprogramming factor family members. Exemplary Reprogramming factor mediated diseases and disorders which can be treated with cell/tissues regenerated from stem cells obtained using the antisense compounds comprise: abnormal Reprogramming factor expression and/or function, cancer, a renal disease, a cardiovascular disease or disorder, inflammation, a neurological disease or disorder, scarring, ophthalmic diseases or disorders (e.g., microphthalmia, coloboma, myopia, anophthalmia/microphthalmia - esophageal atresia syndrome, etc.), Septooptic dysplasia, an autoimmune disease or disorder, diabetes, atherosclerosis, an immunodeficiency disease, a disease caused by infectious agent (e.g., including, viral, bacterial, fungal, protozoan etc.), a genetic disease (e.g. Duchenne muscular dystrophy), a metabolic disease or disorder (e.g. diabetes, obesity, metabolic syndrome, lysosomal disease etc.), trauma (e.g. spinal cord injury, burns, etc), ischemia, a vascular disease, a hepatic disease or disorder, Psoriasis, a disease, a disease or disorder requiring transplantation of cells, tissues, and organs, disorder or condition requiring stem cell therapy and a congenital disease or disorder.

[0080] In some instances, the present disclosure provides a method for preparing an induced pluripotent stem cell by reprogramming of a somatic cell, which comprises the step of contacting the aforementioned reprogramming factor with the somatic cell.

[0081] There is also provided a method, which comprises the step of adding the aforementioned reprogramming factor to a culture of the somatic cell; the aforementioned method, which comprises the step of introducing a gene encoding the aforementioned reprogramming factor into the somatic cell; the aforementioned method, which comprises the step of introducing said gene into the somatic cell by using a recombinant vector containing at least one kind of gene encoding the aforementioned reprogramming factor; and the aforementioned method, wherein a somatic cell isolated from a patient is used as the somatic cell.

[0082] In another aspect, the present disclosure provides an induced pluripotent stem cell obtained by the aforementioned method. The present disclosure also provides a somatic cell derived by inducing differentiation of the aforementioned induced pluripotent stem cell.

[0083] The present disclosure further provides a method for stem cell therapy, which comprises the step of transplanting a somatic cell, wherein said cell is obtained by inducing differentiation of an induced pluripotent stem cell obtained by the aforementioned method using a somatic cell isolated and collected from a patient, into said patient. Several kinds of, preferably approximately 200 kinds of iPS cells prepared from somatic cells derived from healthy humans can be stored in an iPS cell bank as a library of iPS cells, and one kind or more kinds of the iPS cells in the library can be used for preparation of somatic cells, tissues, or organs that are free of rejection by a patient to be subjected to stem cell therapy.

[0084] The present disclosure further provides a method for evaluating a physiological function or toxicity of a compound, a medicament, a poison or the like by using various cells obtained by inducing differentiation of an induced pluripotent stem cell obtained by the aforementioned method.

[0085] The present disclosure also provides a method for improving ability of differentiation and/or growth of a cell, which comprises the step of contacting the aforementioned reprogramming factor with the cell, and further provides a cell obtained by the aforementioned method, and a somatic cell derived by inducing differentiation of a cell obtained by the aforementioned method.

[0086] By using the reprogramming factor provided by the present disclosure, reprogramming of a differentiated cell nucleus can be conveniently and highly reproducibly induced without using embryos or ES cells, and an induced pluripotent stem cell, as an undifferentiated cell having differentiation ability, pluripotency, and growth ability similar to those of ES cells, can be established. For example, an induced pluripotent stem cell having high growth ability and differentiation pluripotency can be prepared from a patient's own somatic cell by using the reprogramming factor of the present disclosure. Cells obtainable by differentiating said cell (for example, cardiac muscle cells, insulin producing cells, nerve cells and the like) are extremely useful, because they can be utilized for stem cell transplantation therapies for a variety of diseases such as cardiac insufficiency, insulin dependent diabetes mellitus, Parkinson's disease and spinal cord injury, thereby the ethical problem concerning the use of human embryo and rejection after transplantation can be avoided. Further, various cells obtainable by differentiating the induced pluripotent stem cell (for example, cardiac muscle cells, hepatic cells and the like) are highly useful as systems for evaluating efficacy or toxicity of compounds, medicaments, poisons and the like.

[0087] In some instances, the antisense oligonucleotides of the present disclosure are used in conjunction with other transcription factors required to induce pluripotent stem cells from a given adult tissue. These cells can be further introduced into a lesion and allowed to differentiate into a required cell type in vitro or in vivo.

[0088] In some instances, the antisense oligonucleotides are administered to a cell or patient to maintain pluripotency and/or self-renewing characteristics and is suitable for any stem cell, progenitor cell, or any cell derived. As an illustrative example, any pluripotent human ESC or a respective cell line may be used in the method. Means of deriving a population of such cells are well established in the art (cf e.g. Thomson, J. A. et al. [1998] Science 282, 1145-1147 or Cowan, C. A. et al. [2004] N. Engl. J. Med. 350, 1353-1356). Furthermore, 71 independent human ESC lines are for example known to exist, of which 11 cell lines are available for research purposes, such as GE01, GE09, BG01, BG02, TE06 or WA09. Adult stem cells may for instance be isolated from blood from the placenta and umbilical cord left over after birth, or from myofibers, to which they are associated as so called "satellite cells"

[0089] Where the method is intended to be used for a progenitor cell, i.e. a cell giving rise to mature somatic cells, any progenitor cell may be used in this method of the disclosure. Examples of suitable progenitor cells include, but are not limited to, neuronal progenitor cells, endothelial progenitor cells, erythroid progenitor cells, cardiac progenitor cells, oligodendrocyte progenitor cells, retinal progenitor cells, or hematopoietic progenitor cells.

[0090] Progenitor cells, such as endothelial progenitor cells obtainable from peripheral blood, have for example been found to possess high expression levels of Nanog and Oct-4. CNS progenitor cells such as retinal progenitor cells have been reported to express high levels of Sox2.

[0091] As indicated above, the antisense oligonucleotides modulate gene expression and/or

function in a cell. In some instances an endogenous SOX gene is functionally active. In some of these instances the respective cell is a stem or a progenitor cell. Examples of stem cells that may be used in the method of the present disclosure include, but are not limited to, embryonic stem cells, trophoblast stem cells and extraembryonic stem cells. In some instances of the methods of the disclosure an ESC (embryonic stem cell) may be used. ESC of human origin (i.e. a human ESC) are disclosed, but not encompassed by the present invention. In other instances, the cell is a progenitor cell. In yet other instances the cell is a cancer cell. An illustrative example of a cancer cell is teratoma cancer cell, such as for example F9, NTERA2, C3H, TES-1, 1246 (including 1246-3A), SuSa (including SuSa/DXRIO and SKOV-3/DXRIO), AT805 (including ATDC5), HTST, HGRT, PC (e.g. PCC3/A/1) or GCT27. Two further illustrative example of a cancer cell are a HeLa cell and an MCF-7 cell. In some instances the cell is a hybrid cell of a stem cell and a somatic cell. In some of these instances a cell of an established eukaryotic cell line is selected, such as for instance HEK, COS, CHO, CRE, MT4, DE (duck embryo), QF (quail fibrosarcoma), NSO, BHK, Sf9, PC12, or High 5. An illustrative example is a HEK 293T cell.

[0092] In one instance, KLF4 antisense oligonucleotides reprogram stem cells and modulate self-renewal. Stem cells are contacted with one or more KLF4 antisense oligonucleotides. For example, overexpression of KLF4 in ES cells inhibited differentiation into erythroid progenitors, and increased their capacity to generate secondary embryoid bodies. In concert with Oct3/4 and Sox2, KLF4 activates expression of Lefty], a gene expressed in ES cells, but lost during differentiation process.

[0093] In another preferred instance, one or more antisense KLF4 oligonucleotides are administered to one or more stem cells. The cells can be stem cells isolated from the bone marrow as a progenitor cell, or cells obtained from any other source, such as for example, ATCC.

[0094] "Bone marrow derived progenitor cell" (BMDC) or "bone marrow derived stem cell" refers to a primitive stem cell with the machinery for self-renewal constitutively active. Included in this definition are stem cells that are totipotent, pluripotent and precursors. A "precursor cell" can be any cell in a cell differentiation pathway that is capable of differentiating into a more mature cell. As such, the term "precursor cell population" refers to a group of cells capable of developing into a more mature cell. A precursor cell population can comprise cells that are totipotent, cells that are pluripotent and cells that are stem cell lineage restricted (i.e. cells capable of developing into less than all hematopoietic lineages, or into, for example, only cells of erythroid lineage). As used herein, the term "totipotent cell" refers to a cell capable of developing into all lineages of cells. Similarly, the term "totipotent population of cells" refers to a composition of cells capable of developing into all lineages of cells. Also as used herein, the term "pluripotent cell" refers to a cell capable of developing into a variety (albeit not all) lineages and are at least able to develop into all hematopoietic lineages (e.g., lymphoid, erythroid, and thrombocytic lineages). Bone marrow derived stem cells contain two well-characterized types of stem cells. Mesenchymal stem cells (MSC) normally form chondrocytes and osteoblasts. Hematopoietic stem cells (HSC) are of mesodermal origin that normally give rise to cells of the blood and immune system (e.g., erythroid, granulocyte/macrophage, megakaryocyte and lymphoid lineages). In addition, hematopoietic stem cells also have been shown to have the potential to differentiate into the cells of the liver (including hepatocytes, bile duct cells), lung, kidney (e.g., renal tubular epithelial cells and renal parenchyma), gastrointestinal tract, skeletal muscle fibers, astrocytes of the CNS, Purkinje neurons, cardiac muscle (e.g., cardiomyocytes), endothelium and skin.

[0095] In another preferred instance, the antisense oligonucleotides modulate the expression and/or function of KLF4 in patients suffering from or at risk of developing diseases or disorders associated with KLF4. Any number of diseases or conditions can be treated by modulation of KLF4, including stem cell renewal and re-programming.

[0096] In other preferred instances, the oligomeric compounds of the present disclosure also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenosine, variants may be produced which contain thymidine, guanosine or cytidine at this position. This may be done at any of the positions of the antisense compound. These compounds are then tested using the methods described herein to determine their ability to inhibit expression of a target nucleic acid. Antisense oligonucleotides which modulate expression, activity or any functions of KLF4 would also modulate factors in cell pathways that are associated with KLF4 activity, function or expression.

[0097] Examples of factors which are modulated by KLF4 are shown in the table below:

Factor/condition Activation targets	Factor/condition Repression targets
1200015N20Rik	Bax
A33 antigen	CD11d
B2R	Cyclin B1
CYP1A1	Cyclin D1
Cytokeratin 4	Cyclin E
EBVED-L2	Fgf5
hSMVT	Histidine decarboxylase
1AP	KLF2
iNOS	Laminin a1
Keratin 4	Nes
Keratin 19	Ornithine decarboxylase
KLF4	p53
Laminin-a 3A	PAI-1
Laminin-y 1	SM22a
Lefty1	SM a-actin
Nanog	Sp1
Oct4	
p21Cip1	
p27Kip1	
p57Kip2	
PKG-Ia	
Rb	
Sox2	
SPRR1A	
SPRR2A	

Factor/condition Activation targets	Factor/condition Repression targets
Tbx3	
u-PAR	

[0098] In one instance, the oligonucleotides are specific for polynucleotides of a Reprogramming factor, which includes, without limitation noncoding regions. The Reprogramming factor targets comprise variants of a Reprogramming factor; mutants of a Reprogramming factor, including SNPs; noncoding sequences of a Reprogramming factor; alleles, fragments and the like. Preferably the oligonucleotide is an antisense RNA molecule.

[0099] In accordance with instances of the disclosure, the target nucleic acid molecule is not limited to a Reprogramming factor polynucleotides alone but extends to any of the isoforms, receptors, homologs, non-coding regions and the like of a Reprogramming factor.

[0100] In another instance, an oligonucleotide targets a natural antisense sequence (natural antisense to the coding and non-coding regions) of a Reprogramming factor targets, including, without limitation, variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense RNA or DNA molecule.

[0101] In another instance, the oligomeric compounds of the present disclosure also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenine, variants may be produced which contain thymidine, guanosine, cytidine or other natural or unnatural nucleotides at this position. This may be done at any of the positions of the antisense compound.

[0102] In some instances, homology, sequence identity or complementarity, between the antisense compound and target is from about 50% to about 60%. In some instances, homology, sequence identity or complementarity, is from about 60% to about 70%. In some instances, homology, sequence identity or complementarity, is from about 70% to about 80%. In some instances, homology, sequence identity or complementarity, is from about 80% to about 90%. In some instances, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

[0103] An antisense compound is specifically hybridizable when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a loss of activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which specific binding is desired. Such conditions include, i.e., physiological conditions in the case of in vivo assays or therapeutic treatment, and conditions in which assays are performed in the case of in vitro assays.

[0104] An antisense compound, whether DNA, RNA, chimeric, substituted etc, is specifically hybridizable when binding of the compound to the target DNA or RNA molecule interferes with the normal function of the target DNA or RNA to cause a loss of utility, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target sequences under conditions in which specific binding is desired, i.e., under physiological conditions in the case

of in vivo assays or therapeutic treatment, and in the case of in vitro assays, under conditions in which the assays are performed.

[0105] In another preferred instance, targeting of a Reprogramming factor including without limitation, antisense sequences which are identified and expanded, using for example, PCR, hybridization etc., one or more of the sequences set forth as SEQ ID NO: 4 to 6, and the like, modulate the expression or function of a Reprogramming factor. In one instance, expression or function is up-regulated as compared to a control. In another preferred instance, expression or function is down-regulated as compared to a control.

[0106] In another preferred instance, oligonucleotides comprise nucleic acid sequences set forth as SEQ ID NOS: 7 to 17 including antisense sequences which are identified and expanded, using for example, PCR, hybridization etc. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like. Examples of modified bonds or internucleotide linkages comprise phosphorothioate, phosphorodithioate or the like. In another preferred instance, the nucleotides comprise a phosphorus derivative. The phosphorus derivative (or modified phosphate group) which may be attached to the sugar or sugar analog moiety in the modified oligonucleotides of the present disclosure may be a monophosphate, diphosphate, triphosphate, alkylphosphate, alkanephosphate, phosphorothioate and the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and oligonucleotides, *per se*, is also known and need not be described here.

[0107] The specificity and sensitivity of antisense is also harnessed by those of skill in the art for therapeutic uses. Antisense oligonucleotides have been employed as therapeutic moieties in the treatment of disease states in animals and man. Antisense oligonucleotides have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that oligonucleotides can be useful therapeutic modalities that can be configured to be useful in treatment regimes for treatment of cells, tissues and animals, especially humans.

[0108] In instances of the present disclosure oligomeric antisense compounds, particularly oligonucleotides, bind to target nucleic acid molecules and modulate the expression and/or function of molecules encoded by a target gene. The functions of DNA to be interfered comprise, for example, replication and transcription. The functions of RNA to be interfered comprise all vital functions such as, for example, translocation of the RNA to the site of protein translation, translation of protein from the RNA, splicing of the RNA to yield one or more mRNA species, and catalytic activity which may be engaged in or facilitated by the RNA. The functions may be up-regulated or inhibited depending on the functions desired.

[0109] The antisense compounds, include, antisense oligomeric compounds, antisense oligonucleotides, external guide sequence (EGS) oligonucleotides, alternate splicers, primers, probes, and other oligomeric compounds that hybridize to at least a portion of the target nucleic acid. As such, these compounds may be introduced in the form of single-stranded, double-stranded, partially single-stranded, or circular oligomeric compounds.

[0110] Targeting an antisense compound to a particular nucleic acid molecule, in the context of this disclosure, can be a multistep process. The process usually begins with the identification of a target nucleic acid whose function is to be modulated. This target nucleic acid may be, for example, a

cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a nucleic acid molecule from an infectious agent. In the present disclosure, the target nucleic acid encodes a Reprogramming factor.

[0111] The targeting process usually also includes determination of at least one target region, segment, or site within the target nucleic acid for the antisense interaction to occur such that the desired effect, e.g., modulation of expression, will result. Within the context of the present disclosure, the term "region" is defined as a portion of the target nucleic acid having at least one identifiable structure, function, or characteristic. Within regions of target nucleic acids are segments. "Segments" are defined as smaller or sub-portions of regions within a target nucleic acid. "Sites," as used in the present disclosure, are defined as positions within a target nucleic acid.

[0112] In one instance, the antisense oligonucleotides bind to the natural antisense sequences of a Reprogramming factor and modulate the expression and/or function of a Reprogramming factor (SEQ ID NO: 1 to 3). Examples of antisense sequences include SEQ ID NOS: 4 to 17.

[0113] In another instance, the antisense oligonucleotides bind to one or more segments of a Reprogramming factor polynucleotide and modulate the expression and/or function of a Reprogramming factor. The segments comprise at least five consecutive nucleotides of a Reprogramming factor sense or antisense polynucleotides.

[0114] In another instance, the antisense oligonucleotides are specific for natural antisense sequences of a Reprogramming factor wherein binding of the oligonucleotides to the natural antisense sequences of a Reprogramming factor modulate expression and/or function of a Reprogramming factor.

[0115] In another instance, oligonucleotide compounds comprise sequences set forth as SEQ ID NOS: 7 to 17, antisense sequences which are identified and expanded, using for example, PCR, hybridization etc. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like. Examples of modified bonds or internucleotide linkages comprise phosphorothioate, phosphorodithioate or the like. In another preferred instance, the nucleotides comprise a phosphorus derivative. The phosphorus derivative (or modified phosphate group) which may be attached to the sugar or sugar analog moiety in the modified oligonucleotides of the present disclosure may be a monophosphate, diphosphate, triphosphate, alkylphosphate, alkanephosphate, phosphorothioate and the like. The preparation of the above-noted phosphate analogs, and their incorporation into nucleotides, modified nucleotides and oligonucleotides, per se, is also known and need not be described here.

[0116] Since, as is known in the art, the translation initiation codon is typically 5'-AUG (in transcribed mRNA molecules; 5'-ATG in the corresponding DNA molecule), the translation initiation codon is also referred to as the "AUG codon," the "start codon" or the "AUG start codon". A minority of genes has a translation initiation codon having the RNA sequence 5'-GUG, 5'-UUG or 5'-CUG; and 5'-AUA, 5'-ACG and 5'-CUG have been shown to function in vivo. Thus, the terms "translation initiation codon" and "start codon" can encompass many codon sequences, even though the initiator amino acid in each instance is typically methionine (in eukaryotes) or formylmethionine (in prokaryotes). Eukaryotic and prokaryotic genes may have two or more alternative start codons, any one of which may be preferentially utilized for translation initiation in a particular cell type or tissue, or under a particular

set of conditions. In the context of the disclosure, "start codon" and "translation initiation codon" refer to the codon or codons that are used in vivo to initiate translation of an mRNA transcribed from a gene encoding a Reprogramming factor, regardless of the sequence(s) of such codons. A translation termination codon (or "stop codon") of a gene may have one of three sequences, i.e., 5'-UAA, 5'-UAG and 5'-UGA (the corresponding DNA sequences are 5'-TAA, 5'-TAG and 5'-TGA, respectively).

[0117] The terms "start codon region" and "translation initiation codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation initiation codon. Similarly, the terms "stop codon region" and "translation termination codon region" refer to a portion of such an mRNA or gene that encompasses from about 25 to about 50 contiguous nucleotides in either direction (i.e., 5' or 3') from a translation termination codon. Consequently, the "start codon region" (or "translation initiation codon region") and the "stop codon region" (or "translation termination codon region") are all regions that may be targeted effectively with the antisense compounds of the present disclosure.

[0118] The open reading frame (ORF) or "coding region," which is known in the art to refer to the region between the translation initiation codon and the translation termination codon, is also a region which may be targeted effectively. Within the context of the present disclosure, a targeted region is the intragenic region encompassing the translation initiation or termination codon of the open reading frame (ORF) of a gene.

[0119] Another target region includes the 5' untranslated region (5'UTR), known in the art to refer to the portion of an mRNA in the 5' direction from the translation initiation codon, and thus including nucleotides between the 5' cap site and the translation initiation codon of an mRNA (or corresponding nucleotides on the gene). Still another target region includes the 3' untranslated region (3'UTR), known in the art to refer to the portion of an mRNA in the 3' direction from the translation termination codon, and thus including nucleotides between the translation termination codon and 3' end of an mRNA (or corresponding nucleotides on the gene). The 5' cap site of an mRNA comprises an N7-methylated guanosine residue joined to the 5'-most residue of the mRNA via a 5'-5' triphosphate linkage. The 5' cap region of an mRNA is considered to include the 5' cap structure itself as well as the first 50 nucleotides adjacent to the cap site. Another target region for this disclosure is the 5' cap region.

[0120] Although some eukaryotic mRNA transcripts are directly translated, many contain one or more regions, known as "introns," which are excised from a transcript before it is translated. The remaining (and therefore translated) regions are known as "exons" and are spliced together to form a continuous mRNA sequence. In one instance, targeting splice sites, i.e., intron-exon junctions or exon-intron junctions, is particularly useful in situations where aberrant splicing is implicated in disease, or where an overproduction of a particular splice product is implicated in disease. An aberrant fusion junction due to rearrangement or deletion is another instance of a target site. mRNA transcripts produced via the process of splicing of two (or more) mRNAs from different gene sources are known as "fusion transcripts". Introns can be effectively targeted using antisense compounds targeted to, for example, DNA or pre-mRNA.

[0121] In another instance, the antisense oligonucleotides bind to coding and/or non-coding regions of a target polynucleotide and modulate the expression and/or function of the target molecule.

[0122] In another instance, the antisense oligonucleotides bind to natural antisense polynucleotides and modulate the expression and/or function of the target molecule.

[0123] In another instance, the antisense oligonucleotides bind to sense polynucleotides and modulate the expression and/or function of the target molecule.

[0124] Alternative RNA transcripts can be produced from the same genomic region of DNA. These alternative transcripts are generally known as "variants". More specifically, "pre-mRNA variants" are transcripts produced from the same genomic DNA that differ from other transcripts produced from the same genomic DNA in either their start or stop position and contain both intronic and exonic sequence.

[0125] Upon excision of one or more exon or intron regions, or portions thereof during splicing, pre-mRNA variants produce smaller "mRNA variants". Consequently, mRNA variants are processed pre-mRNA variants and each unique pre-mRNA variant must always produce a unique mRNA variant as a result of splicing. These mRNA variants are also known as "alternative splice variants". If no splicing of the pre-mRNA variant occurs then the pre-mRNA variant is identical to the mRNA variant.

[0126] Variants can be produced through the use of alternative signals to start or stop transcription. Pre-mRNAs and mRNAs can possess more than one start codon or stop codon. Variants that originate from a pre-mRNA or mRNA that use alternative start codons are known as "alternative start variants" of that pre-mRNA or mRNA. Those transcripts that use an alternative stop codon are known as "alternative stop variants" of that pre-mRNA or mRNA. One specific type of alternative stop variant is the "polyA variant" in which the multiple transcripts produced result from the alternative selection of one of the "polyA stop signals" by the transcription machinery, thereby producing transcripts that terminate at unique polyA sites. Within the context of the disclosure, the types of variants described herein are also instances of target nucleic acids.

[0127] The locations on the target nucleic acid to which the antisense compounds hybridize are defined as at least a 5-nucleotide long portion of a target region to which an active antisense compound is targeted.

[0128] While the specific sequences of certain exemplary target segments are set forth herein, one of skill in the art will recognize that these serve to illustrate and describe particular instances within the scope of the present disclosure. Additional target segments are readily identifiable by one having ordinary skill in the art in view of this disclosure.

[0129] Target segments 5-100 nucleotides in length comprising a stretch of at least five (5) consecutive nucleotides selected from within the illustrative preferred target segments are considered to be suitable for targeting as well.

[0130] Target segments can include DNA or RNA sequences that comprise at least the 5 consecutive nucleotides from the 5'-terminus of one of the illustrative preferred target segments (the remaining nucleotides being a consecutive stretch of the same DNA or RNA beginning immediately upstream of the 5'-terminus of the target segment and continuing until the DNA or RNA contains about 5 to about 100 nucleotides). Similarly preferred target segments are represented by DNA or RNA sequences that comprise at least the 5 consecutive nucleotides from the 3'-terminus of one of

the illustrative preferred target segments (the remaining nucleotides being a consecutive stretch of the same DNA or RNA beginning immediately downstream of the 3'-terminus of the target segment and continuing until the DNA or RNA contains about 5 to about 100 nucleotides). One having skill in the art armed with the target segments illustrated herein will be able, without undue experimentation, to identify further preferred target segments.

[0131] Once one or more target regions, segments or sites have been identified, antisense compounds are chosen which are sufficiently complementary to the target, i.e., hybridize sufficiently well and with sufficient specificity, to give the desired effect.

[0132] In instances of the disclosure the oligonucleotides bind to an antisense strand of a particular target. The oligonucleotides are at least 5 nucleotides in length and can be synthesized so each oligonucleotide targets overlapping sequences such that oligonucleotides are synthesized to cover the entire length of the target polynucleotide. The targets also include coding as well as non coding regions.

[0133] In one instance, , specific nucleic acids are targeted by antisense oligonucleotides. Targeting an antisense compound to a particular nucleic acid, is a multistep process. The process usually begins with the identification of a nucleic acid sequence whose function is to be modulated. This may be, for example, a cellular gene (or mRNA transcribed from the gene) whose expression is associated with a particular disorder or disease state, or a non coding polynucleotide such as for example, non coding RNA (ncRNA).

[0134] RNAs can be classified into (1) messenger RNAs (mRNAs), which are translated into proteins, and (2) non-protein-coding RNAs (ncRNAs). ncRNAs comprise microRNAs, antisense transcripts and other Transcriptional Units (TU) containing a high density of stop codons and lacking any extensive "Open Reading Frame". Many ncRNAs appear to start from initiation sites in 3' untranslated regions (3'UTRs) of protein-coding loci. ncRNAs are often rare and at least half of the ncRNAs that have been sequenced by the FANTOM consortium seem not to be polyadenylated. Most researchers have for obvious reasons focused on polyadenylated mRNAs that are processed and exported to the cytoplasm. Recently, it was shown that the set of non-polyadenylated nuclear RNAs may be very large, and that many such transcripts arise from intergenic regions. The mechanism by which ncRNAs may regulate gene expression is by base pairing with target transcripts. The RNAs that function by base pairing can be grouped into (1) cis encoded RNAs that are encoded at the same genetic location, but on the opposite strand to the RNAs they act upon and therefore display perfect complementarity to their target, and (2) trans-encoded RNAs that are encoded at a chromosomal location distinct from the RNAs they act upon and generally do not exhibit perfect base-pairing potential with their targets.

[0135] Without wishing to be bound by theory, perturbation of an antisense polynucleotide by the antisense oligonucleotides described herein can alter the expression of the corresponding sense messenger RNAs. However, this regulation can either be discordant (antisense knockdown results in messenger RNA elevation) or concordant (antisense knockdown results in concomitant messenger RNA reduction). In these cases, antisense oligonucleotides can be targeted to overlapping or non-overlapping parts of the antisense transcript resulting in its knockdown or sequestration. Coding as well as non-coding antisense can be targeted in an identical manner and that either category is capable of regulating the corresponding sense transcripts - either in a concordant or discordant

manner. The strategies that are employed in identifying new oligonucleotides for use against a target can be based on the knockdown of antisense RNA transcripts by antisense oligonucleotides or any other means of modulating the desired target

[0136] Several nucleotide sequences are relevant to the disclosure, including:

The following nucleotide sequences are relevant to the disclosure:

SEQ ID NO: 1: Homo sapiens SRY (sex determining region Y)-box 2 (SOX2), mRNA (NCBI Accession No.: NM_003106.2).

SEQ ID NO: 2: Homo sapiens Kruppel-like factor 4 (gut) (KLF4), mRNA (NCBI Accession No.: NM_004235.4).

SEQ ID NO: 3: Homo sapiens POU class 5 homeobox 1 (POU5F1), transcript variant 1, mRNA (NCBI Accession No.: NM_002701.4).

SEQ ID NO: 4: Human Natural SOX2 antisense sequence (AI885646)

SEQ ID NO: 5: Mouse Natural KLF4 antisense sequence (DB461753)

SEQ ID NO: 6: Human Natural POU5F1 antisense sequence (AI926793)

SOX2 antisense oligonucleotides, SEQ ID NOs: 7 and 8. 'r' indicates RNA and * indicates phosphothioate bond.

KLF4 antisense oligonucleotides, SEQ ID NOs: 9 to 12. * indicates phosphothioate bond.

POU5F1 antisense oligonucleotides, SEQ ID NOs: 13 to 17. * indicates phosphothioate bond.

Figure 5 shows the SOX2 sense oligonucleotides, SEQ ID NOs: 18 and 19. The sense oligonucleotides SEQ ID NO: 18 and 19 are the reverse complements of the antisense oligonucleotides SEQ ID NO: 7 and 8 respectively. 'r' indicates RNA.

Strategy 1: In the case of discordant regulation, knocking down the antisense transcript elevates the expression of the conventional (sense) gene. Should that latter gene encode for a known or putative drug target, then knockdown of its antisense counterpart could conceivably mimic the action of a receptor agonist or an enzyme stimulant.

Strategy 2: In the case of concordant regulation, one could concomitantly knock down both antisense and sense transcripts and thereby achieve synergistic reduction of the conventional (sense) gene expression. If, for example, an antisense oligonucleotide is used to achieve knockdown, then this strategy can be used to apply one antisense oligonucleotide targeted to the sense transcript and another antisense oligonucleotide to the corresponding antisense transcript, or a single energetically symmetric antisense oligonucleotide that simultaneously targets overlapping sense and antisense transcripts.

[0137] According to the present disclosure, antisense compounds include antisense oligonucleotides, ribozymes, external guide sequence (EGS) oligonucleotides, siRNA compounds,

single- or double-stranded RNA interference (RNAi) compounds such as siRNA compounds, and other oligomeric compounds which hybridize to at least a portion of the target nucleic acid and modulate its function. As such, they may be DNA, RNA, DNA-like, RNA-like, or mixtures thereof, or may be mimetics of one or more of these. These compounds may be single-stranded, doublestranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges, mismatches or loops. Antisense compounds are routinely prepared linearly but can be joined or otherwise prepared to be circular and/or branched. Antisense compounds can include constructs such as, for example, two strands hybridized to form a wholly or partially double-stranded compound or a single strand with sufficient self-complementarity to allow for hybridization and formation of a fully or partially double-stranded compound. The two strands can be linked internally leaving free 3' or 5' termini or can be linked to form a continuous hairpin structure or loop. The hairpin structure may contain an overhang on either the 5' or 3' terminus producing an extension of single stranded character. The double stranded compounds optionally can include overhangs on the ends. Further modifications can include conjugate groups attached to one of the termini, selected nucleotide positions, sugar positions or to one of the internucleoside linkages. Alternatively, the two strands can be linked via a non-nucleic acid moiety or linker group. When formed from only one strand, dsRNA can take the form of a self-complementary hairpin-type molecule that doubles back on itself to form a duplex. Thus, the dsRNAs can be fully or partially double stranded. Specific modulation of gene expression can be achieved by stable expression of dsRNA hairpins in transgenic cell lines, however, in some instances, the gene expression or function is up regulated. When formed from two strands, or a single strand that takes the form of a self-complementary hairpin-type molecule doubled back on itself to form a duplex, the two strands (or duplex-forming regions of a single strand) are complementary RNA strands that base pair in Watson-Crick fashion.

[0138] Once introduced to a system, the compounds of the disclosure may elicit the action of one or more enzymes or structural proteins to effect cleavage or other modification of the target nucleic acid or may work via occupancy-based mechanisms. In general, nucleic acids (including oligonucleotides) may be described as "DNA-like" (i.e., generally having one or more 2'-deoxy sugars and, generally, T rather than U bases) or "RNA-like" (i.e., generally having one or more 2'-hydroxyl or 2'-modified sugars and, generally U rather than T bases). Nucleic acid helices can adopt more than one type of structure, most commonly the A- and B-forms. It is believed that, in general, oligonucleotides which have B-form-like structure are "DNA-like" and those which have A-formlike structure are "RNA-like." In some (chimeric) instances, an antisense compound may contain both A- and B-form regions.

[0139] In another instance, the desired oligonucleotides or antisense compounds, comprise at least one of: antisense RNA, antisense DNA, chimeric antisense oligonucleotides, antisense oligonucleotides comprising modified linkages, interference RNA (RNAi), short interfering RNA (siRNA); a micro, interfering RNA (miRNA); a small, temporal RNA (stRNA); or a short, hairpin RNA (shRNA); small RNA-induced gene activation (RNAa); small activating RNAs (saRNAs), or combinations thereof.

[0140] dsRNA can also activate gene expression, a mechanism that has been termed "small RNA-induced gene activation" or RNAa. dsRNAs targeting gene promoters induce potent transcriptional activation of associated genes. RNAa was demonstrated in human cells using synthetic dsRNAs, termed "small activating RNAs" (saRNAs).

[0141] Small double-stranded RNA (dsRNA), such as small interfering RNA (siRNA) and microRNA (miRNA), have been found to be the trigger of an evolutionary conserved mechanism known as RNA interference (RNAi). RNAi invariably leads to gene silencing. However, in instances described in detail in the examples section which follows, oligonucleotides are shown to increase the expression and/or function of the Reprogramming factor polynucleotides and encoded products thereof. dsRNAs may also act as small activating RNAs (saRNA). Without wishing to be bound by theory, by targeting sequences in gene promoters, saRNAs would induce target gene expression in a phenomenon referred to as dsRNA-induced transcriptional activation (RNAa).

[0142] In a further instance, the "preferred target segments" identified herein may be employed in a screen for additional compounds that modulate the expression of a Reprogramming factor polynucleotide. "Modulators" are those compounds that decrease or increase the expression of a nucleic acid molecule encoding a Reprogramming factor and which comprise at least a 5-nucleotide portion that is complementary to a preferred target segment. The screening method comprises the steps of contacting a preferred target segment of a nucleic acid molecule encoding sense or natural antisense polynucleotides of a Reprogramming factor with one or more candidate modulators, and selecting for one or more candidate modulators which decrease or increase the expression of a nucleic acid molecule encoding a Reprogramming factor polynucleotide, e.g. SEQ ID NOS: 7 to 17. Once it is shown that the candidate modulator or modulators are capable of modulating (e.g. either decreasing or increasing) the expression of a nucleic acid molecule encoding a Reprogramming factor polynucleotide, the modulator may then be employed in further investigative studies of the function of a Reprogramming factor polynucleotide, or for use as a research, diagnostic, or therapeutic agent in accordance with the present disclosure.

[0143] Targeting the natural antisense sequence modulates the function of the target gene. For example, the Reprogramming factor (e.g. accession numbers NM_003106, NM_004235, NM_002701). In a preferred instance, the target is an antisense polynucleotide of the Reprogramming factor. In a preferred instance, an antisense oligonucleotide targets sense and/or natural antisense sequences of a Reprogramming factor polynucleotide (e.g. accession numbers NM_003106, NM_004235, NM_002701), variants, alleles, isoforms, homologs, mutants, derivatives, fragments and complementary sequences thereto. Preferably the oligonucleotide is an antisense molecule and the targets include coding and noncoding regions of antisense and/or sense Reprogramming factor polynucleotides.

[0144] The preferred target segments of the present disclosure may be also be combined with their respective complementary antisense compounds of the present disclosure to form stabilized double-stranded (duplexed) oligonucleotides.

[0145] Such double stranded oligonucleotide moieties have been shown in the art to modulate target expression and regulate translation as well as RNA processing via an antisense mechanism. Moreover, the double-stranded moieties may be subject to chemical modifications. For example, such double-stranded moieties have been shown to inhibit the target by the classical hybridization of antisense strand of the duplex to the target, thereby triggering enzymatic degradation of the target.

[0146] In a preferred instance, an antisense oligonucleotide targets Reprogramming factor polynucleotides (e.g. accession numbers NM_003106, NM_004235, NM_002701), variants, alleles, isoforms, homologs, mutants, derivatives, fragments and complementary sequences thereto.

Preferably the oligonucleotide is an antisense molecule.

[0147] In accordance with instances of the disclosure, the target nucleic acid molecule is not limited to Reprogramming factor alone but extends to any of the isoforms, receptors, homologs and the like of a Reprogramming factor molecule.

[0148] In another instance, an oligonucleotide targets a natural antisense sequence of a Reprogramming factor polynucleotide, for example, polynucleotides set forth as SEQ ID NO: 4 to 6, and any variants, alleles, homologs, mutants, derivatives, fragments and complementary sequences thereto. Examples of antisense oligonucleotides are set forth as SEQ ID NOS: 7 to 17.

[0149] In one instance, the oligonucleotides are complementary to or bind to nucleic acid sequences of a Reprogramming factor antisense, including without limitation noncoding sense and/or antisense sequences associated with a Reprogramming factor polynucleotide and modulate expression and/or function of a Reprogramming factor molecule.

[0150] In another instance, the oligonucleotides are complementary to or bind to nucleic acid sequences of a Reprogramming factor natural antisense, set forth as SEQ ID NO: 4 to 6 and modulate expression and/or function of a Reprogramming factor molecule.

[0151] In a instance, oligonucleotides comprise sequences of at least 5 consecutive nucleotides of SEQ ID NOS: 7 to 17 and modulate expression and/or function of a Reprogramming factor molecule.

[0152] The polynucleotide targets comprise Reprogramming factor, including family members thereof, variants of a Reprogramming factor; mutants of a Reprogramming factor, including SNPs; noncoding sequences of a Reprogramming factor; alleles of a Reprogramming factor; species variants, fragments and the like. Preferably the oligonucleotide is an antisense molecule.

[0153] In another instance, the oligonucleotide targeting Reprogramming factor polynucleotides, comprise: antisense RNA, interference RNA (RNAi), short interfering RNA (siRNA); micro interfering RNA (miRNA); a small, temporal RNA (stRNA); or a short, hairpin RNA (shRNA); small RNA-induced gene activation (RNAa); or, small activating RNA (saRNA).

[0154] In another instance, targeting of a Reprogramming factor polynucleotide, e.g. SEQ ID NO: 4 to 6 modulate the expression or function of these targets. In one instance, expression or function is up-regulated as compared to a control. In another preferred instance, expression or function is down-regulated as compared to a control.

[0155] In another instance, antisense compounds comprise sequences set forth as SEQ ID NOS: 7 to 17. These oligonucleotides can comprise one or more modified nucleotides, shorter or longer fragments, modified bonds and the like.

[0156] In another instance, SEQ ID NOS: 7 to 17 comprise one or more LNA nucleotides.

[0157] Table 1 shows exemplary antisense oligonucleotides useful in the methods of the disclosure.

Sequence ID	Antisense Sequence Name	Sequence
SEQ ID NO:7	CUR-0404	rArUrArArUrArArArUrGrGrArArCrGrUrGrGrCrUrGrGrUrArGrArU
SEQ ID NO:8	CUR-0406	rCrUrGrArGrUrUrUrCrCrArGrUrGrGrGrUrArUrArUrUrUrArGrUrG
SEQ ID NO:9	CUR-0933	T*G*A*G*T*G*G*T*C*A*G*T*G*T*T*T*C*T*T
SEQ ID NO:10	CUR-0931	G*T*G*T*C*T*T*T*G*T*A*C*T*T*G*C*T*C*C*C*T
SEQ ID NO:11	CUR-0930	C*C*C*T*T*T*A*C*T*C*T*C*T*T*C*T*C*T*C*T*C
SEQ ID NO:12	CUR-0932	T*T*G*C*T*A*C*T*T*G*T*T*G*T*C*T*G*A*G
SEQ ID NO:13	CUR-1139	G*C*C*A*T*C*A*T*T*G*T*A*C*T*C*C*A*C*T
SEQ ID NO:14	CUR-1140	A*G*T*T*G*G*G*T*G*T*G*G*T*G*G*C*T*C*A
SEQ ID NO:15	CUR-1141	T*G*G*T*C*C*C*A*G*C*C*A*C*T*T*A*G*G*A*G
SEQ ID NO:16	CUR-1142	G*G*C*G*G*G*A*G*G*A*T*T*T*C*T*T*G*A*G*G*A
SEQ ID NO:17	CUR-1143	A*T*G*N*A*G*G*T*G*T*G*G*A*G*T*G*A*T*T*C

[0158] The modulation of a desired target nucleic acid can be carried out in several ways known in the art. For example, antisense oligonucleotides, siRNA etc. Enzymatic nucleic acid molecules (e.g., ribozymes) are nucleic acid molecules capable of catalyzing one or more of a variety of reactions, including the ability to repeatedly cleave other separate nucleic acid molecules in a nucleotide base sequence-specific manner. Such enzymatic nucleic acid molecules can be used, for example, to target virtually any RNA transcript.

[0159] Because of their sequence-specificity, trans-cleaving enzymatic nucleic acid molecules show promise as therapeutic agents for human disease. Enzymatic nucleic acid molecules can be designed to cleave specific RNA targets within the background of cellular RNA. Such a cleavage event renders the mRNA non-functional and abrogates protein expression from that RNA. In this manner, synthesis of a protein associated with a disease state can be selectively inhibited.

[0160] In general, enzymatic nucleic acids with RNA cleaving activity act by first binding to a target RNA. Such binding occurs through the target binding portion of a enzymatic nucleic acid which is held in close proximity to an enzymatic portion of the molecule that acts to cleave the target RNA. Thus, the enzymatic nucleic acid first recognizes and then binds a target RNA through complementary base pairing, and once bound to the correct site, acts enzymatically to cut the target RNA. Strategic cleavage of such a target RNA will destroy its ability to direct synthesis of an encoded

protein. After an enzymatic nucleic acid has bound and cleaved its RNA target, it is released from that RNA to search for another target and can repeatedly bind and cleave new targets.

[0161] Several approaches such as in vitro selection (evolution) strategies have been used to evolve new nucleic acid catalysts capable of catalyzing a variety of reactions, such as cleavage and ligation of phosphodiester linkages and amide linkages.

[0162] The development of ribozymes that are optimal for catalytic activity would contribute significantly to any strategy that employs RNA-cleaving ribozymes for the purpose of regulating gene expression. The hammerhead ribozyme, for example, functions with a catalytic rate (kcat) of about 1 min⁻¹ in the presence of saturating (10 mM) concentrations of Mg²⁺ cofactor. An artificial "RNA ligase" ribozyme has been shown to catalyze the corresponding self-modification reaction with a rate of about 100 min⁻¹. In addition, it is known that certain modified hammerhead ribozymes that have substrate binding arms made of DNA catalyze RNA cleavage with multiple turn-over rates that approach 100 min⁻¹. Finally, replacement of a specific residue within the catalytic core of the hammerhead with certain nucleotide analogues gives modified ribozymes that show as much as a 10-fold improvement in catalytic rate. These findings demonstrate that ribozymes can promote chemical transformations with catalytic rates that are significantly greater than those displayed in vitro by most natural self-cleaving ribozymes. It is then possible that the structures of certain self-cleaving ribozymes may be optimized to give maximal catalytic activity, or that entirely new RNA motifs can be made that display significantly faster rates for RNA phosphodiester cleavage.

[0163] Intermolecular cleavage of an RNA substrate by an RNA catalyst that fits the "hammerhead" model was first shown in 1987. The RNA catalyst was recovered and reacted with multiple RNA molecules, demonstrating that it was truly catalytic.

[0164] Catalytic RNAs designed based on the "hammerhead" motif have been used to cleave specific target sequences by making appropriate base changes in the catalytic RNA to maintain necessary base pairing with the target sequences. This has allowed use of the catalytic RNA to cleave specific target sequences and indicates that catalytic RNAs designed according to the "hammerhead" model may possibly cleave specific substrate RNAs in vivo.

[0165] RNA interference (RNAi) has become a powerful tool for modulating gene expression in mammals and mammalian cells. This approach requires the delivery of small interfering RNA (siRNA) either as RNA itself or as DNA, using an expression plasmid or virus and the coding sequence for small hairpin RNAs that are processed to siRNAs. This system enables efficient transport of the pre-siRNAs to the cytoplasm where they are active and permit the use of regulated and tissue specific promoters for gene expression.

[0166] In one instance, an oligonucleotide or antisense compound comprises an oligomer or polymer of ribonucleic acid (RNA) and/or deoxyribonucleic acid (DNA), or a mimetic, chimera, analog or homolog thereof. This term includes oligonucleotides composed of naturally occurring nucleotides, sugars and covalent internucleoside (backbone) linkages as well as oligonucleotides having non-naturally occurring portions which function similarly. Such modified or substituted oligonucleotides are often desired over native forms because of desirable properties such as, for example, enhanced cellular uptake, enhanced affinity for a target nucleic acid and increased stability in the presence of nucleases.

[0167] According to the present disclosure, the oligonucleotides or "antisense compounds" include antisense oligonucleotides (e.g. RNA, DNA, mimetic, chimera, analog or homolog thereof), ribozymes, external guide sequence (EGS) oligonucleotides, siRNA compounds, single- or double-stranded RNA interference (RNAi) compounds such as siRNA compounds, saRNA, aRNA, and other oligomeric compounds which hybridize to at least a portion of the target nucleic acid and modulate its function. As such, they may be DNA, RNA, DNA-like, RNA-like, or mixtures thereof, or may be mimetics of one or more of these. These compounds may be single-stranded, double-stranded, circular or hairpin oligomeric compounds and may contain structural elements such as internal or terminal bulges, mismatches or loops. Antisense compounds are routinely prepared linearly but can be joined or otherwise prepared to be circular and/or branched. Antisense compounds can include constructs such as, for example, two strands hybridized to form a wholly or partially double-stranded compound or a single strand with sufficient self-complementarity to allow for hybridization and formation of a fully or partially double-stranded compound. The two strands can be linked internally leaving free 3' or 5' termini or can be linked to form a continuous hairpin structure or loop. The hairpin structure may contain an overhang on either the 5' or 3' terminus producing an extension of single stranded character. The double stranded compounds optionally can include overhangs on the ends. Further modifications can include conjugate groups attached to one of the termini, selected nucleotide positions, sugar positions or to one of the internucleoside linkages. Alternatively, the two strands can be linked via a non-nucleic acid moiety or linker group. When formed from only one strand, dsRNA can take the form of a self-complementary hairpin-type molecule that doubles back on itself to form a duplex. Thus, the dsRNAs can be fully or partially double stranded. Specific modulation of gene expression can be achieved by stable expression of dsRNA hairpins in transgenic cell lines. When formed from two strands, or a single strand that takes the form of a self-complementary hairpin-type molecule doubled back on itself to form a duplex, the two strands (or duplex-forming regions of a single strand) are complementary RNA strands that base pair in Watson-Crick fashion.

[0168] Once introduced to a system, the compounds of the disclosure may elicit the action of one or more enzymes or structural proteins to effect cleavage or other modification of the target nucleic acid or may work via occupancy-based mechanisms. In general, nucleic acids (including oligonucleotides) may be described as "DNA-like" (i.e., generally having one or more 2'-deoxy sugars and, generally, T rather than U bases) or "RNA-like" (i.e., generally having one or more 2'-hydroxyl or 2'-modified sugars and, generally U rather than T bases). Nucleic acid helices can adopt more than one type of structure, most commonly the A- and B-forms. It is believed that, in general, oligonucleotides which have B-form-like structure are "DNA-like" and those which have A-formlike structure are "RNA-like." In some (chimeric) instances, an antisense compound may contain both A- and B-form regions.

[0169] The antisense compounds in accordance with this disclosure can comprise an antisense portion from about 5 to about 80 nucleotides (i.e. from about 5 to about 80 linked nucleosides) in length. This refers to the length of the antisense strand or portion of the antisense compound. In other words, a single-stranded antisense compound of the disclosure comprises from 5 to about 80 nucleotides, and a double-stranded antisense compound of the disclosure (such as a dsRNA, for example) comprises a sense and an antisense strand or portion of 5 to about 80 nucleotides in length. One of ordinary skill in the art will appreciate that this comprehends antisense portions of 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59,

60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, or 80 nucleotides in length, or any range therewithin.

[0170] In one instance, the antisense compounds of the disclosure have antisense portions of 10 to 50 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies oligonucleotides having antisense portions of 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, or 50 nucleotides in length, or any range therewithin. In some instances, the oligonucleotides are 15 nucleotides in length.

[0171] In one instance, the antisense or oligonucleotide compounds of the disclosure have antisense portions of 12 or 13 to 30 nucleotides in length. One having ordinary skill in the art will appreciate that this embodies antisense compounds having antisense portions of 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides in length, or any range therewithin.

[0172] In another instance, the oligomeric compounds of the present disclosure also include variants in which a different base is present at one or more of the nucleotide positions in the compound. For example, if the first nucleotide is an adenosine, variants may be produced which contain thymidine, guanosine or cytidine at this position. This may be done at any of the positions of the antisense or dsRNA compounds. These compounds are then tested using the methods described herein to determine their ability to inhibit expression of a target nucleic acid.

[0173] In some instances, homology, sequence identity or complementarity, between the antisense compound and target is from about 40% to about 60%. In some instances, homology, sequence identity or complementarity, is from about 60% to about 70%. In some instances, homology, sequence identity or complementarity, is from about 70% to about 80%. In some instances, homology, sequence identity or complementarity, is from about 80% to about 90%. In some instances, homology, sequence identity or complementarity, is about 90%, about 92%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99% or about 100%.

[0174] In another instance, the antisense oligonucleotides, such as for example, nucleic acid molecules set forth in SEQ ID NOS: 4 to 17 comprise one or more substitutions or modifications. In one instance, the nucleotides are substituted with locked nucleic acids (LNA).

[0175] In another instance, the oligonucleotides target one or more regions of the nucleic acid molecules sense and/or antisense of coding and/or non-coding sequences associated with Reprogramming factor and the sequences set forth as SEQ ID NOS: 1 to 3 and 4 to 6. The oligonucleotides are also targeted to overlapping regions of SEQ ID NOS: 1 to 3 and 4 to 6.

[0176] Certain oligonucleotides of this disclosure are chimeric oligonucleotides. "Chimeric oligonucleotides" or "chimeras," in the context of this disclosure, are oligonucleotides which contain two or more chemically distinct regions, each made up of at least one nucleotide. These oligonucleotides typically contain at least one region of modified nucleotides that confers one or more beneficial properties (such as, for example, increased nuclease resistance, increased uptake into cells, increased binding affinity for the target) and a region that is a substrate for enzymes capable of cleaving RNA:DNA or RNA:RNA hybrids. By way of example, RNase H is a cellular endonuclease which cleaves the RNA strand of an RNA:DNA duplex. Activation of RNase H,

therefore, results in cleavage of the RNA target, thereby greatly enhancing the efficiency of antisense modulation of gene expression. Consequently, comparable results can often be obtained with shorter oligonucleotides when chimeric oligonucleotides are used, compared to phosphorothioate deoxyoligonucleotides hybridizing to the same target region. Cleavage of the RNA target can be routinely detected by gel electrophoresis and, if necessary, associated nucleic acid hybridization techniques known in the art. In one preferred instance, a chimeric oligonucleotide comprises at least one region modified to increase target binding affinity, and, usually, a region that acts as a substrate for RNase H. Affinity of an oligonucleotide for its target (in this case, a nucleic acid encoding ras) is routinely determined by measuring the T_m of an oligonucleotide/target pair, which is the temperature at which the oligonucleotide and target dissociate; dissociation is detected spectrophotometrically. The higher the T_m , the greater is the affinity of the oligonucleotide for the target.

[0177] Chimeric antisense compounds of the disclosure may be formed as composite structures of two or more oligonucleotides, modified oligonucleotides, oligonucleosides and/or oligonucleotides mimetics as described above. Such; compounds have also been referred to in the art as hybrids or gapmers. Representative United States patents that teach the preparation of such hybrid structures comprise, but are not limited to, US patent nos. 5,013,830; 5,149,797; 5, 220,007; 5,256,775; 5,366,878; 5,403,711; 5,491,133; 5,565,350; 5,623,065; 5,652,355; 5,652,356; and 5,700,922.

[0178] In another instance, the region of the oligonucleotide which is modified comprises at least one nucleotide modified at the 2' position of the sugar, most preferably a 2'-Oalkyl, 2'-O-alkyl-O-alkyl or 2'-fluoro-modified nucleotide. In other preferred instances, RNA modifications include 2'-fluoro, 2'-amino and 2' O-methyl modifications on the ribose of pyrimidines, abasic residues or an inverted base at the 3' end of the RNA. Such modifications are routinely incorporated into oligonucleotides and these oligonucleotides have been shown to have a higher T_m (i.e., higher target binding affinity) than; 2'-deoxyoligonucleotides against a given target. The effect of such increased affinity is to greatly enhance RNAi oligonucleotide inhibition of gene expression. RNase H is a cellular endonuclease that cleaves the RNA strand of RNA:DNA duplexes; activation of this enzyme therefore results in cleavage of the RNA target, and thus can greatly enhance the efficiency of RNAi inhibition. Cleavage of the RNA target can be routinely demonstrated by gel electrophoresis. In another preferred instance, the chimeric oligonucleotide is also modified to enhance nuclease resistance. Cells contain a variety of exo- and endo-nucleases which can degrade nucleic acids. A number of nucleotide and nucleoside modifications have been shown to make the oligonucleotide into which they are incorporated more resistant to nuclease digestion than the native oligodeoxynucleotide. Nuclease resistance is routinely measured by incubating oligonucleotides with cellular extracts or isolated nuclease solutions and measuring the extent of intact oligonucleotide remaining over time, usually by gel electrophoresis. Oligonucleotides which have been modified to enhance their nuclease resistance survive intact for a longer time than unmodified oligonucleotides. A variety of oligonucleotide modifications have been demonstrated to enhance or confer nuclease resistance. Oligonucleotides which contain at least one phosphorothioate modification are presently more preferred. In some cases, oligonucleotide modifications which enhance target binding affinity are also, independently, able to enhance nuclease resistance. Some desirable modifications can be found in De Mesmaeker et al. (1995) *Acc. Chem. Res.*, 28:366-374.

[0179] Specific examples of some preferred oligonucleotides envisioned for this disclosure include those comprising modified backbones, for example, phosphorothioates, phosphotriesters, methyl

phosphonates, short chain alkyl or cycloalkyl intersugar linkages or short chain heteroatomic or heterocyclic intersugar linkages. Most preferred are oligonucleotides with phosphorothioate backbones and those with heteroatom backbones, particularly $\text{CH}_2\text{--NH--O--CH}_2$, $\text{CH}_2\text{--N(CH}_3\text{)--O--CH}_2$ [known as a methylene(methylimino) or MMI backbone], $\text{CH}_2\text{--O--N(CH}_3\text{)--CH}_2$, $\text{CH}_2\text{--N(CH}_3\text{)--N(CH}_3\text{)--CH}_2$ and $\text{O--N(CH}_3\text{)--CH}_2\text{--CH}_2$ backbones, wherein the native phosphodiester backbone is represented as O--P--O--CH_2 . The amide backbones disclosed by De Mesmaeker et al. (1995) *Acc. Chem. Res.* 28:366-374 are also preferred. Also preferred are oligonucleotides having morpholino backbone structures (Summerton and Weller, U.S. Pat. No. 5,034,506). In other preferred instances, such as the peptide nucleic acid (PNA) backbone, the phosphodiester backbone of the oligonucleotide is replaced with a polyamide backbone, the nucleotides being bound directly or indirectly to the aza nitrogen atoms of the polyamide backbone (Nielsen et al. (1991) *Science* 254, 1497). Oligonucleotides may also comprise one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH, SH, SCH₃, F, OCN, OCH₃, OCH₃, O(CH₂)_n CH₃, O(CH₂)_n NH₂ or O(CH₂)_n CH₃ where n is from 1 to about 10; C₁ to C₁₀ lower alkyl, alkoxyalkoxy, substituted lower alkyl, alkaryl or aralkyl; Cl; Br; CN; CF₃; OCF₃; O--, S--, or N-alkyl; O--, S--, or N-alkenyl; SOCH₃; SO₂ CH₃; ONO₂; NO₂; N₃; NH₂; heterocycloalkyl; heterocycloalkaryl; aminoalkylamino; polyalkylamino; substituted silyl; an RNA cleaving group; a reporter group; an intercalator; a group for improving the pharmacokinetic properties of an oligonucleotide; or a group for improving the pharmacodynamic properties of an oligonucleotide and other substituents having similar properties. A preferred modification includes 2'-methoxyethoxy [2'-O-CH₂ CH₂ OCH₃, also known as 2'-O-(2-methoxyethyl)] (Martin et al., (1995) *Helv. Chim. Acta*, 78, 486). Other preferred modifications include 2'-methoxy (2'-O-CH₃), 2'-propoxy (2'-OCH₂ CH₂CH₃) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyls in place of the pentofuranosyl group.

[0180] Oligonucleotides may also include, additionally or alternatively, nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleotides include adenine (A), guanine (G), thymine (T), cytosine (C) and uracil (U). Modified nucleotides include nucleotides found only infrequently or transiently in natural nucleic acids, e.g., hypoxanthine, 6-methyladenine, 5-Me pyrimidines, particularly 5-methylcytosine (also referred to as 5-methyl-2' deoxycytosine and often referred to in the art as 5-Me-C), 5-hydroxymethylcytosine (HMC), glycosyl HMC and gentobiosyl HMC, as well as synthetic nucleotides, e.g., 2-aminoadenine, 2-(methylamino)adenine, 2-(imidazolylalkyl)adenine, 2-(aminoalkylamino)adenine or other heterosubstituted alkyladenines, 2-thiouracil, 2-thiothymine, 5-bromouracil, 5-hydroxymethyluracil, 8-azaguanine, 7-deazaguanine, N⁶ (6-aminohexyl)adenine and 2,6-diaminopurine. (Komberg, A., *DNA Replication*, W. H. Freeman & Co., San Francisco, 1980, pp75-77; Gebeyehu, G., (1987) et al. *Nucl. Acids Res.* 15:4513). A "universal" base known in the art, e.g., inosine, may be included. 5-Me-C substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C. (Sanghvi, Y. S., in Crooke, S. T. and Lebleu, B., eds., *Antisense Research and Applications*, CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions.

[0181] Another modification of the oligonucleotides of the disclosure involves chemically linking to the oligonucleotide one or more moieties or conjugates which enhance the activity or cellular uptake of the oligonucleotide. Such moieties include but are not limited to lipid moieties such as a cholesterol moiety, a cholesteryl moiety, a thioether, e.g., hexyl-S-tritylthiol, a thiocholesterol, an

aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero- 3-H-phosphonate, a polyamine or a polyethylene glycol chain, or adamantane acetic acid. Oligonucleotides comprising lipophilic moieties, and methods for preparing such oligonucleotides are known in the art, for example, U.S. Pat. Nos. 5,138,045, 5,218,105 and 5,459,255.

[0182] It is not necessary for all positions in a given oligonucleotide to be uniformly modified, and in fact more than one of the aforementioned modifications may be incorporated in a single oligonucleotide or even at within a single nucleoside within an oligonucleotide. The present disclosure also includes oligonucleotides which are chimeric oligonucleotides as hereinbefore defined.

[0183] In another instance, the nucleic acid molecule of the present disclosure is conjugated with another moiety including but not limited to abasic nucleotides, polyether, polyamine, polyamides, peptides, carbohydrates, lipid, or polyhydrocarbon compounds. Those skilled in the art will recognize that these molecules can be linked to one or more of any nucleotides comprising the nucleic acid molecule at several positions on the sugar, base or phosphate group.

[0184] The oligonucleotides used in accordance with this disclosure may be conveniently and routinely made through the well-known technique of solid phase synthesis. Equipment for such synthesis is sold by several vendors including Applied Biosystems. Any other means for such synthesis may also be employed; the actual synthesis of the oligonucleotides is well within the talents of one of ordinary skill in the art. It is also well known to use similar techniques to prepare other oligonucleotides such as the phosphorothioates and alkylated derivatives. It is also well known to use similar techniques and commercially available modified amidites and controlled-pore glass (CPG) products such as biotin, fluorescein, acridine or psoralen-modified amidites and/or CPG (available from Glen Research, Sterling VA) to synthesize fluorescently labeled, biotinylated or other modified oligonucleotides such as cholesterol-modified oligonucleotides.

[0185] In accordance with the disclosure, use of modifications such as the use of LNA monomers to enhance the potency, specificity and duration of action and broaden the routes of administration of oligonucleotides comprised of current chemistries such as MOE, ANA, FANA, PS etc (Uhlman, et al. (2000) Current Opinions in Drug Discovery & Development Vol. 3 No 2). This can be achieved by substituting some of the monomers in the current oligonucleotides by LNA monomers. The LNA modified oligonucleotide may have a size similar to the parent compound or may be larger or preferably smaller. It is preferred that such LNA-modified oligonucleotides contain less than about 70%, more preferably less than about 60%, most preferably less than about 50% LNA monomers and that their sizes are between about 5 and 25 nucleotides, more preferably between about 12 and 20 nucleotides.

[0186] Modified oligonucleotide backbones comprise, but are not limited to, phosphorothioates, chiral phosphorothioates, phosphorodithioates, phosphotriesters, aminoalkylphosphotriesters, methyl and other alkyl phosphonates comprising 3'-alkylene phosphonates and chiral phosphonates, phosphinates, phosphoramidates comprising 3'-amino phosphoramidate and aminoalkylphosphoramidates, thionophosphoramidates, thionoalkylphosphonates, thionoalkylphosphotriesters, and boranophosphates having normal 3'-5' linkages, 2'-5' linked analogs of these, and those having inverted polarity wherein the adjacent pairs of nucleoside units

are linked 3'-5' to 5'-3' or 2'-5' to 5'-2'. Various salts, mixed salts and free acid forms are also included.

[0187] Representative United States patents that teach the preparation of the above phosphorus containing linkages comprise, but are not limited to, US patent nos. 3,687,808; 4,469,863; 4,476,301; 5,023,243; 5, 177,196; 5,188,897; 5,264,423; 5,276,019; 5,278,302; 5,286,717; 5,321,131; 5,399,676; 5,405,939; 5,453,496; 5,455, 233; 5,466,677; 5,476,925; 5,519,126; 5,536,821; 5,541,306; 5,550,111; 5,563, 253; 5,571,799; 5,587,361; and 5,625,050.

[0188] Modified oligonucleotide backbones that do not include a phosphorus atom therein have backbones that are formed by short chain alkyl or cycloalkyl internucleoside linkages, mixed heteroatom and alkyl or cycloalkyl internucleoside linkages, or one or more short chain heteroatomic or heterocyclic internucleoside linkages. These comprise those having morpholino linkages (formed in part from the sugar portion of a nucleoside); siloxane backbones; sulfide, sulfoxide and sulfone backbones; formacetyl and thioformacetyl backbones; methylene formacetyl and thioformacetyl backbones; alkene containing backbones; sulfamate backbones; methyleneimino and methylenehydrazino backbones; sulfonate and sulfonamide backbones; amide backbones; and others having mixed N, O, S and CH₂ component parts.

[0189] Representative United States patents that teach the preparation of the above oligonucleosides comprise, but are not limited to, US patent nos. 5,034,506; 5,166,315; 5,185,444; 5,214,134; 5,216,141; 5,235,033; 5,264, 562; 5, 264,564; 5,405,938; 5,434,257; 5,466,677; 5,470,967; 5,489,677; 5,541,307; 5,561,225; 5,596, 086; 5,602,240; 5,610,289; 5,602,240; 5,608,046; 5,610,289; 5,618,704; 5,623, 070; 5,663,312; 5,633,360; 5,677,437; and 5,677,439.

[0190] In other oligonucleotide mimetics, both the sugar and the internucleoside linkage, i.e., the backbone, of the nucleotide units are replaced with novel groups. The base units are maintained for hybridization with an appropriate nucleic acid target compound. One such oligomeric compound, an oligonucleotide mimetic that has been shown to have excellent hybridization properties, is referred to as a peptide nucleic acid (PNA). In PNA compounds, the sugar-backbone of an oligonucleotide is replaced with an amide containing backbone, in particular an aminoethylglycine backbone. The nucleobases are retained and are bound directly or indirectly to aza nitrogen atoms of the amide portion of the backbone. Representative United States patents that teach the preparation of PNA compounds comprise, but are not limited to, US patent nos. 5,539,082; 5,714,331; and 5,719,262. Further teaching of PNA compounds can be found in Nielsen, et al. (1991) Science 254, 1497-1500.

[0191] In another instance of the disclosure the oligonucleotides with phosphorothioate backbones and oligonucleosides with heteroatom backbones, and in particular- CH₂-NH-O-CH₂-, -CH₂-N (CH₃)-O-CH₂-known as a methylene (methylimino) or MMI backbone, - CH₂-O-N (CH₃)-CH₂-, -CH₂N(CH₃)-N(CH₃) CH₂-and-O-N(CH₃)-CH₂-CH₂- wherein the native phosphodiester backbone is represented as-O-P-O-CH₂- of the above referenced US patent no. 5,489,677, and the amide backbones of the above referenced US patent no. 5,602,240. Also preferred are oligonucleotides having morpholino backbone structures of the above-referenced US patent no. 5,034,506.

[0192] Modified oligonucleotides may also contain one or more substituted sugar moieties. Preferred oligonucleotides comprise one of the following at the 2' position: OH; F; O-, S-, or N-alkyl; O-, S-, or N-alkenyl; O-, S-or N-alkynyl; or O alkyl-O-alkyl, wherein the alkyl, alkenyl and alkynyl may

be substituted or unsubstituted C to CO alkyl or C2 to CO alkenyl and alkynyl. Particularly preferred are $O(CH_2)_n OmCH_3$, $O(CH_2)_n OCH_3$, $O(CH_2)_n NH_2$, $O(CH_2)_n CH_3$, $O(CH_2)_n ONH_2$, and $O(CH_2)_n ON(CH_2)_m CH_3$ where n and m can be from 1 to about 10. Other preferred oligonucleotides comprise one of the following at the 2' position: C to CO, (lower alkyl, substituted lower alkyl, alkaryl, aralkyl, O-alkaryl or O-aralkyl, SH, SCH₃, OCN, Cl, Br, CN, CF₃, OCF₃, SOCH₃, SO₂CH₃, ONO₂, NO₂, N₃, NH₂, heterocycloalkyl, heterocycloalkaryl, aminoalkylamino, polyalkylamino, substituted silyl, an RNA cleaving group, a reporter group, an intercalator, a group for improving the pharmacokinetic properties of an oligonucleotide, or a group for improving the pharmacodynamic properties of an oligonucleotide, and other substituents having similar properties. A preferred modification comprises 2'-methoxyethoxy (2'-O-CH₂CH₂OCH₃, also known as 2'-O-(2-methoxyethyl) or 2'-MOE) (Martin et al., (1995) *Helv. Chim. Acta*, 78, 486-504) i.e., an alkoxyalkoxy group. A further preferred modification comprises 2'-dimethylaminoethoxy, i.e., a $O(CH_2)_2 ON(CH_3)_2$ group, also known as 2'-DMAOE, as described in examples herein below, and 2'-dimethylaminoethoxyethoxy (also known in the art as 2'-O-dimethylaminoethoxyethyl or 2'-DMAEOE), i.e., 2'-O-CH₂-O-CH₂-N(CH₂)₂.

[0193] Other modifications comprise 2'-methoxy (2'-O CH₃), 2'-aminopropoxy (2'-O CH₂CH₂CH₂NH₂) and 2'-fluoro (2'-F). Similar modifications may also be made at other positions on the oligonucleotide, particularly the 3' position of the sugar on the 3' terminal nucleotide or in 2'-5' linked oligonucleotides and the 5' position of 5' terminal nucleotide. Oligonucleotides may also have sugar mimetics such as cyclobutyl moieties in place of the pentofuranosyl sugar. Representative United States patents that teach the preparation of such modified sugar structures comprise, but are not limited to, US patent nos. 4,981,957; 5,118,800; 5,319,080; 5,359,044; 5,393,878; 5,446,137; 5,466,786; 5,514, 785; 5,519,134; 5,567,811; 5,576,427; 5,591,722; 5,597,909; 5,610,300; 5,627,053; 5,639,873; 5,646, 265; 5,658,873; 5,670,633; and 5,700,920.

[0194] Oligonucleotides may also comprise nucleobase (often referred to in the art simply as "base") modifications or substitutions. As used herein, "unmodified" or "natural" nucleotides comprise the purine bases adenine (A) and guanine (G), and the pyrimidine bases thymine (T), cytosine (C) and uracil (U). Modified nucleotides comprise other synthetic and natural nucleotides such as 5-methylcytosine (5-me-C), 5-hydroxymethyl cytosine, xanthine, hypoxanthine, 2-aminoadenine, 6-methyl and other alkyl derivatives of adenine and guanine, 2-propyl and other alkyl derivatives of adenine and guanine, 2-thiouracil, 2-thiothymine and 2-thiocytosine, 5-halouracil and cytosine, 5-propynyl uracil and cytosine, 6-azo uracil, cytosine and thymine, 5-uracil (pseudo-uracil), 4-thiouracil, 8-halo, 8-amino, 8-thiol, 8-thioalkyl, 8-hydroxyl and other 8-substituted adenines and guanines, 5-halo particularly 5-bromo, 5-trifluoromethyl and other 5-substituted uracils and cytosines, 7-methylguanine and 7-methyladenine, 8-azaguanine and 8-azaadenine, 7-deazaguanine and 7-deazaadenine and 3-deazaguanine and 3-deazaadenine.

[0195] Further, nucleotides comprise those disclosed in United States Patent No. 3,687,808, those disclosed in 'The Concise Encyclopedia of Polymer Science And Engineering', pages 858-859, Kroschwitz, J.I., ed. John Wiley & Sons, 1990, those disclosed by Englisch et al., 'Angewandte Chemie, International Edition', 1991, 30, page 613, and those disclosed by Sanghvi, Y.S., Chapter 15, 'Antisense Research and Applications', pages 289-302, Crooke, S.T. and Lebleu, B. ea., CRC Press, 1993. Certain of these nucleotides are particularly useful for increasing the binding affinity of the oligomeric compounds of the disclosure. These comprise 5-substituted pyrimidines, 6-azapyrimidines and N-2, N-6 and O-6 substituted purines, comprising 2-aminopropyladenine, 5-

propynyluracil and 5-propynylcytosine. 5-methylcytosine substitutions have been shown to increase nucleic acid duplex stability by 0.6-1.2°C (Sanghvi, Y.S., Crooke, S.T. and Lebleu, B., eds, 'Antisense Research and Applications', CRC Press, Boca Raton, 1993, pp. 276-278) and are presently preferred base substitutions, even more particularly when combined with 2'-O-methoxyethyl sugar modifications.

[0196] Representative United States patents that teach the preparation of the above noted modified nucleotides as well as other modified nucleotides comprise, but are not limited to, US patent nos. 3,687,808, as well as 4,845,205; 5,130,302; 5,134,066; 5,175, 273; 5, 367,066; 5,432,272; 5,457,187; 5,459,255; 5,484,908; 5,502,177; 5,525,711; 5,552,540; 5,587,469; 5,596,091; 5,614,617; 5,750,692, and 5,681,941.

[0197] Another modification of the oligonucleotides of the disclosure involves chemically linking to the oligonucleotide one or more moieties or conjugates, which enhance the activity, cellular distribution, or cellular uptake of the oligonucleotide.

[0198] Such moieties comprise but are not limited to, lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g., hexyl-S-tritylthiol (Manoharan et al., (1992) Ann. N. Y. Acad. Sci., 660, 306-309; Manoharan et al., (1993) Bioorg. Med. Chem. Let., 3, 2765-2770), a thiocholesterol, an aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-H-phosphonate, a polyamine or a polyethylene glycol chain, or adamantane acetic acid, a palmityl moiety, or an octadecylamine or hexylamino-carbonyl-t oxcholesterol moiety.

[0199] Representative United States patents that teach the preparation of such oligonucleotides conjugates comprise, but are not limited to, US patent nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552, 538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486, 603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762, 779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082, 830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5, 245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391, 723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5, 565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599, 928 and 5,688,941.

[0200] *Drug discovery:* The compounds of the present disclosure can also be applied in the areas of drug discovery and target validation. The present disclosure comprehends the use of the compounds and preferred target segments identified herein in drug discovery efforts to elucidate relationships that exist between a Reprogramming factor polynucleotide and a disease state, phenotype, or condition. These methods include detecting or modulating a Reprogramming factor polynucleotide comprising contacting a sample, tissue, cell, or organism with the compounds of the present disclosure, measuring the nucleic acid or protein level of a Reprogramming factor polynucleotide and/or a related phenotypic or chemical endpoint at some time after treatment, and optionally comparing the measured value to a non-treated sample or sample treated with a further compound of the disclosure. These methods can also be performed in parallel or in combination with other experiments to determine the function of unknown genes for the process of target validation or to determine the validity of a particular gene product as a target for treatment or prevention of a particular disease, condition, or phenotype.

Assessing Up-regulation or Inhibition of Gene Expression:

[0201] Transfer of an exogenous nucleic acid into a host cell or organism can be assessed by directly detecting the presence of the nucleic acid in the cell or organism. Such detection can be achieved by several methods well known in the art. For example, the presence of the exogenous nucleic acid can be detected by Southern blot or by a polymerase chain reaction (PCR) technique using primers that specifically amplify nucleotide sequences associated with the nucleic acid. Expression of the exogenous nucleic acids can also be measured using conventional methods including gene expression analysis. For instance, mRNA produced from an exogenous nucleic acid can be detected and quantified using a Northern blot and reverse transcription PCR (RT-PCR).

[0202] Expression of RNA from the exogenous nucleic acid can also be detected by measuring an enzymatic activity or a reporter protein activity. For example, antisense modulatory activity can be measured indirectly as a decrease or increase in target nucleic acid expression as an indication that the exogenous nucleic acid is producing the effector RNA. Based on sequence conservation, primers can be designed and used to amplify coding regions of the target genes. Initially, the most highly expressed coding region from each gene can be used to build a model control gene, although any coding or non coding region can be used. Each control gene is assembled by inserting each coding region between a reporter coding region and its poly(A) signal. These plasmids would produce an mRNA with a reporter gene in the upstream portion of the gene and a potential RNAi target in the 3' non-coding region. The effectiveness of individual antisense oligonucleotides would be assayed by modulation of the reporter gene. Reporter genes useful in the methods of the present disclosure include acetohydroxyacid synthase (AHAS), alkaline phosphatase (AP), beta galactosidase (LacZ), beta glucuronidase (GUS), chloramphenicol acetyltransferase (CAT), green fluorescent protein (GFP), red fluorescent protein (RFP), yellow fluorescent protein (YFP), cyan fluorescent protein (CFP), horseradish peroxidase (HRP), luciferase (Luc), nopaline synthase (NOS), octopine synthase (OCS), and derivatives thereof. Multiple selectable markers are available that confer resistance to ampicillin, bleomycin, chloramphenicol, gentamycin, hygromycin, kanamycin, lincomycin, methotrexate, phosphinothricin, puromycin, and tetracycline. Methods to determine modulation of a reporter gene are well known in the art, and include, but are not limited to, fluorometric methods (e.g. fluorescence spectroscopy, Fluorescence Activated Cell Sorting (FACS), fluorescence microscopy), antibiotic resistance determination.

[0203] Sox2, KLF4, and POU5F1 proteins and mRNA expression can be assayed using methods known to those of skill in the art and described elsewhere herein. For example, immunoassays such as the ELISA can be used to measure protein levels. Reprogramming factor antibodies for ELISAs are available commercially, e.g., from R&D Systems (Minneapolis, MN), Abcam, Cambridge, MA.

[0204] In instances, Sox2, KLF4, and POU5F1 expression (e.g., mRNA or protein) in a sample (e.g., cells or tissues in vivo or in vitro) treated using an antisense oligonucleotide of the disclosure is evaluated by comparison with Reprogramming factor expression in a control sample. For example, expression of the protein or nucleic acid can be compared using methods known to those of skill in the art with that in a mock-treated or untreated sample. Alternatively, comparison with a sample treated with a control antisense oligonucleotide (e.g., one having an altered or different sequence)

can be made depending on the information desired. In another instance, , a difference in the expression of the Reprogramming factor protein or nucleic acid in a treated vs. an untreated sample can be compared with the difference in expression of a different nucleic acid (including any standard deemed appropriate by the researcher, e.g., a housekeeping gene) in a treated sample vs. an untreated sample.

[0205] Observed differences can be expressed as desired, e.g., in the form of a ratio or fraction, for use in a comparison with control. In instances, the level of a Reprogramming factor mRNA or protein, in a sample treated with an antisense oligonucleotide of the present disclosure, is increased or decreased by about 1.25-fold to about 10-fold or more relative to an untreated sample or a sample treated with a control nucleic acid. In instances, the level of a Reprogramming factor mRNA or protein is increased or decreased by at least about 1.25-fold, at least about 1.3-fold, at least about 1.4-fold, at least about 1.5-fold, at least about 1.6-fold, at least about 1.7-fold, at least about 1.8-fold, at least about 2-fold, at least about 2.5-fold, at least about 3-fold, at least about 3.5-fold, at least about 4-fold, at least about 4.5-fold, at least about 5-fold, at least about 5.5-fold, at least about 6-fold, at least about 6.5-fold, at least about 7-fold, at least about 7.5-fold, at least about 8-fold, at least about 8.5-fold, at least about 9-fold, at least about 9.5-fold, or at least about 10-fold or more.

Kits, Research Reagents, Diagnostics, and Therapeutics

[0206] The compounds of the present disclosure can be utilized for diagnostics, therapeutics, and prophylaxis, and as research reagents and components of kits. Furthermore, antisense oligonucleotides, which are able to inhibit gene expression with exquisite specificity, are often used by those of ordinary skill to elucidate the function of particular genes or to distinguish between functions of various members of a biological pathway.

[0207] For use in kits and diagnostics and in various biological systems, the compounds of the present disclosure, either alone or in combination with other compounds or therapeutics, are useful as tools in differential and/or combinatorial analyses to elucidate expression patterns of a portion or the entire complement of genes expressed within cells and tissues.

[0208] As used herein the term "biological system" or "system" is defined as any organism, cell, cell culture or tissue that expresses, or is made competent to express products of the Reprogramming factors. These include, but are not limited to, humans, transgenic animals, cells, cell cultures, tissues, xenografts, transplants and combinations thereof.

[0209] As one non limiting example, expression patterns within cells or tissues treated with one or more antisense compounds are compared to control cells or tissues not treated with antisense compounds and the patterns produced are analyzed for differential levels of gene expression as they pertain, for example, to disease association, signaling pathway, cellular localization, expression level, size, structure or function of the genes examined. These analyses can be performed on stimulated or unstimulated cells and in the presence or absence of other compounds that affect expression patterns.

[0210] Examples of methods of gene expression analysis known in the art include DNA arrays or

microarrays (Brazma and Vilo, (2000) FEBS Lett., 480, 17-24; Celis, et al., (2000) FEBS Lett., 480, 2-16), SAGE (serial analysis of gene expression) (Madden, et al., (2000) Drug Discov. Today, 5, 415-425), READS (restriction enzyme amplification of digested cDNAs) (Prashar and Weissman, (1999) Methods Enzymol., 303, 258-72), TOGA (total gene expression analysis) (Sutcliffe, et al., (2000) Proc. Natl. Acad. Sci. U.S.A., 97, 1976-81), protein arrays and proteomics (Celis, et al., (2000) FEBS Lett., 480, 2-16; Jungblut, et al., Electrophoresis, 1999, 20, 2100-10), expressed sequence tag (EST) sequencing (Celis, et al., FEBS Lett., 2000, 480, 2-16; Larsson, et al., J. Biotechnol., 2000, 80, 143-57), subtractive RNA fingerprinting (SuRF) (Fuchs, et al., (2000) Anal. Biochem. 286, 91-98; Larson, et al., (2000) Cytometry 41, 203-208), subtractive cloning, differential display (DD) (Jurecic and Belmont, (2000) Curr. Opin. Microbiol. 3, 316-21), comparative genomic hybridization (Carulli, et al., (1998) J. Cell Biochem. Suppl., 31, 286-96), FISH (fluorescent in situ hybridization) techniques (Going and Gusterson, (1999) Eur. J. Cancer, 35, 1895-904) and mass spectrometry methods (To, Comb. (2000) Chem. High Throughput Screen, 3, 235-41).

[0211] The compounds of the disclosure are useful for research and diagnostics, because these compounds hybridize to nucleic acids encoding a Reprogramming factor. For example, oligonucleotides that hybridize with such efficiency and under such conditions as disclosed herein as to be effective Reprogramming factor modulators are effective primers or probes under conditions favoring gene amplification or detection, respectively. These primers and probes are useful in methods requiring the specific detection of nucleic acid molecules encoding a Reprogramming factor and in the amplification of said nucleic acid molecules for detection or for use in further studies of a Reprogramming factor. Hybridization of the antisense oligonucleotides, particularly the primers and probes, of the disclosure with a nucleic acid encoding a Reprogramming factor can be detected by means known in the art. Such means may include conjugation of an enzyme to the oligonucleotide, radiolabeling of the oligonucleotide, or any other suitable detection means. Kits using such detection means for detecting the level of a Reprogramming factor in a sample may also be prepared.

[0212] The specificity and sensitivity of antisense are also harnessed by those of skill in the art for therapeutic uses. Antisense compounds have been employed as therapeutic moieties in the treatment of disease states in animals, including humans. Antisense oligonucleotide drugs have been safely and effectively administered to humans and numerous clinical trials are presently underway. It is thus established that antisense compounds can be useful therapeutic modalities that can be configured to be useful in treatment regimes for the treatment of cells, tissues and animals, especially humans.

[0213] For therapeutics, an animal, preferably a human, suspected of having a disease or disorder which can be treated by modulating the expression of a Reprogramming factor polynucleotide is treated by administering antisense compounds in accordance with this disclosure. For example, in one non-limiting instance, the methods comprise the step of administering to the animal in need of treatment, a therapeutically effective amount of a Reprogramming factor modulator. The Reprogramming factor modulators of the present disclosure effectively modulate the activity of a Reprogramming factor or modulate the expression of a Reprogramming factor protein. In one instance, the activity or expression of a Reprogramming factor in an animal is inhibited by about 10% as compared to a control. Preferably, the activity or expression of a Reprogramming factor in an animal is inhibited by about 30%. More preferably, the activity or expression of a Reprogramming factor in an animal is inhibited by 50% or more. Thus, the oligomeric compounds modulate expression of a Reprogramming factor mRNA by at least 10%, by at least 50%, by at least 25%, by

at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, by at least 95%, by at least 98%, by at least 99%, or by 100% as compared to a control.

[0214] In one instance, the activity or expression of a Reprogramming factor and/or in an animal is increased by about 10% as compared to a control. Preferably, the activity or expression of a Reprogramming factor in an animal is increased by about 30%. More preferably, the activity or expression of a Reprogramming factor in an animal is increased by 50% or more. Thus, the oligomeric compounds modulate expression of a Reprogramming factor mRNA by at least 10%, by at least 50%, by at least 25%, by at least 30%, by at least 40%, by at least 50%, by at least 60%, by at least 70%, by at least 75%, by at least 80%, by at least 85%, by at least 90%, by at least 95%, by at least 98%, by at least 99%, or by 100% as compared to a control.

[0215] For example, the reduction of the expression of a Reprogramming factor may be measured in serum, blood, adipose tissue, liver or any other body fluid, tissue or organ of the animal. Preferably, the cells contained within said fluids, tissues or organs being analyzed contain a nucleic acid molecule encoding Reprogramming factor peptides and/or the Reprogramming factor protein itself.

[0216] The compounds of the disclosure can be utilized in pharmaceutical compositions by adding an effective amount of a compound to a suitable pharmaceutically acceptable diluent or carrier. Use of the compounds and methods of the disclosure may also be useful prophylactically.

Conjugates

[0217] Another modification of the oligonucleotides of the disclosure involves chemically linking to the oligonucleotide one or more moieties or conjugates that enhance the activity, cellular distribution or cellular uptake of the oligonucleotide. These moieties or conjugates can include conjugate groups covalently bound to functional groups such as primary or secondary hydroxyl groups. Conjugate groups of the disclosure include intercalators, reporter molecules, polyamines, polyamides, polyethylene glycols, polyethers, groups that enhance the pharmacodynamic properties of oligomers, and groups that enhance the pharmacokinetic properties of oligomers. Typical conjugate groups include cholesterols, lipids, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. Groups that enhance the pharmacodynamic properties, in the context of this disclosure, include groups that improve uptake, enhance resistance to degradation, and/or strengthen sequence-specific hybridization with the target nucleic acid. Groups that enhance the pharmacokinetic properties, in the context of this disclosure, include groups that improve uptake, distribution, metabolism or excretion of the compounds of the present disclosure. Representative conjugate groups are disclosed in International Patent Application No. PCT/LTS92/09196, filed Oct. 23, 1992, and U.S. Pat. No. 6,287,860. Conjugate moieties include, but are not limited to, lipid moieties such as a cholesterol moiety, cholic acid, a thioether, e.g., hexyl-5- tritylthiol, a thiocholesterol, an aliphatic chain, e.g., dodecandiol or undecyl residues, a phospholipid, e.g., di-hexadecyl-rac-glycerol or triethylammonium 1,2-di-O-hexadecyl-rac-glycero-3-Hphosphonate, a polyamine or a polyethylene glycol chain, or adamantane acetic acid, a palmityl moiety, or an octadecylamine or hexylamino-carbonyl-oxycholesterol moiety. Oligonucleotides of the

disclosure may also be conjugated to active drug substances, for example, aspirin, warfarin, phenylbutazone, ibuprofen, suprofen, fenbufen, ketoprofen, (S)-(+)-pranoprofen, carprofen, dansylsarcosine, 2,3,5-triiodobenzoic acid, flufenamic acid, folinic acid, a benzothiadiazide, chlorothiazide, a diazepine, indomethicin, a barbiturate, a cephalosporin, a sulfa drug, an antidiabetic, an antibacterial or an antibiotic.

[0218] Representative United States patents that teach the preparation of such oligonucleotides conjugates include, but are not limited to, U.S. Pat. Nos. 4,828,979; 4,948,882; 5,218,105; 5,525,465; 5,541,313; 5,545,730; 5,552,538; 5,578,717; 5,580,731; 5,580,731; 5,591,584; 5,109,124; 5,118,802; 5,138,045; 5,414,077; 5,486,603; 5,512,439; 5,578,718; 5,608,046; 4,587,044; 4,605,735; 4,667,025; 4,762,779; 4,789,737; 4,824,941; 4,835,263; 4,876,335; 4,904,582; 4,958,013; 5,082,830; 5,112,963; 5,214,136; 5,082,830; 5,112,963; 5,214,136; 5,245,022; 5,254,469; 5,258,506; 5,262,536; 5,272,250; 5,292,873; 5,317,098; 5,371,241; 5,391,723; 5,416,203; 5,451,463; 5,510,475; 5,512,667; 5,514,785; 5,565,552; 5,567,810; 5,574,142; 5,585,481; 5,587,371; 5,595,726; 5,597,696; 5,599,923; 5,599,928 and 5,688,941.

Formulations

[0219] The compounds of the disclosure may also be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, as for example, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. Representative United States patents that teach the preparation of such uptake, distribution and/or absorption-assisting formulations include, but are not limited to, U.S. Pat. Nos. 5,108,921; 5,354,844; 5,416,016; 5,459,127; 5,521,291; 5,543,165; 5,547,932; 5,583,020; 5,591,721; 4,426,330; 4,534,899; 5,013,556; 5,108,921; 5,213,804; 5,227,170; 5,264,221; 5,356,633; 5,395,619; 5,416,016; 5,417,978; 5,462,854; 5,469,854; 5,512,295; 5,527,528; 5,534,259; 5,543,152; 5,556,948; 5,580,575; and 5,595,756.

[0220] Although, the antisense oligonucleotides do not need to be administered in the context of a vector in order to modulate a target expression and/or function, instances of the disclosure relates to expression vector constructs for the expression of antisense oligonucleotides, comprising promoters, hybrid promoter gene sequences and possess a strong constitutive promoter activity, or a promoter activity which can be induced in the desired case.

[0221] In an instance, disclosure practice involves administering at least one of the foregoing antisense oligonucleotides with a suitable nucleic acid delivery system. In one instance, that system includes a non-viral vector operably linked to the polynucleotide. Examples of such nonviral vectors include the oligonucleotide alone (e.g. any one or more of SEQ ID NOS: 7 to 17) or in combination with a suitable protein, polysaccharide or lipid formulation.

[0222] Additionally suitable nucleic acid delivery systems include viral vector, typically sequence from at least one of an adenovirus, adenovirus-associated virus (AAV), helper-dependent adenovirus, retrovirus, or hemagglutinin virus of Japan-liposome (HVJ) complex. Preferably, the viral vector comprises a strong eukaryotic promoter operably linked to the polynucleotide e.g., a cytomegalovirus (CMV) promoter.

[0223] Additionally preferred vectors include viral vectors, fusion proteins and chemical conjugates. Retroviral vectors include Moloney murine leukemia viruses and HIV-based viruses. One preferred HIV-based viral vector comprises at least two vectors wherein the gag and pol genes are from an HIV genome and the env gene is from another virus. DNA viral vectors are preferred. These vectors include pox vectors such as orthopox or avipox vectors, herpesvirus vectors such as a herpes simplex I virus (HSV) vector [Geller, A.I. et al., (1995) J. Neurochem, 64: 487; Lim, F., et al., in DNA Cloning: Mammalian Systems, D. Glover, Ed. (Oxford Univ. Press, Oxford England) (1995); Geller, A.I. et al., (1993) Proc Natl. Acad. Sci.: U.S.A.:90 7603; Geller, A.I., et al., (1990) Proc Natl. Acad. Sci USA: 87:1149], Adenovirus Vectors (LeGal LaSalle et al., Science, 259:988 (1993); Davidson, et al., (1993) Nat. Genet. 3: 219; Yang, et al., (1995) J. Virol. 69: 2004) and Adeno-associated Virus Vectors (Kapliitt, M.G., et al., (1994) Nat. Genet. 8:148).

[0224] The antisense compounds of the disclosure encompass any pharmaceutically acceptable salts, esters, or salts of such esters, or any other compound which, upon administration to an animal, including a human, is capable of providing (directly or indirectly) the biologically active metabolite or residue thereof.

[0225] The term "pharmaceutically acceptable salts" refers to physiologically and pharmaceutically acceptable salts of the compounds of the disclosure: i.e., salts that retain the desired biological activity of the parent compound and do not impart undesired toxicological effects thereto. For oligonucleotides, preferred examples of pharmaceutically acceptable salts and their uses are further described in U.S. Pat. No. 6,287,860.

[0226] The present disclosure also includes pharmaceutical compositions and formulations that include the antisense compounds of the disclosure. The pharmaceutical compositions of the present disclosure may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic and to mucous membranes including vaginal and rectal delivery), pulmonary, e.g., by inhalation or insufflation of powders or aerosols, including by nebulizer; intratracheal, intranasal, epidermal and transdermal), oral or parenteral. Parenteral administration includes intravenous, intraarterial, subcutaneous, intraperitoneal or intramuscular injection or infusion; or intracranial, e.g., intrathecal or intraventricular, administration.

[0227] For treating tissues in the central nervous system, administration can be made by, e.g., injection or infusion into the cerebrospinal fluid. Administration of antisense RNA into cerebrospinal fluid is described, e.g., in U.S. Pat. App. Pub. No. 2007/0117772, "Methods for slowing familial ALS disease progression,".

[0228] When it is intended that the antisense oligonucleotide of the present disclosure be administered to cells in the central nervous system, administration can be with one or more agents capable of promoting penetration of the subject antisense oligonucleotide across the blood-brain barrier. Injection can be made, e.g., in the entorhinal cortex or hippocampus. Delivery of neurotrophic factors by administration of an adenovirus vector to motor neurons in muscle tissue is described in, e.g., U.S. Pat. No. 6,632,427, "Adenoviral-vector-mediated gene transfer into medullary motor neurons,". Delivery of vectors directly to the brain, e.g., the striatum, the thalamus, the hippocampus, or the substantia nigra, is known in the art and described, e.g., in U.S. Pat. No.

6,756,523, "Adenovirus vectors for the transfer of foreign genes into cells of the central nervous system particularly in brain,".

[0229] Administration can be rapid as by injection or made over a period of time as by slow infusion or administration of slow release formulations.

[0230] The subject antisense oligonucleotides can also be linked or conjugated with agents that provide desirable pharmaceutical or pharmacodynamic properties. For example, the antisense oligonucleotide can be coupled to any substance, known in the art to promote penetration or transport across the blood-brain barrier, such as an antibody to the transferrin receptor, and administered by intravenous injection. The antisense compound can be linked with a viral vector, for example, that makes the antisense compound more effective and/or increases the transport of the antisense compound across the blood-brain barrier. Osmotic blood brain barrier disruption can also be accomplished by, e.g., infusion of sugars including, but not limited to, meso erythritol, xylitol, D(+) galactose, D(+) lactose, D(+) xylose, dulcitol, myo-inositol, L(-) fructose, D(-) mannitol, D(+) glucose, D(+) arabinose, D(-) arabinose, cellobiose, D(+) maltose, D(+) raffinose, L(+) rhamnose, D(+) melibiose, D(-) ribose, adonitol, D(+) arabitol, L(-) arabitol, D(+) fucose, L(-) fucose, D(-) lyxose, L(+) lyxose, and L(-) lyxose, or amino acids including, but not limited to, glutamine, lysine, arginine, asparagine, aspartic acid, cysteine, glutamic acid, glycine, histidine, leucine, methionine, phenylalanine, proline, serine, threonine, tyrosine, valine, and taurine. Methods and materials for enhancing blood brain barrier penetration are described, e.g., in U. S. Patent No. 4,866,042, "Method for the delivery of genetic material across the blood brain barrier," 6,294,520, "Material for passage through the blood-brain barrier," and 6,936,589, "Parenteral delivery systems,".

[0231] The subject antisense compounds may be admixed, encapsulated, conjugated or otherwise associated with other molecules, molecule structures or mixtures of compounds, for example, liposomes, receptor-targeted molecules, oral, rectal, topical or other formulations, for assisting in uptake, distribution and/or absorption. For example, cationic lipids may be included in the formulation to facilitate oligonucleotide uptake. One such composition shown to facilitate uptake is LIPOFECTIN (available from GIBCO-BRL, Bethesda, MD).

[0232] Oligonucleotides with at least one 2'-O-methoxyethyl modification are believed to be particularly useful for oral administration. Pharmaceutical compositions and formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, suppositories, sprays, liquids and powders. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Coated condoms, gloves and the like may also be useful.

[0233] The pharmaceutical formulations of the present disclosure, which may conveniently be presented in unit dosage form, may be prepared according to conventional techniques well known in the pharmaceutical industry. Such techniques include the step of bringing into association the active ingredients with the pharmaceutical carrier(s) or excipient(s). In general, the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

[0234] The compositions of the present disclosure may be formulated into any of many possible dosage forms such as, but not limited to, tablets, capsules, gel capsules, liquid syrups, soft gels,

suppositories, and enemas. The compositions of the present disclosure may also be formulated as suspensions in aqueous, non-aqueous or mixed media. Aqueous suspensions may further contain substances that increase the viscosity of the suspension including, for example, sodium carboxymethylcellulose, sorbitol and/or dextran. The suspension may also contain stabilizers.

[0235] Pharmaceutical compositions of the present disclosure include, but are not limited to, solutions, emulsions, foams and liposome-containing formulations. The pharmaceutical compositions and formulations of the present disclosure may comprise one or more penetration enhancers, carriers, excipients or other active or inactive ingredients.

[0236] Emulsions are typically heterogeneous systems of one liquid dispersed in another in the form of droplets usually exceeding 0.1 μm in diameter. Emulsions may contain additional components in addition to the dispersed phases, and the active drug that may be present as a solution in either the aqueous phase, oily phase or itself as a separate phase. Microemulsions are included as an instance of the present disclosure. Emulsions and their uses are well known in the art and are further described in U.S. Pat. No. 6,287,860.

[0237] Formulations of the present disclosure include liposomal formulations. As used in the present disclosure, the term "liposome" means a vesicle composed of amphiphilic lipids arranged in a spherical bilayer or bilayers. Liposomes are unilamellar or multilamellar vesicles which have a membrane formed from a lipophilic material and an aqueous interior that contains the composition to be delivered. Cationic liposomes are positively charged liposomes that are believed to interact with negatively charged DNA molecules to form a stable complex. Liposomes that are pH-sensitive or negatively-charged are believed to entrap DNA rather than complex with it. Both cationic and noncationic liposomes have been used to deliver DNA to cells.

[0238] Liposomes also include "sterically stabilized" liposomes, a term which, as used herein, refers to liposomes comprising one or more specialized lipids. When incorporated into liposomes, these specialized lipids result in liposomes with enhanced circulation lifetimes relative to liposomes lacking such specialized lipids. Examples of sterically stabilized liposomes are those in which part of the vesicle-forming lipid portion of the liposome comprises one or more glycolipids or is derivatized with one or more hydrophilic polymers, such as a polyethylene glycol (PEG) moiety. Liposomes and their uses are further described in U.S. Pat. No. 6,287,860.

[0239] The pharmaceutical formulations and compositions of the present disclosure may also include surfactants. The use of surfactants in drug products, formulations and in emulsions is well known in the art. Surfactants and their uses are further described in U.S. Pat. No. 6,287,860.

[0240] In one instance, the present disclosure employs various penetration enhancers to effect the efficient delivery of nucleic acids, particularly oligonucleotides. In addition to aiding the diffusion of non-lipophilic drugs across cell membranes, penetration enhancers also enhance the permeability of lipophilic drugs. Penetration enhancers may be classified as belonging to one of five broad categories, i.e., surfactants, fatty acids, bile salts, chelating agents, and non chelating nonsurfactants. Penetration enhancers and their uses are further described in U.S. Pat. No. 6,287,860.

[0241] One of skill in the art will recognize that formulations are routinely designed according to their

intended use, i.e. route of administration.

[0242] Preferred formulations for topical administration include those in which the oligonucleotides of the disclosure are in admixture with a topical delivery agent such as lipids, liposomes, fatty acids, fatty acid esters, steroids, chelating agents and surfactants. Preferred lipids and liposomes include neutral (e.g. dioleoyl-phosphatidyl DOPE ethanolamine, dimyristoylphosphatidyl choline DMPC, distearoylphosphatidyl choline) negative (e.g. dimyristoylphosphatidyl glycerol DMPG) and cationic (e.g. dioleoyltetramethylaminopropyl DOTAP and dioleoyl-phosphatidyl ethanolamine DOTMA).

[0243] For topical or other administration, oligonucleotides of the disclosure may be encapsulated within liposomes or may form complexes thereto, in particular to cationic liposomes. Alternatively, oligonucleotides may be complexed to lipids, in particular to cationic lipids. Preferred fatty acids and esters, pharmaceutically acceptable salts thereof, and their uses are further described in U.S. Pat. No. 6,287,860.

[0244] Compositions and formulations for oral administration include powders or granules, microparticulates, nanoparticulates, suspensions or solutions in water or non-aqueous media, capsules, gel capsules, sachets, tablets or minitabets. Thickeners, flavoring agents, diluents, emulsifiers, dispersing aids or binders may be desirable. Preferred oral formulations are those in which oligonucleotides of the disclosure are administered in conjunction with one or more penetration enhancers surfactants and chelators. Preferred surfactants include fatty acids and/or esters or salts thereof, bile acids and/or salts thereof. Preferred bile acids/salts and fatty acids and their uses are further described in U.S. Pat. No. 6,287,860. Also preferred are combinations of penetration enhancers, for example, fatty acids/salts in combination with bile acids/salts. A particularly preferred combination is the sodium salt of lauric acid, capric acid and UDCA. Further penetration enhancers include polyoxyethylene-9-lauryl ether, polyoxyethylene-20-cetyl ether. Oligonucleotides of the disclosure may be delivered orally, in granular form including sprayed dried particles, or complexed to form micro or nanoparticles. Oligonucleotide complexing agents and their uses are further described in U.S. Pat. No. 6,287,860.

[0245] Compositions and formulations for parenteral, intrathecal or intraventricular administration may include sterile aqueous solutions that may also contain buffers, diluents and other suitable additives such as, but not limited to, penetration enhancers, carrier compounds and other pharmaceutically acceptable carriers or excipients.

[0246] Certain instances of the disclosure provide pharmaceutical compositions containing one or more oligomeric compounds and one or more other chemotherapeutic agents that function by a non-antisense mechanism. Examples of such chemotherapeutic agents include but are not limited to cancer chemotherapeutic drugs such as daunorubicin, daunomycin, dactinomycin, doxorubicin, epirubicin, idarubicin, esorubicin, bleomycin, mafosfamide, ifosfamide, cytosine arabinoside, bischloroethyl- nitrosurea, busulfan, mitomycin C, actinomycin D, mithramycin, prednisone, hydroxyprogesterone, testosterone, tamoxifen, dacarbazine, procarbazine, hexamethylmelamine, pentamethylmelamine, mitoxantrone, amsacrine, chlorambucil, methylcyclohexylnitrosurea, nitrogen mustards, melphalan, cyclophosphamide, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-azacytidine, hydroxyurea, deoxycytosine, 4-hydroxyperoxycyclo-phosphoramidate, 5-fluorouracil (5-FU), 5-fluorodeoxyuridine (5-FUdR), methotrexate (MTX), colchicine, taxol, vincristine, vinblastine, etoposide (VP-16), trimetrexate, irinotecan, topotecan, gemcitabine, teniposide, cisplatin and

diethylstilbestrol (DES). When used with the compounds of the disclosure, such chemotherapeutic agents may be used individually (e.g., 5-FU and oligonucleotide), sequentially (e.g., 5-FU and oligonucleotide for a period of time followed by MTX and oligonucleotide), or in combination with one or more other such chemotherapeutic agents (e.g., 5-FU, MTX and oligonucleotide, or 5-FU, radiotherapy and oligonucleotide). Anti-inflammatory drugs, including but not limited to nonsteroidal anti-inflammatory drugs and corticosteroids, and antiviral drugs, including but not limited to ribivirin, vidarabine, acyclovir and ganciclovir, may also be combined in compositions of the disclosure. Combinations of antisense compounds and other non-antisense drugs are also within the scope of this disclosure. Two or more combined compounds may be used together or sequentially.

[0247] In another related instance, compositions of the disclosure may contain one or more antisense compounds, particularly oligonucleotides, targeted to a first nucleic acid and one or more additional antisense compounds targeted to a second nucleic acid target. For example, the first target may be a particular antisense sequence of a Reprogramming factor, and the second target may be a region from another nucleotide sequence. Alternatively, compositions of the disclosure may contain two or more antisense compounds targeted to different regions of the same Reprogramming factor nucleic acid target. Numerous examples of antisense compounds are illustrated herein and others may be selected from among suitable compounds known in the art. Two or more combined compounds may be used together or sequentially.

Dosing:

[0248] The formulation of therapeutic compositions and their subsequent administration (dosing) is believed to be within the skill of those in the art. Dosing is dependent on severity and responsiveness of the disease state to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of the disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual oligonucleotides, and can generally be estimated based on EC₅₀s found to be effective in in vitro and in vivo animal models. In general, dosage is from 0.01 µg to 100 g per kg of body weight, and may be given once or more daily, weekly, monthly or yearly, or even once every 2 to 20 years. Persons of ordinary skill in the art can easily estimate repetition rates for dosing based on measured residence times and concentrations of the drug in bodily fluids or tissues. Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the oligonucleotide is administered in maintenance doses, ranging from 0.01 µg to 100 g per kg of body weight, once or more daily, to once every 20 years.

[0249] In instances, a patient is treated with a dosage of drug that is at least about 1, at least about 2, at least about 3, at least about 4, at least about 5, at least about 6, at least about 7, at least about 8, at least about 9, at least about 10, at least about 15, at least about 20, at least about 25, at least about 30, at least about 35, at least about 40, at least about 45, at least about 50, at least about 60, at least about 70, at least about 80, at least about 90, or at least about 100 mg/kg body weight. Certain injected dosages of antisense oligonucleotides are described, e.g., in U.S. Pat. No. 7,563,884, "Antisense modulation of PTP1B expression,".

[0250] While various instances of the present disclosure have been described above, it should be understood that they have been presented by way of example only, and not limitation. Numerous changes to the disclosed instances can be made in accordance with the disclosure herein without departing from the spirit or scope of the disclosure. Thus, the breadth and scope of the present disclosure should not be limited by any of the above described instances.

[0251] By their citation of various references in this document, Applicants do not admit any particular reference is "prior art" to their invention. Embodiments of inventive compositions and methods are illustrated in the following examples.

EXAMPLES

[0252] The following non-limiting Examples serve to illustrate selected embodiments of the invention. It will be appreciated that variations in proportions and alternatives in elements of the components shown will be apparent to those skilled in the art and are within the scope of instances of the present disclosure.

Example 1: Design of antisense oligonucleotides specific for a nucleic acid molecule antisense to a Reprogramming factor and/or a sense strand of a Reprogramming factor polynucleotide

[0253] As indicated above the term "oligonucleotide specific for" or "oligonucleotide targets" refers to an oligonucleotide having a sequence (i) capable of forming a stable complex with a portion of the targeted gene, or (ii) capable of forming a stable duplex with a portion of an mRNA transcript of the targeted gene.

[0254] Selection of appropriate oligonucleotides is facilitated by using computer programs that automatically align nucleic acid sequences and indicate regions of identity or homology. Such programs are used to compare nucleic acid sequences obtained, for example, by searching databases such as GenBank or by sequencing PCR products. Comparison of nucleic acid sequences from a range of species allows the selection of nucleic acid sequences that display an appropriate degree of identity between species. In the case of genes that have not been sequenced, Southern blots are performed to allow a determination of the degree of identity between genes in target species and other species. By performing Southern blots at varying degrees of stringency, as is well known in the art, it is possible to obtain an approximate measure of identity. These procedures allow the selection of oligonucleotides that exhibit a high degree of complementarity to target nucleic acid sequences in a subject to be controlled and a lower degree of complementarity to corresponding nucleic acid sequences in other species. One skilled in the art will realize that there is considerable latitude in selecting appropriate regions of genes for use in the present disclosure.

[0255] An antisense compound is "specifically hybridizable" when binding of the compound to the target nucleic acid interferes with the normal function of the target nucleic acid to cause a modulation of function and/or activity, and there is a sufficient degree of complementarity to avoid non-specific binding of the antisense compound to non-target nucleic acid sequences under conditions in which

specific binding is desired, i.e., under physiological conditions in the case of in vivo assays or therapeutic treatment, and under conditions in which assays are performed in the case of in vitro assays

[0256] The hybridization properties of the oligonucleotides described herein can be determined by one or more in vitro assays as known in the art. For example, the properties of the oligonucleotides described herein can be obtained by determination of binding strength between the target natural antisense and a potential drug molecules using melting curve assay.

[0257] The binding strength between the target natural antisense and a potential drug molecule (Molecule) can be estimated using any of the established methods of measuring the strength of intermolecular interactions, for example, a melting curve assay.

[0258] Melting curve assay determines the temperature at which a rapid transition from double-stranded to single-stranded conformation occurs for the natural antisense/Molecule complex. This temperature is widely accepted as a reliable measure of the interaction strength between the two molecules.

[0259] A melting curve assay can be performed using a cDNA copy of the actual natural antisense RNA molecule or a synthetic DNA or RNA nucleotide corresponding to the binding site of the Molecule. Multiple kits containing all necessary reagents to perform this assay are available (e.g. Applied Biosystems Inc. MeltDoctor kit). These kits include a suitable buffer solution containing one of the double strand DNA (dsDNA) binding dyes (such as ABI HRM dyes, SYBR Green, SYTO, etc.). The properties of the dsDNA dyes are such that they emit almost no fluorescence in free form, but are highly fluorescent when bound to dsDNA.

[0260] To perform the assay the cDNA or a corresponding oligonucleotide are mixed with Molecule in concentrations defined by the particular manufacturer's protocols. The mixture is heated to 95 °C to dissociate all pre-formed dsDNA complexes, then slowly cooled to room temperature or other lower temperature defined by the kit manufacturer to allow the DNA molecules to anneal. The newly formed complexes are then slowly heated to 95 °C with simultaneous continuous collection of data on the amount of fluorescence that is produced by the reaction. The fluorescence intensity is inversely proportional to the amounts of dsDNA present in the reaction. The data can be collected using a real time PCR instrument compatible with the kit (e.g. ABI's StepOne Plus Real Time PCR System or LightTyper instrument, Roche Diagnostics, Lewes, UK).

[0261] Melting peaks are constructed by plotting the negative derivative of fluorescence with respect to temperature ($-d(\text{Fluorescence})/dT$) on the y-axis) against temperature (x-axis) using appropriate software (for example LightTyper (Roche) or SDS Dissociation Curve, ABI). The data is analyzed to identify the temperature of the rapid transition from dsDNA complex to single strand molecules. This temperature is called T_m and is directly proportional to the strength of interaction between the two molecules. Typically, T_m will exceed 40 °C.

Example 2: Modulation of a Reprogramming factor polynucleotide Treatment of HepG2 cells with antisense oligonucleotides

[0262] HepG2 cells from ATCC (cat# HB-8065) were grown in growth media (MEM/EBSS (Hyclone cat #SH30024, or Mediatech cat # MT-10-010-CV) +10% FBS (Mediatech cat# MT35- 011-CV)+ penicillin/streptomycin (Mediatech cat# MT30-002-CI)) at 37°C and 5% CO₂. One day before the experiment the cells were replated at the density of 1.5×10^5 /ml into 6 well plates and incubated at 37°C and 5% CO₂. On the day of the experiment the media in the 6 well plates was changed to fresh growth media. All antisense oligonucleotides were diluted to the concentration of 20 µM. Two µl of this solution was incubated with 400 µl of Opti-MEM media (Gibco cat#31985-070) and 4 µl of Lipofectamine 2000 (Invitrogen cat# 11668019) at room temperature for 20 min and applied to each well of the 6 well plates with HepG2 cells. A Similar mixture including 2 µl of water instead of the oligonucleotide solution was used for the mock-transfected controls. After 3-18 h of incubation at 37°C and 5% CO₂ the media was changed to fresh growth media. 48 h after addition of antisense oligonucleotides the media was removed and RNA was extracted from the cells using SV Total RNA Isolation System from Promega (cat # Z3105) or RNeasy Total RNA Isolation kit from Qiagen (cat# 74181) following the manufacturers' instructions. 600 ng of RNA was added to the reverse transcription reaction performed using Verso cDNA kit from Thermo Scientific (cat#AB1453B) or High Capacity cDNA Reverse Transcription Kit (cat# 4368813) as described in the manufacturer's protocol. The cDNA from this reverse transcription reaction was used to monitor gene expression by real time PCR using ABI Taqman Gene Expression Mix (cat#4369510) and primers/probes designed by ABI (Applied Biosystems Taqman Gene Expression Assay: Hs00602736_s1, Hs00358836_m1, Hs03005111_g1 by Applied Biosystems Inc., Foster City CA). The following PCR cycle was used: 50°C for 2 min, 95°C for 10 min, 40 cycles of (95°C for 15 seconds, 60°C for 1 min) using Mx4000 thermal cycler (Stratagene).

[0263] Fold change in gene expression after treatment with antisense oligonucleotides was calculated based on the difference in 18S-normalized dCt values between treated and mock-transfected samples.

Results

[0264] Real time PCR results show that the levels of SOX2 mRNA in HepG2 cells are significantly increased 48 h after treatment with two of the siRNAs designed to SOX2 antisense ai885646 (Fig 1A).

[0265] Real time PCR results show that the levels of KLF4 mRNA in HepG2 cells are significantly increased 48 h after treatment with two of the oligos designed to KLF4 antisense. (Fig 1B).

[0266] Real time PCR results show that the levels of POU5F1 mRNA are significantly increased in HepG2 cells 48 h after treatment with some of the antisense oligonucleotides to POU5F1 natural antisense AI926793 (Fig 1C).

[0267] In addition, while a particular feature of the invention may have been disclosed with respect to only one of several implementations, such feature may be combined with one or more other features of the other implementations as may be desired and advantageous for any given or particular application.

[0268] The Abstract of the disclosure will allow the reader to quickly ascertain the nature of the technical disclosure. It is submitted with the understanding that it will not be used to interpret or limit the scope or meaning of the following claims.

SEQUENCE LISTING

[0269]

<110> CuRNA, Inc.

<120> TREATMENT OF REPROGRAMMING FACTOR RELATED DISEASES BY INHIBITION OF NATURAL ANTISENSE TRANSCRIPT TO A REPROGRAMMING FACTOR

<130> SOX2

<150> 61/179,056

<151> 2009-05-18

<150> 61/286,852

<151> 2009-12-16

<150> 61/233,996

<151> 2009-08-14

<160> 17

<170> PatentIn version 3.5

<210> 1

<211> 2518

<212> DNA

<213> Homo sapiens

<400> 1

ctattaactt gttcaaaaaa gtatcaggag ttgtcaaggc agagaagaga gtgtttgcaa	60
aagggggaaa gtagtttgc tgcctctttaa gactaggact gagagaaaga agaggagaga	120
gaaagaaagg gagagaagtt tgagcccccag gcttaagcct ttccaaaaa taataataac	180
aatcatcggc ggccggcagga tcggccagag gaggagggaa gcgctttttt tgatcctgat	240
tccagtttgc ctctctcttt ttttccccc aattattctt cgcctgattt tctctcgcgga	300
gccctgcgct cccgacaccc ccgcccgct cccctcctcc tctccccccg ccgcggggcc	360
ccccaaagtc ccggccgggc cgagggtcgg cggccgccgg cgggcccgggc ccgcgcacag	420
cgcccgcatg tacaacatga tggagacgga gctgaagccg ccgggcccgc agcaaacttc	480
ggggggcggc ggccggcaact ccaccgcggc ggccggccgg gcgaaccaga aaaacagccc	540
ggaccgcgtc aagcggccca tgaatgcctt catggtgtgg tcccgcgggc agcggcgcaa	600
gatggcccag gagaacccca agatgcacaa ctccgagatc agcaagcgcc tgggcgccga	660
gtggaaactt ttgtcggaga cggagaagcg gccgttcac gacgaggcta agcggctgcg	720
agcgtgcac atgaaggagc acccggatta taaataccgg ccccgccgga aaaccaagac	780

```

gctcatgaag aaggataagt acacgctgcc cggcgggctg ctggcccccg ggggcaatag      840
catggcgagc ggggtcgggg tgggcgccgg cctgggcgcg ggcggaacc agcgcatgga      900
cagttacgcg cacatgaacg gctggagcaa cggcagctac agcatgatgc aggaccagct      960
gggctacccg cagcaccgcg gectcaatgc gcacggcgca gcgcagatgc agcccatgca     1020
ccgctacgac gtgagcgccc tgcagtacaa ctccatgacc agctcgcaga cctacatgaa     1080
cggctcgccc acctacagca tgtcctactc gcagcagggc acccctggca tggctcttgg     1140

ctccatgggt teggtggtca agtccgaggg cagctccagc cccctgtgg ttacctcttc     1200
ctcccactcc agggcgccct gccaggccgg ggacctccgg gacatgatca gcatgtatct     1260
ccccggcgcc gaggtgccgg aacccgccgc cccagcaga cttcacatgt cccagcacta     1320
ccagagcgggc cgggtgcccg gcacggccat taacggcaca ctgcccctct cacacatgtg     1380
agggcgggac agcgaactgg aggggggaga aattttcaaa gaaaaacgag ggaaatggga     1440
ggggtgcaaa agaggagagt aagaaacagc atggagaaaa cccggtacgc tcaaaaagaa     1500
aaaggaaaaa aaaaaatccc atcaccacaa gcaaatgaca gctgcaaaag agaaccacaa     1560
tcccatccac actcacgcaa aaaccgcgat gccgacaaga aaacttttat gagagagatc     1620
ctggacttct ttttggggga ctatttttgt acagagaaaa cctggggagg gtggggaggg     1680
cgggggaatg gaccttgtat agatctggag gaaagaaagc tacgaaaaac tttttaaaag     1740
ttctagtggg acggtaggag ctttgaggga agtttgcaaa agtctttacc aataatattt     1800
agagctagtc tccaagcgac gaaaaaatg ttttaatat tgcaagcaac ttttgtacag     1860
tatttatcga gataaacatg gcaatcaaaa tgtccattgt ttataagctg agaatttgcc     1920
aatatttttc aaggagaggc ttcttgctga attttgattc tgcagctgaa atttaggaca     1980
gttgcaaacg tgaaaagaag aaaattattc aaatttggac attttaattg tttaaaaatt     2040
gtacaaaagg aaaaaattag aataagtact ggcgaaacct ctctgtggtc ttgtttaaaa     2100
agggcaaaag ttttagactg tactaaattt tataacttac tgttaaaagc aaaaatggcc     2160
atgcaggttg acaccgttgg taatttataa tagcttttgt tcgatcccaa ctttccattt     2220
tgttcagata aaaaaaacca tgaaattact gtgtttgaaa tattttctta tggtttgtaa     2280
tattttctga aatttattgt gatattttta gggtttcccc cttttatttt ccgtagttgt     2340
attttaaaag attcggctct gtattatttg aatcagctct ccgagaatcc atgtatatat     2400
ttgaactaat atcatcctta taacaggtag attttcaact taagttttta ctccattatg     2460
cacagtttga gataaataaa tttttgaaat atggacactg aaaaaaaaaa aaaaaaaa     2518

```

<210> 2

<211> 2949

<212> DNA

<213> Homo sapiens

<400> 2

```

agtttcccg cagagagaa cgaacgtgtc tgcgggcgcg cggggagcag aggcggtggc      60
gggcggcggc ggcaccggga gccgccgagt gacctcccc cgccctctg gccccacc     120
ctcccacccg ccggtggccc gcgcccatgg ccgcgcgcgc tcacacacaa tcaccggagt     180
ccgcgccttg cgccgcgac cagttcgcag ctccgcgcca cggcagccag tctcacctgg     240
cggcacccgc cgccaccgc cccggccaca gccctgcgc ccacggcagc actcgaggcg     300

```

```

-- - - - - - - - - - -
accgcgacag tgggtggggga cgctgctgag tggaagagag cgcagccggg ccaccggacc 360

tacttactcg ccttgctgat tgtctatddd tgcgtttaca acttttctaa gaacttttgt 420
atacaaagga actttttaaa aaagacgctt ccaagttata tttaatccaa agaagaagga 480
tctcggccaa tttgggggtt tgggttttgg ctctgcttct tctcttcgtt gactttgggg 540
ttcaggtgcc ccagctgctt cgggctgccg aggaccttct gggcccccac attaatgagg 600
cagccacctg gcgagtctga catggctgtc agcgacgcgc tgcctccatc tttctccacg 660
ttcgcgtctg gcccgggcggg aaggagagaag acactgcgtc aagcagggtc cccgaataac 720
cgctggcggg aggagctctc ccacatgaag cgacttcccc cagtgccttc cggccgcccc 780
tatgacctgg cggggcgac cgtggccaca gacctggaga gggcgggagc cggtgccgct 840
tgcggcggtg gcaacctggc gccctacct cggagagaga ccgaggagt caacgatctc 900
ctggacctgg actttatct ctccaattcg ctgacctc ctccggagtc agtgccgcc 960
accgtgtcct cgtcagcgtc agcctcctct tctgtctgc cgtcagcag cggccctgcc 1020
agcgcgccct ccacctgcag ctccacctat ccgatccggg ccgggaacga cccgggcgtg 1080
gcgcggggcg gcacggggcg aggcctctc tatggcaggg agtccgctcc cctccgacg 1140
gctcccttca acctggcgga catcaacgac gtgagccct cggggcggtt cgtggccgag 1200
ctcctgcggc cagaattgga cccggtgtac attccgcgc agcagccga gccgccagt 1260
ggcgggctga tgggcaagt cgtgtgaag gctgtctga gcgcctctgg cagcgagtac 1320
ggcagcccgt cgtcatcag cgtcagcaaa ggcagccct acggcagcca cccggtggtg 1380
gtggcgccct acaacggcg gcgcgcgc acgtgcccc agatcaagca ggagcggtc 1440
tctcgtgca ccaactggg cgctggacct cctctcagca atggccacc gccggtgca 1500
cacgacttcc cctggggcg gcagctcccc agcaggacta ccccgacct ggtcttgag 1560
gaagtgtga gcagcaggga ctgtaccct gccctgcgc ttcctcccg ctccatccc 1620
caccggggc ccaattacc atccttctg ccgatcaga tgcagccga agtccgcgc 1680
ctccattacc aagagctcat gccaccggg tctgcatgc cagaggagcc caagccaaag 1740
agggaagac gatcgtggc ccggaagg accgccccc acacttgtga ttacggggc 1800
tgcgcaaaa cctacacaaa gagttccat ctcaaggcac acctgcgaac ccacacagt 1860
gagaaacctt acctgtga ctgggacggc tgtggatgga aattcgccc ctcatgaa 1920
ctgaccaggc actaccgta acacacggg caccgccgt tccagtcca aaaatgcgac 1980
cgagcatttt ccaggtcgg ccacctgcc ttacacatga agaggcattt ttaaatccca 2040
gacagtggat atgaccaca ctgccagaag agaattcagt attttttact tttcacactg 2100
tcttccgat gagggaagga gccagccag aaagcactac aatcatggtc agttcccaa 2160
ctgagtcac ttgtgagtg ataacagga aaaatgagga atccaaaaga caaaaatcaa 2220
agaacagatg ggtctgtga ctggatctt tatcattcca attctaaatc cgaattgaat 2280

attcctggac ttacaaaatg ccaagggggg gactggaagt tgtggatac agggataaaa 2340
ttatatccgt gagttggggg agggaagacc agaattccct tgaatttgt attgatcaa 2400
tataagcata aaagatcacc ttgtattct tttacctctt aaaagccatt attatgatgt 2460
tagaagaaga ggaagaaatt caggtacaga aaacatgttt aaatagccta aatgatggtg 2520
cttggtgagt cttggttcta aaggtaccaa acaagggaag caaagttttc aaactgctgc 2580
atactttgac aaggaaaatc tataattgtc ttccgatcaa ctttatgac ctaagttagg 2640
taatatacct attttacttc tttaacattt ttatacagac actctattat acactatatt 2700

```

```

-----
ttcagatgtg caataatttg tacaatggtt tattccaag tatgccttaa gcagaacaaa 2760
tgtgtttttc tatatagttc cttgccttaa taaatatgta atataaattt aagcaaacgt 2820
ctatttttga tattttgtaa ctacaaagta aaatgaacat tttgtggagt ttgtattttg 2880
catactcaag gtgagaatta agtttttaaat aaacctataa tatttttatct gaaaaaaaaa 2940
aaaaaaaaa 2949

```

<210> 3

<211> 1411

<212> DNA

<213> Homo sapiens

<400> 3

```

ccttcgcaag ccttcatttc accaggcccc cggtctgggg cgccttcctt ccccatggcg 60
ggacacctgg cttecgattt cgccttcctc cccctccag gtggtggagg tgatgggcca 120
ggggggccgg agccgggctg ggttgatcct cggacctggc taagcttcca aggccctcct 180
ggaggggccag gaatcgggcc ggggggtggg ccaggctctg aggtgtgggg gattccccca 240
tgccccccgc cgtatgagtt ctgtgggggg atggcgtact gtgggccccg ggttgagtg 300
gggctagtgc cccaaggcgg cttggagacc tctcagcctg agggcgaaagc aggagtgggg 360
gtggagagca actccgatgg ggcctccccg gagcctgca ccgtcacccc tggtgccgtg 420
aagctggaga aggagaagct ggagcaaac ccggaggagt ccaggacat caaagctctg 480
cagaaagaac tcgagcaatt tgccaagctc ctgaagcaga agaggatcac cctgggatat 540
acacaggccg atgtggggct caccctgggg gttctatttg ggaaggattt cagccaaacg 600
accatctgcc gctttgaggc tctgcagctt agcttcaaga acatgtgtaa gctgcggccc 660
ttgctgcaga agtgggtgga ggaagctgac aacaatgaaa atcttcagga gatatgcaa 720
gcagaaaccc tcgtgcaggc ccgaaagaga aagcgaacca gtatcgagaa ccgagtgaga 780
ggcaacctgg agaatttggt cctgcagtgc ccgaaaccca cactgcagca gatcagccac 840
atcgcccagc agcttgggct cgagaaggat gtgggtccgag tgtgggtctg taaccggcgc 900
cagaagggca agcgatcaag cagcgactat gcacaacgag aggatthtga ggctgctggg 960
tctcctttct cagggggacc agtgtccttt cctctggccc cagggcccca ttttggtagc 1020
ccaggetatg ggagccctca cttcactgca ctgtactcct cggtcctttt ccctgagggg 1080
gaagcctttc cccctgtctc cgtcaccact ctgggctctc ccatgcattc aaactgaggt 1140
gcctgccctt ctaggaatgg gggacagggg gaggggagga gctagggaaa gaaaacctgg 1200
agtttgtgcc agggtttttg ggattaagtt cttcattcac taaggaagga attgggaaca 1260
caaagggtgg gggcagggga gtttggggca actggttggg gggaaggtga agttcaatga 1320
tgctcttgat tttaatccca catcatgtat cacttttttc ttaaataaag aagcctggga 1380
cacagtagat agacacactt aaaaaaaaaa a 1411

```

<210> 4

<211> 993

<212> DNA

<213> Homo sapiens

<220>

<221> misc_feature

<222> (956)..(956)

<223> n is a, c, g, t or u

<220>

<221> misc_feature

<222> (968)..(968)

<223> n is a, c, g, t or u

<400> 4

```

ccaggactta tcagctggga taggcctcac ttacaagaca gctctgttca gtatttggaa      60
gaaagcctga caggtttctt cggaaaagtg gccatccatg gaatgaatga aatgttctct      120
ttctattcca gggattgcag tggcaaagct aggctaggtc ttggaggctg gtgtaaggcg      180
atgtgggtga aggcaggagg ctgatggaaa gactgggggg aagaaaagcc gaaatggatt      240
cacggtgcct tggatgaagg acgagagggg aactgcaagc tccttcaact ggttctgtcc      300
ggtgagaagt gatcaagctt gggctgacaa gaggctcagg gagccctcac gttctttcgc      360
ttttttacct gccaatcaaa ctgctacaag acaacaccct gatctggcat ggacatgtcc      420
gtgtttacct taatattaca cctggattaa acagaaaaca gtgctgtatt ttgaagagcg      480
ccgaatatgt atgcattttg agaaacccaa tctcaccctt tccgggttcc caagaactaa      540
aacaagccat aacttgagag aaaaaggaga accttcgggg ggcaggaagg ttgattggaa      600
ataacttaag gaaagtctgc agaattcttt tttttacaac ttttctgagt ttccagtggg      660
tatatttagt gtgagtttga cagtaacagg ctagggaggg cagagattgg agaaattggg      720
ggtcggggga gtgattatgg gaagaagggt agtaaggaaac aaaacaatgc accgttttgt      780
aaagataata aatggaacgt ggctggtaga tactattcag tacattttct tagggtgagt      840
aagggtagac caggggagga gggggcggag agagtgttac agaagaaaga aaataagtaa      900
ccctgatggt ttaagccctt tataaaaaga aatggcatca ggtttttttt ctttantccc      960
ccaccancc tttgtagtca agtgcattha aag                                     993

```

<210> 5

<211> 493

<212> DNA

<213> Homo sapiens

<400> 5

```

aaaaagaagg ctgtaagctc agacaacaag tagcaaaaga aacactgacc actcacaggc      60
tggggattca gaacctgcgg atgtggaggg ccagagggct gaatttgcaa gttgggcaat      120
ggggaaagcc ttcacaaagg agaaagcctt tgggctggaa ttggaggagt aggagtcaat      180
agagagagaa gagagtaaag ggcaaggcca tcattccagg tttagggagc aagtacaaag      240
acacagagaa atggaaggct tggcctgttt ggagaatgtc aagctgtctg gggttgagag      300
aaagacattc taacaaggct tccattaaag aggctagatt tgctgcttgg acattttcta      360
gctcttctga gacaactttt tgttgagtta agcgaaaatg aatgaaaatc agatggacca      420
tatttctctt gaatgcacac gggcttaaaa acgaagggga aaaagaactt gaccagaatg      480
tgtcttggct ttt                                     493

```

<210> 6
 <211> 418
 <212> DNA
 <213> Homo sapiens

<220>
 <221> misc_feature
 <222> (359)..(359)
 <223> n is a, c, g, t or u

<400> 6
 ttaagacagg gtctcacttt gttgcccaaga ctggagtgca gtggagtaca atgatggctc 60
 aatgtagcct cgatctactg ggccaaagca atccttctgc tccagcctcc taagtggctg 120
 ggaaccacagg cttggaccac tgtgccctgt tagttttttt attttttagta gagatggggc 180
 cctgccatgt taccagagct ggtcttgaac tctgtctctc aagaaatcct cccgcctctg 240
 ccgcccagtg tcatgattaa aggcgtgagc caccacaccc aactttcaac tccaaccocg 300
 ctccctggca ctctctcagg ctctgcacat cccagctgtc tggaatcact ccacacctnc 360
 atgtttcttca ggaaccacagg tgcttgaccc cctctcccag accttggcac tgtgcctt 418

<210> 7
 <211> 27
 <212> DNA
 <213> Homo sapiens

<400> 7
 auaauaaaug gaacguggcu gguagau 27

<210> 8
 <211> 27
 <212> DNA
 <213> Homo sapiens

<400> 8
 cugaguuuucc aguggguaua uuuagug 27

<210> 9
 <211> 19
 <212> DNA
 <213> Homo sapiens

<400> 9
 tgagtggta gtgtttctt 19

<210> 10
 <211> 21
 <212> DNA
 <213> Homo sapiens

<400> 10

gtgtctttgt acttgctccc t 21

<210> 11

<211> 21

<212> DNA

<213> Homo sapiens

<400> 11

ccctttactc tcttctctc c 21

<210> 12

<211> 19

<212> DNA

<213> Homo sapiens

<400> 12

ttgctacttg ttgtctgag 19

<210> 13

<211> 19

<212> DNA

<213> Homo sapiens

<400> 13

gccatcattg tactccact 19

<210> 14

<211> 19

<212> DNA

<213> Homo sapiens

<400> 14

agttgggtgt ggtggctca 19

<210> 15

<211> 20

<212> DNA

<213> Homo sapiens

<400> 15

tggtcccagc cacttaggag 20

<210> 16

<211> 21

<212> DNA

<213> Homo sapiens

<400> 16

ggcgggagga ttcttgagg a 21

<210> 17

<211> 20
 <212> DNA
 <213> Homo sapiens

 <220>
 <221> misc_feature
 <222> (4)..(4)
 <223> n is a, c, g, t or u

 <400> 17
 atgnaggtgt ggagtgattc 20

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description

- [US61179056A \[0001\]](#)
- [US61233996B \[0001\]](#)
- [US61286852A \[0001\]](#)
- [US2009068742A \[0003\]](#)
- [US5432272A \[0039\] \[0196\]](#)
- [US5013830A \[0177\]](#)
- [US5149797A \[0177\]](#)
- [US5220007A \[0177\]](#)
- [US5256775A \[0177\]](#)
- [US5366878A \[0177\]](#)
- [US5403711A \[0177\]](#)
- [US5491133A \[0177\]](#)
- [US5565350A \[0177\]](#)
- [US5623065A \[0177\]](#)
- [US5652355A \[0177\]](#)
- [US5652356A \[0177\]](#)
- [US5700922A \[0177\]](#)
- [US5034506A \[0179\] \[0189\] \[0191\]](#)
- [US5138045A \[0181\] \[0199\] \[0218\]](#)
- [US5218105A \[0181\] \[0199\] \[0218\]](#)
- [US5459255A \[0181\] \[0196\]](#)
- [US3687808A \[0187\] \[0195\] \[0196\]](#)

- [US4469863A](#) [0187]
- [US4476301A](#) [0187]
- [US5023243A](#) [0187]
- [US5177196A](#) [0187]
- [US5188897A](#) [0187]
- [US5264423A](#) [0187]
- [US5276019A](#) [0187]
- [US5278302A](#) [0187]
- [US5286717A](#) [0187]
- [US5321131A](#) [0187]
- [US5399676A](#) [0187]
- [US5405939A](#) [0187]
- [US5453496A](#) [0187]
- [US5455233A](#) [0187]
- [US5466677A](#) [0187] [0189]
- [US5476925A](#) [0187]
- [US5519126A](#) [0187]
- [US5536821A](#) [0187]
- [US5541306A](#) [0187]
- [US5550111A](#) [0187]
- [US5563253A](#) [0187]
- [US5571799A](#) [0187]
- [US5587361A](#) [0187]
- [US5625050A](#) [0187]
- [US5166315A](#) [0189]
- [US5185444A](#) [0189]
- [US5214134A](#) [0189]
- [US5216141A](#) [0189]
- [US5235033A](#) [0189]
- [US5264562A](#) [0189]
- [US5264564A](#) [0189]
- [US5405938A](#) [0189]
- [US5434257A](#) [0189]
- [US5470967A](#) [0189]
- [US5489677A](#) [0189] [0191]
- [US5541307A](#) [0189]
- [US5561225A](#) [0189]
- [US5596086A](#) [0189]
- [US5602240A](#) [0189] [0189] [0191]
- [US5610289A](#) [0189] [0189]
- [US5608046A](#) [0189] [0199] [0218]
- [US5618704A](#) [0189]
- [US5623070A](#) [0189]
- [US5663312A](#) [0189]
- [US5633360A](#) [0189]
- [US5677437A](#) [0189]
- [US5677439A](#) [0189]

- [US5539082A](#) [0190]
- [US5714331A](#) [0190]
- [US5719262A](#) [0190]
- [US4981957A](#) [0193]
- [US5118800A](#) [0193]
- [US5319080A](#) [0193]
- [US5359044A](#) [0193]
- [US5393878A](#) [0193]
- [US5446137A](#) [0193]
- [US5466786A](#) [0193]
- [US5514785A](#) [0193] [0199] [0218]
- [US5519134A](#) [0193]
- [US5567811A](#) [0193]
- [US5576427A](#) [0193]
- [US5591722A](#) [0193]
- [US5597909A](#) [0193]
- [US5610300A](#) [0193]
- [US5627053A](#) [0193]
- [US5639873A](#) [0193]
- [US5646265A](#) [0193]
- [US5658873A](#) [0193]
- [US5670633A](#) [0193]
- [US5700920A](#) [0193]
- [US4845205A](#) [0196]
- [US5130302A](#) [0196]
- [US5134066A](#) [0196]
- [US5175273A](#) [0196]
- [US5367066A](#) [0196]
- [US5457187A](#) [0196]
- [US5484908A](#) [0196]
- [US5502177A](#) [0196]
- [US5525711A](#) [0196]
- [US5552540A](#) [0196]
- [US5587469A](#) [0196]
- [US5596091A](#) [0196]
- [US5614617A](#) [0196]
- [US5750692A](#) [0196]
- [US5681941A](#) [0196]
- [US4828979A](#) [0199] [0218]
- [US4948882A](#) [0199] [0218]
- [US5525465A](#) [0199] [0218]
- [US5541313A](#) [0199] [0218]
- [US5545730A](#) [0199] [0218]
- [US5552A](#) [0199]
- [US538A](#) [0199]
- [US5578717A](#) [0199] [0218]
- [US5580731A](#) [0199] [0199] [0218] [0218]

- [US5591584A](#) [0199] [0218]
- [US5109124A](#) [0199] [0218]
- [US5118802A](#) [0199] [0218]
- [US5414077A](#) [0199] [0218]
- [US5486603A](#) [0199] [0218]
- [US5512439A](#) [0199] [0218]
- [US5578718A](#) [0199] [0218]
- [US4587044A](#) [0199] [0218]
- [US4605735A](#) [0199] [0218]
- [US4667025A](#) [0199] [0218]
- [US4762779A](#) [0199] [0218]
- [US4789737A](#) [0199] [0218]
- [US4824941A](#) [0199] [0218]
- [US4835263A](#) [0199] [0218]
- [US4876335A](#) [0199] [0218]
- [US4904582A](#) [0199] [0218]
- [US4958013A](#) [0199] [0218]
- [US5082830A](#) [0199] [0199] [0218] [0218]
- [US5112963A](#) [0199] [0199] [0218] [0218]
- [US5214136A](#) [0199] [0199] [0218] [0218]
- [US5245022A](#) [0199] [0218]
- [US5254469A](#) [0199] [0218]
- [US5258506A](#) [0199] [0218]
- [US5262536A](#) [0199] [0218]
- [US5272250A](#) [0199] [0218]
- [US5292873A](#) [0199] [0218]
- [US5317098A](#) [0199] [0218]
- [US5371241A](#) [0199] [0218]
- [US5391723A](#) [0199] [0218]
- [US5416203A](#) [0199] [0218]
- [US5451463A](#) [0199] [0218]
- [US5510475A](#) [0199] [0218]
- [US5512667A](#) [0199] [0218]
- [US5565552A](#) [0199] [0218]
- [US5567810A](#) [0199] [0218]
- [US5574142A](#) [0199] [0218]
- [US5585481A](#) [0199] [0218]
- [US5587371A](#) [0199] [0218]
- [US5595726A](#) [0199] [0218]
- [US5597696A](#) [0199] [0218]
- [US5599923A](#) [0199] [0218]
- [US5599928A](#) [0199] [0218]
- [US5688941A](#) [0199] [0218]
- [USLTS9209196W](#) [0217]
- [US6287860B](#) [0217] [0225] [0236] [0238] [0239] [0240] [0243] [0244] [0244]
- [US5552538A](#) [0218]
- [US5108921A](#) [0219] [0219]

- [US5354844A \[0219\]](#)
- [US5416016A \[0219\] \[0219\]](#)
- [US5459127A \[0219\]](#)
- [US5521291A \[0219\]](#)
- [US5543165A \[0219\]](#)
- [US5547932A \[0219\]](#)
- [US5583020A \[0219\]](#)
- [US5591721A \[0219\]](#)
- [US4426330A \[0219\]](#)
- [US4534899A \[0219\]](#)
- [US5013556A \[0219\]](#)
- [US5213804A \[0219\]](#)
- [US5227170A \[0219\]](#)
- [US5264221A \[0219\]](#)
- [US5356633A \[0219\]](#)
- [US5395619A \[0219\]](#)
- [US5417978A \[0219\]](#)
- [US5462854A \[0219\]](#)
- [US5469854A \[0219\]](#)
- [US5512295A \[0219\]](#)
- [US5527528A \[0219\]](#)
- [US5534259A \[0219\]](#)
- [US5543152A \[0219\]](#)
- [US5556948A \[0219\]](#)
- [US5580575A \[0219\]](#)
- [US5595756A \[0219\]](#)
- [US907603A \[0223\]](#)
- [US20070117772A \[0227\]](#)
- [US6632427B \[0228\]](#)
- [US6756523B \[0228\]](#)
- [US4866042A \[0230\]](#)
- [US6294520B \[0230\]](#)
- [US6936589B \[0230\]](#)
- [US7563884B \[0249\]](#)
- [WO61179056A \[0269\]](#)
- [WO61286852A \[0269\]](#)
- [WO61233996A \[0269\]](#)

Non-patent literature cited in the description

- **XU et al.**Cell, 2009, vol. 137, 4647-658 [\[0003\]](#)
- **RICHARDS et al.**Stem Cells, 2006, vol. 24, 51162-1173 [\[0003\]](#)
- **TAKAHASHI et al.**Cell, 2006, vol. 126, 4663-676 [\[0003\]](#)

- **TAKAHASHI et al.**Cell, 2007, vol. 131, 5861-872 [0003]
- **SACHDEVA et al.**PNAS, 2009, vol. 106, 93207-3212 [0003]
- **MAHERALI et al.**Cell Stem Cell, 2007, vol. 1, 55-7 [0003]
- **CHAMBERLAIN et al.**Neurogenetics, 2008, vol. 9, 4227-235 [0003]
- **LAVORGNA et al.**Trends in Biochem. Sci., 2004, vol. 29, 288-94 [0003]
- **LEHNER et al.**Trends in Genetics, 2002, vol. 18, 263-65 [0003]
- **EGUCHI et al.**Ann. Rev. Biochem., 1991, vol. 60, 631-652 [0024]
- **CAPLEN, N. J. et al.**Proc. Natl. Acad. Sci. USA, 2001, vol. 98, 9742-9747 [0034]
- **BERNSTEIN, E. et al.**Nature, 2001, vol. 409, 363-366 [0034] [0034]
- **BOUTLA, A. et al.**Curr. Biol., 2001, vol. 11, 1776-1780 [0034]
- **CECHJ.**American. Med. Assoc., 1988, vol. 260, 3030-3035 [0036]
- **SULLENGER et al.**Cell, 1990, vol. 63, 601-608 [0037]
- **KOMBERGBAKER**DNA ReplicationFreeman19920000 [0039]
- **SCHEIT**Nucleotide AnalogsJohn Wiley19800000 [0040]
- **FREIERALTMANN**Nucl. Acid. Res., 1997, vol. 25, 224429-4443 [0040]
- **TOULMÉ, J. J.**Nature Biotechnology, 2001, vol. 19, 17-18 [0040]
- **MANOHARAN M.**Biochemica et Biophysica Acta, 1999, vol. 1489, 117-139 [0040]
- **FREIER S. M.**Nucleic Acid Research, 1997, vol. 25, 4429-4443 [0040]
- **UHLMAN, E.**Drug Discovery & Development, 2000, vol. 3, 203-213 [0040]
- **HERDEWIN P.**Antisense & Nucleic Acid Drug Dev., 2000, vol. 10, 297-310 [0040]
- **N. K CHRISTIENSEN. et al.**J. Am. Chem. Soc., 1998, vol. 120, 5458-5463 [0040]
- **PRAKASH TPBHAT B.**Curr Top Med Chem., 2007, vol. 7, 7641-9 [0040]
- **CHO EJ et al.**Annual Review of Analytical Chemistry, 2009, vol. 2, 241-264 [0040]
- **ALTSCHUL et al.**J. Mol. Biol., 1990, vol. 215, 403-410 [0045]
- **ZHANGMADDEN**Genome Res., 1997, vol. 7, 649-656 [0045]
- **SMITHWATERMAN**Adv. Appl. Math., 1981, vol. 2, 482-489 [0045]
- **RYAN AK et al.**Genes Dev, 1997, vol. 11, 1207-1225 [0074]
- **RUVKUN G et al.**Cell, 1991, vol. 64, 475-478 [0074]
- **HERR W et al.**Genes Dev, 1995, vol. 9, 1679-1693 [0074]
- **SCHOLER HR et al.**Nature, 1990, vol. 344, 435-9 [0074]
- **BOIANI M et al.**Nat Rev Mol Cell Biol., 2005, vol. 6, 872-84 [0075]
- **BOYER LA et al.**Cell, 2005, vol. 122, 947-56 [0075]
- **TAKEDA J et al.**Nucleic Acids Res, 1992, vol. 20, 4613-4620 [0076]
- **THOMSON, J. A. et al.**Science, 1998, vol. 282, 11451147- [0088]
- **COWAN, C. A. et al.**N. Engl. J. Med., 2004, vol. 350, 1353-1356 [0088]
- **DE MESMAEKER et al.**Acc. Chem. Res., 1995, vol. 28, 366-374 [0178] [0179]
- **NIELSEN et al.**Science, 1991, vol. 254, 1497- [0179]
- **MARTIN et al.**Helv. Chim. Acta, 1995, vol. 78, 486- [0179]
- **KOMBERG, A.**DNA ReplicationW. H. Freeman & Co.1980000075-77 [0180]
- **GEBEYEHU, G. et al.**Nucl. Acids Res., 1987, vol. 15, 4513- [0180]
- **Antisense Research and Applications**CRC Press19930000 [0180]
- **UHLMAN et al.**Current Opinions in Drug Discovery & Development, 2000, vol. 3, 2 [0185]
- **NIELSEN et al.**Science, 1991, vol. 254, 1497-1500 [0190]
- **MARTIN et al.**Helv. Chim. Acta, 1995, vol. 78, 486-504 [0192]
- **Concise Encyclopedia of Polymer Science And Engineering**John Wiley & Sons19900000858-859 [0195]
- **ENGLISCH et al.**Angewandle Chemie, International Edition, 1991, vol. 30, 613- [0195]

- **SANGHVI, Y.S.** Antisense Research and Applications CRC Press 1993 0000289-302 [0195]
- Antisense Research and Applications CRC Press 1993 0000276-278 [0195]
- **MANOHARAN et al.** Ann. N. Y. Acad. Sci., 1992, vol. 660, 306-309 [0198]
- **MANOHARAN et al.** Bioorg. Med. Chem. Lett., 1993, vol. 3, 2765-2770 [0198]
- **BRAZMAVILOF** FEBS Lett., 2000, vol. 480, 17-24 [0210]
- **CELIS et al.** FEBS Lett., 2000, vol. 480, 2-16 [0210] [0210] [0210]
- **MADDEN et al.** Drug Discov. Today, 2000, vol. 5, 415-425 [0210]
- **PRASHARWEISSMAN** Methods Enzymol., 1999, vol. 303, 258-72 [0210]
- **SUTCLIFFE et al.** Proc. Natl. Acad. Sci. U.S.A., 2000, vol. 97, 1976-81 [0210]
- **JUNGBLUT et al.** Electrophoresis, 1999, vol. 20, 2100-10 [0210]
- **LARSSON et al.** J. Biotechnol., 2000, vol. 80, 143-57 [0210]
- **FUCHS et al.** Anal. Biochem., 2000, vol. 286, 91-98 [0210]
- **LARSON et al.** Cytometry, 2000, vol. 41, 203-208 [0210]
- **JURECICBELMONT** Curr. Opin. Microbiol., 2000, vol. 3, 316-21 [0210]
- **CARULLI et al.** J. Cell Biochem. Suppl., 1998, vol. 31, 286-96 [0210]
- **GOINGGUSTERSON** Eur. J. Cancer, 1999, vol. 35, 1895-904 [0210]
- **TO, COMB.** Chem. High Throughput Screen, 2000, vol. 3, 235-41 [0210]
- **GELLER, A.I. et al.** J. Neurochem, 1995, vol. 64, 487- [0223]
- **LIM, F. et al.** DNA Cloning: Mammalian Systems Oxford Univ. Press 1995 0000 [0223]
- **GELLER, A.I. et al.** Proc Natl. Acad. Sci., 1993, [0223]
- **GELLER, A.I. et al.** Proc Natl. Acad. Sci USA, 1990, vol. 87, 1149- [0223]
- **LEGAL LASALLE et al.** Science, 1993, vol. 259, 988- [0223]
- **DAVIDSON et al.** Nat. Genet., 1993, vol. 3, 219- [0223]
- **YANG et al.** J. Virol., 1995, vol. 69, 2004- [0223]
- **KAPLITT, M.G. et al.** Nat. Genet., 1994, vol. 8, 148- [0223]

Patentkrav

1. Oligonukleotid, der er målrettet et naturligt omprogrammeringsfaktor-antisense-transskript til anvendelse som en terapeutisk forbindelse, hvor oligonukleotidet øger ekspressionen af omprogrammeringsfaktoren, og hvor det naturlige antisense-transskript har nukleinsyresekvensen ifølge en hvilken som helst af SEQ ID NO: 4, 5 eller 6, eller en variant deraf, der bevarer funktionen af det naturlige antisense-transskript ifølge SEQ ID NO: 4, 5 eller 6.

2. Oligonukleotid, der er målrettet et naturligt omprogrammeringsfaktor-antisense-transskript til anvendelse i forebyggelse eller behandling af en omprogrammeringsfaktorassocieret sygdom eller forstyrrelse, hvor oligonukleotidet øger ekspressionen af omprogrammeringsfaktoren, og hvor det naturlige antisense-transskript har nukleinsyresekvensen ifølge en hvilken som helst af SEQ ID NO: 4, 5 eller 6, eller en variant deraf, der bevarer funktionen af det naturlige antisense-transskript ifølge SEQ ID NO: 4, 5 eller 6.

3. Anvendelse af et oligonukleotid, der er målrettet et naturligt omprogrammeringsfaktor-antisense-transskript til fremstilling af et medikament til forebyggelse eller behandling af en omprogrammeringsfaktorassocieret sygdom eller forstyrrelse, hvor oligonukleotidet øger ekspressionen af omprogrammeringsfaktoren, og hvor det naturlige antisense-transskript har nukleinsyresekvensen ifølge en hvilken som helst af SEQ ID NO: 4, 5 eller 6, eller en variant deraf, der bevarer funktionen af det naturlige antisense-transskript ifølge SEQ ID NO: 4, 5 eller 6.

4. Anvendelse ifølge krav 3, eller oligonukleotid til anvendelse ifølge krav 2, hvor sygdommen eller forstyrrelsen er udvalgt fra gruppen bestående af abnorm omprogrammeringsfaktorekspression og/eller funktion, cancer, en nyresygdom, en kardiovaskulær sygdom eller forstyrrelse, inflammation, en neurologisk sygdom eller forstyrrelse, ardannelse, øjensygdomme eller -forstyrrelser (f.eks. mikroftalmi, kolobom, myopi, anoftalmi/mikroftalmi - øsofagealt atresisyndrom mv.), septooptisk dysplasi, en autoimmun sygdom eller forstyrrelse, diabetes, aterosklerose, en immundeficienssygdom, en sygdom forårsaget af et infektiøst middel (f.eks.

indbefattende viralt, bakterielt, fungalt, protozoisk mv.), en genetisk sygdom (f.eks. Duchennes muskeldystrofi), en metabolisk sygdom eller forstyrrelse (f.eks. diabetes, fedme, metabolisk syndrom, lysosomal sygdom etc.), traume (f.eks. rygradslæsion, forbrændinger mv), iskæmi, en karsygdom, en leversygdom eller -forstyrrelse, psoriasis, en sygdom eller forstyrrelse, der kræver transplantation af celler, væv og organer, forstyrrelse eller tilstand, der kræver stamcelleterapi og en medfødt sygdom eller forstyrrelse.

5. Fremgangsmåde *in vitro* til øgning af ekspressionen af en omprogrammeringsfaktor i patientceller eller -væv, der omfatter: etablering af kontakt mellem cellerne og vævet og et oligonukleotid, der er specifikt målrettet et naturligt omprogrammeringsfaktor-antisense-transskript; hvorved ekspressionen af omprogrammeringsfaktoren øges, hvor det naturlige antisense-transskript har nukleinsyresekvensen ifølge en hvilken som helst af SEQ ID NO: 4, 5 eller 6, eller en variant deraf, der bevarer funktionen af det naturlige antisense-transskript ifølge SEQ ID NO: 4, 5 eller 6.

6. Anvendelse ifølge et hvilket som helst af kravene 3 eller 4, eller oligonukleotid til anvendelse ifølge et hvilket som helst af kravene 1, 2 eller 4 eller fremgangsmåde ifølge krav 5, hvor oligonukleotidet er:

- a. enkeltstrengt eller
- b. en siRNA-forbindelse.

7. Anvendelse ifølge et hvilket som helst af kravene 3, 4 eller 6, eller oligonukleotid til anvendelse ifølge et hvilket som helst af kravene 1, 2, 4 eller 6 eller fremgangsmåde ifølge krav 5 eller 6, hvor:

- a. oligonukleotidet omfatter mindst én af SEQ ID NO: 7 til 17; og/eller
- b. ekspressionen af omprogrammeringsfaktoren øges med mindst 10 %.

8. Anvendelse ifølge et hvilket som helst af kravene 3, 4, 6 eller 7, eller oligonukleotid til anvendelse ifølge et hvilket som helst af kravene 1, 2, 4, 6 eller 7 eller fremgangsmåde ifølge et hvilket som helst af kravene 5 til 7, hvor oligonukleotidet endvidere omfatter én eller flere modifikationer, der omfatter:

- a. mindst én modificeret internukleosidbinding udvalgt fra: et phosphorthioat, alkylphosphonat, phosphordithioat, alkylphosphonthioat, phosphoramidat, carbamat, carbonat, en phosphattriester, acetamidat, carboxymethylester og kombinationer deraf;
- b. mindst ét modificeret nukleotid udvalgt fra: en peptidnukleinsyre (PNA), en låst nukleinsyre (LNA), en arabino-nukleinsyre, en analog, et derivat og kombinationer deraf; eller
- c. mindst én modificeret sukkerdel udvalgt fra: en 2'-O-methoxyethylmodificeret sukkerdel, en 2'-methoxymodificeret sukkerdel, en 2'-O-alkylmodificeret sukkerdel, en bicyklisk sukkerdel, en 2'-fluordel og kombinationer deraf.

9. Oligonukleotid, der specifikt kan hybridisere til et naturligt antisense-transskript, der har nukleinsyresekvensen ifølge en hvilken som helst af SEQ ID NO: 4, 5 eller 6, eller en variant deraf, der bevarer funktionen af det naturlige antisense-transskript ifølge SEQ ID NO: 4, 5 eller 6.

10. Oligonukleotid ifølge krav 9, hvor oligonukleotidet:

- a. er enkeltstrenget eller en siRNA-forbindelse;
- b. omfatter mindst én af SEQ ID NO: 7 til 17;
- c. endvidere omfatter én eller flere modifikationer, der omfatter:
 - i. mindst én modificeret internukleosidbinding udvalgt fra: et phosphorthioat, alkylphosphonat, phosphordithioat, alkylphosphonthioat, phosphoramidat, carbamat, carbonat, en phosphattriester, acetamidat, carboxymethylester og kombinationer deraf;
 - ii. mindst ét modificeret nukleotid udvalgt fra: en peptidnukleinsyre (PNA), en låst nukleinsyre (LNA), en arabino-nukleinsyre, en analog, et derivat og kombinationer deraf; eller
 - iii. mindst én modificeret sukkerdel udvalgt fra: en 2'-O-methoxyethylmodificeret sukkerdel, en 2'-methoxymodificeret sukkerdel, en 2'-O-alkylmodificeret sukkerdel, en bicyklisk sukkerdel, en 2'-fluordel og kombinationer deraf; og/eller
- d. er mellem 10 til 30 nukleotider langt.

11. Farmaceutisk sammensætning, der omfatter mindst ét oligonukleotid ifølge krav 9 eller 10 og en farmaceutisk acceptabel excipiens.



12. Fremgangsmåde til fremstilling af en induceret, pluripotent stem- (iPS) celle ved nukleær omprogrammering af somatiske celler, der omfatter: etablering af kontakt mellem de somatiske celler og et oligonukleotid ifølge krav 9 eller 10.

13. Fremgangsmåde *in vitro* til øgning af ekspressionen af en omprogrammeringsfaktor i en induceret, pluripotent stem- (iPS) celle, der omfatter etablering af kontakt mellem en somatisk celle og et oligonukleotid ifølge krav 9 eller 10.

14. Induceret, pluripotent stem- (iPS) celle, der kan opnås ved hjælp af fremgangsmåden ifølge krav 12 eller 13, hvor iPS-cellen omfatter et oligonukleotid ifølge krav 9 eller 10.

15. Induceret, pluripotent stem- (iPS) celle ifølge krav 14 til anvendelse i stamcelleterapi, hvor terapien omfatter induktion af differentiering af iPS-cellen for at opnå en differentieret iPS-celle, og transplantation af den differentierede iPS-celle i patienten.

DRAWINGS

FIGURE 1**A****B**

C

