




(86) Date de dépôt PCT/PCT Filing Date: 2011/10/13
(87) Date publication PCT/PCT Publication Date: 2012/04/19
(45) Date de délivrance/Issue Date: 2018/04/10
(85) Entrée phase nationale/National Entry: 2013/04/04
(86) N° demande PCT/PCT Application No.: JP 2011/074091
(87) N° publication PCT/PCT Publication No.: 2012/050233
(30) Priorité/Priority: 2010/10/14 (JP2010-231369)

(51) Cl.Int./Int.Cl. *A01N 37/36* (2006.01),
A01N 47/12 (2006.01), *A01N 47/18* (2006.01),
A01N 47/22 (2006.01), *A01N 47/24* (2006.01),
A01P 3/00 (2006.01)
(72) Inventeurs/Inventors:
KIGUCHI, SO, JP;
TANAKA, SOICHI, JP
(73) Propriétaire/Owner:
SUMITOMO CHEMICAL COMPANY, LIMITED, JP
(74) Agent: KIRBY EADES GALE BAKER

(54) Titre : COMPOSITION DE CONTROLE DE PARASITES FONGIQUES RENFERMANT DE LA MANDESTROBINE ET UN COMPOSE CARBAMATE ET METHODE DE CONTROLE DE PARASITES FONGIQUES
(54) Title: FUNGAL PEST CONTROLLING COMPOSITION COMPRISING MANDESTROBIN AND A CARBAMATE COMPOUND AND METHOD FOR CONTROLLING FUNGAL PESTS



(1)

(57) Abrégé/Abstract:

The present invention provides a composition having excellent pest controlling activity. The composition comprising a compound represented by Formula (1)

(see formula 1)

and one or more carbamate compound(s) selected from Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb and xylycarb.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(10) International Publication Number
WO 2012/050233 A3(43) International Publication Date
19 April 2012 (19.04.2012)

(51) International Patent Classification:

A01N 37/36 (2006.01) A01N 47/12 (2006.01)
A01N 47/24 (2006.01) A01N 47/18 (2006.01)
A01N 47/22 (2006.01) A01P 3/00 (2006.01)

(21) International Application Number:

PCT/JP2011/074091

(22) International Filing Date:

13 October 2011 (13.10.2011)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2010-231369 14 October 2010 (14.10.2010) JP

(71) Applicant (for all designated States except US): **SUMITOMO CHEMICAL COMPANY, LIMITED** [JP/JP]; 27-1, Shinkawa 2-chome, Chuo-ku, Tokyo, 1048260 (JP).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KIGUCHI, So** [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hyogo, 6658555 (JP). **TANAKA, Soichi** [JP/JP]; c/o SUMITOMO CHEMICAL COMPANY, LIMITED, 2-1, Takatsukasa 4-chome, Takarazuka-shi, Hyogo, 6658555 (JP).(74) Agents: **TANAKA, Mitsuo** et al.; AOYAMA & PARTNERS, IMP Building, 3-7, Shiromi 1-chome, Chuo-ku, Osaka-shi, Osaka, 5400001 (JP).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.


(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

[Continued on next page]

(54) Title: PEST CONTROLLING COMPOSITION AND METHOD FOR CONTROLLING PEST



(57) Abstract: The present invention provides a composition having an excellent controlling activity on a pest. The composition comprising a compound represented by Formula (1) and one or more carbamate compound(s) selected from Group (A) shows an excellent controlling activity on a pest. Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, proprocur, XMC, thiodicarb and xylycarb

WO 2012/050233 A3



— *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(88) Date of publication of the international search report:
7 September 2012

FUNGAL PEST CONTROLLING COMPOSITION COMPRISING MANDESTROBIN
AND A CARBAMATE COMPOUND AND
METHOD FOR CONTROLLING FUNGAL PESTS


5 Technical Field

The present invention relates to a pest controlling composition and a method for controlling a pest.

Background Art

10 Hitherto, there have been provided compounds as an active ingredient for a composition for controlling a pest (see e.g., The Pesticide Manual - 15th edition (BCPC published) ISBN 1901396188; and SHIBUYA INDEX (Index of Pesticides) 13th Edition 2008 (SHIBUYA INDEX RESEARCH GROUP
15 published) ISBN 9784881371435).

Also there has been provided a compound of Formula (1):



(see e.g., WO 95/27693 pamphlet and WO 02/10101 pamphlet).


20 Disclosure of Invention

An object of the present invention is to provide a composition having an excellent pest control effect.

The present inventors have intensively studied to find a composition having an excellent pest control effect. As a result, they have found that a composition comprising the compound represented by Formula (1) and one or more carbamate compound(s) selected from the following group (A) shows a synergistic activity, and thus has an excellent control effect on pests, and therefore the present invention has been completed.

The present invention provides:

- [1] A fungal pest controlling composition comprising a compound represented by Formula (1):



and one or more carbamate compound(s) selected from Group (A):


- Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb and xylylcarb.

- [2] The fungal pest controlling composition according to the above [1], wherein the weight ratio of the compound represented by Formula (1) to the carbamate compound(s) is

from 0.0125/1 to 500/1.

[3] The fungal pest controlling composition according to the above [1] or [2], wherein the compound represented by Formula (1) has R-absolute configuration.


5 [4] A method for controlling a fungal pest, wherein the method comprises applying a compound of Formula (1):



and one or more carbamate compound(s) selected from Group (A) to a plant or a soil for cultivating the plant,

10 Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocab, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb,
 15 propoxur, XMC, thiodicarb and xylylcarb.

[5] A method for controlling a fungal pest, wherein the method comprises applying a compound represented by Formula (1):




and one or more carbamate compound(s) selected from Group (A) to a seed,

Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb and xylylcarb.

[6] The method according to the above [4] or [5], wherein the weight ratio of the compound represented by Formula (1) to the carbamate compound(s) is from 0.0125/1 to 500/1.

[7] The method according to any one of the above [4] to [6], wherein the compound represented by Formula (1) has R-absolute configuration.

[8] Use of a combination of a compound represented by Formula (1):



and one or more carbamate compound(s) selected from Group (A) for controlling a fungal pest,


Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb,

fenothiocarb, fenoxycarb, furathiocarb, isoprocarb,
metolcarb, methomyl, methiocarb, oxamyl, pirimicarb,
propoxur, XMC, thiodicarb and xylylcarb.

5 The present invention enables one to control pests.

Mode for Carrying Out the Invention

A pest controlling composition of the present
invention (hereinafter, referred to as a composition of the
10 present invention) comprises a compound represented by
Formula (1):



(hereinafter, referred to as an amide compound of the present invention) and one or more carbamate compound(s) selected from Group (A) (hereinafter, referred to as a carbamate compound of the present invention),

Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocab, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb and xylylcarb.

The present amide compound is described in, for example, WO 95/27693 pamphlet and WO 02/10101 pamphlet, and thus can be prepared according to the method described therein.

The present amide compound has one asymmetric carbon. Herein, a compound represented by Formula (1) wherein an enantiomer having R-absolute configuration is enriched is referred to as an amide compound having R-absolute configuration.

The present amide compound encompasses the following compounds:

compounds represented by Formula (1) wherein an enantiomer having R-absolute configuration amounts to 70% and more of the total amount thereof;

5 compounds represented by Formula (1) wherein an enantiomer having R-absolute configuration amounts to 90% and more of the total amount thereof;

compounds represented by Formula (1) wherein an enantiomer having R-absolute configuration amounts to 95% and more of the total amount thereof.

10 Alanycarb, aldicarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC and thiodicarb to be used in the present invention are known
15 compounds, which are described in, for example, "The PESTICIDE MANUAL - 15th EDITION (BCPC published) ISBN 1901396188", pages 23, 24, 79, 82, 156, 161, 163, 441, 476, 478, 482, 581, 670, 757, 755, 848, 916, 956, 1194 and 1124 respectively. These compounds are either commercially
20 available, or can be prepared by a known method.

Aldoxycarb, cloethocarb, metolcarb and xylylcarb to be used in the present invention are known compounds, which are described in, for example, "SHIBUYA INDEX (Index of Pesticides) 13th Edition 2008 (SHIBUYA INDEX RESEARCH GROUP
25 published) ISBN 9784881371435", pages 30, 32, 31 and 31

respectively. These compounds are either commercially available, or can be prepared by a known method.

The weight ratio of the present amide compound to the present carbamate compound(s) in the composition of the present invention is usually from 0.0125/1 to 500/1 (the present amide compound/the present carbamate compound(s)), preferably 0.025/1 to 100/1, and more preferably 0.1/1 to 10/1.

Although the composition of the present invention may be a mixture alone of the present amide compound and the present carbamate compound(s), the composition of the present invention is usually prepared by mixing the present amide compound, the present carbamate compound(s) and an inert carrier, and if necessary, adding a surfactant or other pharmaceutical additives, and then formulating into the form of an oil solution, emulsifiable concentrate, flowable formulation, wettable powder, granulated wettable powder, dust formulation, granules and so on. Such formulations can be used alone or with an addition of other inert components as an agent for controlling a pest.

Usually, the composition of the present invention can contain 0.1 to 99 % by weight, preferably 0.2 to 90 % by weight, and more preferably 1 to 80 % by weight of the present amide compound and the present carbamate compound(s) in total.

Examples of a solid carrier used on the formulation include finely-divided powder or particles of clay consisting of minerals (e.g., kaolin clay, attapulgite clay, bentonite, montmorillonite, acid clay, pyrophyllite, talc, diatomaceous earth, or calcite), natural organic substances (e.g., corncob powder, or walnut shell powder), synthetic organic substances (e.g., urea), salts (e.g., calcium carbonate, or ammonium sulfate), synthetic inorganic substances (e.g., synthetic hydrous silicon oxide) and so on. Examples of a liquid carrier include aromatic hydrocarbons (e.g., xylene, alkyl benzene, or methylnaphthalene), alcohols (e.g., 2-propanol, ethylene glycol, propylene glycol, or ethylene glycol monoethyl ether), ketones (e.g., acetone, cyclohexanone, or isophorone), vegetable oils (e.g., soybean oil, or cotton oils), petroleum-derived aliphatic hydrocarbons, esters, dimethylsulfoxide, acetonitrile and water.

Examples of the surfactant include anionic surfactant (e.g., alkyl sulfate salts, alkylaryl sulfate salts, dialkyl sulfosuccinate salts, polyoxyethylene alkylaryl ether phosphates, lignin sulfonate, or naphthalenesulfonate formaldehyde polycondensation), nonionic surfactant (e.g., polyoxyethylene alkylaryl ether, polyoxyethylene alkyl polyoxypropylene block copolymer, or sorbitan fatty acid ester) and cationic surfactant (e.g., alkyltrimethyl

ammonium salts).

Examples of the other pharmaceutical additives include water-soluble polymers (e.g., polyvinyl alcohol, or polyvinyl pyrrolidone), polysaccharides (e.g. arabic gum, 5 alginic acid and salts thereof, CMC (carboxymethyl-cellulose), or xanthan gum), inorganic substances (e.g., aluminum magnesium silicate, or alumina-sol), antiseptic agent, coloring agent, and PAP (isopropyl acid phosphate), and stabilizing agent (e.g., BHT).

10 The composition of the present invention can also be prepared by separately formulating the present amide compound and the present carbamate compound(s) into different formulations by the above procedures, if necessary, further diluting each of them with water, 15 thereafter, mixing the separately prepared different formulations or the dilute solutions.

The composition of the present invention may further contain one or more other fungicide(s) and/or insecticide(s).

20 The composition of the present invention is used to control a pest by applying it to a plant or soil for cultivating the plant.

The arthropod pests on which the composition of the present invention exhibits a controlling effect are 25 exemplified below:

Hemiptera:

Planthoppers (Delphacidae) such as small brown planthopper (*Laodelphax striatellus*), brown rice planthopper (*Nilaparvata lugens*), and white-backed rice
5 planthopper (*Sogatella furcifera*); leafhoppers (Deltocephalidae) such as green rice leafhopper (*Nephotettix cincticeps*) and green rice leafhopper (*Nephotettix virescens*); aphids (Aphididae) such as cotton
10 aphid (*Aphis gossypii*), green peach aphid (*Myzus persicae*), cabbage aphid (*Brevicoryne brassicae*), potato aphid (*Macrosiphum euphorbiae*), foxglove aphid (*Aulacorthum solani*), oat bird-cherry aphid (*Rhopalosiphum padi*), and tropical citrus aphid (*Toxoptera citricidus*); stink bugs (Pentatomidae) such as green stink bug (*Nezara antennata*),
15 bean bug (*Riptortus clavetus*), rice bug (*Leptocorisa chinensis*), white spotted spined bug (*Eysarcoris parvus*), stink bug (*Halyomorpha mista*), and tarnished plant bug (*Lysus lineolaris*); whiteflies (Aleyrodidae) such as greenhouse whitefly (*Trialeurodes vaporariorum*), sweet
20 potato whitefly (*Bemisia tabaci*), and silverleaf whitefly (*Bemisia argentifolii*); scales (Coccidae) such as California red scale (*Aonidiella aurantii*), San Jose scale (*Comstockaspis perniciosus*), citrus north scale (*Unaspis citri*), red wax scale (*Ceroplastes rubens*), cottoncushion
25 scale (*Icerya purchasi*); lace bugs (Tingidae); psyllids

(Psyllidae), etc.;

Lepidoptera:

Pyralid moths (Pyralidae) such as rice stem borer
(*Chilo suppressalis*), yellow rice borer (*Tryporyza*
5 *incertulas*), rice leafroller (*Cnaphalocrocis medinalis*),
cotton leafroller (*Notarcha derogata*), Indian meal moth
(*Plodia interpunctella*), *Micractis nubilalis* (*Ostrinia*
furnacalis), European corn borer (*Ostrinia nubilalis*),
cabbage webworm (*Hellula undalis*), and bluegrass webworm
10 (*Pediasia teterrellus*); owlet moths (Noctuidae) such as
common cutworm (*Spodoptera litura*), beet armyworm
(*Spodoptera exigua*), armyworm (*Pseudaletia separata*),
cabbage armyworm (*Mamestra brassicae*), black cutworm
(*Agrotis ipsilon*), beet semi-looper (*Plusia nigrisigna*),
15 *Thoricoplusia* spp., *Heliothis* spp., and *Helicoverpa* spp.;
white butterflies (Pieridae) such as common white (*Pieris*
rapae); tortricid moths (Tortricidae) such as *Adoxophyes*
spp., oriental fruit moth (*Grapholita molesta*), soybean pod
borer (*Leguminivora glycinivorella*), azuki bean podworm
20 (*Matsumuraeses azukivora*), summer fruit tortrix (*Adoxophyes*
orana fasciata), smaller tea tortrix (*Adoxophyes honmai*),
oriental tea tortrix (*Homona magnanima*), apple tortrix
(*Archips fuscocupreanus*), and codling moth (*Cydia*
pomonella); leafblotch miners (Gracillariidae) such as tea
25 leafroller (*Caloptilia theivora*), and apple leafminer

(*Phyllonorycter ringoniella*); Carposinidae such as peach fruit moth (*Carposina niponensis*); Lyonetiid moths (Lyonetiidae) such as *Lyonetia* spp.; tussock moths (Lymantriidae) such as *Lymantria* spp., and *Euproctis* spp.;
5 yponomeutid moths (Yponomeutidae) such as diamondback (*Plutella xylostella*); gelechiid moths (Gelechiidae) such as pink bollworm (*Pectinophora gossypiella*), and potato tubeworm (*Phthorimaea operculella*); tiger moths and allies (Arctiidae) such as fall webworm (*Hyphantria cunea*); and
10 tineid moths (Tineidae) such as casemaking clothes moth (*Tinea translucens*), etc.;

Thysanoptera:

Yellow citrus thrips (*Frankliniella occidentalis*), melon thrips (*Thrips palmi*), yellow tea thrips
15 (*Scirtothrips dorsalis*), onion thrips (*Thrips tabaci*), flower thrips (*Frankliniella intonsa*), and tobacco thrips (*Frankliniella fusca*), etc.;

Diptera:

Leafminer flies (Agromyzidae) such as onion maggot
20 (*Hylemya antiqua*), seed corn maggot (*Hylemya platura*), rice leafminer (*Agromyza oryzae*), rice leafminer (*Hydrellia griseola*), rice stem maggot (*Chlorops oryzae*), legume leafminer (*Liriomyza trifolii*); melon fly (*Dacus cucurbitae*), and Mediterranean fruit fly (*Ceratitis capitata*), etc.;

25

Coleoptera:

Twenty-eight-spotted ladybird (*Epilachna vigintioctopunctata*), cucurbit leaf beetle (*Aulacophora femoralis*), striped flea beetle (*Phyllotreta striolata*),
 5 rice leaf beetle (*Oulema oryzae*), rice curculio (*Echinocnemus squameus*), rice water weevil (*Lissorhoptrus oryzophilus*), boll weevil (*Anthonomus grandis*), azuki bean weevil (*Callosobruchus chinensis*), hunting billbug (*Sphenophorus venatus*), Japanese beetle (*Popillia japonica*),
 10 cupreous chafer (*Anomala cuprea*), Corn root worms (*Diabrotica* spp.), Colorado beetle (*Leptinotarsa decemlineata*), click beetles (*Agriotes* spp.), and cigarette beetle (*Lasioderma serricorne*), etc.;

Orthoptera:

15 African mole cricket (*Gryllotalpa africana*), rice grasshopper (*Oxya yezoensis*), and rice grasshopper (*Oxya japonica*), etc.;

Hymenoptera:

Cabbage sawfly (*Athalia rosae*), leaf-cutting ant
 20 (*Acromyrmex* spp.), and fire ant (*Solenopsis* spp.), etc.;

Acarina:

Spider mites (Tetranychidae) such as two-spotted spider mite (*Tetranychus urticae*), citrus red mite (*Panonychus citri*), and *Oligonychus* spp.; eriophyid mites
 25 (Eriophyidae) such as pink citrus rust mite (*Aculops*

pelekassi); tarsonemid mites (Tarsonemidae) such as broad
 mite (*Polyphagotarsonemus latus*); false spider mites
 (Tenuipalpidae); tuckerellidae; acarid mites (Acaridae)
 such as mold mite (*Tyrophagus putrescentiae*); house dust
 5 mites (Pyroglyphidae) such as *Dermatophagoides farinae*, and
Dermatophagoides ptrebyssus; cheyletoid mites
 (Cheyletidae) such as *Cheyletus eruditus*, *Cheyletus*
malaccensis, and *Cheyletus moorei*;

Nematodes:

10 White tip nematode (*Aphelenchoides besseyi*), and
 strawberry bud nematode (*Nothotylenchus acris*), etc.

The plant diseases which can be controlled by the present invention are exemplified below:

Rice diseases: blast (*Magnaporthe oryzae*),
 15 helminthosporium leaf spot (*Cochliobolus miyabeanus*),
 sheath blight (*Rhizoctonia solani*) and bakanae disease
 (*Gibberella fujikuroi*);

Diseases of barley, wheat, oats and rye: powdery
 mildew (*Erysiphe graminis*), Fusarium head blight (*Fusarium*
 20 *graminearum*, *F. avenaceum*, *F. culmorum*, *F. asiaticum*,
Microdochium nivale), rust (*Puccinia striiformis*,
P. graminis, *P. recondite*, *P. hordei*), snow blight (*Typhula*
sp., *Micronectriella nivalis*), loose smut (*Ustilago tritici*,
U. nuda), bunt (*Tilletia caries*), eyespot
 25 (*Pseudocercosporella herpotrichoides*), scald

(*Rhynchosporium secalis*), leaf blotch (*Septoria tritici*),
glume blotch (*Leptosphaeria nodorum*) and net blotch
(*Pyrenophora teres Drechsler*);

Citrus diseases: melanose (*Diaporthe citri*), scab
5 (*Elsinoe fawcetti*), green mold (*Penicillium digitatum*) and
blue mold (*Penicillium italicum*);

Apple diseases: blossom blight (*Monilinia mali*),
canker (*Valsa ceratosperma*), powdery mildew (*Podosphaera*
leucotricha), Alternaria leaf spot (*Alternaria alternata*
10 *apple pathotype*), scab (*Venturia inaequalis*), bitter rot
(*Colletotrichum acutatum*) and late blight (*Phytophthora*
cactorum);

Pear diseases: scab (*Venturia nashicola*, *V. pirina*),
black spot (*Alternaria alternata Japanese pear pathotype*),
15 rust (*Gymnosporangium asiaticum*) and late blight
(*Phytophthora cactorum*);

Peach diseases: brown rot (*Monilinia fructicola*), scab
(*Cladosporium carpophilum*) and Phomopsis rot (*Phomopsis sp.*);

Grape diseases: anthracnose (*Elsinoe ampelina*), ripe
20 rot (*Glomerella cingulata*), powdery mildew (*Uncinula*
necator), rust (*Phakopsora ampelopsidis*), black rot
(*Guignardia bidwellii*), downy mildew (*Plasmopara viticola*)
and Gray mold (*Botrytis cinerea*);

Diseases of Japanese persimmon: anthracnose
25 (*Gloeosporium kaki*) and leaf spot (*Cercospora kaki*,

Mycosphaerella nawae);

Diseases of gourd family: anthracnose (*Colletotrichum lagenarium*), powdery mildew (*Sphaerotheca fuliginea*), gummy stem blight (*Mycosphaerella melonis*), Fusarium wilt (5 *Fusarium oxysporum*), downy mildew (*Pseudoperonospora cubensis*), Phytophthora rot (*Phytophthora* sp.), gray mold fungus (*Botrytis cinerea*) and damping-off (*Pythium* sp.);

Tomato diseases: early blight (*Alternaria solani*), leaf mold (*Cladosporium fulvum*) and late blight (10 *Phytophthora infestans*);

Egg plant disease: brown spot (*Phomopsis vexans*) and powdery mildew (*Erysiphe cichoracearum*);

Diseases of Cruciferous Vegetables: *Alternaria* leaf spot (*Alternaria japonica*), white spot (*Cercospora* 15 *brassicae*), clubroot (*Plasmodiophora brassicae*), and downy mildew (*Peronospora parasitica*);

Rapeseed diseases: *Sclerotinia* rot (*Sclerotinia sclerotiorum*), black spot (*Alternaria brassicae*), powdery mildew (*Erysiphe cichoracearum*), blackleg (*Leptosphaeria* 20 *maculans*);

Welsh onion diseases: rust (*Puccinia allii*);

Soybean diseases: purple seed stain (*Cercospora* *kikuchii*), Sphaceloma scad (*Elsinoe glycines*), pod and stem blight (*Diaporthe phaseolorum* var. *sojae*), rust (*Phakopsora* 25 *pachyrhizi*) and phytophthora stem rot (*Phytophthora sojae*);

Adzuki-bean diseases: Gray mold (*Botrytis cinerea*),
Sclerotinia rot (*Sclerotinia sclerotiorum*);

Kidney bean diseases: Gray mold (*Botrytis cinerea*),
Sclerotinia rot (*Sclerotinia sclerotiorum*), anthracnose
5 (*Colletotrichum lindemthianum*);

Peanut diseases: leaf spot (*Cercospora personata*),
brown leaf spot (*Cercospora arachidicola*) and southern
blight (*Sclerotium rolfsii*);

Garden pea diseases: powdery mildew (*Erysiphe pisi*);

10 Potato diseases: early blight (*Alternaria solani*) and
late blight (*Phytophthora infestans*);

Strawberry diseases: powdery mildew (*Sphaerotheca
humuli*);

15 Tea diseases: net blister blight (*Exobasidium
reticulatum*), white scab (*Elsinoe leucospila*), gray blight
(*Pestalotiopsis* sp.) and anthracnose (*Colletotrichum theae-
sinensis*);

Cotton diseases: fusarium wilt (*Fusarium oxysporum*),
damping-off (*Rhizoctonia solani*);

20 Tobacco diseases: brown spot (*Alternaria longipes*),
powdery mildew (*Erysiphe cichoracearum*), anthracnose
(*Colletotrichum tabacum*), downy mildew (*Peronospora
tabacina*) and late blight (*Phytophthora nicotianae*);

25 Sugar beet diseases: Cercospora leaf spot (*Cercospora
beticola*), leaf blight (*Thanatephorus cucumeris*), Root rot

(*Thanatephorus cucumeris*), *Aphanomyces* root rot
(*Aphanidermatum cochlioides*);

Rose diseases: black spot (*Diplocarpon rosae*) and
powdery mildew (*Sphaerotheca pannosa*);

5 Chrysanthemum diseases: leaf blight (*Septoria*
chrysanthemi-indici) and white rust (*Puccinia horiana*);

Various plants diseases: diseases caused by *Pythium*
spp. (*Pythium aphanidermatum*, *Pythium debarianum*, *Pythium*
graminicola, *Pythium irregulare*, *Pythium ultimum*), Gray
10 mold (*Botrytis cinerea*), *Sclerotinia* rot (*Sclerotinia*
sclerotiorum),

Japanese radish diseases: *Alternaria* leaf spot
(*Alternaria brassicicola*);

Turfgrass diseases: dollar spot (*Sclerotinia*
15 *homeocarpa*), brown patch and large patch (*Rhizoctonia*
solani); and

Banana diseases: Sigatoka disease (*Mycosphaerella*
fijiensis, *Mycosphaerella musicola*, *Pseudocercospora musae*).

Examples of the plants to which the composition of the
20 present invention can be applied are as follows:

Crops: corn, rice, wheat, barley, rye, oat, sorghum,
cotton, soybean, adzuki-bean, kidney bean, peanut,
buckwheat, beet, rapeseed, sunflower, sugar cane, and
tobacco, etc.;

Vegetables: solanaceous vegetables (eggplant, tomato, pimento, pepper, and potato, etc.), cucurbitaceous vegetables (cucumber, pumpkin, zucchini, watermelon, melon, and squash, etc.), cruciferous vegetables (Japanese radish, white turnip, horseradish, kohlrabi, Chinese cabbage, cabbage, leaf mustard, broccoli, and cauliflower, etc.), asteraceous vegetables (burdock, crown daisy, artichoke, and lettuce, etc.), liliaceous vegetables (green onion, onion, garlic, and asparagus), ammiaceous vegetables (carrot, parsley, celery, and parsnip, etc.), chenopodiaceous vegetables (spinach, and Swiss chard, etc.), lamiaceous vegetables (Perilla frutescens, mint, and basil, etc.), strawberry, sweet potato, Japanese yam, and taro, etc.;

Flowers;

Foliage plants;

Turfgrass;

Fruits: pomaceous fruits (apple, pear, Japanese pear, Chinese quince, and quince, etc.), stone fleshy fruits (peach, plum, nectarine, Japanese apricot, cherry fruit, apricot, and prune, etc.), citrus fruits (Citrus unshiu, orange, lemon, lime, and grapefruit, etc.), nuts (chestnut, walnuts, hazelnuts, almond, pistachio, cashew nuts, and macadamia nuts, etc.), berries (blueberry, cranberry, blackberry, and raspberry, etc.), grape, kaki persimmon, olive, Japanese plum, banana, coffee, date palm, and

coconuts, etc.; and

Trees other than fruit trees: tea, mulberry, flowering plant, roadside trees (ash, birch, dogwood, Eucalyptus, Ginkgo biloba, lilac, maple, Quercus, poplar, Judas tree, Liquidambar formosana, plane tree, zelkova, Japanese arborvitae, fir wood, hemlock, juniper, Pinus, Picea, and Taxus cuspidate), etc.

The aforementioned "plants" include plants which resistances have been imparted by genetic recombination.

Exemplary embodiments of the composition of the present invention are as follows:

a composition comprising the present amide compound and alanycarb wherein the weight ratio of the present amide compound to alanycarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and alanycarb wherein the weight ratio of the present amide compound to alanycarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and alanycarb wherein the weight ratio of the present amide compound to alanycarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and aldicarb wherein the weight ratio of the present amide compound to aldicarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and aldicarb wherein the weight ratio of the present amide

compound to aldicarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and aldicarb wherein the weight ratio of the present amide compound to aldicarb is from 0.1/1 to 10/1;

5 a composition comprising the present amide compound and aldoxycarb wherein the weight ratio of the present amide compound to aldoxycarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and aldoxycarb wherein the weight ratio of the present amide compound to aldoxycarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and aldoxycarb wherein the weight ratio of the present amide compound to aldoxycarb is from 0.1/1 to 10/1;

15 a composition comprising the present amide compound and bendiocarb wherein the weight ratio of the present amide compound to bendiocarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and bendiocarb wherein the weight ratio of the present amide compound to bendiocarb is from 0.025/1 to 100/1;

20 a composition comprising the present amide compound and bendiocarb wherein the weight ratio of the present amide compound to bendiocarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and benfuracar wherein the weight ratio of the present amide compound to benfuracar is from 0.0125/1 to 500/1;

25

a composition comprising the present amide compound and benfuracar wherein the weight ratio of the present amide compound to benfuracar is from 0.025/1 to 100/1;

5 a composition comprising the present amide compound and benfuracar wherein the weight ratio of the present amide compound to benfuracar is from 0.1/1 to 10/1;

a composition comprising the present amide compound and benfuracarb wherein the weight ratio of the present amide compound to benfuracarb is from 0.0125/1 to 500/1;

10 a composition comprising the present amide compound and benfuracarb wherein the weight ratio of the present amide compound to benfuracarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and benfuracarb wherein the weight ratio of the present amide compound to benfuracarb is from 0.1/1 to 10/1;

15 a composition comprising the present amide compound and carbaryl wherein the weight ratio of the present amide compound to carbaryl is from 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and carbaryl wherein the weight ratio of the present amide compound to carbaryl is from 0.025/1 to 100/1;

a composition comprising the present amide compound and carbaryl wherein the weight ratio of the present amide compound to carbaryl is from 0.1/1 to 10/1;

25 a composition comprising the present amide compound

and carbofuran wherein the weight ratio of the present amide compound to carbofuran is from 0.0125/1 to 500/1;

5 a composition comprising the present amide compound and carbofuran wherein the weight ratio of the present amide compound to carbofuran is from 0.025/1 to 100/1;

a composition comprising the present amide compound and carbofuran wherein the weight ratio of the present amide compound to carbofuran is from 0.1/1 to 10/1;

10 a composition comprising the present amide compound and carbosulfan wherein the weight ratio of the present amide compound to carbosulfan is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and carbosulfan wherein the weight ratio of the present amide compound to carbosulfan is from 0.025/1 to 100/1;

15 a composition comprising the present amide compound and carbosulfan wherein the weight ratio of the present amide compound to carbosulfan is from 0.1/1 to 10/1;

20 a composition comprising the present amide compound and cloethocarb wherein the weight ratio of the present amide compound to cloethocarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and cloethocarb wherein the weight ratio of the present amide compound to cloethocarb is from 0.025/1 to 100/1;

25 a composition comprising the present amide compound and cloethocarb wherein the weight ratio of the present

amide compound to cloethocarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and ethiofencarb wherein the weight ratio of the present amide compound to ethiofencarb is from 0.0125/1 to 500/1;

5 a composition comprising the present amide compound and ethiofencarb wherein the weight ratio of the present amide compound to ethiofencarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and ethiofencarb wherein the weight ratio of the present amide compound to ethiofencarb is from 0.1/1 to 10/1;

10 a composition comprising the present amide compound and fenobucarb wherein the weight ratio of the present amide compound to fenobucarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and fenobucarb wherein the weight ratio of the present amide compound to fenobucarb is from 0.025/1 to 100/1;

15 a composition comprising the present amide compound and fenobucarb wherein the weight ratio of the present amide compound to fenobucarb is from 0.1/1 to 10/1;

20 a composition comprising the present amide compound and fenothiocarb wherein the weight ratio of the present amide compound to fenothiocarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and fenothiocarb wherein the weight ratio of the present amide compound to fenothiocarb is from 0.025/1 to 100/1;

25

a composition comprising the present amide compound and fenothiocarb wherein the weight ratio of the present amide compound to fenothiocarb is from 0.1/1 to 10/1;

5 a composition comprising the present amide compound and fenoxycarb wherein the weight ratio of the present amide compound to fenoxycarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and fenoxycarb wherein the weight ratio of the present amide compound to fenoxycarb is from 0.025/1 to 100/1;

10 a composition comprising the present amide compound and fenoxycarb wherein the weight ratio of the present amide compound to fenoxycarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and furathiocarb wherein the weight ratio of the present amide compound to furathiocarb is from 0.0125/1 to 500/1;

15 a composition comprising the present amide compound and furathiocarb wherein the weight ratio of the present amide compound to furathiocarb is from 0.025/1 to 100/1;

20 a composition comprising the present amide compound and furathiocarb wherein the weight ratio of the present amide compound to furathiocarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and isoprocarb wherein the weight ratio of the present amide compound to isoprocarb is from 0.0125/1 to 500/1;

25 a composition comprising the present amide compound

and isoprocarb wherein the weight ratio of the present amide compound to isoprocarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and isoprocarb wherein the weight ratio of the present amide compound to isoprocarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and metolcarb wherein the weight ratio of the present amide compound to metolcarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and metolcarb wherein the weight ratio of the present amide compound to metolcarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and metolcarb wherein the weight ratio of the present amide compound to metolcarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and methomyl wherein the weight ratio of the present amide compound to methomyl is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and methomyl wherein the weight ratio of the present amide compound to methomyl is from 0.025/1 to 100/1;

a composition comprising the present amide compound and methomyl wherein the weight ratio of the present amide compound to methomyl is from 0.1/1 to 10/1;

a composition comprising the present amide compound and methiocarb wherein the weight ratio of the present

amide compound to methiocarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and methiocarb wherein the weight ratio of the present amide compound to methiocarb is from 0.025/1 to 100/1;

5 a composition comprising the present amide compound and methiocarb wherein the weight ratio of the present amide compound to methiocarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and oxamyl wherein the weight ratio of the present amide
10 compound to oxamyl is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and oxamyl wherein the weight ratio of the present amide compound to oxamyl is from 0.025/1 to 100/1;

a composition comprising the present amide compound
15 and oxamyl wherein the weight ratio of the present amide compound to oxamyl is from 0.1/1 to 10/1;

a composition comprising the present amide compound and pirimicarb wherein the weight ratio of the present amide compound to pirimicarb is from 0.0125/1 to 500/1;

20 a composition comprising the present amide compound and pirimicarb wherein the weight ratio of the present amide compound to pirimicarb is from 0.025/1 to 100/1;

a composition comprising the present amide compound and pirimicarb wherein the weight ratio of the present
25 amide compound to pirimicarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and propoxur wherein the weight ratio of the present amide compound to propoxur is from 0.0125/1 to 500/1;

5 a composition comprising the present amide compound and propoxur wherein the weight ratio of the present amide compound to propoxur is from 0.025/1 to 100/1;

a composition comprising the present amide compound and propoxur wherein the weight ratio of the present amide compound to propoxur is from 0.1/1 to 10/1;

10 a composition comprising the present amide compound and XMC wherein the weight ratio of the present amide compound to XMC is from 0.0125/1 to 500/1;

15 a composition comprising the present amide compound and XMC wherein the weight ratio of the present amide compound to XMC is from 0.025/1 to 100/1;

a composition comprising the present amide compound and XMC wherein the weight ratio of the present amide compound to XMC is from 0.1/1 to 10/1;

20 a composition comprising the present amide compound and thiodicarb wherein the weight ratio of the present amide compound to thiodicarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and thiodicarb wherein the weight ratio of the present amide compound to thiodicarb is from 0.025/1 to 100/1;

25 a composition comprising the present amide compound

and thiodicarb wherein the weight ratio of the present amide compound to thiodicarb is from 0.1/1 to 10/1;

a composition comprising the present amide compound and xylylcarb wherein the weight ratio of the present amide compound to xylylcarb is from 0.0125/1 to 500/1;

a composition comprising the present amide compound and xylylcarb wherein the weight ratio of the present amide compound to xylylcarb is from 0.025/1 to 100/1; and

a composition comprising the present amide compound and xylylcarb wherein the weight ratio of the present amide compound to xylylcarb is from 0.1/1 to 10/1.

The method for controlling a pest of the present invention (hereinafter, referred to as the method for controlling of the present invention) comprises applying an effective amount in total of the present amide compound and the present carbamate compound(s) to the plants or the soil for cultivating the plant. Such plants include foliages of plants, seeds of plants, or bulbs of plants. The bulbs herein are intended to mean bulb, corm, rootstock, tuber, tuberous root and rhizophore.

In the method for controlling of the present invention, the present amide compound and the present carbamate compound(s) may be applied separately around the same time to the plant or the soil for cultivating the plant, but is usually applied as the composition of the present invention

for ease and convenience of application.

In the method for controlling of the present invention, examples of the method of applying the present amide compound and the present carbamate compound(s) include
5 foliage treatment, soil treatment, root treatment and seed treatment.

Such foliage treatment includes a method of applying the composition of the present invention to a surface of the plant to be cultivated by a foliage application or a
10 stem application.

Such root treatment includes a method of soaking a whole or a root of the plant in a medicinal solution comprising the present amide compound and the present carbamate compound(s), and a method of attaching a solid
15 formulation comprising the present amide compound, the present carbamate compound(s) and the solid carrier to a root of the plant.

Such soil treatment includes soil broadcast, soil incorporation, and irrigation of the medicinal solution to
20 a soil.

Such seed treatment includes an applying of the composition of the present invention to a seed or a bulb of the plant to be prevented from the plant disease. Specifically, a spray treatment by spraying a suspension of
25 the composition of the present invention in a mist form to

a surface of a seed or a surface of a bulb, a smear treatment by smearing the wettable powder, the emulsifiable concentrate or the flowable formulation of the composition of the present invention with an addition of small amounts
5 of water or alone to a seed or a bulb, an immersion treatment of a seed into a solution of the composition of the present invention for a given time, a film-coating treatment, and a pellet-coating treatment.

Each dose of the present amide compound and the
10 present carbamate compound(s) in the method for controlling of the present invention may vary depending on a kind of plant to be treated, a kind or a frequency of an occurrence of a plant disease as a control subject, a dosage form, a treatment period, a treatment method, a treatment site, a
15 climate condition, etc. In the case of application to foliage of the plant or soil for cultivating the plant, a total amount of the present amide compound and the present carbamate compound(s) is usually 1 to 500 g, preferably 2 to 200 g, and more preferably 10 to 100 g, per 1000 m².
20 Each dose of the present amide compound and the present carbamate compound(s) in the treatment for seed is usually 0.001 to 10 g, and preferably 0.01 to 1 g, per 1kg of seeds as a total amount of the present amide compound and the present carbamate compound(s).

The emulsifiable concentrate, the wetttable powder or the flowable formulation, etc., is usually applied by diluting them with water, and then spreading them. In this case, usually, each concentration of the present amide compound and the present carbamate compound(s) contains 0.0005 to 2% by weight, and preferably 0.005 to 1% by weight of the present amide compound and the present carbamate compound(s) in total. The dust formulation or the granular formulation, etc, is usually applied alone without dilution.

EXAMPLES

Next, the present invention is described in more detail below by the following examples including formulation examples and test examples, but the present invention should not be construed to be limited thereto.

The formulation examples are given below. It is to be noted that in the formulation examples, the term "'part'" indicates "'part by weight'".

Formulation 1

Five (5) parts of the present amide compound, 5 parts of alanycarb, 35 parts of the mixture of white carbon and polyoxyethylene alkylether sulfate ammonium salts (weight ratio 1:1), and 55 parts of water are mixed and the

resulting solution is then subjected to fine grinding according to wet grinding method, so as to obtain a flowable formulation. The same above operations are carried out using aldicarb, aldoxycarb, bendiocarb, 5 benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb or xylylcarb instead of alanycarb, so as to obtain flowable formulations.

10

Formulation 2

Ten (10) parts of the present amide compound, 5 parts of alanycarb and 1.5 parts of sorbitan trioleate are mixed into 28 parts of an aqueous solution that contains 2 parts 15 of polyvinyl alcohol, and the mixed solution is then subjected to fine grinding according to wet grinding method. Thereafter, 45.50 parts of an aqueous solution that contains 0.05 parts of xanthan gum and 0.1 part of aluminum magnesium silicate is added to the resultant product, and 20 10 parts of propylene glycol is further added thereto. The obtained mixture is blended by stirring, so as to obtain the flowable formulation. The same above operations are carried out using aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, 25 ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb,

furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb or xylylcarb instead of alanycarb, so as to obtain flowable formulations.

5 Formulation 3

Ten (10) parts of the present amide compound, 40 parts of alanycarb, 3 parts of calcium lignosulfonate, 2 parts of sodium lauryl sulfate, and 45 parts of synthetic hydrous silicon oxide are fully crushed and mixed, so as to obtain
10 wetable powders. The same above operations are carried out using aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb,
15 oxamyl, pirimicarb, propoxur, XMC, thiodicarb or xylylcarb instead of alanycarb, so as to obtain wettable powders.

The test examples are given below.

Test Examples 1 to 4

20 True leaf of cucumber was punched out with cork borer to 13mm in diameter to prepare a leaf disk. In a 24 well microwell plate that was dispensed with 1ml 0.8% water agar, the leaf disk was placed such that the upper side of the leaf was in an upward direction. Thereto was spread 20
25 microliters of a testing solution prepared by mixing the

present amide compound and carbofuran to a predetermined concentration (for treated group). Control where 20 microliters of ion-exchange water was spread was prepared (for non-treated group). After confirming that the spray solution was dried, conidium of gray mold fungus (*Botrytis cinerea*) was suspended into potato dextrose broth (DIFCO) in a density of about 10^5 conidia/mL and was then subjected to a spray inoculation. After leaving the leaf disk to stand in a growth chamber set up at 15°C for six days, an onset area on each leaf was measured and a preventive value was then calculated by the following equation 1.

The same above operations were carried out using thiodicarb instead of carbofuran, so as to obtain the preventive values.

(Equation 1)

$$\text{Preventive value (\%)} = 100 \times (A-B)/A$$

wherein

A: an onset area rate of plant belonging to non-treated group

B: an onset area rate of plant belonging to treated group

onset area rate = (onset area of the leaf disk)/(the total area of the leaf disk)

The results are shown in Tables 1 and 2.

Table 1

Ex. No.	treatment concentration (ppm)		preventive value (%)
	the present amide compound	carbofuran	
1	2.5	0.5	100
2	1.0	5.0	100

Table 2

Ex. No.	treatment concentration (ppm)		preventive value (%)
	the present amide compound	thiodicarb	
3	2.5	0.5	100
4	1.0	5.0	100

5 Test Examples 5 to 38 and Comparative Examples 1 to 34

The same above operations as described in Test Examples 1 to 4 were carried out using alanycarb, aldicarb, bendiocarb, benfuracarb, carbaryl, carbosulfan, ethiofencarb, fenobucarb, furathiocarb, isoprocarb, 10 metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, XMC or xylylcarb instead of carbofuran, so as to obtain the respective preventive values.

Also the same operations as described in Test Examples 1 to 4 were carried out except that the testing medicine 15 solution was substituted with a predetermined concentration of a dimethyl sulfoxide solution of the present amide compound, so as to calculate the respective preventive values.

The results are shown in Tables 3 to 19.

Table 3

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	alanycarb	
Ex.No.5	2.5	0.5	100
Ex.No.6	1.0	5.0	100
Comp.Ex.No.1	2.5	-	56
Comp.Ex.No.2	1.0	-	46

Table 4

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	aldicarb	
Ex.No.7	2.5	0.5	100
Ex.No.8	1.0	5.0	100
Comp.Ex.No.3	2.5	-	56
Comp.Ex.No.4	1.0	-	46

5 Table 5

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	bendiocarb	
Ex.No.9	2.5	0.5	100
Ex.No.10	1.0	5.0	100
Comp.Ex.No.5	2.5	-	56
Comp.Ex.No.6	1.0	-	46

Table 6

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	benfuracarb	
Ex.No.11	2.5	0.5	100
Ex.No.12	1.0	5.0	100
Comp.Ex.No.7	2.5	-	56
Comp.Ex.No.8	1.0	-	46

Table 7

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	carbaryl	
Ex.No.13	2.5	0.5	100
Ex.No.14	1.0	5.0	100
Comp.Ex.No.9	2.5	-	56
Comp.Ex.No.10	1.0	-	46

Table 8

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	carbosulfan	
Ex.No.15	2.5	0.5	100
Ex.No.16	1.0	5.0	100
Comp.Ex.No.11	2.5	-	56
Comp.Ex.No.12	1.0	-	46

5 Table 9

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	ethiofencarb	
Ex.No.17	2.5	0.5	100
Ex.No.18	1.0	5.0	100
Comp.Ex.No.13	2.5	-	56
Comp.Ex.No.14	1.0	-	46

Table 10

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	fenobucarb	
Ex.No.19	2.5	0.5	100
Ex.No.20	1.0	5.0	100
Comp.Ex.No.15	2.5	-	56
Comp.Ex.No.16	1.0	-	46

Table 11

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	furathiocarb	
Ex.No.21	2.5	0.5	100
Ex.No.22	1.0	5.0	100
Comp.Ex.No.17	2.5	-	56
Comp.Ex.No.18	1.0	-	46

Table 12

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	isoproc carb	
Ex.No.23	2.5	0.5	100
Ex.No.24	1.0	5.0	100
Comp.Ex.No.19	2.5	-	56
Comp.Ex.No.20	1.0	-	46

5 Table 13

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	metolcarb	
Ex.No.25	2.5	0.5	100
Ex.No.26	1.0	5.0	100
Comp.Ex.No.21	2.5	-	56
Comp.Ex.No.22	1.0	-	46

Table 14

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	methomyl	
Ex.No.27	2.5	0.5	100
Ex.No.28	1.0	5.0	100
Comp.Ex.No.23	2.5	-	56
Comp.Ex.No.24	1.0	-	46

Table 15

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	methiocarb	
Ex.No.29	2.5	0.5	100
Ex.No.30	1.0	5.0	100
Comp.Ex.No.25	2.5	-	56
Comp.Ex.No.26	1.0	-	46

Table 16

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	oxamyl	
Ex.No.31	2.5	0.5	100
Ex.No.32	1.0	5.0	100
Comp.Ex.No.27	2.5	-	56
Comp.Ex.No.28	1.0	-	46

5 Table 17

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	pirimicarb	
Ex.No.33	2.5	0.5	100
Ex.No.34	1.0	5.0	100
Comp.Ex.No.29	2.5	-	56
Comp.Ex.No.30	1.0	-	46

Table 18


	treatment concentration (ppm)		preventive value (%)
	the present amide compound	XMC	
Ex.No.35	2.5	0.5	100
Ex.No.36	1.0	5.0	100
Comp.Ex.No.31	2.5	-	56
Comp.Ex.No.32	1.0	-	46

Table 19

	treatment concentration (ppm)		preventive value (%)
	the present amide compound	xylylcarb	
Ex.No.37	2.5	0.5	100
Ex.No.38	1.0	5.0	100
Comp.Ex.No.33	2.5	-	56
Comp.Ex.No.34	1.0	-	46

CLAIMS

1. A fungal pest controlling composition comprising a compound represented by Formula (1):



5

and one or more carbamate compound(s) selected from Group (A):

Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoprocarb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb and xylylcarb.

10


2. The fungal pest controlling composition according to claim 1, wherein the weight ratio of the compound represented by Formula (1) as defined in claim 1 to the carbamate compound(s) is from 0.0125/1 to 500/1.

15

3. The fungal pest controlling composition according to claim 1 or 2, wherein the compound represented by Formula (1) as defined in claim 1 has R-absolute configuration.

20


4. A method for controlling a fungal pest, wherein the method comprises applying a compound of Formula (1):



and one or more carbamate compound(s) selected from Group (A) to a plant or a soil for cultivating the plant,

Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoproc carb, metolcarb, methomyl, methiocarb, oxamyl, pirimicarb, propoxur, XMC, thiodicarb and xylylcarb.

5. A method for controlling a fungal pest, wherein the method comprises applying a compound represented by Formula (1):



and one or more carbamate compound(s) selected from Group (A) to a seed,


Group (A): a group consisting of alanycarb, aldicarb, aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran, carbosulfan, cloethocarb, ethiofencarb, fenobucarb, fenothiocarb, fenoxycarb, furathiocarb, isoproc carb,

metolcarb, methomyl, methiocarb, oxamyl, pirimicarb,
propoxur, XMC, thiodicarb and xylylcarb.

6. The method according to claim 4 or 5, wherein the
weight ratio of the compound represented by Formula (1) as
5 defined in claim 1 to the carbamate compound(s) is from
0.0125/1 to 500/1.


7. The method according to any one of claims 4 to 6,
wherein the compound represented by Formula (1) as defined
in claim 1 has R-absolute configuration.

10 8. Use of a combination of a compound represented by
Formula (1):



and one or more carbamate compound(s) selected from Group (A)
for controlling a fungal pest,

15 Group (A): a group consisting of alanycarb, aldicarb,
aldoxycarb, bendiocarb, benfuracarb, carbaryl, carbofuran,
carbosulfan, cloethocarb, ethiofencarb, fenobucarb,
fenoethiocarb, fenoxycarb, furathiocarb, isoprocarb,
metolcarb, methomyl, methiocarb, oxamyl, pirimicarb,
20 propoxur, XMC, thiodicarb and xylylcarb.



(1)