



(51) International Patent Classification:

A61K 31/195 (2006.01) A61K 31/4402 (2006.01)
A61K 31/44 (2006.01) A61P 3/00 (2006.01)

(21) International Application Number:

PCT/EP2013/057729

(22) International Filing Date:

12 April 2013 (12.04.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

12425073.9 18 April 2012 (18.04.2012) EP
61/672,931 18 July 2012 (18.07.2012) US

(71) Applicant: **NOGRA PHARMA LIMITED** [IE/IE]; 33,
Sir John Rogerson's Quay, Dublin, 2 (IE).

(72) Inventors: **BARONI, Sergio**; Via Piazzola 3, I-24030
Villa D'Adda (BG) (IT). **BELLINIA, Salvatore**; Via
Brusafiera 12, I-33170 Pordenone (IT). **VITI, Francesca**;
Via Benedetto Croce 44, Sesto San Giovanni (IT).

(74) Agents: **LANE, Cathal Michael** et al.; Tomkins & Co., 5
Dartmouth Road, Dublin, 6 (IE).


(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).


Published:

— with international search report (Art. 21(3))

(54) Title: METHODS OF TREATING LACTOSE INTOLERANCE

FIGURE 1
A

B



(57) Abstract: Disclosed herein are methods for treating lactose intolerance, including compounds that may be specific or modulate PPAR γ receptors.

METHODS OF TREATING LACTOSE INTOLERANCE

RELATED APPLICATIONS

[0001] This application claims priority to EP12425073.9, filed April 18, 2012 and U.S.S.N 61/672,931, filed July 18, 2012, both of which are incorporated by reference in their entireties.

BACKGROUND

[0002] Lactase protein is a disaccharidase (β -galactosidase) expressed on the tips of the villi of the small intestine having the ability to hydrolyze lactose into galactose and glucose. Inadequate lactase-phlorizin hydrolase (LPH) activity is responsible for lactose intolerance/malabsorption leading to diarrhea, abdominal pain or bloating after lactose ingestion. Primary lactase deficiency (or lactase non persistence or hypolactasia) is the main cause of lactose intolerance, due to the relative or absolute absence of lactase expression in the small bowel, occurring in childhood at various ages and in different racial groups. Approximately 70% of the world's population has primary lactase deficiency. The percentage of lactose deficiency varies according to ethnicity and is related to the use of dairy products in the diet reaching up to 20% of North European, 40% of Mediterranean European, 80% of Africans, and 90% of Asian population. Typical treatments for lactose intolerance is lactose exclusion (leading to nutritional impairment) or expansive regimen such as the use of lactose deficient milk or lactase supplementation.

[0003] It has been reported that two particular Single Polymorphisme (SNP) nucleotide are tightly associated with adult-type hypolactasia. A C at position _13910 (C₁₃₉₁₀) upstream of the lactase gene is 100% associated and a G at position _22018 (G₂₂₀₁₈) is more than 95% associated with lactase non-persistence in the Finnish population. SNP. Expression of LPH mRNA in the intestinal mucosa in individuals with T₁₃₉₁₀ and A₂₂₀₁₈ is higher than found in individuals with C₁₃₉₁₀ and G₂₂₀₁₈, suggesting a transcriptional regulation of LPH gene. However, much of the regulation of the LPH gene remains unknown. Accordingly, effective agents, that are useful in the treatment of lactose intolerance and related disorders are needed.

[0004] Peroxisome Proliferator Activated Receptors (PPARs) are members of the nuclear hormone receptor super family, which are ligand-activated transcription factors regulating gene expression. PPARs play a role in the regulation of cell differentiation, development and metabolism of higher organisms.

[0005] Three types of PPAR have been identified: alpha, expressed in the liver, kidney, heart and other tissues and organs, beta/delta expressed for example in the brain, and gamma, expressed in three forms: gamma1, gamma2, and gamma3. PPAR γ receptors have been associated with stimulation of keratinocyte differentiation, and has served as a potential drug target for a number of disease states including skin disorders such as psoriasis and atopic dermatitis.

SUMMARY

[0006] This disclosure is generally directed methods of treating, ameliorating or substantially preventing lactose intolerance in a patient need thereof, comprising administering a composition that includes a PPAR modulator, e.g. PPAR α , δ or γ , e.g., a PPAR γ modulator, e.g., agonist (e.g. such as a compound disclosed herein) to a patient in need thereof. Also provided herein is a method for stimulating lactose gene expression in a patient in need thereof, comprising administering a PPAR agonist, e.g., a PPAR α , δ , or PPAR γ agonist to the patient.

[0007] Also provided herein is a method for treating diarrhea, abdominal pain and/or bloating after lactose ingestion in a patient in need thereof, comprising administering a PPAR agonist, e.g. a PPAR α , PPAR δ , PPAR γ agonist. For example, provided herein is a method for ameliorating lactose intolerance in a patient ingesting a composition that includes lactose, comprising administering a PPAR agonist before, substantially simulatenously with, or after said ingestion.

BRIEF DESCRIPTION OF THE FIGURES

[0008] Figure 1A depicts gene expression evaluated in qPCR in synchronized Caco2 cells after 24hours of treatment by compound 34, 1mM. Confirmation of the 12 up-regulated

genes expression observed in microarrays. Up regulated expression of AQP12, OST α and LZTS1 are not confirmed. Results are expressed in fold compared to control (set to value 1).

[0009] Figure 1B depicts gene expression evaluated in qPCR in synchronized Caco2 cells after 24hours of treatment by compound 34, 1mM. Confirmation of the 12 up-regulated genes expression observed in microarrays. Up regulated expression of AQP12, OST α and LZTS1 are not confirmed. Results are expressed in relative expression.

[0010] Figure 2A depicts gene expression evaluated in qPCR in synchronized Caco2 cells after 24hours of treatment by compound 34, 1mM. Confirmation of all the 15 up-regulated genes expression observed in microarrays. Results are expressed in fold, compared to control (set to value 1).

[0011] Figure 2B depicts gene expression evaluated in qPCR in synchronized Caco2 cells after 24hours of treatment by compound 34 1mM. Confirmation of all the 15 up-regulated genes expression observed in microarrays. Results are expressed in relative expression.

[0012] FIGURE 3 depicts gene expression expressed in fold ,of stimulation 2 are correlated of stimulation 1 microarrays results ($r=0.754$, $p=0.0046$).

[0013] Figure 4 shows that Lactase mRNA expression is induced by PPAR. Caco-2 cells were stimulated with various PPAR γ and PPAR α agonists. Statistical differences were analyzed with GraphPad software using a non-parametric Mann-Whitney test.

[0014] Figure 5: Identification of direct repeats in the promoter (3000pb) of the human Lactase gene. DR1 are in bold, and DR2 are in bold and start with "AG". Also indicated are TATA box and lowercases letters represent the beginning of the first exon.

DETAILED DESCRIPTION

[0015] The features and other details of the disclosure will now be more particularly described. Before further description of the present invention, certain terms employed in the specification, examples and appended claims are collected here. These definitions should be read in light of the remainder of the disclosure and understood as by a person of skill in the art. Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art.

Definitions

[0016] “Treating” includes any effect, e.g., lessening, reducing, modulating, or eliminating, that results in the improvement of the condition, disease, disorder and the like.

[0017] The term “alkenyl” as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon double bond, such as a straight or branched group of 2-12, 2-10, or 2-6 carbon atoms, referred to herein as C₂-C₁₂alkenyl, C₂-C₁₀alkenyl, and C₂-C₆alkenyl, respectively. Exemplary alkenyl groups include, but are not limited to, vinyl, allyl, butenyl, pentenyl, hexenyl, butadienyl, pentadienyl, hexadienyl, 2-ethylhexenyl, 2-propyl-2-butenyl, 4-(2-methyl-3-butene)-pentenyl, etc.

[0018] The term “alkoxy” as used herein refers to an alkyl group attached to an oxygen (-O-alkyl-). Exemplary alkoxy groups include, but are not limited to, groups with an alkyl, alkenyl or alkynyl group of 1-12, 1-8, or 1-6 carbon atoms, referred to herein as C₁-C₁₂alkoxy, C₁-C₈alkoxy, and C₁-C₆alkoxy, respectively. Exemplary alkoxy groups include, but are not limited to methoxy, ethoxy, etc. Similarly, exemplary “alkenoxy” groups include, but are not limited to vinyloxy, allyloxy, butenoxy, etc.

[0019] The term “alkyl” as used herein refers to a saturated straight or branched hydrocarbon, such as a straight or branched group of 1-12, 1-10, or 1-6 carbon atoms, referred to herein as C₁-C₁₂alkyl, C₁-C₁₀alkyl, and C₁-C₆alkyl, respectively. Exemplary alkyl groups include, but are not limited to, methyl, ethyl, propyl, isopropyl, 2-methyl-1-propyl, 2-methyl-2-propyl, 2-methyl-1-butyl, 3-methyl-1-butyl, 2-methyl-3-butyl, 2,2-dimethyl-1-propyl, 2-methyl-1-pentyl, 3-methyl-1-pentyl, 4-methyl-1-pentyl, 2-methyl-2-pentyl, 3-methyl-2-pentyl, 4-methyl-2-pentyl, 2,2-dimethyl-1-butyl, 3,3-dimethyl-1-butyl, 2-ethyl-1-butyl, butyl, isobutyl, t-butyl, pentyl, isopentyl, neopentyl, hexyl, heptyl, octyl, etc. In certain embodiments, alkyl refers to C₁-C₆alkyl. In certain embodiments, cycloalkyl refers to C₃-C₆cycloalkyl.

[0020] Alkyl, alkenyl and alkynyl groups can, in some embodiments, be optionally be substituted with or interrupted by at least one group selected from alkanoyl, alkoxy, alkyl, alkenyl, alkynyl, amido, amidino, amino, aryl, arylalkyl, azido, carbamate, carbonate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, imino, ketone, nitro, phosphate, phosphonato, phosphinato, sulfate, sulfide, sulfonamido, sulfonyl and thiocarbonyl.

[0021] The term “alkynyl” as used herein refers to an unsaturated straight or branched hydrocarbon having at least one carbon-carbon triple bond, such as a straight or branched group

of 2-12, 2-8, or 2-6 carbon atoms, referred to herein as C₂-C₁₂alkynyl, C₂-C₈alkynyl, and C₂-C₆alkynyl, respectively. Exemplary alkynyl groups include, but are not limited to, ethynyl, propynyl, butynyl, pentynyl, hexynyl, methylpropynyl, 4-methyl-1-butynyl, 4-propyl-2-pentynyl, and 4-butyl-2-hexynyl, etc.

[0022] The term “aryl” as used herein refers to a mono-, bi-, or other multi-carbocyclic, aromatic ring system. In certain embodiments, aryl refers to a monocyclic and/or bicyclic, 5 to 6 membered ring. The aromatic ring may be substituted at one or more ring positions with substituents selected from alkanoyl, alkoxy, alkyl, alkenyl, alkynyl, amido, amidino, amino, aryl, arylalkyl, azido, carbamate, carbonate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, imino, ketone, nitro, phosphate, phosphonato, phosphinato, sulfate, sulfide, sulfonamido, sulfonyl and thiocarbonyl. The term “aryl” also includes polycyclic ring systems having two or more cyclic rings in which two or more carbons are common to two adjoining rings (the rings are “fused rings”) wherein at least one of the rings is aromatic, e.g., the other cyclic rings may be cycloalkyls, cycloalkenyls, cycloalkynyls, and/or aryls. Exemplary aryl groups include, but are not limited to, phenyl, tolyl, anthracenyl, fluorenyl, indenyl, azulenyl, and naphthyl, as well as benzo-fused carbocyclic moieties such as 5,6,7,8-tetrahydronaphthyl.

[0023] The term “carboxy” as used herein refers to the radical -COOH or its corresponding salts, e.g. -COONa, etc.

[0024] The term “cycloalkyl” as used herein refers to a monovalent saturated or unsaturated cyclic, bicyclic, or bridged bicyclic hydrocarbon group of 3-12, 3-8, 4-8, or 4-6 carbons, referred to herein, e.g., as “C₄₋₈cycloalkyl,” derived from a cycloalkane. Exemplary cycloalkyl groups include, but are not limited to, cyclohexanes, cyclohexenes, cyclopentanes, cyclopentenenes, cyclobutanes and cyclopropanes. Cycloalkyl groups may be substituted with alkanoyl, alkoxy, alkyl, alkenyl, alkynyl, amido, amidino, amino, aryl, arylalkyl, azido, carbamate, carbonate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, imino, ketone, nitro, phosphate, phosphonato, phosphinato, sulfate, sulfide, sulfonamido, sulfonyl and thiocarbonyl. Cycloalkyl groups can be fused to other cycloalkyl, aryl, or heterocyclyl groups. In certain embodiments, cycloalkyl refers to C₃-C₆ alkyl.

[0025] The terms “halo” or “halogen” or “Hal” as used herein refer to F, Cl, Br, or I.

[0026] The term “haloalkyl” as used herein refers to an alkyl group substituted with one or more halogen atoms.

[0027] The term “phenyl” as used herein refers to a 6-membered carbocyclic aromatic ring. The phenyl group can also be fused to a cyclohexane or cyclopentane ring. Phenyl can be substituted with one or more substituents including alkanoyl, alkoxy, alkyl, alkenyl, alkynyl, amido, amidino, amino, aryl, arylalkyl, azido, carbamate, carbonate, carboxy, cyano, cycloalkyl, ester, ether, formyl, halogen, haloalkyl, heteroaryl, heterocyclyl, hydroxyl, imino, ketone, nitro, phosphate, phosphonato, phosphinato, sulfate, sulfide, sulfonamido, sulfonyl and thiocarbonyl.

[0028] The term “pharmaceutically acceptable carrier” or “pharmaceutically acceptable excipient” as used herein refers to any and all solvents, dispersion media, coatings, isotonic and absorption delaying agents, and the like, that are compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. The compositions may also contain other active compounds providing supplemental, additional, or enhanced therapeutic functions.

[0029] The term “pharmaceutical composition” as used herein refers to a composition comprising at least one compound as disclosed herein formulated together with one or more pharmaceutically acceptable carriers.

[0030] “Individual,” “patient,” or “subject” are used interchangeably and include to any animal, including mammals, preferably mice, rats, other rodents, rabbits, dogs, cats, swine, cattle, sheep, horses, or primates, and most preferably humans. The compounds of the invention can be administered to a mammal, such as a human, but can also be other mammals such as an animal in need of veterinary treatment, *e.g.*, domestic animals (*e.g.*, dogs, cats, and the like), farm animals (*e.g.*, cows, sheep, pigs, horses, and the like) and laboratory animals (*e.g.*, rats, mice, guinea pigs, and the like). The mammal treated in the methods of the invention is desirably a mammal in whom modulation of PPAR receptors is desired. “Modulation” includes antagonism (*e.g.*, inhibition), agonism, partial antagonism and/or partial agonism.

[0031] In the present specification, the term “therapeutically effective amount” means the amount of the subject compound that will elicit the biological or medical response of a tissue, system, animal or human that is being sought by the researcher, veterinarian, medical doctor or other clinician. The compounds of the invention are administered in therapeutically effective amounts to treat a disease. Alternatively, a therapeutically effective amount of a compound is

the quantity required to achieve a desired therapeutic and/or prophylactic effect, such as an amount which results in the prevention of or a decrease in the symptoms associated with a disease associated with PPAR receptors.

[0032] The term "pharmaceutically acceptable salt(s)" as used herein refers to salts of acidic or basic groups that may be present in compounds used in the present compositions. Compounds included in the present compositions that are basic in nature are capable of forming a wide variety of salts with various inorganic and organic acids. The acids that may be used to prepare pharmaceutically acceptable acid addition salts of such basic compounds are those that form non-toxic acid addition salts, i.e., salts containing pharmacologically acceptable anions, including but not limited to malate, oxalate, chloride, bromide, iodide, nitrate, sulfate, bisulfate, phosphate, acid phosphate, isonicotinate, acetate, lactate, salicylate, citrate, tartrate, oleate, tannate, pantothenate, bitartrate, ascorbate, succinate, maleate, gentisinate, fumarate, gluconate, glucuronate, saccharate, formate, benzoate, glutamate, methanesulfonate, ethanesulfonate, benzenesulfonate, *p*-toluenesulfonate and pamoate (i.e., 1,1'-methylene-bis-(2-hydroxy-3-naphthoate)) salts. Compounds included in the present compositions that include an amino moiety may form pharmaceutically acceptable salts with various amino acids, in addition to the acids mentioned above. Compounds included in the present compositions that are acidic in nature are capable of forming base salts with various pharmacologically acceptable cations. Examples of such salts include alkali metal or alkaline earth metal salts and, particularly, calcium, magnesium, sodium, lithium, zinc, potassium, and iron salts.

[0033] The compounds of the disclosure may contain one or more chiral centers and/or double bonds and, therefore, exist as stereoisomers, such as geometric isomers, enantiomers or diastereomers. The term "stereoisomers" when used herein consist of all geometric isomers, enantiomers or diastereomers. These compounds may be designated by the symbols "R" or "S," depending on the configuration of substituents around the stereogenic carbon atom. The present invention encompasses various stereoisomers of these compounds and mixtures thereof. Stereoisomers include enantiomers and diastereomers. Mixtures of enantiomers or diastereomers may be designated "(±)" in nomenclature, but the skilled artisan will recognize that a structure may denote a chiral center implicitly.

[0034] Individual stereoisomers of compounds of the present invention can be prepared synthetically from commercially available starting materials that contain asymmetric or

stereogenic centers, or by preparation of racemic mixtures followed by resolution methods well known to those of ordinary skill in the art. These methods of resolution are exemplified by (1) attachment of a mixture of enantiomers to a chiral auxiliary, separation of the resulting mixture of diastereomers by recrystallization or chromatography and liberation of the optically pure product from the auxiliary, (2) salt formation employing an optically active resolving agent, or (3) direct separation of the mixture of optical enantiomers on chiral chromatographic columns. Stereoisomeric mixtures can also be resolved into their component stereoisomers by well known methods, such as chiral-phase gas chromatography, chiral-phase high performance liquid chromatography, crystallizing the compound as a chiral salt complex, or crystallizing the compound in a chiral solvent. Stereoisomers can also be obtained from stereomerically-pure intermediates, reagents, and catalysts by well known asymmetric synthetic methods.

[0035] Geometric isomers can also exist in the compounds of the present invention. The symbol $\equiv$ denotes a bond that may be a single, double or triple bond as described herein. The present invention encompasses the various geometric isomers and mixtures thereof resulting from the arrangement of substituents around a carbon-carbon double bond or arrangement of substituents around a carbocyclic ring. Substituents around a carbon-carbon double bond are designated as being in the “*Z*” or “*E*” configuration wherein the terms “*Z*” and “*E*” are used in accordance with IUPAC standards. Unless otherwise specified, structures depicting double bonds encompass both the “*E*” and “*Z*” isomers.

[0036] Substituents around a carbon-carbon double bond alternatively can be referred to as “cis” or “trans,” where “cis” represents substituents on the same side of the double bond and “trans” represents substituents on opposite sides of the double bond. The arrangement of substituents around a carbocyclic ring are designated as “cis” or “trans.” The term “cis” represents substituents on the same side of the plane of the ring and the term “trans” represents substituents on opposite sides of the plane of the ring. Mixtures of compounds wherein the substituents are disposed on both the same and opposite sides of plane of the ring are designated “cis/trans.”

[0037] The compounds disclosed herein can exist in solvated as well as unsolvated forms with pharmaceutically acceptable solvents such as water, ethanol, and the like, and it is intended that the invention embrace both solvated and unsolvated forms. In one embodiment, the

compound is amorphous. In one embodiment, the compound is a polymorph. In another embodiment, the compound is in a crystalline form.

[0038] The invention also embraces isotopically labeled compounds of the invention which are identical to those recited herein, except that one or more atoms are replaced by an atom having an atomic mass or mass number different from the atomic mass or mass number usually found in nature. Examples of isotopes that can be incorporated into compounds of the invention include isotopes of hydrogen, carbon, nitrogen, oxygen, phosphorus, fluorine and chlorine, such as ^2H , ^3H , ^{13}C , ^{14}C , ^{15}N , ^{18}O , ^{17}O , ^{31}P , ^{32}P , ^{35}S , ^{18}F , and ^{36}Cl , respectively.

[0039] Certain isotopically-labeled disclosed compounds (*e.g.*, those labeled with ^3H and ^{14}C) are useful in compound and/or substrate tissue distribution assays. Tritiated (*i.e.*, ^3H) and carbon-14 (*i.e.*, ^{14}C) isotopes are particularly preferred for their ease of preparation and detectability. Further, substitution with heavier isotopes such as deuterium (*i.e.*, ^2H) may afford certain therapeutic advantages resulting from greater metabolic stability (*e.g.*, increased *in vivo* half-life or reduced dosage requirements) and hence may be preferred in some circumstances. Isotopically labeled compounds of the invention can generally be prepared by following procedures analogous to those disclosed in the *e.g.*, Examples herein by substituting an isotopically labeled reagent for a non-isotopically labeled reagent.

[0040] The term “prodrug” refers to compounds that are transformed *in vivo* to yield a disclosed compound or a pharmaceutically acceptable salt, hydrate or solvate of the compound. The transformation may occur by various mechanisms, such as through hydrolysis in blood. For example, if a compound of the invention or a pharmaceutically acceptable salt, hydrate or solvate of the compound contains a carboxylic acid functional group, a prodrug can comprise an ester formed by the replacement of the hydrogen atom of the acid group with a group such as (C₁-C₈)alkyl, (C₂-C₁₂)alkanoyloxymethyl, 1-(alkanoyloxy)ethyl having from 4 to 9 carbon atoms, 1-methyl-1-(alkanoyloxy)-ethyl having from 5 to 10 carbon atoms, alkoxycarbonyloxymethyl having from 3 to 6 carbon atoms, 1-(alkoxycarbonyloxy)ethyl having from 4 to 7 carbon atoms, 1-methyl-1-(alkoxycarbonyloxy)ethyl having from 5 to 8 carbon atoms, N-(alkoxycarbonyl)aminomethyl having from 3 to 9 carbon atoms, 1-(N-(alkoxycarbonyl)amino)ethyl having from 4 to 10 carbon atoms, 3-phthalidyl, 4-crotonolactonyl, gamma-butyrolacton-4-yl, di-N,N-(C₁-C₂)alkylamino(C₂-C₃)alkyl (such as β-

dimethylaminoethyl), carbamoyl-(C₁-C₂)alkyl, N,N-di(C₁-C₂)alkylcarbamoyl-(C₁-C₂)alkyl and piperidino-, pyrrolidino- or morpholino(C₂-C₃)alkyl.


[0041] Similarly, if a compound of the invention contains an alcohol functional group, a prodrug can be formed by the replacement of the hydrogen atom of the alcohol group with a group such as (C₁-C₆)alkanoyloxymethyl, 1-((C₁-C₆)alkanoyloxy)ethyl, 1-methyl-1-((C₁-C₆)alkanoyloxy)ethyl (C₁-C₆)alkoxycarbonyloxymethyl, N-(C₁-C₆)alkoxycarbonylaminomethyl, succinoyl, (C₁-C₆)alkanoyl, α -amino(C₁-C₄)alkanoyl, arylacyl and α -aminoacyl, or α -aminoacyl- α -aminoacyl, where each α -aminoacyl group is independently selected from the naturally occurring L-amino acids, P(O)(OH)₂, -P(O)(O(C₁-C₆)alkyl)₂ or glycosyl (the radical resulting from the removal of a hydroxyl group of the hemiacetal form of a carbohydrate).

[0042] If a compound of the invention incorporates an amine functional group, a prodrug can be formed by the replacement of a hydrogen atom in the amine group with a group such as R-carbonyl, RO-carbonyl, NRR'-carbonyl where R and R' are each independently (C₁-C₁₀)alkyl, (C₃-C₇)cycloalkyl, benzyl, or R-carbonyl is a natural α -aminoacyl or natural α -aminoacyl-natural α -aminoacyl, —C(OH)C(O)OY¹ wherein Y¹ is H, (C₁-C₆)alkyl or benzyl, -C(OY²)Y³ wherein Y² is (C₁-C₄) alkyl and Y³ is (C₁-C₆)alkyl, carboxy(C₁-C₆)alkyl, amino(C₁-C₄)alkyl or mono-N— or di-N,N—(C₁-C₆)alkylaminoalkyl, —C(Y⁴)Y⁵ wherein Y⁴ is H or methyl and Y⁵ is mono-N— or di-N,N—(C₁-C₆)alkylamino, morpholino, piperidin-1-yl or pyrrolidin-1-yl.

PPARmodulators

[0043] The disclosure provides, at least in part, compounds having PPAR modulating activity, e.g., PPAR γ modulating activity. Such compounds, may, for example, be represented by formulas depicted below. Also contemplated herein are compositions that include a compound represented by the disclosed formulas and e.g., a pharmaceutically or cosmetically acceptable carrier or excipient.

[0044] For example, in an embodiment, contemplated PPAR modulating compounds (e.g. compounds having PPAR activity) may be represented by:



and pharmaceutically acceptable salts, stereoisomers, prodrugs thereof, wherein:

R_1 and R_2 , are each independently selected from the group consisting of H and C_{1-6} alkyl; or R_1 and R_2 together with the nitrogen atom they are bonded to form an aromatic or aliphatic ring with 5 or 6 atoms which may be optionally substituted;



Y and Z are each independently selected from the group consisting of H, OH, COOH, $-OR_3$, $-CH(OR_3)COOH$; and

R_3 is selected from the group consisting of H, phenyl, benzyl, vinyl, allyl, C_{1-6} alkyl or C_{1-6} alkyl substituted by one, two, three or more halogens.

[0045] In an embodiment, Y may be H or COOH. For example, Y may be H and Z may be $CH(OR_3)COOH$, or Y may be COOH and Z may be $-OR_3$. In some embodiments, R_3 may be methyl, ethyl, n-propyl, or isopropyl.

[0046] In other embodiments, the NR_1R_2 moiety may be in the 4' position or may be in the 3' position. In certain embodiments, R_1 and R_2 are H.

[0047] Exemplary compounds also include those represented by formulas IIa or IIb or a pharmaceutically acceptable salt, enantiomer or stereoisomer of:



wherein:

R_1 and R_2 are each independently selected from the group consisting of H and C_{1-6} alkyl; or R_1 and R_2 together, with the nitrogen atom they are bonded to, form an aromatic or aliphatic ring with 5 or 6 atoms;

R_6 is selected from the group consisting of: $-NHOH$, OH , and $-OR_9$;

R_9 is C_{1-6} alkyl;

R_4 is selected from H, phenyl, benzyl, vinyl, allyl, C_{1-6} alkyl or C_{1-6} alkyl substituted by one or more halogens;

R_5 and R_7 are each independently hydrogen or halo, or ;


or

[0048] R_4 and R_5 , or R_4 and R_6 together, form a fused heterocyclic ring with 5 or 6 atoms, optionally substituted with halo or C_{1-6} alkyl; and A is a fused heterocyclic ring; or a pharmaceutically acceptable salt thereof.

[0049] In certain embodiments, the NR_1R_2 moiety of formula IIa may be in the 4' position or may be in the 3' position. In certain embodiments, R_1 and R_2 are H.


[0050] R_9 , in some embodiments, may be methyl, ethyl, n-propyl, or isopropyl.

[0051] In some embodiments a compound can be represented by



wherein p is 1 or 2, R_6 is OH or $-OR_9$, wherein R_9 is defined above, and R_{10} , independently for each occurrence, is selected from the group consisting of H, halo, or C_{1-6} alkyl, e.g. methyl or ethyl.


[0052] Exemplary compounds contemplated herein include:




acceptable salt thereof.


[0053] In some embodiments, contemplated compounds include: 4-amino-N-hydroxy-2-methoxybenzamide (compound 13); 6-methoxy quinoline-5-carboxylic acid (compound 36); 6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid (compound 37); 5-diisopropylaminosalicylic acid (compound 38).

[0054] Other exemplary compounds include those represented by:





(compound 31):



; (compound 28): .

[0055] Compounds contemplated herein include racemic mixtures, and enantiomers of compounds, for example: (±)-2-hydroxy-3-(3'-aminophenyl) propionic acid (compound 20); (±)-2-methoxy-2-(4'-aminophenyl) acetic acid (compound 23); (±)-2-ethoxy-2-(3'-aminophenyl) acetic acid (compound 32); (±)-2-ethoxy-2-(4'-aminophenyl) acetic acid (compound 33); (±)-2-methoxy-3-(4'-aminophenyl) propionic acid (compound 34) "±34" (racemic form); (±)-2-ethoxy-3-(4'-aminophenyl) propionic acid (compound 39); (±)-2-ethoxy-3-(3'-aminophenyl) propionic acid (compound 40).

[0056] For example, the compounds used in the methods of the present invention can be enantiomers of the following racemic mixtures: (R,S)-2-hydroxy-2-(3-aminophenyl)acetic acid (compound 10); (R,S)-2-hydroxy-2-(4-aminophenyl)acetic acid (compound 11); (R,S)-2-hydroxy-3-(4'-aminophenyl)propionic acid (compound 21); (R,S)-2-methoxy-2-(3'-aminophenyl)acetic acid (compound 22); (R,S)-2-methoxy-3-(3'-aminophenyl)propionic acid (compound 35); (R,S)-2-methoxy-3-(4-aminophenyl)propionic acid (compound 34), as well as enantiomers, e.g.: (+) 2-S-methoxy-3-(4-aminophenyl)propionic acid (compound 34); (-) 2-R-methoxy-3-(4-aminophenyl)propionic acid (compound 34).


[0057] Other racemic type mixtures of compounds contemplated include: e.g. (±)-2-hydroxy-2-(3'-aminophenyl)acetic acid (compound 10); (±)-2-hydroxy-2-(4'-aminophenyl)acetic acid (compound 11); (±)-2-hydroxy-3-(4'-aminophenyl)propionic acid (compound 21) and (±)-2-methoxy-2-(3'-aminophenyl)acetic acid (compound 22).

[0058] Further compounds contemplated for use in the disclosed methods: 5-aminosalicylo-hydroxamic acid (compound 5); 3-dimethylaminosalicylic acid (compound 6); 2-methoxy-4-aminobenzoic acid (compound 7); 2-methoxy-5-aminobenzoic acid (compound 8); 5-methylaminosalicylic acid (compound 9); 4-methylaminosalicylic acid (compound 12); 4-acetylaminosalicylic acid (compound 16); 2-ethoxy-4-aminobenzoic acid (compound 18); 2-ethoxy-5-aminobenzoic acid (compound 19); 4-dimethylaminosalicylic acid (compound 24); 2-ethoxy-4-aminobenzoylhydroxamic acid (compound 25); 6-hydroxyquinoline-5-carboxylic acid

(compound 27); 2-(2-propyl)oxy-4-aminobenzoic acid (compound 30); 4-(1-piperazinyl)salicylic acid (compound 41); (R,S) 5-oxa-quinoline-6-carboxylic acid (compound 15); 6-methoxy quinoline-5-carboxylic acid (compound 36); 6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid (compound 37); 5-diisopropylaminosalicylic acid (compound 38); and 4-diisopropylaminosalicylic acid (compound 42).

[0059] Methods for making contemplated compounds may be found for example in WO2007/010516 and WO2007/010514, each hereby incorporated by reference in their entirety.

[0060] Also provided herein are compounds represented by:



I'

wherein X is C₁-C₃alkylene, optionally substituted with one, two or three substituents selected from halogen or hydroxyl;

R₁ is selected from the group consisting of C₁-C₆alkyl, C₃-C₆cycloalkyl, C₂-C₆alkenyl, and C₂-C₆alkynyl;

R₂ is selected from the group consisting of hydrogen and C₁-C₆alkyl;

R₃ is independently selected, for each occurrence from the group consisting of hydrogen, C₁-C₆alkoxy, C₁-C₆alkyl, cyano, C₃-C₆cycloalkyl, halogen, hydroxyl, and nitro;

R₄ is selected from the group consisting of hydrogen and C₁-C₆alkyl;


R₅ is C₁-C₆alkyl;

or pharmaceutically acceptable salts or N-oxides thereof.

[0061] In one embodiment, R₁ can be C₁-C₆alkyl, such as methyl. In one embodiment, R₂ can be hydrogen. In another embodiment, R₃ can be selected from the group consisting of hydrogen, C₁-C₆alkyl, halogen, and hydroxyl. In a further embodiment, R₃ can be hydrogen. In one embodiment, R₄ and R₅ can each be C₁-C₆alkyl. In another embodiment, R₄ may be


hydrogen and R_5 may be methyl. In one embodiment, X may be $(CH_2)_n$, wherein n is 1 or 2, such as 1.

[0062] In another embodiment, $-NR_2-COR_1$ can be in the meta position relative to X as shown in formula III'.




III'

[0063] In another embodiment, $-NR_2-COR_1$ can be in the para position relative to X as shown in formula IV'.



IV'

[0064] The disclosure provides, at least in part, compounds represented by formula II', as depicted below. Also contemplated herein are compositions that include a compound represented by formula II and e.g., a pharmaceutically acceptable carrier.



II'

wherein R₁ is selected from the group consisting of C₁-C₆alkyl, C₃-C₆cycloalkyl, C₂-C₆alkenyl, and C₂-C₆alkynyl;


R₂ is selected from the group consisting of hydrogen and C₁-C₆alkyl;

R₃ is independently selected, for each occurrence from the group consisting of hydrogen, C₁-C₆alkoxy, C₁-C₆alkyl, cyano, C₃-C₆cycloalkyl, halogen, hydroxyl, and nitro;

R₅ is hydrogen or C₁-C₆alkyl;

or pharmaceutically acceptable salts or N-oxides thereof.

[0065] Compounds of Formula II' are also contemplated as shown below, as well as compositions that include a compound represented by formula II' and e.g., a pharmaceutically acceptable carrier. Contemplated compounds can include:



wherein R₁ is selected from the group consisting of C₁-C₆alkyl, C₃-C₆cycloalkyl, C₂-C₆alkenyl, and C₂-C₆alkynyl;

R₃ is independently selected, for each occurrence from the group consisting of hydrogen, C₁-C₆alkoxy, C₁-C₆alkyl, cyano, C₃-C₆cycloalkyl, halogen, hydroxyl, and nitro;

R₄ is selected from the group consisting of hydrogen and C₁-C₆alkyl;


R₅ is hydrogen or C₁-C₆alkyl; and

A is a fused five or six membered heterocycle;

or pharmaceutically acceptable salts or N-oxides thereof.

[0066] In one embodiment, R₁ can be C₁-C₆alkyl, such as methyl. In another embodiment, R₁ and R₃ can each be C₁-C₆alkyl, such as methyl. In one embodiment, R₂ can be hydrogen.

[0067] In some embodiments, a compound can be represented by




wherein p is 1 or 2;


R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

R_4 and R_8 are each independently selected from the group consisting of hydrogen and C_1 - C_6 alkyl;

or pharmaceutically acceptable salts or N-oxides thereof.

[0068] Contemplated compounds, and pharmaceutical compositions, comprising at least one compound, may be selected from the group consisting of: N-acetyl-(R)-(-)-3-(4-aminophenyl)-2-methoxypropionic acid (Compound A), N-acetyl-(S)-(-)-3-(4-aminophenyl)-2-methoxypropionic acid (Compound B), racemic N-acetyl-(S)-(-)-3-(4-aminophenyl)-2-methoxypropionic acid (compound AB);





4-acetamido-N-hydroxy-2-methoxybenzamide; 1-acetyl-6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid, 5-acetamido-2hydroxybenzoic acid (e.g., acetylated 5-aminosalicylic acid) or pharmaceutically acceptable salts or N-oxides thereof.

[0069] Also provided herein are other PPAR modulator or agonists including 5-amino salicylic acid (mesalamine), sulfasalazine, pioglitazone, rosiglitazone, pomegranate flower extract, non-steroidal anti inflammatories including ibuprofen, and Ψ -baptigenin.

[0070] The present disclosure also provides pharmaceutical compositions comprising compounds as disclosed herein formulated together with one or more pharmaceutically or cosmetically acceptable carriers. These formulations include those suitable for oral, rectal, topical, buccal and parenteral (e.g., subcutaneous, intramuscular, intradermal, or intravenous) administration, or for topical use, e.g. as a cosmetic product. Although the most suitable form of administration in any given case will depend on the degree and severity of the condition being treated and on the nature of the particular compound being used.

Therapeutic Applications

[0071] The disclosure is directed at least in part to treating or ameliorating lactose intolerance using a PPAR modulator, e.g., a PPAR γ modulator, such as a disclosed compound. For example, methods of treating diarrhea, abdominal pain and/or bloating after lactose ingestion is provided, wherein a disclosed compound (or e.g., a composition that includes a disclosed compound) is administered to a subject in need thereof, e.g. orally administered. .

[0072] For example, provided herein is a method for ameliorating lactose intolerance in a patient ingesting a composition that includes lactose, comprising administering a PPAR agonist,

e.g., a PPAR α , PPAR δ , and/or a PPAR γ agonist before, substantially simultaneously with, or after said ingestion.

[0073] Also contemplated herein are compositions used for reducing lactose intolerance, e.g. a disclosed composition may form part of, or is used for making, a low lactose content milk or milk product, comprising a PPAR modulator. Such compositions may be part of a whey, a milk or a cheese.

[0074] The compounds of the invention may be administered to subjects (animals and/or humans) in need of such treatment in dosages that will provide optimal pharmaceutical efficacy. It will be appreciated that the dose required for use in any particular application will vary from patient to patient, not only with the particular compound or composition selected, but also with the route of administration, the nature of the condition being treated, the age and condition of the patient, concurrent medication or special diets then being followed by the patient, and other factors which those skilled in the art will recognize, with the appropriate dosage ultimately being at the discretion of the attendant physician. For treating clinical conditions and diseases noted above, the compound of this invention may be administered orally, topically, parenterally, by inhalation spray or rectally in dosage unit formulations containing conventional non-toxic pharmaceutically acceptable carriers, adjuvants and vehicles. The term parenteral as used herein includes subcutaneous injections, intravenous, intramuscular, intrasternal injection or infusion techniques.

[0075] Generally, a therapeutically effective amount of active component will be in the range of from about 0.1 mg/kg to about 100 mg/kg, optionally from about 1 mg/kg to about 100 mg/kg, optionally from about 1 mg/kg to 10 mg/kg. The amount administered will depend on variables such as the type and extent of disease or indication to be treated, the overall health status of the particular patient, the relative biological efficacy of the compounds, formulation of compounds, the presence and types of excipients in the formulation, and the route of administration. The initial dosage administered may be increased beyond the upper level in order to rapidly achieve the desired blood-level or tissue level, or the initial dosage may be smaller than the optimum and the daily dosage may be progressively increased during the course of treatment depending on the particular situation. Human dosage can be optimized, e.g., in a conventional Phase I dose escalation study designed to run from 0.5 mg/kg to 20 mg/kg. Dosing frequency can vary, depending on factors such as route of administration, dosage amount and the

disease condition being treated. Exemplary dosing frequencies are once per day, once per week and once every two weeks.

[0076] Contemplated formulations or compositions comprise a disclosed compound and typically may also include a pharmaceutically acceptable carrier or excipient.

[0077] Contemplated compositions may be administered by various means, depending on their intended use, as is well known in the art. For example, if compositions of the present invention are to be administered orally, they may be formulated as tablets, capsules, granules, powders or syrups. Alternatively, formulations of the present invention may be administered parenterally as injections (intravenous, intramuscular or subcutaneous), drop infusion preparations or enemas or suppositories. For application by the ophthalmic mucous membrane route, compositions of the present invention may be formulated as eyedrops or eye ointments. These formulations may be prepared by conventional means, and, if desired, the compositions may be mixed with any conventional additive, such as an excipient, a binder, a disintegrating agent, a lubricant, a corrigent, a solubilizing agent, a suspension aid, an emulsifying agent or a coating agent.

[0078] In formulations of the subject invention, wetting agents, emulsifiers and lubricants, such as sodium lauryl sulfate and magnesium stearate, as well as coloring agents, release agents, coating agents, sweetening, flavoring and perfuming agents, preservatives and antioxidants may be present in the formulated agents.

[0079] Subject compositions may be suitable for oral, nasal, topical (including buccal and sublingual), rectal, vaginal, aerosol and/or parenteral administration. The formulations may conveniently be presented in unit dosage form and may be prepared by any methods well known in the art of pharmacy. The amount of composition that may be combined with a carrier material to produce a single dose vary depending upon the subject being treated, and the particular mode of administration.

[0080] Formulations suitable for oral administration may be in the form of capsules, cachets, pills, tablets, lozenges (using a flavored basis, usually sucrose and acacia or tragacanth), powders, granules, or as a solution or a suspension in an aqueous or non-aqueous liquid, or as an oil-in-water or water-in-oil liquid emulsion, or as an elixir or syrup, or as pastilles (using an inert base, such as gelatin and glycerin, or sucrose and acacia), each containing a predetermined

amount of a subject composition thereof as an active ingredient. Compositions of the present invention may also be administered as a bolus, electuary, or paste.

[0081] In solid dosage forms for oral administration (capsules, tablets, pills, film-coated tablets, sugar-coated tablets, powders, granules and the like), the subject composition is mixed with one or more pharmaceutically acceptable carriers, such as sodium citrate or dicalcium phosphate, and/or any of the following: (1) fillers or extenders, such as starches, lactose, sucrose, glucose, mannitol, and/or silicic acid; (2) binders, such as, for example, carboxymethylcellulose, alginates, gelatin, polyvinyl pyrrolidone, sucrose and/or acacia; (3) humectants, such as glycerol; (4) disintegrating agents, such as agar-agar, calcium carbonate, potato or tapioca starch, alginic acid, certain silicates, and sodium carbonate; (5) solution retarding agents, such as paraffin; (6) absorption accelerators, such as quaternary ammonium compounds; (7) wetting agents, such as, for example, acetyl alcohol and glycerol monostearate; (8) absorbents, such as kaolin and bentonite clay; (9) lubricants, such a talc, calcium stearate, magnesium stearate, solid polyethylene glycols, sodium lauryl sulfate, and mixtures thereof; and (10) coloring agents. In the case of capsules, tablets and pills, the compositions may also comprise buffering agents. Solid compositions of a similar type may also be employed as fillers in soft and hard-filled gelatin capsules using such excipients as lactose or milk sugars, as well as high molecular weight polyethylene glycols and the like..

[0082] Liquid dosage forms for oral administration include pharmaceutically acceptable emulsions, microemulsions, solutions, suspensions, syrups and elixirs. In addition to the subject composition, the liquid dosage forms may contain inert diluents commonly used in the art, such as, for example, water or other solvents, solubilizing agents and emulsifiers, such as ethyl alcohol, isopropyl alcohol, ethyl carbonate, ethyl acetate, benzyl alcohol, benzyl benzoate, propylene glycol, 1,3-butylene glycol, oils (in particular, cottonseed, groundnut, corn, germ, olive, castor and sesame oils), glycerol, tetrahydrofuryl alcohol, polyethylene glycols and fatty acid esters of sorbitan, cyclodextrins and mixtures thereof.

[0083] Suspensions, in addition to the subject composition, may contain suspending agents as, for example, ethoxylated isostearyl alcohols, polyoxyethylene sorbitol and sorbitan esters, microcrystalline cellulose, aluminum metahydroxide, bentonite, agar-agar and tragacanth, and mixtures thereof.

[0084] Throughout the description, where compositions are described as having, including, or comprising specific components, it is contemplated that compositions also consist essentially of, or consist of, the recited components. Similarly, where processes are described as having, including, or comprising specific process steps, the processes also consist essentially of, or consist of, the recited processing steps. Except where indicated otherwise, the order of steps or order for performing certain actions are immaterial so long as the invention remains operable. Moreover, unless otherwise noted, two or more steps or actions may be conducted simultaneously.

EXAMPLES

[0085] The compounds disclosed herein can be prepared in a number of ways well known to one skilled in the art of organic synthesis. .

Example A Materials and Methods

Cell culture

[0086] Caco-2 colonic adenocarcinoma cell lines were used. Caco-2 cells grew in Dulbecco's modified Eagle's medium (DMEM, Invitrogen, Cergy-Pontoise, France) supplemented with 20% foetal calf serum (FCS, Dutscher, Brumath, France), 1% penicillin-streptomycin (5 ml/l) (Invitrogen) and 1% non-essential amino acids (5 ml/l) (Invitrogen). All cell lines were cultured as confluent monolayers at 37°C in a controlled, 5% CO₂ atmosphere. Each week, the confluent monolayer were trypsinized (Invitrogen) and then replated after a 1:10 dilution. For the stimulation experiments, cells were placed in 6-well plates at a density of 1x10⁶ cells per well. Serum deprivation was used 16 hours prior to stimulation in order to synchronize the cells.

Treatment of the cells

[0087] The cells were treated with Compound 34 (1 mM and 30mM), 5-ASA (30 mM), or pioglitazone (1 µM). When necessary, the vehicle DMSO (Sigma) is used as control (Ctrl). After 24 hrs of stimulation, the wells were rinsed three times with sterile PBS at 4°C. The cells were then lysed with lysis buffer (RA1, Macherey-Nagel) containing 1% β-mercaptoethanol. The plates were frozen at -80°C for subsequent RNA extraction. A time course experiment

consisting of stimulation of Caco-2 cells by compound 34, 1mM during 6hrs and 12 hrs was also done in order to compare gene expression according treatment duration.

RNA extraction

[0088] Total RNA were extracted with a Nucleospin RNA kit (Macherey-Nagel, Hoerd, France). After RNase inactivation, the total RNA was cleaned of trace genomic DNA via a DNase treatment and eluted in RNase-free, DEPC-free water. The purity of the RNA will be evaluated by UV spectroscopy on a Nanodrop system from 220 to 350 nm. Before microarray experiment, RNAs were also profiled on an Agilent 2100 bioanalyzer. One µg of total RNA were used in the microarray analysis (minimum concentration: 50 ng/µl).

Microarrays

[0089] Dual-colour gene expression microarrays were used compare the cRNA from the samples (cells treated with DMSO (Ctrl) vs. cells treated with an agonist (Ago)). Briefly, the RNA from the samples is first reverse-transcribed into cDNA (Affinity-Script RT, Agilent), which is then used as the substrate for the synthesis and amplification of cRNA by T7 RNA polymerase in the presence of cyanine 3-CTP for the Ctrl sample (green fluorescence) and cyanine 5-CTP for the Ago sample (red fluorescence). The two labelled cRNAs were then mixed and hybridized on the same array (G4851A Agilent 8x44K), which is then scanned (with an Agilent G2505B scanner). The fluorescence was visualised after laser excitation and the relative intensities of the two fluorophores will be expressed as a ratio, in order to yield the over- or under-expression status of each gene (using GeneSpring software (Agilent)). This analysis was performed for each agonist. The transcriptomic profiles obtained for each agonist were then compared. Expression of genes of interest was quantified by quantitative PCR, in order to confirm the microarray results.

Quantitative PCR

[0090] Transcripts of genes were estimated by quantitative PCR (qPCR) in order to confirm microarrays results. 1µg of total RNA was reverse-transcribed into cDNA using the High Capacity cDNA Archive kit (Applied biosystems). qPCR was performed using an ABI PRISM 7000 sequence detection system (Applied Biosystem). Primer pairs for each transcript were chosen with qPrimer depot software. Quantification of qPCR signals was performed using ΔC_t relative quantification method using GAPDH as a reference gene. The values were

represented in terms of relative quantity of mRNA level variation or fold increase compared to control conditions.

Generation of PPAR γ knock down cells

[0091] PPAR γ knock down IECs were obtained using the pSUPER.retro system (OligoEngine). Forward and reverse target sequences corresponding to nucleotides 105-123 of the human PPAR γ mRNA (5'-GCCCTTCACTACTGTTGAC-3') were cloned into the BglII/XhoI restriction sites of the pSUPERretro vector (pRS) giving the ShPPAR construct. A negative control pRS plasmid containing the sequence 5'-ACGCTGAGTACTTCGAAAT-3' targeted against the luciferase gene was also generated (ShLuc construct). Both constructions were transfected in Caco-2 cells using Nucleofector technology from Amaxa Biosystems, according to the manufacturer's protocol. Stably transfected clones were selected 24h post-transfection with complete culture medium supplemented with puromycin (5 μ g/ml). The silencing of PPAR γ expression was checked by quantitative RT-PCR and western-blot analysis. Once established, ShPPAR and ShLuc cell lines were maintained in complete medium supplemented with 2.5 μ g/ml puromycin.

Example 1 Microarray

[0092] In order to evaluate the feasibility of transcriptomic analysis of intestinal epithelial cells stimulated with the PPAR agonist compound 34, the gene expression profile of Caco-2 cells only stimulated with compound 34 at the concentration of 1mM was compared. Cell stimulations were done with 4 replicates and a total of 44 000 genes were screened. 52 genes showed a significant different expression in the compound 34, 1mM condition in comparison with controls ($p < 0.05$) (Table 1). 49 genes were up-regulated in the compound 34, 1mM condition in comparison with controls (one gene (LCT 1) with a fold change superior of 5, four genes with a fold change between 3 and 5, and 13 genes with a fold change between 2 and 3 (Table1). Three genes were down-regulated in the compound 34, 1mM condition in comparison with controls (MYB, OXA13 and ANKRD1).

[0093]

Gene Symbol	Description	Fold change	Regulation
LCT	Homo sapiens lactase	5,2924175	up
DMBT1	Homo sapiens deleted in malignant brain tumors 1, transcript variant 2	3,8939168	up
KRT20	Homo sapiens keratin 20	3,1442842	up
AADAC	Homo sapiens arylacetamide deacetylase (esterase)	3,0116935	up
MBL2	Homo sapiens mannose-binding lectin 2	2,8058631	up

GPA33	Homo sapiens glycoprotein A33	2,662811	up
SLC38A4	Homo sapiens solute carrier family 38, member 4, transcript variant 1	2,6554945	up
C11orf86	Homo sapiens chromosome 11 open reading frame 86	2,583426	up
SGK2	Homo sapiens serum/glucocorticoid regulated kinase 2, transcript variant 1	2,5728035	up
MMP19	Homo sapiens matrix metalloproteinase 19, transcript variant 1	2,481643	up
OSTalpha	Homo sapiens organic solute transporter alpha	2,4348273	up
LZTS1	Homo sapiens leucine zipper, putative tumor suppressor 1	2,3766112	up
HMGCS2	Homo sapiens 3-hydroxy-3-methylglutaryl-Coenzyme A synthase 2, transcript variant 1	2,231923	up
PLIN2	Homo sapiens perilipin 2	2,2110133	up
CREB3L3	Homo sapiens cAMP responsive element binding protein 3-like 3	2,1615057	up
SLC23A3	Homo sapiens solute carrier family 23 (nucleobase transporters), member 3 transcript variant 1	1,9120694	up
ACSL5	Homo sapiens acyl-CoA synthetase long-chain family member 5	1,8756381	up
HIGD1A	Homo sapiens HIG1 hypoxia inducible domain family, member 1A	1,8233925	up
MUC17	Homo sapiens mucin 17, cell surface associated	1,79823	up
AGPAT2	Homo sapiens 1-acylglycerol-3-phosphate O-acyltransferase 2	1,7209687	up
FABP1	Homo sapiens fatty acid binding protein 1, liver	1,7156018	up
AQP12A	Homo sapiens aquaporin 12A	1,7124082	up
DNAH12	Homo sapiens dynein, axonemal, heavy chain 12	1,7119135	up
COL16A1	Homo sapiens collagen, type XVI, alpha 1	1,7069026	up
SLC25A42	Homo sapiens solute carrier family 25, member 42	1,697567	up
AQP12B	Homo-sapiens aquaporin-12B	1,6971904	up
ITI13	Homo sapiens inter-alpha (globulin) inhibitor H3	1,6908818	up
EMP1	Homo sapiens epithelial membrane protein 1	1,6551878	up
CAPS	Homo sapiens calcyphosine, transcript variant 1	1,645898	up
ACSM3	Homo sapiens acyl-CoA synthetase medium-chain family member 3 transcript variant 2	1,6409295	up
SLC46A1	Homo sapiens solute carrier family 46, member 1	1,6296347	up
ABCG2	Homo sapiens ATP-binding cassette, sub-family G, member 2	1,6244398	up
MYB	Homo sapiens v-myb myeloblastosis viral oncogene homolog (avian) transcript variant 2	1,6136038	down
COL12A1	Homo sapiens collagen, type XII, alpha 1, transcript variant long	1,5990913	up
CYP2B6	Homo sapiens cytochrome P450, family 2, subfamily B, polypeptide 6	1,5909243	up
MMP19	Homo sapiens matrix metalloproteinase 19 transcript variant 1	1,5854398	up
CREB3L3	Homo sapiens cAMP responsive element binding protein 3-like 3	1,578399	up
ANKRD1	Homo sapiens ankyrin repeat domain 1	1,5722646	down
ZNF575	Homo sapiens zinc finger protein 575	1,5662627	up
SLC25A20	Homo sapiens solute carrier family 25 (carnitine/acylcarnitine translocase), member 20	1,563665	up
GALNT2	Homo sapiens UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 2	1,5570259	up
RHOU	Homo sapiens ras homolog gene family, member U	1,5535966	up
SLC46A1	Homo sapiens solute carrier family 46, member 1	1,5489862	up
PPARGC1B	Peroxisome proliferator-activated receptor gamma coactivator 1-beta	1,5455792	up
CHST3	Homo sapiens carbohydrate (chondroitin 6) sulfotransferase 3	1,5176716	up
HOXA13	Homo sapiens homeobox A13	1,5115318	down
HIGD1A	Homo sapiens HIG1 hypoxia inducible domain family, member 1A, transcript variant 3	1,5103043	up
ACSL4	Homo sapiens acyl-CoA synthetase long-chain family member 4, transcript variant 1	1,5066991	up
LUM	Homo sapiens lumican	1,5007194	up

[0094] **Table 1.** Comparison of transcriptomic profiles in epithelial cells treated by compound 34 1mM versus control. Results are expressed in fold.

Confirmation by qPCR

[0095] We used quantitative PCR to confirm microarrays results. We quantified by quantitative PCR the expression of the 15 most up regulated genes evaluated by microarrays (SGK2 alpha, SGK2 beta, GPA33, CKT 20, LCT1, AADAC, DMBT1, ACSL5, MMP19, SLC28A4, AQP12, MBL2, OSTalpha, HMGCoAS2, LZTS1) in Cacao2 cells cultured in the same condition. Data obtained by quantitative PCR and presented in the Figures 1a-b confirmed array results for the 12/15 up regulated genes except for AQP12, OST α and LZTS1.

[0096] Genes variations were evaluated with a second and independent experiment in stimulated Caco-2 cells. The same stimulation protocol was used (compound 34 1mM vs control) and gene expressions were studied using qRT-PCR. We confirmed the up-regulation of all the selected genes induced by compound 34 (including AQP12, OSTa and LZTS1 which are weakly expressed but upregulated by compound 34) except for GPA33 (Figure 2A and 2B).

[0097] Interestingly, in this experiment, PPAR γ mRNA expression was not modified following compound 34 treatment supporting that compound 34 mediated upregulated target genes was mainly due to the modulation of PPAR γ activity by compound 34, rather than an increased of the receptor expression.

Correlation between gene expression expressed in fold in microarrays and qPCR

[0098] To evaluate more definitively the reproducibility of our microarray data, we evaluated the correlation between quantitative PCR evaluation and microarray analysis. We confirmed a significant strong correlation between microarrays (stimulation 1) and qPCR (stimulation 2) results expressed in Fold Change ($r=0.754$, $p=0.0046$) (Figure 3).

Example 2 Lactase gene as a (PPAR) target gene

[0099] Among the new 15 new target genes of compound 34 expressed by epithelial cells, the gene encoding for lactase was one with the strongest induction (fold change 5.29). This induction was confirmed in the second stimulation suggesting that the lactase gene is one of the most PPAR γ - sensitive genes in our analysis (figure 2a). Due to the functional consequences of

lactase gene abnormalities in lactose deficiency, a common disorder found in 20% of European population, we decided to deeper characterize this result. Inadequate lactase-phlorizin hydrolase (LPH) activity is responsible of lactose intolerance/malabsorption leading to diarrhea, abdominal pain or bloating after lactose ingestion.

[00100] Figure 4 shows the results of lactase transcript expression in different experiments of Caco-2 cells stimulated with various agonists of PPAR. Each experiment has been done at least twice. The lactase gene expression was first confirmed to be stimulated by compound 34 at two different concentrations (1mM, fold increase 4.78, $p=0.0002$; 30mM, fold increase 7.9, $p=0.0006$) (Fig 4A). Time course experiment showed that 6 hours of stimulation by compound 34 1mM is enough to significantly increase Lactase mRNA expression ($p<0.05$) and that the maximum expression is reached at 12 hours of treatment ($p<0.0001$) (Figure 4A). Similarly to compound 34, 5ASA 30mM and pioglitazone 1 μ M, significantly increased similarly lactase mRNA expression (figure 4B)(respectively 12.69 fold increase, $p=0.0006$, 11.76 fold increase, $p=0.0006$). Fenofibrate, a PPAR α agonist, induced weakly and without significance the lactase gene expression in Caco-2 cells (Figure 4C). In Caco2 ShPPAR cell line, a cell line stably expressing a short hairpin anti-sense RNA against PPAR leading to a specific 70% down expression of PPAR γ expression, and lactase mRNA expression is significantly reduced by 63% ($p=0.0012$) compared to ShLuc control cells (Figure 4D). Altogether, these data demonstrated that the human lactase gene is inducible by PPAR agonists in Caco-2 cells and that Lactase gene could be a new PPAR target gene in intestinal epithelial cells.

Presence of PPRE sequences at 3000 bp upstream the LCT gene

[00101] PPAR γ is able to bind DNA as a heterodimer with another nuclear receptor RXR. The heterodimer PPAR γ -RXR recognizes short dimeric palindromic sequences (consensus AGGTCA or TGACCT) spaced by one or two nucleotides giving the DR1 (direct repeat 1) or DR2 (direct repeat 2) thereby defining the PPAR γ response element (PPRE). To better evaluate the potential regulation of the LCT gene by PPAR, we investigated the potential presence of PPRE sequences upstream the LCT gene. Bioinformatics tools were used to find PPRE in the human Lactase gene. Three different programs were used: NUBIScan, PPRE Search and MatInspector Figure 5 represents the result of the *in silico* analysis of the 3000 bases pair upstream of the transcription start point of the human Lactase gene. Only direct repeats that were predicted by at least two different programs are indicated. This analysis revealed that the 3000 bp

upstream of the transcription start site contain several PPRE which could be used by PPAR to regulate Lactase expression.

[00102] Transcriptomic analysis of intestinal epithelial cells has been found to be stimulated by compound 34 1mM using microarray showing a significant modification of 52 genes. These results were confirmed using quantitative RT-PCR in different and independent experiments. Among these 52 genes, we focused of interest on the gene encoding for lactase and confirmed upregulated LCT gene expression by 5.29. 5-ASA and pioglitazone are also able to enhance LCT gene expression in Caco2 cells, the spontaneous level of LCT gene expression is significantly reduced by 70% in PPAR γ deficient epithelial cells compared to control epithelial cells, 5ASA and pioglitazone kept their upregulatory effects on LCT gene expression in PPAR semi deficient epithelial cells, suggesting that impaired expression of LCT gene may be restored by PPAR agonists. Altogether, these data demonstrate that PPAR may be a new modulator of LCT gene expression and suggest that PPAR agonist and particularly a new class of PPAR γ agonist targeting the gut may be a new strategy to treat patients with lactase deficiency.

Example 3 Reporter gene assay and chromatin immunoprecipitation experiments

[00103] The 3000bp of the lactase gene promoter in the pGL4-Luc reporter vector was cloned. This construction (and the control empty vector) is transiently transfected in Caco-2 cells. The cells are then treated with various PPAR agonists and the luciferase activity is measured in order to evaluate the regulation of lactase promoter activity by PPAR modulators. The identification and localization of the PPAR response element(s) is achieved by successive deletions of the promoter region cloned in the pGL4-Luc reporter vector. The physical binding of PPAR γ on the promoter of the Lactase gene is studied by ChIP experiments in Caco-2 cells stimulated or not with PPAR agonists.

Example 4- PPAR γ agonists ability to modulate (to increase) lactase activity

[00104] Both stimulation of Caco-2 cells and *ex-vivo* stimulation of small pieces of mice intestine are used to test whether PPAR γ agonists are able to increase lactase activity. The lactase activity (disaccharidase activity) is assessed with o-nitrophenyl- β -D galactoside (ONPG) as a substrate. Upon hydrolysis of the β -galactosidic bond, ONPG yields galactose and o-nitrophenol, a yellow compound which can be detected by spectrophotometric methods (absorption max =

450 nm). Alternatively, the expression level of lactase gene in the intestine of mice is evaluated which will receive compound 34 or 5-ASA *per os*.

References

[00105] All publications and patents mentioned herein, including those items listed below, are hereby incorporated by reference in their entirety as if each individual publication or patent was specifically and individually incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.


Equivalents

[00106] While specific embodiments of the subject invention have been discussed, the above specification is illustrative and not restrictive. Many variations of the invention will become apparent to those skilled in the art upon review of this specification. The full scope of the invention should be determined by reference to the claims, along with their full scope of equivalents, and the specification, along with such variations.

[00107] Unless otherwise indicated, all numbers expressing quantities of ingredients, reaction conditions, and so forth used in the specification and claims are to be understood as being modified in all instances by the term “about.” Accordingly, unless indicated to the contrary, the numerical parameters set forth in this specification and attached claims are approximations that may vary depending upon the desired properties sought to be obtained by the present invention.

WHAT IS CLAIMED IS:

1. A method for treating and/or ameliorating lactose intolerance or lactase deficiency in a patient in need thereof, comprising administering a PPAR agonist compound to said patient.
2. A method for stimulating lactase gene expression in a patient in need thereof, comprising administering a PPAR γ agonist to said patient.
3. A method for treating diarrhea, abdominal pain and/or bloating after lactose ingestion in a patient in need thereof, comprising administering a PPAR agonist.
4. A method for ameliorating lactose intolerance in a patient ingesting a composition that includes lactose, comprising administering a PPAR agonist before, substantially simultaneously with, or after said ingestion.
5. The method of any one of claims 1-4, wherein the PPAR agonist is represented by Formula I:



and pharmaceutically acceptable salts, stereoisomers, and/or prodrugs thereof, wherein:

R_1 and R_2 , are each independently selected from the group consisting of H and C_{1-6} alkyl; or R_1 and R_2 together with the nitrogen atom they are bonded to form an aromatic or aliphatic ring with 5 or 6 atoms which may be optionally substituted;

Y and Z are each independently selected from the group consisting of H, OH, COOH, $-OR_3$, $-CH(OR_3)COOH$; and


R_3 is selected from the group consisting of H, phenyl, benzyl, vinyl, allyl, C_{1-6} alkyl or C_{1-6} alkyl substituted by one, two, three or more halogens.

6. The method of any one of claims 1-4, wherein the PPAR agonist is selected from the group consisting of: ($\pm$)-2-hydroxy-3-(3'-aminophenyl) propionic acid; ($\pm$)-2-methoxy-2-(4'-aminophenyl) acetic acid; ($\pm$)-2-ethoxy-2-(3'-aminophenyl) acetic acid; ($\pm$)-2-ethoxy-2-(4'-aminophenyl) acetic acid; ($\pm$)-2-methoxy-3-(4'-aminophenyl) propionic acid; ($\pm$)-2-ethoxy-3-(4'-aminophenyl) propionic acid; ($\pm$)-2-ethoxy-3-(3'-aminophenyl) propionic acid, (R,S)-2-methoxy-3-(4-aminophenyl)propionic acid, 5-aminosalicylo-hydroxamic acid (compound 5); 3-

dimethylaminosalicylic acid; 2-methoxy-4-aminobenzoic acid; 2-methoxy-5-amino-5-methylaminosalicylic acid; 4-methylaminosalicylic acid; 4-acetylaminosalicylic acid; 2-ethoxy-4-aminobenzoic acid; 2-ethoxy-5-aminobenzoic acid; 4-dimethylaminosalicylic acid; 2-ethoxy-4-aminobenzoylhydroxamic acid; 6-hydroxyquinoline-5-carboxylic acid; 2-(2-propyl)oxy-4-aminobenzoic acid; 4-(1-piperazinyl)salicylic acid; (R,S) 5-oxa-quinoline-6-carboxylic acid; 6-methoxy quinoline-5-carboxylic acid; 6-methoxy-1,2,3,4-tetrahydroquinoline-5-carboxylic acid; 5-diisopropylaminosalicylic acid; and 4-diisopropylaminosalicylic acid or pharmaceutically acceptable salts thereof.

7. The method of any one of claims 1-6, wherein the PPAR agonist is (-) 2-R-methoxy-3-(4-aminophenyl)propionic acid.

8. The method of any one of claims 1-4, wherein the PPAR agonist is a compound represented by Formula I':



wherein X is C_1 - C_3 alkylene, optionally substituted with one, two or three substituents selected from halogen or hydroxyl;

R_1 is selected from the group consisting of C_1 - C_6 alkyl, C_3 - C_6 cycloalkyl, C_2 - C_6 alkenyl, and C_2 - C_6 alkynyl;

R_2 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl;

R_3 is independently selected, for each occurrence from the group consisting of hydrogen, C_1 - C_6 alkoxy, C_1 - C_6 alkyl, cyano, C_3 - C_6 cycloalkyl, halogen, hydroxyl, and nitro;

R_4 is selected from the group consisting of hydrogen and C_1 - C_6 alkyl;


R₅ is hydrogen C₁-C₆alkyl; or pharmaceutically acceptable salts or N-oxides thereof; and an excipient.

9. The method of claim 8, wherein the compound is N-acetyl(R)-(-)-3-4-(4-aminophenyl)-2-methoxypropionic acid.
10. The method of any one of claims 1-4, wherein the PPAR agonist is selected from the group consisting of mesalamine, sulfasalazine, pioglitazone and rosiglitazone.
11. A food product comprising a dairy component and an PPAR agonist.
12. The food product of claim 11, wherein the dairy component is whey, milk, cheese or cream.
13. Use of a PPAR agonist compound in the treatment of lactose intolerance or lactase deficiency.
14. Use of a PPAR γ agonist in a method of stimulating lactase gene expression.
15. Use of a PPAR agonist in a method of treatment of diarrhoea, abdominal pain and/or bloating after lactose ingestion.

MCG24101

FIGURE 1

A



B





Figure 2

A



B




FIGURE 3




FIGURE 4




FIGURE 5

GTAATTTATTTTACTTCTGTGTCCTAAGGGTAATTTCTCAGGATTGTTTTCAAATTGCTTTTTAGGGGAAATAGGTCAT
TTGCTATATTACAAGCAATCCCCAAATTTTATGGTCTCCAGGAAAAAGTTATTACCGTTTATGATACTAACAGTTCCTGA
GACTTAGCTATGATCAGTATGTTTCATGAGGTGGAGCAGTTCCTGTGTTGCAGCTTTAACAACAGATGGCATTCAATAAA
TCACAAAGTATGTTAAAGGTCACAAAAGCAAAATAACTGTCTGAGGCTAAGGCCACGTGGGACAGTCTAATACCCATGA
GTA CTCAACTTGCCCTTGATGTCTGAGCTTTCCAGTGCAATGTGAATTTGAGCAGCCAGAAATCTATTAGTAGAAAGCAAG
ACAGATTAATATAGGTTAAACAATGATTTAAATATGTTTCTCCCAATAATTATCTCTTTCCCTGGAATCAACTTGTATG
AAACCTTGTCAAAATGTACTCCACAAGTATGTACAATTAAAGTATTTAAAAATAAATGGCAAACATTAAAAACAAGAGTG
AATACTCAAGTAGATTTGTCATGGGATTTTATAGAAGACTGGTATCAGGTAATGTATCTTTAAAGACTAGGCTGCTCT
GCTGACGTAGTAACCATTTTTTATTCCCTTACTTTCTAATAGCCTTGCTTTCACCTAAGAAAAAAAAGGCTAGGCACG
GTGGCTCACGCCTGTAATCCCAGCACTTTGGGAGGCCAAGGTGAGTAGATCACCTGAGGTCAGGAGTTCGAGACAAGCCT
GGCCAACTGGAGAAAACTGTCTCTACTAAAAGTACAAAAATTAGCCAGGCATGGTGGTGGGCACCTGTAATCCCAGCT
ACTCAGGAGGTTGAGGCAGGAGAATTGCTTGAACCAAGAGGTGGAAGTTCAGTGAGCTGAGATCATGCCACTGCCCTC
CAGCTTGGGCGGCAACAACAACAACAAAAAAGATTCCAGGCTTCACCCTCTACAGTCTATGGAAAGTCATGGGAGTCT
ATTA AAAAAA CAGACACTATAAACCAAAATTAAGATGGGAAAATTTTCTATCACGTTTTAATATGTGATATCCAAA
CTCCCATTAAGAATTTTATATCAGTAATAAGATAAACTCAGAAGAAAAATGCCAATTAAACAAGAGAGCTTCACCTC
TTAAATAATCAGAAAAAGAAAAGCACATAATATTTACGTCAAAGTATCAAAAATGTAAAAGTTTTATAATAGGCATTAGC
AAGGCAGACGGATACAGGCACACATCTAACTCTATGTAGGATTTAACTTGGGAAGACAATTTGCATTTCATCAATCAATCAAAA
ACCATTTTTTTTTTGAGATGATCTCCCTGTGTTGCCAGACTGGCCTCAAGCTCCTGGGCTTGAGCAGTCTACAGTCT
GAGCCTCCAAAAGTACTGGCAAAACAGGCGTGATTGACCATGCCAGCCTGTATATAAGGATATTCATTTCGAGCATTGT
TTGCAACACAAAAATGTAAACCAACCGGTGTTAATATGCAAGATGTTAATTTATGGTACATTCTTGCTGTGGAAAGCT
ATGCTATCTGTAGAGGTGGGTCTATATGTTCTGATACAGGCAGAACTCTAAGAGCTATTAAGTGAAGAAAAGGCTGAAAA
TACATATGGCATAATTCCATATATGTTAAATTGTTCTATATTTTAAATGTTATACACGTTTCCATTTGATTTACGTAAT
AATAATGATCCCTGAGCTGTACTGTTGTATACATGCTACAAGAAAGACCCATTTAATCCCCACAGCCTATGATGAATTAT
CCACATTCTACAGGTGACAAAATAGAGGCACAAAGTTAAGTAATTTTGTACAGGTGAGATTTAAACCCAGGCATTCTGAC
TCCTGTATAACCATTAAGATATGCAGAGAAAGAAAAGTGGAAAGATACATATTGCTGAAGATACTTATTATAGGAAGAGG
AGGGGGGAGGGTGAAGGAATTTGCAAGTTTTTCATAGATGTTTCCATATTGTTGAATCTCTTACAAAAATATGTTTCAGCA
TATTTTTAAAGAGAAAAATTTGGGGCAAAATACTTATTTTTGTATTATGTAACAAATTTTAAAAATAATGTGTGGCTGGGT
GCGCTGGCTCACACCTGTAATCCCAACACTTTAGGAGGCTGAGGCAAGAGGATTGCTTGAGCCCAGGAGTTCAAGACCAG
CCTGGGTGACATGGCAAACTCCATCTCTACTAAAAATACAAAAAATTAGCCAGTCGTGGTGGCGCACACCTATGGTCCC
ACCTACCCAGGATGCTGAGATGGGAGGATCACTTGAGCCCAGGAAGTCAAGGCTGCAGGAAGCTGTGATCGCACCACTGC
ACTCCCACCTGGGCAACAGAGTGAGACCCGGTCACCAAAAAACAAAAAAATTTGTAATCGTTTTCTTCAG
ACATTTTCCGGGTTCTCTGCTTAACTTGATAGGAAGTCTGAGGTTTTGTGTTGGTCTTTACCTTTTTTTTTTTTTT
TTTTTTAAGATGGAGTCTCATTCTGTTGCCAGGCTGGAGTGCAGTGGCATGATCTTGGCTCCTGCAACCTCCGCCTCC
TGGGTTCAAGTGATTCTCTGCCCTCAGCCTCCTGAGTAGCCGGGACTACAGGCGCATGCCACGATGCCTGGCTAATTTTT
TGTATTTTTAGTAGAGATGGGGTTTACCATGTTAGCTAGGACGGTCTCGATCTCCTGACCTCGTGATCCGCCACCTCG
GCCTCCCAAAGTGTGGAATTACAGGTGTGAGCCACCACGCCCGCCCTGATCTTTACATTTTTAAATATTGCATTAGTG
AACCGTGTACTGATTTTGTGATCATAGATAACCCAGTTAAATATTAAGTCTTAATTATCACTTAGTATTTACAACCTCA
GTTGCAATTATAAAGTAAGGGTCCACATACCTCCTAACAGttcctagaaa

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/057729

A. CLASSIFICATION OF SUBJECT MATTER
INV. A61K31/195 A61K31/44 A61K31/4402 A61P3/00
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data, EMBASE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2007/096148 A1 (LODERS CROKLAAN BV [NL]; BASSAGANYA-RIERA JOSEP [US]; O'SHEA MARIANNE) 30 August 2007 (2007-08-30) abstract page 14, line 5 - line 23 -----	11,12
A	WO 2008/104557 A1 (GIULIANI INT LTD [IE]; BARONI SERGIO [IT]; DESREUMAUX PIERRE [FR]; BEL) 4 September 2008 (2008-09-04) page 1 page 8 - page 16 -----	1-15
A	WO 2007/010516 A2 (GIULIANI INT LTD [IE]; NACCARI GIANCARLO [IT]; BARONI SERGIO [IT]) 25 January 2007 (2007-01-25) cited in the application abstract claims ----- -/-	1-15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

24 May 2013

Date of mailing of the international search report

11/06/2013

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Damiani, Federica

INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2013/057729

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 2007/010514 A2 (GIULIANI INT LTD [IE]; NACCARI GIANCARLO [IT]; BARONI SERGIO [IT]) 25 January 2007 (2007-01-25) cited in the application abstract claims -----	1-15
A	HARRINGTON L K ET AL: "A re-appraisal of lactose intolerance.", INTERNATIONAL JOURNAL OF CLINICAL PRACTICE OCT 2008, vol. 62, no. 10, October 2008 (2008-10), pages 1541-1546, XP002697625, ISSN: 1742-1241 the whole document -----	1-15

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2013/057729

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2007096148	A1	30-08-2007	NONE
WO 2008104557	A1	04-09-2008	AR 065525 A1 10-06-2009 AT 553759 T 15-05-2012 AU 2008220762 A1 04-09-2008 CA 2678159 A1 04-09-2008 DK 2131829 T3 29-05-2012 EP 2131829 A1 16-12-2009 ES 2384195 T3 02-07-2012 HR P20120473 T1 31-07-2012 IE 20070129 A1 24-12-2008 JP 2010520166 A 10-06-2010 PT 2131829 E 20-06-2012 RS 52251 B 31-10-2012 SI 2131829 T1 28-09-2012 US 2011152225 A1 23-06-2011 WO 2008104557 A1 04-09-2008
WO 2007010516	A2	25-01-2007	AU 2006271165 A1 25-01-2007 BR PI0613396 A2 11-01-2011 CA 2616146 A1 25-01-2007 CN 101243039 A 13-08-2008 CN 102020577 A 20-04-2011 EA 200800372 A1 30-06-2008 EA 201000708 A1 29-10-2010 EP 1910270 A2 16-04-2008 JP 2009502777 A 29-01-2009 KR 20080039428 A 07-05-2008 NZ 566045 A 27-04-2012 NZ 592228 A 26-10-2012 US 2009048343 A1 19-02-2009 US 2013005813 A1 03-01-2013 WO 2007010516 A2 25-01-2007
WO 2007010514	A2	25-01-2007	AU 2006271163 A1 25-01-2007 BR PI0613398 A2 11-01-2011 CA 2615978 A1 25-01-2007 CN 101243043 A 13-08-2008 EP 1910275 A2 16-04-2008 JP 2009502775 A 29-01-2009 NZ 566044 A 29-06-2012 US 2009118357 A1 07-05-2009 US 2012316230 A1 13-12-2012 WO 2007010514 A2 25-01-2007