



(51) International Patent Classification:

A61K 48/00 (2006.01)

(21) International Application Number:

PCT/US2019/064032

(22) International Filing Date:

02 December 2019 (02.12.2019)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:


62/773,885 30 November 2018 (30.11.2018) US
62/826,462 29 March 2019 (29.03.2019) US

(71) Applicant: **DANA-FARBER CANCER INSTITUTE, INC.** [US/US]; 450 Brookline Avenue, Boston, Massachusetts 02215 (US).(72) Inventors: **MARASCO, Wayne A.**; 43 Rice Street, Wellesley, Massachusetts 02481 (US). **ZHU, Quan Karen**; 9 Carriage Hill Circle, Southborough, Massachusetts 01772 (US). **WANG, Yufei**; 20 Chapel Street, Brookline, Massachusetts 02446 (US). **CHANG, Matthew**; 35 Vernon St, Brookline, Massachusetts 02135 (US).(74) Agent: **ESTRADA DE MARTIN, Paula**; 201 St. Charles Avenue, Suite 3600, New Orleans, Louisiana 70170 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: CHIMERIC ANTIGEN RECEPTOR FACTORIES AND METHODS OF USE THEREOF

**FIG. 28**

(57) **Abstract:** This invention is directed to chimeric antigen receptors and cells comprising the same, wherein the cells further secrete monoclonal antibodies locally at a tumor site.



Published:

- *without international search report and to be republished
upon receipt of that report (Rule 48.2(g))*

CHIMERIC ANTIGEN RECEPTOR FACTORIES
AND METHODS OF USE THEREOF

[0001] This application claims priority from U.S. Provisional Application No. 62/773,885, filed on November 30, 2018, and U.S. Provisional Application No. 62/826,462, filed on March 29, 2019, the entire contents of each of which are incorporated herein by reference in their entireties.

[0002] All patents, patent applications and publications cited herein are hereby incorporated by reference in their entirety. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art as known to those skilled therein as of the date of the invention described and claimed herein.

[0003] This patent disclosure contains material that is subject to copyright protection. The copyright owner has no objection to the facsimile reproduction by anyone of the patent document or the patent disclosure as it appears in the U.S. Patent and Trademark Office patent file or records, but otherwise reserves any and all copyright rights.

INCORPORATION-BY-REFERENCE OF SEQUENCE LISTING

[0004] The contents of the text file named "[]", which was created on [], are hereby incorporated by reference in their entirety.

FIELD OF THE INVENTION

[0005] This invention is directed to chimeric antigen receptors and cells comprising the same, wherein the cells further secrete monoclonal antibodies locally at a tumor site.

BACKGROUND OF THE INVENTION

[0006] Clear cell renal cell carcinoma (ccRCC) is the major type of RCC which is among the 10 most common cancers in both men and women. Other types of renal carcinoma include papillary renal cell carcinoma, chromophobe renal cell carcinoma, and other or unclassified types of renal cell carcinoma. See, for example, Lancet 373 (2009) 1119-32.

SUMMARY OF THE INVENTION

[0007] Other objects and advantages of this invention will become readily apparent from the ensuing description.

[0008] Aspects of the invention are drawn towards an engineered cell comprising a chimeric antigen receptor. In embodiments, the chimeric antigen receptor comprises an extracellular ligand binding domain that is specific for a first antigen and a second antigen on the surface of a cancer cell, wherein the first antigen comprises CAIX and the second antigen comprises CD70.

[0009] In embodiments, the CAR further comprises a transmembrane polypeptide, an intracellular signaling domain, and/or a co-stimulatory domain.

[0010] In embodiments, the extracellular ligand binding domain comprises an antibody or fragment thereof. For example, the antibody comprises a VH and/or VL according to Table 2, or any combination thereof. For example, the antibody comprises a VH and/or VL according to Table 4, or any combination thereof. For example, the extracellular binding domain comprises a a VH and/or a VL of Table 2 and Table 4, or any combination thereof. For example, the antibody comprises a CDR1, CDR2, and/or CDR3 of Table 1, or any combination thereof. For example, the antibody comprises a CDR1, CDR2, and/or CDR3 of Table 3, or any combination thereof. For example, the extracellular binding domain comprises a CDR1, CDR2, and/or CDR3 of Table 1 and Table 3, or any combination thereof.

[0011] In embodiments, the engineered cell expresses and secretes a recombinant polypeptide.

[0012] In embodiments, the recombinant polypeptide comprises an antibody or fragment thereof, or a cytokine. For example, the recombinant polypeptide comprises an antibody or fragment thereof specific for TIGIT, GITR, PD-L1, PD-L2, PD-1, CTLA-4, VISTA, CD70, TIM-3, LAG-3, CD40L, or CCR4. For example, the recombinant polypeptide comprises a cytokine comprising IL-12, IL-15, or IL-18.

[0013] In embodiments, the recombinant polypeptide modulates the immune system of a subject. For example, the recombinant polypeptide is an immune checkpoint blockade antibody.

[0014] In embodiments, the recombinant polypeptide modulates tumor vasculogenesis. For example, the recombinant polypeptide can be specific for VEGF, VEGFR1, VEGFR2, PDGF, Ang-1, or AT1.

[0015] In embodiments, the engineered cell is a T cell, an NK cell, or an NKT cell. For example, the T cell is CD4+, CD8+, CD3+ panT cells, or any combination thereof. For example, the T cell is a mixed population of CD4+ and CD8+ T cells.

[0016] Aspects of the disclosure are further drawn towards a nucleic acid construct encoding a chimeric antigen receptor. In embodiments the chimeric antigen receptor comprises an extracellular ligand binding domain that is specific for a first antigen and a second antigen on the surface of a cancer cell, wherein the first antigen comprises CAIX and the second antigen comprises CD70.

[0017] In embodiments, the nucleic acid construct further encodes for a transmembrane polypeptide, an intracellular signaling domain, and/or a co-stimulatory domain.

[0018] In embodiments, the nucleic acid construct further encodes for a recombinant polypeptide.

[0019] Aspects of the disclosure are further drawn towards a vector comprising the nucleic acid construct described herein.

[0020] Still further, aspects of the disclosure are drawn towards a cell comprising the vector described herein.

[0021] Aspects of the disclosure are also drawn towards a method for treating a subject afflicted with cancer. In embodiments, the method comprises administering to a subject a therapeutically effective amount of the engineered cell described herein.

[0022] Also, aspects of the disclosure are drawn towards a method of reducing progression or promoting regression of a cancer in a subject. In embodiments, the method comprises administering the subject a therapeutically effective amount of the engineered cell described herein.

[0023] Still further, aspects of the disclosure are drawn towards a method of reducing cellular proliferation of a cancer cell in a subject. In embodiments, the method comprises administering the subject a therapeutically effective amount of the engineered cell described herein.

[0024] In embodiments, the cancer comprises renal cell carcinoma.

[0025] Aspects of the disclosure are drawn to a chimeric antigen receptor (CAR) comprising an extracellular ligand binding domain, wherein the extracellular ligand binding domain is specific for a first antigen and a second antigen on the surface of a cancer cell, wherein the first antigen comprises CAIX and the second antigen comprises CD70.

[0026] In embodiments, the CAR further comprises a transmembrane polypeptide, an intracellular signaling domain, and/or a co-stimulatory domain.

[0027] In embodiments, the extracellular ligand binding domain comprises an antibody or fragment thereof.

[0028] Further, aspects of the invention are drawn towards a cell comprising the chimeric antigen receptor (CAR) as described herein.

[0029] Still further, aspects of the invention are drawn towards an engineered cell comprising a first chimeric antigen receptor and a second chimeric antigen receptor, wherein the first chimeric antigen receptor comprises an extracellular ligand binding domain specific for CAIX, and wherein the second chimeric antigen receptor comprises an extracellular ligand binding domain that is specific for CD70.

[0030] In embodiments, the engineered cell expresses and secretes a recombinant polypeptide.

[0031] In embodiments, the first chimeric antigen receptor and the second chimeric antigen receptor are expressed from a single nucleic acid construct.

BRIEF DESCRIPTION OF THE FIGURES

[0032] **FIG. 1** shows a schematic of bispecific tandem CAR T anti-CD70 and anti-CAIX scFvs were combined in different permutations by changing the order of the two targeting scFvs with various linkers and hinges connected to a costimulatory domain (non-limiting examples comprising CD28, 41BB, CD28-41BB, or 41BB-CD28) and an activating domain (such as, CD3). In order to elevate the efficacy (such as, address heterogeneity) and safety (such as, limit on-target off-tumor effect), the 2nd generation CAR T cell factories can be designed with the introduction of the second cassette anti-CD70 scFv.

[0033] **FIG. 2** shows that 2nd generation CAR T cell factories can be established by combining a series of CARs (such as different scFvs, linkers, and hinges) with immune checkpoint blockade as payload. Based on these data, 8 constructs were established by using G36 as the anti-CAIX scFv and B7 as the anti-CD70 scFv. The skilled artisan will recognize that any scFvs can be utilized according to the invention. The dual CAR engineering is the

most important part and the payload was replaced with zsgreen to indicate the transduction efficiency.

[0034] FIG. 3 shows CAIX and CD70 upregulated and co-expressed on ccRCC. IHC staining of ccRCC primary cell lines generated from patient samples indicates that CAIX and CD70 are both highly expressed and co-expressed on ccRCC. Without wishing to be bound by theory, CAIX and CD70 are two potential targets for ccRCC therapy. To validate targets, we conducted IHC for staining of CAIX and CD70 in primary cell line which is generated from ccRCC patients. From these images, we can see that CAIX and CD70 staining are 100% positive, demonstrating that CAIX and CD70 are highly and simultaneously expressed in ccRCC. This is presently being validated in a study which determines CAIX and CD70 expression on ccRCC patients with at least 150 samples.

[0035] FIG. 4 shows CAIX and CD70 upregulated and co-expressed on ccRCC. IHC staining of ccRCC patient samples indicates that CAIX and CD70 are both highly expressed and co-expressed on ccRCC.

[0036] FIG. 5 shows establishment of CRISPR skrc-59 cell lines. For further evaluation *in vitro*, four CRISPR engineered skrc-59 cell lines were established with four different phenotypes, (i) CAIX+ CD70+, (ii) CAIX+ CD70-, (iii) CAIX- CD70+, and (iv) CAIX- CD70-. Four CRISPR skrc-59 cell lines have been established. Four different phenotypes, (i) CAIX+ CD70+, (ii) CAIX+ CD70-, (iii) CAIX- CD70+, and (iv) CAIX- CD70- are used for the *in vitro* assay described herein. The corresponding table shows CAIX and CD70 expression level on 4 different cell lines that were quantified.

[0037] FIG. 6 shows graphs of anti-CD70 minibodies that showed selective binding to CD70+ SKRC59 cells. Phage display (panned against CD70+ skrc-59 cells and subtracted against CD70- skrc-59 cells) indicates that a series of anti-CD70 minibodies showed binding with CD70 positive ccRCC skrc-59 cells. Anti-CD70 minibodies were expressed via Expi293

cells in a 6 well plate. 3 days after transfection, the supernatant was harvested and IgG quantification ELISA (Bethyl) was conducted to approximate the concentration of minibody in the supernatant. This approximate concentration was used to normalize the supernatants for FACS binding curves. Staining was carried out via standard FACS staining protocol using an anti-hFc-APC secondary. As can be seen, one of the hits with killing activity (#9) is very nonspecific. The other two monoCARs which exhibit killing activity (#3, 7) show good specificity to CD70.

[0038] FIG. 7 demonstrates anti-CD70 CAR T cells show killing activity in Celigo killing assay. These anti-CD70 scFv candidates were cloned into vectors and were packaged into lentiviruses to transduce primary T cells. CAR T cells were assessed for antiproliferation activity in Celigo assay. Results showed that #3, #7, #9 have efficacy compared to CD70 ligand CD27 CAR T cells. Also, these hits were cloned into pHAGE vector and conducted Celigo killing assay. These graphs showed that anti-CD70 CAR T cells exhibited killing activity in Celigo killing assay with Effector: Target ratio of 2:1. We compared the cell number in different time points and normalized with untreated cells in corresponding time point. So that Y axis is the percentage of treated cells/ untreated cells and X axis is different treatment. After 24h coculture with T cells, the SKRC59 cell number of treatment group is significantly decreased compared to untreated group and untransduced T cell group, demonstrating the anti-proliferation activity of these CAR T cells. After 48h coculture, we found 3 candidates which are comparable to CD27, a ligand of CD70. So that 3,7,9 were used for the further Chromium 51 killing assay.

[0039] FIG. 8 demonstrates anti-CD70 CAR T cells show killing activity in Chromium 51 killing assay on CAIX+ CD70+ cells. CAR T cells were also assessed for killing activity in Chromium 51 release assay. Results showed that #7 CAR T cells have the enhanced efficacy compared to CD27 (CD70 ligand) CAR T cells. A chromium 51 4h release assay was

conducted. After 4h incubation with chromium 51 labeled target cells, it validated that B7 has killing activity against SKRC59 CD70+ cells.

[0040] **FIG. 9** shows 2nd generation CAR-T cells. CAR T cells can be generated by using zsGreen instead of immune checkpoint blockade. Based on these data, 8 constructs were established by using G36 as the anti-CAIX scFv and B7 as the anti-CD70 scFv. The dual CAR engineering is the most important part and the payload was replaced with zsgreen to indicate the transduction efficiency.

[0041] **FIG. 10** shows transfected 293T cells binding with CAIX-PE. 293T cells were transfected with different constructs of dual CAR and the binding assay with PE labeled CAIX protein was performed. All dual CARs bind to CAIX, and different orientation of two scFvs influence the EC50 of anti-CAIX scFv binding. Anti-CAIX scFv G36 prefer the 2nd cassette after linker. For example, 293T cells were transfected with these 8 bispecific constructs and also corresponding monoCARs and stained with CAIX-PE. After normalization by transfect efficiency, different orientation of two scFvs influence the EC50 of anti-CAIX scFv binding. Anti-CAIX scFv G36 prefer to in the 2nd cassette after linker.

[0042] **FIG. 11** shows transfected 293T cells binding with CD70-APC. 293T cells were transfected with different constructs of dual CAR and the binding assay was performed with PE labeled CAIX protein. All dual CARs bind to CD70, and different orientation of two scFvs does not influence the EC50 of anti-CD70 scFv binding. Anti-CD70 scFv B7 does not have a significant preference. For example, 293T cells were transfected with these 8 bispecific constructs and corresponding monoCARs and stained with CD70-PE. After normalization by transfect efficiency, different orientation of the two scFvs does not influence the EC50 of anti-CD70 scFv binding. Anti-CD70 scFv B7 does not have a significant preference.

[0043] **FIG. 12** shows B7-GGGGS3-G36 killing assay with Celigo. B7-GGGGS3-G36 CAR for example was taken and selective killing assay was performed on mixed CAIX+ CD70+ dual positive cells with CAIX+ or CD70+ single positive or CAIX- CD70- cells. B7-GGGGS3-G36 CAR T cells had more killing activity on the targeted cells (CAIX+ CD70+) than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-).

[0044] **FIG. 13** shows B7-GGGGS3-G36 killing assay with FACS. Four different CRISPR engineered skrc-59 cells were transduced with BFP fluorescence group. Cells were mixed with 1:1:1:1 ratio and were treated with B7-GGGGS3-G36 CAR T cells or culture medium. After treatment, stain cells with PE labeled anti-CD70 antibody and APC labeled anti-CAIX antibody and, perform flow cytometry. It showed that B7-GGGGS3-G36 has selective killing to CAIX+ CD70+ cells whose population is reduced from 26.7% to 18.5%.

[0045] **FIG. 14** is a schematic of fine-tuned anti-CAIX CAR T. anti-CAIX scFvs with various KDs were generated as CARs and corresponding killing activity was assessed. In order to limit on-target off-tumor effect, the 2nd generation CAR T cell factories have been designed with the introduction of the second cassette anti-CD70 scFv.

[0046] **FIG. 15** is a diagram rendering of fine-tuned anti-CAIX CAR-T cells. For validation experiments, CAR T cells were generated by using zsGreen instead of immune checkpoint blockade. 8 constructs were established by using G36 as the anti-CAIX scFv and B7 as the anti-CD70 scFv. The dual CAR engineering is an important part and the payload was replaced with zsGreen to indicate the transduction efficiency. A series of antibodies against CAIX were identified with different KD values from 1.49nM to 99.58nM (See also FIG. 16). 13 antibodies were classified into 4 groups by binding experiment.

[0047] **FIG. 16** depicts table showing 19 anti-CAIX ScFvs identified and the corresponding binding.

[0048] **FIG. 17** shows graphs of anti-CAIX monoCAR T killing on CAIX+CD70+. These anti-CAIX CAR T cells were assessed for antiproliferation activity in Celigo assay. Results showed based on killing activity, 19 CARs can be separated into 4 groups. G37, G39, G125 (++++> G10, G21, G36, G40, G45, G57, G62, G98, G106, G119 (++> G6, G9, G17, G27, G28 (++)> G104 (++)).

[0049] **FIG. 18** shows graphs of Anti-CAIX monoCAR T killing on CAIX+CD70+. These anti-CAIX CAR T cells were assessed for antiproliferation activity in Celigo assay. Results showed based on killing activity, that 19 CARs can be separated into 4 groups. G37, G39, G125 (++++> G10, G21, G36, G40, G45, G57, G62, G98, G106, G119 (++> G6, G9, G17, G27, G28 (++)> G104 (++)).

[0050] **FIG. 19** shows Anti-CAIX monoCAR T killing on CAIX+CD70+. These anti-CAIX CAR T cells were assessed for antiproliferation activity in Celigo assay. Results showed based on killing activity, 19 CARs can be separated into 4 groups. G37, G39, G125 (++++> G10, G21, G36, G40, G45, G57, G62, G98, G106, G119 (++> G6, G9, G17, G27, G28 (++)> G104 (++)).

[0051] **FIG. 20** is a schematic of a selective killing assay (e.g., using either a Celigo assay or FACS assay). In order to explore the selectivity of bispecific CARs on 4 different cell lines, 2 different selectivity assay were performed. For example, CAIX+ CD70+ cells were mixed with mono positive cells and then CART cells were added. After certain time incubation, the number of double positive cells and single positive cells were measured through Celigo. Also, 4 different cell lines were mixed with the ratio of 1:1:1:1, and CART cells were added. After 24h coculture, all the cells are collected and stained with CAIX-APC and CD70-PE by FACS.

[0052] **FIG. 21** are graphs showing tandem CAR killing on skrc-59 mixed cells. Selective killing assay was performed on mixed CAIX+ CD70+ dual positive cells with CAIX+, or

CD70+ single positive, or CAIX- CD70- cells. Bispecific CAR T cells had more killing activity on the targeted cells (CAIX+ CD70+) than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-) with E:T=5:1.

[0053] **FIG. 22** are graphs showing tandem CAR killing on skrc-59 mixed cells. Selective killing assay was performed on mixed CAIX+ CD70+ dual positive cells with CAIX+, or CD70+ single positive, or CAIX- CD70- cells. Bispecific CAR T cells had more killing activity on the targeted cells (CAIX+ CD70+) than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-) with E:T=5:1.

[0054] **FIG. 23** are graphs showing tandem CAR killing on skrc-59 mixed cells. Selective killing assay was performed on mixed CAIX+ CD70+ dual positive cells with CAIX+, or CD70+ single positive, or CAIX- CD70- cells. Bispecific CAR T cells had more killing activity on the targeted cells (CAIX+ CD70+) than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-) with E:T=10:1.

[0055] **FIG. 24** shows CAIX and CD70 upregulated and co-expressed on ccRCC.

[0056] **FIG. 25** shows design of CARs to address unwanted side-effects previously associated with CART cell therapy. To apply CART cell therapy in solid tumors, we elevate the efficacy and safety profiling. First, to restore the effective anti-cancer immunity, CAR-T cell factories were developed, which is able to modulate the tumor microenvironment through secreting human anti-immune checkpoint inhibitor monoclonal antibodies (mAbs) locally at the tumor site to reverse T cell exhaustion. In order to limit on-target off-tumor side effects, fine-tuned CARs were designed with reduced affinity scFv to expand the therapeutic window by limiting the recognition of the tumor associated antigen on normal tissues. Also, bispecific CARs were designed by introducing the second cassette, such as anti-CD70 scFv, to increase the preferential killing on the double positive population, resulting in the elevation of safety profiling.

[0057] **FIG. 26** shows CAR-T cell factories show enhanced killing. We engineered a bicistronic lentiviral vector to express the anti-CAIX scFv linked to CD28 and CD3z signaling domains in the first cassette and anti-PDL1 mAb in the second expression cassette. So that the CAR-T cell factories are able to target CAIX and secrete checkpoint blockade inhibitor at the tumor site to convert suppressive tumor microenvironment. In RCC orthotopic mouse model, 1E7 CAR-T cells were injected by i.v. on Day 0 and 2.5E6 CAR-T cells were injected on Day 17. CAR-T cell factories showed enhanced killing efficacy compared to the CAR-T cells secreting irrelevant antibodies in our orthotopic RCC mouse model.

[0058] **FIG. 27** shows schematic of assessment of CAR-T on humanized orthotopic ccRCC mouse model.

[0059] **FIG. 28** shows MRI imaging of RCC mouse model.

[0060] **FIG. 29** shows a schematic of an experiment designed to establish stress model and compare 2nd generation and 3rd generation CAR and CD8 vs. CD4/8 on orthotopic ccRCC mouse model.

[0061] **FIG. 30** shows 41BB performed superior killing *in vivo*. Efficient tumor regression was observed in G36-41BB and G36-CD28-41BB treated group at the dose of 1E7. Mixed CD4/CD8 G36-41BB CAR-T performed superior killing even at the dose of 3E6. Comparing two 2nd generation CAR constructs and one 3rd generation CAR construct, G36-41BB outperformed than G36-CD28 and G36-CD28-41BB

[0062] **FIG. 31** shows G36-41BB stress model. Dose of 3E6 can be used for G36-41BB stress model.

[0063] **FIG. 32** shows 41BB performed superior killing *in vivo*.

[0064] **FIG. 33** shows 41BB CAR-T cell performed the best expansion *in vivo*.

[0065] **FIG. 34** shows CD8 vs. CD4/CD8.

[0066] **FIG. 35** shows CD4 and CD8 T cells expansion *in vivo*.

[0067] **FIG. 36** shows a schematic of a fine-tuned anti-CAIX CAR-T which has been designed to limit on-target off-tumor effect. Without wishing to be bound by theory, lowering the affinity of anti-CAIX scFv, the CAR only recognizes the high density of CAIX on ccRCC. A series of anti-CAIX scFvs with various KDs were cloned into lentiviral vectors and corresponding CAR-T cells were generated and assessed the killing activity.

[0068] **FIG. 37** is a schematic detailing the "therapeutic window". "Therapeutic window" is a term originally from pharmaceutical toxicology and can refer to a range of doses between efficacy and toxicity, achieving the highest therapeutic benefit without resulting in unacceptable toxicity; it is the range between the minimum effective dose (MED) and the maximum tolerated dose (MTD. This concept was applied to optimize CAR-T therapy. To expand the therapeutic window, we can fine tune the CAR affinity with the antigens by assembling the scFvs with different KDs in the CAR construct, such as from 1nm to 100nm. Ideally, after optimization, the CAR only recognize the high density antigens on tumor cells not the low density antigens on normal cells. Targeting antigens expressed exclusively on tumor cells or antigens that are expressed only on non-critical tissues widens the therapeutic window as direct toxicity on vital tissues would not occur. On the other hand, targeting antigens that are expressed in critical normal tissues/cells narrows the window by decreasing MTD

[0069] **FIG. 38** is a schematic describing expanding the therapeutic window to address on-target off-tumor side effects.

[0070] **FIG. 39** shows IHC double staining of CAIX and CD70 on patient samples. We found that CD70 and CAIX are highly expressed and co-expressed on ccRCC. Thus, we choose CD70 as our second target.

[0071] **FIG. 40** shows IHC double staining of CAIX and CD70 on patient samples.

[0072] **FIG. 41** shows a panel of scFvs against CAIX with different KD values from 1.49nM to 99.58nM as the table shows. Corresponding CAR-T cells were generated and screened by Celigo killing assay

[0073] **FIG. 42** shows that there is correlation between affinity of scFv and killing of CART cells, 19 CARs were divided into 4 groups according to the cytotoxicity and to be tested against skrc-59 cells with different expression levels of CAIX. G37, G39, G125> G10, G21, G36, G40, G45, G57, G62, G98, G106, G119> G6, G9, G17, G27, G28> G104.

[0074] **FIG. 43** shows "Or" gating to capture tumor cell heterogeneity but not kill low target density healthy cells.

[0075] **FIG. 44** shows CD70 can be an ideal target as the second target for our bispecific CAR since it's highly expressed on kidney cancer, especially clear cell renal cell carcinoma.

[0076] **FIG. 45** shows IHC double staining of CAIX and CD70 on ccRCC patient samples. CD70 and CAIX are highly expressed and co-expressed on ccRCC. Thus, we choose CD70 as our second target.

[0077] **FIG. 46** shows IHC double staining of CAIX and CD70 on ccRCC patient samples. CD70 and CAIX are highly expressed and co-expressed on ccRCC. Thus, we choose CD70 as our second target.

[0078] **FIG. 47** shows CAIX and CD70 upregulated and co-expressed on ccRCC.

[0079] **FIG. 48** shows anti-CD70 minibodies showed selective binding to CD70+ SKRC59 cells. By phage display (panned against CD70+ skrc-59 cells and subtracted against CD70-skrc-59 cells), we discovered a series of anti-CD70 minibodies which showed promising binding with CD70 positive ccRCC skrc-59 cells. Anti-CD70 minibodies were expressed via Expi293 cells in a 6 well plate. 3 days after transfection, the supernatant was harvested and IgG quantification ELISA (Bethyl) was conducted to approximate the concentration of minibody in the supernatant. This approximate concentration was used to normalize the

supernatants for FACS binding curves. Staining was carried out via standard FACS staining protocol using an anti-hFc-APC secondary. One of the hits with killing activity (#9) is very nonspecific. The other two monoCARs which exhibit killing activity (#3, 7) show good specificity to CD70.

[0080] FIG. 49 shows anti-CD70 B7 CAR-T cells showed promising killing against CD70+ skrc-59 cells. By phage display (panned against CD70+ skrc-59 cells and subtracted against CD70- skrc-59 cells), we discovered a series of anti-CD70 minibodies which showed promising binding with CD70 positive ccRCC skrc-59 cells. Also, these hits were cloned into lentiviral vector and corresponding CART cells were conducted Celigo killing assay. From the screening, we found B7 as the candidate.

[0081] FIG. 50 shows bispecific CAR constructs. A series of constructs were established by using G36 as the anti-CAIX scFv and B7 as the anti-CD70 scFv in different rotations (or orientations) with different linkers.

[0082] FIG. 51 shows that anti-scFv G36 prefer to be in the 2nd cassette after the linker. Transfect 293T cells with these 8 bispecific constructs and also corresponding monoCARs and stain with CAIX-PE. After normalization by transfect efficiency, it shows that different orientation of two scFvs influence the EC₅₀ of anti-CAIX scFv binding. Anti-CAIX scFv G36 prefer to in the 2nd cassette after linker.

[0083] FIG. 52 shows that anti-CD70 scFv B7 has no preference. Transfect 293T cells with different constructs of dual CAR and perform the binding assay with APC labeled CD70 protein. After normalization by transfection efficiency, it shows that different orientation of two scFvs doesn't influence the EC₅₀ of anti-CD70 scFv binding. Anti-CD70 scFv B7 doesn't have a significant preference.

[0084] **FIG. 53** shows that establishment of 4 CRISPR skrc-59 cell lines. For further evaluation in vitro, 4 CRISPR engineered skrc-59 cell lines were established with 4 different phenotypes, CAIX+ CD70+, CAIX+ CD70-, CAIX- CD70+, and CAIX- CD70-.

[0085] **FIG. 54** shows B7-GGGGS3-G36 shows preferential killing. Take B7-GGGGS3-G36 CAR as example we performed selective killing assay on mixed CAIX+ CD70+ dual positive cells, with CAIX+ or CD70+ single positive cells, or CAIX- CD70- cells. It showed that B7-GGGGS3-G36 CAR-T cells have preferential killing activity against the targeted cells (CAIX+ CD70+) than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-). After certain time incubation, the number of double positive cells and single positive cells were measured through Celigo. And we can see that B7-GGGGS3 has limited selective index on CAIX+CD70+ while mixed with CAIX+ cells.

[0086] **FIG. 55** shows that B7-GGGGS3-G36 shows preferential killing. 4 different CRISPR engineered skrc-59 cells were transduced with BFP fluorescence group. Cells were mixed with 1:1:1:1 ratio and were treated with B7-GGGGS3-G36 CAR-T cells or culture medium. After treatment, stain cells with PE labeled anti-CD70 antibody and APC labeled anti-CAIX antibody and, perform flow cytometry. It showed that B7-GGGGS3-G36 has selective killing to CAIX+ CD70+ cells whose population is reduced from 26.7% to 18.5%. In this selective killing data from FACS, it showed that B7-GGGGS3-G36 has selective killing to CAIX+ CD70+ cells whose population is reduced from 26.7% to 18.5%.

[0087] **FIG. 56** shows a schematic of a bispecific split CAR T. Anti-CD70 and anti-CAIX scFvs were expressed on cell surface with different costimulation domains. See, for example, dual(split) CAR T, Nat Rev Cancer 16(9):566-81.

[0088] **FIG. 57** shows split CAR killing on ccRCC primary cancer cell. Split CAR T cells were assessed killing activity on primary ccRCC cancer cells with monoCAR T and Tandem

CAR T cells. It showed that split CAR achieved a superior killing at low E:T ratio, like 1:1 ratio.

[0089] **FIG. 58** provides multiple sequence alignment of amino acid sequences of anti-carbonic anhydrase IX (G250) scFv clones.

[0090] **FIG. 59** provides alignment of human and mouse CAIX amino acid sequences.

[0091] **FIG. 60** provides structure of carbonic anhydrase IX (G250).

[0092] **FIG. 61** shows CAR T cell killing assay of Skrc-59 transduced cells. Celigo image cytometry is shown.

[0093] **FIG. 62** shows homology study of 10 different species. Tree: distance on each branch equals number of differences between sequences (e.g. 0.1 means 10 % differences between two sequences); distance between 2 species equals sum length of all branches connecting them. homology > 60% → potential cross reactivity.

[0094] **FIG. 63** shows establishment of skrc-59 stable cell lines expressing CAIX or CD70 from different species. CRISPR knock out CAIX-/CD70- skrc-59 cells → transduced with the 10 different constructs (CD70 or CAIX from the 5 different species) → sorted by FACS. Half transduced with BFP (for Celigo). Stained with commercial antibodies.

[0095] **FIG. 64** shows binding assay of anti-CAIX(G36) scFv. Normalized with the binding data of commercial antibodies. Analyzed using a non-linear regression and a log (agonist) vs. response model.

[0096] **FIG. 65** shows killing assay of anti-CAIX (G36) CAR T cells. Monkey 100% (same killing as human, cross reactive); mouse, hamster 50% (significant killing).

[0097] **FIG. 66** shows killing assay of anti-CD70 (B7) CAR T cells.

[0098] **FIG. 67** shows killing assay of 20 different anti-CAIX CAR T cells.

[0099] **FIG. 68** shows fold change killing assay G36 E:T 10:1.

[00100] **FIG. 69** shows killing assay G36 E:T 5:1.

- [00101] **FIG. 70** shows killing assay G36 n=2.
- [00102] **FIG. 71** shows killing assay 20 anti-CAIX scFvs.
- [00103] **FIG. 72** shows all candidates monkey.
- [00104] **FIG. 73** shows all candidates mouse.
- [00105] **FIG. 74** shows all candidates hamster.
- [00106] **FIG. 75** shows fold change B7 killing assay.
- [00107] **FIG. 76** shows the amino acid sequence and germline alignment of anti-cd70 antibodies.
- [00108] **FIG. 77** shows a nucleic acid construct of a split car.
- [00109] **FIG. 78** shows the amino acid sequence and germline alignment of anti-PDL1 and anti-PD1 sequences.
- [00110] **FIG. 79** shows the amino acid sequence and germline alignment of anti-CAIX antibodies.
- [00111] **FIG. 80** shows tandem CAR B7-GGGGS5-G36 cytotoxicity.
- [00112] **FIG. 81** shows the amino acid sequence and germline alignment of anti-TIGIT antibodies.
- [00113] **FIG. 82** shows the anti-PD-L1 amino acid sequences.
- [00114] **FIG. 83** shows the anti-PD1 nucleic acid and amino acid sequences.

DETAILED DESCRIPTION OF THE INVENTION

[00115] Chimeric Antigen Receptor (CAR) T-cell therapies represent an exciting area of discovery that has already revolutionized treatment in several blood-borne cancers. In Acute Lymphoblastic Leukemia (ALL), for example, remission rates were demonstrated to be 80 to 90 percent, leading to FDA approval. The technology can utilize a patient's own immune cells to fight their cancer by engineering them to better recognize particular proteins located

on the cancer cells. For example, after that change has been effected in the lab, the immune cells are grown extensively in the lab, multiplying their cancer-killing potential, and finally infused back to the patient where they can attack cancer wherever it is in the body with increased efficacy and numbers.

[00116] CAR T-cell therapies have been largely elusive in solid tumors such as renal cell carcinoma (RCC) because the cancer cells create a tumor microenvironment that turns off immune penetration. We have successfully created and advanced mouse models that can provide crucial information for this technology. Our approach relies upon the creation of humanized RCC in mice that can be studied to better understand their immunologic fingerprint. Examining how human RCC behaves in mice has allowed the lab to identify antigens, such as Carbonic Anhydrase IX [CAIX] and CD70, that could be targeted using CAR T-cell therapy in humans and other mammals. To large degree, these two antigens, namely CAIX and CD70, are only found together on kidney cancer tumor cells, allowing the immune system to attack locally with minimal impact on healthy tissue. There have also been advances in counteracting the immunosuppressive effects within the microtumor environment by engineering T-cells that produce and/or secrete check-point inhibitor antibodies. Pairing these approaches could allow for dramatically improved efficacy within the tumor microenvironment.

[00117] Described herein are CAR T-cells that target CAIX and CD70, for example, that can be administered to animals with close monitoring for changes in the tumor microenvironment and subsequent anti-tumor effects. The skilled artisan will recognize that these approaches can also be combined with currently available technology, such as the use of immune checkpoint inhibitors, that have already positive impacted the RCC clinical landscape.

[00118] Detailed descriptions of one or more embodiments are provided herein. It is to be understood, however, that the present invention may be embodied in various forms.

Therefore, specific details disclosed herein are not to be interpreted as limiting, but rather as a basis for the claims and as a representative basis for teaching one skilled in the art to employ the present invention in any appropriate manner.

[00119] The singular forms “a”, “an” and “the” include plural reference unless the context clearly dictates otherwise. The use of the word “a” or “an” when used in conjunction with the term “comprising” in the claims and/or the specification may mean “one,” but it is also consistent with the meaning of “one or more,” “at least one,” and “one or more than one.”

[00120] Wherever any of the phrases “for example,” “such as,” “including” and the like are used herein, the phrase “and without limitation” is understood to follow unless explicitly stated otherwise. Similarly “an example,” “exemplary” and the like are understood to be nonlimiting.

[00121] The term “substantially” allows for deviations from the descriptor that do not negatively impact the intended purpose. Descriptive terms are understood to be modified by the term “substantially” even if the word “substantially” is not explicitly recited.

[00122] The terms “comprising” and “including” and “having” and “involving” (and similarly “comprises”, “includes,” “has,” and “involves”) and the like are used interchangeably and have the same meaning. Specifically, each of the terms is defined consistent with the common United States patent law definition of “comprising” and is therefore interpreted to be an open term meaning “at least the following,” and is also interpreted not to exclude additional features, limitations, aspects, etc. Thus, for example, “a process involving steps a, b, and c” means that the process includes at least steps a, b and c. Wherever the terms “a” or “an” are used, “one or more” is understood, unless such interpretation is nonsensical in context.

[00123] As used herein, the term “about” can refer to approximately, roughly, around, or in the region of. When the term “about” is used in conjunction with a numerical range, it modifies that range by extending the boundaries above and below the numerical values set forth. In general, the term “about” is used herein to modify a numerical value above and below the stated value by a variance of 20 percent up or down (higher or lower).

[00124] **Chimeric Antigen Receptor (CAR) T-cell Therapies**

[00125] Chimeric-antigen receptor (CAR) T-cell therapies redirect a patient’s T-cells to kill tumor cells by the exogenous expression of a CAR. Referring to FIG. 1, for example, a CAR is a membrane spanning fusion protein that links the antigen recognition domain of an antibody or fragment to the intracellular signaling domains of the T-cell receptor and co-receptor. For example, chimeric antigen receptors fuse antigen-specific antibody fragments to T-cell co-stimulatory domains and the CD3 zeta intracellular signaling domain, allowing for the re-direction of T-cells towards an antigen presented on a cell of interest, for example, onto tumor cells.

[00126] The term “antibody” herein is used in the broadest sense and can refer to immunoglobulin molecules and immunologically active portions of immunoglobulin (Ig) molecules, i.e., molecules that contain an antigen binding site that specifically binds (immunoreacts with) an antigen. “Specifically binds” or “immunoreacts with” can refer to the antibody reacting with one or more antigenic determinants of the desired antigen and does not react with other polypeptides. Antibodies of the disclosure can include, but are not limited to, polyclonal, monoclonal, humanized, fully human, bispecific, multispecific, chimeric, dAb (domain antibody), single chain antibodies, Fab, Fab’ and F(ab’)2 fragments, scFvs, diabodies, minibodies, scFv-Fc fusions, and Fab expression libraries. Unless specified to the contrary, any reference to “antibody” or “antibodies” made herein encompasses, for

example, any (or all) of these molecules so long as they exhibit the desired antigen-binding activity.

[00127] A single chain Fv (“scFv”) polypeptide molecule is a covalently linked VH::VL heterodimer, which can be expressed from a gene fusion including VH- and VL-encoding genes linked by a peptide-encoding linker. (See Huston et al. (1988) Proc Nat Acad Sci USA 85(16):5879-5883). A number of methods have been described to discern chemical structures for converting the naturally aggregated, but chemically separated, light and heavy polypeptide chains from an antibody V region into an scFv molecule, which will fold into a three dimensional structure substantially similar to the structure of an antigen-binding site. See, e.g., U.S. Pat. Nos. 5,091,513; 5,132,405; and 4,946,778.

[00128] Solid tumors offer unique challenges for CAR-T therapies. Unlike blood cancers, tumor-associated target proteins are overexpressed between the tumor and healthy tissue resulting in on-target/off-tumor T-cell killing of healthy tissues. Furthermore, immune repression in the tumor microenvironment (TME) limits the activation of CAR-T cells towards killing the tumor. Aspects of the disclosure address these problems. For example, embodiments comprise T cells comprising a bispecific CAR that (a) targets two antigens on a cancer cell to mitigate on-target/off-tumor T-cell killing. Referring to FIG. 12, for example, B7-GGGGS-G36 CAR T cells had more killing activity on the targeted renal carcinoma cells (CAIX+ CD70+) than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-). In embodiments, the bispecific CAR can also (b) secrete a checkpoint blockade antibody that removes repression in the tumor microenvironment. Other embodiments comprise a fine-tuned CAR that only recognizes the high density antigens on tumor cells, not the low density antigens on normal cells. Without wishing to be bound by theory, this can be accomplished by, for example, lowering the affinity of the antibody or antibodies associated with the CAR.

[00129] FIG 1, for example, provides a schematic of a bispecific CAR that targets two antigens, such as CAIX and CD70. Bispecific CARs can refer to CARs that has binding specificities for at least two different antigens. For example, bispecific CARs can comprise monoclonal antibodies, such as human or humanized antibodies, or fragments thereof. In the present case, one of the binding specificities is CAIX and/or CD70. The second binding target is any other antigen, and advantageously is a cell-surface protein or receptor or receptor subunit. For example, one of the binding specificities is for CAIX and the second binding specificity is for CD70.

[00130] As reported (see, for example, J Clin Oncol 24 (2006) 20-22; Molecular Therapy 21 (2013) 4), the first clinical trial of the first generation anti-CAIX G250 CAR-T cells on renal cell carcinoma RCC (RCC) patients failed due to on-target, off-tumor side effects. 2 of the first 3 patients developed hepatitis due to CAIX expression on bile ducts. The introduction of a second antibody, such as a second scFv, can result in elevated safety to the subject, because the dual targeted CAR T can reduce or eliminate on-target/off-tumor effects. Referring to FIG. 24, for example, CAIX and CD70 expression is upregulated and co-expressed on ccRCC. On the other hand, CAIX is expressed on bile duct (mostly in cytoplasm), and CD70 is not expressed on bile duct. See, for example, British Journal of Cancer 103 (2010) 676-684.

[00131] Bispecific antibodies can be prepared as full length antibodies or antibody fragments (e.g. F(ab')₂ bispecific antibodies). Methods for making bispecific antibodies are known in the art. See for example U.S. patent 8,329,178, which is incorporated herein by reference in its entirety.

[00132] Antibody molecules obtained from humans relate to any of the classes IgG, IgM, IgA, IgE and IgD, which differ from one another by the nature of the heavy chain present in

the molecule. Certain classes have subclasses as well, such as IgG1, IgG2, and others.

Furthermore, in humans, the light chain may be a kappa chain or a lambda chain.

[00133] The term “antigen-binding site,” or “binding portion” can refer to the part of the immunoglobulin molecule that participates in antigen binding. The antigen binding site is formed by amino acid residues of the N-terminal variable (“V”) regions of the heavy (“H”) and light (“L”) chains. Three highly divergent stretches within the V regions of the heavy and light chains, referred to as “hypervariable regions,” are interposed between more conserved flanking stretches known as “framework regions,” or “FRs”. Thus, the term “FR” refers to amino acid sequences which are naturally found between, and adjacent to, hypervariable regions in immunoglobulins. In an antibody molecule, the three hypervariable regions of a light chain and the three hypervariable regions of a heavy chain are disposed relative to each other in three dimensional space to form an antigen-binding surface. The antigen-binding surface is complementary to the three-dimensional surface of a bound antigen, and the three hypervariable regions of each of the heavy and light chains are referred to as “complementarity-determining regions,” or “CDRs.” See, for example, Tables 1 and 3, which provide the CDRs for anti-CAIX and anti-CD70 antibodies.

[00134] An emerging mechanism associated with the progression of tumors is the immune checkpoint pathway, which include cellular interactions that prevent excessive activation of T cells under normal conditions, allowing T cell function in a self-limited manner. As an evasion mechanism, many tumors are able to stimulate the expression of immune checkpoint molecules, resulting in an anergic phenotype of T cells that cannot restrain tumor progression. For example, emerging clinical data highlight the importance of one inhibitory ligand and receptor pair as an immune checkpoint: the programmed death-ligand 1 (PD-L1; B7-H1 and CD274) and programmed death receptor-1 (PD-1; CD279), in preventing killing of cancer cells by cytotoxic T-lymphocytes. PD1 receptor is expressed by many cell types

like T cells, B cells, Natural Killer cells (NK) and host tissues. Tumors and Antigen-presenting cells (APC) expressing PD-L1 can block T cell receptor (TCR) signaling of cytotoxic T-lymphocytes through binding to receptor PD-1, decreasing the production of cytokines and T cell proliferation. PD-L1 overexpression can be found in many tumor types and may also mediate an immunosuppressive function through its interaction with other proteins, including CD80 (B7.1), blocking its ability to activate T cells through binding to CD28.

[00135] Genetic engineering of human lymphocytes to express tumor-directed chimeric antigen receptors (CAR) can produce antitumor effector cells that bypass tumor immune escape mechanisms that are due to abnormalities in protein-antigen processing and presentation. Moreover, these transgenic receptors can be directed to tumor-associated antigens that are not protein-derived. In certain embodiments of the disclosure, there are lymphocytes (CARTS) that are modified to comprise at least a CAR, and in particular embodiments of the invention, a single CAR targeting two or more antigens (e.g., a bispecific CAR). In some embodiments, the cells comprise a split CAR, such as anti-CD70 and anti-CAIX scFvs expressed on cell surface with different costimulation domains. Further, some embodiments comprise a fine-tuned CAR. In some embodiments, the CARTS are further modified to express and secrete one or more polypeptides, such as for example an antibody or a cytokine, such as IL-12, IL-15, or IL-18. Such CARTS are referred to herein as armed CARTS or CAR factories. Armed CARTS allow for simultaneous secretion of the polypeptide locally at the targeted site (i.e., tumor site).

[00136] Referring to FIG. 56, for example, a split CAR comprises two or more CARs on the surface of a cell, such as a T cell or NK cell. The CARs can be specific for two or more antigens, such as CD70 and CAIX. FIG. 77 provides an example of a nucleic acid construct encoding split CARs. In this example, the first CAR is specific for CAIX, and the second

CAR is specific for CD70. As described herein, the CARs can be in any orientation desired. For example, the first CAR can be specific for CD70 and the second CAR can be specific for CAIX. As shown in the example, first and the second CARs can be expressed from a single nucleic acid construct. In such an example, a nucleic acid encoding a cleavable linker can be located between the nucleic acids encoding the first and the second CAR. In other embodiments, the two CARs can be expressed in the same cell but from two separate nucleic acid constructs.

[00137] A modified TCR called chimeric antigen receptor (CAR), such as a CAR containing single chain variable antibody fragment (scFv) previously selected by high affinity against a specific tumor associated antigen, is a powerful new approach against cancer. The scFv presented in the CAR is linked to an intracellular signaling block that includes CD3 ζ to induce T cell activation followed by antigen binding. This structure is characteristic for first-generation CARs, which were improved to second- and generation CARs that link the signaling co-stimulatory endodomains of CD28, 4-1BB, or OX40 to CD3 or 3rd-generation CARs that links two elements to CD3 ζ in tandem. These endodomains are required for complete T cell activation during TCR recognition by antigen-presenting cells (APCs), improving cytokine production and proliferation of CAR-T cells. The effect of CART cells has heretofore been modest for the treatment of solid tumors, due to difficulty in finding unique tumor associated antigens, inefficient homing of T cells to tumor locations, low persistence of T cells in the body and the immunosuppressive microenvironment of solid tumors.

[00138] In particular cases, the lymphocytes can include a receptor that is chimeric, non-natural and engineered at least in part by the hand of man. In particular cases, the engineered chimeric antigen receptor (CAR) has one, two, three, four, or more components, and in some

embodiments the one or more components facilitate targeting or binding of the lymphocyte to one or more tumor antigen-comprising cancer cells.

[00139] The CAR according to the disclosure comprises at least one transmembrane polypeptide comprising at least one extracellular ligand-binding domain and; one transmembrane polypeptide comprising at least one intracellular signaling domain; such that the polypeptides assemble together to form a Chimeric Antigen Receptor. Exemplary CARS useful in aspects of the disclosure include those disclosed in for example PCT/US2006/046350, PCT/US2015/067178, PCT/US2015/067225, and PCT/US2019/022272, each of which are incorporated by reference herein in their entireties..

[00140] The term "extracellular ligand-binding domain" as used herein can refer to an oligo- or polypeptide that can bind a ligand. The domain can interact with a cell surface molecule. For example, the extracellular ligand-binding domain can be chosen to recognize a ligand that acts as a cell surface marker on target cells associated with a particular disease state.

[00141] In particular, the extracellular ligand-binding domain can comprise an antigen binding domain or antigen recognition domain derived from an antibody against an antigen of the target. The antigen binding domain or antigen recognition domain can be an antibody fragment. An "antibody fragment" can be a molecule other than an intact antibody that comprises a portion of an intact antibody that binds the antigen to which the intact antibody binds. Examples of antibody fragments include but are not limited to Fv, Fab, Fab', Fab'-SH, F(ab')₂; diabodies; linear antibodies; single-chain antibody molecules (e.g. scFv); and multispecific antibodies formed from antibody fragments. Referring to FIG. 1, for example, one embodiment comprises a CAR with two scFvs as the antigen recognition domains. Referring to FIG. 14, for example, one embodiment comprises a CAR with one scFv as the antigen recognition domain.

[00142] The antigen recognition domain can be directed towards any antigen target of interest. In embodiments, the antigen target of interest is on the surface of a cell, such as the surface of a cancer cell. Non-limiting examples of antigen targets comprise CAIX and/or CD70.

[00143] In some embodiments, the CAR is specific for CAIX and/or CD70.

[00144] In embodiments, said extracellular ligand-binding domain is a single chain antibody fragment (scFv) comprising the light (VL) and the heavy (VH) variable fragment of a target antigen specific monoclonal antibody joined by a flexible linker. The skilled artisan will recognize that embodiments can comprise different linkers which are typically known in the art. See, for example, *Chen, et al. "Fusion protein linkers: property, design and functionality." Advanced drug delivery reviews 65.10 (2013): 1357-1369*, which is incorporated by reference herein in its entirety. For example, using different linkers will allow one to fine tune the dual-targeting CAR construct. The linker length can vary depending on the antibodies of the dual-targeting CAR construct, their angle of approach to the target epitope, topography of the target on the tumor cell membrane. Referring to FIG. 2, for example, the flexible linkers can include GGGG1, GGGG3, GGGG5, or IgG1 hinge. In some embodiments, the number of Gs in the linker can be 2, 3, 4, 5, 6, or 7 in combination with either S1, S2, S3, S4, S5, or S6. For example, the scFv antibody is specific for CAIX and/or CD70. As shown in FIG. 10 and FIG. 50, for example, the orientations of the scFvs to the linker can vary. In one nucleic acid construct displayed in FIG. 50, the anti-CAIX scFv can be in the first cassette (i.e., before the linker), and the anti-CD70 cassette can be in the second cassette (i.e., after the linker). Alternatively, the anti-CAIX scFv can be in the second cassette, and the anti-CD70 scFv can be in the first cassette. Referring to FIG. 51, for example, anti-CAIX scFv G36 can be in the first cassette, and anti-CD70 B7 can be in the second cassette. Alternatively, anti-CAIX scFv G36 can be in the second cassette, and anti-

CD70 B7 can be in the first cassette. Linkers of various lengths and flexibilities can be utilized as described herein. As shown, different orientations of the two scFvs can influence binding. For example, G36 has higher binding what engineered as the second cassette. On the other hand, as shown in FIG. 11, anti-CD70 scFv B7 does not have significant preference.

[00145] Examples of antibodies useful in constructing the CARs according to the disclosure include those detailed in Table 1, 2, 3 or 4 herein. See also, for example, WO/2007/065027 and WO/2016/100985, the contents of which are hereby incorporated by reference in their entireties.

[00146] The antigen recognition domain useful in constructing the CAR-Ts, for example scFVs directed toward CAIX and/or CD70, can be synthesized, engineered, and/or produced using nucleic acids (e.g., DNA). The DNA encoding the antigen recognition domain can be cloned in frame to DNA encoding necessary CAR-T elements such as, but not limited to, CD8 hinge regions, transmembrane domains, co-stimulatory domains of molecules of immunological interest such as, but not limited to, CD28 and 41BB and CD3-zeta intracellular signaling domains. See FIG. 2, for example.

[00147] Binding domains other than scFv can also be used for predefined targeting of lymphocytes, such as camelid single-domain antibody fragments or receptor ligands, antibody binding domains, antibody hypervariable loops or CDRs as non-limiting examples.

[00148] In an embodiment, the transmembrane domain further comprises a stalk region between said extracellular ligand-binding domain and said transmembrane domain. The term "stalk region" used herein can mean any oligo- or polypeptide that functions to link the transmembrane domain to the extracellular ligand-binding domain. In particular, stalk region are used to provide more flexibility and accessibility for the extracellular ligand-binding domain. A stalk region may comprise up to 300 amino acids, for example, 10 to 100 amino acids or, for example, 25 to 50 amino acids. Stalk region may be derived from all or part of

naturally occurring molecules, such as from all or part of the extracellular region of CD8, CD4 or CD28, or from all or part of an antibody constant region. Alternatively the stalk region may be a synthetic sequence that corresponds to a naturally occurring stalk sequence, or may be an entirely synthetic stalk sequence. In an embodiment said stalk region is a part of human CD8 alpha chain.

[00149] The signal transducing domain or intracellular signaling domain of the CAR of the invention is responsible for intracellular signaling following the binding of extracellular ligand binding domain to the target resulting in the activation of the immune cell and immune response. In other words, the signal transducing domain is responsible for the activation of at least one of the normal effector functions of the immune cell in which the CAR is expressed. For example, the effector function of a T cell can be a cytolytic activity or helper activity including the secretion of cytokines. Thus, the term "signal transducing domain" can refer to the portion of a protein which transduces the effector signal function signal and directs the cell to perform a specialized function.

[00150] Signal transduction domain can comprise two distinct classes of cytoplasmic signaling sequence, those that initiate antigen-dependent primary activation, and those that act in an antigen-independent manner to provide a secondary or co-stimulatory signal. Primary cytoplasmic signaling sequence can comprise signaling motifs which are known as immunoreceptor tyrosine-based activation motifs of ITAMs. ITAMs are well defined signaling motifs found in the intracytoplasmic tail of a variety of receptors that serve as binding sites for syk/zap70 class tyrosine kinases. Examples of ITAM used in the invention can include as non-limiting examples those derived from TCR zeta, FcR gamma, FcR beta, FcR epsilon, CD3 gamma, CD3 delta, CD3 epsilon, CD5, CD22, CD79a, CD79b and CD66d. In another embodiment, the signaling transducing domain of the CAR can comprise the CD3 zeta signaling domain, or the intracytoplasmic domain of the Fc epsilon RI beta or gamma

chains. In another embodiment, the signaling is provided by CD3 zeta together with co-stimulation provided by CD28 and a tumor necrosis factor receptor (TNFr), such as 4-1BB or OX40), for example.

[00151] In an embodiment the intracellular signaling domain of the CAR of the disclosure comprises a co-stimulatory signal molecule. In some embodiments the intracellular signaling domain contains 2, 3, 4 or more co-stimulatory molecules in tandem. A co-stimulatory molecule is a cell surface molecule other than an antigen receptor or their ligands that is required for an efficient immune response.

[00152] "Co-stimulatory ligand" can refer to a molecule on an antigen presenting cell that specifically binds a cognate co-stimulatory molecule on a T-cell, thereby providing a signal which, in addition to the primary signal provided by, for instance, binding of a TCR/CD3 complex with an MHC molecule loaded with peptide, mediates a T cell response, including, but not limited to, proliferation activation, differentiation and the like. A co-stimulatory ligand can include but is not limited to CD7, B7-1 (CD80), B7-2 (CD86), PD-L1, PD-L2, 4-1BBL, OX40L, inducible costimulatory ligand (ICOS-L), intercellular adhesion molecule (ICAM, CD30L, CD40, CD70, CD83, HLA-G, MICA, M1CB, HVEM, lymphotoxin beta receptor, 3/TR6, ILT3, ILT4, an agonist or antibody that binds Toll ligand receptor and a ligand that specifically binds with B7-H3. A co-stimulatory ligand also encompasses, inter alia, an antibody that specifically binds with a co-stimulatory molecule present on a T cell, such as but not limited to, CD27, CD28, 4-1BB, OX40, CD30, CD40, PD-1, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LTGHT, NKG2C, B7-H3, a ligand that specifically binds with CD83.

[00153] A "co-stimulatory molecule" can refer to the cognate binding partner on a T-cell that specifically binds with a co-stimulatory ligand, thereby mediating a co-stimulatory response by the cell, such as, but not limited to proliferation. Co-stimulatory molecules

include, but are not limited to an MHC class 1 molecule, BTLA and Toll ligand receptor. Examples of costimulatory molecules include CD27, CD28, CD8, 4-1BB (CD137), OX40, CD30, CD40, PD-1, ICOS, lymphocyte function-associated antigen-1 (LFA-1), CD2, CD7, LIGHT, NKG2C, B7-H3 and a ligand that specifically binds with CD83 and the like.

[00154] In another particular embodiment, said signal transducing domain is a TNFR-associated Factor 2 (TRAF2) binding motifs, intracytoplasmic tail of costimulatory TNFR member family. Cytoplasmic tail of costimulatory TNFR family member contains TRAF2 binding motifs consisting of the major conserved motif (P/S/A)X(Q/E)E or the minor motif (PXQXXD), wherein X is any amino acid. TRAF proteins are recruited to the intracellular tails of many TNFRs in response to receptor trimerization.

[00155] Chimeric antigen receptors fuse antigen-recognition domains to signaling domains (also referred to as stimulatory domains) that modulate (i.e., stimulate) cell signaling. Non-limiting examples of such stimulatory domains comprise those of CD28, 41BB, and/or CD3-zeta intracellular signaling domains. See FIG. 2, for example.

[00156] The distinguishing features of appropriate transmembrane polypeptides comprise the ability to be expressed at the surface of an immune cell, in particular lymphocyte cells or Natural killer (NK) cells, and to interact together for directing cellular response of immune cell against a predefined target cell. The different transmembrane polypeptides of the CAR of the present invention comprising an extracellular ligand-binding domain and/or a signal transducing domain interact together to take part in signal transduction following the binding with a target ligand and induce an immune response. The transmembrane domain can be derived either from a natural or from a synthetic source. The transmembrane domain can be derived from any membrane-bound or transmembrane protein.

[00157] The term "a part of" used herein can refer to any subset of the molecule, that is a shorter peptide. Alternatively, amino acid sequence functional variants of the polypeptide

can be prepared by mutations in the DNA which encodes the polypeptide. Such variants or functional variants include, for example, deletions from, or insertions or substitutions of, residues within the amino acid sequence. Any combination of deletion, insertion, and substitution may also be made to arrive at the final construct, provided that the final construct possesses the desired activity, especially to exhibit a specific anti-target cellular immune activity. The functionality of the CAR of the invention within a host cell is detectable in an assay suitable for demonstrating the signaling potential of said CAR upon binding of a particular target. Such assays are available to the skilled person in the art. For example, this assay allows the detection of a signaling pathway, triggered upon binding of the target, such as an assay involving measurement of the increase of calcium ion release, intracellular tyrosine phosphorylation, inositol phosphate turnover, or interleukin (IL) 2, interferon .gamma., GM-CSF, IL-3, IL-4 production thus effected.

[00158] Carbonic Anhydrase IX (CAIX)

[00159] A number of mAbs have been identified that react with surface antigens on RCC. These include mAbs that recognize differentiation and overexpressed antigens as well as mAbs that identify RCC-associated antigens not expressed in normal kidney (Michael, 2003; Yang, 2003). The gene for CAIX, also known as G250 and MN is located on chromosomes 9p12 to 13 and encodes a transmembrane protein that binds zinc and has CA activity (Zavada, 1997; Grabmaier, 2000). In HeLa cells derived from human carcinoma of cervix uteri and in RCC cell lines, CAIX/G250/MN/ is found both at the plasma membrane and as a nuclear protein with apparent molecular weights of 58 and 54 kDa. It is N-glycosylated, and in the non-reduced state it forms oligomers (Pastorekova, 1992). Sequence analysis of the predicted CAIX protein shows that it contains a signal peptide (aa 1-37), an extracellular (EC) part (aa 38-414), a hydrophobic transmembrane region of 20 amino acids

(aa 415-434) and a small C-terminus cytoplasmic portion of 25 amino acids (aa 435-459). The human and murine CAIX amino acid sequences are shown in Figure 59. The extracellular portion is composed of two distinct domains. The region between the signal peptide and the CA domain (aa 53-111) shows significant homology (38% identity) with a keratin sulfate attachment domain of a human large aggregating proteoglycan, aggrecan (Doege, 1991). In the PG-like domain of CAIX, a hexapeptide motif with consensus E-E-D-L-P-E (SEQ ID NO: []) is repeated 7 times. The carbonic anhydrase domain is located close to the plasma membrane (aa 135-391). The CAIX antigen appears at malignant transformation and stains positive in about 95% of clear cell RCC specimens as well as in most renal cell metastases.

[00160] Epitopes expressed on the cell surface of tumor cells are superior targets for humoral anti-cancer therapy since, unlike intracellular antigens, they are accessible to circulating antibodies in vivo. Human monoclonal antibodies (mAbs) have become a well-tolerated treatment option in an increasing number of cancers. The concept of selective tumor targeting with antibodies is based on the avid interaction between the antibody and an antigen that is expressed on malignant cells, but not on normal tissues. Many mechanisms have been proposed for the ability of antibodies against tumors to mediate their effects in vivo. For example, engagement of the antibody Fc domain with effector cell FcγRs leads to antibody-dependent cell-mediated cytotoxicity (ADCC). Some (antagonist or inhibitory) antibodies can block the signaling on tumor cells and in this way may act synergistically with immune effector responses by rendering the tumor cells more susceptible to immune effector cell triggered apoptosis or lytic cell death (Baselga, 1998). Another way that antibodies can be utilized is through the construction and functional expression of chimeric-immune receptors or “T-bodies” on T-lymphocytes otherwise known as “designer T-cells”. The antigen binding domain of the chimeric receptor can consist of an antigen-binding domain,

for example, a single-chain antibody (scFv), while the intracellular signaling domain is derived from the cytoplasmic part of a membrane-bound receptor that can induce cellular activation (Maher, 2002; Pinthus, 2003). T-lymphocytes grafted with a chimeric receptor have the combined advantages of MHC-independence and antibody-based antigen binding with efficient T-cell activation upon specific binding to the receptor ligand. This activation results in the production and secretion of cytokines such as IL-2, interferon, GM-CSF and TNF- α . Antigen-specific lysis of tumor cells both in vitro and in vivo have been reported. T-lymphocytes can be permanently grafted with antigen-specific chimeric receptors by retroviral transduction of vector constructs encoding the receptor molecule of choice (reviewed by Rivière, 2004).

[00161] Embodiments of the present invention can comprise an isolated human monoclonal antibody or fragment thereof that immunospecifically binds to a carbonic anhydrase IX (G250) protein. Such antibodies can reduce carbonic anhydrase activity of said protein. For example, such anti-CAIX antibodies can include those described in WO 2007/065027 and U.S. 8,466,263, each of which are incorporated herein by reference in their entireties. See, for example, Figure 58, which provides multiple sequence alignment of amino acid sequences of anti-carbonic anhydrase IX (G250) scFv clones. For example, embodiments of the invention comprise single chain antibodies, such as scFvs G6, G9, G10, G17, G21, G27, G28, G36, G37, G39, G40, G45, G57, G62, G98, G104, G119, or G125 as well as any other scFvs identified according to the methods disclosed herein.

[00162] For example, the amino acids of the CDRs of the anti-CAIX antibodies are provided in Table 1:

CLONE	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
G6 (VH)	TYAMT ([])	AVSGSGGSTYYADSVKG ([])	GPVLRYGFDI ([])
G6 (VL)	TGSRNIGADYDVH ([])	ANNNRPS ([])	QSYDSSLRAWV ([])
G9 (VH)	SYAMS ([])	AISGSGGSTYYADSVKG ([])	SHSSGG FDY ([])
G9 (VL)	TGSSSNIGRGYNVH ([])	GNTNRPS ([])	QSYDSSLRAWV ([])
G10 (VH)	SYAMS ([])	AISANGGTYYADSVKG ([])	NGNYRGAFDI ([])
G10 (VL)	TGSSSNIGAGYDVH ([])	GNSNRPS ([])	QSYDRSLSWV ([])
G17 (VH)	GFTFSSYA ([])	ISGSGGST ([])	ATYGDYGLSDY ([])
G17 (VL)	SSNIGAGYD ([])	ANN ([])	QSYDSSLRAWV ([])
G18 (VH)	SYAMS ([])	AISGSGGSTYYADSVKG ([])	AAAGFDY ([])
G18 (VL)	TGSSSNIGRGYNVH ([])	DD'INRPS ([])	QSYDSSLRAWV ([])
G21 (VH)	SYAMS ([])	AISGSGGSTYYADSVKG ([])	SHSSGGFDY ([])
G21 (VL)	TGSSSNIGRGThTVH ([])	GNTNRPS ([])	QSYDSSLRAWV ([])
G27 (VH)	NYAMT ([])	LISYDGSVTHYTDSVKG ([])	GSGYQE ([])
G27 (VL)	GGNNIGSKSVE ([])	YDSRPS ([])	QVWDSSSDHHVV ([])
G28 (VH)	GFTFSNYA ([])	ISYDGSVT ([])	ARGSGYQEH ([])
G28 (VL)	NIGSKS ([])	YDS ([])	QVWDSSSDHHVV ([])
G36 (VH)	SYAMS ([])	AISANGGTYYADSVKG ([])	NGNYRGAFDI ([])
G36 (VL)	TGSSSNIGAGFDVH ([])	GNTNRPS ([])	QSYDSRLSAWV ([])
G37 (VH)	SYAMS ([])	AISANGGTYYADSVKG ([])	NGNYRGAFDI ([])
G37 (VL)	TGSRNIGADYDVH ([])	ANNNRPS ([])	QSYDSSLRAWV ([])
G39 (VH)	SYAMS ([])	AISGSGGSTYYADSVKG ([])	IGRYSSSLGY ([])
G39 (VL)	TGSSSNIGRGYNVH ([])	DNTNRPS ([])	QSYDSGLRWV ([])
G40 (VH)	SYAMS ([])	AISGSGGSTYYADSVKG ([])	YGDYGLSDY ([])
G40 (VL)	TGSSSNIGAGYDVH ([])	ANNNRPS ([])	QSYDSSLRAWV ([])
G45 (VH)	SYAMS ([])	AISANGGTYYADSVKG ([])	NGNYRGAFDI ([])
G45 (VL)	TGTSSNIGAGYDVH ([])	GNNNRPS ([])	QSYDKSLSWV ([])
G57 (VH)	SYAMS ([])	AISGSGVSTYYADSVKG ([])	YCSSTSCYRGMDV ([])
G57 (VL)	TGSSSNIGAGYDVH ([])	ANNNRPS ([])	QSYDSSLRAWV ([])
G62 (VH)	SYAMS ([])	AISANGGTYYADSVKG ([])	NGNYRGAFDI ([])
G62 (VL)	TGSSSNIGAGYDVH ([])	GNNNRPS ([])	QSYDKSLTWV ([])
G82 (VH)	SYGMH ([])	VISYDGSNKYYADSVKG ([])	GRAARPPFDY ([])
G82 (VL)	SGSSSNIGSNYVY ([])	RNNQRPS ([])	AAWDDSLNGVV ([])
G94 (VH)	SYGMH ([])	VISYDGSNKYYADSVKG ([])	EAPYSSSLDAFDI ([])
G94 (VL)	TGSSSNIGRGYNVH ([])	GNSNRPS ([])	HSRDNNGHHI ([])
G98 (VH)	SYAMS ([])	AISANGGTYYADSVKG ([])	NGNYRGAFDI ([])

G98 (VL)	TGSSSNIGAGYDVH ([])	GNSNRPS ([])	QSYDSSL SAWV ([])
G104 (VH)	IYAMS ([])	AISGSGGGTYHADS VKG ([])	FSAYSGYDL ([])
G104 (VL)	TGSSSNIGRGYNVH ([])	DNTNRPS ([])	QSYDSGLRWV ([])
G106 (VH)	SYAMS ([])	AISANGGTTYADSVKG ([])	NGNYRGAFDI ([])
G106 (VL)	TGSSSNIGAGFDVH ([])	GNNNRPS ([])	QSYDSSL SAWV ([])
G119 (VH)	SYAMS ([])	AISANGGTTYADSVKG ([])	NGNYRGAFDI ([])
G119 (VL)	TGSSSNIGAGYDVH ([])	GNTNRPS ([])	QSYDSTLRVWM ([])
G124 (VH)	KYAMS ([])	GISGSGGSTYYADSVKG ([])	SSRSGYFLP-LDY ([])
G124 (VL)	QGNLSRYYYPS ([])	GKNNRPS ([])	SSRDNTDNRVV ([])
G125 (VH)	SYGMH ([])	AISGSGGSTYYADSVKG ([])	AAVTGGFDP ([])
G125 (VL)	GGDNIGRKSVH ([])	DDRDRPS ([])	QVWDSSSKHYV ([])

[00163] For example, the amino acid sequences of the VH and VL regions of the anti-CAIX antibodies are provided in Table 2:

CLONE	VH/VL	SEQ ID NO:
G6 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASEFTFGTYAMTWVRQAPGKGLEW VSAVSGSGGSTYYADSVKGRFTISRDNRSNTLYLQMNSLRADDTAVYY CAR GPVLRYGFDI WGQGTMTIVSS	([])
G6 (VL)	QSVLTQPPSVSGAPGQRITISCTGSRSNIGADYDVHWYQQLPGTAPKLLI YANNNRPSGVPGRFSASKSGTSASLAISGLQAEDEADYYCQSYDSSLRA WVFGGGTKLAVLG	([])
G9 (VH)	QVQLQESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ARSHSSGGFDYWGQGTMTVTVSS	([])
G9 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGRGYNVHWYQQLPGTAPKLLI YGNTNRPSGVPDRFSGSKSGTSASLAITGLQAEDEGDYYCQSYDSSL SA WVFGGGTKLTVLG	([])
G10 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDI WGQGTMTVTVSS	([])
G10 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLI YGNSNRPSGVPDRFSGSKSGSSASLAITGLQAEDEAHYYCQSYDRSLSW VFGGGTKLTVLG	([])
G17 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ATYGDYGS LDY	([])
G17 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLI YANNNRPSGVP.DRFSGSK.SGTSASLAITGLQAEDEADYYCQSYDSSLR AWV	([])
G18 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ARAAAGFDYWGQGTMTVTVSS	([])

G18 (VL)	QSVLTQPPSVSGAPGQRTISCTGSSSNIGRGYNVHWYQQLPGTAPKLLI YDDINRPSGVPHRFSGSGKSGTSASLAITGLQAEDEADYYCQSYDSSLRA WVFGGGTKLAVLG	()
G21 (VH)	QVQLQESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ARSHSSGGFDYWGGQGLVTVSS	()
G21 (VL)	QSVLTQPPSVSGAPGQRTISCTGSSSNIGRGYNVHWYQQLPGTAPKLLI YGNTNRPSGVPDRFSGSKSGTSASLAITGLQAXDEGDYYCQSYDSSLRA WVFGGGTKLTVLG	()
G27 (VH)	QVTLKESGGGVVQPGTSLRLSCAASGFTFSNYAMTWVRQAPGKGLEW VGLISYDGSVTHYTDSVKGRFTISRDN AKNSLYLQMNTLRADDTAVYY CAR GSGYQEHWGQGLVTVSS	()
G27 (VL)	LPVLTQPPSVSVAPGQTARITCGGNNIGSKSVEWYQQKPGQAPVLVIYY DSDRPSGIPERFSGSNSGNTATLTISRVEAGDEADYYCQVWDSSSDHHV VFGGGTKLTVLG	()
G28 (VH)	QVTLKESGG.GVVQPGTSLRLSCAASGFTFSNYAMTWVRQAPGKGLEW VGLISYDGSVTHYTDSVK.GRFTISRDN AKNSLYLQMNTLRADDTAVYY CARGSGYQEH	()
G28 (VL)	LPVLTQPPS.VSVAPGQTARITCGGNNIGSKSVHWYQQKPGQAPVLVIY YDSDRPSGIP.ERFSGSNSGNTATLTISRVEAGDEADYYCQVWDSSSDHH VV	()
G36 (VH)	EVQLVQSGGGVVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDIWGQGTMTVSS	()
G36 (VL)	QSVLTQPPSVSGAPGQRTISCTGSSSNIGAGFDVHWYQQLPGTAPKLLI YGNTNRPSGVPDRFSGSKSGTSASLAITGLQAEDETDYYCQSYDSRLSA WVFGGGTKLTVLG	()
G37 (VH)	QVQLVQSGGGVVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDI WGQGTMTVSS	()
G37 (VL)	QSVLTQPPSVSGAPGQRTISCTGSRNIGADYDVHWYQQLPGTAPKLLI YANNRPSGVPDRFSGSKSGTSASLAITGLQAEDETDYFCQSYDSSLRA WVFGGGTKVTVLG	()
G39 (VH)	QVQLQESGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC AKIGRYSSSLGYWGQGLVTVSS	()
G39 (VL)	QSVLTQPPSVSGAPGQRTISCTGSSSNIGRGYNVHWYQQLPGTAPKLLI YDNTNRPSGVPARFSGSKSATSASLAITGLQADDEADYYCQSYDSGLR WVFGGGTKLTLLR	()
G40 (VH)	QVQLVQSGGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC ATYGDYGSLDYWGGQGLVTVSS	()
G40 (VL)	QSVLTQPPSVSGAPGQRTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLI YANNRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLRA WVFGGGTKLAVLG	()

G45 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDI WGQGTMTVTVSS	(1)
G45 (VL)	QSVLTQPPSVSGAPGQRITISCTGTSSNIGAGYDVHWYQQLPGAAPRVLI YGNNNRPSGVPDRFSGSKSGTSASLAITGLQSEDEADYYCQSYDKSLS- WVFGGGTKLTVLR	(1)
G57 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFTFS SYAMS WVRQAPGKGLEWVS AISGSGVSTYYADSVKG RFTISRDN SKNTLYLQMNSLRAEDTAVYYCAK YCSSTSCYRGMDV WGKGTLVTVSS	(1)
G57 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLI YANNNRPSGVPDRFSGSKSGTSASLAITGLQAEDEADYYCQSYDSSLRA WVFGGGTKLAVLG	(1)
G62 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDI WGQGTMTVTVSS	(1)
G62 (VL)	QSVLTQPPSVSGAPGQRITISCTGSSSNIGAGYDVHWYQQVPGKAPKVV IYGNNNRPSGVPDRFSGSKSGASASLAITGLQTEDEADYYCQSYDKSLT WVFGGGTKVTVLG	(1)
G82 (VH)	QVQLVQSGGGVQVQGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEW VAVISYDGSNKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CARGRAARPPFDYWGQGTMTVTVSS	(1)
G82 (VL)	QPVLTPPPSASGTPGQRVTISCSGSSSNIGSNYVYVYQQLPGTAPKLPY RNNQRPSGVPDRFSGSKSGTSASLAISGLRSEDEADYYCAAWDDSLNG VWFGGGTKLTVLR	(1)
G94 (VH)	QVQLVQSGGGVQVQGRSLRLSCAASGFTFSSYGMHWVRQAPGKGLEW VAVISYDGSNKYYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CAREAPYSSSLDAFDIWGQGTMTVTVSS	(1)
G94 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGRGYNVHWYQQLPGTAPKLLI YGNSNRPSGVPDRFSGSSGNTASLTITGAQAEDEADYYCHSRDNNGH HIFGGGTCLTVLS	(1)
G98 (VH)	QVQLVQSGGGVQVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMN'SLRAEDTAVYY CANNGNYRGAFDI WGQGTMTVTVSS	(1)
G98 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQHLPGTAPKLLI YGNSNRPSGVPDRFSGSKSGTSASLAITGLQAEDETDYFCQSYDSSLSA WVFGGGTKVTVLG	(1)
G104 (VH)	QVQLQESGGGLVQPGGSLRLSCAASGFTFSIYAMSWVRQAPGKGLEWV SAISGSGGGTYHADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYC AK FSAYSGYDLWGQGTMTVTVSS	(1)
G104 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGRGYNVHWYQQLPGTAPKLLI YDNTNRPSGVPARFSGSKSATSASLTITGLQADDEADYYCQSYDSGLR WVFGGGTKLTLLG	(1)
G106 (VH)	EVQLVQSGGGLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDI WGQGTMTVTVSS	(1)

G106 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGFDVHWYQQLPGTAPRLLI YGNNNRPSGVPDRFSGSKSGTSASLAITGLQAEDETDYFCQSYDSSL SAWVFGGGTKVTVLR	()
G119 (VH)	QVQLVQSGGGLVQPGGSLRLSCAASGFPFSSYAMSWVRQAPGKGLEW VSAISANGGTTYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYY CANNGNYRGAFDIWGQGMVIVSS	()
G119 (VL)	QSVLTQPPSVSGAPGQRVTISCTGSSSNIGAGYDVHWYQQLPGTAPKLLI YGNTNRPSGVPDRFSGSKSGTSASLAIIIGLQADDEADYYCQSYDSTLRV WMFGGGTKLTVLG	()
G124 (VH)	QVQLVQSGGGLVQPGGSLRLSCAAPEFTFSKYAMSWVRQAPGKGLEW VSGISGSGGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYYC AKSSRSGYFLPLDYWGQGLTVTVSS	()
G124 (VL)	SSELTQDPAVSVALGQTVRITCQGNLSRYYYPSWYQKPGQAPVLVIY GKNNRPSGIPDRFSGSSSGNTASLTITGTQAEDEADYYCSSRDNTDNRV VFGGGTKLTVLG	()
G125 (VH)	EVQLVESGGGVVQPGRLRLSCAASGFTFSSYGMHWVRQAPGKGLEW VSAISGSGGSTYYADSVKGRFTISRDNKNTLYLQMNSLRAEDTAVYY CARAAVTGGFDPWGQGLTVTVSS	()
G125 (VL)	QPGLTQPPSVSVAPGQTARITCGGDNIGRKSVDHWYQQRPGQAPILVIR DDRDRPSGIPERFSGSSSVNTATLIISRVEAGDEADYYCQVWDSSSKHY VFGPGTKVTALG	()

[00164] Embodiments can also comprise a consensus sequence of any of the amino acid sequences described herein. For example, when four or more clones have the same amino acid at a given position, that position in the consensus is designated by that amino acid.

[00165] **CD70**

[00166] CD70 is found on the surface of tumors cells of the kidney (clear cell carcinoma and papillary carcinoma, for example), pancreas, larynx or pharynx, melanoma, ovary, lung adenocarcinoma, colon, breast, and brain. See, for example, British Journal of Cancer 103 (2010) 676-684.

[00167] Embodiments of the present invention can comprise an isolated human monoclonal antibody or fragment thereof that immunospecifically binds to CD70. For example, embodiments of the invention comprise single chain antibodies, such as scFvs A20,

B2, B3, B5, B7, B8, or B9 as well as any other scFvs identified according to the methods disclosed herein.

[00168] For example, the amino acids of the CDRs of the anti-CD70 antibodies are provided in Table 3:

CLONE	CDR1 (SEQ ID NO)	CDR2 (SEQ ID NO)	CDR3 (SEQ ID NO)
A20 (VH)	GFTFSSYA([])	ISGSGGSR([])	ARGRGGHGMDV([])
A20 (VL)	SSNIGSNY([])	RNN([])	AAWDDSLNGLV([])
B2 (VH)			
B2 (VL)			
B3 (VH)	GGTFSSQA ([])	IIPFFGVP ([])	AVLKGRGNFDF ([])
B3 (VL)	YSVFHSPNNKNY([])	WAS ([])	QQRSNWPLT ([])
B5 (VH)	GFTVSNYA([])	KSGSDGRT([])	AKGIYDVTGSSFDS([])
B5 (VL)	ALPKKY([])	EDS([])	YSTDSSGNHKV([])
B7 (VH)	GFTVSNYA ([])	KSGSDGRT ([])	AKGIYDVTGSSFDS ([])
B7 (VL)	SGSIASNY ([])	EDN ([])	QSYDSGNRRV ([])
B8 (VH)	GFTVSTSH([])	KDSGGKT([])	ARARSPDPYDGS GFDAFDI([])
B8 (VL)	SNNVGNQG([])	RNN([])	SAWDSSLSAWV([])
B9 (VH)	GFIFSDYY ([])	IRSRGET ([])	ARHRKSFTDLDAFDL ([])
B9 (VL)	QDIGTD ([])	KAS ([])	QHFNNYPAT ([])

[00169] For example, the amino acid sequences of the VH and VL regions of the anti-cd70 antibodies are provided in Table 4:

CLONE	VH/VL	SEQ ID NO:
A20 (VH)	QVQLVQSGG.GLVQPGGSLRLSCAASGFTFSSYAMSWVRQAPGKGLE WVSLISGSGGSRYADSVK.GRFTISRDN SKNTLYLQMNNLRAEDTAV YYCARGRGGHGMDV	([])
A20 (VL)	QPGLTQPPSASGTPGQRVTISCSGSSNIGSNYVYWYQQLPGTAPKLLI YRNNQRPSGVPDRFSGSKSGTSASLAISGLQSEDEADYYCAAWDDSLN GLV	([])
B2 (VH)		
B2 (VL)		
B3 (VH)	QVQLVQSGAEVKKPGSSVKVSCRSSGGTFSSQAFSWVRQAPGQGLEW MGR IIPFFGVPTYAQR FQGRVTITADKSP TAYMELTSLRSDDTAVYYC AVLKGRGNFDF	([])
B3 (VL)	DIVMTQSPDSLAVSLGERATINCKSSYSVFHSPNNKNYLAWYQQRPGQ PPKLLIYWASTRGSGVP.DRFSGSGSGTDFTLTISLEPEDFAVYYCQQR SNWPLT	([])
B5 (VH)	QVQLVQSGGGLVQPRGSLRLSCAASGFTVSNYAMSWVRQAPGKGLE WVATKSGSDGR TYYADSVKGRFTIARDNSKNSLYLQMNSLRAADTA VYYCAKGIYDVTGSSFDS	([])
B5 (VL)	SYELTQPPS.VSVSPGQTARITCSGDALPKKYAYWYQQKSGQAPVLVM FEDSKRPSGIPERFSGSSSGTMTLTISGAQVEDEADYYCYSTDSSGNH KV	([])

B7 (VH)	QVQLVQSGG.GL VQPRGSLRLSCAASGFTVSNYAMSWVRQAPGKGLE WVATKSGSDGRYYADSVKGRFTIARDNSKNSLYLQMNSLRAADTA VYYCAKGIYDVTGSSFDS	(I)
B7 (VL)	NFMLTQPHS.VSESPGKTVTISCTRSSGSIASNYVQWYQQRPGSAPTTVI YEDNQRPSPGVDRFSGSIDSSNSASLTISGLKTEDEADYYCQSYDSGN RRV	(I)
B8 (VH)	EVQLVESGG.GVVQPGRSLRLSCAASGFTVSTSHMSWVRQAPGKGLE WLSGKDSGGKTYADSVR.GRFTIARDDSLNTVFLQMNNMRDEDSGV YYCARARPSDPYDGSFGDAFDI	(I)
B8 (VL)	SYELTQPPS.VSKGLRQTATLTCTGNSNNVGNQGAAWLQQHQGHPPKL LSYRNNNRPSGIS.ERFSASRSGNTASLTITGLQPEDEADYYCSAWDSSL SAWV	(I)
B9 (VH)	QVQLVQSGG.GL VKPRGSLRLSCAASGFIFSDYYMSWIRQAPGKGLQW VASIRSRRGETNYADSVK.GRFTIARDNAEKSLYLQMNSLRAEDAAVY YCARHRKSFTDLDAFDL	(I)
B9 (VL)	DIVMTQSPSTLSASVGDRVTITCRASQDIGTDLSWYQQKPGKAPKLLIY KASSLESGVPSRFSGSGSGTDFLTISLQPDFFATYYCQHFNNYPAT	(I)

[00170] Embodiments can also comprise a consensus sequence of any of the amino acid sequences described herein. For example, when four or more clones have the same amino acid at a given position, that position in the consensus is designated by that amino acid.

[00171] *Cells*

[00172] Embodiments of the disclosure include cells that express a CAR (i.e., CARTS). The cell can be of any kind, including an immune cell that can express the CAR for cancer therapy or a cell, such as a bacterial cell, that harbors an expression vector that encodes the CAR. As used herein, the terms "cell," "cell line," and "cell culture" can be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. For example, all progeny may not be identical due to deliberate or inadvertent mutations. In the context of expressing a heterologous nucleic acid sequence, "host cell" can refer to a eukaryotic cell that is can replicate a vector and/or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors. A host cell can be "transfected" or "transformed," which can refer to a process by

which exogenous nucleic acid is transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny. As used herein, the terms "engineered" and "recombinant" cells or host cells can refer to a cell into which an exogenous nucleic acid sequence, such as, for example, a vector, has been introduced. Therefore, recombinant cells are distinguishable from naturally occurring cells which do not contain a recombinantly introduced nucleic acid. In embodiments of the invention, a host cell is a T cell, including a cytotoxic T cell (also known as TC, Cytotoxic T Lymphocyte, CTL, T-Killer cell, cytolytic T cell, CD8⁺ T-cells or killer T cell); NK cells and NKT cells are also encompassed in the disclosure.

[00173] Some vectors can employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to maintain them and to permit replication of a vector. Also understood and known are techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and their cognate polypeptides, proteins, or peptides.

[00174] The cells can be autologous cells, syngeneic cells, allogenic cells and even in some cases, xenogeneic cells.

[00175] In many situations one may wish to be able to kill the modified CTLs, where one wishes to terminate the treatment, the cells become neoplastic, in research where the absence of the cells after their presence is of interest, or other event. For this purpose one can provide for the expression of certain gene products in which one can kill the modified cells under controlled conditions, such as inducible suicide genes.

[00176] *Armed CARTS*

[00177] The invention further includes CARTs that are modified to secrete one or more polypeptides. Such CARTs can be referred to as CART factories, CAR T cell factories, or armed CARTs. The polypeptide can be, for example, an antibody or fragment thereof as described herein. For example, the polypeptide can be an antibody or cytokine. In embodiments, the antibody is specific for TIGIT, CAIX, GITR, PD-L1, PD-L2, PD-1, CCR4, CTLA-4, VISTA, or CD70. For example, CAR T cell factories can secrete PD-L1 mAbs locally at the tumor site to restore the effective anti-cancer immunity and/or reverse T cell exhaustion. In embodiments, the armed CART secretes IL-12, IL-15, or IL-18.

[00178] For example, a second expression construct, which can be in the same DNA vector as that which encodes the CAR (e.g. the antigen-recognition domain) or in a second separate vector, can be used to encode a mini body (scFv-Fc) or antibody, or a fragment thereof, that is directed against a single or multiple antigens of interest, and can be cloned after an internal ribosomal entry site (IRES). Referring to the figures, the second expression cassette comprises either a fluorescent molecule or an immune-modulating minibody.

[00179] Armed CARTS have the advantage of simultaneously secreting a polypeptide at the targeted site, e.g. tumor site. For example, armed CARTS can secrete anti-TIGIT antibodies or fragments thereof. TIGIT is a T-cell coinhibitory receptor that limits antitumor and other T-cell dependent chronic immune responses, such as CD8⁺ T cell-dependent immune responses. TIGIT is expressed on subsets of activated T cells and natural killer (NK) cells. For example, TIGIT is highly expressed on tumor-infiltrating T-cells. In cancer models, antibody blockade of TIGIT contributed to enhanced CD8⁺ T cell effector function and tumor clearance.

[00180] In embodiments, the anti-TIGIT antibody of the armed CART comprises one or more of the anti-TIGIT antibody clones (or fragments thereof, such as FR1, FR2, FR3,

FR4, CDR1, CDR2, CDR3, or any combinations of the framework and/or CDR regions therein) described in FIG.81.

[00181] For example, the anti-TIGIT antibody can comprise a CDR1 of the VH region of having the sequence: GYTF...TSYG (SEQ ID NO:[]); CDR2 of the VH region having the sequence: ISAY..NGNT (SEQ ID NO:[]); CDR3 of the VH region having the sequence: ARDPGLWFGLTHDYFDY (SEQ ID NO:[]); CDR1 of the VL region having the sequence SSNI....GSNT (SEQ ID NO:[]); CDR2 of the VL region having the sequence: RN.....N (SEQ ID NO:[]); and CDR3 of the VL region having the sequence: AAWDDSRSGPV (SEQ ID NO:[]).

[00182] For example, the anti-TIGIT antibody can comprise a CDR1 of the VH region of having the sequence: GFTF....SDYS (SEQ ID NO:[]); CDR2 of the VH region having the sequence: INSD..GSRT (SEQ ID NO:[]); CDR3 of the VH region having the sequence: ARGPGFFGFDI (SEQ ID NO:[]); CDR1 of the VL region having the sequence RSNI....GRNS (SEQ ID NO:[]); CDR2 of the VL region having the sequence: SN.....N (SEQ ID NO:[]); and CDR3 of the VL region having the sequence: AAWDARLTGPL (SEQ ID NO:[]).

[00183] For example, the anti-TIGIT antibody can comprise a CDR1 of the VH region of having the sequence: GYSF....TNYW (SEQ ID NO:[]); CDR2 of the VH region having the sequence: INPV..NSRT (SEQ ID NO:[]); CDR3 of the VH region having the sequence: ARYYYYAMEV (SEQ ID NO:[]); CDR1 of the VL region having the sequence SSNI....GSNT (SEQ ID NO:[]); CDR2 of the VL region having the sequence: RN.....N (SEQ ID NO:[]); and CDR3 of the VL region having the sequence: EAWDDSLNGPV (SEQ ID NO:[]).

[00184] For example, the anti-TIGIT antibody can comprise a CDR1 of the VH region of having the sequence: GYTF....TNYG (SEQ ID NO:[]); CDR2 of the VH region having

the sequence: VDNN..NGNI (SEQ ID NO:[]); CDR3 of the VH region having the sequence: ARGLFSSRWYLVWFDLP (SEQ ID NO:[]); CDR1 of the VL region having the sequence SSDVG...GYNY (SEQ ID NO:[]); CDR2 of the VL region having the sequence: EV.....T (SEQ ID NO:[]); and CDR3 of the VL region having the sequence: SSYTRSSTSYVV (SEQ ID NO:[]).

[00185] For example, the anti-TIGIT antibody can comprise a CDR1 of the VH region of having the sequence: GGTF....SSYA (SEQ ID NO:[]); CDR2 of the VH region having the sequence: ILPM..FGST (SEQ ID NO:[]); CDR3 of the VH region having the sequence: ARGRDIVAPSN SGFDV (SEQ ID NO:[]); CDR1 of the VL region having the sequence SNNV....GNQG (SEQ ID NO:[]); CDR2 of the VL region having the sequence: RN.....D (SEQ ID NO:[]); and CDR3 of the VL region having the sequence: SAYDRSLNAWV (SEQ ID NO:[]).

[00186] In other embodiments, armed CARTS can secrete anti-PDL1 antibodies or fragments thereof. For example, the armed CARTS can secrete anti-PDL1 antibodies or fragments thereof disclosed in provisional patent application no. 62/624,455, which is incorporated by reference herein in its entirety. Exemplary anti-PDL1 antibodies include antibodies having a VH amino acid sequence having SEQ ID NO: [] and/or a VL amino acid sequence having SEQ ID NO: []. See FIG. 78 and FIG. 81, for example.

[00187] In embodiments, the anti-PDL1 antibodies of the armed CART comprises one or more of the anti-PDL1 antibody clones (or fragments thereof, such as FR1, FR2, FR3, FR4, CDR1, CDR2, CDR3, or any combinations of the framework and/or CDR regions therein) described in FIG. 78 and/or FIG. 81.

[00188] For example, the amino acid sequences of the heavy and light chain complementary determining regions of the PDL-1 antibodies are below:

Sequence ID	V _H CDR1	V _H CDR2	V _H CDR3
42 mut	GGTFSSYA (SEQ ID NO: [])	IIPFGTA (SEQ ID NO: [])	ARGRQMFGAGIDF (SEQ ID NO: [])
50-6B6.1 mut	GYTLSSHG (SEQ ID NO: [])	ISAHNGHA (SEQ ID NO: [])	ARVHAALYYGMDV (SEQ ID NO: [])
50-6B6.2	GYTLSSHG (SEQ ID NO: [])	ISAHNGHA (SEQ ID NO: [])	ARVHAALYYGMDV (SEQ ID NO: [])
50-7B3	GYTLSSHG (SEQ ID NO: [])	ISAHNGHA (SEQ ID NO: [])	ARVHAALYYGMDV (SEQ ID NO: [])
50-5B9	GYTLSSHG (SEQ ID NO: [])	ISAHNGHA (SEQ ID NO: [])	ARVHAALYYGMDV (SEQ ID NO: [])

Sequence ID	V _L CDR1	V _L CDR2	V _L CDR3
42 mut	SGSIDSNY (SEQ ID NO: [])	EDN (SEQ ID NO: [])	QSYDSNNRHVI (SEQ ID NO: [])
50-6B6.1 mut	NIGSKG (SEQ ID NO: [])	DDR (SEQ ID NO: [])	QVWDSGSDHWV (SEQ ID NO: [])
50-6B6.2	NIGDKG (SEQ ID NO: [])	DDS (SEQ ID NO: [])	QVWDSSSDHWV (SEQ ID NO: [])
50-7B3	NIGNKG (SEQ ID NO: [])	DDS (SEQ ID NO: [])	QVWDSSSDHWV (SEQ ID NO: [])
50-5B9	NIGGKG (SEQ ID NO: [])	DDY (SEQ ID NO: [])	QVWDSSSDHWV (SEQ ID NO: [])

[00189] For example, the anti-PDL1 antibodies have a heavy chain with three CDRs including the amino acid sequences SEQ ID NO: [], [], and/or [], respectively and a light chain with three CDRs including the amino acid sequence.

[00190] In other embodiments, armed CARTS can secrete anti-PD1 antibodies or fragments thereof. Exemplary anti-PD1 antibodies include antibodies having a V_H amino acid sequence having SEQ ID NO: [] and a V_L amino acid sequence having SEQ ID NO: []. See FIG. 78 and FIG. 83, for example.

[00191] In embodiments, the anti-PD1 antibodies of the armed CART comprises on or more of the anti-PD1 antibody clones (or fragments thereof, such as FR1, FR2, FR3, FR4, CDR1, CDR2, CDR3, or any combinations of the framework and/or CDR regions therein) described in FIG. 78 and FIG. 83. For example, the anti-PD1 antibodies have a heavy chain with three CDRs including the amino acid sequences SEQ ID NO: [], [], and/or [], respectively and a light chain with three CDRs including the amino acid sequences SEQ ID NO: [], [], and/or [], respectively. See FIG. 78 and FIG. 83, for example.

[00192] In embodiments, the amino acid sequences of the heavy and light chain complementary determining regions of the PD-1 antibodies are shown below:

Heavy chain (V_H) complementary determining regions (CDRs) of the PD-1 antibodies

Sequence ID	V _H CDR1	V _H CDR2	V _H CDR3
P4-B3	GFTFDDYA (SEQ ID NO: [])	ISWNSGSI (SEQ ID NO: [])	ASDYGDKYYYYGMDV (SEQ ID NO: [])
P4-B7	GYTFTTYW (SEQ ID NO: [])	IYPDDSDT (SEQ ID NO: [])	AFWGASGAPVNGFDI (SEQ ID NO: [])
PD1#2	GDSVSSDNYF (SEQ ID NO: [])	VYYNGNT (SEQ ID NO: [])	ATETPPTSYPFNSGPFDS (SEQ ID NO: [])
PD1#3	GYTFNRFG (SEQ ID NO: [])	TNPYNGNT (SEQ ID NO: [])	ARVVAVNGMDV (SEQ ID NO: [])
PD1#13	GFTFSSYA (SEQ ID NO: [])	ISYDGSNK (SEQ ID NO: [])	ASQTVAGSDY (SEQ ID NO: [])
HL-7	GFTFDDYA (SEQ ID NO: [])	ISWNSGSI (SEQ ID NO: [])	ASDYGDKYYYYGMDV (SEQ ID NO: [])
HL-14	GFTFDDYA (SEQ ID NO: [])	ISWNSGSI (SEQ ID NO: [])	ASDYGDKY\$YYGMDV (SEQ ID NO: [])
HLkin-1	GFTFDDFA (SEQ ID NO: [])	ISWNSGSI (SEQ ID NO: [])	ASDYGDKYYYYGMDV (SEQ ID NO: [])
HLkin-1 HL-7 mut2	GFTFDDFA (SEQ ID NO: [])	ISWNSGSI (SEQ ID NO: [])	ASDYGDKYYYYGMDV (SEQ ID NO: [])
HLkin-1 HL-7 HL-14 mut3	GFTFDDFA (SEQ ID NO: [])	ISWNSGSI (SEQ ID NO: [])	ASDYGDKY\$YYGMDV (SEQ ID NO: [])

Light chain (V_L) complementary determining regions (CDRs) of the PD-1 antibodies

Sequence ID	V _L CDR1	V _L CDR2	V _L CDR3
P4-B3	SSNIGSNT (SEQ ID NO: [])	NDN (SEQ ID NO: [])	AAWDGGLNGRGV (SEQ ID NO: [])
P4-B7	SSNIGAGYV (SEQ ID NO: [])	SNN (SEQ ID NO: [])	AAWDDSLNAPV (SEQ ID NO: [])
PD1#2	SNNVGAHG (SEQ ID NO: [])	RNN (SEQ ID NO: [])	SSWDSSLSGYV (SEQ ID NO: [])
PD1#3	SGSIAAYY (SEQ ID NO: [])	EDN (SEQ ID NO: [])	QSYDSSNLWV (SEQ ID NO: [])
PD1#13	NIGSKS (SEQ ID NO: [])	DDS (SEQ ID NO: [])	QVWHSVSDQGV (SEQ ID NO: [])
HL-7	SSNIGSNT (SEQ ID NO: [])	NDN (SEQ ID NO: [])	AAWDGGLNGRGV (SEQ ID NO: [])
HL-14	SSNIGSNT (SEQ ID NO: [])	NDN (SEQ ID NO: [])	AAWDGGLNGRGV (SEQ ID NO: [])
HLkin-1	SSNIGSNT (SEQ ID NO: [])	NDN (SEQ ID NO: [])	AAWDGGLNGRGV (SEQ ID NO: [])
HLkin-1 HL-7 mut2	SSNIGSNT (SEQ ID NO: [])	NDN (SEQ ID NO: [])	AAWDGGLNGRGV (SEQ ID NO: [])
HLkin-1 HL-7 HL-14 mut3	SSNIGSNT (SEQ ID NO: [])	NDN (SEQ ID NO: [])	AAWDGGLNGRGV (SEQ ID NO: [])

[00193] In other embodiments, armed CARTS can secrete anti-CCR4 antibodies or fragments thereof. For example, armed CARTS can secrete anti-CCR4 antibodies or fragments as described in WO2009/086514, WO 2013/166500, PCT/US2015/054202, or PCT/US2016/026232.

[00194] For example, the anti-CCR4 protein antibody or fragment thereof of the armed CART can comprise an antibody having a VH CDR1 region having the amino acid sequence GYTFASY (SEQ ID NO: []); a VH CDR2 region having the amino acid sequence WINPGNVNTKYNEKFKG (SEQ ID NO: []); a VH CDR3 region having the amino acid sequence STYYRPLDY (SEQ ID NO: []); and/or VL CDR1 region having amino acid sequence KSSQSILYSSNQKNYLA (SEQ ID NO: []); a VL CDR2 region having the amino

acid sequence WASTRES (SEQ ID NO: []) and/or a VL CDR3 region having the amino acid sequence HQYLSSYT (SEQ ID NO: []).

[00195] For example, the anti-CCR4 protein or fragment thereof of the armed CAR can comprise an antibody having a VH amino acid sequence

QVQLVQSGAEVKKPGASVKVSCKASGYTF

ASYMHWMRQAPGQGLEWIGWINPGNVNTKYNEKFKGRATLTVDSTNTAYMEL

SSLRSEDVAVYYCARSTYYRPLDYWGQGTLVTVSS (SEQ ID NO: []) and/or a VL amino acid sequence

DIVMTQSPDSLAVSLGERATINCKSSQSILYSSNQKNYLAWYQQKPGQSPKLLIYWA

STRESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCHQYLSSYTFGQGTKLEIK

(SEQ ID NO: []).

[00196] For example, the anti-CCR4 protein antibody or fragment thereof of the armed CART can comprise an antibody having a:

[00197] VH chain of antibody 1-44: (SEQ ID NO: [])

QVQLVQSGAEVKKPGASVKVSCKASGYTFASQWMHWMRQAPGQGLEWIGWINPG

NVNTKYNEKFKGRATLTVDSTNTAYMELSSLRSEDVAVYYCARSTWYRPLDYWG

QGTLVTVSS

[00198] VL chain of antibody 1-44 (SEQ ID NO: [])

DIVMTQSPDSLAVSLGERATINCKSSQSILYSSNQKNYLAWYQQKPGQSPKLLIYWA

STR ESGVPDRFSGSGSGTDFTLTISLQAEDVAVYYCHQYISSYTFGQGTKLEIK

[00199] VH chain of antibody 1-49 (SEQ ID NO: [])

QVQLVQSGAEVKKPGASVKVSCKASGYTFASSWMHWMRQAPGQGLEWIGWINPG

NVNTKYNEKFKGRATLTVDSTNTAYMELSSLRSEDVAVYYCARSTWYRPNDYWG

QGTLVTVSS

[00200] VL chain of antibody 1-49 (SEQ ID NO: [])

DIVMTQSPDSLAVSLGERATINCKSSQSILYSSNQKNYLAWYQQKPGQSPKLLIYW
ASTRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCHQYKSSYTFGQGGTKLEIK

[00201] VH chain of antibody 2-1 (SEQ ID NO: [])

QVQLVQSGAEVKKPGASVKVSCKASGYTFASSWMHWMRQAPGQGLEWIGWINPG
NVNTKYNEKFKGRATLTVDSTNTAYMELSSLRSEDTAVYYCARTTRYRPLDYWG
QGTLVTVSS

[00202] VL chain of antibody 2-1 (SEQ ID NO: [])

DIVMTQSPDSLAVSLGERATINCKSSQSILYSSNQKNYLAWYQQKPGQSPKLLIYWA
ST RESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCHQYRSSYTFGQGGTKLEIK

[00203] VH chain of antibody 2-2 (SEQ ID NO: [])

QVQLVQSGAEVKKPGASVKVSCKASGYTFASQYMHWMRQAPGQGLEWIGWINPG
NVNTKYNEKFKGRATLTVDSTNTAYMELSSLRSEDTAVYYCARLTYYRPPDYWG
QGTLVTVSS

[00204] VL chain of antibody 2-2 (SEQ ID NO: [])

DIVMTQSPDSLAVSLGERATINCKSSQSILYSSNQKNYLAWYQQKPGQSPKLLIYWA
ST RESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCHQYYSSYTFGQGGTKLEIK

[00205] VH chain of antibody 2-3 (SEQ ID NO: [])

QVQLVQSGAEVKKPGASVKVSCKASGYTFASAWMHWMRQAPGQGLEWIGWINPG
NVNTKYNEKFKGRATLTVDSTNTAYMELSSLRSEDTAVYYCARSTYYRPLDYWG
QGTLV TVSS

[00206] VL chain of antibody 2-3 (SEQ ID NO: [])

DIVMTQSPDSLAVSLGERATINCKSSQSILYSSNQKNYLAWYQQKPGQSPKLLIYWA
STRESGVPDRFSGSGSGTDFTLTISSLQAEDVAVYYCHQYMSSYTFGQGGTKLEIK

[00207] The amino acid sequences of the heavy and light chain complementary determining regions of the anti-CCR4 antibodies are shown in the Table below.

Antibody	Variable region	CDR1	CDR2	CDR3
Mouse 1567	VH	GYTFASY {SEQ ID NO: 21}	INPGNVNT {SEQ ID NO: 27}	STYYRPLDY {SEQ ID NO: 29}
Humanized 1567	VH	GYTFASY {SEQ ID NO: 21}	INPGNVNT {SEQ ID NO: 27}	STYYRPLDY {SEQ ID NO: 29}
Ab1-44	VH	GYTFASQW {SEQ ID NO: 22}	INPGNVNT {SEQ ID NO: 27}	STWYRPLDY {SEQ ID NO: 30}
Ab1-49	VH	GYTFASSW {SEQ ID NO: 23}	INPGNVNT {SEQ ID NO: 27}	STWYRPNDY {SEQ ID NO: 31}
Ab2-1	VH	GYTFASSW {SEQ ID NO: 23}	INPGNVNT {SEQ ID NO: 27}	TTRYRPLDY {SEQ ID NO: 32}
Ab2-2	VH	GYTFASQY {SEQ ID NO: 24}	INPGNVNT {SEQ ID NO: 27}	LTYYRPPDY {SEQ ID NO: 33}
Ab2-3	VH	GYTFASAW {SEQ ID NO: 25}	INPGNVNT {SEQ ID NO: 27}	STYYRPLDY {SEQ ID NO: 29}
Mouse 1567	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYLSSYT {SEQ ID NO: 34}
Humanized 1567	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYLSSYT {SEQ ID NO: 34}
Ab1-44	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYISSYT {SEQ ID NO: 35}
Ab1-49	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYKSSYT {SEQ ID NO: 36}
Ab2-1	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYRSSYT {SEQ ID NO: 37}
Ab2-2	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYYSST {SEQ ID NO: 38}
Ab2-3	VL	QSILYSSNQKNY {SEQ ID NO: 26}	WASTRE {SEQ ID NO: 28}	HQYMSSYT {SEQ ID NO: 39}

[00208] Armed CART can be constructed by including a nucleic acid encoding the polypeptide of interest after the intracellular signaling domain. In embodiments, there is an internal ribosome entry site, (IRES), positioned between the intracellular signaling domain and the polypeptide of interest. One skilled in the art can appreciate that more than one polypeptide can be expressed by employing multiple IRES sequences in tandem.

[00209] In one embodiment, the methods and compositions presented herein provide a target-specific T cell, such as a T cell with specificity for CAIX and/or CD70, armed with the

power to secrete polypeptides in the tumor microenvironment, for example to combat T cell exhaustion. For example, myeloid-derived suppressor cells (MDSCs) are a heterogeneous population of early myeloid progenitors, immature granulocytes, macrophages, and dendritic cells at different stages of differentiation that comprise the tumor microenvironment.

MDSCs are induced by pro-inflammatory cytokines and are found in increased numbers in infectious and inflammatory disease conditions. Without wishing to be bound by theory, their presence in the tumor microenvironment indicates a causative role in promoting tumor-associated immune suppression. Human MDSCs express Siglec-3/CD33 (GENBANK Accession No. NM_001772.4 (for nucleotide sequence) and NP_001763.3 (for amino acid sequence) and have heterogeneous expression of CD14 (GENBANK Accession No. NM_000591.4 (for nucleotide sequence) and NP_000582.1 (for amino acid sequence)) and CD15 (GENBANK Accession No. NM_002033.3 (for nucleotide sequence) and NP_002024.1 (for amino acid sequence)) which show that multiple subsets exist. Other MDSC markers useful for identifying MDSCs include, but are not limited to B7-1/CD8 (NM_001145873.1 (for nucleotide sequence), NP_001139345.1 (for amino acid sequence)), CCR2(NM_001123041.2 (for nucleotide sequence), NP_001116513.2 (for amino acid sequence)), CD1d (NM_001766.3 (for nucleotide sequence), NP_001757.1 (for amino acid sequence)), CD2 (NM_001328609.2 (for nucleotide sequence), NP_001315538.1 (for amino acid sequence)), CD31/PECAM-1 (NM_000442.5 (for nucleotide sequence), NP_000433.4 (for amino acid sequence)), CD43 (NM_001030288.3 (for nucleotide sequence), NP_001025459.1 (for amino acid sequence)), CD44(NM_000610.4 (for nucleotide sequence), NP_000601.3 (for amino acid sequence)), gp130 (NM_002184.4 (for nucleotide sequence), NP_002175.2 ((for amino acid sequence)), PD-L1 (NM_014143.4 (for nucleotide sequence), NP_054862.1 (for amino acid sequence)), and CD162 (NM_001206609.2 (for nucleotide sequence), NP_001193538.1 (for amino acid sequence)).

[00210] In one embodiment, the methods and compositions presented herein provide a target-specific T cell, such as a T cell with specificity for CAIX and/or CD70, armed with the power to secrete polypeptides that target MDSCs in the tumor microenvironment. For example the secreted polypeptides can target one or more MDSC markers (e.g., CD33, CD14, and/or CD15, or other MDSC markers listed herein).

[00211] **Introduction of Constructs into CTLs**

[00212] Expression vectors that encode the CARs can be introduced as one or more DNA molecules or constructs, where there may be at least one marker that will allow for selection of host cells that contain the construct(s).

[00213] The constructs can be prepared in conventional ways, where the genes and regulatory regions can be isolated, as appropriate, ligated, cloned in an appropriate cloning host, analyzed by restriction or sequencing, or other convenient means. Particularly, using PCR, individual fragments including all or portions of a functional unit can be isolated, where one or more mutations may be introduced using "primer repair", ligation, in vitro mutagenesis, etc., as appropriate. The construct(s) once completed and demonstrated to have the appropriate sequences may then be introduced into the CTL by any convenient means. The constructs may be integrated and packaged into non-replicating, defective viral genomes like Adenovirus, Adeno-associated virus (AAV), or Herpes simplex virus (HSV) or others, including retroviral vectors or lentiviral vectors, for infection or transduction into cells. The constructs may include viral sequences for transfection. Alternatively, the construct may be introduced by fusion, electroporation, biolistics, transfection, lipofection, or the like. The host cells can be grown and expanded in culture before introduction of the construct(s), followed by the appropriate treatment for introduction of the construct(s) and integration of the construct(s). The cells are then expanded and screened by virtue of a marker present in the

construct. Various markers that may be used successfully include hppt, neomycin resistance, thymidine kinase, hygromycin resistance, etc.

[00214] In some instances, one can have a target site for homologous recombination, where a construct be integrated at a particular locus. For example,) can knock-out an endogenous gene and replace it (at the same locus or elsewhere) with the gene encoded for by the construct using materials and methods as are known in the art for homologous recombination. For homologous recombination, one may use either .OMEGA. or O-vectors. See, for example, Thomas and Capecchi, Cell (1987) 51, 503-512; Mansour, et al., Nature (1988) 336, 348-352; and Joyner, et al., Nature (1989) 338, 153-156.

[00215] The constructs can be introduced as a single DNA molecule encoding at least the CAR and optionally another gene, or different DNA molecules having one or more genes. Other genes include genes that encode therapeutic molecules or suicide genes, for example. The constructs may be introduced simultaneously or consecutively, each with the same or different markers.

[00216] Vectors containing useful elements such as bacterial or yeast origins of replication, selectable and/or amplifiable markers, promoter/enhancer elements for expression in prokaryotes or eukaryotes, etc. that may be used to prepare stocks of construct DNAs and for carrying out transfections are well known in the art, and many are commercially available.

[00217] Methods of Use

[00218] Aspects of the disclosure are directed towards methods of treating a subject afflicted with a cancer.

[00219] For example, aspects of the disclosure are directed towards methods of killing a cancer cell, such as a renal cancer cell. Referring to FIG. 12, for example, B7-GGGGS-G36 CAR T cells had more killing activity on the targeted renal carcinoma cells (CAIX+ CD70+)

than non-targeted cells (CAIX+ CD70-, CAIX- CD70+, CAIX- CD70-). Further, referring to FIG. 57, for example, bispecific split CAR achieved superior killing when compared to mono CAR or bispecific CARs.

[00220] Aspects of the disclosure are further directed towards methods of stopping or reducing progression or promoting regression of a cancer in a subject.

[00221] Still further, aspects of the disclosure are directed towards a method of reducing cellular proliferation of a cancer cell in a subject. See, for example, FIG. 80.

[00222] "Cancer" and "cancerous" can refer to or describe, for example, the physiological condition in mammals that is characterized by unregulated cell growth. Examples of cancer include but are not limited to, carcinoma, lymphoma, blastoma, sarcoma, and leukemia. More particular examples of such cancers include squamous cell cancer, smallcell lung cancer, non-small cell lung cancer, adenocarcinoma of the lung, squamous carcinoma of the lung, cancer of the peritoneum, hepatocellular cancer, gastrointestinal cancer, pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer, bladder cancer, hepatoma, breast cancer, colon cancer, colorectal cancer, endometrial or uterine carcinoma, salivary gland carcinoma, kidney cancer, liver cancer, prostate cancer, vulval cancer, thyroid cancer, hepatic carcinoma and various types of head and neck cancer. For example, the cancer is renal cell carcinoma, such as ccRCC.

[00223] In cancer, the normal intercellular interactions in tissues are disrupted, and the tumor microenvironment evolves to accommodate the growing tumor. The tumor microenvironment (TME) can refer to the cellular environment in which a tumor exists, including components such as surrounding blood vessels, immune cells, fibroblasts, bone marrow-derived inflammatory cells, lymphocytes, signaling molecules and the extracellular matrix (ECM). Tumor microenvironment is complex and is heavily influenced by immune system.

[00224] This invention provides CAR-T cell therapy for renal cell carcinoma, among others (such as those described herein). The secretion of a mono, bi-, or tri-specific minibody, antibody or minibody/antibody fusion protein by the CAR-T cell at the tumor site could provide additional benefit by altering (i.e., modulating) the immune-repressive tumor microenvironment.

[00225] In embodiments, the method comprises administering to a subject afflicted with a cancer a therapeutically effective amount of an engineered cell as described herein. Therapeutically effective amounts can depend on the severity and course of the cancer, previous therapy, the subject's health status, weight, and response to the drugs, and the judgment of the treating physician.

[00226] The subject can be afflicted with cancer such as liquid cancers (i.e., blood cancers) and/or solid cancers (i.e., tumors). The cancer can be benign or malignant, and can be one that is influenced by the immune system.

[00227] Embodiments as described herein can modulate the immune system so as to treat the subject afflicted with cancer. "Modulating" can refer to up-regulation, induction, stimulation, potentiation, and/or relief of inhibition, as well as inhibition, attenuation and/or down-regulation or suppression. In embodiments, the activity of the subject's immune system is modulated, the microenvironment surround the cancer cell and/or tumor is modulated, or both. For example, embodiments as described herein can alter the immune-repressive tumor microenvironment, reducing the microenvironment-dependent immune suppression, so as to modulate (or allow) the immune system to kill tumor cells.

[00228] One embodiment is directed towards methods of treating a subject afflicted with renal cell carcinoma. Immune therapies, such as those described herein, offer an exciting therapeutic option for RCC. For example, embodiments comprise engineering a chimeric-antigen receptor (CAR) T-cell for RCC.

[00229] An "individual" or "subject" can be a mammal. Mammals include, but are not limited to, domesticated animals (e.g., cows, sheep, cats, dogs, and horses), primates (e.g., humans and non-human primates such as monkeys), rabbits, and rodents (e.g., mice and rats). In certain embodiments, the individual or subject is a human.

[00230] The cells according to the disclosure can be used for treating cancer in a patient in need thereof. In another embodiment, said isolated cell according to the invention can be used in the manufacture of a medicament for treatment of a cancer, viral infections of autoimmune disorders, in a patient in need thereof.

[00231] The present disclosure can rely on methods for treating patients in need thereof, said method comprising at least one of the following steps: (a) providing a chimeric antigen receptor cells according to the invention and (b) administering the cells to said patient.

[00232] Said treatment can be ameliorating, curative or prophylactic. It can be either part of an autologous immunotherapy or part of an allogenic immunotherapy treatment. By autologous, it is meant that cells, cell line or population of cells used for treating patients are originating from said patient or from a Human Leucocyte Antigen (HLA) compatible donor. By allogeneic is meant that the cells or population of cells used for treating patients are not originating from said patient but from a donor.

[00233] The invention is particularly suited for allogenic immunotherapy, insofar as it allows the transformation of T-cells, obtained from donors, into non-alloreactive cells. This may be done under standard protocols and reproduced as many times as needed. The resulted modified T cells may be pooled and administrated to one or several patients, being made available as an "off the shelf" therapeutic product.

[00234] Cells that can be used with the disclosed methods are described in the previous section. Said treatment can be used to treat patients diagnosed with cancer. Cancers that may

be treated include tumors that are not vascularized, or not yet substantially vascularized, as well as vascularized tumors. The cancers may comprise nonsolid tumors (such as hematological tumors, for example, leukemias and lymphomas) or may comprise solid tumors. Types of cancers to be treated with the CARs of the invention include, but are not limited to, carcinoma, blastoma, and sarcoma, and certain leukemia or lymphoid malignancies, benign and malignant tumors, and malignancies e.g., sarcomas, carcinomas, and melanomas. Adult tumors/cancers and pediatric tumors/cancers are also included.

[00235] It can be a treatment in combination with one or more therapies against cancer selected from the group of antibodies therapy, chemotherapy, cytokines therapy, dendritic cell therapy, gene therapy, hormone therapy, laser light therapy and radiation therapy.

[00236] According to an embodiment of the invention, said treatment can be administrated into patients undergoing an immunosuppressive treatment. The present invention uses cells or population of cells, which have been made resistant to at least one immunosuppressive agent due to the inactivation of a gene encoding a receptor for such immunosuppressive agent. In this aspect, the immunosuppressive treatment should help the selection and expansion of the T-cells according to the invention within the patient.

[00237] In a further embodiment, the cell compositions of the present invention are administered to a patient in conjunction with (e.g., before, simultaneously or following) bone marrow transplantation, T cell ablative therapy using either chemotherapy agents such as, fludarabine, external-beam radiation therapy (XRT), cyclophosphamide, or antibodies such as OKT3 or CAM PATH. In another embodiment, the cell compositions of the present invention are administered following B-cell ablative therapy such as agents that react with CD20, e.g., Rituxan. For example, in one embodiment, subjects may undergo standard treatment with high dose chemotherapy followed by peripheral blood stem cell transplantation. In certain embodiments, following the transplant, subjects receive an infusion

of the expanded immune cells of the present invention. In an additional embodiment, expanded cells are administered before or following surgery. Said modified cells obtained by any one of the methods described here can be used in a particular aspect of the invention for treating patients in need thereof against Host versus Graft (HvG) rejection and Graft versus Host Disease (GvHD); therefore in the scope of the present invention is a method of treating patients in need thereof against Host versus Graft (HvG) rejection and Graft versus Host Disease (GvHD) comprising treating said patient by administering to said patient an effective amount of modified cells comprising inactivated TCR alpha and/or TCR beta genes.

[00238] *Administration of Cells*

[00239] The disclosure is particularly suited for allogenic immunotherapy, insofar as it allows the transformation of T-cells, obtained from donors, into non-alloreactive cells. This can be done under standard protocols and reproduced as many times as needed. The resulted modified T cells may be pooled and administrated to one or several patients, being made available as an "off the shelf" therapeutic product.

[00240] Depending upon the nature of the cells, the cells can be introduced into a host organism, e.g. a mammal, in a wide variety of ways. The cells can be introduced at the site of the tumor, in specific embodiments, although in alternative embodiments the cells hone to the cancer or are modified to hone to the cancer. The number of cells that are employed will depend upon a number of circumstances, the purpose for the introduction, the lifetime of the cells, the protocol to be used, for example, the number of administrations, the ability of the cells to multiply, the stability of the recombinant construct, and the like. The cells may be applied as a dispersion, for example, being injected at or near the site of interest. The cells may be in a physiologically-acceptable medium.

[00241] In some embodiments, the cells are encapsulated to inhibit immune recognition and placed at the site of the tumor.

[00242] The cells can be administered as desired. Depending upon the response desired, the manner of administration, the life of the cells, the number of cells present, various protocols may be employed. The number of administrations will depend upon the factors described above at least in part.

[00243] The administration of the cells or population of cells according to the present invention can be carried out in any convenient manner, including by aerosol inhalation, injection, ingestion, transfusion, implantation or transplantation. The compositions described herein may be administered to a patient subcutaneously, intradermally, intratumorally, intranodally, intramedullary, intramuscularly, by intravenous or intralymphatic injection, or intraperitoneally. In one embodiment, the cell compositions of the present invention are administered by intravenous injection.

[00244] The administration of the cells or population of cells can consist of the administration of 10^4 - 10^9 cells per kg body weight, for example, 10^5 to 10^6 cells/kg body weight including all integer values of cell numbers within those ranges. The cells or population of cells can be administered in one or more doses. In another embodiment, said effective amount of cells are administered as a single dose. In another embodiment, said effective amount of cells are administered as more than one dose over a period time. Timing of administration is within the judgment of managing physician and depends on the clinical condition of the patient. The cells or population of cells may be obtained from any source, such as a blood bank or a donor. While individual needs vary, determination of optimal ranges of effective amounts of a given cell type for a particular disease or conditions within the skill of the art. An effective amount means an amount which provides a therapeutic or prophylactic benefit. The dosage administered will be dependent upon the age, health and

weight of the recipient, kind of concurrent treatment, if any, frequency of treatment and the nature of the effect desired.

[00245] It should be appreciated that the system is subject to many variables, such as the cellular response to the ligand, the efficiency of expression and, as appropriate, the level of secretion, the activity of the expression product, the particular need of the patient, which may vary with time and circumstances, the rate of loss of the cellular activity as a result of loss of cells or expression activity of individual cells, and the like. Therefore, for each individual patient, even if there were universal cells which could be administered to the population at large, each patient would be monitored for the proper dosage for the individual, and such practices of monitoring a patient are routine in the art.

[00246] *Nucleic Acid-Based Expression Systems*

[00247] The CARs of the disclosure can be expressed from an expression vector. Recombinant techniques to generate such expression vectors are well known in the art.

[00248] DNA constructs, which can also be referred to as "DNA vectors", as described herein, can be cloned into a vector which will be used to transduce and produce chimeric-antigen receptor T-cells that secrete polypeptides and/or fragments thereof. For example, DNA constructs can be cloned into a lentiviral vector for production of lentivirus, which will be used to transduce and produce chimeric-antigen receptor T-cells that secrete a mono, bi- or tri-specific immune-modulating antibody/minibody and/or antibody-fusion protein at the tumor site.

[00249] *Vectors*

[00250] The term "vector" can refer to a carrier nucleic acid molecule into which a nucleic acid sequence can be inserted for introduction into a cell where it can be replicated. A

nucleic acid sequence can be "exogenous," which means that it is foreign to the cell into which the vector is being introduced or that the sequence is homologous to a sequence in the cell but in a position within the host cell nucleic acid in which the sequence is ordinarily not found. Vectors include plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (see, for example, Maniatis et al., 1988 and Ausubel et al., 1994, both incorporated herein by reference).

[00251] The term "expression vector" refers to any type of genetic construct comprising a nucleic acid coding for a RNA that can be transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. In other cases, these sequences are not translated, for example, in the production of antisense molecules or ribozymes. Expression vectors can contain a variety of "control sequences," which refer to nucleic acid sequences necessary for the transcription and translation of an operably linked coding sequence in a particular host cell. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described infra.

[00252] *Promoters and Enhancers*

[00253] A "promoter" is a control sequence that is a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It can contain genetic elements at which regulatory proteins and molecules may bind, such as RNA polymerase and other transcription factors, to initiate the specific transcription a nucleic acid sequence. The phrases "operatively positioned," "operatively linked," "under control," and "under transcriptional control" mean that a promoter is in a correct functional location and/or orientation in relation

to a nucleic acid sequence to control transcriptional initiation and/or expression of that sequence.

[00254] A promoter comprises a sequence that functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as, for example, the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation. Additional promoter elements regulate the frequency of transcriptional initiation. These are located in the region 30 110 bp upstream of the start site, although a number of promoters have been shown to contain functional elements downstream of the start site as well. To bring a coding sequence "under the control of" a promoter, one positions the 5' end of the transcription initiation site of the transcriptional reading frame "downstream" of (i.e., 3' of) the chosen promoter. The "upstream" promoter stimulates transcription of the DNA and promotes expression of the encoded RNA.

[00255] The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either cooperatively or independently to activate transcription. A promoter may or may not be used in conjunction with an "enhancer," which refers to a cis-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[00256] A promoter can be one naturally associated with a nucleic acid sequence, as may be obtained by isolating the 5 prime' non-coding sequences located upstream of the coding segment and/or exon. Such a promoter can be referred to as "endogenous." Similarly,

an enhancer may be one naturally associated with a nucleic acid sequence, located either downstream or upstream of that sequence. Alternatively, certain advantages will be gained by positioning the coding nucleic acid segment under the control of a recombinant or heterologous promoter, which refers to a promoter that is not normally associated with a nucleic acid sequence in its natural environment. A recombinant or heterologous enhancer refers also to an enhancer not normally associated with a nucleic acid sequence in its natural environment. Such promoters or enhancers may include promoters or enhancers of other genes, and promoters or enhancers isolated from any other virus, or prokaryotic or eukaryotic cell, and promoters or enhancers not "naturally occurring," i.e., containing different elements of different transcriptional regulatory regions, and/or mutations that alter expression. For example, promoters that are most commonly used in recombinant DNA construction include the lactamase (penicillinase), lactose and tryptophan (trp) promoter systems. In addition to producing nucleic acid sequences of promoters and enhancers synthetically, sequences may be produced using recombinant cloning and/or nucleic acid amplification technology, including PCR.TM., in connection with the compositions disclosed herein (see U.S. Pat. Nos. 4,683,202 and 5,928,906, each incorporated herein by reference). Furthermore, the control sequences that direct transcription and/or expression of sequences within non-nuclear organelles such as mitochondria, chloroplasts, and the like, can be employed as well.

[00257] Naturally, it will be important to employ a promoter and/or enhancer that effectively directs the expression of the DNA segment in the organelle, cell type, tissue, organ, or organism chosen for expression. Those of skill in the art of molecular biology generally know the use of promoters, enhancers, and cell type combinations for protein expression, (see, for example Sambrook et al. 1989, incorporated herein by reference). The promoters employed may be constitutive, tissue-specific, inducible, and/or useful under the appropriate conditions to direct high level expression of the introduced DNA segment, such

as is advantageous in the large-scale production of recombinant proteins and/or peptides. The promoter may be heterologous or endogenous.

[00258] Additionally any promoter/enhancer combination could also be used to drive expression. Use of a T3, T7 or SP6 cytoplasmic expression system is another embodiment. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

[00259] The identity of tissue-specific promoters or elements, as well as assays to characterize their activity, is well known to those of skill in the art.

[00260] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily determine this and providing the necessary signals

[00261] In certain embodiments of the disclosure, the use of internal ribosome entry sites (IRES) elements are used to create multigene, or polycistronic, messages, and these may be used in the invention.

[00262] Vectors can include a multiple cloning site (MCS), which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector. "Restriction enzyme digestion" refers to catalytic cleavage of a nucleic acid molecule with an enzyme that functions only at specific locations in a nucleic acid molecule. Many of these restriction enzymes are commercially available. Use of such enzymes is widely understood by those of skill in the art. Frequently, a vector is linearized or fragmented using a restriction enzyme that cuts within the MCS to allow exogenous sequences to be ligated to the vector. "Ligation"

refers to the process of forming phosphodiester bonds between two nucleic acid fragments, which may or may not be contiguous with each other. Techniques involving restriction enzymes and ligation reactions are well known to those of skill in the art of recombinant technology.

[00263] Splicing sites, termination signals, origins of replication, and selectable markers may also be employed.

[00264] *Plasmid Vectors*

[00265] In certain embodiments, a plasmid vector can be used to transform a host cell. Plasmid vectors containing replicon and control sequences which are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences which can provide phenotypic selection in transformed cells. In a non-limiting example, *E. coli* is often transformed using derivatives of pBR322, a plasmid derived from an *E. coli* species. pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, for example, promoters which can be used by the microbial organism for expression of its own proteins.

[00266] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM.TM. 11 may be utilized in making a recombinant phage vector which can be used to transform host cells, such as, for example, *E. coli* LE392.

[00267] Further useful plasmid vectors include pIN vectors (Inouye et al., 1985); and pGEX vectors, for use in generating glutathione S transferase (GST) soluble fusion proteins

for later purification and separation or cleavage. Other suitable fusion proteins are those with galactosidase, ubiquitin, and the like.

[00268] Bacterial host cells, for example, *E. coli*, comprising the expression vector, are grown in any of a number of suitable media, for example, LB. The expression of the recombinant protein in certain vectors may be induced, as would be understood by those of skill in the art, by contacting a host cell with an agent specific for certain promoters, e.g., by adding IPTG to the media or by switching incubation to a higher temperature. After culturing the bacteria for a further period, for example, between 2 and 24 h, the cells are collected by centrifugation and washed to remove residual media.

[00269] *Viral Vectors*

[00270] The ability of certain viruses to infect cells or enter cells via receptor mediated endocytosis, and to integrate into host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign nucleic acids into cells (e.g., mammalian cells). Components of the present invention may be a viral vector that encodes one or more CARs of the invention. Non-limiting examples of virus vectors that may be used to deliver a nucleic acid of the present invention are described below.

[00271] *Adenoviral Vectors*

[00272] A particular method for delivery of the nucleic acid involves the use of an adenovirus expression vector. Although adenovirus vectors are known to have a low capacity for integration into genomic DNA, this feature is counterbalanced by the high efficiency of gene transfer afforded by these vectors. "Adenovirus expression vector" is meant to include those constructs containing adenovirus sequences sufficient to (a) support packaging of the construct and (b) to ultimately express a tissue or cell specific construct that has been cloned

therein. Knowledge of the genetic organization of adenovirus, a 36 kb, linear, double stranded DNA virus, allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kb (Grunhaus and Horwitz, 1992).

[00273] *AAV Vectors*

[00274] The nucleic acid may be introduced into the cell using adenovirus assisted transfection. Increased transfection efficiencies have been reported in cell systems using adenovirus coupled systems (Kelleher and Vos, 1994; Cotten et al., 1992; Curiel, 1994). Adeno associated virus (AAV) is an attractive vector system for use in the cells of the present invention as it has a high frequency of integration and it can infect nondividing cells, thus making it useful for delivery of genes into mammalian cells, for example, in tissue culture (Muzyczka, 1992) or in vivo. AAV has a broad host range for infectivity (Tratschin et al., 1984; Laughlin et al., 1986; Lebkowski et al., 1988; McLaughlin et al., 1988). Details concerning the generation and use of rAAV vectors are described in U.S. Pat. Nos. 5,139,941 and 4,797,368, each incorporated herein by reference.

[00275] *Retroviral Vectors*

[00276] Retroviruses are useful as delivery vectors because of their ability to integrate their genes into the host genome, transferring a large amount of foreign genetic material, infecting a broad spectrum of species and cell types and of being packaged in special cell lines (Miller, 1992).

[00277] In order to construct a retroviral vector, a nucleic acid (e.g., one encoding the sequence of interest) is inserted into the viral genome in the place of certain viral sequences to produce a virus that is replication defective. In order to produce virions, a packaging cell line containing the gag, pol, and env genes but without the LTR and packaging components is

constructed (Mann et al., 1983). When a recombinant plasmid containing a cDNA, together with the retroviral LTR and packaging sequences is introduced into a special cell line (e.g., by calcium phosphate precipitation for example), the packaging sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann et al., 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration and stable expression require the division of host cells (Paskind et al., 1975).

[00278] Lentiviruses are complex retroviruses, which, in addition to the common retroviral genes gag, pol, and env, contain other genes with regulatory or structural function. Lentiviral vectors are well known in the art (see, for example, Naldini et al., 1996; Zufferey et al., 1997; Blomer et al., 1997; U.S. Pat. Nos. 6,013,516 and 5,994,136). Some examples of lentivirus include the Human Immunodeficiency Viruses: HIV-1, HIV-2 and the Simian Immunodeficiency Virus: SIV. Lentiviral vectors have been generated by multiply attenuating the HIV virulence genes, for example, the genes env, vif, vpr, vpu and nef are deleted making the vector biologically safe.

[00279] Recombinant lentiviral vectors can infect non-dividing cells and can be used for both in vivo and ex vivo gene transfer and expression of nucleic acid sequences. For example, recombinant lentivirus can infect a non-dividing cell wherein a suitable host cell is transfected with two or more vectors carrying the packaging functions, namely gag, pol and env, as well as rev and tat is described in U.S. Pat. No. 5,994,136, incorporated herein by reference. One may target the recombinant virus by linkage of the envelope protein with an antibody or a particular ligand for targeting to a receptor of a particular cell-type. By inserting a sequence (including a regulatory region) of interest into the viral vector, along with another

gene which encodes the ligand for a receptor on a specific target cell, for example, the vector is now target-specific.

[00280] **Other Viral Vectors**

[00281] Other viral vectors may be employed as vaccine constructs in the present invention. Vectors derived from viruses such as vaccinia virus (Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar et al., 1988), sindbis virus, cytomegalovirus and herpes simplex virus may be employed. They offer several attractive features for various mammalian cells (Friedmann, 1989; Ridgeway, 1988; Baichwal and Sugden, 1986; Coupar et al., 1988; Horwich et al., 1990).

[00282] **Delivery Using Modified Viruses**

[00283] A nucleic acid to be delivered may be housed within an infective virus that has been engineered to express a specific binding ligand. The virus particle will thus bind specifically to the cognate receptors of the target cell and deliver the contents to the cell. A new approach designed to allow specific targeting of retrovirus vectors was developed based on the chemical modification of a retrovirus by the chemical addition of lactose residues to the viral envelope. This modification can permit the specific infection of hepatocytes via sialoglycoprotein receptors.

[00284] Another approach to targeting of recombinant retroviruses was designed in which biotinylated antibodies against a retroviral envelope protein and against a specific cell receptor were used. The antibodies were coupled via the biotin components by using streptavidin (Roux et al., 1989). Using antibodies against major histocompatibility complex class I and class II antigens, they demonstrated the infection of a variety of human cells that bore those surface antigens with an ecotropic virus in vitro (Roux et al., 1989).

[00285] *Vector Delivery and Cell Transformation*

[00286] Suitable methods for nucleic acid delivery for transfection or transformation of cells are known to one of ordinary skill in the art. Such methods include, but are not limited to, direct delivery of DNA such as by ex vivo transfection, by injection, and so forth. Through the application of techniques known in the art, cells may be stably or transiently transformed.

[00287] *Ex Vivo Transformation*

[00288] Methods for transfecting eukaryotic cells and tissues removed from an organism in an ex vivo setting are known to those of skill in the art. Thus, cells or tissues may be removed and transfected ex vivo using nucleic acids of the present invention. In some embodiments, the transplanted cells or tissues may be placed into an organism. In other embodiments, a nucleic acid is expressed in the transplanted cells.

[00289] *Kits of the Invention*

[00290] Any of the compositions described herein may be comprised in a kit. In a non-limiting example, one or more cells for use in cell therapy and/or the reagents to generate one or more cells for use in cell therapy that harbors recombinant expression vectors may be comprised in a kit. The kit components are provided in suitable container means.

[00291] Some components of the kits may be packaged either in aqueous media or in lyophilized form. The container means of the kits can include at least one vial, test tube, flask, bottle, syringe or other container means, into which a component may be placed, and suitably aliquoted. Where there are more than one component in the kit, the kit also can contain a second, third or other additional container into which the additional components

may be separately placed. However, various combinations of components may be comprised in a vial. The kits of the present invention also can include a means for containing the components in close confinement for commercial sale. Such containers may include injection or blow molded plastic containers into which the desired vials are retained.

[00292] When the components of the kit are provided in one and/or more liquid solutions, the liquid solution is an aqueous solution, with a sterile aqueous solution being particularly useful. In some cases, the container means may itself be a syringe, pipette, and/or other such like apparatus, from which the formulation may be applied to an infected area of the body, injected into an animal, and/or even applied to and/or mixed with the other components of the kit.

[00293] However, the components of the kit may be provided as dried powder(s). When reagents and/or components are provided as a dry powder, the powder can be reconstituted by the addition of a suitable solvent. For example, the solvent may also be provided in another container means. The kits may also comprise a second container means for containing a sterile, pharmaceutically acceptable buffer and/or other diluent.

[00294] In particular embodiments of the invention, cells that are to be used for cell therapy are provided in a kit, and in some cases the cells are essentially the sole component of the kit. The kit may comprise reagents and materials to make the desired cell. In specific embodiments, the reagents and materials include primers for amplifying desired sequences, nucleotides, suitable buffers or buffer reagents, salt, and so forth, and in some cases the reagents include vectors and/or DNA that encodes a CAR as described herein and/or regulatory elements therefor.

[00295] In particular embodiments, there are one or more apparatuses in the kit suitable for extracting one or more samples from an individual. The apparatus may be a syringe, scalpel, and so forth.

[00296] In some cases of the invention, the kit, in addition to cell therapy embodiments, also includes a second cancer therapy, such as chemotherapy, hormone therapy, and/or immunotherapy, for example. The kit(s) may be tailored to a particular cancer for an individual and comprise respective second cancer therapies for the individual.

[00297] **Combination Therapy**

[00298] In certain embodiments of the invention, methods of the present invention for clinical aspects are combined with other agents effective in the treatment of hyperproliferative disease, such as anti-cancer agents. An "anti-cancer" agent can negatively affect cancer in a subject, for example, by killing cancer cells, inducing apoptosis in cancer cells, reducing the growth rate of cancer cells, reducing the incidence or number of metastases, reducing tumor size, inhibiting tumor growth, reducing the blood supply to a tumor or cancer cells, promoting an immune response against cancer cells or a tumor, preventing or inhibiting the progression of cancer, or increasing the lifespan of a subject with cancer. For example,, these other compositions would be provided in a combined amount effective to kill or inhibit proliferation of the cell. This process may involve contacting the cancer cells with the expression construct and the agent(s) or multiple factor(s) at the same time. This may be achieved by contacting the cell with a single composition or pharmacological formulation that includes both agents, or by contacting the cell with two distinct compositions or formulations, at the same time, wherein one composition includes the expression construct and the other includes the second agent(s).

[00299] Tumor cell resistance to chemotherapy and radiotherapy agents represents a major problem in clinical oncology. One goal of current cancer research is to find ways to improve the efficacy of chemo- and radiotherapy by combining it with other therapies. In one embodiment, cell therapy could be used similarly in conjunction with chemotherapeutic,

radiotherapeutic, or immunotherapeutic intervention, as well as pro-apoptotic or cell cycle regulating agents.

[00300] Alternatively, the present inventive therapy may precede or follow the other agent treatment by intervals ranging from minutes to weeks. In embodiments where the other agent and present invention are applied separately to the individual, one would generally ensure that a significant period of time did not expire between the time of each delivery, such that the agent and inventive therapy would still be able to exert an advantageously combined effect on the cell. In such instances, one may contact the cell with both modalities within about 12-24 h of each other (for example, within about 6-12 h of each other). In some situations, it may be desirable to extend the time period for treatment significantly, however, where several days (2, 3, 4, 5, 6 or 7) to several weeks (1, 2, 3, 4, 5, 6, 7 or 8) lapse between the respective administrations.

[00301] The treatment cycles would be repeated as necessary. For example, various standard therapies, as well as surgical intervention, may be applied in combination with the inventive cell therapy.

[00302] **Chemotherapy**

[00303] Cancer therapies also include a variety of combination therapies with both chemical and radiation based treatments. Combination chemotherapies include, but are not limited to, for example, abraxane, altretamine, docetaxel, herceptin, methotrexate, novantrone, zoladex, cisplatin (CDDP), carboplatin, procarbazine, mechlorethamine, cyclophosphamide, camptothecin, ifosfamide, melphalan, chlorambucil, busulfan, nitrosurea, dactinomycin, daunorubicin, doxorubicin, bleomycin, plicomycin, mitomycin, etoposide (VP16), tamoxifen, raloxifene, estrogen receptor binding agents, taxol, gemcitabine, navelbine, farnesyl-protein transferase inhibitors, transplatin, 5-fluorouracil, vincristin,

vinblastin and methotrexate, or any analog or derivative variant of the foregoing and also combinations thereof.

[00304] In specific embodiments, chemotherapy for the individual is employed in conjunction with the invention, for example before, during and/or after administration of the invention

[00305] **Radiotherapy**

[00306] Other factors that cause DNA damage and have been used extensively include what are commonly known as .gamma.-rays, X-rays, and/or the directed delivery of radioisotopes to tumor cells. Other forms of DNA damaging factors are also useful such as microwaves and UV-irradiation. It is most likely that all of these factors effect a broad range of damage on DNA, on the precursors of DNA, on the replication and repair of DNA, and on the assembly and maintenance of chromosomes. Dosage ranges for X-rays range from daily doses of 50 to 200 roentgens for prolonged periods of time (3 to 4 wk), to single doses of 2000 to 6000 roentgens. Dosage ranges for radioisotopes vary widely, and depend on the half-life of the isotope, the strength and type of radiation emitted, and the uptake by the neoplastic cells.

[00307] The terms "contacted" and "exposed," when applied to a cell, are used herein to describe the process by which a therapeutic construct and a chemotherapeutic or radiotherapeutic agent are delivered to a target cell or are placed in direct juxtaposition with the target cell. To achieve cell killing or stasis, both agents are delivered to a cell in a combined amount effective to kill the cell or prevent it from dividing.

[00308] **Immunotherapy**

[00309] Immunotherapeutics rely on the use of immune effector cells and molecules to target and destroy cancer cells. The immune effector may be, for example, an antibody specific for some marker on the surface of a tumor cell. The antibody alone may serve as an effector of therapy or it may recruit other cells to actually effect cell killing. The antibody also may be conjugated to a drug or toxin (chemotherapeutic, radionuclide, ricin A chain, cholera toxin, pertussis toxin, etc.) and serve merely as a targeting agent. Alternatively, the effector may be a lymphocyte carrying a surface molecule that interacts, either directly or indirectly, with a tumor cell target. Various effector cells include cytotoxic T cells and NK cells.

[00310] Immunotherapy other than the inventive therapy described herein could thus be used as part of a combined therapy, in conjunction with the present cell therapy. The approach for combined therapy is discussed herein. For example, the tumor cell must bear some marker that is amenable to targeting, i.e., is not present on the majority of other cells. Many tumor markers exist and any of these may be suitable for targeting in the context of the present invention. Common tumor markers include PD-1, PD-L1, CTLA4, carcinoembryonic antigen, prostate specific antigen, urinary tumor associated antigen, fetal antigen, tyrosinase (p97), gp68, TAG-72, HMFG, Sialyl Lewis Antigen, MucA, MucB, PLAP, estrogen receptor, laminin receptor, erb B and p155.

[00311] **Genes**

[00312] In yet another embodiment, the secondary treatment is a gene therapy in which a therapeutic polynucleotide is administered before, after, or at the same time as the present invention clinical embodiments. A variety of expression products are encompassed within the

invention, including inducers of cellular proliferation, inhibitors of cellular proliferation, or regulators of programmed cell death.

[00313] **Surgery**

[00314] Approximately 60% of persons with cancer will undergo surgery of some type, which includes preventative, diagnostic or staging, curative and palliative surgery. Curative surgery is a cancer treatment that may be used in conjunction with other therapies, such as the treatment of the present invention, chemotherapy, radiotherapy, hormonal therapy, gene therapy, immunotherapy and/or alternative therapies.

[00315] Curative surgery includes resection in which all or part of cancerous tissue is physically removed, excised, and/or destroyed. Tumor resection refers to physical removal of at least part of a tumor. In addition to tumor resection, treatment by surgery includes laser surgery, cryosurgery, electrosurgery, and microscopically controlled surgery (Mohs' surgery). For example, the present invention may be used in conjunction with removal of superficial cancers, precancers, or incidental amounts of normal tissue.

[00316] Upon excision of part of all of cancerous cells, tissue, or tumor, a cavity may be formed in the body. Treatment may be accomplished by perfusion, direct injection or local application of the area with an additional anti-cancer therapy. Such treatment may be repeated, for example, every 1, 2, 3, 4, 5, 6, or 7 days, or every 1, 2, 3, 4, and 5 weeks or every 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months. These treatments may be of varying dosages as well.

[00317] **Other Agents**

[00318] In some embodiments, other agents may be used in combination with the present invention to improve the therapeutic efficacy of treatment. These additional agents

include immunomodulatory agents, agents that affect the upregulation of cell surface receptors and GAP junctions, cytostatic and differentiation agents, inhibitors of cell adhesion, or agents that increase the sensitivity of the hyperproliferative cells to apoptotic inducers. Immunomodulatory agents include tumor necrosis factor; interferon alpha, beta, and gamma; IL-2 and other cytokines; F42K and other cytokine analogs; or MIP-1, MIP-1beta, MCP-1, RANTES, and other chemokines. In some embodiments, upregulation of cell surface receptors or their ligands such as Fas/Fas ligand, DR4 or DR5/TRAIL would potentiate the apoptotic inducing abilities of the present invention by establishment of an autocrine or paracrine effect on hyperproliferative cells. Increases intercellular signaling by elevating the number of GAP junctions would increase the anti-hyperproliferative effects on the neighboring hyperproliferative cell population. In other embodiments, cytostatic or differentiation agents can be used in combination with the present invention to improve the anti-hyperproliferative efficacy of the treatments. Inhibitors of cell adhesion can also be used to improve the efficacy of the present invention. Examples of cell adhesion inhibitors are focal adhesion kinase (FAKs) inhibitors and Lovastatin. In some embodiments, other agents that increase the sensitivity of a hyperproliferative cell to apoptosis, such as the antibody c225, could be used in combination with the present invention to improve the treatment efficacy.

[00319] Methods of Assessing Activity of Engineered CAR T cells

[00320] Aspects of the disclosure are further directed towards methods of and kits for assessing the killing capabilities of engineered CAR T cells. Specifically, embodiments are directed towards an immune complex analysis method and kits to determine the CAR T cell activity during co-culture with cancer cells. First, different target cancer cells (for example, HEK293T, MDA-MB-231, MDA-MB-468, HCC38, and skrc59) are stained with a dye, (for

example, ViaStain™ Tracer Blue dye), seeded in a plate (for example a 96-well plate), and incubated for a period of time (for example 12 hours, or overnight). Next, different T cell types (for example, two different T cell types) are added to the wells (for example at a ratio of 20:1 effector-to-target (E:T) ratios) and allowed to co-culture for a period of time (for example, 24 hours). Finally, the plate is scanned and analyzed (for example, using the bright field and blue fluorescent channels). The immune complexes were analyzed by confluence measurement and compared to the negative control of untransduced T cells. As a result, data plots displayed the CAR T cell activities for all of tested target and effector cells combinations. Utilization of an image cytometry platform can visually confirm interactions between effector and target cells, thus making results highly accurate and robust.

EXAMPLES

[00321] Examples are provided below to facilitate a more complete understanding of the invention. The following examples illustrate the exemplary modes of making and practicing the invention. However, the scope of the invention is not limited to specific embodiments disclosed in these Examples, which are for purposes of illustration only, since alternative methods can be utilized to obtain similar results.

Example 1

[00322] Clear cell renal cell carcinoma (ccRCC) is the major type of RCC which is among the 10 most common cancers in both men and women. Chimeric Antigen Receptor (CAR) T cells have proven to be a powerful, clinically translatable immunotherapy for hematologic malignancies. However, these results have not been translatable to solid tumors due to inefficient homing of CAR T cells, the suppressive tumor microenvironment, and on-target off-tumor toxicities resulted from the sharing of CAR T targeting epitopes on healthy tissues. To combat the suppressive microenvironment, immune checkpoint blockade has shown an enhanced effect on antitumor response by restoring the local antitumor immunity. CAR T cell factories were designed to empower CAR T cells through the secretion of human anti-immune checkpoint inhibitor monoclonal antibodies (mAbs) locally at the tumor site. Our results show a dramatic improvement in CAR T killing of ccRCC *in vitro* and *in vivo* by reversing CAR T cell and tumor infiltrating lymphocyte (TIL) exhaustion.

[00323] CAIX is an ideal target for ccRCC therapy and used as a CAR target for the first clinical trial (see, for example, Lamers, Sleijfer et al. 2006; Lamers, Willemsen et al. 2011). However, it led to serious side effects due to CAIX expression on the bile duct. Therefore, it is crucial to develop a CAR with elevated efficacy and safety (such as limited on-target off-tumor effect). To achieve that, second generation CARs were developed by introducing a 2nd targeting scFv (for example, anti-CD70 scFv) in the CAR T cell factory

together with the first targeting scFv (for example, anti-CAIX scFv) to allow the CAR to target two unique antigens simultaneously. See, for example, FIG. 1. By IHC staining of ccRCC patient samples, we found that target CD70 is an ideal target to be utilized as the 2nd target since it is highly expressed on ccRCC and co-expressed with CAIX.

[00324] Our 27 billion-member human scFv-phage display library was panned against the antigen expressing skrc-59 ccRCC cells and subtracted with the antigen absent skrc-59 ccRCC cells to identify new scFvs. Their binding kinetics (K_{on}/K_{off}) and affinity (K_d) were then measured via an OctetRed 96 instrument and scFvs with desirable kinetics were evaluated for their ability to bind with antigen expressing cells. Candidates were cloned into vectors where anti-CD70 and anti-CAIX scFvs were combined in different permutations by changing the order of the two targeting scFvs with various linkers connected to a costimulatory domain (CD28, 41BB) and an activating domain (CD3). The 4th generation lentivirus packaging system was used to obtain CAR lentiviruses and primary T cells isolated from PBMCs were transduced to express dual CAR and were tested against different cell lines *in vitro*. For further evaluation *in vivo*, a humanized orthotopic ccRCC mouse model was established by injecting luciferized ccRCC cells under the kidney capsule of NSG-SGM3 mice with a reconstituted human immune system.

[00325] In summary, utilizing a dual CAR T discovery platform, we generated a series of CARs with different scFvs, linkers, and hinges. The skilled artisan will recognize that embodiments can comprise different combinations of scFvs, linkers, and hinges. For example, difference combinations of scFvs, linkers, and hinges can be used to treat different cancers.

[00326] By combining the best dual CAR with a payload, such as an immune checkpoint blockade payload, the 2nd generation CAR T was discovered. Without wishing to

be bound by theory, the 2nd generation CAR T cell factories can be used for the treatment of cancers, such as ccRCC, and are able to eliminate side effects on normal tissues.

[00327] References cited in this Example:

[00328] Lamers, C. H., S. Sleijfer, A. G. Vulto, W. H. Kruit, M. Kliffen, R. Debets, J. W. Gratama, G. Stoter and E. Oosterwijk (2006). "Treatment of metastatic renal cell carcinoma with autologous T-lymphocytes genetically retargeted against carbonic anhydrase IX: first clinical experience." J Clin Oncol 24(13): e20-22.

[00329] Lamers, C. H., R. Willemsen, P. van Elzakker, S. van Steenbergen-Langeveld, M. Broertjes, J. Oosterwijk-Wakka, E. Oosterwijk, S. Sleijfer, R. Debets and J. W. Gratama (2011). "Immune responses to transgene and retroviral vector in patients treated with ex vivo-engineered T cells." Blood 117(1): 72-82.

Example 2

[00330] Chimeric Antigen Receptor (CAR) T cells have proven to be a powerful immunotherapy for hematologic malignancies, however, these have not been translated to solid tumors. Our CAR T cell factories empower CAR T cells through secreting antibodies, such as human anti-immune checkpoint inhibitor monoclonal antibodies (mAbs), locally at the tumor site to restore tumor microenvironment to achieve solid tumor cure. With the introduction of one or more additional scFvs, the 2nd generation CAR has elevated efficacy and safety and holds great promise in clinic.

[00331] The bispecific tandem CAR was designed to target antigens associated with tumors, such as CAIX and CD70 (see FIG. 1 and FIG. 2, for example) which are both highly expressed and co-expressed on ccRCC primary cell by IHC (FIG. 3). Successful panning of a 27 billion member phage display library has led to the identification of several anti-CD70 scFvs (see FIG. 6, for example). The anti-CAIX scFvs we discovered before and anti-CD70

scFvs were cloned into pHAGE vector with different linkers (See FIG. 9, for example). Then lentiviruses were packaged and transduced with primary T cells. CAR T cells were obtained and evaluated on 4 different CRISPR engineered cell lines (see FIG. 5, for example). The killing activity was assessed in Celigo and Cr51 release assay (See FIG. 12 and 13, for example). Selectivity was shown as described herein.

[00332] This CAR T cell factories can be used in therapy of CAIX and/or CD70 over-expressed cancers or in combination with other therapies.

EXAMPLE 3

[00333] CAIX and CD70 dual IHC staining quantification

	Scan #	Tissue	RCCT numbers	T	N	M	Stage	% CD70 Positive Cells	% CAIX Positive Cells	% CD70 & CAIX Positive Cells	% CD70 or CAIX Positive Cells
1	477491	Normal Kidney	RCCT0461	3a	2		III				0.0
2	477495	Normal Kidney	RCCT0487	3a	0	1	IV				0.0
3	477471 & 477472	Fat cells and liver with bile duct									
5	477473 & 477474	Bile duct									
7	477465	PRCC	RCCT0165	2			II				0.0
8	477466	PRCC	RCCT0115	3b	0	0	III	59.8	96.5	59.4	96.9
9	477470	PRCC	RCCT0564	3		0	III				0.0
10	477477	ccRCC	RCCT0015	T1a	x	x	I	38.5	93.3	37.2	94.6
11	477501	ccRCC	RCCT0257	1a	0	0	I	25.5	98.4	25.5	98.4
12	477469	ccRCC	RCCT0434	1b			I	88.1	93.3	85.7	95.7
13	477496	ccRCC	RCCT0574	1b		0	I	93.4	93.7	93.2	93.9
14	477498	ccRCC	RCCT0581	2	0	0	II	15.7	92.4	15.2	92.9
15	477504	ccRCC	RCCT0402	2	0	0	II	89.5	89.6	82.6	96.4
16	477475	ccRCC	RCCT0520	2a		0	II	84.8	94.8	84.0	95.6
17	477481	ccRCC	RCCT0565	2a	0	0	II	0.8	95.1	0.8	95.1
18	477482	ccRCC	RCCT0566	2a	0		II	25.6	86.9	18.5	94.0
19	477483	ccRCC	RCCT0573	2a	0	0	II	81.0	97.0	81.0	97.1
20	477479	ccRCC	RCCT0049	2a	0	0	II	0.2	91.3	0.2	91.3
21	477476	ccRCC	RCCT0533	3	0	0	III	0.5	95.7	0.5	95.7
22	477487	ccRCC	RCCT0129	3	1		III	68.0	95.7	68.0	95.7
23	477488	ccRCC	RCCT0010	3a	0		III	94.2	95.0	93.9	95.3
24	477497	ccRCC	RCCT0411	3a	1		III	31.1	98.6	30.8	98.9
25	477499	ccRCC	RCCT0330	T3b	0	X	III	80.2	81.3	65.6	96.0
26	477500	ccRCC	RCCT0334	3b		0	III	83.6	90.8	81.0	93.3
27	477490	ccRCC	RCCT0317			1	IV				0.0
28	477478	ccRCC	RCCT0444	1b	0	1	IV				0.0
29	477484	ccRCC	RCCT0156	2	0	1	IV				0.0
30	477502	ccRCC	RCCT0394	2	0	1	IV	70.1	91.3	64.4	97.0
31	477485	ccRCC	RCCT0056	3	0	1	IV	42.7	93.0	41.9	93.8
32	477486	ccRCC	RCCT0170	3a	0	1	IV	46.0	17.7	12.6	51.0
33	477468	ccRCC	RCCT0400	3b		1	IV	10.8	62.2	9.7	63.3
34	477489	ccRCC	RCCT0447	3b	1	1	IV	21.8	95.3	21.8	95.3
35	477494	ccRCC	RCCT0119	3b	1	1	IV	0.6	96.7	0.6	96.7
36	477503	ccRCC	RCCT0383	3b	0	1	IV	89.1	93.9	88.5	94.5
37	477493	ccRCC	RCCT0016	4	0	1	IV	3.4	94.7	3.1	95.0
38	477467	ccRCC	RCCT0384					45.3	97.3	45.2	97.3
39	477480	ccRCC						56.7	97.1	56.5	97.4
40	477492	ccRCC	RCCT0563			0		30.5	95.8	29.9	96.5

EQUIVALENTS

[00334] Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such equivalents are considered to be within the scope of this invention, and are covered by the following claims.

What is claimed:

1. An engineered cell comprising a chimeric antigen receptor, wherein the chimeric antigen receptor comprises an extracellular ligand binding domain that is specific for a first antigen and a second antigen on the surface of a cancer cell, wherein the first antigen comprises CAIX and the second antigen comprises CD70.
2. The engineered cell of claim 1, wherein the CAR further comprises a transmembrane polypeptide and an intracellular signaling domain.
3. The engineered cell of claim 2, wherein the CAR further comprises a co-stimulatory domain.
4. The engineered cell of claim 1, wherein the extracellular ligand binding domain comprises an antibody or fragment thereof.
5. The engineered cell of claim 4, wherein the antibody comprises a VH and/or VL according to Table 2, or any combination thereof.
6. The engineered cell of claim 4, wherein the antibody comprises a VH and/or VL according to Table 4, or any combination thereof.
7. The engineered cell of claim 4, wherein the extracellular binding domain comprises a VH and/or a VL of Table 2 and Table 4, or any combination thereof.
8. The engineered cell of claim 4, wherein the antibody comprises a CDR1, CDR2, and/or CDR3 of Table 1, or any combination thereof.
9. The engineered cell of claim 4, wherein the antibody comprises a CDR1, CDR2, and/or CDR3 of Table 3, or any combination thereof.
10. The engineered cell of claim 4, wherein the extracellular binding domain comprises a CDR1, CDR2, and/or CDR3 of Table 1 and Table 3, or any combination thereof.
11. The engineered cell of claim 1, wherein the engineered cell expresses and secretes a recombinant polypeptide.

12. The engineered cell of claim 11, wherein the recombinant polypeptide comprises an antibody or fragment thereof, or a cytokine.
13. The engineered cell of claim 11, wherein the recombinant polypeptide modulates the immune system of a subject.
14. The engineered cell of claim 11, wherein the recombinant polypeptide is an immune checkpoint blockade antibody.
15. The engineered cell of claim 11, wherein the recombinant polypeptide modulates tumor vasculogenesis.
16. The engineered cell of claim 15, wherein the recombinant polypeptide is specific for VEGF, VEGFR1, VEGFR2, PDGF, Ang-1, or AT1.
17. The engineered cell of claim 11, wherein the recombinant polypeptide comprises an antibody or fragment thereof specific for TIGIT, GITR, PD-L1, PD-L2, PD-1, CTLA-4, VISTA, CD70, TIM-3, LAG-3, CD40L, or CCR4.
18. The engineered cell of claim 12, wherein the cytokine comprises IL-12, IL-15, or IL-18.
19. The engineered cell of claim 1, wherein the cell comprises a T cell, an NK cell, or an NKT cell.
20. The engineered cell of claim 19, wherein the T cell is CD4+, CD8+, CD3+ panT cells, or any combination thereof.
21. The engineered cell of claim 19, wherein the T cell is a mixed population of CD4+ and CD8+ T cells.
22. A nucleic acid construct encoding a chimeric antigen receptor, wherein the chimeric antigen receptor comprises an extracellular ligand binding domain that is specific for a first antigen and a second antigen on the surface of a cancer cell, wherein the first antigen comprises CAIX and the second antigen comprises CD70.

23. The nucleic acid construct of claim 22, wherein the chimeric antigen receptor further comprises a transmembrane polypeptide and an intracellular signaling domain.
24. The nucleic acid construct of claim 23, wherein the chimeric antigen receptor further comprises a co-stimulatory domain.
25. The nucleic acid construct of claim 23, wherein the nucleic acid construct further encodes a recombinant polypeptide.
26. The nucleic acid construct of claim 25, wherein the recombinant polypeptide can be secreted from an engineered cell.
27. The nucleic acid construct of claim 25, wherein the recombinant polypeptide comprises an antibody or fragment thereof, or a cytokine.
28. The nucleic acid construct of claim 25, wherein the recombinant polypeptide modulates the immune system of a subject.
29. The nucleic acid construct of claim 25, wherein the recombinant polypeptide is an immune checkpoint blockade antibody.
30. The nucleic acid construct of claim 25, wherein the recombinant polypeptide modulates tumor vasculogenesis.
31. The nucleic acid construct of claim 25, wherein the recombinant polypeptide comprises an antibody or fragment thereof specific for TIGIT, GITR, PD-L1, PD-L2, PD-1, CTLA-4, VISTA, CD70, TIM-3, LAG-3, CD40L, or CCR4.
32. The nucleic acid of claim 27, wherein the cytokine comprises IL12, IL15 or IL18.
33. The nucleic acid of claim 30, wherein the recombinant polypeptide is specific for VEGF, VEGFR1, VEGFR2, PDGF, Ang-1, or AT1
34. A vector comprising the nucleic acid construct of claim 22.
35. A cell comprising the vector of claim 34.

36. A method for treating a subject afflicted with cancer, the method comprising administering the subject a therapeutically effective amount of the engineered cell of claim 1.
37. A method of reducing progression or promoting regression of a cancer in a subject, the method comprising administering the subject a therapeutically effective amount of the engineered cell of claim 1.
38. A method of reducing cellular proliferation of a cancer cell in a subject, the method comprising administering the subject a therapeutically effective amount of the engineered cell of claim 1.
39. The method of any one of claims 36-38, wherein cancer comprises renal cell carcinoma.
40. A chimeric antigen receptor (CAR) comprising an extracellular ligand binding domain, wherein the extracellular ligand binding domain is specific for a first antigen and a second antigen on the surface of a cancer cell, wherein the first antigen comprises CAIX and the second antigen comprises CD70.
41. The CAR of claim 40, wherein the CAR further comprises a transmembrane polypeptide and an intracellular signaling domain.
42. The CAR of claim 41, wherein the CAR further comprises a co-stimulatory domain.
43. The CAR of claim 40, wherein the extracellular ligand binding domain comprises an antibody or fragment thereof.
44. A cell comprising the chimeric antigen receptor (CAR) of claim 40.
45. An engineered cell comprising a first chimeric antigen receptor and a second chimeric antigen receptor, wherein the first chimeric antigen receptor comprises an extracellular ligand binding domain specific for CAIX, and wherein the second

chimeric antigen receptor comprises an extracellular ligand binding domain that is specific for CD70.

46. The engineered cell of claim 45, wherein the engineered cell expresses and secretes a recombinant polypeptide.
47. The engineered cell of claim 46, wherein the recombinant polypeptide comprises an antibody or fragment thereof or a cytokine.
48. The engineered cell of claim 46, wherein the recombinant polypeptide modulates the immune system of a subject.
49. The engineered cell of claim 46, wherein the recombinant polypeptide is an immune checkpoint blockade antibody.
50. The engineered cell of claim 46, wherein the recombinant polypeptide modulates tumor vasculogenesis.
51. The engineered cell of claim 46, wherein the recombinant polypeptide comprises an antibody or fragment thereof specific for TIGIT, GITR, PD-L1, PD-L2, PD-1, CTLA-4, VISTA, CD70, TIM-3, LAG-3, CD40L, or CCR4.
52. The engineered cell of claim 47, wherein the cytokine comprises IL12, IL15 or IL18.
53. The nucleic acid of claim 50, wherein the recombinant polypeptide is specific for VEGF, VEGFR1, VEGFR2, PDGF, Ang-1, or AT1.
54. The engineered cell of claim 45, wherein the cell comprises a T cell or an NK cell.
55. The engineered cell of claim 54, wherein the T cell is T cell is CD4+, CD8+, CD3+ panT cells, or any combination thereof.
56. The engineered cell of claim 54, wherein the T cell is a mixed population of CD4+ and CD8+ T cells.
57. The engineered cell of claim 45, wherein the first chimeric antigen receptor and the second chimeric antigen receptor are expressed from a single nucleic acid construct.




FIG. 1

2nd generation CAR-T cell factories








FIG. 2

FIG. 3

ccRCC-Patient 9-T
primary cell line



ccRCC-Patient 9-D
primary cell line



CAIX

CD70





FIG. 4



	CAIX	CD70
Skrc-59 CAIX+ CD70+	2658.93	43.43
Skrc-59 CAIX+ CD70-	1403.36	3.98
Skrc-59 CAIX- CD70+	30.37	132.66
Skrc-59 CAIX- CD70-	35.38	3.57

FIG. 5 – CONT.




FIG. 6




FIG. 7




FIG. 8

2nd generation CAR-T cells





FIG. 9



	G36-GGGGS1-7	7-GGGGS1-G36	7-GGGGS3-G36	G36-GGGGS5-CD27	7-GGGGS5-G36	G36-hinge-7	7-hinge-G36	G36
EC ₅₀ (ug/ml)	0.7004	0.1093	0.1177	0.4107	0.09124	0.5921	0.1138	0.1315

FIG. 10





FIG. 11

	G36-GGGGS1-B7	B7-GGGGS1-G36	B7-GGGGS3-G36	G36-GGGGS5-CD27	B7-GGGGS5-G36	G36-hinge-B7	B7-hinge-G36	CD27	B7
EC ₅₀ (ug/ml)	0.1548	0.1348	0.1928	0.6188	0.1251	0.1766	0.2344	0.9166	0.2267



skrc-59 CA9+ CD70+ mCardinal CA9+ CD70- BFP skrc-59 CA9+ CD70+ mCardinal CA9- CD70+ BFP 13/92

Selectivity index = BFP / mCardinal = non-target cells / target cells

FIG. 12




FIG. 13




FIG. 14




FIG. 15

FIG. 16

19 anti-CAIX ScFvs

CAIX Abs	Affinity (KD) (nM)	Mapping* (CA-GST) (OD450)	Cross-competition Grouping	% CA Inhibition by anti-CAIX scFvFc	Internalization
G119	1.49	—	1	25	+++***
G10	1.62	3	1	15	+
G37	1.89	3	1	40**	—***
G106	3.20	2.4	1	20	++
G36	3.22	3	1	10	+++***
G45	12.50	—	1	—	+
G39	3.43	0.25	1	50**	17/92
G57	4.25	—	1	—	—
G40	21.78	—	1a	15	—
G6	25.90	0.9	1a	50**	—
G27	25.12	—	2	—	+
G125	40.32	—	3	40**	—
G9	99.58	0.4	3	—	—

*All anti-CAIX antibodies recognize the CA domain but only a selected group interacts with bacterially produced CA-GST (Figure 3D &E).

**Percentage of carbonic anhydrase inhibition by CAIX antibodies was estimated from Figure 4B. The rest data are from Figure 4A.

***Internalization measured by both flow cytometry and ImageStream. + and — denotes degree of internalization as measured by regular flow cytometry (Figure 5).

doi:10.1371/journal.pone.0009625.t004

Group 1, group 1a, group 2, group 3.

FIG. 16 – CONT.

Group 1, group 1a, group 2, group 3.

	scFv-Fc	KD(nM)	Internalization	Killing activity	Sequencing validation	pHAGE	Lentivirus
1	G6	25.90	-	++	✓	✓	✓
2	G9	99.58	-	++	✓	✓	✓
3	G10	1.62	+	+++	✓	✓	✓
4	G17			++	✓	✓	✓
5	G21			+++	✓	✓	✓
6	G27	25.12	+	++	✓	✓	✓
7	G28			++	✓	✓	✓
8	G36	3.22	++	+++	✓	✓	✓
9	G37	1.89	-	++++	✓	✓	✓
10	G39	3.43	-	++++	✓	✓	✓
11	G40	21.78	-	+++	✓	✓	✓
12	G45	12.50	+	+++	✓	✓	✓
13	G57	4.25	-	+++	✓	✓	✓
14	G62			+++	✓	✓	✓
15	G98			+++	✓	✓	✓
16	G104	Non-binder		+	✓	✓	✓
17	G106	3.20	++	+++	✓	✓	✓
18	G119	1.49	+++	+++	✓	✓	✓
19	G125	40.32	-	++++	✓	✓	✓

Anti-CAIX scFvs were cloned into pHAGE vector and packaged as lentivirus. CAR T cells were generated and tested killing activity. We proved that there is correlation between affinity of scFv and killing of CART, G37, G39, G125> G10, G21, G36, G40, G45, G57, G62, G98, G106, G119> G6, G9, G17, G27, G28> G104.




FIG. 17

20/92




FIG. 18




FIG. 19

- Celigo assay
- FACS assay





FIG. 20


E:T=5:1



Selectivity index = BFP / mCardinal = non-target cells / target cells


FIG. 21

E:T=5:1



Selectivity index = BFP / mCardinal = non-target cells / target cells

FIG. 22



Selectivity index = BFP / mCardinal = non-target cells / target cells

FIG. 23




FIG. 24




FIG. 25




FIG. 26

Oncotarget 7:23 (2016)




FIG. 27




FIG. 28




FIG. 29




FIG. 30




FIG. 31




FIG. 32




FIG. 33




FIG. 34




FIG. 35





FIG. 36



Front Immunol. 2018 Oct 26;9:2486

FIG. 37





FIG. 38



477495 T3a M1
ccRCC
Normal kidney

FIG. 39




FIG. 40

CAIX Abs	Affinity (KD) (nM)	Mapping* (CA-GST) (OD450)	Cross-competition Grouping	% CA Inhibition by anti-CAIX scFvFc	Internalization
G119	1.49	—	1	25	+++**
G10	1.62	3	1	15	+
G37	1.89	3	1	40**	+++**
G106	3.20	2.4	1	20	++
G36	3.22	3	1	10	+++**
G45	12.50	—	1	—	+
G39	3.43	0.25	1	50**	—
G57	4.25	—	1	—	—
G40	21.78	—	1a	15	—
G6	25.90	0.9	1a	50**	—
G27	25.12	—	2	—	+
G125	40.32	—	3	40**	—
G9	99.58	0.4	3	—	—


*All anti-CAIX antibodies recognize the CA domain but only a selected group interacts with bacterially produced CA-GST (Figure 3D &E).
**Percentage of carbonic anhydrase inhibition by CAIX antibodies was estimated from Figure 4B. The rest data are from Figure 4A.
***Internalization measured by both flow cytometry and ImageStream. + and — denotes degree of internalization as measured by regular flow cytometry (Figure 5).
doi:10.1371/journal.pone.0209625.t004

FIG. 41

	scFv-Fc	KD(nM)	Internalization	MonoCAR killing	MonoCAR-293T binding
1	G6	25.90	-	++	21.3
2	G9	99.58	-	++	24.9
3	G10	1.62	+	+++	39.6
4	G17			++	29.6
5	G21			+++	29.5
6	G27	25.12	+	++	15.5
7	G28			++	8.4
8	G36	3.22	++	+++	32.1
9	G37	1.89	-	+++	33.7
10	G39	3.43	-	+++	25.7
11	G40	21.78	-	+++	29.6
12	G45	12.50	+	+++	26
13	G57	4.25	-	+++	19.4
14	G62			+++	35.1
15	G98			+++	35
16	G104	Non-binder		+	0.9
17	G106	3.20	++	+++	47.7
18	G119	1.49	+++	+++	36
19	G125	40.32	-	++++	34.5
20	G250				

FIG. 42

“Or” Gating to Capture Tumor Cell Heterogeneity



But not Kill Low Target Density Healthy Cells



FIG. 43

Tumor type	CD70+ of total	%CD70+
Kidney ^a	204 of 283	72
Pancreas	33 of 140	23
Larynx or pharynx ^b	18 of 82	22
Melanoma	15 of 98	16
Ovary	37 of 241	15
Lung adenocarcinoma	17 of 172	10
Colon	17 of 194	9
Breast	5 of 204	2
Brain	6 of 59	10
6189 of 230 (82%) clear cell carcinoma and 4 of 6 (50%) papillary carcinoma.		
^a Includes nasopharyngeal cancer.		

British Journal of Cancer 103 (2010) 676-684

FIG. 44





FIG. 45



Stage IV
477490 M1
ccRCC

FIG. 46




FIG. 47




FIG. 48




FIG. 49




FIG. 50





FIG. 51



	G36-GGGGS1-B7	B7-GGGGS1-G36	G36-GGGGS3-CD27	B7-GGGGS5-G36	G36-hinge-B7	B7-hinge-G36	CD27	B7
EC ₅₀ (ug/ml)	0.1348	0.1348	0.1928	0.6188	0.1251	0.1766	0.2344	0.2267

FIG. 52





FIG. 53



Selectivity index = BFP / mCardinal = non-target cells / target cells

FIG. 54




FIG. 55




FIG. 56




FIG. 57

VH:

	1	FW1	FW2	FW3	FW4	FW5	FW6	FW7	FW8	FW9	FW10	FW11	FW12	FW13	FW14	FW15	FW16	FW17	FW18	FW19	FW20	FW21	FW22	FW23	FW24	FW25	FW26	FW27	FW28	FW29	FW30	FW31	FW32	FW33	FW34	FW35	FW36	FW37	FW38	FW39	FW40	FW41	FW42	FW43	FW44	FW45	FW46	FW47	FW48	FW49	FW50	FW51	FW52	FW53	FW54	FW55	FW56	FW57	FW58	FW59	FW60	FW61	FW62	FW63	FW64	FW65	FW66	FW67	FW68	FW69	FW70	FW71	FW72	FW73	FW74	FW75	FW76	FW77	FW78	FW79	FW80	FW81	FW82	FW83	FW84	FW85	FW86	FW87	FW88	FW89	FW90	FW91	FW92	FW93	FW94	FW95	FW96	FW97	FW98	FW99	FW100	FW101	FW102	FW103	FW104	FW105	FW106	FW107	FW108	FW109	FW110	FW111	FW112	FW113	FW114	FW115	FW116	FW117	FW118	FW119	FW120	FW121	FW122	FW123	FW124	FW125	FW126	FW127	FW128	FW129	FW130	FW131	FW132	FW133	FW134	FW135	FW136	FW137	FW138	FW139	FW140	FW141	FW142	FW143	FW144	FW145	FW146	FW147	FW148	FW149	FW150	FW151	FW152	FW153	FW154	FW155	FW156	FW157	FW158	FW159	FW160	FW161	FW162	FW163	FW164	FW165	FW166	FW167	FW168	FW169	FW170	FW171	FW172	FW173	FW174	FW175	FW176	FW177	FW178	FW179	FW180	FW181	FW182	FW183	FW184	FW185	FW186	FW187	FW188	FW189	FW190	FW191	FW192	FW193	FW194	FW195	FW196	FW197	FW198	FW199	FW200	FW201	FW202	FW203	FW204	FW205	FW206	FW207	FW208	FW209	FW210	FW211	FW212	FW213	FW214	FW215	FW216	FW217	FW218	FW219	FW220	FW221	FW222	FW223	FW224	FW225	FW226	FW227	FW228	FW229	FW230	FW231	FW232	FW233	FW234	FW235	FW236	FW237	FW238	FW239	FW240	FW241	FW242	FW243	FW244	FW245	FW246	FW247	FW248	FW249	FW250	FW251	FW252	FW253	FW254	FW255	FW256	FW257	FW258	FW259	FW260	FW261	FW262	FW263	FW264	FW265	FW266	FW267	FW268	FW269	FW270	FW271	FW272	FW273	FW274	FW275	FW276	FW277	FW278	FW279	FW280	FW281	FW282	FW283	FW284	FW285	FW286	FW287	FW288	FW289	FW290	FW291	FW292	FW293	FW294	FW295	FW296	FW297	FW298	FW299	FW300	FW301	FW302	FW303	FW304	FW305	FW306	FW307	FW308	FW309	FW310	FW311	FW312	FW313	FW314	FW315	FW316	FW317	FW318	FW319	FW320	FW321	FW322	FW323	FW324	FW325	FW326	FW327	FW328	FW329	FW330	FW331	FW332	FW333	FW334	FW335	FW336	FW337	FW338	FW339	FW340	FW341	FW342	FW343	FW344	FW345	FW346	FW347	FW348	FW349	FW350	FW351	FW352	FW353	FW354	FW355	FW356	FW357	FW358	FW359	FW360	FW361	FW362	FW363	FW364	FW365	FW366	FW367	FW368	FW369	FW370	FW371	FW372	FW373	FW374	FW375	FW376	FW377	FW378	FW379	FW380	FW381	FW382	FW383	FW384	FW385	FW386	FW387	FW388	FW389	FW390	FW391	FW392	FW393	FW394	FW395	FW396	FW397	FW398	FW399	FW400	FW401	FW402	FW403	FW404	FW405	FW406	FW407	FW408	FW409	FW410	FW411	FW412	FW413	FW414	FW415	FW416	FW417	FW418	FW419	FW420	FW421	FW422	FW423	FW424	FW425	FW426	FW427	FW428	FW429	FW430	FW431	FW432	FW433	FW434	FW435	FW436	FW437	FW438	FW439	FW440	FW441	FW442	FW443	FW444	FW445	FW446	FW447	FW448	FW449	FW450	FW451	FW452	FW453	FW454	FW455	FW456	FW457	FW458	FW459	FW460	FW461	FW462	FW463	FW464	FW465	FW466	FW467	FW468	FW469	FW470	FW471	FW472	FW473	FW474	FW475	FW476	FW477	FW478	FW479	FW480	FW481	FW482	FW483	FW484	FW485	FW486	FW487	FW488	FW489	FW490	FW491	FW492	FW493	FW494	FW495	FW496	FW497	FW498	FW499	FW500	FW501	FW502	FW503	FW504	FW505	FW506	FW507	FW508	FW509	FW510	FW511	FW512	FW513	FW514	FW515	FW516	FW517	FW518	FW519	FW520	FW521	FW522	FW523	FW524	FW525	FW526	FW527	FW528	FW529	FW530	FW531	FW532	FW533	FW534	FW535	FW536	FW537	FW538	FW539	FW540	FW541	FW542	FW543	FW544	FW545	FW546	FW547	FW548	FW549	FW550	FW551	FW552	FW553	FW554	FW555	FW556	FW557	FW558	FW559	FW560	FW561	FW562	FW563	FW564	FW565	FW566	FW567	FW568	FW569	FW570	FW571	FW572	FW573	FW574	FW575	FW576	FW577	FW578	FW579	FW580	FW581	FW582	FW583	FW584	FW585	FW586	FW587	FW588	FW589	FW590	FW591	FW592	FW593	FW594	FW595	FW596	FW597	FW598	FW599	FW600	FW601	FW602	FW603	FW604	FW605	FW606	FW607	FW608	FW609	FW610	FW611	FW612	FW613	FW614	FW615	FW616	FW617	FW618	FW619	FW620	FW621	FW622	FW623	FW624	FW625	FW626	FW627	FW628	FW629	FW630	FW631	FW632	FW633	FW634	FW635	FW636	FW637	FW638	FW639	FW640	FW641	FW642	FW643	FW644	FW645	FW646	FW647	FW648	FW649	FW650	FW651	FW652	FW653	FW654	FW655	FW656	FW657	FW658	FW659	FW660	FW661	FW662	FW663	FW664	FW665	FW666	FW667	FW668	FW669	FW670	FW671	FW672	FW673	FW674	FW675	FW676	FW677	FW678	FW679	FW680	FW681	FW682	FW683	FW684	FW685	FW686	FW687	FW688	FW689	FW690	FW691	FW692	FW693	FW694	FW695	FW696	FW697	FW698	FW699	FW700	FW701	FW702	FW703	FW704	FW705	FW706	FW707	FW708	FW709	FW710	FW711	FW712	FW713	FW714	FW715	FW716	FW717	FW718	FW719	FW720	FW721	FW722	FW723	FW724	FW725	FW726	FW727	FW728	FW729	FW730	FW731	FW732	FW733	FW734	FW735	FW736	FW737	FW738	FW739	FW740	FW741	FW742	FW743	FW744	FW745	FW746	FW747	FW748	FW749	FW750	FW751	FW752	FW753	FW754	FW755	FW756	FW757	FW758	FW759	FW760	FW761	FW762	FW763	FW764	FW765	FW766	FW767	FW768	FW769	FW770	FW771	FW772	FW773	FW774	FW775	FW776	FW777	FW778	FW779	FW780	FW781	FW782	FW783	FW784	FW785	FW786	FW787	FW788	FW789	FW790	FW791	FW792	FW793	FW794	FW795	FW796	FW797	FW798	FW799	FW800	FW801	FW802	FW803	FW804	FW805	FW806	FW807	FW808	FW809	FW810	FW811	FW812	FW813	FW814	FW815	FW816	FW817	FW818	FW819	FW820	FW821	FW822	FW823	FW824	FW825	FW826	FW827	FW828	FW829	FW830	FW831	FW832	FW833	FW834	FW835	FW836	FW837	FW838	FW839	FW840	FW841	FW842	FW843	FW844	FW845	FW846	FW847	FW848	FW849	FW850	FW851	FW852	FW853	FW854	FW855	FW856	FW857	FW858	FW859	FW860	FW861	FW862	FW863	FW864	FW865	FW866	FW867	FW868	FW869	FW870	FW871	FW872	FW873	FW874	FW875	FW876	FW877	FW878	FW879	FW880	FW881	FW882	FW883	FW884	FW885	FW886	FW887	FW888	FW889	FW890	FW891	FW892	FW893	FW894	FW895	FW896	FW897	FW898	FW899	FW900	FW901	FW902	FW903	FW904	FW905	FW906	FW907	FW908	FW909	FW910	FW911	FW912	FW913	FW914	FW915	FW916	FW917	FW918	FW919	FW920	FW921	FW922	FW923	FW924	FW925	FW926	FW927	FW928	FW929	FW930	FW931	FW932	FW933	FW934	FW935	FW936	FW937	FW938	FW939	FW940	FW941	FW942	FW943	FW944	FW945	FW946	FW947	FW948	FW949	FW950	FW951	FW952	FW953	FW954	FW955	FW956	FW957	FW958	FW959	FW960	FW961	FW962	FW963	FW964	FW965	FW966	FW967	FW968	FW969	FW970	FW971	FW972	FW973	FW974	FW975	FW976	FW977	FW978	FW979	FW980	FW981	FW982	FW983	FW984	FW985	FW986	FW987	FW988	FW989	FW990	FW991	FW992	FW993	FW994	FW995	FW996	FW997	FW998	FW999	FW1000	FW1001	FW1002	FW1003	FW1004	FW1005	FW1006	FW1007	FW1008	FW1009	FW1010	FW1011	FW1012	FW1013	FW1014	FW1015	FW1016	FW1017	FW1018	FW1019	FW1020	FW1021	FW1022	FW1023	FW1024	FW1025	FW1026	FW1027	FW1028	FW1029	FW1030	FW1031	FW1032	FW1033	FW1034	FW1035	FW1036	FW1037	FW1038	FW1039	FW1040	FW1041	FW1042	FW1043	FW1044	FW1045	FW1046	FW1047	FW1048	FW1049	FW1050	FW1051	FW1052	FW1053	FW1054	FW1055	FW1056	FW1057	FW1058	FW1059	FW1060	FW1061	FW1062	FW1063	FW1064	FW1065	FW1066	FW1067	FW1068	FW1069	FW1070	FW1071	FW1072	FW1073	FW1074	FW1075	FW1076	FW1077	FW1078	FW1079	FW1080	FW1081	FW1082	FW1083	FW1084	FW1085	FW1086	FW1087	FW1088	FW1089	FW1090	FW1091	FW1092	FW1093	FW1094	FW1095	FW1096	FW1097	FW1098	FW1099	FW1100	FW1101	FW1102	FW1103	FW1104	FW1105	FW1106	FW1107	FW1108	FW1109	FW1110	FW1111	FW1112	FW1113	FW1114	FW1115	FW1116	FW1117	FW1118	FW1119	FW1120	FW1121	FW1122	FW1123	FW1124	FW1125	FW1126	FW1127	FW1128	FW1129	FW1130	FW1131	FW1132	FW1133	FW1134	FW1135	FW1136	FW1137	FW1138	FW1139	FW1140	FW1141	FW1142	FW1143	FW1144	FW1145	FW1146	FW1147	FW1148	FW1149	FW1150	FW1151	FW1152	FW1153	FW1154	FW1155	FW1156	FW1157	FW1158	FW1159	FW1160	FW1161	FW1162	FW1163	FW1164	FW1165	FW1166	FW1167	FW1168	FW1169	FW1170	FW1171	FW1172	FW1173	FW1174	FW1175	FW1176	FW1177	FW1178	FW1179	FW1180	FW1181	FW1182	FW1183	FW1184	FW1185	FW1186	FW1187	FW1188	FW1189	FW1190	FW1191	FW1192	FW1193	FW1194	FW1195	FW1196	FW1197	FW1198	FW1199	FW1200	FW1201	FW1202	FW1203	FW1204	FW1205	FW1206	FW1207	FW1208	FW1209	FW1210	FW1211	FW1212	FW1213	FW1214	FW1215	FW1216	FW1217	FW1218	FW1219	FW1220	FW1221	FW1222	FW1223	FW1224	FW1225	FW1226	FW1227	FW1228	FW1229	FW1230	FW1231	FW1232	FW1233	FW1234	FW1235	FW1236	FW1237	FW1238	FW1239	FW1240	FW1241	FW1242	FW1243	FW1244	FW1245	FW1246	FW1247	FW1248	FW1249	FW1250	FW1251	FW1252	FW1253	FW1254	FW1255	FW1256	FW1257	FW1258	FW1259	FW1260	FW1261	FW1262	FW1263	FW1264	FW1265	FW1266	FW1267	FW1268	FW1269	FW1270	FW1271	FW1272	FW1273	FW1274	FW1275	FW1276	FW1277	FW1278	FW1279	FW1280	FW1281	FW1282	FW1283	FW1284	FW1285	FW1286	FW1287	FW1288	FW1289	FW1290	FW1291	FW1292	FW1293	FW1294	FW1295	FW1296	FW1297	FW1298	FW1299	FW1300	FW1301	FW1302	FW1303	FW1304	FW1305	FW1306	FW1307	FW1308	FW1309	FW1310	FW1311	FW1312	FW1313	FW1314	FW1315	FW1316	FW1317	FW1318	FW1319	FW1320	FW1321	FW1322	FW1323	FW1324	FW1325	FW1326	FW1327	FW1328	FW1329	FW
--	---	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	-------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	--------	----

1	10	20	30	40	50	59
HCA IX	NAPLCSPWLPELLIPAPAPGLTVQILL SLLLMPVHPQLPRMQ-EDSPLGGSSSGEDDP					
MCA IX	NASLGPSWAPLESTPAP-- --TAQUILLFLIQVSAQPGQLSGMQGEPS-LGDSSSSGEDE-					
61	70	80	90	100	110	
HCA IX	LGEED-LPSEEDSPREE-DPPGEEDLPGEEDLPGEEDLPGEEDLPGEEDLP					
MCA IX	LGV-DVLPSEEDAP-EEADPP--D--GE-D-P-- --PEVN--SEDRMEE-SLGLLEDLS					
120	130	140	150	160	170	
HCA IX	TVEAPGDPQEP-QNNAHRD-KEGDDQSHWRYGGDPWPRVSPACAGRFQSPVDIRPQLA-					
MCA IX	TPEAP-- --EHSQGS-HGDEKGGGH-SHWSYGGTLLWPQVSPACAGRFQSPVDIR--LER					
180	190	200	210	220	230	
HCA IX	-AFCPALRPLELLGFLPPLPELRUNNGHSVQLTLPGLLENALGPGRREYRALQLHLHWG					
MCA IX	TAFCRTLOPLELLGYELQPLPELSLSNNGHTVQLTLPGLKNALGPQGEYRALQLHLHWG					
240	250	260	270	280	290	
HCA IX	AAGRPGSENTVEGHRFPAAEIHVVHLSLTAFAFVDEALGRPGGLAVLAALFLEEGPEENSAYE					
MCA IX	TSDHPGSENTVNGHRFPAAEIHVVHLSLTAFAFVDEALGRPGGLAVLAALFLEEGPEENSAYE					
300	310	320	330	340	348	
HCA IX	QLLSRL EEIAEEGS--ETQVPGLDISALLPSDFSRYFQYEGSLTTPPCAQGVITWTFENQT					
MCA IX	QLLSHLEETS EEGSKIE--IPGLDVSAALLPSDFSRYRYRYEGSLTTPPCSQGVITWTFENET					
360	370	380	390	400	408	
HCA IX	VMLS AKQLHTLSDTLWGPGRDSRLQLNFRATQPLNGRVIEASFPAGVDS SPRAAEPVQLNS					
MCA IX	VKLS AKQLHTLSVSLWGPGRDSRLQLNFRATQPLNGRVIEASFPAGVDS SP-- --EPVHVNS					
420	430	440	450	459		
HCA IX	CLAGDILALVFGLLFAVTSVAFVLMRRQHR-- --GTKGGVSYRPAEVAETGA					
HCA IX	CFTAGDILALVFGLLFAVTSIAFLQLRRQHRHS GTKDRVSYS PAEMTETGA					

FIG. 59




FIG. 60





FIG. 61



- Homology > 60%
- 5 species: cynomolgus monkey, rat, mouse, guinea pig, hamster

FIG. 62




FIG. 63





FIG. 64


- Vs Skrc-59 engineered cell lines (6 species)
- Effector: target ratio = 10:1



- Human, monkey > hamster, mouse > rat > guinea pig
- More sensitive than binding assay


FIG. 65

- Vs Skrc-59 engineered cell lines (4 species)
- Effector: target ratio = 10:1



- **Mouse > monkey > rat, human**
→ All potential models for B7
→ Candidate model for B7-GGGGS3-G36 = cynomolgus monkey

FIG. 66



- B7-linker-G36: safety optimization required
→ fine tuning anti-CAIX affinity
→ 20 scFvs: different epitopes + affinities

- Effector: target ratio = 10:1 (- = 2:1)

FIG. 67




FIG. 68





FIG. 69




FIG. 70




FIG. 71




FIG. 72




FIG. 73




FIG. 74




FIG. 75

78/92

HC+L2

	FR1-IMGT (1-26)	CDR3-IMGT (27-38)	FR2-IMGT (39-55)	CDR2-IMGT (56-85)	FR3-IMGT (66-104)	CDR3-IMGT
IGKV3-21*05 F B9_R	QVGLVSSGGLVSPGSLRLSCAAS QVGLVSSGGLVSPGSLRLSCAAS	QPTF...SDYF QPTF...SDYF	KSHIQAPGKGLHRYVY KSHIQAPGKGLHRYVY	KSSS...SSYF KPSR...HRET	NYADSVK.GRFTTIRGMKNSILYLDNLSLRAEDTAVYIC NYADSVK.GRFTTIRGMKNSILYLDNLSLRAEDTAVYIC	AK AKHKESTLLDAFEL
IGKV1-6*03 F B9_L	DIQMTQSPSTLSASVGRVTITCKAS DIQMTQSPSTLSASVGRVTITCKAS	QSI...SSN QSI...SSN	LAWYQKPGKGLHRYVY LAWYQKPGKGLHRYVY	KA...SS KA...SS	SLDSGVP.SRPSSGG...SGTDTFLTSSLOPDPFRTYIC SLDSGVP.SRPSSGG...SGTDTFLTSSLOPDPFRTYIC	QQNSYS QHPNRYFAT
IGKV3-23*04 F B7_R	EVQLVSSGGLVSPGSLRLSCAAS EVQLVSSGGLVSPGSLRLSCAAS	QPTF...SDYA QPTF...SDYA	KSHYQAPGKGLHRYVY KSHYQAPGKGLHRYVY	KSSS...GSET KSSS...DRET	YVADSVK.GRFTTIRGMKNSILYLDNLSLRAEDTAVYIC YVADSVK.GRFTTIRGMKNSILYLDNLSLRAEDTAVYIC	AK AKGIIIDVTGSEFCE
IGLV6-57*01 F B7_L	KRWLTAPHS.VSESPKRTVTISCTRS KRWLTAPHS.VSESPKRTVTISCTRS	SGSI...ASNY SGSI...ASNY	VQWYQKPGKGLHRYVY VQWYQKPGKGLHRYVY	ED...N ED...N	QPSGVP.DRFSGLSSGSSNSLSLIGLNTVEEDAVYIC QPSGVP.DRFSGLSSGSSNSLSLIGLNTVEEDAVYIC	QSYGSSN QSLDSCHRRV
IGKV1-63*06 F B3_R	QVQLVSSGGLVSPGSLRLSCAAS QVQLVSSGGLVSPGSLRLSCAAS	QPTF...SDYA QPTF...SDYA	KSHYQAPGKGLHRYVY KSHYQAPGKGLHRYVY	KSSS...GSET KSSS...DRET	YVADSVK.GRFTTIRGMKNSILYLDNLSLRAEDTAVYIC YVADSVK.GRFTTIRGMKNSILYLDNLSLRAEDTAVYIC	AK AKHKESTLLDAFEL
IGKV4-1*01 F B3_L	DIQMTQSPSTLSASVGRVTITCKAS DIQMTQSPSTLSASVGRVTITCKAS	QSI...SSN QSI...SSN	LAWYQKPGKGLHRYVY LAWYQKPGKGLHRYVY	KA...SS KA...SS	SLDSGVP.SRPSSGG...SGTDTFLTSSLOPDPFRTYIC SLDSGVP.SRPSSGG...SGTDTFLTSSLOPDPFRTYIC	QQNSYS QHPNRYFAT

FIG. 76

79/92

80+LC

	FR1-IMGT (1-26)	CDR1-IMGT (27-38)	FR2-IMGT (39-55)	CDR2-IMGT (56-65)	FR3-IMGT (66-104)	CDR3-IMGT
IGHV3-23*04 F A20_H	EYGLVESGG, GLVQPGSSLRISCAS QVQLVQSGG, GLVQPGCELRLSCAS	GFVF, ..., SSYA GFVF, ..., SSYA	MSVYSQAPGKGLRWVA MSVYQAPGKGLRWEL	TSGS, ...GGST TSGS, ...GGSR	YFADGVK, GRFTISGRKSNNTLYIQMWSLRASDTAVVYC YFADSVK, GRFTISGRKSNNTLYIQMWSLRASDTAVVYC	AZ ARCGSGHGMIV
IGLV1-4*01 F A20_L	QSVLTQPPS, ASGTPGKNTICGSS QVQLTQPPS, ASGTPGKNTICGSS	SSNI, ..., GSNY SSNI, ..., GSNY	VYVYQQLPGTAPKLLY VYVYQQLPGTAPKLLY	RS, ..., ...N RS, ..., ...N	QRPSGVP, DRFGSK, ...SGTASLAIHSLRSEDEADYYC QRPSGVP, DRFGSK, ...SGTASLAIHSLRSEDEADYYC	AAMHSLSG AAMDSELNLY
IGHV3-23*04 F B5_H	EYGLVESGG, GLVQPGSSLRISCAS QVQLVQSGG, GLVQPHSSLRISCAS	GFVF, ..., SSYA GFVF, ..., SSYA	MSVYSQAPGKGLRWVA MSVYQAPGKGLRWAT	TSGS, ...GGST XSGS, ...GHT	YFADSVK, GRFTISGRKSNNTLYIQMWSLRASDTAVVYC YFADSVK, GRFTIADNRKSNNTLYIQMWSLRASDTAVVYC	AZ ARSLYGTGSGFDS
IGLV3-10*01 F B5_L	SYELTQPPS, VSVSPQTRITCGSD SYELTQPPS, VSVSPQTRITCGSD	ALP, ..., ...KRY ALP, ..., ...KRY	AYWYQSGQAPVLYY AYWYQSGQAPVLYME	ED, ..., ...S ED, ..., ...S	KRPSGIP, ERFGSS, ...SGTATLITISCAQVEADYYC KRPSGIP, ERFGSS, ...SGTATLITISCAQVEADYYC	YSTDSGSH YSTDSGSHIV
IGHV3-56*02 F B8_H	EYGLVESGG, GLVQPGSSLRISCAS EYGLVESGG, GNVQPGSLRLSCAS	GFVF, ..., GSNY GFVF, ..., STSH	MSVYSQAPGKGLRWVY MSVYQAPGKGLRWLSG	IYSS, ...GST EDSG, ...GKT	YFADSVK, GRFTISGRKSNNTLYIQMWSLRASDTAVVYC YFADSVK, GRFTIADNRKSNNTLYIQMWSLRASDTAVVYC	AZ ARAPSPFDSSGFDATCI
IGLV10-54*01 F B8_L	QSVLTQPPS, VSKGLAQTRITCTON SYELTQPPS, VSKGLAQTRITCTON	SNWV, ..., GRQG SNWV, ..., GRQG	AAMVQSGHPPKLLSY AAMVQSGHPPKLLSY	RS, ..., ...N RS, ..., ...N	KRPSGIS, EEELASR, ...SGTASLITITGLQPSDEADYYC KRPSGIS, ERFGSK, ...SGTASLITITGLQPSDEADYYC	SAWDSLSA SAWDSLSAW

FIG. 76 con't

80/92




FIG. 77

HC+LC

	FRI-1MST (1-26)	CUR1-1MST (27-38)	FRI2-1MST (39-55)	CUR2-1MST (56-65)	FRI3-1MST (66-104)	CUR3-1MST
IGHV1-69*05_F FDL1_42_H	QVQLVQSGA.EVKKPQSSVYSCKAS QVQLVQSGA.EVKKPQSSVYSCKAS	GGTF...SSYA GGTF...SSYA	ISWVRQAPQGLEMMGG ISWVRQAPQGLEMMGG	IIPI..PSTA IIPI..PSTA	NYAQRQ.GRVTTADKSTSTAYMELSLRSEDTAVYIC NYAQRQ.GRVTTADKSTSTAYMELSLRSEDTAVYIC	NR ARQRQFQACIDF
IGHV6-57*02_F FDL1_42_L	HPMLTQPHS.VSEERKNTVTISCTGS HPMLTQPHS.VSEERKNTVTISCTGS	SGSI...ASNY SGSI...DSNI	VQWQQRPASAPTVIY VQWQQRPASAPTVIY	ED.....N ED.....N	QRFSGVF.DRFSGHDSUSASLTISGNTEDPADYIC QRFSGVF.DRFSGHDSUSASLTISGNTEDPADYIC	QSYQEN QSYDNNRHVI
IGHV3-2*01_F FDL14P3_H	EVQLVESGG.SLVQPSRSLRLSCAAS QVQAVQSGG.GLVQPSRSLRLSCAAS	GGTF...DQYA GGTF...DQFA	MHWVRQAPKGLKLVVCG MHWVRQAPKGLKLVVCG	ISNN..SGSI ISNN..SGSI	SYAIVSYK.GRFTTIRDAKASLIYDQMSLRARQETALLYIC GLMDSVK.GRFTVSRDWAASLIYQWMSLRARQETAVYIC	AKD ASTDSKISYYGHHV
IGHV1-24*01_F FDL14P3_L	QSVLTQPHS.ASGTPQRTVTISCSGS QSVLTQPHS.ASGTPQRTVTISCSGS	SGNI...GSNI SGNI...GSNI	VHWYQGLPCTAPKLLIY VHWYQGLPCTAPKLLIY	SN.....N ED.....N	QRFSGVF.DRFSGK..SGTSSASLAIISGLQSEDEADYIC QRFSGVF.DRFSGK..SGTSSASLAIISGLQSEDEADYIC	AAWDSLNG AAWDSLNGRQV

FIG. 78

80*LC

	461-1MGT (1-26)	461-1MGT (27-38)	462-1MGT (39-55)	462-1MGT (56-66)	463-1MGT (66-104)	463-1MGT
IGHV3-23*04_F G17_H	EVQIVESGG,CLVQPGSLRLSCAAS QVQLVQSGG,CLVQPGSLRLSCAAS	QPTF...SGLA QPTF...SSYA	MHWVQAPFGKLEWVSA MSWVAQAPFGKLEWVSA	ISGS...GGST ISGS...GGST	YYADSVK,GRFTISRNSKNTLYIQMNSLAEETAVYYC YYADSVK,GRFTISRNSKNTLYIQMNSLAEETAVYYC	AK ATYGDHGSLEDS
IGHV1-40*01_F G17_L	QSVLTQPPS,VSGAPGQRTTISCOS QSVLTQPPS,VSGAPGQRTTISCOS	SSNIG...AGVD SSNIG...AGVD	VHWYQLPGRAPLLLY VHWYQLPGRAPLLLY	GR.....S AK.....M	MRPSGVP,DRFSOSK...SOTSRSLAITGLQAEDEAATYC MRPSGVP,DRFSOSK...SOTSRSLAITGLQAEDEAATYC	QSYDSGLSS QSYDSGLRHW
IGHV3-30*04_F G25_H	QVQLVESGG,GVVQPGSLRLSCAAS QVTLKESGG,GVVQPGSLRLSCAAS	QPTF...SSYA QPTF...SNLA	MHWVQAPFGKLEWVAV MTWVQAPFGKLEWVGL	ISYC...GSNK ISYD...GSPT	YYADSVK,GRFTISRNSKNTLYIQMNSLAEETAVYYC HYTDSVR,GRFTISRDRAKSLYIQMTLRAQDTAVYYC	KA ARSSGYQRH
IGHV3-21*02_F G28_L	SYVLTAQPPS,VSVAPGQRTARTCSN LEVLTQPPS,VSVAPGQRTARTCSN	NIG.....SKS NIG.....SKS	VHWYQAPFGQAPLVVY VHWYQAPFGQAPLVVY	ED.....S ED.....S	DRPSGIP,DRFSNSN...SMTPLTITSHVPAAGDEAATYC DRPSGIP,DRFSNSN...SMTPLTITSHVPAAGDEAATYC	QVWDSSSSK QVWDSSTCHRY

FIG. 79




FIG. 80

84/92

209964 Homoap IGV1-44*01 F HTGTH18-E1-C3 (T161)	FR1-1MGT (1-26)	CDR1-1MGT (27-38)	FR2-1MGT (39-55)	CDR2-1MGT (56-65)	FR3-1MGT (66-104)	CDR3	FW4
QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC AP QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC APDGLMFLTQVTFDY MCGGLATVPS							
217654 Homoap IGV1-44*01 F HTGTH18-E1-C3 (T161)	FR1-1MGT (1-26)	CDR1-1MGT (27-38)	FR2-1MGT (39-55)	CDR2-1MGT (56-65)	FR3-1MGT (66-104)	CDR3	FW4
QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC AP QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC APDGLMFLTQVTFDY MCGGLATVPS							
217652 Homoap IGV1-44*02 F HTGTH18-E2-S7 (T166)	FR1-1MGT (1-26)	CDR1-1MGT (27-38)	FR2-1MGT (39-55)	CDR2-1MGT (56-65)	FR3-1MGT (66-104)	CDR3	FW4
QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC AP QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC APDGLMFLTQVTFDY MCGGLATVPS							
217654 Homoap IGV1-44*01 F HTGTH18-E2-S7 (T166)	FR1-1MGT (1-26)	CDR1-1MGT (27-38)	FR2-1MGT (39-55)	CDR2-1MGT (56-65)	FR3-1MGT (66-104)	CDR3	FW4
QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC AP QVQLVQSGEGLVVPGRGASVSRCAE GYTF...SYG LHWYQAFGQSDLEMMH ISAV...NNT NYRQSLQ.GEVNTTTFHTSTAYRELSLSHSHITAVYC APDGLMFLTQVTFDY MCGGLATVPS							

FIG. 81

[illegible]

Sequence ID	V-GENE and allele	J-GENE and allele	D-GENE and allele	FR1-MST	COR1-MST	FR2-MST	CDK2-MST	FR3-MST	COR3-MST	FR4-MST
NTGTHS-E1C3 (T131)	HY1-18701 F	HJ702 F	HE3-10701 F	QVQLVQ33GA EVKHPGASVKVSQAS	GYTF...TSYG	ISWYRQAPGGGLEWNGW	ISAY...AGNT	NYAQNLQ GRVTRITDTISTATYAEIRLSRSDDTAVYYC	ARQPGIWFGLTHETFFDY	WGQGTLLTVSS
	HY3-74702 F	HJ702 F	HQ3-9701 F	EVQLVQ33GG GLVKPGGSLRLSCAS	QYTF...SDYS	ASDVRQAPGGGLEWVSR	WSD...GSRT	NYADSVK SPFTSRDMMNNLLYLIRNSLRADETAMYYC	ARQPSFFFDI	WGQGTLLTVSS
	HY5-51701 F	HJ702 F		EVQLVQ33GA EVKHPGGSLSKRCAS	QYSF...TNYW	ISWYRQAPGGGLEWMAI	HPY...NSRT	YSPSFQGSQVTSVQKS/ITAYLQWLSLKASDTAMYYC	ARYYYVAMEV	WGQGTLLTVSS
P5-H12	HY1-18701 F	HJ702 F	HQ6-13701 F	EVQLVQ33GA EVKHPGASVKVSQAS	GYTF...TNYG	ISWYRQAPGGGLEWNGW	YUNN...NGWI	NYAQNL GRVTRITDTISTATYAEIRLSRSDDTAVYYC	ARGLFSRVPILWDP	WGQGTLLTVSS
P6-G7	HY1-59701 F	HJ702 F	HQ6-12701 F	QVQLDQ33GA EVKHPGSSVKVSQAS	GGTF...SSVA	ISWYRQAPGGGLEWMAIG	ILPL...FGST	NYACKFQGRLLIADESTRTVYLEHSLTSEDITAVYYC	ARGEDVAPSNSSGFQV	WGQGTLLTVSS
86/92										
NTGTHS-E1C3 (T131)	LV1-44701 F	LJ702 F		SYELTOPPSASSTPGQRTVTHSCGS	SSNI...GSNT	VSWYQGLPGTAPKLLIY	RNI...N	QRPSQIP DRFSGSK SGTSAQLAKGLDSEDEADYYC	AAWDSSRSQVY	FGGGRLLTVL
	LV1-44701 F	LJ701 F		SYELTOPPSASGTPGQRTVTHSCGS	RSM...GRNS	VWYQGLPGTAPKLLIY	SN...N	QRPSNP GRFSGR SGTSAQLAKGLDSEDEDTYYC	AAWDARLTQPL	F33
	LV1-44701 F	LJ702 F		LPALTOPPSASGTPGQRTVTHSCGS	SSNI...GSNT	VWYQGLPGTAPKLLIY	RRI...N	QRPSQIP DRFSGST SGTSAQLAKGLDSEDEADYYC	EAKWDSLNGPV	FGG
P5-E10	LV2-14701 F	LJ701 F		QSALTQPPASVSGSPGSHITCTST	SSDVG...GVNY	VSWYQDHPGAPKLLIY	EV...T	ERPSQNS NAFSSSK SONTASLTSGLOADEDESDYYC	SSYTRSSSTSYVY	F33
P6-H12	LV10-59701 F	LJ702 F		SYELTOPPSVSKGLKQTATLTCTGN	SNHV...QMGG	AAWYQDHPGAPKLLIY	RRI...D	NRPSSISERRSASR SONTASLTSGLOPEDEADYYC	SAYDRELNAVY	FGG

FIG. 81 CON'T

The anti-PD-L1 amino acid sequences

VH

Sequence ID	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
40 mut	QVQLVQSGAEVKKPGSSVKVCKAS	GGTFSSYA	ISWVRQAPGQGLEWMGG	IIPIFGTA	NYAQKFGGRVTITADKSTSTAYMELSLRSED	ARGRMFGAGIDF	WGPGTLTVSS
50-6B6.1 mut	QVQLVQSGGEVKKPGASVKVCKAS	GYTLSSHG	ITWVRQAPGQGLEWMGW	ISAHNGHA	SNAQKVEDRVTMTTDTSTNTAYMELRSLT	ARVHAALYYGMDV	WGQGTLLTVSS
50-6B6.2	QVQLVQSGGEVKKPGASVKVCKAS	GYTLSSHG	ITWVRQAPGQGLEWMGW	ISAHNGHA	SNAQKVEDRVTMTTDTSTNTAYMELRSLT	ARVHAALYYGMDV	WGQGTLLTVSS
50-7B3	QVQLVQSGGEVKKPGASVKVCKAS	GYTLSSHG	ITWVRQAPGQGLEWMGW	ISAHNGHA	SNAQKVEDRVTMTTDTSTNTAYMELRSLT	ARVHAALYYGMDV	WGQGTLLTVSS
50-5B9	QVQLVQSGGEVKKPGASVKVCKAS	GYTLSSHG	ITWVRQAPGQGLEWMGW	ISAHNGHA	SNAQKVEDRVTMTTDTSTNTAYMELRSLT	ARVHAALYYGMDV	WGQGTLLTVSS

VL

Sequence ID	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
40 mut	RFRLTQPHSVSESPGKTVTISCTRS	SGSIDSNY	VQWYQQRPQGSAPTIVY	EDN	QRPSGVDPDRFSGSIDSSNSASLTISGLKTE	QSYDSNNRHVI	FGGGTKLTVL
50-6B6.1 mut	SYELTQPPSVSLAPGQSARISCGGD	NIGSKG	VHWYQKPGQAPVAVVY	DDR	DRPSGIPERFSGSNSGNTATLTISRVEAGDE	QVWDSGSDHWV	FGGGTKLTVL
50-6B6.2	LPVLTQPPSVSAAPGQTARISCGGS	NIGDKG	VHWYQKPGQAPVAVVY	DDS	DRPSGIPERFSGSNSGNTATLTISRVEAGDE	QVWDSGSDHWV	FGGGTKLTVL
50-7B3	SYELTQPPSVSVAPGQTARITCGGN	NIGNKG	VHWYQKPGQAPVAVVY	DDS	DRPSGIPERFSGSNSGNTATLTISRVEAGDE	QVWDSGSDHWV	FGGGTKLTVL
50-5B9	LPVLTQPPSVSVVAGQTARITCRGN	NIGGKG	VHWYQKPGQAPVAVVY	DDY	SRRSGIPERFSGSHSGSAATLTISRVEAGDE	QVWDSGSDHWV	FGGGTKLTVL

FIG. 82

Ab P4-B3 Variable Region nucleic acid sequences
V _H chain of Ab P4-B3 VH (IGHV3-9*01)
CAGGTGCAGCTGGTGCAGTCTGGGGGAGGCTTGGTACAGCCTGGCAGGTCCCTGAGAC TCTCCTGTGCAGCCTCTGGATTACCTTTGATGATTATGCCATGCACTGGGTCCGGCAAG CTCCAGGGAAGGGCCTGGAGTGGGTCTCAGGTATTAGTTGGAATAGTGGTAGCATAGG CTATGCGGACTCTGTGAAGGGCCGATTACCGTCTCCAGAGACAACGCCAAGAAGTCA CTGTATCTGCAAATGAACAGCCTGAGAGCCGAGGACACGGCTGTGTATTACTGTGCGA GTGACTACGGTGACAAATACTACTACTACGGTATGGACGTCTGGGGCAAAGGGACCAC GGTCACCGTCTCCTCA
V _L chain of Ab P4-B3 VL (IGLV1-44*01)
CAGCCTGGGCTGACTCAGCCACCCTCAGCGTCTGGGACCCCCGGGCAGAGGGTCACCA TCTCTTGTCTGGAAGCAGCTCCAACATCGGAAGTAATACTGTCAACTGGTATCAGCAA TTCCCCGGAAAGGCCCCCAAACCTCCTCATCTTTAATGATAATCAGCGGCCCTCAGGGGT CCCTGACCGCTTCTCTGCTTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATTAGTGGCCT CCAGTCTGAGGATGAGGCTGACTATTACTGTGCGGCATGGGATGGCGGTCTGAATGGTC GAGGGGTGTTTCGCGGAGGGACCAAACCTGACCGTCTCA
Ab P4-B3 Variable Region amino acid sequences
V _H chain of Ab P4-B3 VH (IGHV3-9*01)
QVQLVQSGGGLVQPGRSLRLSCAASGFTFDDYAMHWVRQAPGKGLEWVSGISWNSGSIGY ADSVKGRFTVSRDNAKNSLYLQMNSLRAEDTAVYYCASDYGDKYYYYGMDVWGKGTTV TVSS
V _L chain of Ab P4-B3 VL (IGLV1-44*01)
QPGLTQPPSASGTPGQRTVISCSSSSNIGSNTVNWYQQFPGKAPKLLIFNDNQRPSGVPDRF SASKSGTSASLAISGLQSEDEADYYCAAWDGGLNGRGVFGGGTKLTVL
Ab P4-B3 Constant Region nucleic acid sequences – wild type IgG1 monomer
CH1
ACCAAGGGCCCATCGGTCTTCCCCCTGGCACCCCTCCTCCAAGAGCACCTCTGGGGGGCAC AGCGGCCCTGGGCTGCCTGGTCAAGGACTACTTCCCCGAACCGGTGACGGTGTCTGTGGA ACTCAGGCGCCCTGACCAGCGGCGTGACACCTTCCCGGCTGTCTACAGTCCTCAGGA CTCTACTCCCTCAGCAGCGTGGTGACCGTGCCCTCCAGCAGCTTGGGCACCCAGACCTA CATCTGCAACGTGAATCACAAGCCCAGCAACACCAAGGTGGACAAGAAA
Hinge
GCAGAGCCCAAATCTTGTGACAAAACCTCACACATGCCCACCGTGCCCA
CH2
GCACCTGAACTCCTGGGGGGGACCGTCAGTCTTCCTCTTCCCCCAAACCCAAGGACAC CCTCATGATCTCCCGGACCCCTGAGGTACATGCGTGGTGGTGGACGTGAGCCACGAAG ACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAGAC AAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCCTC CTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCC TCCAGCCCCCATCGAGAAAACCATCTCCAAGCCAAA
CH3
GGGCAGCCCCGAGAACCACAGGTGTACACCCTGCCCCCATCCCGGGATGAGCTGACCA AGAACCAGGTACGCCTGACCTGCCTGGTCAAAGGCTTCTATCCAGCGACATCGCCGTG GAGTGGGAGAGCAATGGGCAGCCGGAGAACAACTACAAGACCACGCCTCCCGTGGCTGG ACTCCGACGGCTCCTTCTTCTCTACAGCAAGCTCACCGTGGACAAGAGCAGGTGGCAG

FIG. 83

CAGGGGAACGTCTTCTCATGCTCCGTGATGCATGAGGCTCTGCACAACCACTACACGCA GAAGAGCCTCTCCCTGTCTCCGGGTAAATGA
C _L GGTCAGCCCAAGGCTGCCCCCTCGGTCACTCTGTTCCCGCCCTCCTCTGAGGAGCTTCA AGCCAACAAGGCCACACTGGTGTGTCTCATAAGTGACTTCTACCCGGGAGCCGTGACA GTGGCCTGGAAGGCAGATGGCAGCCCCGTCAAGGCGGGAGTGGAGACCACACACCCT CCAAACAAAGCAACAACAAGTACGCGGCCAGCAGCTATCTGAGCCTGACGCCTGAGCA GTGGAAGTCCCACAGAAGCTACAGCTGCCAGGTACGCATGAAGGGAGCACCGTGGAG AAGACAGTGGCCCCCTACAGAATGTTTCATGA

Ab P4-B3 Constant Region amino acid sequences – wild type IgG1 monomer
CH1 ASTKGPSVFPLAPSSKSTSGGTAALGLCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGL YSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKK
Hinge AEPKSCDKTHTCPPCP
CH2 APELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKP REEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAK
CH3 GQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLDSD GSFFLYSKLTVDKSRWQQGNVFSVMHEALHNHYTQKSLSLSPGK
C _L GOPKAAPSVTLFPPSSEELQANKATLVCLISDFYPGAVTVAWKADGSPVKAGVETTTPSKQ SNNKYAASSYLSLTPEQWKSHRSYSCQVTHEGSTVEKTVAPTECS

Ab P4-B7 Variable Region nucleic acid sequences
V _H chain of Ab P4-B7 VH (IGHV5-51*01) CAGGTGCAGCTGGTGCAGTCTGGAGCAGAGGTGAAGAAGCCCCGGGGAGTCTCTGAAGA TCTCCTGTAAGGATTCTGGATACACCTTTACCACCTACTGGATCGGCTGGGTGCGCCAG CTGCCCCGGGAAAGGCCTGGAGTTGATGGGGATCATCTATCCTGATGACTCTGATACCAC ATACAGCCCGTCCTTCCAAGGCCATGTCACCATCTCAGCCGACAAGTCCATCAACACCG CCTACCTGCAGTGGAGCAGCCTGAAGGCCTCGGACACCGCCATGTATTACTGTGCGTTT TGGGGTGCAGTGGAGCGCCAGTGAATGGTTTGTATCTGGGGCCAAGGCACCCTGG TCACCGTCTCCTCA
V _L chain of Ab P4-B7 VL (IGLV1-44*01) CTGCCTGTGCTGACTCAGCCACCCCTCAGCGTCTGGGACCCCCGGGCAGAGGGTCACCAT CTCCTGCACTGGGAGCAGCTCCAACATCGGGGCAGGTTATGTTGTACACTGGTACCAGC AGCTCCAGGAACGGCCCCCAACTCCTCATCTATAGTAATAATCAGCGGCCCTCAGGG GTCCCTGACCGATTCTCTGGCTCCAAGTCTGGCACCTCAGCCTCCCTGGCCATCAGTGG GCTCCAGTCTGAGGATGAGGCTGATTACTGTGCAGCATGGGATGACAGCCTGAATG CTCCGGTGTTCGGCGGAGGGACCAAGCTGACCGTCTCA

Ab P4-B7 Variable Region amino acid sequences
V _H chain of Ab P4-B7 VH (IGHV5-51*01)

FIG. 83 con't

90/92

QVQLVQSGAEVKKPAGESLKISCKDSGYTFTTYWIGWVRQLPGKGLELMGIIYPDDSDTTYSPSFQGHVTISADKSINTAYLQWSSLKASDTAMYYCAFWGASGAPVNGFDIWGQGLTVTVSS
V_L chain of Ab P4-B7 VL (IGLV1-44*01)
LPVLTQPPSASGTPGQRVTISCTGSSSNIGAGYVVHWYQQLPGTAPKLLIYSNNQRPSGVPRFSGSKSGTSASLAISGLQSEDEADYYCAAWDDSLNAPVFGGGTKLTVLL

PD1#2 Variable Region nucleic acid sequences
V_H chain of Ab PD1#2 VH (IGHV4-61*01)
CAGGTACAGCTGCAGCAGTCAGGCCAGGACTGGTGAGGCCTTCGGCGACCTGTCCCTCACCTGCACTGTCTCTGGTGACTCCGTCAGCAGTGATAATTACTTCTGGAGTTGGATTCCGCAGCCCCCAGGGAAGCCACTGGAGTGGATTGGCTATGTCTATTACAATGGGAACACCAACTACAACCCCTCCTTCAACAGTCGAGTCACCATGTCACCTGACACGTCCAAGAACCA GTTCTCCTTGAAGCTGAGGTCTGTGACCGCCGCGGACACGGCCTTTTATTACTGTGCGA CAGAGACGCCCCCAACCAGCTATTTTAATAGTGGACCTTTGACTCCTGGGGCCAGGGC ACCCTGGTCACCGTCTCCTCG
V_L chain of Ab PD1#2 VL (IGLV10-54*01)
CAGCCTGGGCTGACTCAGCCACCCTCGGTGTCCAAGGGCTTGAGACAGACCGCCACACTCACCTGCACTGGGAGCAGCAACAATGTAGGCGCCACGGAGCAGCTTGGCTGCAGCAGCACCAGGGCCACCCTCCCAAACCTCCTTGCTACAGGAATAACAACCGGCCCTCAGGGATCTCAGAGAGATTCTCTGCATCCAGGTCAGGAAACACAGCCTCCCTGACCATTATTGGAC TCCAGCCTGAGGACGAGGGTGACTATTACTGCTCATCATGGGACAGCAGCCTCAGTGGT TATGTCTTCGGACCTGGGACCAAAGTCACCGTCCTA

Ab PD1#2 Variable Region amino acid sequences
V_H chain of Ab PD1#2 VH (IGHV4-61*01)
QVQLQQSGPGLVRPSATLSLTCTVSGDSVSSDNYFWSWIRQPPGKPLEWIGYVYYNGNTNYPNPSFNSRVTMSLDTSKNQFSKLRSVTAADTAFYYCATETPPTSYPFNSGPFDSWGQGLTVTVSS
V_L chain of Ab PD1#2 VL (IGLV10-54*01)
QPGLTQPPSVSKGLRQTATLTCTGSSNNVGAHGAAWLQQHQGHPPKLLAYRNNNRPSGISE RFSASRSGNTASLTIIGLQPEDEGDYYCSSWDSSLSGYVFGPGTKVTVL

PD1#3 Variable Region nucleic acid sequences
V_H chain of Ab PD1#3 VH (IGHV1-18*01)
CAGGTGCAGCTGGTGCAGTCTGGGGCTGAGGTGAAGAAGCCTGGGTCCTCAGTGAAGG TCTCCTGCAAGACTTCTGGCTACACCTTTAACAGGTTTGGTCTCACCTGGGTGCGACAG GCCCCTGGACAAGGGCTTGAGTGGATGGGATGGACCAACCCTTACAATGGTAACACAA GGTATGCACAGAAGTTCCAGGGCAGAGTCACCATGACCACAGACACATCCACGAGCAC AGCCTACATGGAGCTGAGGAGCCTGAGATCTGACGACACGGCCATGTATTTCTGTGCGA GAGTCGTAGCCGTAAACGGTATGGACGTCTGGGGCCAAGGGACCACGGTCACCGTCTC CTCA
V_L chain of Ab PD1#3 VL (IGLV6-57*01)
AATTTTATGCTGACTCAGCCCCACTCTGTGTCTGGAGTCTCCGGGGAAGACGGTTACCAT

FIG. 83 con't

CTCCTGCACCCGCAACAGTGGCAGCATTGCCGCTACTATGTGCAGTGGTACCAGCAGC
 GCCCCGGGCAGTTCCCCCACCCTGTGATCTATGAAGATAACCAAAGACCCTCTGGGGTC
 CCTGATCGGTTCTCTGGCTCCATCGACAGCTCCTCCAACCTCTGCCTCCCTCACCATCTCT
 GGACTGAAGACTGAGGACGAGGCTGACTACTACTGTGAGTCTTATGATAGCAGCAATCT
 TTGGGTGTTCCGGCGGAGGGACCAAGCTGACCGTCCTA

Ab PD1#3 Variable Region amino acid sequences

V_H chain of Ab PD1#3 VH (IGHV1-18*01)

QVQLVQSGAEVKKPGSSVKVSKTSGYTFNRFGLTWVRQAPGQGLEWMGWNPYNGNT
 RYAQKFQGRVTMTTDTSTSTAYMELRSLRSDDTAMYFCARVAVNGMDVWGQGTITVTV
 SS

V_L chain of Ab PD1#3 VL (IGLV6-57*01)

NFMLTQPHSVSESPGKTVTISCIRNSGSIAAYYVQWYQQRPGSSPTTVIYEDNQRPSGVPDR
 FSGSIDSSSNSASLTISGLKTEADYYCQSYDSSNLWVFGGGTKLTVL

Ab PD1#13 Variable Region nucleic acid sequences

V_H chain of Ab PD1#13 VH (IGHV3-30*01)

GAGGTGCAGCTGGTGCAGTCTGGGGGAGGCGTGGTCCAGCCTGGGAGGTCCCTGAGAC
 TCTCCTGTGCAGCCTCTGGATTACCTTCAGTAGCTATGCTATGCACTGGGTCCGCCAGG
 CTCCAGGCAAGGGGCTGGAGTGGGTGGCAGTTATATCATATGATGGAAGCAATAAATA
 CTACGCAGACTCCGTGAAGGGCCGATTACCATCTCCAGAGACAATTCCAAGAACACG
 CTGTATCTGCAAATGAACAGCCTGAGAGCTGAGGACACGGCTGTGTATTACTGTGCGAG
 CCAAACAGTGGCTGGAAGTGACTACTGGGGCCAGGGCACCTGGTCACCGTCTCCTCA

V_L chain of Ab PD1#13 VL (IGLV1-44*01)

CAGCCTGGGCTGACTCAGCCACCCTCGGTGCCAGTGGCCCCAGGACAGACGGCCAGGA
 TTACCTGTGGGGGAAACAACATTGGAAGTAAAAGTGTGCACTGGTACCAGCAGAAGCC
 AGGCCAGGCCCTGTGCTGGTCGTCTATGATGATAGCGACCGGCCCTCAGGGATCCCTG
 AGCGATTCTCTGGCTCCAACCTCTGGGAACACGGCCACCCTGACCATCAGCAGGGTCGAA
 GCCGGGGATGAGGCCGACTATTACTGTGAGGTGTGGCATAGTGTTAGTGATCAAGGGG
 TCTTCGGAAGTGGGACCAAAGTCACCGTCCTA

Ab PD1#13 Variable Region amino acid sequences

V_H chain of Ab PD1#13 VH (IGHV3-30*01)

EVQLVQSGGGVVPGRSLRLSCAASGFTFSSYAMHWVRQAPGKGLEWVAVISYDGSNKY
 YADSVKGRFTISRDN SKNTLYLQMNSLRAEDTAVYYCASQTVAGSDYWGQGLTVTVSS

V_L chain of Ab PD1#13 VL (IGLV1-44*01)

QPGLTQPPSPVPAPGQTARITCGGNNIGSKSVHWYQQKPGQAPVLVYDDSDRPSGIPERFS
 GSNSGNTATLTISRVEAGDEADYYCQVWHSVSDQGQVFGTGTKVTVL

FIG. 83 con't




FIG. 83 con't