



(86) Date de dépôt PCT/PCT Filing Date: 2006/02/02
 (87) Date publication PCT/PCT Publication Date: 2006/12/07
 (45) Date de délivrance/Issue Date: 2012/09/25
 (85) Entrée phase nationale/National Entry: 2007/11/08
 (86) N° demande PCT/PCT Application No.: US 2006/004011
 (87) N° publication PCT/PCT Publication No.: 2006/130188
 (30) Priorités/Priorities: 2005/05/31 (US60/686,016);
 2005/06/21 (US60/692,439)

(51) Cl.Int./Int.Cl. *A61K 35/74* (2006.01),
A23K 1/00 (2006.01), *A23K 1/18* (2006.01),
A23L 1/30 (2006.01)
 (72) Inventeurs/Inventors:
 BOILEAU, THOMAS WILLIAM-MAXWELL, US;
 KIELY, BARRY PIUS, IE;
 O'MAHONY, LIAM DIARMUID, IE;
 MACSHARRY, JOHN, IE;
 SUNVOLD, GREGORY DEAN, US
 (73) Propriétaires/Owners:
 THE IAMS COMPANY, US;
 ALIMENTARY HEALTH LTD., IE
 (74) Agent: MBM INTELLECTUAL PROPERTY LAW LLP

(54) Titre : BIFIDOBACTERIES DE PROBIOTIQUES FELINS
 (54) Title: FELINE PROBIOTIC BIFIDOBACTERIA

(57) **Abrégé/Abstract:**

According to the invention there is provided a strain of lactic acid bacteria of the genus Bifidobacteria obtainable by isolation from resected and washed feline gastrointestinal tract having a probiotic activity in animals. Methods of use and compositions comprising the Bifidobacteria of the present invention are also provided.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
7 December 2006 (07.12.2006)

PCT

(10) International Publication Number
WO 2006/130188 A1

(51) International Patent Classification:

A61K 35/74 (2006.01) A23L 1/30 (2006.01)
A23K 1/00 (2006.01) A23K 1/18 (2006.01)(74) Common Representative: THE IAMS COMPANY; c/o
Eileen L. Hughett, The Procter & Gamble Company, Win-
ton Hill Business Center, 6110 Center Hill Road, Cincin-
nati, Ohio 45224 (US).

(21) International Application Number:

PCT/US2006/004011

(22) International Filing Date: 2 February 2006 (02.02.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/686,016 31 May 2005 (31.05.2005) US
60/692,439 21 June 2005 (21.06.2005) US(81) Designated States (*unless otherwise indicated, for every
kind of national protection available*): AE, AG, AL, AM,
AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN,
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI,
GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE,
KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV,
LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI,
NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG,
SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US,
UZ, VC, VN, YU, ZA, ZM, ZW.(71) Applicants (*for all designated States except US*): THE
IAMS COMPANY [US/US]; 7250 Poe Avenue, Dayton,
Ohio 45414 (US). ALIMENTARY HEALTH LTD.
[IE/IE]; 7 Pearse Street, Kinsale, County Cork (IE).(84) Designated States (*unless otherwise indicated, for every
kind of regional protection available*): ARIPO (BW, GH,
GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM,
ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM),
European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,
FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT,
RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA,
GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): BOILEAU,
Thomas, William-Maxwell [US/US]; 6072 Rothchester
Drive, Galloway, Ohio 43119 (US). KIELY, Barry,
Pius [IE/IE]; Simla Villa, Passage West, Cork (IE).
O'MAHONY, Liam, Diarmuid [IE/IE]; 41 Maryville Es-
tate, Ballintemple, Cork (IE). MACSHARRY, John
[IE/IE]; 11 Abbotswood Mews, Monastery Road,
Rochestown, Cork (IE). SUNVOLD, Gregory, Dean
[US/US]; 3158 Minnich-wysong Road, Lewisburg, Ohio
45338 (US).

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guid-
ance Notes on Codes and Abbreviations" appearing at the begin-
ning of each regular issue of the PCT Gazette.

(54) Title: FELINE PROBIOTIC BIFIDOBACTERIA

(57) Abstract: According to the invention there is provided a strain of lactic acid bacteria of the genus Bifidobacteria obtainable by isolation from resected and washed feline gastrointestinal tract having a probiotic activity in animals. Methods of use and compositions comprising the Bifidobacteria of the present invention are also provided.



WO 2006/130188 A1

FELINE PROBIOTIC BIFIDOBACTERIA

FIELD OF THE INVENTION

The present invention relates to the field of probiotic micro-organisms, more specifically feline probiotic lactic acid bacteria and methods of use.

BACKGROUND OF THE INVENTION

The defense mechanisms to protect the mammalian gastrointestinal (GI) tract from colonisation by bacteria are highly complex. The GI tract of most mammals are colonised by native microflora, and invasive pathogenic micro-organisms. In a healthy state, these competing microflora are in a state of equilibrium. Modification of the intestinal microflora equilibrium may lead to or prevent many GI disorders, both in humans, and other mammalian species, such as companion animals including cats, dogs and rabbits. The well being of companion animals is closely related to their feeding and GI health, and maintenance of the intestinal microflora equilibrium in these animals may result in healthier companion animals.

The number and composition of the intestinal microflora tend to be stable, although age and diet may modify it. Gastric acidity, bile, intestinal peristalsis and local immunity are factors thought to be important in the regulation of bacterial flora in the small intestine of human beings and various other mammals. Often companion animal GI disorders, including those found in felines, are linked to bacterial overgrowth and the production of enterotoxins by pathogenic bacteria. These factors disrupt the intestinal microflora equilibrium and can promote inflammation and aberrant immune responses.

During the last few years, research has begun to highlight some valuable strains of bacteria and their potential use as probiotic agents. Probiotics are considered to be preparations of bacteria, either viable or dead, their constituents such as proteins or carbohydrates, or purified fractions of bacterial ferments that promote mammalian health by preserving the natural microflora in the GI tract, and reinforcing the normal controls on aberrant immune responses. It is believed by some that probiotic bacteria are more effective when derived from the species, or closely related species, intended to be treated.

Therefore, there is a need for probiotic strains derived from companion animals to be used for companion animals that are different to those derived from humans.

WO 01/90311 discloses probiotic micro-organisms isolated from faecal samples obtained from cats having probiotic activity. However, these bacteria were obtained from faecal samples, and may not form part of the natural intestinal microflora present in the upper portion of the GI tract.

Consequently, there is a need to provide strains of bacteria obtainable by isolation from the natural intestinal microflora present in the upper portion of the GI tract that are particularly adapted for cats, and have been selected for their probiotic properties and ability to survive processing, and to incorporate these strains into compositions that are suitable for their use.

SUMMARY OF THE INVENTION

An object of the present invention is to provide feline probiotic bifidobacteria.

According to the invention there is provided strains of lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract having a probiotic activity in animals. The lactic acid bacterial strains are preferably selected from the species comprising *Bifidobacterium longum*, *Bifidobacterium animalis*, *Bifidobacterium adolescentis*, *Bifidobacterium bifidum*, *Bifidobacterium infantis*, or *Bifidobacterium thermophilum*.

In accordance with an aspect of the present invention, there is provided, a method of maintaining or improving the health of a companion animal comprising orally administering lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract.

In accordance with another aspect of the present invention, there is provided, a composition comprising a strain of lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract having probiotic activity, and a carrier.

In accordance with another aspect of the present invention, there is provided, a method of reducing odor associated with the feces or urine of a companion animal comprising orally administering lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract.

2a

In accordance with another aspect of the present invention, there is provided, a method of reducing odor associated with a litter box of a companion animal comprising orally administering lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract.

In a preferred embodiment, the lactic acid bacterial strain is a strain selected from the group comprising *Bifidobacterium longum* having a 16s-23s spacer region DNA sequence having greater than 95% homology to SEQ. ID NO. 1, or greater than 94% homology to SEQ. ID NO. 2.

In a further preferred embodiment, the lactic acid bacterial strain is selected from the group comprising *Bifidobacterium longum* NCIMB 41290 (*AHF5340*), *Bifidobacterium longum* NCIMB 41291 (*AHF1231*) and mixtures thereof.

Furthermore, the present invention is directed towards providing uses of lactic acid bacteria obtainable by isolation from resected and washed feline gastrointestinal tract for maintaining and improving companion animal health, and compositions comprising the lactic acid bacteria.

These and other features, aspects, and advantages of the present invention will become evident to those skilled in the art from a reading of the present disclosure.

Sequences

SEQ. ID NO. 1 – 16s-23s intergenic spacer nucleotide sequence from *Bifidobacterium longum* NCIMB 41290 (AHF5340).

SEQ. ID NO. 2 – 16s-23s intergenic spacer nucleotide sequence from *Bifidobacterium longum* NCIMB 41291 (AHF1231).

SEQ. ID NO. 3 – Left 16s-23s PCR primer sequence for sequence analysis.

SEQ. ID NO. 4 – Right 16s-23s PCR primer sequence for sequence analysis.

Bacterial Deposit Numbers

The table below indicates *Bifidobacterium* species and strain number for strains that are examples of the present invention. The bacterial strains are deposited with the National Collections of Industrial Food and Marine Bacteria (NCIMB), Aberdeen, UK.

Strain	Deposit Number	16 s-23s Sequence
Bifidobacterium AHF534D	NCIMB 41290	SEQ. ID NO. 1
Bifidobacterium AHF123R	NCIMB 41291	SEQ. ID NO. 2

DETAILED DESCRIPTION OF THE INVENTION

All weights, measurements and concentrations herein are measured at 25°C on the composition in its entirety, unless otherwise specified.

Unless otherwise indicated, all percentages of compositions referred to herein are weight percentages and all ratios are weight ratios.

Unless otherwise indicated, all molecular weights are weight average molecular weights.

Unless otherwise indicated, the content of all literature sources referred to within this text are incorporated herein in full by reference.

Except where specific examples of actual measured values are presented, numerical values referred to herein should be considered to be qualified by the word “about”.

Within the following description, the abbreviation CFU (“colony-forming unit”) designates the number of bacterial cells revealed by microbiological counts on agar plates, as will be commonly understood in the art.

As used herein, the term “mutants thereof” includes derived bacterial strains comprising DNA mutations in other DNA sequences in the bacterial genome excluding the 16s-23s intergenic sequence.

As used herein, the term “DNA mutations” includes natural or induced mutations comprising at least single base alterations including deletions, insertions, transversions, and other DNA modifications known to those skilled in the art, including genetic modification introduced into a parent nucleotide or amino acid sequence whilst maintaining at least 50% homology to the parent sequence. Preferably, the sequence comprising the DNA mutation or mutations has at least 60%, more preferably at least 75%, more preferably still 85% homology with the parental sequence. As used herein, sequence “homology” can be determined using standard techniques known to those skilled in the art. For example, homology may be determined using the on-line homology algorithm “BLAST” program, publicly available at <http://www.ncbi.nlm.nih.gov/BLAST/>.

As used herein “genetic modification” includes the introduction of exogenous and/or endogenous DNA sequences into the genome of an organism either by insertion into the genome of said organism or by vectors including plasmid DNA or bacteriophage as known by one skilled in the art, said DNA sequence being at least two deoxyribonucleic acid bases in length.

As used herein, “companion animal” means a domestic animal. Preferably, “companion animal” means a domestic feline (cat), canine (dog), rabbit, ferret, horse, cow, or the like. More preferably, “companion animal” means a domestic feline.

Lactic Acid Bifidobacteria Strains

The first aspect of the present invention comprises a strain of lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract having probiotic activity in animals. Probiotics are micro-organisms, either viable or dead, processed compositions of micro-organisms, their constituents such as proteins or carbohydrates, or purified fractions of bacterial ferments that beneficially affect a host. The general use of probiotic bacteria is in the form of viable cells. However, it can be extended to non-viable cells such as killed cultures or compositions containing beneficial factors expressed by the probiotic bacteria. This may

include thermally killed micro-organisms, or micro-organisms killed by exposure to altered pH or subjected to pressure. For the purpose of the present invention, "probiotics" is further intended to include the metabolites generated by the micro-organisms of the present invention during fermentation, if they are not separately indicated. These metabolites may be released to the medium of fermentation, or they may be stored within the micro-organism. As used herein "probiotic" also includes bacteria, bacterial homogenates, bacterial proteins, bacterial extracts, bacterial ferment supernatants, and mixtures thereof, which perform beneficial functions to the host animal when given at a therapeutic dose.

It has been found that lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation directly from resected and washed GI tract of mammals are adherent to the GI tract following feeding of viable bacterial cells, and are also significantly immunomodulatory when fed to animals in viable, non-viable or fractionated form. Without being bound by theory, it is believed that *Bifidobacteria* obtainable by isolation from resected and washed GI tract closely associate with the gut mucosal tissues. Without further being bound by theory, this is believed to result in the probiotic *Bifidobacteria* of the present invention generating alternative host responses that result in its probiotic action. It has been found that probiotic bacteria obtainable by isolation from resected and washed GI tract can modulate the host's immune system via direct interaction with the mucosal epithelium, and the host's immune cells. This immunomodulation, in conjunction with the traditional mechanism of action associated with probiotic bacteria, i.e. the prevention of pathogen adherence to the gut by occlusion and competition for nutrients, results in the *Bifidobacteria* of the present invention being highly efficacious as a probiotic organism.

The *Bifidobacteria* of the present invention, obtainable by isolation from resected and washed feline GI tract, have *in vitro* anti-microbial activity against a number of pathogenic bacterial strains/species. Without being bound by theory, it is believed that this *in vitro* anti-microbial activity is indicative of potential probiotic activity *in vivo* in animals, preferably companion animals such as felines. The lactic acid bacteria of the present invention preferably have *in vitro* anti-microbial activity against *Salmonella*

typhimurium, *Listeria monocytogenes*, *Listeria innocua* or *Eschericia coli*, more preferably a mixture of these strains, more preferably still, all of these strains.

Without being bound by theory, it is believed that the anti-microbial activity of the lactic acid bacteria of the present invention may be the result of a number of different actions by the lactic acid bacteria herein. It has previously been suggested in the art that several strains of bacteria isolated from faecal samples exert their probiotic effect in the GI tract following oral consumption by preventing the attachment of pathogenic organisms to the gut mucosa by occlusion. This requires oral consumption of "live" or viable bacterial cells in order for a colony of bacteria to be established in the gut. However, it is believed that the *Bifidobacteria* of the present invention, obtainable by isolation from resected and washed feline GI tract, whilst exerting some probiotic effect due to occlusion if given in a viable form, may deliver a substantial probiotic effect in either the viable or non-viable form due to the production during fermentation *in vitro* of a substance or substances that either inhibit the growth of or kill pathogenic micro-organisms, and/or alter the host animal's immune competence. This form of probiotic activity is desirable, as the bacteria of the present invention can be given as either viable or non-viable cultures or purified fermentation products and still deliver a beneficial therapeutic effect to the host animal.

Preferably, the lactic acid bacteria of the present invention are able to maintain viability following transit through the GI tract. This is desirable in order for live cultures of the bacteria to be taken orally, and for colonisation to occur in the intestines and bowel following transit through the oesophagus and stomach. Colonisation of the intestine and bowel by the lactic acid bacteria of the present invention is desirable for long-term probiotic benefits to be delivered to the host. Oral dosing of non-viable cells or purified isolates thereof induces temporary benefits, but as the bacteria are not viable, they are not able to grow, and continuously deliver a probiotic effect *in situ*. As a result this may require the host to be dosed regularly in order to maintain the health benefits. In contrast, viable cells that are able to survive gastric transit in the viable form, and subsequently colonise by adhering to and proliferating on the gut mucosa are able to deliver probiotic effects continuously *in situ*.

Therefore, it is preferable that the lactic acid bacteria of the present invention maintain viability after suspension in a media having a pH of 2.5 for 1 hour. As used herein, "maintain viability" means that at least 25% of the bacteria initially suspended in the test media are viable using the plate count method known to those skilled in the art. Preferably, "maintain viability" means that at least 50% of the bacteria initially suspended are viable. It is desirable for the lactic acid bacteria of the present invention to maintain viability following exposure to low pH as this mimics the exposure to gastric juices in the stomach and upper intestine *in vivo* following oral consumption in animals.

Furthermore, it is preferable that the lactic acid bacteria of the present invention have a growth of at least 66% when in the presence of at least 0.3% porcine bile salts. Growth, as used herein is described in further detail in example 3. More preferably, the bacteria of the present invention have a growth of at least 66% when in the presence of at least 1% feline bile salts. Without being bound by theory it is believed that the lactic acid bacteria of the present invention, capable of maintaining viability in the presence of at least 0.3% porcine bile salts, are able to survive the conditions present in the intestine. This is thought to be a result of the addition of porcine bile to the culture medium mimicking the conditions of the intestine.

Further still, it is preferable that the lactic acid bacteria of the present invention have significant adhesion to gut epithelial cells *in vitro*. As used herein, "significant adhesion" means at least 1% of the total number of lactic acid bacteria co-incubated with the epithelial cells *in vitro* adhere to the epithelial cells. More preferably, at least 1.5% of bacterial cells co-incubated adhere to epithelial cells *in vitro*. Without wishing to be bound by theory, it is believed that gut epithelial cell adherence *in vitro* is indicative of the lactic acid bacteria's ability to colonise the GI tract of an animal *in vivo*.

Preferably, the strain of lactic acid bacteria according to the present invention is of a species selected from the group comprising *Bifidobacterium longum*, *Bifidobacterium animalis*, *Bifidobacterium adolescentis*, *Bifidobacterium bifidum*, *Bifidobacterium infantis*, or *Bifidobacterium thermophilum*, preferably *Bifidobacterium longum*.

The 16s-23s intergenic polynucleotide sequence is known to those skilled in the art as the sequence of DNA in the bacterial genome that can be used in order to identify

different species and strains of bacteria. This intergenic polynucleotide sequence can be determined by the method detailed below in example 4.

In a preferred embodiment of the present invention, the strain of lactic acid bacteria is selected from the group comprising *Bifidobacteria* having a 16s-23s intergenic polynucleotide sequence that has greater than 95% homology with SEQ. ID NO. 1 or greater than 94% homology with SEQ. ID NO. 2. More preferably the strain of lactic acid bacteria is selected from the group comprising Lactobacilli having a 16s-23s polynucleotide sequence having greater than 95%, more preferably still greater than 97%, homology to SEQ. ID NO. 1, or SEQ. ID NO. 2. More preferably, the strain of lactic acid bacteria according to the present invention is selected from the group comprising *Bifidobacteria* having a 16s-23s polynucleotide sequence according to SEQ. ID NO. 1 or SEQ. ID NO. 2. More preferably still, the strain of lactic acid bacteria according to the present invention is selected from *Bifidobacterium longum* NCIMB 41290 (*AHF5340*), *Bifidobacterium longum* NCIMB 41291 (*AHF1231*) or a mutant thereof.

The strain of lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline gastrointestinal tract can be used to deliver probiotic benefit following oral consumption in animals, preferably companion animals or humans. This probiotic benefit generally maintains and improves the overall health of the animal. Non-limiting elements of animal health and physiology that benefit, either in therapeutically relieving the symptoms of, or disease prevention by prophylaxis include inflammatory disorders, immunodeficiency, inflammatory bowel disease, irritable bowel syndrome, cancer (particularly those of the gastrointestinal and immune systems), diarrhoeal disease, antibiotic associated diarrhoea, appendicitis, autoimmune disorders, multiple sclerosis, Alzheimer's disease, amyloidosis, rheumatoid arthritis, arthritis, joint mobility, diabetes mellitus, insulin resistance, bacterial infections, viral infections, fungal infections, periodontal disease, urogenital disease, surgical associated trauma, surgical-induced metastatic disease, sepsis, weight loss, weight gain, excessive adipose tissue accumulation, anorexia, fever control, cachexia, wound healing, ulcers, gut barrier infection, allergy, asthma, respiratory disorders, circulatory disorders, coronary heart disease, anaemia, disorders of the blood coagulation system, renal disease, disorders of the central nervous system, hepatic disease, ischaemia, nutritional disorders, osteoporosis,

endocrine disorders, and epidermal disorders. Preferred are treatment of the gastrointestinal tract, including treatment or prevention of diarrhoea; immune system regulation, preferably the treatment or prevention of autoimmune disease and inflammation; maintaining or improving the health of the skin and/or coat system, preferably treating or preventing atopic disease of the skin; ameliorating or reducing the effects of aging, including mental awareness and activity levels; preventing disorders associated with the hypothalamus-pituitary-adrenal axis, and improving joint health whereby improving mobility.

The treatment of the disorders disclosed above may be measured using techniques known to those skilled in the art. For example, inflammatory disorders including autoimmune disease and inflammation may be detected and monitored using *in vivo* immune function tests such as lymphocyte blastogenesis, natural killer cell activity, antibody response to vaccines, delayed-type hypersensitivity, and mixtures thereof. Such methods are briefly described herein, but well known to those skilled in the art.

1. Lymphocyte blastogenesis: This assay measures the proliferative response *in vitro* of lymphocytes isolated from fresh whole blood of test and control animals to various mitogens and is a measure of overall T- and B-cell function. Briefly, peripheral blood mononucleocytes (PBMC) are isolated from whole blood by Ficoll-Hypaque density centrifugation methods known to those skilled in the art. The isolated PBMCs are washed twice in RPMI 1640 cell media supplemented with HEPES, L-glutamine and penicillin/streptomycin. The washed cells are resuspended in RPMI 1640, counted, and the cell density adjusted appropriately. The 2×10^5 cells are exposed to a range of concentrations (0.1 $\mu\text{g/ml}$ to 100 $\mu\text{g/ml}$) of various mitogens, some examples of which include pokeweed mitogen (Gibco), phytohaemagglutinin (Gibco) and concanavalin A (Sigma) in triplicate for 72 hours at 37°C and 5% CO₂ with 10% foetal bovine serum (Sigma). At 54 hours the cells are pulsed with 1 μCi ³H-thymidine, and the cells harvested and scintillation counts read on a TopCount NXT at 72 hours.
2. Natural killer cell activity: As described in US 6,310,090, this assay measures the *in vitro* effector activity of natural killer cells isolated from fresh whole blood of test and control animals. Natural killer cells are a component of the innate

immune function of a mammal. Feline thyroid adenocarcinoma cells were used as target cells in assessing NK cell cytotoxic activity. This cell line was previously shown to be susceptible to killing by feline NK cell. Target cells were cultured in a T75 flask with 20 mL minimum essential medium (MEM; Sigma Chem. Co., St. Louis, Mo.) supplemented with 10% fetal calf serum (FCS), 100 U/mL of penicillin and 100 µg/mL of streptomycin. When confluent, target cells were trypsinized, washed 3 times and resuspended to 5×10^5 cells/mL in complete medium (RPMI-1640+10% FCS+100 U/mL of penicillin+100 µg/mL of streptomycin). Triplicate 100 . µL aliquots of the target cells were pipetted into 96-well U-bottom plates (Costar, Cambridge, Mass.) and incubated for 8 hours to allow cell adherence. Lymphocytes (effector cells; 100 . µL) isolated by Ficoll-Hypaque separation (as described above) were then added to the target cells to provide an effector/target cell (E:T) ratio of 10:1. After 10 hours of incubation at 37°C., 20 . µl of a substrate containing 5 . µg of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) was added. The mixture was incubated for 4 hours at 37°C. after which the unmetabolized MTT was removed by aspiration. The formazan crystals were dissolved by adding 200 µL of 95% ethanol. Optical density was measured at 570 nm using a microplate reader. The percentage of NK cell-specific lysis was calculated as follows:

$$\text{Specific Cytotoxicity (\%)} = 100 \times \{1 - [(\text{OD of target cells and effector cells} \\ - \text{OD of effector cells}) / (\text{OD of target cells})]\}$$

3. Antibody response to vaccines: The test subjects are given an array (up to 5) of vaccines after at least 12 weeks of probiotic or control feeding. The vaccines may be a mixture of novel and redundant vaccines. Non-limiting examples of vaccine arrays that may be used include mixtures of vaccines prepared by Fort Dodge Animal Health. Non-limiting examples of vaccines suitable for use herein include Feline distemper, adenovirus, coronavirus, parainfluenza, and parvovirus. The test subject's vaccine history will determine the vaccines to be used. The specific antibodies to the vaccines given are measured in blood for 3 weeks and the length and strength of response in control and probiotic feeding groups compared.

4. Delayed-type hypersensitivity: An *in vivo*, non-invasive method of assessing immune system status. This test comprises an intradermal injection of the polyclonal mitogen Phytohemmagglutinin (PHA) in combination with sheep red blood cells a multivalent vaccine, histamine (100 μ L of 0.0275 g/L Histamine Phosphate; Greer, Lenoir, NC), or PBS (100 μ L of Phosphate Buffered Saline, 8.5 g/L; Sigma). The immune response to the antigen is recorded as skinfold thickness using calipers at time intervals of 0, 24, 48 and 72 hours post-injection. An increase in skinfold thickness is indicative of a greater hypersensitivity response that should be decreased by treatment with the bacteria of the present invention.

Additional methods for determining the effect of the *Bifidobacteria* bacteria of the present invention are described in US 6,133,323 and US 6,310,090.

Furthermore, ameliorating the effects of age may be determined using dual x-ray absorptometry or CT scan for measuring body composition, including body fat mass, fat-free mass and bone mineral content. Similarly, this method may be used to determine anatomy changes such as weight loss or bone density in subjects following infection.

The *Bifidobacteria* of the present invention may also be used in a method for reducing stress levels in companion animals. Concentrations of blood stress hormones including epinephrine, norepinephrine, dopamine, Cortisol, C-reactive protein and other acute phase proteins may be measured to determine stress levels and their reduction or maintenance. These hormones are recognized biomarkers of stress and can be readily measured using techniques known to those skilled in the art. Additionally, direct measure of adrenal size as an *in vivo* marker of activity of the hypothalamus-pituitary-adrenal axis may be measured by CT imaging.

Further still, maintenance or improvement of the health of the skin and/or coat system of companion animals, including atopic disease of the skin, may be measured using skin and coat assessments conducted by two trained individuals. Examples of criteria examined during such assessments include:

- a) Shedding index: A shedding index is assigned to each test subject by collecting hair produced during a standardized brushing session. The hair is retained and weighed, and control and test subjects compared.

- b) Subjective skin/coat evaluations: Trained panelists subjectively evaluate skin and coat condition by assessing shedding, dander, shine, uniformity, softness and density.
- c) Skin functional assessment: The barrier function of the skin may be assessed by wiping the skin surface with an acetone-soaked gauze. This technique effectively disrupts the skin barrier by removing single cell layers and associated lipid fractions of the stratum corneum. Barrier disruption is quantified by measuring the increase in transepidermal water loss (TEWL) and the degree of redness of the insulted site using methods known to those skilled in the art. Redness (erythema) scores are obtained using the previously described camera and lighting system. TEWL readings and redness scores are obtained immediately before and after disruption, and at five and 24-hour endpoints to assess the protective and healing properties of skin.

The treatment or prevention of diarrhoea in companion animals may be measured using stool scores. Stools scores may be recorded daily according to the following guidelines and control and test groups compared before and after feeding with the bacteria according to the present invention.

Score: 5 Extremely Dry

This stool is hard and does not stick to surfaces. Stool will roll when pushed. No indentations are made when stool is picked up. Stool is often defecated in groups of individual stools instead of one complete unit. The stool maintains original shape after collection.

Score: 4 Firm (Ideal stool)

This stool is firm, well shaped, and cylindrical. This stool does not break apart easily when picked up. This stool may leave residue on surfaces and gloves. This stool is often defecated as one unit. The stool maintains original shape after collection.

Score: 3 Soft, with shape

This stool is soft, however there are definite shapes. This stool will break apart easily and will definitely leave residue on surfaces and gloves. The stool often loses original shape after collection. This stool is often present with another score but can comprise whole stool sample.

Score: 2 Soft, without shape

This stool is soft and will have no cylindrical shape. The shape often associated with a "2" is a "cow patty" shape. This stool will lose the original shape when collected and will definitely leave residue on surfaces and gloves. This stool score is often present with another score but can comprise the whole stool sample. This stool sample may spread over an area of several inches.

Score: 1 Liquid

This stool score will always resemble liquid and there may or may not be particulate matter present. This stool will often be defecated in groups of piles instead of one complete unit. Mucous is often present with this stool sample. This stool sample is very difficult to collect and residue is always left on surfaces and gloves. This stool sample may spread over an area of several inches.

In addition, other observations are also recorded, including: blood in stool; foreign object in stool; or mucous in stool.

Furthermore, the treatment of gastrointestinal infection in companion animals may comprise improving microbial ecology of companion animals. Improving the microbial ecology of companion animals preferably comprises reducing the levels of pathogenic bacteria in the faeces of companion animals. The levels of pathogenic bacteria present in the faeces of companion animals may be enumerated using the standard plate count method known to those skilled in the art. More preferably, the pathogenic bacteria are selected from the group consisting of *Clostridia*, *Escherichia*, *Salmonella*, bacteriodes and mixtures thereof. Non-limiting examples of suitable strains of pathogenic bacteria include *C. perfringens*, *C. difficile*, *Escherichia coli*, *Salmonella typhimurium* and mixtures thereof.

The method of use of the bacteria of the present invention may also include the treatment, either prophylactic or therapeutic of the urinary tract of mammals, preferably companion animals. Non-limiting examples of urinary tract treatment include treatment or prevention of urinary tract infections, treatment or prevention of kidney disease, including urinary tract stones, treatment or prevention of bladder infections and the like. Without being bound by theory, it is believed that the *Bifidobacteria* bacteria of the present invention are useful in preventing these ailments as a result of their ability to

degrade oxalic acid, as demonstrated *in vitro*. Oxalic acid is a by-product of urinary metabolism that can form insoluble precipitates that result in kidney, bladder and other urinary tract infections. By degrading oxalic acid, and therefore potentially preventing its precipitation and build up in the urinary tract, the bacteria of the present invention may treat and prevent infections and other ailments of the urinary tract. Oxalic acid degradation may be measured *in vitro* using the Oxalic acid test kit cat # 755699 commercially available from Boehringer Mannheim/R-Biopharm.

The *Bifidobacteria* of the present invention may be used in a method for improving or maintaining the health of companion animals comprising improving fibre digestion. Improving fibre digestion is desirable as it promotes the growth of said probiotic bacteria, as well as beneficial endogenous microflora, which aid in the suppression of some potentially pathogenic bacteria. In addition, a decrease in the amount of toxic metabolites and detrimental enzymes that result from colonic fermentation has been documented in humans (Tomomatsu, H. "Health effects of oligosaccharides", (1994) *Food Technol*, **48**, 61-65). Fibre digestion may be determined using the method described in Vickers et al. (2001), "Comparison of fermentation of selected fructooligosaccharides and other fiber substrates by feline colonic microflora", *Am. J. Vet. Res.* **61** (4), 609-615, with the exception that instead of inoculating using diluted fecal samples each experiment used pure cultures of the bacterial strains of interest.

The feline probiotic strains of the present invention may be used to reduce the odor of the feces and urine and concomitantly in the litterbox by reducing the production of compounds in the feces and urine that cause odor. Non-limiting examples of odor-causing compounds include ammonia, indoles, phenols, amines, branched chain fatty acids, and volatile sulphur-containing compounds. Without wishing to be bound by theory it is believed that reducing the levels of these compounds in the feces or urine of a companion animal reduces the odor associated with the feces or urine. Furthermore, for companion animals that use a litter box, there is a concomitant decrease in litter box odor.

The method of use of the lactic acid bacteria of the present invention typically involves oral consumption by the animal. Oral consumption may take place as part of the normal dietary intake, or as a supplement thereto. The oral consumption typically occurs

at least once a month, preferably at least once a week, more preferably at least once per day. The lactic acid bacteria of the present invention may be given to the companion animal in a therapeutically effective amount to maintain or improve the health of the animal, preferably a companion animal. As used herein, the term "therapeutically effective amount" with reference to the lactic acid bacteria, means that amount of the bacteria sufficient to provide the desired effect or benefit to a host animal in need of treatment, yet low enough to avoid adverse effects such as toxicity, irritation, or allergic response, commensurate with a reasonable benefit/risk ratio when used in the manner of the present invention. The specific "therapeutically effective amount" will vary with such factors as the particular condition being treated, the physical condition of the user, the duration of the treatment, the nature of concurrent therapy (if any), the specific dosage form to be used, the carrier employed, the solubility of the dose form, and the particular dosing regimen.

Preferably, the lactic acid bacteria are given to the companion animal at a dose of from $1.0E+04$ to $1.0E+14$ CFU per day, more preferably from $1.0E+06$ to $1.0E+12$ CFU per day. The composition preferably may contain at least 0.001% of from $1.0E+04$ to $1.0E+12$ CFU/g of the lactic acid bacteria of the genus *Bifidobacteria* obtainable by isolation from resected and washed feline GI tract. The lactic acid bacteria can be given to the animal in either viable form, or as killed cells, or distillates, isolates or other fractions of the fermentation products of the lactic acid bacteria of the present invention, or any mixture thereof.

Preferably, the lactic acid bacteria, or a purified or isolated fraction thereof, are used to prepare a composition intended to maintain or improve the health of an animal. As indicated above, the composition may be part of the normal dietary intake, or a supplement. Where the composition comprises part of the normal dietary intake, the composition may be in the form of a dried animal food such as biscuits or kibbles, a processed grain feed, a wet animal food, yoghurts, gravies, chews, treats and the like.

Such compositions may comprise further components. Other components are beneficial for inclusion in the compositions used herein, but are optional for purposes of the invention. For example, food compositions are preferably nutritionally balanced. In one embodiment, the food compositions may comprise, on a dry matter basis, from about

20% to about 50% crude protein, preferably from about 22% to about 40% crude protein, by weight of the food composition. The crude protein material may comprise any material having a protein content of at least about 15% by weight, non-limiting examples of which include vegetable proteins such as soybean, cotton seed, and peanut, animal proteins such as casein, albumin, and meat tissue. Non-limiting examples of meat tissue useful herein include fresh meat, and dried or rendered meals such as fish meal, poultry meal, meat meal, bone meal and the like. Other types of suitable crude protein sources include wheat gluten or corn gluten, and proteins extracted from microbial sources such as yeast.

Furthermore, the food compositions may comprise, on a dry matter basis, from about 5% to about 35% fat, preferably from about 10% to about 30% fat, by weight of the food composition. Further still, food compositions comprising the lactic acid bacteria of the present invention may also comprise from about 4% to about 25% total dietary fibre. The compositions may also comprise a multiple starch source as described in WO99/51108.

The compositions of the present invention may further comprise a source of carbohydrate. Grains or cereals such as rice, corn, milo, sorghum, barley, alfalfa, wheat, and the like are illustrative sources. In addition, the compositions may also contain other materials such as dried whey and other dairy by products.

The compositions comprising the bacteria of the present invention may also comprise a prebiotic. "Prebiotic" includes substances or compounds that are fermented by the intestinal flora of the companion animal and hence promote the growth or development of lactic acid bacteria in the gastro-intestinal tract of the companion animal at the expense of pathogenic bacteria. The result of this fermentation is a release of fatty acids, in particular short-chain fatty acids in the colon. This has the effect of reducing the pH value in the colon. Non-limiting examples of suitable prebiotics include oligosaccharides, such as inulin and its hydrolysis products commonly known as fructooligosaccharides, galacto- oligosaccharides, xylo-oligosaccharides or oligo derivatives of starch. The prebiotics may be provided in any suitable form. For example, the prebiotic may be provided in the form of plant material which contains the fiber. Suitable plant materials include asparagus, artichokes, onions, wheat or chicory, or

residues of these plant materials. Alternatively, the prebiotic fiber may be provided as an inulin extract, for example extracts from chicory are suitable. Suitable inulin extracts may be obtained from Orafit SA of Tirlemont 3300, Belgium under the trade mark "Raftiline". For example, the inulin may be provided in the form of Raftiline (g) ST which is a fine white powder which contains about 90 to about 94% by weight of inulin, up to about 4% by weight of glucose and fructose, and about 4 to 9% by weight of sucrose. Alternatively, the fiber may be in the form of a fructooligosaccharide such as obtained from Orafit SA of Tirlemont 3300, Belgium under the trade mark "Raftilose". For example, the inulin may be provided in the form of Raftilose (g) P95. Otherwise, the fructooligosaccharides may be obtained by hydrolyzing inulin, by enzymatic methods, or by using micro-organisms.

For dried companion animal foods a suitable process is extrusion cooking, although baking and other suitable processes may be used. When extrusion cooked, the dried companion animal food is usually provided in the form of a kibble. If a prebiotic is used, the prebiotic may be admixed with the other ingredients of the dried companion animal food prior to processing. A suitable process is described in European patent application No 0850569. If a probiotic micro-organism is used, the organism is best coated onto or filled into the dried companion animal food. A suitable process is described in European patent publication Number EP 0 862 863.

For wet foods, the processes described in US patents 4,781,939 and 5,132,137 may be used to produce simulated meat products. Other procedures for producing chunk type products may also be used; for example cooking in a steam oven. Alternatively, loaf type products may be produced by emulsifying a suitable meat material to produce a meat emulsion, adding a suitable gelling agent, and heating the meat emulsion prior to filling into cans or other containers. Typical wet food compositions may comprise from about 5% to about 15% protein, from about 1% to about 10% fat, and from about 1% to about 7% fibre. Non-limiting ingredients that may be used in wet food compositions include chicken, turkey, beef, whitefish, chicken broth, turkey broth, beef broth, chicken liver, brewers rice, corn grits, fish meal, egg, beet pulp, chloride, flax meal, lamb, beef by-products, chicken by-products and mixtures thereof.

In another embodiment, supplement compositions such as biscuits, chews, and other treats may comprise, on a dry matter basis, from about 20% to about 60% protein, or from about 22% to about 40% protein, by weight of the supplement composition. As another example, the supplement compositions may comprise, on a dry matter basis, from about 5% to about 35% fat, or from about 10% to about 30% fat, by weight of the supplement composition. Food and supplement compositions intended for use by felines or felines are commonly known in the art.

The companion animal foods may contain other active agents such as long chain fatty acids and zinc. Suitable long chain fatty acids include alpha-linoleic acid, gamma linolenic acid, linoleic acid, eicosapentanoic acid, and docosahexanoic acid. Fish oils are a suitable source of eicosapentanoic acids and docosahexanoic acid.

Borage oil, blackcurrent seed oil and evening primrose oil are suitable sources of gamma linolenic acid. Safflower oils, sunflower oils, corn oils and soy bean oils are suitable sources of linoleic acid. These oils may also be used in the coating substrates referred to above. Zinc may be provided in various suitable forms, for example as zinc sulfate or zinc oxide. Further, many ingredients commonly used in companion animal foods are sources of fatty acids and zinc. It has been observed that the combination of chicory, as a source of prebiotic, with a linoleic-acid rich oil, such as soy bean oil, provides unexpected benefits, suggestive of a synergistic effect.

Where the composition is in the form of a gravy, the composition preferably comprises at least 10% of a broth, or stock, non-limiting examples of which include vegetable beef, chicken or ham stock. Typical gravy compositions may comprise from about 0.5% to about 5% crude protein, from about 2% to about 5% crude fat, and from about 1% to about 5% fibre.

Further non-limiting examples of supplements suitable for use herein include powders, oil suspensions, milk-based suspensions, cheeses, cocoa-butter-based compositions and pills or capsules. Where the composition is in the form of a pill, suitable binding agents are required to maintain the pill in a solid, pressed form. Non-limiting examples of suitable binding agents include the natural gums such as xanthan gum, pectins, lecithins, alginates and others known to those skilled in the art. Where the composition is in the form of a capsule, the composition is preferably encapsulated using

technologies known to those skilled in the art. Non-limiting examples of suitable encapsulation materials include polyvinyl alcohol (PVA), polyvinylpyrrolidone (PVP), alginates, and gelatin. Yoghurt-based compositions may comprise from about 1% to about 5% protein, from about 10% to about 20% carbohydrate, from about 1% to about 5% fibre, from about 1% to about 5% fat and from about 50% to about 90% liquid carrier such as milk.

Examples

The following examples are provided to illustrate the invention and are not intended to limit the scope thereof in any manner.

Example 1: Isolation of Lactic Acid bacteria from feline GI tracts

Feline intestinal samples were obtained from healthy cats presenting at the local veterinarians for owner initiated and approved euthanasia. All animals were healthy and disease-free. The colon, mid-colon, caecum and ileum of each cat were dissected in order to expose the mucosa.

Supernatants were removed following agitation of the mucosal tissue (vortexed for 1 minute) and following mechanical homogenisation of the tissue. Each supernatant was plated on de Mann Rogosa Sharpe (MRS) agar. These were incubated anaerobically, using the Anerocult GasPak system, for 48 hours at 37°C. Isolated colonies from the plates were re-streaked onto either MRS and again grown anaerobically under the same conditions. Isolated colonies were re-streaked a further 4 times in order to purify a single strain. Colony morphology and microscopic appearance were assessed. Suitable isolates were tested for Gram reaction and catalase activity. Identification of gram positive, catalase negative rods was performed using API testing (API 50CHL, BioMerieux). Harvested cells were washed twice with 0.05M phosphate buffer (pH 6.5) and cysteine-HCl (500 mg/l) followed by sonication. Centrifugation removed cellular debris. Supernatants were incubated with NaF (6 mg/ml) and Na iodoacetate (10 mg/ml) for 30 minutes at 37°C. The reaction was stopped by incubation with hydroxylamine HCl (pH6.5) for 10 minutes at room temperature. Colour development was monitored following the addition of HCl (4M), FeCl₃.6H₂O (5% (w/v) in 0.1M HCl) and fructose-6-phosphate (Na salt). Formation of acetyl phosphate from fructose-6-phosphate was evidenced by the reddish colour formed by the ferric chelate of its hydroxymate.

Example 2: Screening for Anti-Microbial Activity

Each of the isolated lactic acid bacterial strains was incubated anaerobically in MRS broth. 2 μ l of each culture were spotted onto MRS agar plates and incubated anaerobically overnight. *Salmonella typhimurium* and Enteropathogenic *E. Coli* (ExPEC) were pre-grown overnight and 100 μ l inoculated into molten agar (1% v/v). This indicator culture was poured onto the surface of the inoculated MRS plates. Following overnight incubation, zones of inhibition around the probiotic colony were measured. All experiments were performed in duplicate on three separate occasions. In addition, incorporating the buffer 2% betaglycerophosphate into the agar enabled assessment of the contribution of acid production to the observed pathogen inhibition *in vitro*.

The data presented in Table 2 clearly demonstrate that the lactic acid bacteria strains of the present invention obtainable by isolation from resected and washed feline GI tract have significant anti-microbial activity *in vitro*, indicative of potential probiotic activity.

Table 2

	AHF534D	1231
<i>S. typhimurium</i>	5.5	3.67
ExPEC	6.17	6

Example 3: *In Vitro* Measures of Survival and Colonisation

pH Tolerance

Bacterial cells were harvested from overnight cultures, washed twice in phosphate buffer (pH 6.5) and resuspended in MRS/TPY broth adjusted with 1M HCl to pH 2.5. The cells were incubated anaerobically at 37°C and their survival measured at intervals of 0, 30, 60, 120, 240 and 360 minutes using the plate count method known to those skilled in the art. Table 3 summarises this data per strain.

Table 3

Survival of strains in a low pH environment (pH 2.5). Data are log CFU counts.

STRAIN	TIME (min)					
	0	30	60	120	180	360
AHF5340	8.34	8.22	8.29	8.19	8.26	8.12
AHF1231	9.10	9.06	9.07	9.04	8.97	8.91

Bile Resistance

The bacterial strains were streaked onto MRS agar supplemented with porcine bile (Sigma) at 0.5%, 1% and 5% (w/v). Plates were incubated at 37°C under anaerobic conditions and the growth recorded after 48 hours. Growth was compared with control plates by an experienced observer, and the growth of colonies described as:

Negative (0) – no growth;

+ (1) – Hazy translucent growth (<33% control-plates with 0% bile);

++ (2) – Definite growth but not as good as controls (>33% but <66%);

+++ (3) – Growth equivalent to controls (>66%).

Once the growth of the colonies in the presence of bile salts is compared with the controls, the growth descriptors are given numerical values of 0, 1, 2 or 3 (-; +; ++, +++ respectively), and then expressed as a percentage, where 3 represents 100%.

Table 4 demonstrates that the *Bifidobacterium* of the present invention clearly demonstrate a resistance to bile salts, being able to grow and form colonies at a level of at least 66% in most instances when exposed to 0.3% porcine bile salts.

Table 4

Survival of strains in various concentrations of porcine bile

STRAIN	PERCENTAGE BILE (%)							
	0	0.3	0.5	1	2	5	7.5	10
AHF5340	+++	+++	+++	++	++	++	++	++
AHF1231	+++	++	+	-	-	-	-	-

Furthermore, in order to assess any differences in the ability of the strains to colonise the GI tract of cats, the bacterial strains were streaked onto MRS agar

supplemented with feline bile at 0.5%, 1% and 2% (w/v). Feline bile was obtained from cats undergoing endoscopy in a clinical setting during a non-terminal procedure. Plates were incubated at 37°C under anaerobic conditions and the growth recorded after 48 hours. Growth was compared with control plates by an experienced observer, and the growth of colonies described as:

Negative (0) – no growth;

+ (1) – Hazy translucent growth (<33% control-plates with 0% bile);

++ (2) – Definite growth but not as good as controls (>33% but <66%);

+++ (3) – Growth equivalent to controls (>66%).

Once the growth of the colonies in the presence of bile salts is compared with the controls, the growth descriptors are given numerical values of 0, 1, 2 or 3 (-; +; ++, +++ respectively), and then expressed as a percentage, where 3 represents 100%.

Table 5 demonstrates that the *Bifidobacterium* of the present invention clearly demonstrate a resistance to feline bile salts, being able to grow and form colonies at a level of at least 66% in most instances when exposed to 1% feline bile salts.

Table 5

Survival of strains in various concentrations of feline bile

STRAIN	PERCENTAGE BILE (%)			
	0	0.5	1	2
AHF5340	+++	+++	++	++
AHF1231	+++	++	++	+

Gut Epithelial Cell Adhesion

The human epithelial cell line, HT-29, was used to assess the adhesion properties of selected strains. Epithelial cells were routinely cultured as a monolayer in 75 cm² tissue culture flasks at 37°C in a humidified atmosphere containing 5% CO₂ in Dulbecco's Minimal Essential Media (DMEM) containing 10% foetal calf serum (FCS), pen/strep, glutamine and fungizone. For experimental purposes, the epithelial cells were seeded at a concentration of 5 x 10⁵ cells/ml (3 mls total volume) per well in 6 well culture plates (Sarstedt). Following incubation for 7 days, to allow differentiation, the epithelial monolayers were washed with antibiotic-free medium containing 10% FCS. Bacterial suspensions plus/in antibiotic-free DMEM were added to each well and the cells

incubated for 90 minutes at 37°C. Following incubation, the monolayers were washed three times with PBS. The epithelial cells were lysed in deionised H₂O and the number of adherent bacteria enumerated using the plate count method known to those skilled in the art. Adhesion was expressed as a percentage of the number of bacteria initially plated. *Bifidobacteria longum* AHF534D had an adhesion level of 1.7%, whilst *Bifidobacteria longum* AHF1231 had an adhesion level of 44.1%.

Example 4: 16s-23s Intergenic Polynucleotide Sequencing

The feline *Bifidobacterium* isolates were centrifuged in stock tubes and the resulting pellet lysed in 100 µl of Extraction Solution and 25 µl of Tissue Preparation solution (Sigma, XNAT2 Kit), incubated for 10 minutes at room temperature for 10 minutes. The samples were then incubated for 5 minutes at 95°C and then 100 µl of Neutralization Solution (XNAT2 kit) was added. The genomic DNA solution was then, quantified using a Nanodrop spectrophotometer and stored at 4°C.

PCR was performed using the intergenic spacer (IGS) primers, IGS L: 5'-GCTGGATCACCTCCTTTC-3' and IGS R: 5'-CTGGTGCCAAGGCATCCA-3' (Bridgidi et al 2000, System Appl. Microbiol., 23, 391-399 (2000)). The cycling conditions were 94°C for 3 min (1 cycle), 94°C for 30 sec, 53°C for 30 sec, 72°C for 30 sec (28 cycles). The PCR reaction contained 4 µl (50ng) of DNA, PCR mix (XNAT2 kit), 0.4 µM IGS L and R primer (MWG Biotech, Germany). The PCR reactions were performed on an Eppendorf thermocycler. The PCR products (10 µl) were ran alongside a molecular weight marker (100 bp Ladder, Roche) on a 2 % agarose EtBr stained gel in TAE, to determine the IGS profile.

PCR products of *Bifidobacterium* (single band) were purified using the Promega Wizard PCR purification kit.

The purified PCR products were sequenced using the primer sequences (above) for the intergenic spacer region. Sequence data was the searched against the NCBI nucleotide database to determine the identity of the strain by nucleotide homology.

Following sequencing, the obtained sequences for the four deposited strains were compared with the on-line sequence database "BLAST", available at <http://www.ncbi.nlm.nih.gov/BLAST/> for homology with other deposited bacterial 16s-23s sequences. The closest matches for *Bifidobacterium longum* NCIMB 41290

(AHF5340) and *Bifidobacterium longum* NCIMB 41291 (AHF1231) was the strain *Bifidobacterium longum* NCC2705, having a percent homology score of 95% and 94% respectively.

Example 5: Example Compositions

Examples 1 to 4 are examples of dried kibble compositions comprising the probiotic *Bifidobacteria* of the present invention.

Ingredient	Percentage on a weight Basis			
	Ex. 1	Ex. 2	Ex. 3	Ex. 4
Cereal grains	To 100	To 100	To 100	To 100
Poultry by-product meal	43.5	40	45	35
Poultry fat	1.28	1.02	1.16	1.35
Egg product	2.4	2.1	2.5	2.2
Chicken liver meal	1.0	1.0	1.0	1.0
Brewer's dried yeast	1.0	1.0	1.0	1.0
Monosodium phosphate	1.0	1.0	1.0	1.0
Calcium carbonate	0.8	0.8	0.8	0.8
Potassium chloride	0.6	0.6	0.6	0.6
Vitamins	0.4	0.4	0.4	0.4
Choline chloride	0.3	0.3	0.3	0.3
Minerals	0.3	0.3	0.3	0.3
DL-Methionine	0.1	0.1	0.1	0.1
Sodium Chloride	0.03	0.03	0.03	0.03
Probiotic (1 x 10 ¹⁰ cfu/g NCIMB 41290 in sunflower oil)	1	0.5	-	0.6
Probiotic (1 x 10 ¹⁰ cfu/g NCIMB 41291 in sunflower oil)	-	0.5	1	0.4

Examples 5 to 7 are examples of wet companion animal food compositions comprising the probiotic *Bifidobacteria longum* of the present invention.

Ingredient	Percentage on a weight Basis		
	Ex. 5	Ex. 6	Ex. 7
Water	To 38	To 47	To 50
Poultry Liver	To 25	To 20	To 15
Poultry Products	25	20	20
Brewers Rice	5	7	10
Egg Product	3	2.5	1.5
Poultry Fat	2.9	3.0	3.2
Chicken Stock	0.6	0.7	0.9
Taurine	0.1	0.1	0.1
Vitamins	0.05	0.1	0.1
Minerals	0.05	0.1	0.1
Probiotic (1 x 10 ¹⁰ cfu/g NCIMB 41290)	4	5	6

Examples 8 to 10 are examples of yoghurt supplement compositions comprising the probiotic *Bifidobacteria longum* of the present invention.

Ingredient	Percentage on a weight Basis		
	Ex. 8	Ex. 9	Ex. 10
Milk	38	42	48
Sugar	12	12	10
Modified Starch	1.0	0.8	0.8
Prebiotic	0.25	0.3	0.5
Probiotic (1 x 10 ¹⁰ cfu/g NCIMB 41291)	4	5	6

All documents cited in the Detailed Description of the Invention are, in relevant part, incorporated herein by reference; the citation of any document is not to be construed as an admission that it is prior art with respect to the present invention.

While particular embodiments of the present invention have been illustrated and described, it would be obvious to those skilled in the art that various other changes and modifications can be made without departing from the spirit and scope of the invention. It is therefore intended to cover in the appended claims all such changes and modifications that are within the scope of this invention.

PCT

Original (for SUBMISSION)

0-1	Form PCT/RO/134 (SAFE) Indications Relating to Deposited Microorganism(s) or Other Biological Material (PCT Rule 13bis)	
0-1-1	Prepared Using	PCT-SAFE [EASY mode] Version 3.51.001.176 MT/FOP 20060101/0.20.4rc.2.7
0-2	International Application No.	
0-3	Applicant's or agent's file reference	P-171&+/CB
1	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
1-1	page	2
1-2	line	24
1-3	Identification of deposit	
1-3-1	Name of depository institution	NCIMB NCIMB Ltd.
1-3-2	Address of depository institution	Ferguson Building, Craibstone Estate, Bucksburn, Aberdeen AB21 9YA, United Kingdom
1-3-3	Date of deposit	07 December 2005 (07.12.2005)
1-3-4	Accession Number	NCIMB 41290
1-4	Additional Indications	Bifidobacterium longum (AHF5340)
1-5	Designated States for Which Indications are Made	all designations
1-6	Separate Furnishing of Indications	Receipt of Deposit and Viability Proof from the Depository Institution to be supplied at National Phase Entry as required.
	These indications will be submitted to the International Bureau later	
2	The indications made below relate to the deposited microorganism(s) or other biological material referred to in the description on:	
2-1	page	2
2-2	line	25
2-3	Identification of deposit	
2-3-1	Name of depository institution	NCIMB NCIMB Ltd.
2-3-2	Address of depository institution	Ferguson Building, Craibstone Estate, Bucksburn, Aberdeen AB21 9YA, United Kingdom
2-3-3	Date of deposit	20 October 2005 (20.10.2005)
2-3-4	Accession Number	NCIMB 41291
2-4	Additional Indications	Bifidobacterium longum (AHF1231)
2-5	Designated States for Which Indications are Made	all designations
2-6	Separate Furnishing of Indications	Receipt of Deposit and Viability Proof from the Depository Institution to be supplied at National Phase Entry as required.
	These indications will be submitted to the International Bureau later	

Original (for SUBMISSION)

FOR RECEIVING OFFICE USE ONLY

0-4	This form was received with the international application: (yes or no)	
0-4-1	Authorized officer	

FOR INTERNATIONAL BUREAU USE ONLY

0-5	This form was received by the international Bureau on:	
0-5-1	Authorized officer	

SEQUENCE LISTING

<110> Alimentary Health Ltd.
The IAMS Company

<120> Feline Probiotic Bifidobacteria

<130> P171P2&

<140> 2,607,949

<141> 2006-02-02

<160> 4

<170> PatentIn version 3.3

<210> 1

<211> 500

<212> DNA

<213> Bifidobacterium longum

<220>

<221> misc_feature

<222> (3)..(3)

<223> n i s a , c , g , o r t

<220>

<221> misc_feature

<222> (6)..(6)

<223> n i s a , c , g , o r t

<220>

<221> misc_feature

<222> (475)..(475)

<223> n i s a , c , g , o r t

<400> 1

tcnggnngtgc tgggatacct cctttctacg gagaatgcag gatgccgttc ggcgtccggt 60

gcgaaccctc ccgcgcgatg gtcgcgtgtg gacgggttgc tgggtgtggaa gatgtcgttg 120

gctttgcccc gccggtcgtg cgggtgggtgc cgggggtggg cggatgcgct ttggggctcc 180

cggatcgcca cccaggtt ttgcctggcg cgattcgatg cccgtcgtgc ctgcgggccg 240

gccgtgtgcc ggtctggtcg gcgtggcggg gcgtggtggc ttgagaactg gatagtggac 300

gcgagcaaaa caagatTTTT tttgtgtcct tgttttgctg ttgatttcga atcgaactct 360
attgttcggt tcgatcggtt tgtgatcatt tttagtgtga tgatttgcg tctgggaatt 420
tgctagagga atcttgccgc gcatgcatgt tgcgtgtgtg ttgcttgcaa gggcntatgg 480
tggatgcctt gccacccaga 500

<210> 2
<211> 500
<212> DNA
<213> Bifidobacterium longum

<220>
<221> misc_feature
<222> (3)..(3)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (6)..(6)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (441)..(441)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (452)..(452)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (475)..(475)
<223> n is a, c, g, or t

<220>
<221> misc_feature
<222> (485)..(485)
<223> n is a, c, g, or t

<400> 2
tcngngtgc tggggatacc tcctttctac ggagaatgca ggatgctgtt cggcgtccgg 60

tgccaaccct cccgcgcggt ggtcgcgtgt ggacggggtg ctggtgtgga agatgtcgtt	120
ggctttgccc cgccggtcgt gcggtgggtg ccgggggtgg gcggatgcgc ttttgggctc	180
ccggatcgcc accccaggct tttgcctggc gcgattcgat gcccgtcgtg cctgcgggcc	240
ggccgtgtgc cggctctggc ggcgtggcgg tgcgtggtgg cttgagaact ggatagtgga	300
cgcgagcaaa acaagggttt tttgtgtctt tgttttgctg ttgatttcga atcgaactct	360
attgttcggt tcgatcgttt tgtgatcatt tttagtgtga tgatttgcg tctgggaatt	420
tgctagagga atcttgccgc ncatgcatgt tncgtgtgtg ttgcttgcaa gggcntatgg	480
tggangcctt gccacccaga	500

<210> 3
 <211> 18
 <212> DNA
 <213> Artificial

<220>
 <223> Artificial PCR primer

<400> 3	
gctggatcac ctcctttc	18

<210> 4
 <211> 18
 <212> DNA
 <213> Artificial

<220>
 <223> Artificial PCR sequence

<400> 4	
ctggtgcca ggcattcca	18

THE EMBODIMENTS OF THE INVENTION FOR WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A strain which is *Bifidobacterium longum* NCIMB 41290 (AHF5340) or *Bifidobacterium longum* NCIMB 41291 (AHF1231).
2. A composition comprising a strain of lactic acid bacteria of the genus *Bifidobacteria* and a carrier, wherein said strain is *Bifidobacterium longum* NCIMB 41290 (AHF5340) or *Bifidobacterium longum* NCIMB 41291 (AHF1231) or mixtures thereof.
3. The composition according to claim 2, wherein said composition is a companion animal food.
4. The composition according to claim 3, wherein the composition is a cat food.
5. The composition according to claim 3, wherein the composition is a wet animal food or a dry animal food.
6. The composition according to claim 5, wherein the composition is in a form selected from the group consisting of kibbles, chews, and biscuits.
7. The composition according to claim 2, wherein said composition is a companion animal food supplement.
8. The composition according to claim 7, wherein the composition is in the form of a gravy.
9. The composition according to claim 7, wherein the composition is in the form of a yoghurt, cheese, or other dairy produce.

10. The composition according to claim 7, wherein the composition is in a form selected from the group consisting of capsules, tablets, and pills.
11. The composition according to claim 2 comprising at least 0.001% of from about $1.0E+04$ CFU/g to about $1.0E+12$ CFU/g of the lactic acid bacteria.
12. The composition according to claim 2, wherein the lactic acid bacteria are in the form of viable cells.
13. The composition according to claim 2, wherein the lactic acid bacteria are in the form of non-viable cells, or the constituent active fractions thereof.
14. Use of a therapeutically effective amount of a strain of lactic acid bacteria of the genus *Bifidobacteria* for maintenance or improvement of health of a companion animal, wherein the maintenance or improvement of health is selected from the group consisting of regulation of or improvement of the immune system, maintenance or improvement of the health of the skin and/or coat system, amelioration or reduction of the effects of aging, prevention of weight loss during and following infection, treatment or prevention of gastrointestinal infection, treatment or prevention of urinary tract ailments, increase fibre digestion, and reduction of stress levels, wherein said strain is formulated for oral administration, and wherein said strain is *Bifidobacterium longum* NCIMB 41290 (AHF5340) or *Bifidobacterium longum* NCIMB 41291 (AHF1231) or mixtures thereof.
15. The use according to claim 14, for the regulation of or improvement of the immune system of the companion animal.
16. The use according to claim 15, for treatment or prevention of autoimmune disease in said companion animal.

17. The use according to claim 15, for treatment or prevention of inflammation in said companion animal.
18. The use according to claim 14, for maintenance or improvement of the health of the skin and/or coat system of said companion animal.
19. The use according to claim 18, for treatment or prevention of atopic disease of the skin of said companion animal.
20. The use according to claim 14, for amelioration or reduction of the effects of aging in said companion animal.
21. The use according to claim 14, for prevention of weight loss during and following infection in said companion animal.
22. The use according to claim 14, for treatment or prevention of gastrointestinal infection in said companion animal.
23. The use according to claim 22, for improvement of microbial ecology of companion animals.
24. The use according to claim 23, wherein said improvement comprises reduction of the levels of pathogenic bacteria in the faeces of said companion animal.
25. The use according to claim 24, wherein said pathogenic bacteria are selected from the group consisting of *Clostridia*, *Escherichia*, *Salmonella*, and mixtures thereof.
26. The use according to claim 14, for treatment or prevention of urinary tract ailments in said companion animal.

27. The use according to claim 26, wherein said urinary tract ailment comprises urinary tract infection.
28. The use according to claim 26, wherein said urinary tract ailment comprises urinary tract stones.
29. The use according to claim 26, wherein said *Bifidobacteria* increase the degradation of oxalic acid *in vitro*.
30. The use according to claim 14, to increase fibre digestion in said companion animal.
31. The use according to claim 14, to reduce stress levels in said companion animal.
32. The use according to claim 31, wherein said stress levels are measured by determining the level of stress hormones selected from the group consisting of epinephrine, norepinephrine, dopamine, Cortisol, C-reactive protein and mixtures thereof.
33. The use according to claim 14, wherein the strain is for feeding to said companion animal in any amount from 1.0E+04 to 1.0E+12 CFU/animal per day.
34. The use according to claim 14, wherein the companion animal is a cat.
35. Use of an effective amount of a strain of lactic acid bacteria of the genus *Bifidobacteria* to reduce odor associated with the feces or urine of a companion animal, wherein said strain is formulated for oral administration, and wherein said strain is *Bifidobacterium longum* NCIMB 41290 (AHF5340) or *Bifidobacterium longum* NCIMB 41291 (AHF1231) or mixtures thereof.

36. Use of an effective amount of a strain of lactic acid bacteria of the genus *Bifidobacteria* to reduce odor associated with a litter box of a companion animal, wherein said strain is formulated for oral administration, wherein said strain is *Bifidobacterium longum* NCIMB 41290 (AHF5340) or *Bifidobacterium longum* NCIMB 41291 (AHF1231) or mixtures thereof.
37. Use of a strain of lactic acid bacteria of the genus *Bifidobacteria* in the manufacture of a composition for maintenance or improvement of health of a companion animal, wherein the maintenance or improvement of health is selected from the group consisting of regulation of or improvement of the immune system, maintenance or improvement of the health of the skin and/or coat system, amelioration or reduction of the effects of aging, prevention of weight loss during and following infection, treatment or prevention of gastrointestinal infection, treatment or prevention of urinary tract ailments, increase fibre digestion, and reduction of stress levels, wherein said strain is formulated for oral administration, and wherein said strain is *Bifidobacterium longum* NCIMB 41290 (AHF5340) or *Bifidobacterium longum* NCIMB 41291 (AHF1231) or mixtures thereof.
38. The use according to claim 37, wherein said composition is for the regulation of or improvement of the immune system of the companion animal.
39. The use according to claim 38, wherein said composition is for treatment or prevention of autoimmune disease in said companion animal.
40. The use according to claim 38, wherein said composition is for treatment or prevention of inflammation in said companion animal.
41. The use according to claim 37, wherein said composition is for maintenance or improvement of the health of the skin and/or coat system of said companion animal.

42. The use according to claim 41, wherein said composition is for treatment or prevention of atopic disease of the skin of said companion animal.
43. The use according to claim 37, wherein said composition is for amelioration or reduction of the effects of aging in said companion animal.
44. The use according to claim 37, wherein said composition is for prevention of weight loss during and following infection in said companion animal.
45. The use according to claim 37, wherein said composition is for treatment or prevention of gastrointestinal infection in said companion animal.
46. The use according to claim 45, wherein said composition is for improvement of microbial ecology of companion animals.
47. The use according to claim 46, wherein said composition is for reduction of the levels of pathogenic bacteria in the faeces of said companion animal.
48. The use according to claim 47, wherein said pathogenic bacteria are selected from the group consisting of *Clostridia*, *Escherichia*, *Salmonella*, and mixtures thereof.
49. The use according to claim 37, wherein said composition is for treatment or prevention of urinary tract ailments in said companion animal.
50. The use according to claim 49, wherein said urinary tract ailment comprises urinary tract infection.
51. The use according to claim 49, wherein said urinary tract ailment comprises urinary tract stones.

52. The use according to claim 49, wherein said *Bifidobacteria* increase the degradation of oxalic acid *in vitro*.
53. The use according to claim 37, wherein said composition is to increase fibre digestion in said companion animal.
54. The use according to claim 37, wherein said composition is to reduce stress levels in said companion animal.
55. The use according to claim 54, wherein said stress levels are measured by determining the level of stress hormones selected from the group consisting of epinephrine, norepinephrine, dopamine, Cortisol, C-reactive protein and mixtures thereof.
56. The use according to claim 37, wherein said composition is for feeding to an animal in any amount from 1.0E+04 to 1.0E+12 CFU/animal per day.
57. The use according to claim 37, wherein the companion animal is a cat.
58. Use of a strain of lactic acid bacteria of the genus *Bifidobacteria* for in the manufacture of a composition to reduce odor associated with the feces or urine of a companion animal, wherein said strain is formulated for oral administration, wherein said strain is *Bifidobacterium longum* NCIMB 41290 (*AHF5340*) or *Bifidobacterium longum* NCIMB 41291 (*AHF1231*) or mixtures thereof.
59. Use of an effective amount of a strain of lactic acid bacteria of the genus *Bifidobacteria* for in the manufacture of a composition to reduce odor associated with a litter box of a companion animal, wherein said strain is formulated for oral administration, wherein said strain is *Bifidobacterium longum* NCIMB 41290 (*AHF5340*) or *Bifidobacterium longum* NCIMB 41291 (*AHF1231*) or mixtures thereof.