




(86) **Date de dépôt PCT/PCT Filing Date:** 2015/08/13
(87) **Date publication PCT/PCT Publication Date:** 2016/02/18
(45) **Date de délivrance/Issue Date:** 2024/06/11
(85) **Entrée phase nationale/National Entry:** 2017/02/13
(86) **N° demande PCT/PCT Application No.:** US 2015/045142
(87) **N° publication PCT/PCT Publication No.:** 2016/025764
(30) **Priorité/Priority:** 2014/08/13 (US62/036,936)

(51) **Cl.Int./Int.Cl. C12N 15/12** (2006.01),
A61K 31/7088 (2006.01), **A61K 38/37** (2006.01),
A61K 48/00 (2006.01), **A61P 7/04** (2006.01),
C07K 14/755 (2006.01), **C12N 15/85** (2006.01),
C12N 15/86 (2006.01), **C12N 15/864** (2006.01)
(72) **Inventeurs/Inventors:**
SABATINO, DENISE, US;
ELKOUBY, LIRON, US;
HIGH, KATHERINE A., US
(73) **Propriétaire/Owner:**
THE CHILDREN'S HOSPITAL OF PHILADELPHIA, US
(74) **Agent:** MBM INTELLECTUAL PROPERTY AGENCY

(54) **Titre : CASSETTE D'EXPRESSION AMELIOREE POUR LE CONDITIONNEMENT ET L'EXPRESSION DE VARIANTES DU
FACTEUR VIII POUR LE TRAITEMENT DES TROUBLES DE L'HEMOSTASE**
(54) **Title: AN IMPROVED EXPRESSION CASSETTE FOR PACKAGING AND EXPRESSION OF VARIANT FACTOR VIII FOR THE
TREATMENT OF HEMOSTASIS DISORDERS**



(57) **Abrégé/Abstract:**

Factor VIII variants and methods of use thereof are disclosed. In particular embodiments, Factor VIII variants are expressed more efficiently by cells over wild-type Factor VIII proteins, are secreted at increased levels by cells over wild-type Factor VIII proteins, exhibit enhanced activity over wild-type Factor VIII proteins and are packaged more efficiently into viral vectors.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2016/025764 A3

(43) International Publication Date
18 February 2016 (18.02.2016)

WIPO | PCT

(51) International Patent Classification:

A61K 38/37 (2006.01) C12N 15/00 (2006.01)
C07K 14/755 (2006.01)

(21) International Application Number:

PCT/US2015/045142

(22) International Filing Date:

13 August 2015 (13.08.2015)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

62/036,936 13 August 2014 (13.08.2014) US

(71) Applicant: THE CHILDREN'S HOSPITAL OF PHILADELPHIA [US/US]; 34th Street & Civic Center Boulevard, Philadelphia, Pennsylvania 19104 (US).

(72) Inventor; and

(71) Applicant : SABATINO, Denise [US/US]; 28 Llandillo Road, Havertown, Pennsylvania 19083 (US).

(74) Agent: BEDGOOD, Robert M.; Pillsbury Winthrop Shaw Pittman LLP, P.O. Box 10500, McLean, Virginia 22101 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,

BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))

(88) Date of publication of the international search report:

7 April 2016

(54) Title: AN IMPROVED EXPRESSION CASSETTE FOR PACKAGING AND EXPRESSION OF VARIANT FACTOR VIII FOR THE TREATMENT OF HEMOSTASIS DISORDERS




Fig. 6B

(57) Abstract: Factor VIII variants and methods of use thereof are disclosed. In particular embodiments, Factor VIII variants are expressed more efficiently by cells over wild-type Factor VIII proteins, are secreted at increased levels by cells over wild-type Factor VIII proteins, exhibit enhanced activity over wild-type Factor VIII proteins and are packaged more efficiently into viral vectors.

WO 2016/025764 A3

AN IMPROVED EXPRESSION CASSETTE FOR PACKAGING AND EXPRESSION OF VARIANT FACTOR VIII FOR THE TREATMENT OF HEMOSTASIS DISORDERS

5 [0001]

Field of the Invention

[0002] This invention relates to the fields of recombinant coagulation factor production and
10 the treatment of medical disorders associated with aberrant hemostasis. More particularly, the
invention provides an improved expression cassette which more efficiently packages the Factor
VIII variant sequence of the invention, the improved variant also exhibiting enhanced activity
over wild-type Factor VIII proteins.

Introduction

15 [0003] Several publications and patent documents are cited throughout the specification in
order to describe the state of the art to which this invention pertains. Each of these citations is
incorporated herein by reference as though set forth in full.

[0004] Hemophilia is an X-linked bleeding disorder present in 1 in 5,000 males worldwide.
Therapies aimed at increasing clotting factor levels just above 1% of normal are associated with
20 substantial improvement of the severe disease phenotype. Recent clinical trials for AAV-
mediated gene transfer for hemophilia B (HB) have demonstrated sustained long-term expression
of therapeutic levels of factor IX (FIX) but established that the AAV vector dose may be limiting
due to anti-AAV immune responses to the AAV capsid. While these data relate the hemophilia
B, 80% of all hemophilia is due to FVIII deficiency, hemophilia A (HA).

25 [0005] Current treatment for this disease is protein replacement therapy that requires frequent
infusion of the factor VIII protein. There is an immediate need to achieve sustained therapeutic
levels of factor VIII expression so that patients no longer require such frequent protein
treatments. Indeed, continuous factor VIII expression would prevent bleeding episodes and may
ensure that immune tolerance to the protein is established.

30 [0006] In summary, gene therapy for HA presents *3 distinct challenges*: (1) intrinsic
properties of *human* FVIII (hFVIII) make it difficult to express compared to other proteins of
similar size (2) the large size of the FVIII cDNA and sequence specific effects are correlated with
rearrangements which hamper AAV production and (3) high rates of anti-FVIII antibody

(inhibitors) formation in response to protein therapy that occurs in 25-30% of severe (<1% FVIII) HA patients.

[0007] The invention provides improved Factor VIII variants useful for treatment in patients in need thereof, such as a patient with HA.

5

Summary

[0008] In accordance with the invention, codon-optimized Factor VIII (FVIII) encoding nucleic acid variants distinct from wild-type nucleic acid encoding wild-type FVIII are provided. Such codon-optimized FVIII encoding nucleic acid variants encode FVIII protein, optionally without a B domain, such as FVIII protein (e.g., human FVIII protein) without a B domain (see, e.g., Fig. 1). Such codon-optimized FVIII encoding nucleic acid variants exhibit increased expression, 4-5 fold for a particular codon-optimized sequence (CO/CO3), when transferred into a cell (see, e.g., Fig. 6B), leading to increased FVIII protein secretion and therefore activity (see, e.g., Fig. 7). Such codon-optimized FVIII encoding nucleic acid variants also are packaged more efficiently in different AAV vector serotypes, leading to increased production of AAV vectors across AAV serotypes (see, e.g., Tables 3-5).

[0009] Surprisingly, these features dramatically increase FVIII expression, as compared to expression of wild-type nucleic acid encoding wild-type FVIII, or expression of wild type FVIII without B domain, and also increase packaging efficiency into an AAV vector, leading to higher vector yields. Codon-optimized FVIII encoding sequences may exhibit a reduction in rearrangement events that impact the integrity of the FVIII transgene after AAV packaging that is observed with wild type FVIII and other codon-optimized hFVIII transgenes, e.g., CO/CO3 is superior to either of CO1 and CO2. In addition, non-codon optimized and codon-optimized FVIII variants that have a single amino acid change or small deletion (1-4 amino acids) at a PACE-furin cleavage site can exhibit increased secretion *in vivo* (non-codon optimized Fig. 3B, and codon-optimized Fig. 6B), and specific activity of the FVIII protein. Invention FVIII encoding nucleic acid variants, that encode FVIII with or without B-domain, and/or with or without mutated PACE-furin cleavage recognition site can increase expression of FVIII, increase production of AAV vector, as well as provide increased efficacy in the context of gene transfer by increased circulating levels of FVIII and achieving hemostasis with lower FVIII thereby reducing the effective dose required for beneficial therapeutic outcomes.

[0010] In one embodiment, a Factor VIII (FVIII) encoding nucleic acid variant exhibits greater expression when compared to wild type FVIII expression. In another embodiment, a Factor VIII (FVIII) encoding nucleic acid variant comprises a B domain deletion and exhibits greater expression when compared to wild type FVIII comprising a B domain deletion

expression. In an additional embodiment, FVIII encoding nucleic acid variant is more efficiently packaged into a viral vector such as an adenovirus-associated virus (AAV) vector.

[0011] In a further embodiment, a FVIII encoded by said nucleic acid variant exhibits greater biological activity when compared to wild type FVIII or when compared to wild type FVIII comprising a B domain deletion (e.g., as determined by a clotting assay or reduced bleeding in a FVIII assay or FVIII deficiency model).

[0012] In still further embodiments, codon optimized FVIII encoding nucleic acid variants encode FVIII protein with a mutated PACE-furin cleavage recognition site, referred to as a variant FVIII or FVIII protein. Such codon-optimized FVIII nucleic acid variants that encode FVIII protein with mutated PACE-furin cleavage recognition site exhibit increased secretion and blood clotting activity (see, e.g., Figs. 2, 6 and 7). In particular aspects, FVIII encoding nucleic acid variants have 1, 2, 3 or all 4 of the codons encoding the PACE/furin cleavage site of FVIII substituted or deleted. In such aspects, variant FVIII proteins encoded by such FVIII nucleic acid variants have 1, 2, 3 or all 4 of the PACE/furin cleavage site of FVIII substituted or deleted. In more particular aspects, 1, 2, 3 or all 4 of the codons encoding the PACE/furin cleavage site set forth as HHQR or RHQR from positions 1645-1648 is/are deleted in FVIII encoding nucleic acid variants.

[0013] In additional aspects, FVIII encoding nucleic acid variants have one or more leucine codons changed to CTG compared to TTA, TTG, CTT, CTC or CTA in wild type FVIII encoding nucleic acid. In further aspects, FVIII encoding nucleic acid variants have 2-5, 5-10, 10-20, 20-50, 50-100, 100-250, 250-500, 500-750 or 750-850 CTG leucine codons modified from TTA, TTG, CTT, CTC or CTA leucine codons in wild type FVIII encoding nucleic acid. In yet additional aspects, FVIII encoding nucleic acid variants have greater than 85% CTG leucine codons modified from TTA, TTG, CTT, CTC or CTA leucine codons in wild type FVIII encoding nucleic acid. In still further aspects, FVIII encoding nucleic acid variants have all CTG leucine codons modified from TTA, TTG, CTT, CTC or CTA leucine codons in wild type FVIII encoding nucleic acid. In particular aspects, FVIII encoding nucleic acid variants have between about 50-59%, or 50-56%, or 50-53% GC content. In other aspects, FVIII encoding nucleic acid variants have one or more AAG lysine codons compared to AAA lysine codons in wild type FVIII encoding nucleic acid.

[0014] In still additional embodiments, FVIII encoding nucleic acid variant are at least 75% identical to wild type human FVIII nucleic acid or wild type human FVIII nucleic acid comprising a B domain deletion. In more particular aspects, FVIII encoding nucleic acid variant are about 75-85% identical (e.g., about 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%

or 85% identical) to wild type human FVIII nucleic acid or wild type human FVIII nucleic acid comprising a B domain deletion.

[0015] In particular aspects, FVIII encoding nucleic acid variants are set forth in any of SEQ ID NOs:1-7 and 9.

5 [0016] In various embodiments, FVIII encoding nucleic acid variants are mammalian, such as human. Such mammalian FVIII encoding nucleic acid variants including human forms may be based upon wild type FVIII or wild type FVIII comprising a B domain deletion.

[0017] In accordance with the invention, also provided are expression vectors that include FVIII encoding nucleic acid variants as set forth herein. In particular embodiments, an
10 expression vector comprises an adenovirus-associated virus (AAV) vector, a retroviral vector, an adenoviral vector, a plasmid, or a lentiviral vector. In particular aspects, an AAV vector comprises an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 or AAV-2i8 AAV serotype.

[0018] Expression vectors can include additional components or elements. In particular
15 embodiments, an expression vector such as AAV vector further includes an intron, an expression control element, one or more AAV inverted terminal repeats (ITRs) (e.g., any of: AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 or AAV-2i8 AAV serotypes, or a combination thereof) and/or a filler polynucleotide sequence. In particular aspects, an intron is within or flanks a FVIII encoding nucleic acid variant, and/or an
20 expression control element is operably linked to the FVIII encoding nucleic acid variant, and/or an AAV ITR(s) flanks the 5' or 3' terminus of the FVIII encoding nucleic acid variant, and/or a filler polynucleotide sequence flanks the 5' or 3' terminus of the FVIII encoding nucleic acid variant.

[0019] In particular aspects, an expression control element comprises a constitutive or
25 regulatable control element, or a tissue-specific expression control element or promoter. In more particular aspects, an expression control element comprises an element that confers expression in liver. In additional particular aspects, an expression control element comprises a TTR promoter or mutant TTR promoter, such as SEQ ID NO:8.

[0020] In accordance with the invention, additionally provided are host cells expressing the
30 FVIII encoded by the nucleic acid variants as set forth herein. In particular embodiments, a host cell includes FVIII encoding nucleic acid variant or an expression vector comprising a FVIII encoding nucleic acid variant. In particular aspects, such host cells produce FVIII protein encoded by the nucleic acid variants and FVIII protein produced is recovered. Such FVIII protein produced by the cells, optionally isolated and/or purified, can be administered to a subject
35 as set forth herein.

[0021] In accordance with the invention, further provided are virus vectors that include the FVIII encoding nucleic acid variants or the expression vectors comprising the FVIII encoding nucleic acid variants. In particular embodiments, a virus vector comprises an AAV vector. In particular aspects, an AAV vector comprises a VP1, VP2 and/or VP3 capsid sequence having
 5 75% or more sequence identity (*e.g.*, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, 99.6%, 99.7%, 99.8%, etc.) to AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 or AAV-2i8 VP1, VP2 and/or VP3 sequences. In more particular aspects, an AAV vector comprises a VP1, VP2 and/or VP3 capsid sequence selected from any of: AAV1, AAV2, AAV3, AAV4, AAV5, AAV6,
 10 AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 and AAV-2i8 AAV serotypes.

[0022] In accordance with the invention, yet additionally provided are compositions comprising FVIII encoding nucleic acid variants set forth herein. In particular embodiments, pharmaceutical compositions include an expression vector, or a virus or AAV vector, in a biologically compatible carrier or excipient. Such pharmaceutical compositions optionally
 15 include empty capsid AAV (*e.g.*, lack vector genome comprising FVIII encoding nucleic acid variant). In additional particular embodiments, FVIII encoding nucleic acid variants, expression vectors, or virus or AAV vectors are encapsulated in a liposome or mixed with phospholipids or micelles.

[0023] In accordance with the invention, still further provided are methods for delivering or
 20 transferring FVIII encoding nucleic acid variants into a mammal or a mammalian cell. In one embodiment, a method includes administering or contacting a FVIII encoding nucleic acid variant, an expression vector comprising FVIII encoding nucleic acid variant, or a virus or AAV vector comprising a FVIII encoding nucleic acid variant to a mammal or mammalian cell, thereby delivering or transferring the nucleic acid sequence into the mammal or mammalian cell. Such
 25 methods introduce FVIII encoding nucleic acid variants into a mammalian cell in culture or in a subject (*e.g.*, a patient).

[0024] Methods of the invention also include treating mammalian subjects (*e.g.*, patients) such as humans in need of Factor VIII (the human produces an insufficient amount of Factor VIII protein, or a defective or aberrant Factor VIII protein). In one embodiment, a method of treating
 30 a mammal in need of Factor VIII, includes: providing a FVIII encoding nucleic acid variant, or an expression vector comprising FVIII encoding nucleic acid variant, or a virus or AAV vector comprising a FVIII encoding nucleic acid variant; and administering an amount of the FVIII encoding nucleic acid variant, or an expression vector comprising FVIII encoding nucleic acid variant, or a virus or AAV vector comprising a FVIII encoding nucleic acid variant to the

mammalian subject such that Factor VIII encoded by the FVIII encoding nucleic acid variant, is expressed in the mammalian subject.

[0025] In another embodiment, a method for treatment of a hemostasis related disorder in a patient in need thereof (e.g., the patient produces an insufficient amount of Factor VIII protein, or a defective or aberrant Factor VIII protein) includes administration of a therapeutically effective amount of a FVIII encoding nucleic acid variant, or an expression vector comprising FVIII encoding nucleic acid variant, or a virus or AAV vector comprising a FVIII encoding nucleic acid variant in a biologically acceptable carrier to the patient.

[0026] In particular aspects of the invention methods, Factor VIII is expressed at levels having a beneficial or therapeutic effect on the mammal; and/or Factor VIII is expressed in a cell, tissue or organ of the mammal. Such aspects include introduction of FVIII encoding nucleic acid variant into a tissue or organ such as liver. Such aspects also include introduction of FVIII encoding nucleic acid variant into a secretory cell. Such aspects further include introduction of FVIII encoding nucleic acid variant into an endocrine cell or an endothelial cell. Such aspects additionally include introduction of FVIII encoding nucleic acid variant into an hepatocyte, a sinusoidal endothelial cell, a megakaryocyte, a platelet or hematopoietic stem cell.

[0027] Candidate subjects (e.g., a patient) and mammals (e.g., humans) for administration (e.g., delivery) of a FVIII encoding nucleic acid variant, or an expression vector comprising FVIII encoding nucleic acid variant, or a virus or AAV vector comprising a FVIII encoding nucleic acid variant include those having or those at risk of having a disorder such as: hemophilia A, von Willebrand diseases and bleeding associated with trauma, injury, thrombosis, thrombocytopenia, stroke, coagulopathy, disseminated intravascular coagulation (DIC) or over-anticoagulation treatment disorder.

[0028] Candidate subjects (e.g., a patient) and mammals (e.g., humans) for administration (e.g., delivery) of a FVIII encoding nucleic acid variant, or an expression vector comprising FVIII encoding nucleic acid variant, or a virus or AAV vector comprising a FVIII encoding nucleic acid variant include those or sero-negative for AAV antibodies, as well as those having or those at risk of developing AAV antibodies. Such subjects (e.g., a patient) and mammals (e.g., humans) may be sero-negative or sero-positive for an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV-Rh10 or AAV-Rh74 serotype.

[0029] Methods of the invention therefore further include administering empty capsid AAV to said mammal or said patient. In particular embodiments, empty capsid of AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV-12, AAV-Rh10 and/or AAV-Rh74 serotype is further administered to the mammal or patient.

[0030] Methods of administration (e.g., delivery) in accordance with the invention include any mode of contact or delivery, *ex vivo* or *in vivo*. In particular embodiments administration (e.g., delivery) is: intravenously, intraarterially, intramuscularly, subcutaneously, intra-cavity, intubation, or via catheter.

5 [0031] The invention also provide methods for testing the improved FVIII variants in small and large animal models that are tolerant to human FVIII in order to assess dosing and monitor immunogenicity of the variants. The HB dog model is superior for predicting the effective dose required for human FIX gene therapy. Moreover, use of this model provides a setting that allows assessment of humans currently receiving protein replacement therapy with hFVIII-BDD without
10 evidence of an anti-hFVIII antibody response who are likely to develop an immune response to such variants.

Description of Drawings

[0032] **Figure 1 shows** proteolytic cleavages of factor VIII. FVIII has a domain structure consisting of three A domains (A1, A2, and A3), two C domains (C1 and C2), and a large B
15 domain encoded by exon 14 of the gene that is dispensable for FVIII function. FVIII functions as a cofactor in the activation of factor X to factor Xa in a reaction catalyzed by factor IXa. The FVIII is secreted as a heterodimer composed of FVIII heavy and light chains that are stabilized in the circulation by von Willebrand factor (vWF). Upon thrombin cleavage, FVIII forms the activated heterotrimer.

20 [0033] **Figure 2 shows** Factor VIII Expression after AAV Delivery of codon-optimized factor VIII. The codon-optimized human factor VIII gene (CO) was delivered in an adeno-associated viral vector serotype 8 to hemophilia A/CD4 knockout mice. The mice were delivered 1×10^{11} vector genomes (vg)/ mouse of hFVIII-SQ (wild type factor VIII cDNA), TTR-hFVIII-SQ-CO, TTRmut-hFVIII-SQ-CO or TTRmut-hFVIII-SQ-CO-delP/F. ELISA was performed on
25 plasma collected at 2, 4, 8 and 12 weeks post vector administration to determine the factor VIII protein levels in the circulation.

[0034] **Figure 3A shows** non-codon optimized AAV-hFVIII constructs. The AAV-hFVIII vectors have the wild type B-domain deleted (BDD) human factor VIII cDNA. The enhancer/promoter element is a minimal hepatic control region-human α -1 antitrypsin
30 enhancer/promoter (HCR-hAAT) that has a shortened version of the hepatic control region of the human apolipoprotein E gene locus (Nathwani, et al. 2006, Sabatino et al. 2011). A 65-bp SV40 intron and a 134-bp SV40 polyadenylation are also included. The only difference between these constructs is that one utilizes the wild type hFVIII-BDD (top construct) and the other

utilizes the wild type hFVIII-BDD with a PACE-furin deletion variant (bottom construct). All five P/F deletion variants (Table 1) were introduced into this expression cassette.

[0035] **Figure 3B shows** FVIII expression after AAV8 delivery of hFVIII PACE-furin deletion variants. AAV (5×10^{11} vg/mouse) was delivered to hemophilia A/CD4 knockout (immunodeficient) mice. The hFVIII expression (antigen and activity) was followed in the plasma through 12 weeks after AAV administration. * = $p < 0.05$

[0036] **Figure 4A shows** human wild-type and codon optimized FVIII transgene constructs. TTR promoter drives expression of codon-optimized FVIII. SQ, the B-domain deleted form that has a S743 and Q1638 at the junction; ITR, inverted terminal repeat sequences required for packaging of the transgene into the AAV vectors.

[0037] **Figure 4B shows** that codon optimization of hFVIII leads to higher expression levels. A comparison of CO1 FVIII, CO2 FVIII, and wild-type FVIII AAV delivered FVIII expression and activity levels at 4 and 8 weeks post AAV vector administration.

[0038] **Figure 4C shows** that codon optimization of hFVIII leads to higher expression levels. A comparison of CO1 FVIII, CO3/CO FVIII, and wild-type FVIII AAV delivered FVIII expression and activity levels at 4 and 8 weeks post AAV vector administration.

[0039] **Figure 5A shows** FVIII expression after AAV delivery of CO3. Comparison of the TTR promoter and the TTR mutant (TTRm) promoter with FVIII-CO3. The TTR mutant promoter with the PACE-furin deletion $\Delta 4$ (delP/F) introduced into CO3.

[0040] **Figure 5B shows** data indicating that the TTR mutant promoter results in increased FVIII-CO3 expression after AAV delivery.

[0041] **Figure 6A shows** human FVIII transgene constructs. The transgene constructs utilize a modified transthyretin promoter (TTRm), synthetic intron (Intron), and polyadenylation signal (Poly A). The factor VIII transgenes are the wild type B-domain deleted human factor VIII cDNA (WT FVIII) or the codon-optimized B-domain deleted human factor VIII cDNA (CO FVIII). The PACE-furin deletion variants $\Delta 3$ or $\Delta 4$ were introduced into the hFVIII transgenes. $\Delta 3$, deletion of three residues (amino acid 1645-1647) of the PACE-furin cleavage recognition site. $\Delta 4$, deletion of four residues (amino acid 1645-1648) of the PACE-furin cleavage recognition site. SQ, the B-domain deleted form that has a S743 and Q1638 at the junction; ITR, inverted terminal repeat sequences required for packaging of the transgene into the AAV vectors.

[0042] **Figure 6B shows** Factor VIII expression after AAV delivery of wild type and codon-optimized FVIII constructs. The hFVIII antigen levels were determined by a human FVIII specific ELISA (Affinity Biologicals). FVIII activity determined by Coatest assay correlated with the FVIII antigen levels. These data are presented as fold difference compared to the TTRmut-hFVIII-wtSQ at the 2 and 4 week time points.

[0043] **Figure 7** shows *in vivo* hemostatic challenge of mice after AAV delivery of wild type and codon-optimized hFVIII constructs. At 6 weeks post vector administration (1x10¹¹vg/mouse), the mice treated with AAV-hFVIII constructs were challenged using a tail clip assay (Ivanciu L et al. Nature Biotech 2011, 29(11):1028-1033). Mice were anesthetized with isoflurane and the tail was pre-warmed to 37°C followed by tail transection at a diameter of 3mm. Blood was collected for 10 minutes in 14ml of saline at 37°C. Total blood loss (μl) was determined by measuring total hemoglobin by absorbance at 575nm following red cell lysis. Blood loss of the hemophilia A mice that were treated with AAV-hFVIII constructs was compared to the blood loss from untreated hemophilia A mice and wild type mice.

Detailed Description

[0044] Previous reports have shown that AAV liver gene therapy of canine FVIII (cFVIII) results in long-term expression of therapeutic levels of FVIII in naïve HA dogs (Sabatino et al. Mol Ther 19, 442-449, 2011). This strategy also eradicated inhibitors to cFVIII in HA dogs with pre-existing neutralizing cFVIII antibodies (Finn, J.D. et al., Blood 116, 5842-5848, 2010). These data demonstrate that AAV-FVIII can significantly improve the disease phenotype and eliminate FVIII inhibitors. Interestingly, cFVIII is inherently more stable than human FVIII (hFVIII), resulting in increased biological activity (Sabatino, D.E. et al. Blood 114, 4562-4565, 2009). Thus, the AAV dose of cFVIII does not predict the therapeutic dose of hFVIII. This is in contrast to factor IX that has similar biological activity in both species and predicted the therapeutic AAV dose in humans. Notably, the hemophilia dog model has been an excellent predictor of both efficacy and immune response to the transgene in liver and muscle trials for HB.

[0045] To understand the biochemical basis for enhanced function of cFVIII, we identified a PACE-furin cleavage recognition site in FVIII that is unique in cFVIII compared to all other species, including hFVIII. Introduction of a single amino acid change at this site in hFVIII confers increased stability and higher biological activity to hFVIII (Siner, J.I., et al. Blood 121, 4396-4403, 2013).

[0046] Disclosed herein are nucleic acid variants that encode human FVIII, distinct from wild-type nucleic acid that encode FVIII. Such nucleic acid variants that encode human FVIII are expressed at increased levels in cells and/or animals, which in turn can provide increased FVIII protein levels *in vivo*. Also disclosed herein are nucleic acid variants that encode human FVIII that can be packaged more efficiently by AAV across different serotypes. Further disclosed herein are nucleic acid variants that encode human FVIII protein variants having higher stability and/or biological activity *in vitro* and/or *in vivo*.

[0047] Still further disclosed herein are data showing efficacy and safety of AAV delivery of several variants of human FVIII in animal models of hemophilia A. In particular, for example, codon-optimized nucleic acid that encode human FVIII with and without deletions in the intracellular protease cleavage recognition site exhibit one or more of the following: 1) increased expression in cells and/or animals; 2) increased activity, as reflected by increased clotting, for example; 3) increased stability; and 4) achieve therapeutic effect at lower AAV doses than native hFVIII, with no apparent increase in immunogenicity over the currently used B-domain deleted FVIII constructs.

[0048] The terms “polynucleotide” and “nucleic acid” are used interchangeably herein to refer to all forms of nucleic acid, oligonucleotides, including deoxyribonucleic acid (DNA) and ribonucleic acid (RNA). Polynucleotides include genomic DNA, cDNA and antisense DNA, and spliced or unspliced mRNA, rRNA tRNA and inhibitory DNA or RNA (RNAi, *e.g.*, small or short hairpin (sh)RNA, microRNA (miRNA), small or short interfering (si)RNA, trans-splicing RNA, or antisense RNA). Polynucleotides include naturally occurring, synthetic, and intentionally modified or altered polynucleotides (*e.g.*, variant nucleic acid). Polynucleotides can be single, double, or triplex, linear or circular, and can be of any length. In discussing polynucleotides, a sequence or structure of a particular polynucleotide may be described herein according to the convention of providing the sequence in the 5' to 3' direction.

[0049] As used herein, the terms “modify” or “variant” and grammatical variations thereof, mean that a nucleic acid, polypeptide or subsequence thereof deviates from a reference sequence. Modified and variant sequences may therefore have substantially the same, greater or less expression, activity or function than a reference sequence, but at least retain partial activity or function of the reference sequence. A particular example of a modification or variant is a codon-optimized nucleic acid sequence that encodes FVIII.

[0050] A “nucleic acid” or “polynucleotide” variant refers to a modified sequence which has been genetically altered compared to wild-type. The sequence may be genetically modified without altering the encoded protein sequence. Alternatively, the sequence may be genetically modified to encode a variant protein. A nucleic acid or polynucleotide variant can also refer to a combination sequence which has been codon modified to encode a protein that still retains at least partial sequence identity to a reference sequence, such as wild-type protein sequence, and also has been codon-modified to encode a variant protein. For example, some codons of such a nucleic acid variant will be changed without altering the amino acids of the protein (FVIII) encoded thereby, and some codons of the nucleic acid variant will be changed which in turn changes the amino acids of the protein (FVIII) encoded thereby.

[0051] The term “variant Factor VIII (FVIII)” refers to a modified FVIII which has been genetically altered such that the encoded protein exhibits increased expression when compared to when compared to unmodified wild-type FVIII or FVIII-BDD. Such a variant can be referred to as a “Factor VIII (FVIII) encoding nucleic acid variant.” An increase of at least 30-fold was attained with the TTR_{mut}-hFVIII-SQ-CO delP/F expression cassette compared to HCR/hAAT promoter driving wild-type hFVIII (Fig. 2). Particular examples of such FVIII modifications are codon-optimized FVIII encoding variants which exhibit increased expression and/or increased packaging efficiency into AAV vectors compared to non codon-optimized wild-type FVIII or FVIII-BDD, and do so without modifications to the nucleic acid that result in amino acid changes to the encoded FVIII protein. When comparing expression, if the variant FVIII protein retains the B-domain, it is appropriate to compare it to wild-type FVIII expression; and if the variant FVIII protein has a B-domain deletion, it be compared to expression of wild-type FVIII that also has a B-domain deletion.

[0052] A “variant Factor VIII (FVIII)” can also mean a modified FVIII protein such that the modified protein has an amino acid alteration compared to wild-type FVIII and exhibits an increase in activity and/or stability compared to wild-type FVIII. Examples of such particular FVIII protein modifications are genetic modifications that lead to PACE-furin cleavage recognition site mutations, deletions or substitutions in the FVIII protein, referred to herein as a variant FVIII protein. When comparing activity and/or stability, if the variant FVIII protein retains the B-domain, it is appropriate to compare it to wild-type FVIII; and if the variant FVIII protein has a B-domain deletion, it be compared to wild-type FVIII that also has a B-domain deletion.

[0053] The nucleotide sequences described herein are readily obtainable from GenBank. For human FVIII, see Accession No. NG-011403.1. For canine FVIII, see Accession No. NM-001003212-1. FVIII-BDD, hFVIII-BDD, cFVIII-BDD and the like refer to a FVIII variant which lacks the B domain (see, e.g., Fig. 1).

[0054] The “polypeptides,” “proteins” and “peptides” encoded by the “nucleic acid” or “polynucleotide” sequences,” include full-length native (FVIII) sequences, as with naturally occurring wild-type proteins, as well as functional subsequences, modified forms or sequence variants so long as the subsequence, modified form or variant retain some degree of functionality of the native full-length protein. For example, a FVIII protein can have a B-domain deletion as set forth herein and retain clotting function. In methods and uses of the invention, such polypeptides, proteins and peptides encoded by the nucleic acid sequences can be but are not required to be identical to the endogenous protein that is defective, or whose expression is insufficient, or deficient in the treated mammal.

[0055] Non-limiting examples of modifications include one or more nucleotide or amino acid substitutions (*e.g.*, 1-3, 3-5, 5-10, 10-15, 15-20, 20-25, 25-30, 30-40, 40-50, 50-100, 100-150, 150-200, 200-250, 250-500, 500-750, 750-850 or more nucleotides or residues). An example of a nucleic acid modification is codon optimization, *e.g.*, for a leucine codon that is not CTG to be modified to CTG, or a lysine codon that is not AAG to be modified to AAG. Another example of a nucleic acid codon optimization modification is increasing GC content. In particular aspects, a variant nucleic acid sequence encoding human FVIII protein has 1-5% more GC content than native sequence encoding human Factor FVIII (*e.g.*, 1, 2, 3, 4 or 5% more GC content); or has 5-10% more GC content than native (wild-type) sequence encoding human Factor FVIII (*e.g.*, 5, 6, 7, 8, 9 or 10% more GC content); or has 10-15% more GC content than native (wild-type) sequence encoding human Factor FVIII (*e.g.*, 10, 11, 12, 13, 14 or 15% more GC content).

[0056] An example of an amino acid modification is a conservative amino acid substitution or a deletion (*e.g.*, subsequences or fragments, or deletion of PACE/furin cleavage site) of a reference sequence, *e.g.* FVIII. In particular embodiments, a modified or variant sequence retains at least part of a function or activity of unmodified sequence.

[0057] All mammalian and non-mammalian forms of nucleic acid encoding proteins, including other mammalian forms of the FVIII nucleic acid and FVIII proteins disclosed herein are expressly included, either known or unknown. Thus, the invention includes genes and proteins from non-mammals, mammals other than humans, and humans, which genes and proteins function in a substantially similar manner to the FVIII (*e.g.*, human) genes and proteins described herein.

[0058] The term “vector” refers to small carrier nucleic acid molecule, a plasmid, virus (*e.g.*, AAV vector), or other vehicle that can be manipulated by insertion or incorporation of a nucleic acid. Such vectors can be used for genetic manipulation (*i.e.*, “cloning vectors”), to introduce/transfer polynucleotides into cells, and to transcribe or translate the inserted polynucleotide in cells. An “expression vector” is a specialized vector that contains a gene or nucleic acid sequence with the necessary regulatory regions needed for expression in a host cell. A vector nucleic acid sequence generally contains at least an origin of replication for propagation in a cell and optionally additional elements, such as a heterologous polynucleotide sequence, expression control element (*e.g.*, a promoter, enhancer), intron, ITR(s), selectable marker (*e.g.*, antibiotic resistance), polyadenylation signal.

[0059] A viral vector is derived from or based upon one or more nucleic acid elements that comprise a viral genome. Particular viral vectors include lentivirus, pseudo-typed lentivirus and parvo-virus vectors, such as adeno-associated virus (AAV) vectors. Also provided are vectors comprising a nucleic acid sequence encoding a variant FVIII polypeptide.

[0060] The term “recombinant,” as a modifier of vector, such as recombinant viral, e.g., lenti- or parvo-virus (*e.g.*, AAV) vectors, as well as a modifier of sequences such as recombinant polynucleotides and polypeptides, means that the compositions have been manipulated (*i.e.*, engineered) in a fashion that generally does not occur in nature. A particular example of a recombinant vector, such as an AAV vector would be where a polynucleotide that is not normally present in the wild-type viral (*e.g.*, AAV) genome is inserted within the viral genome. An example of a recombinant polynucleotide would be where a nucleic acid (*e.g.*, gene) encoding a FVIII protein is cloned into a vector, with or without 5', 3' and/or intron regions that the gene is normally associated within the viral (*e.g.*, AAV) genome. Although the term “recombinant” is not always used herein in reference to vectors, such as viral and AAV vectors, as well as sequences such as polynucleotides, recombinant forms including polynucleotides, are expressly included in spite of any such omission.

[0061] A recombinant viral “vector” or “AAV vector” is derived from the wild type genome of a virus, such as AAV by using molecular methods to remove the wild type genome from the virus (*e.g.*, AAV), and replacing with a non-native nucleic acid, such as a FVIII encoding nucleic acid variant sequence. Typically, for AAV one or both inverted terminal repeat (ITR) sequences of AAV genome are retained in the AAV vector. A “recombinant” viral vector (*e.g.*, AAV) is distinguished from a viral (*e.g.*, AAV) genome, since all or a part of the viral genome has been replaced with a non-native sequence with respect to the viral (*e.g.*, AAV) genomic nucleic acid such as FVIII encoding nucleic acid variant sequence. Incorporation of a non-native sequence therefore defines the viral vector (*e.g.*, AAV) as a “recombinant” vector, which in the case of AAV can be referred to as a “rAAV vector.”

[0062] A recombinant vector (*e.g.*, lenti-, parvo-, AAV) sequence can be packaged- referred to herein as a “particle” for subsequent infection (transduction) of a cell, *ex vivo*, *in vitro* or *in vivo*. Where a recombinant vector sequence is encapsidated or packaged into an AAV particle, the particle can also be referred to as a “rAAV.” Such particles include proteins that encapsidate or package the vector genome. Particular examples include viral envelope proteins, and in the case of AAV, capsid proteins.

[0063] A vector “genome” refers to the portion of the recombinant plasmid sequence that is ultimately packaged or encapsidated to form a viral (*e.g.*, AAV) particle. In cases where recombinant plasmids are used to construct or manufacture recombinant vectors, the vector genome does not include the portion of the “plasmid” that does not correspond to the vector genome sequence of the recombinant plasmid. This non vector genome portion of the recombinant plasmid is referred to as the “plasmid backbone,” which is important for cloning and amplification of the plasmid, a process that is needed for propagation and recombinant virus

production, but is not itself packaged or encapsidated into virus (*e.g.*, AAV) particles. Thus, a vector “genome” refers to the nucleic acid that is packaged or encapsidated by virus (*e.g.*, AAV).

[0064] A “transgene” is used herein to conveniently refer to a nucleic acid that is intended or has been introduced into a cell or organism. Transgenes include any nucleic acid, such as a gene that encodes a polypeptide or protein (*e.g.*, Factor VIII).

[0065] In a cell having a transgene, the transgene has been introduced/transferred by way of vector, such as AAV, “transduction” or “transfection” of the cell. The terms “transduce” and “transfect” refer to introduction of a molecule such as a nucleic acid into a cell or host organism. The transgene may or may not be integrated into genomic nucleic acid of the recipient cell. If an introduced nucleic acid becomes integrated into the nucleic acid (genomic DNA) of the recipient cell or organism it can be stably maintained in that cell or organism and further passed on to or inherited by progeny cells or organisms of the recipient cell or organism. Finally, the introduced nucleic acid may exist in the recipient cell or host organism extrachromosomally, or only transiently.

[0066] A “transduced cell” is a cell into which the transgene has been introduced. Accordingly, a “transduced” cell (*e.g.*, in a mammal, such as a cell or tissue or organ cell), means a genetic change in a cell following incorporation of an exogenous molecule, for example, a nucleic acid (*e.g.*, a transgene) into the cell. Thus, a “transduced” cell is a cell into which, or a progeny thereof in which an exogenous nucleic acid has been introduced. The cell(s) can be propagated and the introduced protein expressed, or nucleic acid transcribed. For gene therapy uses and methods, a transduced cell can be in a subject.

[0067] An “expression control element” refers to nucleic acid sequence(s) that influence expression of an operably linked nucleic acid. Control elements, including expression control elements as set forth herein such as promoters and enhancers, Vector sequences including AAV vectors can include one or more “expression control elements.” Typically, such elements are included to facilitate proper heterologous polynucleotide transcription and if appropriate translation (*e.g.*, a promoter, enhancer, splicing signal for introns, maintenance of the correct reading frame of the gene to permit in-frame translation of mRNA and, stop codons etc.). Such elements typically act in cis, referred to as a “cis acting” element, but may also act in trans.

[0068] Expression control can be effected at the level of transcription, translation, splicing, message stability, etc. Typically, an expression control element that modulates transcription is juxtaposed near the 5’ end (*i.e.*, “upstream”) of a transcribed nucleic acid. Expression control elements can also be located at the 3’ end (*i.e.*, “downstream”) of the transcribed sequence or within the transcript (*e.g.*, in an intron). Expression control elements can be located adjacent to or at a distance away from the transcribed sequence (*e.g.*, 1-10, 10-25, 25-50, 50-100, 100 to 500,

or more nucleotides from the polynucleotide), even at considerable distances. Nevertheless, owing to the length limitations of certain vectors, such as AAV vectors, expression control elements will typically be within 1 to 1000 nucleotides from the transcribed nucleic acid.

[0069] Functionally, expression of operably linked nucleic acid is at least in part controllable by the element (*e.g.*, promoter) such that the element modulates transcription of the nucleic acid and, as appropriate, translation of the transcript. A specific example of an expression control element is a promoter, which is usually located 5' of the transcribed sequence *e.g.*, Factor VIII (FVIII) encoding nucleic acid variant. A promoter typically increases an amount expressed from operably linked nucleic acid as compared to an amount expressed when no promoter exists.

10 [0070] An "enhancer" as used herein can refer to a sequence that is located adjacent to the heterologous polynucleotide. Enhancer elements are typically located upstream of a promoter element but also function and can be located downstream of or within a sequence (*e.g.*, Factor VIII (FVIII) encoding nucleic acid variant). Hence, an enhancer element can be located 100 base pairs, 200 base pairs, or 300 or more base pairs upstream or downstream of Factor VIII (FVIII) encoding nucleic acid variant. Enhancer elements typically increase expressed of an operably

15 linked nucleic acid above expression afforded by a promoter element.

[0071] An expression construct may comprise regulatory elements which serve to drive expression in a particular cell or tissue type. Expression control elements (*e.g.*, promoters) include those active in a particular tissue or cell type, referred to herein as a "tissue-specific expression control elements/promoters." Tissue-specific expression control elements are typically active in specific cell or tissue (*e.g.*, liver). Expression control elements are typically active in particular cells, tissues or organs because they are recognized by transcriptional activator proteins, or other regulators of transcription, that are unique to a specific cell, tissue or organ type. Such regulatory elements are known to those of skill in the art (see, *e.g.*, Sambrook et

20 al. (1989) and Ausubel et al. (1992)).

[0072] The incorporation of tissue specific regulatory elements in the expression constructs of the invention provides for at least partial tissue tropism for the expression of the variant FVIII or functional fragments thereof. Examples of promoters that are active in liver are the TTR promoter, human alpha 1-antitrypsin (hAAT) promoter; albumin, Miyatake, et al. *J. Virol.*, 71:5124-32 (1997); hepatitis B virus core promoter, Sandig, et al., *Gene Ther.* 3:1002-9 (1996);

30 alpha-fetoprotein (AFP), Arbuthnot, et al., *Hum. Gene Ther.*, 7:1503-14 (1996)], among others. An example of an enhancer active in liver is apolipoprotein E (apoE) HCR-1 and HCR-2 (Allan et al., *J. Biol. Chem.*, 272:29113-19 (1997)).

[0073] Expression control elements also include ubiquitous or promiscuous promoters/enhancers which are capable of driving expression of a polynucleotide in many

35

different cell types. Such elements include, but are not limited to the cytomegalovirus (CMV) immediate early promoter/enhancer sequences, the Rous sarcoma virus (RSV) promoter/enhancer sequences and the other viral promoters/enhancers active in a variety of mammalian cell types, or synthetic elements that are not present in nature (*see, e.g.*, Boshart et al, *Cell*, 41:521-530 (1985)),

5 the SV40 promoter, the dihydrofolate reductase promoter, the cytoplasmic β -actin promoter and the phosphoglycerol kinase (PGK) promoter.

[0074] Expression control elements also can confer expression in a manner that is regulatable, that is, a signal or stimuli increases or decreases expression of the operably linked heterologous polynucleotide. A regulatable element that increases expression of the operably

10 linked polynucleotide in response to a signal or stimuli is also referred to as an "inducible element" (*i.e.*, is induced by a signal). Particular examples include, but are not limited to, a hormone (*e.g.*, steroid) inducible promoter. Typically, the amount of increase or decrease conferred by such elements is proportional to the amount of signal or stimuli present; the greater the amount of signal or stimuli, the greater the increase or decrease in expression. Particular non-

15 limiting examples include zinc-inducible sheep metallothioneine (MT) promoter; the steroid hormone-inducible mouse mammary tumor virus (MMTV) promoter; the T7 polymerase promoter system (WO 98/10088); the tetracycline-repressible system (Gossen, et al., *Proc. Natl. Acad. Sci. USA*, 89:5547-5551 (1992)); the tetracycline-inducible system (Gossen, et al., *Science*, 268:1766-1769 (1995); see also Harvey, et al., *Curr. Opin. Chem. Biol.* 2:512-518 (1998)); the

20 RU486-inducible system (Wang, et al., *Nat. Biotech.* 15:239-243 (1997) and Wang, et al., *Gene Ther.* 4:432-441 (1997)]; and the rapamycin-inducible system (Magari, et al., *J. Clin. Invest.* 100:2865-2872 (1997); Rivera, et al., *Nat. Medicine*. 2:1028-1032 (1996)). Other regulatable control elements which may be useful in this context are those which are regulated by a specific physiological state, *e.g.*, temperature, acute phase, development.

25 [0075] Expression control elements also include the native element(s) for the heterologous polynucleotide. A native control element (*e.g.*, promoter) may be used when it is desired that expression of the heterologous polynucleotide should mimic the native expression. The native element may be used when expression of the heterologous polynucleotide is to be regulated temporally or developmentally, or in a tissue-specific manner, or in response to specific

30 transcriptional stimuli. Other native expression control elements, such as introns, polyadenylation sites or Kozak consensus sequences may also be used.

[0076] The term "operably linked" means that the regulatory sequences necessary for expression of a coding sequence are placed in the appropriate positions relative to the coding sequence so as to effect expression of the coding sequence. This same definition is sometimes

35 applied to the arrangement of coding sequences and transcription control elements (*e.g.*

promoters, enhancers, and termination elements) in an expression vector. This definition is also sometimes applied to the arrangement of nucleic acid sequences of a first and a second nucleic acid molecule wherein a hybrid nucleic acid molecule is generated.

[0077] In the example of an expression control element in operable linkage with a nucleic acid, the relationship is such that the control element modulates expression of the nucleic acid. More specifically, for example, two DNA sequences operably linked means that the two DNAs are arranged (cis or trans) in such a relationship that at least one of the DNA sequences is able to exert a physiological effect upon the other sequence.

[0078] Accordingly, additional elements for vectors include, without limitation, an expression control (*e.g.*, promoter/enhancer) element, a transcription termination signal or stop codon, 5' or 3' untranslated regions (*e.g.*, polyadenylation (polyA) sequences) which flank a sequence, such as one or more copies of an AAV ITR sequence, or an intron.

[0079] Further elements include, for example, filler or stuffer polynucleotide sequences, for example to improve packaging and reduce the presence of contaminating nucleic acid. AAV vectors typically accept inserts of DNA having a size range which is generally about 4 kb to about 5.2 kb, or slightly more. Thus, for shorter sequences, inclusion of a stuffer or filler in order to adjust the length to near or at the normal size of the virus genomic sequence acceptable for AAV vector packaging into virus particle. In various embodiments, a filler/stuffer nucleic acid sequence is an untranslated (non-protein encoding) segment of nucleic acid. For a nucleic acid sequence less than 4.7 Kb, the filler or stuffer polynucleotide sequence has a length that when combined (*e.g.*, inserted into a vector) with the sequence has a total length between about 3.0-5.5Kb, or between about 4.0-5.0Kb, or between about 4.3-4.8Kb.

[0080] An intron can also function as a filler or stuffer polynucleotide sequence in order to achieve a length for AAV vector packaging into a virus particle. Introns and intron fragments that function as a filler or stuffer polynucleotide sequence also can enhance expression.

[0081] The phrase "hemostasis related disorder" refers to bleeding disorders such as hemophilia A, hemophilia A patients with inhibitory antibodies, deficiencies in coagulation Factors, VII, VIII, IX and X, XI, V, XII, II, von Willebrand factor, combined FV/FVIII deficiency, vitamin K epoxide reductase C1 deficiency, gamma-carboxylase deficiency; bleeding associated with trauma, injury, thrombosis, thrombocytopenia, stroke, coagulopathy, disseminated intravascular coagulation (DIC); over-anticoagulation associated with heparin, low molecular weight heparin, pentasaccharide, warfarin, small molecule antithrombotics (*i.e.* FXa inhibitors); and platelet disorders such as, Bernard Soulier syndrome, Glanzman thrombasthenia, and storage pool deficiency.

[0082] The term "isolated," when used as a modifier of a composition, means that the compositions are made by the hand of man or are separated, completely or at least in part, from their naturally occurring *in vivo* environment. Generally, isolated compositions are substantially free of one or more materials with which they normally associate with in nature, for example, one or more protein, nucleic acid, lipid, carbohydrate, cell membrane.

[0083] With reference to nucleic acids of the invention, the term "isolated " refers to a nucleic acid molecule that is separated from one or more sequences with which it is immediately contiguous (in the 5' and 3' directions) in the naturally occurring genome (genomic DNA) of the organism from which it originates. For example, the "isolated nucleic acid" may comprise a DNA or cDNA molecule inserted into a vector, such as a plasmid or virus vector, or integrated into the DNA of a prokaryote or eukaryote.

[0084] With respect to RNA molecules of the invention, the term "isolated " primarily refers to an RNA molecule encoded by an isolated DNA molecule as defined above. Alternatively, the term may refer to an RNA molecule that has been sufficiently separated from RNA molecules with which it would be associated in its natural state (i.e., in cells or tissues), such that it exists in a "substantially pure" form (the term "substantially pure" is defined below).

[0085] With respect to protein, the term "isolated protein" or "isolated and purified protein" is sometimes used herein. This term refers primarily to a protein produced by expression of an isolated nucleic acid molecule. Alternatively, this term may refer to a protein which has been sufficiently separated from other proteins with which it would naturally be associated, so as to exist in "substantially pure" form.

[0086] The term "isolated" does not exclude combinations produced by the hand of man, for example, a recombinant vector (*e.g.*, rAAV) sequence, or virus particle that packages or encapsidates a vector genome and a pharmaceutical formulation. The term "isolated" also does not exclude alternative physical forms of the composition, such as hybrids/chimeras, multimers/oligomers, modifications (*e.g.*, phosphorylation, glycosylation, lipidation) or derivatized forms, or forms expressed in host cells produced by the hand of man.

[0087] The term "substantially pure" refers to a preparation comprising at least 50-60% by weight the compound of interest (*e.g.*, nucleic acid, oligonucleotide, protein, etc.). The preparation can comprise at least 75% by weight, or about 90-99% by weight, of the compound of interest. Purity is measured by methods appropriate for the compound of interest (*e.g.* chromatographic methods, agarose or polyacrylamide gel electrophoresis, HPLC analysis, and the like).

[0088] The phrase "consisting essentially of" when referring to a particular nucleotide sequence or amino acid sequence means a sequence having the properties of a given SEQ ID

NO: For example, when used in reference to an amino acid sequence, the phrase includes the sequence per se and molecular modifications that would not affect the basic and novel characteristics of the sequence.

[0089] The term "oligonucleotide," as used herein refers to primers and probes, and is defined as a nucleic acid molecule comprised of two or more ribo- or deoxyribonucleotides, such as more than three. The exact size of the oligonucleotide will depend on various factors and on the particular application for which the oligonucleotide is used.

[0090] The term "probe" as used herein refers to an oligonucleotide, polynucleotide or nucleic acid, either RNA or DNA, whether occurring naturally as in a purified restriction enzyme digest or produced synthetically, which is capable of annealing with or specifically hybridizing to a nucleic acid with sequences complementary to the probe. A probe may be either single-stranded or double-stranded. The exact length of the probe will depend upon many factors, including temperature, source of probe and method of use. For example, for diagnostic applications, depending on the complexity of the target sequence, the oligonucleotide probe typically contains 15-25 or more nucleotides, although it may contain fewer nucleotides.

[0091] The probes herein are selected to be "substantially" complementary to different strands of a particular target nucleic acid sequence. This means that the probes must be sufficiently complementary so as to be able to "specifically hybridize" or anneal with their respective target strands under a set of pre-determined conditions. Therefore, the probe sequence need not reflect the exact complementary sequence of the target. For example, a non-complementary nucleotide fragment may be attached to the 5' or 3' end of the probe, with the remainder of the probe sequence being complementary to the target strand. Alternatively, non-complementary bases or longer sequences can be interspersed into the probe, provided that the probe sequence has sufficient complementarity with the sequence of the target nucleic acid to anneal therewith specifically.

[0092] The term "specifically hybridize" refers to the association between two single-stranded nucleic acid molecules of sufficiently complementary sequence to permit such hybridization under pre-determined conditions generally used in the art (sometimes termed "substantially complementary"). In particular, the term refers to hybridization of an oligonucleotide with a substantially complementary sequence contained within a single-stranded DNA or RNA molecule of the invention, to the substantial exclusion of hybridization of the oligonucleotide with single-stranded nucleic acids of non-complementary sequence.

[0093] The term "primer" as used herein refers to an oligonucleotide, either RNA or DNA, either single-stranded or double-stranded, either derived from a biological system, generated by restriction enzyme digestion, or produced synthetically which, when placed in the proper

environment, is able to act functionally as an initiator of template-dependent nucleic acid synthesis. When presented with an appropriate nucleic acid template, suitable nucleoside triphosphate precursors of nucleic acids, a polymerase enzyme, suitable cofactors and conditions such as a suitable temperature and pH, the primer may be extended at its 3' terminus by the
 5 addition of nucleotides by the action of a polymerase or similar activity to yield a primer extension product.

[0094] The primer may vary in length depending on the particular conditions and requirements of the application. For example, in diagnostic applications, the oligonucleotide primer is typically 15-25 or more nucleotides in length. The primer must be of sufficient
 10 complementarity to the desired template to prime the synthesis of the desired extension product, that is, to be able to anneal with the desired template strand in a manner sufficient to provide the 3' hydroxyl moiety of the primer in appropriate juxtaposition for use in the initiation of synthesis by a polymerase or similar enzyme. It is not required that the primer sequence represent an exact complement of the desired template. For example, a non-complementary nucleotide sequence
 15 may be attached to the 5' end of an otherwise complementary primer. Alternatively, non-complementary bases may be interspersed within the oligonucleotide primer sequence, provided that the primer sequence has sufficient complementarity with the sequence of the desired template strand to functionally provide a template-primer complex for the synthesis of the extension product.

20 [0095] The term "identity," "homology" and grammatical variations thereof, mean that two or more referenced entities are the same, when they are "aligned" sequences. Thus, by way of example, when two polypeptide sequences are identical, they have the same amino acid sequence, at least within the referenced region or portion. Where two polynucleotide sequences are identical, they have the same polynucleotide sequence, at least within the referenced region or
 25 portion. The identity can be over a defined area (region or domain) of the sequence. An "area" or "region" of identity refers to a portion of two or more referenced entities that are the same. Thus, where two protein or nucleic acid sequences are identical over one or more sequence areas or regions they share identity within that region. An "aligned" sequence refers to multiple polynucleotide or protein (amino acid) sequences, often containing corrections for missing or
 30 additional bases or amino acids (gaps) as compared to a reference sequence.

[0096] The identity can extend over the entire length or a portion of the sequence. In particular aspects, the length of the sequence sharing the percent identity is 2, 3, 4, 5 or more contiguous nucleic acids or amino acids, *e.g.*, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, etc. contiguous nucleic acids or amino acids. In additional particular aspects, the length of the
 35 sequence sharing identity is 21 or more contiguous nucleic acids or amino acids, *e.g.*, 21, 22, 23,

24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, etc. contiguous nucleic acids or amino acids. In further particular aspects, the length of the sequence sharing identity is 41 or more contiguous nucleic acids or amino acids, *e.g.* 42, 43, 44, 45, 45, 47, 48, 49, 50, etc., contiguous nucleic acids or amino acids. In yet further particular aspects, the length of the

5 sequence sharing identity is 50 or more contiguous nucleic acids or amino acids, *e.g.*, 50-55, 55-60, 60-65, 65-70, 70-75, 75-80, 80-85, 85-90, 90-95, 95-100, 100-110, etc. contiguous nucleic acids or amino acids.

[0097] As set forth herein, Factor VIII (FVIII) encoding nucleic acid variants will be distinct from (e.g, non wild-type) but will exhibit sequence identity with wild-type FVIII encoding

10 nucleic acid. In codon optimized FVIII encoding nucleic acid variants, denoted CO1, CO2 and CO/CO3, at the nucleotide sequence level, a codon optimized FVIII encoding nucleic acid variant will typically be at least about 70% identical, more typically about 75% identical, even more typically about 80%-85% identical to wild-type FVIII encoding nucleic acid. Thus, for example, a codon optimized FVIII encoding nucleic acid variant may have 75%-85% identity to wild-type

15 FVIII encoding gene, or to each other, i.e., CO1 vs. CO2, CO1 vs. CO3, CO2 vs. CO3, etc. as set forth herein.

[0098] At the amino acid sequence level, a variant such as a variant FVIII protein will be at least about 70% identical, more typically about 80% identical, even more typically about 90% or more identity. In other embodiments, a variant such as a variant FVIII protein has at least 90%,

20 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more identity to a reference sequence, *e.g.* wild-type FVIII protein with or without B-domain.

[0099] To determine identity, if the variant FVIII (FVIII encoding nucleic acid variant or FVIII protein) retains the B-domain, it is appropriate to compare identity to wild-type FVIII. If the variant FVIII (FVIII encoding nucleic acid variant or FVIII protein) has a B-domain deletion,

25 it is appropriate to compare identity to wild-type FVIII that also has a B-domain deletion.

[0100] The terms “homologous” or “homology” mean that two or more referenced entities share at least partial identity over a given region or portion. “Areas, regions or domains” of homology or identity mean that a portion of two or more referenced entities share homology or are the same. Thus, where two sequences are identical over one or more sequence regions they

30 share identity in these regions. “Substantial homology” means that a molecule is structurally or functionally conserved such that it has or is predicted to have at least partial structure or function of one or more of the structures or functions (*e.g.*, a biological function or activity) of the reference molecule, or relevant/corresponding region or portion of the reference molecule to which it shares homology.

[0101] The extent of identity (homology) or "percent identity" between two sequences can be ascertained using a computer program and/or mathematical algorithm. For purposes of this invention comparisons of nucleic acid sequences are performed using the GCG Wisconsin Package version 9.1, available from the Genetics Computer Group in Madison, Wisconsin. For convenience, the default parameters (gap creation penalty = 12, gap extension penalty = 4) specified by that program are intended for use herein to compare sequence identity. Alternately, the Blastn 2.0 program provided by the National Center for Biotechnology Information(found on the world wide web at ncbi.nlm.nih.gov/blast/; Altschul et al., 1990, J Mol Biol 215:403-410) using a gapped alignment with default parameters, may be used to determine the level of identity and similarity between nucleic acid sequences and amino acid sequences. For polypeptide sequence comparisons, a BLASTP algorithm is typically used in combination with a scoring matrix, such as PAM100, PAM 250, BLOSUM 62 or BLOSUM 50. FASTA (*e.g.*, FASTA2 and FASTA3) and SSEARCH sequence comparison programs are also used to quantitate extent of identity (Pearson et al., *Proc. Natl. Acad. Sci. USA* 85:2444 (1988); Pearson, *Methods Mol Biol.* 132:185 (2000); and Smith et al., *J. Mol. Biol.* 147:195 (1981)). Programs for quantitating protein structural similarity using Delaunay-based topological mapping have also been developed (Bostick et al., *Biochem Biophys Res Commun.* 304:320 (2003)).

[0102] Nucleic acid molecules, expression vectors (*e.g.*, vector genomes), plasmids, including Factor VIII (FVIII) encoding nucleic acid variants of the invention may be prepared by using recombinant DNA technology methods. The availability of nucleotide sequence information enables preparation of isolated nucleic acid molecules of the invention by a variety of means. For example, Factor VIII (FVIII) encoding nucleic acid variants can be made using various standard cloning, recombinant DNA technology, via cell expression or *in vitro* translation and chemical synthesis techniques. Purity of polynucleotides can be determined through sequencing, gel electrophoresis and the like. For example, nucleic acids can be isolated using hybridization or computer-based database screening techniques. Such techniques include, but are not limited to: (1) hybridization of genomic DNA or cDNA libraries with probes to detect homologous nucleotide sequences; (2) antibody screening to detect polypeptides having shared structural features, for example, using an expression library; (3) polymerase chain reaction (PCR) on genomic DNA or cDNA using primers capable of annealing to a nucleic acid sequence of interest; (4) computer searches of sequence databases for related sequences; and (5) differential screening of a subtracted nucleic acid library.

[0103] Nucleic acids of the invention may be maintained as DNA in any convenient cloning vector. In a one embodiment, clones are maintained in a plasmid cloning/expression vector, such as pBluescript (Stratagene, La Jolla, CA), which is propagated in a suitable *E. coli* host cell.

Alternatively, nucleic acids may be maintained in vector suitable for expression in mammalian cells. In cases where post-translational modification affects coagulation function, nucleic acid molecule can be expressed in mammalian cells.

[0104] Factor VIII (FVIII) encoding nucleic acid variants of the invention include cDNA,
5 genomic DNA, RNA, and fragments thereof which may be single- or double-stranded. Thus, this invention provides oligonucleotides (sense or antisense strands of DNA or RNA) having sequences capable of hybridizing with at least one sequence of a nucleic acid of the invention. Such oligonucleotides are useful as probes for detecting FVIII expression.

[0105] A B-domain deleted, codon optimized FVIII encoding nucleic acid variant of the
10 invention, optionally encoding FVIII polypeptide variant with or without a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment of either as described herein, may be prepared in a variety of ways, according to known methods. The protein may be purified from appropriate sources, e.g., transformed bacterial or animal cultured cells or tissues which express engineered FVIII by immunoaffinity purification.

[0106] The availability of invention variant nucleic acid molecules encoding FVIII,
15 optionally encoding FVIII polypeptide variant with or without a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment of either as described herein, enables production of FVIII using in vitro expression methods known in the art. For example, a cDNA or gene may be cloned into an appropriate in vitro transcription vector, such as pSP64 or pSP65 for
20 in vitro transcription, followed by cell-free translation in a suitable cell-free translation system, such as wheat germ or rabbit reticulocyte lysates. In vitro transcription and translation systems are commercially available, e.g., from Promega Biotech, Madison, Wisconsin or BRL, Rockville, Maryland.

[0107] Alternatively, according to an embodiment, larger quantities of FVIII may be
25 produced by expression in a suitable prokaryotic or eukaryotic expression system. For example, part or all of a Factor VIII (FVIII) encoding nucleic acid variant for example, may be inserted into a plasmid vector adapted for expression in a bacterial cell, such as E. coli or a mammalian cell line such as baby hamster kidney (BHK), CHO or Hela cells. Alternatively, in an embodiment, tagged fusion proteins comprising FVIII can be generated. Such FVIII-tagged
30 fusion proteins are encoded by part or all of a DNA molecule, ligated in the correct codon reading frame to a nucleotide sequence encoding a portion or all of a desired polypeptide tag which is inserted into a plasmid vector adapted for expression in a bacterial cell, such as E. coli or a eukaryotic cell, such as, but not limited to, yeast and mammalian cells.

[0108] Vectors such as those described herein comprise the regulatory elements necessary
35 for expression of the DNA in the host cell positioned in such a manner as to permit expression of

the encoded protein in the host cell. Such regulatory elements required for expression include, but are not limited to, promoter sequences, enhancer sequences and transcription initiation sequences as set forth herein and known to the skilled artisan.

[0109] Factor VIII (FVIII) encoding nucleic acid variant optionally encoding variant FVIII proteins as set forth herein, produced by gene expression in a recombinant prokaryotic or eukaryotic system, may be purified according to methods known in the art. In an embodiment, a commercially available expression/secretion system can be used, whereby the recombinant protein is expressed and thereafter secreted from the host cell, to be easily purified from the surrounding medium. If expression/secretion vectors are not used, an alternative approach involves purifying the recombinant protein by affinity separation, such as by immunological interaction with antibodies that bind specifically to the recombinant protein or nickel columns for isolation of recombinant proteins tagged with 6-8 histidine residues at their N-terminus or C-terminus. Alternative tags may comprise the FLAG epitope, GST or the hemagglutinin epitope. Such methods are commonly used by skilled practitioners.

[0110] FVIII proteins, prepared by the aforementioned methods, may be analyzed according to standard procedures. For example, such proteins may be assessed for altered coagulation properties according to known methods.

[0111] As disclosed herein, a convenient way of producing a polypeptide according to the invention is to express nucleic acid encoding it, by use of the nucleic acid in an expression system. A variety of expression systems of utility for the methods of the invention are well known to those of skill in the art.

[0112] Accordingly, the invention also provides methods of making a polypeptide (as disclosed), the method including expression from nucleic acid encoding the polypeptide (generally nucleic acid). This may conveniently be achieved by culturing a host cell, containing such a vector, under appropriate conditions which cause or allow production of the polypeptide. Polypeptides may also be produced in *in vitro* systems.

[0113] Methods and uses of the invention of the invention include delivering (transducing) nucleic acid (transgene) into host cells, including dividing and/or non-dividing cells. The nucleic acids, recombinant vector (*e.g.*, rAAV), methods, uses and pharmaceutical formulations of the invention are additionally useful in a method of delivering, administering or providing a protein to a subject in need thereof, as a method of treatment. In this manner, the nucleic acid is transcribed and the protein may be produced *in vivo* in a subject. The subject may benefit from or be in need of the protein because the subject has a deficiency of the protein, or because production of the protein in the subject may impart some therapeutic effect, as a method of treatment or otherwise.

[0114] Vectors including lenti- or parvo-virus vector (*e.g.*, AAV) sequences, recombinant virus particles, methods and uses may be used to deliver a Factor VIII (FVIII) encoding nucleic acid variant with a biological effect to treat or ameliorate one or more symptoms associated with a FVIII deficiency or abnormality. Recombinant lenti- or parvo-virus vector (*e.g.*, AAV)

5 sequences, plasmids, recombinant virus particles, methods and uses may be used to provide therapy for various disease states involving or due to a FVIII deficiency or abnormality.

[0115] Invention nucleic acids, vectors, recombinant vectors (*e.g.*, rAAV), and recombinant virus particles, methods and uses permit the treatment of genetic diseases, *e.g.*, a FVIII deficiency. For deficiency state diseases, gene transfer can be used to bring a normal gene into
10 affected tissues for replacement therapy, as well as to create animal models for the disease using antisense mutations. For unbalanced disease states, gene transfer could be used to create a disease state in a model system, which could then be used in efforts to counteract the disease state. The use of site-specific integration of nucleic acid sequences to correct defects is also possible.

[0116] In particular embodiments, Factor VIII (FVIII) encoding nucleic acid variants (*e.g.*,
15 codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (*e.g.*, codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either, may be used, for example, as therapeutic and/or prophylactic agents (protein or nucleic acid) which modulate the blood coagulation cascade or as a transgene in gene. For example, Factor VIII (FVIII)
20 encoding nucleic acid variants may have similar coagulation activity as wild-type FVIII, or altered coagulation activity compared to wild-type FVIII, such as in the case of a variant FVIII protein (*e.g.*, a codon optimized nucleic acid variant encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment. Cell-based strategies allow continuous expression of Factor VIII (FVIII) encoding nucleic acid variant
25 or FVIII protein variants in hemophilia A patients. As disclosed herein, certain modifications of FVIII molecules (nucleic acid and protein) result in increased expression at the nucleic acid level, increased packaging efficiency by virus (*e.g.*, AAV) vector, and increased coagulation activity and greater stability at the protein level thereby effectively improving hemostasis.

[0117] Factor VIII (FVIII) encoding nucleic acid variants (*e.g.*, codon optimized variants
30 encoding FVIII), variant FVIII protein encoding nucleic acid variants (*e.g.*, codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either may be used for a variety of purposes in accordance with the invention. In one embodiment, a nucleic acid delivery vehicle (*i.e.*, an expression vector) for modulating blood coagulation is provided wherein the expression vector
35 comprises a Factor VIII (FVIII) encoding nucleic acid variants (*e.g.*, codon optimized variants

encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either as described herein. Administration of FVIII-encoding expression vectors to a patient results in the expression of FVIII protein which serves to alter the coagulation cascade. In accordance with the invention, a Factor VIII (FVIII) encoding nucleic acid variant may encode a FVIII polypeptide as described herein (e.g., a variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment, whose expression increases hemostasis. In one embodiment, a Factor VIII (FVIII) encoding nucleic acid variant encodes a FVIII polypeptide variant, such as a FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment thereof.

[0118] In additional embodiments of the invention, compositions and methods are provided for administration of a viral vector comprising a Factor VIII (FVIII) encoding nucleic acid variants (e.g., codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either. In one embodiment, the expression vector comprising Factor VIII (FVIII) encoding nucleic acid variants (e.g., codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either, is a viral vector.

[0119] Expression vectors comprising Factor VIII (FVIII) encoding nucleic acid variant (e.g., codon optimized variant encoding FVIII), a variant FVIII protein encoding nucleic acid variant (e.g., codon optimized variant encoding variant FVIII with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either may be administered alone, or in combination with other molecules useful for modulating hemostasis. According to the invention, the expression vectors or combination of therapeutic agents may be administered to the patient alone or in a pharmaceutically acceptable or biologically compatible compositions.

[0120] Viral vectors such as lenti- and parvo-virus vectors, including AAV serotypes and variants thereof provide a means for delivery of nucleic acid into cells *ex vivo*, *in vitro* and *in vivo*, which encode proteins such that the cells express the encoded protein. AAV are viruses useful as gene therapy vectors as they can penetrate cells and introduce nucleic acid/genetic material so that the nucleic acid/genetic material may be stably maintained in cells. In addition, these viruses can introduce nucleic acid/genetic material into specific sites, for example. Because AAV are not associated with pathogenic disease in humans, AAV vectors are able to deliver

heterologous polynucleotide sequences (*e.g.*, therapeutic proteins and agents) to human patients without causing substantial AAV pathogenesis or disease.

[0121] Viral vectors which may be used in the invention include, but are not limited to, adeno-associated virus (AAV) vectors of multiple serotypes (*e.g.*, AAV-1 to AAV-12, and others) and hybrid/chimeric AAV vectors, lentivirus vectors and pseudo-typed lentivirus vectors [e.g., Ebola virus, vesicular stomatitis virus (VSV), and feline immunodeficiency virus (FIV)], herpes simplex virus vectors, adenoviral vectors (with or without tissue specific promoters/enhancers), vaccinia virus vectors, retroviral vectors, lentiviral vectors, non-viral vectors and others.

10 [0122] AAV and lentiviral particles may be used to advantage as vehicles for effective gene delivery. Such virions possess a number of desirable features for such applications, including tropism for dividing and non-dividing cells. Early clinical experience with these vectors also demonstrated no sustained toxicity and immune responses were minimal or undetectable. AAV are known to infect a wide variety of cell types *in vivo* and *in vitro* by receptor-mediated
15 endocytosis or by transcytosis. These vector systems have been tested in humans targeting retinal epithelium, liver, skeletal muscle, airways, brain, joints and hematopoietic stem cells. Non-viral vectors, for example, based on plasmid DNA or minicircles, are also suitable gene transfer vectors for a large gene as that encoding FVIII.

[0123] It may be desirable to introduce a vector that can provide, for example, multiple
20 copies of a desired gene and hence greater amounts of the product of that gene. Improved AAV and lentiviral vectors and methods for producing these vectors have been described in detail in a number of references, patents, and patent applications, including: Wright J.F. (*Hum Gene Ther* 20:698-706, 2009) a technology used for the production of clinical grade vector at Children's Hospital of Philadelphia. Lentiviral vector can also be produced at CHOP and the other vectors
25 are available through the Lentivirus vector production core laboratory by NHLBI Gene Therapy Resource Program (GTRP) - Lentivirus Vector Production Core Laboratory.

[0124] Accordingly, in various embodiments of the invention a vector includes a lenti- or parvo-viral vector, such as an adeno-viral vector. In particular embodiments, a recombinant vector is a parvovirus vector. Parvoviruses are small viruses with a single-stranded DNA
30 genome. "Adeno-associated viruses" (AAV) are in the parvovirus family.

[0125] Accordingly, the invention provides viral vectors that include Factor VIII (FVIII) encoding nucleic acid variants (*e.g.*, codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (*e.g.*, codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a
35 functional fragment of either. For example, a recombinant AAV vector can include a nucleic acid

encoding a desired protein (*e.g.*, Factor VIII), such as a codon optimized Factor VIII (FVIII) encoding nucleic acid variant (*e.g.*, codon optimized variant encoding FVIII), where the encoded FVIII protein optionally has B-domain deletion, variant FVIII protein encoding nucleic acid variants (*e.g.*, codon optimized nucleic acid variants encoding variant FVIII protein with a

5 PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either. Vector delivery or administration to a subject (*e.g.*, mammal) therefore provides FVIII (via *e.g.*, codon optimized nucleic acid variant encoding FVIII), variant FVIII protein encoding nucleic acid variants (*e.g.*, codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either,

10 to a subject such as a mammal (*e.g.*, human).

[0126] Direct delivery of vectors or ex-vivo transduction of human cells followed by infusion into the body will result in FVIII expression thereby exerting a beneficial therapeutic effect on hemostasis. In the context of invention Factor VIII described herein, such administration enhances pro-coagulation activity.

15 **[0127]** AAV vectors and lentiviral vectors do not typically include viral genes associated with pathogenesis. Such vectors typically have one or more of the wild type AAV genes deleted in whole or in part, for example, rep and/or cap genes, but retain at least one functional flanking ITR sequence, as necessary for the rescue, replication, and packaging of the recombinant vector into an AAV vector particle. For example, only the essential parts of vector *e.g.*, the ITR and

20 LTR elements, respectively are included. An AAV vector genome would therefore include sequences required in cis for replication and packaging (*e.g.*, functional ITR sequences)

[0128] Recombinant AAV vector, as well as methods and uses thereof, include any viral strain or serotype. As a non-limiting example, a recombinant AAV vector can be based upon any AAV genome, such as AAV-1, -2, -3, -4, -5, -6, -7, -8, -9, -10, -11, -12, -rh74, -rh10 or AAV-2i8,

25 for example. Such vectors can be based on the same strain or serotype (or subgroup or variant), or be different from each other. As a non-limiting example, a recombinant AAV vector based upon one serotype genome can be identical to one or more of the capsid proteins that package the vector. In addition, a recombinant AAV vector genome can be based upon an AAV (*e.g.*, AAV2) serotype genome distinct from one or more of the AAV capsid proteins that package the vector.

30 For example, the AAV vector genome can be based upon AAV2, whereas at least one of the three capsid proteins could be a AAV1, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8 or variant thereof, for example.

[0129] In particular embodiments, adeno-associated virus (AAV) vectors include AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10,

35 Rh74 and AAV-2i8, as well as variants (*e.g.*, capsid variants, such as amino acid insertions,

additions and substitutions) thereof as set forth in WO 2013/158879 (International Application PCT/US2013/037170) and WO 2015/013313 (International Application PCT/US2014/047670).

AAV variants include AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 and AAV-2i8 variants. Accordingly, AAV vectors and

5 AAV variants (*e.g.*, capsid variants) that include (encapsidate or package) Factor VIII (FVIII) encoding nucleic acid variants (*e.g.*, codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (*e.g.*, codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either, are provided.

10 **[0130]** AAV and AAV variants (*e.g.*, capsid variants) serotypes (*e.g.*, VP1, VP2, and/or VP3 sequences) may or may not be distinct from other AAV serotypes, including, for example, AAV1-AAV12, Rh74 or or Rh10 (*e.g.*, distinct from VP1, VP2, and/or VP3 sequences of any of AAV1-AAV12, Rh74 or Rh10 serotypes).

[0131] As used herein, the term “serotype” is a distinction used to refer to an AAV having a capsid that is serologically distinct from other AAV serotypes. Serologic distinctiveness is determined on the basis of the lack of cross-reactivity between antibodies to one AAV as compared to another AAV. Such cross-reactivity differences are usually due to differences in capsid protein sequences/antigenic determinants (*e.g.*, due to VP1, VP2, and/or VP3 sequence differences of AAV serotypes). Despite the possibility that AAV variants including capsid variants may not be serologically distinct from a reference AAV or other AAV serotype, they differ by at least one nucleotide or amino acid residue compared to the reference or other AAV serotype.

[0132] Under the traditional definition, a serotype means that the virus of interest has been tested against serum specific for all existing and characterized serotypes for neutralizing activity and no antibodies have been found that neutralize the virus of interest. As more naturally occurring virus isolates of are discovered and/or capsid mutants generated, there may or may not be serological differences with any of the currently existing serotypes. Thus, in cases where the new virus (*e.g.*, AAV) has no serological difference, this new virus (*e.g.*, AAV) would be a subgroup or variant of the corresponding serotype. In many cases, serology testing for neutralizing activity has yet to be performed on mutant viruses with capsid sequence modifications to determine if they are of another serotype according to the traditional definition of serotype. Accordingly, for the sake of convenience and to avoid repetition, the term “serotype” broadly refers to both serologically distinct viruses (*e.g.*, AAV) as well as viruses (*e.g.*, AAV) that are not serologically distinct that may be within a subgroup or a variant of a given serotype.

[0133] AAV vectors therefore include gene/protein sequences identical to gene/protein sequences characteristic for a particular serotype. As used herein, an “AAV vector related to AAV1” refers to one or more AAV proteins (*e.g.*, VP1, VP2, and/or VP3 sequences) that has substantial sequence identity to one or more polynucleotides or polypeptide sequences that
 5 comprise AAV1. Analogously, an “AAV vector related to AAV8” refers to one or more AAV proteins (*e.g.*, VP1, VP2, and/or VP3 sequences) that has substantial sequence identity to one or more polynucleotides or polypeptide sequences that comprise AAV8. An “AAV vector related to AAV-Rh74” refers to one or more AAV proteins (*e.g.*, VP1, VP2, and/or VP3 sequences) that has substantial sequence identity to one or more polynucleotides or polypeptide sequences that
 10 comprise AAV-Rh74. Such AAV vectors related to another serotype, *e.g.*, AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8, can therefore have one or more distinct sequences from AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 and AAV-2i8, but can exhibit substantial sequence identity to one or more genes and/or proteins, and/or have one or
 15 more functional characteristics of AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8 (*e.g.*, such as cell/tissue tropism). Exemplary non-limiting AAV variants include capsid variants of any of VP1, VP2, and/or VP3.

[0134] In various exemplary embodiments, an AAV vector related to a reference serotype has a polynucleotide, polypeptide or subsequence thereof that includes or consists of a sequence
 20 at least 80% or more (*e.g.*, 85%, 90%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, etc.) identical to one or more AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8 (*e.g.*, such as VP1, VP2, and/or VP3 sequences).

[0135] Compositions, methods and uses of the invention include AAV sequences
 25 (polypeptides and nucleotides), and subsequences thereof that exhibit less than 100% sequence identity to a reference AAV serotype such as AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, or AAV-2i8, but are distinct from and not identical to known AAV genes or proteins, such as AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8, genes or
 30 proteins, etc. In one embodiment, an AAV polypeptide or subsequence thereof includes or consists of a sequence at least 75% or more identical, *e.g.*, 80%, 85%, 85%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.1%, 99.2%, 99.3%, 99.4%, 99.5%, etc., up to 100% identical to any reference AAV sequence or subsequence thereof, such as AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12,

Rh10, Rh74 or AAV-2i8 (*e.g.*, VP1, VP2 and/or VP3). In particular aspects, an AAV variant has 1, 2, 3, 4, 5, 5-10, 10-15, 15-20 or more amino acid substitutions.

[0136] Recombinant AAV vectors, including AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8 and variant, related, hybrid and chimeric sequences, can be constructed using recombinant techniques that are known to the skilled artisan, to include one or more nucleic acid sequences (transgenes) flanked with one or more functional AAV ITR sequences.

[0137] In one embodiment of the invention, FVIII polypeptide variants, such as a human FVIII variant with a PACE-furin cleavage recognition site mutation, deletion or substitution, may be administered to a patient via infusion in a biologically compatible carrier, for example, via intravenous injection. The FVIII polypeptide variants, such as a human FVIII variant with a PACE-furin cleavage recognition site mutation, deletion or substitution of the invention may optionally be encapsulated into liposomes or mixed with other phospholipids or micelles to increase stability of the molecule. FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment, may be administered alone or in combination with other agents known to modulate hemostasis (*e.g.*, Factor V, Factor Va or derivatives thereof).

[0138] An appropriate composition in which to deliver FVIII polypeptide variants, such as a human FVIII variant with a PACE-furin cleavage recognition site mutation, deletion or substitution may be determined by a medical practitioner upon consideration of a variety of physiological variables, including, but not limited to, the patient's condition and hemodynamic state. A variety of compositions well suited for different applications and routes of administration are well known in the art and are described hereinbelow.

[0139] A preparation containing purified FVIII protein, such as FVIII variant with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment, contains a physiologically acceptable matrix and may be formulated as a pharmaceutical preparation. The preparation can be formulated using substantially known prior art methods, it can be mixed with a buffer containing salts, such as NaCl, CaCl₂, and amino acids, such as glycine and/or lysine, and in a pH range from 6 to 8. Until needed, the purified preparation containing FVIII variant can be stored in the form of a finished solution or in lyophilized or deep-frozen form.

[0140] A preparation can be stored in lyophilized form and is dissolved into a visually clear solution using an appropriate reconstitution solution. Alternatively, the preparation according to the invention can also be made available as a liquid preparation or as a liquid that is deep-frozen. The preparation according to the invention may optionally be especially stable, *i.e.*, it can be allowed to stand in dissolved form for a prolonged time prior to administration or delivery.

[0141] The preparation according to the invention can be made available as a pharmaceutical preparation with FVIII activity in the form of a one-component preparation or in combination with other factors in the form of a multi-component preparation. Prior to processing the purified protein into a pharmaceutical preparation, the purified protein is subjected to the conventional
 5 quality controls and fashioned into a therapeutic form of presentation. In particular, during the recombinant manufacture, the purified preparation is tested for the absence of cellular nucleic acids as well as nucleic acids that are derived from the expression vector, such as is described in EP 0 714 987.

[0142] The pharmaceutical protein preparation may be used at dosages of between 30-100
 10 IU/kg (One I.U is 100 ng/ml) at as single daily injection or up to 3 times/day for several days. Patients may be treated immediately upon presentation at the clinic with a bleed. Alternatively, patients may receive a bolus infusion every eight to twelve hours, or if sufficient improvement is observed, a once daily infusion of the variant FVIII described herein.

[0143] The invention also provides compositions, such as compositions including Factor VIII
 15 (FVIII) encoding nucleic acid variants (e.g., codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), and functional fragments of either, having one or more of the following attributes:
 1) exhibits increased expression by cells or in animals; 2) exhibits increased secretion by cells; 3)
 20 exhibits increased activity, as reflected by increased clotting, for example; 4) exhibits increased stability; and 5) exhibits increased packaging by AAV vector.

[0144] Accordingly, invention nucleic acids, vectors, recombinant vectors (e.g., rAAV), and recombinant virus particles and other compositions, agents, drugs, biologics (proteins) can be incorporated into pharmaceutical compositions. Such pharmaceutical compositions are useful
 25 for, among other things, administration and delivery to a subject *in vivo* or *ex vivo*.

[0145] In particular embodiments, pharmaceutical compositions also contain a pharmaceutically acceptable carrier or excipient. Such excipients include any pharmaceutical agent that does not itself induce an immune response harmful to the individual receiving the composition, and which may be administered without undue toxicity.

[0146] As used herein the term “pharmaceutically acceptable” and “physiologically acceptable” mean a biologically acceptable formulation, gaseous, liquid or solid, or mixture thereof, which is suitable for one or more routes of administration, *in vivo* delivery or contact. A
 30 “pharmaceutically acceptable” or “physiologically acceptable” composition is a material that is not biologically or otherwise undesirable, e.g., the material may be administered to a subject
 35 without causing substantial undesirable biological effects. Thus, such a pharmaceutical

composition may be used, for example in administering a nucleic acid, vector, viral particle or protein to a subject.

[0147] Pharmaceutically acceptable excipients include, but are not limited to, liquids such as water, saline, glycerol, sugars and ethanol. Pharmaceutically acceptable salts can also be included therein, for example, mineral acid salts such as hydrochlorides, hydrobromides, phosphates, sulfates, and the like; and the salts of organic acids such as acetates, propionates, malonates, benzoates, and the like. Additionally, auxiliary substances, such as wetting or emulsifying agents, pH buffering substances, and the like, may be present in such vehicles.

[0148] The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding, free base forms. In other cases, a preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

[0149] Pharmaceutical compositions include solvents (aqueous or non-aqueous), solutions (aqueous or non-aqueous), emulsions (*e.g.*, oil-in-water or water-in-oil), suspensions, syrups, elixirs, dispersion and suspension media, coatings, isotonic and absorption promoting or delaying agents, compatible with pharmaceutical administration or *in vivo* contact or delivery. Aqueous and non-aqueous solvents, solutions and suspensions may include suspending agents and thickening agents. Such pharmaceutically acceptable carriers include tablets (coated or uncoated), capsules (hard or soft), microbeads, powder, granules and crystals. Supplementary active compounds (*e.g.*, preservatives, antibacterial, antiviral and antifungal agents) can also be incorporated into the compositions.

[0150] Pharmaceutical compositions can be formulated to be compatible with a particular route of administration or delivery, as set forth herein or known to one of skill in the art. Thus, pharmaceutical compositions include carriers, diluents, or excipients suitable for administration by various routes.

[0151] Compositions suitable for parenteral administration comprise aqueous and non-aqueous solutions, suspensions or emulsions of the active compound, which preparations are typically sterile and can be isotonic with the blood of the intended recipient. Non-limiting illustrative examples include water, buffered saline, Hanks' solution, Ringer's solution, dextrose, fructose, ethanol, animal, vegetable or synthetic oils. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran.

[0152] Additionally, suspensions of the active compounds may be prepared as appropriate oil injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

[0153] Cosolvents and adjuvants may be added to the formulation. Non-limiting examples of cosolvents contain hydroxyl groups or other polar groups, for example, alcohols, such as isopropyl alcohol; glycols, such as propylene glycol, polyethyleneglycol, polypropylene glycol, glycol ether; glycerol; polyoxyethylene alcohols and polyoxyethylene fatty acid esters. Adjuvants include, for example, surfactants such as, soya lecithin and oleic acid; sorbitan esters such as sorbitan trioleate; and polyvinylpyrrolidone.

[0154] After pharmaceutical compositions have been prepared, they may be placed in an appropriate container and labeled for treatment. For administration of FVIII-containing vectors or polypeptides, such labeling would include amount, frequency, and method of administration.

[0155] Pharmaceutical compositions and delivery systems appropriate for the compositions, methods and uses of the invention are known in the art (*see, e.g.,* Remington: The Science and Practice of Pharmacy (2003) 20th ed., Mack Publishing Co., Easton, PA; Remington's Pharmaceutical Sciences (1990) 18th ed., Mack Publishing Co., Easton, PA; The Merck Index (1996) 12th ed., Merck Publishing Group, Whitehouse, NJ; Pharmaceutical Principles of Solid Dosage Forms (1993), Technomic Publishing Co., Inc., Lancaster, Pa.; Ansel and Stoklosa, Pharmaceutical Calculations (2001) 11th ed., Lippincott Williams & Wilkins, Baltimore, MD; and Poznansky et al., Drug Delivery Systems (1980), R. L. Juliano, ed., Oxford, N.Y., pp. 253-315).

[0156] The invention also provides methods for introducing a Factor VIII (FVIII) encoding nucleic acid variants (e.g., codon optimized variant encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variant encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), and functional fragments of either into a cell or an animal. In a particular embodiment, the invention provides methods for modulating hemostasis. In one embodiment, a method includes contact or administration of an individual (patient or subject such as a mammal) with a nucleic acid delivery vehicle (e.g., an AAV vector) comprising a Factor VIII (FVIII) encoding nucleic acid variant (e.g., codon optimized variant encoding FVIII), or variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variant encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either, under conditions wherein the FVIII polypeptide is expressed in the individual. In another

embodiment, a method includes providing cells of an individual (patient or subject such as a mammal) with a nucleic acid delivery vehicle (e.g., an AAV vector) comprising a Factor VIII (FVIII) encoding nucleic acid variant (e.g., codon optimized variant encoding FVIII), or variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variant encoding
 5 variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either, under conditions wherein the FVIII polypeptide is expressed in the individual.

[0157] From the foregoing, it can be seen that Factor VIII (FVIII) encoding nucleic acid variants (e.g., codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic
 10 acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either, may be used in the treatment of disorders associated with deficient, insufficient or aberrant blood coagulation.

[0158] Compositions of Factor VIII (FVIII) encoding nucleic acid variants (e.g., codon
 15 optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or functional fragments, of either, including vectors, recombinant vectors (e.g., rAAV), and recombinant virus particles can be administered, and methods and uses of the invention can be provided, in a sufficient or effective amount to a
 20 subject in need thereof. An “effective amount” or “sufficient amount” refers to an amount that provides, in single or multiple doses, alone or in combination, with one or more other compositions (therapeutic agents such as a drug), treatments, protocols, or therapeutic regimens agents, a detectable response of any duration of time (long or short term), an expected or desired outcome in or a benefit to a subject of any measurable or detectable degree or for any duration of
 25 time (e.g., for minutes, hours, days, months, years, or cured).

[0159] Doses can vary and depend upon the type, onset, progression, severity, frequency, duration, or probability of the disease to which treatment is directed, the clinical endpoint desired, previous or simultaneous treatments, the general health, age, gender, race or immunological competency of the subject and other factors that will be appreciated by the skilled artisan. The
 30 dose amount, number, frequency or duration may be proportionally increased or reduced, as indicated by any adverse side effects, complications or other risk factors of the treatment or therapy and the status of the subject. The skilled artisan will appreciate the factors that may influence the dosage and timing required to provide an amount sufficient for providing a therapeutic or prophylactic benefit.

[0160] The dose to achieve a therapeutic effect, *e.g.*, the dose in vector genomes/per kilogram of body weight (vg/kg), will vary based on several factors including, but not limited to: route of administration, the level of heterologous polynucleotide expression required to achieve a therapeutic effect, the specific disease treated, any host immune response to the viral vector, a host immune response to the heterologous polynucleotide or expression product (protein), and the stability of the protein expressed. One skilled in the art can determine a rAAV/vector genome dose range to treat a patient having a particular disease or disorder based on the aforementioned factors, as well as other factors. Generally, doses will range from at least 1×10^8 , or more, for example, 1×10^9 , 1×10^{10} , 1×10^{11} , 1×10^{12} , 1×10^{13} or 1×10^{14} , or more, vector genomes per kilogram (vg/kg) of the weight of the subject, to achieve a therapeutic effect. AAV dose in the range of 1×10^{10} - 1×10^{11} in mice, and 1×10^{12} - 1×10^{13} in dogs have been effective.

[0161] Using hemophilia B as an example, generally speaking, it is believed that, in order to achieve a therapeutic effect, a blood coagulation factor concentration that is greater than 1% of factor concentration found in a normal individual is needed to change a severe disease phenotype to a moderate one. A severe phenotype is characterized by joint damage and life-threatening bleeds. To convert a moderate disease phenotype into a mild one, it is believed that a blood coagulation factor concentration greater than 5% of normal is needed. With respect to treating such a hemophilic subject, a typical dose is at least 1×10^{10} vector genomes (vg) per kilogram (vg/kg) of the weight of the subject, or between about 1×10^{10} to 1×10^{11} vg/kg of the weight of the subject, or between about 1×10^{11} to 1×10^{12} vg/kg of the weight of the subject, or between about 1×10^{12} to 1×10^{13} vg/kg of the weight of the subject, to achieve a desired therapeutic effect.

[0162] The doses of an “effective amount” or “sufficient amount” for treatment (*e.g.*, to ameliorate or to provide a therapeutic benefit or improvement) typically are effective to provide a response to one, multiple or all adverse symptoms, consequences or complications of the disease, one or more adverse symptoms, disorders, illnesses, pathologies, or complications, for example, caused by or associated with the disease, to a measurable extent, although decreasing, reducing, inhibiting, suppressing, limiting or controlling progression or worsening of the disease is a satisfactory outcome.

[0163] An effective amount or a sufficient amount can but need not be provided in a single administration, may require multiple administrations, and, can but need not be, administered alone or in combination with another composition (*e.g.*, agent), treatment, protocol or therapeutic regimen. For example, the amount may be proportionally increased as indicated by the need of the subject, type, status and severity of the disease treated or side effects (if any) of treatment. In addition, an effective amount or a sufficient amount need not be effective or sufficient if given in

single or multiple doses without a second composition (*e.g.*, another drug or agent), treatment, protocol or therapeutic regimen, since additional doses, amounts or duration above and beyond such doses, or additional compositions (*e.g.*, drugs or agents), treatments, protocols or therapeutic regimens may be included in order to be considered effective or sufficient in a given subject.

- 5 Amounts considered effective also include amounts that result in a reduction of the use of another treatment, therapeutic regimen or protocol, such as administration of recombinant clotting factor protein (*e.g.*, FVIII) for treatment of a clotting disorder (*e.g.*, hemophilia A).

[0164] Accordingly, methods and uses of the invention also include, among other things, methods and uses that result in a reduced need or use of another compound, agent, drug,
10 therapeutic regimen, treatment protocol, process, or remedy. For example, for a blood clotting disease, a method or use of the invention has a therapeutic benefit if in a given subject a less frequent or reduced dose or elimination of administration of a recombinant clotting factor protein to supplement for the deficient or defective (abnormal or mutant) endogenous clotting factor in the subject. Thus, in accordance with the invention, methods and uses of reducing need or use of
15 another treatment or therapy are provided.

[0165] An effective amount or a sufficient amount need not be effective in each and every subject treated, nor a majority of treated subjects in a given group or population. An effective amount or a sufficient amount means effectiveness or sufficiency in a particular subject, not a group or the general population. As is typical for such methods, some subjects will exhibit a
20 greater response, or less or no response to a given treatment method or use.

[0166] The term “ameliorate” means a detectable or measurable improvement in a subject’s disease or symptom thereof, or an underlying cellular response. A detectable or measurable improvement includes a subjective or objective decrease, reduction, inhibition, suppression, limit or control in the occurrence, frequency, severity, progression, or duration of the disease, or
25 complication caused by or associated with the disease, or an improvement in a symptom or an underlying cause or a consequence of the disease, or a reversal of the disease. For HemA, an effective amount would be an amount that reduces frequency or severity of acute bleeding episodes in a subject, for example, or an amount that reduces clotting time as measured by a clotting assay, for example.

30 [0167] Accordingly, pharmaceutical compositions of the invention include compositions wherein the active ingredients are contained in an effective amount to achieve the intended therapeutic purpose. Determining a therapeutically effective dose is well within the capability of a skilled medical practitioner using the techniques and guidance provided in the invention.

[0168] Therapeutic doses will depend on, among other factors, the age and general condition
35 of the subject, the severity of the aberrant blood coagulation phenotype, and the strength of the

control sequences regulating the expression levels of Factor VIII (FVIII) encoding nucleic acid variants (e.g., codon optimized variants encoding FVIII), variant FVIII protein encoding nucleic acid variants (e.g., codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution), or a functional fragment of either.

5 Thus, a therapeutically effective amount in humans will fall in a relatively broad range that may be determined by a medical practitioner based on the response of an individual patient to vector-based FVIII treatment.

[0169] Compositions such as pharmaceutical compositions may be delivered to a subject, so as to allow production of a biologically active protein (e.g., Factor VIII (FVIII) encoded by nucleic acid variants such as codon optimized variants encoding FVIII, or variant FVIII protein encoded by nucleic acid variants such as codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment of either) or by inducing continuous expression of the FVIII transgene in vivo by gene- and or cell-based therapies or by ex-vivo modification of the patient's or donor's cells. In a particular embodiment, pharmaceutical compositions comprising sufficient genetic material to enable a recipient to produce a therapeutically effective amount of a FVIII polypeptide can influence hemostasis in the subject. Alternatively, as disclosed herein, an effective amount of a variant Factor VIII polypeptide, such as FVIII with one or more mutations, deletions or substitutions in the intracellular protease cleavage recognition site (PACE/Furin), may be directly infused into a patient in need thereof.

[0170] The compositions may be administered alone or in combination with at least one other agent, such as a stabilizing compound, which may be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be administered to a patient alone, or in combination with other agents (e.g., co-factors) which influence hemostasis.

[0171] Variant Factor VIII polypeptides, alone or in combination with other agents may be administered or contacted or directly infused into a patient in an appropriate biological carrier as described herein. Expression vectors of the invention comprising FVIII encoding variant nucleic acid sequences, such as codon optimized variants encoding FVIII, or variant FVIII protein encoded by nucleic acid variants such as codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment of either, may be administered to a patient by a variety of means to achieve and optionally maintain for a period of time a prophylactically and/or therapeutically effective level of FVIII polypeptide. One of skill in the art could readily determine specific protocols for using

the FVIII encoding expression vectors of the invention for the therapeutic treatment of a particular patient.

[0172] Protocols for the generation of adenoviral vectors and administration to patients have been described in U.S. Patent Nos. 5,998,205; 6,228,646; 6,093,699; 6,100,242; and International Patent Application Nos. WO 94/17810 and WO 94/23744.

In particular, for example, AAV vectors are employed to deliver Factor VIII (FVIII) encoded by nucleic acid variants such as codon optimized variants encoding FVIII, or variant FVIII protein encoded by nucleic acid variants such as codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment of either, to a patient in need thereof.

[0173] Factor VIII (FVIII) encoded by nucleic acid variants such as codon optimized variants encoding FVIII, or variant FVIII protein encoded by nucleic acid variants such as codon optimized nucleic acid variants encoding variant FVIII protein with a PACE/Furin cleavage site mutation, deletion or substitution, or a functional fragment of either delivered by way of AAVvectors of the invention may be administered to a patient by any means known.

[0174] Methods and uses of the invention include delivery and administration systemically, regionally or locally, or by any route, for example, by injection or infusion. Delivery of the pharmaceutical compositions in vivo may generally be accomplished via injection using a conventional syringe, although other delivery methods such as convection-enhanced delivery are envisioned (See e.g., U.S. Pat. No. 5,720,720). For example, compositions may be delivered subcutaneously, epidermally, intradermally, intrathecally, intraorbitally, intramucosally, intraperitoneally, intravenously, intra-pleurally, intraarterially, orally, intrahepatically, via the portal vein, or intramuscularly. Other modes of administration include oral and pulmonary administration, suppositories, and transdermal applications. A clinician specializing in the treatment of patients with blood coagulation disorders may determine the optimal route for administration of the adenoviral-associated vectors comprising FVIII nucleic acid sequences based on a number of criteria, including, but not limited to: the condition of the patient and the purpose of the treatment (e.g., enhanced or reduced blood coagulation).

[0175] Invention methods and uses can be combined with any compound, agent, drug, treatment or other therapeutic regimen or protocol having a desired therapeutic, beneficial, additive, synergistic or complementary activity or effect. Exemplary combination compositions and treatments include second actives, such as, biologics (proteins), agents and drugs. Such biologics (proteins), agents, drugs, treatments and therapies can be administered or performed prior to, substantially contemporaneously with or following any other method or use of the

invention, for example, a therapeutic method of treating a subject for a blood clotting disease such as HemA.

[0176] The compound, agent, drug, treatment or other therapeutic regimen or protocol can be administered as a combination composition, or administered separately, such as concurrently or
5 in series or sequentially (prior to or following) delivery or administration of a nucleic acid, vector, recombinant vector (*e.g.*, rAAV), or recombinant virus particle. The invention therefore provides combinations in which a method or use of the invention is in a combination with any compound, agent, drug, therapeutic regimen, treatment protocol, process, remedy or composition, set forth herein or known to one of skill in the art. The compound, agent, drug, therapeutic
10 regimen, treatment protocol, process, remedy or composition can be administered or performed prior to, substantially contemporaneously with or following administration of a nucleic acid, vector, recombinant vector (*e.g.*, rAAV), or recombinant virus particle of the invention, to a subject.

[0177] The invention is useful in animals including human and veterinary medical
15 applications. Suitable subjects therefore include mammals, such as humans, as well as non-human mammals. The term “subject” refers to an animal, typically a mammal, such as humans, non-human primates (apes, gibbons, gorillas, chimpanzees, orangutans, macaques), a domestic animal (dogs and cats), a farm animal (poultry such as chickens and ducks, horses, cows, goats, sheep, pigs), and experimental animals (mouse, rat, rabbit, guinea pig). Human subjects include
20 fetal, neonatal, infant, juvenile and adult subjects. Subjects include animal disease models, for example, mouse and other animal models of blood clotting diseases such as HemA and others known to those of skill in the art.

[0178] Subjects appropriate for treatment in accordance with the invention include those having or at risk of producing an insufficient amount or having a deficiency in a functional gene
25 product (*e.g.*, FVIII protein), or produce an aberrant, partially functional or non-functional gene product (*e.g.*, FVIII protein), which can lead to disease. Subjects appropriate for treatment in accordance with the invention also include those having or at risk of producing an aberrant, or defective (mutant) gene product (protein) that leads to a disease such that reducing amounts, expression or function of the aberrant, or defective (mutant) gene product (protein) would lead to
30 treatment of the disease, or reduce one or more symptoms or ameliorate the disease. Target subjects therefore include subjects having aberrant, insufficient or absent blood clotting factor production, such as hemophiliacs (*e.g.*, hemophilia A).

[0179] Subjects appropriate for treatment in accordance with the invention also include those having or at risk of producing antibodies against AAV. AAV vectors can be administered or
35 delivered to such subjects using several techniques. For example, empty capsid AAV (*i.e.*, AAV

lacking a FVIII nucleic acid) can be delivered to bind to the AAV antibodies in the subject thereby allowing the AAV vector bearing the FVIII nucleic acid to transform cells of the subject. Amounts of empty capsid AAV to administer can be calibrated based upon the amount of AAV antibodies produced in a particular subject. Empty capsid can be of any AAV serotype, for example, AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV12, Rh10, Rh74 or AAV-2i8.

[0180] Alternatively or in addition to, AAV vector can be delivered by direct intramuscular injection (*e.g.*, one or more slow-twitch fibers of a muscle). In another alternative, a catheter introduced into the femoral artery can be used to delivery AAV vectors to liver via the hepatic artery. Non-surgical means can also be employed, such as endoscopic retrograde cholangiopancreatography (ERCP), to deliver AAV vectors directly to the liver, thereby bypassing the bloodstream and AAV antibodies. Other ductal systems, such as the ducts of the submandibular gland, can also be used as portals for delivering AAV vectors into a subject that develops or has preexisting anti-AAV antibodies.

[0181] Administration or *in vivo* delivery to a subject can be performed prior to development of an adverse symptom, condition, complication, etc. caused by or associated with the disease. For example, a screen (*e.g.*, genetic) can be used to identify such subjects as candidates for invention compositions, methods and uses. Such subjects therefore include those screened positive for an insufficient amount or a deficiency in a functional gene product (*e.g.*, FVIII protein), or that produce an aberrant, partially functional or non-functional gene product (*e.g.*, FVIII protein).

[0182] Administration or *in vivo* delivery to a subject in accordance with the methods and uses of the invention as disclosed herein can be practiced within 1-2, 2-4, 4-12, 12-24 or 24-72 hours after a subject has been identified as having the disease targeted for treatment, has one or more symptoms of the disease, or has been screened and is identified as positive as set forth herein even though the subject does not have one or more symptoms of the disease. Of course, methods and uses of the invention can be practiced 1-7, 7-14, 14-21, 21-48 or more days, months or years after a subject has been identified as having the disease targeted for treatment, has one or more symptoms of the disease, or has been screened and is identified as positive as set forth herein.

[0183] A “unit dosage form” as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity optionally in association with a pharmaceutical carrier (excipient, diluent, vehicle or filling agent) which, when administered in one or more doses, is calculated to produce a desired effect (*e.g.*, prophylactic or therapeutic effect). Unit dosage forms may be within, for example, ampules and

vials, which may include a liquid composition, or a composition in a freeze-dried or lyophilized state; a sterile liquid carrier, for example, can be added prior to administration or delivery *in vivo*. Individual unit dosage forms can be included in multi-dose kits or containers. Recombinant vector (*e.g.*, rAAV) sequences, recombinant virus particles, and pharmaceutical compositions thereof can be packaged in single or multiple unit dosage form for ease of administration and uniformity of dosage.

[0184] The invention provides kits with packaging material and one or more components therein. A kit typically includes a label or packaging insert including a description of the components or instructions for use *in vitro*, *in vivo*, or *ex vivo*, of the components therein. A kit can contain a collection of such components, *e.g.*, a nucleic acid, recombinant vector, virus (*e.g.*, AAV) vector, or virus particle and optionally a second active, such as another compound, agent, drug or composition.

[0185] A kit refers to a physical structure housing one or more components of the kit. Packaging material can maintain the components sterilely, and can be made of material commonly used for such purposes (*e.g.*, paper, corrugated fiber, glass, plastic, foil, ampules, vials, tubes, etc.).

[0186] Labels or inserts can include identifying information of one or more components therein, dose amounts, clinical pharmacology of the active ingredient(s) including mechanism of action, pharmacokinetics and pharmacodynamics. Labels or inserts can include information identifying manufacturer, lot numbers, manufacture location and date, expiration dates. Labels or inserts can include information identifying manufacturer information, lot numbers, manufacturer location and date. Labels or inserts can include information on a disease for which a kit component may be used. Labels or inserts can include instructions for the clinician or subject for using one or more of the kit components in a method, use, or treatment protocol or therapeutic regimen. Instructions can include dosage amounts, frequency or duration, and instructions for practicing any of the methods, uses, treatment protocols or prophylactic or therapeutic regimes described herein.

[0187] Labels or inserts can include information on any benefit that a component may provide, such as a prophylactic or therapeutic benefit. Labels or inserts can include information on potential adverse side effects, complications or reactions, such as warnings to the subject or clinician regarding situations where it would not be appropriate to use a particular composition. Adverse side effects or complications could also occur when the subject has, will be or is currently taking one or more other medications that may be incompatible with the composition, or the subject has, will be or is currently undergoing another treatment protocol or therapeutic

regimen which would be incompatible with the composition and, therefore, instructions could include information regarding such incompatibilities.

[0188] Labels or inserts include “printed matter,” *e.g.*, paper or cardboard, or separate or affixed to a component, a kit or packing material (*e.g.*, a box), or attached to an ampule, tube or vial containing a kit component. Labels or inserts can additionally include a computer readable medium, such as a bar-coded printed label, a disk, optical disk such as CD- or DVD-ROM/RAM, DVD, MP3, magnetic tape, or an electrical storage media such as RAM and ROM or hybrids of these such as magnetic/optical storage media, FLASH media or memory type cards.

[0189] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described herein.

[0190]

[0191] Various terms relating to the biological molecules of the invention are used hereinabove and also throughout the specification and claims.

[0192] All of the features disclosed herein may be combined in any combination. Each feature disclosed in the specification may be replaced by an alternative feature serving a same, equivalent, or similar purpose. Thus, unless expressly stated otherwise, disclosed features (*e.g.*, nucleic acid variant, vector, plasmid, recombinant vector (*e.g.*, rAAV) sequence, or recombinant virus particle) are an example of a genus of equivalent or similar features.

[0193] As used herein, the singular forms “a”, “and,” and “the” include plural referents unless the context clearly indicates otherwise. Thus, for example, reference to “a nucleic acid” includes a plurality of such nucleic acids, reference to “a vector” includes a plurality of such vectors, and reference to “a virus” or “particle” includes a plurality of such viruses/particles.

[0194] As used herein, all numerical values or numerical ranges include integers within such ranges and fractions of the values or the integers within ranges unless the context clearly indicates otherwise. Thus, to illustrate, reference to 80% or more identity, includes 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94% etc., as well as 81.1%, 81.2%, 81.3%, 81.4%, 81.5%, etc., 82.1%, 82.2%, 82.3%, 82.4%, 82.5%, etc., and so forth.

[0195] Reference to an integer with more (greater) or less than includes any number greater or less than the reference number, respectively. Thus, for example, a reference to less than 100,

includes 99, 98, 97, etc. all the way down to the number one (1); and less than 10, includes 9, 8, 7, etc. all the way down to the number one (1).

[0196] As used herein, all numerical values or ranges include fractions of the values and integers within such ranges and fractions of the integers within such ranges unless the context clearly indicates otherwise. Thus, to illustrate, reference to a numerical range, such as 1-10 includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, as well as 1.1, 1.2, 1.3, 1.4, 1.5, etc., and so forth. Reference to a range of 1-50 therefore includes 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, etc., up to and including 50, as well as 1.1, 1.2, 1.3, 1.4, 1.5, etc., 2.1, 2.2, 2.3, 2.4, 2.5, etc., and so forth.

[0197] Reference to a series of ranges includes ranges which combine the values of the boundaries of different ranges within the series. Thus, to illustrate reference to a series of ranges, for example, of 1-10, 10-20, 20-30, 30-40, 40-50, 50-60, 60-75, 75-100, 100-150, 150-200, 200-250, 250-300, 300-400, 400-500, 500-750, 750-850, includes ranges of 1-20, 1-30, 1-40, 1-50, 1-60, 10-30, 10-40, 10-50, 10-60, 10-70, 10-80, 20-40, 20-50, 20-60, 20-70, 20-80, 20-90, 50-75, 50-100, 50-150, 50-200, 50-250, 100-200, 100-250, 100-300, 100-350, 100-400, 100-500, 150-250, 150-300, 150-350, 150-400, 150-450, 150-500, etc.

[0198] The invention is generally disclosed herein using affirmative language to describe the numerous embodiments and aspects. The invention also specifically includes embodiments in which particular subject matter is excluded, in full or in part, such as substances or materials, method steps and conditions, protocols, or procedures. For example, in certain embodiments or aspects of the invention, materials and/or method steps are excluded. Thus, even though the invention is generally not expressed herein in terms of what the invention does not include aspects that are not expressly excluded in the invention are nevertheless disclosed herein.

[0199] A number of embodiments of the invention have been described. Nevertheless, one skilled in the art, without departing from the spirit and scope of the invention, can make various changes and modifications of the invention to adapt it to various usages and conditions. Accordingly, the following examples are intended to illustrate but not limit the scope of the invention claimed in any way.

EXAMPLE 1

[0200] Hemophilia A (HA) is an X-linked bleeding disease characterized by deficiency in factor VIII (FVIII), a key component of the coagulation cascade (Kazazian, H.H., et al.

Hemophilia A: Deficiency of coagulation factor VIII. in *The Metabolic and Molecular Bases of Inherited Disease* (eds. Scriver, C.R., Beaudet, A.L., Sly, W.S. & Valle, D.) 4367-4392

(McCraw-Hill, New York, 2001). The FVIII gene contains 26 exons that span 186 kb and is

synthesized as a large precursor molecule (2332 amino acids) (**Figure 1**) (Kaufman, R.J., et al., The biology and genetics of factor VIII deficiency in hemophilia A. in *Hemostasis and Thrombosis: Basic principles and clinical practice* (eds. Colman, R.W., Hirsch, J., Mander, V.J., Clowes, A.W. & George, J.N.) (Lippincott-Raven, Philadelphia, 1998).

5 [0201] Affected individuals commonly suffer joint, muscle, as well as intracranial and intraperitoneal hemorrhages that can be lethal. The normal plasma FVIII level is 100-200 ng/ml, but small amounts of circulating FVIII (~1-2 ng/ml) are sufficient to have a substantial effect on the clinical course of patients with severe disease. The current treatment for HA patients is protein replacement therapy using recombinant or plasma-derived FVIII. However, these
10 products are only available to ~20% of the HA population worldwide. The major complication of this therapy is the development of neutralizing antibodies (inhibitors) to FVIII that occurs in 25-30% of patients with severe HA. Since inhibitors render the FVIII protein therapy ineffective, bypass agents (FVIIa) are used to achieve hemostasis, however, these products are very expensive alternatives.

15 [0202] Disclosed herein are gene constructs for use in gene therapy methods to treat hemophilia. In addition, these factor VIII (FVIII) gene constructs may be useful *in vitro* in the setting of protein expression systems. Each gene construct can optionally include one or more of a expression control (e.g., promoter) element, factor VIII gene and other regulatory features required for expression of the gene, such as introns, ITRs, stop codons, poly A signals, etc.

20 **EXAMPLE 2**

[0203] *Codon-optimization of the factor VIII DNA sequence*

[0204] It has previously been described that codon-optimization of the DNA sequence can improve factor VIII expression in the setting of gene therapy (Ward et al. 2011, Blood 117(3):798-807). Codon-optimization of a DNA sequence is based on the fact that each DNA
25 codon is a series of three nucleotides that encodes for a specific amino acid residue in the protein chain during the protein synthesis. There are 64 different codons but only 20 amino acids thus there are many codons that can code for the same amino acid. There is codon usage bias observed among different species that reflects a selection process for optimizing the translation process. Importantly, while the DNA sequence is modified in codon optimization, the protein sequence is
30 identical to the wild type sequence.

[0205] Codon optimization was performed for the B-domain deleted factor VIII (hFVIII-SQ) cDNA sequence. Four different codon optimized versions of the factor VIII cDNA were assessed in expression assays and surprisingly, several of the DNA sequences expressed higher levels of factor VIII.

[0206] One of the sequences not only expressed higher levels of factor VIII but also had a unique feature (SEQ ID NO:1). This sequence could be packaged into an adeno-associated viral (AAV) vector better than the other codon optimized sequences or the wild type sequence. That is, the factor VIII gene, along with its regulatory elements, are flanked by inverted terminal repeats (ITRs) that provide the signal required to package the sequence into the AAV viral capsid. Data suggests that the size of the DNA sequence between the ITRs must be similar to the size of the normal virus genome (4.7 Kb) for optimal packaging efficiency. In the case of factor VIII, we have gene sequences that are 5.0 Kb in length but still appear to be able to be packaged, however, it may not be efficiently packaged as a complete gene.

[0207] Another factor that may contribute to the ability to package a particular gene is the DNA sequence itself. Some DNA sequences may be more prone to rearrangements, or may have folding structures based on charge, that affect the ability to package into the viral capsid. That is, it appears that not only the size but also the DNA sequence affects the packaging ability of a transgene construct. In our studies we identified novel factor VIII DNA sequences that do not occur in nature that express higher levels of factor VIII (see, e.g., **Figures 2, 4A and 6A**) but also are packaged more efficiently into the AAV vector. This improved packaging feature generates higher vector yields (Tables 3-5) across serotypes and will allow production of homogenous AAV vector that meets stringent quality control criteria for use in human gene therapy.

[0208] To further improve expression levels, codon-optimized human FVIII cDNA constructs were generated with a transthyretin (TTR) or mutant TTR promoter (see description below) and a variant of factor VIII that deletes the PACE-furin cleavage recognition site from amino acid position 1645-1648 (TTRmut-hFVIII-SQ-CO-delP/F) (see description of FVIII variants below). These FVIII transgene constructs were packaged into adeno-associated viral (AAV) vectors and delivered to hemophilia A/CD4 knockout mice. The levels of the FVIII expression were higher than the expression from the wild type hFVIII-SQ as follows: TTR-hFVIII-SQ-CO was 5-fold higher, TTRmut-hFVIII-SQ-CO was 16-fold higher and TTRmut-hFVIII-SQ-CO-delP/F was 33-fold higher.

EXAMPLE 3

[0209] *Variants of factor VIII*

[0210] Modifications were introduced into the protein sequence of factor VIII. The modifications introduced into the codon optimized DNA sequence changes a PACE-furin cleavage recognition site. Modification of this site improves the stability and biological activity of the protein. The amino acid modifications introduced into a codon-optimized factor VIII of PACE-furin cleavage recognition site are shown in Table 1.

[0211] Deletion variants of the PACE-furin (P/F) cleavage recognition site at amino acid position 1645-1648 were introduced into the wild type B-domain deleted human factor VIII (hFVIII) gene. Stable baby hamster kidney (BHK) cells lines were generated that expressed each variant and recombinant protein was purified for analysis.

- 5 [0212] cFVIII, canine factor VIII; hFVIII-RH, a variant that has a single amino acid substitution at amino acid position 1645 from an arginine (R) to a histidine (H) which generates a P/F site that is identical to the cFVIII sequence; Δ 1645, deletion of residue R1645, Δ 2, deletion of residues R1645 and H1646; Δ 3, deletion of residues R1645, H1646 and Q1647; Δ 4, deletion of residues R1645, H1646, Q1647 and R1648; and Δ 1648, deletion of residue R1648.

10 **Table 1: Human Factor VIII variants of the PACE-Furin recognition site**

hFVIII Variant	Amino acids at PACE-furin recognition site						
	1642	1643	1644	1645	1646	1647	1648
hFVIII	V	L	K	R	H	Q	R
hFVIII-RH	V	L	K	H	H	Q	R
Δ 4: hFVIII-del1645-1648	V	L	K	-	-	-	-
Δ 1645: hFVIII-del1645	V	L	K	-	H	Q	R
Δ 2: hFVIII-del1645-1646	V	L	K	-	-	Q	R
Δ 3: hFVIII-del1645-1647	V	L	K	-	-	-	R
Δ 1648: hFVIII-del1648	V	L	K	R	H	Q	-

- [0213] Recombinant variant proteins were purified and have been compared in different assays both in vitro and in vivo to characterize procoagulant function. In vitro the P/F variants have higher biological activity than wild type hFVIII with the Δ 3 having the highest biological activity followed by the Δ 4 and Δ 2 variants.
- 15

[0214] In vivo, all of these variant hFVIII proteins were infused into hemophilia A mice in a tail clip challenge assay to measure blood loss after an injury. The Δ 3 and Δ 4 variants showed improved ability to reduce blood loss in this challenge model.

- [0215] All five of the P/F deletion variants were introduced into an adeno-associated viral (AAV) vector expression cassette containing the wild type factor VIII gene (**Figure 3A**). AAV-hFVIII was delivered to hemophilia A/CD4KO mice (5×10^{11} vg/mouse) and expression of FVIII was determined by antigen and activity (**Figure 3B**). The expression levels of Δ 3 were 4-fold higher than wild type FVIII-BDD while Δ 4 and Δ 1645 were two and three-fold higher, respectively. The Δ 2 and Δ 1648 were similar to wild type FVIII. These results support the
- 20

introduction of the $\Delta 3$ and $\Delta 4$ PACE-furin deletion variants into the improved codon-optimized FVIII constructs.

EXAMPLE 4

5 [0216] *TTR Promoter*

[0217] The improved promoter element that utilized in these studies is the transthyretin (TTR) promoter with four nucleotide changes that improves factor VIII expression. The characterization of this promoter was originally described in Costa and Grayson 1991, Nucleic Acids Research 19(15):4139-4145. These in vitro studies suggested that a modification of four
10 nucleotides increases the affinity of the hepatocyte nuclear factor (HNF) transcription factor for its binding site in the DNA sequence. Thus, in a novel synthetic promoter, we introduced the four nucleotide changes beginning at nucleotide 67 of the TTR promoter sequence that is not known to occur in nature. It was modified from TAmGTGTAG to TATTGACTTAG (SEQ ID NO:8).

15 **EXAMPLE 5**

[0218] *Comparison of wild type and codon-optimized (CO) transgenes*

[0219] AAV transgene constructs were generated to demonstrate the contribution of each FVIII sequence or sequence variant to the improved FVIII expression. These constructs contain the same regulatory elements as the construct described in **Figure 2**; however, their FVIII
20 sequence is different. Based on the studies described above, the best performing PACE-furin deletion variants ($\Delta 3$ and $\Delta 4$) were introduced into these hFVIII transgenes. Also, all of these FVIII transgenes are a B-domain deleted (BDD) form. All of AAV vectors in this study were produced at the Research Vector Core at CCMT and were titered side by side by quantitative PCR and silver staining.

25

Table 2A shows hFVIII Sequence Identities between WT hFVIII and 3 Codon Optimized Variants, denoted CO1, CO2 and CO3

hFVIII 4374 nucleotides

Sequence 1	Sequence 2	% Identity	# NT Identities	# NT Differences
WT	CO3	77.34%	3386	992
WT	CO1	77.14%	3374	1000
WT	CO2	75.74%	3315	1062
CO3	CO1	82.20%	3597	779
CO3	CO2	81.91%	3590	793

Table 2B shows nucleotide frequencies for each codon optimized hFVIII variant denoted CO1, CO2 and CO3

	WT hFVIII	CO3	CO2	CO1
Nucleotide Frequency	Frequency, %	Frequency, %	Frequency, %	Frequency, %
A	A: 1,266 28.9%	A: 1,145 26.2%	A: 1,072 24.5%	A: 1,063 24.3%
C	C: 970 22.2%	C: 1,108 25.3%	C: 1,197 27.4%	C: 1,371 31.3%
G	G: 964 22.0%	G: 1,133 25.9%	G: 1,149 26.3%	G: 1,191 27.2%
T	T: 1,174 26.8%	T: 988 22.6%	T: 956 21.9%	T: 749 17.1%
GC	GC: 1,934 44.2%	GC: 2,241 51.2%	GC: 2,346 53.6%	GC: 2,562 58.6%

[0220] In this study, the five human FVIII transgenes were: (1) wild type human FVIII-BDD (hFVIIIwtSQ); (2) wild type human FVIII-BDD with the deletion of four residues in the PACE-furin recognition sequence (hFVIIIwtSQΔ4); (3) codon-optimized (CO) hFVIII-BDD (hFVIIIICO); (4) codon-optimized with deletion of three residues of the PACE-furin recognition

site (hFVIII Δ 3); and (5) codon-optimized hFVIII with deletion of four residues of the PACE-furin recognition site (hFVIII Δ 4)(**Figure 6**).

[0221] This data is an improvement of the data in **Figure 2**, in which the wild type FVIII construct used a different promoter (HCR-hAAT) and was not in the same expression cassette as the codon optimized (CO) constructs making it difficult to demonstrate the individual contribution of each FVIII transgene. In addition, the wild type FVIII with the PACE/furin variant that deletes four amino acid residues at this site (hFVIIIwtSQ Δ 4) is included to demonstrate the contribution of the PACE-furin deletion in the context of the wild type hFVIII (**Figure 4**).

10 [0222] As described, AAV-hFVIII was delivered to hemophilia A/CD4KO mice (1×10^{11} vg/mouse). However, a different AAV serotype that was developed based upon an AAV variant, denoted Rh74_{VV}. Previous data was generated with AAV serotype 8 (AAV8).

[0223] The hFVIII expression was measured by antigen levels (ELISA) as well as activity (Coatest assay). For Rh74_{VV}, levels of hFVIII expression at 4 weeks post vector administration
15 were 9.4+1.5ng/ml (6.3% of normal FVIII levels) for (hFVIIIwtSQ), 32.4+13.0ng/ml (21.6%) for (hFVIIIwtSQ Δ 4), 42.2+7.1ng/ml (28.1%) (hFVIII Δ CO), 177.8+8.5ng/ml (118.5%) (hFVIII Δ CO Δ 3) and 114.3+60.2ng/ml (76.2%) (hFVIII Δ CO Δ 4) (**Figure 6**).

[0224] At 6 weeks post vector administration, mice after AAV delivery of wild type and codon-optimized hFVIII constructs were assayed by in vivo hemostatic challenge. The results
20 (**Figure 7**) show that clotting activity in the hemophilia A mice, as reflected by reduced blood loss, is greater for the CO optimized FVIII with or without the P/F deletions, and is comparable to wild type mice.

[0225] Since the levels of FVIII protein expression obtained with the AAV variant (Rh74_{VV}) are lower than the FVIII levels with AAV8, the data is presented using fold difference compared
25 to the wild type FVIII construct. Importantly, the overall conclusions are the same. Levels of expression with the introduction of the Δ 4 P/F variant is about 2-4 fold higher compared to the wild type sequence alone. The CO transgene results in expression that is 5-fold higher than the wild type hFVIII (hFVIII-wtSQ). The introduction of the P/F deletion variants into the codon-optimized hFVIII (hFVIII-CO) results in expression that is 3-4 fold higher (Δ 3) or 2-3 fold higher
30 (Δ 4) than the CO transgene alone. Thus, the hFVIII Δ CO Δ 3 construct is 16-19-fold better than hFVIIIwtSQ, which is substantially higher than either the P/F deletion variants or codon optimization alone.

[0226] Significantly, the contribution of the P/F deletion variants is consistent with both the wild type and the codon-optimized hFVIII sequences (**Figure 4**). Furthermore, this contribution

was also observed using another expression cassette (**Figure 3**). That is, the $\Delta 3$ P/F variant results in 3-4 fold higher expression levels and $\Delta 4$ results in 2-3 fold higher expression levels regardless of whether they are in the wild type DNA sequence or the codon-optimized DNA sequence. Although not wishing to be bound by any particular theory, the combined synergistic effect is believed due to the different mechanisms that are responsible for the codon-optimization (improved translation of the RNA sequence) and the PACE-furin variants (improved secretion and higher biological activity of the protein).

[0227] As disclosed (Figure 2), the TTRm promoter increases expression 2-fold compared to the TTR promoter alone. While not included in this study, a reasonable estimate based on the other studies is that the optimized construct—TTRm-hFVIII Δ CO3 is ~30-40-fold better than TTR-hFVIIIwtSQ.

EXAMPLE 6

[0228] *Vector Yields of Codon-Optimized and Non-Codon Optimized Constructs*

[0229] Adeno-associated viral (AAV) vector yields for human factor VIII preparations are shown in Tables 3-5. The adeno-associated viral (AAV) vector genomes per roller bottle for wild type human FVIII-SQ (hFVIII) (3 preparations, denoted Prep1-Prep3) and the codon-optimized human FVIII (hFVIII-CO, also referred to as CO3) (5 separate preparations of CO3, denoted Prep1-Prep5) are shown. All vector preparations of FVIII-CO3 were made using the same production protocol and purification scheme. The average yield of AAV8-hFVIII-SQ-CO was 4.60×10^{12} vg/rollerbottle (13 vector preparations) compared to the average yield of the AAV8-hFVIII-SQ of 2.77×10^{12} vg/rollerbottle (3 vector preparations). The yields are 2-fold higher with the CO FVIII sequence compared to the wild type FVIII sequence.

Table 3: AAV-8 vector yields for human factor VIII preparations

hFVIII Transgene	AAV Vector	Vector Genomes/ Roller Bottle
hFIX	AAV8-hFIX-19	1.00×10^{13}
Wild type	AAV8-hFVIII-Prep 1	2.20×10^{12}
	AAV8-hFVIII-Prep 2	3.66×10^{12}
	AAV8-hFVIII-Prep 3	2.46×10^{12}
	AAV8-hFVIII-CO-Prep 1	4.54×10^{12}
Codon Optimized	AAV8-hFVIII-CO-Prep 2	3.62×10^{12}
	AAV8-hFVIII-CO-Prep 3	4.71×10^{12}
	AAV8-hFVIII-CO-Prep 4	5.00×10^{12}
	AAV8-hFVIII-CO-Prep 5	4.30×10^{12}

[0230] The mean yield of all vector preparations of FVIII with the standard deviation are shown in Tables 4 and 5. Again, while AAV preparations produced by a different vector core were observed to have a lower yield, a comparison of both the wild type vectors and the CO vectors in a different serotype (Rh74_{VV}) shows that the yields are consistently 2-fold higher. This demonstrates that yield of FVIII vector preparations is independent of the vector core and the AAV serotype.

Table 4: AAV-8 vector yields for human factor VIII preparations with standard deviation

hFVIII Transgene	AAV Vector	Number of Preparations	Mean Vector Genomes/ Roller Bottle
hFIX	AAV8-hFIX-19		1.00×10^{13}
Wild type	AAV8-hFVIII	3	$2.67 \times 10^{12} \pm 9.29 \times 10^{11}$
Codon Optimized	AAV8-hFVIII-CO	13	$4.60 \times 10^{12} \pm 6.85 \times 10^{11}$

Table 5: Rh74_{VV} AAV vector yields for human factor VIII preparations with standard deviation

hFVIII Transgene	AAV Vector	Number of Preparations	Mean Vector Genomes/ Roller Bottle
Wild type	Rh74 _{VV} -hFVIII	7	$9.58 \times 10^{11} \pm 5.07 \times 10^{11}$
Codon Optimized	Rh74 _{VV} -hFVIII-CO	5	$2.39 \times 10^{12} \pm 6.85 \times 10^{11}$

Codon-optimized factor VIII cDNA, aka CO/CO3 (SEQ ID NO:1)

5 ATGCAGATTGAGCTGTCAACTTGCTTTTTCTGTGCCTGCTGAGATTTTGTTTTTCCGCTACTAGAAG
 ATACTACCTGGGGGCTGTGGAACGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 ACGCAGCATTTCCACCTAGAGTCCCTAAATCATTCCTTCAACACCAGCGTGGTCTATAAGAAAAACA
 CTGTTCTGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGATGGGACTGCT
 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCATTACACTGAAAAACATGGCCTCACACC
 10 CCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCGAAAGGGGCAGAGTATGACGATCAG
 ACTTCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTGGCGGTCTCATACCTATGTGTGGCAGGT
 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCCTCTGTGCCTGACCTACTCTTATCTGAGTCACGTGG
 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGAAGGGAGCCTGGCT
 AAGGAGAAAACCCAGACACTGCATAAGTTTCATTCTGCTGTTTCGCCGTGTTTGACGAAGGAAAAATCATG
 15 GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCTTGGCCCAAAA
 TGCACACTGTGAACGGCTACGTCAATCGTCACTGCCTGGACTGATCGGCTGCCACCGAAAGAGCGTG
 TATTGGCATGTATCGGAATGGGCACCACACCTGAAGTGCACTCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCTCCCTGGAGATCTCTCCAATTACCTTCTTGACAGCTCAGACTC
 TGCTGATGGATCTGGGACAGTTCCTGCTGTTTTGCCACATCAGCTCCCACCAGCATGATGGCATGGAG
 20 GCCTACGTGAAAGTGGACAGCTGTCCCAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTTCGATGACGATAACAGCCCCCT
 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCACTGGTGTGGCACCAGACGATCGATCCTACAAAATCTCAGTATCT
 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAGGTTTCATGGCTTATACCGATGAAA
 25 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 GGGGACACCTGCTGATCATTTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 TACAGATGTGCGCCCTCTGTACAGCCGAGACTGCCAAAGGGCGTCAAACACCTGAAGGACTTCCCAA
 TCCTGCCCCGGGAAATTTTTAAGTATAAATGGACTGTCACCGTCGAGGATGGCCCCACTAAGAGCGAC
 CCTAGGTGCCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAAGGGATCTGGCCAGCGGACTGAT
 30 CGGCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA
 GGAATGTGATTCTGTTCAGTGTCTTTGACGAAAACCGGTTCATGGTATCTGACCGAGAACATCCAGAGA
 TTCTTGCCATAATCCAGCCGGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 TATTAATGGCTACGTGTTTCGACAGTCTGCAGCTGTCACTGTGTCTGCACGAGGTTCGCTTACTGGTATA
 TCCTGAGCATTGGAGCACAGACAGATTTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAA
 35 ATGGTGTATGAGGACACACTGACTCTGTTTCCCTTCAGCGGCGAAACCGTGTATGTCCATGGAGAA
 TCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTCAGGAATCGCGGGATGACTGCCCTGCTGA
 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTATACGAGGACATCAGCGCATAT
 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCTTTTAGTCAGAATCCTCCAGTGCTGAAGAGGCA
 CCAGCGCGAGATCACCCGCACTACCCTGCAGAGTGATCAGGAAGAGATCGACTACGACGATACAATTT
 40 CTGTGGAATGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCTCGATCATTC
 CAGAAGAAAACCCGGCATTACTTTATTGCTGCAGTGGAGCGCTGTGGGATTATGGCATGTCTCTAG
 TCCTCACGTGCTGCGAAATCGGGCCAGTCAGGGAGCGTCCACAGTTCAAGAAAGTGGTCTTCAGG
 AGTTTACAGACGGATCCTTTACTCAGCCACTGTACCGGGCGAACTGAACGAGCACCTGGGGCTGCTG

GGACCTATATCAGAGCTGAAGTGGAGGATAACATTATGGTCACCTTCAGAAATCAGGCATCTAGGCC
 TTACAGTTTTTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGGAGCAGAACCACGAAAAA
 ACTTCGTGAAGCCTAATGAGACCAAAACATACTTTTGGAAAGGTGCAGCACCATATGGCCCCAACAAAA
 GACGAATTCGATTGCAAGGCATGGGCCTATTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGG
 5 CCTGATCGGGCCACTGCTGGTGTGTCATACTAACACCCTGAATCCCGCACACGGCAGGCAGGTCACTG
 TCCAGGAATTCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAGCTGGTACTTCACCGAAAACATG
 GAGCGAAATTGCCGGGCTCCATGTAATATTAGATGGAAGACCCACATTCAAGGAGAATACCGCTT
 TCATGCCATCAATGGGTATATTATGGATACTCTGCCCGGACTGGTCATGGCTCAGGACCAGAGAATCA
 GGTGGTACCTGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTCAATTCAGCGGACACGTGTTT
 10 ACTGTCCGGAAGAAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGGCGTGTTCGAAACCGT
 CGAGATGCTGCCTAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCTGATTGGGGAGCACCTGCATGCCG
 GAATGTCTACCCTGTTTCTGGTGTACAGTAATAAGTGTGACACACCCCTGGGGATGGCTTCCGGACAT
 ATCCGGGATTTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCA
 CTATTCCGGGTCTATCAACGCTTGGTCCACAAAAGAGCCCTTCTCTTGGATTAAAGTGGACCTGCTGG
 15 CACCAATGATCATTGATGGCATCAAACTCAGGGGGCCAGGCAGAAAGTCTCCTCTCTGTACATCTCA
 CAGTTTATCATCATGTACAGCCTGGATGGCAAGAAATGGCAGACATAACCGGCAATAGCACAGGGAC
 TCTGATGGTGTCTTTGGCAACGTGGACAGTTCAGGGATCAAGCACAAACATTTTCAATCCCCCTATCA
 TTGCTAGATACATCAGGCTGCACCCAACCCATTATTCTATTGAAGTACACTGCGGATGGAACGTATG
 GGGTGCATCTGAACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTGACGCCAGAT
 20 TACCGCTAGCTCCTACTTCACTAATATGTTTGCTACCTGGAGCCCTCCAAAGCACGACTGCATCTGC
 AGGGACGAAGCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTGCGATTTTCAG
 AAACTATGAAGGTGACCGGAGTCACAACCTCAGGGCGTGAAAAGTCTGCTGACCTCAATGTACGTCAA
 GGAGTTCCTGATCTCTAGTTCACAGGACGGCCACCAAGTGGACACTGTTCTTTCAGAACGGAAAGGTGA
 AAGTCTTCCAGGGCAATCAGGATTCTTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACT
 25 CGCTACCTGCGAATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTGGGCTG
 CGAGGCCAGGACCTGTATTGA

Codon-optimized factor VIII cDNA, aka CO/CO3, with R1645H variant (SEQ ID NO:2)

ATGCAGATTGAGCTGTCAACTTGCTTTTTTCTGTGCTGCTGAGATTTTGTTTTTCCGCTACTAGAAG
 ATACTACCTGGGGGCTGTGGAACGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 30 ACGCACGATTTCCACCTAGAGTCCCTAAATCATTTCCCTTCAACACCAGCGTGGTCTATAAGAAAAACA
 CTGTTTCGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCTCGGCCACCCCTGGATGGGACTGCT
 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCAATACACTGAAAAACATGGCCTCACACC
 CCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCCGAAGGGGCAGAGTATGACGATCAG
 ACTTCCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTTGGCGGGTCTCATACCTATGTGTGGCAGGT
 35 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCTCTGTGCTGACCTACTCTTATCTGAGTCACGTGG
 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGAAGGGAGCCTGGCT
 AAGGAGAAAACCCAGACACTGCATAAGTTCATTCTGCTGTTCCCGTGTGTTGACGAAGGAAAATCATG
 GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCTTGGCCCCAAA
 TGCACACTGTGAACGGCTACGTCAATCGTCACTGCTGCTGGACTGATCGGCTGCCACCGAAAAGAGCGTG
 40 TATTGGCATGTATCGGAATGGGCACCACACCTGAAGTGCATCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCTCCCTGGAGATCTCTCCAATTACCTTCTGACAGCTCAGACTC
 TGCTGATGGATCTGGGACAGTTCCTGCTGTTTTGCCACATCAGCTCCCACCAGCATGATGGCATGGAG
 GCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTGATGACGATAACAGCCCCCT
 45 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCACTGGTGTGGCACCAGACGATCGATCTACAAATCTCAGTATCT
 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAGGTTTCATGGCTTATACCGATGAAA
 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 GGGGACACCCTGCTGATCATTTTTTAAGAACAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 50 TACAGATGTGCGCCCTCTGTACAGCCGGAGACTGCCAAAGGGCGTCAAAACACCTGAAGGACTTCCCAA
 TCCTGCCCGGGGAAATTTTTAAGTATAAATGGACTGTCACCGTTCGAGGATGGCCCCACTAAGAGCGAC
 CCTAGGTGCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAGGGATCTGGCCAGCGGACTGAT
 CGGCCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA

GGAATGTGATTCTGTTTCAGTGTCTTTGACGAAAACCGGTCATGGTATCTGACCGAGAACATCCAGAGA
 TTCCTGCCTAATCCAGCCGGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 TATTAATGGCTACGTGTTTCGACAGTCTGCAGCTGTCAGTGTGTCTGCACGAGGTCGCTTACTGGTATA
 TCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAA
 5 ATGGTGTATGAGGACACACTGACTCTGTTCCCTTCAGCGGCGAAAACCGTGTATGTCCATGGAGAA
 TCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTAGGAATCGCGGGATGACTGCCCTGCTGA
 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTCATACGAGGACATCAGCGCATAT
 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCTTTTAGTCAGAATCCTCCAGTGTCTGAAGCACC
 CCAGCGCGAGATCACCCGCACTACCCTGCAGAGTGATCAGGAAGAGATCGACTACGACGATACAATTT
 10 CTGTGGAATGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCTCTGATCATTC
 CAGAAGAAAACCCGGCATTACTTTATTGCTGCAGTGGAGCGCTGTGGGATTATGGCATGTCTCTAG
 TCCTCACGTGCTGCGAAATCGGGCCAGTCAGGGAGCGTCCACAGTTCAAGAAAGTGGTCTTCCAGG
 AGTTTACAGACGGATCCTTTACTCAGCCACTGTACCGGGGCGAACTGAACGAGCACCTGGGGCTGCTG
 GGACCCATATCAGAGCTGAAGTGGAGGATAACATTATGGTCACCTTCAGAAATCAGGCATCTAGGCC
 15 TTACAGTTTTTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGAGCAGAACCACGAAAAA
 ACTTCGTGAAGCCTAATGAGACCAAAACATACTTTTGGAAAGGTGCAGCACCATATGGCCCCAACAAAA
 GACGAATTCGATTGCAAGGCATGGGCCATTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGG
 CCTGATCGGGCCACTGCTGGTGTGTCTACTAACACCCCTGAATCCCGCACACGGCAGGCAGGTCACTG
 TCCAGGAATTCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAAGCTGGTACTTCACCGAAAAATG
 20 GAGCGAAATTGCCGGGCTCCATGTAATATTAGATGGAAGACCCACATTCAAGGAGAACTACCGCTT
 TCATGCCATCAATGGGTATATTATGGATACTCTGCCCGGACTGGTCATGGCTCAGGACCAGAGAATCA
 GGTGGTACCTGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTCATTTCAGCGGACACGTGTTT
 ACTGTCCGGAAGAAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGGCGTGTTCGAAACCGT
 CGAGATGCTGCCTAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCTGATTGGGGAGCACCTGCATGCCG
 25 GAATGTCTACCCTGTTTCTGGTGTACAGTAATAAGTGTGAGACACCCCTGGGGATGGCTTCCGGACAT
 ATCCGGGATTTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCA
 CTATTCGGGTCTATCAACGCTTGGTCCACAAAAGAGCCTTTCTCTTGGATTAAGGTGGACCTGCTGG
 CACCAATGATCATTATGGCATCAAACTCAGGGGGCCAGGCAGAAAGTCTCTCTCTGTACATCTCA
 CAGTTTATCATCTGATGACGCTGGATGGCAAGAAATGGCAGACATACCGCGGCAATAGCAGAGGGAC
 30 TCTGATGGTGTCTTTGGCAACGTGGACAGTTCAAGGATCAAGCACAACATTTTCAATCCCCCTATCA
 TTGCTAGATACATCAGGCTGCACCCAACCCATTATTCTATTGAAGTACACTGCGGATGGAAGTATG
 GGGTGCATCTGAACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTGACGCCAGAT
 TACCGCTAGCTCCTACTTCACTAATATGTTTGCTACCTGGAGCCCTCCAAAGCACGACTGCATCTGC
 AGGGACGAAGCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTGCATTTTCAG
 35 AAACTATGAAGGTGACCGGAGTCAACAACCTCAGGGCGTGAAAAGTCTGCTGACCTCAATGTACGTCAA
 GGAGTTCCTGATCTCTAGTTCACAGGACGGCCACCAGTGGACACTGTTCTTTCAGAACGGAAAGGTGA
 AAGTCTTCCAGGGCAATCAGGATTCTTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACT
 CGCTACCTGCGAATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTCTGGGCTG
 CGAGGCCCAGGACCTGTATTGA

40 **Codon-optimized factor VIII cDNA, aka CO/CO3, with deletion of amino acid 1645-1648
 variant, Δ4 (SEQ ID NO:3)**

ATGCAGATTGAGCTGTCAACTTGCTTTTTTCTGTGCTGCTGAGATTTTGTTTTTCCGCTACTAGAAG
 ATACTACCTGGGGGCTGTGGAAGTGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 ACGCACGATTTCCACCTAGAGTCCCTAAATCATTCCCTTCAACACCAGCGTGGTCTATAAGAAAAACA
 45 CTGTTTCGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGATGGGACTGCT
 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCATTACACTGAAAAACATGGCCTCACACC
 CCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCCGAAGGGGCAGAGTATGACGATCAG
 ACTTCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTGGCGGGTCTCATACCTATGTGTGGCAGGT
 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCCTCTGTGCCTGACCTACTCTTATCTGAGTCACGTGG
 50 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACCTGCTGGTGTGTAGGGAAGGGAGCCTGGCT
 AAGGAGAAAACCCAGACACTGCATAAGTTTCTGCTGTTTCCCGTGTGTTGACGAAGGAAAAATCATG
 GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCTTGGCCAAAAA
 TGCACACTGTGAACGGCTACGTCAATCGTCACTGCCCTGGACTGATCGGCTGCCACCGAAAGAGCGTG

TATTGGCATGTCATCGGAATGGGCACCACACCTGAAGTGCACTCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCCTCCCTGGAGATCTCTCCAATTACCTTCTGACAGCTCAGACTC
 TGCTGATGGATCTGGGACAGTTCCTGCTGTTTTGCCACATCAGCTCCCACCAGCATGATGGCATGGAG
 GCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 5 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTTCGATGACGATAACAGCCCCCT
 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCCTGGTGTGGCACCAGACGATCGATCCTACAAATCTCAGTATCT
 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAGGTTTCATGGCTTATACCGATGAAA
 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 10 GGGGACACCCTGCTGATCATTTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 TACAGATGTGCGCCCTCTGTACAGCCGGAGACTGCCAAAGGGCGTCAAACACCTGAAGGACTTCCCAA
 TCCTGCCCGGGGAAATTTTTAAGTATAAATGGACTGTACCGTTCGAGGATGGCCCCACTAAGAGCGAC
 CTTAGGTGCCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAGGGATCTGGCCAGCGGACTGAT
 CGGCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA
 15 GGAATGTGATTCTGTTCAGTGTCTTTGACGAAAACCGGTTCATGGTATCTGACCGAGAACATCCAGAGA
 TTCTGCTAATCCAGCCGGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 TATTAATGGCTACGTGTTCGACAGTCTGCAGCTGTCACTGTGTCTGCACGAGGTTCGCTTACTGGTATA
 TCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAA
 ATGGTGTATGAGGACACACTGACTCTGTTCCCTTCAGCGGCGAAACCGTGTTCATGTCCATGGAGAA
 20 TCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTTCAAGGAATCGCGGGATGACTGCCCTGCTGA
 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTCATACGAGGACATCAGCGCATAT
 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCTTTTAGTCAGAATCCTCCAGTGTCTGAAGGAGAT
 CACCCGCACTACCCTGCAGAGTGATCAGGAAGAGATCGACTACGACGATACAATTTCTGTGGAAATGA
 AGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCCCTCGATCATTCCAGAAGAAAAACC
 25 CGGCATTACTTTATTTGCTGCAGTGGAGCGCTGTGGGATTATGGCATGTCTCTAGTCTCACGTGCT
 GCGAAATCGGGCCCAGTCAGGGAGCGTCCACAGTTCAAGAAAGTGGTCTTCCAGGAGTTTACAGACG
 GATCCTTTACTCAGCCACTGTACCGGGGCGAAGTGAACGAGCACCTGGGGCTGCTGGGACCCCTATATC
 AGAGCTGAAGTGGAGGATAACATTATGGTCACTTCAGAAATCAGGCATCTAGGCCTTACAGTTTTTA
 TTCAAGCCTGATCTCTTACGAAGAGGACAGAGGCGAGGAGCAGAACACGAAAAAACTTCGTGAAGC
 30 CTAATGAGACCAAAACATACTTTTGGAAAGTGCAGCACCATATGGCCCCAACAAAAAGACGAATTCGAT
 TGCAAGGCATGGGCCTATTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGGCCTGATCGGGCC
 ACTGCTGGTGTGTACTACTAACACCCTGAATCCCGCACACGGCAGGCAGGTCACTGTCCAGGAATTTCG
 CCCTGTTCTTTACCATCTTTGATGAGACAAAAAGCTGGTACTTCACCGAAAAACATGGAGCGAAATTGC
 CGGGCTCCATGTAATATTCAGATGGAAGACCCACATTCAAGGAGAACTACCGCTTTCATGCCATCAA
 35 TGGGTATATTATGGATACTCTGCCCCGACTGGTCAITGGCTCAGGACCAGAGAATCAGGTGGTACCTGC
 TGAGCATGGGGTCCAACGAGAATATCCACTCAATTCATTTCAGCGGACACGTGTTTACTGTCCGGAAG
 AAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGCGTGTTCGAAACCGTCGAGATGCTGCC
 TAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCTGATTGGGGAGCACCTGCATGCCGAATGTCTACCC
 TGTTTCTGGTGTACAGTAATAAGTGTGACACACCCCTGGGGATGGCTTCCGGACATATCCGGGATTTT
 40 CAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCCCTAAGCTGGCTAGACTGCACTATTCCGGGTC
 TATCAACGCTTGGTCCACAAAAGAGCCTTTCTCTTGGATTAAAGGTGGACCTGCTGGCACCAATGATCA
 TTCATGGCATCAAACTCAGGGGGCCAGGCAGAAGTTCTCCTCTCTGTACATCTCACAGTTTATCATC
 ATGTACAGCCTGGATGGCAAGAAATGGCAGACATACCGCGCAATAGCACAGGGACTCTGATGGTGT
 CTTTGGCAACGTGGACAGTTCAGGGATCAAGCACAACTTTTCAATCCCCCTATCATTGCTAGATACA
 45 TCAGGCTGCACCCAACCCATTATTCTATTGCAAGTACACTGCGGATGGAAGTATGAGGCTGCGATCTG
 AACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTGACGCCAGATTACCGCTAGCTC
 CTACTTCACTAATATGTTTGCTACCTGGAGCCCCCTCCAAAGCACGACTGCATCTGCAGGGACGAAGCA
 ACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTTCGATTTTCAGAAAACTATGAAG
 GTGACCGGAGTCACTCACTCAGGGCGTGAAGTCTGCTGACCTCAATGTACGTCAAGGAGTTCTTGAT
 50 CTCTAGTTTACAGGACGGCCACCAGTGGACACTGTCTTTTCAAGACGGAAGGTGAAAGTCTTCCAGG
 GCAATCAGGATTCTTTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACTCGCTACCTGCGA
 ATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTGGGCTGCGAGGCCAGGA
 CCTGATTGA

Codon-optimized factor VIII cDNA, aka CO/CO3, with deletion of amino acid 1645 variant, Δ1645 (SEQ ID NO:4)

ATGCAGATTGAGCTGTCAACTTGCTTTTTCTGTGCCTGCTGAGATTTTGTTTTTCCGCTACTAGAAG
 A TACTACCTGGGGGCTGTGGAAGTGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 5 ACGCACGATTTCCACCTAGAGTCCCTAAATCATTCCCCTTCAACACCAGCGTGGTCTATAAGAAAAACA
 CTGTTTCGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGATGGGACTGCT
 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCATTACACTGAAAAACATGGCCTCACACC
 CCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCCGAAGGGGCAGAGTATGACGATCAG
 ACTTCCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTGGCGGGTCTCATACCTATGTGTGGCAGGT
 10 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCCCTGTGCGCTGACCTACTCTTATCTGAGTCACGTGG
 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGAAGGGAGCCTGGCT
 AAGGAGAAAACCCAGACACTGCATAAGTTTATTCTGCTGTTTCGCCGTGTTTGACGAAGGAAAATCATG
 GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCTTGGCCCAAAA
 TGCACACTGTGAACGGCTACGTCAATCGCTCACTGCCCTGGACTGATCGGCTGCCACCGAAAGAGCGTG
 15 TATTGGCATGTGATCGGAATGGGCACCACACCTGAAGTGCATCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCCTCCCTGGAGATCTCTCCAATTACCTTCTGACAGCTCAGACTC
 TGCTGATGGATCTGGGACAGTTCCTGCTGTTTTGCCACATCAGCTCCCACCAGCATGATGGCATGGAG
 GCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTTCGATGACGATAACAGCCCCCT
 20 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCACTGGTGTGGCACCAGACGATCGATCCTACAAATCTCAGTATCT
 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAGGTTTATGGCTTATACCGATGAAA
 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 GGGGACACCCTGCTGATCATTTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 25 TACAGATGTGCGCCCTCTGTACAGCCGGAGACTGCCAAAGGGCGTCAAAACACCTGAAGGACTTCCCAA
 TCCTGCCCGGGGAAATTTTAAAGTATAAATGGACTGTCAACCGTCGAGGATGGCCCCACTAAGAGCGAC
 CCTAGGTGCCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAAGGGATCTGGCCAGCGGACTGAT
 CGGCCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA
 GGAATGTGATTCTGTTTCAGTGTCTTTGACGAAAACCGGTGATGGTATCTGACCGAGAACATCCAGAGA
 30 TTCTGCTAATCCAGCCGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 TATTAATGGCTACGTGTTTCGACAGTCTGCAGTGTGCTGTCGACGAGGTGCGTTACTGGTATA
 TCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAAA
 ATGGTGTATGAGGACACACTGACTCTGTTCCCTTCAGCGGCGAAAACCGTGTATTATGTCCATGGAGAA
 TCCCGGGCTGTGGATCTGGGATGCCACAACAGCGATTTTCAAGGAATCGCGGGATGACTGCCCTGCTGA
 35 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTCATACGAGGACATCAGCGCATAT
 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCTTTTAGTCAGAATCCTCCAGTGTGTAAGCACC
 GCGCGAGATCACCCGCACTACCCTGCAGAGTGATCAGGAAGAGATCGACTACGACGATACAATTTCTG
 TGGAAATGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCCCTCGATCATTCCAG
 AAGAAAACCCGGCATTACTTTATTGCTGCAGTGGAGCGCCTGTGGGATTATGGCATGTCTCTAGTCC
 40 TCACGTGCTGCGAAATCGGGCCAGTCAGGGAGCGTCCCACAGTTCAAGAAAGTGGTCTTCCAGGAGT
 TTACAGACGGATCCTTTACTCAGCCACTGTACCGGGCGAACTGAACGAGCACCTGGGGCTGCTGGGA
 CCCTATATCAGAGCTGAAGTGGAGGATAACATTATGGTCACCTTCAGAAATCAGGCATCTAGGCCTTA
 CAGTTTTTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGGAGCAGAACCACGAAAAAACT
 TCGTGAAGCCTAATGAGACCAAAACATACTTTTGAAGGTGCAGCACCATATGGCCCCAACAAAAGAC
 45 GAATTGATTGCAAGGCATGGGCCATTTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGGCCT
 GATCGGGCCACTGCTGGTGTGTCATACTAACACCTGAATCCCGCACACGCGCAGGCAGGTCACTGTCC
 AGGAATTCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAAGCTGGTACTTCACCGAAAAACATGGAG
 CGAAATTGCCGGGCTCCATGTAATATTAGATGGAAGACCCACATTCAAGGAGAACTACCGCTTTTCA
 TGCCATCAATGGGTATATTATGGATACTCTGCCCGGACTGGTCATGGCTCAGGACCAGAGAATCAGGT
 50 GGTACCTGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTCATTTACGCGGACACGTGTTTACT
 GTCCGGAAGAAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGGCGTGTTCGAAACCGTCTGA
 GATGCTGCCTAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCTGATTGGGGAGCACCTGCATGCCGAA
 TGTCTACCCTGTTTCTGGTGTACAGTAATAAGTGTGACACACCCCTGGGGATGGCTTCCGGACATATC
 CGGGATTTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCACTA

TTCCGGGTCTATCAACGCTTGGTCCACAAAAGAGCCTTTCTCTTGGATTAAAGGTGGACCTGCTGGCAC
 CAATGATCATTTCATGGCATCAAACTCAGGGGGCCAGGCAGAAATTCTCTCTCTGTACATCTCACAG
 TTTATCATCATGTACAGCCTGGATGGCAAGAAATGGCAGACATACCGCGGCAATAGCACAGGGACTCT
 GATGGTGTCTTTGGCAACGTGGACAGTTCAGGGATCAAGCACAACATTTTCAATCCCCCTATCATTG
 5 CTAGATACATCAGGCTGCACCCAACCCATTATTCTATTTCGAAGTACACTGCGGATGGAAGTATGGGG
 TCGATCTGAACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTGACGCCCAGATTAC
 CGCTAGCTCCTACTTCACTAATATGTTTGCTACCTGGAGCCCCCCTCAAAGCACGACTGCATCTGCAGG
 GACGAAGCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTCGATTTTCAGAAA
 ACTATGAAGGTGACCGGAGTCACAACCTCAGGGCGTGAAAAGTCTGCTGACCTCAATGTACGTCAAGGA
 10 GTTCCTGATCTCTAGTTCACAGGACGGCCACAGTGGACACTGTTCTTTTCAGAACGGAAAGGTGAAAG
 TCTTCCAGGGCAATCAGGATTCTTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACTCGC
 TACCTGCGAATCCACCCACAGTCCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTGGGCTGCGA
 GGCCAGGACCTGTATTGA

Codon-optimized factor VIII cDNA, aka CO/CO3, with deletion of amino acids 1645 and 1646 variant, $\Delta 2$ (SEQ ID NO:5)

15 ATGCAGATTGAGCTGTCAACTTGCTTTTTTCTGTGCTGCTGAGATTTTGTTTTTCCGCTACTAGAAG
 ATACTACCTGGGGGCTGTGGAACGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 ACGCACGATTTCCACCTAGAGTCCCTAAATCATTCCCCTTCAACACCAGCGTGGTCTATAAGAAAACA
 CTGTTTCGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGATGGGACTGCT
 20 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCATTACACTGAAAAACATGGCCTCACACC
 CCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCCGAAGGGGCAGAGTATGACGATCAG
 ACTTCCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTGGCGGGTCTCATACCTATGTGTGGCAGGT
 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCTCTGTGCTGACCTACTCTTATCTGAGTCACGTGG
 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGAAGGGAGCCTGGCT
 25 AAGGAGAAAAACCCAGACACTGCATAAGTTTATTCTGCTGTTCCCGTGTGTTGACGAAGGAAAAATCATG
 GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCTTGGCCCCAAA
 TGCACACTGTGAACGGCTACGTCAATCGTCTACCTGCGCTGGACTGATCGGCTGCCACCGAAAGAGCGTG
 TATTGGCATGTATCGGAATGGGCACCACACCTGAAGTGCACCTCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCCTCCCTGGAGATCTCTCCAATTACCTTCTGACAGCTCAGACTC
 30 TGCTGATGGATCTGGGACAGTTCTGTGTTTTGCCACATCAGCTCCCACCAGCATGATGGCATGGAG
 GCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTTCGATGACGATAACAGCCCCCT
 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCACCTGGTGTGGCACCAGACGATCGATCTACAAATCTCAGTATCT
 35 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAAGTTTCATGGCTTATACCGATGAAA
 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 GGGGACACCCTGCTGATCATTTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 TACAGATGTGCGCCCTCTGTACAGCCGGAGACTGCCAAAGGGCGTCAAAACCTGAAGGACTTCCCAA
 TCCTGCCCGGGGAAATTTTAAAGTATAAATGGACTGTACCGTTCGAGGATGGCCCCACTAAGAGCGAC
 40 CCTAGGTGCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAGGGATCTGGCCAGCGGACTGAT
 CGGCCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA
 GGAATGTGATTCTGTTCAGTGTCTTTGACGAAAACCGTTCATGGTATCTGACCGAGAACATCCAGAGA
 TTCTGCGCTAATCCAGCCGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 TATTAATGGCTACGTGTTGACAGTCTGCAGCTGTCACTGTGTCTGCACGAGGTCGCTTACTGGTATA
 45 TCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAA
 ATGGTGTATGAGGACACACTGACTCTGTTTCCCCTTCAGCGCGCAAACCGTGTATGTCCATGGAGAA
 TCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTCAAGGAATCGCGGGATGACTGCCCTGCTGA
 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTCATACGAGGACATCAGCGCATAT
 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCTTTTAGTCAGAATCCTCCAGTGCTGAAGCAGCG
 50 CGAGATCACCCGCACTACCTGCAGAGTGATCAGGAAGAGATCGACTACGACGATACAATTTCTGTGG
 AAATGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCCTCGATCATTCCAGAAG
 AAAACCCGGCATTACTTTATTGCTGCAGTGGAGCGCTGTGGGATTATGGCATGTCTCTAGTCTCTCA
 CGTGCTGCGAAATCGGGCCAGTCAGGGAGCGTCCACAGTTCAAGAAAGTGGTCTTCCAGGAGTTTA

CAGACGGATCCTTTACTCAGCCACTGTACCGGGGCGAACTGAACGAGCACCTGGGGCTGCTGGGACCC
 TATATCAGAGCTGAAGTGGAGGATAACATTATGGTCACCTTCAGAAATCAGGCATCTAGGCCTTACAG
 TTTTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGGAGCAGAACCACGAAAAAACTTCG
 TGAAGCCTAATGAGACCAAAACATACTTTTGGAAAGGTGCAGCACCATATGGCCCCAACAAAAGACGAA
 5 TTCGATTGCAAGGCATGGGCCTATTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGGCCTGAT
 CGGGCCACTGCTGGTGTGTCATACTAACACCCTGAATCCCGCACACGGCAGGCAGGTCACTGTCCAGG
 AATTCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAGCTGGTACTTCACCGAAAAACATGGAGCGA
 AATTGCCGGGCTCCATGTAATATTTCAGATGGAAGACCCACATTCAAGGAGAACTACCGCTTTTCATGC
 CATCAATGGGTATATTATGGATACTCTGCCCCGACTGGTCATGGCTCAGGACCAGAGAATCAGGTGGT
 10 ACCTGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTTCATTCAGCGGACACGTGTTTACTGTC
 CGGAAGAAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGCGCTGTTTCGAAACCGTCGAGAT
 GCTGCCCTAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCTGATTGGGGAGCACCTGCATGCCGGAATGT
 CTACCCTGTTTCTGGTGTACAGTAATAAGTGTGCAGACACCCCTGGGGATGGCTTCCGGACATATCCGG
 GATTTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCATATTTC
 15 CGGGTCTATCAACGCTTGGTCCACAAAAGAGCCTTTCTCTTGGATTAAAGGTGGACCTGCTGGCACC
 TGATCATTTCATGGCATCAAACTCAGGGGGCCAGGCAGAAAGTTCTCCTCTCTGTACATCTCACAGTTT
 ATCATCATGTACAGCCTGGATGGCAAGAAATGGCAGACATAACCGCGGCAATAGCACAGGGACTCTGAT
 GGTGTTCTTTGGCAACGTGGACAGTTTCAGGGATCAAGCACAAACATTTTCAATCCCCCTATCATTGCTA
 GATACATCAGGCTGCACCCAACCCATTATTCTATTTCGAAGTACACTGCGGATGGAAGTATGGGGTGC
 20 GATCTGAACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTGACGCCAGATTACCGC
 TAGCTCCTACTTCACATAATATGTTTGCTACCTGGAGCCCTCCAAAGCACGACTGCATCTGCAGGGAC
 GAAGCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTGATTTTCAGAAAACCT
 ATGAAGGTGACCGGAGTCACAACCTCAGGGCGTGAAAAGTCTGCTGACCTCAATGTACGTCAAGGAGTT
 CCTGATCTCTAGTTCACAGGACGGCCACCAGTGGACACTGTTCTTTTCAGAACGGAAAGGTGAAAGTCT
 25 TCCAGGGCAATCAGGATTCTTTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACTCGCTAC
 CTGCGAATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTCTGGGCTGCGAGGC
 CCAGGACCTGTATTGA

Codon-optimized factor VIII cDNA, aka CO/CO3, with deletion of amino acids 1645-1647 variant, Δ3 (SEQ ID NO:6)

ATGCAGATTGAGCTGTCAACTTGCTTTTTTCTGTGCCTGCTGAGATTTTGTTTTTCCGCTACTAGAAG
 AACTACCTGGGGGCTGTGGAACGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 ACGCACGATTTCCACCTAGAGTCCCTAAATCATTCCCCTTCAACACCAGCGTGGTCTATAAGAAAAACA
 CTGTTCTGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGATGGGACTGCT
 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCATTACACTGAAAAACATGGCCTCACACC
 35 CCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCCGAAGGGGCAGAGTATGACGATCAG
 ACTTCCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTTGGCGGGTCTCATACCTATGTGTGGCAGGT
 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCTCTGTGCCTGACCTACTCTTATCTGAGTCACGTGG
 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGAAGGGAGCCTGGCT
 AAGGAGAAAACCCAGACACTGCATAAGTTCATTCTGCTGTTCCCGTGTTTGACGAAGGAAAATCATG
 40 GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGAATGCCGCTTCAGCCAGAGCTTGGCCAAAA
 TGCACACTGTGAACGGCTACGTCAATCGCTCACTGCTGGACTGATCGGCTGCCACCGAAAGAGCGTG
 TATTGGCATGTCTCGGAATGGGCACCACACCTGGAAGTGCATCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCCTCCCTGGAGATCTCTCCAATTACCTTCCCTGACAGCTCAGACTC
 TGCTGATGGATCTGGGACAGTTCTGTGTTTTGCCACATCAGCTCCCAACAGCATGATGGCATGGAG
 45 GCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTTCGATGACGATAACAGCCCT
 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCACTGGTGTGGCACCAGACGATCGATCCTACAAATCTCAGTATCT
 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAAGTTTCATGGCTTATACCGATGAAA
 50 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 GGGGACACCCTGCTGATCATTTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 TACAGATGTGCGCCCTCTGTACAGCCGGAGACTGCCAAAGGGCGTCAAAACCTGAAGGACTTCCCAA
 TCCTGCCCGGGGAAATTTTTAAGTATAAATGGACTGTCAACCGTCGAGGATGGCCCCACTAAGAGCGAC

CCTAGGTGCCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAAAGGGATCTGGCCAGCGGACTGAT
 CGGCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA
 GGAATGTGATTCTGTTCAAGTGTCTTTGACGAAAACCGGTCATGGTATCTGACCGAGAACATCCAGAGA
 TTCTGCGCTAATCCAGCCGGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 5 TATTAATGGCTACGTGTTTCGACAGTCTGCAGCTGTCAAGTGTCTGACGAGGTCGCTTACTGGTATA
 TCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAA
 ATGGTGTATGAGGACACACTGACTCTGTTCCCTTCAGCGGCGAAACCGTGTATGTCCATGGAGAA
 TCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTTCAGGAATCGCGGGATGACTGCCCTGCTGA
 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTCATACGAGGACATCAGCGCATAT
 10 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCCTTTAGTCAGAATCCTCCAGTGTGTAAGCGCGA
 GATCACCCGCACTACCTGTCAGAGTGTACAGGAAGAGATCGACTACGACGATACAATTTCTGTGGAAA
 TGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCTCTGATCATTCCAGAAGAAA
 ACCCGGCATTACTTTATGCTGCAGTGGAGCGCCTGTGGGATTATGGCATGTCTCTAGTCTCTACGT
 GCTCGGAAATCGGGCCAGTCAGGGAGCGTCCACAGTTCAGAAAGTGGTCTTCCAGGAGTTTACAG
 15 ACGGATCCTTTACTCAGCCACTGTACCGGGCGAAGTGAACGAGCACCTGGGGCTGTCTGGGACCTAT
 ATCAGAGCTGAAGTGGAGGATAACATTATGGTCACCTTCAGAAATCAGGCATCTAGGCCTTACAGTTT
 TTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGAGCAGAAACCACGAAAAAACTTCGTGA
 AGCCTAATGAGACCAAAACATACTTTTGAAGGTGCAGCACCATATGGCCCAACAAAAGACGAATTC
 GATTGCAAGGCATGGGCTATTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGGCCTGATCGG
 20 GCCACTGCTGGTGTGTCTATACTAACACCTGAATCCCGCACACGGCAGGCAGGTCAGTGTCCAGGAAT
 TCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAGCTGGTACTTCACCGAAAACATGGAGCGAAAT
 TGCCGGGCTCCATGTAATATTCAGATGGAAGACCCACATTCAGGAGAACTACCGCTTTCATGCCAT
 CAATGGGTATATTATGGATACTCTGCCCGGACTGGTCATGGCTCAGGACCAGAGAATCAGGTGGTACC
 TGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTCATTTACGCGGACACGTGTTTACTGTCCGG
 25 AAGAAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGGCGTGTTCGAAACCGTCGAGATGCT
 GCCTAGCAAGGCAGGGATCTGGAGAGTGGATGCCGTATGGGGAGCACCTGCATGCCGGAATGTCTA
 CCCTGTTTCTGGTGTACAGTAATAAGTGTGACACACCCCTGGGGATGGCTTCCGGACATATCCGGGAT
 TTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCACTATTCCGG
 GTCTATCAACGCTTGGTCCACAAAAGACCTTTCTCTTGGATTAAGGTGGACCTGCTGGCACCAATGA
 30 TCATTTCATGGCATCAAAACTCAGGGGGCCAGGCAGAAAGTTCTCTCTGTACATCTCACAGTTTATC
 ATCATGTACAGCCTGGATGGCAAGAAATGGCAGACATACC GCGGCAATAGCACAGGGACTCTGATGGT
 GTTCTTTGGCAACGTGGACAGTTTCAGGGATCAAGCACAAACATTTTCAATCCCCCTATCATTGCTAGAT
 ACATCAGGCTGCACCAACCCATTATTCTATTTCGAAGTACACTGCGGATGGAACCTGATGGGGTGCAT
 CTGAACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTGACGCCCAGATTACCGCTAG
 35 CTCTACTTCACTAATATGTTTGCTACCTGGAGCCCCCCTCAAAGCACGACTGCATCTGCAGGGACGAA
 GCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTGATTTTCAGAAAACATATG
 AAGGTGACCGGAGTCACAACCTCAGGGCGTGAAGAGTCTGCTGACCTCAATGTACGTCAAGGAGTTCCT
 GATCTCTAGTTCACAGGACGGCCACAGTGGACACTGTTCTTTCAGAACGGAAGGTGAAAGTCTTCC
 AGGGCAATCAGGATTCCCTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACTCGCTACCTG
 40 CGAATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTGGGCTGCGAGGCCCA
 GGACCTGTATTGA

**Codon-optimized FVIII cDNA, aka CO/CO3, with deletion of amino acid 1648 variant,
 Δ1648 (SEQ ID NO:7)**

ATGCAGATTGAGCTGTCAACTTGCTTTTTTCTGTGCTGCTGAGATTTTGTTTTTTCCGCTACTAGAAG
 45 ATACTACCTGGGGGCTGTGGAAGTGTCTTGGGATTACATGCAGAGTGACCTGGGAGAGCTGCCAGTGG
 ACGCACGATTTCCACCTAGAGTCCCTAAATCATTCCCCCTCAACACCAGCGTGGTCTATAAGAAAAACA
 CTGTTCTGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGATGGGACTGCT
 GGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCAATTACACTGAAAAACATGGCCTCACACC
 CCGTGAGCCTGCATGCTGTGGGCGTCAGTACTGGAAGGCTTCCGAAGGGGCGAGATATGACGATCAG
 50 ACTTCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTGGCGGGTCTCATACCTATGTGTGGCAGGT
 CCTGAAAGAGAATGGCCCCATGGCTTCCGACCCCTGTGCGCTGACCTACTCTTATCTGAGTCACGTGG
 ACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGGAAGGGAGCCTGGCT
 AAGGAGAAAACCCAGACACTGCATAAGTTTCTGCTGTTTCGCCGTGTTTGACGAAGGAAAAATCATG

GCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCTTGGCCCCAAAA
 TGCACACTGTGAACGGCTACGTCAATCGCTCACTGGCTGATCGGCTGCCACCGAAAGAGCGTG
 TATTGGCATGTATCGGAATGGGCACCACACCTGAAGTGCATCCATTTTCTGGAGGGGCATACCTT
 TCTGGTCCGCAACCACCGACAGGCCTCCCTGGAGATCTCTCCAATTACCTTCTGACAGCTCAGACTC
 5 TGCTGATGGATCTGGGACAGTTCCTGCTGTTTTGCCACATCAGCTCCCACCAGCATGATGGCATGGAG
 GCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATGAGGAAGCTGA
 AGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTTCGATGACGATAACAGCCCCCT
 CCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCTAAGACATGGGTCCATTACATCGCAGCCGAG
 GAAGAGGACTGGGATTATGCACCACTGGTGTGGCACCAGACGATCGATCCTACAAATCTCAGTATCT
 10 GAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAGGTTTCATGGCTTATACCGATGAAA
 CCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTACGGAGAAGTG
 GGGGACACCCCTGCTGATCATTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATCCACATGGAAT
 TACAGATGTGCGCCCTCTGTACAGCCGGAGACTGCCAAAGGGCGTCAAACACCTGAAGGACTTCCCAA
 TCCTGCCCGGGGAAATTTTTAAGTATAAATGGACTGTCACCGTCGAGGATGGCCCCCACTAAGAGCGAC
 15 CCTAGGTGCCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAAGGGATCTGGCCAGCGGACTGAT
 CGGCCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATGTCCGACAAGA
 GGAATGTGATTCTGTTCAGTGTCTTTGACGAAAACCGGTCATGGTATCTGACCGAGAACATCCAGAGA
 TTCTTGCTAATCCAGCCGGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAACATCATGCATAG
 TATTAATGGCTACGTGTTCGACAGTCTGCAGCTGTCAGTGTGTCTGCACGAGGTCGCTTACTGGTATA
 20 TCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTTTAAGCATAAA
 ATGGTGTATGAGGACACACTGACTCTGTTCCCTTCAGCGGCAGAACCGTGTATATGTCCATGGAGAA
 TCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTCAAGGAATCGCGGGATGACTGCCCTGCTGA
 AAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTATACGAGGACATCAGCGCATAT
 CTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCCTTTAGTCAGAATCCTCCAGTGCTGAAGAGGCA
 25 CCAGGAGATCACCCGCACTACCTTGCAGAGTGATCAGGAAGAGATCGACTACGACGATACAATTTCTG
 TGGAAATGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCCTCGATCATTCCAG
 AAGAAAACCCGGCATTACTTTATGCTGCAGTGGAGCGCCTGTGGGATTATGGCATGTCTCTAGTCC
 TCACGTGCTGCGAAATCGGGCCAGTCAGGGAGCGTCCACAGTTCAGAAAGTGGTCTTCCAGGAGT
 TTACAGAGTGCCTTTACTCAGCCACTGTACCGGGGCGAACTGAACGAGCACCTGGGGCTGCTGGGA
 30 CCCTATATCAGAGCTGAAGTGGAGGATAACATTATGTTGGTCACTTCAGAAATCAGGCATCTAGGCCTTA
 CAGTTTTTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGGAGCAGAACCACGAAAAAACT
 TCGTGAAGCCTAATGAGACCAAAACATACTTTTGGAAAGTGCAGCACCATATGGCCCCAACAAAAGAC
 GAATTGATTGCAAGGCATGGGCCTATTTTTCTGACGTGGATCTGGAGAAGGACGTCCACAGTGGCCT
 GATCGGGCCACTGCTGGTGTGTCTATACTAACACCTGAATCCCGCACACGGCAGGCAGGTCACTGTCC
 35 AGGAATTCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAAGCTGGTACTTCACCGAAAACATGGAG
 CGAAATTGCCGGCTCCATGTAATATTAGATGGAAGACCCACATTCAAGGAGAACTACCGCTTTCA
 TGCCATCAATGGGTATATTATGGATACTCTGCCCGGACTGGTCATGGCTCAGGACCAGAGAATCAGGT
 GGTACCTGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTCAATTCAGCGGACACGTGTTTACT
 GTCCGGAAGAAAGAAGATATAAAATGGCCCTGTACAACCTGTATCCCGGCGTGTTCGAAAACCGTCGA
 40 GATGCTGCCTAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCGTATTGGGGAGCACCTGCATGCCGGAA
 TGTCTACCTGTTTCTGGTGTACAGTAATAAGTGTGACACACCCCTGGGGATGGCTTCCGGACATATC
 CGGGATTTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCACTA
 TTCCGGGTCTATCAACGCTTGGTCCACAAAAGAGCCTTTCTCTTGGATTAAGGTGGACCTGCTGGCAC
 CAATGATCATTATGGCATCAAACTCAGGGGGCCAGGCAGAACTTCTCCTCTGTACATCTCACAG
 45 TTTATCATCATGTACAGCCTGGATGGCAAGAAATGGCAGACATACCGCGGCAATAGCACAGGGACTCT
 GATGGTGTCTTTGGCAACGTGGACAGTTTCAAGGATCAAGCACAACATTTTCAATCCCCCTATCATTG
 CTAGATACATCAGGCTGCACCAACCCATTATTCTATTTCGAAGTACACTGCGGATGGAACCTGATGGGG
 TGCGATCTGAACAGTTGTTCAATGCCCCGGAATGGAGTCCAAGGCAATCTCTGACGCCAGATTAC
 CGCTAGCTCCTACTTACTAATATGTTTGTACCTGGAGCCCCCTCAAAGCACGACTGCATCTGCAGG
 50 GACGAAGCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGTCGATTTTCAGAAA
 ACTATGAAGGTGACCGGAGTCACAACCTCAGGGCGTGAAGAGTCTGCTGACCTCAATGTACGTCAAGGA
 GTTCCTGATCTCTAGTTTACAGGACGGCCACCAGTGGACACTGTCTTTTCAAGACGAAAGGTGAAAG
 TCTTCCAGGGCAATCAGGATTCTTTTACACCTGTGGTCAACTCTCTGGACCCACCCCTGCTGACTCGC
 TACCTGCGAATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAGTCTGGGCTGCGA
 55 GGCCAGGACCTGTATTGA

TTR promoter with 4 nucleotide mutation (TTRm), SEQ ID NO:8

GTCTGTCTGCACATTTTCGTAGAGCGAGTGTTCCGATACTCTAATCTCCCTAGGCAAGGTTTCATATTGA
 CTAGGTTACTTATTCTCCTTTTGTGACTAAGTCAATAATCAGAATCAGCAGGTTTGGAGTCAGCTT
 GGCAGGGATCAGCAGCCTGGGTTGGAAGGAGGGGGTATAAAAGCCCTTCACCAGGAGAAGCCGTCAC
 ACAGATCCACAAGCTCCT

[0231] The DNA sequence of the entire full-length factor VIII construct including the TTR promoter with the four nucleotide mutation, synthetic intron, codon-optimized factor VIIIcDNA and polyadenylation (Poly A) signal sequence are provided (SEQ ID NO:9). Also highlighted in SEQ ID NO: 9 are the codons coding for the 4 amino acid PACE/furin cleavage site, wherein 1, 2, 3 or all 4 amino acids are optionally deleted.

Full length construct including TTR promoter with 4 nucleotide mutation (TTRm), synthetic intron, codon-optimized factor VIII cDNA (PACE/furin underlined), poly A (SEQ ID NO:9)

GTCTGTCTGCACATTTTCGTAGAGCGAGTGTTCCGATACTCTAATCTCCCTAGGCAAGGTTTCATATTga
 ctTAGGTTACTTATTCTCCTTTTGTGACTAAGTCAATAATCAGAATCAGCAGGTTTGGAGTCAGCTT
 GGCAGGGATCAGCAGCCTGGGTTGGAAGGAGGGGGTATAAAAGCCCTTCACCAGGAGAAGCCGTCAC
 ACAGATCCACAAGCTCCTGCTAGCAGGTAAGTGCCGTGTGTGGTTCCCGCGGGCCTGGCCTCTTTACG
 GGTTATGGCCCTTGCGTGCCCTTGAATTACTGACACTGACATCCACTTTTTCTTTTCTCCACAGGTTT
 AAACGCCACCATGCAGATTGAGCTGTCAACTTGCTTTTTCTGTGCTGCTGAGATTTTGTTTTCCG
 CTACTAGAAGATACTACCTGGGGGCTGTGGAAGTGTCTTGGGATTACATGCAGAGTGACCTGGGAGAG
 CTGCCAGTGGACGCACGATTTCCACCTAGAGTCCCTAAATCATTCCTTCAACACCAGCGTGGTCTA
 TAAGAAAACACTGTTTCGTGGAGTTTACTGATCACCTGTTCAACATCGCTAAGCCTCGGCCACCCTGGA
 TGGGACTGCTGGGACCAACAATCCAGGCAGAGGTGTACGACACCGTGGTCATTACACTGAAAAACATG
 GCCTCACACCCCGTGAGCCTGCATGCTGTGGGCGTCAGCTACTGGAAGGCTTCCGAAGGGGCAGAGTA
 TGACGATCAGACTTCCCAGAGAGAAAAAGAGGACGATAAGGTGTTTCTGGCGGGTCTCATACCTATG
 TGTGGCAGGTCTGAAAGAGAATGGCCCCATGGCTTCCGACCTCTGTGCTGACCTACTCTTATCTG
 AGTCACGTGGACCTGGTCAAGGATCTGAACAGCGGACTGATCGGAGCACTGCTGGTGTGTAGGGAAGG
 GAGCCTGGCTAAGGAGAAAACCCAGACACTGCATAAGTTCATTCTGCTGTTCGCCGTGTTTGACGAAG
 GAAAATCATGGCACAGCGAGACAAAGAATAGTCTGATGCAGGACCGGGATGCCGCTTCAGCCAGAGCT
 TGGCCCAAAATGCACACTGTGAACGGCTACGTCAATCGCTCACTGCCTGGACTGATCGGCTGCCACCG
 AAAGAGCGTGTATTGGCATGTATCGGAATGGGCACCACACCTGAAGTGCATCCATTTTCTGGAGG
 GGCATACCTTTCTGGTCCGCAACCACCGACAGGCCCTCCCTGGAGATCTCTCCAATTACCTTCTTGACA
 GCTCAGACTCTGCTGATGGATCTGGGACAGTTTCTGCTGTTTTGCCACATCAGCTCCCACCAGCATGA
 TGGCATGGAGGCCTACGTGAAAGTGGACAGCTGTCCCGAGGAACCTCAGCTGAGGATGAAGAACAATG
 AGGAAGCTGAAGACTATGACGATGACCTGACCGACTCCGAGATGGATGTGGTCCGATTGATGACGAT
 AACAGCCCCCTCCTTTATCCAGATTAGATCTGTGGCCAAGAAACACCCTAAGACATGGGTCCATTACAT
 CGCAGCCGAGGAAGAGGACTGGGATTATGCACCACTGGTGTGGCACCAGACGATCGATCCTACAAAT
 CTCAGTATCTGAACAATGGACCACAGCGGATTGGCAGAAAGTACAAGAAAGTGAGGTTTCATGGCTTAT
 ACCGATGAAACCTTCAAGACTCGCGAAGCAATCCAGCACGAGAGCGGGATTCTGGGACCACTGCTGTA
 CGGAGAAGTGGGGGACACCCTGCTGATCATTTTTAAGAACCAGGCCAGCAGGCCTTACAATATCTATC
 CACATGGAATTACAGATGTGCGCCCTCTGTACAGCCGAGACTGCCAAAGGGCGTCAAACACCTGAAG
 GACTTCCCAATCCTGCCCGGGGAAATTTTTAAGTATAAATGGACTGTCACCGTCGAGGATGGCCCCAC
 TAAGAGCGACCCTAGGTGCCTGACCCGCTACTATTCTAGTTTCGTGAATATGGAAGGGATCTGGCCA
 GCGGACTGATCGGCCCACTGCTGATTTGTTACAAAGAGAGCGTGGATCAGAGAGGCAACCAGATCATG
 TCCGACAAGAGGAATGTGATTCTGTTCAAGTGTCTTTGACGAAAACCGGTCATGGTATCTGACCGAGAA
 CATCCAGAGATTCTGCCTAATCCAGCCGAGTGCAGCTGGAAGATCCTGAGTTTCAGGCTTCTAAACA
 TCATGCATAGTATTAATGGCTACGTGTTTCGACAGTCTGCAGCTGTCAAGTGTGTCTGCACGAGGTGCGT
 TACTGGTATATCCTGAGCATTGGAGCACAGACAGATTTCTGAGCGTGTCTTTTCCGGCTACACTTT
 TAAGCATAAAATGGTGTATGAGGACACACTGACTCTGTTCCCTTCAGCGGCGAAACCGTGTATTATGT

CCATGGAGAATCCCGGGCTGTGGATCCTGGGATGCCACAACAGCGATTTTCAGGAATCGCGGGATGACT
 GCCCTGCTGAAAGTGTCAAGCTGTGACAAGAACACCGGAGACTACTATGAAGATTCATACGAGGACAT
 CAGCGCATATCTGCTGTCCAAAAACAATGCCATTGAACCCAGGTCTTTTAGTCAGAATCCTCCAGTGC
 TGAAG**AGGCACCAGCGC**GAGATCACCCGCACTACCCTGCAGAGTGATCAGGAAGAGATCGACTACGAC
 5 GATACAATTTCTGTGGAAATGAAGAAAGAGGACTTCGATATCTATGACGAAGATGAGAACCAGAGTCC
 TCGATCATTCCAGAAGAAAACCCGGCATTACTTTATTGCTGCAGTGGAGCGCCTGTGGGATTATGGCA
 TGTCTCTAGTCTCTACGTGCTGCGAAATCGGGCCCCAGTCAGGGAGCGTCCACAGTTCAAGAAAAGTG
 GTCTTCCAGGAGTTTACAGACGGATCCTTTACTCAGCCACTGTACCGGGGCGAACTGAACGAGCACCT
 GGGGCTGCTGGGACCCTATATCAGAGCTGAAGTGGAGGATAACATTATGGTCACTTTCAGAAATCAGG
 10 CATCTAGGCCTTACAGTTTTTATTCAAGCCTGATCTCTTACGAAGAGGACCAGAGGCAGGGAGCAGAA
 CCACGAAAAAACTTCGTGAAGCCTAATGAGACCAAAAACATACTTTTGGAAGGTGCAGCACCATATGGC
 CCCAACAAAAGACGAATTCGATTGCAAGGCATGGGCCATTTTTCTGACGTGGATCTGGAGAAGGACG
 TCCACAGTGGCCTGATCGGGCCACTGCTGGTGTGTCTACTAACAACCTGAATCCCGCACACGGCAGG
 CAGGTCACTGTCCAGGAATTCGCCCTGTTCTTTACCATCTTTGATGAGACAAAAAGCTGGTACTTCAC
 15 CGAAAACATGGAGCGAAATTGCCGGGCTCCATGTAATATTCAGATGGAAGACCCACATTCAAGGAGA
 ACTACCGCTTTTCATGCCATCAATGGGTATATTATGGATACTCTGCCCGGACTGGTCATGGCTCAGGAC
 CAGAGAATCAGGTGGTACCTGCTGAGCATGGGGTCCAACGAGAATATCCACTCAATTCATTTCAGCGG
 ACACGTGTTTACTGTCCGGAAGAAAGAAGAGTATAAAATGGCCCTGTACAACCTGTATCCCGGCGTGT
 TCGAAACCGTCGAGATGCTGCCTAGCAAGGCAGGGATCTGGAGAGTGGAAATGCCTGATTGGGGAGCAC
 20 CTGTCATGCCGGAATGTCTACCCTGTTTCTGGTGTACAGTAATAAGTGTGCAGACACCCCTGGGGATGGC
 TTCCGGACATATCCGGGATTTCCAGATTACCGCATCTGGACAGTACGGCCAGTGGGCCCTTAAGCTGG
 CTAGACTGCATATTCCGGGTCTATCAACGCTTGGTCCACAAAAGAGCCTTTCTCTTGGATTAAGGTG
 GACCTGCTGGCACCAATGATCATTTCATGGCATCAAACTCAGGGGGCCAGGCAGAAAGTTCTCTCTCT
 GTACATCTCACAGTTTATCATCATGTACAGCCTGGATGGCAAGAAATGGCAGACATACCGCGGCAATA
 25 GCACAGGGACTCTGATGGTGTCTTTGGCAACGTGGACAGTTCAGGGATCAAGCACAACTTTTCAAT
 CCCCCTATCATTGCTAGATACATCAGGCTGCACCAACCCATTATTCTATTCGAAGTACACTGCGGAT
 GGAACCTGATGGGGTCCGATCTGAACAGTTGTTCAATGCCCTGGGAATGGAGTCCAAGGCAATCTCTG
 ACGCCACAGATTACCGCTAGCTCCTACTTCACTAATATGTTTGCTACCTGGAGCCCCCTCAAAGCAGCA
 CTGCATCTGCAGGAGCAAGCAACGCATGGCGACCACAGGTGAACAATCCCAAGGAGTGGCTGCAGGT
 30 CGATTTTCAGAAAACATGAAGGTGACCGGAGTACCAACTCAGGGCGTGAAAAGTCTGCTGACCTCAA
 TGTACGTCAAGGAGTTCTGATCTCTAGTTACAGGACGGCCACCAGTGGACACTGTTCTTTTCAGAAC
 GGAAAGGTGAAAGTCTTCCAGGGCAATCAGGATTCCCTTTACACCTGTGGTCAACTCTCTGGACCCACC
 CCTGCTGACTCGCTACCTGCGAATCCACCCACAGTCTGGGTGCATCAGATTGCACTGAGAATGGAAG
 TCCTGGGCTGCGAGGCCAGGACCTGTATTGAGCGGCCGCAATAAAAAGATCAGAGCTCTAGAGATCTG
 35 TGTGTTGGTTTTTTGTGT

Codon-optimized factor VIII cDNA, aka CO1 (SEQ ID NO:10)

ATGCAGATCGAGCTGTCTACCTGCTTCTTCCCTGTGCTGCTGCGGTTCTGCTTCAGCGCCACCAGACG
 GTACTATCTGGGCGCCGTGGAACCTGAGCTGGGACTACATGCAGAGCGACCTGGGCGAGCTGCCCGTGG
 ATGCCAGATTCCCTCCAAGAGTGCCCAAGAGCTTCCCTTCAACACCTCCGTGGTGTACAAGAAAACC
 40 CTGTTCTGTGGAATTCACCGACCACCTGTTCAATATCGCCAAGCCCAGACCCCCCTGGATGGGCCTGCT
 GGGACCTACAATTCAGGCCGAGGTGTACGACACCGTCGTGATCACCTGAAGAACATGGCCAGCCACC
 CCGTGTCTCTGTCATGCCGTGGGAGTGTCTTACTGGAAGGCCCTGAGGGCGCCGAGTACGACGATCAG
 ACCAGCCAGCGCGAGAAAGAGGACGACAAGGTGTTCCCTGGCGGCAGCCACACCTACGTGTGGCAGGT
 GCTGAAAGAAAACGGCCCCATGGCCTCCGACCTCTGTGCTGACATACAGCTACCTGAGCCACGTGG
 45 ACCTCGTGAAGGACCTGAACAGCGGCCTGATCGGAGCCCTGCTCGTGTGTAGAGAGGGCAGCCTGGCC
 AAAGAGAAAACCCAGACCCTGCACAAGTTCATCCTGCTGTTTCGCCGTGTTTCGACGAGGGCAAGAGCTG
 GCACAGCGAGACAAAGAACAGCCTGATGCAGGACCGGGACGCCGCTCTGCTAGAGCCTGGCCAAAAA
 TGCACACCGTGAACGGCTACGTGAACAGAAGCCTGCCGGGACTGATCGGCTGCCACCGGAAGTCTGTG
 TACTGGCACGTGATCGGCATGGGCACCACCCCTGAGGTGCACAGCATCTTTCTGGAAGGACACACCTT
 50 TCTCGTGCAGAACACCGGCAGGCCAGCTGGAAATCAGCCCTATCACCTTCTGACCGCCAGACAC
 TGCTGATGGACCTGGGCCAGTTTCTGCTGTTCTGCCACATCAGCTCCCAACAGCACGACGGCATGGAA
 GCCTACGTGAAGGTGGACAGCTGCCCCGAGGAACCCAGCTGCGGATGAAGAACAACGAGGAAGCCGA
 GGACTACGACGACGACCTGACCGACAGCGAGATGGACGTGGTGCCTTCGACGACGATAACAGCCCCA

GCTTCATCCAGATCAGAAGCGTGGCCAAGAAGCACCCCAAGACCTGGGTGCACTATATCGCCGCCGAG
 GAAGAGGACTGGGATTACGCCCCCTCTGGTGTGGCCCCGACGACAGAAAGCTACAAGAGCCAGTACCT
 GAACAATGGCCCCCAGCGGATCGGCCGGAAGTATAAGAAAAGTGCGGTTCATGGCCTACACCGACGAGA
 CATTCAAGACCAGAGAGGCCATCCAGCACGAGAGCGGCATCCTGGGCCCTCTGCTGTATGGCGAAGTG
 5 GGCAGACACCCTGCTGATCATCTTCAAGAACAGGCCAGCAGACCCTACAACATCTACCCCTACGGCAT
 CACCGACGTGCGGCCCTGTACTCTAGAAGGCTGCCAAGGGCGTGAAACACCTGAAGGACTTCCCCA
 TCCTGCCCCGCGAGATCTTCAAGTACAAGTGGACCGTGACCGTGGAAGATGGCCCCACCAAGAGCGAC
 CCCAGATGCCTGACACGGTACTACAGCAGCTTCGTGAACATGGAACGGGACCTGGCCTCCGGCCTGAT
 TGGCCCCACTGCTGATCTGCTACAAAGAAAGCGTGGACCAGCGGGGCAACCAGATCATGAGCGACAAGC
 10 GGAACGTGATCCTGTTTAGCGTGTTCGATGAGAACCGGTCTCTGGTATCTGACCGAGAATATCCAGCGG
 TTCTTGCCCAACCCTGCCGGCGTGCAGCTGGAAGATCCTGAGTTCCAGGCCCTCAACATCATGCACTC
 CATCAATGGCTATGTGTTCGACAGCCTGCAGCTGAGCGTGTGCCTGCACGAGGTGGCCTACTGGTACA
 TCCTGAGCATCGGGGCCAGACCGACTTCCTGTCCGTGTTCTTCTCCGGCTACACCTTCAAGCAAAA
 ATGGTGTACGAGGATACCCCTGACCCTGTTCCCTTTAGCGGCGAAACCGTGTTCATGAGCATGGAAAA
 15 CCCCCGCCCTGTGGATCCTGGGCTGCCACAACAGCGACTTCCGGAACAGAGGCATGACCGCCCTGCTGA
 AGGTGTCCAGCTGCGACAAGAACACCGGCGACTACTACGAGGACAGCTATGAGGACATCAGCGCCTAC
 CTGCTGAGCAAGAACAATGCCATCGAGCCCAGAAGCTTCAGCCAGAACCCCCCGTGCTGAAGCGGCA
 CCAGAGAGAGATCACCCGGACCACCCTGCAGTCCGACCAGGAAGAGATCGATTACGACGACACCATCA
 GCGTGGAATGAAGAAAGAAGATTTTCGACATCTACGACGAGGACGAGAACCAGAGCCCCCGGTCTTT
 20 CAGAAAAAGACCCGGCACTACTTCATTGCCGCTGTGGAACGGCTGTGGGACTACGGCATGAGCAGCAG
 CCCTCACGTGCTGAGAAACAGGGCCCAGAGCGGCAGCGTGCCCCAGTTCAAGAAAGTGGTGTTCAGG
 AATTACAGACGGCAGCTTCACCCAGCCTCTGTACCGCGGCGAGCTGAATGAGCACCTGGGACTGCTG
 GGCCCCATATCAGAGCCGAAGTGAAGATAACATCATGGTCACCTTCCGGAATCAGGCCTCCCGGCC
 CTACAGCTTCTACAGCTCCCTGATCAGCTACGAAGAGGACCAGAGACAGGGCGCTGAGCCCCGGAAGA
 25 ACTTCGTGAAGCCCAACGAGACTAAGACCTACTTTTGGAAAGGTGCAGCACCATGGCCCCCTACAAAG
 GACGAGTTCGACTGCAAGGCCCTGGGCCACTTCTCCGATGTGGACCTGGAAAAGGACGTGCACTCTGG
 GCTGATCGGGCCCCCTGCTCGTGTGCCACACCAACACCCTGAATCCCGCCACGGCAGACAAGTGACAG
 TGCAGGAATTCGCCCTGTTCTTACCATCTTCGACGAAACAAAGAGCTGGTACTTCACCGAAAACATG
 GAAAGAACTGCCGGCTCCCTGCAACATCCAGATGGAAGATCCACCTTCAAAGAGAACTACCGGTT
 30 CCACGCCATCAACGGCTACATCATGGACACACTGCCCGGCCCTCGTGATGGCTCAGGATCAGCGGATCC
 GGTGGTATCTGCTGTCCATGGGCTCCAACGAGAACATCCACAGCATCCACTTCAGCGGCCACGTGTTT
 ACCGTGCGGAAAAAAGAAGAGTACAAAATGGCCCTGTACAACCTGTACCCTGGGGTGTTCGAGACAGT
 GGAAATGCTGCCCAGCAAGGCCGGCATCTGGCGGGTGGAAATGTCTGATCGGCGAGCATCTGCACGCTG
 GGATGAGCACACTGTTTCTGGTGTACAGCAACAAGTGCCAGACACCTCTGGGCATGGCCTCTGGCCAC
 35 ATCCGGGACTTTTCAGATCACAGCCAGCGGCCAGTAIGGCCAGTGGGCCCCAAAAGTGGCCAGACTGCA
 CTACAGCGGCAGCATCAACGCCTGGTCCACCAAAGAGCCCTTCAGCTGGATCAAGGTGGACCTGCTGG
 CTCCCATGATCATCCACGGAATCAAGACCCAGGGCGCCAGACAGAAGTTACGACAGCCTGTACATCAGC
 CAGTTCATCATCATGTACAGCCTGGACGGCAAGAAGTGGCAGACCTACCGGGGCAATAGCACCGGCAC
 CCTGATGGTGTCTTCTCGGCAACGTGGACTCCAGCGGCATTAAGCACAAACATCTTCAACCCCCCATCA
 40 TTGCCCGGTACATCCGGCTGCACCCACCCACTACAGCATCCGGTCCACCCTGAGAATGGAAGTATG
 GGCTGCGACCTGAACCTCTGCAGCATGCCCTGGGGATGGAAAGCAAGGCCATCTCCGACGCCAGAT
 CACCGCCTCCAGCTACTTCACCAACATGTTCCGCCACCTGGTCCCCATCCAAGGCCCGGCTGCATCTGC
 AGGGCAGAAGCAATGCTTGGAGGCCCAAGTGAACAACCCCAAAGAATGGCTGCAGGTGGACTTCCAG
 AAAACCATGAAAGTGACCGGCTGACCACCCAGGGCGTGAAGTCTCTGCTGACCTCTATGTACGTGAA
 45 AGAGTTCCTGATCTCCAGCAGCCAGGACCGCCACCAGTGGACCTGTTTTTCCAGAACGGCAAGTGA
 AAGTGTTCAGGGGAACCAGGACTCCTTACCCCCGTCTGTAATAGCCTGGACCTCCACTGCTGACC
 AGATACCTGCGGATCCACCCTCAGAGTTGGGTGCACCAGATTGCTCTGCGGATGGAAGTGTGGGATG
 CGAGGCCAGGACCTGTACTGA

Codon-optimized factor VIII cDNA, aka CO2 (SEQ ID NO:11)

50 ATGCAGATCGAGCTGTCTACCTGCTTCTTCCCTGTGCCTGCTGCGGTTCTGCTTTAGCGCTACTAGACG
 CTACTACCTGGGCGCCGTGGAAGTGAAGTGGGACTATATGCAGTCAGACCTGGGCGAGCTGCCCGTGG
 ACGCTAGATTTCCACCTAGAGTGCCTAAGAGCTTCCCTTTAACACCTCCGTGGTCTATAAGAAAACC
 CTGTTCTGTCGAGTTACCGATCACCTGTTTAATATCGCTAAGCCTAGACCCCCCTGGATGGGCCTGCT

GGGCCCTACTATTTCAGGCCGAGGTCTACGACACCGTCGTGATCACCCCTGAAGAATATGGCTAGTCACC
 CCGTCAGCCTGCACGCCGTGGGCGTCAGCTACTGGAAGGCTAGCGAGGGCGCCGAGTACGACGATCAG
 ACTAGTCAGCGCGAGAAAGAGGACGACAAAGTCTTTCCTGGCGGCTCTCACACCTACGTGTGGCAGGT
 CCTGAAAGAAAACGGCCCTATGGCTAGCGACCCCTGTGCTGACCTATAGCTACCTGAGTCACGTGG
 5 ACCTGGTCAAGGACCTGAATAGCGGCCTGATCGGCCCTGCTCGTGTGTAGAGAGGGCTCACTGGCT
 AAAGAGAAAACCTCAGACCCTGCACAAGTTTATCCTGCTGTTTCGCCGTGTTTCGACGAGGGCAAGAGCTG
 GCACTCAGAGACTAAGAATAGCCTGATGCAGGATAGGGACGCCCTAGCGCTAGAGCCTGGCCTAAGA
 TGCACACCGTGAACGGCTACGTGAACAGATCACTGCCCGGACTGATCGGCTGTCAACGGAAGTCCGTC
 TACTGGCACGTGATCGGAATGGGCACTACCCCCGAGGTGCACTCTATCTTTCTGGAAGGCCACACCTT
 10 CCTCGTCAGAAATCACCGGCAGGCTAGCCTCGAGATTAGCCCTATCACCTTCTTGACCGCTCAGACAC
 TGCTGATGGACCTGGGCCAGTTCTGCTGTTTTGTACATTAGCTCTCACACGACGACGGGATGGAA
 GCCTACGTGAAAGTGGATAGCTGCCCGAGGAACCTCAGCTGAGAATGAAGAACAACGAGGAAGCCGA
 GGATTACGACGACGACCTGACCGATAGCGAGATGGACGTGTCAGATTTCGACGACGATAACTCACCTA
 GCTTTATTTCAGATTAGATCAGTGGCTAAGAAGCACCCCTAAGACCTGGGTGCACTATATCGCCCGCGAG
 15 GAAGAGGACTGGGACTACGCCCTCTGGTGCTGGCCCCGACGATAGAAGCTATAAGTCTCAGTACCT
 GAACAACGGCCCTCAGCGGATCGGCCGGAAGTATAAGAAAAGTGCAGTTTATGGCCTACACCGACGAAA
 CCTTCAAGACTAGAGAGGCTATTTCAGCACGAGTCAGGGATCCTGGGCCCCCTGCTCTACGGCGAAGTG
 GCGGACACCCCTGCTGATTATCTTTAAGAATCAGGCTAGTAGGCCCTATAATATCTACCCCCACGGAAT
 CACCGACGTGCGGCCCTCTACTCTAGACGGCTGCCTAAGGGCGTGAAGCACCTGAAGGACTTCCCTA
 20 TTCTGCCCGGCGAGATCTTTAAGTACAAGTGGACCGTGACCGTCGAGGACGGCCCTACTAAGTCCGAC
 CCTCGGTGCCTGACTAGGTACTACTCTAGCTTCGTGAATATGGAACGGGACCTGGCTAGCGGACTGAT
 TGGCCCTCTGCTGATCTGCTACAAAGAATCAGTGGATCAGCGGGGCAATCAGATTATGAGCGATAAGC
 GGAACGTGATCCTGTTTAGTGTGTTTGACGAGAATAGGTCTCGGTATCTGACCGAGAATATCCAGCGG
 TTCTGCTTAACCCCTGCCGGCGTGACGCTGGAAGATCCCGAGTTTCAGGCTAGCAATATTATGCACTC
 25 TATTAACGGATACGTGTTTCGATAGCCTGCAGCTGAGCGTGTGCCTGCACGAGGTGGCCTACTGGTATA
 TCCTGTCTATCGGCGCTCAGACCGACTTCCTGAGCGTGTTCCTTAGCGGCTACACCTTTAAGCACAAAG
 ATGGTCTACGAGGATACCCTGACCTGTTCCCTTTAGCGGCGAAACCGTGTTTATGTCTATGGAAAA
 CCCC GGCTGTGGATCCTGGGGTGTACAATAGCGACTTTAGGAATAGAGGAATGACCGCCCTGCTGA
 AAGTGTCTAGCTGCGATAAGAACACCGGCGACTATTACGAGGACTCTTACGAGGATATTAGCGCCTAC
 30 CTGCTGTCTAAGAACAACGCTATCGAGCCTAGAAGCTTCAGTCAGAACCCCCCGTGCTGAAGCGGCA
 CCAGAGAGAGATCACTAGAACTACCCTGCAGAGCGACCAGGAAGAGATCGACTACGACGACACTATTA
 GCGTCGAGATGAAGAAAGAGGATTTTCGATATCTACGACGAGGACGAGAAACAGTCACCTAGATCCTTC
 CAGAAGAAAAC TAGGCACTACTTTATTGCCGCCGTGAGCGGCTGTGGGACTACGGAATGAGTTCTAG
 CCCTCACGTGCTGAGAAATAGGGCTCAGTCAGGCTCAGTGCCTCAGTTCAAGAAAGTGGTGTTCAGG
 35 AATTCACCGACGGCAGCTTCACTCAGCCCTCTATAGGGGCGAGCTGAACGAGCACCTGGGACTGCTG
 GGACCTTATATTAGAGCCGAAGTCGAGGACAATATTATGGTCACCTTTAGGAACGAGCCCTCTAGGCC
 CTACAGCTTCTACTCTAGCCTGATCAGCTACGAGGAAGATCAGCGGCAGGGGGCCGAACCTAGAAAGA
 ACTTCGTGAAGCCTAACGAGACTAAGACCTACTTCTGGAAGGTGCAGCACCACATGGCCCCCTACTAAG
 GACGAGTTCGACTGTAAAGCCTGGGCCTACTTTAGCGACGTGGACCTCGAGAAGGACGTGCACTCAGG
 40 GCTGATCGGACCTCTGCTCGTCTGTACACTAACACCCCTGAACCCCGCTCACGGCCGGCAGGTACAG
 TGCAGGAATTCGCCCTGTTCTTCACTATCTTCGACGAACTAAGAGCTGGTACTTCACAGAGAATATG
 GAAAGAACTGTAGGGCCCCCTGTAATATTCAGATGGAAGATCCTACCTTTAAAGAGAACTATAGGTT
 TCACGCTATTAACGGCTATATTATGGACACCCCTGCCCGGCTCGTGATGGCTCAGGATCAGCGGATTA
 45 ACCGTCCGGAAGAAAGAGATATAAGATGGCCCTCTATAACCTCTACCCCGGCGTGTTCGAGACAGT
 GGAAATGCTGCCTAGTAAAGCCGGAATCTGGCGGGTTCGAGTGTCTGATCGGCGAGCACCTCCACGCCG
 GAATGAGCACCCCTGTTTCTGGTCTACTCTAACAAAGTGTACAGACCCCTCTGGGAATGGCTAGCGGCCAC
 ATTAGAGACTTTTCAGATCACCGCTAGCGGACAGTACGGCCAGTGGGCCCTAAGCTGGCTAGACTGCA
 CTATAGCGGATCTATCAACGCCCTGGTCTACCAAAGAGCCCTTTAGCTGGATTAAGGTGGACCTGCTGG
 50 CCCCTATGATTATTCACGGGATTAAGACTCAGGGCGCTAGGCAGAAGTTTAGTAGCCTCTATATTAGT
 CAGTTTATCATTATGTATAGCCTGGACGGCAAGAAGTGGCAGACCTATAGAGGCAATAGCACCGGCAC
 CCTGATGGTGTCTTCGGCAACGTGGACTCTAGCGGGATCAAGCACAAATATCTTTAACCCCCCTATTA
 TCGCTAGATATATTAGGCTGCACCCTACTCACTACTCTATTAGGTCTACCTGAGGATGGAAGTATG
 55 GGCTGCGATCTGAATAGCTGCTCTATGCCCTGGGGATGGAATCTAAGGCTATTAGCGACGCTCAGAT
 CACAGCCTCTAGCTACTTCACTAATATGTTTCGTACCTGGTCCCCCTAGCAAGGCCCGGCTGCACCTCC

AGGGCAGATCTAACGCTTGGCGGCCTCAGGTCAACAACCCTAAAGAGTGGCTGCAGGTCGACTTTTCAG
 AAAACTATGAAGGTCACCGGCGTGACCACTCAGGGCGTGAAATCACTGCTGACCTCTATGTACGTGAA
 AGAGTTCCTGATTAGCTCTAGCCAGGACGGCCACCAGTGGACCCTGTTCTTTCAGAACGGCAAAGTGA
 AAGTGTTCAGGGCAATCAGGATAGCTTCACCCCCGTGGTCAATAGCCTGGATCCCCACTGCTGACT
 5 AGATACCTGAGAATTCACCCTCAGTCTTGGGTGCACCAGATCGCCCTGAGAATGGAAGTCCTGGGCTG
 TGAAGCTCAGGACCTCTACTAA

Certain Definitions/Abbreviations Used

SQ and BDD: FVIII with B domain deletion

TTRm: TTR promoter with 4 mutations, from TAmGTGTAG to TATTGACTTAG

10 CO and CO3: codon optimized FVIII nucleic acid variant, set forth as SEQ ID NO:1

CO1: codon optimized FVIII nucleic acid variant, set forth as SEQ ID NO:10

CO2: codon optimized FVIII nucleic acid variant, set forth as SEQ ID NO:11

Prep designations in Table 3 refer to 5 separate vector preparations of CO/CO3.

hFVIII-RH: Arg to His substitution at position 1645 of human FVIII

15 Δ4: deletion of amino acids 1645 to 1648 of FVIII

Δ1645: deletion of amino acid 1645 (Arg) of FVIII

Δ2: deletion of amino acids 1645 and 1646 of FVIII

Δ3: deletion of amino acids 1645 to 1647 of FVIII

Δ1648: deletion of amino acid 1648 of FVIII

20 [0232] While certain of the embodiments of the invention have been described and
 specifically exemplified above, it is not intended that the invention be limited to such
 embodiments. Various modifications may be made thereto without departing from the scope and
 spirit of the invention, as set forth in the following claims.

THE EMBODIMENTS OF THE INVENTION FOR WHICH AN EXCLUSIVE PROPERTY OR PRIVILEGE IS CLAIMED ARE DEFINED AS FOLLOWS:

1. A Factor VIII (FVIII) encoding nucleic acid variant, wherein the FVIII encoded by said nucleic acid variant exhibits greater expression when compared to expression of wild type FVIII encoded by GenBank sequence accession number NG 011403.1, wherein said nucleic acid variant has one or more leucine codons changed to CTG compared to TTA, TTG, CTT, CTC or CTA in said wild type FVIII, has 50% to 59% GC content, and is at least 94% identical to the full length sequence of SEQ ID NO:1.
2. A Factor VIII (FVIII) encoding nucleic acid variant, wherein the FVIII encoded by said nucleic acid variant comprises a B domain deletion and exhibits greater expression when compared to expression of wild type FVIII encoded by GenBank sequence accession number NG 011403.1 comprising a B domain deletion, wherein said nucleic acid variant has one or more leucine codons changed to CTG compared to TTA, TTG, CTT, CTC or CTA in said wild type FVIII, has 45.2% to 59.2% GC content, and is at least 94% identical to the full length sequence of SEQ ID NO:1.
3. The variant FVIII as claimed in claim 1 or 2, wherein said nucleic acid variant is more efficiently packaged into an adeno-associated virus (AAV) vector.
4. The variant FVIII as claimed in any one of claims 1-3, wherein said FVIII encoded by said nucleic acid variant exhibits greater biological activity when compared to wild type FVIII or when compared to wild type FVIII comprising a B domain deletion.
5. The variant FVIII as claimed in claim 4, wherein said biological activity is determined by a clotting assay or reduced bleeding in a FVIII assay or FVIII deficiency model.
6. The variant FVIII as claimed in any one of claims 1-4, wherein said nucleic acid variant has 2-5, 5-10, 10-20, 20-50, 50-100, 100-250, 250-500, 500-750 or 750-850 CTG leucine codons modified from TTA, TTG, CTT, CTC or CTA leucine codons in wild type FVIII.

7. The variant FVIII as claimed in any one of claims 1-4, wherein said nucleic acid variant has greater than 85% CTG leucine codons modified from TTA, TTG, CTT, CTC or CTA leucine codons in wild type FVIII.

8. The variant FVIII as claimed in any one of claims 1-4, wherein said nucleic acid variant has all CTG leucine codons modified from TTA, TTG, CTT, CTC or CTA leucine codons in wild type FVIII.

9. The variant FVIII as claimed in any one of claims 1-4, wherein said nucleic acid variant has one or more AAG lysine codons compared to AAA lysine codons in wild type FVIII.

10. The variant FVIII as claimed in claim 1, wherein said nucleic acid variant has between 50-56%, or 50-53% GC content.

11. The variant FVIII as claimed in any one of claims 1-4, wherein said nucleic acid variant is at least 95% identical to the full length sequence of SEQ ID NO:1.

12. The variant FVIII as claimed in any one of claims 1-4, wherein said nucleic acid variant is human FVIII, and/or wherein said wild type FVIII or wild type FVIII comprising a B domain deletion is human FVIII.

13. The variant FVIII as claimed in claim 1, wherein said nucleic acid variant comprises any of SEQ ID NOs:1-7 and 9.

14. The variant FVIII as claimed in any one of claims 1-13, wherein 1, 2, 3 or all 4 codons encoding the PACE/furin cleavage site is/are deleted.

15. The variant FVIII as claimed in any one of claims 1-13, wherein 1, 2, 3 or all 4 codons encoding the PACE/furin cleavage site set forth as HHQR or RHQR from positions 1645-1648 is/are deleted.

16. An expression vector comprising the variant FVIII of any one of claims 1-15.
17. The expression vector of claim 16, selected from the group consisting of an adeno associated virus (AAV) vector, a retroviral vector, an adenoviral vector, a plasmid, and a lentiviral vector.
18. The expression vector of claim 17, wherein said AAV vector comprises an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 or AAV-2i8 AAV serotype.
19. The expression vector of claim 16, further comprising an intron, an expression control element, one or more adeno-associated virus (AAV) inverted terminal repeats (ITRs) and/or a filler polynucleotide sequence.
20. The expression vector of claim 19, wherein the intron is within or flanks the variant FVIII, or wherein the expression control element is operably linked to the variant FVIII, or wherein the AAV ITR(s) flanks the 5' or 3' terminus of the variant FVIII, or wherein the filler polynucleotide sequence flanks the 5' or 3' terminus of the variant FVIII.
21. The expression vector of claim 19, wherein the expression control element comprises a constitutive or regulatable control element, or a tissue-specific expression control element or promoter.
22. The expression vector of claim 19, wherein the expression control element comprises an element that confers expression in liver.
23. The expression vector of claim 19, wherein the expression control element comprises a TTR promoter or mutant TTR promoter.
24. The expression vector of claim 23, wherein the mutant TTR promoter comprises SEQ ID NO:8.

25. The expression vector of claim 19, wherein the ITR comprises one or more ITRs of any of: AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 or AAV-2i8 AAV serotypes, or a combination thereof.
26. A host cell expressing the FVIII encoded by said nucleic acid variant of any one of claims 1-15.
27. A host cell comprising the nucleic acid variant of any one of claims 1-15 or the expression vector of any one of claims 16-25.
28. A virus vector comprising the nucleic acid variant of any one of claims 1-15 or the expression vector of any one of claims 16-25.
29. An AAV vector comprising the nucleic acid variant of any one of claims 1-15 or the expression vector of any one of claims 16-25.
30. The AAV vector of claim 29, wherein the AAV vector comprises a VP1, VP2 and/or VP3 capsid sequence having 75% or more sequence identity to AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 or AAV-2i8 VP1, VP2 and/or VP3 sequences.
31. The AAV vector of claim 29, wherein the AAV vector comprises a VP1, VP2 and/or VP3 capsid sequence selected from any of: AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, Rh10, Rh74 and AAV-2i8 AAV serotypes.
32. A pharmaceutical composition comprising the nucleic acid variant of any one of claims 1-15, the expression vector of any one of claims 16-25, or the virus vector of claim 28 or AAV vector of any one of claims 29-31 in a biologically compatible carrier or excipient.
33. The pharmaceutical composition of claim 32, further comprising empty capsid AAV.

34. The nucleic acid variant of any one of claims 1-15, the expression vector of any one of claims 16-25, or the virus or AAV vector of any one of claims 28-31 encapsulated in a liposome or mixed with phospholipids or micelles.

35. A use of the nucleic acid variant of any one of claims 1-15, the expression vector of any one of claims 16-25, or the virus vector of claim 28 or AAV vector of any one of claims 29-31 to deliver or transfer a nucleic acid sequence into a mammal or a mammalian cell.

36. The use of claim 35, wherein said mammalian cell is in culture or is present in a subject.

37. A use of the nucleic acid variant of any one of claims 1-15, the expression vector of any one of claims 16-25, or the virus vector of claim 28 or AAV vector of any one of claims 29-31 to treat a mammal in need of Factor VIII.

38. The use of claim 35 or 37, wherein said Factor VIII provides a beneficial or therapeutic effect on the mammal.

39. The use of claim 35 or 37, wherein expression of said Factor VIII is in a cell, tissue or organ of said mammal.

40. The use of claim 35 or 37, wherein, the cell comprises a secretory cell.

41. The use of claim 35 or 37, wherein the cell comprises an endocrine cell or an endothelial cell.

42. The use of claim 35 or 37, wherein the cell comprises a hepatocyte, a sinusoidal endothelial cell, a megakaryocyte, a platelet or hematopoietic stem cell.

43. The use of claim 39, wherein the tissue or organ of said mammal comprises liver.

44. The use of claim 35 or 37, wherein the mammal produces an insufficient amount of Factor VIII protein, or a defective or aberrant Factor VIII protein.
45. The use of claim 35 or 37, wherein the mammal has hemophilia A.
46. A use of the nucleic acid variant of any one of claims 1-15, the expression vector of any one of claims 16-25, or the virus of claim 28 or AAV vector of any of claims 29-31 for treatment of a hemostasis related disorder in a patient in need thereof.
47. The use of claim 35, 37 or 46, wherein said mammal or said patient has a disorder selected from the group consisting of hemophilia A, von Willebrand diseases and bleeding associated with trauma, injury, thrombosis, thrombocytopenia, stroke, coagulopathy, disseminated intravascular coagulation (DIC) and over-anticoagulation treatment disorders.
48. The use of any one of claims 35-47, wherein the nucleic acid variant of any one of claims 1-15, the expression vector of any one of claims 16-25, or the virus of claim 28 or AAV vector of any one of claims 29-31 is for delivery to said mammal or said patient intravenously, intraarterially, intramuscularly, subcutaneously, intra-cavity, or by intubation, or via catheter.
49. The use of any one of claims 35 and 37-45, wherein said mammal or said patient is human.
50. The use of any one of claims 35-45, wherein said mammal or said patient is sero-positive or sero-negative for an AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV-Rh10 or AAV-Rh74 serotype.
51. The use of any one of claims 35-45, further comprising use of an empty capsid AAV.
52. The use of any one of claims 35-45, further comprising use of an empty capsid of AAV1, AAV2, AAV3, AAV4, AAV5, AAV6, AAV7, AAV8, AAV9, AAV10, AAV11, AAV-12, AAV-Rh10 and/or AAV-Rh74 serotype.

53. A method of producing FVIII protein comprising expressing in a cell the variant FVIII as claimed in any one of claims 1-15, or the expression vector of claim 16 and recovering said FVIII protein produced by the cells.

54. The method of claim 53, further comprising purifying or isolating said FVIII protein produced by the cells.

55. A Factor VIII (FVIII) encoding nucleic acid variant according to the sequence of any one of SEQ ID NO:1, SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:5, SEQ ID NO:6, SEQ ID NO:7, SEQ ID NO:9, SEQ ID NO:10 and SEQ ID NO:11.

Fig. 1




Fig. 2




Fig. 3A




Fig. 3B




Fig. 4A

HCR/HAAT-hFVIIIwtSQ

EC	HCR/	WT	WT	WT
EC	HAAT	WT	WT	WT

TTR-hFVIII CO

EC	TTR	CO	CO	CO
EC	TTR	CO	CO	CO

Fig. 4B
AAV delivery of hFVIII in hemophilia A/CD4 KO mice




Fig. 4C
AAV delivery of hFVIII in hemophilia A/CD4 KO mice




Fig. 5A




Fig. 5B




Fig. 6A




Fig. 6B




Fig. 7

