

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
22 June 2017 (22.06.2017)

(10) International Publication Number
WO 2017/103850 A1

(51) International Patent Classification:

A61K 31/702 (2006.01) A61P 31/04 (2006.01)
A61K 31/7012 (2006.01) A61P 31/12 (2006.01)

(21) International Application Number:

PCT/IB2016/057675

(22) International Filing Date:

15 December 2016 (15.12.2016)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

15200067.5 15 December 2015 (15.12.2015) EP
16154144.6 3 February 2016 (03.02.2016) EP

(71) Applicant: GLYCOM A/S [DK/DK]; Kogle Allé 4, 2970
Hørsholm (DK).

(72) Inventors: MCCONNELL, Bruce; 2 Chemin du Crêt à
l'Aigle, 1814 La Tour de Peilz (CH). VIGSNÆS, Louise
Kristine; Ravneholmsvej 20, 2400 København NV (DK).

(74) Agent: BECK GREENER; Fulwood House, 12 Fulwood
Place, London WC1V 6HR (GB).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KH, KN,
KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA,
MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG,
NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS,
RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY,
TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN,
ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— with international search report (Art. 21(3))

(54) Title: MIXTURE OF HMOS

(57) Abstract: A synthetic mixture of human milk oligosaccharides (HMOs) consisting essentially of lacto -N-neotetraose (LNnT), lacto-N-tetraose (LNT), 2'-fucosyllactose (2'FL), 3'-O-sialyllactose (3'-SL), 6'-O-sialyllactose (6'-SL) either difucosyllactose (DFL) or 3-fucosyllactose (3-FL), preferably DFL, and optionally lactose. Said synthetic composition is useful for: treating or preventing viral and/or bacterial infection in a non-infant human; modulating the microbiota of a non-infant human; and/or improving the cognitive function of a non-infant human and as a pharmaceutical or nutritional composition.



WO 2017/103850 A1

MIXTURE OF HMOs

FIELD OF THE INVENTION

The present invention relates to synthetic mixtures of human milk oligosaccharides
5 (“HMOs”), particularly of LNT, 2'-FL, 3'-SL, 6'-SL and either DFL or 3-FL and
applications of the mixtures in human health.

BACKGROUND OF THE INVENTION

HMOs have become the subject of much interest in recent years due to their roles in
numerous biological processes occurring in the human organism. Mammalian milk contains
10 at least 130 of these complex oligosaccharides (Urashima et al, Milk Oligosaccharides, Nova
Biomedical Books, New York, 2011, ISBN: 978-1-61122-831-1).

Previously, the only source of HMOs had been mammalian milk which contains mostly
water, together with 55-70 g/l lactose, 24-59 g/l lipids, ca. 13 g/l proteins, 5-15 g/l HMOs and
ca. 1.5 g/l minerals.

15 However, several processes for synthesizing HMOs have been developed in recent
years due to their roles in numerous human biological processes. In this regard, processes
have been developed for producing HMOs by microbial fermentations, enzymatic processes,
chemical syntheses, or combinations of these technologies. For example, by chemical
processes, Galp β 1-4GlcNAcp β 1-3Galp β 1-4Glc or lacto-N-neotetraose (“LNT”) can be
20 made as described in WO 2011/100980 and WO 2013/044928; Galp β 1-3GlcNAcp β 1-
3Galp β 1-4Glc or lacto-N-tetraose (“LNT”) can be synthesized as described in WO
2012/155916 and WO 2013/044928; 6'-O-sialyllactose (“6'-SL”) can be synthesized as
described in WO 2011/100979; a mixture of LNT and LNT can be made as described in
WO 2013/091660; 2'-O-fucosyllactose (“2'-FL”) can be made as described in WO
25 2010/115934 and WO 2010/115935; 3-fucosyllactose (“3-FL”) can be made as described in
WO 2013/139344; and 6'-SL can be made as described in WO 2010/100979. As examples
of biotechnological processes, WO 01/04341 and WO 2007/101862 describe how to make
core human milk oligosaccharides optionally substituted by fucose or sialic acid, including
LNT, 6'-SL and 3'-O-sialyllactose (“3'-SL”) using genetically modified *E. coli*; and WO
30 2015/032412 describes making 2'-FL and difucosyllactose or Fuc(α 1-2)Gal(β 1-4)[Fuc(α 1-
3)]Glc (“DFL”) using genetically modified *E. coli*. As an example of enzymatic processes,
sialylated oligosaccharides can be made as described in EP-A-577580.

Efforts have also been made to develop processes for synthesizing enzymatically
mixtures of HMO oligosaccharides, without having to synthesize all of the component

oligosaccharides of the mixture as described in WO 2012/156897 and WO 2012/156898. Such processes have provided reaction mixtures containing a plurality of different oligosaccharides.

Evidence is accumulating that the resident community of microbes, called the microbiota, in the human digestive tract plays a major role in health and disease. When the composition of the intestinal microbiota is thrown off balance, the human host can suffer consequences. Recent research has implicated intestinal microbiota imbalances in individual disorders as diverse as cancer, obesity, inflammatory bowel disease, psoriasis, asthma, and possibly even autism. Individual non-digestible fibres, including HMOs, are believed to beneficially modulate the microbiota, and they are of increasing interest for treating one or more of such disorders. However, many digestible fibres non-specifically modulate the microbiota while others are not able to provide sufficiently broad, but specific, modulation.

Therefore, there has been a need to specifically modulate the microbiota, so as to address individual disorders in different ways and also to address simultaneously multiple disorders. In particular, there has been a need for a composition that can be used for, *inter alia*, treating and/or preventing bacterial and viral infections, particularly in the intestinal and respiratory tracts, improving cognitive function and/or increasing the efficacy of anticancer agents against tumors.

SUMMARY OF THE INVENTION

A first aspect of this invention relates to a synthetic mixture of HMOs consisting essentially of LNnT, LNT, 2'-FL, 3'-SL, 6'-SL and either DFL or 3-FL, preferably DFL. This mixture can optionally include lactose. The mixture of HMOs preferably consists essentially of:

- i. about 55 wt % to about 75 wt % of 2'-FL, more preferably about 60 wt % to about 70 wt % of 2'-FL;
- ii. about 2 wt % to about 10 wt % of LNnT, more preferably about 3 wt % to about 7 wt % of LNnT;
- iii. about 10 wt % to about 20 wt % of LNT, more preferably about 12 wt % to about 18 wt % of LNT;
- iv. about 1 wt % to about 15 wt % of DFL or 3-FL, more preferably about 1 wt % to about 10 wt %, even more preferably about 2 wt % to about 10 wt % of DFL or 3-FL such as e.g. about 2 wt % to about 8 wt %;

- v. about 1 wt % to about 10 wt % of 3'-SL, more preferably about 2 wt % to about 8 wt % of 3'-SL; and
- vi. about 1 wt % to about 15 wt % of 6'-SL, preferably about 5 wt % to about 15 wt % of 6'-SL, more preferably about 7 wt % to about 13 wt % of 6'-SL.

5 A second aspect of the invention relates to a nutritional or pharmaceutical composition comprising a synthetic mixture of HMOs according to the first aspect of the invention.

A third aspect of this invention relates to a synthetic mixture of HMOs or a composition comprising a synthetic mixture of HMOs for use in: i) preventing and/or treating viral and/or bacterial infections in a non-infant human; ii) specifically modulating the indigenous
10 microbiota of a non-infant human; and/or iii) improving the cognitive function of a non-infant human. The synthetic mixture of HMOs consists essentially of LNnT, LNT, 2'-FL, 3'-SL, 6'-SL, either DFL or 3-FL, preferably DFL, and optionally lactose, as described above. The synthetic HMO mixture or the nutritional or pharmaceutical composition comprising it contains a plurality of different HMOs with novel combinations of properties and biological
15 activities. The composition is especially useful against viral and bacterial, intestinal infections through specific modulation of the intestinal microbiota by an increase in *Bifidobacterium*, modulation of intestinal binding of viruses and pathogenic bacteria to intestinal epithelial cells, and improvement of intestinal barrier function. The composition is also especially useful against viral and bacterial, respiratory tract infections by inhibiting pathogen binding to
20 human epithelial cells.

A fourth aspect of this invention relates to a method of modulating the indigenous microbiota of a non-infant human to increase the abundance of *Bifidobacterium* in order to increase the efficacy of anticancer agents against tumors in a non-infant human patient. The method involves administering to the non-infant human a synthetic mixture of HMOs
25 according to the first aspect of this invention or a nutritional or pharmaceutical composition according to the second aspect of this invention, as described above. *Bifidobacterium* can act as an immune booster, hence the increase in its abundance can strengthen a cancer patient's response to an anticancer agent. This property makes the mixture suitable as an aid in cancer therapy.

30 A fifth aspect of this invention relates to a method of modulating the indigenous intestinal microbiota of a non-infant human to increase *Bifidobacterium* and/or *Barnesiella* abundance and also to reduce the abundance of *Ruminococcus gnavus*. The method involves administering to the non-infant human a synthetic mixture of HMOs according to the first aspect of this invention or a nutritional or pharmaceutical composition according to the
35 second aspect of this invention, as described above. The increased *Bifidobacterium* and/or *Barnesiella* abundance and reduced abundance of *Ruminococcus gnavus* render the non-

infant human's intestinal milieu less prone to inflammation and provide improved intestinal barrier function. These effects can prevent and/or treat conditions such as inflammatory bowel disease, irritable bowel syndrome, and other conditions associated with inflammation and impaired gut barrier function.

5 A sixth aspect of this invention relates to a method of modulating the indigenous microbiota of a non-infant human to increase *Bifidobacterium* abundance and to at least maintain the abundance of *Faecalibacterium*. The method involves administering to the non-infant human a synthetic mixture of HMOs according to the first aspect of this invention or a nutritional or pharmaceutical composition according to the second aspect of this invention,
10 as described above. Increasing the *Bifidobacterium* abundance and at least maintaining the abundance of *Faecalibacterium* render the non-infant human's intestinal milieu less prone to inflammation and provide improved intestinal barrier function. Preferably the abundance of *Ruminococcus gnavus* is reduced. These effects can prevent and/or treat conditions such as inflammatory bowel disease, irritable bowel syndrome, and other conditions associated
15 with inflammation and impaired gut barrier function.

In both the fifth and sixth aspects of the invention, the abundance of Proteobacteria is also preferably reduced.

A seventh aspect of this invention relates to a method of preventing or treating viral and/or bacterial infections in a non-infant human, especially intestinal infections and
20 infections of the respiratory tract. The method comprises administering, to the non-infant human, a synthetic mixture of HMOs according to the first aspect of this invention or a nutritional or pharmaceutical composition according to the second aspect of this invention, as described above.

An eighth aspect of this invention relates to a method of improving the cognitive
25 function of a non-infant human. The method comprises administering, to the non-infant human, a synthetic mixture of HMOs according to the first aspect of this invention or a nutritional or pharmaceutical composition according to the second aspect of this invention, as described above.

DETAILED DESCRIPTION OF THE INVENTION

30 In accordance with this invention, the following terms preferably have the following meanings:

"Non-infant human" or "non-infant" preferably means a human of 3 years of age and older. Accordingly, a non-infant human is a human of any age above 3 years old, e.g. it can be a child, a teenager, an adult or an elderly person.

"Synthetic mixture of HMOs" means a mixture which is artificially prepared and preferably means a mixture in which at least one HMO is produced *ex vivo* chemically and/or biologically, e.g. by means of chemical reaction, enzymatic reaction or recombinantly. More preferably, all of the HMOs in the mixture are produced *ex vivo* chemically and/or biologically. In some embodiments, a synthetic mixture of the invention may be, but preferably is not, identical with a naturally occurring mixture

"Microbiota", "microflora" and "microbiome" mean a community of living microorganisms that typically inhabits a bodily organ or part, particularly the gastro-intestinal organs of non-infant humans. The most dominant members of the gastrointestinal microbiota include microorganisms of the phyla of *Firmicutes*, *Bacteroidetes*, *Actinobacteria*, *Proteobacteria*, *Synergistetes*, *Verrucomicrobia*, *Fusobacteria*, and *Euryarchaeota*; at genus level *Bacteroides*, *Faecalibacterium*, *Bifidobacterium*, *Roseburia*, *Alistipes*, *Collinsella*, *Blautia*, *Coprococcus*, *Ruminococcus*, *Eubacterium* and *Dorea*; at species level *Bacteroides uniformis*, *Alistipes putredinis*, *Parabacteroides merdae*, *Ruminococcus bromii*, *Dorea longicatena*, *Bacteroides caccae*, *Bacteroides thetaiotaomicron*, *Eubacterium hallii*, *Ruminococcus torques*, *Faecalibacterium prausnitzii*, *Ruminococcus lactaris*, *Collinsella aerofaciens*, *Dorea formicigenerans*, *Bacteroides vulgatus* and *Roseburia intestinalis*. The gastrointestinal microbiota includes the mucosa-associated microbiota, which is located in or attached to the mucus layer covering the epithelium of the gastrointestinal tract, and luminal-associated microbiota, which is found in the lumen of the gastrointestinal tract.

"Modulating of microbiota" means exerting a modifying or controlling influence on microbiota, in particular an influence leading to an increase in the indigenous intestinal abundance of *Bifidobacterium*, *Barnesiella* and/or *Faecalibacterium*, and reduction of the intestinal abundance of *Ruminococcus gnavus* and/or *Proteobacteria*.

"Proteobacteria" are a phylum of Gram-negative bacteria and include a wide variety of pathogenic bacteria, such as *Escherichia*, *Salmonella*, *Vibrio*, *Helicobacter*, *Yersinia* and many other notable genera.

"Therapy" means treatment given or action taken to reduce or eliminate symptoms of a disease or pathological condition.

"Preventive treatment" or "prevention" in the present context means a treatment given or an action taken to diminish the risk of the onset or recurrence of a disease.

"Viral infection" in the present context means an infection of the stomach and intestines, or any other parts of the gastro-intestinal tract, by a virus often resulting in inflammatory response, commonly known as the stomach flu. The symptoms may include one or more of the following: watery diarrhoea, nausea and vomiting, headache, muscle aches, joint aches fever, chills, sweating, clammy skin, abdominal cramps and pain, loss of

appetite, weight loss. Examples of viral gastroinfections include, but not limited to, viral gastroinfections caused by Rotavirus or Norovirus.

“Bacterial infection” in the present context means an infection of the stomach and intestines, or any other parts of the gastro-intestinal tract, by a pathogenic bacteria, commonly known as bacterial enteritis. The symptoms may include one or more of the following: loss of appetite, nausea and vomiting, diarrhoea, abdominal pains and cramps, blood in the stool, fever. Examples of bacterial infections include, but not limited to, gastroinfections by pathogenic strains of *Yersinia*, *Staphylococcus*, *Shigella*, *Salmonella*, *Campylobacter* and *E. Coli*.

“Infections of the respiratory tract” refers to any viral and/or bacterial infections of upper and lower respiratory tract, i.e. infections of the sinuses, throat, airways or lungs.

“Cancer” preferably refers to a neoplasm; for example of any part of one or more of the following: respiratory, gastro-intestinal, urinary tracts or liver.

“Oral administration” means any conventional form for the delivery of a composition through the mouth. Accordingly, oral administration is a form of enteral administration.

“Cognitive function” refers to cerebral activities that lead to knowledge, including all means and mechanisms of acquiring information. Cognitive functions encompass reasoning, memory, attention, and language and lead directly to the attainment of information and, thus, knowledge.

In accordance with this invention, it has been surprisingly discovered that a synthetic HMO mixture consisting essentially of LNnT, LNT, 2'-FL, 3'-SL, 6'-SL and either DFL or 3-FL, preferably DFL, and optionally lactose, can provide an anti-infective composition for preventing or treating, bacterial or viral infections through specific modulation of the intestinal microbiota, binding of viruses, reduction in pathogenic translocation and improvement of intestinal barrier function in a non-infant human. Further, the HMO mixture of this invention acts as a decoy receptor and binds to rotaviruses to prevent the rotaviruses from adhering to human intestinal cells. These properties, coupled with an improvement in intestinal barrier function, make the HMO mixture suitable for preventing and treating intestinal infections.

It has also been found that the HMO mixture of this invention can increase the brain ganglioside and glycoprotein sialic acid concentrations, and enhance hippocampal long term potentiation through the vagus nerve, leading to increased synaptogenesis and neurodevelopment in a non-infant human. This makes the HMO mixture suitable for administration to non-infant humans to improve their cognitive function.

The HMO mixture of this invention can also:

i) increase the indigenous intestinal abundance of *Bifidobacterium*, and

- ii) increase the intestinal abundance of *Barnesiella* and/or at least maintain the intestinal abundance of *Faecalibacterium*, and
- iii) reduce the intestinal abundance of *Ruminococcus gnavus* and/or Proteobacteria

5 in non-infant humans.

These effects can render an intestinal milieu less prone to inflammation. Coupled with an improvement in intestinal barrier function, these effects of the HMO mixture can prevent and/or treat conditions such as inflammatory bowel disease, irritable bowel syndrome, and other conditions associated with inflammation and impaired barrier function.

10 Surprisingly, the increase of bifidobacteria in the gut, induced by the HMO mixture of this invention, increases the efficacy of anticancer agents against tumors. The bifidobacteria act as immune helpers, strengthening a cancer patient's response to an anticancer agent. This property makes the HMO mixture suitable as an aid in cancer therapy.

Preferably the HMO mixture of this invention contains: i) about 55 wt % to about 75 wt
15 % of 2'-FL, more preferably about 60 wt % to about 70 wt %; ii) about 2 wt % to about 10 wt % of LNnT, more preferably about 3 wt % to about 7 wt %; iii) about 10 wt % to about 20 wt % of LNT, more preferably about 12 wt % to about 18 wt %; iv) about 1 wt % to about 15 wt % of DFL or 3-FL, preferably about 1 wt % to about 10 wt %, even more preferably about 2 wt % to about 10 wt %; v) about 1 wt % to about 10 wt % of 3'-SL, more preferably about 2
20 wt % to about 8 wt %; and vi) about 1 wt % to about 15 wt % of 6'-SL, preferably about 5 wt % to about 15 wt %, more preferably about 7 wt % to about 13 wt %. The HMO mixture of this invention can also contain lactose, but it is not considered an active ingredient of the mixture.

The synthetic HMO mixture can be administered to a non-infant human in any suitable
25 form such as a nutritional or pharmaceutical composition, for example, a unit dosage form (for example, a tablet, a capsule, a sachet of powder, etc.).

The nutritional or pharmaceutical composition consists of the synthetic mixture of HMOs consisting essentially of LNnT, LNT, 2'-FL, 3'-SL, 6'-SL, either DFL or 3-FL, and optionally lactose, and suitable excipients. The nutritional or pharmaceutical composition
30 does not contain any other HMOs than those mentioned above.

In one embodiment, the HMO mixture of this invention can be in the form of a nutritional composition. For example, the nutritional composition can be a food composition, a rehydration solution, or a dietary maintenance or supplement for elderly individuals or immunocompromised individuals. The nutritional composition can contain sources of protein,
35 lipids and/or digestible carbohydrates and can be in powdered or liquid forms. The composition can be designed to be the sole source of nutrition or a nutritional supplement.

Suitable protein sources include milk proteins, soy protein, rice protein, pea protein and oat protein, or mixtures thereof. Milk proteins can be in the form of milk protein concentrates, milk protein isolates, whey protein or casein, or mixtures of both. The protein can be whole protein or hydrolysed protein, either partially hydrolysed or extensively hydrolysed. Hydrolysed protein offers the advantage of easier digestion which can be important for non-infants with inflamed GI tracts. The protein can also be provided in the form of free amino acids. The protein can comprise about 5 % to about 30 % of the energy of the nutritional composition, normally about 10 % to 20 %.

The protein source can be a source of glutamine, threonine, cysteine, serine, proline, or a combination of these amino acids. The glutamine source can be a glutamine dipeptide and/or a glutamine enriched protein. Glutamine can be included due to the use of glutamine by enterocytes as an energy source. Threonine, serine and proline are important amino acids for the production of mucin. Mucin coats the GI tract and can improve mucosal healing. Cysteine is a major precursor of glutathione, which is key for the antioxidant defences of the body.

Suitable digestible carbohydrates include maltodextrin, hydrolysed or modified starch or corn starch, glucose polymers, corn syrup, corn syrup solids, high fructose corn syrup, rice-derived carbohydrates, pea-derived carbohydrates, potato-derived carbohydrates, tapioca, sucrose, glucose, fructose, sucrose, lactose, honey, sugar alcohols (e.g., maltitol, erythritol, sorbitol), or mixtures thereof. Preferably the composition is free from added lactose. Generally digestible carbohydrates provide about 35 % to about 55 % of the energy of the nutritional composition. A particularly suitable digestible carbohydrate is a low dextrose equivalent (DE) maltodextrin.

Suitable lipids include medium chain triglycerides (MCT) and long chain triglycerides (LCT). Preferably the lipid is a mixture of MCTs and LCTs. For example, MCTs can comprise about 30 % to about 70 % by weight of the lipids, more specifically about about 50 % to about 60 % by weight. MCTs offer the advantage of easier digestion which can be important for non-infants with inflamed GI tracts. Generally, the lipids provide about 35 % to about 50 % of the energy of the nutritional composition. The lipids can contain essential fatty acids (omega-3 and omega-6 fatty acids). Preferably these polyunsaturated fatty acids provide less than about 30 % of total energy of the lipid source. Decreasing the levels of these polyunsaturated fatty acids is believed to decrease sensitivity to peroxidation; which can be beneficial for non-infants having inflammatory conditions.

Suitable sources of long chain triglycerides are rapeseed oil, sunflower seed oil, palm oil, soy oil, milk fat, corn oil, high oleic oils, and soy lecithin. Fractionated coconut oils are a suitable source of medium chain triglycerides. The lipid profile of the nutritional composition

is preferably designed to have a polyunsaturated fatty acid omega-6 (n-6) to omega-3 (n-3) ratio of about 4:1 to about 10:1. For example, the n-6 to n-3 fatty acid ratio can be about 6:1 to about 9:1.

The nutritional composition may also include vitamins and minerals. If the nutritional composition is intended to be a sole source of nutrition, it preferably includes a complete vitamin and mineral profile. Examples of vitamins include vitamins A, B-complex (such as B1, B2, B6 and B12), C, D, E and K, niacin and acid vitamins such as pantothenic acid, folic acid and biotin. Examples of minerals include calcium, iron, zinc, magnesium, iodine, copper, phosphorus, manganese, potassium, chromium, molybdenum, selenium, nickel, tin, silicon, vanadium and boron.

The nutritional composition can also include a carotenoid such as lutein, lycopene, zeaxanthin, and beta-carotene. The total amount of carotenoid included can vary from about 0.001 µg/ml to about 10 µg/ml. Lutein can be included in an amount of from about 0.001 µg/ml to about 10 µg/ml, preferably from about 0.044 µg/ml to about 5 µg/ml of lutein. Lycopene can be included in an amount from about 0.001 µg/ml to about 10 µg/ml, preferably about 0.0185 µg/ml to about 5 µg/ml of lycopene. Beta-carotene can comprise from about 0.001 µg/ml to about 10 mg/ml, for example about 0.034 µg/ml to about 5 µg/ml of beta-carotene.

The nutritional composition preferably also contains reduced concentrations of sodium; for example, from about 300 mg/l to about 400 mg/l. The remaining electrolytes can be present in concentrations set to meet needs without providing an undue renal solute burden on kidney function. For example, potassium is preferably present in a range of about 1180 to about 1300 mg/l; and chloride is preferably present in a range of about 680 to about 800 mg/l.

The nutritional composition can also contain various other conventional ingredients such as preservatives, emulsifying agents, thickening agents, buffers, fibres and prebiotics (e.g. fructooligosaccharides, galactooligosaccharides), probiotics (e.g. *B. animalis* subsp. *lactis* BB-12, *B. lactis* HN019, *B. lactis* Bi07, *B. infantis* ATCC 15697, *L. rhamnosus* GG, *L. rhamnosus* HNOOI, *L. acidophilus* LA-5, *L. acidophilus* NCFM, *L. fermentum* CECT5716, *B. longum* BB536, *B. longum* AH1205, *B. longum* AH1206, *B. breve* M-16V, *L. reuteri* ATCC 55730, *L. reuteri* ATCC PTA-6485, *L. reuteri* DSM 17938), antioxidant/anti-inflammatory compounds including tocopherols, carotenoids, ascorbate/vitamin C, ascorbyl palmitate, polyphenols, glutathione, and superoxide dismutase (melon), other bioactive factors (e.g. growth hormones, cytokines, TFG-β), colorants, flavours, and stabilisers, lubricants, and so forth.

The nutritional composition can be formulated as a soluble powder, a liquid concentrate, or a ready-to-use formulation. The composition can be fed to a human in need via a nasogastric tube or orally. Various flavours, fibres and other additives can also be present.

5 The nutritional compositions can be prepared by any commonly used manufacturing techniques for preparing nutritional compositions in solid or liquid form. For example, the composition can be prepared by combining various feed solutions. A protein-in-fat feed solution can be prepared by heating and mixing the lipid source and then adding an emulsifier (e.g. lecithin), fat soluble vitamins, and at least a portion of the protein source
10 while heating and stirring. A carbohydrate feed solution is then prepared by adding minerals, trace and ultra trace minerals, thickening or suspending agents to water while heating and stirring. The resulting solution is held for 10 minutes with continued heat and agitation before adding carbohydrates (e.g. the HMOs and digestible carbohydrate sources). The resulting feed solutions are then blended together while heating and agitating and the pH adjusted to
15 6.6-7.0, after which the composition is subjected to high-temperature short-time processing during which the composition is heat treated, emulsified and homogenized, and then allowed to cool. Water soluble vitamins and ascorbic acid are added, the pH is adjusted to the desired range if necessary, flavours are added, and water is added to achieve the desired total solid level.

20 For a liquid product, the resulting solution can then be aseptically packed to form an aseptically packaged nutritional composition. In this form, the nutritional composition can be in ready-to-feed or concentrated liquid form. Alternatively, the composition can be spray-dried and processed and packaged as a reconstitutable powder.

 When the nutritional product is a ready-to-feed nutritional liquid, it may be preferred
25 that the total concentration of HMOs in the liquid, by weight of the liquid, is from about 0.0001 % to about 2.0 %, including from about 0.001 % to about 1.5 %, including from about 0.01 % to about 1.0 %. When the nutritional product is a concentrated nutritional liquid, it may be preferred that the total concentration of HMOs in the liquid, by weight of the liquid, is from about 0.0002 % to about 4.0 %, including from about 0.002 % to about 3.0 %, including
30 from about 0.02 % to about 2.0 %.

 In other embodiment, the HMO mixture of this invention can be comprised in a pharmaceutical composition. The pharmaceutical composition can contain a pharmaceutically acceptable carrier, e.g. phosphate buffered saline solution, mixtures of ethanol in water, water and emulsions such as an oil/water or water/oil emulsion, as well as
35 various wetting agents or excipients. The pharmaceutical composition can also contain other materials that do not produce an adverse, allergic or otherwise unwanted reaction when

administered to non-infants. The carriers and other materials can include solvents, dispersants, coatings, absorption promoting agents, controlled release agents, and one or more inert excipients, such as starches, polyols, granulating agents, microcrystalline cellulose, diluents, lubricants, binders, and disintegrating agents. If desired, tablet dosages of the anti-infective compositions can be coated by standard aqueous or non-aqueous techniques.

The pharmaceutical compositions can be administered orally, e.g. as a tablet, capsule, or pellet containing a predetermined amount, or as a powder or granules containing a predetermined concentration or a gel, paste, solution, suspension, emulsion, syrup, bolus, electuary, or slurry, in an aqueous or non-aqueous liquid, containing a predetermined concentration. Orally administered compositions can include binders, lubricants, inert diluents, flavouring agents, and humectants. Orally administered compositions such as tablets can optionally be coated and can be formulated so as to provide sustained, delayed or controlled release of the mixture therein.

The pharmaceutical compositions can also be administered by rectal suppository, aerosol tube, naso-gastric tube or direct infusion into the GI tract or stomach.

The pharmaceutical compositions can also include therapeutic agents such as antiviral agents, antibiotics, probiotics, analgesics, and anti-inflammatory agents. The proper dosage of these compositions for a non-infant human can be determined in a conventional manner, based upon factors such immune status, body weight and age. In some cases, the dosage will be at a concentration similar to that found for the HMOs in human breast milk. The required amount would generally be in the range from about 200 mg to about 20 g per day, in certain embodiments from about 300 mg to about 15 g per day, from about 400 mg to about 10 g per day, in certain embodiments from about 500 mg to about 10 g per day, in certain embodiments from about 1 g to about 10 g per day. Appropriate dose regimes can be determined by conventional methods.

Pharmaceutical compositions of the invention can be used for treatment of a concerned disease in combination with other medicaments prescribed for said disease, e.g. in combination with an anti-diabetes medicine or antibiotic therapy.

Yet in other embodiment, the composition comprising the HMO mixture of this invention is a unit dosage form. The unit dosage form can contain an acceptable carrier, e.g. phosphate buffered saline solution, mixtures of ethanol in water, water and emulsions such as an oil/water or water/oil emulsion, as well as various wetting agents or excipients. The unit dosage form can also contain other materials that do not produce an adverse, allergic or otherwise unwanted reaction when administered to a patient. The carriers and other materials can include solvents, dispersants, coatings, absorption promoting agents,

controlled release agents, and one or more inert excipients, such as starches, polyols, granulating agents, microcrystalline cellulose, diluents, lubricants, binders, and disintegrating agents. If desired, tablet dosages of the composition can be coated by standard aqueous or nonaqueous techniques.

5 A unit dosage form of this invention can be administered orally, e.g. as a tablet, capsule, or pellet containing a predetermined amount of the mixture, or as a powder or granules containing a predetermined concentration of the mixture or a gel, paste, solution, suspension, emulsion, syrup, bolus, electuary, or slurry, in an aqueous or nonaqueous liquid, containing a predetermined concentration of the mixture. An orally administered composition
10 can include one or more binders, lubricants, inert diluents, flavouring agents, and humectants. An orally administered composition such as a tablet can optionally be coated and can be formulated so as to provide sustained, delayed or controlled release of the HMO mixture therein.

A unit dosage form of this invention can also be administered by rectal suppository,
15 aerosol tube, naso-gastric tube or direct infusion into the GI tract or stomach.

A unit dosage form of this invention can also include therapeutic agents such as antiviral agents, antibiotics, probiotics, analgesics, and anti-inflammatory agents. The proper dosage of such a composition for a patient can be determined in a conventional manner, based upon factors such as the patient's immune status, body weight and age. In some
20 cases, the dosage will be at a concentration similar to that found for the HMOs of the composition in human breast milk. The required amount would generally be in the range from about 200 mg to about 20 g per day, in certain embodiments from about 300 mg to about 15 g per day, from about 400 mg to about 10 g per day, in certain embodiments from about 500 mg to about 10 g per day, in certain embodiments from about 1 g to about 10 g
25 per day. Appropriate dose regimes can be determined by methods known to those skilled in the art.

EXAMPLES

Example 1

A total of 50 healthy male and female subjects are recruited to participate in the study.
30 After a screening visit and run-in period of 1-2 weeks, the subjects are selected and randomized into 2 groups, each of 25 subjects. One group is administered a treatment product containing 5 g of the following HMO mixture of this invention:

- i) 14.2 wt % of LNT
- ii) 5.3 wt % of LNT

- iii) 63.7 wt % of 2'-FL
- iv) 4.2 wt % of DFL,
- v) 3.7 wt % of 3'-SL and
- vi) 8.9 wt % of 6'-SL

5 The other group is administered a placebo (containing 2 grams of glucose). The treatment product and the placebo are in powder form in a unit dosage container.

The subjects are eligible to participate if they are at least 18 years of age. All recruited subjects are able and willing to understand and comply with the study procedures. Subjects are excluded if: they have participated in a clinical study one month prior to screening visit;
10 they have abnormal results in the screening tests which are clinically relevant for study participation; they are suffering for a severe disease such as malignancy, diabetes, severe coronary disease, kidney disease, neurological disease, or severe psychiatric disease or any condition which can confound the results of the study; used highly dosed probiotic supplements (yoghurt allowed) for 3 months prior to the study; consumed antibiotic drugs 3
15 months prior to the study; consumed on a regular basis any medication that might interfere with symptom evaluation 2 weeks prior to the study; and pregnant or lactating.

At the screening visit, medical history and concomitant medication is registered and a blood sample for safety analyses is collected. A faecal sample kit is distributed. Subjects are instructed to keep their samples in the freezer until the next visit.

20 At the second visit, eligibility criteria are checked and eligible subjects are randomised to the two arms in the trial. The faecal samples are collected and equipment for new samples are distributed. Subjects are familiarised with an interactive internet enabled system which records data daily and are provided with either treatment or placebo products. Subjects are reminded not to change their usual diet during the study. Blood samples are
25 collected for biomarker studies. The faecal samples are stored at -80 °C until analysis. Faecal samples are subjected to 16S rRNA sequencing analysis.

The study runs for 8 weeks with the subjects consuming either a placebo or a treatment product daily. Subjects are instructed to consume the products in the morning with breakfast. Compliance is monitored through the interactive internet enabled system. The
30 subjects also use the system to record:

- Bristol Stool Form (BSF) scale information,
- symptom information such as abdominal pain, abdominal discomfort, abdominal cramping, abdominal bloating, and abdominal fullness,
- additional Gastrointestinal Symptom Rating Scale (GSRS) information.

This questionnaire includes 15 items covering five dimensions (abdominal pain, indigestion, reflux, diarrhoea, constipation) and uses a seven-graded Likert scale.

At the end of the study, each subject has an exit visit with the medical team. Faecal samples and blood samples are collected and analysed as before.

5 The faecal analysis indicates that the subjects treated with the HMO mixture of this invention have increased abundance of *Bifidobacterium* and *Barnesiella* and reduced abundance of Firmicutes, especially *Clostridia*, and *Ruminococcus gnavus*. The abundance of *Faecalibacterium* is unchanged in these subjects. The abundance of Proteobacteria is decreased in these subjects.

10 Example 2

A total of 50 healthy male and female subjects are recruited to participate in the study. After a screening visit and run-in period of 1-2 weeks, the subjects are selected and randomized into 2 groups, each of 25 subjects. One group is administered a treatment product containing 5 g of the following HMO mixture of this invention:

- 15 i) 14.2 wt % of LNT
 ii) 5.3 wt % of LNT
 iii) 63.7 wt % of 2'-FL
 iv) 4.2 wt % of 3-FL,
 v) 3.7 wt % of 3'-SL and
20 vi) 8.9 wt % of 6'-SL

The other group is administered a placebo (containing 2 grams of glucose). The treatment product and the placebo are in powder form in a unit dosage container.

The subjects are eligible to participate if they are at least 18 years of age. All recruited subjects are able and willing to understand and comply with the study procedures. Subjects
25 are excluded if: they have participated in a clinical study one month prior to screening visit; they have abnormal results in the screening tests which are clinically relevant for study participation; they are suffering for a severe disease such as malignancy, diabetes, severe coronary disease, kidney disease, neurological disease, or severe psychiatric disease or any condition which can confound the results of the study; used highly dosed probiotic
30 supplements (yoghurt allowed) for 3 months prior to the study; consumed antibiotic drugs 3 months prior to the study; consumed on a regular basis any medication that might interfere with symptom evaluation 2 weeks prior to the study; and pregnant or lactating.

At the screening visit, medical history and concomitant medication is registered and a blood sample for safety analyses is collected. A faecal sample kit is distributed. Subjects are
35 instructed to keep their samples in the freezer until the next visit.

At the second visit, eligibility criteria are checked and eligible subjects are randomised to the two arms in the trial. The faecal samples are collected and equipment for new samples are distributed. Subjects are familiarised with an interactive internet enabled system which records data daily and are provided with either treatment or control products. Subjects are reminded not to change their usual diet during the study. Blood samples are collected for biomarker studies. The faecal samples are stored at -80° C until analysis. Faecal samples are subjected to 16S rRNA sequencing analysis.

The study runs for 8 weeks with the subjects consuming either a placebo or a treatment product daily. Subjects are instructed to consume the products in the morning with breakfast. Compliance is monitored through the interactive internet enabled system. The subjects also use the system to record:

- Bristol Stool Form (BSF) scale information,
- symptom information such as abdominal pain, abdominal discomfort, abdominal cramping, abdominal bloating, and abdominal fullness,
- additional Gastrointestinal Symptom Rating Scale (GSRS) information.

This questionnaire includes 15 items covering five dimensions (abdominal pain, indigestion, reflux, diarrhoea, constipation) and uses a seven-graded Likert scale.

At the end of the study, each subject has an exit visit with the medical team. Faecal samples and blood samples are collected and analysed as before.

The faecal analysis indicates that the subjects treated with the HMO mixture of this invention have increased abundance of *Bifidobacterium* and *Barnesiella* and reduced abundance of Firmicutes, especially *Clostridia*, and *Ruminococcus gnavus*. The abundance of *Faecalibacterium* is unchanged in these subjects. The abundance of Proteobacteria is decreased in these subjects.

Example 3

Twenty 7-week-old C57BL/6J female mice are individually housed to avoid contamination between mice and provided with irradiated food and water. The mice are separated into 2 groups, each of 10 mice.

The mice are treated with ampicillin (0.5 g/liter) in their drinking water, which is changed every 3 days. After 1 week, the ampicillin addition to the drinking water is terminated. Thereafter, 1 group is administered a treatment product containing the following HMO mixture of this invention:

14.2 wt % of LNT

5.3 wt % of LNT

63.7 wt % of 2'-FL

4.2 wt % of DFL,
3.7 wt % of 3'-SL and
8.9 wt % of 6'-SL.

The treatment product is added to the drinking water of 1 group at a total concentration of 40 mg/ml. The other group receives drinking water with 40 mg/ml of glucose. Fresh water is administered daily, and all mice have free access to the drinking water. The mice are fed a rodent chow and are given fresh chow daily.

Two days after termination of the ampicillin treatment, mice of each group are infected by oral gavage with a vancomycin-resistant *Enterococcus faecium* strain (VRE). Fresh faecal pellets are collected at different time points to determine the VRE levels. VRE is quantified by plating serial dilutions of faecal pellets on Enterococcosel agar plates with vancomycin. VRE colonies are identified by appearance and confirmed by Gram staining. PCR of the vanA gene, which confers resistance to vancomycin, is used to confirm the presence of VRE in infected mice.

The mice are monitored for 2 weeks and are then euthanized. Luminal contents from the ileum, cecum and colon are collected and immediately frozen and stored at -80 °C. DNA is extracted using a 96-well PowerSoil DNA Isolation Kit (MO-BIO). A minimum of one sample-well per plate is kept empty to serve as a negative control during PCR. PCR is done with the forward primer S-D-Bact-0341-b-S-17 and reverse primer S-D-Bact-0785-a-A-21 (Klindworth et al. *Nucleic Acids Res.* **41**, e1 (2013)) with Illumina adapters attached. These are universal bacterial 16S rDNA primers, which target the V3-V4 region. The following PCR program is used: 98 °C for 30 sec, 25x (98 °C for 10 s, 55 °C for 20 s, 72 °C for 20 s), 72 °C for 5 min. Amplification is verified by running the products on a 1% agarose gel. Barcodes are added in a nested PCR using the Nextera Index Kit V2 (Illumina) with the following PCR program: 98 °C for 30 sec, 8x (98 °C for 10 s, 55 °C for 20 s, 72 °C for 20 s), 72 °C for 5 min. Attachment of primers is verified by running the products on a 1 % agarose gel.

Products from the nested PCR are normalized using the SequelPrep Normalization Plate Kit and pooled. Pooled libraries are concentrated by evaporation and the DNA concentration of pooled libraries was measured on a Qubit fluorometer using the Qubit High Sensitivity Assay Kit (Thermo Fisher Scientific). Sequencing is done on a MiSeq desktop sequencer using the MiSeq Reagent Kit V3 (Illumina) for 2x 300 bp paired-end sequencing. The 64-bit version of USEARCH (Edgar, 2013) is used for bioinformatical analysis of the sequence data.

In the mice treated with the HMO mixture of this invention, VRE colonisation is reduced to undetectable levels within 14 days. The density of VRE reduces within 5 days. The mice treated with the HMO mixture also showed a higher abundance of

Porphyromonadaceae, especially *Barnesiella*. The untreated mice continue to harbour large numbers of VRE throughout the intestine.

Example 4

Twenty 7-week-old C57BL/6J female mice are individually housed to avoid
5 contamination between mice and provided with irradiated food and water. The mice are separated into 2 groups, each of 10 mice.

The mice are treated with ampicillin (0.5 g/liter) in their drinking water, which is changed every 3 days. After 1 week, the ampicillin addition to the drinking water is terminated. Thereafter, 1 group is administered a treatment product containing the following

10 HMO mixture of this invention:

14.2 wt % of LNT

5.3 wt % of LNT

63.7 wt % of 2'-FL

4.2 wt % of 3'-FL,

15 3.7 wt % of 3'-SL and

8.9 wt % of 6'-SL.

The treatment product is added to the drinking water of 1 group at a total concentration of 40 mg/ml. The other group receives drinking water with 40 mg/ml of glucose. Fresh water is administered daily, and all mice have free access to the drinking water. The mice are fed a
20 rodent chow and are given fresh chow daily.

Two days after termination of the ampicillin treatment, mice of each group are infected by oral gavage with a vancomycin-resistant *Enterococcus faecium* strain (VRE). Fresh faecal pellets are collected at different time points to determine the VRE levels. VRE is quantified by plating serial dilutions of faecal pellets on Enterococcosel agar plates with vancomycin.

25 VRE colonies are identified by appearance and confirmed by Gram staining. PCR of the vanA gene, which confers resistance to vancomycin, is used to confirm the presence of VRE in infected mice.

The mice are monitored for 2 weeks and are then euthanized. Luminal contents from the ileum, cecum and colon are collected and immediately frozen and stored at -80 °C. DNA
30 is extracted using a 96-well PowerSoil DNA Isolation Kit (MO-BIO). A minimum of one sample-well per plate is kept empty to serve as a negative control during PCR. PCR is done with the forward primer S-D-Bact-0341-b-S-17 and reverse primer S-D-Bact-0785-a-A-21 (Klindworth et al. *Nucleic Acids Res.* **41**, e1 (2013)) with Illumina adapters attached. These are universal bacterial 16S rDNA primers, which target the V3-V4 region. The following PCR
35 program is used: 98 °C for 30 sec, 25x (98 °C for 10 s, 55 °C for 20 s, 72 °C for 20 s), 72 °C

for 5 min. Amplification is verified by running the products on a 1% agarose gel. Barcodes are added in a nested PCR using the Nextera Index Kit V2 (Illumina) with the following PCR program: 98 °C for 30 sec, 8x (98 °C for 10 s, 55 °C for 20 s, 72 °C for 20 s), 72 °C for 5 min. Attachment of primers is verified by running the products on a 1 % agarose gel.

5 Products from the nested PCR are normalized using the SequalPrep Normalization Plate Kit and pooled. Pooled libraries are concentrated by evaporation and the DNA concentration of pooled libraries was measured on a Qubit fluorometer using the Qubit High Sensitivity Assay Kit (Thermo Fisher Scientific). Sequencing is done on a MiSeq desktop sequencer using the MiSeq Reagent Kit V3 (Illumina) for 2x 300 bp paired-end
10 sequencing. The 64-bit version of USEARCH (Edgar, 2013) is used for bioinformatical analysis of the sequence data.

In the mice treated with the HMO mixture of this invention, VRE colonisation is reduced to undetectable levels within 14 days. The density of VRE reduces within 5 days. The mice treated with the HMO mixture also showed a higher abundance of

15 *Porphyromonadaceae*, especially *Barnesiella*. The untreated mice continue to harbour large numbers of VRE throughout the intestine.

Example 5

Forty-five 6-week-old female C57BL/6 mice are injected subcutaneously with B16 melanoma cells. The mice are then randomly assigned to three groups (fifteen mice in each
20 group), and individually housed to avoid contamination between mice. The mice are fed a pelleted semi-synthetic AIN 76 diet (Research Diets Inc., New Brunswick, NJ) with: 5% of the following HMO mixture of this invention in the feed for group A:

14.2 wt % of LNT

5.3 wt % of LNT

25 63.7 wt % of 2'-FL

4.2 wt % of DFL,

3.7 wt % of 3'-SL and

8.9 wt % of 6'-SL;

5% of the following HMO mixture of this invention in the feed for group B:

30 14.2 wt % of LNT

5.3 wt % of LNT

63.7 wt % of 2'-FL

4.2 wt % of 3-FL,

3.7 wt % of 3'-SL and

35 8.9 wt % of 6'-SL; and

5% corn-starch in the feed for group C (control).

Fresh water is administered daily and all mice have free access to drinking water and feed. Three days after the tumor implantation, mice are injected with 200 µg of CTLA-4 antibodies. Additional antibody treatments are given every 3 days until the end of the experiment (14 days). Food consumption, bodyweight and tumor volume (diameter) are measured throughout the study. Fresh faecal samples are collected at day 0, 4, 7, 10, and 14. Samples are immediately frozen and stored at -80 °C until further analysis.

Fourteen days after implantation of the B16 cells, mice are euthanized by cervical dislocation. Tumors are excised and weighed. Sections of tumors are fixed in 4% paraformaldehyde and stored at -80 °C. Mucosal and intestinal contents from caecum and colon are removed and stored at -80 °C.

For analysis of tumor infiltrating T cells, tumors are minced into small pieces and digested with 0.2 mg/ml DNase and 1.67 Wünsch U/ml Liberase (Roche). Obtained cell suspensions are filtered through a 40-µm nylon cell strainer and red blood cells lysed. CD8⁺ T cells are purified from B16 tumor-derived cell suspensions using CD8α (Ly-2) MicroBeads (Miltenyi Biotec), following the manufacturer's protocol. Obtained cells are incubated for 2 hours in the presence of 1 µl/ml of brefeldin A, washed and incubated with rat anti-mouse CD16/CD32 mAb (2.4G2) to block nonspecific binding, and then stained with CD8α-PE-Cy5 and CD69-PE, followed by intracellular staining with IFN-γ-PE-Cy7 or isotype control antibodies according to the manufacturer's instructions (BD Biosciences — Pharmingen). The density of CD8 GFP⁺ and GFP⁻ T cells in the tumors are calculated by dividing the total number of obtained cells by the tumor weight.

To assess the microbiota profile, DNA is extracted from the faecal samples, mucosal and intestinal content using a 96-well PowerSoil DNA Isolation Kit (MO-BIO). A minimum of one sample-well per plate is kept empty to serve as a negative control during PCR. PCR is done with the forward primer S-D-Bact-0341-b-S-17 and reverse primer S-D-Bact-0785-a-A-21 (Klindworth et al. Nucleic Acids Res. 41, e1 (2013)) with Illumina adapters attached. These are universal bacterial 16S rDNA primers, which target the V3-V4 region. The following PCR program is used: 98 °C for 30 sec, 25x (98° C for 10 s, 55 °C for 20 s, 72 °C for 20 s), 72 °C for 5 min. Amplification is verified by running the products on a 1 % agarose gel. Barcodes are added in a nested PCR using the Nextera Index Kit V2 (Illumina) with the following PCR program: 98 °C for 30 sec, 8x (98 °C for 10 s, 55 °C for 20 s, 72 °C for 20 s), 72 °C for 5 min. Attachment of primers is verified by running the products on a 1 % agarose gel. Products from the nested PCR are normalized using the SequalPrep Normalization Plate Kit and pooled. Pooled libraries are concentrated by evaporation and the DNA concentration of pooled libraries is measured on a Qubit fluorometer using the Qubit High

Sensitivity Assay Kit (Thermo Fisher Scientific). Sequencing is done on a MiSeq desktop sequencer using the MiSeq Reagent Kit V3 (Illumina) for 2 x 300 bp paired-end sequencing. The 64-bit version of USEARCH (Edgar, 2013) is used for bioinformatical analysis of the sequence data.

5 The results show that ingestion of both HMO mixtures beneficially modulates the microbiota by increasing the abundance of bifidobacteria both in the mucosal and luminal intestinal environment. Additionally, the results show that the abundance of bifidobacteria positively correlates with the priming and accumulation of infiltrating T cells in the tumor microenvironment and negatively correlates with tumor size. Collectively, these results
10 indicate that both HMO mixtures can increase the abundance of bifidobacteria and through this increase the efficacy of anti-cancer agents against tumor outgrowth.

Example 6

Mice (5 weeks old; n=30) are randomly allocated to one of the following groups: A) control (n=10); B) 1% mixture of HMOs (n=10); C) 5% mixture of HMOs (n=10). The
15 composition of the HMO blend is given in the table below.

	g/100 ml	
	1%	5%
2'-FL (g)	0.64	3.20
LNT (g)	0.05	0.25
3'-SL (g)	0.05	0.25
6'-SL (g)	0.09	0.45
DFL (g)	0.05	0.25

The HMOs are provided in drinking water. All animals have free access to same diet (KLIBA 2122).

After 2 weeks, the mice are challenged with Influenza strain PR8 at a dose of 100
20 PFU per mouse by intranasal inoculation. The mice are monitored for the next 14 days to assess clinical score of illness symptoms and body weight loss.

The results show that the HMO mixture protects against sickness symptoms and body weight loss caused by the influenza virus infection.

CLAIMS

1. A synthetic mixture of HMOs consisting essentially of LNnT, LNT, 2'-FL, 3'-SL, 6'-SL, either DFL or 3-FL, preferably DFL, and optionally lactose.
- 5 2. The synthetic mixture of claim 1 that consists essentially of:
 - i. about 55 wt % to about 75 wt %, preferably about 60 wt % to about 70 wt %, of 2'-FL ;
 - ii. about 2 wt % to about 10 wt %, preferably about 3 wt % to about 7 wt %, of LNnT;
 - 10 iii. about 10 wt % to about 20 wt %, preferably about 12 wt % to about 18 wt %, of LNT;
 - iv. about 1 wt % to about 15 wt %, preferably about 2 wt % to about 10 wt %, of DFL or 3-FL;
 - v. about 1 wt % to about 10 wt %, preferably about 2 wt % to about 8 wt %, of 3'-SL; and
 - 15 vi. about 1 wt % to about 15 wt %, preferably about 5 wt % to about 15 wt %, more preferably about 7 wt % to about 13 wt %, of 6'-SL.
3. The synthetic mixture of claim 1 or 2 for use in: i) preventing and/or treating viral and/or bacterial infections in a non-infant human; ii) modulating the microbiota of a non-
 20 infant human; and/or iii) improving the cognitive function of a non-infant human.
4. A pharmaceutical or nutritional composition comprising a synthetic mixture of claim 1 or 2.
5. The composition of claim 4 for use in: i) preventing and/or treating viral and/or bacterial infections in a non-infant human; ii) modulating the microbiota of a non-infant
 25 human; and/or iii) improving the cognitive function of a non-infant human.
6. A method of modulating the microbiota of a non-infant human to increase the abundance of *Bifidobacterium* in order to increase the efficacy of anticancer agents against tumors in a non-infant human cancer patient, the method comprising:
 - administering, to the non-infant human, a synthetic mixture of claim 1 or 2, or a
 30 nutritional or pharmaceutical composition of claim 4.

7. A method of modulating the microbiota of a non-infant human to increase the abundance of *Bifidobacterium* and/or *Barnesiella* and to reduce the abundance of *Ruminococcus gnavus*, and preferably also to reduce the abundance of Proteobacteria, the method comprising:

- 5 - administering, to the non-infant human, a synthetic mixture of claim 1 or 2, or a nutritional or pharmaceutical composition of claim 4.

8. A method of modulating the microbiota of a non-infant human to increase *Bifidobacterium* abundance and to at least maintain the abundance of *Faecalibacterium*, and preferably also to reduce the abundance of Proteobacteria, the method comprising:

- 10 - administering, to the human, a synthetic mixture of claim 1 or 2, or a nutritional or pharmaceutical composition of claim 4.

9. A method of preventing or treating viral and/or bacterial infections in a non-infant human, especially intestinal infections and infections of the respiratory tract, the method comprising:

- 15 - administering, to the non-infant human, a mixture of claim 1 or 2, or a nutritional or pharmaceutical composition of claim 4.

10. A method of improving the cognitive function of a non-infant human, the method comprising:

- 20 - administering, to the non-infant human, a mixture of claim 1 or 2, or a nutritional or pharmaceutical composition of claim 4.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2016/057675

A. CLASSIFICATION OF SUBJECT MATTER A61K 31/702 (2006.01), A61K 31/7012 (2006.01), A61P 31/04 (2006.01), A61P 31/12 (2006.01)		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) IPC & CPC: A61K, A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched DK, NO, SE, FI: Classes as above.		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) EPODOC, WPI, FULL TEXT: ENGLISH, GERMAN; NPL, MEDLINE, XPESP, XPOAC		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 2013/185780 A1 (GLYCOM A/S) 2013.12.19 Claims 11-17; Page 4, lines 12-23; Page 8, lines 14-20; Page 10, lines 14-21; Page 11, lines 5-23; Example 5	1-5
X	EP 2465509 A1 (NESTEC S.A.) 2012.06.20 Claims 1-4, 9, 11-12; Paragraphs [0003], [0023], [0074]	1-5
X	WO 2015/071402 A1 (NESTEC S.A.) 2015.05.21 Claims 1-2, 5-7, 9, 14 ; Page 4, lines 17-18; Page 9, lines 18-29; Page 18, lines 13-14	1-5
A	EP 2768314 B1 (NESTEC S.A.) 2015.11.25 Claims 1-4	-
☐ Further documents are listed in the continuation of Box C. ☒ See patent family annex.		
* Special categories of cited documents: “A” document defining the general state of the art which is not considered to be of particular relevance “E” earlier application or patent but published on or after the international filing date “L” document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) “O” document referring to an oral disclosure, use, exhibition or other means “P” document published prior to the international filing date but later than the priority date claimed “T” later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention “X” document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone “Y” document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art “&” document member of the same patent family		
Date of the actual completion of the international search 19/01/2017		Date of mailing of the international search report 20/01/2017
Name and mailing address of the ISA Nordic Patent Institute Helgeshøj Allé 81 DK - 2630 Taastrup, Denmark. Facsimile No. + 45 43 50 80 08		Authorized officer E-ri Maria Sol Telephone No. +45 43 50 85 88

INTERNATIONAL SEARCH REPORT

International application No.

PCT/IB2016/057675

Box No. II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 6-10

because they relate to subject matter not required to be searched by this Authority, namely:

The subject matter of claim 6 concerns a method of modulating the microbiota of a non-infant human to increase the abundance of *Bifidobacterium*, wherein said method comprises administering to the non-infant human, a synthetic mixture, nutritional or a pharmaceutical composition. As the subject matter of claim 6 concerns a method for treatment of the human body by therapy, said subject matter of claim 6 is excluded from search and examination under PCT Rules 39.1 and 67.1.

The same argument applies mutatis mutandis, for the subject matter of claims 7-10.

2. ☐ Claims Nos.:

because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:

because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box No. III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.

2. ☐ As all searchable claims could be searched without effort justifying additional fees, this Authority did not invite payment of additional fees.

3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:

4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest and, where applicable, the payment of a protest fee.

☐ The additional search fees were accompanied by the applicant's protest but the applicable protest fee was not paid within the time limit specified in the invitation.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT
Information on patent family members

International application No.

PCT/IB2016/057675

Patent document cited in search report / Publication date	Patent family member(s) / Publication date
WO 2013185780 A1 2013.12.19	CN 104428307 A 2015.03.18 EP 2861609 A1 2015.04.22 EP 2861609 A4 2015.11.04 US 2015183814 A1 2015.07.02
EP 2465509 A1 2012.06.20	CA 2818012 A1 2012.06.14 WO 2012076323 A1 2012.06.14 TW 201302205 A 2013.01.16 AU 2011340882 A1 2013.05.30 AU 2011340882B B2 2016.10.20 SG 190323 A1 2013.06.28 MX 2013005727 A 2013.07.03 US 2013236424 A1 2013.09.12 EP 2643006 A1 2013.10.02 CL 2013001433 A1 2013.12.13 CN 103582485 A 2014.02.12 RU 2013128570 A 2014.12.27 RU 2593321 C2 2016.08.10 BR 112013012598 A2 2016.08.09
WO 2015071402 A1 2015.05.21	AU 2014350157 A1 2016.04.14 CN 105744938 A 2016.07.06 EP 3071208 A1 2016.09.28 US 2016296540 A1 2016.10.13
EP 2768314 A1 2014.08.27	WO 2013057061 A1 2013.04.25 AU 2012325072 A1 2014.03.13 AU 2012325072 A2 2014.05.29 CN 103889240 A 2014.06.25 CN 103889240B B 2016.09.14 CL 2014000832 A1 2014.08.22 EP 2768314 B1 2015.11.25 US 2014286909 A1 2014.09.25 MX 2014004724 A 2015.02.05 RU 2014119903 A 2015.11.27 PT 2768314E E 2016.02.17 ES 2560023T T3 2016.02.17