

(19) **DANMARK**



Patent- og
Varemærkestyrelsen

(10) **DK/EP 3551633 T3**

(12) Oversættelse af
europæisk patentskrift

-
- (51) Int.Cl.: **C 07 D 491/20 (2006.01)** **A 61 K 31/438 (2006.01)** **A 61 P 17/00 (2006.01)**
- (45) Oversættelsen bekendtgjort den: **2021-06-28**
- (80) Dato for Den Europæiske Patentmyndigheds
bekendtgørelse om meddelelse af patentet: **2021-03-31**
- (86) Europæisk ansøgning nr.: **16809388.8**
- (86) Europæisk indleveringsdag: **2016-12-12**
- (87) Den europæiske ansøgnings publiceringsdag: **2019-10-16**
- (86) International ansøgning nr.: **EP2016080548**
- (87) Internationalt publikationsnr.: **WO2018108230**
- (84) Designerede stater: **AL AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HR HU IE IS IT LI LT LU LV
MC MK MT NL NO PL PT RO RS SE SI SK SM TR**
- (73) Patenthaver: **UNION therapeutics A/S, Tuborg Havnevej 18, 2900 Hellerup, Danmark**
- (72) Opfinder: **LIANG, Xifu, c/o LEO Pharma A/S, Industriparken 55, 2750 Ballerup, Danmark**
LARSEN, Jens, c/o LEO Pharma A/S, Industriparken 55, 2750 Ballerup, Danmark
ANDERSEN, Peter, c/o LEO Pharma A/S, Industriparken 55, 2750 Ballerup, Danmark
NIELSEN, Simon Feldbæk, c/o LEO Pharma A/S, Industriparken 55, 2720 Vanløse, Danmark
- (74) Fuldmægtig i Danmark: **NORDIC PATENT SERVICE A/S, Bredgade 30, 1260 København K, Danmark**
- (54) Benævnelse: **SUBSTITUERED E PYRAZOLOAZEPIN-4-ONER OG ANVENDELSE DERA F SOM
PHOSPHODIESTERASEHÆMMERE**
- (56) Fremdragne publikationer:
WO-A1-2007/040435
WO-A2-2008/104175

Description

FIELD OF THE INVENTION

5 **[0001]** The present invention relates to novel substituted pyrazoloazepin-4-ones with phosphodiesterase inhibitory activity, and to their use as therapeutic agents in the treatment of inflammatory diseases and conditions.

BACKGROUND OF THE INVENTION

10 **[0002]** Phosphodiesterases are enzymes that catalyse the hydrolysis of cyclic AMP and/or cyclic GMP in cells to 5-AMP and 5-GMP, respectively, and as such they are critical to cellular regulation of cAMP or cGMP levels. Of the 11 phosphodiesterases identified so far, phosphodiesterase (PDE) 4, PDE7 and PDE8 are selective for cAMP. PDE4 is the most important modulator of cAMP expressed in immune and inflammatory cells such as neutrophils, macrophages and T-lymphocytes. As cAMP is a key second messenger in the modulation of inflammatory responses, PDE4 has been
 15 found to regulate inflammatory responses of inflammatory cells by modulating proinflammatory cytokines such as TNF- α , IL-2, IFN- γ , GM-CSF and LTB₄. Inhibition of PDE4 has therefore become an attractive target for the therapy of inflammatory diseases such as asthma, chronic obstructive pulmonary disease (COPD), rheumatoid arthritis, atopic dermatitis, psoriasis, inflammatory bowel disease such as Crohn's disease etc. (M.D. Houslay et al., Drug Discovery Today 10 (22), 2005, pp. 1503-1519). As atopic dermatitis (AD) patients have increased PDE-activity, PDE4-inhibition
 20 would also appear to be a viable treatment of AD (Journal of Investigative Dermatology (1986), 87(3), 372-6).

[0003] The PDE4 gene family consists at least of four genes, A, B, C and D, which have a high degree of homology (V. Boswell Smith and D. Spina, Curr. Opinion Investig. Drugs 6(11), 2006, pp. 1136-1141). The four PDE4 isoforms are differentially expressed in different tissues and cell types. Thus, PDE4B is predominantly expressed in monocytes and neutrophils, but not in cortex and epithelial cells, while PDE4D is expressed in lung, cortex, cerebellum and T-cells
 25 (C. Kroegel and M. Foerster, Exp. Opinion Investig. Drugs 16(1), 2007, pp. 109-124). It has been speculated that inhibition of PDE4D in the brain is associated with the adverse effects found when administering PDE4 inhibitors clinically, primarily nausea and emesis, whereas inhibition of PDE4B is associated with anti-inflammatory effects (B. Lipworth, Lancet 365, 2005, pp. 167-175).

[0004] Numerous PDE4 inhibitors have been studied for their therapeutic effect on inflammatory diseases, primarily
 30 asthma and COPD.

[0005] WO 2007/040435 (Astrazeneca AB) discloses 5,6-dihydropyrazolo[3,4-e][1,4]-diazepin-4(1H)-one derivatives for the treatment of asthma and chronic obstructive pulmonary disease. The compounds are stated to be selective inhibitors of PDE4 over other PDEs.

[0006] WO 2001/049689 (Warner-Lambert Company) discloses pyrazolo[3,4-e]diazepines. The compounds are stated
 35 to inhibit the PDE4 enzyme.

[0007] WO2008/060597 (Vertex Pharmaceuticals Inc) relates to compounds as protein kinase inhibitors.


[0008] WO2004/098520 (IRM LLC) relates to compounds as protein kinase inhibitors.

[0009] There is a continuous need for developing novel PDE4 inhibitors which have a more favourable therapeutic window, i.e. fewer adverse effects, while retaining their therapeutic effect.

SUMMARY OF THE INVENTION

[0010] It is an object of the present invention to provide novel compounds. In one aspect the present invention relates
 45 to PDE4 inhibitors that could have a stability profile in biological tissue that implies that a low systemic exposure of the compounds to be observed upon e.g. topical administration, indicating that the compounds of the present invention could have high clearance in human liver microsomes, that they could hydrolyse in human whole blood and could display stability towards enzymatic hydrolyses in human keratinocytes.

[0011] In one aspect the invention provides a compound of general formula (I)



(I)

wherein

R_1 and R_4 are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl;

5 R_2 and R_3 are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl; or

R_2 and R_3 together with the carbon atom to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, tetrahydrofuranyl, or tetrahydropyranyl ring;

10 $n = 0, 1$ or 2 ;

Q is -O-C(O)- R_5 ;

15 R_5 is selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from R_6 , or wherein said aryl is optionally benzodioxole; and wherein said (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl are optionally substituted with one or more substituents independently selected from R_7 ;

20 R_6 consists of halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂ R_x , -S(O)₂ NR_aR_b , -C(O) R_x , -C(O) NR_aR_b and -OR_x;

R_7 consists of halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O) NR_aR_b , -C(O)OR_a, -S(O)₂ R_x , -S(O)₂ NR_aR_b , -OR_x, and -SR_x;

25 R_x consists of (C₁-C₄)alkyl and (C₃-C₆)cycloalkyl;

R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, or

30 R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)alkyl or -C(O) R_x ; or

pharmaceutically acceptable salts, hydrates or solvates thereof.

35 **[0012]** In another aspect, the invention provides pharmaceutical compositions comprising a compound of general formula (I) as defined above together with a pharmaceutically acceptable vehicle or excipient or pharmaceutically acceptable carrier(s), optionally together with one or more other therapeutically active compound(s).


[0013] In another aspect, the invention provides the use of a compound of the invention, for the manufacture of pharmaceutical compositions for the prophylaxis, treatment, prevention or amelioration of a disease, disorder or condition responsive to PDE4 inhibitory activity.

40 **[0014]** In yet another aspect the invention provides a method for treatment, prevention or alleviation of diseases, disorders or conditions responsive to PDE4 inhibitory activity, and which method comprises the step of administering to a living animal body a therapeutically effective amount of the compound of formula (I) of the invention. Other objects of the invention will be apparent to the person skilled in the art from the following detailed description and examples.

45 DETAILED DESCRIPTION OF THE INVENTION

[0015] In one aspect the invention provides a compound of Formula (I)

50



55

(I)

wherein

R₁ and R₄ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl;

5 R₂ and R₃ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl; or

R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, tetrahydrofuranyl, or tetrahydropyranyl ring; n = 0, 1 or 2;

10 Q is -O-C(O)-R₅;

R₅ is selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from R₆, or wherein said aryl is benzodioxole; and wherein said (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl are optionally substituted with one or more substituents independently selected from R₇; R₆ consists of halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, C(O)NR_aR_b and -OR_x;

R₇ consists of halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O)NR_aR_b, -C(O)OR_a, -S(O)₂R_x, -S(O)₂NR_aR_b, -OR_x, and -SR_x;

20 R_x consists of (C₁-C₄)alkyl and (C₃-C₆)cycloalkyl;

R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, or

25 R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)alkyl or -C(O)R_x; and

pharmaceutically acceptable salts, hydrates or solvates thereof.

30 **[0016]** In another aspect the invention provides a compound of Formula (I) wherein R₁ and R₄ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl;

R₂ and R₃ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl; or

R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, tetrahydrofuranyl, or tetrahydropyranyl ring; n = 0, 1 or 2;

35 Q is -O-C(O)-R₅;

R₅ is selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from R₆; and wherein said (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl are optionally substituted with one or more substituents independently selected from R₇;

40 R₆ consists of halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, C(O)NR_aR_b and -OR_x;

R₇ consists of halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O)NR_aR_b, -C(O)OR_a, -S(O)₂R_x, -S(O)₂NR_aR_b, -OR_x, and -SR_x;

R_x consists of (C₁-C₄)alkyl and (C₃-C₆)cycloalkyl;

45 R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, or

R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)alkyl or -C(O)R_x; or

a pharmaceutically acceptable salt, hydrate or solvate thereof.

[0017] In another aspect the invention provides a compound of Formula (I) wherein

50 R₁ and R₄ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl;

R₂ and R₃ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl; or

R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl or oxetanyl ring; n = 0, 1 or 2;

Q is -O-C(O)-R₅;

55 R₅ is selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from R₆, or wherein said aryl is optionally benzodioxole; and wherein said (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl are optionally substituted with one or more substituents independently selected from R₇;

R₆ consists of halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, C(O)NR_aR_b and -OR_x;

R₇ consists of halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O)NR_aR_b, -C(O)OR_a, -S(O)₂R_x, -S(O)₂NR_aR_b, -OR_x, and -SR_x;

5 R_x consists of (C₁-C₄)alkyl and (C₃-C₆)cycloalkyl;

R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, or

R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)alkyl or -C(O)R_x; or

a pharmaceutically acceptable salts, hydrates or solvates thereof.

10 **[0018]** In another aspect the invention provides a compound of Formula (I) wherein

R₁ and R₄ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl;

R₂ and R₃ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl; or

R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl or oxetanyl ring;

n = 0, 1 or 2;

15 Q is -O-C(O)-R₅;

R₅ is selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, phenyl; wherein said phenyl is optionally substituted with one or more substituents independently selected from R₆; and wherein said (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl are optionally substituted with one or more substituents independently selected from R₇;

20 R₆ consists of halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, C(O)NR_aR_b and -OR_x;

R₇ consists of halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O)NR_aR_b, -C(O)OR_a, -S(O)₂R_x, -S(O)₂NR_aR_b, -OR_x, and -SR_x;

R_x consists of (C₁-C₄)alkyl and (C₃-C₆)cycloalkyl;

25 R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, or

R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)alkyl or -C(O)R_x; or

a pharmaceutically acceptable salt, hydrate or solvate thereof.

[0019] In another aspect the invention provides a compound of Formula (I) wherein

30 R₁ and R₄ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl;

R₂ and R₃ are independently selected from the group consisting of hydrogen and (C₁-C₄)alkyl; or

R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, tetrahydrofuranyl, or tetrahydropyranyl ring; n = 0, 1 or 2;

Q is -O-C(O)-R₅;

35 R₅ is selected from the group consisting of (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from R₆;

R₆ consists of halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b and -OR_x;

40 R_x is (C₁-C₄)alkyl;

R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, or

R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)alkyl; and

pharmaceutically acceptable salts, hydrates or solvates thereof.

45 **[0020]** In one embodiment of the present invention, R₁ and R₄ are both hydrogen.

[0021] In another embodiment of the present invention, one of R₂ and R₃ is hydrogen and the other one of R₂ and R₃ is (C₁-C₄)alkyl.

[0022] In another embodiment of the present invention, R₂ and R₃ are both hydrogen.

[0023] In another embodiment of the present invention, R₂ and R₃ are both (C₁-C₄)alkyl, e.g. both methyl.

50 **[0024]** In another embodiment of the present invention, R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl, cyclobutyl, cyclopentyl, oxetanyl, tetrahydrofuranyl, or tetrahydropyranyl ring.

[0025] In another embodiment of the present invention, R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl or oxetanyl ring.

[0026] In another embodiment of the present invention, n is 1.

55 **[0027]** In another embodiment of the present invention, R₅ is (C₁-C₆)alkyl.

[0028] In another embodiment of the present invention, R₅ is (C₃-C₆)cycloalkyl, e.g. cyclopentyl.

[0029] In another embodiment of the present invention, R₅ is aryl, optionally substituted with one or more substituents independently selected from R₆.

[0030] In another embodiment of the present invention, R_5 is phenyl, optionally substituted with one or more substituents independently selected from R_6 .

[0031] In another embodiment of the present invention, R_5 is phenyl, substituted with one substituent selected from R_6 .

[0032] In another embodiment of the present invention, R_5 is phenyl, substituted with two substituents independently selected from R_6 .

[0033] In another embodiment of the present invention, R_6 consists of halogen, cyano, hydroxyl, (C_1-C_4) alkyl, halo (C_1-C_4) alkyl and (C_1-C_4) alkyloxy.

[0034] In another embodiment of the present invention, R_6 is (C_1-C_4) alkyl, e.g. methyl, ethyl.

[0035] In another embodiment of the present invention, R_6 is $-S(O)_2R_x$.

[0036] In another embodiment of the present invention, R_x is methyl.

[0037] In another embodiment of the present invention, R_6 is $-C(O)R_a$.

[0038] In another embodiment of the present invention, R_6 is $C(O)NR_aR_b$.

[0039] In another embodiment of the present invention, R_6 is $-S(O)_2NR_aR_b$.

[0040] In another embodiment of the present invention R_7 consists of halogen, hydroxyl, (C_1-C_4) alkoxy, $-C(O)NR_aR_b$, $-C(O)OR_a$, $-S(O)_2R_x$, $-OR_x$, and $-SR_x$.

[0041] In another embodiment of the present invention, R_a and R_b are independently selected from the group consisting of hydrogen and (C_1-C_4) alkyl.

[0042] In another embodiment of the present invention, R_a and R_b are both (C_1-C_4) alkyl, e.g. both methyl.

[0043] In another embodiment of the present invention, when R_5 is substituted with one of R_6 consisting of $-S(O)_2R_x$ and $-S(O)_2NR_aR_b$, the substituent is in the para position.

[0044] In another embodiment of the present invention, R_1 and R_4 are both hydrogen, one of R_2 and R_3 is hydrogen and the other one of R_2 and R_3 is (C_1-C_4) alkyl, n is 1, R_5 is phenyl, R_6 is $-S(O)_2R_x$ or $-S(O)_2NR_aR_b$, R_x is (C_1-C_4) alkyl and R_a and R_b are independently selected from the group consisting of hydrogen and (C_1-C_4) alkyl.

[0045] In another embodiment of the present invention, all of R_1 , R_2 , R_3 , R_4 are hydrogen, n is 1, R_5 is phenyl, and R_6 is (C_1-C_4) alkyl.

[0046] In another embodiment of the present invention, all of R_1 , R_2 , R_3 , R_4 are hydrogen, n is 1, R_5 is selected from the group consisting of (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl, (C_1-C_6) alkoxy, and aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from halogen, cyano, hydroxyl, (C_1-C_4) alkyl, halo (C_1-C_4) alkyl, (C_1-C_4) alkyloxy, $-S(O)_2R_x$, $-S(O)_2NR_aR_b$, $-C(O)R_x$, $C(O)NR_aR_b$, or wherein said aryl is optionally benzodioxole; and wherein said (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl are optionally substituted with one or more substituents independently selected from halogen, hydroxyl, (C_1-C_4) alkoxy, $-C(O)NR_aR_b$, $-C(O)OR_a$, $-S(O)_2R_x$, $-OR_x$, and $-SR_x$; R_x consists of (C_1-C_4) alkyl and (C_3-C_6) cycloalkyl; R_a and R_b are independently selected from the group consisting of hydrogen, (C_1-C_4) alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C_1-C_4) alkyl or $-C(O)R_x$.

[0047] In another embodiment of the present invention, all of R_1 , R_2 , R_3 , R_4 are hydrogen, n is 1, R_5 is phenyl which is optionally substituted with one or more substituents independently selected from halogen, cyano, (C_1-C_4) alkyl, halo (C_1-C_4) alkyl, and (C_1-C_4) alkyloxy.

[0048] In another embodiment of the present invention, all of R_1 , R_2 , R_3 , R_4 are hydrogen, n is 1, R_5 is phenyl which is optionally substituted with one or more substituents independently selected from the group consisting of $-S(O)_2R_x$, $-S(O)_2NR_aR_b$, $-C(O)R_x$, $C(O)NR_aR_b$; R_x consists of (C_1-C_4) alkyl and (C_3-C_6) cycloalkyl; R_a and R_b are independently selected from the group consisting of hydrogen and (C_1-C_4) alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C_1-C_4) alkyl or $-C(O)R_x$.

[0049] In another embodiment of the present invention, all of R_1 , R_2 , R_3 , R_4 are hydrogen, n is 1, R_5 is phenyl substituted with $C(O)NR_aR_b$, wherein R_a and R_b are independently selected from the group consisting of hydrogen and (C_1-C_4) alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C_1-C_4) alkyl or $-C(O)R_x$; and R_x is (C_1-C_4) alkyl.

[0050] In another embodiment of the present invention, R_1 and R_4 are both hydrogen, one of R_2 and R_3 is hydrogen and the other one of R_2 and R_3 is (C_1-C_4) alkyl, n is 1, R_5 is selected from the group consisting of (C_1-C_6) alkyl, (C_3-C_6) cycloalkyl and aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from halogen, cyano, (C_1-C_4) alkyl, halo (C_1-C_4) alkyl, (C_1-C_4) alkyloxy, $-S(O)_2R_x$, $-S(O)_2NR_aR_b$ and $-C(O)R_x$; R_x consists of (C_1-C_4) alkyl and (C_3-C_6) cycloalkyl; R_a and R_b are independently selected from the group consisting of hydrogen, (C_1-C_4) alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C_1-C_4) alkyl or $-C(O)R_x$.

[0051] In another embodiment of the present invention, R_1 and R_4 are both hydrogen, one of R_2 and R_3 is hydrogen and the other one of R_2 and R_3 is (C_1-C_4) alkyl, n is 1, R_5 is phenyl which is optionally substituted with one or more substituents independently selected from halogen, cyano, (C_1-C_4) alkyl, halo (C_1-C_4) alkyl, (C_1-C_4) alkyloxy, $-S(O)_2R_x$, $-S(O)_2NR_aR_b$, and $-C(O)R_x$; R_x consists of (C_1-C_4) alkyl and (C_3-C_6) cycloalkyl; R_a and R_b are independently selected

from the group consisting of hydrogen, (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)-alkyl or -C(O)R_x.

[0052] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, one of R₂ and R₃ is hydrogen and the other one of R₂ and R₃ is (C₁-C₄)-alkyl, n is 1, R₅ is phenyl substituted with C(O)NR_aR_b, wherein R_a and R_b are independently selected from the group consisting of hydrogen and (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)-alkyl or -C(O)R_x; and R_x is (C₁-C₄)-alkyl.

[0053] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, one of R₂ and R₃ is hydrogen and the other one of R₂ and R₃ is (C₁-C₄)-alkyl, n is 1, R₅ is phenyl optionally substituted with halogen, (C₁-C₄)-alkyl and (C₁-C₄)-alkyloxy.

[0054] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, R₂ and R₃ are both (C₁-C₄)-alkyl, n is 1, R₅ is selected from the group consisting of (C₁-C₆)-alkyl, (C₃-C₆)-cycloalkyl and aryl; wherein said aryl is optionally substituted with one or more substituents independently selected from halogen, cyano, (C₁-C₄)-alkyl, halo(C₁-C₄)-alkyl, (C₁-C₄)-alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, C(O)NR_aR_b and -OR_x; R_x consists of (C₁-C₄)-alkyl and (C₃-C₆)-cycloalkyl; R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more of -C(O)R_x.

[0055] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, R₂ and R₃ are both (C₁-C₄)-alkyl, n is 1, R₅ is phenyl which is optionally substituted with one or more substituents independently selected from halogen, cyano, (C₁-C₄)-alkyl, halo(C₁-C₄)-alkyl, (C₁-C₄)-alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, C(O)NR_aR_b and -OR_x; R_x consists of (C₁-C₄)-alkyl and (C₃-C₆)-cycloalkyl; R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more of -C(O)R_x.

[0056] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, R₂ and R₃ are both (C₁-C₄)-alkyl, n is 1, R₅ is phenyl substituted with C(O)NR_aR_b, wherein R_a and R_b are independently selected from the group consisting of hydrogen and (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl, wherein said heterocycloalkyl is optionally substituted with one or more (C₁-C₄)-alkyl or -C(O)R_x; and R_x is (C₁-C₄)-alkyl.

[0057] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, R₂ and R₃ together with the carbon atom to which they are attached form a cyclopropyl ring, n is 1, R₅ is phenyl substituted with one or more substituents independently selected from halogen, (C₁-C₄)-alkyl, and (C₁-C₄)-alkyloxy.

[0058] In another embodiment of the present invention, R₁ and R₄ are both hydrogen, R₂ and R₃ together with the carbon atom to which they are attached form a oxetanyl ring, n is 1, R₅ is phenyl substituted with one or more substituents independently selected from halogen, (C₁-C₄)-alkyl, and (C₁-C₄)-alkyloxy.

[0059] In another embodiment of the present invention, R₁ is (C₁-C₄)-alkyl, all of R₂, R₃, R₄ are hydrogen, n is 1, R₅ is phenyl optionally substituted with one or more substituents independently selected from halogen, (C₁-C₄)-alkyl, and (C₁-C₄)-alkyloxy.

[0060] In another embodiment of the present invention, all of R₁, R₂, R₃, R₄ are hydrogen, n is 2, R₅ is phenyl which is optionally substituted with one or more substituents independently selected from halogen, (C₁-C₄)-alkyl, -S(O)₂R_x, -S(O)₂NR_aR_b, and -C(O)R_x; R_x consists of (C₁-C₄)-alkyl and (C₃-C₆)-cycloalkyl; R_a and R_b are independently selected from the group consisting of hydrogen, (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl.

[0061] In another embodiment of the present invention, all of R₁, R₂, R₃, R₄ are hydrogen, n is 2, R₅ is phenyl substituted with C(O)NR_aR_b, wherein R_a and R_b are independently selected from the group consisting of hydrogen and (C₁-C₄)-alkyl, or R_a and R_b together with the nitrogen to which they are attached form a 4-6 membered heterocycloalkyl. In another embodiment of the present invention, all of R₁, R₂, R₃ are hydrogen, n is 0, R₅ is phenyl which is optionally substituted with one or more substituents independently selected from halogen, (C₁-C₄)-alkyl, and (C₁-C₄)-alkyloxy.

[0062] Specific examples of compounds of formula (I) may be selected from the group consisting of:

3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methylbenzoate (compound 101);

3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-ethylbenzoate (compound 102);

[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(methylsulfamoyl)benzoate (compound 103); [(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(dimethylsulfamoyl)benzoate (compound 104); [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methylsul-

fonylbenzoate (compound 105); and pharmaceutically acceptable salts, hydrates or solvates thereof.

Definitions

5 **[0063]** As used throughout the present specification and appended claims, the following terms have the indicated meaning:

The term "alkyl" is intended to indicate a radical obtained when one hydrogen atom is removed from a branched or linear hydrocarbon. Said alkyl comprises 1-6, such as 1-4, such as 1-3, such as 2-3 or such as 1-2 carbon atoms. The term includes the subclasses normal alkyl (*n*-alkyl), secondary and tertiary alkyl, such as methyl, ethyl, *n*-propyl, isopropyl, 10 *n*-butyl, isobutyl, *sec*.-butyl, *tert*.-butyl, *n*-pentyl, isopentyl, neopentyl, *n*-hexyl and isohexyl.

[0064] The terms "alkyloxy" and "alkoxy" are intended to indicate a radical of the formula -OR', wherein R' is alkyl as indicated herein, wherein the alkyl group is appended to the parent molecular moiety through an oxygen atom, e.g. methoxy (-OCH₃), ethoxy (-OCH₂CH₃), *n*-propoxy, isopropoxy, butoxy, *tert*-butoxy, and the like.

15 **[0065]** The term alkoxyalkyl is intended to indicate an alkyl group as defined above substituted with one or more alkoxy groups as defined above, e.g. methoxymethyl, ethoxymethyl, 2-methoxyethyl, 2-ethoxyethyl, 3-methoxyprop-1-yl, and the like. The term "haloalkyl" is intended to indicate an alkyl group as defined herein substituted with one or more halogen atoms as defined herein, e.g. fluoro or chloro, such as fluoromethyl, difluoromethyl or trifluoromethyl.

[0066] The terms "haloalkyloxy" and "haloalkoxy" are intended to indicate a haloalkyl group as defined herein which is appended to the parent molecular moiety through an oxygen atom, such as difluoromethoxy or trifluoromethoxy.

20 **[0067]** The term "halogen" is intended to indicate a substituent from the 7th main group of the periodic table, such as fluoro, chloro and bromo.

[0068] The term "alkylthio" is intended to indicate a radical of the formula -S-R', wherein R' is alkyl as indicated herein, wherein the alkyl group is appended to the parent molecular moiety through a sulphur atom, e.g. -S-CH₃ (methylthio) or -S-CH₂CH₃ (ethylthio).

25 **[0069]** The term "cyano" is intended to indicate a -CN group attached to the parent molecular moiety through the carbon atom.

[0070] The term "cycloalkyl" is intended to indicate a saturated cycloalkane hydrocarbon radical, comprising 3-7 carbon atoms, 3-6 carbon atoms, 3-5 carbon atoms, 3-4 carbon atoms, e.g. cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl.

30 **[0071]** The term "aryl" is intended to indicate a radical of aromatic carbocyclic rings comprising 6-13 carbon atoms, 6-9 carbon atoms, such as 6 carbon atoms, in particular 5- or 6-membered rings, including fused carbocyclic rings with at least one aromatic ring. If the aryl group is a fused carbocyclic ring, the point of attachment of the aryl group to the parent molecular moiety may be through an aromatic or through an aliphatic carbon atom within the aryl group. Representative examples of aryl include, but are not limited to phenyl, naphthyl, indenyl, indanyl, dihydronaphthyl, tetrahydro-naphthyl and fluorenyl.

35 **[0072]** The term "heteroaryl" is intended to indicate radicals of monocyclic heteroaromatic rings comprising 5- or 6-membered ring which contains from 1-5 carbon atoms and from 1-4 heteroatoms selected from oxygen, sulphur and nitrogen. The heteroaryl radical may be connected to the parent molecular moiety through a carbon atom or a nitrogen atom contained anywhere within the heteroaryl group. Representative examples of heteroaryl groups include, but are not limited to, furanyl, imidazolyl, isothiazolyl, isoxazolyl, oxadiazolyl, oxazolyl, pyrazinyl, pyrazolyl, pyridazinyl, pyridyl, 40 pyrimidinyl, pyrrolyl, tetrazolyl, thiadiazolyl, thiazolyl, thienyl, triazolyl.

[0073] The term "heterocycloalkyl" is intended to indicate a cycloalkane radical as described herein, wherein one or more carbon atoms are replaced by heteroatoms, comprising 1-6 carbon atoms, e.g. 2-5 or 2-4 carbon atoms, further comprising 1-3 heteroatoms, preferably 1 to 2 heteroatoms, selected from O, N, or S, S(=O) or S(=O)₂. The heterocycloalkyl radical may be connected to the parent molecular moiety through a carbon atom or a nitrogen atom contained 45 anywhere within the heterocycloalkyl group. Representative examples of heterocycloalkyl groups include, but are not limited to azepanyl, azetidiny, aziridinyl, dioxolanyl, dioxolyl, imidazolidinyl, morpholinyl, oxetanyl, piperazinyl, piperidinyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothiophenyl, tetrahydrothiopyranyl.

[0074] The term "(4-6) membered heterocycloalkyl" is intended to indicate a heterocycloalkyl as defined herein, comprising 4-6 ring-atoms, and comprising 1-5 carbon atoms, e.g. 2-5 or 2-4 carbon atoms, further comprising 1-3 heteroatoms, preferably 1 to 2 heteroatoms, selected from O, N, S, S(=O) or S(=O)₂. Representative examples of (4-6) 50 membered heterocycloalkyl groups include azetidiny, dioxanyl, dioxolanyl, imidazolidinyl, morpholinyl, thiomorpholinyl, thiomorpholinyl-1-oxide, thiomorpholinyl-1,1-dioxide, oxetanyl, piperazinyl, piperidinyl, pyrrolidinyl, tetrahydrofuranyl, tetrahydropyranyl, tetrahydrothiopyranyl, tetrahydrothiophenyl, thietanyl.

[0075] The term "hydrocarbon radical" is intended to indicate a radical containing only hydrogen and carbon atoms, it may contain one or more double and/or triple carbon-carbon bonds, and it may comprise cyclic moieties in combination with branched or linear moieties. Said hydrocarbon comprises 1-6 carbon atoms, and preferably comprises 1-5, e.g. 1-4, e.g. 1-3, e.g. 1-2 carbon atoms. The term includes alkyl, cycloalkyl and aryl, as indicated herein.

[0076] In some instances, the number of carbon atoms in a hydrocarbon radical (e.g. alkyl, cycloalkyl and aryl) is

indicated by the prefix "(C_a-C_b)", wherein a is the minimum number and b is the maximum number of carbons in the hydrocarbon radical. Thus, for example (C₁-C₄)alkyl is intended to indicate an alkyl radical comprising from 1 to 4 carbon atoms, and (C₃-C₆)cycloalkyl is intended to indicate a cycloalkyl radical comprising from 3 to 6 carbon ring atoms.

[0077] The term "hydroxyalkyl" is intended to indicate an alkyl group as defined above substituted with one or more hydroxy, e.g. hydroxymethyl, hydroxyethyl, hydroxypropyl.

The term "hydroxyl" is intended to indicate an -OH group.

The term "oxo" is intended to indicate an oxygen atom which is connected to the parent molecular moiety via a double bond (=O).

The term "thioxo" is intended to indicate a sulfur atom which is connected to the parent molecular moiety via a double bond (=S).

The group C(O) is intended to represent a carbonyl group (C=O).

The term "pharmaceutically acceptable salt" is intended to indicate salts prepared by reacting a compound of formula I, which comprise a basic moiety, with a suitable inorganic or organic acid, such as hydrochloric, hydrobromic, hydroiodic, sulfuric, nitric, phosphoric, formic, acetic, 2,2-dichloroacetic, adipic, ascorbic, L-aspartic, L-glutamic, galactaric, lactic, maleic, L-malic, phthalic, citric, propionic, benzoic, glutaric, gluconic, D-glucuronic, methanesulfonic, salicylic, succinic, malonic, tartaric, benzenesulfonic, ethane-1,2-disulfonic, 2-hydroxy ethanesulfonic acid, toluenesulfonic, sulfamic or fumaric acid. Further examples of pharmaceutical acceptable salts are listed in Berge, S.M.; J. Pharm. Sci.; (1977), 66(1), 1-19.

[0078] The term "solvate" is intended to indicate a species formed by interaction between a compound, e.g. a compound of formula I, and a solvent, e.g. alcohol, glycerol or water, wherein said species are in a crystalline form. When water is the solvent, said species is referred to as a hydrate.

[0079] The term "treatment" as used herein means the management and care of a patient for the purpose of combating a disease, disorder or condition. The term is intended to include the delaying of the progression of the disease, disorder or condition, the amelioration, alleviation or relief of symptoms and complications, and/or the cure or elimination of the disease, disorder or condition. The term may also include prevention of the condition, wherein prevention is to be understood as the management and care of a patient for the purpose of combating the disease, condition or disorder and includes the administration of the active compounds to prevent the onset of the symptoms or complications. Nonetheless, prophylactic (preventive) and therapeutic (curative) treatments are two separate aspects.

[0080] The terms "disease", "condition" and "disorder" as used herein are used interchangeably to specify a state of a patient which is not the normal physiological state of man.

[0081] The term "medicament" as used herein means a pharmaceutical composition suitable for administration of the pharmaceutically active compound to a patient.

[0082] The term "pharmaceutically acceptable" as used herein means suited for normal pharmaceutical applications, i.e. giving rise to no adverse events in patients etc.

The compounds of the invention may be obtained in crystalline form either directly by concentration from an organic solvent or by crystallisation or re-crystallisation from an organic solvent or mixture of said solvent and a co-solvent that may be organic or inorganic, such as water. The crystals may be isolated in essentially solvent-free form or as a solvate, such as a hydrate. The invention covers all crystalline modifications and forms and also mixtures thereof.

[0083] Compounds of formula (I) may or may not comprise asymmetrically substituted (chiral) carbon atoms which give rise to the existence of isomeric forms, e.g. enantiomers and possibly diastereomers. The present invention relates to all such isomers, either in optically pure form or as mixtures thereof (e.g. racemic mixtures or partially purified optical mixtures). Pure stereoisomeric forms of the compounds and the intermediates of this invention may be obtained by the application of procedures known in the art. The various isomeric forms may be separated by physical separation methods such as selective crystallization and chromatographic techniques, e.g. high pressure liquid chromatography using chiral stationary phases. Enantiomers may be separated from each other by selective crystallization of their diastereomeric salts which may be formed with optically active amines, such as l-ephedrine, or with optically active acids. Optically purified compounds may subsequently be liberated from said purified diastereomeric salts. Enantiomers may also be resolved by the formation of diastereomeric derivatives. Alternatively, enantiomers may be separated by chromatographic techniques using chiral stationary phases. Pure stereoisomeric forms may also be derived from the corresponding pure stereoisomeric forms of the appropriate starting materials, provided that the reaction occur stereoselectively or stereospecifically. If a specific stereoisomer is desired, said compound will be synthesized by stereoselective or stereospecific methods of preparation. These methods will advantageously employ chiral pure starting materials.

Furthermore, when a double bond or a fully or partially saturated ring system is present in the molecule geometric isomers may be formed. It is intended that any geometric isomer, as separated, pure or partially purified geometric isomers or mixtures thereof are included within the scope of the invention.

[0084] Deuterated analogues. Any formula given herein is also intended to represent unlabeled forms as well as isotopically labeled forms of the compounds. For example, any hydrogen represented by "H" in any of the formulae herein is intended to represent all isotopic forms of hydrogen, e.g. ¹H, ²H or D, ³H. Enrichment with heavier isotopes, particularly

deuterium (i.e. ^2H or D) may afford certain therapeutic advantages due to for example an increased metabolic skin stability or an increased, systemic, in vivo clearance. Changes that would result in reduced dosage requirements or an improvement in therapeutic index. It is understood that deuterium in this context is regarded as a substituent of a compound of the formula (I).

[0085] Isotopically-enriched compounds of formula (I) can generally be prepared by conventional techniques known to those skilled in the art or by processes analogous to those described in the accompanying Preparations and Examples using any appropriate isotopically enriched reagent in place of the non-enriched reagent previously employed.

Medical use

[0086] As the compounds of the invention could exhibit PDE4 inhibitory activity, the compounds could be useful as therapeutic agents for inflammatory allergic diseases such as bronchial asthma, COPD, allergic rhinitis, and nephritis; autoimmune diseases such as rheumatoid arthritis, multiple sclerosis, Crohn's disease, and systemic lupus erythematosus; acute or chronic cutaneous wound disorders; diseases of the central nervous system such as depression, amnesia, and dementia; organopathy associated with ischemic reflux caused by cardiac failure, shock, and cerebrovascular diseases, and the like; insulin-resistant diabetes; wounds; AIDS, and the like.

[0087] In one embodiment, the compounds of the present invention are considered useful for the treatment, prevention or alleviation of dermal diseases or conditions.

[0088] In another embodiment, the compounds of the present invention are considered useful for the treatment, prevention or alleviation of dermal diseases or conditions selected from the group consisting of proliferative and inflammatory skin disorders, dermatitis, atopic dermatitis, seborrheic dermatitis, contact dermatitis, psoriasis, cancer, epidermal inflammation, alopecia, alopecia areata, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, urticaria, pruritis, and eczema.

[0089] In another embodiment, the compounds of the present invention are considered useful for the treatment or alleviation of atopic dermatitis.

[0090] In another embodiment, the compounds of the present invention are considered useful for the treatment or alleviation of psoriasis.

[0091] In another embodiment, the compounds of the present invention are considered useful for the treatment or alleviation of alopecia areata.

[0092] In another embodiment, the compounds of the present invention are considered useful for the treatment or alleviation of acne.

[0093] In another embodiment, the compounds of the present invention are considered useful for the treatment or alleviation of pruritis.

[0094] In another embodiment, the compounds of the present invention are considered useful for the treatment or alleviation of eczema.

[0095] Compounds of the invention, optionally in combination with other active compounds, may be useful for the treatment of dermal diseases or conditions, in particular for the treatment of proliferative and inflammatory skin disorders, dermatitis, atopic dermatitis, seborrheic dermatitis, contact dermatitis, psoriasis, cancer, epidermal inflammation, alopecia, alopecia areata, skin atrophy, steroid induced skin atrophy, skin ageing, photo skin ageing, acne, urticaria, pruritis, and eczema.

Besides being useful for human treatment, the compounds of the present invention may also be useful for veterinary treatment of animals including mammals such as horses, cattle, sheep, pigs, dogs, and cats.

[0096] For use in therapy, compounds of the present invention are typically in the form of a pharmaceutical composition. The invention therefore relates to a pharmaceutical composition comprising a compound of formula (I), optionally together with one or more other therapeutically active compound(s), together with a pharmaceutically acceptable excipient or vehicle. The excipient must be "acceptable" in the sense of being compatible with the other ingredients of the composition and not deleterious to the recipient thereof.

[0097] In the form of a dosage unit, the compound may be administered one or more times a day at appropriate intervals, always depending, however, on the condition of the patient, and in accordance with the prescription made by the medical practitioner. Conveniently, a dosage unit of a topical formulation contain between 0.001 mg and 1000 mg, preferably between 0.01 mg and 100 mg, such as 0.1-50 mg of a compound of formula (I). Also, conveniently, a dosage unit of a topical formulation contain between 0.01 mg and 10 g mg, preferably between 0.1 mg and 1000 mg, such as 1-500 mg of a compound of formula (I).

[0098] A suitable dosage of the compound of the invention will depend, inter alia, on the age and condition of the patient, the severity of the disease to be treated and other factors well known to the practising physician. The compound may be administered either orally, parenterally or topically according to different dosing schedules, e.g. daily or with weekly intervals. In general a single dose will be in the range from 0.0001 to 10 mg/kg body weight, e.g. in the range from 0.001 to 5 mg/kg body weight. Also, in general a single dose will be in the range from 0.001 to 100 mg/kg body

weight, e.g. in the range from 0.01 to 10 mg/kg body weight. The compound may be administered as a bolus (i.e. the entire daily dose is administered at once) or in divided doses two or more times a day.

[0099] In the context of topical treatment it may be more appropriate to refer to a "usage unit", which denotes unitary, i.e. a single dose which is capable of being administered to a patient, and which may be readily handled and packed, remaining as a physically and chemically stable unit dose comprising either the active material as such or a mixture of it with solid or liquid pharmaceutical diluents or carriers. A "usage unit" is capable of being administered topically to a patient in an application per square centimetre of the skin of from 0.1 mg to 50 mg and preferably from 0.2 mg to 5 mg of the final formulation in question.

[0100] It is also envisaged that in certain treatment regimes, administration with longer intervals, e.g. every other day, every week, or even with longer intervals may be beneficial.

[0101] If the treatment involves administration of another therapeutically active compound it is recommended to consult Goodman & Gilman's *The Pharmacological Basis of Therapeutics*, 9th Ed., J.G. Hardman and L.E. Limbird (Eds.), McGraw-Hill 1995, for useful dosages of said compounds.

[0102] The administration of a compound of the present invention with one or more other active compounds may be either concomitantly or sequentially.

[0103] The formulations include e.g. those in a form suitable for oral (including sustained or timed release), rectal, parenteral (including subcutaneous, intraperitoneal, intramuscular, intraarticular and intravenous), transdermal, ophthalmic, topical, dermal, nasal or buccal administration. Topical administration of the claimed formulation is particularly suitable.

[0104] The formulations may conveniently be presented in dosage unit form and may be prepared by but not restricted to any of the methods well known in the art of pharmacy, e.g. as disclosed in Remington, *The Science and Practice of Pharmacy*, 21ed ed., 2005. All methods include the step of bringing the active ingredient into association with the carrier, which constitutes one or more accessory ingredients. In general, the formulations are prepared by uniformly and intimately bringing the active ingredient into association with a liquid carrier, semisolid carrier or a finely divided solid carrier or combinations of these, and then, if necessary, shaping the product into the desired formulation.

[0105] Formulations of the present invention suitable for oral and buccal administration may be in the form of discrete units as capsules, sachets, tablets, chewing gum or lozenges, each containing a predetermined amount of the active ingredient; in the form of a powder, granules or pellets; in the form of a solution or a suspension in an aqueous liquid or non-aqueous liquid, such as ethanol or glycerol; or in the form of a gel, a nano- or microemulsion, an oil-in-water emulsion, a water-in-oil emulsion or other dispensing systems. The oils may be edible oils, such as but not restricted to e.g. cottonseed oil, sesame oil, coconut oil or peanut oil. Suitable dispersing or suspending agents for aqueous suspensions include synthetic or natural surfactants and viscosifying agents such as but not restricted to tragacanth, alginate, acacia, dextran, sodium carboxymethylcellulose, gelatin, methylcellulose, hydroxypropylmethylcellulose, hydroxypropylcellulose, carbomers, polyvinylpyrrolidone, polysorbates, sorbitan fatty acid esters,. The active ingredients may also be administered in the form of a bolus, electuary or paste.

[0106] A tablet may be made by compressing, moulding or freeze drying the active ingredient optionally with one or more accessory ingredients. Compressed tablets may be prepared by compressing, in a suitable machine, the active ingredient(s) in a free-flowing form such as a powder or granules, optionally mixed by a binder and/or filler, such as e.g. lactose, glucose, mannitol starch, gelatine, acacia gum, tragacanth gum, sodium alginate, calcium phosphates, microcrystalline cellulose, carboxymethylcellulose, methylcellulose, hydroxypropylcellulose, hydroxypropylmethylcellulose, ethylcellulose, hydroxyethylcellulose, polyethylene glycol, waxes or the like; a lubricant such as e.g. sodium oleate, sodium stearate, magnesium stearate, sodium benzoate, sodium acetate, sodium chloride or the like; a disintegrating agent such as e.g. starch, methylcellulose, agar, bentonite, croscarmellose sodium, sodium starch glycollate, crospovidone or the like or a dispersing agent, such as polysorbate 80. Moulded tablets may be made by moulding, in a suitable machine, a mixture of the powdered active ingredient and suitable carrier moistened with an inert liquid diluent. Freeze dried tablets may be formed in a freeze-dryer from a solution of the drug substance. A suitable filler can be included.

[0107] Formulations for rectal administration may be in the form of suppositories in which the compound of the present invention is admixed with low melting point, water soluble or insoluble solids such as cocoa butter, hydrogenated vegetable oils, polyethylene glycol or fatty acids esters of polyethylene glycols, while elixirs may be prepared using myristyl palmitate.

[0108] Formulations suitable for parenteral administration conveniently comprise a sterile oily or aqueous preparation of the active ingredients, which is preferably isotonic with the blood of the recipient, e.g. isotonic saline, isotonic glucose solution or buffer solution. Furthermore, the formulation may contain cosolvent, solubilising agent and/or complexation agents. The formulation may be conveniently sterilised by for instance filtration through a bacteria retaining filter, addition of sterilising agent to the formulation, irradiation of the formulation or heating of the formulation. Liposomal formulations as disclosed in e.g. *Encyclopedia of Pharmaceutical Technology*, vol.9, 1994, are also suitable for parenteral administration.

[0109] Alternatively, the compounds of formula (I) may be presented as a sterile, solid preparation, e.g. a freeze-dried powder, which is readily dissolved in a sterile solvent immediately prior to use.

[0110] Transdermal formulations may be in the form of a plaster, patch, microneedles, liposomal or nanoparticulate delivery systems or other cutaneous formulations applied to the skin.

[0111] Formulations suitable for ophthalmic administration may be in the form of a sterile aqueous preparation of the active ingredients, which may be in microcrystalline form, for example, in the form of an aqueous microcrystalline suspension. Liposomal formulations or biodegradable polymer systems e.g. as disclosed in Encyclopedia of Pharmaceutical Technology, vol.2, 1989, may also be used to present the active ingredient for ophthalmic administration.

[0112] Formulations suitable for topical, such as dermal, intradermal or ophthalmic administration include liquid or semi-solid preparations such as liniments, lotions, gels, applicants, sprays, foams, film forming systems, microneedles, micro- or nano-emulsions, oil-in-water or water-in-oil emulsions such as creams, ointments or pastes; or solutions or suspensions such as drops. Compositions for ophthalmic treatment may additionally contain cyclodextrin.

[0113] For topical administration, the compound of formula (I) may typically be present in an amount of from 0.001 to 20% by weight of the composition, such as 0.01% to about 10 %, but may also be present in an amount of up to about 100% of the composition.

[0114] Formulations suitable for nasal or buccal administration include powder, self-propelling and spray formulations, such as aerosols and atomisers. Such formulations are disclosed in greater detail in e.g. Modern Pharmaceutics, 2nd ed., G.S. Banker and C.T. Rhodes (Eds.), page 427-432, Marcel Dekker, New York; Modern Pharmaceutics, 3th ed., G.S. Banker and C.T. Rhodes (Eds.), page 618-619 and 718-721, Marcel Dekker, New York and Encyclopedia of Pharmaceutical Technology, vol. 10, J. Swarbrick and J.C. Boylan (Eds), page 191-221, Marcel Dekker, New York.

[0115] In addition to the aforementioned ingredients, the formulations of a compound of formula (I) may include one or more additional ingredients such as diluents, buffers, flavouring agents, colourant, surface active agents, thickeners, penetration enhancing agents, solubility enhancing agents preservatives, e.g. methyl hydroxybenzoate (including antioxidants), emulsifying agents and the like.

[0116] When the active ingredient is administered in the form of salts with pharmaceutically acceptable non-toxic acids or bases, preferred salts are for instance easily watersoluble or slightly soluble in water, in order to obtain a particular and appropriate rate of absorption.

[0117] The pharmaceutical composition may additionally comprise one or more other active components conventionally used in the treatment of dermal disease or conditions, e.g. selected from the group consisting of glucocorticoids, vitamin D and vitamin D analogues, antihistamines, platelet activating factor (PAF) antagonists, anticholinergic agents, methylxanthines, β -adrenergic agents, COX-2 inhibitors, JAK inhibitors, other PDEs, salicylates, indomethacin, flufenamate, naproxen, timegadine, gold salts, penicillamine, serum cholesterol lowering agents, retinoids, zinc salts, salicylazosulfa-pyridine and calcineurin inhibitors.

METHODS OF PREPARATION

[0118] The compounds of the present invention can be prepared in a number of ways well known to those skilled in the art of synthesis. The compounds of formula (I) may for example be prepared using the reactions and techniques outlined below together with methods known in the art of synthetic organic chemistry, or variations thereof as appreciated by those skilled in the art. Preferred methods include, but are not limited to, those described below. The reactions are carried out in solvents appropriate to the reagents and materials employed and suitable for the transformations being effected. Also, in the synthetic methods described below, it is to be understood that all proposed reaction conditions, including choice of solvent, reaction atmosphere, reaction temperature, duration of experiment and work-up procedures, are chosen to be conditions of standard for that reaction, which should be readily recognized by one skilled in the art. Not all compounds falling into a given class may be compatible with some of the reaction conditions required in some of the methods described. Such restrictions to the substituents which are compatible with the reaction conditions will be readily apparent to one skilled in the art and alternative methods can be used.

[0119] The compounds of the present invention or any intermediate may be purified if required using standard methods well known to a synthetic organic chemist, e.g. methods described in "Purification of Laboratory Chemicals", 6th ed. 2009, W. Amarego and C. Chai, Butterworth-Heinemann.

[0120] Starting materials are either known or commercially available compounds or can be prepared by routine synthetic methods well known to a person skilled in the art.

[0121] ^1H nuclear magnetic resonance (NMR) spectra were recorded at 300 or 600 MHz unless otherwise specified. Chemical shift values (δ , in ppm) are quoted relative to internal tetramethylsilane ($\delta = 0.00$) standards. The value of a multiplet, either defined doublet (d), triplet (t), quartet (q) or (m) at the approximate midpoint is given unless a range is quoted. (br) indicates a broad peak, whilst (s) indicates a singlet. The organic solvents used were usually anhydrous. Chromatography was performed on Merck silica gel 60 (0.040 - 0.063 mm). The solvent ratios indicated refer to v:v unless otherwise noted. All NMR spectra are recorded in DMSO- d_6 unless another solvent is stated.

Analytical UPLC/MS

5 **[0122]** Analytical UPLC/MS is performed on a Waters Acquity UPLC-system and SQD-MS. Column: Waters Acquity HSS T3 1.8 μ m, 2.1 x 50mm; solvent system: A = 10mM Ammonium acetate in water + 0.1% HCOOH and B = acetonitrile + 0.1% HCOOH; flow rate = 1.2 mL/min; method (1.4 min): Linear gradient method from 5 % B to 95 % B over 0.9 minutes then 95 % B for 0.3 minutes. Column temperature is 60°C.

Preparative purification HPLC/MS:

10 **[0123]** Preparative HPLC/MS was performed on a Waters AutoPurification system with a Waters SQD2 mass spectrometer. This includes three steps, pre-analysis, preparative purification and re-analysis on the purified compound.

[0124] Solvent: A = 0.1% formic acid and solvent B = acetonitrile with 0,1% formic acid.

Analytical pre-analysis using the following method:

15 **[0125]** Column: Waters SUNFIRE C-18, 100 mm x 4.6 mm, 5 μ m
Flow rate = 1.2 mL/min. (method 10 min)
[0126] Method: Linear gradient method going from 10 % B to 95 % B in 6.5 minutes and staying at 95 % B for another 1.5 minutes to obtain the retention time of the compounds provides the following four different preparative gradient
20 methods:

Preparative methods:

25 **[0127]** Column: Waters SUNFIRE C-18, 100 mm x 19 mm, 5 μ m
Flow rate = 20 mL/min. (method 8 min) 0-3 min: 5% B for 2 minutes followed by a linear gradient method going from 5% B to 35 % B in 3 minutes and going to 100% B and staying at 100 % B for another 1.4 minutes.
3.01-5 min: 15% B for 1 minutes followed by a linear gradient method going from 15% B to 55% B in 4 minutes and going to 100% B and staying at 100% B for another 1.4 minutes.
5.01-7.5 min: 30% B for 1 minutes followed by a linear gradient method going from 30% B to 70% B in 4 minutes and
30 going to 100% B and staying at 100 % B for another 1.4 minutes.
7.51-10 min: 50% B for 1 minutes followed by a linear gradient method going from 50% B to 100% B in 4 minutes and staying at 100% B for another 1.5 minutes.

[0128] The fractions were collected based on ion traces of relevant ions and PDA signal (240-400 nm).

35 Re-analysis method for fractions:

[0129] Column: Waters Xselect C18; 50x3.0mm 5 μ m

Flow rate = 1.2 mL/min. (method 5 min)

40 **[0130]** Method: Linear gradient method going from 10 % B to 95 % B in 3 minutes and staying at 95 % B for another 0.5 min.

Instruments:

45 **[0131]** Waters 2767 Sample Manager
Waters 2545 Binary Gradient Module
Waters SFO System Fluidics Organizer
Waters 515 HPLC Pump
Waters 2998 Photodiode Array Detector
Waters SQDetector 2

50 **[0132]** LCMS Method "XE Metode 7 CM"

A quality check was performed on a Waters LCT Premier MS instrument and a Waters Aquity UPLC.

Column: Waters Aquity UPLC HSS T3 1.8 μ m, 2.1 x 50 mm, at 40°C.

Solvents: A = 10 mM ammonium acetate + 0.1% HCOOH, B = MeCN + 0.1% HCOOH. Flow: 0.7 ml/min. Injection volume 2 μ l. UV detection range 240 - 400 nm.

55 **[0133]** Gradient:

Time	% A	% B
0.00 min	99	1
0.50 min	94	6
1.00 min	94	6
2.60 min	5	95
3.80 min	5	95
3.81 min	99	1
4.80 min	99	1


[0134] The MW confirmation and purity was extracted and checked with OpenLynx.

[0135] The following abbreviations have been used throughout:

ACN	acetonitrile
DCE	1,2-dichloroethane
DCM	dichloromethane
DMAP	<i>N,N</i> -dimethylpyridin-4-amine
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
EDAC	(3-dimethylamino-propyl)-ethyl-carbodiimide
EtOH	ethanol
MeOH	methanol
EtOAc	ethyl acetate
L	litre
LAH	lithium aluminium hydride
Me	methyl
NMR	nuclear magnetic resonance
PG	protecting group
PPTS	pyridinium <i>p</i> -toluenesulfonate
RT	room temperature
THF	tetrahydrofuran
TEA	triethylamine

General Methods


[0136] Compounds of general formula (I) of the invention may for example be prepared according to the following non-limiting general methods and examples. R_1 , R_2 , R_3 , R_4 , Q and n are as previously defined for the compounds of general formula (I):



a) Synthesis of Compound 004 is outlined in Scheme 1

[0137] Compound 004 is prepared as described in WO 2008/110308 in three steps.


Scheme 1



b) Synthesis of compound 006 is outlined in Scheme 2

[0138] Compound 006 is prepared according to standard procedures known to a chemist skilled in the art of organic synthesis. Reaction of compound 004 with propionyl chloride or propionic anhydride in the presence of a base such as pyridine, triethylamine, or diisopropylethylamine in a solvent such as DCM, THF gives compound 005. Treatment of acetone cyanohydrin, Et₃N delivers Compound 006.

Scheme 2



c) Synthesis of a compound of general formula (V) is outlined in Scheme 3

[0139] A compound of general formula (II) may be prepared by treatment of compound 006 with a compound of general formula (III) in the presence of a suitable acid such as acetic acid and 4-toluenesulphonic acid or in the absence of an acid in a suitable solvent such as methanol, ethanol, propanol, isopropanol, butanol, and THF at suitable temperature between rt and 150 °C (preferably 50 °C to 120 °C).

[0140] A compound of general formula (II) may also be prepared by treatment of compound 006 with a compound of formula (III) in the presence of a suitable base such as triethylamine, K₂CO₃, Bu₄NOH, KOH in a suitable solvent such as methanol, ethanol, propanol, isopropanol, butanol, and THF at suitable temperature between rt and 150 °C (preferably 50 °C to 120 °C), especially when a compound of general formula (II) is in a salt form such as hydrochloride and oxalate.


[0141] A compound of general formula (III) may be commercially available or prepared according to standard procedures known to a chemist skilled in the art of organic synthesis (for examples, see: Gever, Gabriel, J. Am. Chem. Soc. (1954), 76, 1283-1285; Ghali, N. I. J. Org. Chem. (1981), 46 (26), 5413-14; Kim, Yongju, ACS Medicinal Chemistry Letters (2012), 3 (2), 151-154). The protecting group may be acyl such as acetyl and benzoyl or tetrahydropyranyl, but not limited to these.

[0142] A compound of general formula (IV) may be prepared by procedures similar to those described, for example, by Bardakos, Vasilios et al. Chem Ber. (1976), 109, 1898-1910; Wang, Xiao-Feng et al. J. Med. Chem. (2014), 57 (4), 1390-1402. Typically, a compound of general formula (IV) may be prepared by treatment of formula (II) with hydrazoic acid, sodium azide or trimethylsilyl azide in the presence of a suitable Bronsted-Lowry acid such as hydrochloride, sulfuric acid, methanesulfonic acid, trifluoroacetic acid, phosphoric acid, trifluoromethanesulfonic acid, and trifluoroborane ether-

ate or a mixture of two or more of these or in the presence of a suitable Lewis acid such as trifluoroborane etherate in a suitable solvent such as water, dichloromethane, chloroform, acetic acid, methanesulfonic acid, toluene, and benzene or a mixture of two or more of these at a suitable temperature between 0 °C to 100 °C.

[0143] A compound of formula (V) represents a protected alcohol. A compound of general formula (IV) may be prepared by removal of a protecting group according to standard procedures known to a chemist skilled in the art of organic synthesis (for a review on protecting group, see: Kocienski, Philip J. Protecting groups, Georg Thieme Verlag, Stuttgart, New York, 2004).

Scheme 3




d) Synthesis of a compound of general formula I where Q represents -O-C(O)-R₅ (R₅ is as previous defined) is outlined in Scheme 4

[0144] A compound of general formula (I) where Q represents -O-C(O)-R₅ may be prepared from a compound of formula (IV) according to standard procedures known to a chemist skilled in the art of organic synthesis (for ester formation, see: Junzo Otera, Esterification: Methods, Reactions, and Applications, Wiley-VCH, Weinheim (2004)).

[0145] For example, a compound of general formula (I) may be prepared by reaction of a compound of general formula (IV) with HO-(CO)-R₅ in the presence of a suitable couplings reagent such as DCC, EDAC and a suitable catalyst such as DMAP in a suitable solvent such dichloromethane, dichloroethane, acetonitrile and ethyl acetate.

Scheme 4




15 Preparations and Examples

20 Preparation 1 (Compounds 001 and 002)

1-Tetrahydropyran-4-ylidenepropan-2-one and 1-(3,6-dihydro-2H-pyran-4-yl)propan-2-one

25 **[0146]**




30 **[0147]** To a solution of KOH (40 g, 722.9 mmol) in H₂O (200 mL) and EtOH (800 mL) at 0 °C, 1-dimethoxyphosphorylpropan-2-one (100 g, 602.04 mmol) in EtOH (100 mL) and tetrahydropyran-4-one (60 g, 602.04 mmol) in EtOH (100 mL) were added and stirred at 26 °C for 4 h. On completion of the reaction, excess solvent was evaporated under vacuum and the resulting residue was treated with water (200 mL) and extracted with diethyl ether (2 x 200 mL). The combined organic layers were washed with brine (200 mL), dried over Na₂SO₄, and concentrated in vacuum to afford the title compounds as pale yellow liquid (crude). The crude was used for the next step without any further purification.

35 Preparation 2 (Compound 003)

Ethyl 8-hydroxy-10-oxo-3-oxaspiro[5.5]undec-8-ene-11-carboxylate


40 **[0148]**



55 **[0149]** To a solution of EtONa (97.14 g, 1428 mmol) in ethanol (800 mL), diethyl malonate (114 g, 714.28 mmol) and 1-tetrahydropyran-4-ylidenepropan-2-one and 1-(3,6-dihydro-2H-pyran-4-yl)propan-2-one (100 g, 713.3 mmol) in EtOH (200 mL) were added slowly and stirred at reflux temperature for 16 h. On completion excess solvent was evaporated under vacuum to afford the title compound as brown solid (crude). The crude was used for the next step without any further purification.

Preparation 3 (Compound 004)

8-Hydroxy-3-oxaspiro[5.5]undec-8-en-10-one


[0150]

[0151] To ethyl 8-hydroxy-10-oxo-3-oxaspiro[5.5]undec-8-ene-11-carboxylate (100 g, 393.3 mmol), 6N NaOH (1 L) was added and stirred for 16 h at 26 °C. Then conc.HCl (500 mL) was added slowly at 0 °C to adjust the pH to ~2 and refluxed for 4h. On completion of the reaction, reaction mixture was treated with water and extracted with EtOAc (2 x 500 mL). The combined organic layers were washed with brine (500 mL), dried over Na₂SO₄, concentrated in vacuo, and purified by washed with diethyl ether to afford the title compound as a brown solid.

¹H NMR (300 MHz, DMSO-d₆) δ 11.02 (s, 1H), 5.19 (s, 1H), 3.71 - 3.45 (m, 4H), 2.29 (s, 4H), 1.45 (t, J = 5.4 Hz, 4H).

Preparation 4 (Compound 005)

(10-Oxo-3-oxaspiro[5.5]undec-8-en-8-yl) propanoate


[0152]

[0153] To a solution of 8-hydroxy-3-oxaspiro[5.5]undec-8-en-10-one (90 g, 494.5 mmol) in DCM (1.5 L), Pyridine (58.6 g, 741.7 mmol) and propionyl chloride (45.5 g, 494.5 mmol) were added slowly at 0 °C and it was stirred for 4 h at 26 °C. On completion the reaction mixture was treated with 1N.HCl and the organic layer was washed with brine (700 mL), dried over Na₂SO₄, and concentrated to afford the title compound as a brown liquid.

¹H NMR (300 MHz, CDCl₃) δ 5.91 (d, J = 1.2 Hz, 1H), 3.68 (t, J = 5.4 Hz, 4H), 2.56 (d, J = 1.3 Hz, 2H), 2.51 (q, J = 7.5, 2H), 2.44 (s, 2H), 1.73 - 1.49 (m, 4H), 1.22 (t, J = 7.5, 3H).

Preparation 5 (Compound 006)

9-Propanoyl-3-oxaspiro[5.5]undecane-8,10-dione

[0154]

[0155] To a solution of (10-oxo-3-oxaspiro[5.5]undec-8-en-8-yl) propanoate (90 g, 378.15 mmol) in ACN (1500 mL),


TEA (38.2 g, 378.15 mmol) and acetone cyanohydrin (48 g, 567.22 mmol) were added at 0 °C and stirred for 3 h at 26 °C. On completion excess solvent was evaporated under vacuum and the resulting residue was purified by silica gel column chromatography (1% MeOH in DCM as eluent) to afford the title compound as a pale yellow liquid.

¹H NMR (300 MHz, CDCl₃) δ 3.77 - 3.58 (m, 4H), 3.06 (q, J = 7.2 Hz, 2H), 2.67 (s, 2H), 2.51 (s, 2H), 1.62 - 1.47 (m, 4H), 1.13 (t, J = 7.3 Hz, 3H).

Preparation 6 (Compound 007)

3-Hydrazinopropan-1-ol

[0156]




[0157] A solution of NaOH (42.5 g, 1063 mmol) and N₂H₄·H₂O (269.5 g, 5319 mmol) was heated up to 100 °C, 3-chloropropan-1-ol (100 g, 1063 mmol) was added at same temperature and stirred for 5 h. On completion excess solvent was evaporated under vacuum, the resulting residue was treated with EtOH and the resulting solid was filtered. Filtrate was concentrated and the excess N₂H₄·H₂O was removed under downward distillation, to afford the title compound as a colorless liquid (crude), which was used for the next step without any further purification.

Preparation 7 (Compound 008)

3-Ethyl-1-(3-hydroxypropyl)spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one

[0158]




[0159] To a solution of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (160 g, 671.7 mmol) in ethanol (1.6 Lit), TEA (135.7 g, 1344.5 mmol) and 3-hydrazinopropan-1-ol (72.6 g, 806 mmol) were added slowly and stirred at 80 °C for 16 h. On completion excess solvent was evaporated under vacuum and the resulting residue was purified by silica gel (100-200 mesh) column chromatography (5% MeOH in DCM as eluent) to afford the title compound as a yellow liquid. ¹H NMR (300 MHz, CDCl₃) δ 4.24 - 4.13 (m, 2H), 3.71 (t, J = 5.4 Hz, 4H), 3.61 (td, J = 5.8, 2.2 Hz, 2H), 2.95 - 2.76 (m, 4H), 2.52 (s, 2H), 2.13 - 1.95 (m, 2H), 1.73 - 1.50 (m, 4H), 1.23 (t, J = 7.5 Hz, 3H).

HPLC-Retention time (XE Methode 7 CM): 1.59 minutes. Detected "M+1"-mass: 308.19.

Preparation 8 (Compound 009)

3-(3-Ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)propyl benzoate

[0160]




[0161] To a solution of 3-ethyl-1-(3-hydroxypropyl)spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one (160 g, 547.9 mmol) in DCM (1.6 L), TEA (138.3 g, 1369.8 mmol) and Benzoyl chloride (153.4 g, 1095.9 mmol) were added slowly at 0 °C and stirred for 16 h at 26 °C. On completion of the reaction, volatiles were evaporated under vacuum and the resulting residue was treated with water. The organic layer was washed with brine (500 mL), dried over Na₂SO₄, concentrated and purified by silica gel (100-200 mesh) column chromatography (70% EtOAc in PE as eluent) to afford the title compound as a pale yellow liquid, which was used directly in the next step.

Preparation 9 (Compound 010)

3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl benzoate

[0162]



[0163] To a solution of 3-(3-ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)propyl benzoate (160 g, 404.0 mmol) in CH₃SO₃H (775.8 g, 8080.8 mmol), NaN₃ (131.3 g, 2020.2 mmol) was added portion wise at 0 °C and stirred for 16h at 26 °C. On completion, the reaction mixture was diluted with water at 0 °C and extracted with 10% MeOH: DCM (2 x 250 mL). The combined organic layers were dried over Na₂SO₄, concentrated, and purified by silica gel (100-200 mesh) column chromatography (5% MeOH in DCM as eluent) to afford the title compound as pale yellow liquid.

¹H NMR (600 MHz, DMSO-d₆) δ 7.99 - 7.86 (m, 2H), 7.75 - 7.59 (m, 1H), 7.52 (t, J = 7.8 Hz, 2H), 7.40 (t, J = 5.8 Hz, 1H), 4.27 (t, J = 5.9 Hz, 2H), 4.16 (t, J = 6.7 Hz, 2H), 3.57 (dt, J = 12.0, 4.9 Hz, 2H), 3.49 (ddd, J = 11.7, 6.8, 4.7 Hz, 2H), 2.98 (d, J = 5.8 Hz, 2H), 2.75 (s, 2H), 2.72 (q, J = 7.5 Hz, 2H), 2.20 (p, J = 6.5 Hz, 2H), 1.44 - 1.30 (m, 4H), 1.09 (t, J = 7.5 Hz, 3H).


HPLC-Retention time (XE Metode 7 CM): 2.04 minutes.

Detected "M+1"-mass: 412.23.

Preparation 10 (Compound 011)

3-Ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0164]




[0165] To a solution of 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl benzoate (85 g, 206.81 mmol) in MeOH (850 mL), K_2CO_3 (57 g, 413.62 mmol) was added at 0 °C and stirred for 3 h at 26 °C. On completion, the reaction was filtered and it was washed with MeOH. The filtrate was concentrated and purified by Neutral Alumina column chromatography (5% MeOH in DCM as eluent) to afford the title compound as off-white solid. ^1H NMR (400 MHz, $DMSO-d_6$) δ 7.39 (t, J = 5.7 Hz, 1H), 4.58 (s, 1H), 4.01 (t, J = 7.1 Hz, 2H), 3.77 - 3.46 (m, 4H), 3.38 (t, J = 6.0 Hz, 2H), 3.01 (d, J = 5.7 Hz, 2H), 2.85 - 2.66 (m, 4H), 1.84 (p, J = 6.5 Hz, 2H), 1.44 (t, J = 5.6 Hz, 4H), 1.11 (t, J = 7.5 Hz, 3H).
HPLC-Retention time (XE Metode 7 CM): 1.59 minutes.
Detected "M+1"-mass: 308.19.

Preparation 11 (Compound 012)

Methyl (2S)-2-methyl-3-tetrahydropyran-2-yloxy-propanoate

[0166]




[0167] Methyl (2S)-3-hydroxy-2-methyl-propanoate (30 g, 356 mmol) was added to a solution of 3,4-dihydro-2H-pyran (16 g, 135 mmol) and PPTS (2 g, 8 mmol) in DCM (50 mL) at rt. After 2 h, the reaction was washed with H_2O (100 mL), saturated aqueous $NaHCO_3$ (100 mL) and with brine (100 mL), dried ($MgSO_4$), and concentrated in vacuo to afford the title compound as a colorless liquid, which was used in the next step without characterization.

Preparation 12 (Compound 013)

(2R)-2-Methyl-3-tetrahydropyran-2-yloxy-propan-1-ol

[0168]




[0169] Methyl (2S)-2-methyl-3-tetrahydropyran-2-yloxy-propanoate (31 g, 153 mmol) was cooled to 0 °C and treated dropwise with 200 mL (200 mmol) of LAH (1 M in THF). The solution was stirred at rt for 1 h. Under ice-bath cooling, to the solution was dropwise added 2N NaOH (40 mL) (very slow addition). The mixture was filtered. The filtrate was concentrated. The residue was purified by chromatography (heptane/ethyl acetate 4:1 (R_f = 0.1) to heptane/ethyl acetate 0:1), giving the title compound as a colorless oil. ^1H NMR (300 MHz, CDCl_3) δ 4.58 (1H, br), 3.93-3.29 (m, 6H), 2.67-2.49 (m, 1H), 2.13-1.94 (m, 1H), 1.90-1.42 (m, 6H), 0.98-0.83 (m, 3H).

Preparation 13 (Compound 014)

[(2S)-2-Methyl-3-tetrahydropyran-2-yloxy-propyl] methanesulfonate

[0170]




[0171] To a solution of (2R)-2-methyl-3-tetrahydropyran-2-yloxy-propan-1-ol (18 g, 103.3 mmol) and Et_3N (20 g) in DCM (100 mL) was dropwise added methanesulfonyl chloride (14.2 g, 124 mmol) at 0 °C. The obtained mixture was stirred at rt for 0.5 h and washed with H_2O , 1 N HCl and NaHCO_3 solution. The organic phase was dried over Na_2SO_4 and concentrated in vacuo, giving the title compound as a brown syrup.

^1H NMR (300 MHz, CDCl_3) δ 4.62 - 4.50 (m, 1H), 4.35 - 4.09 (m, 2H), 3.83 (ddt, J = 11.1, 7.4, 3.7 Hz, 1H), 3.73 (dd, J = 9.8, 5.2 Hz, 0.5H), 3.67 (dd, J = 9.9, 7.0 Hz, 0.5H), 3.56 - 3.46 (m, 1H), 3.37 (dd, J = 9.8, 5.0 Hz, 0.5H), 3.30 (dd, J = 9.8, 7.1 Hz, 0.5H), 3.01/3.01 (s, 3H), 2.31 - 2.10 (m, 1H), 1.93 - 1.45 (m, 6H), 1.04 (dd, J = 6.9, 2.8 Hz, 3H).

Preparation 14 (Compound 015)

[(2R)-2-Methyl-3-tetrahydropyran-2-yloxy-propyl]hydrazine

[0172]




[0173] A solution of [(2S)-2-methyl-3-tetrahydropyran-2-yloxy-propyl] methanesulfonate From Preparation 13 (103 mmol) and hydrazine monohydrate (30 mL) in isopropanol (100 mL) was heated to reflux for 2 h. The mixture was then concentrated in vacuo. The residue was taken up in DCM and washed with brine. The aqueous phase was extracted twice with DCM. The combined organic phases were dried over Na_2SO_4 and concentrated in vacuo, giving the title compound (crude). The crude was used directly in the next step without further purification.

Preparation 15 (Compound 016)

3-Ethyl-1-[(2R)-2-methyl-3-tetrahydropyran-2-yloxy-propyl]spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one

[0174]




15 **[0175]** A solution of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (15 g, 63 mmol) and [(2R)-2-Methyl-3-tetrahydropyran-2-yloxy-propyl]hydrazine (15 g, 80 mmol) in isopropanol (100 mL) was heated reflux for 5 h. The solution was concentrated in vacuo. The residue was purified by chromatography (ethyl acetate, $R_f = 0.41$), giving the as an oil. The compound was used in the next step without characterization.

20 Preparation 16 (Compound 017)

3-Ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one

25 **[0176]**




40 **[0177]** To a solution of 3-ethyl-1-[(2R)-2-methyl-3-tetrahydropyran-2-yloxy-propyl]spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one (27 g, 69 mmol) in MeOH/H₂O (3:1, 200 mL) was added methanesulfonic acid (2 mL) at rt. The solution was stirred at 50 °C for 1 h and concentrated in vacuo in order to remove MeOH. The aqueous mixture was diluted with water and extracted three times with DCM. The combined organic phases were dried and concentrated in vacuo. The residue was purified by chromatography (ethyl acetate/MeOH 20:1), giving the title compound as an oil.

45 ¹H NMR (300 MHz, DMSO-d₆) δ 4.63 (t, $J = 5.2$ Hz, 1H), 4.05 (dd, $J = 13.7, 6.5$ Hz, 1H), 3.81 (dd, $J = 13.7, 7.7$ Hz, 1H), 3.68-3.50 (m, 4H), 3.37 - 3.15 (m, 2H), 2.95 - 2.82 (m, 2H), 2.80-2.65 (m, 2H), 2.41 (s, 2H), 2.15-2.00 (m, 1H), 1.53-1.39 (m, 4H), 1.13 (t, $J = 7.5$ Hz, 3H), 0.80 (d, $J = 6.8$ Hz, 3H).

50 Preparation 17 (Compound 018)

3-Ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

55 **[0178]**



[0179] To a solution of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one (17 g, 55.5 mmol) in methanesulfonic acid (100 mL) was added sodium azide (7.2 g, 111 mmol) in portion at rt (exothermic). The obtained reaction mixture was stirred at rt for 2 h. To the mixture was added 400 mL of water. The mixture was neutralized with NaOH and concentrated. The residue was taken up in EtOH (400 mL). The precipitate was filtered off and washed with EtOH. The filtrate was concentrated in vacuo. The residue was taken up in DCM. The precipitate was filtered off and washed with DCM. The filtrate was concentrated in vacuo. The residue was purified twice by chromatography (DCM/MeOH 10:1), giving the title compound as a white solid.

^1H NMR (600 MHz, DMSO- d_6) δ 7.41 (t, J = 5.7 Hz, 1H), 4.64 (t, J = 5.1 Hz, 1H), 4.00 (dd, J = 13.9, 6.5 Hz, 1H), 3.73 (dd, J = 13.9, 7.8 Hz, 1H), 3.69 - 3.60 (m, 2H), 3.58-3.50 (m, 2H), 3.30-3.20 (m, 2H), 3.01 (d, J = 5.8 Hz, 2H), 2.81-2.79 (m, 4H), 2.16 - 2.01 (m, 1H), 1.50-1.40 (m, 4H), 1.11 (t, J = 7.5 Hz, 3H), 0.81 (d, J = 6.8 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.67 minutes.

Detected "M+1"-mass: 321.21.

General Procedure 1: esterification

[0180] To a solution of an acid (2 equiv) in DMF (0.1 mL) were added a solution of an alcohol (0.013 mmol) in DMF (0.1 mL) and a solution of DMAP (2 equiv) in DMF (0.1 mL). To this resulting solution was added a suspension of EDAC (2.7 equiv) in DMF (0.1 mL). The mixture was shaken at 50 °C overnight. The crude was subjected to preparative LCMS purification, giving an ester.


General Procedure 2: esterification

[0181] An alcohol (0.013 mmol) was dissolved in DCE (0.15 mL). A solution of an acid (2 equiv) in DCE (0.2 mL) and a solution of DMAP (1 equiv) in DCE (0.1 mL) were added. To the resulting mixture was added EDAC (2 equiv). The mixture was then shaken at 50 °C overnight and concentrated in vacuo. The residue was redissolved in DMF (0.3 mL) and subjected to preparative LCMS purification, giving an ester.

Example 1 (Compound 101)

3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methylbenzoate

[0182]



[0183] 4-Methylbenzoic acid and 3-ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one were treated as described in the general procedure 1, giving the title compound.


¹H NMR (600 MHz, DMSO-d₆) δ 7.92 - 7.72 (m, 2H), 7.41 (t, J = 5.7 Hz, 1H), 7.32 (d, J = 8.1 Hz, 2H), 4.24 (t, J = 6.0 Hz, 2H), 4.15 (t, J = 6.8 Hz, 2H), 3.63 - 3.53 (m, 2H), 3.54 - 3.45 (m, 2H), 2.98 (d, J = 5.8 Hz, 2H), 2.79 - 2.67 (m, 4H), 2.38 (s, 3H), 2.18 (p, J = 6.5 Hz, 2H), 1.45 - 1.32 (m, 4H), 1.10 (t, J = 7.5 Hz, 3H). HPLC-Retention time (XE Metode 7 CM): 2.12 minutes.

Detected "M+1"-mass: 426.24.

Example 2 (Compound 102)

3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-ethylbenzoate

[0184]



[0185] 3-Ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one and 4-ethylbenzoic acid were treated as described in General procedure 2, giving the title compound.


¹H NMR (300 MHz, DMSO-d₆) δ 7.85 (d, J = 8.3 Hz, 2H), 7.48 - 7.29 (m, 3H), 4.24 (t, J = 6.0 Hz, 2H), 4.14 (t, J = 6.7 Hz, 2H), 3.69 - 3.40 (m, 4H), 2.98 (d, J = 5.7 Hz, 2H), 2.83 - 2.59 (m, 6H), 2.19 (p, J = 6.4 Hz, 2H), 1.36 (t, J = 5.4 Hz, 4H), 1.19 (t, J = 7.6 Hz, 3H), 1.10 (t, J = 7.5 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 2.23 minutes.

Detected "M+1"-mass: 440.25.

Example 3 (Compound 103)

[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2-methyl-propyl]
4-(methylsulfamoyl)benzoate

[0186]

[0187] 3-Ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one and 4-(methylsulfamoyl)benzoic acid were treated as described in General procedure 2, giving the title compound.

¹H NMR (600 MHz, DMSO-d₆) δ 8.17 - 8.06 (m, 2H), 7.94 - 7.84 (m, 2H), 7.71 (s, 1H), 7.44 (t, J = 5.8 Hz, 1H), 4.24 (dd, J = 11.0, 6.2 Hz, 1H), 4.20 - 4.07 (m, 2H),


3.99 (dd, J = 14.2, 6.7 Hz, 1H), 3.60 (dt, J = 11.8, 4.9 Hz, 1H), 3.57 - 3.42 (m, 3H), 3.02 - 2.93 (m, 2H), 2.79 (d, J = 16.6 Hz, 1H), 2.73 - 2.65 (m, 3H), 2.57 - 2.52 (m, 1H), 2.43 (s, 3H), 1.38 (t, J = 5.5 Hz, 2H), 1.37 - 1.27 (m, 2H), 1.07 (t, J = 7.5 Hz, 3H), 1.01 (d, J = 6.9 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 2.02 minutes.

Detected "M+1"-mass: 519.23.

Example 4 (Compound 104)

[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2-methyl-propyl]
4-(dimethylsulfamoyl)benzoate

[0188]

[0189] 3-Ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one and 4-(dimethylsulfonyl)benzoic acid were treated as described in General procedure 2, giving the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 8.19 - 8.07 (m, 2H), 7.90 - 7.82 (m, 2H), 7.36 (t, J = 5.7 Hz, 1H), 4.34 - 3.91 (m, 4H), 3.65 - 3.39 (m, 4H), 2.98 (d, J = 5.7 Hz, 2H), 2.89 - 2.61 (m, 10H), 2.75-2.60 (m, 1H), 1.48 - 1.24 (m, 4H), 1.06 (t, J = 7.5 Hz, 3H), 1.02 (d, J = 7.0 Hz, 3H).


HPLC-Retention time (XE Metode 7 CM): 2.12 minutes.

Detected "M+1"-mass: 533.24.

Example 5 (Compound 105)

[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2-methyl-propyl] 4-methylsulfonylbenzoate

[0190]



[0191] 3-Ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one and 4-methylsulfonylbenzoic acid were treated as described in General procedure 2, giving the title compound.

¹H NMR (600 MHz, DMSO-d₆) δ 8.17 - 8.10 (m, 2H), 8.08 - 8.03 (m, 2H), 7.40 (t, J = 5.8 Hz, 1H), 4.25 (dd, J = 11.0, 6.3 Hz, 1H), 4.18 (dd, J = 11.0, 5.1 Hz, 1H), 4.11 (dd, J = 14.1, 7.8 Hz, 1H), 3.99 (dd, J = 14.1, 6.6 Hz, 1H), 3.60 (dt, J = 11.8, 4.9 Hz, 1H), 3.58 - 3.43 (m, 3H), 3.28 (s, 3H), 3.04 - 2.91 (m, 2H), 2.78 (d, J = 16.5 Hz, 1H), 2.74 - 2.64 (m, 3H), 2.60 - 2.51 (m, 1H), 1.39 (t, J = 5.4 Hz, 2H), 1.37 - 1.27 (m, 2H), 1.08 (t, J = 7.5 Hz, 3H), 1.01 (d, J = 6.8 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.99 minutes.

Detected "M+1"-mass: 504.21.

PDE4 assay

[0192] The human PDE4D catalytic domain (UniProt no. Q08499 [S380-L740]) was incubated with a mixture of non-labelled cAMP (cyclic adenosine monophosphate) and fluorescein amidite (FAM) conjugated cAMP and titrated test or reference compound. Following brief incubation the enzymatic reaction was stopped by addition of binding buffer containing nanoparticles with immobilized trivalent metal ions capable of binding 1) AMP phospho groups and 2) terbium (Tb) donor fluorophores. Subsequent excitation of the Tb donor triggers time-resolved FRET to adjacent FAM acceptor molecules resulting in light emission. In the presence of a PDE4 inhibitor, AMP generation was reduced resulting in a lower fluorescence signal. The cAMP phosphodiester is not bound by the detection system.

[0193] The results were calculated as the molar concentrations resulting in 50% inhibition of the substrate cleavage compared to controls samples, and are expressed as a range of IC₅₀ (nM).

[0194] The results are shown in Table 1 below.

PDE4 IC₅₀ ranges

[0195]

* indicates that IC₅₀ values are > 500 nM
** indicates that IC₅₀ values are > 100 and < 500 nM
*** indicates that IC₅₀ values are < 100 nM

Table 1

Example	Compound	PDE4 IC ₅₀ range
1	101	***
2	102	***
3	103	***
4	104	***
5	105	***

[0196] The Examples 6 - 139 shown in Table 2 were prepared by reacting Compound 011 as described in General Procedure (G.P.) 1 or 2 with the appropriate acid:








Table 2

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
6	106		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl acetate	2	1.87	**
7	107		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methylpropanoate	2	1.96	**
8	108		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxypropanoate	1	1.80	*
9	109		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl cyclopentanecarboxylate	2	2.08	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
10	110		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-hydroxycyclobutanecarboxylate	1	1.71	**
15	111		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-ethylbutanoate	2	2.15	***
25	112		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-isopropoxyacetate	1	1.89	*
30	113		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methylsulfanylpropanoate	1	1.94	**
40	114		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3,3-difluorocyclobutanecarboxylate	1	2.00	**
50	115		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methyl benzoate	1	2.13	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
16	116		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methyl benzoate	1	2.13	***
17	117		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-hydroxybenzoate	2	2.08	***
18	118		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluorobenzoate	2	2.08	***
19	119		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluorobenzoate	2	2.04	***
20	120		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluorobenzoate	1	2.08	***
21	121		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl cis-4-hydroxycyclohexanecarboxylate	1	1.79	**
22	122		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl trans-4-hydroxycyclohexanecarboxylate	1	1.79	**

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
23	123		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-cyanobenzoate	2	2.00	***
24	124		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-cyanobenzoate	2	2.02	***
25	125		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-cyanobenzoate	1	1.98	***
26	126		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3,4-dimethylbenzoate	2	2.20	***
27	127		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2,4-dimethylbenzoate	2	2.23	***
28	128		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxybenzoate	1	2.05	***
29	129		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxybenzoate	1	2.07	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
30	130		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methoxybenzoate	1	1.98	***
31	131		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methylsulfonylpropanoate	1	1.72	*
32	132		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-4-methylbenzoate	2	2.17	***
33	133		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-4-methylbenzoate	2	2.12	***
34	134		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-fluoro-2-methylbenzoate	2	2.17	***
35	135		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluoro-3-methylbenzoate	1	2.17	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
36	136		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-2-methylbenzoate	2	2.17	***
37	137		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-5-methylbenzoate	1	2.13	***
38	138		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluoro-2-methylbenzoate	2	2.17	***
39	139		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-6-methylbenzoate	2	2.13	***
40	140		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chlorobenzoate	1	2.10	***
41	141		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chlorobenzoate	1	2.16	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
42	142		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chlorobenzoate	1	2.18	***
43	143		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3,5-difluorobenzoate	2	2.13	***
44	144		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3,4-difluorobenzoate	2	2.12	***
45	145		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2,3-difluorobenzoate	2	2.08	***
46	146		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2,6-difluorobenzoate	2	2.05	***
47	147		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2,5-difluorobenzoate	2	2.07	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
48	148		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxycyclohexanecarboxylate	1	1.95	**
49	149		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4,4-difluorocyclohexanecarboxylate	1	2.09	**
50	150		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-acetyl benzoate	2	2.00	***
51	151		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-acetyl benzoate	2	1.99	***
52	152		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-cyano-3-fluorobenzoate	2	2.07	***
53	153		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-cyano-5-fluorobenzoate	2	2.06	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
54	154		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-cyano-2-fluorobenzoate	1	2.04	***
55	155		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-cyano-4-fluorobenzoate	2	2.06	***
56	156		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-cyano-2-fluorobenzoate	2	2.00	***
57	157		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 1,3-benzodioxole-5-carboxylate	1	2.03	***
58	158		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxy-3-methylbenzoate	1	2.15	***
59	159		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxy-4-methylbenzoate	2	2.19	***
60	160		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methoxy-4-methylbenzoate	2	2.05	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
61	161		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-ethoxybenzoate	2	2.16	***
62	162		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-4-methoxybenzoate	2	2.08	***
63	163		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-5-methoxybenzoate	2	2.13	***
64	164		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-4-methoxybenzoate	2	2.07	***
65	165		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluoro-3-methoxybenzoate	2	2.10	***
66	166		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-3-methoxybenzoate	2	2.04	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
67	167		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-5-methoxybenzoate	2	2.07	***
68	168		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-fluoro-2-methoxybenzoate	1	2.02	***
69	169		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-6-methoxybenzoate	1	2.04	***
70	170		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-2-methoxybenzoate	2	2.08	***
71	171		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluoro-2-methoxybenzoate	2	2.02	***
72	172		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-4-methylbenzoate	2	2.26	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
73	173		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-4-methylbenzoate	2	2.19	***
74	174		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-2-methylbenzoate	2	2.26	***
75	175		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-chloro-2-methylbenzoate	2	2.25	***
76	176		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(difluoromethyl)benzoate	2	2.12	***
77	177		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-4-fluorobenzoate	2	2.15	***
78	178		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-2-fluorobenzoate	2	2.15	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
79	179		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-5-fluorobenzoate	2	2.21	***
80	180		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-6-fluorobenzoate	2	2.12	***
81	181		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chloro-3-fluorobenzoate	2	2.21	***
82	182		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-5-fluorobenzoate	2	2.14	***
83	183		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-chloro-2-fluorobenzoate	2	2.15	***
84	184		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-4-fluorobenzoate	2	2.20	***







(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
85	185		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chloro-2-fluorobenzoate	2	2.18	***
86	186		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-4-methoxybenzoate	2	2.14	***
87	187		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chloro-3-methoxybenzoate	2	2.19	***
88	188		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-5-methoxybenzoate	2	2.22	***
89	189		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-4-methoxybenzoate	2	2.14	***
90	190		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-5-methoxybenzoate	2	2.14	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
91	191		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-3-methoxybenzoate	2	2.08	***
92	192		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chloro-2-methoxybenzoate	2	2.12	***
93	193		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-chloro-2-methoxybenzoate	2	2.10	***
94	194		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-2-methoxybenzoate	2	2.16	***
95	195		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-(trifluoromethyl)benzoate	1	2.16	***
96	196		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(trifluoromethyl)benzoate	2	2.22	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
97	197		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(trifluoromethyl)benzoate	2	2.24	***
98	198		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methylsulfonylbenzoate	1	1.91	***
99	199		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methylsulfonylbenzoate	2	1.90	***
100	200		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methylsulfonylbenzoate	2	1.91	***
101	201		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-sulfamoylbenzoate	2	1.84	***
102	202		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-sulfamoylbenzoate	1	1.85	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
103	203		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methyl-3-(trifluoromethyl)benzoate	2	2.30	***
104	204		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methyl-3-(trifluoromethyl)benzoate	2	2.29	***
105	205		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-methyl-5-(trifluoromethyl)benzoate	2	2.30	***
106	206		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-2-(trifluoromethyl)benzoate	2	2.19	***
107	207		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-6-(trifluoromethyl)benzoate	2	2.18	***
108	208		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-5-(trifluoromethyl)benzoate	2	2.27	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
109	209		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluoro-2-(trifluoromethyl)benzoate	2	2.21	***
110	210		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-fluoro-4-(trifluoromethyl)benzoate	2	2.27	***
111	211		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-fluoro-3-(trifluoromethyl)benzoate	2	2.25	***
112	212		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-5-(trifluoromethyl)benzoate	2	2.21	***
113	213		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-fluoro-4-(trifluoromethyl)benzoate	2	2.24	***
114	214		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-ethylsulfonylbenzoate	2	1.96	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
115	215		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-ethylsulfonfylbenzoate	2	1.97	***
116	216		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(methysulfamoyl)benzoate	2	1.93	***
117	217		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(methysulfamoyl)benzoate	2	1.93	***
118	218		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxy-3-(trifluoromethyl)benzoate	2	2.22	***
119	219		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-methoxy-2-(trifluoromethyl)benzoate	2	2.21	***
120	220		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxy-5-(trifluoromethyl)benzoate	2	2.27	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
121	221		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chloro-2-(trifluoromethyl)benzoate	2	2.30	***
122	222		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-chloro-3-(trifluoromethyl)benzoate	2	2.33	***
123	223		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-3-(trifluoromethyl)benzoate	2	2.25	***
124	224		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 5-chloro-2-(trifluoromethyl)benzoate	2	2.28	***
125	225		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-4-(trifluoromethyl)benzoate	2	2.33	***
126	226		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-4-(trifluoromethyl)benzoate	2	2.30	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
127	227		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-chloro-5-(trifluoromethyl)benzoate	2	2.35	***
128	228		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2-chloro-5-(trifluoromethyl)benzoate	2	2.28	***
129	229		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-isopropylsulfonylbenzoate	2	2.02	***
130	230		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-isopropylsulfonylbenzoate	2	2.02	**
131	231		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(dimethylsulfonyl)benzoate	2	2.03	***
132	232		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-cyclopentylsulfonylbenzoate	2	2.13	**

(continued)


Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
133	233		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-pyrrolidin-1-ylsulfonylbenzoate	2	2.11	
134	234		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-pyrrolidin-1-ylsulfonylbenzoate	2	2.10	
135	235		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 2,4-bis(trifluoromethyl)benzoate	2	2.34	***
136	236		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3,4-bis(trifluoromethyl)benzoate	2	2.37	***
137	237		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-morpholinosulfonylbenzoate	2	2.02	
138	238		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-morpholinosulfonylbenzoate	2	2.03	

(continued)

Ex.	Cmpd	Structure	IUPAC Name	G.P.	HPLC Rt (min)	PDE4 IC ₅₀ range
139	239		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(4-acetylpiperazin-1-yl)sulfonylbenzoate	2	1.95	

Preparation 18 (Compound 240)


3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-formylbenzoate

[0197]

[0198] To a solution of 3-ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one (250 mg, 0.81 mmol) in MeCN (3 mL) was added 4-formylbenzoic acid (147 mg, 0.98 mmol), EDAC (187 mg, 0.98 mmol) and DMAP (10 mg, 0.082 mmol). The mixture was stirred overnight at 50 °C before it was evaporated to dryness under vacuum. Silica gel (100-200 mesh) column chromatography (0 to 15% MeOH in DCM as eluent) afforded the title compound.

Preparation 19 (Compound 241)

4-[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propoxycarbonyl]benzoic acid

[0199]

[0200] Jones Reagent (5 mL) was at 0 °C added to a solution of 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-formylbenzoate (308 mg, 0.701 mmol) in acetone/water (45 mL, 2:1). The solution was stirred at room temperature for 18 hours before isopropanol (5 mL) was added. The obtained mixture was stirred for another 30 min, diluted with water and extracted three times with DCM. The combined organic phases were

concentrated in vacuo. Silica gel (100-200 mesh) column chromatography (0 to 15% MeOH in DCM as eluent) afforded the title compound.

¹H NMR (DMSO-d₆) δ: 13.34 (s, 1H), 8.12 - 7.91 (m, 4H), 7.39 (t, J = 5.7 Hz, 1H), 4.31 (t, J = 5.9 Hz, 2H), 4.17 (t, J = 6.6 Hz, 2H), 3.65 - 3.40 (m, 4H), 2.99 (d, J = 5.7 Hz, 2H), 2.80 - 2.64 (m, 4H), 2.29 - 2.15 (m, 2H), 1.38 (t, J = 5.4 Hz, 4H), 1.09 (t, J = 7.5 Hz, 3H).

General Procedure 3: Amide formation

[0201] DMF (0.2 mL) solutions of the acid (0.022 mmol), DIPEA (3 equiv.) and HATU (1.2 equiv.) were added to a vial containing an amine (2 equiv.) and the mixture was shaken at room temperature overnight. The crude was subjected to preparative LCMS purification, giving an amide.

[0202] The Examples 140 - 149 in Table 3 were prepared by reacting Compound 241 as described in General Procedure 3 with the appropriate amine:

Table 3


Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
140	242		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)propyl 4-carbamoylbenzoate	1.78	***
141	243		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-cazepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(methylcarbamoyl)benzoate	1.83	***
142	244		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-cazepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(ethylcarbamoyl)benzoate	1.89	***
143	245		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-cazepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(dimethylcarbamoyl)benzoate	1.86	**

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
144	246		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(isopropylcarbamoyl)benzoate	1.97	***
145	247		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(pyrrolidine-1-carbonyl)benzoate	1.93	**
146	248		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(piperidine-1-carbonyl)benzoate	2.03	***
147	249		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(morpholine-4-carbonyl)benzoate	1.86	**
148	250		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(4-methylpiperazine-1-carbonyl)benzoate	1.67	**
149	251		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 4-(4-acetylpiperazine-1-carbonyl)benzoate	1.80	**

Preparation 20 (Compound 252)


3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-formylbenzoate

[0203]

[0204] To a solution of 3-ethyl-1-(3-hydroxypropyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one (250 mg, 0.81 mmol) in MeCN (3 mL) was added 3-formylbenzoic acid (147 mg, 0.98 mmol), EDAC (187 mg, 0.98 mmol) and DMAP (10 mg, 0.082 mmol). The mixture was stirred overnight at 50 °C before it was evaporated to dryness under vacuum. Silica gel (100-200 mesh) column chromatography (0 to 15% MeOH in DCM as eluent) afforded the title compound.

Preparation 21 (Compound 253)

3-[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propoxycarbonyl]benzoic acid

[0205]

[0206] Jones Reagent (5 mL) was at 0 °C added to a solution of 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-formylbenzoate (341 mg, 0.776 mmol) in acetone/water (45 mL, 2:1). The solution was stirred at room temperature for 18 hours before isopropanol (5 mL) was added.

The obtained mixture was stirred for another 30 min, diluted with water and extracted three times with DCM. The combined organic phases were concentrated in vacuo. Silica gel (100-200 mesh) column chromatography (0 to 15% MeOH in DCM as eluent) afforded the title compound.

¹H NMR (DMSO-d₆) δ: 13.31 (s, 1H), 8.51 - 8.42 (m, 1H), 8.23 - 8.10 (m, 2H), 7.71 - 7.61 (m, 1H), 7.44 - 7.32 (m, 1H), 4.31 (t, J = 6.0 Hz, 2H), 4.16 (t, J = 6.7 Hz, 2H), 3.66 - 3.38 (m, 4H), 2.99 (d, J = 5.7 Hz, 2H), 2.79 - 2.63 (m, 4H), 2.31 - 2.14 (m, 2H), 1.38 (t, J = 5.3 Hz, 4H), 1.09 (t, J = 7.5 Hz, 3H).

[0207] The Examples 150 - 159 in Table 4 were prepared by reacting Compound 253 as described in General Procedure 3 with the appropriate amine:

Table 4

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
150	254		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-carbamoylbenzoate	1.79	***
151	255		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(methylecarbamoyl)benzoate	1.83	***
152	256		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(ethylcarbamoyl)benzoate	1.89	***
153	257		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(dimethylecarbamoyl)benzoate	1.86	***
154	258		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(isopropylcarbamoyl) benzoate	1.96	***
155	259		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(pyrrolidine-1-carbonyl)benzoate	1.93	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
156	260		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(piperidine-1-carbonyl)benzoate	2.03	***
157	261		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(morpholine-4-carbonyl)benzoate	1.86	***
158	262		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(4-methylpiperazine-1-carbonyl)benzoate	1.67	**
159	263		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)propyl 3-(4-acetylpiperazine-1-carbonyl)benzoate	1.80	***

[0208] The following Examples 160 - 251 in Table 5 were prepared by reacting Compound 018 as described in General Procedure 2 with the appropriate acid:







Table 5

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
160	264		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methylpropanoate	2.05	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
161	265		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] cyclopenta ne ca rboxylate	2.18	***
162	266		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-ethylbutanoate	2.24	***
163	267		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] benzoate	2.13	***
164	268		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methyl benzoate	2.22	***
165	269		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluorobenzoate	2.17	***
166	270		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluorobenzoate	2.16	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
167	271		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-cyanobenzoate	2.09	***
168	272		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-cyanobenzoate	2.10	***
169	273		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3,4-dimethylbenzoate	2.29	***
170	274		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,4-dimethylbenzoate	2.32	***
171	275		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxybenzoate	2.15	***
172	276		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-4-methylbenzoate	2.22	***







(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
173	277		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-fluoro-2-methylbenzoate	2.26	***
174	278		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-6-methylbenzoate	2.21	***
175	279		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlorobenzoate	2.27	***
176	280		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,6-difluorobenzoate	2.13	***
177	281		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,5-difluorobenzoate	2.16	***
178	282		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3,4-difluorobenzoate	2.21	***







(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
179	283		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,3-difluorobenzoate	2.17	***
180	284		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-acetylbenzoate	2.09	***
181	285		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-acetylbenzoate	2.07	***
182	286		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-cyano-5-fluoro-benzoate	2.14	***
183	287		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-cyano-2-fluoro-benzoate	2.08	***
184	288		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-cyano-4-fluoro-benzoate	2.14	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
185	289		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-cyano-3-fluoro-benzoate	2.16	***
186	290		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methoxy-3-methylbenzoate	2.24	***
187	291		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methoxy-4-methylbenzoate	2.14	***
188	292		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluoro-4-methoxy-benzoate	2.16	***
189	293		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-4-methoxy-benzoate	2.16	***
190	294		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-3-methoxy-benzoate	2.13	***







(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
191	295		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-5-methoxy-benzoate	2.16	***
192	296		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluoro-2-methoxy-benzoate	2.16	***
193	297		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluoro-3-methoxy-benzoate	2.18	***
194	298		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-6-methoxy-benzoate	2.12	***
195	299		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluoro-2-methoxy-benzoate	2.11	***
196	300		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-2-methyl-benzoate	2.37	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
197	301		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(difluoromethyl)benzoate	2.20	***
198	302		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chloro-2-fluoro-benzoate	2.24	***
199	303		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-3-fluoro-benzoate	2.30	***
200	304		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-5-fluoro-benzoate	2.22	***
201	305		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-3-fluoro-benzoate	2.22	***
202	306		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-chloro-2-fluoro-benzoate	2.25	***







(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
203	307		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-6-fluoro-benzoate	2.20	***
204	308		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chloro-4-fluoro-benzoate	2.29	***
205	309		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-2-fluoro-benzoate	2.27	***
206	310		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chloro-4-methoxy-benzoate	2.23	***
207	311		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-3-methoxy-benzoate	2.16	***
208	312		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-3-methoxy-benzoate	2.27	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
209	313		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chloro-2-methoxybenzoate	2.25	***
210	314		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-chloro-2-methoxybenzoate	2.19	***
211	315		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-2-methoxybenzoate	2.21	***
212	316		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(trifluoromethyl)benzoate	2.32	***
213	317		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-l-yl)-2-methyl-propyl] 3-methylsulfonylbenzoate	1.97	***
214	318		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methylsulfonylbenzoate	1.98	***







(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
215	319		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-sulfamoylbenzoate	1.92	***
216	320		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-sulfamoylbenzoate	1.92	***
217	321		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methyl-3-(trifluoromethyl)benzoate	2.38	***
218	322		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methyl-5-(trifluoromethyl)benzoate	2.38	***
219	323		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluoro-4-(trifluoromethyl)benzoate	2.35	***
220	324		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-5-(trifluoromethyl)benzoate	2.30	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
221	325		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluoro-4-(trifluoromethyl)benzoate	2.32	***
222	326		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluoro-2-(trifluoromethyl)benzoate	2.29	***
223	327		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-fluoro-2-(trifluoromethyl)benzoate	2.28	***
224	328		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluoro-3-(trifluoromethyl)benzoate	2.34	***
225	329		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluoro-5-(trifluoromethyl)benzoate	2.35	***
226	330		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-ethylsulfonylbenzoate	2.04	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
227	331		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-ethylsulfonylbenzoate	2.05	***
228	332		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(methylsulfamoyl)benzoate	2.00	***
229	333		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methoxy-2-(trifluoromethyl)benzoate	2.28	***
230	334		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methoxy-3-(trifluoromethyl)benzoate	2.30	***
231	335		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-methoxy-2-(trifluoromethyl)benzoate	2.28	***
232	336		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxy-5-(trifluoromethyl)benzoate	2.36	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
233	337		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxy-4-(trifluoromethyl)benzoate	2.35	***
234	338		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chloro-4-(trifluoromethyl)benzoate	2.42	***
235	339		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-3-(trifluoromethyl)benzoate	2.33	***
236	340		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-2-(trifluoromethyl)benzoate	2.38	***
237	341		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-4-(trifluoromethyl)benzoate	2.38	***
238	342		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chloro-3-(trifluoromethyl)benzoate	2.42	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
239	343		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-chloro-2-(trifluoromethyl)benzoate	2.36	***
240	344		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chloro-5-(trifluoromethyl)benzoate	2.44	***
241	345		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chloro-5-(trifluoromethyl)benzoate	2.36	***
242	346		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-isopropylsulfonylbenzoate	2.10	***
243	347		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-isopropylsulfonylbenzoate	2.11	***
244	348		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-cyclopentylsulfonylbenzoate	2.21	***

(continued)


Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
245	349		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-pyrrolidin-1-ylsulfonylbenzoate	2.19	***
246	350		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-pyrrolidin-1-ylsulfonylbenzoate	2.17	***
247	351		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3,4-bis(trifluoromethyl)benzoate	2.45	***
248	352		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-morpholinosulfonylbenzoate	2.09	***
249	353		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-morpholinosulfonylbenzoate	2.11	***
250	354		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(4-acetylpiperazin-1-yl)sulfonylbenzoate	2.02	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
251	355		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(4-acetylpiperazin-1-yl)sulfonylbenzoate	2.00	***

Preparation 22 (Compound 356)

[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-formylbenzoate


[0209]

[0210] To a solution of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one (128 mg, 0.40 mmol) in MeCN (3 mL) was added 4-formylbenzoic acid (72 mg, 0.48 mmol), EDAC (92 mg, 0.48 mmol) and DMAP (5 mg, 0.04 mmol). The mixture was shaken at 50 °C for ½ hour and then overnight at room temperature before it was evaporated to dryness under vacuum. Silica gel (100-200 mesh) column chromatography (0 to 15% MeOH in DCM as eluent, R_f(DCM/MeOH : 20/1) = 0.13) afforded the title compound as an oil.

¹H NMR (Chloroform-d) δ: 10.11 (s, 1H), 8.22 - 8.08 (m, 2H), 8.02 - 7.88 (m, 2H), 6.31 - 5.90 (m, 1H), 4.34 - 4.21 (m, 2H), 4.11 (dd, J = 14.0, 7.5 Hz, 1H), 3.96 (dd, J = 14.0, 6.9 Hz, 1H), 3.74 - 3.54 (m, 4H), 3.13 (d, J = 5.8 Hz, 2H), 2.88 (q, J = 7.5 Hz, 2H), 2.79 - 2.60 (m, 3H), 1.64 - 1.40 (m, 4H), 1.24 (t, J = 7.5 Hz, 3H), 1.10 (d, J = 7.0 Hz, 3H).

Preparation 23 (Compound 357)

4-[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propoxy]carbonylbenzoic acid

[0211]

[0212] Jones Reagent (1 mL) was added at room temperature to a solution of [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-formylbenzoate (60 mg, 0.13 mmol) in acetone/water (9 mL, 2:1). The solution was stirred at room temperature for 18 hours before isopropanol (1 mL) was added. The obtained mixture was stirred for another 30 min, diluted with water and extracted three times with DCM. The combined organic phases were concentrated in vacuo. Silica gel (100-200 mesh) column chromatography (R_f(MeOH/AcOH/DCM : 1/0.02/10) = 0.31) afforded the title compound.

¹H NMR (DMSO-d₆) δ: 13.45 (s, 1H), 8.08 - 7.99 (m, 4H), 7.45 (t, J = 5.8 Hz, 1H), 4.23 (dd, J = 11.0, 6.3 Hz, 1H), 4.16 (dd, J = 11.0, 5.0 Hz, 1H), 4.12 (dd, J = 14.1, 7.8 Hz, 1H), 3.99 (dd, J = 14.1, 6.7 Hz, 1H), 3.60 (dt, J = 11.8, 4.9 Hz, 1H), 3.57 - 3.44 (m, 3H), 3.04 - 2.92 (m, 2H), 2.80 (d, J = 16.6 Hz, 1H), 2.75 - 2.66 (m, 3H), 2.57 - 2.48 (m, 1H), 1.39 (t, J = 5.4 Hz, 2H), 1.37 - 1.30 (m, 2H), 1.08 (t, J = 7.5 Hz, 3H), 1.01 (d, J = 6.9 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.94 minutes.

Detected "M+1"-mass: 470.21

[0213] The Examples 252 - 262 in Table 6 were prepared by reacting Compound 357 as described in General Procedure 3 with the appropriate amine:


Table 6

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
252	358		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7 ,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-carbamoyl benzoate	1.86	***
253	359		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7 ,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(methylcarbamoyl)benzoate	1.90	***
254	360		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7 ,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(ethylcarbamoyl)benzoate	1.97	***
255	361		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7 ,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(dimethylcarbamoyl)benzoate	1.93	***

(continued)


Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
256	362		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(isopropylcarbamoyl)benzoate	2.04	***
257	363		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(pyrrolidine-1-carbonyl)benzoate	2.00	***
258	364		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(piperidine-1-carbonyl)benzoate	2.11	***
259	365		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(piperazine-1-carbonyl)benzoate	1.72	***
260	366		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(morpholine-4-carbonyl)benzoate	1.93	***
261	367		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(4-methyl piperazine-1-carbonyl)benzoate	1.73	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
262	368		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(4-acetyl piperazine-1-carbonyl)benzoate	1.86	***

Preparation 24 (Compound 369)

[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-formylbenzoate


[0214]

[0215] To a solution of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one (64 mg, 0.20 mmol) in MeCN (2 mL) was added 3-formylbenzoic acid (36 mg, 0.24 mmol), EDAC (46 mg, 0.24 mmol) and DMAP (5 mg, 0.04 mmol). The mixture was shaken at 50 °C for 30 minutes and then overnight at room temperature before it was evaporated to dryness under vacuum. Silica gel (100-200 mesh) column chromatography (R_f (DCM/MeOH : 20/1) = 0.13) afforded the title compound as an oil.

¹H NMR (Chloroform-d) δ : 10.09 (s, 1H), 8.53 - 8.45 (m, 1H), 8.26 (dt, 1H), 8.10 (dt, J = 7.7, 1.5 Hz, 1H), 7.66 (t, J = 7.7 Hz, 1H), 5.86 (s, 1H), 4.36 - 4.20 (m, 2H), 4.12 (dd, J = 14.0, 7.6 Hz, 1H), 3.97 (dd, J = 14.0, 7.0 Hz, 1H), 3.74 - 3.53 (m, 4H), 3.13 (d, J = 5.8 Hz, 2H), 2.86 (q, J = 7.5 Hz, 2H), 2.76 - 2.68 (m, 2H), 1.69 (s, 1H), 1.64 - 1.43 (m, 4H), 1.22 (t, J = 7.5 Hz, 3H), 1.10 (d, J = 6.9 Hz, 3H).

Preparation 25 (Compound 370)

3-[(2R)-3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propoxy]carbonylbenzoic acid

[0216]

[0217] Jones Reagent (1 mL) was added at room temperature to a solution of [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-formylbenzoate (30 mg, 0.065 mmol) in acetone/water (9 mL, 2:1). The solution was stirred at room temperature for 18 hours before isopropanol (1 mL) was added. The obtained mixture was stirred for another 30 min, diluted with water and extracted three times with DCM . The combined organic phases were concentrated in vacuo. Silica gel (100-200 mesh) column chromatography (R_f(MeOH/AcOH/DCM : 1/0.02/10) = 0.31) afforded the title compound.

¹H NMR (DMSO-d₆) δ: 13.46 (s, 1H), 8.47 (t, J = 1.8 Hz, 1H), 8.20 (dt, J = 7.7, 1.5 Hz, 1H), 8.14 (dt, J = 7.8, 1.5 Hz, 1H), 7.66 (t, J = 7.8 Hz, 1H), 7.44 (t, J = 5.8 Hz, 1H), 4.24 (dd, J = 11.0, 6.2 Hz, 1H), 4.15 (dd, J = 11.0, 5.3 Hz, 1H), 4.10 (dd, J = 14.1, 7.5 Hz, 1H), 3.99 (dd, J = 14.1, 6.9 Hz, 1H), 3.60 (dt, J = 11.8, 4.9 Hz, 1H), 3.57 - 3.44 (m, 3H), 3.03 - 2.93 (m, 2H), 2.78 (d, J = 16.5 Hz, 1H), 2.74 - 2.66 (m, 3H), 2.57 - 2.52 (m, 1H), 1.39 (t, J = 5.4 Hz, 2H), 1.37 - 1.32 (m, 2H), 1.08 (t, J = 7.5 Hz, 3H), 1.00 (d, J = 6.9 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.94 minutes.

Detected "M+1"-mass: 470.21

[0218] The Examples 263 - 273 in Table 7 were prepared by reacting Compound 370 as described in General Procedure 3 with the appropriate amine:

Table 7

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
263	371		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7 ,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-carbamoylbenzoate	1.85	***
264	372		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(methylcarbamoyl)benzoate	1.90	***
265	373		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(ethylcarbamoyl)benzoate	1.96	***
266	374		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(dimethylcarbamoyl)benzoate	1.93	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
267	375		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(isopropylcarbamoyl)benzoate	2.04	***
268	376		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(pyrrolidine-1-carbonyl)benzoate	2.00	***
269	377		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(piperidine-1-carbonyl)benzoate	2.10	***
270	378		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(piperazine-1-carbonyl)benzoate	1.72	***
271	379		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(morpholine-4-carbonyl)benzoate	1.93	***
272	380		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(4-methyl piperazine-1-carbonyl)benzoate	1.73	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
273	381		[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(4-acetylpiperazine-1-carbonyl)benzoate	1.86	***

Preparation 26 (Compound 382)

3-Ethyl-1-[(2S)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0219]



[0220] 3-Ethyl-1-[(2S)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-4-one was prepared in a manner similar to that of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one described in Preparation 11 to Preparation 17 using methyl (2R)-3-hydroxy-2-methyl-propanoate instead of methyl (2S)-3-hydroxy-2-methyl-propanoate as starting material.
¹H NMR (DMSO-d₆) δ: 7.39 (t, J = 5.6 Hz, 1H), 4.62 (t, J = 5.1 Hz, 1H), 4.01 (dd, J = 13.9, 6.5 Hz, 1H), 3.74 (dd, J = 13.9, 7.7 Hz, 1H), 3.68 - 3.47 (m, 4H), 3.27 (t, J = 5.4 Hz, 2H), 3.01 (d, J = 5.7 Hz, 2H), 2.79 - 2.68 (m, 4H), 2.16 - 1.95 (m, 1H), 1.54 - 1.35 (m, 4H), 1.11 (t, J = 7.5 Hz, 3H), 0.81 (d, J = 6.8 Hz, 3H).
HPLC-Retention time (XE Metode 7 CM): 1.66 minutes.
Detected "M+1"-mass: 321.21.
[0221] The following Examples 274 - 279 in Table 8 were prepared by reacting Compound 382 as described in General Procedure 2 with the appropriate acid:

Table 8

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
274	383		[(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] benzoate	2.13	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
275	384		[(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl]4-methyl benzoate	2.22	***
276	385		[(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluorobenzoate	2.17	***
277	386		[(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluorobenzoate	2.16	***
278	387		[(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl]3-methoxybenzoate	2.15	***
279	388		[(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-clazepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlorobenzoate	2.27	***

Preparation 27 (Compound 389)

(3-Hydroxy-2,2-dimethyl-propyl) benzoate

[0222]




[0223] 3-Hydroxy-2,2-dimethylpropanol (93.4 g, 900 mmol) was dissolved in DCM (600 mL), cooled in an ice bath under argon. DMAP (7.3 g, 60 mmol) and pyridine (48.3 mL, 600 mmol) was added, followed by dropwise addition of benzoyl chloride (42.2 g, 34.9 mL, 300 mmol) over 15 min. The mixture was stirred at rt for 3 days and quenched with 4N HCl (90 mL). After phase separation, the organic phase was washed with sat. NaHCO₃ (50 mL), water (50 mL) and brine (50 mL), dried over MgSO₄, filtered and concentrated in vacuo. Column chromatography (10% EtOAc in heptane) afforded the title compound.

Preparation 28 (Compound 390)

(2,2-Dimethyl-3-oxo-propyl) benzoate

[0224]




[0225] DMSO (17.9 mL, 251 mmol) in DCM (229 mL) was cooled to -70 °C in a 2 necked bottle equipped with thermometer, addition funnel and argon flow. Oxalyl chloride (25.5 mL, 297 mmol) in DCM (114 mL) was added slowly while maintaining temperature between -60 to -70 °C. The mixture was stirred at -70 °C for 0.5 hour before (3-hydroxy-2,2-dimethyl-propyl) benzoate (47.6 g, 229 mmol) in DCM (229 mL) was added slowly while maintaining temperature between -60 and -70 °C. The reaction mixture was stirred at -70 °C for 0.5 h. Triethylamine (127 mL, 914 mmol) was added slowly while maintaining temperature between -60 and -70 °C. The mixture was stirred for 2 hours. Water (100 mL) was added and the mixture was phase separated. The aqueous phase was extracted with three times with DCM (3 x 50 mL) and the combined organic phases were washed with water (50 mL), brine (50 mL), dried over MgSO₄ and filtered. Concentrated in vacuo afforded approx. 60 g of reddish oil with precipitation. This mixture was suspended in DCM (20 mL) and filtered. Column chromatography (10 to 35% EtOAc in heptane) afforded the title compound.

¹H NMR (300 MHz, CDCl₃) δ 9.64 (s, 1H), 8.22 - 7.79 (m, 2H), 7.64 - 7.50 (m, 1H), 7.50 - 7.35 (m, 2H), 4.37 (s, 2H), 1.21 (s, 6H).

Preparation 29 (Compound 391)

[3-(2-Tert-butoxycarbonylhydrazino)-2,2-dimethyl-propyl] benzoate

[0226]




[0227] BOC-NHNH₂ (21.8 g, 165 mmol) and AcOH (28.3 mL, 495 mmol) was added to a solution of (2,2-dimethyl-3-oxo-propyl) benzoate (34.0 g, 165 mmol) in MeOH (200 mL). The reaction mixture was stirred at rt for 30 minutes before it was cooled to 0 °C. Sodium cyanoborohydride (15.5 g, 247 mmol) was added portion wise over 5 minutes and the mixture was stirred for an additional 45 minutes at 0 °C. The reaction was quenched with water (100 mL), and the mixture was extracted twice with DCM (2 x 200 mL). The combined organic phases were washed with water (100 mL), dried over MgSO₄, filtered and concentrated in vacuo. Column chromatography (heptane/ethyl acetate 9:1 to 1:1) afforded the title compound as an oil.

¹H NMR (300 MHz, CDCl₃) δ 8.20 - 7.99 (m, 2H), 7.63 - 7.50 (m, 1H), 7.50 - 7.37 (m, 2H), 6.13 (s, 1H), 4.23 - 4.15 (m, 1H), 4.14 (s, 2H), 2.86 (s, 2H), 1.42 (s, 9H), 1.04 (s, 6H).

Preparation 30 (Compound 392)

(3-Hydrazino-2,2-dimethyl-propyl) benzoate ditrifluoroacetic acid salt


[0228]

[0229] [3-(2-Tert-butoxycarbonylhydrazino)-2,2-dimethyl-propyl] benzoate (40 g, 124.1 mmol) was dissolved in DCM. TFA (50 mL) was added and the obtained solution was stirred at rt for 1 hour. Concentrated in vacuo gave the title compound.

^1H NMR (300 MHz, CDCl_3) δ 8.03 - 7.96 (m, 2H), 7.64 - 7.53 (m, 1H), 7.50 - 7.40 (m, 2H), 4.20 (s, 2H), 3.09 (s, 2H), 1.16 (s, 6H).

Preparation 31 (Compound 393)

[3-(3-Ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] benzoate

[0230]


[0231] A mixture of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (15 g, 63.0 mmol), (3-hydrazino-2,2-dimethyl-propyl) benzoate.2TFA (33.6 g, 74.6 mmol) and DIPEA (33.6 g, 251.8 mmol) in ethanol (200 mL) was stirred at rt for 10 minutes before it was heated to reflux for 2 hours. The mixture was then concentrated in vacuo. Water (100 mL) was added and the aqueous mixture was extracted three times with DCM (3 x 100 mL). The combined organic phases were dried over MgSO_4 , filtered and concentrated in vacuo. Chromatography (heptane/ethyl acetate 9:1 to 1:9) gave the title product as an oil.

^1H NMR (300 MHz, CDCl_3) δ : 8.07 - 8.00 (m, 2H), 7.63 - 7.56 (m, 1H), 7.50 - 7.43 (m, 2H), 4.13 (s, 2H), 4.00 (s, 2H), 3.73 - 3.53 (m, 4H), 2.83 (q, $J = 7.5$ Hz, 2H), 2.68 (s, 2H), 2.46 (s, 2H), 1.66 - 1.52 (m, 3H), 1.52 - 1.40 (m, 2H), 1.22 (t, $J = 7.5$ Hz, 3H), 1.13 (s, 6H).

Example 280 (Compound 394)

[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] benzoate

[0232]




[0233] To a solution of [3-(3-ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] benzoate (15 g, 35.3 mmol) in methanesulfonic acid (30 mL) and DCM (50 mL) was added sodium azide (11.5 g, 176.7 mmol) in portions at rt over 1 hour. Then the mixture was stirred at rt for 2 hours. The reaction was not complete. Ethanol (500 mL) was added to the mixture before it was neutralized with solid sodium hydroxide. The solid material was filtered off. The filtrate was concentrated in vacuo. The residue was taken up in chloroform (100 mL). The mixture was left standing for 30 minutes. The solid material was filtered off again. The filtrate was concentrated in vacuo. The residue was purified three times by chromatography (CHCl₃/MeOH 10:1, R_f = 0.17), giving the title compound as a solid.

¹H NMR (600 MHz, DMSO-d₆) δ 7.98 (t, J = 7.7 Hz, 2H), 7.67 (t, J = 7.7 Hz, 1H), 7.54 (t, J = 7.7 Hz, 2H), 7.42 (t, J = 5.7 Hz, 1H), 4.10 (s, 2H), 4.01 (s, 2H), 3.60-3.55 (m, 2H), 3.52-3.45 (m, 2H), 2.97 (d, J = 5.7 Hz, 2H), 2.75 (s, 2H), 2.71 (q, J = 7.5 Hz, 2H), 1.37 (t, J = 5.4 Hz, 4H), 1.09 (t, J = 7.5 Hz, 3H), 1.04 (s, 6H). HPLC-Retention time (XE Metode 7 CM): 2.25 minutes.

Detected "M+1"-mass: 440.25.

Preparation 32 (Compound 395)

3-Ethyl-1-(3-hydroxy-2,2-dimethyl-propyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one



[0235] To a mixture of [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] benzoate (7.0 g, 15.9 mmol) in MeOH (50 mL) was added NaOH (1 g) at rt. The mixture was stirred at 50 °C for 0.5 h until starting material disappeared. Then the solution was neutralized by using 4N HCl and concentrated in vacuo. The residue was taken up in EtOH. The insoluble was filtered off. The filtrate was concentrated in vacuo. The residue was purified by chromatography (MeOH/DCM 1:10, R_f = 0.36). Recrystallized in TBME/heptane gave the title compound as colorless crystals.

¹H NMR (300 MHz, DMSO-d₆) δ 7.40 (t, J = 5.7 Hz, 1H), 4.75 (t, J = 5.3 Hz, 1H), 3.82 (s, 2H), 3.74 - 3.45 (m, 4H), 3.13 (d, J = 5.2 Hz, 2H), 2.99 (d, J = 5.6 Hz, 2H), 2.79 (s, 2H), 2.72 (d, J = 7.5 Hz, 2H), 1.50-1.35 (m, 4H), 1.11 (t, J = 7.5 Hz, 3H), 0.83 (s, 6H).

HPLC-Retention time (XE Metode 7 CM): 1.78 minutes.







Detected "M+1"-mass: 336.22.

[0236] The following Examples 281 - 325 in Table 9 were prepared by reacting Compound 395 as described in General Procedure 2 with the appropriate acid:

Table 9

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
281	396		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] cyclopentanecarboxylate	2.31	***
282	397		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-ethylbutanoate	2.37	***
283	398		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-methyl benzoate	2.34	***
284	399		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-fluorobenzoate	2.29	***
285	400		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-fluorobenzoate	2.28	***
286	401		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-fluorobenzoate	2.25	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
287	402		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-cyanobenzoate	2.20	***
288	403		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-cyanobenzoate	2.22	***
289	404		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-methoxybenzoate	2.27	***
290	405		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-chlorobenzoate	2.39	***
291	406		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-chlorobenzoate	2.38	***
292	407		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-chlorobenzoate	2.31	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
293	408		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3,5-difluorobenzoate	2.34	***
294	409		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3,4-difluorobenzoate	2.33	***
295	410		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-acetyl benzoate	2.20	***
296	411		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-acetyl benzoate	2.18	***
297	412		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-cyano-3-fluoro-benzoate	2.27	***
298	413		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-cyano-2-fluoro-benzoate	2.23	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
299	414		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-ethoxybenzoate	2.37	***
300	415		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-chloro-6-fluoro-benzoate	2.32	***
301	416		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-chloro-4-fluoro-benzoate	2.36	***
302	417		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(trifluoromethyl)benzoate	2.42	***
303	418		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-(trifluoromethyl)benzoate	2.35	***
304	419		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(trifluoromethyl)benzoate	2.44	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
305	420		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-methylsulfonylbenzoate	2.07	***
306	421		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-methylsulfonylbenzoate	2.08	***
307	422		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-methylsulfonylbenzoate	2.07	***
308	423		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-sulfamoylbenzoate	2.01	***
309	424		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-sulfamoylbenzoate	2.01	***
310	425		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-fluoro-5-(trifluoromethyl)benzoate	2.47	***
311	426		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-ethylsulfonylbenzoate	2.14	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
312	427		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-ethylsulfonylbenzoate	2.15	***
313	428		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(methylsulfamoyl)benzoate	2.09	***
314	429		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-chloro-3-(trifluoromethyl) benzoate	2.54	***
315	430		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-chloro-5-(trifluoromethyl) benzoate	2.48	***
316	431		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-isopropylsulfonylbenzoate	2.20	***
317	432		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-isopropylsulfonylbenzoate	2.21	***
318	433		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(dimethylsulfamoyl)benzoate	2.21	***

(continued)


Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
319	434		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-cyclopentylsulfonylbenzoate	2.31	***
320	435		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-pyrrolidin-1-ylsulfonylbenzoate	2.27	***
321	436		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-pyrrolidin-1-ylsulfonylbenzoate	2.29	***
322	437		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-morpholinosulfonylbenzoate	2.21	***
323	438		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-morpholinosulfonylbenzoate	2.19	***
324	439		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(4-acetypiperazin-1-yl)sulfonylbenzoate	2.10	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
325	440		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(4-acetylpiperazin-1-yl) sulfonylbenzoate	2.08	***

Example 326 (Compound 441)

[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2,2-dimethyl-propyl] 4-[benzyl(methyl)sulfamoyl]benzoate


[0237]

[0238] To a mixture of 3-ethyl-1-(3-hydroxy-2,2-dimethyl-propyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one (8.9 mmol, 3 g), 4-[benzyl(methyl)sulfamoyl]benzoic acid (3.2 g, 10.7 mmol) and DMAP (110 mg, 0.9 mmol) in MeCN (70 mL) was added EDAC (10.7 mmol, 2.1 g) at rt. The mixture was stirred at rt for 18 hours. The mixture gradually became a solution. After 18 h, the product was precipitated and filtered off, giving 3.6 g of product as a white solid. The mother liquor was concentrated in vacuo. The residue was purified by chromatography (ethyl acetate/MeOH 20:1, R_f = 0.21), giving 1.3 g of product as a white foam.

¹H NMR (DMSO-d₆) δ: 8.21 - 8.15 (m, 2H), 8.01 - 7.96 (m, 2H), 7.46 (t, J = 5.8 Hz, 1H), 7.41 - 7.36 (m, 2H), 7.35 - 7.28 (m, 3H), 4.17 (d, J = 5.3 Hz, 4H), 4.02 (s, 2H), 3.58 (dt, J = 11.8, 4.9 Hz, 2H), 3.52 - 3.45 (m, 2H), 2.97 (d, J = 5.8 Hz, 2H), 2.75 (s, 2H), 2.68 (q, J = 7.5 Hz, 2H), 2.57 (s, 3H), 1.44 - 1.32 (m, 4H), 1.13 - 1.02 (m, 9H).

Example 327 (Compound 442)

[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(methylsulfamoyl)benzoate

[0239]

[0240] A mixture of [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-[benzyl(methyl)sulfamoyl]benzoate (2.4 g) in methanesulfonic acid (15 mL) was heated at 80 °C for 3 hours. The reaction was cooled to rt and diluted with DCM (50 mL). A solution of NaHCO₃ (25 g) in water (250 mL)

was cooled to approximately 5 °C. To this solution was dropwise added the reaction solution. After phase separation, the aqueous phase was extracted twice with DCM. The combined organic phases were dried over Na₂SO₄ and concentrated in vacuo. The residue was purified by chromatography (ethyl acetate/MeOH), giving the product as a foam. Crystallized of the foam from n-butyl acetate (15 mL), gave the title compound as colorless crystals.

¹H NMR (DMSO-d₆) δ: 8.21 - 8.13 (m, 2H), 7.95 - 7.87 (m, 2H), 7.67 (s, 1H), 7.40 (t, J = 5.6 Hz, 1H), 4.15 (s, 2H), 4.01 (s, 2H), 3.66 - 3.42 (m, 4H), 2.97 (d, J = 5.7 Hz, 2H), 2.74 (s, 2H), 2.70 (q, J = 7.7 Hz, 2H), 2.44 (s, 3H), 1.38 (t, J = 5.4 Hz, 4H), 1.11 - 1.01 (m, 9H).


HPLC-Retention time (XE Metode 7 CM): 2.11 minutes.

Detected "M+1"-mass: 533.23.

Preparation 33 (Compound 443)

4-[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2,2-dimethyl-propoxy]carbonylbenzoic acid

[0241]



[0242] 4-[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propoxy]carbonylbenzoic acid was prepared in a manner similar to that of 4-[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propoxy]carbonylbenzoic acid described in Preparation 22 & 23 using 3-ethyl-1-(3-hydroxy-2,2-dimethyl-propyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one instead of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one as starting material.

[0243] The Examples 328 - 338 in Table 10 were prepared by reacting Compound 443 as described in General Procedure 3 with the appropriate amine:

Table 10

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
328	444		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-carbamoyl benzoate	1.94	***
329	445		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(methylcarbamoyl)benzoate	1.99	***

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
330	446		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 4-(ethylcarbamoyl)benzoate	2.06	***
331	447		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 4-(dimethylcarbamoyl)benzoate	2.02	***
332	448		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 4-(isopropylcarbamoyl)benzoate	2.13	***
333	449		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 4-(pyrrolidine-1-carbonyl) benzoate	2.10	***
334	450		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 4-(piperidine-1-carbonyl) benzoate	2.21	***
335	451		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 4-(piperazine-1-carbonyl) benzoate	1.79	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
336	452		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(morpholine-4-carbonyl) benzoate	2.02	***
337	453		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(4-methyl piperazine-1-carbonyl)benzoate	1.79	***
338	454		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 4-(4-acetyl piperazine-1-carbonyl)benzoate	1.94	***

Preparation 34 (Compound 455)

3-[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2,2-dimethyl-propoxy]carbonylbenzoic acid

[0244]



[0245] 3-[3-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)-2,2-dimethyl-propoxy]carbonylbenzoic acid






was prepared in a manner similar to that of 3-[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propoxy]-carbonylbenzoic acid described in Preparation 24 & 25 using 3-ethyl-1-(3-hydroxy-2,2-dimethyl-propyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one instead of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one as starting material.

[0246] The Examples 339 - 349 in Table 11 were prepared by reacting Compound 455 as described in General Procedure 3 with the appropriate amine:

Table 11

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
339	456		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-carbamoylbenzoate	1.94	
340	457		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(methylcarbamoyl)-benzoate	1.98	***
341	458		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(ethylcarbamoyl)-benzoate	2.05	***
342	459		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(dimethylcarbamoyl)-benzoate	2.02	***
343	460		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(isopropylcarbamoyl)-benzoate	2.13	***
344	461		[3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 3-(pyrrolidine-1-carbonyl) benzoate	2.10	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
345	462		[3-(3-ethyl-4-oxo- spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)- 2,2 -dimethyl-propyl] 3-(piperidine-1-carbonyl) benzoate	2.20	***
346	463		[3-(3-ethyl-4-oxo- spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)- 2,2- dimethyl-propyl] 3-(piperazine-1-carbonyl) benzoate	1.78	***
347	464		[3-(3-ethyl-4-oxo- spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 3-(morpholine-4-carbonyl) benzoate	2.01	***
348	465		[3-(3-ethyl-4-oxo- spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)- 2,2dimethyl-propyl] 3-(4-methylpiperazine-1- carbonyl)benzoate	1.79	***
349	466		[3-(3-ethyl-4-oxo- spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)-2,2- dimethyl-propyl] 3-(4-acetylpiperazine-1- carbonyl)benzoate	1.93	***

Preparation 35 (Compound 467)

[1-[[Tert-butyl(diphenyl)silyl]oxymethyl]cyclopropyl]methanol

[0247]




[0248] To a solution of [1-(hydroxymethyl)cyclopropyl]methanol (5.4 g, 52.94 mmol) and diisopropylethylamine (5.3 g, 105.96 mmol) in DCM (53 mL) was tert-butylchlorodiphenylsilane (4.3 g, 15.88 mmol) added dropwise. The mixture was stirred at 26 °C for 16 hours. On completion, the reaction mixture was concentrated under reduced pressure and the residue was diluted with water (150 mL) and extracted twice with ethyl acetate (2 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over anhydrous Na₂SO₄ and concentrated under reduced pressure to afford the title compound as yellow oil.

Preparation 36 (Compound 468)

[1-[[Tert-butyl(diphenyl)silyl]oxymethyl]cyclopropyl]methyl benzoate

[0249]




[0250] Benzoyl chloride (4.5 mL, 38.8 mmol) was slowly added at 0 °C to a solution of [1-[[tert-butyl(diphenyl)silyl]oxymethyl]cyclopropyl]methanol (11 g, 32.3 mmol) and pyridine (4.0 mL, 48.5 mmol) in DCM (150 mL). The mixture was stirred at 26 °C for 16 hours. On completion, the reaction mixture was diluted with water (200 mL) and extracted twice with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (5% EtOAc in Pet ether as eluent) to afford the title compound as gummy liquid.

¹H NMR (400MHz, CDCl₃): δ 8.02 - 7.96 (m, 2H), 7.65 (dd, J = 1.5, 7.9 Hz, 4H), 7.60 - 7.50 (m, 1H), 7.46 - 7.28 (m, 8H), 4.35 (s, 2H), 3.66 (s, 2H), 1.05 (s, 9H), 0.62 - 0.57 (m, 2H), 0.56 - 0.50 (m, 2H).

Preparation 37 (Compound 469)

[1-(Hydroxymethyl)cyclopropyl]methyl benzoate

[0251]




[0252] TBAF (12.9 g, 49.5 mmol) was added slowly at 0 °C to a solution of [1-[[tert-butyl-(diphenyl)silyl]oxymethyl]cyclopropyl]methyl benzoate (11 g, 24.7 mmol) in THF (250 mL). The reaction mixture was stirred at 26 °C for 16 hours. On completion the reaction was quenched with water (150 mL) and the mixture was extracted twice with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, and concentrated under vacuum to afford the crude title compound as gummy liquid.

Preparation 38 (Compound 470)

(1-Formylcyclopropyl)methyl benzoate

[0253]




[0254] PCC (14.6 g, 67.9 mmol) was added at 0 °C to a solution of [1-(hydroxymethyl)-cyclopropyl]methyl benzoate (7 g, 24.7 mmol) in DCM (500 mL). The mixture was stirred at 26 °C for 2 hours. On completion the reaction mixture was filtered through celite. The filtrate was dried over Na₂SO₄, concentrated and the resulting residue was purified by silica gel column chromatography (15% EtOAc in Pet ether as eluent) to afford the title compound as gummy liquid.

¹H NMR (300MHz, CDCl₃): δ 9.06 (d, J = 1.8 Hz, 1H), 8.02 (br d, J = 7.3 Hz, 2H), 7.61 - 7.50 (m, 1H), 7.48 - 7.37 (m, 2H), 4.57 (d, J = 1.5 Hz, 2H), 1.41 - 1.31 (m, 2H), 1.30 - 1.24 (m, 2H).

Preparation 39 (Compound 471)

[1-[(2-Tert-butoxycarbonylhydrazino)methyl]cyclopropyl]methyl benzoate

[0255]



[0256] Boc-NH-NH₂ (5.6 g, 86.8 mmol) was added at 0 °C to a solution of (1-formyl-cyclopropyl)methyl benzoate (5 g, 24.5 mmol) in MeOH (150 mL). The mixture was stirred at 26 °C for 30 minutes. NaBH₃CN (3.2 g, 36.7 mmol) and AcOH (1 mL) were added to the reaction mixture at 0 °C which was then stirred for another 16 hours at 26 °C. On completion, the reaction was treated with aq. NaHCO₃ solution, and then extracted twice with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (10% EtOAc in Pet ether as eluent) to afford the title compound as gummy liquid.


¹H NMR (400MHz, CDCl₃): δ 8.13 - 7.99 (m, 2H), 7.59 - 7.53 (m, 1H), 7.48 - 7.41 (m, 2H), 6.06 (br s, 1H), 4.29 (s, 2H), 4.05 (br s, 1H), 2.91 (s, 2H), 1.45 (s, 9H), 0.66 - 0.61 (m, 2H), 0.61 - 0.55 (m, 2H).

Preparation 40 (Compound 472)

[1-(Hydrazinomethyl)cyclopropyl]methyl benzoate hydrochloride

[0257]

5



10

[0258] Dioxane.HCl (50 mL) was added to a solution of [1-[(2-tert-butoxycarbonylhydrazino)methyl]cyclopropyl]methyl benzoate (2 g, 6.2 mmol) in DCM (10 mL) and the mixture was stirred at 26 °C for 16 hours. On completion excess solvent was evaporated under vacuum and washed with diethyl ether to afford the crude title compound as brown solid, which was used in the next step without further purification.

Preparation 41 (Compound 473)


15

[1-[(3-Ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)methyl]cyclopropyl]methyl benzoate

[0259]

20

25



30

35

[0260] [1-(Hydrazinomethyl)cyclopropyl]methyl benzoate hydrochloride (1.3 g, 5.0 mmol) were added to a solution of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (1 g, 4.2 mmol) and TEA (1.27 g, 12.6 mmol) in EtOH (150 mL). The mixture was stirred at 85 °C for 3 h. On completion, excess solvent was evaporated under vacuum and the residue was diluted with water and extracted twice with EtOAc (2 x 50 mL). The combined organic layer were washed with brine (50 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (60% EtOAc in Pet ether as eluent) to afford the title compound as brown gummy liquid.

40

¹H NMR (400MHz, CDCl₃): δ 8.06 - 7.93 (m, 2H), 7.65 - 7.54 (m, 1H), 7.51 - 7.38 (m, 2H), 4.13 (d, J=11.7 Hz, 4H), 3.68 - 3.54 (m, 4H), 2.83 (q, J = 7.7 Hz, 2H), 2.71 (s, 2H), 2.41 (s, 2H), 1.59 - 1.54 (m, 2H), 1.50 - 1.40 (m, 2H), 1.30 - 1.19 (m, 3H), 0.89 - 0.82 (m, 2H), 0.81 - 0.76 (m, 2H).

Example 350 (Compound 474)


45

[1-[(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydro-pyran]-1-yl)methyl]cyclopropyl]methyl benzoate

[0261]

50

55



[0262] NaN₃ (0.383 g, 5.9 mmol) was added at 0 °C to a solution of [1-[(3-ethyl-4-oxo-spiro[5,7]-dihydroindazole-6,4'-tetrahydropyran]-1-yl)methyl]cyclopropylmethyl benzoate (1.0 g, 1.18 mmol) in MeSO₃H (2.27 g, 5.9 mmol). The mixture was stirred at 26 °C for 16 hours. On completion, the reaction was treated with aq. NaHCO₃ solution at 0 °C, and then extracted twice with EtOAc (2 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (2% MeOH in DCM as eluent) to afford the title compound as colorless solid.

¹H NMR (300MHz, DMSO-*d*₆): δ 8.00 - 7.82 (m, 2H), 7.69 - 7.61 (m, 1H), 7.56 - 7.45 (m, 2H), 7.37 (br t, *J* = 5.7 Hz, 1H), 4.11 (d, *J* = 9.9 Hz, 4H), 3.58 - 3.38 (m, 4H), 2.96 (br d, *J* = 5.5 Hz, 2H), 2.80 - 2.64 (m, 4H), 1.39 - 1.22 (m, 4H), 1.07 (t, *J* = 7.5 Hz, 3H), 0.80 - 0.73 (m, 2H), 0.73 - 0.66 (m, 2H).


HPLC-Retention time (XE Metode 7 CM): 2.14 minutes.

Detected "M+1"-mass: 438.23.

Preparation 42 (Compound 475)

3-Ethyl-1-[[1-(hydroxymethyl)cyclopropyl]methyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0263]



[0264] K₂CO₃ (0.235 g, 1.71 mmol) was added at 0 °C to a solution of [1-[(3-ethyl-4-oxo-spiro[6,8]-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]cyclopropylmethyl benzoate (0.25 g, 0.57 mmol) in MeOH (5 mL). The mixture was stirred at 26 °C for 1 hour. On completion, excess solvent was evaporated under vacuum and the residue was diluted with water and extracted twice with EtOAc (2 x 100 mL). The combined organic layers were washed with brine (25 mL), dried over Na₂SO₄, concentrated and the residue was purified by Neutral Alumina column chromatography (5% MeOH in DCM as eluent) to afford the title compound as colorless solid.

¹H NMR (300MHz, DMSO-*d*₆): δ 7.40 (br t, *J* = 5.7 Hz, 1H), 4.73 (br s, 1H), 3.99 (s, 2H), 3.74 - 3.45 (m, 4H), 3.14 (s, 2H), 3.01 (br d, *J* = 5.8 Hz, 2H), 2.84 (s, 2H), 2.72 (q, *J* = 7.3 Hz, 2H), 1.44 (br t, *J* = 5.3 Hz, 4H), 1.10 (t, *J* = 7.5 Hz, 3H), 0.55 - 0.46 (m, 2H), 0.45 - 0.37 (m, 2H).

HPLC-Retention time (XE Metode 7 CM): 1.70 minutes.

Detected "M+1"-mass: 334.21.

[0265] The following Examples 351 - 355 in Table 12 were prepared by reacting Compound 475 as described in General Procedure 2 with the appropriate acid:


Table 12

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
351	476		[1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl) methyl]cyclopropyl]methyl 4-methyl benzoate	2.22	***
352	477		[1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl) methyl]cyclopropyl]methyl 3-fluorobenzoate	2.18	***
353	478		[1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl) methyl]cyclopropyl]methyl 4-fluorobenzoate	2.17	***
354	479		[1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl) methyl]cyclopropyl]methyl 3-methoxybenzoate	2.17	***
355	480		[1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl) methyl]cyclopropyl]methyl 4-chlorobenzoate	2.28	***

Preparation 43 (Compound 481)

[3-(Bromomethyl)oxeta n-3-yl] methanol

[0266]




[0267] KOH (1.07 g, 19.08 mmol) was added slowly to a solution of 2,2-bis(bromomethyl)-propane-1,3-diol (5 g, 19.08 mmol) in MeOH (50 mL). The mixture was stirred at 60 °C for 16 hours. On completion, excess solvent was evaporated under vacuum and the residue was diluted with water (100 mL) and extracted twice with diethyl ether (2 x 100 mL). The combined organic layers were washed with brine (100 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (50% EtOAc in Pet ether as eluent) to afford the title compound as pale yellow liquid.

¹H NMR (300MHz, DMSO-*d*₆): δ 5.07 (t, *J* = 5.5 Hz, 1H), 4.37 - 4.20 (m, 4H), 3.83 (s, 2H), 3.66 (d, *J* = 5.5 Hz, 2H).

Preparation 44 (Compound 482)

3-Ethylspiro[5,7-dihydro-1H-indazole-6,4'-tetrahydropyran]-4-one

[0268]




[0269] NH₂NH₂·H₂O (0.47 g, 9.45 mmol) was added to a solution of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (1.5 g, 6.3 mmol) and Et₃N (1.6 mL, 12.6 mmol) in EtOH (30 mL). The mixture was stirred at 80 °C for 2 hours. On completion, excess solvent was evaporated under vacuum and the residue was diluted with water and extracted twice with EtOAc (2 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (2% MeOH in DCM as eluent) to afford the title compound as yellow solid.

¹H NMR (400MHz, CDCl₃): δ 3.71 (t, *J* = 5.4 Hz, 4H), 3.04 - 2.91 (m, 2H), 2.86 (s, 2H), 2.53 (s, 2H), 1.68 - 1.53 (m, 4H), 1.33 - 1.21 (m, 4H).

Preparation 45 (Compound 483)

3-Ethylspiro[1,5,6,8-tetrahydropyrazolo[4,3-*c*]azepine-7,4'-tetrahydropyran]-4-one

[0270]




[0271] NaN₃ (0.305 g, 4.7 mmol) were added at 26 °C to a solution of 3-ethylspiro[5,7-dihydro-1H-indazole-6,4'-tetrahydropyran]-4-one (1 g, 4.27 mmol) in CHCl₃ (15 mL) and H₂SO₄ (3 mL). The mixture was stirred for 16 hours. On completion, excess solvent was evaporated under vacuum and the residue was treated with aq. Na₂CO₃ solution, extracted twice with EtOAc (2 x 50 mL). The combined organic layers were washed with brine (50 mL), dried over Na₂SO₄, concentrated and the residue was purified by silica gel (100-200 mesh) column chromatography (3% MeOH in DCM as eluent) to afford the title compound as gummy liquid.

¹H NMR (300MHz, DMSO-*d*₆): δ 12.69 - 12.39 (m, 1H), 12.26 - 12.00 (m, 1H), 9.06 (br s, 1H), 7.39 (br s, 1H), 3.66 - 3.50 (m, 4H), 3.06 - 2.84 (m, 1H), 2.83 - 2.54 (m, 3H), 2.36 (brs, 1H), 2.14 (brs, 1H), 1.53 - 1.37 (m, 4H), 1.18 - 1.04 (m, 2H).

Preparation 46 (Compound 484)

3-Ethyl-1-[[3-(hydroxymethyl)oxetan-3-yl]methyl]spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0272]



[0273] [3-(Bromomethyl)oxetan-3-yl]methanol (0.39 g, 2.2 mmol) and CS₂CO₃ (1.3 g, 4.01 mmol) were added to a solution of 3-ethylspiro[1,5,6,8-tetrahydropyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one (0.5 g, 2.008 mmol) in MeCN (10 mL). The mixture was stirred at 80 °C for 16 hours. On completion, excess solvent was evaporated under vacuum and the residue was dissolved in DCM and filtered through celite. The filtrate was concentrated and the residue was purified by prep HPLC (separations of isomers), to afford the title compound as colorless solid.

¹H NMR (300MHz, DMSO-*d*₆): δ 7.43 (br t, *J* = 5.6 Hz, 1H), 5.06 (br s, 1H), 4.55 (d, *J* = 6.2 Hz, 2H), 4.27 (d, *J* = 6.2 Hz, 2H), 4.21 (s, 2H), 3.68 - 3.49 (m, 4H), 3.42 (br s, 2H), 3.01 (br d, *J* = 5.4 Hz, 2H), 2.81 - 2.66 (m, 4H), 1.42 (br t, *J* = 5.2 Hz, 4H), 1.09 (t, *J* = 7.4 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.62 minutes.

Detected "M+1"-mass: 350.20.

[0274] The following Examples 356 - 361 in Table 13 were prepared by reacting Compound 484 as described in General Procedure 2 with the appropriate acid:

Table 13

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
356	485		[3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl]methyl benzoate	2.02	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
357	486		[3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl]methyl 4-methylbenzoate	2.10	***
358	487		[3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl]methyl 3-fluorobenzoate	2.06	***
359	488		[3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl]methyl 4-fluorobenzoate	2.06	***
360	489		[3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl]methyl 3-methoxybenzoate	2.05	***
361	490		[3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl]methyl 4-chlorobenzoate	2.15	***

Preparation 47 (Compound 491)

(R/S)-Ethyl 3-(3-ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)butanoate

[0275]




[0276] A mixture of 3-ethylspiro[5,7-dihydro-1H-indazole-6,4'-tetrahydropyran]-4-one (220 mg, 0.94 mmol), (R/S)-ethyl 3-bromobutanoate (220 mg, 1.12 mmol), CS_2CO_3 (612 mg, 1.87 mmol) in DMF (2 mL) was stirred at rt for 2 hours. The mixture was diluted with water (10 mL) and EtOAc (10 mL). After phase separation, the aqueous phase was extracted twice with EtOAc (2 x 10 mL). The combined organic phases were dried over MgSO_4 , filtered, and concentrated in vacuo. The residue was purified by prep acidic LCMS to afford the title compound.

^1H NMR (300 MHz, CDCl_3) δ 4.76 - 4.61 (m, 1H), 4.12 - 3.97 (m, 2H), 3.82 - 3.64 (m, 4H), 3.17 (dd, J = 16.8, 9.5 Hz, 1H), 3.03 (d, J = 16.3 Hz, 1H), 2.86 (d, J = 7.5 Hz, 1H), 2.81 (d, J = 7.5 Hz, 1H), 2.75 - 2.71 (m, 1H), 2.70 - 2.65 (m, 1H), 2.50 (s, 2H), 1.70 - 1.56 (m, 4H), 1.50 (d, J = 6.8 Hz, 3H), 1.25 - 1.13 (m, 6H).

Preparation 48 (Compound 492)

Ethyl 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butanoate

[0277]




[0278] To a mixture of ethyl 3-(3-ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)butanoate (135 mg, 0.39 mmol) and sodium azide (76 mg, 1.16 mmol) in chloroform (2 mL) was added methanesulfonic acid (0.50 mL, 7.7 mmol) at rt. The mixture was stirred at rt for 1h. Additional sodium azide (76 mg, 1.16 mmol) and methanesulfonic acid (0.50 mL, 7.7 mmol) was added. The mixture was stirred at rt for another hour and quenched by slow addition of the reaction mixture to saturated, aqueous NaHCO_3 (25 mL). The crude was extracted three times with DCM (3 x 15 mL). The combined organic phases were dried over MgSO_4 , filtered and concentrated in vacuo. Column chromatography (DCM to DCM:MeOH:AcOH 100:10:1 as eluent) afforded the title compound.

^1H NMR (300 MHz, $\text{DMSO}-d_6$) δ 7.40 (t, J = 5.8 Hz, 1H), 4.77 - 4.61 (m, 1H), 3.96 (q, J = 7.1 Hz, 2H), 3.73 - 3.49 (m, 4H), 3.06 - 2.93 (m, 3H), 2.87 - 2.65 (m, 5H), 1.46 (t, J = 5.6 Hz, 4H), 1.36 - 1.30 (m, 3H), 1.13 - 1.01 (m, 6H).

Preparation 49 (Compound 493)

(R/S)-3-Ethyl-1-(3-hydroxy-1-methyl-propyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0279]



[0280] To a solution of ethyl 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butanoate (50 mg, 0.14 mmol) in EtOH (dry) (1.5 mL) under argon was added sodium borohydride (104.1 mg, 2.8 mmol) at rt. The obtained mixture was stirred at this temperature for 2 hours. The mixture was diluted with water (2 mL) and extracted three times with DCM (3 x 5 mL). The combined organic phases were dried over MgSO_4 , filtered and concentrated in vacuo, giving the crude title compound.

^1H NMR (300 MHz, DMSO-d_6) δ 7.36 (t, $J = 5.7$ Hz, 1H), 4.44 (dt, $J = 9.0, 6.3$ Hz, 1H), 3.75 - 3.47 (m, 4H), 3.40 - 3.30 (m, 2H), 3.11 - 2.91 (m, 3H), 2.93 - 2.62 (m, 4H), 1.97 (ddt, $J = 13.9, 9.7, 5.1$ Hz, 1H), 1.79 (ddt, $J = 13.7, 8.3, 5.6$ Hz, 1H), 1.44 (t, $J = 4.6$ Hz, 4H), 1.34 (d, $J = 6.6$ Hz, 3H), 1.10 (t, $J = 7.5$ Hz, 3H). HPLC-Retention time (XE Metode 7 CM): 1.65 minutes.

Detected "M+1"-mass: 322.21.

[0281] Example 362 in Table 14 was prepared by reacting Compound 493 as described in General Procedure 1 with benzoic acid:


Table 14

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC_{50} range
362	494		3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl benzoate	2.12	***

Preparation 50 (Compound 495)

4-Hydrazinobutan-1-ol

[0282]




[0283] To hydrazine monohydrate (20 mL) was added 4-chlorobutan-1-ol (10.8 g, 99.5 mmol) at 0 °C. The solution was

stirred at rt for 2 days. To the solution was added NaOH (4 g, 99.5 mmol). The mixture was stirred for 1 hour and concentrated in vacuo. The residue was taken up in isopropanol (50 mL). The precipitate was filtered off over celite. The filtrate was concentrated in vacuo, giving an oil, which was used without further purification.

Preparation 51 (Compound 496)

3-Ethyl-1-(4-hydroxybutyl)spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one

[0284]




[0285] A reaction solution of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (1.2 g, 5.0 mmol), 4-hydrazinobutan-1-ol (0.8 g, 8 mmol) and triethylamine (2 mL) in EtOH (20 mL) was heated to 100 °C for 1 hour. The solution was concentrated in vacuo. Column chromatography (ethyl acetate/MeOH 9:1, R_f = 0.2), gave the title compound as an orange oil.

¹H NMR (300 MHz, CDCl₃) δ 4.07 (t, J = 7.0 Hz, 2H), 3.78 - 3.59 (m, 6H), 2.85 (q, J = 7.5 Hz, 2H), 2.75 (s, 2H), 2.52 (s, 2H), 2.01 - 1.86 (m, 2H), 1.75 - 1.45 (m, 7H), 1.24 (t, J = 7.5 Hz, 3H).

Preparation 52 (Compound 497)

3-Ethyl-1-(4-hydroxybutyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0286]



[0287] To a mixture of 3-ethyl-1-(4-hydroxybutyl)spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-4-one (1.09 g, 3.56 mmol) and sodium azide (1.2 g, 18 mmol) in CHCl₃ (5 mL) was added methanesulfonic acid (3.4 g, 35.6 mmol). The obtained mixture was stirred at rt overnight. The mixture was then diluted with water and neutralized by addition of NaOH. The mixture was extracted three times with ethyl acetate. The combined organic phases were dried and concentrated in vacuo. Column chromatography (ethyl acetate/MeOH 10:1) afforded the title compound as an oil.

¹H NMR (DMSO-d₆) δ: 7.39 (t, J = 5.7 Hz, 1H), 4.43 (t, J = 5.1 Hz, 1H), 3.96 (t, J = 7.3 Hz, 2H), 3.69 - 3.60 (m, 2H), 3.60 - 3.49 (m, 2H), 3.43 - 3.36 (m, 2H), 3.01 (d, J = 5.7 Hz, 2H), 2.77 - 2.67 (m, 4H), 1.80 - 1.66 (m, 2H), 1.51 - 1.35 (m, 6H), 1.10 (t, J = 7.5 Hz, 3H).

HPLC-Retention time (XE Methode 7 CM): 1.63 minutes.

Detected "M+1"-mass: 322.21.
[0288] The following Examples 363 - 378 in Table 15 were prepared by reacting Compound 497 as described in General Procedure 2 with the appropriate acid:

5

Table 15

10

15

20

25

30

35

40

45

50

55

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
363	498		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl benzoate	2.12	***
364	499		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 3-methylbenzoate	2.22	***
365	500		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-methylbenzoate	2.21	***
366	501		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-fluorobenzoate	2.15	***
367	502		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 3-acetylbenzoate	2.06	***
368	503		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-acetylbenzoate	2.07	**

(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
369	504		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 3-methylsulfonylbenzoate	1.97	**
370	505		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-methylsulfonylbenzoate	1.97	**
371	506		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-ethylsulfonylbenzoate	2.03	**
372	507		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 3-(methylsulfamoyl) benzoate	1.99	***
373	508		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-(methylsulfamoyl) benzoate	2.00	**
374	509		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 3-(dimethylsulfamoyl)benzoate	2.09	***


(continued)

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
375	510		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-(dimethylsulfamoyl)benzoate	2.10	**
376	511		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-cyclopentylsulfonylbenzoate	2.19	**
377	512		4-(3-ethyl-4-oxo-spi ro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-pyrrolidin-1-ylsulfonylbenzoate	2.17	**
378	513		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-morpholinosulfonylbenzoate	2.09	**

Preparation 53 (Compound 514)

4-[4-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butoxycarbonyl]benzoic acid

[0289]



[0290] 4-[4-(3-Ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butoxycarbonyl]benzoic acid was prepared in a manner similar to that of 4-[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-

c]azepine-7,4'-tetrahydropyran]-1-yl)-2-methyl-propoxy]carbonylbenzoic acid described in Preparation 22 & 23 using 3-ethyl-1-(4-hydroxybutyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one instead of 3-ethyl-1-[(2R)-3-hydroxy-2-methyl-propyl]-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one as starting material.




¹H NMR (DMSO-d6) δ: 8.05 (s, 4H), 7.40 (t, J = 5.8 Hz, 1H), 4.32 (t, J = 6.3 Hz, 2H), 4.05 (t, J = 7.2 Hz, 2H), 3.65 - 3.58 (m, 2H), 3.54 - 3.47 (m, 2H), 3.00 (d, J = 5.7 Hz, 2H), 2.75 (s, 2H), 2.72 (q, J = 7.5 Hz, 2H), 1.91 - 1.82 (m, 2H), 1.79 - 1.70 (m, 2H), 1.47 - 1.35 (m, 4H), 1.10 (t, J = 7.5 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.94 minutes.

Detected "M+1"-mass: 470.23.

[0291] The Examples 379 - 381 in Table 16 were prepared by reacting Compound 514 as described in General Procedure 3 with the appropriate amine:


Table 16

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
379	515		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-(methylcarbamoyl)benzoate	1.88	**
380	516		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-(dimethylcarbamoyl)benzoate	1.92	**
381	517		4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo [4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)butyl 4-(morpholine-4-carbonyl)benzoate	1.92	**

Preparation 54 (Compound 518)

Ethyl 2-(3-ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)acetate

[0292]




[0293] A mixture of 9-propanoyl-3-oxaspiro[5.5]undecane-8,10-dione (218 mg, 0.91 mmol), triethylamine (0.255 mL, 1.83 mmol) and ethyl 2-hydrazinoacetate hydrochloride (141 mg, 0.91 mmol) was stirred at 80°C for 1 hour. The reaction mixture was concentrated *in vacuo*. Column chromatography (heptane:EtOAc (3:1) -> heptane:EtOAc (1:2) as eluent) afforded the title compound as pale yellow oil.

¹H NMR (300 MHz, Chloroform-d) δ 4.81 (s, 2H), 4.25 (q, J = 7.1 Hz, 2H), 3.79 - 3.59 (m, 4H), 2.86 (q, J = 7.5 Hz, 2H), 2.73 (s, 2H), 2.53 (s, 2H), 1.72 - 1.52 (m, 4H), 1.38 - 1.16 (m, 6H).

Preparation 55 (Compound 519)

Ethyl 2-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)acetate

[0294]




[0295] Methanesulfonic acid (1 mL) was added to a mixture of ethyl 2-(3-ethyl-4-oxo-spiro[5,7-dihydroindazole-6,4'-tetrahydropyran]-1-yl)acetate (1.4 g, 4.37 mmol) and sodium azide (0.85 g, 13.1 mmol) in dry chloroform. The mixture was stirred at rt for 15 minutes before additional methanesulfonic acid (2 mL) was added. The mixture was stirred at rt for another 15 minutes before additional methanesulfonic acid (2 mL) was added. The mixture was stirred at rt for 45 minutes. Finally, additional sodium azide (852 mg, 13.1 mmol) and methanesulfonic acid (2 mL) was added. The mixture was stirred at rt for 1 hour. Reaction was quenched with saturated, aqueous NaHCO₃ (120 mL) and extracted three times with DCM (3 x 50 mL). The combined organic phases were concentrated *in vacuo*. Column chromatography (20-100% EtOAc in heptane as eluent) gave the title compound.

¹H NMR (300 MHz, CDCl₃) δ 5.74 (t, J = 5.8 Hz, 1H), 4.81 (s, 2H), 4.24 (q, J = 7.1 Hz, 2H), 3.79 - 3.60 (m, 4H), 3.21 (d, J = 5.8 Hz, 2H), 2.91 (q, J = 7.5 Hz, 2H), 2.69 (s, 2H), 1.68 - 1.54 (m, 4H), 1.33 - 1.21 (m, 6H).

Preparation 56 (Compound 520)

3-Ethyl-1-(2-hydroxyethyl)spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-4-one

[0296]



[0297] To a solution of ethyl 2-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)acetate (546 mg, 1.63 mmol) in THF:H₂O (2:1, 12 mL) was added lithium borohydride (185 mg, 4.88 mmol). The mixture was stirred at rt for 1 hour. Saturated, aqueous NH₄Cl (10 mL) was added. The obtained mixture was stirred at rt for 5 minutes and extracted 10 times with DCM (10 x 20 mL) (Note that the product was very soluble in water). The combined organic phases were dried over MgSO₄ and filtered. Evaporation to dryness afforded the title compound. ¹H NMR (300 MHz, DMSO-d₆) δ 7.37 (t, J = 5.7 Hz, 1H), 4.01 (t, J = 5.6 Hz, 2H), 3.75 - 3.46 (m, 6H), 3.00 (d, J = 5.7 Hz, 2H), 2.82 (s, 2H), 2.73 (q, J = 7.5 Hz, 2H), 1.56 - 1.32 (m, 4H), 1.12 (t, J = 7.5 Hz, 3H).

HPLC-Retention time (XE Metode 7 CM): 1.57 minutes.

Detected "M+1"-mass: 294.18.

[0298] Example 382 in Table 17 was prepared by reacting Compound 520 as described in General Procedure 1 with benzoic acid:

Table 17

Ex.	Cmpd	Structure	IUPAC Name	HPLC Rt (min)	PDE4 IC ₅₀ range
382	521		2-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepine-7,4'-tetrahydropyran]-1-yl)ethyl benzoate	1.99	***

REFERENCES CITED IN THE DESCRIPTION

This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.

Patent documents cited in the description


- WO 2007040435 A [0005]
- WO 2001049689 A [0006]
- WO 2008060597 A [0007]
- WO 2004098520 A [0008]
- WO 2008110308 A [0137]

Non-patent literature cited in the description

- M.D. HOUSLAY et al. *Drug Discovery Today*, 2005, vol. 10 (22), 1503-1519 [0002]
- *Journal of Investigative Dermatology*, 1986, vol. 87 (3), 372-6 [0002]
- V. BOSWELL SMITH ; D. SPINA. *Curr. Opinion Investig. Drugs*, 2006, vol. 6 (11), 1136-1141 [0003]
- C. KROEGEL ; M. FOERSTER. *Exp. Opinion Investig. Drugs*, 2007, vol. 16 (1), 109-124 [0003]
- B. LIPWORTH. *Lancet*, 2005, vol. 365, 167-175 [0003]
- BERGE, S.M. *J. Pharm. Sci.*, 1977, vol. 66 (1), 1-19 [0077]
- Goodman & Gilman's *The Pharmacological Basis of Therapeutics*. McGraw-Hill, 1995 [0101]
- REMINGTON. *The Science and Practice of Pharmacy*. 2005 [0104]
- *Encyclopedia of Pharmaceutical Technology*, 1994, vol. 9 [0108]
- *Encyclopedia of Pharmaceutical Technology*. 1989, vol. 2 [0111]
- *Modern Pharmaceutics*. Marcel Dekker, 427-432 [0114]
- *Modern Pharmaceutics*. Marcel Dekker [0114]
- *Encyclopedia of Pharmaceutical Technology*. Marcel Dekker, vol. 10, 191-221 [0114]
- W. AMAREGO ; C. CHAI. *Purification of Laboratory Chemicals*. Butterworth-Heinemann, 2009 [0119]
- GEVER, GABRIEL. *J. Am. Chem. Soc.*, 1954, vol. 76, 1283-1285 [0141]
- GHALI, N. I. *J. Org. Chem.*, 1981, vol. 46 (26), 5413-14 [0141]
- KIM, YONGJU. *ACS Medicinal Chemistry Letters*, 2012, vol. 3 (2), 151-154 [0141]
- BARDAKOS, VASILIOS et al. *Chem Ber.*, 1976, vol. 109, 1898-1910 [0142]
- WANG, XIAO-FENG et al. *J. Med. Chem.*, 2014, vol. 57 (4), 1390-1402 [0142]
- KOCIENSKI ; PHILIP J. *Protecting groups*. Georg Thieme Verlag, 2004 [0143]
- JUNZO OTERA. *Esterification: Methods, Reactions, and Applications*. Wiley-VCH, 2004 [0144]

PATENTKRAV

1. Forbindelse med den almene formel (I)



5 hvor

R₁ og R₄ uafhængigt er udvalgt fra gruppen bestående af hydrogen og (C₁-C₄)alkyl;

R₂ og R₃ uafhængigt er udvalgt fra gruppen bestående af hydrogen og (C₁-C₄)alkyl; eller

R₂ og R₃ sammen med det carbonatom, hvortil de er bundet, danner en cyclopropyl-, cyclobutyl-, cyclopentyl-, oxetanyl-, tetrahydrofuranyl- eller tetrahydropyranylring;

10 n = 0, 1 eller 2;

Q er -O-C(O)-R₅;

R₅ er udvalgt fra gruppen bestående af (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; hvor aryl eventuelt er substitueret med én eller flere substituenten uafhængigt udvalgt fra R₆, eller hvor aryl er eventuelt benzodioxol; og hvor
15 (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl eventuelt er substitueret med én eller flere substituenten uafhængigt udvalgt fra R₇;

R₆ består af halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, -C(O)NR_aR_b og -OR_x;

R₇ består af halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O)NR_aR_b, -C(O)OR_a, -S(O)₂R_x, -
20 S(O)₂NR_aR_b, -OR_x og -SR_x;

R_x består af (C₁-C₆)alkyl og (C₃-C₆)cycloalkyl;

R_a og R_b uafhængigt er udvalgt fra gruppen bestående af hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, eller

R_a og R_b sammen med det nitrogen, hvortil de er bundet, danner et 4-6-leddet heterocycloalkyl,

25 hvor heterocycloalkyl eventuelt er substitueret med én eller flere (C₁-C₄)alkyl, -C(O)R_x; eller farmaceutisk acceptable salte, hydrater eller solvater deraf.

2. Forbindelse ifølge krav 1, hvor

R₁ og R₄ uafhængigt er udvalgt fra gruppen bestående af hydrogen og (C₁-C₄)alkyl;

30 R₂ og R₃ uafhængigt er udvalgt fra gruppen bestående af hydrogen og (C₁-C₄)alkyl; eller

R₂ og R₃ sammen med det carbonatom, hvortil de er bundet, danner en cyclopropyl- eller oxetanylring;

n = 0, 1 eller 2;

Q er -O-C(O)-R₅;

- 5 R₅ er udvalgt fra gruppen bestående af (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, phenyl; hvor phenyl eventuelt er substitueret med én eller flere substituenten uafhængigt udvalgt fra R₆, eller hvor R₅ er benzodioxol; og hvor (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl eventuelt er substitueret med én eller flere substituenten uafhængigt udvalgt fra R₇;
- 10 R₆ består af halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b, -C(O)R_x, -C(O)NR_aR_b og -OR_x;
R₇ består af halogen, hydroxyl, cyano, (C₁-C₄)alkoxy, -C(O)NR_aR_b, -C(O)OR_a, -S(O)₂R_x, -S(O)₂NR_aR_b, -OR_x, og -SR_x;
R_x består af (C₁-C₆)alkyl og (C₃-C₆)cycloalkyl;
- 15 R_a og R_b uafhængigt er udvalgt fra gruppen bestående af hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-C₄)alkyl, eller
R_a og R_b sammen med det nitrogen, hvortil de er bundet, danner et 4-6-leddet heterocycloalkyl, hvor heterocycloalkylet eventuelt er substitueret med én eller flere (C₁-C₄)alkyl, -C(O)R_x; eller et farmaceutisk acceptabelt salt, hydrat eller solvat deraf.

20

3. Forbindelse ifølge krav 1, hvor

R₁ og R₄ uafhængigt er udvalgt fra gruppen bestående af hydrogen og (C₁-C₄)alkyl;

R₂ og R₃ uafhængigt er udvalgt fra gruppen bestående af hydrogen og (C₁-C₄)alkyl; eller

eller R₂ og R₃ sammen med det carbonatom, hvortil de er bundet, danner en cyclopropyl-,

- 25 cyclobutyl-, cyclopentyl-, oxetanyl-, tetrahydrofuranyl- eller tetrahydropyranylring;

n = 0, 1 eller 2;

Q er -O-C(O)-R₅;

R₅ er udvalgt fra gruppen bestående af (C₁-C₆)alkyl, (C₃-C₆)cycloalkyl, halo(C₁-C₆)alkyl, (C₁-C₆)alkoxy, (C₁-C₆)alkoxy(C₁-C₆)alkyl, aryl; hvor aryl eventuelt er substitueret med én eller

- 30 flere substituenten uafhængigt udvalgt fra R₆;

R₆ består af halogen, cyano, hydroxyl, (C₁-C₄)alkyl, halo(C₁-C₄)alkyl, (C₁-C₄)alkyloxy, -S(O)₂R_x, -S(O)₂NR_aR_b og -OR_x;

R_x er (C₁-C₆)alkyl;

R_a og R_b uafhængigt er udvalgt fra gruppen bestående af hydrogen, (C₁-C₄)alkyl, phenyl-(C₁-

C₄)alkyl, eller

R_a og R_b sammen med det nitrogen, hvortil de er bundet, danner et 4-6-leddet heterocycloalkyl, hvor heterocycloalkylet eventuelt er substitueret med én eller flere (C₁-C₄)alkyl; eller et farmaceutisk acceptabelt salt, hydrat eller solvat deraf.

5

4. Forbindelse ifølge et hvilket som helst af de foregående krav, hvor samtlige af R₁, R₂, R₃, R₄ er hydrogen, n er 1, og R₅ er phenyl, eventuelt substitueret med én eller flere substituenten uafhængigt udvalgt fra R₆.

10 5. Forbindelse ifølge et hvilket som helst af de foregående krav udvalgt fra gruppen bestående af

3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methylbenzoat (forbindelse 101);

15 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-ethylbenzoat (forbindelse 102);

[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(methylsulfamoyl)benzoat (forbindelse 103);

[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(dimethylsulfamoyl)benzoat (forbindelse 104);

20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methylsulfonylbenzoat (forbindelse 105); eller farmaceutisk acceptable salte, hydrater eller solvater deraf.

25 6. Forbindelse ifølge et hvilket som helst af kravene 1 til 4 udvalgt fra gruppen bestående af:

3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propylacetat

3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methylpropanoat

30 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxypropanoat

3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl cyclopentancarboxylat

3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 3-hydroxycyclobutancarboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-ethylbutanoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-isopropoxyacetat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methylsulfanylpropanoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3,3-difluorcyclobutancarboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-hydroxybenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl cis-4-hydroxycyclohexancarboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl trans-4-hydroxycyclohexancarboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-cyanobenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-cyanobenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-cyanobenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3,4-dimethylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 2,4-dimethylbenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxybenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxybenzoat
5 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methoxybenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methylsulfonylpropanoat
10 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-4-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-4-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-fluor-2-methyl-benzoat
15 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluor-3-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-2-methyl-benzoat
20 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-5-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluor-2-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-6-methyl-benzoat
25 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlorbenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlorbenzoat
30 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlorbenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3,5-difluorbenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 3,4-difluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2,3-difluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2,6-difluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2,5-difluorbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxycyclohexancarboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4,4-difluorcyclohexancarboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-acetylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-acetylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-cyano-3-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-cyano-5-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-cyano-2-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-cyano-4-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-cyano-2-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 1,3-benzodioxol-5-carboxylat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxy-3-methyl-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxy-4-methyl-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methoxy-4-methyl-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 3-ethoxybenzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-4-methoxy-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-5-methoxy-benzoat
5 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-4-methoxy-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluor-3-methoxy-benzoat
10 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-3-methoxy-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-5-methoxy-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-fluor-2-methoxy-benzoat
15 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-6-methoxy-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-2-methoxy-benzoat
20 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluor-2-methoxy-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-4-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-4-methyl-benzoat
25 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-2-methyl-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-chlor-2-methyl-benzoat
30 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(difluormethyl)benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-4-fluor-benzoat
3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 3-chlor-2-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-5-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-6-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlor-3-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-5-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-chlor-2-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-4-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlor-2-fluor-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-4-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlor-3-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-5-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-4-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-5-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-3-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlor-2-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-chlor-2-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-2-methoxy-benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 2-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(trifluormethyl)benzoat
 5 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methylsulfonylbenzoat
 10 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-sulfamoylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-sulfamoylbenzoat
 15 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methyl-3-(trifluormethyl) benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methyl-3-(trifluormethyl) benzoat
 20 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-methyl-5-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-2-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-6-(trifluormethyl)benzoat
 25 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-5-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluor-2-(trifluormethyl)benzoat
 30 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-fluor-4-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-fluor-3-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 2-fluor-5-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-fluor-4-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-ethylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-ethylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(methylsulfamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(methylsulfamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methoxy-3-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-methoxy-2-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-methoxy-5-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlor-2-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-chlor-3-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-3-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 5-chlor-2-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-4-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-4-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-chlor-5-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2-chlor-5-(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 3-isopropylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-isopropylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(dimethylsulfamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-cyclopentylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-pyrrolidin-1-ylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-pyrrolidin-1-ylsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 2,4-bis(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3,4-bis(trifluormethyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-morpholinsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-morpholinsulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(4-acetylpiperazin-1-yl)sulfonylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-carbamoylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(methylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(ethylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(dimethylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(isopropylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(pyrrolidin-1-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)propyl 4-(piperidin-1-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(morpholin-4-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(4-methylpiperazin-1-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-(4-acetylpiperazin-1-carbonyl)benzoat;
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-carbamoylbenzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(methylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(ethylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(dimethylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(isopropylcarbamoyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(pyrrolidin-1-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(piperidin-1-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(morpholin-4-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(4-methylpiperazin-1-carbonyl)benzoat
 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 3-(4-acetylpiperazin-1-carbonyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methylpropanoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] cyclopentancarboxylat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-ethylbutanoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluorbenzoat
 5 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluorbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-cyanobenzoat
 10 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-cyanobenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3,4-dimethylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,4-dimethylbenzoat
 15 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxybenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-4-methyl-benzoat
 20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-fluor-2-methyl-benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-6-methyl-benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlorbenzoat
 25 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,6-difluorbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,5-difluorbenzoat
 30 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3,4-difluorbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2,3-difluorbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] 4-acetylbenzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-acetylbenzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-cyano-5-fluor-benzoat
5 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-cyano-2-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-cyano-4-fluor-benzoat
10 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-cyano-3-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methoxy-3-methyl-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methoxy-4-methyl-benzoat
15 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluor-4-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-4-methoxy-benzoat
20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-3-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-5-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluor-2-methoxy-benzoat
25 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluor-3-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-6-methoxy-benzoat
30 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluor-2-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-2-methyl-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] 4-(difluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chlor-2-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-3-fluor-benzoat
5 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-5-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-3-fluor-benzoat
10 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-chlor-2-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-6-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chlor-4-fluor-benzoat
15 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-2-fluor-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chlor-4-methoxy-benzoat
20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-3-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-3-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chlor-2-methoxy-benzoat
25 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-chlor-2-methoxy-benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-2-methoxy-benzoat
30 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methylsulfonylbenzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] 2-methylsulfonylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-sulfamoylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-sulfamoylbenzoat
 5 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methyl-3-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-methyl-5-(trifluormethyl)benzoat
 10 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluor-4-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-5-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-fluor-4-(trifluormethyl)benzoat
 15 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluor-2-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-fluor-2-(trifluormethyl)benzoat
 20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluor-3-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluor-5-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-ethylsulfonylbenzoat
 25 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-ethylsulfonylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(methylsulfamoyl)benzoat
 30 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methoxy-2-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methoxy-3-(trifluormethyl)benzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] 5-methoxy-2-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxy-5-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxy-4-(trifluormethyl)benzoat
5 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chlor-4-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-3-(trifluormethyl)benzoat
10 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-2-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-4-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlor-3-(trifluormethyl)benzoat
15 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 5-chlor-2-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-chlor-5-(trifluormethyl)benzoat
20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 2-chlor-5-(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-isopropylsulfonylbenzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-isopropylsulfonylbenzoat
25 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-cyclopentylsulfonylbenzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-pyrrolidin-1-ylsulfonylbenzoat
30 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-pyrrolidin-1-ylsulfonylbenzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3,4-bis(trifluormethyl)benzoat
[(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] 3-morpholinsulfonylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-morpholinsulfonylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(4-acetylpiperazin-1-yl)sulfonylbenzoat
 5 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(4-acetylpiperazin-1-yl)sulfonylbenzoat
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-carbamoylbenzoat;
 10 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(methylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(ethylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(dimethylcarbamoyl)benzoat;
 15 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(isopropylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(pyrrolidin-1-carbonyl)benzoat;
 20 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(piperidin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(piperazin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(morpholin-4-carbonyl)benzoat;
 25 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(4-methylpiperazin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-(4-acetylpiperazin-1-carbonyl)benzoat;
 30 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-carbamoylbenzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(methylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)-2-methyl-propyl] 3-(ethylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(dimethylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(isopropylcarbamoyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(pyrrolidin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(piperidin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(piperazin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(morpholin-4-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(4-methylpiperazin-1-carbonyl)benzoat;
 [(2R)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-(4-acetylpiperazin-1-carbonyl)benzoat
 [(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] benzoat;
 [(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-methylbenzoat;
 [(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-fluorbenzoat;
 [(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-fluorbenzoat;
 [(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 3-methoxybenzoat;
 [(2S)-3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2-methyl-propyl] 4-chlorbenzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] cyclopentancarboxylat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-2,2-dimethyl-propyl] 2-ethylbutanoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-

- 2,2-dimethyl-propyl] 4-methylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-fluorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 5 2,2-dimethyl-propyl] 4-fluorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 2-fluorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-cyanobenzoat
 10 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-cyanobenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-methoxybenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 15 2,2-dimethyl-propyl] 4-chlorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-chlorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 2-chlorbenzoat
 20 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3,5-difluorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3,4-difluorbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 25 2,2-dimethyl-propyl] 4-acetylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-acetylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-cyano-3-fluor-benzoat
 30 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-cyano-2-fluor-benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-ethoxybenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-

- 2,2-dimethyl-propyl] 2-chlor-6-fluor-benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 2-chlor-4-fluor-benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 5 2,2-dimethyl-propyl] 3-(trifluormethyl)benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 2-(trifluormethyl)benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(trifluormethyl)benzoat
 10 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-methylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-methylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 15 2,2-dimethyl-propyl] 2-methylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-sulfamoylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-sulfamoylbenzoat
 20 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-fluor-5-(trifluormethyl)benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-ethylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 25 2,2-dimethyl-propyl] 4-ethylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(methylsulfamoyl)benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-chlor-3-(trifluormethyl)benzoat
 30 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 2-chlor-5-(trifluormethyl)benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-isopropylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-

- 2,2-dimethyl-propyl] 4-isopropylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(dimethylsulfamoyl)benzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 5 2,2-dimethyl-propyl] 4-cyclopentylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-pyrrolidin-1-ylsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-pyrrolidin-1-ylsulfonylbenzoat
 10 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-morpholinsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-morpholinsulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 15 2,2-dimethyl-propyl] 4-(4-acetylpiperazin-1-yl)sulfonylbenzoat
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(4-acetylpiperazin-1-yl)sulfonylbenzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-carbamoylbenzoat;
 20 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(methylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(ethylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 25 2,2-dimethyl-propyl] 4-(dimethylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(isopropylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(pyrrolidin-1-carbonyl)benzoat;
 30 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(piperidin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(piperazin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-

- 2,2-dimethyl-propyl] 4-(morpholin-4-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 4-(4-methylpiperazin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 5 2,2-dimethyl-propyl] 4-(4-acetylpiperazin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-carbamoylbenzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(methylcarbamoyl)benzoat;
 10 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(ethylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(dimethylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 15 2,2-dimethyl-propyl] 3-(isopropylcarbamoyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(pyrrolidin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(piperidin-1-carbonyl)benzoat;
 20 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(piperazin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(morpholin-4-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 25 2,2-dimethyl-propyl] 3-(4-methylpiperazin-1-carbonyl)benzoat;
 [3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)-
 2,2-dimethyl-propyl] 3-(4-acetylpiperazin-1-carbonyl)benzoat;
 [1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-
 yl)methyl]cyclopropyl]methyl 4-methylbenzoat;
 30 [1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-
 yl)methyl]cyclopropyl]methyl 3-fluorbenzoat;
 [1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-
 yl)methyl]cyclopropyl]methyl 4-fluorbenzoat;
 [1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)methyl]cyclopropyl)methyl 3-methoxybenzoat;
 [1-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]cyclopropyl)methyl 4-chlorbenzoat;
 [3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl)methyl benzoat;
 5 [3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl)methyl 4-methylbenzoat;
 [3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl)methyl 3-fluorbenzoat;
 10 [3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl)methyl 4-fluorbenzoat;
 [3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl)methyl 3-methoxybenzoat;
 [3-[(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)methyl]oxetan-3-yl)methyl 4-chlorbenzoat;
 15 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl benzoat
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butylbenzoat;
 20 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 3-methylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-methylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-fluorbenzoat;
 25 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 3-acetylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-acetylbenzoat;
 30 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 3-methylsulfonylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-methylsulfonylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-

- yl)butyl 4-ethylsulfonylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 3-(methylsulfamoyl)benzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-(methylsulfamoyl)benzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 3-(dimethylsulfamoyl)benzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-(dimethylsulfamoyl)benzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-cyclopentylsulfonylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-pyrrolidin-1-ylsulfonylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-morpholinsulfonylbenzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-(methylcarbamoyl)benzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-(dimethylcarbamoyl)benzoat;
 4-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)butyl 4-(morpholin-4-carbonyl)benzoat;
 2-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)ethyl benzoat; eller
 et farmaceutisk acceptabelt salt, hydrat eller solvat deraf.

25

7. Forbindelse ifølge et hvilket som helst af kravene 1 til 5, hvor forbindelsen er 3-(3-ethyl-4-oxo-spiro[6,8-dihydro-5H-pyrazolo[4,3-c]azepin-7,4'-tetrahydropyran]-1-yl)propyl 4-methylbenzoat (forbindelse 101) eller et farmaceutisk acceptabelt salt, hydrat eller solvat deraf.
8. Farmaceutisk sammensætning, der omfatter en forbindelse ifølge et hvilket som helst af kravene 1-6 sammen med et farmaceutisk acceptabelt bæremedium eller hjælpestof eller en eller flere farmaceutisk acceptable bærere.
9. Anvendelse af en forbindelse ifølge et hvilket som helst af kravene 1-7 til fremstilling

af en farmaceutisk sammensætning.

5 **10.** Anvendelse af en forbindelse ifølge krav 9 i fremstillingen af en farmaceutisk sammensætning til behandling eller lindring af en sygdom, lidelse eller tilstand der er en hudsygdom eller -tilstand.

10 **11.** Anvendelse ifølge krav 10, hvor sygdommen, lidelsen eller tilstanden er proliferative og inflammatoriske hudlidelser, dermatitis, atopisk dermatitis, seboroisk dermatitis, kontaktdermatitis, psoriasis, cancer, epidermal inflammation, alopecia, alopecia areata, hudatrofi, steroid-induceret hudatrofi, hudældning, fotohudældning, acne, urticaria, pruritis eller eksem.

12. **12.** Forbindelse ifølge et hvilket som helst af kravene 1-7 til anvendelse som et lægemiddel.

15 **13.** Forbindelse til anvendelse ifølge krav 12 til anvendelse i behandling eller lindring af en sygdom, lidelse eller tilstand, der er en hudsygdom eller -tilstand.

20 **14.** Forbindelse til anvendelse ifølge krav 13 til anvendelse i behandling af proliferative og inflammatoriske hudlidelser, dermatitis, atopisk dermatitis, seboroisk dermatitis, kontaktdermatitis, psoriasis, cancer, epidermal inflammation, alopecia, alopecia areata, hudatrofi, steroid-induceret hudatrofi, hudældning, fotohudældning, acne, urticaria, pruritis og eksem.