



US 20030158086A1

(19) **United States**

(12) **Patent Application Publication**

Klett-Loch

(10) **Pub. No.: US 2003/0158086 A1**

(43) **Pub. Date: Aug. 21, 2003**

(54) **SYNTHETIC, STATISTIC THYMIC PEPTIDE COMBINATION AND ITS USE AS A PREPARATION WITH IMMUNOLOGICAL AND/OR ENDOCRINOLOGICAL EFFECT**

(76) Inventor: **Lore Maria Klett-Loch**, Mannheim (DE)

Correspondence Address:
ALSTON & BIRD LLP
BANK OF AMERICA PLAZA
101 SOUTH TRYON STREET, SUITE 4000
CHARLOTTE, NC 28280-4000 (US)

(21) Appl. No.: **10/224,283**

(22) Filed: **Aug. 20, 2002**

Related U.S. Application Data

(63) Continuation of application No. 09/766,686, filed on Jan. 22, 2001, now Pat. No. 6,458,761, which is a

continuation of application No. 09/424,625, filed on Nov. 24, 1999, now abandoned, filed as 371 of international application No. PCT/DE97/01061, filed on May 26, 1997.

Publication Classification

(51) **Int. Cl.⁷** **A61K 38/24**
(52) **U.S. Cl.** **514/2**

(57) **ABSTRACT**

Immunologically and/or endocrinologically active preparation containing as an active ingredient short-chain peptides with a weighted quantity ratio and a pattern of amino acids characteristic of thymus tissue.




FIG. 1




FIG. 2

Mitogenic Costimulation
Statistic thymus peptide combination in the test




FIG. 3

Mitogenic Costimulation

Statistic thymus peptide combination in the test




FIG. 4

**SYNTHETIC, STATISTIC THYMIC PEPTIDE
COMBINATION AND ITS USE AS A
PREPARATION WITH IMMUNOLOGICAL
AND/OR ENDOCRINOLOGICAL EFFECT**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

[0001] This application is a continuation of U.S. application Ser. No. 09/424,625, filed Nov. 24, 1999 which is a U.S. National Phase application of PCT/DE97/01061, filed May 26, 1997. The contents of the above applications are incorporated herein by reference in their entirety.

FIELD OF THE INVENTION

[0002] The invention relates to a synthetic, statistic thymic peptide combination and its use as an immunologically and endocrinologically active preparation.

BACKGROUND OF THE INVENTION

[0003] From an increasing number of scientific publications, it becomes obvious that a communication exists by means of low-molecular signaling substances between the brain, the organs of the immune system (bone marrow, thymus, lymph nodes, spleen, intestinal mucosa, pulmonary epithelium, and body skin) and the endocrine system of the hormonal control. From the thymic tissue, low-molecular proteins, so-called signaling peptides, were identified, which function both on a neuronal level as neurotransmitters and on an immunological level as differentiation factors.

[0004] It was possible to prove in animal experiments that low-molecular signaling peptides are capable of inducing a differentiation cascade in T and B lymphocytes from pluripotent parent cells, when such active peptide substances are subcutaneously supplied to the organism. Moreover, it has been found that such signaling molecules, whose composition corresponds to the proteinic elements of the thymus gland, are capable of protecting the organism against opportunistic infections, when same has previously been harmed, for example, by X-rays, cancer chemotherapy, or a latent virus infection. Such early damage to the regulatory equilibrium of the organism is accompanied by hormonal disorders, which express themselves by changed fertility, reduced mental and physical performance, sleep disorders, and loss of appetite. Moreover, humans exhibit a reduced protective function of the skin as well as the quality of hair and nail growth.

[0005] From experience, medical science has countered this syndrome for a long time in that preparations of organic extracts are used as medicines or food supplements. Such preparations of the thymic tissue from animals, in particular thymus of the calf, have found wide use, even when a molecularly defined principle of action of such preparations has not been described until now. Positive data from a series of biological studies in vivo with such thymus preparations permit concluding that substances contained in the thymus tissue exert synergistic effects both on the neuronal and on the immune system. However, in the last five years the risk has come to the fore that such preparations of organic extracts from animals are contaminated with prion proteins. These prion proteins are associated with transmitting the symptoms of BSE (bovine spongiform encephalopathy, "bovine madness").

[0006] It is therefore the object of the present invention to make available an immunologically and/or endocrinologically active preparation, which permits on the one hand extrapolating the positive experience with thymic peptide combinations, but totally excludes on the other hand the risk of BSE. Furthermore, it is the object to specify a synthetic, statistic thymic peptide combination and its use as an immunologically and/or endocrinologically active preparation with corresponding advantages.

SUMMARY OF THE INVENTION

[0007] The present invention accomplishes the foregoing object by the characteristic features of claim 1. Accordingly, an immunologically and/or endocrinologically active preparation contains as an active ingredient short-chain peptides with a weighted quantity ratio and pattern of amino acids characteristic of thymic tissue.

[0008] Claim 2 relates to corresponding method, namely the stepwise lineup of individual, chemically suitable derivatized amino acids to short-chain peptides, and release of these synthetic peptides from their N-terminal and form derivatized on side functions, while maintaining the weighted quantity ratio and pattern of amino acids characteristic of the thymus tissue.

[0009] Claim 3 relates to the use of corresponding preparations, i.e., synthesized, short-chain peptides with a weighted quantity ratio and pattern of amino acids characteristic of thymus tissue with immunologically and/or endocrinologically active properties, namely preparations containing synthetic, statistic thymic peptide combinations. The further dependent claims 4 to 8 relate to quite particular uses, namely in the case of weak immunity and immunodeficiencies, hormonal disorders, as a nutrient for skin, hair, and nails, in the cosmetic field, and finally in the field of food supplements.

[0010] Proceeding from partial hydrolysates of thymic protein, protein mixtures from short-chain thymic peptides and amino acids, which result from an enzymatic and/or chemical partial hydrolysis of thymus tissue, such combinations were clarified by way of chromatography and amino-acid analysis to that extent that the such partial hydrolysates consist of short-chain peptides with a quantity ratio and pattern of amino acid characteristic of thymus tissue. When such amino acids are reacted with one another in chemically suitably derivatized form, with the quantity ratio of the amino acids characteristic of the thymus tissue being maintained, when such synthetic combinations are released from their N-terminal protective groups and coupled again and repeatedly with such weighted combinations of amino acids, which are characteristic of thymus tissues, one will obtain synthetic, statistic thymic peptide combinations to the total exclusion of the prion protein risk.

[0011] Thus, the invention discloses relatively simple ways of purposefully producing immunologically and endocrinologically active preparations with an effect of the preparation that is standardizable for the first time.

[0012] It is possible to produce such thymic peptide combinations in a pharmaceutically reproducible sterile manner, so that a preparation results in this manner, which depending on the number of repeated statistic synthesis operations consists of a family of signaling thymic peptides character-

istic of neurotransmitters and immunologically active signaling molecules. Based on analytical tests in comparison with partial hydrolysis mixtures from natural thymus tissue, it was possible to demonstrate that, within the margin of error of applied analytical methods, synthetic, statistic thymic peptide combinations are comparable in their composition with the natural preparation. From this follows that it was possible to accomplish the object of the present invention to eliminate the BSE risk by synthesizing a thymic peptide combination.

[0013] Surprisingly, it has shown that the synthetic preparation of the present invention exhibits in the biological test on human lymphocytes a superior activity with respect to mitogenic co-stimulation (induction of cell proliferation) in comparison with a commercial thymic preparation produced by partial hydrolysis.

[0014] Furthermore, it comes as a surprise that the synthetic preparation is capable of binding constituents of the extracellular matrix, a property that had previously to be ascribed only to scleroproteins.

BRIEF DESCRIPTION OF THE DRAWINGS

[0015] Having thus described the invention in general terms, reference will now be made to the accompanying drawings, which are not necessarily drawn to scale, and wherein:

[0016] **FIG. 1** is a graphic representation of data from the mitogenic co-stimulation of human peripheral blood lymphocytes (PBL's) obtained from a first donor with phytohemagglutinin (PHA) in the presence of a positive control (REF01), a commercial thymic peptide combination (GKL01) and the statistic thymic peptide combination (SP/01/21 1096).

[0017] **FIG. 2** is a graphic representation of data from the mitogenic co-stimulation of human PBLs obtained from a second donor with PHA in the presence of a positive control (REF01), a commercial thymic peptide combination (GKL01) and the statistic thymic peptide combination (SP/01/211096).

[0018] **FIG. 3** is a graphic representation of data from the mitogenic co-stimulation of human PBLs obtained from a third donor with PHA in the presence of a positive control thymosin fraction 5(TF5), a commercial thymic peptide combination (GKL01) and the statistic thymic peptide combination (SP/01/211096).

[0019] **FIG. 4** is a graphic representation of data from a second mitogenic co-stimulation of human PBLs obtained from the third donor with PHA in the presence of a positive control thymosin fraction 5(TF5), a commercial thymic peptide combination (GKL01) and the statistic thymic peptide combination (SP/01/211096).

DETAILED DESCRIPTION OF THE INVENTION

[0020] The present invention now will be described more fully hereinafter with reference to the accompanying drawings, in which preferred embodiments of the invention are shown. This invention may, however, be embodied in many different forms and should not be construed as limited to the embodiments set forth herein; rather, these embodiments are

provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art. Like numbers refer to like elements throughout.

[0021] Many modifications and other embodiments of the invention will come to mind to one skilled in the art to which this invention pertains having the benefit of the teachings presented in the foregoing descriptions and the associated drawings. Therefore, it is to be understood that the invention is not to be limited to the specific embodiments disclosed and that modifications and other embodiments are intended to be included within the scope of the appended claims. Although specific terms are employed herein, they are used in a generic and descriptive sense only and not for purposes of limitation.

[0022] The invention is described in greater detail with reference to the following examples.

EXAMPLE 1

[0023] The statistic thymic peptide combinations of the present invention were examined for bioactivity in the test system of the mitogenic co-stimulation in comparison with a commercial thymic preparation from partial hydrolysis. Tested were end concentrations of 100, 50, and 25 $\mu\text{g/ml}$.

[0024] In the case of two donors (KFG and WH) the statistic thymic peptide combination shows a statically significantly higher bioactivity than a commercial thymic preparation from partial hydrolysis. While at the high concentration of 100 $\mu\text{g/ml}$, an inhibition of cell proliferation was observed for a commercial thymic preparation from partial hydrolysis, the statistic thymic peptide combination led to a dose-dependent increase in the proliferation of human lymphocytes.

[0025] From two blood donations, human peripheral blood lymphocytes were isolated, and the proliferation rate of these cells was measured, after a suboptimal prestimulation (0.5 $\mu\text{g/ml}$ phytohemagglutinin (PHA)) under the action of different concentrations of a commercial thymic preparation from partial hydrolysis, and the statistic thymic peptide combination in comparison with the positive control REF 01 (thymosin fraction 5).

[0026] In terms of the measuring technique, the proliferation rate was determined by the following method: During the active-ingredient dependent induction of the T-cell proliferation, the DNA synthesis is stimulated for purposes of cell division. Therefore, it is possible to measure via incorporation of a radioactive DNA component (3H-thymidine) the T-cell growth with reference to incorporated radioactivity.

[0027] **FIGS. 1 and 2** show results from the stimulation of the blood lymphocytes of donor KFG and donor WH. The test data are to be related to the result of the control formulation after adding phytohemagglutinin (Kontr.+PHA). In the test, the reference thymus preparation was used as positive control in a concentration of 100 $\mu\text{g/ml}$; a commercial thymic preparation from partial hydrolysis and the statistic thymic peptide combination were tested in the three concentrations of 100, 50, and 25 $\mu\text{g/ml}$.

[0028] **FIG. 1** shows the results for donor KFG. The reference thymus preparation used a positive control led to

a very obvious increase in the proliferation rate. For a commercial thymus preparation produced by partial hydrolysis, however, it was possible to observe a dose-dependent inhibition. Only in the case of the lowest concentration of 25 $\mu\text{g/ml}$ was the value of the control formulation reached. By the action of the statistic thymic peptide preparation, the cell proliferation rate was statistically significantly higher in the case of the three tested concentrations than in the control formulation.

[0029] According to FIG. 2, the lymphocytes of donor WH showed comparable results. Again, the reference thymus preparation used as positive control resulted in a considerable increase in the proliferation of human lymphocytes. A commercial thymus preparation produced by partial hydrolysis led in the concentrations of 50 and 25 $\mu\text{g/ml}$ to a slight increase in the cell proliferation rate, but not in the high concentration of 100 $\mu\text{g/ml}$. However, for the statistic thymic peptide combination SP/01/211096, it was possible to observe a clear and dose-dependent increase in the DNA synthesis rate.

[0030] The object of these tests according to Example 1 was the comparative testing of the synthetic, statistic thymic peptide preparation with a commercial thymus preparation produced by partial hydrolysis in the test system of the mitogenic co-stimulation. The intent was to test the capability of the test substance of increasing the cell proliferation rate of human blood lymphocytes.

[0031] For the statistic thymic peptide combination, it was possible to find in the case of two donors a clearly higher bioactivity in comparison with the commercial thymus preparation from partial hydrolysis. In addition, donor WH showed a dose-dependent increase in the DNA synthesis rate. Surprisingly, the addition of a commercial thymus preparation from partial hydrolysis to the cell culture formulations led rather to an inhibition of the cell proliferation. Only in the case of low concentrations of 50 or 25 $\mu\text{g/ml}$ was the cell proliferation rate slightly above the value of the control formulation in the case of donor WH.

EXAMPLE 2

[0032] The following example relates to the stimulation of the blood lymphocytes of donor DB. The test data are to be related with reference to FIGS. 3 and 4 to the result of the control formulation after adding phytohemagglutinin at PHA. The thymosin fraction No. 5 (TF5) was used as positive control in two concentrations (third and fourth columns from the left). For a comparison, a commercial partial thymic hydrolysate (fifth and sixth columns) was also used in the same concentration. As a control PHA was used alone.

[0033] In the mitogenic co-stimulation with suboptimal stimulation of the blood cells with PHA of 0.05 $\mu\text{g/ml}$, the synthetic-statistic thymic peptide combination shows a surprisingly distinct activity both in the case of a dose of the claimed synthesized product of 100 $\mu\text{g/ml/cell}$ culture formulation and in the case of 0.4 $\mu\text{g/ml/cell}$ culture formulation.

[0034] Likewise, in the case of a slight prestimulation of the blood cells with PHA in an amount of 0.025 $\mu\text{g/ml}$ /formulation, one can clearly notice the effects of the synthetic-statistic thymic peptide combination according to the invention, even though less distinct than in the case of a prestimulation of the blood cells with 0.05 $\mu\text{g/ml}$ PHA/formulation.

[0035] Lastly, one has in this instance a typical, in vitro, two-phase activity, which shows a maximum respectively in the range of 100 $\mu\text{g/ml}$ and 0.4 $\mu\text{g/ml}$, and is in each case better than the commercial partial thymic hydrolysate that is used as comparison preparation. In any event, this two-phase activity is a proof for the wide scope of the immunological, in vitro activity, wherein the great extent of activity in the case of the extremely low concentration of 0.4 $\mu\text{g/ml}$ comes no doubt as a surprise and is of great economic interest due to its really very low concentration.

[0036] The analytical data of the synthetic-statistic thymic peptide combination are as follows:

Certificate of Analysis Data Sheet Synthetic-Statistic Thymic Peptides		
	Specifications Specified	#SP/01/211096 Found
Description:	yellowish powder	yellowish powder
Residual Solvents (GC):	<10%	
Water	<0.1%	9% \pm 0.5%
DMF	<0.1%	n.d.* < 200 ppm**
Methanol	<0.1%	130 ppm < 130 ppm
Toluol	<0.1%	n.d. < 40 ppm
Ethylisopropylamine	<0.1%	n.d. < 200 ppm
Acetic ester	<0.1%	n.d. < 20 ppm
Acetates:	<3%	1.6% \pm 0.2%
Inorganic salts:	<0.5%	<0.2%
Specific rotation: (20° C. c = 0.2; H ₂ O)	-30.0° \pm 5%	-30.2° \pm 0.2°
Free amino acid content: (%, w/w)	GKL01 \pm 5%	complies
Total amino acid composition: (%, w/w)	GKL01 \pm 5%	complies
Peptide content:	50% \pm 5%	complies

That which is claimed:

1. Preparation containing synthetic thymic peptide combinations without the risk of BSE, available by lining up in steps individual, chemically suitably derivatized amino acids to short-chain peptides, and releasing these synthetic peptides from their N-terminal group and form derivatized on side functions, while maintaining the weighted quantity ratio and pattern of amino acids characteristic of thymus tissue.
2. Method of producing an immunologically and/or endocrinologically active preparation containing as an active ingredient short-chain peptides with a weighted quantity ratio and pattern of amino acids characteristic of thymus tissue, characterized by lining up in steps individual, chemically suitably derivatized amino acids to short-chain peptides, and releasing these synthetic peptides from their N-terminal group and form derivatized on side functions, while maintaining the weighted quantity ratio and pattern of amino acids characteristic of thymus tissue.
3. Use of a preparation of claim 1 for producing a substance with immunologically and/or endocrinologically active properties.
4. Use of claim 3 in the case of weak immunity and immunodeficiencies.
5. Use of claim 3 in the case of hormonal disorders.
6. Use of claim 3 as a nutrient for skin, hair, nails.
7. Use of claim 3 in the field of cosmetics.
8. Use of claim 3 in the field of food supplements.

* * * * *