Office de la Propriété Intellectuelle du Canada
Un organisme d'Industrie Canada

Canadian Intellectual Property Office
An agency of Industry Canada

CA 2683887 A1 2008/11/06

(21) 2 683 887

(12) DEMANDE DE BREVET CANADIEN
CANADIAN PATENT APPLICATION

(13) A1


(86) Date de dépôt PCT/PCT Filing Date: 2008/04/23
(87) Date publication PCT/PCT Publication Date: 2008/11/06
(85) Entrée phase nationale/National Entry: 2009/10/14
(86) N° demande PCT/PCT Application No.: JP 2008/057847
(87) N° publication PCT/PCT Publication No.: 2008/133274
(30) Priorités/Priorities: 2007/04/24 (JP2007-114288); 2007/11/08 (JP2007-290589)

(51) Cl.Int./Int.Cl. *C07D 279/06* (2006.01), *A61K 31/497* (2006.01), *A61K 31/54* (2006.01), *A61K 31/541* (2006.01), *A61P 25/28* (2006.01), *A61P 43/00* (2006.01), *C07D 417/12* (2006.01), *C07D 417/14* (2006.01), *C07D 471/04* (2006.01), *C07D 487/04* (2006.01), *C07D 495/04* (2006.01), *C07D 513/04* (2006.01)

(71) Demandeur/Applicant:
SHIONOGI & CO., LTD., JP

(72) Inventeurs/Inventors:
TAMURA, YUUSUKE, JP;
SUZUKI, SHINJI, JP;
TADA, YUKIO, JP; ...

(54) Titre : DERIVES D'AMINODIHYDROTHIAZINE SUBSTITUES PAR DES GROUPES CYCLIQUES
(54) Title: AMINODIHYDROTHIAZINE DERIVATIVES SUBSTITUTED WITH A CYCLIC GROUP



(I)

(57) Abrégé/Abstract:
The invention provides compounds represented by the general formula (I) which are useful as remedies for diseases induced by the production, secretion, or deposition of amyloid β protein. Compounds represented by the general formula (I); pharmaceutically acceptable salts thereof; or solvates of both: (I) wherein A is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group; R^1 is optionally substituted lower alkyl, optionally substituted lower alkenyl, or optionally substituted lower alkynyl; R^{2a} and R^{2b} are each independently hydrogen, optionally substituted lower alkyl, or optionally substituted acyl; and R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogeno, hydroxy, optionally substituted lower alkyl, or the like.

(21) 2 683 887


(13) A1

(72) Inventeurs(suite)/Inventors(continued): YONEZAWA, SHUJI, JP; FUJIKOSHI, CHIAKI, JP; MATSUMOTO, SAE, JP; KOORIYAMA, YUUJI, JP; UENO, TATSUHIKO, JP

(74) Agent: KIRBY EADES GALE BAKER

ABSTRACT

This invention provides a compound of the formula (I):



(I)

wherein the ring A is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group; R^1 is optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl; R^{2a} and R^{2b} are each independently hydrogen, optionally substituted lower alkyl or optionally substituted acyl; R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogen, hydroxy or optionally substituted lower alkyl etc.; a pharmaceutically acceptable salt or solvate thereof,

which is useful for treating diseases induced by production, secretion and/or deposition of amyloid β protein.

DESCRIPTION

AMINODIHYDROTHIAZINE DERIVATIVES SUBSTITUTED WITH A CYCLIC GROUP

TECHNICAL FIELD

This invention relates to a compound having an inhibitory activity against production of amyloid β protein and useful for treating diseases induced by production, secretion and/or deposition of amyloid β protein.

BACKGROUND ART

In the brain of Alzheimer's patient many insoluble spots (senile plaque) are found, which is formed by extraneuronal accumulation of a peptide called amyloid β protein comprised of about 40 amino acids. Neuronal death caused by the senile plaque is considered to develop Alzheimer's Disease and an enhancer of amyloid β protein decomposition or vaccine of amyloid β protein etc. are extensively studied as a remedy for Alzheimer's Disease.

Secretase is an enzyme producing amyloid β protein by intracellular cleavage of a protein called amyloid β protein precursor(APP). An enzyme playing a role for producing an N-terminal of the amyloid β protein is called BACE1 (beta-site APP-cleaving enzyme) and an inhibitor of the BACE1, which will reduce production of amyloid β protein, could be a remedy for treating Alzheimer's disease.

Patent literature 1 discloses a compound with a chemical structure similar to that of the compound of the present invention, having an inhibitory activity of NO synthetase and effective for treating dementia.

Patent literature 2-5 and non-patent literature 1-2 disclose compounds with a chemical structure similar to those of the compound of the present invention, and describe that each compound is useful as an anti-hypotensive agent, morphine-like analgesic or

tranquilizer, intermediate of a therapeutic agent, NPYY5 antagonist, analgesic and the like.

Patent literatures 6-14 disclose BACE-1inhibitors having a chemical structure different from that of the compound of the present invention. Also, patent literature 15 discloses a BACE-1 inhibitor.

Patent literature 1: WO 96/014842 Pamphlet

Patent literature 2: US 3,235,551

Patent literature 3: US 3,227,713

Patent literature 4: JP H09-067355

Patent literature 5: WO 2005/111031 Pamphlet

Patent literature 6: WO 02/96897 Pamphlet

Patent literature 7: WO 04/043916 Pamphlet

Patent literature 8: WO 2005/058311 Pamphlet

Patent literature 9: WO 2005/097767 Pamphlet

Patent literature 10: WO 2006/041404 Pamphlet

Patent literature 11: WO 2006/041405 Pamphlet

Patent literature 12: US 2007/0004786A

Patent literature 13: US 2007/0004730A

Patent literature 14: US 2007/27199A

Patent literature 15: WO 2007/049532 Pamphlet

Non-patent literature 1: Journal of Heterocyclic Chemistry, 14, 717-723(1977)

Non-patent literature 2: Journal of Organic Chemistry, 33(8), 3126-3132(1968).

DISCLOSURE OF INVENTION


PROBLEM TO BE SOLVED

This invention provides with a compound having an inhibitory activity against BACE-1 and useful for treating diseases induced by production, secretion and/or deposition of amyloid β protein.

MEANS TO SOLVE THE PROBLEM

The present invention provides with

1) a compound of the formula (I):



(I)

wherein the ring A is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group,

R^1 is optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group,

R^{2a} and R^{2b} are each independently hydrogen, optionally substituted lower alkyl or optionally substituted acyl,

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted acyl, optionally substituted lower alkoxy, optionally substituted carbocyclyl lower alkyl, optionally substituted heterocyclyl lower alkyl, optionally substituted carbocyclyl lower alkoxy, optionally substituted heterocyclyl lower alkoxy, optionally substituted aralkyl, optionally substituted heteroaralkyl, optionally substituted aralkyloxy, optionally substituted heteroaralkyloxy, optionally substituted lower alkylthio, carboxy, optionally substituted lower alkoxycarbonyl, optionally substituted amino, optionally substituted carbamoyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group, or R^{3a} and R^{3b} or R^{3c} and R^{3d} may form a carbocyclic ring together with a linked carbon atom or may form oxo,

provided the following compounds i) and ii) are excluded;


i) a compound in which R^{2a} is hydrogen, R^{2b} is hydrogen, acetyl or phenyl, R^1 is methyl, and the ring A is phenyl or 4-methoxyphenyl;

ii) a compound in which R^{2a} is hydrogen, R^{2b} is hydrogen, acetyl or phenyl, R^1 is

ethyl and the ring A is 3,4-dimethoxyphenyl,

a pharmaceutically acceptable salt or solvate thereof;

1') a compound of the formula (I):



(I)

wherein the ring A is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group,

R^1 is optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group,

R^{2a} and R^{2b} are each independently hydrogen, optionally substituted lower alkyl or optionally substituted acyl,

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted acyl, optionally substituted lower alkoxy, optionally substituted lower alkylthio, carboxy, optionally substituted lower alkoxycarbonyl, optionally substituted amino, optionally substituted carbamoyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group, or

R^{3a} and R^{3b} or R^{3c} and R^{3d} may form a carbocyclic ring together with a linked carbon atom,

provided the following compounds i) and ii) are excluded:


i) a compound in which R^{2a} is hydrogen, R^{2b} is hydrogen, acetyl or phenyl, R^1 is methyl, and the ring A is phenyl or 4-methoxyphenyl;

ii) a compound in which R^{2a} is hydrogen, R^{2b} is hydrogen, acetyl or phenyl, R^1 is ethyl and the ring A is 3,4-dimethoxyphenyl,

a pharmaceutically acceptable salt or solvate thereof;


2) the compound of 1) or 1') described above,

wherein the ring A is




wherein the ring A' is a carbocyclic group or a heterocyclic group,


G is




(ii')




(ii")




(iv')




(v')



(x')




(xiii')



(xiv')

or



(xixi')

wherein R^5 is hydrogen, lower alkyl or acyl,

R^6 is optionally substituted lower alkyl, optionally substituted lower alkenyl or optionally substituted lower alkynyl,

W^1 is O or S,

W^2 is O, S or NR^5,

Ak is optionally substituted lower alkylene, optionally substituted lower alkenylene or optionally substituted lower alkynylene,


the ring B is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group and each R^5 may be independent:

R^4 is halogen, hydroxyl, mercapto, halogeno lower alkyl, lower alkoxy, amino, lower alkylamino, acylamino or lower alkylthio and each R^4 may be independent;

A pharmaceutically acceptable salt or solvate thereof;


2') the compound of 1) or 1') described above

wherein the ring A is



wherein the ring A' is a carbocyclic group or a heterocyclic group,

G is



wherein R^5 is hydrogen, lower alkyl or acyl,

R^6 is optionally substituted lower alkyl, optionally substituted lower alkenyl or optionally substituted lower alkynyl,

W^1 is O or S,

W^2 is O, S or NR^5,

Ak is optionally substituted lower alkylene, optionally substituted lower alkenylene or optionally substituted lower alkynylene,

the ring B is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group and each R^5 may be independent:

R^4 is halogen, hydroxyl, mercapto, halogeno lower alkyl, lower alkyl, lower alkoxy, amino, lower alkylamino, acylamino or lower alkylthio and each R^4 may be independent;

A pharmaceutically acceptable salt or solvate thereof;

3) the compound of 2) or 2') described above wherein the ring A' is phenyl or a nitrogen-containing aromatic heterocyclic group, a pharmaceutically acceptable salt or solvate thereof;

3') the compound of 2) or 2') described above wherein the ring A' is phenyl, a pharmaceutically acceptable salt or solvate thereof;

3'') the compound of 2) or 2') described above wherein the ring A' is a nitrogen-containing aromatic heteromonocyclic group, a pharmaceutically acceptable salt or solvate thereof;

3''') the compound of 2) or 2') described above wherein the ring A' is pyridyl, a pharmaceutically acceptable salt or solvate thereof;

4) the compound of 1) – 3), 1'), 2'), 3'), 3'') or 3''') described above wherein R^1 is C1-C3 alkyl, a pharmaceutically acceptable salt or solvate thereof;

4') the compound of 1) – 3), 1'), 2'), 3'), 3'') or 3''') described above wherein R^1 is optionally substituted lower alkynyl, a pharmaceutically acceptable salt or solvate thereof;

5) the compound of 1) – 4), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein R^{2a} and R^{2b} are both hydrogen, a pharmaceutically acceptable salt or solvate thereof;

6) the compound of 1) – 5), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein all of R^{3a}, R^{3b}, R^{3c} and R^{3d} are hydrogen, a pharmaceutically acceptable salt or solvate thereof;

6') the compound of 1) – 5), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein R^{3a} and R^{3b} are the same substituent selected from halogen and optionally substituted lower alkyl, a pharmaceutically acceptable salt or solvate thereof;

6'') the compound of 1) – 5), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein R^{3c} and R^{3d} are the same substituent selected from halogen and optionally substituted lower alkyl, a pharmaceutically acceptable salt or solvate thereof;

6''') the compound of 1) – 5), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein R^{3a} and R^{3b} or R^{3c} and R^{3d} form a carbocyclic ring together with a linked carbon atom a pharmaceutically acceptable salt or solvate thereof;

7) the compound of 1) – 5), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein R^{3c} or R^{3d} is optionally substituted carbocyclic ring lower alkoxy or optionally substituted heterocyclyl lower alkoxy, a pharmaceutically acceptable salt or solvate thereof;

7') the compound of 1) – 5), 1'), 2'), 3'), 3''), 3''') or 4') described above wherein R^{3c} and R^{3d} form oxo together with a linked carbon atom, a pharmaceutically acceptable salt or solvate thereof;

8) a pharmaceutical composition comprising a compound of 1) – 7), 1'), 2'), 3'), 3''), 3'''), 4') 6'), 6''), 6''') or 7') described above, a pharmaceutically acceptable salt or solvate thereof as an active ingredient; and

9) a pharmaceutical composition having a BACE 1 inhibitory activity comprising a compound of 1) – 7), 1'), 2'), 3'), 3''), 3'''), 4') 6'), 6''), 6''') or 7') described above, a pharmaceutically acceptable salt or solvate thereof as an active ingredient;

The present invention also provides with

10) the pharmaceutical composition having a BACE 1 inhibitory activity of 9) described above, which is a composition having inhibitory activity of amyloid β protein production;

11) the pharmaceutical composition having a BACE 1 inhibitory activity of 9) described above, which is a medicine for treating diseases induced by production, secretion and/or deposition of amyloid β protein;

12) the pharmaceutical composition having a BACE 1 inhibitory activity of 9) described above, which is a medicine for treating Alzheimer's disease;

13) a method for treating diseases induced by production, secretion and/or deposition of amyloid β protein, characterized in administering a compound of the formula (I) described in 1) above, a pharmaceutically acceptable salt or solvate thereof;

14) use of a compound of the formula (I) described in 1) above, a pharmaceutically acceptable salt or solvate thereof described in 1) above, in manufacturing a medicine for treating diseases induced by production, secretion and/or deposition of amyloid β protein;

15) a method for treating diseases induced by BACE 1 characterized in administering a compound of the formula (I) described in 1) above, a pharmaceutically acceptable salt or solvate thereof;

16) use of a compound of the formula (I) described in 1) above, a pharmaceutically acceptable salt or solvate thereof, in manufacturing a medicine for treating diseases induced by BACE 1;

17) a method for treating Alzheimer's disease characterized in administering a

compound of the formula (I) described in 1) above, a pharmaceutically acceptable salt or solvate thereof; and

18) use of a compound of the formula (I) described in 1) above, a pharmaceutically acceptable salt or solvate thereof in manufacturing a medicine for treating Alzheimer's disease.

EFFECT OF INVENTION

A compound of the present invention is useful for treating diseases induced by production, secretion and/or deposition of amyloid β protein (Alzheimer's disease etc.).

BEST MODE FOR CARRYING OUT THE INVENTION

In this description, "halogen" includes fluorine, chlorine, bromine and iodine.

A moiety of halogen in "halogeno lower alkyl" and "halogeno lower alkoxycarbonyl" is the same as "halogen" above.

"Lower alkyl" includes C1-C15, preferably C1-C10, more preferably C1-C6 and further more preferably C1-C3 straight or branched alkyl, and for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, n-pentyl, isopentyl, neopentyl, hexyl, isohexyl, n-heptyl, isoheptyl, n-octyl, isooctyl, n-nonyl and n-decyl are exemplified.

A moiety of alkyl in "lower alkoxy", "halogeno lower alkyl", "hydroxyl lower alkyl", "hydroxyl lower alkoxy", "lower alkoxycarbonyl", "halogeno lower alkoxycarbonyl", "lower alkoxycarbonyl lower alkyl", "lower alkylamino", "lower alkoxy lower alkyl", "hydroxyimino lower alkyl", "lower alkoxyimino lower alkyl", "amino lower alkyl" "lower alkoxy lower alkoxy", "lower alkoxy lower alkenyl", "lower alkoxycarbonyl lower alkenyl", "lower alkoxy lower alkynyl", "lower alkoxycarbonyl lower alkynyl", "lower alkyl carbamoyl", "hydroxyl lower alkyl carbamoyl", "lower alkoxyimino", "lower alkylthio", "lower alkylsulfonyl", "lower alkylsulfonyloxy", "lower alkyl sulfamoyl", "lower alkyl sulfinyl", "carbocyclyl lower alkyl", "carbocyclyl lower alkyl", "carbocyclyl lower alkoxy", "carbocyclyl lower alkoxycarbonyl", "carbocyclyl lower alkylamino", "carbocyclyl lower alkyl carbamoyl", "cycloalkyl lower alkyl", "cycloalkyl lower alkoxy", "cycloalkyl

lower alkylamino", "cycloalkyl lower alkoxycarbonyl", "cycloalkyl lower alkylcarbamoyl", "aryl lower alkyl", "aryl lower alkoxy", "aryl lower alkylamino", "aryl lower alkoxycarbonyl", "aryl lower alkoxycarbamoyl", "heterocyclyl lower alkyl", "heterocyclyl lower alkoxy", "heterocyclyl lower alkylamino", "heterocyclyl lower alkoxycarbonyl" and "heterocyclyl lower alkylcarbamoyl" is the same as "alkyl" above.

"Optionally substituted lower alkyl" may be substituted with one or more of substituent(s) selected from a substituent group α.

Group α is a group consisting of halogen, hydroxy, lower alkoxy, hydroxy lower alkoxy, lower alkoxy lower alkoxy, acyl, acyloxy, carboxy, lower alkoxycarbonyl, amino, acylamino, lower alkylamino, imino, hydroxyimino, lower alkoxyimino, lower alkylthio, carbamoyl, lower alkylcarbamoyl, hydroxy lower alkylcarbamoyl, sulfamoyl, lower alkylsulfamoyl, lower alkylsulfonyl, cyano, nitro, a carbocyclic group and a heterocyclic group.

One or more of substituent(s) selected from the substituent group α is exemplified as a substituent of "optionally substituted lower alkoxy", "optionally substituted lower alkoxycarbonyl" and "optionally substituted lower alkylthio"

"Lower alkylidene" includes a divalent group derived from the "lower alkyl" above, and methylidene, ethylidene, propylidene, isopropylidene, butylidene, pentylidene and hexylidene etc. are exemplified.

"Lower alkenyl" includes C2-C15, preferably C2-C10, more preferably C2-C6 and further more preferably C2-C4 straight or branched alkenyl having one or more double bond(s) at any position thereof. Examples of lower alkenyl include vinyl, allyl, propenyl, isopropenyl, butenyl, isobutenyl, prenyl, butadienyl, pentenyl, isopentenyl, pentadienyl, hexenyl, isohexenyl, hexadienyl, heptenyl, octenyl, nonenyl, decenyl, undecenyl, dodecenyl, tridecenyl, tetradecenyl, pentadecenyl and the like.

"Lower alkynyl" includes C2-C10, preferably C2-C8, more preferably C3-C6 straight or branched alkynyl having one or more triple bond(s) at any position thereof. Examples of lower alkynyl include ethynyl, propynyl, butynyl, pentynyl, hexynyl, heptynyl,

octynyl, nonynyl, decynyl and the like. Lower alkynyl may additionally have a double bond at any position thereof.

One or more of substituent(s) selected from the substituent group α is exemplified as a substituent of "optionally substituted lower alkenyl" and "optionally substituted lower alkynyl".

A moiety of lower alkenyl in "hydroxyl lower alkenyl", "lower alkoxy lower alkenyl", "lower alkoxycarbonyl lower alkenyl", "carbocyclyl lower alkenyl", "lower alkenyloxy", "lower alkenylthio" and "lower alkenylamino" is the same as that of "lower alkenyl".

A moiety of lower alkynyl in "hydroxyl lower alkynyl", "lower alkoxy lower alkynyl", "lower alkoxycarbonyl lower alkynyl", "carbocyclyl lower alkynyl", "lower alkynyloxy", "lower alkenylamino" and "lower alkynylamino" is the same as that of "lower alkynyl" above.

One or more substituents selected from lower alkyl, acyl, hydroxyl, lower alkoxy, lower alkoxycarbonyl, a carbocyclic group and a heterocyclic group etc. is exemplified as a substituent of "optionally substituted amino" and "optionally substituted carbamoyl".

"Acyl" includes C1-C10 aliphatic acyl, carbocyclyl carbonyl and heterocyclic carbonyl, and examples of acyl include formyl, acetyl, propionyl, butyryl, isobutyryl, valeryl, pivaloyl, hexanoyl, acryloyl, propioloyl, methacryloyl, crotonoyl, benzoyl, cyclohexanecarbonyl, pyridinecarbonyl, furancarbonyl, thiophenecarbonyl, benzothiazolecarbonyl, pyrazinecarbonyl, piperidinecarbonyl, thiomorpholino and the like.

A moiety of acyl in "acylamino" and "acyloxy" is the same as described above.

One or more substituents selected from the substituent group α is exemplified as a substituent in "optionally substituted acyl" and a moiety of the ring in carbocyclyl carbonyl and heterocyclylcarbonyl is optionally substituted with one or more substituent(s) selected from lower alkyl, the substituent group α and lower alkyl substituted with one or more substituent(s) selected from the substituent group α.

"A carbocyclic group" includes cycloalkyl, cycloalkenyl, aryl, and non-aromatic

fused carbocyclic group etc.

"Cycloalkyl" includes C3-C10, preferably C3-C8 and more preferably C4-C8 carbocyclic group and examples of cycloalkyl include cyclopropyl, cyclobutyl, cyclopentyl cyclohexyl, cycloheptyl, cyclooctyl, cyclononyl, cyclodecyl and the like.

A moiety of cycloalkyl in "cycloalkyl lower alkyl", "cycloalkyloxy", "cycloalkyl lower alkoxy", "cycloalkylthio", "cycloalkylamino", "cycloalkyl lower alkylamino", "cycloalkylsulfamoyl", "cycloalkylsulfonyl", "cycloalkylcarbamoyl", "cycloalkyl lower alkylcarbamoyl", "cycloalkyl lower alkoxycarbonyl" and "cycloalkylcarbonyl" is the same as "cycloalkyl" described above

"Cycloalkenyl" includes the above cycloalkyl having one or more double bond(s) at any position on the ring, and examples of the cycloalkenyl include cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptynyl, cyclooctynyl, and cyclohexadienyl etc.

Examples of "aryl" include phenyl, naphthyl, anthryl and phenanthryl etc. and especially phenyl is preferable.

"Non-aromatic fused carbocyclic group" includes a group in which two or more cyclic groups selected from "cycloalkyl", "cycloalkenyl" and "aryl" described above fused, and examples of "Non-aromatic fused carbocyclic group" include indanyl, indenyl, tetrahydronaphthyl and fluorenyl etc.

"Forming a carbocyclic ring together with a linked carbon atom" means that two substituents jointly form "cycloalkyl" above.

A moiety of the carbocyclic ring in "carbocyclyloxy", "carbocyclyl lower alkyl", "carbocyclyl lower alkenyl", "carbocyclyl lower alkynyl", "carbocyclyl lower alkoxy", "carbocyclyl lower alkoxycarbonyl", "carbocyclylthio", "carbocyclyl amino", "carbocyclyl lower alkylamino", "carbocyclylcarbonyl", "carbocyclylsulfamoyl", "carbocyclylsulfonyl", "carbocyclylcarbamoyl", "carbocyclyl lower alkyl carbamoyl", "carbocyclyloxycarbonyl" is the same as the "carbocyclic group".

A moiety of aryl in "aryl lower alkyl", "aryloxy", "aryloxycarbonyl",

"aryloxycarbonyloxy", "aryl lower alkoxycarbonyl", "arylthio", "arylamino", "aryl lower alkoxy", "aryl lower alkylamino", "arylsulfonyl", "arylsulfonyloxy", "arylsulfinyl", "arylsulfamoyl", "arylcarbamoyl" and "aryl lower alkylcarbamoyl" is the same as the "aryl" above.

"Heterocyclic group" includes a heterocyclic group containing one or more heteroatom(s) each independently selected from O, S and N, and examples of "heterocyclic group" include 5- or 6-membered heteroaryl such as pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazolyl, triazinyl, tetrazolyl, furyl, thienyl, isoxazolyl, oxazolyl, oxadiazolyl, isothiazolyl, thiazolyl and thiadiazolyl etc.; a non-aromatic heterocyclic group such as dioxanyl, thiiranyl, oxiranyl, oxetanyl, oxathiolanyl, azetidinyl, thianyl, thiazolidinyl, pyrrolidinyl, pyrrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, pirazolinyl, piperidyl, piperazinyl, morpholinyl, morpholino, thiomorpholinyl, thiomorpholino, dihydropyridyl, tetrahydropyridyl, tetrahydrofuryl, tetrahydropyranyl, dihydrothiazolyl, tetrahydrothiazolyl, tetrahydroisothiazolyl, dihydrooxazinyl, hexahydroazepinyl, tetrahydrodiazepinyl and tetrahydropyridazinyl etc.;

a fused bicyclic heterocyclic group such as indolyl, isoindolyl, indazolyl, indolidinyl, indolinyl, isoindolinyl, quinolyl, isoquinolyl, cinnolinyl, phthaladinyl, quinazolinyl, naphthilidinyl, quinoxalinyl, purinyl, pteridinyl, benzopyranyl, benzimidazolyl, benzotriazolyl, benzisoxazolyl, benzoxazolyl, benzoxadiazolyl, benzoisothiazolyl, benzothiazolyl, benzothiadiazolyl, benzofuryl, isobenzofuryl, benzothienyl, benzotriazolyl, thienopyridyl, thienopyrrolyl, thienopyrazolyl, thienopyrazinyl, furopyrrolyl, thienothienyl, imidazopyridyl, pyrazolopyridyl, thiazolopyridyl, pyrazolopyrimidinyl, pyrazolotriazinyl, pyridazolopyridyl, triazolopyridyl, imidazothiazolyl, pyrazinopyridazinyl, quinazolinyl, quinolyl, isoquinolyl, naphthilidinyl, dihydrothiazolopyrimidinyl, tetrahydroquinolyl, tetrahydroisoquinolyl, dihydrobenzofuryl, dihydrobenzoxazinyl, dihydrobenzimidazolyl, tetrahydrobenzothienyl, tetrahydrobenzofuryl, benzodioxolyl, benzodioxonyl, chromanyl, chromenyl, octahydrochromenyl, dihydrobenzodioxinyl, dihydrobenzooxedinyl, dihydrobenzooxepinyl and dihydothienodioxinyl etc.; and a fused

tricyclic heterocyclic group such as carbazolyl, acridinyl, xanthenyl, phenothiazinyl, phenoxathiinyl, phenoxadinyl, dibenzofuryl, imidazoquinolyl and tetrahydrocarbazolyl etc.; and preferably includes 5- or 6-membered heteroaryl and a non-aromatic heterocyclic group.

A moiety of the heterocyclic group in "heterocyclyl lower alkyl", "heterocyclyloxy", "heterocyclylthio", "heterocyclylcarbonyl", "heterocyclyl lower alkoxy", "heterocyclyl amino", "heterocyclyl carbonylamino", "heterocyclyl sulfamoyl", "heterocyclylsulfonyl", "heterocyclylcarbamoyl", "heterocyclyloxycarbonyl", "heterocyclyl lower alkylamino", "heterocyclyl lower alkoxycarbonyl" and ""heterocyclyl lower alkylcarbamoyl" is the same as the "heterocyclic group" above.

"A nitrogen-containing aromatic heterocyclic group" means a group of the "heterocyclic group" above containing at least one nitrogen atom, and examples of the "nitrogen-containing aromatic heterocyclic group" include 5- or 6-membered heteroaryl such as pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazolyl, triazinyl, tetrazolyl, isoxazolyl, oxazolyl, oxadiazolyl, isothiazolyl, thiazolyl and thiadiazolyl etc.;

a fused bicyclic heterocyclo group such as indolyl, isoindolyl, indazolyl, indolidinyl, indolinyl, isoindolinyl, quinolyl, isoquinolyl, cinnolinyl, phthaladinyl, quinazolinyl, naphthilidinyl, quinoxalinyl, purinyl, pteridinyl, benzopyranyl, benzimidazolyl, benzotriazolyl, benzisoxazolyl, benzoxazolyl, benzoxadiazolyl, benzoisothiazolyl, benzothiazolyl, benzothiadiazolyl, benzotriazolyl, imidazopyridyl, pyrazolopyridine, triazolopyridyl, imidazothiazolyl, pyrazinopyridazinyl, quinazolinyl,quinolyl, isoquinolyl, naphthylidinyl, dihydobenzofuryl, tetrahydroquinolyl, tetrahydroisoquinolyl, dihydobenzoxazine etc.; and

a fused tricyclic heterocyclo group such as carbazolyl, acridinyl, xanthenyl and imidazoquinolyl etc.; and pyrrolidinyl, pyrolinyl, imidazolidinyl, imidazolinyl, pyrazolidinyl, pyrazolinyl, piperidyl, piperazinyl, morpholinyl, morpholino, thiomorpholinyl, thiomorpholino, dihydropiridyl, dihydrobenzimidazolyl, tetrahydropyridyl, tetrahydrothiazolyl and tetrahydroisothiazolyl etc.

"The heterocyclic group" or "nitrogen-containing aromatic heterocyclic group" above may be linked to other group at any position on the ring.

"Nitrogen-containing aromatic monocyclic heterocyclic group" means a monocyclic group in the "nitrogen-containing aromatic heterocyclic group" and examples of the "Nitrogen-containing aromatic monocyclic heterocyclic group" include 5- or 6-membered heteroaryl such as pyrrolyl, imidazolyl, pyrazolyl, pyridyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazolyl, triazinyl, tetrazolyl, isoxazolyl, oxazolyl, oxadiazolyl, isothiazolyl, thiazolyl and thiadiazolyl etc.

"The nitrogen-containing aromatic monocyclic heterocyclic group" above may be linked to other group at any carbon atom on the ring.

Examples of a substituent in the "optionally substituted carbocyclic group" and "optionally substituted heterocyclic group" of the ring A and B include

the substituent group α (preferably halogen, hydroxyl, acyl, acyloxy, carboxy, lower alkoxycarbonyl, carbamoyl, amino, cyano, lower alkylamino, lower alkylthio etc.);

lower alkyl optionally substituted with one or more substituent(s) selected from the substituent group α, hydroxyimino and lower alkoxyimino, wherein examples of preferable substituents include halogen, hydroxyl, lower alkoxy, lower alkoxycarbonyl etc.;

amino lower alkyl substituted with one or more substituent(s) selected from the substituent group α, wherein examples of preferable substituents include acyl, lower alkyl and/or lower alkoxy etc.;

hydroxyimino lower alkyl, lower alkoxyimino lower alkyl;

lower alkenyl optionally substituted with one or more substituent(s) selected from the substituent group α, wherein examples of preferable substituents include lower alkoxycarbonyl, halogen and/or halogeno lower alkoxycarbonyl;

lower alkynyl optionally substituted with one or more substituent(s) selected from the substituent group α, wherein examples of preferable substituents include lower alkoxycarbonyl etc;

lower alkoxy optionally substituted with one or more substituent(s) selected from the

substituent group α, wherein examples of preferable substituents include halogenocarbamoyl, oxetane, lower alkylcarbamoyl, hydroxyl lower alkylcarbamoyl;

lower alkoxy lower alkoxy optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkenyloxy optionally substituted with one or more substituent(s) selected from the substituent group α wherein examples of preferable substituents include halogen, hydroxyl, amino, lower alkyl etc.;

lower alkoxy lower alkenyloxy optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkynyloxy optionally substituted with one or more substituent(s) selected from the substituent group α, wherein examples of preferable substituents include halogen, hydroxyl etc.;

lower alkoxy lower alkynyloxy optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkylthio optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkenylthio optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkynylthio optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkylamino substituted with one or more substituent(s) selected from the substituent group α;

lower alkenylamino substituted with one or more substituent(s) selected from the substituent group α;

lower alkynylamino substituted with one or more substituent(s) selected from the substituent group α;

aminooxy optionally substituted with one or more substituent(s) selected from lower alkylidene and the substituent group α;

acyl substituted with one or more substituent(s) selected from the substituent group α;

lower alkylsulfonyl optionally substituted with one or more substituent(s) selected from the substituent group α;

lower alkylsulfinyl optionally substituted with one or more substituent(s) selected from the substituent group α;

sulfamoyl;

lower alkylsulfamoyl optionally substituted with one or more substituent(s) selected from the substituent group α;

a carbocyclic group (preferably cycloalkyl, aryl etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

a heterocyclic group optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclyl lower alkyl (preferably cycloalkyl lower alkyl, aryl lower alkyl etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl ;

heterocyclyl lower alkyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclyloxy (preferably cycloalkyloxy, aryloxy etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclyloxy optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclyl lower alkoxy (preferably cycloalkyl lower alkoxy, aryl lower alkoxy, etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclyl lower alkoxy (preferably cycloalkyl lower alkoxycarbonyl, aryl lower alkoxycarbonyl etc.) optionally substituted with one or more substituent(s) selected from the

substituent group α, azide, lower alkyl and halogeno lower alkyl,

carbocyclyl lower alkoxycarbonyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclyl lower alkoxycarbonyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclylthio (preferably cycloalkylthio, arylthio etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclylthio optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclyl amino (preferably cycloalkylamino, arylamino etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl,;

heterocyclyl amino optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclyl lower alkylamino (preferably cycloalkyl lower alkylamino, aryl lower alkylamino etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclyl lower alkylamino optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

lower alkylsulfamoyl optionally substituted with one or more substituent(s) selected from the substituent group α;

carbocyclylsulfamoyl (preferably cycloalkyl sulfamoyl, arylsulfamoyl etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclylsulfamoyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclylsulfonyl (preferably cycloalkyl sulfonyl, arylsulfonyl etc.) optionally substituted

with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclylsulfonyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclylcarbamoyl (preferably cycloalkyl carbamoyl, aryl carbamoyl etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclyl carbamoyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

carbocyclyl lower alkylcarbamoyl (preferably cycloalkyl lower alkylcarbamoyl, aryl lower alkylcarbamoyl etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

heterocyclyl lower alkylcarbamoyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl,

carbocyclyloxycarbonyl (preferably cycloalkyloxycarbonyl, aryloxycarbonyl etc.) optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;


heterocyclyloxycarbonyl optionally substituted with one or more substituent(s) selected from the substituent group α, azide, lower alkyl and halogeno lower alkyl;

lower alkylenedioxy optionally substituted with halogen;


oxo, azide and the like.

These may be substituted with one or more substituents selected from these groups.


Also the ring A may be substituted with one or more group(s) selected from




(i)




(ii)




(iii)




(iv)




(v)




(vi)




(vii)




(viii)




(ix)




(x)




(xi)




(xii)




(xiii)




(xiv)




(xv)




(xvi)



(xvii)




(xviii)



(xix)

or



(xixi)

wherein Ak^1, Ak^2 and Ak^3 are each independently a single bond, optionally substituted lower alkylene, optionally substituted lower alkenylene or optionally substituted lower alkynylene;

Ak^4 is optionally substituted lower alkylene, optionally substituted lower alkenylene or optionally substituted lower alkynylene;

W^1 and W^3 are each independently O or S,

W^2 is O, S or NR^5,

R^5 and R^6 are each independently hydrogen, lower alkyl, hydroxy lower alkyl, lower alkoxy lower alkyl, lower alkoxycarbonyl lower alkyl, carbocyclyl lower alkyl, lower alkenyl,


hydroxyl lower alkenyl, lower alkoxy lower alkenyl, lower alkoxycarbonyl lower alkenyl, carbocyclyl lower alkenyl, lower alkynyl, hydroxyl lower alkynyl, lower alkoxy lower alkynyl, lower alkoxycarbonyl lower alkynyl, carbocyclyl lower alkynyl or acyl;

R^7 is hydrogen or lower alkyl;


the ring B is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group; and

p is 1 or 2; W^1, W^3 or W^5 may be independent when it is pluralized. Additionally the oxygen atom of (xii) may be cis or trans to the substituent R^7.


Preferable examples of (i) to (xixi) above include




(ii')




(ii")




(iv')



(v')




(x')



(xiii')

or



(xiv')

wherein Ak is optionally substituted lower alkylene, optionally substituted lower alkenylene or optionally substituted lower alkynylene, and the other symbols are the same as described above.

In other cases of "an optionally substituted carbocyclic group" and "an optionally substituted heterocyclic group", one or more substituent(s) selected from a group of lower alkyl and the substituent group α may be exemplified as a substituent of "an optionally substituted carbocyclic group" and "an optionally substituted heterocyclic group"

"Heteroaryl" includes an aromatic cyclic group among the "heterocyclic group" above.

"Lower alkylene" includes C1-C10, preferably C1-C6, more preferably C1-C3 straight or branched divalent carbon chain , and for example, methylene, dimethylene, trimethylene, tetramethylene and methyl trimethylene are exemplified.

A moiety of lower alkylene in "lower alkylenedioxy" is the same as the "lower alkylene" described above.

"Lower alkenylene" includes C2-C10, preferably C2-C6, more preferably C2-C4 straight or branched divalent carbon chain having a double bond at any arbitrary position thereof, and vinylene, propenylene, butenylene, butadienylene, methyl propenylene, pentenylene and hexenylene are exemplified.

"Lower alkynylene" includes C2-C10, preferably C2-C6, more preferably C2-C4 straight or branched divalent carbon chain having a triple bond and also a double bond at any arbitrary position thereof, and for example, ethynylene, propynylene, butynylene, pentynylene and hexynylene are exemplified.

Examples of a substituent in "optionally substituted lower alkylene", "optionally substituted lower alkenylene" and "optionally substituted lower alkynylene" include the substituent group α、 and preferably halogen and hydroxyl etc. are exemplified.


Examples of a substituent in "optionally substituted carbocyclyl lower alkyl", "optionally substituted heterocyclyl lower alkyl", "optionally substituted carbocyclyl lower alkoxy", and "optionally substituted heterocyclyl lower alkoxy" include one or more substituent(s) selected from lower alkyl and the substituent group α.

In this specification, "solvate" includes a solvate with an organic solvent and a hydrate etc. and hydrate may be coordinated with optional number of water molecule.

The compound (I) includes pharmaceutical acceptable salt thereof. Examples of the pharmaceutical acceptable salt include a salt with an alkali metal such as lithium, sodium and potassium etc., an alkali earth metal such as magnesium, calcium etc., ammonium, an organic base and an amino acid; a salt with an inorganic acid such as hydrochloric acid, sulfuric acid, nitric acid, hydrobromic acid, phosphoric acid or hydroiodic acid etc., and an organic acid such as acetic acid, trifluoroacetic acid, citric acid, lactic acid, tartaric acid, oxalic acid, maleic acid, fumaric acid, mandelic acid, glutaric acid, malic acid, benzoic acid, phthalic acid, benzensulfonic acid, p-toluenesulfonic acid, methanesulfonic acid, ethanesulfonic acid, etc. Especially hydrochloric acid, phosphoric acid, tartaric acid or


methane sulfonic acid is preferable. These salts can be prepared by a method usually carried out.

The compound (I) is not construed to be limited to a specific isomer but to include all possible isomers such as a keto-enol isomer, an imine-enamine isomer, a diastereoisomer, an optical isomer and a rotational isomer etc. For example, a compound (I) in which R^{2a} is hydrogen includes a tautomer as follows;



The compound (I) of the present invention can be prepared, for example, according to the non-patent literature 1 or a method described below;

Preparation of an aminodihydrothiazine ring (I-1) or (I-2):



(In the scheme above, at least one of R^{2b} and R^{2c} is hydrogen, R^{3b} and R^{3d} are each independently hydrogen, halogen, hydroxyl, optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted acyl, optionally substituted lower alkoxy, optionally substituted lower alkylthio, carboxy, optionally substituted lower alkoxycarbonyl, optionally substituted amino, optionally substituted carbamoyl, an optionally substituted


carbocyclic group, or an optionally substituted heterocyclic group, and the other symbols are the same as described above.)

The 1st step: A Grignard reagent having a corresponding substituent of the objective compound such as vinyl magnesium chloride, vinyl magnesium bromide and propenyl magnesium bromide etc. is added to a compound a, which is commercially available or can be prepared by a known method, in a solvent such as ether, tetrahydrofuran etc. or a mixed solvent of ether-tetrahydrofuran etc., at -100°C to 50°C, preferably -80°C to 0°C and the mixture is stirred for 0.2 to 24 hours, preferably 0.2 to 5 hours to give a compound b.

The 2nd step: To a compound b in an acid such as acetic acid, trifluoroacetic acid, hydrochloric acid, sulfuric acid etc. or a mixture thereof under the presence of a solvent such as toluene etc. or without a solvent, is added a substituted thiourea having a corresponding substituent of the objective compound such as thiourea, N-methylthiourea, N,N'-dimethylthiourea etc., and the mixture is stirred at -20°C to 100°C, preferably 0°C to 80°C for 0.5 hours to 120 hours, preferably 1 hour to 72 hours to give a compound c.

The 3rd step: To a compound c in a solvent such as toluene etc. or without a solvent, is added an acid such as trifluoroacetic acid, methanesulfonic acid, trifluoromethanesulfonic acid, sulfuric acid etc. or a mixture thereof and reacted at -20°C to 100°C, preferably 0°C to 50°C for 0.5 hours to 120 hours, preferably 1 hour to 72 hours to give a compound (I-2) when R^{2b} is hydrogen or a compound (I-1) when R^{2c} is hydrogen.

Preparation of an aminodihydrothiazine ring (I-3):



(In the scheme above, L is a leaving group such as halogen or lower alkylsulfonyl etc. and

the other symbols are the same as described above.)


The 1st step: Thiocyanate such as sodium thiocyanate or ammonium thiocyanate etc. is reacted with a compound d, which is commercially available or can be prepared by a known method, in a solvent such as toluene, chloroform, tetrahydrofuran etc. under the presence of water and an acid such as hydrochloric acid or sulfuric acid etc. at 0°C to 150°C, preferably at 20°C to 100°C for 0.5 to 24 hours, preferably 1 to 12 hours to give a compound e.

The 2nd step: A reducing agent such as sodium borohydride etc. is added to and reacted with a compound e in a solvent such as tetrahydrofuran, methanol, ethanol, water etc. or a mixture of ethanol-water etc. under the presence of buffering agent such as sodium dihydrogen phosphate at -80°C to 50°C, preferably at -20°C to 20°C for 0.1 to 24 hours, preferably 0.5 to 12 hours to give a compound f.

The 3rd step: A compound f is reacted with a halogenating agent such as thionyl chloride, phosphoryl chloride, carbon tetrachloride-triphenylphosphine etc. in a solvent such as toluene, dichloromethane etc. or without a solvent at -80°C to 50°C, preferably at -20°C to 20°C for 0.1 to 24 hours, preferably 0.5 to 12 hours; or it is reacted with a sulfonating agent such as methanesulfonyl chloride, p-toluenesulfonyl chloride etc. in a solvent such as toluene, dichloromethane etc. under the presence of a base such as triethylamine etc. at -80°C to 50°C, preferably at -20°C to 20°C for 0.1 to 24 hours, preferably 0.5 to 12 hours to give a compound g.

The 4th step: A compound g is reacted with ammonia or a primary amine such as methylamine etc. in a solvent such as methanol, ethanol, water etc. or a mixture of methanol-water etc. at -20°C to 80°C, preferably at 0°C to 40°C for 0.5 to 48 hours, preferably 1 to 24 hours to give the compound (I-3).

Preparation of an aminodihydrothiazine ring (I-6) or an aminotetrahydrothiazine ring(I-7)



o p q


(In the scheme above, at least one of R^{2b} and R^{2c} is hydrogen and the other symbols are the same as described above.)

The 1st step: Thiourea or a substituted thiourea corresponding to the objective compound such as N-methyl thiourea, N,N-dimethylthiouers, N,N'-dimethylthiouera etc. is reacted with a compound o, which is commercially available or can be prepared by a known method, in a solvent such as ethanol, methanol, tetrahydrofuran, toluene etc. at -20°C to 200°C, preferably at 0°C to 150°C for 0.5 to 200 hours, preferably 1 to 120 hours to give a compound p.

The 2nd step: A Grignard reagent corresponding to the objective compound such as methyl magnesium chloride, ethyl magnesium bromide and benzyl magnesium bromide etc. is added to a compound p in a solvent such as ether, tetrahydrofuran etc. or a mixed solvent thereof at -100°C to 50°C, preferably -80°C to 30°C and the mixture is stirred for 0.2 to 24 hours, preferably 0.5 to 5 hours to give a compound q.

The 3rd step: To a compound q in a solvent such as toluene etc. or without a solvent, is added an acid such as trifluoroacetic acid, methanesulfonic acid, trifluoromethanesulfonic acid, sulfuric acid etc. or a mixture thereof and reacted at -20°C to 100°C, preferably 0°C to 50°C for 0.5 hours to 200 hours, preferably 1 hour to 150 hours to give a compound (I-6) (R^{2c} = H) or a compound (I-7) (R^{2b} = H).

Preparation of an aminodihydrothiazine ring (I-8)



(In the scheme, each symbol is the same as described above)

The 1st step: Ammonium chloride is added to a compound r which can be prepared by a known method in a solvent such as acetic acid etc. at 0°C to 200°C, preferably 10°C to 100°C for 0.1 hours to 100 hours, preferably 0.5 hour to 24 hours to give a compound.


The 2nd step: A reducing agent such as lithium aluminium hydride, diisobutyl aluminium hydride etc. is reacted with a compound s in a solvent such as tetrahydrofuran, diethyl ether etc. at -80°C to 150°C, preferably 0°C to 100°C for 0.1 hours to 24 hours, preferably 0.5 hour to 12 hours to give a compound t.

The 3rd step: Isothiocyanate corresponding to the objective compound such as 4-methoxybenzyl isothiocyanate, t-butyl isothiocyanate etc. or carbamoyl halide corresponding to the objective compound such as N,N-dimethyl thiocarbamoyl chloride, N,N-diethyl thiocarbamoyl chloride etc. is reacted with a compound t in a solvent such as toluene, chloroform, tetrahydrofuran etc. under the presence of a base such as diisopropylethylamine, triethylamine, pyridine, sodium hydroxide etc. or without a base at 0°C to 150°C, preferably 20°C to 100°C for 0.5 hours to 120 hours, preferably 1 hour to 72 hours to give a compound u.

The 4th step: A halogenating agent such as thionyl chloride, phosphoryl oxychloride, carbon tetrachloride-triphenyl phosphine etc. is reacted with a compound u in a solvent such as acetonitrile, toluene, dichloromethane etc. at -80°C to 50°C, preferably -20°C to 20°C for 0.1 hours to 24 hours, preferably 0.5 hour to 12 hours, or a sulfonylating agent such as

methanesulfonyl chloride, p-toluenesulfonyl chloride is reacted with a compound u in a solvent such as toluene, dichloromethane etc. under the presence of a base such as triethylamine at -80°C to 50°C, preferably -20°C to 20°C for 0.1 hours to 24 hours, preferably 0.5 hour to 12 hours. The resulting halogenated compound or sulfonate ester derivative is reacted with a base such as diisopropylethylamine, potassium carbonate, sodium bicarbonate, sodium hydride, sodium hydroxide etc. at 0°C to 150°C, preferably 20°C to 100°C for 0.5 hours to 120 hours, preferably 1 hour to 72 hours to give a compound (I-8).


Preparation of an acylamino derivative (I-13) and/or (I-14)



(In the scheme, R^{17} is optionally substituted lower alkyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group and other symbols are the same as described above)

An acylating agent corresponding to the objective compound such as benzoyl chloride, 2-furoyl chloride, acetic anhydride etc. is reacted with a compound (I-12) in which R^{2b} is hydrogen under the presence of a solvent such as tetrahydrofuran, dichloromethane etc. or without a solvent and under the presence of a base such as pyridine or triethylamine etc. or without a solvent at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours, or a compound (I-12) is reacted with a carboxylic acid having a substituent corresponding to the objective compound such as amino acid or glycolic acid etc. in a solvent such as dimethylformamide, tetrahydrofuran, dichloromethane etc. under the presence of a condensation agent such as dicyclohexylcarbodiimide, carbonyldiimidazole etc. at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours to give a compound (I-13) and/or (I-14)(when R^{2a} is hydrogen).


Preparation of a carbamoyl derivative(I-17)



(In the scheme above, $CONR^{18}R^{19}$ is optionally substituted carbamoyl and the other symbols are the same as described above)

A compound (I-16) having a carboxyl group as a substituent on the ring A is reacted with a primary or secondary amine having a substituent corresponding to the objective compound (e.g., aniline, 2-aminopyridine, dimethylamine etc.) in a solvent such as dimethylformamide, tetrahydrofuran, dichloromethane etc. under the presence of a condensation agent such as dicyclohexylcarbodiimide, carbonyldiimidazole, dicyclohexylcarbodiimide-N-hydroxybenzotriazole etc. at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours to give a compound (I-17).

Preparation of an acylamino derivative (I-19)




(In the scheme, NHR^{20} is optionally substituted amino, $NR^{20}COR^{21}$ is optionally substituted acylamino, optionally substituted ureido or carboxyamino having a substituent on the oxygen atom and the other symbols are the same as described above.)

A compound (I-18) having an optionally substituted amino group on the ring A is reacted with a reagent having a substituent corresponding to the objective compound such as acid chlorides, acid anhydrides, chlorocarbonate esters, isocyanates etc. under the presence of a solvent such as tetrahydrofuran, dichloromethane etc. or without a solvent under the presence of a base such as pyridine, triethylamine etc. or without a base at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours, or a

compound (I-18) is reacted with a carboxylic acid having a substituent corresponding to the objective compound such as benzoic acid, 2-pyridinecarboxylic acid etc. in a solvent such as dimethylformamide, tetrahydrofuran, dichloromethane etc. under the presence of a condensation agent such as dicyclohexylcarbodiimide, carbonyldiimidazole, dicyclohexylcarbodiimide-N-hydroxybenzotriazole etc. at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours to give a compound (I-19).


Preparation of an alkylamino derivative (I-20)



(In the scheme, NHR^{20} is optionally substituted amino and R^{22} is lower alkyl.)

A compound (I-18) having an amino group on the ring A is reacted with an aldehyde having a substituent corresponding to the objective compound such as benzaldehyde, pyridine-2-carboaldehyde etc. and a reducing agent such as sodium cyanoborohydride, sodium triacetoxyborohydride etc. in a solvent such as dichloromethane, tetrahydrofuran etc. under the presence of an acid such as acetic acid etc. or without an acid at -80°C to 100°C, preferably 0°C to 40°C for 0.5 hours to 150 hours, preferably 1 hour to 24 hours to give a compound (I-20).

Preparation of a substituted alkoxy derivative (I-22).




(In the scheme above, R^{23} is optionally substituted lower alkyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group and the other symbols are

the same as described above.)

A compound (I-21) having a hydroxy group on the ring A is reacted with an alkylating agent having a substituent corresponding to the objective compound such as benzyl chloride, methyl iodide etc. in a solvent such as dimethylformamide, tetrahydrofuran etc. under the presence of a base such as potassium carbonate, sodium hydroxide, sodium hydride etc. at -80°C to 100°C, preferably 0°C to 40°C for 0.5 hours to 150 hours, preferably 1 hour to 24 hours, or a compound (I-18) is reacted with an alcohol such as 2-aminoethanol etc. in a solvent such as dimethylformamide, tetrahydrofuran etc. under the presence of a Mitsunobu reagent such as triphenylphosphine-azodicarboxylic acid diethyl ester etc. at -80°C to 100°C, preferably 0°C to 40°C for 0.5 hours to 72 hours, preferably 1 hour to 24 hours to give a compound (I-22).

Introduction of a substituent by palladium coupling




(In the scheme above, Hal is halogen, G is optionally substituted lower alkenyl, optionally substituted alkynyl, optionally substituted alkoxycarbonyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group etc. and the other symbols are the same as described above.)

A compound (I-23) having halogen as a substituent on the ring A is reacted with a compound having a substituent corresponding to the objective compound (e.g., styrene, propargyl alcohol, aryl boronic acid, carbon monoxide etc.) in a solvent such as tetrahydrofuran, dimethylformamide, 1,2-dimethoxyethane, methanol etc. under the presence of a base such as triethylamine, sodium carbonate etc., a palladium catalyst such as palladium acetate, palladium chloride etc. and a ligand such as triphenylphosphine etc. and under irradiation of microwave or without the irradiation, at -80°C to 150°C, preferably 0°C

to 100°C for 0.5 hours to 72 hours, preferably 1 hour to 24 hours to give a compound (I-24).

Preparation of an oxime derivative (I-26)




(I-25) (I-26)

(In the scheme above, R^{24} is hydrogen, optionally substituted lower alkyl etc., R^{25} is hydrogen, optionally substituted lower alkyl, optionally substituted lower alkenyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group etc., and the other symbols are the same as described above.)

A compound (I-25) having an acyl group as a substituent of the ring A is reacted with a hydroxylamine having a substituent corresponding to the objective compound such as hydroxylamine, methoxylamine, O-benzylhydroxylamine etc. or a salt thereof in a solvent such as methanol, ethanol etc. under the presence of an additive such as potassium acetate etc. or without an additive at -80°C to 100°C, preferably 0°C to 40°C for 0.5 hours to 150 hours, preferably 1 hour to 72 hours to give a compound (I-26).

Coupling reaction



(I-27)

(In the scheme above, R^{26} is a substituent corresponding to each objective compound)

The 1st step:

A compound v is reacted with a reagent having a substituent corresponding to the objective compound such as acyl halide, acid anhydride, chlorocarbonate ester, isocyanate etc. (e.g., benzoyl chloride, 2-furoyl chloride, acetic anhydride, benzyl chloroformate, di-tert-butyl dicarbonate, phenyl isocyanate etc.) in a solvent such as tetrahydrofuran,

dichloromethane, dimethylformamide etc. or without a solvent under the presence of a base such as pyridine, triethylamine etc. or without a base at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours, or a compound A is reacted with a carboxylic acid having a substituent corresponding to the objective compound such as benzoic acid, 2-pyridinecarboxylic acid etc. in a solvent such as dimethylformamide, tetrahydrofuran, dichloromethane, methanol etc. under the presence of a condensation agent such as dicyclohexylcarbodiimide, carbonyl diimidazole, dicyclohexylcarbodiimide-N-hydroxybenzotriazole, 4-(4,6-dimethoxy-1,3,5-triazine-2-yl)-4-methylmorpholinium chloride, 2-(7-aza-1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate etc. at -80°C to 100°C, preferably -20°C to 40°C for 0.1 hours to 24 hours, preferably 1 hour to 12 hours to give a compound w.

When the substituent R has a functional group which disturb the said reaction, it can be carried out by protecting the functional group with a suitable protecting group and then deprotecting it at a subsequent appropriate step.


The 2nd step:

A compound w is reacted in a solvent such as methanol, ethanol, ether, tetrahydrofuran, 1,4-dioxane, dichloromethane, ethyl acetate etc. containing trifluoroacetic acid etc. or in neat, or in neat trifluoroacetic acid at -30°C to 100°C, preferably 0°C to 90°C for 0.5 to 12 hours to give a compound (I-27). Alternatively, the objective compound can be synthesized according to the method described in Protective Groups in Organic Synthesis, Theodora W Green (John Wiley & Sons) etc.

Preparation of an optically active isomer

1) Preparation of an optically active isomer ae

For example, an optically active isomer ae, one embodiment of the compounds of the present invention, can be prepared according to the following scheme:



(In the scheme above, R^1 is optionally substituted lower alkyl, optionally substituted lower alkenyl or optionally substituted lower alkynyl; R^{27} is a chiral sulfoxide having optionally substituted lower alkyl, optionally substituted lower alkenyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group, or a chiral auxiliary group such as α-methyl benzyl etc.; R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogen, hydroxyl, optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted acyl, carboxy, optionally substituted lower alkoxycarbonyl, optionally substituted amino, optionally substituted carbamoyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group; R^{28} is optionally substituted lower alkyl or optionally substituted lower alkenyl; R^{2a} and R^{2b} are each independently hydrogen, optionally substituted lower alkyl or optionally substituted acyl; and the other symbols are the same as described above.)

The compounds y and z above can be prepared by a method described in

(1) T. Fujisawa et al., Tetrahedron Lett., 37, 3881-3884 (1996),

(2) D. H. Hua et al, Sulfur Reports, vol. 21, pp. 211-239 (1999)

(3) Y. Koriyama et al., Tetrahedron, 58, 9621-9628 (2002), or

(4) T. Vilavan et al, Cuuent Organic Chemistry, 9, 1315-1392 (2005).

Alternatively, these compounds can be prepared by optical resolution of each

intermediate or the final product, or according to methods described below. Examples of the optical resolution method include a separation of optical isomers using an optically active column, kinetic resolution by an enzyme reaction etc., crystallization of diastereomers by salt formulation using a chiral acid or chiral base, and a preferential crystallization etc.

The 1st step: Compound y can be obtained by reacting Compound x, which is commercially available or can be prepared by a known method, with a chiral reagent having a substituent corresponding to the objective compound such as α-methylbenzylamine, para-toluene, tert-butylsulfine amide etc. at 60 °C to 120 °C, preferably 80 °C to 100 °C in a solvent such as ether, tetrahydrofuran, toluene, benzene etc. or a mixed solvent such as ether-tetrahydrofuran etc. for 0.5 to 24 hours, preferably 0.5 to 5 hours, in the presence of molecular sieves or magnesium sulfate etc., under continuous evaporation by Dean-Stark apparatus, or according to the method described in the above literatures.

The 2nd step: A compound z can be diastereo-selectively obtained by reacting an enolate of lithium, aluminium, zinc, titan etc. prepared form a reagent having a substituent corresponding to the objective compound such as acetate ester etc., which is commercially available or can be prepared by a known method, or ketenesilyl acetate prepared from a reagent having a substituent corresponding to the objective compound such as ethyl acetate etc. with a compound a in a solvent such as ether, tetrahydrofuran, toluene, dichloromethane etc. or a mixed solvent such as ether-tetrahydrofuran etc. under the presence of a Lewis acid such as titanium tetrachloride, ether-trifluoroborane complex etc. or without a Lewis acid at -100°C to 50°C, preferably -80°C to -30°C for 0.5 to 24 hours, preferably 0.5 to 5 hours. Alternatively, the compound z can be diastereo-selectively prepared by the method described in the literature (1) or (3).

The 3rd step: A compound z is reacted with a compound c in a solvent such as methanol, ethanol, ether, tetrahydrofuran, 1,4-dioxane, dichloromethane, ethyl acetate etc. containing hydrogen chloride, trifluoroacetic acid etc. or in neat trifluoroacetic acid at -30°C to 100°C, preferably -10°C to 90°C for 0.5 to 12 hours, preferably 0.5 to 5 hours to give a compound aa.

The 4th step: A reducing agent such as borane-tetrahydrofuran complex, borane-dimethyl sulfoxide complex, borane-triethylamine complex, borane-pyridine complex etc. or ether- or tetrahydrofuran-solution thereof is reacted with a compound aa in a solvent such as ether, tetrahydrofuran, toluene etc. or a mixed solvent such as ether-tetrahydrofuran etc. at -30°C to 30°C, preferably -10°C to 20°C for 0.5 to 12 hours, preferably 0.5 to 5 hours to give a compound ab.

The 5th step: Calcium carbonate or potassium carbonate etc. is added to a compound ab in a solvent such as dichloromethane, toluene etc. or a mixed solvent such as dichloromethane-water etc. and thiophosgene is added at -30°C to 50°C, preferably -10°C to 25°C and the mixture is reacted for 0.5 to 12 hours, preferably 0.5 to 5 hours to give a compound ac.


The 6th step: Oxalyl chloride or thionyl chloride etc. and a catalytic amount of N,N-dimethylformamide are added to a compound ac in a solvent such as dichloromethane, tetrahydrofuran, toluene etc. at -30°C to 50°C, preferably -10°C to 20°C and the mixture is reacted at 0 °C to 100 °C, preferably 20 °C to 90 °C for 0.5 to 12 hours, preferably 0.5 to 5 hours to give a compound ad. Alternatively, it is obtained by a method described in Comprehensive Organic Transformations, Richard C Larock (Mcgraw-Hill).

The 7th step: 15% to 30% Ammonia water or a reagent having a substituent corresponding to the objective compound such as tert-butylamine etc. is added to a compound ad in a solvent such as ethyl acetate, dichloromethane, tetrahydrofuran, toluene etc. at -30°C to 50°C, preferably -10°C to 30°C and the mixture is reacted at -10°C to 30°C, preferably 0°C to 30°C for 0.5 to 72 hours to give a compound ae-i or a compound ae-ii.

When R^{2a} and/or R^{2b} is hydrogen in the resulting compound ae-i or ae-ii, a substituent of the objective compound, R^{2a} and/or R^{2b}, may be further introduced by a conventional method if it is necessary.

1') Method for preparing an optically active isomer Method B

An optically active compound ah of the present invention can be also prepared by a method below:



(In the scheme , the symbols are the same as described above.)


The 1st step to the 4th step: the same as described in 1) above.

The 5th step: Isothiocyanate having a protecting group which is commercially available or can be prepared by a known method is added to a compound ab in a solvent such as dichloromethane, toluene, acetone etc. or a mixed solvent at -30°C to 50°C, preferably -10°C to 25°C and the mixture is reacted for 0.5 to 12 hours, preferably 0.5 to 5 hours to give a compound ag.

The 6th step: Oxalyl chloride or thionyl chloride etc. and a catalytic amount of N,N-dimethylformamide are added to a compound ag in a solvent such as dichloromethane, tetrahydrofuran, toluene etc. at -30°C to 50°C, preferably -10°C to 25°C, or 1-chloro-N,N-2-trimethyl-1-propenenylamine is added to a compound ag, and reacted at 0°C to 100°C, preferably 20°C to 90°C for 0.5 to 72 hours to give a compound ah-i or ah-ii.

2) Introduction of R^{3a} and R^{3b}

An optically active compound ae-iii or ae-iv of the present invention can be also prepared by introducing R^{3a} and R^{3b} as shown below:



(In the scheme above, each symbol is the same as described above)


When preparing a compound ae-iii or ae-iv in which R^{3a} and R^{3b} are substituted on the carbon atom next to S atom, a compound z is processed through the 3'rd and 4'th steps in place of the 3rd and 4th steps of 1) described above, and R^{3a} and R^{3b} are introduced in advance.

The 3'rd step: A Grignard reagent having a substituent corresponding to the objective compound such as methyl magnesium chloride, ethyl magnesium bromide etc. is added to a compound z in a solvent such as ether, tetrahydrofuran etc. or a mixed solvent such as ether-tetrahydrofuran etc. at -100°C to 50°C, preferably -80°C to 30°C, or a compound z is converted to Weinreb Amide and reacted with a Grignard reagent having a substituent corresponding to the objective compound such as $R^{3a}MgBr$, $R^{3b}MgBr$. The reaction mixture is reacted for 0.2 to 24 hours, preferably 0.2 to 5 hours to give a compound aa'.

The 4'th step: A compound aa' is reacted in a solvent such as methanol, ethanol, ether, tetrahydrofuran, 1,4-dioxane, dichloromethane, ethyl acetate etc. containing hydrogen chloride, trifluoroacetic acid etc. or in neat trifluoroacetic acid at -30°C to 100°C, preferably -10°C to 90°C for 0.5 to 12 hours, preferably 0.5 to 5 hours to give a compound ab'.

The compound ab' is processed in the same reactions as the 5th to 7th steps of 1) above to give the objective compound ae-iii or ae-iv.


When the substituent L of a compound ad' is eliminated to give a compound ad'' shown below, the objective compound ae'-iii or ae'-iv is obtained by processing the compound ad'' it through the 7'th step in place of the 7th step described in 1) above.



The 7'th step: A compound ad'' is dissolved in conc. sulfuric acid, trifluoroacetic acid, trifluoromethanesulfonic acid etc. and reacted at -30°C to 100°C, preferably -10°C to 40°C for 0.1 to 12 hours, preferably 0.5 to 5 hours to give a compound ae'.

3) Conversion of a substituent (1)

A preparation of a compound af-1 by conversion of the substituent is illustrated below:



(In the scheme, R^{8a} and R^{8b} are an amino-protecting group, and the other symbols are the same as described above.)


Trisdibenzylideneacetonedipalladium, palladium acetate, palladium(0) prepared in situ etc. and a phosphine ligand such as tri-tert-butylphosphine, dicyclohexylbiphenylphosphine etc. are added to a compound ae-1 in a solvent such as tetrahydrofuran, toluene, xylene etc. and further a reagent having a substituent corresponding to the objective compound such as lithium hexamethylenedisilazide, benzophenonimine etc. is added thereto at -10°C to 30°C, then the reaction mixture is reacted at 30°C to 120°C, preferably 50°C to 100°C for 0.5 to 48 hours, preferably 3 to 20 hours to give a compound

af-1.

Any amino-protecting group which is deprotected by a method described in Protective Groups in Organic Synthesis, Theodora W Green(John Wiley & Sons) etc. can be used and examples of the protecting group include lower alkoxycarbonyl, lower alkenyloxycarbonyl, trialkylsilyl, acyl, methanesulfonyl, trifluoromethanesulfonyl and toluenesulfonyl etc.

4) Conversion of a substituent (2)

A preparation of a compound af-2 by conversion of the substituent is illustrated below:




(In the scheme, each symbol is the same as described above.)

A catalyst of catalytic reduction such as 10% palladium-carbon etc. is added to a compound ae-2 in a solvent such as tetrahydrofuran, ethyl acetate, methanol etc. and it is reacted under the pressure of normal to 5 atom, preferably normal to 2 atom of hydrogen atmosphere at 30°C to 120°C, preferably 50°C to 80°C for 0.5 to 48 hours, preferably 6 to 20 hours to give a compound af-2. Alternatively, the compound af-2 is obtained by a method described in Comprehensive Organic Transformations, Richard C Larock (Mcgraw-Hill).

5) Conversion of a substituent (3)

A preparation of a compound af-3 by conversion of the substituent is illustrated below:



(In the scheme, R^9 is hydroxyl, optionally substituted lower alkyl, optionally substituted lower alkoxy, optionally substituted lower alkylthio, optionally substituted lower alkyl amino, optionally substituted aromatic carbocyclyloxy, optionally substituted heterocyclyloxy, optionally substituted aromatic carbocyclylthio, optionally substituted heterocyclylthio, optionally substituted carbocyclylamino, optionally substituted heterocyclylamino, cyano, azide, an optionally substituted carbocyclic group, an optionally substituted heterocyclic group, optionally substituted carbamoyl etc. and the other symbols are the same as described above.)

A reagent having a substituent corresponding to the objective compound such as ethanol, methanthiol, dimethylamine etc. is added to a compound ae-3 in a solvent such as tetrahydrofuran, ethanol etc. under the presence of a base such as sodium methoxide, potassium tert-butoxide, sodium hydroxide, sodium hydride etc. or without a base at -10°C to 50°C and it is reacted for 0.5 to 12 hours, preferably 1 to 8 hours to give a compound af-3. If necessary, a coupling reaction may be carried out in the same manner as the method for preparing a compound (I-19) described above.


In every step described above, if a starting compound has a functional group which disturb the reaction (e.g., hydroxyl, mercapto, amino, formyl, carbonyl, carboxyl etc.), it is recommended to protect the functional group and deprotect it at a subsequent appropriate step with a method described in Protective Groups in Organic Synthesis, Theodora W Green(John Wiley & Sons) etc.

Further the order of steps may be changed and each reaction intermediate may be isolated and used in the subsequent step.

Examples of a preferable compound in the present invention include the

followings:

In a formula (I')



(I')

1) a compound in which the ring A' is phenyl or a nitrogen-containing aromatic heterocyclic group (hereinafter called a compound in which the ring A' is A'1),

a compound in which the ring A' is benzene, pyridine, indole, benzisoxazole, benzopyrazole, benzofuran, benzothiophene, benzodioxole, or dihydrobenzodioxolane (hereinafter called a compound in which the ring A' is A'2),

a compound in which the ring A' is benzene (hereinafter called a compound in which the ring A' is A'3),

a compound in which the ring A' is pyridine (hereinafter called a compound in which the ring A' is A'4)

2) a compound in which R^1 is optionally substituted lower alkyl (hereinafter called a compound in which R^1 is R1-1),

a compound in which R^1 is methyl (hereinafter called a compound in which R^1 is R1-2),

3) a compound in which R^{2a} and R^{2b} are each independently hydrogen, lower alkyl or acyl(hereinafter called a compound in which R^{2a} and R^{2b} are R2-1),

a compound in which both of R^{2a} and R^{2b} are hydrogens (hereinafter called a compound in which R^{2a} and R^{2b} are R2-2),

4) a compound in which R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogen, hydroxyl, lower alkyl or amino (hereinafter called a compound in which R^{3a}, R^{3b}, R^{3c} and R^{3d} are R3-1),

a compound in which R^{3a} and R^{3b}, or R^{3c} and R^{3d} taken together form cyclopropyl,

cyclobutyl, cyclopentyl or cyclohexyl together (hereinafter called a compound in which R^{3a}, R^{3b}, R^{3c} and R^{3d} are R3-2),

a compound in which R^{3a} and R^{3b},or R^{3c} and R^{3d} are the same substituent selected from halogen and lower alkyl (hereinafter called a compound in which R^{3a}, R^{3b}, R^{3c} and R^{3d} are R3-3),

a compound in which all of R^{3a}, R^{3b}, R^{3c} and R^{3d} are hydrogens (hereinafter called a compound in which R^{3a}, R^{3b}, R^{3c} and R^{3d} are R3-4),

5) a compound in which n is 0 to 2, R^4 is each independently halogen, lower alkoxy, lower alkylamino, lower alkylthio, oxo or lower alkylenedioxy (hereinafter called a compound in which R^4 is R4-1),

a compound in which n is 0 to 2, R^4 is each independently halogen (hereinafter called a compound in which R^4 is R4-2),

6) a compound in which G is (ii), (iv), (v), (x), (xiii) or (xiv) above (hereinafter called a compound in which G is G1),

a compound in which G is (ii'), (ii''), (iv'), (v'), (x'), (xiii') or (xiv') above (hereinafter called a compound in which G is G2),

a compound in which G is (ii'), (ii''), (iv'), (v'), (x'), (xiii') or (xiv') above, and the ring B is optionally substituted pyridyl, optionally substituted pyrazinyl, optionally substituted thiazolyl, optionally substituted isoxazolyl, optionally substituted benzothiazolyl, optionally substituted thiazolopyridyl, optionally substituted quinolyl, optionally substituted isoquinolyl or optionally substituted naphthylidinyl, optionally substituted quinazolinyl, or optionally substituted pyridopyrimidinyl (hereinafter called a compound in which G is G3),

a compound in which G is (ii') above (hereinafter called a compound in which G is G4),

a compound in which a combination of the ring A', R^1, R^{2a} and R^{2b}, R^{3a}, R^{3b}, R^{3c} and R^{3d}, n and R^4, and G is as follows;

(A'1,R1-1,R2-1,R3-1,R4-1,G1),(A'1,R1-1,R2-1,R3-1,R4-1,G2),(A'1,R1-1,R2-1,R3-1,R4-1,G3),(A'1,R1-1,R2-1,R3-1,R4-1,G4),(A'1,R1-1,R2-1,R3-1,R4-2,G1),(A'1,R1-1,R2-1,R3-1,R

4-2,G2),(A'1,R1-1,R2-1,R3-1,R4-2,G3),(A'1,R1-1,R2-1,R3-1,R4-2,G4),(A'1,R1-1,R2-1,R3-2,R4-1,G1),(A'1,R1-1,R2-1,R3-2,R4-1,G2),(A'1,R1-1,R2-1,R3-2,R4-1,G3),(A'1,R1-1,R2-1,R3-2,R4-1,G4),(A'1,R1-1,R2-1,R3-2,R4-2,G1),(A'1,R1-1,R2-1,R3-2,R4-2,G2),(A'1,R1-1,R2-1,R3-2,R4-2,G3),(A'1,R1-1,R2-1,R3-2,R4-2,G4),(A'1,R1-1,R2-1,R3-3,R4-1,G1),(A'1,R1-1,R2-1,R3-3,R4-1,G2),(A'1,R1-1,R2-1,R3-3,R4-1,G3),(A'1,R1-1,R2-1,R3-3,R4-1,G4),(A'1,R1-1,R2-1,R3-3,R4-2,G1),(A'1,R1-1,R2-1,R3-3,R4-2,G2),(A'1,R1-1,R2-1,R3-3,R4-2,G3),(A'1,R1-1,R2-1,R3-3,R4-2,G4),(A'1,R1-1,R2-1,R3-4,R4-1,G1),(A'1,R1-1,R2-1,R3-4,R4-1,G2),(A'1,R1-1,R2-1,R3-4,R4-1,G3),(A'1,R1-1,R2-1,R3-4,R4-1,G4),(A'1,R1-1,R2-1,R3-4,R4-2,G1),(A'1,R1-1,R2-1,R3-4,R4-2,G2),(A'1,R1-1,R2-1,R3-4,R4-2,G3),(A'1,R1-1,R2-1,R3-4,R4-2,G4),(A'1,R1-1,R2-2,R3-1,R4-1,G1),(A'1,R1-1,R2-2,R3-1,R4-1,G2),(A'1,R1-1,R2-2,R3-1,R4-1,G3),(A'1,R1-1,R2-2,R3-1,R4-1,G4),(A'1,R1-1,R2-2,R3-1,R4-2,G1),(A'1,R1-1,R2-2,R3-1,R4-2,G2),(A'1,R1-1,R2-2,R3-1,R4-2,G3),(A'1,R1-1,R2-2,R3-1,R4-2,G4),(A'1,R1-1,R2-2,R3-2,R4-1,G1),(A'1,R1-1,R2-2,R3-2,R4-1,G2),(A'1,R1-1,R2-2,R3-2,R4-1,G3),(A'1,R1-1,R2-2,R3-2,R4-1,G4),(A'1,R1-1,R2-2,R3-2,R4-2,G1),(A'1,R1-1,R2-2,R3-2,R4-2,G2),(A'1,R1-1,R2-2,R3-2,R4-2,G3),(A'1,R1-1,R2-2,R3-2,R4-2,G4),(A'1,R1-1,R2-2,R3-3,R4-1,G1),(A'1,R1-1,R2-2,R3-3,R4-1,G2),(A'1,R1-1,R2-2,R3-3,R4-1,G3),(A'1,R1-1,R2-2,R3-3,R4-1,G4),(A'1,R1-1,R2-2,R3-3,R4-2,G1),(A'1,R1-1,R2-2,R3-3,R4-2,G2),(A'1,R1-1,R2-2,R3-3,R4-2,G3),(A'1,R1-1,R2-2,R3-3,R4-2,G4),(A'1,R1-1,R2-2,R3-4,R4-1,G1),(A'1,R1-1,R2-2,R3-4,R4-1,G2),(A'1,R1-1,R2-2,R3-4,R4-1,G3),(A'1,R1-1,R2-2,R3-4,R4-1,G4),(A'1,R1-1,R2-2,R3-4,R4-2,G1),(A'1,R1-1,R2-2,R3-4,R4-2,G2),(A'1,R1-1,R2-2,R3-4,R4-2,G3),(A'1,R1-1,R2-2,R3-4,R4-2,G4),(A'1,R1-2,R2-1,R3-1,R4-1,G1),(A'1,R1-2,R2-1,R3-1,R4-1,G2),(A'1,R1-2,R2-1,R3-1,R4-1,G3),(A'1,R1-2,R2-1,R3-1,R4-1,G4),(A'1,R1-2,R2-1,R3-1,R4-2,G1),(A'1,R1-2,R2-1,R3-1,R4-2,G2),(A'1,R1-2,R2-1,R3-1,R4-2,G3),(A'1,R1-2,R2-1,R3-1,R4-2,G4),(A'1,R1-2,R2-1,R3-2,R4-1,G1),(A'1,R1-2,R2-1,R3-2,R4-1,G2),(A'1,R1-2,R2-1,R3-2,R4-1,G3),(A'1,R1-2,R2-1,R3-2,R4-1,G4),(A'1,R1-2,R2-1,R3-2,R4-2,G1),(A'1,R1-2,R2-1,R3-2,R4-2,G2),(A'1,R1-2,R2-1,R3-2,R4-2,G3),(A'1,R1-2,R2-1,R3-2,R4-2,G4),(A'1,R1-2,R2-1,R3-3,R4-1,G1),(A'1,R1-2,R2-1,R3-3,R4-1,G2),(A'1,R1-2,R2-1,R3-3,R4-1,G3),(A'1,R1-2,R2-1,R3-3,R4-1,G4),(A'1,R1-2,R2-1,R3-3,R4-2,G1),(A'1,R1-2,R2-1,R3-3,R4-2,G2),(A'1,R1-2,R2-

1,R3-3,R4-2,G3),(A'1,R1-2,R2-1,R3-3,R4-2,G4),(A'1,R1-2,R2-1,R3-4,R4-1,G1),(A'1,R1-2,
R2-1,R3-4,R4-1,G2),(A'1,R1-2,R2-1,R3-4,R4-1,G3),(A'1,R1-2,R2-1,R3-4,R4-1,G4),(A'1,R
1-2,R2-1,R3-4,R4-2,G1),(A'1,R1-2,R2-1,R3-4,R4-2,G2),(A'1,R1-2,R2-1,R3-4,R4-2,G3),(A'
1,R1-2,R2-1,R3-4,R4-2,G4),(A'1,R1-2,R2-2,R3-1,R4-1,G1),(A'1,R1-2,R2-2,R3-1,R4-1,G2),
(A'1,R1-2,R2-2,R3-1,R4-1,G3),(A'1,R1-2,R2-2,R3-1,R4-1,G4),(A'1,R1-2,R2-2,R3-1,R4-2,
G1),(A'1,R1-2,R2-2,R3-1,R4-2,G2),(A'1,R1-2,R2-2,R3-1,R4-2,G3),(A'1,R1-2,R2-2,R3-1,R
4-2,G4),(A'1,R1-2,R2-2,R3-2,R4-1,G1),(A'1,R1-2,R2-2,R3-2,R4-1,G2),(A'1,R1-2,R2-2,R3-
2,R4-1,G3),(A'1,R1-2,R2-2,R3-2,R4-1,G4),(A'1,R1-2,R2-2,R3-2,R4-2,G1),(A'1,R1-2,R2-2,
R3-2,R4-2,G2),(A'1,R1-2,R2-2,R3-2,R4-2,G3),(A'1,R1-2,R2-2,R3-2,R4-2,G4),(A'1,R1-2,R
2-2,R3-3,R4-1,G1),(A'1,R1-2,R2-2,R3-3,R4-1,G2),(A'1,R1-2,R2-2,R3-3,R4-1,G3),(A'1,R1-
2,R2-2,R3-3,R4-1,G4),(A'1,R1-2,R2-2,R3-3,R4-2,G1),(A'1,R1-2,R2-2,R3-3,R4-2,G2),(A'1,
R1-2,R2-2,R3-3,R4-2,G3),(A'1,R1-2,R2-2,R3-3,R4-2,G4),(A'1,R1-2,R2-2,R3-4,R4-1,G1),(
A'1,R1-2,R2-2,R3-4,R4-1,G2),(A'1,R1-2,R2-2,R3-4,R4-1,G3),(A'1,R1-2,R2-2,R3-4,R4-1,G
4),(A'1,R1-2,R2-2,R3-4,R4-2,G1),(A'1,R1-2,R2-2,R3-4,R4-2,G2),(A'1,R1-2,R2-2,R3-4,R4-
2,G3),(A'1,R1-2,R2-2,R3-4,R4-2,G4),

(A'2,R1-1,R2-1,R3-1,R4-1,G1),(A'2,R1-1,R2-1,R3-1,R4-1,G2),(A'2,R1-1,R2-1,R3-1,R4-1,
G3),(A'2,R1-1,R2-1,R3-1,R4-1,G4),(A'2,R1-1,R2-1,R3-1,R4-2,G1),(A'2,R1-1,R2-1,R3-1,R
4-2,G2),(A'2,R1-1,R2-1,R3-1,R4-2,G3),(A'2,R1-1,R2-1,R3-1,R4-2,G4),(A'2,R1-1,R2-1,R3-
2,R4-1,G1),(A'2,R1-1,R2-1,R3-2,R4-1,G2),(A'2,R1-1,R2-1,R3-2,R4-1,G3),(A'2,R1-1,R2-1,
R3-2,R4-1,G4),(A'2,R1-1,R2-1,R3-2,R4-2,G1),(A'2,R1-1,R2-1,R3-2,R4-2,G2),(A'2,R1-1,R
2-1,R3-2,R4-2,G3),(A'2,R1-1,R2-1,R3-2,R4-2,G4),(A'2,R1-1,R2-1,R3-3,R4-1,G1),(A'2,R1-
1,R2-1,R3-3,R4-1,G2),(A'2,R1-1,R2-1,R3-3,R4-1,G3),(A'2,R1-1,R2-1,R3-3,R4-1,G4),(A'2,
R1-1,R2-1,R3-3,R4-2,G1),(A'2,R1-1,R2-1,R3-3,R4-2,G2),(A'2,R1-1,R2-1,R3-3,R4-2,G3),(
A'2,R1-1,R2-1,R3-3,R4-2,G4),(A'2,R1-1,R2-1,R3-4,R4-1,G1),(A'2,R1-1,R2-1,R3-4,R4-1,G
2),(A'2,R1-1,R2-1,R3-4,R4-1,G3),(A'2,R1-1,R2-1,R3-4,R4-1,G4),(A'2,R1-1,R2-1,R3-4,R4-
2,G1),(A'2,R1-1,R2-1,R3-4,R4-2,G2),(A'2,R1-1,R2-1,R3-4,R4-2,G3),(A'2,R1-1,R2-1,R3-4,
R4-2,G4),(A'2,R1-1,R2-2,R3-1,R4-1,G1),(A'2,R1-1,R2-2,R3-1,R4-1,G2),(A'2,R1-1,R2-2,R

3-1,R4-1,G3),(A'2,R1-1,R2-2,R3-1,R4-1,G4),(A'2,R1-1,R2-2,R3-1,R4-2,G1),(A'2,R1-1,R2-2,R3-1,R4-2,G2),(A'2,R1-1,R2-2,R3-1,R4-2,G3),(A'2,R1-1,R2-2,R3-1,R4-2,G4),(A'2,R1-1,R2-2,R3-2,R4-1,G1),(A'2,R1-1,R2-2,R3-2,R4-1,G2),(A'2,R1-1,R2-2,R3-2,R4-1,G3),(A'2,R1-1,R2-2,R3-2,R4-1,G4),(A'2,R1-1,R2-2,R3-2,R4-2,G1),(A'2,R1-1,R2-2,R3-2,R4-2,G2),(A'2,R1-1,R2-2,R3-2,R4-2,G3),(A'2,R1-1,R2-2,R3-2,R4-2,G4),(A'2,R1-1,R2-2,R3-3,R4-1,G1),(A'2,R1-1,R2-2,R3-3,R4-1,G2),(A'2,R1-1,R2-2,R3-3,R4-1,G3),(A'2,R1-1,R2-2,R3-3,R4-1,G4),(A'2,R1-1,R2-2,R3-3,R4-2,G1),(A'2,R1-1,R2-2,R3-3,R4-2,G2),(A'2,R1-1,R2-2,R3-3,R4-2,G3),(A'2,R1-1,R2-2,R3-3,R4-2,G4),(A'2,R1-1,R2-2,R3-4,R4-1,G1),(A'2,R1-1,R2-2,R3-4,R4-1,G2),(A'2,R1-1,R2-2,R3-4,R4-1,G3),(A'2,R1-1,R2-2,R3-4,R4-1,G4),(A'2,R1-1,R2-2,R3-4,R4-2,G1),(A'2,R1-1,R2-2,R3-4,R4-2,G2),(A'2,R1-1,R2-2,R3-4,R4-2,G3),(A'2,R1-1,R2-2,R3-4,R4-2,G4),(A'2,R1-2,R2-1,R3-1,R4-1,G1),(A'2,R1-2,R2-1,R3-1,R4-1,G2),(A'2,R1-2,R2-1,R3-1,R4-1,G3),(A'2,R1-2,R2-1,R3-1,R4-1,G4),(A'2,R1-2,R2-1,R3-1,R4-2,G1),(A'2,R1-2,R2-1,R3-1,R4-2,G2),(A'2,R1-2,R2-1,R3-1,R4-2,G3),(A'2,R1-2,R2-1,R3-1,R4-2,G4),(A'2,R1-2,R2-1,R3-2,R4-1,G1),(A'2,R1-2,R2-1,R3-2,R4-1,G2),(A'2,R1-2,R2-1,R3-2,R4-1,G3),(A'2,R1-2,R2-1,R3-2,R4-1,G4),(A'2,R1-2,R2-1,R3-2,R4-2,G1),(A'2,R1-2,R2-1,R3-2,R4-2,G2),(A'2,R1-2,R2-1,R3-2,R4-2,G3),(A'2,R1-2,R2-1,R3-2,R4-2,G4),(A'2,R1-2,R2-1,R3-3,R4-1,G1),(A'2,R1-2,R2-1,R3-3,R4-1,G2),(A'2,R1-2,R2-1,R3-3,R4-1,G3),(A'2,R1-2,R2-1,R3-3,R4-1,G4),(A'2,R1-2,R2-1,R3-3,R4-2,G1),(A'2,R1-2,R2-1,R3-3,R4-2,G2),(A'2,R1-2,R2-1,R3-3,R4-2,G3),(A'2,R1-2,R2-1,R3-3,R4-2,G4),(A'2,R1-2,R2-1,R3-4,R4-1,G1),(A'2,R1-2,R2-1,R3-4,R4-1,G2),(A'2,R1-2,R2-1,R3-4,R4-1,G3),(A'2,R1-2,R2-1,R3-4,R4-1,G4),(A'2,R1-2,R2-1,R3-4,R4-2,G1),(A'2,R1-2,R2-1,R3-4,R4-2,G2),(A'2,R1-2,R2-1,R3-4,R4-2,G3),(A'2,R1-2,R2-1,R3-4,R4-2,G4),(A'2,R1-2,R2-2,R3-1,R4-1,G1),(A'2,R1-2,R2-2,R3-1,R4-1,G2),(A'2,R1-2,R2-2,R3-1,R4-1,G3),(A'2,R1-2,R2-2,R3-1,R4-1,G4),(A'2,R1-2,R2-2,R3-1,R4-2,G1),(A'2,R1-2,R2-2,R3-1,R4-2,G2),(A'2,R1-2,R2-2,R3-1,R4-2,G3),(A'2,R1-2,R2-2,R3-1,R4-2,G4),(A'2,R1-2,R2-2,R3-2,R4-1,G1),(A'2,R1-2,R2-2,R3-2,R4-1,G2),(A'2,R1-2,R2-2,R3-2,R4-1,G3),(A'2,R1-2,R2-2,R3-2,R4-1,G4),(A'2,R1-2,R2-2,R3-2,R4-2,G1),(A'2,R1-2,R2-2,R3-2,R4-2,G2),(A'2,R1-2,R2-2,R3-2,R4-2,G3),(A'2,R1-2,R2-2,R3-2,R4-2,G4),(A'2,R1-2,R2-2,R3-3,R4-1,G1),(A'2,R1-2,R2-2,R3-3,R4-1,G2),(A'2,R1-2,R2-2,R3-3,R4-1,G3),(A'2,R1-

2,R2-2,R3-3,R4-1,G4),(A'2,R1-2,R2-2,R3-3,R4-2,G1),(A'2,R1-2,R2-2,R3-3,R4-2,G2),(A'2,R1-2,R2-2,R3-3,R4-2,G3),(A'2,R1-2,R2-2,R3-3,R4-2,G4),(A'2,R1-2,R2-2,R3-4,R4-1,G1),(A'2,R1-2,R2-2,R3-4,R4-1,G2),(A'2,R1-2,R2-2,R3-4,R4-1,G3),(A'2,R1-2,R2-2,R3-4,R4-1,G4),(A'2,R1-2,R2-2,R3-4,R4-2,G1),(A'2,R1-2,R2-2,R3-4,R4-2,G2),(A'2,R1-2,R2-2,R3-4,R4-2,G3),(A'2,R1-2,R2-2,R3-4,R4-2,G4),

(A'3,R1-1,R2-1,R3-1,R4-1,G1),(A'3,R1-1,R2-1,R3-1,R4-1,G2),(A'3,R1-1,R2-1,R3-1,R4-1,G3),(A'3,R1-1,R2-1,R3-1,R4-1,G4),(A'3,R1-1,R2-1,R3-1,R4-2,G1),(A'3,R1-1,R2-1,R3-1,R4-2,G2),(A'3,R1-1,R2-1,R3-1,R4-2,G3),(A'3,R1-1,R2-1,R3-1,R4-2,G4),(A'3,R1-1,R2-1,R3-2,R4-1,G1),(A'3,R1-1,R2-1,R3-2,R4-1,G2),(A'3,R1-1,R2-1,R3-2,R4-1,G3),(A'3,R1-1,R2-1,R3-2,R4-1,G4),(A'3,R1-1,R2-1,R3-2,R4-2,G1),(A'3,R1-1,R2-1,R3-2,R4-2,G2),(A'3,R1-1,R2-1,R3-2,R4-2,G3),(A'3,R1-1,R2-1,R3-2,R4-2,G4),(A'3,R1-1,R2-1,R3-3,R4-1,G1),(A'3,R1-1,R2-1,R3-3,R4-1,G2),(A'3,R1-1,R2-1,R3-3,R4-1,G3),(A'3,R1-1,R2-1,R3-3,R4-1,G4),(A'3,R1-1,R2-1,R3-3,R4-2,G1),(A'3,R1-1,R2-1,R3-3,R4-2,G2),(A'3,R1-1,R2-1,R3-3,R4-2,G3),(A'3,R1-1,R2-1,R3-3,R4-2,G4),(A'3,R1-1,R2-1,R3-4,R4-1,G1),(A'3,R1-1,R2-1,R3-4,R4-1,G2),(A'3,R1-1,R2-1,R3-4,R4-1,G3),(A'3,R1-1,R2-1,R3-4,R4-1,G4),(A'3,R1-1,R2-1,R3-4,R4-2,G1),(A'3,R1-1,R2-1,R3-4,R4-2,G2),(A'3,R1-1,R2-1,R3-4,R4-2,G3),(A'3,R1-1,R2-1,R3-4,R4-2,G4),(A'3,R1-1,R2-2,R3-1,R4-1,G1),(A'3,R1-1,R2-2,R3-1,R4-1,G2),(A'3,R1-1,R2-2,R3-1,R4-1,G3),(A'3,R1-1,R2-2,R3-1,R4-1,G4),(A'3,R1-1,R2-2,R3-1,R4-2,G1),(A'3,R1-1,R2-2,R3-1,R4-2,G2),(A'3,R1-1,R2-2,R3-1,R4-2,G3),(A'3,R1-1,R2-2,R3-1,R4-2,G4),(A'3,R1-1,R2-2,R3-2,R4-1,G1),(A'3,R1-1,R2-2,R3-2,R4-1,G2),(A'3,R1-1,R2-2,R3-2,R4-1,G3),(A'3,R1-1,R2-2,R3-2,R4-1,G4),(A'3,R1-1,R2-2,R3-2,R4-2,G1),(A'3,R1-1,R2-2,R3-2,R4-2,G2),(A'3,R1-1,R2-2,R3-2,R4-2,G3),(A'3,R1-1,R2-2,R3-2,R4-2,G4),(A'3,R1-1,R2-2,R3-3,R4-1,G1),(A'3,R1-1,R2-2,R3-3,R4-1,G2),(A'3,R1-1,R2-2,R3-3,R4-1,G3),(A'3,R1-1,R2-2,R3-3,R4-1,G4),(A'3,R1-1,R2-2,R3-3,R4-2,G1),(A'3,R1-1,R2-2,R3-3,R4-2,G2),(A'3,R1-1,R2-2,R3-3,R4-2,G3),(A'3,R1-1,R2-2,R3-3,R4-2,G4),(A'3,R1-1,R2-2,R3-4,R4-1,G1),(A'3,R1-1,R2-2,R3-4,R4-1,G2),(A'3,R1-1,R2-2,R3-4,R4-1,G3),(A'3,R1-1,R2-2,R3-4,R4-1,G4),(A'3,R1-1,R2-2,R3-4,R4-2,G1),(A'3,R1-1,R2-2,R3-4,R4-2,G2),(A'3,R1-1,R2-2,R3-4,R4-2,G3),(A'3,R1-1,R

2-2,R3-4,R4-2,G4),(A'3,R1-2,R2-1,R3-1,R4-1,G1),(A'3,R1-2,R2-1,R3-1,R4-1,G2),(A'3,R1-2,R2-1,R3-1,R4-1,G3),(A'3,R1-2,R2-1,R3-1,R4-1,G4),(A'3,R1-2,R2-1,R3-1,R4-2,G1),(A'3,R1-2,R2-1,R3-1,R4-2,G2),(A'3,R1-2,R2-1,R3-1,R4-2,G3),(A'3,R1-2,R2-1,R3-1,R4-2,G4),(A'3,R1-2,R2-1,R3-2,R4-1,G1),(A'3,R1-2,R2-1,R3-2,R4-1,G2),(A'3,R1-2,R2-1,R3-2,R4-1,G3),(A'3,R1-2,R2-1,R3-2,R4-1,G4),(A'3,R1-2,R2-1,R3-2,R4-2,G1),(A'3,R1-2,R2-1,R3-2,R4-2,G2),(A'3,R1-2,R2-1,R3-2,R4-2,G3),(A'3,R1-2,R2-1,R3-2,R4-2,G4),(A'3,R1-2,R2-1,R3-3,R4-1,G1),(A'3,R1-2,R2-1,R3-3,R4-1,G2),(A'3,R1-2,R2-1,R3-3,R4-1,G3),(A'3,R1-2,R2-1,R3-3,R4-1,G4),(A'3,R1-2,R2-1,R3-3,R4-2,G1),(A'3,R1-2,R2-1,R3-3,R4-2,G2),(A'3,R1-2,R2-1,R3-3,R4-2,G3),(A'3,R1-2,R2-1,R3-3,R4-2,G4),(A'3,R1-2,R2-1,R3-4,R4-1,G1),(A'3,R1-2,R2-1,R3-4,R4-1,G2),(A'3,R1-2,R2-1,R3-4,R4-1,G3),(A'3,R1-2,R2-1,R3-4,R4-1,G4),(A'3,R1-2,R2-1,R3-4,R4-2,G1),(A'3,R1-2,R2-1,R3-4,R4-2,G2),(A'3,R1-2,R2-1,R3-4,R4-2,G3),(A'3,R1-2,R2-1,R3-4,R4-2,G4),(A'3,R1-2,R2-2,R3-1,R4-1,G1),(A'3,R1-2,R2-2,R3-1,R4-1,G2),(A'3,R1-2,R2-2,R3-1,R4-1,G3),(A'3,R1-2,R2-2,R3-1,R4-1,G4),(A'3,R1-2,R2-2,R3-1,R4-2,G1),(A'3,R1-2,R2-2,R3-1,R4-2,G2),(A'3,R1-2,R2-2,R3-1,R4-2,G3),(A'3,R1-2,R2-2,R3-1,R4-2,G4),(A'3,R1-2,R2-2,R3-2,R4-1,G1),(A'3,R1-2,R2-2,R3-2,R4-1,G2),(A'3,R1-2,R2-2,R3-2,R4-1,G3),(A'3,R1-2,R2-2,R3-2,R4-1,G4),(A'3,R1-2,R2-2,R3-2,R4-2,G1),(A'3,R1-2,R2-2,R3-2,R4-2,G2),(A'3,R1-2,R2-2,R3-2,R4-2,G3),(A'3,R1-2,R2-2,R3-2,R4-2,G4),(A'3,R1-2,R2-2,R3-3,R4-1,G1),(A'3,R1-2,R2-2,R3-3,R4-1,G2),(A'3,R1-2,R2-2,R3-3,R4-1,G3),(A'3,R1-2,R2-2,R3-3,R4-1,G4),(A'3,R1-2,R2-2,R3-3,R4-2,G1),(A'3,R1-2,R2-2,R3-3,R4-2,G2),(A'3,R1-2,R2-2,R3-3,R4-2,G3),(A'3,R1-2,R2-2,R3-3,R4-2,G4),(A'3,R1-2,R2-2,R3-4,R4-1,G1),(A'3,R1-2,R2-2,R3-4,R4-1,G2),(A'3,R1-2,R2-2,R3-4,R4-1,G3),(A'3,R1-2,R2-2,R3-4,R4-1,G4),(A'3,R1-2,R2-2,R3-4,R4-2,G1),(A'3,R1-2,R2-2,R3-4,R4-2,G2),(A'3,R1-2,R2-2,R3-4,R4-2,G3),(A'3,R1-2,R2-2,R3-4,R4-2,G4),

(A'4,R1-1,R2-1,R3-1,R4-1,G1),(A'4,R1-1,R2-1,R3-1,R4-1,G2),(A'4,R1-1,R2-1,R3-1,R4-1,G3),(A'4,R1-1,R2-1,R3-1,R4-1,G4),(A'4,R1-1,R2-1,R3-1,R4-2,G1),(A'4,R1-1,R2-1,R3-1,R4-2,G2),(A'4,R1-1,R2-1,R3-1,R4-2,G3),(A'4,R1-1,R2-1,R3-1,R4-2,G4),(A'4,R1-1,R2-1,R3-2,R4-1,G1),(A'4,R1-1,R2-1,R3-2,R4-1,G2),(A'4,R1-1,R2-1,R3-2,R4-1,G3),(A'4,R1-1,R2-1,

R3-2,R4-1,G4),(A'4,R1-1,R2-1,R3-2,R4-2,G1),(A'4,R1-1,R2-1,R3-2,R4-2,G2),(A'4,R1-1,R2-1,R3-2,R4-2,G3),(A'4,R1-1,R2-1,R3-2,R4-2,G4),(A'4,R1-1,R2-1,R3-3,R4-1,G1),(A'4,R1-1,R2-1,R3-3,R4-1,G2),(A'4,R1-1,R2-1,R3-3,R4-1,G3),(A'4,R1-1,R2-1,R3-3,R4-1,G4),(A'4,R1-1,R2-1,R3-3,R4-2,G1),(A'4,R1-1,R2-1,R3-3,R4-2,G2),(A'4,R1-1,R2-1,R3-3,R4-2,G3),(A'4,R1-1,R2-1,R3-3,R4-2,G4),(A'4,R1-1,R2-1,R3-4,R4-1,G1),(A'4,R1-1,R2-1,R3-4,R4-1,G2),(A'4,R1-1,R2-1,R3-4,R4-1,G3),(A'4,R1-1,R2-1,R3-4,R4-1,G4),(A'4,R1-1,R2-1,R3-4,R4-2,G1),(A'4,R1-1,R2-1,R3-4,R4-2,G2),(A'4,R1-1,R2-1,R3-4,R4-2,G3),(A'4,R1-1,R2-1,R3-4,R4-2,G4),(A'4,R1-1,R2-2,R3-1,R4-1,G1),(A'4,R1-1,R2-2,R3-1,R4-1,G2),(A'4,R1-1,R2-2,R3-1,R4-1,G3),(A'4,R1-1,R2-2,R3-1,R4-1,G4),(A'4,R1-1,R2-2,R3-1,R4-2,G1),(A'4,R1-1,R2-2,R3-1,R4-2,G2),(A'4,R1-1,R2-2,R3-1,R4-2,G3),(A'4,R1-1,R2-2,R3-1,R4-2,G4),(A'4,R1-1,R2-2,R3-2,R4-1,G1),(A'4,R1-1,R2-2,R3-2,R4-1,G2),(A'4,R1-1,R2-2,R3-2,R4-1,G3),(A'4,R1-1,R2-2,R3-2,R4-1,G4),(A'4,R1-1,R2-2,R3-2,R4-2,G1),(A'4,R1-1,R2-2,R3-2,R4-2,G2),(A'4,R1-1,R2-2,R3-2,R4-2,G3),(A'4,R1-1,R2-2,R3-2,R4-2,G4),(A'4,R1-1,R2-2,R3-3,R4-1,G1),(A'4,R1-1,R2-2,R3-3,R4-1,G2),(A'4,R1-1,R2-2,R3-3,R4-1,G3),(A'4,R1-1,R2-2,R3-3,R4-1,G4),(A'4,R1-1,R2-2,R3-3,R4-2,G1),(A'4,R1-1,R2-2,R3-3,R4-2,G2),(A'4,R1-1,R2-2,R3-3,R4-2,G3),(A'4,R1-1,R2-2,R3-3,R4-2,G4),(A'4,R1-1,R2-2,R3-4,R4-1,G1),(A'4,R1-1,R2-2,R3-4,R4-1,G2),(A'4,R1-1,R2-2,R3-4,R4-1,G3),(A'4,R1-1,R2-2,R3-4,R4-1,G4),(A'4,R1-1,R2-2,R3-4,R4-2,G1),(A'4,R1-1,R2-2,R3-4,R4-2,G2),(A'4,R1-1,R2-2,R3-4,R4-2,G3),(A'4,R1-1,R2-2,R3-4,R4-2,G4),(A'4,R1-2,R2-1,R3-1,R4-1,G1),(A'4,R1-2,R2-1,R3-1,R4-1,G2),(A'4,R1-2,R2-1,R3-1,R4-1,G3),(A'4,R1-2,R2-1,R3-1,R4-1,G4),(A'4,R1-2,R2-1,R3-1,R4-2,G1),(A'4,R1-2,R2-1,R3-1,R4-2,G2),(A'4,R1-2,R2-1,R3-1,R4-2,G3),(A'4,R1-2,R2-1,R3-1,R4-2,G4),(A'4,R1-2,R2-1,R3-2,R4-1,G1),(A'4,R1-2,R2-1,R3-2,R4-1,G2),(A'4,R1-2,R2-1,R3-2,R4-1,G3),(A'4,R1-2,R2-1,R3-2,R4-1,G4),(A'4,R1-2,R2-1,R3-2,R4-2,G1),(A'4,R1-2,R2-1,R3-2,R4-2,G2),(A'4,R1-2,R2-1,R3-2,R4-2,G3),(A'4,R1-2,R2-1,R3-2,R4-2,G4),(A'4,R1-2,R2-1,R3-3,R4-1,G1),(A'4,R1-2,R2-1,R3-3,R4-1,G2),(A'4,R1-2,R2-1,R3-3,R4-1,G3),(A'4,R1-2,R2-1,R3-3,R4-1,G4),(A'4,R1-2,R2-1,R3-3,R4-2,G1),(A'4,R1-2,R2-1,R3-3,R4-2,G2),(A'4,R1-2,R2-1,R3-3,R4-2,G3),(A'4,R1-2,R2-1,R3-3,R4-2,G4),(A'4,R1-2,R2-1,R3-4,R4-1,G1),(A'4,R1-2,R2-1,R3-4,R4-1,G2),(A'4,R1-2,R2-1,R3-4,R4-1,G3),(A'4,R1-2,R2-1,R3-4,R4-1,G4),(A'4,R

1-2,R2-1,R3-4,R4-2,G1),(A'4,R1-2,R2-1,R3-4,R4-2,G2),(A'4,R1-2,R2-1,R3-4,R4-2,G3),(A'4,R1-2,R2-1,R3-4,R4-2,G4),(A'4,R1-2,R2-2,R3-1,R4-1,G1),(A'4,R1-2,R2-2,R3-1,R4-1,G2),(A'4,R1-2,R2-2,R3-1,R4-1,G3),(A'4,R1-2,R2-2,R3-1,R4-1,G4),(A'4,R1-2,R2-2,R3-1,R4-2,G1),(A'4,R1-2,R2-2,R3-1,R4-2,G2),(A'4,R1-2,R2-2,R3-1,R4-2,G3),(A'4,R1-2,R2-2,R3-1,R4-2,G4),(A'4,R1-2,R2-2,R3-2,R4-1,G1),(A'4,R1-2,R2-2,R3-2,R4-1,G2),(A'4,R1-2,R2-2,R3-2,R4-1,G3),(A'4,R1-2,R2-2,R3-2,R4-1,G4),(A'4,R1-2,R2-2,R3-2,R4-2,G1),(A'4,R1-2,R2-2,R3-2,R4-2,G2),(A'4,R1-2,R2-2,R3-2,R4-2,G3),(A'4,R1-2,R2-2,R3-2,R4-2,G4),(A'4,R1-2,R2-2,R3-3,R4-1,G1),(A'4,R1-2,R2-2,R3-3,R4-1,G2),(A'4,R1-2,R2-2,R3-3,R4-1,G3),(A'4,R1-2,R2-2,R3-3,R4-1,G4),(A'4,R1-2,R2-2,R3-3,R4-2,G1),(A'4,R1-2,R2-2,R3-3,R4-2,G2),(A'4,R1-2,R2-2,R3-3,R4-2,G3),(A'4,R1-2,R2-2,R3-3,R4-2,G4),(A'4,R1-2,R2-2,R3-4,R4-1,G1),(A'4,R1-2,R2-2,R3-4,R4-1,G2),(A'4,R1-2,R2-2,R3-4,R4-1,G3),(A'4,R1-2,R2-2,R3-4,R4-1,G4),(A'4,R1-2,R2-2,R3-4,R4-2,G1),(A'4,R1-2,R2-2,R3-4,R4-2,G2),(A'4,R1-2,R2-2,R3-4,R4-2,G3) or (A'4,R1-2,R2-2,R3-4,R4-2,G4).

Compounds of the present invention are useful for treating diseases induced by production, secretion or deposition of amyloid β protein, and effective for the treatment and/or prophylaxis, or improvement of conditions for Alzheimer's dementia (Alzheimer's disease, senile dementia of Alzheimer type etc.), Down's disease, disturbance of memory, prion disease (Creutzfeldt-Jakob disease etc.), mild cognitive impairment(MCI), Dutch-type hereditary cerebral hemorrhage with amyloidosis, cerebral amyloid angiopathy, other degenerated dementia, vascular degenerated mixed dementia, dementia associated with Parkinson's disease, dementia associated with progressive supranuclear paralysis, dementia associated with corticobasal degeneration, diffuse Lewy Bodies Alzheimer's disease, age-related macular degeneration, Parkinson's disease, or amyloid angiopathy etc.

Since compounds of the present invention have several efficacies such as having a potent inhibitory activity against BACE-1 having a high selectivity against other enzymes etc., they can be a drug with less side effects. Moreover they can be a drug having a wide margin of safety by choosing an optically active isomer of appropriate stereochemistry. Further they have a lot of merits such as good metabolic stability, high solubility, high

absorbability of oral administration, high bioavailability, good clearance and high transitivity to brain, long half-life, high ratio of non protein binding, lower inhibition of hERG channel and CYP, and/or negative result of Ames Test, and, therefore, they can be superior drugs.

A compound of the present invention may be administrated together with other agent(e.g., other agent for treating Alzheimer's disease such as acetylcholine esterase etc.). The compound can be given in combination with an antidementia drug such as donepezil hydrochloride, tacrine, galantamine, rivastigmine, zanapezil, memantine or vinpocetine, for example.

A compound of the present invention may be orally administrated as powder, granule, tablet, capsule, pill or liquid formulation, or parentally administrated as injection, suppository, formulation of transdermal absorption or inhalation. Also, an effective amount of the compound may be formulated together with medicinal additives suitable for the formulation such as an excipient, binder, moistening agent, disintegrant and/or lubricant etc.

Dose of a compounds of the present invention depends on condition of diseases, route of administration, age and body weight of a patient, but in the case of oral administration to an adult, the dose range is usually 0.1 μg to 1 g/day, preferably 0.01 to 200 mg/day and in the case of parenteral administration the dose range is usually 1 μg to 10 g/day, preferably 0.1 to 2 g/day.

EXAMPLES

The present invention is illustrated in details by examples and test examples but the present invention is not limited to these examples.

In EXAMPLES, each abbreviation has the following meaning:

Me: methyl

Et: ethyl

iPr, Pri isopropyl

tBu: t-butyl

Ph: phenyl

Bn: benzyl

Boc: tert-butoxycarbonyl

TFA: trifluroacetic acid


THF: tetrahydrofuran

DMT-MM: 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride

n-hydrate

DMF: N,N-dimethylformamide

Reference Example 1



The 1st step: Compound (1) (101.5 g) was cooled to -18°C and conc. sulfuric acid

(400 ml) was added dropwise in 65 minutes while the inner temperature was kept at -15°C or below. Separately fuming nitric acid (60 ml) was added to conc. sulfuric acid (180 ml) chilled to 4°C in 45 minutes while the temperature was kept at 10°C or below and the resulted mixed acid was added dropwise to the solution of (1) prepared before in an hour while the temperature was kept at -30°C or below. The mixture was stirred at -20°C or below for 1.5 hours, poured into 2.5 kg of ice-water and stirred for an hour. The precipitated crystals were filtered to give compound (2) (121.5 g).

^{1}H-NMR($CDCl_3$): 2.71(3H, d, J=5.1Hz), 7.35(1H, dd, J=9.3, 9.0Hz), 8.41(1H , ddd, J=9.0, 3.9, 3.0Hz), 8.78(1H, dd, J=6.3, 3.0Hz).

The 2nd step: Compound (2) (20 g) was dissolved in ethanol (400 ml), Pd-C (10% dry)(2.0 g) was added thereto and the mixture was stirred in a hydrogen atmosphere at room temperature for 2 hours. Then Pd-C (10% dry)(1.0 g) was added and the mixture was stirred in a hydrogen atmosphere at room temperature for 1.5 hours, and further Pd-C (10% dry)(1.0 g) was added and the mixture was stirred in a hydrogen atmosphere at room temperature for 15 minutes. Pd-C was filtered off, the solvent was evaporated under reduced pressure and the residue of compound (3) (15.9 g) was obtained.

^{1}H-NMR(DMSO-d_6) : 2.50(3H, d, J=4.8Hz), 5.21(2H, brs, 1H), 6.78(1H , ddd, J=8.7, 4.2, 3.0Hz), 6.94(1H, dd, J=6.3, 3.0Hz), 6.99(1H, dd, J=11.4, 8.7Hz).

The 3rd step: Compound (3) (15.8 g) was dissolved in THF (79 ml), anhydrous trifluoroacetic acid (16.1 ml) and triethylamine (20.2 ml) were added under ice cooling and the mixture was stirred for 20 minutes. After the addition of water (30 ml), it was stirred under ice cooling for 20 minutes and the precipitated crystals were filtered. The filtrate was extracted with ethyl acetate (80 ml) and 50 ml, the organic layer was washed with water (60 ml), saturated brine. The crystals filtered previously were dissolved in the organic layer and dried over sodium sulfate. Sodium sulfate was filtered, the filtrate was concentrated under reduced pressure and the residue was dissolved in ethyl acetate under warming. After the addition of hexane (50 ml) and stirring under ice cooling for 20 minutes, the precipitated crystals were filtered. The mother liquid was concentrated again under reduced pressure,

crystallized by the addition of ethyl acetate (8 ml) and hexane (12 ml) and compound (4) (totally 20.4 g) was obtained.

^{1}H-NMR($CDCl_3$) : 2.70(3H, d, J=5.1Hz), 7.24(1H, dd, J=10.5, 9.3Hz), 8.00(1H, dd, J=6.2, 2.9Hz), 8.21(1H, m), 8.78(1H, brs).

The 4th step: 1.6 M Vinyl magnesium chloride-THF solution (122 ml) was dissolved in THF (161 ml), cooled to -40°C in a nitrogen atmosphere and a THF (81 ml) solution of compound (4) (16.1 g) was added dropwise thereto. The reaction solution was stirred at -40°C for 20 minutes, 1.6 M vinyl magnesium chloride-THF solution (20 ml) was further added and the mixture was stirred at -40°C for 15 minutes. The reaction solution was poured into a mixture of chilled ethyl acetate (480 ml), a saturated aqueous solution of ammonium chloride (80 ml) and water (80 ml) with stirring, and the organic layer was separated. The aqueous layer was further extracted with ethyl acetate (200 ml), the organic layers were combined, washed with water (80 ml), and saturated brine successively, and dried over sodium sulfate. Sodium sulfate was filtered, the filtrate was concentrated under reduced pressure and the residue of compound (5) (22.4 g) was obtained.

^{1}H-NMR($CDCl_3$): 1.74(3H, d, J=1.2Hz), 5.16(1H, dd, J=10.5, 0.9Hz), 5.27(1H, d, J=17.3, Hz), 6.26(1H, ddd, J=17.3, 10.5, 1.7Hz), 7.07(1H, dd, J=11.1, 9.6Hz), 7.64-7.69(2H, m), 7.94(1H, brs).

The 5th step: The residue of compound (5) (22.3 g) and thiourea (5.17 g) were dissolved in acetic acid (112 ml), 1 M HCl-ethyl acetate (97 ml) was added thereto and the mixture was stirred at 40°C for 18 hours. The solvent was evaporated under reduced pressure, toluene (150 ml) was added and again concentrated under reduced pressure. After repeating the same procedure, crystals were precipitated. Ethyl acetate (100 ml) was added to the crystalline residue, the mixture was stirred under ice cooling for an hour and the crystals were filtered to give compound (6) (15.1 g).

^{1}H-NMR(DMSO-d_6): 2.08(3H, s), 4.10(2H, d, J=7.8Hz), 5.72(1H, t, J=7.8Hz), 7.23-7.32(1H, m), 7.60-7.69(2H, m), 9.25(3H, brs), 11.39(1H, brs).

The 6th step: Compound (6) (10.0 g) was dissolved in THF (50 ml), conc. sulfuric


acid (5.74 ml) was added thereto and stirred at 60°C for 2 hours. After evaporation of TFA under reduced pressure, ice-water (100 ml) was added. The mixture was stirred under ice-cooling for an hour, and the precipitated crystals were filtered to give compound (7) (11.2 g).

^{1}H-NMR($CDCl_3$): 1.72(3H, s), 2.02-2.18(1H, m), 2.54-2.76(2H, m), 3.14-3.28(1H, m), 7.37(1H, dd, J=11.9, 8.8Hz), 7.62(1H, dd, J=7.5, 3.0Hz), 7.80(1H, ddd, J=8.8, 3.9, 3.0Hz), 8.77(1H, brs), 9.38(1H, brs), 10.66(1H, brs), 11.50(1H, brs).

The 7th step: MeOH (28 ml), THF (35 ml) and 5 N NaOH (10.9 ml) were added to compound (7) (7.00 g) and stirred at 50°C for 4 hours. Toluene (50 ml) was added and extracted, and the aqueous layer was further extracted with toluene (50 ml) and ethyl acetate (60 ml). All the organic layers were combined, washed with water and saturated brine, and dried over sodium sulfate. The solvent was concentrated under reduced pressure, the resulted crystalline residue was washed with hexane (20 ml) to give compound (8) (3.45 g).

^{1}H-NMR($CDCl_3$): 1.60(3H, d, J=1.5Hz), 1.76-1.87(1H, m), 2.44-2.54(1H, m), 2.66-2.76(1H, m), 2.86-2.94(1H, m), 6.50(1H, ddd, J=8.7, 3.6, 3.0Hz), 6.66(1H, dd, J=7.1, 3.0Hz), 6.81(1H, dd, J=12.0, 8.7Hz).

Reference Example 2



The 1st step: Compound (3)(15.6 g) was dissolved in ethyl acetate(78 ml), acetic anhydride(10.6 ml) and pyridine(9.07 ml) were added under ice cooling, and the mixture was stirred for 15 minutes. Ethyl acetate (100 ml) and water (50 ml) were added, extracted, and

the aqueous layer was extracted with ethyl acetate(50 ml). The organic layers were combined, washed with 2 M HCl (50 ml), a saturated solution of sodium bicarbonate(50 ml) and saturated brine, dried over sodium sulfate. Sodium sulfate was filtered, the filtrate was concentrated under reduced pressure, and ethyl acetate (50 ml) and hexane(50 ml) were added to the residue. The mixture was stirred under ice cooling for 30 minutes and the precipitated crystals were filtered to give compound (9)(total 14.9 g).

^{1}H-NMR($CDCl_3$) : 2.20(3H, s), 2.66(3H, d, J=5.1Hz), 7.13(1H, dd, J=10.5, 9.0Hz), 7.70(1H, dd, J=6.3, 3.0Hz), 7.79(1H, brs), 8.11(1H, ddd, J=9.0, 4.1, 3.0Hz).

The 2nd step: Compound (9)(10.0 g) was dissolved in THF(50 ml), cooled in ice and sodium hydride(2.25 g) was added in a nitrogen atmosphere. After stirring for 15 minutes, the resulted mixture was added dropwise to a solution of 1.6 M vinyl magnesium chloride (86 ml)/THF(70 ml) cooled to -40°C. After stirring at -40°C for 15 minutes and then 0°C for 20 minutes, a saturated aqueous solution of ammonium chloride (50 ml)/water (50 ml) was chilled and added. The layers were separated and the aqueous layer was extracted with ethyl acetate(100 ml). Organic layers were combined, washed with water and saturated brine, dried over sodium sulfate. Sodium sulfate was filtered, the filtrate was concentrated under reduced pressure to give a residue of compound (10)(13.7 g).

^{1}H-NMR($CDCl_3$): 1.69(3H, s), 2.16(3H, s), 5.12(1H, d, J=10.5Hz), 5.24(1H, d, J=17.4Hz), 6.26(1H, ddd, J=17.4, 10.5, 1.5Hz), 6.98(1H, dd, J=11.1, 8.7Hz), 7.33(1H, brs), 7.50-7.59(2H, m).

The 3rd step: The residue of compound (10)(6.56 g) and thiourea(1.88 g) were dissolved in acetic acid(33 ml), 1 M HCl-acetic acid(37 ml) was added and stirred at 40°C for 7 hours. The solvent was evaporated under reduced pressure, toluene(50 ml) was added and concentrated again under reduced pressure. The same procedure was repeated again, ethyl acetate(30 ml) was added to the residue and stirred at room temperature overnight. The precipitate was filtered to give compound (11)(5.77 g).


^{1}H-NMR(DMSO-d_6): 2.03(3H, s), 2.06(3H, s), 4.09(2H, d, J=7.5Hz), 5.67(1H, t, J=7.5Hz), 7.12(1H, dd, J=10.7, 8.9Hz), 7.46-7.59(2H, m), 9.24(4H, brs), 10.11(1H, s).

The 4th step: compound (11)(5.16 g) was dissolved in conc. sulfuric acid(15.5 ml) and stirred at room temperature for an hour. It was poured into ice-water (100 ml), adjusted to pH 10 by the addition of an aqueous solution of potassium hydroxide and extracted with ethyl acetate(200 ml) and a little amount of MeOH. The organic layer was washed with water and saturated brine, dried over sodium sulfate. Sodium sulfate was filtered, the filtrate was concentrated under reduced pressure, ethyl acetate(20 ml) and hexane(15 ml) were added to the residue and the precipitate was filtered. The filtrate was concentrated, ethyl acetate (5 ml) and hexane (5 ml) were added and the precipitate was filtered to give compound (12)(total 3.16 g).

^{1}H-NMR($CDCl_3$): 1.62(3H, d, J=0.9Hz), 1.80-1.91(1H, m), 2.16(3H, s), 2.47-2.58(1H, m), 2.62-2.73(1H, m), 2.87-2.98(1H, m), 4.36(2H, brs), 6.99(1H, dd, J=11.7, 8.7Hz), 7.14(1H, dd, J=7.1, 3.0Hz), 7.80(1H, ddd, J=8.7, 4.2, 3.0Hz), 7.97(1H, brs),

The 5th step: Compound (12)(2.50 g) was suspended in ethanol(25 ml), 6 M HCl(10.2 ml) was added and the mixture was stirred at 90°C for 3 hours. 2 M NaOH(35 ml) was added, the organic solvent was evaporated and the residue was extracted with ethyl acetate (70 ml). The aqueous layer was further extracted with ethyl acetate (30 ml), organic layers were combined, washed with water and saturated brine, and dried over sodium sulfate. Sodium sulfate was filtered, the filtrate was concentrated under reduced pressure, and the crystalline residue was washed with ethyl acetate (3 ml) and hexane (10 ml). The crystals were filtered to give compound (8)(total 1.22 g).

Reference Example 3



The 1st step: A ethanol solution of 20% sodium ethoxide (5.12 ml, 16.2 mmol, 40 eq.) was added to compound (13)(150 mg, 406 μmol) and stirred at room temperature for 6 hours. The reaction solvent was evaporated under reduced pressure, 2 M hydrochloric


acid(8.12 ml, 16.2 mmol, 40 eq.) was added to the resulted residue and extracted with chloroform. The extracting solution was washed with water and dried over anhydrous sodium sulfate. The crude product (189 mg) was obtained by evaporation of the solvent under reduced pressure, to which 4 M HCl-ethyl acetate solution (1.89 ml) was added and the mixture was stirred at room temperature for 14 hours. A saturated aqueous solution of sodium bicarbonate was added to the reaction solution, extracted with ethyl acetate and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (14)(90.8 mg, 76% yield) as a yellow powder.

^{1}H NMR ($CDCl_3$)δ1.52 (3H, t, J = 6.8 Hz), 1.67 (3H, s), 1.93-2.00 (1H, m), 2.60-2.67 (2H, m), 2.94-3.00 (1H, m), 4.19 (2H, q, J = 6.8 Hz), 6.93 (1H, d, J = 9.3 Hz), 8.14 (1H, dd, J = 8.7, 2.4Hz), 8.31(1H, d, J = 2.5 Hz).

The 2nd step: A powder of 10% palladium-carbon (45.4 mg) was added to a methanol(908 μl) solution of compound (14)(90.8 mg, 307 μmol) and the mixture was stirred in a hydrogen atmosphere for 22 hours. The reaction mixture was filtered through a Celite pad and the filtrate was evaporated under reduced pressure. The residue was washed with ethyl acetate to give compound (15)(65.8 mg, 81% yield) as an yellow powder.

^{1}H NMR (DMSO-d_6)δ 1.29 (3H, t, J = 6.9 Hz), 1.45 (3H, s), 1.51-1.58 (1H, m), 2.46-2.48 (1H, m), 2.61-2.64 (1H, m), 2.80-2.83 (1H, m), 3.85-3.91 (2H, m), 6.38 (1H, dd, J = 8.3, 2.5 Hz), 6.52 (1H, d, J = 2.4Hz), 6.67 (1H, d, J = 8.6 Hz)

Reference Example 4



The 1st step: 3'-Bromoacetophenone (15.0 g) and compound (16)(9.13 g) were dissolved in tetrahydrofuran (250 ml), and tetraethoxytitanium (39.5 ml) was added thereto at room temperature with stirring. Then the reaction mixture was stirred at 75°C for 5 hours, and saturated brine was added after disappearance of compound (1) was confirmed. Titanium oxide formed in the reaction was filtered off, the filtrate was extracted with ethyl acetate, the organic layer was dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced pressure. The residue was purified with a column chromatography to give compound (17)(20.1 g).

^{1}H-NMR($CDCl_3$) : 1.33 (9H, s), 2.75 (3H, s), 7.30 (1H, t. J=7.8) 7.59-7.63 (1H, m), 7.79 (1H, d, J=7.8) 8.0 (1H, s)

The 2nd step: A 2.64 M hexane solution of n-butyllithium (79.5 ml) is added dropwise to a tetrahydrofuran (100 ml) solution of diisopropylamine (42.1 ml) in a nitrogen atmosphere at -78°C. After stirring at 0°C for 30 minutes, the reaction solution was again cooled to -78°C and tert-butyl acetate (26.9 ml) dissolved in tetrahydrofuran (100 ml) is added dropwise. After stirring at -78°C for 30 minutes, chlorotriisopropoxytitanium dissolved in tetrahydrofuran (150 ml) is added dropwise. After stirring at the same

temperature for 70 minutes, compound (2)(20.1 g) dissolved in tetrahydrofuran (100 ml) was added dropwise. After then, the reaction solution was stirred at -78°C for 3 hours, and an aqueous solution of ammonium chloride was added after disappearance of compound (2) was confirmed. Titanium oxide formed in the reaction was filtered off, the filtrate was extracted with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give a crude product of compound (18)(26.4 g).

The 3rd step: The crude product of compound (18)(26.4 g) was dissolved in toluene (80 ml), the solution was added dropwise to a 1 M solution of aluminium diisobutyl hydride in toluene (253 ml) with stirring at 0°C. The reaction solution was stirred at room temperature for 1.5 hours and 1 N hydrochloric acid solution was added after disappearance of compound (3) was confirmed. The mixture was extracted with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified by crystallization to give compound (19)(18.1 g).

^{1}H-NMR($CDCl_3$) : 1.28 (9H, s,), 1.71 (3H, s), 2.19-2.24 (2H, m), 3.27-3.32(1H, m), 3.54-3.66 (1H, m), 3.87-3.97 (1H, m), 5.10-5.11(1H, m), 7.22 (1H, t. J=8.1) 7.32-7.41 (2H, m), 7.56-7.58 (1H, m)

The 4th step: Compound (19)(18. g) was dissolved in methanol(30 ml), and 10% hydrochloric acid in methanol(130 ml) was added dropwise therein at room temperature. The reaction solution was stirred at room temperature for 4 hours and 1 N hydrochloric acid was added after disappearance of compound (4) was confirmed. The mixture was extracted with ethyl acetate, the aqueous layer was neutralized with a 2 N aqueous solution of sodium hydroxide and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced pressure to give a crude product of compound (20)(14.1 g).

The 5th step: The crude product of compound (20)(32.8 g) and potassium carbonate (37.1 g) were dissolved in a mixed solvent of toluene (450 ml) and water (225 ml) and

thiophosgene (15.3 ml) was added dropwise cooled at 0°C with stirring. After then, the reaction solution was stirred at 0°C for an hour and water was added when disappearance of compound (5) was confirmed. The mixture was extracted with ethyl acetate and the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give a crude product of compound (21)(38.4 g).

The 6th step: The crude product of (21)(38.4 g) was dissolved in toluene (384 ml), and thionyl chloride (29.4 ml) and N,N-dimethylformamide (1.04 ml) were added dropwise at 0°C with stirring. After then, the reaction solution was stirred at 80°C for 5 hours, and after disappearance of compound (6) was confirmed, the reaction solution was evaporated under reduced pressure to give a crude product of compound (22)(40.9 g).

The 7th step: The crude product of compound (22)(40.9 g) was dissolved in tetrahydrofuran (250 ml) and 25% ammonia-water (250 ml) was added with stirring at 0°C. After then the reaction solution was stirred at room temperature for 16 hours, and a saturated aqueous solution of sodium bicarbonate was added after disappearance of compound (21) was confirmed. The organic layer was separated and the aqueous solution was extracted with dichloromethane. The organic layers were combined, dried over anhydrous magnesium sulfate and evaporated under reduced pressure to give a crude product of compound (23)(38.3 g).

The 8th step: The crude product of compound (23)(38.3 g) is dissolved in tetrahydrofuran (383 ml), di-tert-butyl dicarbonate (61.5 g) and N,N-dimethylaminopyridine (1.64 g) are added and the mixture was stirred at room temperature for 72 hours. After disappearance of compound (23) was confirmed, the solvent was evaporated under reduced pressure. The residue was purified with a silicagel column chromatography to give compound (24)(45.3 g)


^{1}H-NMR($CDCl_3$) : 1.54 (9H, s,), 1.57 (3H, s), 1.96 (2H, t, J=6.0), 2.80-2.92(1H, m), 3.00-3.13 (1H, m), 7.21 (1H, t. J=8.1) 7.28-7.41 (2H, m), 7.52-7.55 (1H, m)

The 9th step: In a nitrogen atmosphere, compound (24)(12.1 g),

trisdibenzylideneacetonedipalladium(1.14 g) and dicyclohexylbiphenylphosphine (0.88 g) were dissolved in toluene (125 ml), and a 1.6 M solution of lithium hexamethyldisilazide in tetrahydrofuran (46.9 ml) was added with stirring at room temperature. The reaction solution was warmed up to 80°C and stirred for 16 hours. After disappearance of compound (21) was confirmed, the reaction solution was cooled at 0°C and diethyl ether and 1 N hydrochloric acid were added. After stirring at 0°C for 10 minutes, the solution was neutralized with a saturated aqueous solution of sodium carbonate. It was extracted with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (25)(6.84 g).

^{1}H-NMR($CDCl_3$) : 1.51 (9H, s,), 1.69 (3H, s), 2.01-2.12 (1H, m), 2.40-2.51(1H, m), 2.67-2.76 (2H, m), 6.55-6.67 (3H, m), 7.15 (1H, t. J=8.1).

Reference Example 5



The 1st step: After the addition of compound (1)(70.00 g) to conc. sulfuric acid(279 ml) cooled in an acetonitrile /dry ice bath with stirring, a mixture of fuming nitric acid(42 ml)

and conc. sulfuric acid(98 ml) were added dropwise. After stirring for 16 minutes, the mixture was gradually poured into ice, the precipitated crystals were filtered and dried to give compound (2)(77.79 g).

^{1}H-NMR ($CDCl_3$) δ: 2.71 (3H, d, J = 4.9 Hz), 7.34 (1H, t, J = 9.3 Hz), 8.40 (1H, ddd, J = 9.3, 6.2, 3.0 Hz), 8.78 (1H, dd, J = 6.2, 3.0 Hz).

The 2nd step: A solution of compound (2)(73.94 g), (R)-(+)-2-methyl-2-propanesulfinamide (53.82 g) and tetraethyl orthotitanate (230.20 g) in tetrahydrofuran (500 ml) were reacted for 2.5 hours under heating to reflux, and then the reaction mixture was gradually poured into ice and the resulted insoluble materials were filtered. It was extracted with ethyl acetate, the organic layer was dried over anhydrous magnesium sulfate and evaporated under reduced pressure. The residue was crystallized from ethyl acetate/n-hexane to give compound (26)(85.44 g).

^{1}H-NMR ($CDCl_3$) δ: 1.34 (9H, s), 2.81 (3H, d, J = 3.5 Hz), 7.29 (1H, t, J = 8.9 Hz), 8.31 (1H, dt, J = 8.9, 2.9 Hz), 8.55 (1H, dd, J = 6.3, 2.9 Hz).

The 3rd step: A solution of tert-butyl acetate (6.08 g) in tetrahydrofuran (10 ml) was added dropwise to a 2 M solution of lithium diisopropylamide/tetrahydrofuran/n-heptane/ethylbenzene (27.9 ml) cooled in an acetone/dry ice bath with stirring. After stirring for 20 minutes, a solution of chlorotitanium isopropoxide (17.5 ml) in tetrahydrofuran (30 ml) was added dropwise, the mixture was stirred for an hour and a solution of compound (26)(5.00 g) in tetrahydrofuran (10 ml) was added dropwise. After reacting for an hour, the reaction solution was gradually poured into an aqueous solution of ammonium chloride cooled in ice with stirring and the resulted insoluble materials were filtered. It was extracted with ethyl acetate, the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (27)(5.49 g).

^{1}H-NMR ($CDCl_3$) δ: 1.30 (9H, s), 1.35 (9H, s), 1.86 (3H, s), 3.11 (1H, dd, J = 16.2, 2.1 Hz), 3.26 (1H, dd, J = 16.2, 2.1 Hz), 5.55 (1H, s), 7.18 (1H, dd, J = 11.1, 8.9 Hz), 8.18 (1H, ddd, J

= 8.9, 4.0, 2.9 Hz), 8.53 (1H, dd, J = 7.0, 2.9 Hz).

Ratio of diastereomers (3S:3R=97:3) HPLC Column : CHIRALPAK AS-RH, Detection :254 nm : Column temp. : 25°C, Mobile phase : 40% MeCNaq., Flow rate : 0.5ml/min.

Note: As to the stereochemistry of compound (27) obtained above, it is known that 3S-isomer is preferentially prepared as written in the literature A etc. and it is also possible to prepare each diastereomer selectively by choosing appropriate metal species and/or a reaction condition.

Literature A: (1) T. Fujisawa et al., *Tetrahedron Lett.*, 37, 3881-3884 (1996), (2) D. H. Hua et al, *Sulfur Reports*, vol. 21, pp. 211-239 (1999), (3) Y. Koriyama et al., *Tetrahedron*, 58, 9621-9628 (2002), (4) Yong Qin et al., *J. Org. Chem.*, 71, 1588-1591 (2006).

The 4th step: A solution of 4 M HCl/1,4-dioxane (50 ml) was added to compound (27)(12.74 g) and the mixture was stirred at 80°C for an hour, diethyl ether (50 ml) was added, the precipitated crystals were filtered and dried to give compound (28)(7.67 g).

^{1}H-NMR (DMSO-d_6) δ: 1.76 (3H, s), 3.25 (2H, s), 7.62 (1H, dd, J = 11.4, 9.4 Hz), 8.33-8.48 (2H, m).

The 5th step: A solution of 1 M tetrahydrofuran-borane in tetrahydrofuran (2029 ml) was added dropwise to a solution of compound (28)(141.32 g) in tetrahydrofuran (707 ml) cooled in ice with stirring and it was reacted for 3 hours and 6 minutes. The reaction mixture was poured into a mixture of sodium bicarbonate (511 g), ice (1500 g) and ethyl acetate (3000 ml) stirred at room temperature, extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (29)(115.46 g) as a crude product.

The 6th step: Toluene (25 ml) and water (12.5 ml) were added to the compound (29) (3.76 g) obtained in the 5th step and stirred under ice cooling. After the addition of potassium carbonate (7.97 g), thiophosgene (2.85 g) was added dropwise. After reacting for 3 hours, water was added, extracted with toluene and the organic layer was dried over

anhydrous magnesium sulfate. A part of the solvent was evaporated under reduced pressure to give compound (30) as a crude product.

The 7th step: Compound (30) obtained in the 6th step was dissolved in toluene (17.4 ml) and thionyl chloride (6.67 g) and N,N-dimethylformamide (0.128 ml) were added with stirring at room temperature. The mixture was stirred at 80°C for 2 hours, water was added, extracted with toluene and concentrated under reduced pressure to give compound (31)(4.03 g) as a crude product.


The 8th step: Compound (31)(4.03 g) obtained in the 7th step was dissolved in tetrahydrofuran (23.8 ml) and 28% ammonia-water (23.8 ml)was added under ice cooling with stirring. The mixture was stirred at room temperature for 3 days, the reaction solution was concentrated under reduced pressure and ethyl acetate was added therein. Conc. hydrochloric acid(6 ml) was added under ice cooling with stirring, the precipitated crystals were washed with ethyl acetate and water and dried to give compound (32)(2.14 g).

^{1}H-NMR (DMSO-d_6) δ: 1.76 (3H, s), 2.13-2.24 (1H, m), 2.68-2.74 (2H, m), 3.19-3.25 (1H, m), 7.63 (1H, dd, J = 11.4, 8.9 Hz), 8.07 (1H, dd, J = 7.0, 3.5 Hz), 8.36 (1H, dt, J = 8.9, 3.5 Hz), 11.22 (1H, s).

The 9th step: Compound (32) (100 mg) was dissolved in methanol(2 ml), 10% palladium-carbon powder (50 mg) was added, and the mixture was stirred in a hydrogen atmosphere at room temperature for 18 hours. Insoluble materials were filtered off, the filtrate was evaporated under reduced pressure, sodium carbonate and water were added therein and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and the solvent was evaporated under reduced pressure to give compound (33)(68 mg).

^{1}H-NMR ($CDCl_3$)δ: 1.59 (3H, s), 1.81 (1H, ddd, J = 14.1, 10.9, 3.5 Hz), 2.47 (1H, ddd, J = 14.1, 5.9, 3.5 Hz), 2.71 (1H, td, J = 10.9, 3.5 Hz), 2.89 (1H, ddd, J = 10.9, 5.9, 3.5 Hz), 3.57 (2H, br s), 6.49 (1H, dt, J = 8.5, 3.3 Hz), 6.67 (1H, dd, J = 6.9, 3.3 Hz), 6.80 (1H, dd, J = 11.8, 8.5 Hz).

Reference Example 6



The 1st step: A solution of compound (38)(5.00 g), (R)-(+)-2-methyl-2-propanesulfinamide (3.33 g) and tetraethyl orthotitanate (17.11 g) in tetrahydrofuran (50 ml) was reacted under heating to reflux for 7 hours, and then, it was poured portionwise into saturated brine and the resulted insoluble materials were filtered off. It was extracted with ethyl acetate, the organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure. The residue was purified with a silicagel column chromatography to give compound (39)(6.37 g).

^{1}H-NMR ($CDCl_3$) δ: 1.34 (9H, s), 2.79 (3H, s), 8.26 (1H, t, J = 2.3 Hz), 8.76 (1H, d, J = 2.3 Hz), 8.96 (1H, d, J = 2.3 Hz).

The 2nd Step: A solution of 2.66 M n-butyllithium/n-hexane (32.4 ml) was added dropwise to a solution of diisopropylamine (9.36 g) in tetrahydrofuran (39 ml) cooled in an acetone/dry ice bath with stirring and the mixture was stirred under ice cooling for 30 minutes. The reaction solution was stirred again in an acetone/dry ice bath and a solution of tert-butyl acetate (4.88 g) in tetrahydrofuran (8 ml) was added dropwise. After stirring for 40 minutes, a solution of chlorotitanium triisopropoxide (23.00 g) in tetrtahydrofuran (88 ml)

was added dropwise. After stirring for 10 minutes, a solution of compound (39)(6.37 g) in tetrahydrofuran (65 ml) was added dropwise. After reacting for 30 minutes, the reaction solution was poured portionwise into an aqueous solution of ammonium chloride and the resulted insoluble materials were filtered off. It was extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (40)(8.03 g) as a crude product.

The 3rd step: Lithium aluminium hydride (2.85 g) was added portionwise to a solution of the compound (40)(8.03 g) obtained in the 2nd step in tetrahydrofuran (100 ml) cooled in ice with stirring and the mixture was stirred for 2 hours. Acetone, water, and a 1N aqueous solution of sodium hydroxide were added portionwise and the mixture was stirred at room temperature for 30 minutes. The insoluble materials were filtered off and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate and concentrated under reduced pressure to give compound (41)(5.83 g) as a crude product.

The 4th step: A solution of 10% HCl/methanol (60 ml) was added to a solution of the compound (41)(5.83 g) obtained in the 3rd step in methanol (60 ml) cooled in ice with stirring and stirred at room temperature for 16 hours. The reaction solution was made alkaline by the addition of water and potassium carbonate, extracted with chloroform, the organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give compound (42)(5.07 g) as a crude product.

The 5th step: Imidazole (2.24 g) and t-butyldimethylsilyl chloride (3.77 g) were added to a solution of the compound (42)(5.07 g) obtained in the 4th step in N,N-dimethylformamide (26 ml) with stirring at room temperature and the mixture was stirred for 1 hour and 40 minutes. After extraction with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (43)(3.82 g).

^{1}H-NMR ($CDCl_3$) δ: -0.04 (3H, s), -0.01 (3H, s), 0.85 (9H, s), 1.51 (3H, s), 1.98 (2H, t, J=

6.0 Hz), 3.49-3.54 (1H, m), 3.65 (1H, dt, J = 11.1, 6.0 Hz), 8.02 (1H, t, J = 2.2 Hz), 8.53 (1H, d, J = 2.2 Hz), 8.63 (1H, d, J = 2.2 Hz).

The 6th step: Toluene (25 ml) and water (13 ml) were added to compound (43)(3.82 g) and stirred under ice cooling. After the addition of potassium carbonate (5.14 g), thiophosgene (1.83 g) was added dropwise. After reacting for 2 hours, water was added, extracted with chloroform and the organic layer was dried over anhydrous magnesium sulfate. A part of the solvent was evaporated under reduced pressure to give compound (44) as a crude product.

The 7th step: Thionyl chloride (4.43 g) and N,N-dimethylformamide (0.08 ml) were added to a solution of the compound (7) obtained in the 6th step in toluene (25 ml) with stirring at room temperature. The mixture was stirred at 80°C for 5 hours, concentrated under reduced pressure to give compound (45)(5.03 g) as a crude product.

The 8th step: 28% Ammonia water (60 ml) was added to a solution of the compound (45)(5.03 g) obtained in the 7th step in tetrahydrofuran (60 ml) stirred under ice cooling and the mixture was stirred at room temperature for 14 hours. The reaction solution was concentrated under reduced pressure to give compound (46)(4.92 g) as a crude product.

The 9th step: A mixture of the compound (46)(4.92 g) obtained in the 8th step, di-t-butyl dicarbonate (9.28 g), triethylamine (3.23 g), 4-dimethylaminopyridine (0.13 g) and tetrahydrofuran (106 ml) was stirred at room temperature for 3 days. The insoluble materials were filtered off, water was added to the filtrate and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give compound (47)(8.31 g) as a crude product.


The 10th step: A mixture of the compound (47)(8.31 g) obtained in the 9th step, di-t-butyl dicarbonate (6.96 g), triethylamine (3.23 g), 4-dimethylaminopyridine (0.13 g) and tetrahydrofuran (50 ml) was stirred at room temperature for an hour. After the addition of water, it was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (48)(1.23 g).


^{1}H-NMR ($CDCl_3$) δ: 1.53 (18H, s), 1.60 (3H, s), 1.93 (1H, ddd, J = 13.8, 9.4, 3.9 Hz), 2.06 (1H, ddd, J = 13.8, 3.9, 1.9 Hz), 2.91 (1H, ddd, J = 12.9, 3.9, 1.9 Hz), 3.15 (1H, ddd, J = 12.9, 9.4, 3.9 Hz), 7.89 (1H, t, J = 2.1 Hz), 8.55-8.57 (2H, m).

The 11th step: Compound (48)(190 mg), trisdibenzylideneacetonedipalladium (54 mg), dicyclohexylbiphenylphosphine (41 mg) were dissolved in toluene (5 ml), stirred at room temperaturea, and 1.6 M solution of lithium hexamethyldisilazide in tetrahydrofuran (0.73 ml) was added therein. The reaction solution was warmed up to 85°C and stirred for 9 hours, then, it was cooled in ice and diethyl ether and a 1 N solution of hydrochloric acid were added. After stirring for 10 minutes, it was neutralized by the addition of a saturated aqueous solution of sodium carbonate and extracted with dichloromethane. The organic layer was dried over anhydrous sodium sulfate, concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (49)(27 mg).

^{1}H-NMR ($CDCl_3$) δ: 1.51 (9H, s), 1.68 (3H, s), 2.12 (1H, ddd, J = 14.8, 11.0, 3.0 Hz), 2.38-2.47 (1H, m), 2.64-2.70 (1H, m), 2.78-2.82 (1H, m), 3.80 (2H, br s), 6.90 (1H, t, J = 2.4 Hz), 7.98 (1H, dd, J = 10.4, 2.4 Hz).

Reference Example 7





The 1st step: A solution of compound (50)(38.93 g), (R)-(+)-2-methyl-2-propanesulfinamide (13.20 g) and tetraethyl orthotitanate (67.76 g) in tetrahydrofuran (389 ml) was reacted under heating to reflux for 4 hours. A saturated aqueous solution of ammonium chloride was added portionwise therein and the resulted insoluble materials were filtered off. The filtrate was concentrated and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give compound (51)(30.52 g) as a crude product.

^{1}H-NMR ($CDCl_3$) δ: 1.32 (9H, s), 2.83 (3H, s), 7.55-7.65 (2H, m), 8.06 (1H, d, J = 8.5 Hz).

The 2nd step: A solution of tert-butyl acetate (22.99 g) in tetrahydrofuran (148 ml) was added dropwise to a 2.0 M solution of lithium diisoprorylamide/n-heptane/ethylbenzene/tetrahydrofuran (202.5 ml) cooled in an acetone/dry ice bath with stirring. After stirring for 45 minutes, a solution of chlorotitanium triisopropoxide (108.36 g) in tetrahydrofuran (342 ml) was added dropwise and stirred for 40 minutes. A solution of the compound (51)(30.52 g) in tetrahydrofuran (342 ml) was added dropwise and reacted for an hour. The reaction solution was poured portionwise into an

aqueous solution of ammonium chloride with stirring under ice cooling and the resulted insoluble materials were filtered off. It was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (52)(27.40 g).

^{1}H-NMR ($CDCl_3$) δ: 1.30 (9H, s), 1.35 (9H, s), 1.65 (3H, s), 3.01 (1H, d, J = 16.5 Hz), 3.38 (1H, d, J = 16.5 Hz), 5.60 (1H, s), 7.31 (1H, dd, J = 5.9, 2.7 Hz), 7.48-7.50 (2H, m).

The 3rd step: Lithium aluminium hydride (5.67 g) was added portionwise to a solution of the compound (52)(22.40 g) in tetrahydrofuran (336 ml) stirred in an ice salt bath and stirred for 7 hours. After the addition of acetone, water and a 1 N aqueous solution of sodium hydroxide, the insoluble materials were filtered off and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give compound (53)(18.75 g) as a crude product.

The 4th step: A solution of 10% HCl/methanol(94 ml) was added to a solution of the compound (53)(18.75 g) obtained in the 3rd step in methanol(94 ml) stirred under ice cooling and stirred at room temperature for 1.5 hours. The reaction solution was made alkaline by the addition of water and potassium carbonate and extracted with chloroform. The organic layer was dried over anhydrous sodium sulfate and concentrated under reduced pressure to give compound (54)(21.03 g) as a crude product.

The 5th step: Imidazole (5.49 g) and tert-butyldimethylsilyl chloride (10.53 g) were added to a solution of the compound (54)(21.03 g) in N,N-dimethylformamide (210 ml) stirred at room temperature and the mixture was stirred for an hour. After extraction with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (55)(20.12 g).

^{1}H-NMR ($CDCl_3$) δ: -0.04 (3H, s), -0.02 (3H, s), 0.84 (9H, s), 1.47 (3H, s), 1.95-2.15 (2H, m), 3.54-3.63 (2H, m), 7.29 (1H, dd, J = 6.1, 2.6 Hz), 7.45-7.48 (2H, m).

The 6th step: Toluene (66 ml) and water (33 ml) were added to compound (55)(10.06

g) and stirred under ice cooling. After the addition of potassium carbonate (11.13 g), thiophosgene (2.86 ml) was added dropwise. After reacting for an hours, water was added, extracted with chloroform and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (56)(9.43 g).

^{1}H-NMR ($CDCl_3$) δ: -0.03 (6H, s), 0.82 (9H, s), 1.80 (3H, s), 2.21-2.24 (1H, m), 2.44-2.48 (1H, m), 3.57 (1H, ddd, J = 12.0, 5.8, 4.8 Hz), 3.71 (1H, ddd, J = 12.0, 5.8, 4.8 Hz), 7.37 (1H, dd, J = 7.5, 1.2 Hz), 7.48-7.58 (2H, m).

The 7th step: 28% Ammonia water (47 ml) was added to compound (56)(9.43 g) dissolved in tetrahydrofuran (94 ml) stirred at room temperature. After stirring for 16 hours, water was added, extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (57)(6.35 g) as a crude product.

The 8th step: Acetic acid(1.09 g) and a 1.0 M solution of tetrabutylammonium fluoride/tetrahydrofuran (18.20 ml) were added to a solution of the compound (57)(6.35 g) obtained in the 7th step in tetrahydrofuran (127 ml) stirred under ice cooling. After stirring at room temperature for 3 hours, water and potassium carbonate were added and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (58)(4.47 g).

^{1}H-NMR ($CDCl_3$) δ: 1.85 (3H, s), 2.27-2.31 (2H, br m), 3.73-3.83 (2H, m), 5.86 (2H, br s), 7.43 (1H, d, J = 7.8 Hz), 7.52 (1H, d, J = 7.8 Hz), 7.61 (1H, t, J = 7.8 Hz), 7.81 (1H, br s).

The 9th step: 1-Chloro-N,N,2-trimethyl-1-propenylamine (2.16 g) was added to compound (58)(4.47 g) dissolved in dichloromethane (89 ml) stirred under ice cooling. After stirring at room temperature for 1.5 hours, water was added and extracted with dichloromethane. The organic layer was dried over anhydrous sodium sulfate, concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (59)(2.91 g).

^{1}H-NMR ($CDCl_3$) δ: 1.53 (3H, s), 1.88 (1H, ddd, J = 13.9, 10.1, 3.8 Hz), 2.40 (1H, ddd, J = 13.9, 6.6, 3.8 Hz), 2.71 (1H, ddd, J = 13.9, 10.1, 3.8 Hz), 2.95 (1H, tt, J = 6.6, 3.8 Hz), 4.33 (2H, br s), 7.29 (1H, dd, J = 7.5, 1.2 Hz), 7.41-7.50 (1H, m).


The 10th step: A mixture of compound (59)(2.91 g), di-tert-butyl dicarbonate (5.52 g), 4-dimethylaminopyridine (0.12 g) and tetrahydrofuran (29 ml) was stirred at room temperature for 2.5 hours. The reaction solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (60)(1.23 g).

^{1}H-NMR ($CDCl_3$) δ: 1.53 (23H, s), 1.60 (3H, s), 1.93 (1H, ddd, J = 13.8, 9.4, 3.9 Hz), 2.06 (1H, ddd, J = 13.8, 3.7, 1.8 Hz), 2.91 (1H, ddd, J = 12.7, 3.7, 1.9 Hz), 3.15 (1H, ddd, J = 12.9, 9.2, 3.7 Hz), 7.89 (1H, t, J = 2.1 Hz), 8.55-8.57 (2H, m).

The 11th step: Compound (60)(3.30 g), trisdibenzylideneacetonedipalladium(0.93 g), dicyclohexylbiphenylphosphine (0.73 g) were dissolved in toluene (66 ml), stirred at room temperature, and a 1.6 M solution of lithium hexamethyldisilazide in tetrahydrofuran (12.7 ml) was added therein. The reaction solution was warmed up to 80°C and stirred for 8 hours, then, it was cooled in ice and diethyl ether and a 1 N solution of hydrochloric acid were added. After stirring for 5 minutes, it was neutralized by the addition of a saturated aqueous solution of sodium carbonate and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (61)(1.55 g).

^{1}H-NMR ($CDCl_3$) δ: 1.61 (3H, s), 1.74-1.80 (1H, m), 1.96-2.11 (1H, m), 2.64-2.82 (2H, m), 4.41 (2H, br s), 6.39 (1H, dd, J = 8.1, 0.6 Hz), 6.71 (1H, dd, J = 8.1, 0.6 Hz), 7.42 (1H, t, J = 8.1 Hz).

Reference Example 8



The 1st step: Compound (63)(3.31 g) prepared in the same manner as Reference Example 6 and 7 described above was dissolved in dichloromethane (16.5 ml), bis(2,4-dimethoxybenzyl)amine (4.45 g)was added and the solvent was evaporated under reduced pressure after stirring at room temperature for an hour and standing for 15 hours. The residue was purified with silicagel chromatography to give compound (63)(5.77 g).

^{1}H-NMR($CDCl_3$) : -0.10(3H, s), -0.07(3H, s), 0.77(9H, s), 1.93(3H, s), 2.08-2.27(1H, m), 3.06-3.28(1H, m), 3.38(1H, ddd, J=10.8, 6.8, 6.8 Hz), 3.55(1H, ddd, J=10.8, 6.8, 6.8 Hz), 3.78(6H, s), 3.79(6H, s), 4.81-5.05(1H, br), 6.43-6.50(4H, m), 7.07(1H, d, J=1.9 Hz), 7.17(2H, d, J=7.3 Hz), 8.05-8.16(2H, m).

The 2nd step: Compound (63)(5.77 g) was dissolved in tetrahydrofuran (60 ml) and acetic acid(1.01 g) and 1M tetrabutylammonium fluoride/tetrahydrofuran-solution (15 ml) were added and the mixture was stirred at room temperature for 150 minutes. Water was added, extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel chromatography to give compound(64)(3.94 g).


^{1}H-NMR($CDCl_3$) : 1.99(3H, s), 1.91-2.02(1H, m), 3.10(1H, ddd, J=14.6, 6,6, 5.2), 3.36-3.51(2H, m), 3.78(6H, s), 3.80(6H, s), 4.58(2H, d, J=15.8 Hz), 4.72(2H, d, J=15.8 Hz), 6.43-6.51(4H, m), 7.12(1H, dd, J=5.3, 2.0 Hz), 7.18-7.29(3H, m), 8.20(1H, dd, J=5.3, 0.6 Hz), 8.28-8.31(1H, br).

The 3rd step: Compound (64)(3.94 g) was dissolved in dichloromethane (20 ml) and

1-chloro-N,N',2-trimethyl-1-propenylamine (1.86 ml) was added under ice cooling with stirring. After stirring at room temperature for 2 hours, water was added, extracted with chloroform and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound(65)(3.41 g).

^{1}H-NMR($CDCl_3$) : 1.49(3H, s), 1.99(1H, ddd, J=13.3, 7.6, 3.2 Hz), 2.31(1H, ddd, J=13.3, 8.6, 3.7), 2.78(1H, ddd, J=12.2, 8.6, 3.2 Hz), 3.04(1H, ddd, J=12.2, 7.6, 3.7 Hz), 3.77(6H, s), 3.79(6H, s), 4.60(2H, d, J=15.8 Hz), 4.76(2H, d, J=15.8 Hz), 6.45-6.52(4H, m), 7.08(1H, dd, J=5.3, 2.1 Hz), 7.17-7.27(3H, m), 8.40(1H, dd, J=5.3, 0.5 Hz).

Reference Example 9



The 1st step: Compound (66)(4.72 g) derived by a conventional method from an intermediate prepared in the same manner as the compound (27) described above was dissolved in tetrahydrofuran (150 ml) and a diethylether solution of methyl magnesium bromide (3M, 37 ml) was added dropwise with stirring under ice cooling in a nitrogen stream for 12 minutes. After stirring 3 hours, a saturated aqueous solution of ammonium chloride (190 ml) was added dropwise, extracted with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was

evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (67)(2.11 g).

^{1}H-NMR(DMSO-d_6) : 0.75(3H, s), 1.09(3H, s), 1.21(9H, s), 1.79(3H, s), 2.06(1H, m), 2.29(1H, m), 4.97(1H, s), 6.57(1H, s), 7.17(1H, dd, J=8.7, 12.0Hz), 7.48-7.53(1H, m), 7.99-8.03(1H, m), 11.26(1H, bs).

The 2nd step: Compound (67)(2.11 g) was dissolved in methanol(7.8 ml) and hydrochloric acid-methanol solution (5-10%)(15.6 ml) was added with stirring at room temperature, and the mixture was stirred for 1.5 hours. Then the reaction solution was poured into ice water and ethyl acetate (100 ml), a saturated aqueous solution of sodium bicarbonate (50 ml) was added and extracted with ethyl acetate. The aqueous layer was further extracted with ethyl acetate (50 ml), organic layers were combined, washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was crystallized from n-hexane to give compound (68)(1.42 g).

^{1}H-NMR(DMSO-d_6) : 0.65(3H, s), 1.10(3H, s), 1.43(3H, s), 1.85(1H, d, J=14.4Hz), 2.17(1H, dd, J=1.5, 14.4Hz), 7.12(1H, dd, J=2.7, 12.0Hz), 7.60-7.64(1H, m), 7.90(1H, dd, J=2.7, 7.5Hz), 11.35(1H, bs).

The 3rd step: Toluene (9.6 ml) and water (4.8 ml) were added to compound (68)(1.42 g) and suspended, potassium carbonate (2.13 g) was added with stirring under ice cooling and 2 minutes later thiophosgene (0.51 ml) was added at once and the stirring was continued. The temperature was back to room temperature 40 minutes later, toluene (40 ml) and water were added and extracted an hour later. The aqueous layer was further extracted with toluene, organic layers were combined, washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give a crude product (69)(2.02 g).

The 4th step: Tetrahydrofuran (17 ml) was added to triphenylphosphine (1.735 g) and N-chlorosuccinimide (833 mg), suspended in a nitrogen stream and stirred at room temperature for 10 minutes. A tetrahydrofuran (21 ml) solution of the crude product

(69)(2.02 g) was added dropwise using a dropping funnel for 2 minutes. After stirring for 6 hours, the mixture was left stand at room temperature overnight. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (70)(828 mg).

^{1}H-NMR(DMSO-d_6) : 1.54(3H, s), 1.86(3H, s), 2.81(1H, d, J=13.8Hz), 2.92(1H, d, J=13.8Hz), 4.73(1H, s), 4.85(1H, m), 7.28-7.35(1H, m), 7.77-7.82(2H, m),11.39(1H, bs).


The 5th step: Compound (70)(828 mg) was dissolved in tetrahydrofuran (4 ml), conc. ammonia water (28%)(4 ml) was added with stirring under ice cooling and the temperature was back to room temperature after stirring for 5 minutes. After 25 hours, the reaction mixture was poured into ice water and extracted with ethyl acetate (50 ml). The organic layer was washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel chromatography to give a compound (71)(260 mg).

^{1}H-NMR(DMSO-d_6) : 1.47(3H, bs), 1.66(3H, bs), 2.58(1H, d, J=12.3Hz), 4.71(1H, s), 4.87(3H, bs), 6.42(1H, bs), 6.51(1H, dd, J=2.7, 7.2Hz), 6.75(2H, bs), 7.54(1H, bs).

The 6th step: Compound (71)(245 mg) was dissolved in chilled conc. sulfuric acid(4.9 ml) and stirred under ice cooling for 2 hours. The reaction solution was poured into ice water with stirring and pH was adjusted to 2-3 by the addition of a 5N aqueous solution of sodium hydroxide. Ethyl acetate (100 ml) and an aqueous solution of potassium carbonate were added and extracted under alkaline condition. The alkaline layer was further extracted with ethyl acetate (50 ml). Organic layers were combined, washed with saturated brine (50 ml) and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (72)(101 mg) as a crystal.

^{1}H-NMR(DMSO-d_6) : 0.83(3H, s), 1.27(3H, s), 1.44(3H, s), 1.54(1H, d, J=14.1Hz), 2.45(1H, d, J=14.1Hz), 4.79(2H, s), 5.89(2H, bs), 6.32-6.37(1H, m), 6.58(1H, dd, J=2.7, 7.2Hz), 6.72(1H, dd, J=8.7, 12.3Hz).

Reference Example 10



(73) (74) (75) (76) (77) (78) (79) (80) (81)

The 1st step: Ethyl bromodifluoroacetate (0.77 ml) was added to a suspension of zinc dust (392 mg) in tetrahydrofuran (4 ml) with stirring in a nitrogen stream at room temperature, stirred for 15 minutes, ethyl bromodifluoroacetate (0.29 ml) was added, stirred for 30 minutes to prepare a solution of ethyl bromozincdifluoroacetate. This solution was added to a solution of compound (73) in tetrahydrofuran (3 ml) in a nitrogen stream and stirred for 8 hours. 3% Ammonia water was added to the reaction mixture with stirring under ice cooling, extracted with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (74)(696 mg).

^{1}H-NMR(DMSO-d_6)δ: 1.17 (3H, t, J=7.2 Hz), 1.18 (9H, s), 2.00 (3H, brs), 4.24 (2H, q, J=7.2 Hz), 5.56 (1H, brs), 7.56 (dd, J=9.0, 11.7 Hz), 8.36 (1H, m), 8.49 (1H, dd, J=3.0, 6.6 Hz).

The 2nd step: Compound (74)(670 mg) was dissolved in tetrahydrofuran (6.7 ml) and lithium borohydride (71 mg) was added in a nitrogen stream with stirring under ice cooling. After stirring for 30 minutes, acetic acid(198 mg) and ice water were added to the reaction mixture, extracted with ethyl acetate, the organic layer was washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (75)(401 mg).

^{1}H-NMR(DMSO-d_6)δ: 1.20(9H, s), 2.00 (3H, d, J=3.6 Hz), 3.80 (1H, m), 4.00 (1H, m), 5.99 (1H, s), 6.34 (1H, t, J=5.7 Hz), 7.53 (1H, dd, J=9.0, 12.0 Hz), 8.31 (1H, m), 8.50 (1H, dd, J=2.7, 6.6 Hz).

The 3rd step: Compound (75)(394 mg) was dissolved in methanol(3 ml), and 4N-HCl/1,4-dioxane (1.35 ml) was added with stirring under ice cooling. After stirring for 30 minutes, the mixture was stirred at room temperature for 1.5 hours. Ice water was added to the reaction solution and washed with ethyl acetate. The aqueous layer was made alkaline by the addition of a 2M aqueous solution of potassium carbonate and extracted with ethyl acetate. The organic layer was dried over anhydrous magnesium sulfate, the solvent was evaporated under reduced pressure to give compound (76)(293 mg).

^{1}H-NMR(DMSO-d_6)δ: 1 .62 (3H, d, J=2.7 Hz), 2.62 (2H, brs), 3.65-3.83 (2H, m), 5.31 (1H, brt), 7.44 (1H, dd, J=9.0, 11.4 Hz), 8.23 (1H, m), 8.59 (1H, dd, J=3.0, 6.9 Hz).

The 4th step: Compound (76)(266 mg) was dissolved in acetone (3 ml) and benzoyl isothiocyanate (164 mg) was added in a nitrogen stream with stirring under ice cooling. After stirring for an hour, the mixture was stirred at room temperature for an hour. The reaction solution was concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (77)(353 mg).

^{1}H-NMR(DMSO-d_6)δ: 2.30 (3H, brs), 3.65-3.96 (2H, m), 5.90 (1H, brt), 7.42-7.68 (4H, m), 7.93-7.96 (2H, m), 8.17-8.33 (2H, m), 11.42 (1H, brs), 12.31 (1H, brs).

The 5th step: Compound (77)(348 mg) was dissolved in dichloromethane (4 ml) and 1-chloro-N,N-2-trimethyl-1-propenylamine (131 mg) was added in a nitrogen stream with stirring under ice cooling. After stirring for 15 hours at room temperature, ice water was added and neutralized by the addition of potassium carbonate. It was extracted with ethyl acetate, washed with brine and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (78)(308 mg).

^{1}H-NMR($CDCl_3$)δ: 1.89 (3H, d, J=3.0 Hz), 3.17 (1H, ddd, J=8.4, 10.2, 13.2 Hz), 3.51 (1H, ddd, J=6.0, 13.2, 19.2 Hz), 7.23 (1H, dd, J=9.0, 10.8 Hz), 7.49-7.64 (3H, m), 7.91 (2H, d,

J=7.2 Hz), 8.24 (1H, m), 8.43 (1H, dd, J=3.0, 6.6 Hz), 8.57 (1H, br).

The 6th step: Compound (78)(297 mg) was dissolved in ethanol(4 ml), water (1.5 ml) and conc. hydrochloric acid(1.5 ml) were added and the mixture was stirred at 90°C for 2.5 hours. Water was added to the reaction solution, washed with ethyl acetate and the aqueous layer was made alkaline by the addition of a 2M aqueous solution of potassium carbonate. It was extracted with ethyl acetate, washed with brine and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (79)(89 mg).

^{1}H-NMR($CDCl_3$)δ: 1.85 (3H, d, J=3.6 Hz), 3.15 (1H, ddd, J=8.7, 10.5, 12.9 Hz), 3.50 (1H, ddd, J=5.4, 12.9, 18.3 Hz), 4.51 (2H, brs), 7.19 (1H, dd, J=9.0, 11.1 Hz), 8.20 (1H, ddd, J=3.0, 6.9, 9.0 Hz), 8.54 (1H, dd, J=3.0, 6.9 Hz).


The 7th step: Compound (79)(82 mg) was dissolved in dichloromethane (1 ml), di-tert-butyldicarbonate (176 mg) and 4-dimethylaminopyridine (4 mg) were added and the mixture was stirred at room temperature for 30 minutes. The reaction solution was concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (80)(101 mg).

^{1}H-NMR($CDCl_3$)δ: 1.56 (18H, S), 1 .90 (3H, d, J=3.6 Hz), 3.27 (1H, ddd, J=6.6, 9.3, 12.9 Hz), 3.69 (1H, ddd, J=4.2, 12.9, 17.4 Hz), 7.23 (1H, dd, J=9.0, 12.0 Hz), 8.24 (1H, ddd, J=3.0, 9.0, 12.0 Hz), 8.41 (1H, ddd, J=2.4, 3.0, 6.0 Hz).

The 8th step: Compound (80)(4.76 g) was dissolved in methanol(70 ml), 10% Pd-C(containing 50% water)(2.38 g) was added and the mixture was stirred in a hydrogen atmosphere for 20 hours. The catalyst was filtered off, the solvent was evaporated under reduced pressure to give compound (81)(4.43 g)

^{1}H-NMR($CDCl_3$)δ: 1.54 (18H, S), 1 .85 (3H, d, J=2.4 Hz), 3.24 (1H, m), 3.44 (1H, m), 3.53 (2H, brs), 6.61 (1H, m), 6.82-6.89 (2H, m).

Reference Example 11



The 1st step: A solution of 2.0M lithium diisopropylamide/n-heptane/ethyl benzene (182 ml) in tetrahydrofuran (150 ml) was cooled in a dry ice-acetone bath, and a solution of methyl isobutyrate (27.17 g) in tetrahydrofuran (90 ml) was added dropwise with stirring. After stirring for 40 minutes, a solution of chlorotitanium triisopropoxide (97.07 g) in tetrahydrofuran (300 ml) was added dropwise. After stirring for 15 minutes, a solution of compound (86)(25.39 g) in tetrahydrofuran (150 ml) was added dropwise. After the reaction for 2.5 hours, the reaction mixture was poured portionwise into an aqueous solution of ammonium chloride stirred under ice cooling and the formed insoluble materials were filtered. It was extracted with ethyl acetate, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (87)(23.98 g). ^{1}H-NMR ($CDCl_3$) δ: 1.12 (3H, s), 1.22 (3H, s), 1.35 (9H, s), 1.99 (3H, d, J = 5.8 Hz), 3.75 (3H, s), 5.65 (1H, s), 7.20 (0H, dd, J = 11.5, 8.9 Hz), 8.18-8.21 (1H, m), 8.45 (1H, dd, J = 6.9, 2.9 Hz).

The 2nd step: Compound (87)(391 mg) was dissolved in tetrahydrofuran (4 ml) and lithium borohydride (44 mg) was added in 3 minutes in a nitrogen stream with stirring at room temperature. After stirring for 2 hours, lithium borohydride (22 mg) was further added and the stirring was continued. After stirring for 2 hours, a saturated aqueous

solution of ammonium chloride was slowly added to the reaction solution with stirring under ice cooling, extracted with ethyl acetate 5 minutes later, the organic layer was washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (88)(175 mg).

^{1}H-NMR(DMSO-d_6)δ: 0.65 (3H, d, J=1.8Hz), 0.93 (3H, s), 1.22 (9H, s), 1.93 (3H, d, J=6.6Hz), 3.24 (1H, d, J=8.4Hz), 3.74 (1H, d, J=8.4Hz), 5.96 (1H, bs), 6.75 (1H, s), 7.47 (1H, dd, J=9.0, 12.0Hz), 8.23 (1H, ddd, J=3.0, 3.0, 9.0Hz), 8.39 (1H, dd, J=3.0, 6.9Hz).

The 3rd step: Compound (88)(331 mg) was dissolved in methanol(1.5 ml), and a hydrogen chloride-methanol solution (5-10%)(3 ml) was added with stirring at room temperature. After stirring for 1.5 hours, the reaction solvent was evaporated under reduced pressure. The residue was dissolved in ethyl acetate-methanol(9:1), poured into ice water, and a saturated aqueous solution of sodium bicarbonate (4 ml) was added, extracted , washed with saturated brine and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the powder obtained by the addition of n-hexane to the solid was filtered to give a compound (89)(207 mg).

1H-NMR(DMSO-d_6) δ: 0.80 (6H, s), 1.59 (3H, d, J=4.5Hz), 3.16 (1H, d, J=10.8Hz), 7.38 (1H, dd, J=9.0, 12.0Hz), 8.17 (1H, ddd, J=3.0, 3.0, 9.0Hz), 8.64 (1H, dd, J=3.0, 6.9Hz)

The 4th step: Compound (89)(150 mg) was dissolved in acetone (3 ml) and benzoyl isothiocyanate (0.079 ml) was added in a nitrogen stream with stirring under ice cooling. After stirring for 2 hours, the reaction solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (90)(236 mg).

LCMS: 420 m/z$[M+H]^+$

The 5th step: Compound (90)(233 mg) was dissolved in dichloromethane (4 ml) and chloropropenylamine (0.081 ml) was added at once in a nitrogen stream with stirring at room temperature. After stirring for 23 hours, the reaction solution was poured into ice water, extracted with ethyl acetate, washed with saturated brine and the organic layer was dried

over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (91)(128 mg).

^{1}H-NMR(DMSO-d_6)δ: 0.83 (3H, s), 1.12 (3H, s), 1.72 (3H, s), 2.69 (1H, d, J=13.2Hz), 2.90-3.10(1H, m), 7.44-7.58 (4H, m), 8.00 (2H,d, J=7.5Hz), 8.23-8.35 (2H, m), 10.75 (1H, bs).

The 6th step: Compound (91)(20 mg) was suspended in 99.5% ethanol(0.4 ml), 6N hydrochloric acid(0.2 ml) was added and the mixture was stirred in a oil bath heated to 90°C. After stirring for 17 hours, the reaction solution was poured into water, and extracted with ethyl acetate. The aqueous layer was made alkaline by the addition of a saturated aqueous solution of potassium carbonate (pH=11), extracted with ethyl acetate, washed with saturated brine and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (92)(14 mg).

^{1}H-NMR(DMSO-d_6)δ: 0.72 (3H, s), 1.00 (3H, d, J=3.6Hz), 1.54 (3H, d, J=4.8Hz), 2.61 (1H, d, J=12.3Hz), 3.09 (1H, d, J=12.3Hz), 5.98 (2H, s), 7.41 (1H, dd, J=9.0, 11.7Hz), 8.16-8.21 (1H, m), 8.42 (1H, dd, J=3.0, 6.9Hz).

The 7th step: Compound (92)(12 mg) was dissolved in dichloromethane (0.1 ml) and a di-tert-butyldicarbonate-dichloromethane solution (0.0966M, 1.2 ml) was added with stirring at room temperature. After stirring for 2 hours, the reaction solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (93)(15 mg).


^{1}H-NMR(DMSO-d_6)δ: 0.70 (3H, s), 1.02 (3H, s), 1.43 (9H, s), 1.56 (3H, bs), 2.61 (1H, d, J=12.9Hz), 3.16 (1H, m), 7.45 (1H, dd, J=9.0, 11.4Hz), 8.20-8.24 (1H, m), 8.35 (1H, m), 9.87 (1H, bs).

The 8th step: Methanol(4.1 ml) was added to compound (93)(823 mg), suspended, and 10% Pd-C(50% wet)(412 mg) was added. A catalytic reduction was carried out at normal pressure, and methanol(8.2 ml) was added when a solid was precipitated and the reduction was further continued. After 23 hours, the catalyst was filtered through a Celite

pad, washed with warm methanol, and the washings were combined. The solvent was evaporated under reduced pressure and the powder precipitated by the addition of diisopropylether to the residue was filtered to give compound (94)(638 mg).

^{1}H-NMR(DMSO-d_6)δ: 0.87 (3H, bs), 1.06 (3H, bs), 1.39 (9H, s), 1.57 (3H, bs), 2.66-2.72 (2H, m), 4.97 (2H, bs), 6.45-6.47 (2H, m), 6.78 (1H, m), 9.65 (1H, bs).

Reference Example 12



(104) (105) (106)


The 1st step: 3-(Trifluoromethyl)-1H-pyrazole (591 mg) was dissolved in dimethylformamide (7 ml), potassium carbonate (601 mg) and compound (104)(500 mg) were added thereto and stirred at room temperature overnight. The reaction was quenched by an addition of water. The insoluble materials were filtered and washed with diisopropylether. The resulted solid was dried under reduced pressure to give compound (105)(644 mg).

^{1}H-NMR ($CDCl_3$) δ: 4.08 (3H, s), 6.81 (1H, d, J = 2.5 Hz), 8.65 (1H, s), 9.14 (1H, s), 9.45 (1H, s).


The 2nd step: Compound (105)(640 mg) was added to a mixed solvent of water-methanol(6 ml, 1:1), lithium hydroxide (84 mg) was added and the mixture was stirred at room temperature for 4 hours. The reaction solution was acidified by the addition of 2N hydrochloric acid, the insoluble materials were filtered off and washed with diisopropylether. The resulted solid was dried under reduced pressure to give compound (106)(343 mg).

^{1}H-NMR (DMSO-d_6) δ: 7.20 (1H, d, J = 2.5 Hz), 8.93 (1H, s), 9.12 (1H, s), 9.33 (1H, s).

Reference Example 13



(107) (108) (109)



(110) (111) (112)

The 1st step: A mixture of compound (107)(1000 mg), dioxane (2 ml), and 28% ammonia water (2 ml) was stirred at 50°C for 19 hours. The reaction solution was concentrated under reduced pressure. Water was added to the residue, extracted with ethyl acetate and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give a compound (108)(476 mg).

^{1}H-NMR ($CDCl_3$) δ: 1.63 (9H, s), 5.04 (2H, br s), 8.03 (1H, s), 8.69 (1H, s).

The 2nd step: 3-Bromo-2-oxopropanoic acid ethyl ester (1582 mg) was added to compound (108)(475 mg) in dimethoxyethane (4 ml) and the mixture was stirred at 75°C for 2.5 hours. The reaction solution was diluted with diisopropylether, the insoluble materials were filtered, washed with diisopropylether and hexane, and dried under reduced pressure. The residue was stirred in tert-butyl alcohol(7.5 ml) at 95°C for 2 hours. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (109)(709 mg).

^{1}H-NMR ($CDCl_3$) δ: 1.46 (3H, t, J = 7.1 Hz), 1.66 (9H, s), 4.50 (2H, q, J = 7.1 Hz), 8.35 (1H, s), 8.89 (1H, s), 9.24 (1H, s).

The 3rd step: A mixture of compound (09)(270 mg), dioxane (3 ml) and 28% ammonia water (2.5 ml) was stirred in a pressure bottle at 50°C for 6 hours. The reaction solution was concentrated under reduced pressure to give a crude product of compound (110)(249 mg).


^{1}H-NMR of the crude product ($CDCl_3$) δ: 1.67 (9H, s), 5.79 (1H, br s), 8.35 (1H, s), 8.90 (1H, s), 9.15 (1H, s).

The 4th step: 2,2,2-Trichloroacetyl chloride (253 mg) was added at 0°C to a mixture

of compound (10)(146 mg), triethylamine (282 mg) and dimethylaminopyridine (6.8 mg) in tetrahydrofuran (9 ml), and the mixture was stirred at room temperature for 2 hours. The reaction solution was diluted with ethyl acetate and the reaction was quenched by the addition of a saturated aqueous solution of sodium bicarbonate. It was extracted with ethyl acetate, dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced pressure to give compound (111)(99 mg) as a crude product.

The 5th step: Compound (111)(95 mg) was dissolved in chloroform (3 ml), trifluoroacetic acid (1330 mg) was added and the mixture was stirred at room temperature for 4 hours. The reaction solution was concentrated under reduced pressure to give a crude product. The residue was suspended with ethyl acetate and diisopropylether and the insoluble materials were filtered and washed with diisopropylether. The residue was dried under reduced pressure to give a composition including compound (112).

Reference Example 14



The 1st step: A 2.6M n-butyl lithium/hexane solution (9.38 ml) was added dropwise for 10 minutes to diisopropylamine (2.75 g) dissolved in tetrahydrofuran (25 ml) under stirring in a dry ice/acetone bath. After stirring in a ice bath for 10 minutes and in a dry ice/acetone bath for 10 minutes, tert-butyl α-benzyloxyacetate (5.21 g) dissolved in tetrahydrofuran (25 ml) was added dropwise for 30 minutes. After stirring for 40 minutes,

chlorotitaniumtriisopropoxide (6.60 g) dissolved in tetrahydrofuran (50 ml) was added dropwise. After stirring for 30 minutes, compound (73)(2.68 g) dissolved in tetrahydrofuran (50 ml) was added dropwise for 10 minutes and stirred for 90 minutes. A suspension of ammonium chloride (7.52 g) in tetrahydrofuran-water (1:1, 40 ml) was stirred at room temperature, and the reaction mixture was added thereto at once and the precipitated insoluble materials were filtered. The filtrate was extracted with ethyl acetate, the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (4.49 g).

^{1}H-NMR($CDCl_3$)δ : 1.14 (3.6H, s), 1.22 (3.6H, s),1.27 (5.4H, s), 1.39 (5.4H, s), 1.96 (1.2H, s), 1.99 (1.8H, s), 4.31 (0.4H, s), 4.34 (0.6H, d, J=11.6Hz), 4.41 (0.4H, d, J=11.6Hz), 4.45 (0.6H, s), 4.56 (0.4H, s), 4.68 (0.6H, d, J=11.6Hz), 4.81 (0.4H, d, J=11.6Hz), 5.01 (0.6H, s), 7.06-7.38 (6H, m), 8.18 (0.6H, d, J=8.8Hz), 8.24 (0.4H, d, J=9.1Hz), 8.42-8.47 (1H, m).

The 2nd step: Compound (113)(4.49 g) was dissolved in trifluoroacetic acid (44 ml), stirred at room temperature for an hour and the solvent was evaporated under reduced pressure. The resulted residue was dissolved in 10% hydrochloric acid-methanol (44 ml), stirred at room temperature overnight and the reaction solution was concentrated under reduced pressure. The residue was dissolved in tetrahydrofuran (22 ml) and a solution of 1M-borane-tetrahydrofuran complex in tetrahydrofuran (44.1 ml) was added dropwise for 15 minutes under ice cooling, and the mixture was stirred at room temperature for 2.5 hours. Water (50 ml) was added therein with stirring under ice cooling, stirred for 15 minutes and ethyl acetate (50 ml) and potassium carbonate (16 g) were added. It was extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the obtained compound (114)(3.27 g) was used in the next step without purification.

The 3rd step: Benzoyl isothiocyanate (1.41 ml) was added to compound (114)(3.27 g) in methylenechloride (16.5 ml), stirred at room temperature for an hour and the solvent was evaporated under reduced pressure. The residue was purified with a silicagel column

chromatography to give compound (115)(3.14 g).

^{1}H-NMR($CDCl_3$)δ: 2.14 (1.35H, s), 2.21 (1.65H, s), 3.73-4.07 (3H, m), 4.43 (0.55H, d, J=11.5 Hz), 4.63 (0.55H, d, J=11.5Hz), 4.74 (0.45H, d, J=11.5Hz), 4.78 (0.45H, d, J=11.5Hz), 7.20-7.38 (4H, m), 7.43-7.51 (2H, m), 7.56-7.63 (1H, m), 7.75-7.86 (2H, m), 8.08-8.17 (1H, m), 8.24-8.34 (1H, m), 8.91-9.01 (1H, m), 11.81 (0.55H, s), 11.90 (0.45H, s).

The 4th step: α-Chlorotetramethylenamine (1.67 ml) was added to compound (115)(3.14 g) in methylenechloride (15.5 ml), stirred at room temperature for 30 minutes and pH was adjusted to over 11 by the addition of water (15 ml) and potassium carbonate. It was extracted with chloroform and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (116)(2.66 g).

^{1}H-NMR($CDCl_3$)δ: 1.58 (3bH, s), 1.81 (3aH, s), 2.76 (bH, dd, J=13.4, 1.8Hz), 3.09 (bH, dd, J=13.4, 6.1Hz), 3.16 (aH, dd, J=13.8, 3.9Hz), 3.35 (aH, dd, J=13.8, 1.8Hz), 4.21-4.25 (aH, m), 4.28 (aH, d, J=12.4Hz), 4.33-4.38 (bH, m), 4.49-4.56 (a+bH, m), 4.73 (bH, d, J=11.9Hz), 6.83-7.60 (10H, m), 7.91-8.23 (3H, m), 8.25-8.30 (bH, m), 8.74 (aH, m).

The 5th step: Hydrazine monohydrate (0.73 ml) was added to compound (116)(1.44 g) in ethanol(7.2 ml) and stirred at room temperature for 2 hours. Water was added, extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the obtained compound (117)(1.14 g) was used in the next step without purification.

The 6th step: A solution of di-tert-butyl dicarbonate (1.65 g) in methylenechloride (5.5 ml) and 4-dimethylaminopyridine (37 mg) were added to compound (117)(1.14 g) in methylenechloride (5.5 ml) and stirred at room temperature for an hour. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (118)(1.52 g).


^{1}H-NMR($CDCl_3$)δ: 1.47 (8.1H, s), 1.51 (9.9H, s), 1.53 (1.35H, s), 1.75 (1.65H, s), 3.01-3.49 (2H, m), 3.81-3.86 (0.55H, m), 4.07-4.09 (0.45H, m), 4.17 (0.45H, d, J=12.1Hz), 4.25 (0.55H, d, J=11.6Hz), 4.41 (0.45H, d, J=12.1Hz), 4.49 (0.55H, d, J=11.6Hz), 6.73-6.78 (1H,

m), 6.94-7.23 (5H, m), 8.11-8.18 (1H, m), 8.22-8.27 (0.55H, m), 8.51-8.55 (0.45H, m).

The 7th step: 20w/w% Palladium hydroxide supported by carbon (40 mg) was added to a solution of compound (118)(211.3 mg) in ethanol (2 ml), stirred in a hydrogen atmosphere of 1 atom at room temperature for 22 hours and filtered through a Celite pad. The filtrate was concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (119)(149.1 mg).

^{1}H-NMR($CDCl_3$)δ: 1.46 (8.1H, s), 1.51 (9.9H, s), 1.54 (1.35H, s), 1.74 (1.65H, s), 2.91 (0.55H, d, J=12.9Hz), 3.02 (0.55H, dd, J=12.9, 6.3Hz), 3.15 (0.45H, dd, J=13.3, 3.0Hz), 3.37-3.73 (2H, br), 3.43 (0.45H, d, J=13.3Hz), 4.13-4.18 (1H, m), 4.22 (0.45H, d, J=11.9Hz), 4.34 (0.45H, d, J=11.9Hz), 4.49 (0.55H, d, J=11.6Hz), 4.59 (0.55H, d, J=11.6Hz), 6.45-6.61 (1H, m), 6.71-7.39 (7H, m).

Example 1 Preparation of compound 46




(34) (33) 46

Compound (34)(125 mg) and DMT-MM (162 mg) were suspended in methanol (1.2 ml), stirred at room temperature for 30 minutes and compound (33)(117 mg) was added therein. After stirring for 5 hours, the product was isolated by a silicagel thin-layer chromatography to give the objective compound (46)(13.5 mg).

^{1}H-NMR (DMSO-d_6) δ: 1.47 (3H, s), 1.81 (1H, d, J = 11.6 Hz), 2.12 (1H, bs), 2.54-2.59 (1H, m), 2.97 (1H, bs), 3.28 (2H, d, 6.4 Hz), 3.52 (2H, d, 6.5 Hz), 3.88 (3H, s), 5.69 (2H, s), 7.09 (1H, dd, J = 11.8, 6.8 Hz), 7.50 (1H, s), 7.60 (1H, d, J = 7.8 Hz), 7.67 (1H, s), 10.06 (1H, s).

Example 2 Preparation of compound 86




The 1st step: Sodium hydride (302 mg) is added to DMF (3.0 ml) and 3-butene alcohol (3.0 ml) under ice cooling in a nitrogen atmosphere. After stirring at room temperature for 1.0 hour, compound (35)(300 mg) was added and stirred under heating at 65°C. After 7 hours, the reaction solution was neutralized by the addition of 2M hydrochloric acid and concentrated under reduced pressure. Water was added to the resulted residue and filtered to give compound (36)(87 mg, 23.7 %).

The 2nd step: Compound (36)(65.8 mg) and compound (37)(50 mg) were dissolved in methanol (2.0 ml), DMT-MM (93.7 mg) was added and the mixture was stirred at room temperature. After 6 hours, the solvent was evaporated under reduced pressure and the residue was purified with a column chromatography using chloroform/methanol to give compound (86)(40 mg, 44.5 %).

^{1}H-NMR (DMSO-d_6) δ: 1.65 (3H, s), 2.03-2.09 (1H, m), 2.34-2.38 (1H, m), 2.51-2.61 (2H, m), 3.10-3.13 (1H, m), 3.57 (2H, t, J = 4.4 Hz), 4.45 (2H, t, J = 6.4 Hz), 5.13 (2H, dd, J = 29.1, 13.9 Hz), 5.83-5.92 (1H, m), 7.08 (1H, d, J = 7.8 Hz), 7.40 (1H, t, J = 8.0 Hz), 7.84 (1H, s), 7.91 (1H, d, J = 8.1 Hz), 8.36 (1H, s), 8.87 (1H, s), 10.56 (1H, s).


Example 3



A carboxylic acid, R-COOH corresponding to the objective compound (0.115 mmol) and 4-(4,6-dimethoxy-1,3,5-triazin-2-yl)-4-methylmorpholinium chloride (0.106 mmol) were dissolved in methanol (0.4 ml) and stirred by shaking at room temperature for 1.5 hours. A solution of compound A (0.0884 mmol) in methanol (0.4 ml) was added and

the mixture was stirred for 8 hours. The reaction solvent was concentrated, dissolved in ethyl acetate (1 ml) and dimethylsulfoxide (0.5 ml), a 2N aqueous solution of sodium hydroxide (1 ml) was added and stirred by shaking for 2 hours. The organic layer was separated and concentrated to give a crude product of compound B. Trifluoroacetic acid(0.3 ml) was added and stirred by shaking at room temperature for 14 hours, dimethylsulfoxide (0.4 ml) was added and the product was purified with preparative LC/MS to give the objective compound C.

Example 4 Preparation of compound 668



(82) (83) (84)

(85) **668**

The 1st step: Compound (82)(506 mg) was dissolved in chloroform (30.0 ml), an aqueous solution (10.0 ml) of sodium bicarbonate (851 mg) and thiophosgene (0.111 ml) were added and stirred under ice cooling for 40 minutes. The organic layer was separated from the reaction solution and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (83)(457 mg).

^{1}H-NMR(DMSO-d_6)δ: 1.44 (9H, s), 1.51 (3H, s), 1.60 (1H, s), 2.17 (1H, s), 2.68 (1H, s), 3.05 (1H, s), 7.30 (1H, t, J= 10.1 Hz), 7.42 (1H, s), 7.58 (1H, s).

The 2nd step: Compound (83)(240 mg) was dissolved in methylelechloride (3.60 ml), pyridine-2-ylmethanamine (74.8 mg) and triethylamine (0.192 ml) were added and the mixture was stirred at room temperature for 40 minutes. The reaction solution was washed with distilled water, the separated organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (84)(210 mg).

^{1}H-NMR(DMSO-d_6)δ: 1.46 (9H, s), 1.65 (3H, s), 2.05 (2H, s), 2.57 (1H, s), 2.97 (1H, s), 4.86 (2H, s), 7.29 (3H, m), 7.40 (1H, d, J=7.3 Hz), 7.66 (1H, s), 7.84 (1H, s), 8.27 (1H, s),

8.58 (1H, s), 9.96 (1H, s).


The 3rd step: Compound (84)(95.1 ml) was dissolved in toluene (1.50 ml), dicyclohexylcarbodiimide (40.1 mg) was added and the mixture was stirred under irradiation of microwave at 100°C for 20 minutes. The reaction solution was concentrated under reduced pressure and the residue was purified with a column chromatography to give compound (85)(38.0 mg).

^{1}H-NMR(DMSO-d_6)δ: 1.40 (9H, s), 1.61 (3H, s), 1.94 (2H,s), 2.57 (1H,s), 2.88 (1H, s), 6.55 (1H, d, J= 6.3 Hz), 6.59 (1H, d, J= 8.6 Hz), 7.07 (1H, d, 8.6 Hz), 7.13 (2H, s), 7.29 (1H, s), 7.41 (1H, s, J= 9.3 H), 7.96 (1H, d, J= 6.8 Hz), 8.85 (1H, s).

The 4th step: Compound (85)(38.0 mg) was dissolved in chloroform (0.50 ml), trifluoroacetic acid (1.00 ml) was added and stirred at room temperature for 2 hours. The reaction solution was extracted with a mixture of chloroform/methanol and washed with an aqueous solution of potassium carbonate and distilled water. The separated organic layer was dried over anhydrous sodium sulfate and the solvent was evaporated under reduced pressure. Diisopropylether was added to the residue and the precipitated powder was filtered to give compound (668)(9.39 mg).

^{1}H-NMR(DMSO-d_6)δ: 1.58 (3H, s), 1.90 (1H, s), 2.44 (1H, s), 2.62 (1H, t, J= 9.7 Hz), 3.06 (1H, s), 6.57 (2H, td, J-15.0, 6.3Hz), 7.05 (1H, dd, J= 12.1, 10.6 Hz), 7.15 (1H, s), 7.24 (1H, d, J= 5.3 Hz), 7.31 (1H, dd, J= 7.7, 3.7 Hz), 7.42 (1H, d, J- 8.8 Hz), 7.99 (1H, d, J= 6.8 Hz), 8.85 (1H, s).

Example 5 Preparation of compound 674



(73) (95) (96) (97) (98) (99) (100) (101) (102)

(103) 674

The 1st step: A solution of 2.0M lithium diisopropylamide/n-heptane/ethylbenzene (172 ml) in tetrahydrofuran (280 ml) was cooled in a dry ice/acetone bath and a solution of tert-butyl cyclopropancarboxylate (36.6 g) in tetrahydrofuran (60 ml) was added dropwise with stirring. After stirring for an hour, a solution of chlorotitanium tirisopropoxide (92 g) in tetrahydrofuran (190 ml)was added dropwise, stirred for 10 minutes and a solution of compound (73)(24.56 g) in tetrahydrofuran (120 ml) was added dropwise. After reaction for 2 hours, the reaction solution was added portionwise to an aqueous solution of ammonium chloride with stirring under ice cooling and the precipitated insoluble materials were filtered. It was extracted with ethyl acetate, the organic layer was dried over anhydrous sodium sulfate.

The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (95)(15.49 g).

^{1}H-NMR ($CDCl_3$) δ: 1.16-1.19 (2H, m), 1.24 (9H, s), 1.28-1.32 (2H, m), 1.36 (9H, s), 1.46 (3H, s), 1.50-1.55 (2H, m), 1.64-1.72 (2H, m), 5.45 (1H, s), 7.11-7.16 (1H, m), 8.11-8.16 (1H, m), 8.67 (1H, dd, J = 6.9, 2.9 Hz).

The 2nd step: 2.0 M Hydrochloric acid/ethyl acetate (30 ml) was added to compound (95)(2.48 g) and stirred at 65°C for 5.5 hours. Diisopropylether was added and the precipitated solid was filtered to give a crude product of compound (96)(1.66 g).

The 3rd step: A solution of compound (96)(1.66 g) in tetrahydrofuran (8.3 ml) was stirred under ice cooling and a solution of 1M borane/tetrahydrofuran (21.8 ml) was added and the mixture was stirred at room temperature for 2 hours and 45 minutes. Ice and sodium bicarbonate were added, extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give a crude product of compound (97)(1.36 g).

The 4th step: A solution of compound (97)(1.36 g) in acetone (20 ml) was stirred under ice cooling, a solution of benzoyl isothiocyanate (0.92 g) in acetone (6 ml) was added and the mixture was stirred for 40 minutes. After the addition of water, the reaction solution was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (98)(1.68 g).

^{1}H-NMR ($CDCl_3$) δ: 0.67-0.73 (3H, m), 0.84-0.88 (1H, m), 1.73 (1H, t, J = 5.6 Hz), 2.29 (3H, d, J = 2.0 Hz), 3.44 (1H, dd, J = 12.2, 5.1 Hz), 3.82 (1H, dd, J = 12.2, 5.1 Hz), 7.14 (1H, dd, J = 11.0, 9.0 Hz), 7.52 (2H, t, J = 7.6 Hz), 7.63 (1H, t, J = 7.6 Hz), 7.87 (2H, d, J = 7.6 Hz), 8.17 (1H, ddd, J = 9.0, 3.9, 2.9 Hz), 8.27 (1H, dd, J = 6.8, 2.9 Hz), 8.82 (1H, s), 11.75 (1H, s).

The 5th step: Compound (98)(1.68 g) was dissolved in dichloromethane (17 ml), stirred under ice cooling and 1-chloro-N,N,2-trimethyl-1-propenylamine (0.60 g) was added. After stirring at room temperature for an hour, water was added, the reaction solution was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified

with a silicagel column chromatography to give compound (99)(1.34 g).

^{1}H-NMR ($CDCl_3$) δ: 0.77-0.82 (1H, m), 0.95-1.07 (2H, m), 1.38-1.40 (1H, m), 1.52 (3H, d, *J* = 1.1 Hz), 2.25 (1H, d, *J* = 13.0 Hz), 3.05 (1H, d, *J* = 13.0 Hz), 7.27 (1H, dd, *J* = 10.8, 8.9 Hz), 7.40-7.54 (3H, m), 8.18-8.27 (3H, m), 8.36 (1H, dd, *J* = 6.7, 2.7 Hz).

The 6th step: Hydrazine monohydrate (038 g) was added to a solution of compound (99)(1.00 g) in ethanol (10 ml) under stirring at room temperature. After stirring for 4 hours, it was stirred under heating at 50°C for 2 hours. The reaction solution was concentrated under reduced pressure, water was added, and the mixture was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give compound (100)(0.69 g) as a crude product.

The 7th step: A mixture of compound (100)(0.91 g), di-tert-butyldicarbonate (1.55 g), 4-dimethylaminopyridine (0.04 g) and tetrahydrofuran (9.1 ml) was stirred at room temperature for 1 hour and 15 minutes. Water was added, the reaction solution was extracted with ethyl acetate and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (101)(1.28 g).

^{1}H-NMR ($CDCl_3$) δ: 0.37-0.41 (1H, m), 0.50-0.54 (1H, m), 0.68 (2H, t, *J* = 7.7 Hz), 1.56 (18H, s), 1.78 (3H, d, *J* = 4.0 Hz), 2.35 (1H, d, *J* = 12.7 Hz), 3.57 (1H, dd, *J* = 12.7, 1.8 Hz), 7.12-7.21 (1H, m), 8.15 (1H, ddd, *J* = 8.9, 3.9, 3.0 Hz), 8.39 (1H, dd, *J* = 6.7, 3.0 Hz).

The 8th step: Compound (101)(1.28 g) was dissolved in ethyl acetate (13 ml), 10% Pd-C(0.64 g) was added and the mixture was stirred at room temperature for 13 hours and 30 minutes. The insoluble materials were filtered, the filtrate was concentrated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (102)(1.07 g).

^{1}H-NMR ($CDCl_3$) δ: 0.51-0.58 (2H, m), 0.81-0.86 (2H, m), 1.54 (18H, s), 1.64 (3H, d, *J* = 3.0 Hz), 2.60 (1H, d, *J* = 12.4 Hz), 3.08 (1H, d, *J* = 12.4 Hz), 3.50 (2H, s), 6.51 (1H, ddd, *J* = 8.6, 3.7, 3.0 Hz), 6.78-6.84 (2H, m), 7.18-7.21 (1H, m).


The 9th step: A soluiton of 5-methylpyrazine-2-carboxylic acid(59 mg) in

N,N-dimethylformamide (1.5 ml) was stirred under ice cooling, and 1–[bis(dimethylamino)methylene]-1H-1,2,3-triazolo(4,5-b)pyridinium-3-oxide hexafluorophosphate (196 mg) and triethylamine (61 mg) were added and the mixture was stirred for 10 minutes. A solution of compound (102)(200 mg) in N,N-dimethylformamide (3 ml) was added and the mixture was stirred at room temperature for 4 hours. Water was added, extracted with ethyl acetate and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (103)(170 mg).

The 10th step: Compound (103)(170 mg) was dissolved in dichloromethane (0.75 ml), stirred under ice cooling, trifluoroacetic acid(0.75 ml) was added and the mixture was stirred at room temperature for 3 hours. After concentration of the reaction solution under reduced pressure, ice water was added, potassium carbonate was added with stirring under ice cooling and extracted with ethyl acetate. The organic layer was dried over anhydrous sodium sulfate, the solvent was evaporated under reduced pressure and ether/hexane was added to the residue. The precipitated solid was filtered to give compound (674)(104 mg)

^{1}H-NMR ($CDCl_3$) δ: 0.53-0.59 (1H, m), 0.65-0.72 (1H, m), 0.85-0.91 (1H, m), 1.14-1.17 (1H, m), 1.47 (3H, d, J = 2.0 Hz), 2.46 (1H, d, J = 12.1 Hz), 2.69 (3H, s), 2.89 (1H, dd, J = 12.1, 1.3 Hz), 7.06 (1H, dd, J = 11.5, 8.8 Hz), 7.45 (1H, dd, J = 6.8, 2.8 Hz), 7.94 (1H, ddd, J = 8.8, 4.0, 2.8 Hz), 8.44 (1H, d, J = 1.3 Hz), 9.36 (1H, d, J = 1.3 Hz), 9.60 (1H, s).

Example 6 preparation of compound 687




(120) 687

The 1st step: Dichloromethane-trifluoroacetic acid (1:1, 1 ml) was added to compound (120)(49 mg) and stirred at room temperature for an hour. The reaction solution was concentrated under reduced pressure, dimethyl sulfoxide-acetic anhydride (1:1, 1 ml) was added to the residue, stirred at 50°C for 1.5 hours and the solvent was evaporated under

reduced pressure. Hydrochloric acid (1M, 0.5 ml) was added to the residue and stirred at 50°C for 1 hours. A saturated aqueous solution of sodium bicarbonate was added, extracted with chloroform, and the organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography. A mixture of chloroform-diethyl ether/ethyl acetate was added and the precipitated slid was filtered to give compound 687(17 mg)

^{1}H-NMR($CDCl_3$)δ: 1.78 (3H, s), 3.52 (1H, d, J=15.1Hz), 3.73 (1H, d, 15.1Hz), 7.06 (1H, dd, J=10.4, 8.6Hz), 7.73 (1H, dd, J=6.6, 1.3Hz), 7.82-7.86 (1H, m), 7.90 (1H, d, J=1.3Hz), 8.48 (1H, d, J=1.3Hz), 9.79 (1H, s).

Example 7 Preparation of compound 680,681 and 682



The 1st step: In a nitrogen atmosphere, diisopropylamine (20.3 ml) and tetrahydrofuran (83.5 ml) were added and cooled to -60°C in an acetone/dry ice bath and a solution of 1.63M n-butyl lithium/n-hexane was added dropwise while stirring was continued until the temperature rose to 0°C. After stirring for 30 minutes, the reaction

solution was cooled to -60°C in an acetone/dry ice bath and a solution of tert-butyl acetate (16.8 ml) in tetrahydrofuran (22.2 ml) was added dropwise with stirring. After stirring for 45 minutes, a solution of compound (121)(11.1 g) in tetrahydrofuran (22.2 ml) was added dropwise. After 2.5 hours, a saturated aqueous solution of ammonium chloride (100 ml) was stirred under ice cooling and the reaction solution was poured portionwise therein, extracted with ethyl acetate, and the organic layer was dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (122)(8.83 g).

^{1}H-NMR ($CDCl_3$) δ: 1.32 (17H, s), 1.93 (3H, s), 3.15 (1H, d, J = 16.4 Hz), 3.66 (1H, d, J = 16.2 Hz), 5.50 (1H, s), 8.12 (1H, s), 8.36 (1H, s).

The 2nd step: Lithium aluminium hydride (0.76 g) and tetrahydrofuran (15 ml) were cooled in an ice-salt bath in a nitrogen atmosphere and a solution of compound (122)(3.03 g) in tetrahydrofuran (10 ml) was added dropwise with stirring. After stirring for 15 minutes, acetone (4 ml) was added, insoluble materials were filtered, extracted with ethyl acetate, and the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give compound (123)(2.30 g) as a crude product.

MS: 385 m/z$[M+H]^+$

The 3rd step: 10% Hydrochloric acid/methanol solution (30 ml) was added to compound (123)(2.2 g) and stirred at room temperature for 1.5 hours. The solvent was evaporated under reduced pressure, the residue was basified with a 2.0M aqueous solution of potassium carbonate and extracted with ethyl acetate (100 ml). The organic layer was washed with water, dried over anhydrous magnesium sulfate and the solvent was evaporated under reduced pressure. Toluene (30 ml) and water (15 ml) were added to the resulted crude product(2.25 g), cooled in a ice bath and potassium carbonate (1.58 g) and thiophosgene (0.656 ml) were added with stirring. After stirring at room temperature for 30 minutes, the reaction solution was extracted with toluene, the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under

reduced pressure, toluene (30 ml), thionyl chloride (1.25 ml) and N,N-dimethylformamide (0.044 ml) were added to the resulted residue and the mixture was stirred under heating at 80°C for 1.5 hours. The solvent was evaporated under reduced pressure, ice water was added, extracted with ethyl acetate, and the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (124)(1.26 g).

^{1}H-NMR ($CDCl_3$) δ:1.56 (3H, s), 2.54-2.64 (1H, m), 3.07-3.17 (1H, m), 3.29-3.38 (1H, m), 3.50-3.57 (1H, m), 8.13 (1H, d, J = 2.4 Hz), 8.44 (1H, d, J = 2.4 Hz).

The 4th step: Tetrahydrofuran (12.6 ml) and 28% ammonia water (6.3 ml) were added to compound (124)(1.26 g) and stirred at room temperature for 1.5 hours. The reaction solution was extracted with ethyl acetate, the organic layer was washed with saturated brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to give compound (125)(1.13 g) as a crude product.

^{1}H-NMR ($CDCl_3$) δ: 1.70 (3H, s), 2.15-2.21 (1H, m), 2.52-2.58 (1H, m), 2.70-2.77 (1H, m), 3.05-3.11 (1H, m), 4.44 (2H, br s), 8.12 (1H, s), 8.34 (1H, s).

The 5th step: Tetrahydrofuran (11.3 ml) and di-tert-butyldicarbonate (0.89 ml) were added to compound (125)(1.13 g) and stirred at room temperature for an hour. Di-tert-butyldicarbonate (1.13 ml) and 4-dimethylaminopyridine (0.086 g) were added and further stirred at room temperature for 2 hours. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (126)(1.59 g).

^{1}H-NMR ($CDCl_3$) δ: 1.53 (18H, s), 1.73 (3H, s), 1.90-1.97 (1H, m), 2.63-2.69 (1H, m), 2.93-2.99 (1H, m), 3.21-3.28 (1H, m), 8.24 (1H, d, *J* = 2.3 Hz), 8.36 (1H, d, *J* = 2.3 Hz).

The 6th step: N, N-Dimethylformamide (40 ml) was added to compound (126)(2.00 g) in a nitrogen stream, cooled in an ice bath with stirring and sodium methoxide (2.074 g) was added therein. After stirring at room temperature for 1.5 hours, the reaction solution was warmed up to 60°C and stirred for 2 hours. It was cooled in a ice bath, neutralized by

the addition of 2N hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with 2M aqueous solution of potassium carbonate and brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (127)(1.69 g).

^{1}H-NMR ($CDCl_3$) δ: 1.52 (9H, s), 1.70 (3H, s), 1.96-2.03 (1H, m), 2.54-2.61 (1H, m), 2.80-2.85 (1H, m), 2.97-3.00 (1H, m), 3.97 (3H, s), 7.62 (1H, d, J = 1.5 Hz), 8.15 (1H, d, J = 1.5 Hz).

The 7th step: Compound (127)(1.571 g), trisdibenzylideneacetonedipalladium(0.414 g) and butynyl-1-adamantylphosphine (0.324 g) were dissolved in toluene under a nitrogen stream, and a solution of 1.6M lithium hexamethyldisilazide/tetrahydrofuran (5.66 ml) was added at room temperature with stirring. The reaction solution was warmed up to 80°C and stirred for 3 hours. Then diethyl ether and 1N hydrochloric acid were added with stirring under ice cooling. After stirring for 5 minutes, it was neutralized by the addition of a saturated aqueous solution of sodium carbonate, extracted with ethyl acetate and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (128)(1.55 g).

^{1}H-NMR ($CDCl_3$) δ: 1.52 (9H, s), 1.72 (3H, s), 1.86-1.93 (1H, m), 2.02 (2H, s), 2.52-2.59 (1H, m), 2.74-2.79 (1H, m), 3.13-3.18 (1H, m), 3.90 (3H, s), 6.96 (1H, d, J = 2.3 Hz), 7.59 (1H, d, J = 1.8 Hz).

The 8th step: Compound (128)(0.20 g), 5-methylpyridine-2-carboxylic acid(0.10 g) and O-(7-azabenzotriazole-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate (HATU)(0.28 g) were dissolved in N,N-dimethylformamide (2 ml), triethylamine (0.119 ml) was added and the mixture was stirred at room temperature for 1.0 hours. A 2M aqueous solution of potassium carbonate was added, extracted with ethyl acetate, and the organic layer was washed with brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure and the resulted residue was dissolved in chloroform(4.0

ml), trifluoroacetic acid(1.0 ml) was added and the mixture was stirred at room temperature for 3 hours. The solvent was evaporated under reduced pressure, the residue was made basic by the addition of a 2.0M aqueous solution of potassium carbonate, extracted with ethyl acetate and the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (680)(0.096 g).

1H-NMR (DMSO-d6) δ: 1.47 (3H, s), 1.77-1.83 (1H, m), 2.34-2.39 (1H, m), 2.48-2.53 (1H, m), 2.63 (3H, s), 2.89-2.96 (1H, m), 3.90 (3H, s), 5.86 (2H, br s), 8.10 (1H, d, J = 2.3 Hz), 8.47 (1H, d, J = 2.5 Hz), 8.69 (1H, s), 9.14 (1H, s), 10.69 (1H, s).

The 9th step: Compound (680)(0.096 g) and sodium iodide (0.193 g) were dissolved in acetonitrile (5.0 ml), trimethylsilylchloride (0.164 ml) was added and the mixture was stirred at room temperature for 2.5 hours. Sodium iodide (0.193 g) and trimethyl silylchloride (0.164 ml) were added and stirring was continued at room temperature for 12 hours. A 2.0M aqueous solution of potassium carbonate, extracted with ethyl acetate and the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure to give compound (681)(0.073 g) as a crude product.

^{1}H-NMR (DMSO-d_6) δ: 1.52 (3H, s), 1.80-1.85 (1H, m), 2.62 (3H, s), 2.64-2.69 (2H, m), 2.96-3.01 (1H, m), 7.77 (1H, d, J = 2.5 Hz), 7.96 (1H, d, J = 2.3 Hz), 8.67 (1H, s), 9.10 (1H, s), 10.58 (1H, s).






The 10th step: Compound (681)(0.031 g) was dissolved in tetrahydrofuran (2.0 ml), di-tert-butyldicarbonate (0.030 ml) was added and the mixture was stirred at room temperature for 1.5 hours. Di-tert-butyldicarbonate (0.030 ml) was further added and the stirring was continued at room temperature for 2.0 hours. The reaction solution was concentrated under reduced pressure, the resulted residue was dissolved in N, N-dimethylformamide (0.5 ml) and potassium carbonate (23.9 mg) was added. A solution of methyl iodide (12.2 mg) in N, N-dimethylformamide (0.5 ml) was added with stirring at room temperature. After stirring at room temperature for 3 hours, methyl iodide (11.05 mg)

was added and the mixture was stirred at room temperature for 2 hours. Brine was added, extracted with ethyl acetate and the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure, the resulted residue was dissolved in chloroform (2.0 ml), trifluoroacetic acid (0.5 ml) was added and the mixture was stirred at room temperature for 3 hours. The reaction solution was concentrated under reduced pressure, the resulted residue was made alkaline by the addition of a 2.0M aqueous solution of potassium carbonate, extracted with ethyl acetate and the organic layer was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated under reduced pressure and the residue was purified with a silicagel column chromatography to give compound (682)(4.2 mg).






^{1}H-NMR (DMSO-d_6) δ: 1.46 (3H, s), 1.95-2.01 (1H, m), 2.33-2.39 (1H, m), 2.62 (3H, s), 2.64-2.69 (1H, m), 2.74 (3H, s), 2.92-2.98 (1H, m), 7.90 (1H, d, J = 2.5 Hz), 7.94-7.95 (1H, m), 8.67 (1H, s), 9.09 (1H, s), 10.57 (1H, s).

The other compounds are prepared in the same manner. Chemical structures and physical constants are shown below.





[Table 1]

Compound No.	Structure
1	
2	
3	
4	
5	






[Table 2]

Compound No.	Structure
6	
7	
8	
9	
10	






[Table 3]

Compound No.	Structure
11	
12	
13	
14	
15	






[Table 4]

Compound No.	Structure
16	
17	
18	
19	
20	






[Table 5]

Compound No.	Structure
21	
22	
23	
24	
26	






[Table 6]

Compound No.	Structure
27	
28	
29	
30	
31	






[Table 7]

Compound No.	Structure
32	
33	
34	
35	
36	






[Table 8]

Compound No.	Structure
37	
38	
39	
40	
41	





[Table 9]

Compound No.	Structure
42	 Chiral
43	 Chiral
44	 Chiral
45	 Chiral
46	 Chiral






[Table 10]

Compound No.	Structure
47	
48	
49	
50	
51	






[Table 11]

Compound No.	Structure
52	
53	
54	
55	
56	






[Table 12]

Compound No.	Structure
57	
58	
59	
60	
61	






[Table 13]

Compound No.	Structure
62	
63	
64	
65	
66	






[Table 14]

Compound No.	Structure
67	
68	
69	
70	
71	






[Table 15]

Compound No.	Structure
72	
73	
74	
75	
76	




[Table 16]

Compound No.	Structure
77	
78	
79	
80	
81	






[Table 17]

Compound No.	Structure
82	
83	
84	
85	
86	






[Table 18]

Compound No.	Structure
87	
88	
89	
90	
91	





[Table 19]

Compound No.	Structure
92	
93	
94	
95	
96	






[Table 20]

Compound No.	Structure
97	
98	
99	
100	
101	





[Table 21]

Compound No.	Structure
102	
103	
104	
105	
106	






[Table 22]

Compound No.	Structure
107	
108	
109	
110	
111	





[Table 23]

Compound No.	Structure
112	
113	
114	
115	
116	






[Table 24]

Compound No.	Structure
117	
118	
119	
120	
121	






[Table 25]

Compound No.	Structure
122	
123	
124	
125	
126	





[Table 26]

Compound No.	Structure
127	
128	
129	
130	
131	






[Table 27]

Compound No.	Structure
132	
133	
134	
135	
136	






[Table 28]

Compound No.	Structure
137	
138	
139	
140	






[Table 29]

Compound No.	Structure
141	
142	
143	
144	
145	






[Table 30]

Compound No.	Structure
146	 Chiral
147	 Chiral
148	
149	 Chiral
150	 Chiral






[Table 31]

Compound No.	Structure
151	
152	
153	
154	
155	







[Table 32]

Compound No.	Structure
156	
157	
158	
159	
160	

[Table 33]

Compound No.	Structure
161	
162	
163	
164	
165	

[Table 34]

Compound No.	Structure
166	
167	
168	
169	
170	
171	

[Table 35]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
1			1H-NMR (CDCl3) δ : 8.69 (1.0H, br s), 8.57 (1.0H, s), 8.53-8.41 (1.0H, m), 8.36-8.33 (1.0H, m), 7.90-7.81 (1.0H, m), 7.56 (1.0H, br s), 7.41-7.29 (2.0H, m), 7.00-6.88 (2.0H, m), 2.88-2.63 (2.0H, m), 2.50-2.38 (1.0H, m), 2.06-2.00 (1.0H, m), 1.66 (3.0H,	
2	344			
3			1H-NMR (CDCl3) δ : 7.79 (1.0H, br s), 7.65-7.64 (1.0H, m), 7.48-7.41 (1.0H, m), 7.31 (1.0H, t, J = 8.01 Hz), 7.04-7.01 (1.0H, m), 6.23 (1.0H, br s), 2.93-2.65 (2.0H, m), 2.57 (3.0H, br s), 2.40 (1.0H, ddd, J = 14.11, 5.34, 3.43 Hz), 2.27 (3.0H, br s), 2.09-1.92 (1.0H, m), 1.67 (3.0H, s).	
4			1H-NMR (CDCl3) δ : 7.86-7.83 (1.0H, m), 7.45-7.42 (1.0H, m), 7.35 (1.0H, t, J = 12.96 Hz), 7.21 (1.0H, br s), 7.04-7.01 (1.0H, m), 4.23 (3.0H, s), 2.90-2.86 (1.0H, m), 2.77-2.61 (1.0H, m), 2.38-2.30 (1.0H, m), 1.99-1.89 (1.0H, m), 1.60 (3.0H, s).	
5			1H-NMR (DMSO-d6) δ : 9.81 (1.0H, br s), 7.70-7.65 (2.0H, m), 7.22 (1.0H, t, J = 7.85 Hz), 7.06-7.03 (1.0H, m), 6.53 (1.0H, s), 2.92-2.85 (1.0H, m), 2.61-2.52 (1.0H, m), 2.28 (3.0H, s), 2.02-1.97 (1.0H, m), 1.73-1.67 (1.0H, m), 1.39 (3.0H, s).	
6	425			
7	415			
8	361			
9	331			
10	347			
11	360			
12	379			
13	367			
14	331			
15			1H-NMR (DMSO-d6) δ : 10.02 (1.0H, s), 7.61-7.55 (2.0H, m), 7.25 (1.0H, t, J = 7.93 Hz), 7.09 (1.0H, d, J = 7.78 Hz), 6.26 (1.0H, s), 3.86 (3.0H, s), 2.91-2.87 (1.0H, m), 2.59-2.54 (1.0H, m), 2.00-1.96 (1.0H, m), 1.75-1.62 (1.0H, m), 1.39 (3.0H, s).	
16	404			
17	422			
18	360			
19	349			
20	349			
21	388			
22	365			
23	392			
24	385			
26			1H-NMR (MeOD) δ :1.73(1H, s), 1.98(1H, s), 2.29(1H, s), 2.76(1H, s), 6.58(1H, s), 6.79(1H, s), 6.92(1H, s), 7.13(1H, s), 8.01(1H, s), 8.55(1H, s)	
27			1H-NMR (DMSO-d6) δ : 1.65(1H, s), 2.07(1H, t, J = 13.1 Hz), 2.57(1H, d, J = 11.6 Hz), 3.10(1H, s), 7.10(1H, d, J = 7.1 Hz), 7.42(1H, t, J = 7.5 Hz), 7.72(2H, s), 7.84(1H, d, J =8.3 Hz), 7.92(1H, t, J = 9.5 Hz), 8.55(1H, s), 10.74 (1H, s)	
28	358			
29			1H-NMR (CDCl3) δ : 1.71(s3H, s), 2.00(1H, d, J =8.8 Hz), 2.45(1H, d, J = 12.4 Hz), 2.78(1H, t, J = 12.5 Hz), 2.88(aH, s, J = 13.6 Hz), 3.94(3H, s), 5.30(1H, s) 7.05(1H, d, J = 7.8 Hz), 7.35(1H, t, J = 8.2 Hz), 7.48(1H, s), 7.56(1H, s), 7.65(1H, d, J = 7.8 Hz), 9.58(1H, s)	

[Table 36]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
30	376			
31			1H-NMR (MeOD) δ :1.75(3H, s), 2.10-2.13(1H, m), 2.49-2.62(1H, m), 2.65-2.71(2H, m), 2.80(3H, s), 7.02(1H, d, J = 8.6 Hz), 7.41(1H, t, J = 7.8 Hz), 7.53(1H, s), 8.12(1H, d, J = 8.1 Hz), 9.34(2H, s), 9.79(1H, s)	
32	372			
33	373			
34			1H-NMR (DMSO-d6) δ : 1.63(3H, s), 1.70(3H, s, J =5.3 Hz), 1.99-2.02(1H, br m), 2.28(1H, s), 2.57-2.60(1H, dr m), 3.07(1H, d, J = 10.1 Hz), 4.65(2H, s), 5.70(1H, d, 14.1 Hz), 5,91(1H, d, 7.3 Hz), 7.06(1H,s), 7.36(1H, s), 7.59(1H, d, J =5.1 Hz), 7.82 (1H, s), 7.89(1H, s, J 5.1 Hz), 8.36(1H, s), 10.48(1H, s)	
35	395			
36	428			
37	412			
38	396			
39	238			
40	408			
41	428			
42	379			
43	390			
44	386			
45	420			
46	435			
47			1H-NMR (DMSO-d6) δ : 1.41 (3H, s), 1.69-1.75 (1H, m), 2.01-2.05 (1H, m), 2.55-2.58 (1H, m), 2.74-2.76 (2H, m), 2.88-2.91 (1H, m), 7.06 (1H, d, J = 7.8 Hz), 7.26 (1H, t, J = 7.8 Hz), 7.70-7.73 (2H, m), 7.91 (1H, br s), 7.99 (1H, s), 8.63 (1H, s), 9.97 (1H, s).	
48			1H-NMR (DMSO-d6) δ : 1.42 (3H, s), 1.71-1.85 (3H, m), 2.04-2.08 (1H, m), 2.56-2.58 (1H, m), 2.82 (2H, t, J = 7.2 Hz), 2.88-2.93 (1H, m), 3.41-3.43 (2H, m), 7.06 (1H, d, J = 7.8 Hz), 7.26 (1H, t, J = 7.6 Hz), 7.70-7.73 (2H, m), 7.99 (2H, s), 8.65 (1H, s), 10.00 (1H, s).	
49			1H-NMR (DMSO-d6) δ : 1.57 (3H, s), 1.62 (3H, s), 1.94-1.97 (1H, m), 2.37-2.40 (1H, m), 2.56-2.60 (1H, m), 2.80-2.82 (2H, m), 3.03-3.06 (1H, m), 7.04 (1H, d, J = 7.6 Hz), 7.34 (1H, t, J = 7.6 Hz), 7.78-7.82 (2H, m), 8.01 (1H, s), 8.06 (1H, br s), 8.64 (1H, s), 10.13 (1H, s).	
50			1H-NMR (DMSO-d6) δ : 1.45 (3H, s), 1.76-1.79 (1H, m), 2.09-2.13 (1H, m), 2.40 (3H, br s), 2.57-2.60 (1H, m), 2.83 (2H, t, J = 5.9 Hz), 2.93-2.94 (1H, m), 3.49-3.51 (2H, m), 5.74 (1H, s), 7.06 (1H, d, J = 7.6 Hz), 7.27 (1H, t, J = 7.8 Hz), 7.74 (1H, br s), 7.77 (1H, s), 7.95 (2H, br s), 8.02 (1H, s), 8.66 (1H, s), 10.02 (1H, s).	
51			1H-NMR (DMSO-d6) δ : 1.40 (3H, s), 1.70-1.75 (1H, m), 1.85-1.90 (2H, m), 1.93-2.02 (2H, m), 2.14-2.21 (1H, m), 2.53-2.60 (1H, m), 2.86-2.94 (1H, m), 3.82 (1H, q, J = 7.1 Hz), 3.98 (1H, q, J = 7.2 Hz), 4.35-4.39 (1H, m), 7.06 (1H, d, J = 7.6 Hz), 7.23 (1H, t, J = 7.8 Hz), 7.56-7.60 (2H, m), 9.57 (1H, s).	

[Table 37]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
52			1H-NMR (DMSO-d6) δ: 1.38 (3H, s), 1.61-1.81 (5H, m), 1.95-2.08 (2H, m), 2.53-2.58 (1H, m), 2.88 (4H, t, J = 6.6 Hz), 3.65-3.68 (1H, m), 5.67-5.85 (2H, m), 7.04 (1H, d, J = 7.8 Hz), 7.22 (1H, t, J = 7.8 Hz), 7.48 (1H, s), 7.59 (1H, d, J = 7.8 Hz), 9.87 (1H, s).	
53			1H-NMR (DMSO-d6) δ: 1.39 (3H, s), 1.62-1.80 (5H, m), 1.99-2.07 (2H, m), 2.52-2.58 (1H, m), 2.87-2.91 (3H, m), 3.68 (2H, dd, J = 8.7, 5.9 Hz), 7.04 (1H, d, J = 7.6 Hz), 7.23 (1H, t, J = 8.0 Hz), 7.47 (1H, s), 7.61 (1H, d, J = 7.8 Hz), 9.90 (1H, s).	
54			1H-NMR (DMSO-d6) δ: 1.40 (3H, s), 1.70-1.75 (1H, m), 1.95-2.05 (2H, m), 2.09-2.37 (2H, m), 2.52-2.57 (1H, m), 2.87-2.94 (1H, m), 4.03 (1H, q, J = 7.1 Hz), 4.19 (1H, dd, J = 8.6, 4.3 Hz), 7.06 (1H, d, J = 7.3 Hz), 7.25 (1H, t, J = 8.0 Hz), 7.49 (1H, s), 7.56 (1H, d, J = 7.8 Hz), 7.86 (1H, s), 10.01 (1H, s).	
55			1H-NMR (DMSO-d6) δ: 1.40 (3H, s), 1.69-1.75 (1H, m), 1.97-2.03 (2H, m), 2.09-2.36 (4H, m), 2.52-2.57 (1H, m), 2.87-2.93 (1H, m), 4.17-4.20 (1H, m), 7.06 (1H, d, J = 7.8 Hz), 7.25 (1H, t, J = 8.0 Hz), 7.54 (2H, t, J = 8.0 Hz), 7.87 (1H, s), 10.00 (1H, s).	
56			1H-NMR (DMSO-d6) δ: 1.40 (3H, s), 1.71-1.76 (1H, m), 2.00-2.03 (1H, m), 2.19-2.26 (1H, m), 2.46-2.58 (5H, m), 2.88-2.94 (1H, m), 5.03-5.06 (1H, m), 7.09 (1H, d, J = 7.1 Hz), 7.27 (1H, t, J = 8.1 Hz), 7.50 (1H, s), 7.56 (1H, d, J = 8.1 Hz), 10.22 (1H, s).	
57			1H-NMR (DMSO-d6) δ: 1.40 (3H, s), 1.69-1.75 (1H, m), 2.00-2.04 (1H, m), 2.19-2.27 (1H, m), 2.46-2.57 (6H, m), 2.87-2.94 (1H, m), 5.03-5.06 (1H, m), 7.09 (1H, d, J = 8.3 Hz), 7.26 (1H, t, J = 8.0 Hz), 7.51 (1H, s), 7.56 (1H, d, J = 8.3 Hz), 10.22 (1H, s).	
58			1H-NMR (DMSO-d6) δ: 1.48 (3H, s), 1.78-1.84 (1H, m), 2.11-2.18 (1H, m), 2.55-2.61 (1H, m), 2.93-2.99 (1H, m), 7.14 (1H, d, J = 7.8 Hz), 7.34 (1H, t, J = 8.0 Hz), 7.77-7.82 (2H, m), 8.93 (1H, s), 9.11 (1H, s).	
59			1H-NMR (DMSO-d6) δ: 1.43 (3H, s), 1.73-1.78 (1H, m), 2.01-2.08 (1H, m), 2.54-2.59 (1H, m), 2.89-2.96 (1H, m), 7.16 (1H, d, J = 8.1 Hz), 7.34 (1H, t, J = 8.0 Hz), 7.57 (1H, s), 7.68 (1H, d, J = 7.1 Hz), 8.70 (1H, d, J = 2.5 Hz), 8.77 (1H, d, J = 2.3 Hz), 10.80 (1H, s).	
60			1H-NMR (DMSO-d6) δ: 1.42 (3H, s), 1.70-1.76 (1H, m), 2.01-2.06 (1H, m), 2.54-2.60 (1H, m), 2.88-2.95 (1H, m), 3.97 (3H, s), 7.12 (1H, d, J = 7.8 Hz), 7.30 (1H, t, J = 7.8 Hz), 7.62-7.68 (2H, m), 8.29 (1H, d, J = 2.3 Hz), 8.40 (1H, d, J = 2.5 Hz), 10.53 (1H, s).	
61			1H-NMR (DMSO-d6) δ: 0.91 (3H, t, J = 7.2 Hz), 1.32-1.40 (5H, m), 1.51-1.58 (2H, m), 1.68-1.73 (1H, m), 1.97-2.05 (0H, m), 2.55-2.60 (1H, m), 2.85-2.92 (1H, m), 5.77 (2H, br s), 7.07 (1H, d, J = 7.6 Hz), 7.25 (1H, t, J = 8.0 Hz), 7.73 (2H, t, J = 6.7 Hz), 7.83-7.87 (1H, m), 7.96 (1H, s), 8.64 (1H, s), 9.95 (1H, s).	
62			1H-NMR (DMSO-d6) δ: 1.43 (3H, s), 1.71-1.77 (1H, m), 2.02-2.09 (1H, m), 2.55-2.61 (1H, m), 2.87-2.95 (1H, m), 3.31 (3H, s), 3.70-3.74 (2H, m), 4.51-4.54 (2H, m), 7.11 (1H, d, J = 7.1 Hz), 7.29 (1H, t, J = 7.7 Hz), 7.75 (1H, d, J = 8.3 Hz), 7.80 (1H, s), 8.43 (1H, s), 8.87 (1H, s), 10.35 (1H, s).	

[Table 38]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
63			1H-NMR (DMSO-d6) δ : 1.29 (3H, t, J = 7.5 Hz), 1.43 (3H, s), 1.71-1.77 (1H, m), 2.01-2.09 (1H, m), 2.55-2.61 (1H, m), 2.90-2.96 (3H, m), 7.12 (1H, d, J = 7.8 Hz), 7.30 (1H, t, J = 7.8 Hz), 7.76 (1H, d, J = 8.1 Hz), 7.82 (1H, s), 8.70 (1H, s), 9.18 (1H, s), 10.53 (1H, s).	
64			1H-NMR (DMSO-d6) δ : 1.41 (3H, s), 1.70-1.75 (1H, m), 1.99-2.06 (1H, m), 2.52-2.57 (1H, m), 2.87-2.94 (1H, m), 7.11 (1H, d, J = 7.6 Hz), 7.28 (1H, t, J = 7.8 Hz), 7.47-7.59 (5H, m), 7.65 (2H, d, J = 7.6 Hz), 8.30 (0H, s).	
65			1H-NMR (DMSO-d6) δ : 1.38 (3H, s), 1.67-1.72 (1H, m), 1.94-2.00 (1H, m), 2.03 (3H, s), 2.50-2.55 (1H, m), 2.85-2.92 (1H, m), 7.06 (1H, d, J = 7.1 Hz), 7.23 (1H, t, J = 7.7 Hz), 7.48 (2H, t, J = 8.3 Hz), 10.55 (1H, s).	
66	707.0(2M+1)			
67	743.1(2M+1)			
68			1H-NMR (DMSO-d6) δ : 1.45 (3H, s), 1.74-1.80 (1H, m), 2.06-2.13 (1H, m), 2.56-2.61 (1H, m), 2.90-2.97 (1H, m), 7.15 (1H, d, J = 7.6 Hz), 7.33 (1H, t, J = 7.7 Hz), 7.79 (1H, d, J = 7.8 Hz), 7.87 (1H, s), 8.04 (1H, s), 8.45 (1H, s), 9.29 (2H, d, J = 8.3 Hz), 10.79 (1H, s).	
69			1H-NMR (DMSO-d6) δ : 1.43 (3H, s), 1.71-1.76 (1H, m), 2.02-2.09 (1H, m), 2.55-2.63 (4H, m), 2.88-2.94 (1H, m), 7.12 (1H, d, J = 8.1 Hz), 7.29 (1H, t, J = 7.8 Hz), 7.74 (1H, d, J = 8.1 Hz), 7.81 (1H, s), 8.70 (1H, s), 9.08 (1H, s), 10.45 (1H, s).	
70			1H-NMR (DMSO-d6) δ : 1.43 (3H, s), 1.71-1.77 (1H, m), 2.02-2.09 (1H, m), 2.55-2.61 (1H, m), 2.87-2.95 (1H, m), 3.31 (3H, s), 3.70-3.74 (2H, m), 4.51-4.54 (2H, m), 7.11 (1H, d, J = 7.1 Hz), 7.29 (1H, t, J = 7.7 Hz), 7.75 (1H, d, J = 8.3 Hz), 7.80 (1H, s), 8.43 (1H, s), 8.87 (1H, s), 10.35 (1H, s).	
71			1H-NMR (DMSO-d6) δ : 0.94 (3H, t, J = 7.3 Hz), 1.43-1.47 (4H, m), 1.72-1.79 (3H, m), 2.02-2.09 (1H, m), 2.58 (1H, t, J = 9.7 Hz), 2.91 (1H, s), 4.40 (2H, t, J = 6.6 Hz), 7.11 (1H, d, J = 7.8 Hz), 7.29 (1H, t, J = 8.1 Hz), 7.75 (1H, d, J = 7.3 Hz), 7.80 (1H, s), 8.37 (1H, s), 8.86 (1H, s), 10.34 (1H, s).	
72	771.1(2M+1)			
73			1H-NMR (DMSO-d6) δ : 1.40 (3H, s), 1.68-1.73 (1H, m), 1.98-2.05 (1H, m), 2.11 (3H, s), 2.55-2.60 (1H, m), 2.69 (2H, t, J = 6.8 Hz), 2.86-2.92 (1H, m), 3.55-3.59 (2H, m), 5.77 (2H, br s), 7.07 (1H, d, J = 8.1 Hz), 7.25 (1H, t, J = 8.0 Hz), 7.71-7.75 (2H, m), 8.00 (2H, s), 8.65 (1H, s), 9.98 (1H, s).	
74			1H-NMR (DMSO-d6) δ : 0.99 (3H, t, J = 7.5 Hz), 1.39 (3H, s), 1.56 (2H, td, J = 14.3, 7.2 Hz), 1.67-1.73 (1H, m), 1.95-2.02 (1H, m), 2.38 (2H, t, J = 6.8 Hz), 2.50-2.55 (1H, m), 2.86-2.93 (1H, m), 7.06 (1H, d, J = 7.8 Hz), 7.23 (1H, t, J = 8.3 Hz), 7.47-7.51 (2H, m).	
75	839.1(2M+1)			
76	835.1(2M+1)			
77	782.9(2M+1)			
78			1H-NMR (DMSO-d6) δ : 1.50 (3H, s), 1.79-1.84 (1H, m), 2.23-2.30 (1H, m), 2.55-2.60 (1H, m), 2.96-3.02 (1H, m), 7.16 (1H, dd, J = 11.6, 8.8 Hz), 7.75-7.78 (2H, m), 8.89 (1H, s), 9.07 (1H, s), 10.74 (1H, br s).	

[Table 39]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
79			1H-NMR (DMSO-d6) δ: 1.49 (4H, s), 1.77-1.83 (1H, m), 2.16-2.23 (1H, m), 2.56-2.62 (1H, m), 2.95-3.01 (1H, m), 5.87 (2H, br s), 7.17 (1H, dd, J = 11.7, 8.5 Hz), 7.76-7.82 (2H, m), 9.96 (2H, d, J = 3.8 Hz), 10.82 (1H, s).	
80			1H-NMR (DMSO-d6) δ: 1.33 (3H, t, J = 7.2 Hz), 1.42 (3H, s), 1.71-1.76 (1H, m), 2.03-2.07 (1H, m), 2.55-2.59 (1H, m), 2.89-2.92 (1H, m), 3.25 (2H, q, J = 7.3 Hz), 7.11 (1H, d, J = 7.8 Hz), 7.29 (1H, t, J = 7.8 Hz), 7.74 (1H, d, J = 8.1 Hz), 7.79 (1H, s), 8.66 (1H, s), 9.07 (1H, s), 10.45 (1H, s).	
81	831.1(2M+1)			
82	850.9(2M+1)			
83	795.0(2M+1)			
84	758.8(2M+1)			
85	750.9(2M+1)			
86	795.1(2M+1)			
87			1H-NMR (DMSO-d6) δ: 1.01 (6H, d, J = 6.8 Hz), 1.44 (3H, s), 1.73-1.78 (1H, m), 2.05-2.13 (2H, m), 2.56-2.61 (1H, m), 2.89-2.95 (1H, m), 4.19 (2H, d, J = 6.6 Hz), 7.12 (1H, d, J = 8.1 Hz), 7.29 (1H, t, J = 7.8 Hz), 7.76 (1H, d, J = 7.8 Hz), 7.82 (1H, s), 8.41 (1H, d, J = 1.3 Hz), 8.87 (1H, s), 10.36 (1H, s).	
88			1H-NMR (DMSO-d6) δ: 1.70 (3H, s), 2.02-2.08 (1H, m), 2.58-2.64 (2H, m), 3.15-3.19 (1H, m), 5.16 (2H, q, J = 8.8 Hz), 7.27 (1H, dd, J = 11.9, 8.8 Hz), 7.85-7.98 (2H, m), 8.62 (1H, s), 8.92 (1H, s), 10.83 (1H, s).	
89			1H-NMR (DMSO-d6) δ: 1.65 (3H, s), 2.04-2.11 (1H, m), 2.54-2.62 (3H, m), 2.83-2.95 (2H, m), 3.11-3.14 (1H, m), 4.65 (2H, t, J = 5.8 Hz), 7.09 (1H, d, J = 7.6 Hz), 7.42 (1H, t, J = 8.0 Hz), 7.87-7.92 (2H, m), 8.43 (1H, s), 8.91 (1H, s), 10.62 (1H, s).	
90			1H-NMR (DMSO-d6) d: 1.41 (3H, s), 1.69-1.74 (1H, m), 2.00-2.04 (1H, m), 2.56-2.61 (1H, m), 2.87-2.92 (1H, m), 3.24 (6H, s), 3.37-3.52 (12H, m), 3.59 (2H, t, J = 4.5 Hz), 3.80 (2H, t, J = 4.3 Hz), 4.52 (2H, t, J = 4.4 Hz), 5.81 (2H, br s), 7.12 (1H, d, J = 7.6 Hz), 7.28 (1H, t, J = 8.0 Hz), 7.75 (1H, d, J = 8.3 Hz), 7.81 (1H, s), 8.42 (1H, s), 8.87 (1H, s), 10.33 (1H, s).	
91			1H-NMR (DMSO-d6) δ: 1.41 (3H, s), 1.69-1.75 (1H, m), 1.85 (3H, s), 1.98-2.05 (1H, m), 2.55-2.61 (1H, m), 2.86-2.93 (1H, m), 5.09 (2H, d, J = 2.0 Hz), 5.79 (2H, br s), 7.12 (1H, d, J = 7.8 Hz), 7.28 (1H, t, J = 7.8 Hz), 7.74 (1H, d, J = 8.3 Hz), 7.80 (1H, s), 8.45 (1H, s), 8.90 (1H, s), 10.36 (1H, s).	
92			1H-NMR (DMSO-d6) δ: 1.41 (4H, s), 1.69-1.74 (1H, m), 1.98-2.05 (1H, m), 2.56-2.61 (1H, m), 2.87-2.93 (1H, m), 4.74 (2H, td, J = 15.0, 3.1 Hz), 5.79 (2H, br s), 6.34-6.61 (1H, m), 7.12 (1H, d, J = 7.8 Hz), 7.29 (1H, t, J = 8.0 Hz), 7.74 (1H, d, J = 8.1 Hz), 7.81 (1H, s), 8.54 (1H, s), 8.90 (1H, s), 10.40 (1H, s).	
93	837.0(2M+1)			
94			1H-NMR (DMSO-d6) δ: 1.41 (4H, s), 1.69-1.74 (1H, m), 1.98-2.05 (1H, m), 2.55-2.61 (2H, m), 2.71-2.75 (1H, m), 2.87-2.93 (1H, m), 4.49-4.61 (4H, m), 5.05-5.11 (1H, m), 5.79 (2H, br s), 7.12 (1H, d, J = 7.3 Hz), 7.28 (1H, t, J = 8.0 Hz), 7.74 (1H, d, J = 8.6 Hz), 7.81 (1H, s), 8.47 (1H, s), 8.88 (1H, s), 10.36 (1H, s).	
95	801.0(2M+1)			

[Table 40]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
96			1H-NMR (DMSO-d6) δ : 1.42 (3H, s), 1.70-1.78 (1H, m), 1.97-2.04 (3H, m), 2.55-2.60 (1H, m), 2.87-2.93 (1H, m), 3.26 (3H, s), 3.49 (2H, t, J = 6.2 Hz), 4.20 (2H, t, J = 6.4 Hz), 5.86 (2H, br s), 7.10 (1H, d, J = 7.8 Hz), 7.28 (1H, t, J = 8.1 Hz), 7.61 (1H, dd, J = 8.8, 2.8 Hz), 7.77-7.78 (2H, m), 8.11 (1H, d, J = 8.6 Hz), 8.38 (1H, d, J = 2.8 Hz), 10.32 (1H, s).	
97			1H-NMR (DMSO-d6) δ : 1.41 (3H, s), 1.70-1.75 (1H, m), 1.99-2.06 (1H, m), 2.56-2.61 (1H, m), 2.87-2.93 (1H, m), 4.40-4.50 (2H, m), 4.74-4.88 (2H, m), 5.81 (2H, br s), 7.10 (1H, d, J = 7.6 Hz), 7.28 (1H, t, J = 8.2 Hz), 7.66 (1H, dd, J = 8.8, 2.8 Hz), 7.77-7.78 (2H, m), 8.13 (1H, d, J = 8.8 Hz), 8.43 (1H, d, J = 2.5 Hz), 10.33 (1H, s).	
98			1H-NMR (DMSO-d6) δ : 1.41 (3H, s), 1.74-1.76 (7H, m), 1.99-2.06 (1H, m), 2.56-2.61 (1H, m), 2.86-2.93 (1H, m), 4.71 (2H, d, J = 6.3 Hz), 5.44-5.49 (1H, m), 5.81 (2H, br s), 7.10 (1H, d, J = 7.3 Hz), 7.28 (1H, t, J = 8.1 Hz), 7.60 (1H, d, J = 8.8 Hz), 7.77 (2H, s), 8.11 (1H, d, J = 8.6 Hz), 8.36 (1H, s), 10.30 (1H, s).	
99	799.0(2M+1)			
100	827.0(2M+1)			
101	867.1(2M+1)			
102	865.1(2M+1)			
103	382			
104	412			
105			1H-NMR (DMSO-d6) δ : 1.41 (4H, s), 1.69-1.74 (1H, m), 1.98-2.05 (1H, m), 2.55-2.60 (1H, m), 2.69-2.75 (2H, m), 2.86-2.93 (2H, m), 4.49 (2H, t, J = 6.4 Hz), 5.82 (2H, br s), 7.12 (1H, d, J = 7.3 Hz), 7.28 (1H, t, J = 7.7 Hz), 7.74 (1H, d, J = 7.8 Hz), 7.80 (1H, s), 8.42 (1H, s), 8.88 (1H, s), 10.34 (1H, s).	
106			1H-NMR (DMSO-d6) δ : 1.41 (3H, s), 1.69-1.76 (3H, m), 1.98-2.05 (1H, m), 2.56-2.61 (1H, m), 2.64-2.69 (2H, m), 2.87-2.93 (1H, m), 4.45 (2H, t, J = 6.4 Hz), 5.80 (2H, br s), 7.12 (1H, d, J = 7.8 Hz), 7.28 (1H, t, J = 7.8 Hz), 7.74 (1H, d, J = 8.3 Hz), 7.80 (1H, s), 8.42 (1H, s), 8.88 (1H, s), 10.34 (1H, s).	
107			1H-NMR (DMSO-d6) δ : 1.47-1.54 (1H, m), 1.86 (3H, s), 2.03-2.09 (1H, m), 2.88-2.94 (1H, m), 3.09-3.15 (1H, m), 4.43-4.47 (1H, m), 5.08-5.11 (2H, m), 5.76 (2H, br s), 7.04-7.06 (1H, m), 7.27-7.31 (1H, m), 7.68-7.70 (1H, m), 7.79 (1H, s), 8.45 (1H, s), 8.89 (1H, s), 10.39 (1H, s).	
108	412			
109	398			
110			1H-NMR (DMSO-d6) δ : 1.45 (3H, s), 1.75-1.81 (1H, m), 2.08-2.14 (1H, m), 2.56-2.61 (1H, m), 2.90-2.97 (1H, m), 7.08 (1H, d, J = 6.8 Hz), 7.27-7.36 (2H, m), 7.77-7.79 (2H, m), 8.01 (1H, d, J = 8.6 Hz), 8.22 (1H, d, J = 2.8 Hz), 10.27 (1H, s).	
111			1H-NMR (DMSO-d6) δ : 1.45 (3H, s), 1.75-1.80 (1H, m), 2.07-2.14 (1H, m), 2.56-2.61 (1H, m), 2.90-2.97 (1H, m), 5.28 (2H, s), 7.09 (1H, d, J = 7.8 Hz), 7.23-7.32 (3H, m), 7.57 (2H, dd, J = 8.3, 5.6 Hz), 7.70 (1H, dd, J = 8.7, 2.7 Hz), 7.78-7.81 (2H, m), 8.13 (1H, d, J = 8.6 Hz), 8.45 (1H, d, J = 2.8 Hz), 10.36 (1H, s).	
112	441			






[Table 41]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
113			1H-NMR (DMSO-d6) δ: 1.43 (3H, s), 1.71-1.77 (1H, m), 2.05-2.13 (7H, m), 2.56-2.61 (1H, m), 2.88-2.94 (1H, m), 5.75 (2H, s), 7.12 (1H, d, J = 8.1 Hz), 7.30 (1H, t, J = 7.8 Hz), 7.74-7.80 (2H, m), 8.66 (1H, s), 8.93 (1H, s), 10.44 (1H, s).	
114			1H-NMR (DMSO-d6) δ: 1.47 (3H, s), 1.77-1.82 (1H, m), 2.13-2.20 (1H, m), 2.57-2.62 (1H, m), 2.93-3.00 (1H, m), 3.75-3.79 (2H, m), 4.18 (2H, t, J = 4.4 Hz), 5.00 (1H, br s), 5.89 (2H, br s), 7.12 (1H, dd, J = 11.4, 8.8 Hz), 7.58-7.63 (1H, m), 7.72-7.82 (2H, m), 8.09 (1H, d, J = 8.6 Hz), 8.39 (1H, s), 10.34 (1H, s).	
115	416			
116			1H-NMR (DMSO-d6) δ: 1.47 (4H, s), 1.76-1.81 (1H, m), 2.15-2.22 (1H, m), 2.55-2.60 (1H, m), 2.93-2.99 (1H, m), 4.47 (3H, s), 5.87 (2H, br s), 7.13 (1H, dd, J = 12.0, 8.7 Hz), 7.79 (2H, ddd, J = 18.3, 8.0, 3.1 Hz), 8.89 (1H, s), 9.23 (1H, s), 10.72 (1H, s).	
117			1H-NMR (DMSO-d6) δ: 1.48 (3H, s), 1.77-1.82 (1H, m), 2.14-2.21 (1H, m), 2.57-2.62 (1H, m), 2.93-3.00 (1H, m), 3.32 (3H, s), 4.56-4.59 (2H, m), 5.87 (2H, br s), 6.27-6.33 (1H, m), 6.75 (1H, d, J = 12.4 Hz), 7.13 (1H, dd, J = 11.9, 8.8 Hz), 7.77-7.83 (2H, m), 8.77 (1H, s), 9.25 (1H, s), 10.64 (1H, s).	
118	418			
119	361			
120	825.1(2M+1)			
121	416			
122			1H-NMR (DMSO-d6) δ: 1.41 (3H, s), 1.68-1.74 (1H, m), 1.98-2.05 (1H, m), 2.21 (6H, s), 2.55-2.60 (1H, m), 2.67 (2H, t, J = 5.6 Hz), 2.86-2.93 (1H, m), 4.48 (2H, t, J = 5.4 Hz), 5.79 (2H, br s), 7.11 (1H, d, J = 7.8 Hz), 7.28 (1H, t, J = 7.8 Hz), 7.73 (1H, d, J = 7.6 Hz), 7.79 (1H, s), 8.39 (1H, s), 8.86 (1H, s), 10.33 (1H, s).	
123	395			
124	414			
125	417			
126			1H-NMR (DMSO-d6) δ: 1.45 (3H, s), 1.75-1.81 (1H, m), 2.13-2.20 (1H, m), 2.55-2.60 (1H, m), 2.92-2.99 (1H, m), 3.88 (6H, s), 5.81 (2H, br s), 7.08 (1H, dd, J = 11.6, 8.8 Hz), 7.50-7.55 (1H, m), 7.66-7.69 (1H, m), 10.07 (1H, s).	
127	380			
128	372			
129	418			
130	412			
131			1H-NMR (DMSO-d6) δ: 1.48 (3H, s), 1.77-1.82 (1H, m), 2.18-2.25 (1H, m), 2.55-2.60 (1H, m), 2.94-3.01 (1H, m), 4.72 (2H, t, J = 13.8 Hz), 6.45 (1H, t, J = 53.9 Hz), 7.10-7.15 (1H, m), 7.74-7.79 (2H, m), 8.50 (1H, s), 8.87 (1H, s), 10.47 (1H, s).	
132			1H-NMR (DMSO-d6) δ: 1.49 (3H, s), 1.78-1.83 (1H, m), 2.19-2.26 (1H, m), 2.56-2.60 (1H, m), 2.94-3.01 (1H, m), 4.61-4.86 (4H, m), 7.09-7.14 (1H, m), 7.75-7.79 (2H, m), 8.43 (1H, s), 8.84 (1H, s), 10.43 (1H, s).	
133	400			






[Table 42]

Compound No.	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
134			1H-NMR (DMSO-d6) δ : 1.37 (3H, t, J = 7.1 Hz), 1.49 (3H, s), 1.78-1.83 (1H, m), 2.19-2.26 (1H, m), 2.56-2.61 (1H, m), 2.95-3.01 (1H, m), 4.44 (2H, q, J = 7.0 Hz), 7.13 (1H, dd, J = 11.6, 9.1 Hz), 7.73-7.78 (2H, m), 8.35 (1H, s), 8.83 (1H, s), 10.41 (1H, s)	
135	411			
136	412			
137			1H-NMR (CDCl3) δ : 1.63 (3H, s), 1.81-1.91 (1H, m), 2.21-2.32 (1H, m), 2.56-2.67 (1H, m), 2.75-2.83 (1H, m), 3.77 (3H, s), 5.24 (2H, s), 6.47 (1H, dd, J = 3.2, 0.6Hz), 6.83 (2H, d, J = 8.9Hz), 7.02 (1H, dd, 8.0, 1.8Hz), 7.07 (1H,d, 3.2Hz), 7.12 (2H, d, J = 8.9Hz), 7.26 (1H,dd, 1.8, 0.6Hz), 7.57 (1H, d, J = 8.0Hz).	
138	400			
139	246			
140	356			
141	376			
142	410			
143	378			
144	398			
145	432			
146	529			
147	377			
148	438			
149	390			
150				212.2
151				211.0, 266.3, 301.8
152				285.2
153	403			
154	403			
155	404			
156	388			
157	389			
158	412			
159	380			
160	381			
161			1H-NMR (DMSO-d6) δ : 1.47 (3H, s), 1.77-1.82 (1H, m), 2.15-2.21 (8H, m), 2.56-2.67 (3H, m), 2.93-3.00 (1H, m), 4.21 (2H, t, J = 5.4 Hz), 5.88 (2H, br s), 7.11 (1H, dd, J = 11.6, 9.3 Hz), 7.59-7.61 (1H, m), 7.73-7.80 (2H, m), 8.09 (1H, d, J = 8.6 Hz), 8.37 (1H, s), 10.33 (1H, s).	
162	402			
163	408			
164	464			
165	459			
166	404			
167	420			
168	375			
169	432			
170	380			
171	376			





[Table 43]

Compound No.	Structure	MS(M+1)
172		
173		459
174		403
175		426
176		393




[Table 44]

Compound No.	Structure	MS(M+1)
177		359
178		402
179		447
180		435
181		396






[Table 45]

Compound No.	Structure	MS(M+1)
182		376
183		385
184		375
185		378
186		412






[Table 46]

Compound No.	Structure	MS(M+1)
187		366
188		429
189		364
190		404
191		439






[Table 47]

Compound No.	Structure	MS(M+1)
192		412
193		426
194		393
195		352
196		414






[Table 48]

Compound No.	Structure	MS(M+1)
197		
198		414
199		364
200		397
201		428






[Table 49]

Compound No.	Structure	MS(M+1)
202		
203		398
204		410
205		422
206		395






[Table 50]

Compound No.	Structure	MS(M+1)
207		414
208		410
209		402
210		
211		





[Table 51]

Compound No.	Structure	MS(M+1)
212		433
213		466
214		464
215		427
216		400






[Table 52]

Compound No.	Structure	MS(M+1)
217		442
218		386
219		402
220		362
221		





[Table 53]

Compound No.	Structure	MS(M+1)
222		399
223		
224		352
225		402
226		395






[Table 54]

Compound No.	Structure	MS(M+1)
227		362
228		375
229		380
230		
231		400






[Table 55]

Compound No.	Structure	MS(M+1)
232		
233		
234		422
235		395
236		364





[Table 56]

Compound No.	Structure	MS(M+1)
237		362
238		427
239		455
240		420
241		406






[Table 57]

Compound No.	Structure	MS(M+1)
242		471
243		406
244		420
245		383
246		






[Table 58]

Compound No.	Structure	MS(M+1)
247		455
248		435
249		416
250		416





[Table 59]

Compound No.	Structure	MS(M+1)
251		402
252		388
253		420
254		
255		524






[Table 60]

Compound No.	Structure	MS(M+1)
256		348
257		
258		395
259		395
260		402






[Table 61]

Compound No.	Structure	MS(M+1)
261		336
262		
263		
264		334
265		384






[Table 62]

Compound No.	Structure	MS(M+1)
266		402
267		402
268		
269		396
270		427






[Table 63]

Compound No.	Structure	MS(M+1)
271		444
272		416
273		429
274		
275		376






[Table 64]

Compound No.	Structure	MS(M+1)
276		425
277		425
278		429
279		430
280		






[Table 65]

Compound No.	Structure	MS(M+1)
281		448
282		411
283		
284		438
285		





[Table 66]

Compound No.	Structure	MS(M+1)
286		437
287		437
288		348
289		429
290		448






[Table 67]

Compound No.	Structure	MS(M+1)
291		398
292		
293		419
294		
295		422






[Table 68]

Compound No.	Structure	MS(M+1)
296		430
297		410
298		410
299		
300		401






[Table 69]

Compound No.	Structure	MS(M+1)
301		
302		400
303		349
304		426
305		363






[Table 70]

Compound No.	Structure	MS(M+1)
306		415
307		
308		424
309		406
310		383






[Table 71]

Compound No.	Structure	MS(M+1)
311		470
312		422
313		476
314		401
315		428






[Table 72]

Compound No.	Structure	MS(M+1)
316		413
317		442
318		442
319		411
320		434






[Table 73]

Compound No.	Structure	MS(M+1)
321		
322		463
323		
324		
325		410






[Table 74]

Compound No.	Structure	MS(M+1)
326		390
327		410
328		410
329		410
330		384






[Table 75]

Compound No.	Structure	MS(M+1)
331		479
332		429
333		427
334		427
335		410






[Table 76]

Compound No.	Structure	MS(M+1)
336		428
337		426
338		401
339		400
340		






[Table 77]

Compound No.	Structure	MS(M+1)
341		441
342		442
343		442
344		430
345		428






[Table 78]

Compound No.	Structure	MS(M+1)
346		430
347		411
348		413
349		478
350		






[Table 79]

Compound No.	Structure	MS(M+1)
351		384
352		443
353		403
354		
355		421





[Table 8]

Compound No.	Structure	MS(M+1)
356		422
357		421
358		369
359		430
360		424






[Table 81]

Compound No.	Structure	MS(M+1)
361		416
362		429
363		
364		
365		398





[Table 82]

Compound No.	Structure	MS(M+1)
366		425
367		425
368		
369		424






[Table 83]

Compound No.	Structure	MS(M+1)
370		413
371		430
372		408
373		426
374		437






[Table 84]

Compound No.	Structure	MS(M+1)
375		424
376		
377		427
378		424



[Table 85]

Compound No.	Structure	MS(M+1)
379		424
380		493
381		458
382		395
383		407






[Table 86]

Compound No.	Structure	MS(M+1)
384		416
385		364
386		
387		
388		






[Table 87]

Compound No.	Structure	MS(M+1)
389		
390		
391		
392		413
393		446






[Table 88]

Compound No.	Structure	MS(M+1)
394		445
395		428
396		413
397		494
398		428





[Table 89]

Compound No.	Structure	MS(M+1)
399		404
400		375
401		444
402		444
403		448






[Table 90]

Compound No.	Structure	MS(M+1)
404		440
405		365
406		414
407		443
408		385





[Table 91]

Compound No.	Structure	MS(M+1)
409		423
410		410
411		
412		393
413		348






[Table 92]

Compound No.	Structure	MS(M+1)
414		414
415		438
416		410
417		
418		464





[Table 93]

Compound No.	Structure	MS(M+1)
419		461
420		462
421		412
422		466
423		437






[Table 94]

Compound No.	Structure	MS(M+1)
424		411
425		411
426		351
427		478
428		462






[Table 95]

Compound No.	Structure	MS(M+1)
429		
430		443
431		470
432		
433		378





[Table 96]

Compound No.	Structure	MS(M+1)
434		451
435		355
436		351
437		509
438		420





[Table 97]

Compound No.	Structure	MS(M+1)
439		429
440		406
441		494
442		458
443		483




[Table 98]

Compound No.	Structure	MS(M+1)
444		457
445		452
446		550
447		437






[Table 99]

Compound No.	Structure	MS(M+1)
448		495
449		455
450		481
451		426







[Table 100]

Compound No.	Structure	MS(M+1)
452		454
453		480
454		404
455		441







[Table 101]

Compound No.	Structure	MS(M+1)
456		417
457		395
458		362
459		393
460		








[Table 102]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
461		346			
462		349		1H-NMR (DMSO-d6) d: 10.02 (1.0H, s), 8.59 (1.0H, s), 7.73-7.66 (2.0H, m), 7.09 (1.0H, dd, J = 12.00, 8.97 Hz), 5.83 (2.0H, br s), 2.97-2.95 (1.0H, m), 2.59-2.56 (1.0H, m), 2.17-2.16 (1.0H, m), 1.79-1.76 (1.0H, m), 1.47 (3.0H, s).	
463		362			
464		441			
465		456			
466				1H-NMR (DMSO-d6) d: 10.70 (1.0H, s), 8.76 (1.0H, s), 8.36 (1.0H, s), 8.03 (1.0H, s), 6.44 (1.0H, s), 5.93 (2.0H, br s), 3.00-2.97 (1.0H, m), 2.63-2.61 (1.0H, m), 2.19 (3.0H, s), 2.00-1.98 (1.0H, m), 1.82-1.80 (1.0H, m), 1.60 (9.0H, s), 1.43 (3.0H, s).	








[Table 103]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value: ascending order)	UV
467		369			
468		396			
469		450			
470		383			
471		417			
472		364			








[Table 104]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
473		361			
474		332			
475		378			
476		345			
477		392			
478		365			
479		359			








[Table 105]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
480		360			
481	 ClH	366			
482		345			
483		394			
484	 ClH	385			
485		347			
486		347			








[Table 106]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
487		347			
488	 ClH	362			
489		405			
490		381			
491	 ClH	379			
492		421			
493		379			








[Table 107]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
494		426			
495	 ClH	363			
496		378			
497		426			
498	 ClH	374			
499	 ClH	374			
500	 ClH	363			








[Table 108]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value: ascending order)	UV
501		400			
502	 ClH	384			
503	 ClH	359			
504		367			
505	 ClH	365			
506	 ClH	365			
507	 ClH	365			







[Table 109]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value: ascending order)	UV
508	 ClH	365			
509		411			
510		363			
511	 ClH	363			
512		393			
513	 ClH	408			
514		413			







[Table 110]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
515		411			
516		413			
517		441			
518	 ClH	348			
519		429			
520	 ClH	394			
521	 ClH	402			







[Table 111]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
522		378			
523		441			
524	 ClH	380			
525	 ClH	379			
526		414			
527		428			







[Table 112]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value: ascending order)	UV
528		433			
529	 ClH	362			
530	 ClH	392			
531		426			
532	 ClH	364			
533	 ClH	364			







[Table 113]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
534	 ClH	404			
535	 ClH	394			
536	 ClH	383			
537		428			
538		404			
539	 ClH	401			








[Table 114]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
540	 ClH	384			
541		442			
542	 ClH	401			
543	 ClH	404			
544		511			
545	 ClH	400			








[Table 115]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
546	 ClH			1H-NMR (DMSO-d6) d: 10.92 (1H, s), 10.45 (1H, s), 8.45 (1H, s), 8.42 (1H, s), 8.30 (1H, d, J = 8.8 Hz), 7.79-7.78 (3H, m), 7.31 (1H, dd, J = 12.3, 9.2 Hz), 3.22 (1H, d, J = 13.4 Hz), 2.72-2.65 (2H, m), 2.11 (1H, t, J = 11.5 Hz), 1.73 (3H, s).	
547		359			
548		359			
549		403			
550		343			
551		343			








[Table 116]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
552		363			
553		348			
554		363			
555		374			
556		383			
557				1H-NMR (DMSO-d6) δ: 10.72 (1H, s), 8.93 (1H, s), 8.90 (1H, s), 8.49 (1H, s), 8.36 (1H, s), 8.24 (1H, s), 4.88-4.64 (4H, m), 3.03-2.97 (1H, m), 2.65-2.58 (1H, m), 2.13-2.07 (1H, m), 1.89-1.81 (1H, m), 1.48 (3H, s).	
558		402			








[Table 117]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
559		418			
560		387			
561		411			
562		431			
563		342			
564		372			
565		390			








[Table 118]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
566		428			
567		429			
568		419			
569		442			
570		456			
571		443			
572		396			








[Table 119]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
573		447			
574		430			
575		458			
576		412			
577		426			
578		426			
579		440			








[Table 120]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
580		480			
581		363			
582		393			
583		437			
584		366			
585		360			
586		380			







[Table 121]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
587		363			
588		323			
589					250.9, 288.7
590					298.2
591					252.1, 305.3
592					250.9, 288.7
593					216.9, 292.3







[Table 122]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
594					214.5, 289.9
595					297
596					250.9, 302.9
597					289.9
598					297
599					214.5, 289.9
600		404, 807 (2M+1)			







[Table 123]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
601		448, 895 (2M+1)			
602		389			
603		391			
604		391			
605		436			
606		388			







[Table 124]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
607					
608					
609		377			
610					
611					
612		332			







[Table 125]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
613		346			
614					
615					
616					
617					
618					







[Table 126]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
619					
620					
621					
622					
623					
624					







[Table 127]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	uv
625					
626					
627					
628					
629					
630					





[Table 128]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
631					
632					
633					
634					
635					
636					








[Table 129]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value:ascending order)	UV
637		387			
638					
639					
640					
641					
642					








[Table 130]

Compound No.	Structure	MS (M+1)	MP	NMR (solvent, shift value: ascending order)	UV
643					
644		475			
645		397			
646		414			








[Table 131]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
647		404	
648		377	
649		388	
650		389	
651		453	
652		399	
653		371	








[Table 132]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
654		360	
655		374	
656		458	
657		411	
658		419	
659		383	
660			1H-NMR (CDCl3) δ: 1.84 (3H, d-like), 3.16 (1H; ddd, J = 6.9, 12.6, 14.4 Hz), 3.36 (1H, ddd, J = 6.0, 12.6, 18.9 Hz), , 4.61 (2H, br), 7.07 (1H, dd, J = 8.7, 11.7 Hz), 7.49-7.63 (2H, m), 7.67 (1H, dd, J = 3.0, 6.9 Hz), 7.95 (1H, ddd, J = 3.0, 6.9, 8.7 Hz), 8.41 (1H, m), 9.85 (1H, brs).








[Table 133]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
661			1H-NMR (CDCl3) δ: 0.89 (3H, s), 1.11 (3H, d, J = 3.0 Hz), 1.67 (3H, d, J = 4.2 Hz), 2.63 (1H, d, J = 12.0 Hz), 3.12 (1H, d, J = 12.0 Hz), 4.29 (2H, br), 7.02 (1H, dd, J = 8.7, 12.3 Hz), 7.49-7.64 (3H, m), 7.96 (1H, ddd, J = 3.0, 6.6, 8.7 Hz), 8.45 (1H, m), 9.81 (1H, brs).
662			1H-NMR (CDCl3) δ: 1.85 (3H, d-like), 2.69 (3H, s), 3.17 (1H, ddd, J = 6.9, 12.6, 14.4 Hz), 3.37 (1H, ddd, J = 6.3, 12.9, 18.9 Hz), 4.54 (2H, brs), 7.08 (1H, dd, J = 8.7, 11.7 Hz), 7.69 (1H, dd, J = 2.7, 6.9 Hz), 7.87 (1H, dd, J = 2.4, 8.4 Hz), 7.93 (1H, ddd, J = 2.7, 6.6, 8.7 Hz), 8.24 (1H, dd, J = 0.6, 8.4 Hz), 8.55 (1H, dd, J = 0.6, 2.4 Hz), 9.82 (1H, brs).
663			1H-NMR (CDCl3) δ: 1.85 (3H, d-like), 3.18 (1H, ddd, J = 7.2, 12.9, 15.0 Hz), 3.37 (1H, ddd, J = 6.0, 12.6, 18.9 Hz), 4.60 (2H, br), 7.08 (1H, dd, J = 8.7, 11.7 Hz), 7.71 (1H, dd, J = 3.0, 6.6 Hz), 7.90 (1H, ddd, J = 3.0, 6.6, 8.7 Hz), 8.41 (1H, d, J = 0.9 Hz), 9.35 (1H, d, J = 0.9 Hz), 9.63 (1H, brs).
664			1H-NMR (CDCl3) δ: 1.83 (3H, d-like), 2.51 (3H, s), 3.16 (1H, ddd, J = 6.9, 12.9, 15.3 Hz), 3.34 (1H, ddd, J = 6.3, 12.9, 19.2 Hz), 4.53 (2H, brs), 7.05 (1H, dd, J = 8.7, 11.4 Hz), 7.62 (1H, dd, J = 2.7, 6.9 Hz), 7.82 (1H, ddd, J = 2.7, 6.9, 8.7 Hz), 8.16 (1H, s), 8.70 (1H, brs).
665			1H-NMR (CDCl3) δ: 0.95 (3H, t, J = 7.5 Hz), 1.32-1.44 (2H, m), 1.59-1.69 (2H, m), 1.85 (3H, d-like), 2.70 (2H, t, J = 7.5 Hz), 3.17 (1H, ddd, J = 6.9, 12.9, 15.3 Hz), 3.34 (1H, ddd, J = 6.3, 12.9, 19.2 Hz), 4.53 (2H, brs), 7.07 (1H, dd, J = 8.7, 11.7 Hz), 7.67-7.70 (2H, m), 7.94 (1H, ddd, J = 2.7, 6.6, 8.7 Hz), 8.18 (1H, dd, J = 0.6, 7.8 Hz), 8.40 (1H, dd, J = 0.6, 1.8 Hz), 9.97 (1H, brs).
666		396	
667		365	








[Table 134]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
668		356	
669		403	
670			1H-NMR (CDCl3) δ: 1.38 (3H, t, J = 7.6 Hz), 1.63 (3H, s), 1.88-1.97 (1H, m), 2.41-2.50 (1H, m), 2.69-2.78 (1H, m), 2.84 (2H, q, J = 7.6 Hz), 2.93-3.01 (1H, m), 7.02 (1H, dd, J = 11.8, 8.8 Hz), 7.34 (1H, dd, J = 7.1, 2.6 Hz), 7.89 (1H, ddd, J = 8.8, 4.3, 2.8 Hz), 8.16 (1H, s), 8.69 (1H, s).
671		370	
672		432	
673		412	
674			1H-NMR ($CDCl_3$) δ: 0.53-0.59 (1H, m), 0.65-0.72 (1H, m), 0.85-0.91 (1H, m), 1.14-1.17 (1H, m), 1.47 (3H, d, J = 2.0 Hz), 2.46 (1H, d, J = 12.1 Hz), 2.69 (3H, s), 2.89 (1H, dd, J = 12.1, 1.3 Hz), 7.06 (1H, dd, J = 11.5, 8.8 Hz), 7.45 (1H, dd, J = 6.8, 2.8 Hz), 7.94 (1H, ddd, J = 8.8, 4.0, 2.8 Hz), 8.44 (1H, d, J = 1.3 Hz), 9.36 (1H, d, J = 1.3 Hz), 9.60 (1H, s).








[Table 135]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
675		402	
676		426	
677		396	
678		430	
679		372	
680			^{1}H-NMR (DMSO-d_6) δ: 1.47 (3H, s), 1.77-1.83 (1H, m), 2.34-2.39 (1H, m), 2.48-2.53 (1H, m), 2.63 (3H, s), 2.89-2.96 (1H, m), 3.90 (3H, s), 5.86 (2H, br s), 8.10 (1H, d, J = 2.3 Hz), 8.47 (1H, d, J = 2.5 Hz), 8.69 (1H, s), 9.14 (1H, s), 10.69 (1H, s).
681			^{1}H-NMR (DMSO-d_6) δ: 1.52 (3H, s), 1.80-1.85 (1H, m), 2.62 (3H, s), 2.64-2.69 (2H, m), 2.96-3.01 (1H, m), 7.77 (1H, d, J = 2.5 Hz), 7.96 (1H, d, J = 2.3 Hz), 8.67 (1H, s), 9.10 (1H, s), 10.58 (1H, s).








[Table 136]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
682			^{1}H-NMR (DMSO-d_6) δ: 1.46 (3H, s), 1.95-2.01 (1H, m), 2.33-2.39 (1H, m), 2.62 (3H, s), 2.64-2.69 (1H, m), 2.74 (3H, s), 2.92-2.98 (1H, m), 7.90 (1H, d, *J* = 2.5 Hz), 7.94-7.95 (1H, m), 8.67 (1H, s), 9.09 (1H, s), 10.57 (1H, s).
683		482	
684		482	
685		400	
686		424	
687		427	
688		402	

[Table 137]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
689		390	
690		413	
691		374	
692		428	
693			1H-NMR (DMSO-d6) δ: 0.77 (3H, s), 1.30 (3H, s), 1.43 (3H, s), 1.71 (1H, d, J = 13.8 Hz), 2.33 (1H, d, J = 13.8 Hz), 3.65 (1H, t, J = 2.4 Hz), 5.13 (2H, d, J = 2.4 Hz), 6.05 (2H, br), 7.19 (1H, d-like), 7.25 (1H, t, J = 7.8 Hz), 7.71 (1H, d-like), 7.87 (1H, s-like), 8.47 (1H, d, J = 1.2 Hz), 8.90 (1H, d, J = 1.2 Hz), 10.37 (1H, brs).
694			1H-NMR (DMSO-d6) δ: 0.77 (3H, s), 1.28 (3H, s), 1.42 (3H, s), 1.67 (1H, d, J = 14.1 Hz), 2.23 (1H, d, J = 14.1 Hz), 4.02 (3H, s), 5.82 (2H, brs), 7.19 (1H, d-like), 7.24 (1H, t, J = 7.8 Hz), 7.70 (1H, d-like), 7.86 (1H, s-like), 8.40 (1H, d, J = 1.2 Hz), 8.89 (1H, d, J = 1.2 Hz), 10.31 (1H, brs).
695			1H-NMR (DMSO-d6) δ: 0.79 (3H, s), 1.32 (3H, s), 1.46 (3H, s), 1.78 (1H, d, J = 14.1 Hz), 1.86 (3H, t, J = 2.4 Hz), 2.39 (1H, d, J = 14.1 Hz), 5.90 (1H, q, J = 2.4 Hz), 6.49 (2H, br), 7.18 (1H, d-like), 7.27 (1H, t, J = 7.8 Hz), 7.73 (1H, d-like), 7.87 (1H, s-like), 8.44 (1H, d, J = 1.2 Hz), 8.89 (1H, d, J = 1.2 Hz), 10.38 (1H, brs).

[Table 138]

Compound No.	Structure	MS [M+1]	NMR (solvent, shift value)
696			1H-NMR (DMSO-d6) δ: 0.77 (3H, s), 1.30 (3H, s), 1.43 (3H, s), 1.71 (1H, d, J = 14.1 Hz), 2.32 (1H, d, J = 14.1 Hz), 5.95 (2H, br), 7.22 (1H, d-like), 7.27 (1H, t, J = 7.8 Hz), 7.75 (1H, d-like), 7.88 (1H, s-like), 8.29 (1H, dd, J = 0.6, 8.1 Hz), 8.58 (1H, dd, J = 2.1, 8.1 Hz), 9.19 (1H, dd, J = 0.6, 2.1 Hz), 10.65 (1H, brs).
697		410	
698			
699			
700			
701		400	
702		443	

Test Example Assay of β-secretase-inhibiting activity

Forty eight point five μL of substrate peptide solution (Biotin-XSEVNLDAEFRHDSGC-Eu: X=ε-amino-n-capronic acid, Eu=Europium cryptate) was added to each well of 96-hole half-area plate (a black plate: Corning Incorporated), and after addition of 0.5 μl of the test sample (dissolved in N,N'-dimethylformaldehyde) and 1 μl of Recombinant human BACE-1(R&D Systems), the reaction mixture was incubated at 30°C for 3 hours. The substrate peptide was synthesized by reacting Cryptate TBPCOOH mono SMP (CIS bio international) with Biotin-XSEVNLDAEFRHDSGC (Peptide Institute, Inc.). The final concentrations of the substrate peptide and Recombinant human BACE-1 were adjusted to 18 nM and 7.4 nM respectively, and the reaction was performed in sodium acetate buffer (50 mM sodium acetate, pH 5.0, 0.008% Triton X-10).

After the incubation for reaction, 50 μl of 8.0 μg/ml Streptavidin-XL665(CIS bio international) dissolved in phosphate buffer (150 mM K_2HPO_4 -KH_2PO_4, pH 7.0, 0.008 % Triton X-100, 0.8 M KF) was added to each well and left stand at 30°C for an hour. After then, fluorescence intensity was measured (excitation wavelength: 320 nm, measuring wavelength: 620 nm and 665 nm) using Wallac 1420 multilabel counter (Perkin Elmer life sciences). Enzymatic activity was determined from counting ratio of each wavelength (10,000 x Count 665/Count 620) and 50% inhibitory concentration against the enzymatic activity was calculated. IC_{50} values of the test compounds are indicated in Table 139 - 141.

[Table 139]

Compound No.	IC50 (uM)
3	0.08
11	0.17
12	0.16
26	4.85
34	0.10
38	0.14
41	0.15
62	0.17
65	0.72
66	0.15
70	0.09
71	0.16
72	0.11
76	0.18
80	0.07
86	0.19
87	0.09

Compound No.	IC50 (uM)
92	0.08
93	0.08
94	0.17
101	0.08
105	0.13
106	0.12
109	0.10
111	0.18
114	0.16
126	2.14
136	0.11
141	0.12
149	9.25
150	2.48
151	6.77
155	5.96
163	6.79
164	0.08

[Table 140]

Compound No.	IC50(μM)
576	0.02
577	0.03
579	0.14
580	0.13
585	0.06
589	0.06
591	0.10
593	0.28
594	0.17
597	0.07
598	0.03
599	0.03
645	0.20

Compound No.	IC50(μM)
508	0.04
514	0.24
517	0.13
521	0.18
526	0.41
530	0.09
537	0.02
542	0.02
544	0.17
551	0.02
559	0.42
563	0.02
566	0.18

Compound No.	IC50(μM)
461	0.03
464	0.43
468	0.02
473	0.17
478	0.08
481	0.05
484	0.29
488	0.26
491	0.09
494	0.03
497	0.09
502	0.04
505	0.03

The following compounds have shown IC_{50} values equal to or under 1 μM in the same assay;

compounds 4, 5, 6, 8, 10, 18, 19, 20, 21, 22, 29, 32, 33, 35, 43, 45, 46, 58, 59, 63, 64, 68, 69, 75, 77, 78, 79, 81, 82, 83, 84, 85, 88, 89, 90, 91, 95, 96, 97, 98, 100, 102, 103, 104, 107, 108, 110, 112, 113, 115, 116, 117, 118, 119, 120, 121, 123, 124, 125, 127, 131, 132, 133, 134, 135, 142, 143, 144, 145, 148, 152, 157, 158, 162 and 165.

Also, compounds 462, 463, 465, 466, 467, 469, 470, 471, 472, 474, 475, 476, 477, 479, 480, 482, 483, 485, 486, 487, 489, 490, 492, 493, 495, 496, 499, 500, 501, 503, 504, 506, 507, 508, 509, 510, 511, 512, 515, 516, 518, 519, 520, 522, 523, 524, 527, 528, 529, 531, 532, 533, 536, 538, 539, 540, 541, 542, 545, 546, 547, 548, 549, 550, 552, 553, 554, 555, 556, 557, 558, 560, 561, 562, 564, 565, 567, 568, 569, 570, 571, 572, 573, 574, 575,

578, 581, 582, 583, 584, 586, 587, 590, 592, 595, 596, 600, 601, 602, 603, 604, 605, 606, 609, 612, 613, 637, 644 and 646 showed IC_{50} values equal to or under 1 μM in the same assay;

[Table 141]

Compound No.	IC50 (μM)
652	0.06
655	0.01
660	0.11
662	0.17
664	0.02
665	0.87
667	0.02
668	0.51
669	0.01
674	0.01
680	0.05
689	0.04
701	0.20
702	0.16

The following compounds also showed IC_{50} values equal to or under 1 μM in the same assay;

compounds 647, 648, 649, 650, 651, 653, 654, 656, 657, 658, 659, 661, 663, 666, 670, 671, 672, 673, 675, 676, 677, 678, 679, 683, 684, 685, 686, 687, 688, 690, 691, 692, 693, 694, 695, 696 and 697, 698, 699, 700, 701 and 702.

Formulation Example 1

Granular formulation is prepared with the following ingredients;

Ingredient	compound of the formula (I)	10 mg
	lactose	700 mg
	corn starch	274 mg
	HPC-L	16 mg
		1000 mg

Compound of the formula (I) and lactose are put through a sieve of No. 60 mesh.

Corn starch is put through a sieve of No.120 mesh and these are mixed with V-shaped mixer. An aqueous solution of HPC-L (Hydroxypropyl cellulose of Low viscosity) is added to the mixed powder, kneaded, granulated (extrusion granulation; pore diameter 0.5-1 mm) and put into a drying process. The resulted dried granule is sieved with vibrating screen (12/60 mesh) to give a granular formulation.

Formulation example 2

Granular formulation for capsule filling is prepared with the following ingredients;

Ingredient	compound of the formula (I)	15 mg
	lactose	90 mg
	corn starch	42 mg
	HPC-L	3 mg
		150 mg

Compound of the formula (I) and lactose are put through a sieve of No. 60 mesh. Corn starch is put through a sieve of No.120 mesh and these are mixed. An aqueous solution of HPC-L is added to the mixed powder, kneaded, granulated and dried. Particle size of the resulted dried granule is regulated and each of 150 mg is filled in No.5 hard-gelatin capsule.

Formulation Example 3

Tablet is prepared with the following ingredients;

Ingredient	compound of the formula (I)	10 mg
	lactose	90 mg
	microcrystalline cellulose	30 mg
	CMC-Na	15 mg
	magnesium stearate	5 mg
		150 mg

Compound of the formula (I), lactose, microcrystalline cellulose and CMC-Na (sodium salt of carboxymethylcellulose) are put through a sieve of No.60 mesh and mixed. Magnesium stearate is mixed with the mixed granule above to give a mixed powder for

tablet, which is compressed by a tabletting machine to give a tablet of 150 mg.

Formulation Example 4

The following ingredients were warmed, mixed and sterilized to give an injection.


Ingredient	compound of the formula (I)	3 mg
	non-ionic surfactant	15 mg
	purified water for injection	1 ml

INDUSTRIAL APPLICABILITY

A compound of the present invention can be a useful drug for treating diseases induced by production, secretion and/or deposition of amyloid β protein.

CLAIMS

1. A compound of the formula (I):



(I)

wherein the ring A is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group,

R^1 is optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted lower alkynyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group,

R^{2a} and R^{2b} are each independently hydrogen, optionally substituted lower alkyl or optionally substituted acyl,

R^{3a}, R^{3b}, R^{3c} and R^{3d} are each independently hydrogen, halogen, hydroxy, optionally substituted lower alkyl, optionally substituted lower alkenyl, optionally substituted acyl, optionally substituted lower alkoxy, optionally substituted carbocyclyl lower alkyl, optionally substituted heterocyclyl lower alkyl, optionally substituted carbocyclyl lower alkoxy, optionally substituted heterocyclyl lower alkoxy, optionally substituted aralkyl, optionally substituted heteroaralkyl, optionally substituted aralkyloxy, optionally substituted heteroaralkyloxy, optionally substituted lower alkylthio, carboxy, optionally substituted lower alkoxycarbonyl, optionally substituted amino, optionally substituted carbamoyl, an optionally substituted carbocyclic group or an optionally substituted heterocyclic group, or R^{3a} and R^{3b} or R^{3c} and R^{3d} may form a carbocyclic ring together with a linked carbon atom or may form oxo,

provided the following compounds i) and ii) are excluded;

i) a compound in which R^{2a} is hydrogen, R^{2b} is hydrogen, acetyl or phenyl, R^1 is


methyl, and the ring A is phenyl or 4-methoxyphenyl; and

ii) a compound in which R^{2a} is hydrogen, R^{2b} is hydrogen, acetyl or phenyl, R^1 is ethyl and the ring A is 3,4-dimethoxyphenyl;

a pharmaceutically acceptable salt or solvate thereof.


2. The compound of claim 1,

wherein the ring A is




wherein the ring A' is a carbocyclic group or a heterocyclic group,


G is




(ii')




(ii'')




(iv')




(v')



(x')




(xiii')



(xiv')

or



(xixi')

wherein R^5 is hydrogen, lower alkyl or acyl,

R^6 is optionally substituted lower alkyl, optionally substituted lower alkenyl or optionally substituted lower alkynyl,

W^1 is O or S,

W^2 is O, S or NR^5,

Ak is optionally substituted lower alkylene, optionally substituted lower alkenylene or optionally substituted lower alkynylene,

the ring B is an optionally substituted carbocyclic group or an optionally substituted heterocyclic group and each R^5 may be independent:

R^4 is halogen, hydroxyl, mercapto, halogeno lower alkyl, lower alkoxy, amino, lower alkylamino, acylamino or lower alkylthio and each R^4 may be independent;

a pharmaceutically acceptable salt or solvate thereof.

3. The compound of claim 2, wherein the ring A' is phenyl or a nitrogen-containing aromatic heterocyclic group; a pharmaceutically acceptable salt or solvate thereof.

4. The compound of claim 2, wherein the ring A' is a nitrogen-containing aromatic heterocyclic group; a pharmaceutically acceptable salt or solvate thereof.

5. The compound of any of claims 1-4, wherein R^1 is C1-C3 alkyl; a pharmaceutically acceptable salt or solvate thereof.

6. The compound of any of claims 1-5, wherein both of R^{2a} and R^{2b} are hydrogen; a pharmaceutically acceptable salt or solvate thereof.

7. The compound of any of claims 1-6, wherein all of R^{3a}, R^{3b}, R^{3c} and R^{3d} are hydrogen; a pharmaceutically acceptable salt or solvate thereof.

8. The compound of any of claims 1-6, wherein R^{3a} and R^{3b} are the same substituent selected from halogen and optionally substituted lower alkyl; a pharmaceutically acceptable salt or solvate thereof.

9. The compound of any of claims 1-6 and 8, wherein R^{3c} and R^{3d} are the same substituent selected from halogen and optionally substituted lower alkyl; a pharmaceutically acceptable salt or solvate thereof.

10. The compound of any of claims 1-6, wherein R^{3a} and R^{3b}, or R^{3c} and R^{3d} form a carbocyclic ring together with a linked carbon atom; a pharmaceutically acceptable salt or solvate thereof.


11. A pharmaceutical composition comprising a compound of any of claims 1-10, a pharmaceutically acceptable salt or solvate thereof as an active ingredient.

12. A pharmaceutical composition having a BACE1-inhibitory activity comprising a compound of any of claims 1-10, a pharmaceutically acceptable salt or solvate thereof as an active ingredient.

13. Method for treating diseases induced by BACE 1 characterized in administrating a compound of any of claims 1-10, a pharmaceutically acceptable salt or solvate thereof.

14 Use of a compound of any of claims 1-10, a pharmaceutically acceptable salt or

solvate thereof, in manufacturing a medicine for treating diseases induced by BACE 1.



(I)