



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification ⁶ : A61K 45/05, 39/395, 38/16, 38/17, 38/19, C07K 16/00</p>	<p>A1</p>	<p>(11) International Publication Number: WO 99/52552</p> <p>(43) International Publication Date: 21 October 1999 (21.10.99)</p>
<p>(21) International Application Number: PCT/US99/08430</p> <p>(22) International Filing Date: 15 April 1999 (15.04.99)</p> <p>(30) Priority Data: 60/081,895 15 April 1998 (15.04.98) US</p> <p>(71) Applicant: BRIGHAM & WOMEN'S HOSPITAL, INC. [US/US]; 75 Francis Street, Boston, MA 02115 (US).</p> <p>(72) Inventor: BLUMBERG, Richard, S.; 34 Lagrane Street, Chestnut Hill, MA 02167 (US).</p> <p>(74) Agent: VAN AMSTERDAM, John, R.; Wolf, Greenfield & Sacks, P.C., 600 Atlantic Avenue, Boston, MA 02210 (US).</p>	<p>(81) Designated States: CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</p> <p>Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	
<p>(54) Title: T CELL INHIBITORY RECEPTOR COMPOSITIONS AND USES THEREOF</p>		
<p>(57) Abstract</p> <p>The invention relates to compositions which bind T cell inhibitory receptor molecules and modulate T cell activity, and methods of using such compositions. Such compositions include biliary glycoprotein binding agents. Methods for modulating killer T cell activities, including cytotoxicity and proliferation also are provided.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

T CELL INHIBITORY RECEPTOR COMPOSITIONS AND USES THEREOF

Field of the Invention

The invention relates to compositions which bind T cell inhibitory receptor molecules
5 and modulate T cell activity.

Background of the Invention

The biologic role of human intestinal intraepithelial lymphocytes (iIEL) and their
functional relationship with the intestinal epithelial cell (IEC) remains incompletely
10 characterized. Human iIELs have been variably shown to exhibit cytolytic function and possibly
immunoregulatory function through the secretion of a variety of cytokines (Ebert, *Science*,
82:81-85, 1990; Lundqvist et al., *J. Immunol.*, 157:1926-1934, 1996; Balk et al., *Science*,
253:1411-1415, 1991). Whether these functional activities are related to processes that may be
unique to the gut associated lymphoid tissue such as oral tolerance and local immunosurveillance
15 against IEC injury and neoplastic transformation is, however, unclear. Moreover, the molecules
on the cell surface of iIELs and their IEC counterreceptors which regulate the functional
activation of iIELs and which may be utilized in this special microenvironment are only
beginning to be elucidated.

A significant fraction of human iIELs of both the small and large intestine are CD8- $\alpha\beta^+$
20 and CD45RO⁺ T cells which express a limited array of $\alpha\beta$ and, to a lesser extent, $\gamma\delta$ -T cell
receptors (TCR) (Balk et al, 1991; Jarry et al., *Eur. J. Immunol.*, 20:1097-1103, 1990; Blumberg
et al., *J. Immunol.*, 150:5144-5153, 1993; Van Kerckhove et al., *J. Exp. Med.*, 175:57-63, 1992;
Chowers et al., *J. Exp. Med.*, 180:183-190, 1994). These phenotypic properties indicate that
iIELs are memory cells which localize to the basolateral surface of IECs for the recognition of a
25 limited number of antigens in the context of major histocompatibility complex (MHC) class I or
class I-like molecules on the IEC. However, the majority of iIELs in mouse and human are
CD28⁻ suggesting that other costimulatory molecules for TCR/CD3 complex-mediated activation
may be important in providing necessary secondary signals for iIEL activation (Gelfanov et al.,
J. Immunol., 155:76-82, 1995; Gramzinski et al., *Int. Immunol.* 5:145-153, 1993; Russell et al., *J.*
30 *Immunol.*, 157:3366-3374, 1996). Candidate costimulatory molecules for human iIELs include
CD2 (Ebert, *Gastroenterology*, 97:1372-1381, 1989), CD101 (Russell et al., 1996), BY-55
(Anumanthan et al., *J. Immunol.*, 161:2780-2790, 1998) and $\alpha^E\beta_7$ (Parker et al., *Proc. Natl. Acad.*

Sci., 89:1924-1928, 1992) which are expressed by the majority of iIELs.

It has also become increasingly evident that in addition to activating costimulatory molecules, T cells can express a variety of molecules that deliver an inhibitory signal such that either the initial activation of the T cell is prevented or the activated state is downregulated. The former include the killer inhibitory receptors (KIR) which are expressed on a subset of T cells and bind specific types of major histocompatibility complex (MHC) class I molecules on the target cells (Lanier et al., *Immunology*, 7:75-82, 1995). The latter includes CTLA-4 (CD152) which, when expressed after T cell activation, binds either CD80 (B7.1) or CD86 (B7.2) on antigen presenting cells (Walunas et al., *J. Exp. Med.*, 183:2541-2550, 1996; Krummel et al., *J. Exp. Med.* 183:2533-2540, 1996). These inhibitory receptors characteristically contain immunoglobulin-like domains extracellularly and one or more immune receptor tyrosine-based inhibitory motifs (ITIM) in their cytoplasmic tails which consists of the consensus sequence I/L/VxYxxL/V (Vély et al., *J. Immunol.*, 159:2075-2077, 1997). In the case of CTLA-4, the ITIM is slightly modified to GxYxxM (Cambier, *Proc. Natl. Acad. Sci.*, 94:5993-5995, 1997). ITIM-containing receptors function in the recruitment of either the *Src* homology domain-containing protein tyrosine phosphatases, SHP-1 and SHP-2, or the SH2 domain-containing inositol polyphosphate 5-phosphatase, SHIP (Isakov, *Immunol. Res.*, 16:85-100, 1997). These phosphatases function in the dephosphorylation of signaling molecules recruited by immune receptor tyrosine-based activation motif (ITAM) bearing receptors like those contained in the CD3- γ , δ , ϵ and ζ chains that associate with the TCR. As such, ITIM bearing receptors on T cells are predicted to downregulate activation events elicited by ITAM bearing receptors if both are ligated in close proximity to one another. Importantly, neither CTLA-4 nor CD80/CD86 have been observed on human iIELs or IECs of the small intestine, respectively.

25

Summary of the Invention

It has now been discovered that biliary glycoprotein (BGP; also known as CD66a and C-CAM), a member of the carcinoembryonic antigen family (CEA), is an inhibitory receptor for activated T cells contained within the human intestinal epithelium. These studies suggest that, in a regional microenvironment that is predominantly CD28/CTLA4-CD80/CD86 negative, other receptor-ligand interactions may provide necessary downregulatory signals to limit T cell activation and immunopathology.

30

According to one aspect of the invention, methods for enhancing specifically the

cytotoxicity or proliferation of killer T cells in a subject are provided. The methods include administering to a subject in need of such treatment an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance the cytotoxicity or proliferation of killer T cells in the subject. In certain embodiments the agent is an antibody or antibody fragment which binds only a single biliary glycoprotein polypeptide. Preferred antibody fragments include Fab fragments. In other embodiments the agent comprises a ligand for the biliary glycoprotein polypeptide, wherein the ligand binds only a single biliary glycoprotein polypeptide. In preferred embodiments the ligand is fused to an immunoglobulin molecule or a fragment thereof, or is a soluble biliary glycoprotein molecule or fragment thereof.

According to another aspect of the invention, methods for suppressing specifically the cytotoxicity or proliferation of killer T cells in a subject are provided. The methods include administering to a subject in need of such treatment an agent that selectively increases cross-linking of biliary glycoprotein polypeptides in an amount effective to suppress the activity of killer T cells in the subject. In certain embodiments the agent is an antibody, preferably a monoclonal antibody. In other embodiments the agent comprises a ligand for the biliary glycoprotein polypeptide which binds two or more biliary glycoprotein polypeptides. In preferred embodiments, the ligand is fused to an immunoglobulin molecule or a fragment thereof, or the ligand includes a biliary glycoprotein polypeptide or fragment thereof.

According to another aspect of the invention, a composition is provided. The composition includes an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance cytotoxicity or proliferation of killer T cells in a subject, and a pharmaceutically-acceptable carrier. In certain embodiments the agent is an antibody or antibody fragment which binds only a single biliary glycoprotein molecule. Preferred antibody fragments include Fab fragments. In preferred embodiments, the agent comprises a ligand for the biliary glycoprotein polypeptide which binds only a single biliary glycoprotein polypeptide. Preferably such a ligand is fused to an immunoglobulin molecule or a fragment thereof. In certain embodiments ligand is biliary glycoprotein or a fragment thereof. Preferably the compositions are pharmaceutical compositions.

According to still another aspect of the invention, a composition is provided which includes an agent that selectively increases cross-linking of biliary glycoprotein polypeptides in an amount effective to suppress cytotoxicity or proliferation of killer T cells in a subject. The composition also include a pharmaceutically-acceptable carrier. In some embodiments the agent

is an antibody, preferably a monoclonal antibody. In other embodiments the agent includes a ligand for the biliary glycoprotein polypeptide which binds two or more biliary glycoprotein polypeptides. Preferably the ligand is fused to an immunoglobulin molecule or a fragment thereof. In other preferred embodiments the ligand is biliary glycoprotein or a fragment thereof.

5 Preferably the compositions are pharmaceutical compositions.

According to still other aspects of the invention, methods for enhancing specifically the cytotoxicity or proliferation of killer T cells also are provided. The methods include contacting a population of killer T cells with an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance the cytotoxicity or proliferation of
10 the killer T cells. The biliary glycoprotein binding agents are as described above in methods for enhancing killer T cell activity in a subject.

In another aspect of the invention, methods for suppressing specifically cytotoxicity or proliferation of killer T cells are provided. The methods include contacting a population of killer T cells with an agent that selectively increases cross-linking of biliary glycoprotein polypeptides
15 in an amount effective to suppress the cytotoxicity or proliferation of the killer T cells. The biliary glycoprotein binding agents are as described above in methods for suppressing killer T cell activity in a subject.

According to another aspect of the invention, an isolated fusion protein is provided. The isolated fusion protein includes a biliary glycoprotein polypeptide or a fragment thereof fused to
20 an immunoglobulin molecule or a fragment thereof. The components of the fusion protein can be fused directly, or a linker molecule such as a peptide can be interposed between the biliary glycoprotein component and the immunoglobulin component. Other polypeptides can be substituted for the immunoglobulin component as will be apparent to one of ordinary skill in the art. In certain embodiments, biliary glycoprotein (or fragment thereof) component of the fusion
25 protein selectively binds a monoclonal antibody selected from the group consisting of 34B1, 5F4 and 26H7. Preferably the fragment of biliary glycoprotein is selected from the group consisting of the N-domain of CD66a, NA1B1 domains of CD66a, the NA1B1A2 domains of CD66a. The fragment of the immunoglobulin molecule preferably is the Fc portion of the immunoglobulin molecule.

30 According to another aspect of the invention, an isolated fusion protein is provided which includes two or more biliary glycoprotein polypeptides, or fragments thereof which bind biliary glycoprotein. The fusion protein is useful for selecting biliary glycoprotein binding agents which

bind two (or more) biliary glycoprotein molecules, particularly those agents which cross-link biliary glycoprotein molecules.

According to yet another aspect of the invention, methods for identifying compounds which enhance or suppress killer T cell activity are provided. The methods include contacting a population of killer T cells which express biliary glycoprotein with a compound that binds biliary glycoprotein, and determining the cytotoxicity or proliferation of the population of killer T cells relative to a control. Compounds which increase the cytotoxicity or proliferation are compounds which enhance the killer T cell activity, and compounds which decrease the cytotoxicity or proliferation are compounds which suppress the killer T cell activity. In certain embodiments, the methods includes the steps of providing a biliary glycoprotein polypeptide or a fragment thereof, contacting the biliary glycoprotein polypeptide or a fragment thereof with a compound, and determining the binding of the compound to the biliary glycoprotein polypeptide or a fragment thereof. The compound is used in the foregoing methods for testing the increase or decrease of killer T cell cytotoxicity or proliferation.

In another aspect of the invention, methods for selectively treating a subject having a condition characterized by aberrant killer T cell activity are provided. The methods include administering to a subject in need of such treatment a pharmacological agent which is selective for biliary glycoprotein, in an amount effective to normalize the aberrant killer T cell activity.

In the foregoing aspects and embodiments of the invention, preferred killer T cells include CD4⁺ T cells, CD8⁺ T cells, NK cells, intestinal intraepithelial lymphocytes and peripheral blood T cells. Particularly preferred killer T cells are CD8⁺ T cells.

The use of the foregoing compositions in the preparation of medicament also is provided. In preferred embodiments, the medicament is useful in the treatment of conditions related to immune system function, including autoimmune disease, cancer and transplantation.

These and other objects of the invention will be described in further detail in connection with the detailed description of the invention.

Brief Description of the Drawings

Fig. 1 depicts identification of three mAbs (34B1, 5F4 and 26H7) which recognize IECs but not resting iIELs.

Fig. 2 shows that iIELs and PBTs express an antigen that is recognized by the 34B1, 5F4 and 26H7 mAbs after activation.

Fig. 3 shows that the N-domain of BGP is the cognate antigen of the 34B1-related mAbs.

Fig. 4 shows that the 34B1-related mAbs specifically immunoprecipitate BGP on COS cell transfectants and activated iIELs.

Fig. 5 depicts the specificity of the anti-BGP mAbs for other CD66 family members.

5 Fig. 6 shows an analysis of CD66 isoform expression by activated human iIELs.

Fig. 7 depicts inhibition of anti-CD3 directed and lymphokine activated killer activity of iIELs by anti-BGP antibodies.

Fig. 8 shows inhibition of human allo-mixed lymphocyte reaction by anti-BGP monoclonal antibodies.

10

Detailed Description of the Invention

Evidence for a role of human biliary glycoprotein (BGP, CD66a) as an inhibitory molecule on activated iIELs has been provided through the characterization of three monoclonal antibodies raised against an activated iIEL cell line. These data are especially relevant to iIELs
15 as they suggest that other molecules such as biliary glycoprotein may contribute to downregulation of T cell activation in the absence of CTLA-4. These studies are also relevant to extending the function of biliary glycoprotein to an important role in immunoregulation of T lymphocytes in general given the observation that biliary glycoprotein is also expressed by activated human peripheral blood T cells.

20 Human biliary glycoprotein is a member of the CEA family of glycoproteins, part of the immunoglobulin supergene family, and encoded in a large cluster on chromosome 19 (Watt et al., *Blood*, 84:200-210, 1994; Daniel et al., *Int. J. Cancer*, 55:303-310, 1993; Teixeira et al., *Blood*, 84:211-219, 1994; Teixeira, 1996; Barnett et al., *Molec. Cell. Biol.*, 13:1273-1282, 1993). The CEA-cluster is highly related to the genetically linked, pregnancy specific gene cluster
25 (Thompson et al., *J. Clin. Lab Anal.*, 5:344-366, 1991; Öbrink, *Curr Opin Cell Biol.*, 9:616-626, 1997). The CEA-subgroup of this family is serologically defined as CD66a (BGP or C-CAM), CD66b (CGM6), CD66c (NCA), CD66d (CGM1) and CD66e (CEA). These structurally related glycoproteins consist of a highly homologous membrane distal amino terminal IgV-like N-domain and variable numbers of membrane distal IgC2-like domains in the case of BGP, NCA,
30 CGM6 and CEA. In contrast to human CEA, CGM6 and NCA, which are linked to the membrane by a glycosyl phosphatidyl-inositol anchor, CGM1 and BGP are type 1 transmembrane glycoproteins. Both of the latter exist as isoforms containing short or long

cytoplasmic tails.

Biliary glycoprotein, and its mouse and rat homologues C-CAM (Rosenberg et al., *Cancer Res.*, 53:4938-4945, 1993; Lin et al., *J. Biol. Chem.*, 264:14408-14414, 1989; Öbrink, *BioEssays*, 13:227-234, 1991), have been regarded mainly as molecules which function in cell-cell adhesion that are expressed primarily by epithelial cells of the gastrointestinal tract and biliary tree, neutrophils and, more recently, B cells. Biliary glycoprotein also serves as a receptor for mouse hepatitis virus (Williams et al., *Proc. Natl. Acad. Sci.*, 88:5533-5536, 1991) and for *Opa* proteins of *Neisseria* species of bacteria (Virji et al., *Mol. Microbiol.*, 5:941-950, 1996). It is of interest that ligation of biliary glycoprotein on epithelial cells may deliver a negative growth signal which may be decreased during tumor formation due to diminished expression of biliary glycoprotein (Rosenberg et al., 1993; Kunath et al., *Oncogene*, 11:2375-2382, 1995; Brümmer et al., *Oncogene*, 11:1649-1655, 1995). Biliary glycoprotein also exhibits a high degree of alternate transcriptional processing resulting in at least eight potential alternate transmembrane transcripts. Two of these transcripts, BGP_a and BGP_b, encode a long cytoplasmic tail of 67 amino acids containing two ITIM motifs which suggest a role as inhibitory receptors (Öbrink, 1997). Indeed, this cytoplasmic tail, when tyrosine phosphorylated, is capable of binding SHP-1 in a mouse colon carcinoma cell line (Beauchemin et al., *Oncogene*, 783-790, 1996). Such interactions may account for the inhibitory growth effect of this molecule on epithelial cells.

The studies contained herein describe the unexpected finding that, whereas biliary glycoprotein is constitutively expressed by IECs, it is an activation molecule on T cells adjacent to the epithelium. The study of peripheral blood T cells, on the other hand, show the unexpected result that biliary glycoprotein is constitutively expressed at low levels and upregulated by T-cell activation. This difference between iIELs and PBTs suggests that biliary glycoprotein expression may be actively suppressed in the epithelium under normal conditions. The association of biliary glycoprotein with the activation state also resembles CTLA-4 expression.

Using cytotoxicity, which is a major function of iIELs, as a measure, it appears that biliary glycoprotein on activated iIELs functions as an inhibitory molecule for CD3-directed cytotoxic activity. In this manner, biliary glycoprotein should be considered as a killer inhibitory receptor. Although the ligand for biliary glycoprotein on the IEC is unknown, a candidate ligand is biliary glycoprotein itself or another CD66 family member in view of the known homophilic and heterophilic interactions between the CD66 group members (Watt et al., 1994; Oikawa et al.,

Biochem. Biophys. Res., 186:881-887, 1992; Teixeira et al., 1994; Öbrink, 1997). It therefore can be hypothesized that ligation of biliary glycoprotein on the IEC by an activated iIEL can serve to function in the inhibition of IEC growth. Corollarily, the binding of biliary glycoprotein on the activated iIEL by BGP on the IEC can limit the activation of the T cell. In tumors of the
5 epithelium where biliary glycoprotein expression has been observed to be diminished, the growth inhibition effect of the iIEL on the IEC might be lost (Brümmer, 1995).

Thus the invention involves the finding that molecules which bind to biliary glycoprotein (i.e., "biliary glycoprotein binding agents") on killer T cells, such as antibodies, can inhibit or enhance the activity of killer T cells, such as cytotoxicity and/or proliferation. As used herein,
10 "killer T cells" includes CD4⁺ T cells, CD8⁺ T cells, and NK cells. Biliary glycoprotein binding agents which increase cross-linking of biliary glycoprotein polypeptides increase the inhibitory signal of biliary glycoprotein, thereby suppressing the activity of killer T cells. Biliary glycoprotein binding agents which decrease cross-linking of biliary glycoprotein polypeptides decrease the inhibitory signal of biliary glycoprotein, thereby enhancing the activity of killer T
15 cells. The invention also embraces molecules which enhance or suppress killer T cell activity but which do not function according to the cross-linking properties described above. For example, a particular biliary glycoprotein binding agent which suppresses the activity of killer T cells may bind biliary glycoprotein and increase the inhibitory signal of biliary glycoprotein without increasing cross-linking (e.g., by inducing a conformational change in biliary
20 glycoprotein).

Modulation of killer T cell activity by molecules which bind biliary glycoprotein expressed on the surface of killer T cells is useful for specifically enhancing or suppressing an immune response *in vivo*, which may be useful for the treatment of conditions related to immune function including autoimmune disease, cancer, and transplantation (e.g., bone marrow or
25 organs). Modulation of killer T cell activity also is useful in *in vitro* and/or non-therapeutic applications including determining whether T cells of a subject are functional (e.g. proliferation and/or cytotoxic functions), to determine if a treatment has rendered killer T cells non-functional, in experimental models of cancer, autoimmune disease, and transplantation, e.g., to determine the effects of increases or decreases in killer T cell function on particular organs or physiological
30 processes, and to test for agents which increase or decrease killer T cell activity. Other uses will be apparent to one of ordinary skill in the art.

The molecules which bind biliary glycoprotein and modulate killer T cell activity (biliary

glycoprotein binding agents) include antibodies and fragments thereof, ligands for biliary glycoprotein, fragments thereof and fusion proteins containing ligands or other biliary glycoprotein binding molecules. Still other biliary glycoprotein binding agents can be identified by screening compounds for the ability to enhance or suppress killer T cell activity, using assays
5 described herein and those assays of T cell activity which are standard in the art. Exemplary methods for preparing fusion proteins useful according to the invention for modulating killer T cell activity are described herein; additional exemplary methods for preparing such fusion proteins are described in U.S. Patent 5,434,131. The molecules which bind biliary glycoprotein can be used alone as a primary therapy or in combination with other therapeutics as a
10 combination therapy to enhance the therapeutic benefits of other medical treatments.

The invention also involves agents which bind to biliary glycoprotein and/or fragments of the biliary glycoprotein and induce or suppress killer T cell activity. In addition to the uses described herein, such binding agents can be used in screening assays to detect the presence or absence of a biliary glycoprotein polypeptide, the presence or location of iIELs and in
15 purification protocols to isolate iIELs and other killer T cells which express biliary glycoprotein. Likewise, such binding agents can be used to selectively target drugs, toxins or other molecules to killer T cells which express biliary glycoprotein. In this manner, killer T cells which express biliary glycoprotein can be treated with cytotoxic compounds, thereby reducing unwanted immune responses.

The biliary glycoprotein binding agents useful according to the invention, including
20 antibodies and other polypeptides, are isolated agents. As used herein, with respect to biliary glycoprotein binding agents, the term "isolated" means that the agents are substantially pure and are essentially free of other substances with which they may be found in nature or *in vivo* systems to an extent practical and appropriate for their intended use. In particular, the agents are
25 sufficiently pure and are sufficiently free from other biological constituents of their hosts cells so as to be useful in, for example, producing pharmaceutical preparations. Because an isolated biliary glycoprotein binding agent may be admixed with a pharmaceutically acceptable carrier in a pharmaceutical preparation, the biliary glycoprotein binding agents may comprise only a small percentage by weight of the preparation. A biliary glycoprotein binding agent is nonetheless
30 substantially pure in that it has been substantially separated from the substances with which it may be associated in living systems.

According to one embodiment, the biliary glycoprotein binding agent used in the

invention is an intact anti-biliary glycoprotein monoclonal antibody in an isolated form, preferably in a soluble form, or in a pharmaceutical preparation. An intact monoclonal antibody, as is well known in the art, is an assembly of polypeptide chains linked by disulfide bridges. Two principle polypeptide chains, referred to as the light chain and heavy chain, make up all major structural classes (isotypes) of antibody. Both heavy chains and light chains are further divided into subregions referred to as variable regions and constant regions. As used herein the term "monoclonal antibody" refers to a homogenous population of immunoglobulins which specifically bind to an epitope (i.e. antigenic determinant) of human biliary glycoprotein.

The invention, therefore, includes the use of antibodies or fragments of antibodies having the ability to selectively bind to biliary glycoprotein, particularly as expressed on the cell surface of killer T cells, such as intestinal intraepithelial lymphocytes and activated peripheral blood lymphocytes. Antibodies include polyclonal and monoclonal antibodies, prepared according to conventional methodology. Examples include the monoclonal antibodies 34B1, 5F4 and 26H7 described in the Examples. Additional antibodies which are reactive with biliary glycoprotein, particularly those raised against biliary glycoprotein expressed on killer T cells, can be prepared according to standard methods.

Antibodies can be prepared by any of a variety of methods, including administering protein, fragments of protein, cells expressing the protein or fragments thereof and the like to an animal to induce polyclonal antibodies. The production of monoclonal antibodies is according to techniques well known in the art. As detailed herein, such antibodies may be used, for example, to identify tissues expressing protein or to purify protein. Antibodies also may be coupled to specific labeling agents for imaging or to cytotoxic agents, including, but not limited to, methotrexate, radioiodinated compounds, toxins such as ricin, other cytostatic or cytolytic drugs, and so forth.

Significantly, as is well-known in the art, only a small portion of an antibody molecule, the paratope, is involved in the binding of the antibody to its epitope (see, in general, Clark, W.R. (1986) The Experimental Foundations of Modern Immunology Wiley & Sons, Inc., New York; Roitt, I. (1991) Essential Immunology, 7th Ed., Blackwell Scientific Publications, Oxford). The pFc' and Fc regions, for example, are effectors of the complement cascade but are not involved in antigen binding. An antibody from which the pFc' region has been enzymatically cleaved, or which has been produced without the pFc' region, designated an F(ab')₂ fragment, retains both of the antigen binding sites of an intact antibody. Similarly, an antibody from which

the Fc region has been enzymatically cleaved, or which has been produced without the Fc region, designated an Fab fragment, retains one of the antigen binding sites of an intact antibody molecule. Proceeding further, Fab fragments consist of a covalently bound antibody light chain and a portion of the antibody heavy chain denoted Fd. The Fd fragments are the major
5 determinant of antibody specificity (a single Fd fragment may be associated with up to ten different light chains without altering antibody specificity) and Fd fragments retain epitope-binding ability in isolation.

Within the antigen-binding portion of an antibody, as is well-known in the art, there are complementarity determining regions (CDRs), which directly interact with the epitope of the
10 antigen, and framework regions (FRs), which maintain the tertiary structure of the paratope (see, in general, Clark, 1986; Roitt, 1991). In both the heavy chain Fd fragment and the light chain of IgG immunoglobulins, there are four framework regions (FR1 through FR4) separated respectively by three complementarity determining regions (CDR1 through CDR3). The CDRs, and in particular the CDR3 regions, and more particularly the heavy chain CDR3, are largely
15 responsible for antibody specificity.

It is now well-established in the art that the non-CDR regions of a mammalian antibody may be replaced with similar regions of conspecific or heterospecific antibodies while retaining the epitopic specificity of the original antibody. This is most clearly manifested in the development and use of "humanized" antibodies in which non-human CDRs are covalently
20 joined to human FR and/or Fc/pFc' regions to produce a functional antibody. Thus, for example, PCT International Publication Number WO 92/04381 teaches the production and use of humanized murine RSV antibodies in which at least a portion of the murine FR regions have been replaced by FR regions of human origin. Such antibodies, including fragments of intact antibodies with antigen-binding ability, are often referred to as "chimeric" antibodies.

Thus, as will be apparent to one of ordinary skill in the art, the present invention also
25 provides for F(ab')₂, Fab, Fv and Fd fragments; chimeric antibodies in which the Fc and/or FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; chimeric F(ab')₂ fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human
30 or non-human sequences; chimeric Fab fragment antibodies in which the FR and/or CDR1 and/or CDR2 and/or light chain CDR3 regions have been replaced by homologous human or non-human sequences; and chimeric Fd fragment antibodies in which the FR and/or CDR1 and/or

CDR2 regions have been replaced by homologous human or non-human sequences. The present invention also includes so-called single chain antibodies. Thus, the invention involves polypeptides of numerous size and type that bind specifically to biliary glycoprotein. These polypeptides may be derived also from sources other than antibody technology. For example, such polypeptide binding agents can be provided by degenerate peptide libraries which can be readily prepared in solution, in immobilized form or as phage display libraries. Combinatorial libraries also can be synthesized of peptides containing one or more amino acids. Libraries further can be synthesized of peptoids and non-peptide synthetic moieties.

Phage display can be particularly effective in identifying binding peptides useful according to the invention, using methods such as those described in Hart et al., *J. Biol. Chem.* 269:12468 (1994). Briefly, one prepares a phage library (using e.g. m13, fd, or lambda phage), displaying inserts from 4 to about 80 amino acid residues using conventional procedures. The inserts may represent a completely degenerate or biased array. One then can select phage-bearing inserts which bind to biliary glycoprotein or a fragment thereof. This process can be repeated through several cycles of reselection of phage that bind to biliary glycoprotein or a fragment thereof. Repeated rounds lead to enrichment of phage bearing particular sequences. DNA sequence analysis can be conducted to identify the sequences of the expressed polypeptides. The minimal linear portion of the sequence that binds to biliary glycoprotein or a fragment thereof can be determined. One can repeat the procedure using a biased library containing inserts containing part or all of the minimal linear portion plus one or more additional degenerate residues upstream or downstream thereof. Thus, biliary glycoprotein can be used to screen peptide libraries, including phage display libraries, to identify and select peptide biliary glycoprotein binding agents for modulating killer T cell activity. Preferably, the biliary glycoprotein binding agents are characterized as to their ability to cross-link biliary glycoprotein. Such binding molecules can also be used, as described, for screening assays, for diagnostic assays, for purification protocols or for targeting drugs, toxins and/or labeling agents (e.g. radioisotopes, fluorescent molecules, etc.) to cells, especially killer T cells, which express biliary glycoprotein on the cell surface. Drug molecules that would disable or destroy cells which express such biliary glycoprotein or a fragment thereof are known to those skilled in the art and are commercially available. For example, the immunotoxin art provides examples of toxins which are effective when delivered to a cell by an antibody or fragment thereof. Examples of toxins include ribosome-damaging toxins derived from plants or bacterial such as ricin, abrin,

saporin, *Pseudomonas* endotoxin, diphtheria toxin, A chain toxins, blocked ricin, etc.

Additionally small polypeptides including those containing the biliary glycoprotein binding fragment (CDR3 region) may easily be synthesized or produced by recombinant means to produce a biliary glycoprotein binding agent useful according to the invention. Such methods are well known to those of ordinary skill in the art. Peptides can be synthesized for example, using automated peptide synthesizers which are commercially available. The peptides can be produced by recombinant techniques by incorporating the DNA expressing the peptide into an expression vector and transforming cells with the expression vector to produce the peptide.

The sequence of the CDR regions, for use in synthesizing peptide biliary glycoprotein binding agents, may be determined by methods known in the art. The heavy chain variable region is a peptide which generally ranges from 100 to 150 amino acids in length. The light chain variable region is a peptide which generally ranges from 80 to 130 amino acids in length. The CDR sequences within the heavy and light chain variable regions which include only approximately 3-25 amino acid sequences may easily be sequenced by one of ordinary skill in the art. The peptides may even be synthesized by commercial sources.

To determine whether a peptide binds to biliary glycoprotein any known binding assay may be employed. For example, the peptide may be immobilized on a surface and then contacted with labeled biliary glycoprotein. The amount of biliary glycoprotein which interacts with the peptide or the amount which does not bind to the peptide may then be quantitated to determine whether the peptide binds to biliary glycoprotein. A surface having the aforementioned anti-biliary glycoprotein monoclonal antibodies immobilized thereto may serve as a positive control.

Screening of biliary glycoprotein binding agents also can be carried out utilizing a competition assay. If the biliary glycoprotein binding agent being tested competes with an anti-biliary glycoprotein monoclonal antibody, as shown by a decrease in binding of the monoclonal antibody, then it is likely that the agent and the anti-biliary glycoprotein monoclonal antibody bind to the same, or a closely related, epitope. Still another way to determine whether an agent has the specificity of the anti-biliary glycoprotein monoclonal antibodies described above is to pre-incubate the monoclonal antibody with biliary glycoprotein with which it is normally reactive (i.e., binds), and then add the agent being tested to determine if the agent being tested is inhibited in its ability to bind biliary glycoprotein. If the agent being tested is inhibited then, in all likelihood, it has the same or a functionally equivalent epitope and specificity as the anti-

biliary glycoprotein monoclonal antibodies.

Using routine procedures known to those of ordinary skill in the art, one can determine whether a biliary glycoprotein binding agent is useful according to the invention by determining whether the agent is one which modulates killer T cell proliferation or cytotoxicity in an *in vitro* assay such as measuring release of TNF from killer T cells or by ⁵¹Cr release assay (see, e.g., 5 Herin et al., *Int. J. Cancer* 39:390-396, 1987). Other assays are described in the Examples and elsewhere herein.

The polypeptides (e.g. antibodies) and other biliary glycoprotein binding agents described above can also be used immunotherapeutically for killer T cell sensitive disorders in humans. 10 The term "immunotherapeutically" or "immunotherapy" as used herein in conjunction with the biliary glycoprotein binding agents denotes both prophylactic as well as therapeutic administration. Thus, the peptides can be administered to high-risk subjects in order to lessen the likelihood and/or severity of a killer T cell sensitive disease, such as a tumor, transplant rejection or autoimmune disease, or administered to subjects already evidencing such diseases.

15 In certain aspects the invention encompasses methods for modulating specifically the cytotoxicity or proliferation of killer T cells *in situ*. The method includes administering to a subject in need of such treatment an agent which binds selectively a biliary glycoprotein polypeptide in an amount effective to enhance or suppress the cytotoxicity or proliferation of the killer T cells in the subject. As shown in the Examples, the activity of killer T cells is subject to 20 specific modulation because killer T cells express biliary glycoprotein. Methods for modulating specifically the cytotoxicity or proliferation of killer T cells also are provided wherein a population of killer T cells is contacted with a biliary glycoprotein binding agent. When a biliary glycoprotein binding agent is administered to a subject or contacted to a population of killer T cells, the inhibitory activity of biliary glycoprotein is modulated. Biliary glycoprotein binding 25 agents which increase or decrease killer T cell activity can be selected using the assays described herein and according to standard killer T cell cytotoxicity and proliferation assays, such as mixed lymphocyte reactions, chromium release assays, TNF release assays, and thymidine incorporation assays. It is believed that a monovalent biliary glycoprotein binding agent will inhibit the inhibitory signal of biliary glycoprotein by reducing cross-linking of biliary 30 glycoprotein polypeptides expressed by killer T cells, and that a multivalent biliary glycoprotein binding agent (having two or more biliary glycoprotein binding sites) will increase the inhibitory signal of biliary glycoprotein in killer T cells by increasing cross-linking of biliary glycoprotein

polypeptides expressed by killer T cells.

By definition, the term “*in situ*” encompasses and includes the terms *in vivo*, *ex vivo* and *in vitro*. The compositions of the invention are useful for many *in vitro* purposes. For example, the compositions of the invention are useful for screening compounds which inhibit killer T cell proliferation or cytotoxicity. Such a screening assay may be performed *in vitro* by setting up cell proliferation or cytotoxicity assays including a biliary glycoprotein binding agent which increases killer T cell proliferation or cytotoxicity and a population of killer T cells. Potential killer T cell proliferation or cytotoxicity inhibitors may be added to the mixture and the effect on proliferation or cytotoxicity may be measured. Agents which increase proliferation or cytotoxicity of killer T cell can be screened using similar assays. Thus methods for identifying compounds which bind biliary glycoprotein (or modulate activation of biliary glycoprotein by other molecules such as natural ligands) and enhance or suppress killer T cell activity are provided according to the invention. Other *in vitro* uses, such as research purposes, are known to those of ordinary skill in the art.

Ex vivo uses also will be readily identified by those of skill in the art. *Ex vivo* uses include, for example, the stimulation of proliferation or cytotoxicity of killer T cells which have been removed from a mammalian subject and which are subsequently returned to the body of the mammalian subject.

The present invention also includes methods for treating a condition characterized by aberrant killer T cell activity, such as cytotoxicity or proliferation. The methods involve the step of administering to a subject having such a condition a pharmacological agent which selectively binds biliary glycoprotein and modulates a killer T cell activity, in an amount effective to increase or decrease T-cell proliferation or cytotoxicity.

A “condition characterized by aberrant killer T cell activity” as used herein is any condition associated with adverse physiological consequences in which an increase or decrease of killer T cell function, embodied by an increase or decrease in killer T cell proliferation or cytotoxicity, results in an improvement of the adverse physiological consequences. Such conditions include disorders of the immune system, such as immunodeficiency, autoimmunity and transplant rejection, as well as disorders involving undesirable cellular invasion by microorganisms or undesirable cell growth such as tumors.

The biliary glycoprotein binding agents are administered in effective amounts. As used herein, an “effective amount” of a biliary glycoprotein binding agent is an amount which is

sufficient to modulate (increase or decrease) biliary glycoprotein inhibitory function, resulting in a modulation of killer T cell proliferation or cytotoxicity. Modulating biliary glycoprotein inhibition of killer T cell activity is sufficient to produce the desired effect in which the symptoms associated with the conditions characterized by aberrant killer T cell activity are ameliorated or decreased. Preferably an effective amount of the peptide is a therapeutically effective amount for modulating killer T cell proliferation or cytotoxicity *in vivo*. Generally, a therapeutically effective amount may vary with the subject's age, condition, weight and gender, as well as the extent of the disease in the subject and can be determined by one of skill in the art as a matter of routine experimentation. The dosage may be adjusted by the individual physician in the event of any complication. A therapeutically effective amount typically will vary from about 0.01 mg/kg to about 500 mg/kg, were typically from about 0.1 mg/kg to about 200 mg/kg, and often from about 0.2 mg/kg to about 20 mg/kg, in one or more dose administrations daily, for one or several days (depending of course of the mode of administration and the factors discussed above).

One of skill in the art can determine what an effective amount of a biliary glycoprotein binding agent is by determining the ability of the agent to modulate killer T cell proliferation or cytotoxicity in an *in vitro* assay. Exemplary assays for measuring the ability of a biliary glycoprotein binding agent to modulate killer T cell proliferation or cytotoxicity are provided in the Examples and have been discussed above. The exemplary assays are predictive of the ability of a biliary glycoprotein binding agent to modulate killer T cell activity *in vivo* and/or *ex vivo* and, hence, can be used to select agents for therapeutic applications.

According to the invention, a biliary glycoprotein binding agent may be administered in a pharmaceutically acceptable composition. In general, pharmaceutically-acceptable carriers for antibodies, antibody fragments, and other biliary glycoprotein binding agents (including small molecules such as those derived from combinatorial libraries) are well-known to those of ordinary skill in the art. As used herein, a pharmaceutically-acceptable carrier means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredients, i.e., the ability of the agent to modulate killer T cell activity. Pharmaceutically acceptable carriers include diluents, fillers, salts, buffers, stabilizers, solubilizers and other materials which are well-known in the art. Exemplary pharmaceutically acceptable carriers for peptides are described in U.S. Patent No. 5,211,657. The agents of the invention may be formulated into preparations in solid, semi-solid, liquid or gaseous forms such as tablets,

capsules, powders, granules, ointments, solutions, depositories, inhalants and injections, for oral, parenteral or surgical administration. The invention also embraces pharmaceutical compositions which are formulated for local administration, such as by implants.

According to the methods of the invention the agents can be administered by injection, by
5 gradual infusion over time or by any other medically acceptable mode. The administration may, for example, be intravenous, intraperitoneal, intramuscular, intracavity, subcutaneous or transdermal. Preparations for parenteral administration includes sterile aqueous or nonaqueous solutions, suspensions and emulsions. Examples of nonaqueous solvents are propylene glycol, polyethylene glycol, vegetable oil such as olive oil, an injectable organic esters such as
10 ethylolate. Aqueous carriers include water, alcoholic/aqueous solutions, emulsions or suspensions, including saline and buffered media. Parenteral vehicles include sodium chloride solution, Ringer's dextrose, dextrose and sodium chloride, lactated Ringer's or fixed oils. Intravenous vehicles include fluid and nutrient replenishers, electrolyte replenishers (such as those based on Ringer's dextrose), and the like. Preservatives and other additives may also be
15 present such as, for example, antimicrobials, antioxidants, chelating agents, and inert gases and the like. Those of skill in the art can readily determine the various parameters for preparing these alternative pharmaceutical compositions without resort to undue experimentation.

The methods of the invention also encompass administering biliary glycoprotein binding agents in conjunction with conventional therapies for treating immune system disorders. For
20 example, the methods of the invention may be practiced simultaneously with conventional treatments. The particular conventional treatment depends, of course, on the nature of the disorder. When, for example, the condition related to aberrant killer T cell activity is a tumor, a conventional mode of treatment is chemotherapy. The agents of the invention which increase killer T cell activity (e.g., decrease biliary glycoprotein inhibitory activity) may be administered
25 in conjunction with chemotherapy in the treatment of the tumor in order to provide enhanced tumoricidal effects. Other immune system diseases can be treated concurrently with the biliary glycoprotein binding agents described herein and other molecules which bind T cells and affect T cell function, such as CTLA4Ig fusion proteins, as described in U.S. Patent 5,434,131.

The following examples are provided to illustrate specific instances of the practice of the
30 present invention and are not to be construed as limiting the present invention to these examples. As will be apparent to one of ordinary skill in the art, the present invention will find application in a variety of compositions and methods.

Examples

Materials and Methods

Antibodies: The 34B1, 26H7 and 5F4 monoclonal antibodies (mAbs) were produced by immunizing BALB/c mice with the activated human mucosal lymphocyte line, 191E, as previously described (Russell et al., 1996). Three intraperitoneal injections and a final intravenous injection of 5×10^6 lymphocytes were given at two-week intervals. Three days after the intravenous immunization, splenocytes were isolated and fused with NS1 murine myeloma cells in the presence of PEG (m.w. 1450) as described previously. Hybridomas were selected with aminopterin-containing medium, and hybridoma supernatants were screened by indirect immunoperoxidase staining of frozen intestinal and tonsillar tissue sections. Positive hybridomas were subcloned twice by limiting dilution, and ascites containing the antibody was produced by intraperitoneal injection of the hybridoma cells into pristane-treated BALB/c mice. The isotypes of 34B1 (IgG1), 26H7 (IgG1) and 5F4 (IgG1) were determined by ELISA using murine isotype-specific mAb (Boehringer Mannheim, Indianapolis, IN). W6/32 is a mouse IgG2a mAb specific for human MHC class I (kindly provided by Dr. Jack Strominger, Dana-Farber Cancer Institute, Boston, MA). OKT3 (IgG2a is a mouse anti-human CD3 mAb (kindly provided by Dr. Robert Finberg, Dana-Farber Cancer Institute, Boston, MA). TS2/18 (kindly provided by Dr. Llyod Klickstein, Brigham and Women's Hospital) is an anti-CD2 mAb (mouse IgG2a). OKT11 (kindly provided by Dr. Ellis Reionherz, Dana-Farber Cancer Institute) is an anti-CD2 mAb (mouse IgG29). OKT4 and OKT8 are mouse IgG2a mAbs specific for human CD4 and CD8- α , respectively (obtained from American Type Culture Collection, Bethesda, MD). UCHT1, directly conjugated to phycoerythrin, is a mouse IgG1 mAb specific for human CD3- ϵ (Dako, Denmark). The MA22 (CD66a; clone YG-C94G7; IgG1), MA26 (CD66a; clone 4.3.17; IgG1), MA27 (CD66e; clone 26/5/1; IgG2a), MA28 (CD66e; clone 26/3/13; IgG1), MA30 (CD66c; clone 9A6; IgG1), MA41 (CD66b; clone BIRMA17c; IgG1), MA61 (CD66b; clone 80H3; IgG1), MA76 (CD66ae; clone 12-140-4; IgG1), MA79 (CD66b; clone B13.9; IgG1), MA81 (CD66b; clone G10F5; IgG1), MA83 (CD66e; clone b7.8.5; IgG1), MA84 (CD66de; clone COL-1; IgG2a), MA86 (CD66acde; clone B6.2; IgG1) and MA91 (CD66e; clone T84.66; IgG1) are mouse mAbs which were obtained from the VIth Leukocyte Typing Workshop, Osaka, Japan. The isotype matched mouse IgG1 negative control mAb was purchased from Kakopatts (Copenhagen, Denmark) or Cappel (West Chester, PA). mAbs were purified by affinity purification and protein-A or G sepharose columns by standard methods.

Cells and Cell Lines: Peripheral blood mononuclear cells (PBMC) were obtained by Ficoll-Hypaque gradient centrifugation using standard methods. Peripheral blood T cells were stimulated by cultivating PBMCs for 72 hours at 37°C in RPMI-1640 (Gibco, Grand Island, NY) containing penicillin/streptomycin (100 units/ml), 10 mM Hepes pH7.4, 10% fetal calf serum and 1 µg/ml phytohemagglutinin-P (PHA-P) (Murex Diagnostics, Dartford, England). Human iIEL cell lines EEI-10 (small intestine), EEI-5 (small intestine) and CLI (large intestine) were generated as previously described (Christ et al., *Immunol. Let.*, 58:159-165, 1997) and maintained by stimulation with 1 µg/ml PHA-P cells in RPMI-1640 containing 10% human serum (type AB, Sigma, St. Louis, MO), 5 units/ml rIL-4 (Genzyme, Cambridge, MA) and 2 nM rIL-2 (a kind gift from Ajinomoto Co., Ltd., Japan) and irradiated PBMC as feeder cells. HT29 is a human intestinal epithelial cell (IEC) line obtained from the ATCC. COS is a monkey kidney fibroblast cell line. These latter cell lines were maintained in RPMI-1640 containing 10% heat-inactivated fetal calf serum (Gibco), penicillin and streptomycin, nonessential amino acids and 10 mM HEPES (complete medium) at 37°C in 5% CO₂.

Transfectants: The BGPx' molecule was constructed as follows. The N terminal domain and the transmembrane/ cytoplasmic domains of human BGPc were each amplified separately by PCR with the primer pairs BGPAMP-S: caucaucaucauaagettatggggcacctc (SEQ ID NO:1) and NTM-AS: gccattttctggggcabctccgggtatac (SEQ ID NO:2); NTM-S: gtatacccgagactgcccccaagaaaatggc (SEQ ID NO:3) and BGPTRANS-CYT-AS: cuacuacuacuaagactatgaagttggttg (SEQ ID NO:4), respectively, where the NTM primers were hybrids of the 3' end of the N terminal domain and the 5' end of the transmembrane domain. The PCR reaction consisted of 5 µl 10xTaq buffer (10 mM Tris-HCl pH8.3, 50 mM KCl, 0.1% gelatin), 3 µl 1.5 mM MgCl₂, 1µl 200 µM of each dNTP, 1 µM of each primer, 1 Unit Taq polymerase and 1 µg cDNA in a final volume of 50 µl. The PCR reaction was carried out at conditions of 94°C for 10 min, followed by 25 cycles of 94°C for 1 min, 55°C for 1 min, and 72°C for 2 min, plus a final extension time of 10 min at 72°C. After passing the PCR products through S-300 columns, 5 µl of each PCR product were used in a second PCR. After the PCR products had annealed, the BGPAMP and TRANS-CYT AS primers were added to the reaction mix and the PCR reaction carried out as described above. The resulting PCR product was cloned into the pAMP 1 vector using the CloneAMP system as detailed by the manufacturer (Gibco-BRL, Gaithersburg, MD), transformed into DH5α competent bacteria and positive transformants selected by PCR. The resulting BGPx' cDNA was extracted and sequenced by standard

methods. The BGPx' cDNA was digested with *EcoRI* and *NotI* restriction enzymes and subcloned into the pcDNA1/Amp vector (Invitrogen, Carlsbad, CA). The BGPx' cDNA in this vector and the pSV2neo plasmid (Clontech, Palo Alto, CA) were linearized with *XhoI* and *BamHI*, respectively, and electroporated into CHO cells at a ratio of 15:1, selected in G418 and on the FACS cell sorter to create a stable CHO-BGPx' cell line as described earlier (Watt et al., 1994). CHO cells stably transfected with BGPx', Neomycin (Neo), BGPc (Watt et al., 1994) and BGPa (Oikawa et al., *Biochem. Biophys. Res. Commun.*, 186:881-887, 1992) and HeLa cells stably transfected with CEA, CGM1, NCA and CGM6 have been previously described (Daniel et al., 1993).

10 *Flow Cytometry:* For one color immunofluorescence, approximately 1×10^6 cells were stained with 1 μg of the primary mAb for 30 minutes on ice. Cells were then washed with phosphate buffered saline (PBS) containing 2% fetal calf serum or 0.2% bovine serum albumin and 0.02% sodium azide (WB) followed by incubation for 30 minutes with 1 μg of a goat-anti-mouse fluorescein isothiocyanate labeled secondary antibody (Zymed, San Francisco, CA) 15 diluted in WB. The cells were washed with WB and either examined fresh or resuspended in 1% paraformaldehyde in PBS. The stained and fixed cells were analyzed on either an Epics V flow cytometer (Coulter, Hialeah, FL) or a FACSCalibur (Becton-Dickinson, Sunnyvale, CA). For two color immunofluorescence, staining was performed as described above except that after incubation with primary conjugate, the stained cells were blocked for 20 minutes in PBS 20 containing 20% normal mouse serum followed by incubation with a directly conjugated mAb.

Immunohistochemistry: Tissue samples were mounted in OCT compound (Ames Co., Elkart, IN), frozen in liquid nitrogen or in a cryostat and stored at -70°C . Frozen tissue sections 4 μm thick were fixed in acetone for 5 minutes, air dried, and stained by an indirect immunoperoxidase method (Canchis et al., *Immunology*, 80:561-569, 1994) using avidin-biotin- 25 peroxidase complex (Vector Laboratories, Burlingame, CA) and 3-amino-9-ethylcarbazole (Aldrich Chemical Co., Inc., Milwaukee, WI) as the chromogen.

COS cell expression cloning: A cDNA library was constructed in the pCDM8 vector using poly(A)⁺ RNA from resting and activated human PBT and from NK cells in the vector pAEXF (Hall et al., *Proc. Natl. Acad. Sci.*, 93:11780-11785, 1996). For the first round of 30 selection, COS cells were transfected via the DEAE-Dextran procedure (Seed et al., *Proc. Natl. Acad. Sci.*, 84:3365-3369, 1987) with 0.2 μg of library DNA per 100 mm dish. After 40 hr, cells were harvested, incubated with 34B1 mAb (1:500 dilution of ascites), washed, and panned on

anti-IgG1 coated plates as previously described (Seed et al., 1987; Freeman et al., *J. Immunol.*, 143:2714-2722, 1989). Episomal DNA was prepared from adherent cells, re-introduced into *E. coli*, transfected into COS cells by polyethylene glycol-mediated fusion of spheroplasts (Seed et al., 1987), and the panning with 34B1 mAb repeated. Individual plasmid DNAs were transfected
5 into COS cells via the DEAE-Dextran procedure and analyzed after 72 hr for cell surface expression by indirect immunofluorescence and flow cytometry.

Radiolabeling, immunoprecipitation and electrophoresis: COS cells, 96 hours after transient transfection, were removed nonenzymatically from plastic Petri dishes and labeled with Na-[¹²⁵I] by the lactoperoxidase catalyzed method as previously described (Balk et al.,
10 *Science*, 265:259-262, 1994). After washing, radiolabeled cells were lysed in immunoprecipitation buffer (IPB) containing 150 mM sodium chloride, 50 mM Tris pH 7.8, 10 mM Iodoacetamide, 1 mM EDTA (1 mM PMSF and 1 µg/ml each of leupeptin, pepstatin, aprotinin, and chymostatin with 1% Nonidet P40 as a detergent). Cells were lysed on ice for 30-60 minutes followed by centrifugation at 14,000g for 15 minutes at 4°C. The supernatant from
15 the centrifugation was then centrifuged at 100,000g at 4°C in a TL-100 ultracentrifuge. The supernatant from this centrifugation was incubated with 20 µl of packed protein-A Sepharose beads and rocked overnight at 4°C. After preclearing with either an irrelevant isotype matched mAb or normal mouse serum coupled to either protein-A or protein-G Sepharose beads, specific immunoprecipitations were performed by rocking overnight at 4°C with mAbs coupled to
20 protein-A or G sepharose (Pharmacia, Piscataway, NJ) and the beads washed with IPB containing detergent. For N-glycanase digestions, immunoprecipitates were suspended in 0.25 M Na-HP0₄ and 1 mM EDTA after boiling in 0.8% β-mercaptoethanol and 0.5% SDS. Immunoprecipitates were then either treated with 1 unit of N-glycanase (Genzyme, Boston, MA) or mock treated at 37°C overnight and resuspended in Laemmli buffer containing reducing
25 agents. Solubilized immunoprecipitates were then resolved in 12.5% polyacrylamide gels in the presence of SDS.

Production of soluble recombinant proteins: Details of the pIG plus vector (R&D Systems Europe Ltd., Abingdon, UK) containing the Fc genomic fragment of human IgG1 and incorporating the hinge (H), CH2 and CH3 domains of the construction of the CD66a-Fc soluble
30 proteins containing the N, NA1B1 and NA1B1A2 extracellular domains, Muc-18-Fc and NCAM-Fc have been described previously (Teixeira et al., 1994; Buckley et al., *J. Cell. Sci.*, 109:437, 1996; Teixeira, 1996). NCAM-Fc, Muc-18-Fc (R&D Systems) and the CD66-Fc (N-

Fc, NA1B1-Fc and NA1B1A2-Fc) cDNAs were transfected into COS cells and the secreted soluble protein purified on protein-A Sepharose as previously described (Watt et al., 1994; Teixeira et al., 1994).

Analysis of antibody binding to soluble recombinant proteins: As previously described (Teixeira et al., 1994), ninety-six well flat bottom microtiter plates (Immulon 3) were coated with 100 µl anti-human Fc (Sigma, St. Louis, MO) at a final concentration of 1 µg/ml in 10 mM Tris-HCl pH 8 overnight at 4°C. Wells were washed four times, blocked for 1 hr at room temperature with 0.25% BSA, 0.05% Tween 20 in PBS (pH 7.4), and coated overnight at 4°C with 50 µl of soluble Fc construct at a final concentration of 10 µg/ml in PBS. After washing the wells four times with PBS, 50 µl of mAb at varying dilutions in PBS were added per well for 1-2 h at room temperature. Wells were washed with PBS four times and 50 µl of 1:4000 dilution of alkaline phosphatase conjugated goat anti-mouse Ig (Boehringer-Mannheim) in PBS were added per well for 1 h at room temperature. The wells were washed four times with PBS and 200 µl paranitrophenyl phosphate substrate were added (Sigma) and developed for 15-45 min at room temperature. The absorbance at 405 nm was determined. All experiments were carried out in triplicate and repeated at least twice.

Redirected Lysis: Cytotoxicity was evaluated as previously described (Probert et al., *J Immunol.*, 158:1941-1948, 1997). Briefly, the P815 mouse mastocytoma cell line was labeled with 100 µCi [⁵¹Cr] (New England Nuclear, Boston, MA) at 37°C for 30 minutes. 2x10³ radiolabeled cells, in 100 µl complete medium, were added to 100 µl of varying concentrations of effector T cells in 100 µl of complete medium in triplicate in a 96 well V-bottom plate. Prior to addition of target cells, the effector cells were incubated for 20 minutes at room temperature with the OKT3 mAb (200 ng/ml of purified antibody) and/or 34B1, 26H7 or 5F4 mAbs (either 1:600 dilution of ascites or varying concentrations of purified antibody). After 5 hours, 100 µl of supernatant were removed for analysis in a γ-counter (LKB Wallac Clini Gamma 1272, Finland). Spontaneous and maximal release were measured by incubating target cells with medium or 1% Nonidet-P40, respectively. Percent cytotoxicity was calculated using the formula (experimental release - spontaneous release) x 100/ (maximal release - spontaneous release).

Statistics: Differences between samples were evaluated with a non-paired, Student's T-test using the Sigma Stat (Jandel Scientific, San Rafael, CA) program.

Example 1: Constitutive expression of the 34B1-related antigen on the cell surface of normal

human IECs.

During the development of iIEL specific mAbs, which were obtained by immunizing mice with an iIEL T-cell line from human small intestine propagated *in vitro*, it was noted that a certain fraction of the mAbs stained IECs as shown by immunohistochemistry of normal human small intestine. Three of these mAbs and the characterization of the antigen that they recognized were of particular interest, as described below.

Fig. 1 depicts identification of three mAbs (34B1, 5F4 and 26H7) which recognize IECs but not resting iIELs. Panels A-C show immunohistology of normal human large intestine stained with the 34B1 (panel A), 5F4 (panel B) and 26H7 (panel C) mAbs with binding detected by subsequent incubation with a goat anti-mouse horseradish peroxidase conjugated antibody as described in the Materials and Methods. The precipitated brown reaction product indicates specific staining on the enterocyte. [Magnification: 20X]. Staining with normal mouse serum was negative (data not shown).

Staining of human intestinal tissue sections showed that these three mAbs (34B1, 26H7 and 5F4) only stained IECs (Fig. 1). This *in vivo* tissue staining with these antibodies appeared to be on the cell surface as confirmed by flow cytometry analysis of a normal human IEC line, HT29. Since these three antibodies did not stain iIELs *in situ*, as determined by immunohistochemistry (Fig. 1), or immediately after isolation as determined by flow cytometry (data not shown), it was suspected that iIELs, activated during the process of *in vitro* cultivation, expressed neoantigens that were constitutively expressed by IECs.

Example 2: The 34B1-related antigen is an activation antigen on normal human iIELs and peripheral blood T Cells.

To determine whether iIEL cell lines expressed neoantigens that were shared with IECs, the staining of iIELs was examined after *in vitro* cultivation with PHA-P. One color flow cytometry analysis was performed of an activated iIEL cell line derived from the small intestine, EEI-5, as described in Fig. 1. As shown in Fig. 1A, iIELs *in situ* and freshly isolated iIELs (data not shown) did not stain with the 34B1, 26H7 and 5F4 mAbs. However, after maintenance *in vitro* as continuous cell lines with PHA-P activation every 10-14 days, all of the iIELs expressed the antigen recognized by these three mAbs (Fig. 2A). Staining of an iIEL T-cell line established from the small intestine, EEI-5, which was 90% CD8⁺ and 10% CD4⁺ as shown in Fig. 1B indicates that all iIEL expressed the antigen recognized by the three mAbs after *in vitro*

activation. The three mAbs exhibited slightly different staining patterns suggesting that they either recognized a different molecule or a different epitope on the same molecule. Similar observations were made with an iIEL T cell line prepared from the large intestine, CLI, which was 40% CD8⁺, 30% CD4⁺ and 30% double negative (CD4⁻CD8⁻) consistent with the *in vivo* phenotype of iIELs in this tissue site (Lundqvist et al., *Int. Immunol.* 7:1473-1480, 1996; data not shown).

These characteristics were, however, not confined to iIELs since PBTs expressed the 34B1-related antigen and upregulated this expression after stimulation with PHA-P *in vitro*. Fig. 2B shows the two-color fluorescence analysis of normal PBTs, before and after three days of stimulation with 1 µg/ml PHA-P, with the 34B1, 5F4 and 26H7 mAbs after gating on the CD3-positive lymphocytes and after subtracting background staining with directly conjugated, isotype matched control antibody. The thick black line shows staining of PBTs without PHA-P stimulation and the thin black line (arrow) with PHA-P stimulation. Prior to PHA-P stimulation, a discrete population of T cells exhibited increased staining with the 34B1, 5F4 and 26H7 mAbs (Fig. 2B). After PHA-P stimulation, a small but significant shift in the intensity of staining of the entire population with all three mAbs was observed. Thus, the 34B1-related antigen is an activation antigen on both normal human T cells in the intestinal epithelium and peripheral blood.

Example 3: The 34B1-related antigen is expressed on epithelial cells, B cells and granulocytes in a wide variety of organs.

The data above suggested that the 34B1-related antigen was expressed by epithelial cells of the intestine and activated T cells. A limited organ survey of the distribution of this cellular expression was therefore examined. As can be seen in Table 1, the antigen(s) recognized by the 34B1, 26H7 and 5F4 mAbs were similarly expressed within a wide variety of tissues suggesting that the molecule(s) of interest had a functional role in diverse organ systems. All three mAbs consistently recognized an antigen on epithelial cells of the small and large intestine, biliary tree, kidney, skin and thymus. In addition, scattered granulocytes in several organs and cells within germinal centers of tonsils, which were consistent with B cells, also stained positive. The staining of granulocytes was confirmed by immunohistochemical analysis of peripheral blood granulocytes (data not shown). Thus, these results, together with the phenotypic studies described above, suggested that the 34B1-related antigen(s) was primarily expressed by a wide

variety of epithelial cell types, B and T lymphocytes, granulocytes and natural killer (NK) cells, based upon staining of an NK cell line (data not shown).

Table I: Tissue Staining of the 34B1-Related mAbs

5	<u>Tissue</u>	<u>Staining Pattern</u>
10	<u>Kidney</u>	Proximal tubules (+) Glomeruli (+) Endothelium (+)
15	<u>Liver</u>	Biliary canaliculi (+) Bile ducts (+): luminal surfaces
20	<u>Lymph Node</u>	Sinusoids (+): Granulocytes/Platelets
25	<u>Skin</u>	Epidermis (-) Eccrine/Sweat glands (+)
30	<u>Small Intestine</u>	Enterocyte (+): Villous > Crypt Goblet cells (-) Granulocytes (+)
35	<u>Thymus</u>	Hassal corpuscles (+)
40	<u>Tonsil</u>	Germinal centers (+) Epithelium (+)

Staining of all tissues is shown as described in the Materials and Methods. For each tissue, cellular staining was graded as either absent (-) or present (+) as defined by a pathologist.

Example 4: Identification of the 34B1-related antigen as biliary glycoprotein (BGP).

To identify the molecule recognized by the 34B1-related mAbs, the 34B1 mAb was used to clone the cDNA which coded for the cognate antigen of the 34B1 mAb by COS cell expression cloning after transfection with a mixture of three cDNA libraries from resting and activated human PBTs and NK cells. These cDNA libraries were utilized because previous studies showed expression of the antigen recognized by the 34B1, 26H7 and 5F4 mAbs in these cell types. Transiently transfected COS cells were subjected to three rounds of immunoselection and panning with the 34B1 mAb. After the third round of panning, 17 of 50 random-selected *E. coli* transformants contained plasmids with a 3.3kb-insert. The inserts in these plasmids were

similar by restriction digest analysis. COS cells transfected with these plasmids were specifically stained with the 34B1 mAb. One of these plasmids, pPAN3.1, was selected for further characterization.

This plasmid directed the translation, when transfected into COS cells, of a 120-kD glycoprotein which was specifically recognized by the 34B1 and 5F4 mAbs and that resolved as a major band of approximately 70-kD and several minor bands of lower molecular weight after digestion with N-glycanase (Fig. 4). Cell surface proteins of COS cells transiently transfected with the pPAN3.1 vector encoding BGPb (lanes a-c) or the pCDM8 vector (lanes d and e) and the activated iIEL cell line, EEI-10 (lanes f and g) were radiolabeled with [¹²⁵I] and immunoprecipitated with either the 34B1 mAb (lanes b, c, d, e, f and g) or normal mouse serum (lane a) and the immunoprecipitates resolved under reducing conditions with (lanes c, e and g) or without (lanes a, b, d and f) prior N-glycanase treatment. Identical observations were made with the 5F4 and 26H7 mAbs (data not shown). A similar glycoprotein was immunoprecipitated from radiolabeled cells surface iIEL proteins by all three mAbs (Fig. 4). Complete DNA sequencing of this cDNA on both strands revealed a sequence that was 97% identical to the 'b' splice variant of BGP or CD66a (Gen Bank accession #X14831) with all the differences occurring outside the coding region. Since the cDNA predicted a polypeptide backbone of 58kD, the data in Fig. 4 suggest that several of the carbohydrate modifications were relatively resistant to N-glycanase digestion.

BGPs are members of the immunoglobulin supergene family that consists of an N-terminal immunoglobulin V- (IgV) related domain, that is highly homologous to the N-domains of other carcinoembryonic antigen (CEA) or CD66 family members, followed by several IgC2-related domains A1 and B1, and the A2, Y or Z domains which are unique to BGP isoforms (Watt et al., 1994; Oikawa et al., 1992; Teixeira, 1994; Barnett et al., 1993; Thompson, 1991).

In order to confirm that the 34B1-related mAbs were reactive with BGP and to define the specific protein domain to which these mAbs were directed, the antibodies were tested in a binding assay with Fc-fusion proteins containing either the N-domain of CD66a, NA1B1 domains of CD66a, the NA1B1A2 domains of CD66a, and N-CAM (CD56) as a negative control. Fig. 3A is a schematic diagram of the Fc-fusion proteins used in the ELISA to test the mAbs as described in the Materials and Methods. Fc-fusion proteins containing the N, NA1B1 and NA1B1A2 domains of CD66a or N-CAM (CD56) as a negative control were tested in an ELISA as described in the Materials and Methods with the 34B1, 5F4 and 26H7 mAbs in

comparison to the positive control antibodies, MA22, MA76 and MA26 (Fig. 3B). As can be seen in Fig. 3, these studies confirmed the recognition of BGP (CD66a) by the three mAbs and showed that all three mAbs reacted with the N-domain.

In order to further confirm that the cognate antigen of the 34B1-related mAbs was BGP, the three mAbs were tested for their ability to stain CHO cells stably transfected with several splice variants of BGP (BGP_a, BGP_c and BGP_{x'}) and HeLa cells transfected with other members of the CD66 serologic cluster including CD66b (CEA gene related member 1, CGM1), CD66c (CEA gene related member 6, CGM6), CD66d (Nonspecific cross reacting antigen, NCA) and CD66e (CEA). Fig. 5 shows the flow cytometric analysis of BGP_a, BGP_c and BGP_{x'} transfectants of CHO, and CEA, NCA, CCGM6 and CGM1 transfectants of HeLa cells in comparison to the mock (Neo) transfectants after staining with either the 34B1, 5F4 and 26H7 mAbs or isotype matched IgG1 antibody as a negative control. All transfectants were positively stained with control mAbs specific for the transfected cDNA (data not shown). Except for CGM1 (CD66b), the 34B1 mAb stained all the CD66 family members tested including all of the CD66a splice variants. The 26H7 and 5F4 mAbs, however, only stained the CD66a splice variants suggesting they were likely specific for the N-domain of this molecule. These results, together with those from phenotyping and the distribution of BGP on epithelial cells, granulocytes, B cells, T cells and NK cells described above and previously reported (Thompson et al., 1991; Möller et al., 1996), clearly identify the N-domain of BGP as the cognate antigen of the 34B1, 26H7 and 5F4 mAbs and that, in addition, the 34B1 mAb recognizes other CD66 forms.

Example 5: iIELs express only CD66a isoform.

Activated human iIELs were stained with a panel of mAbs that have been presented at the Sixth International Workshop on Monoclonal Antibodies and which are specific for all of the CD66 isoforms. The human iIEL cell line from small intestine, EEI-10, was stained 8 days after activation with a series of anti-CD66 mAbs as described in the Materials and Methods. Each panel shows an overlay of the CD66 specific mAbs with the staining obtained with normal mouse serum as a negative control. The specificity of the mAbs for the CD66 isoforms is indicated in the panels. Activated iIELs do not express any CD66 isoforms other than CD66a based upon staining with a large panel of mAbs specific for CD66a-e (mAbs MA27, MA28, MA30, MA41, MA61, MA76, 79, MA81, MA83, MA84, MA86 and MA91 (Fig. 6).

To draw a further similarity between BGP and ITIM containing KIRs, the effects of the CD66a specific mAbs on the pattern of tyrosine phosphorylated proteins after ligation of CD3 was assessed with the Eei-10 cell line. iIELs (1×10^6) were incubated in 100 μ l RPMI-1640 containing 0.1% bovine serum albumin (Sigma) with 100 ng/ml OKT3 mAb and either 10 μ g/ml of the 5F4 mAb or 10 μ g/ml normal IgG1 in the presence of 20 μ g/ml goat anti-mouse IgG antibody (Pierce, Rockford, IL) as crosslinker. After either 2 or 8 minutes incubation at 37°C, the reaction was stopped with 1 ml PBS containing 10mM Na_2VO_3 (Sigma), the pellet incubated on ice for 30 minutes in 100 μ l lysis buffer containing 0.2X PBS, 100 μ M Na_2VO_3 , 1 mM PMSF, 5 mM iodoacetamide and 20 μ g/ml aprotinin and Laemmli buffer added with reducing agents. The lysates were boiled for 5 minutes, resolved by SDS-PAGE on a 10% gel, Western transferred to Immobilon filters, immunoblotted with the PY20 mAb (Zymed, San Francisco, CA) and developed with a horseradish peroxidase conjugated goat anti-mouse antibody (Zymed) and enhanced chemiluminescence. Cross-linking of CD3 with the OKT3 mAb in the presence of the 5F4 mAb, in comparison to an isotype controlled antibody, resulted in a significant alteration in the kinetics and level of phosphorylation suggesting that a phosphatase was activated as a consequence of BGP cross-linking.

Example 6: BGP functions as a killer inhibitory receptor for iIELs.

The observation that BGP was expressed on activated iIELs as defined by staining with the BGP-specific mAbs, 34B1, 26H7 and 5F4 was unique and unexpected since BGP has previously been primarily viewed as a molecule expressed on epithelial cells and granulocytes and involved in cell-cell adhesion and regulation of epithelial cell growth. The function of BGP on iIELs and T cells in general was, however, unknown. Importantly, the cytoplasmic tail of the BGP_a and BGP_b splice variants, but not CD66b-e, contain two ITIM domains within the cytoplasmic tail separated by 21 amino acids raising the possibility that BGP might function as an inhibitory molecule on T cells (Beauchemin et al., 1996; Öbrink, 1997). In order to define the role of the BGP antigen in T cell function, the effects of the three mAbs on the function of iIELs was examined.

A major function of activated iIELs is as cytolytic effector cells. Therefore, the effects of the three mAbs on the cytolytic function of iIELs was examined in a redirected lysis assay, the results of which are depicted in Fig. 7. The anti-CD3 directed lysis (OKT3) of the KJ-3 iIEL cell line was examined as described in the Materials and Methods in the absence or presence of either

the CD66a specific mAbs, 5F4 or 26H7, or an isotype matched IgG1 antibody at a concentration of 100.0 $\mu\text{g/ml}$ at effector:target ratios of 100:1, 50:1 and 25:1. The cytolysis in the absence of added antibodies (medium) and presence of the test antibodies alone (26H7, 5F4 and control IgG1) are also shown. The standard error of the mean for each measurement is indicated. The data presented in Fig. 7 is representative of six experiments. When an iIEL cell line, EEI-10, derived from normal human small intestine, was examined in a redirected lysis assay with the P815 mouse mastocytoma cell line as a target cell, significant cytotoxicity was elicited with, but not without, anti-CD3 cross-linking using the OKT3 mAb (Fig. 7). Neither the 34B1, 26H7 nor 5F4 mAbs at a variety of different concentrations were able to stimulate cytolysis of the P815 cell line indicating that cross-linking BGP did not directly activate iIELs. However, cross-linking BGP with all three anti-BGP mAbs resulted in a significant inhibition of the anti-CD3 directed cytolysis of the P815 cell line in comparison to an isotype matched IgG1 control antibody which exhibited no inhibition. The 5F4 mAb inhibited the lysis by 22%, 35% and 38% at effector:target ratios of 100:1, 50:1 and 25:1, respectively. The inhibition by the control antibody at similar effector:target ratios was -4%, 5% and 5%. Moreover, the anti-CD3 directed cytolysis was not inhibited by a CD2-specific mAb, TS 2/18, which would be expected to inhibit CD58-like interactions with the P815 cell line, suggesting that the inhibition by the anti-CD66a mAbs was not likely due simply to an effect on adhesion. Thus, cross-linking of BGP inhibited the anti-CD3 directed cytolytic activity of iIELs.

When iIELs were harvested early after cytokine treatment, a significant amount of cytolysis was observed against the P815 cell line consistent with cytokine induced killer activity (lymphokine activated killer activity, LAK), a property previously described for iIEL. To test whether this cytolytic activity was also subject to inhibition by the anti-BGP mAbs, the iIEL cell line was exposed to the P815 cell line as target in the presence or absence of the BGP specific mAbs, a pool of the BGP specific mAbs or normal IgG1. The effects of the anti-BGP specific mAbs 34B1, 26H7 and 5F4 on cytolysis of the p815 cell line by the EEI-10 iIEL cell line was examined at an effector target ratio of 25:1. The cytolysis of the p815 cell line was not affected by the irrelevant IgG1 antibody at either 2 $\mu\text{g/ml}$ or 6 $\mu\text{g/ml}$. The inhibition of the cytotoxicity was significant for all anti-BGP antibodies (5F4, $p = 0.15$; 26H7, $p = 0.048$, 34B1, $p = 0.004$; pool, $p = 0.0016$). The anti-BGP mAbs inhibited the cytolytic activity of the iIEL cell line by as much as 70% using the pool of mAbs at 2 $\mu\text{g/ml}$ each, and up to 50% inhibition at doses of individual mAbs of 2 $\mu\text{g/ml}$, indicating the lymphokine activated killer activity of iIELs was also

subject to inhibition by the anti-BGP antibodies.

Example 7: Anti-BGP antibodies can inhibit the BGP inhibitory signal in an allogenic mixed lymphocyte reaction.

5 Human peripheral blood mononuclear cells (donor A) were irradiated (5000 Rads) and cocultivated with an equivalent number of nonirradiated peripheral blood mononuclear cells prepared by Ficoll-Hypaque gradient centrifugation at a total concentration of 2×10^5 cells per well of a 96-well U-bottom plate in a total volume of 200 μ l in quadruplicate. After 96 hours, 0.5 μ Ci of [3 H]-thymidine was added per well for 18 hours of incubation and the plates harvested
10 and counted. The culture conditions contained no additives (medium) or various concentration of diluted ascites (mouse anti-human BGP monoclonal antibodies 34B1 and 5F4) or normal mouse serum. The experimental data depicted in Table II are representative of 5 experiments. The mean \pm SE is presented. The augmentation of T cell proliferation in the presence of anti-BGP antibodies is consistent with the inhibition of a BGP inhibitory signal.

15 *Table II: Allogeneic Mixed Lymphocyte Reaction*

	<u>Treatment</u>	<u>CPM</u>
	Medium	16,330 \pm 2,131
	mouse anti-human BGP (34B1)	
20	1:125	41,246 \pm 1,779
	1:250	54,578 \pm 2,437
	1:500	34,987 \pm 1,545
	1:750	33,978 \pm 1,036
	1:1000	31,522 \pm 1,087
25	mouse anti-human BGP (5F4)	
	1:125	35,330 \pm 1,615
	1:250	33,191 \pm 883
	1:500	27,862 \pm 2,635
30	1:750	23,511 \pm 2,365
	1:1000	38,826 \pm 3,103

Normal mouse serum (negative control)

1:125	18,491 ± 544
1:500	17,957 ± 772
1:1000	19,343 ± 471

5

Fig. 8 shows the results of similar experiments. Stimulator (irradiated peripheral blood mononuclear cells) and responder (nonirradiated peripheral blood mononuclear cells) from two unrelated human study subjects were cocultivated in at 2×10^5 cells per well each in a 96-well flat bottomed plate in the presence or absence of either phytohemagglutinin-P (PHA; 1 $\mu\text{g}/\text{ml}$), irrelevant IgG1 at various concentrations (1-100 $\mu\text{g}/\text{ml}$) or the anti-CD66a specific monoclonal antibodies 34B1, 26H7 or 5F4 at various concentrations (1-100 $\mu\text{g}/\text{ml}$). After four days, 1 μCurie of ^3H -thymidine was added per well for the last 18 hours of incubation and the plates harvested for assessment of proliferation. The y-axis shows the counts per minute. The S.E.M. is shown for each measurement.

15

The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.

20

All of the references described herein are incorporated by reference.

What is claimed is:

Claims

1. A method for enhancing specifically the cytotoxicity or proliferation of killer T cells in a subject, comprising:

administering to a subject in need of such treatment an agent that selectively reduces
5 cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance the cytotoxicity or proliferation of killer T cells in the subject.

2. The method of claim 1, wherein the agent is an antibody or antibody fragment which binds only a single biliary glycoprotein polypeptide.

10

3. The method of claim 2, wherein the antibody fragment is a Fab fragment.

4. The method of claim 1, wherein the agent comprises a ligand for the biliary glycoprotein polypeptide, wherein the ligand binds only a single biliary glycoprotein polypeptide.

15

5. The method of claim 4, wherein the ligand is fused to an immunoglobulin molecule or a fragment thereof.

6. The method of claim 4, wherein the ligand is a soluble biliary glycoprotein molecule or
20 fragment thereof.

7. The method of claim 1, wherein the killer T cells are selected from the group consisting of CD4⁺ T cells, CD8⁺ T cells and NK cells.

25 8. The method of claim 1, wherein the killer T cells are intestinal intraepithelial lymphocytes.

9. The method of claim 1, wherein the killer T cells are peripheral blood T cells.

30 10. A method for suppressing specifically the cytotoxicity or proliferation of killer T cells in a subject, comprising:

administering to a subject in need of such treatment an agent that selectively increases

cross-linking of biliary glycoprotein polypeptides in an amount effective to suppress the activity of killer T cells in the subject.

11. The method of claim 10, wherein the agent is an antibody.

5

12. The method of claim 11, wherein the antibody is a monoclonal antibody.

13. The method of claim 10, wherein the agent comprises a ligand for the biliary glycoprotein polypeptide, wherein the ligand binds two or more biliary glycoprotein polypeptides.

10

14. The method of claim 13, wherein the ligand is fused to an immunoglobulin molecule or a fragment thereof.

15. The method of claim 13, wherein the ligand comprises a biliary glycoprotein polypeptide or fragment thereof.

15

16. The method of claim 10, wherein the killer T cells are selected from the group consisting of CD4⁺ T cells, CD8⁺ T cells and NK cells.

20 17. The method of claim 10, wherein the killer T cells are intestinal intraepithelial lymphocytes.

18. The method of claim 10, wherein the killer T cells are peripheral blood T cells.

25 19. A composition comprising:

an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance cytotoxicity or proliferation of killer T cells in a subject, and

a pharmaceutically-acceptable carrier.

30

20. The composition of claim 19, wherein the agent is an antibody or antibody fragment which binds only a single biliary glycoprotein molecule.

21. The composition of claim 20, wherein the antibody fragment is a Fab fragment.
22. The composition of claim 19, wherein the agent comprises a ligand for the biliary glycoprotein polypeptide, wherein the ligand binds only a single biliary glycoprotein polypeptide..
5
23. The composition of claim 22, wherein the ligand is fused to an immunoglobulin molecule or a fragment thereof.
- 10 24. The composition of claim 22, wherein the ligand is biliary glycoprotein or a fragment thereof.
25. A composition comprising:
an agent that selectively increases cross-linking of biliary glycoprotein polypeptides in an
15 amount effective to suppress cytotoxicity or proliferation of killer T cells in a subject, and
a pharmaceutically-acceptable carrier.
26. The composition of claim 25, wherein the agent is an antibody.
- 20 27. The composition of claim 26, wherein the antibody is a monoclonal antibody.
28. The composition of claim 25, wherein the agent comprises a ligand for the biliary glycoprotein polypeptide, wherein the ligand binds two or more biliary glycoprotein polypeptides.
25
29. The composition of claim 28, wherein the ligand is fused to an immunoglobulin molecule or a fragment thereof.
30. The composition of claim 28, wherein the ligand is biliary glycoprotein or a fragment
30 thereof.
31. A method for enhancing specifically cytotoxicity or proliferation of killer T cells,

comprising:

contacting a population of killer T cells with an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance the cytotoxicity or proliferation of the killer T cells.

5

32. The method of claim 31, wherein the agent is an antibody or antibody fragment that binds one biliary glycoprotein molecule.

33. The method of claim 32, wherein the antibody fragment is a Fab fragment.

10

34. The method of claim 31, wherein the agent comprises a ligand for the biliary glycoprotein polypeptide which binds only a single biliary glycoprotein polypeptide.

35. The method of claim 34, wherein the ligand is fused to an immunoglobulin molecule or a
15 fragment thereof.

36. The method of claim 34, wherein the ligand is a soluble biliary glycoprotein molecule or a fragment thereof.

20 37. The method of claim 31, wherein the killer T cells are selected from the group consisting of CD4⁺ T cells, CD8⁺ T cells and NK cells.

38. The method of claim 31, wherein the killer T cells are intestinal intraepithelial lymphocytes.

25

39. The method of claim 31, wherein the killer T cells are peripheral blood T cells.

40. A method for suppressing specifically cytotoxicity or proliferation of killer T cells, comprising:

30

contacting a population of killer T cells with an agent that selectively increases cross-linking of biliary glycoprotein polypeptides in an amount effective to suppress the cytotoxicity or proliferation of the killer T cells.

41. The method of claim 40, wherein the agent is an antibody.
42. The method of claim 41, wherein the antibody is a monoclonal antibody.
- 5 43. The method of claim 40, wherein the agent comprises a ligand for the biliary glycoprotein polypeptide, wherein the ligand binds two or more biliary glycoprotein polypeptides.
44. The method of claim 43, wherein the ligand is fused to an immunoglobulin molecule or a fragment thereof.
- 10 45. The method of claim 43, wherein the ligand comprises a soluble biliary glycoprotein molecule or a fragment thereof.
46. The method of claim 40, wherein the killer T cells are selected from the group consisting
15 of CD4⁺ T cells, CD8⁺ T cells and NK cells.
47. The method of claim 40, wherein the killer T cells are intestinal intraepithelial lymphocytes.
- 20 48. The method of claim 40, wherein the killer T cells are peripheral blood T cells.
49. An isolated fusion protein comprising a biliary glycoprotein polypeptide or a fragment thereof fused to an immunoglobulin molecule or a fragment thereof.
- 25 50. The isolated fusion protein of claim 49, wherein the biliary glycoprotein or fragment thereof selectively binds a monoclonal antibody selected from the group consisting of 34B1, 5F4 and 26H7.
51. The isolated fusion protein of claim 50, wherein the fragment of biliary glycoprotein is
30 selected from the group consisting of the N-domain of CD66a, NA1B1 domains of CD66a, the NA1B1A2 domains of CD66a.

52. The isolated fusion protein of claim G1, wherein the fragment of the immunoglobulin molecule is the Fc portion of the immunoglobulin molecule.

53 An isolated fusion protein comprising two or more biliary glycoprotein polypeptides or
5 fragments thereof which bind biliary glycoprotein.

54. A method for identifying compounds which enhance or suppress killer T cell activity, comprising,

(a) contacting a population of killer T cells which express biliary glycoprotein with a
10 compound that binds biliary glycoprotein, and

(b) determining the cytotoxicity or proliferation of the population of killer T cells relative to a control, wherein compounds which increase the cytotoxicity or proliferation are compounds which enhance the killer T cell activity, and wherein compounds which decrease the cytotoxicity or proliferation are compounds which suppress the killer T cell activity.

15

55. The method of claim 54, further comprising the steps of

(a) providing a biliary glycoprotein polypeptide or a fragment thereof,

(b) contacting the biliary glycoprotein polypeptide or a fragment thereof with a
compound,

20 (c) determining the binding of the compound to the biliary glycoprotein polypeptide or a fragment thereof, wherein the compound is used in step (a) of claim H1.

56. A method for selectively treating a subject having a condition characterized by aberrant killer T cell activity comprising,

25 administering to a subject in need of such treatment a pharmacological agent which is selective for biliary glycoprotein, in an amount effective to normalize the aberrant killer T cell activity.




Fig. 1A



Fig. 1B



Fig. 1C

2/9




Fig. 2A




Fig. 2B





Fig. 3A



MONOCLONAL ANTIBODIES

Fig. 3B




Fig. 4




Fig. 5




Fig. 6





Fig. 7



LEGEND

(#) = ANTIBODY CONCENTRATION, μ /ml

STIMULATORS = IRRADIATED FEEDER CELLS ALONE

RESPONDERS = RESPONDER CELLS ALONE

No Ab = STIMULATOR AND RESPONDER CELLS MIXED WITH NO TREATMENT

CPM = COUNTS PER MINUTE

Fig. 8

SEQUENCE LISTING

<110> BRIGHAM & WOMEN'S HOSPITAL, INC.
 5 <120> T CELL INHIBITORY RECEPTOR COMPOSITIONS AND USES THEREOF
 <130> B0801/7141WO
 <150> US 60/081,895
 10 <151> 1998-04-15
 <160> 4
 <170> FastSEQ for Windows Version 3.0
 15 <210> 1
 <211> 30
 <212> DNA
 <213> Homo sapiens
 20 <400> 1
 caucauac auaagcttat ggggcacctc 30
 <210> 2
 25 <211> 30
 <212> DNA
 <213> Homo sapiens
 <400> 2
 30 gccattttct tggggcabct cgggtatac 30
 <210> 3
 <211> 30
 <212> DNA
 35 <213> Homo sapiens
 <400> 3
 gtatacccg g agctgcccc a gaaaatggc 30
 40 <210> 4
 <211> 30
 <212> DNA
 <213> Homo sapiens
 45 <400> 4
 cuacuacuac uaagactatg aagttggttg 30

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US99/08430

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 45/05, 39/395, 38/16, 38/17, 38/19; C07K 16/00

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/85.1, 130.1, 133.1, 134.1, 144.1, 154.1, 172.1; 514/8, 885

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EDLUND et al. Evidence for calmodulin binding to the cytoplasmic domains of two C-CAM isoforms. FEBS. July 1993, vol. 327, pages 90-94, especially Abstract and Results, paragraph 1.	1, 25, 28-30
Y	US 5,688,690 A (VALIANTE ET AL) 18 November 1997, see entire document, especially col 7, lines 11-20.	1-9
Y	CICCONE et al. Opposing functions of activatory T-cell receptors and inhibitory NK-cell receptors on cytotoxic T cells. Immunology Today. October 1996, Vol. 17, pages 450-453, see entire article.	1-9



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

09 JULY 1999

Date of mailing of the international search report

30 AUG 1999

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer


 MARIANNE DIBRINO

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/08430

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LUENGPAILIN et al. Demonstration of a biliary-associated glycoprotein in human serum. Clinica Chimica Acta. 1996, Vol. 244, pages 237-240, see entire article.	4-6
Y	SOPP et al. Cross-reactivity of monoclonal antibodies to defined human leucocyte differentiation antigens with bovine cells. Veterinary Immunology and Immunopathology. 1997, Vol. 56, pages 11-25.	1-9, 25-27
Y	US 5,288,854 A (DIAMOND ET AL) 22 February 1994, see entire document, especially column 11, lines 7-26.	1, 4-6
Y	WATT et al. CD66 Identifies the biliary glycoprotein (BGP) adhesion molecule: cloning, expression, and adhesion functions of the BGPc Splice Variant. Blood. 01 July 1994, vol. 84, pages 200-210.	25-27, 28-30
Y	TEIXEIRA et al. The N-domain of the biliary glycoprotein (BGP) adhesion molecule mediates homotypic binding: domain interactions and epitope analysis of BGPc. Blood. 01 July 1994, vol. 84, pages 211-219.	25, 28-30
Y	OIKAWA et al. Homotypic and heterotypic Ca ⁺⁺ -independent cell adhesion activities of biliary glycoprotein, a member of carcinoembryonic antigen family, expressed on cho cell surface. biochem. Biophys. Res. Comm. 31 July, 1992, Vol. 186, pages 881-887.	25, 28-30

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/08430

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-9, 31-39, 25-30

4. No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- The additional search fees were accompanied by the applicant's protest.
No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US99/08430

A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/130.1, 133.1, 134.1, 144.1, 154.1, 173.1, 178.1, 193.1; 514/8, 885

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

STN(BIOSIS, MEDLINE, EMBASE); APS; Blumberg, biliary glycoprotein, bgp, cd66, cd66a, enhance cytotoxicity, cea, enhance proliferation, antibody, IEL, intraepithelial lymphocytes, monoclonal, ccam c-cam, t cell, inhibitory receptors NK cell, Natural Killer Cells, T cell, CD4, CD8, fusion protein, cross-link, cross link, prevent cross linking, homotypic adhesion

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1. In order for all inventions to be searched, the appropriate additional search fees must be paid.

Group I, claims 1-9 and 31-39, drawn to a method for *enhancing* cytotoxicity or proliferation of killer T cells.

Group II, claims 10-18 and 40-48, drawn to a method for *suppressing* cytotoxicity or proliferation of killer T cells.

Group III, claims 49-53, drawn to an isolated fusion protein comprising a biliary glycoprotein polypeptide fused to an immunoglobulin molecule.

Group IV, claims 54-55, drawn to a method for identifying compounds which enhance or suppress killer T cell activity.

Group V, claim 56, drawn to a method for selectively treating a subject having a condition characterized by aberrant killer T cell activity.

Group VI, claims 19-24, drawn to a composition comprising an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to enhance cytotoxicity or proliferation of killer T cells in a subject.

Group VII, claims 25-30, drawn to a composition comprising an agent that selectively reduces cross-linking of biliary glycoprotein polypeptides in an amount effective to suppress cytotoxicity or proliferation of killer T cells in a subject.

The inventions listed as Groups I-VII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

1. The products of Groups III, VI and VII (compositions comprising agents) are structurally different.
2. The putative special technical feature linking Groups I-VII appears to be the fusion protein of Group V, claim 49. However, Edlund et al (FEBS vol 327 (1), pp 90-94, especially Abstract and Results, paragraph 1) teach fragments of fusion proteins of biliary glycoproteins C-CAM 1 and C-CAM 2 with unrelated proteins. It would be obvious to use antibody as the unrelated protein. Therefore, the technical feature linking the inventions of Groups I-VII does not constitute a special technical feature as defined by PCT Rule 13.2, as it does not define a contribution over the prior art.

Therefore, there is no special technical feature which provides for unity of invention.