



(86) Date de dépôt PCT/PCT Filing Date: 2006/04/06  
 (87) Date publication PCT/PCT Publication Date: 2006/10/26  
 (45) Date de délivrance/Issue Date: 2011/06/14  
 (85) Entrée phase nationale/National Entry: 2007/09/07  
 (86) N° demande PCT/PCT Application No.: US 2006/012999  
 (87) N° publication PCT/PCT Publication No.: 2006/113166  
 (30) Priorité/Priority: 2005/04/15 (US11/106,798)

(51) Cl.Int./Int.Cl. *A01N 59/00* (2006.01),  
*A01N 25/30* (2006.01), *A01P 1/00* (2006.01),  
*A61K 33/20* (2006.01), *A61L 2/18* (2006.01)  
 (72) Inventeurs/Inventors:  
 MORELLI, JOSEPH, US;  
 LAPORTE, KARLA, US;  
 LI, JUNZHONG, US  
 (73) Propriétaire/Owner:  
 ECOLAB INC., US  
 (74) Agent: CASSAN MACLEAN

(54) Titre : COMPOSITIONS DESINFECTANTES A BASE DE CHLORITE ACIDIFIE PRESENTANT DES STABILISANTS OLEFINIQUES

(54) Title: ACIDIFIED CHLORITE DISINFECTANT COMPOSITIONS WITH OLEFIN STABILIZERS

(57) **Abrégé/Abstract:**

A two-part disinfecting systems, as well as disinfecting compositions and methods for making and using the same. The two-part disinfecting system contains a first part and a second part adapted to be mixed to yield an aqueous disinfecting composition, wherein the first part comprises a chlorite and the second part comprises an acid, and wherein the first part, the second part, or both the first and second parts comprise an olefin compound.

## (12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
26 October 2006 (26.10.2006)

PCT

(10) International Publication Number  
**WO 2006/113166 A1**

## (51) International Patent Classification:

A01N 59/00 (2006.01) A61K 33/20 (2006.01)  
A01N 25/30 (2006.01) A61L 2/18 (2006.01)  
A01P 1/00 (2006.01)

## (21) International Application Number:

PCT/US2006/012999

(22) International Filing Date: 6 April 2006 (06.04.2006)

(25) Filing Language: English

(26) Publication Language: English

## (30) Priority Data:

11/106,798 15 April 2005 (15.04.2005) US

(71) Applicant (for all designated States except US): **ECOLAB INC.** [US/US]; Ecolab Center, 370 N. Wabasha Street, St. Paul, MN 55102 (US).

## (72) Inventors; and

(75) Inventors/Applicants (for US only): **MORELLI, Joseph** [US/US]; 10622 NE 173rd Place, Bothell, WA 98011 (US). **LAPORTE, Karla** [US/US]; 550 Mount Olympus Drive Sw, Redmond, WA 98027 (US). **LI, Junzhong** [CN/US]; 12700 Durham Way, Apple Valley, MN 55124 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: ACIDIFIED CHLORITE DISINFECTANT COMPOSITIONS WITH OLEFIN STABILIZERS

(57) Abstract: A two-part disinfecting systems, as well as disinfecting compositions and methods for making and using the same. The two-part disinfecting system contains a first part and a second part adapted to be mixed to yield an aqueous disinfecting composition, wherein the first part comprises a chlorite and the second part comprises an acid, and wherein the first part, the second part, or both the first and second parts comprise an olefin compound.

WO 2006/113166 A1

**ACIDIFIED CHLORITE DISINFECTANT**  
**COMPOSITIONS WITH OLEFIN STABILIZERS**

**FIELD OF THE INVENTION**

The present invention relates to two-part disinfecting systems, as  
5 well as disinfecting compositions and methods for making and using the  
same, and in a particular embodiment to a two-part disinfecting system that,  
when mixed, yields a disinfecting composition having reduced chlorine  
dioxide generation and extended color longevity.

**BACKGROUND OF THE INVENTION**

10 Many diseases arise from the growth and spread of microorganisms  
that can affect all aspects of life, from human health, to animal health, to  
food and water safety, to the safety of the environments we live in.  
Disinfectants have found wide spread application in all these areas.  
Hospitals perform rigorous programs to disinfect and sterilize their  
15 environments. Consumer homes are replete with disinfectant hand cleaners,  
sprays, hard surface cleaners, disinfectant wipes, and fruit and vegetable  
washes. Disinfectants are widely used on farms where the difference  
between healthy and sick animals can mean the difference between  
profitability and loss.

20 Mastitis is one of the most common and economically costly  
diseases confronting milk producers. Economic losses result from poorer  
milk quality, lower milk production, and potential culling of chronically  
infected animals. The use of disinfectant solutions both before and after  
milking has found great success in preventing mastitis, particularly

disinfectants based on acidified chlorite such as those commercially available from Alcide Corporation (Redmond, WA) and Ecolab Inc. (St. Paul, MN), and disclosed in U.S. Pat. Nos. 6,524,624, and 6,699,510.

Acidified chlorite (AC) disinfectants are commonly used as two-  
5 part products having a first part containing a chlorite (such as sodium chlorite) and a second part containing an acid. The AC disinfectant is formed upon mixing the first and second parts, and typically only in amounts sufficient for a given use period. Depending upon the desired characteristics and/or intended use of the AC disinfectant, either the first or  
10 second part, or both parts, may contain one or more optional ingredients such as skin conditioners, healing agents, surfactants, thickeners, builders, film-forming agents, and/or preservatives. Also, depending upon the two-part system, the AC disinfecting composition may be formed by simply mixing the first and second parts, often in approximately equal volumes, or  
15 may involve some additional dilution step before or after mixing.

Color has proved to be an important attribute for teat disinfectants, allowing farmers to visually confirm that the disinfectant has been properly applied to the teat. This is particularly advantageous for confirming application to large herds when multiple farm workers are applying the  
20 disinfectant to many different animals. Unfortunately, many such colorants used to impart the color are subject to chemical degradation upon formation of the AC disinfectant. Once the chlorite-containing part and acid part are combined, they form chlorous acid. Over time, the chlorous acid undergoes complex chemical transformations to form chlorine dioxide.

Formation of too much chlorine dioxide is undesirable because chlorine dioxide is a gas in its natural state, which is noxious; chlorine dioxide can be corrosive to metal surfaces; and chlorine dioxide degrades colorants. Colorant degradation leads to ineffective coloration and teat marking as the  
5 disinfectant ages--that is, from the point in time following mixing of the first and second parts to form the AC disinfectant.

Teat disinfectants are generally considered animal "drugs" in most countries, and thus controlled by the regulatory agencies overseeing the same. Most often, the only colorants that can be used in a teat disinfectant  
10 are those dyes that have been approved for use in food and/or drugs. For example, in the United States approved dyes can be found in 21 C.F.R. § 70.3. When present in an AC disinfectant, these dyes are susceptible to chemical oxidation and rapidly lose their color following formation of the disinfectant.

15 Previous attempts to address this problem have largely focused on use of pigments as opposed to dyes (see e.g., WO 99/16418, WO 99/16309 and EP 0 904 693 A1). Pigments are insoluble colorants and less susceptible to chemical degradation within the AC disinfectant. However, pigments are plagued by problems associated with settling out of solution,  
20 staining parlor floors, and clogging milk filters. In addition, such pigments are not approved in some countries for use in teat disinfectants since they are not approved for food or drug use by their regulatory agencies.

Accordingly, there remains a need in the art for improved AC disinfectants generally, as well as a need for controlling the formation of



wherein there are at least two saturated carbon atoms between  $(R_1)CH=CH$  and a heteroatom and a saturated carbon is a carbon free of double and triple bonds.

The olefin compound may be in a protonated form (i.e., sulfonic acid), a salt form, or a mixture thereof, and generally contains from 6 to 60 carbon atoms, although it is understood that more or fewer carbons would fit within the invention.

Alpha olefin sulfonates have been previously discussed as compositions that aid in reducing the generation of chlorine dioxide. See U.S. Patent No. 6,524,624 and U.S. Patent Application Publication No. 20030180247.

Now, it has been discovered that a specific olefin chemical structure has been found to be especially effective at reducing chlorine dioxide generation. In particular, it has been discovered that an olefin compound having a hydrocarbon chain with at least one double bond that is at least two saturated carbon atoms away from a hydrophilic group having at least four heteroatoms are especially effective.

These olefin compounds are advantageous for three reasons. First, they provide a formulator with a broader range of materials to select from where there are broader choices for foaming, wetting, and cleaning characteristics. Second, some are effective at reducing chlorine dioxide generation and consequently color degradation in smaller quantities than alpha olefin sulfonates. Third, some alpha olefin sulfonates can be drying

to skin at elevated levels. Some olefin compounds of the present invention are less drying than the alpha olefin sulfonates.

When combined, the first part and second part form a disinfecting composition having utility over a wide range of applications. The olefin  
5 compound has surprisingly been found to reduce the generation of chlorine dioxide, providing a disinfecting composition that is longer lasting, and has less odors. When the optional oxidizable colorant is present, this results in a disinfecting composition having extended color longevity. Such compositions are particularly useful as teat dips, as well as for other  
10 disinfecting applications where reduction in the generation of chlorine dioxide or extended color longevity is desirable.

In a further embodiment, a method for making a disinfecting composition is disclosed by combining the first part and the second part of the two-part disinfecting system. Such combination may involve mixing  
15 liquid forms of the first part and second part, or may involve diluting or dissolving the first part and/or second part prior to mixing, at the time of mixing, and/or after mixing.

In yet another embodiment, a method for disinfecting a substrate is disclosed by contacting the substrate with an effective amount of a  
20 disinfecting composition of this invention. Such substrates include any surface, material, or fluid that would benefit from being disinfected, including the skin or tissue of a warm-blooded animal, in particular the teat of a dairy cow, goat or sheep, as well as hard surfaces generally and food surfaces such as meat and meat parts (including cooked and uncooked beef,

poultry, pork, other generally recognized red meats, and fish), fruits and vegetables, and process waters, such as flume waters, cooling tower waters, equipment, and facility cleaning solutions, etc.

These and other aspects of this invention will be evident upon  
5 reference to the following detailed description of the invention.

### DETAILED DESCRIPTION OF THE INVENTION

#### Definitions

For the following defined terms, these definitions shall be applied,  
unless a different definition is given in the claims or elsewhere in this  
10 specification.

All numeric values are herein assumed to be modified by the term  
“about,” whether or not explicitly indicated. The term “about” generally  
refers to a range of numbers that one of skill in the art would consider  
equivalent to the recited value (i.e., having the same function or result). In  
15 many instances, the term “about” may include numbers that are rounded to  
the nearest significant figure.

Weight percent, percent by weight, % by weight, wt %, and the like  
are synonyms that refer to the concentration of a substance as the weight of  
that substance divided by the weight of the composition and multiplied by  
20 100.

The recitation of numerical ranges by endpoints includes all  
numbers subsumed within that range (e.g. 1 to 5 includes 1, 1.5, 2, 2.75, 3,  
3.80, 4 and 5).

As used in this specification and the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the content clearly dictates otherwise. Thus, for example, reference to a composition containing “a compound” includes a mixture of two or more compounds.

5 As used in this specification and the appended claims, the term “or” is generally employed in its sense including “and/or” unless the content clearly dictates otherwise.

The use of the terms “antimicrobial” and “biocide” in this application does not mean that any resulting products are approved for use  
10 as an antimicrobial agent or biocide.

As noted above, in one embodiment a two-part disinfecting system is disclosed comprising a first part and a second part adapted to be combined to yield an aqueous disinfecting composition. The first part comprises a chlorite and the second part comprises an acid. In addition, the  
15 first part, the second part, or both the first and second parts, further comprise an olefin compound.

Acidified chlorite compositions may be generated by combining chlorite (i.e.,  $\text{ClO}_2^-$ ), typically in the form of a metal salt such as sodium chlorite, with an acid activator. Such compositions are effective  
20 disinfectants due to the generation of antimicrobial oxidants, particularly chlorous acid (i.e.,  $\text{HClO}_2$ ). Chlorous acid is formed very rapidly upon acidification of chlorite in an equilibrium process governed by the solution pH. Chlorous acid can subsequently undergo a series of chemical reactions to form chlorine dioxide. Although not wishing to be limited by the

following theory, it is believed that the olefin compound reduces generation of chlorine dioxide upon formation of the disinfecting composition by affecting the rate by which chlorous acid is converted to chlorine dioxide. When an oxidizable colorant is present, it is degraded in significant part by the chlorine dioxide generated within the disinfectant. Thus, the olefin compound, by controlling chlorine dioxide generation, imparts extended color longevity to the disinfecting composition by limiting oxidation of the colorant. Chlorine dioxide is a particularly pungent gas that can be unpleasant at excessive levels in air. Unlike chlorous acid, which stays in solution at the surface being disinfected, chlorine dioxide can escape into the air around the user. Slowing the rate of chlorine dioxide formation leads to a longer lasting disinfectant composition with less odors to the user.

The first and second parts may both be in the form of an aqueous composition, emulsion, microemulsion, cream or gel, or one or both may be in a concentrated, non-aqueous or solid form. For example, the first and second parts may be aqueous compositions or gels to be mixed in approximately equal volumes to form the disinfecting composition, or may be concentrates or solids to be diluted by or dissolved in water, and then mixed to yield the disinfecting composition. Alternatively, the first and/or second parts may be in a non-aqueous or solid form (such as a powder or tablet) to be mixed with or dissolved in water prior to combination. To avoid excessive generation of chlorine dioxide which may occur upon

combination of concentrated forms, it is preferable to mix the first and second parts after the parts are diluted with or dissolved in water.

The chlorite of the first part is typically an alkali or alkaline earth metal chlorite, such as potassium or sodium chlorite, and more typically sodium chlorite. The chlorite is present in the first part in an amount such  
5 that, when combined with the second part, it is present within the disinfecting composition at a concentration ranging from about 0.005% to about 3% by weight, from about 0.05% to about 0.5% by weight, and from about 0.1% to about 0.4%.

10 The acid of the second part is any compound or mixture of compounds that will acidify the chlorite of the first part. In one embodiment, the acid has a  $pK_a$  ranging from 2 to 5. The acid can be an organic acid, inorganic acid, or mixture thereof. Organic acids include (but are not limited to) formic acid, acetic acid, glycolic acid, lactic acid,  
15 pyruvic acid, malic acid, mandelic acid, citric acid, tartaric acid, adipic acid, succinic acid, malonic acid, propionic acid, heptanoic acid, octanoic acid, nonanoic acid, salicylic acid, benzoic acid, gluconic acid, or mixtures thereof. The organic acid can also be alkyl-, alkylaryl-, and arylsulfonic acids such as octanesulfonic acid, toluenesulfonic acid, cumenesulfonic  
20 acid, dodecylbenzenesulfonic acid, and homo- & copolymers containing poly(styrenesulfonic acid) and poly(acrylamidopropylsulfonic acid). Inorganic acids include (but are not limited to) sulfuric acid, sulfamic acid, phosphoric acid, hydrochloric acid, nitric acid, boric acid, or mixtures thereof. Other acids that may be used include (but are not limited to)

hydrated metals salts of iron, aluminum, zirconium, vanadium, and gadolinium as described in U.S. Pat. No. 5,820,822. Acids also include (but are not limited to) solid acid exchange resins, such as Amberlite™, Diaion™, Dowex™ and Duolite™, as well as aluminum silicate zeolites.

5 Alternatively, the acid may be any organic acid precursor which forms an acid upon contact with water, such as acid anhydrides, esters, and sulfonate esters. Examples of organic acid precursors are described in U.S. Pat. No. 4,585,482.

The acid is present in the second part in an amount such that, when  
10 combined with the first part, it is present within the disinfecting composition at a concentration ranging from about 0.05% to about 10% by weight, from about 0.5% to about 5.0% by weight, and from about 1.0% to about 3.0% by weight.

Alternatively, the amount of acid in the second part may be  
15 characterized by the pH of the disinfecting composition. In this regard, the acid is present in the second part in an amount such that, when combined with the first part, the pH of the disinfecting composition is below about 5, from about 2 to about 5, and from about 2.3 to about 3.5.

The optional oxidizable colorant is a colorant that undergoes color  
20 loss upon contact with chlorine dioxide at concentrations generally encountered in acidified chlorite disinfectants. The colorant may be present in the first or second part and is preferably present in the second part. Such colorants are typically soluble in vehicles which may be used as carriers for the first or second part, including (but not limited to) water, alcohol,

glycerin and/or oil. In the practice of this invention, either a single oxidizable colorant, or a mixture of two or more oxidizable colorants, may be present in the second part. The amount of oxidizable colorant present in the first or second part is an amount that, upon combination with the other  
5 part, will impart the desired color and/or color intensity to the disinfecting composition.

In the United States, colorants for use in foods and/or drugs are generally classified by the Food and Drug Administration (FDA) as either (1) a food, drug and cosmetic (FD&C) colorant, (2) a drug and cosmetic  
10 (D&C) colorant, or (3) an externally applied drug and cosmetic (Ext. D&C) colorant. These colorants may be identified using Colour Index Numbers (CI#) established by the Society of Dyers and Colourists (UK) and the American Association of Textile Chemists & Colorists (Color Index, Society of Dyers and Colorists and American Association of Textile  
15 Chemists & Colorists, Rev. 3.sup.rd ed, Branford, 1975).

Representative FD&C colorants include (but are not limited to) FD&C Blue #1 (CI#42090), FD&C Blue #2 (CI#73015), FD&C Green #3 (CI# 42053), FD&C Red #3, FD&C Red #4 (CI# 14700), FD&C Red #40 (CI# 16035), FD&C Yellow #5 (CI# 19140), FD&C Yellow #6 (CI#  
20 15980), Orange B, and Citrus Red #2.

Representative D&C colorants include (but are not limited to) D&C Violet #2 (CI# 61565), D&C Green #5 (CI# 61570), D&C Green #6 (CI# 61565), D&C Green #8 (CI# 59040), D&C Orange #4 (CI# 15510), D&C Yellow #7, D&C Yellow #8 (CI# 45350), D&C Yellow #10 (CI# 47005),

D&C Yellow #11 (CI# 47000), D&C Red #6 (CI# 15850), D&C Red #17 (CI# 26100), D&C Red #22 (CI# 45380), D&C Red #28 (CI# 45410), and D&C Red #33 (CI# 17200).

Representative Ext. D&C colorants include (but are not limited to)

5 Ext. D&C: Violet #2 (CI# 60730), Ext. Yellow #7 (CI# 10316),

Other representative food colorants include (but are not limited to)

Acid Green 1 (CI# 10020), Food Yellow 2 (CI# 13015), Acid Yellow 36 (CI# 13065), Food Yellow 8 (CI# 14720), Acid Orange 20 (CI# 14600), Food Red 3 (CI# 14720), Food Red 2 (CI# 14815), Acid Red 88 (CI# 15620), Food Orange 2 (CI# 15980), Acid Red 26 (CI# 16150), Food Red 7 (CI# 16155), Food Red 9 (CI# 16135), Acid Orange 10 (CI# 16230), Acid Red 18 (CI# 16255), Acid Red (CI# 16290), Acid Red 1 (CI# 18050), Acid Red 155 (CI# 18130), Acid Yellow 121 (CI# 18690), Acid Red 180 (CI# 18736), Acid Yellow 11 (CI# 18820), Acid Yellow 40 (CI# 18950), Acid Yellow 5 (CI# 18965), Acid Black 1 (CI# 20470), Acid Red 163 (CI# 24790), Acid Red 73 (CI# 27290), Food Black 2 (CI# 27755), Food Black 1 (CI# 28440), Direct Orange 34 (CI# 40215), Acid Blue 3 (CI# 42051), Acid Blue 5 (CI# 42052), Green S (CI# 44090), and Brown HT (CI# 20285).

20 In addition, oxidizable colorants also include naturally occurring colorants such as red cabbage extract, beet root extract, carminic acid, curcumin, beta carotene, annatto extract, grape skin extract, astaxanthin, canthaxanthin, henna, guaiazulene, and mixtures thereof.

Oxidizable colorants of this invention also include any combination of two or more of the above FD&C, D&C, Ext. D&C, food colorants, and naturally occurring colorants. Furthermore, the oxidizable colorant may, upon contact with the other part, undergo a change in color. Such color change maybe attributable, for example, to a change in pH going from the pH of one part to the pH of the resulting composition. Alternatively, either part may optionally contain a colorant such that, when combined with the other part, the resulting composition has a color different from either the first and second parts.

As noted above, it has been surprisingly found that the presence of an olefin compound in either the first part, the second part, or both the first and second parts, reduces the generation of chlorine dioxide in the resulting disinfecting composition. As a result, when the optional oxidizable colorant is present, the olefin compound imparts enhanced color longevity to the disinfection composition. As used herein, an "olefin compound" refers to a hydrocarbon having at least one double bond that is at least two saturated carbon atoms away from a hydrophilic group having at least four heteroatoms selected from the group consisting of oxygen, nitrogen, sulfur, chlorine, bromine, fluorine, iodine, phosphorous and mixtures thereof. The term "hydrophilic" refers to the portion of the molecule that tends to be "water-loving" or more soluble. The term "saturated carbon" refers to a carbon atom that is free of double or triple bonds. The olefin compound of the present invention has the following structure:



where  $R_1 = H-$ ,  $CH_3-$ ,  $CH_3(CH_2)_n-$ ,  $CH_3(CH_2)_nCH(OH)-$ , and  $MO_2C(CH_2)_n-$ ;

$n = 1-20$  and preferably  $1-10$ ;

5  $R_2 = --(CH_2)_m-$ ,  $--CH(CH_2)_mCO_2M$ ;  
|

$m = 2-20$  and preferably  $2-10$ ;

$M = H, Na, K, Ca, Mg$ , ammonia, or an organic cation such as monoethanolamine, or triethanolamine; and


10  $X =$  a head group having at least four heteroatoms selected from the group consisting of oxygen, nitrogen, sulfur, chlorine, bromine, fluorine, iodine, and phosphorous and mixtures thereof,

wherein there are at least two saturated carbon atoms between  $(R_1)CH=CH$  and a heteroatom and a saturated carbon is a carbon free of  
15 double and triple bonds.

Because of the presence of the hydrophobic group  $(R_1)CH=CH(R_2)$  and the hydrophilic group  $(X)$ , the olefin compounds of the present invention exhibit surfactant properties, and consequently can be characterized in terms of their HLB (Hydrophile – Lipophile Balance)  
20 value. The HLB is an expression of the relationship between the hydrophobic group and the hydrophilic group of the molecule. Generally, the higher the HLB value, the more water-soluble the molecule. Hydrophilicity increases with increasing number of heteroatoms. HLB values greater than 10 are considered water-soluble. For the present  
25 invention, the olefin compound preferably has a HLB value equal to or greater than 8, and more preferably equal to or greater than 12.

Some non-limiting examples of suitable olefin compounds include those compositions that are oleyl-based, ricinoleyl-based, and undecylenyl-based. Oleyl based compounds are compounds derived from oleic acid. This includes addition products in the form of esters, amides, oxazolines, and ene synthesis (hydro-allyl-addition) adducts. It also includes derivatives formed by converting oleic acid into oleyl alcohol or oleyl amine and further chemically modifying the alcohol and amine groups. Ricinoleyl- and undecylenyl-based compounds refer to the analogous array of compounds derived from the corresponding ricinoleic acid and undecylenic acid, respectively.

Some non-limiting examples of oleyl-based olefin compounds include those having the following structure:





Examples of substructures that may be used as the -X group are provided in Table 1.

**Table 1** Oleyl-Based -X Groups

Substructure -X	INCI Name	Example Trade Names
(CO)NCH <sub>3</sub> CH <sub>2</sub> CH <sub>2</sub> SO <sub>3</sub> Na	Sodium Methyl Oleoyl Taurate	Geroon™ T-77 (Rhodia) Hostapon™ T (Hoechst Celanese)
(CO)N(CH <sub>2</sub> CH <sub>2</sub> OH) <sub>2</sub>	Oleamide DEA	Incromide™ OD (Croda) Mackamide™ NOA (McIntyre)
(CO)NH(CH <sub>2</sub> ) <sub>3</sub> N(CH <sub>3</sub> ) <sub>2</sub> →O	Oleamidopropylamine oxide	Incromine™ Oxide O (Croda) Mackamine™ OAO (McIntyre)
(CO)NH(CH <sub>2</sub> ) <sub>3</sub> N(CH <sub>3</sub> ) <sub>2</sub> <sup>+</sup> CH <sub>2</sub> COO <sup>-</sup>	Oleamidopropyl betaine	Mackam™ HV (McIntyre) Incronam™ OP30 (Croda)

Substructure -X	INCI Name	Example Trade Names
$(CO)NH(CH_2)_3N(CH_3)_2CH_2CH(OH)CH_2R$ R = hydrolyzed collagen moiety	Oleamidopropyl dimonium hydroxypropyl hydrolyzed collagen	Mackpro™ OLP (McIntyre)
$(CO)NH(CH_2)_3N(CH_3)_2^+CH_2CH(OH)CH_2SO_3^-$	Oleamidopropyl hydroxysultaine	
$(CO)NH(CH_2)_3N(CH_3)_2^+CHCH(OH)CH_2OH \cdot Cl^-$	Oleamidopropyl PG-dimonium chloride	Lexquat™ AMG-O (Inolex)
$(CO)NCH_3CH_2COOH$	Oleoyl sarcosine	Crodasinic™ O (Croda) Hamposyl™ O (Hampshire)
$(CO)NH(CH_2)_2N(CH_2CH_2OH)CH_2CH(OH)CH_2SO_3^-Na^+$	Sodium oleoamphohydroxypropyl sulfonate	Miranol™ OS-D (Rhodia)
$(CO)(OCH_2CH_2)_nO(CO)-$ n is average of polydisperse compositions	PEG-# Dioleate Where # is 4 thru 150 and correspond to n	Kessco™ PEG 400 Dioleate™ (Stepan) Kessco PEG 4000 Dioleate (Stepan) Kessco PEG 6000 Dioleate (Stepan)
$(CO)NH(CH_2CH_2O)_nH$ n is average of polydisperse compositions	PEG-7 Oleamide	Ethomid™ O/17 (Akzo)
$N(CH_2CH_2O)_xH$   $(CH_2CH_2O)_yH$	PEG-# Oleamine Where # is 2 thru 30 and corresponds to x+y	Ethomeen™ O/12 (Akzo) Ethomeen O/15
$(CO)(OCH_2CH_2)_nOH$ n is average of polydisperse compositions	PEG-# Oleate Where # is 2 through 150 and correspond to n	Kessco Diglycol Oleate (Stepan) Ethofat™ O/15 (Akzo) Kessco PEG 600 Monooleate (Stepan)
$NCH_3(CH_2CH_2O)_xH^+ \cdot Cl^-$   $(CH_2CH_2O)_yH$	PEG-# Oleammonium chloride Where # is 3 through 15 and correspond to x+y	Ethoquad™ O/12H (Akzo) Ethoquad O/25
$(CO)NHCH_2CH_2N(CH_2CH_2OH)(CH_2COONa)$	Sodium Oleoamphoacetate	Miranol OM (Rhodia)
$(CO)NHCH_2CH_2N(CH_2CH_2OH)(CH_2CH_2COONa)$	Sodium Oleoamphopropionate	Miranol OM-SF (Rhodia)
$(CO)OCH_2CH_2SO_3Na$	Sodium Oleoyl isethionate	
$OSO_3Na$	Sodium Oleyl Sulfate	
$(CO)NHCH_2CH_2OCOCH(SO_3Na)CH_2COONa$	Disodium oleamido MEA sulfosuccinate	Mackanate™ OM (McIntyre) Rewopol™ SB E 280 (Rewo)
$NHCH_2CH(CH_3)OCOCH(SO_3Na)CH_2COONa$	Disodium oleamido MIPA sulfosuccinate	Mackanate OP (McIntyre) Emcol™ 4161L (Witco)
$NH(CH_2CH_2O)_nCOCH(SO_3Na)CH_2COONa$	Disodium Oleamido PEG-2 Sulfosuccinate (also diammonium oleamido PEG-2 sulfosuccinate)	Mackanate OD-35 (McIntyre) Geropon SBG-280 (Rhodia)
$(CO)OCH_2CH(OH)CH_2OH$	Glyceryl Oleate	Aldol™ MO (Lonza) Witconol™ 2421 (Witco)

Substructure -X	INCI Name	Example Trade Names
	Sorbitan Oleate	Crill™ 4 (Croda) Witconol 2500 (Witco)
PEG derivatized 	PEG-# sorbitan oleate PEG-# sorbitan tetraoleate PEG-# sorbitan hexaoleate Where # varies from 2 to 50 and corresponds to the level of polyethylene glycol derivatization	Nikkol™ TO 106 (Nikko) Rheodol™ 460 (Kao) Atlas™ G-1096 (ICI)
$N^+(CH_2CH_2OH)_2CH_2CO_2^-$	Dihydroxyethyl oleyl glycinate	
$OCH_2CH(OH)CH_2(OCH_2CH_2)_nOH$	PEG-# Glyceryl Oleate Where # ranges from 10-30 and corresponds to the average n	Nikkol™ TMGO-15 (Nikko) Tagat™ O (Goldschmidt)
Ester of polyglyceryl-X and oleic acid	Polyglyceryl-# oleate Where # is the average repeating units of a glycerin polymer	Abil™ WS 08 (Goldschmidt) Emulsunt™ 1055 (ISP) Nikkol Decaglyn 1-O (Nikko)
$-(OCH_2CH_2)_nOH$ n is average of polydisperse compositions	Oleth-# where # is 3,4,5,6,7,8,9,10,12,15,16, 20, 23, 25, 30, 40, 44, 50 and corresponds to n	Volpo™ 10 (Croda) Marlowet™ OA-5 (Huls) Volpo 20 (Croda)
$-(OCH_2CH_2)_nOCH_2COOH$ n is average of polydisperse compositions	Oleth-# Carboxylic Acid Where # is 3, 6, 10 and corresponds to n	Akypo™ RO-20 (Chem Y) Akypo RO-50 (Chem Y)
$-(OCH_2CH_2)_nOPO_2OH$ n is average of polydisperse compositions	Oleth-# Phosphate Where # is 3, 4, 5, 10, 20 and corresponds to n	Crodafos™ N-3 (Croda) Crodafos N-10 (Croda)
$-OCOCH(SO_3Na)CH_2COONa$	Disodium oleyl sulfosuccinate	
$-(OCH_2CH_2)_nO-COCH(SO_3Na)CH_2CO_2Na$	Disodium Oleth-3 Sulfosuccinate	Incrossul™ OTS (Croda)
$-(OCH_2CH_2)_nOSO_3^- \frac{1}{2}Mg^{2+}$	Magnesium Oleth sulfate n = 1-4 on average	Mixture in Texapon™ ASV (Henkel)

Some non-limiting examples of ricinoleyl-based olefin compounds

include those having the following structure:



- 5 Examples of substructures that may be used as the -X group are provided in Table 2.

**Table 2 Ricinoleyl-Based -X Groups**

Substructure -X	INCI Name	Example Trade Names
(CO)N(CH <sub>2</sub> CH <sub>2</sub> OH) <sub>2</sub>	Ricinoleamide DEA	Mackamide R (McIntyre)
(CO)NH(CH <sub>2</sub> ) <sub>3</sub> N(CH <sub>3</sub> ) <sub>2</sub> <sup>+</sup> CH <sub>2</sub> COO <sup>-</sup>	Oleamidopropyl betaine	Rewoteric <sup>TM</sup> AM R 40 (Rewo Chemische) Mackam RA (McIntyre)
(CO)NH(CH <sub>2</sub> ) <sub>3</sub> N(CH <sub>3</sub> ) <sub>3</sub> <sup>+</sup> CH <sub>3</sub> OSO <sub>3</sub> <sup>-</sup>	Ricinamidopropyltrimonium methosulfate	Rewoquat <sup>TM</sup> RTM 50 (Rewo)
(CO)NH(CH <sub>2</sub> CH <sub>2</sub> O) <sub>n</sub> H n is average of polydisperse compositions	PEG-40 Ricinoleamide	
(CO)O(CH <sub>2</sub> CH <sub>2</sub> O) <sub>n</sub> H n is average of polydisperse compositions	PEG-# Ricinoleate Where # is 2 through 9 and correspond to n	Atlas G4929 (ICI)
(CO)NHCH <sub>2</sub> CH <sub>2</sub> OCOCH(SO <sub>3</sub> Na)CH <sub>2</sub> COONa	Disodium ricinoleamido MEA sulfosuccinate	Mackanate RM (McIntyre) Monamate <sup>TM</sup> RMEA-40 (Mona)
(CO)OCH <sub>2</sub> CH(OH)CH <sub>2</sub> OH	Glyceryl ricinoleate	Aldo <sup>TM</sup> MR (Lonza) Tegin <sup>TM</sup> RZ (Goldschmidt)
(CO)OCH <sub>2</sub> CH(OH)CH <sub>2</sub> (OCH <sub>2</sub> CH <sub>2</sub> ) <sub>n</sub> OH	PEG-# Glyceryl Ricinoleate	

Some non-limiting examples of undecylenyl-based olefin compounds

- 5 include those having the following structure:



Examples of substructures that may be used as the -X group are provided in Table 3.


**Table 3 Undecylenyl-Based -X Groups**

10


Substructure -X	INCI Name	Example Trade Names
(CO)N(CH <sub>2</sub> CH <sub>2</sub> OH) <sub>2</sub>	Undecylenamide DEA	Rewocid <sup>TM</sup> DU 185 (Rewo Chemische)
(CO)NH(CH <sub>2</sub> ) <sub>3</sub> N(CH <sub>3</sub> ) <sub>2</sub> →O	Undecylenamidopropylamine oxide	
(CO)N(CH <sub>3</sub> ) <sub>2</sub> <sup>+</sup> CH <sub>2</sub> COO <sup>-</sup>	Undecylenamidopropyl betaine	
(CO)NHCH <sub>2</sub> CH <sub>2</sub> OCOCH(SO <sub>3</sub> Na)CH <sub>2</sub> COONa	Disodium undecylenamido MEA sulfosuccinate	Emcol 5330 (Witco) Rewocid SB U 185 (Rewo)

In addition to the above-described olefin compounds, the following olefin compounds are examples of olefin compounds that may be used in the present invention:

1. Disodium Dihydroxyethyl Sulfosuccinylundecylenate  
5 (commercially available as Grillosan DS 7911 from RITA).




2. Disodium PEG-8 Ricinosuccinate (commercially available as Grillosan 8C12 from RITA).



10

3. Sodium Bisglycol Ricinosulfosuccinate (commercially available as Grillosol 8C from RITA).



15

In some preferred embodiments, the olefin compound is water soluble or water dispersible. In some preferred embodiments, the olefin compound has an HLB equal to or greater than 8. Preferred olefin compounds have the hydrophilic group connected to the olefin group via an amide, ether, quaternary ammonium, or amine group. Most preferably, the  
20

olefin compound is an oleyl-, ricinoleyl, or undecylenyl-based derivative where the hydrophilic group is connected to the olefin group via an amide, ether, quaternary ammonium group, or amine group.

Some examples of preferred olefin compounds include sodium  
5 oleoyl N-methyl taurate, oleth-X where X is 3-20, oleth-X phosphate where X is 3-20, oleth-X carboxylate where X is 3-20, oleth sulfate, oleamidopropyl betaine, ricinoleamidopropyl betaine, and oleylmethylbis(hydroxyethyl)ammonium chloride.

The olefin compound is preferably an oleyl-based compound, and in  
10 particular sodium N-methyl oleoyl taurate commercially available as Geropon T-77 from Rhodia; oleamidopropylbetaine commercially available as Mackam HV from McIntyre; Oleth-10 and 20 commercially available as Volpo-10 and Volpo-20 from Croda; Oleth-10 phosphate commercially available as Crodofos-10 from Croda; and  
15 oleylmethylbis(hydroxyethyl)ammonium chloride commercially available as Ethoquad O/12 PG from Akzo Nobel.

The olefin compound is present in the first part, the second part, or both the first and second parts in an amount such that when the first part and second part are combined, it is present within the disinfecting  
20 composition at a concentration ranging from about 0.005% to about 50% by weight, generally from about 0.05% to about 10% by weight, and from about 0.1% to about 5% by weight. Mixtures of olefin compounds are also encompassed within this invention. For example, one type of olefin

compound may be present in the first part, with a different type present in the second part.

Various optional ingredients may also be present in the first part, the second part, or both first and second parts of the two-part system. Such ingredients include (but are not limited to) wetting agents, builders, textural modifiers, film-forming polymers, surfactants, colorants, preservatives and mixtures thereof. The wetting agents facilitate contact of the disinfecting composition with the skin or surface, and can be selected from those materials recognized to provide this effect, in both identity and amount. Builders help boost the cleaning performance of surfactant systems and are typically associated with materials that are capable of complexing with polyvalent cations, such as calcium and magnesium. Textural modifiers are those materials which primarily affect the body of the mixed disinfecting composition in terms of retention, flow and lubricity. These include thickening agents such as alkyl celluloses, alkoxy celluloses, xanthan gum, guar gum, and polyacrylamide derivatives, of which the polymer of 2-acrylamido-2-methylpropane sulfonic acid is a preferred example. Inorganic thickening agents include hectorite, synthetic hectorite, magnesium aluminum silicate, bentonite, montmorillonite, and amorphous silicon dioxide. Thickening can also be achieved by the combination of selective surfactant classes. Other textural modifiers include lanolin derivatives, acyl lactylates, polyethylene glycol, glyceryl esters, and mixtures thereof. Skin conditioning and skin healing agents include glycerin, sorbitol, pyrrolidone carboxylic acid, mineral oils, silicone oils, protein hydrolysates, petrolatum, hydrocarbon

emollient alcohols and esters, allantoin, aloe vera extracts, and urea. Film-forming polymers include the above-referenced polyacrylamides, as well as the class of poly(vinyl alcohols/vinyl acetates), polyurethanes, chitosan, polyvinyl pyrrolidone, and polyvinyl pyrrolidone copolymers.

5           In a further embodiment, a method for disinfecting a substrate is disclosed, wherein the method comprises contacting the substrate with an effective amount of the disinfecting composition formed by combining the first part and the second part of the two-part disinfecting system of this invention. In this context, the substrate may be any surface or material in need of, or that  
10           would benefit from, such disinfection, including (but not limited to) skin or tissue, as well as body fluids and mucosal membranes. For example, the substrate may be a wound where disinfection would aid healing. The substrate may be the inside of an animal's mouth where disinfection would help prevent gingivitis and halitosis. The substrate may include any item that is intimately  
15           placed in, on, or around the body of an animal, such as dentures, braces, and contact lenses. In a specific application, the substrate is the teat of a dairy cow, goat or sheep. In addition, the substrate may be any surface of a food product, such as meat, fish, seafood, fruits and vegetables. The substrate may also include food contact surfaces, and nonfood contact surfaces in food processing plants.  
20           The substrate may include any hard surface, such as (but not limited to) floors, walls, countertops, containers, instruments and/or equipment found in homes, hospitals, and manufacturing facilities. In a specific application, the

hard surfaces may include housing and equipment surfaces in animal rearing and production environments. Materials that may benefit from disinfection include, for example, process waters, such as flume waters, cooling tower waters, livestock drinking waters, equipment, and facility  
5 cleaning solutions.

In a further aspect of this invention, this invention is directed to a method for making a disinfecting composition comprising combining the first part and the second part of the two-part disinfecting system. In one embodiment, the first and second parts are both aqueous solutions,  
10 emulsions, microemulsions, creams or gels, and may be adapted to be combined in equal or different volumes. In another embodiment, at least one of the first or second parts is in a concentrated, non-aqueous or solid form, and the concentrated, non-aqueous or solid form is first diluted with or dissolved in water, and then combined with the other part. Alternatively,  
15 the dilution or dissolving step can occur prior to combination with the other part, or simultaneous with combination.

The following examples are provided for the purpose of illustration, not limitation.

### EXAMPLES

#### 20 Example 1

A series of two-part acidified chlorite products were prepared by combining an acid part comprising 2.64% lactic acid, 10.00% glycerin, 0.30% FD&C Yellow #5, 0.30% xanthan gum, 0.04% sodium benzoate and an olefin compound formulated at 0.045M (or no olefin compound for the

control) with a chlorite part containing 0.64% sodium chlorite and sufficient quantities of sodium hydroxide to yield a pH of 2.75 when mixed 1:1. The rate of color fading at 35°C was quantified by UV/Vis spectroscopy at 425nm.

5 **Table 4** Olefin Compounds and Effect on Color Fading at 35°C

Olefin Compound (0.045 M)	Weight Percent*	Time After Mixing (** >50% of Initial Color Intensity Remaining)				
		½ hour	1 hour	2 hour	3 hour	4 hour
Sodium N-methyl oleoyl taurate	1.91	**	**	**	**	**
Oleth-10	3.19	**	**	**	**	**
Oleth-10 phosphate	3.55	**	**	**	**	**
Oleamidopropyl betaine	1.91	**	**	**	**	**
Oleymethylbis (hydroxyethyl) ammonium chloride	1.81	**	**	**	**	**
Disodium oleamido MEA sulfosuccinate	2.48	**	**	**	**	**
Control (no olefin)	----	**	-	-	-	-

\*The weight percent corresponds to the active weight percent of olefin derivative in the acid part. This corresponds to an equal molar comparison of the olefin derivative at 0.0225M in the mixed product composition.

10

The olefin compounds in Table 4 significantly extended the color marking longevity vs the control.

### Example 2

15 Three compositions were prepared according to example 1 where the olefin compound was 0.4% sodium oleoyl N-methyl taurate (Geropon T-77), or a saturated homolog of 0.4% sodium cocoyl N-methyl taurate (Geropon TC 270), or a no olefin control. All products possessed a mixed

product pH of 2.75. The rate of color fading at 35°C was quantified by UV/Vis spectroscopy at 425nm.

**Table 5** Comparison of a Olefin vs Non-Olefin Compound and the Effect of Color Fading at 35°C

5

Component	Time After Mixing (** >50% of Initial Color Intensity Remaining)				
	½ hour	1 hour	2 hour	3 hour	4 hour
Sodium N-methyl oleoyl taurate	**	**	**	**	-
Sodium N-methyl cocoyl taurate	**	-	-	-	-
Control (no olefin)	**	-	-	-	-

Table 5 illustrates that the olefin compound (sodium N-methyl oleoyl taurate) extended the colorant intensity longer while the saturated cocoyl homolog did not.

10 **Example 3**

A series of two-part acidified chlorite products were prepared that when combined 1:1 produced compositions containing 1.32% lactic acid, 0.32% sodium chlorite, an olefin compound formulated at a 0.0225M, and sufficient quantities of sodium hydroxide to produce a mixed product pH of

15 2.75. The rate of chlorine dioxide generation was determined by UV/Vis spectroscopy at 360 nm (54.31 ppm/A.U. cm).

**Table 6** Olefin Compounds and the Rate of Chlorine Dioxide Generation

Olefin Compound	Weight Percent*	Time After Mixing (ppm Chlorine Dioxide)				
		1 hour	2 hour	3 hour	4 hour	5 hour
Sodium N-methyl oleoyl taurate	1.91	4	5	5	6	7
Oleth-10	3.19	10	16	21	25	28
Oleth-10 phosphate	3.55	7	14	17	21	25
Control (no olefin)	----	40	62	83	96	110

5 \*Weight percents correspond to the active weight percent of the olefin derivative in the acid part. This corresponds to an equal molar comparison of the olefin derivative at 0.0225M in the mixed product composition.

Table 6 illustrates that the olefin compounds significantly slowed the rate of chlorine dioxide generation relative to the control.

#### 10 Example 4

Three acidified chlorite compositions were prepared according to Example 1 where the olefin compound was 1.4% sodium oleoyl taurate (Geropon T-77), or 1.4% alpha olefin sulfonate (Bioterge AS-40, Stepan), or a no olefin control, with a chlorite part containing 0.64% sodium chlorite and sufficient quantities of sodium hydroxide to achieve a pH of 2.75 when  
15 mixed 1:1. The rate of sodium chlorite consumption was quantified using a standard iodometric titration method.

**Table 7** Olefin Compounds and the Rate of Sodium Chlorite Consumption

20

Olefin Compound	Time After Mixing (** >50% Initial NaClO <sub>2</sub> Level)				
	1 hour	2 hour	3 hour	4 hour	5 hour
Sodium N-methyl oleoyl taurate	**	**	**	**	**
Alpha olefin sulfonate	**	**	**	-	-
Control (no olefin)	**	-	-	-	-

Results in Table 7 illustrate that olefin compounds significantly slowed the rate of sodium chlorite consumption. Sodium N-methyl oleoyl taurate (Geroxon T-77) exhibited a stronger effect than alpha olefin sulfonate (Bioterge AS-40) on an equal weight-base comparison.

5 **Example 5**

The germicidal performance of a two-part acidified chlorite composition was evaluated using the AOAC Germicidal and Detergent Sanitizing Action test method with a 15 second exposure period. A disinfectant composition was prepared according to Example 1 with 0.4% sodium oleoyl N-methyl taurate as the olefin compound. Ninety-nine mL samples were taken initially, at 2.5 hours, and at 5 hours after mixing and inoculated with 1 mL of the bacterial challenge. Samples from the resulting solutions were neutralized, plated, and enumerated using standard methods. In parallel, a water control was tested in place of the disinfectant composition.

**Table 8** Germicidal Testing of Acidified Chlorite Composition (15 sec. Exposure Time)

Bacteria	Time After Mixing (** no surviving bacteria observed)			Control Bacterial Count (Log CFU/mL)
	Initial	2.5 hours	5 hours	
<i>Pseudomonas aeruginosa</i> ATCC 15442	**	**	**	6.92
<i>Escherichia coli</i> ATCC 11229	**	**	**	6.97
<i>Klebsiella pneumoniae</i> ATCC 4352	**	**	**	6.81
<i>Enterobacter aerogenes</i> ATCC 13048	**	**	**	7.20
<i>Streptococcus uberis</i> ATCC 27958	**	**	**	6.51
<i>Streptococcus dysgalactiae</i> ATCC 27957	**	**	**	5.93

The results in Table 8 show that the present invention is effective at killing a variety of microorganisms compared to a control having a bacterial count of almost at least six logs in every example.

**Example 6**

- 5 **Table 9** A two part teat disinfectant product to mixed 36.57:1 by volume Acid Part + Chlorite Part.

Acid Part	wt%
Lactic Acid	2.95%
Propylene Glycol	10.00%
Lanolin Ethoxylate	1.00%
FD&C Blue #1	0.05%
FD&C Yellow #5	0.10%
Linear Dodecylbenzene	2.00%
Sulfonic Acid	
Mackam HV <sup>1</sup>	2.00%
Xanthan Gum	0.30%
Sodium Benzoate	0.18%
Potassium Hydroxide	0.18%
DI Water	q.s.
Chlorite Part	wt%
Sodium Chlorite, 25%	15.00%
DI Water	q.s.

1. Mackam HV: Oleamidopropyl Betaine (McIntyre)

**Example 7****Table 10** Two-part teat disinfectant product to be mixed 1:1 by volume

Acid Part	wt%
Mandelic Acid	3.00%
Glycerin	5.00%
FD&C Yellow #5	0.30%
Hydroxyethyl Cellulose	1.00%
Sodium Benzoate	0.20%
Bioterger <sup>TM</sup> AS-40 <sup>2</sup>	0.50%
Crodafos 10 <sup>3</sup>	0.50%
DI Water	q.s
Chlorite Part	wt%
Sodium Chlorite	0.64%
Polysulfonic Acid <sup>4</sup>	15.00%
Edetate Disodium	0.17%
Sodium Hydroxide	0.56%
DI Water	q.s.

2. Bioterger AS-40: C14-16 Olefin Sulfonate (Stapan)

3. Crodafos 10: Oleth -10 Phosphate (Croda)

4. Polysulfonic Acid: Cosmedia<sup>TM</sup> HSP (Cognis)

5

**Example 8****Table 11** Two-part teat disinfectant product to be mixed 1:1 by volume

Acid Part	wt%
Lactic Acid	2.64%
Glycerin	10.00%
FD&C Yellow #5	0.30%
Xanthan Gum	0.30%
Sodium Benzoate	0.04%
Volpo-10 <sup>5</sup>	2.00%
Octanoic Acid	0.20%
DI Water	q.s.
Chlorite Part	wt%
Sodium Chlorite	0.64%
Edetate Disodium	0.04%
Sodium Hydroxide	0.032%
DI Water	q.s.

5. Volpo-10, Oleth-10 (Croda)

10

**Example 9**

**Table 12** A two part hard surface cleaner/sanitizer to be mixed 1:1 by volume.

Acid Part	wt%
Citric Acid	2.00%
Neodol <sup>TM</sup> 1-5 <sup>6</sup>	1.50%
Volpo 10	1.00%
Dowanol <sup>TM</sup> DnB <sup>7</sup>	3.00%
DI Water	q.s.
Chlorite Part	wt%
Sodium Chlorite, 25%	2.00%
DI Water	q.s.

6. Neodol 1-5: C11 Akyl Ethoxylate EO=5 (Shell)

5 7. Dowanol DnB: Propylene Glycol n-Butyl Ether (Dow)

**Example 10**

**Table 13** A two part mouth rinse to be mixed 1:1 by volume:


Acid Part	wt%
Malic Acid	2.70%
Propylene Glycol	20.0%
Menthol	0.07%
Sodium Benzoate	0.05%
Sodium Saccharin	0.50%
FD&C Blue #1	1.5 ppm
Sodium Oleth Sulfate	0.10%
DI Water	q.s.
Chlorite Part	wt%
Sodium Chlorite	0.27%
Sodium Fluoride	0.10%
Sodium Hydroxide	0.07%
DI Water	q.s.

10 From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the

spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.

## WHAT IS CLAIMED IS:

1. A two-part disinfecting system comprising a first part and a second part adapted to be mixed to yield an aqueous disinfecting composition, wherein the first part comprises a chlorite and the second part comprises an acid, and wherein the first part, the second part, or both the first and second parts contain an olefin compound having the formula:



where  $R_1 = H-$ ,  $CH_3-$ ,  $CH_3(CH_2)_n-$ ,  $CH_3(CH_2)_nCH(OH)-$ ,  
or  $MO_2C(CH_2)_n-$ ;

10  $n = 1-20$ ;

$$R_2 = -(CH_2)_m-, \text{ or } -\underset{\substack{| \\ \text{CO}_2M}}{CH}(CH_2)_m-$$

$m = 2-20$ ;

15  $M = H, Na, K, Ca, Mg, \text{ ammonia, or an organic cation; and}$

$X = \text{a hydrophilic group having at least four heteroatoms selected from the group consisting of oxygen, nitrogen, sulfur, chlorine, bromine, fluorine, iodine, phosphorous and mixtures thereof,}$

20 wherein there are at least two saturated carbon atoms between  $(R_1)CH=CH$  and a heteroatom and a saturated carbon is a carbon free of double and triple bonds and the olefin compound is present in the first part, the second part, or both the first part and second part so that, when the first part and second part are

combined, it is present within the disinfecting composition at a concentration ranging from about 0.005% to about 50% by weight.

2. The system of claim 1, further comprising an oxidizable colorant present in the first part, the second part, or both the first part and second  
5 part.
3. The system of claim 1, wherein the olefin compound is selected from the group consisting of oleyl-based, ricinoleyl-based, and undecylenyl-based olefin compounds, and mixtures thereof.
4. The system of claim 1, wherein the olefin compound is selected  
10 from the group consisting of sodium oleoyl N-methyl taurate, Oleth-X, Oleth-X phosphate, Oleth-X carboxylate where X is 3-20, oleth sulfate, oleamidopropyl betaine, ricinoleamidopropyl betaine, oleylmethylbis(hydroxyethyl)ammonium chloride and mixtures thereof.
5. The system of claim 1, wherein the hydrophilic group is connected  
15 to the olefin compound via an amide, ether, quaternary ammonium or amine group.
6. The system of claim 1, wherein the olefin compound is present in the first part, the second part, or both the first and second parts in an amount so that, when the first part and second part are combined, it is  
20 present within the disinfecting composition at a concentration ranging from 0.05% to 10% by weight.
7. The system of claim 1, wherein the chlorite is a metal chlorite.
8. The system of claim 7, wherein the metal chlorite is an alkali or alkaline earth metal chlorite.

9. The system of claim 7, wherein the metal chlorite is sodium chlorite or potassium chlorite.
10. The system of claim 1, wherein the chlorite is present in the first part in an amount so that, when combined with the second part, it is present  
5 within the disinfecting composition at a concentration ranging from about 0.005% to about 3% by weight.
11. The system of claim 1, wherein the acid has a  $pK_a$  ranging from 2 to 5.
12. The system of claim 1, wherein the acid is an organic acid, an  
10 inorganic acid, or mixtures thereof.
13. The system of claim 12, wherein the organic acid is glycolic acid, lactic acid, malic acid, mandelic acid, citric acid, tartaric acid, adipic acid, succinic acid, malonic acid, heptanoic acid, octanoic acid, nonanoic acid, benzoic acid, gluconic acid, or a mixture thereof.
- 15 14. The system of claim 12, wherein the inorganic acid is sulfuric acid, sulfamic acid, phosphoric acid, hydrochloric acid, nitric acid, or a mixture thereof.
- 15 15. The system of claim 1, wherein the acid is present in the second part in an amount so that, when combined with the first part, it is present within  
20 the disinfecting composition at a concentration ranging from about 0.05% to about 10% by weight.
16. The system of claim 1, wherein the acid is present in the second part in an amount so that, when combined with the first part, the pH of the disinfecting composition is below about 5.


17. The system of claim 2, wherein the oxidizable colorant is (a) a food, drug and cosmetic colorant, (b) a drug and cosmetic colorant, or (c) an externally applied drug and cosmetic colorant.

18. The system of claim 1, wherein both the first part and the second part are independently in the form of an aqueous solution, emulsion, microemulsion, cream or gel.

19. The system of claim 1, wherein the first part, the second part, or both the first and second parts further comprise a textural modifier, wetting agent, thickening agent, skin conditioner, healing agent, film-forming polymer, surfactant, preservative, or a mixture thereof.


20. Use of the resulting disinfecting composition of claim 1, the use comprising the application of the composition to a surface selected from the group consisting of a hard surface, animal surface, human surface, food surface, meat surface, poultry surface, seafood surface, and process waters.

21. A disinfecting composition comprising a chlorite, an acid and an olefin compound having the formula:



where  $R_1 = H-$ ,  $CH_3-$ ,  $CH_3(CH_2)_n-$ ,  $CH_3(CH_2)_nCH(OH)-$ , or  $MO_2C(CH_2)_n-$ ;

$n = 1-20$ ;



$m = 2-20$ ;

$M = H, Na, K, Ca, Mg, \text{ ammonia, or an organic cation; and}$

X = a hydrophilic group having at least four heteroatoms selected from the group consisting of oxygen, nitrogen, sulfur, chlorine, bromine, fluorine, iodine, phosphorous and mixtures thereof,

5 wherein there are at least two saturated carbon atoms between  $(R_1)CH=CH$  and a heteroatom and a saturated carbon is a carbon free of double and triple bonds.


22. The composition of claim 21, wherein the olefin compound is present from about 0.005 wt.% to about 50 wt.%.

10 23. The composition of claim 21, further comprising an oxidizable colorant.

24. The composition of claim 21, wherein the olefin compound is selected from the group consisting of oleyl-based, ricinoleyl-based, and undecylenyl-based olefin compounds, and mixtures thereof.

15 25. The composition of claim 21, wherein the olefin compound is selected from the group consisting of sodium oleoyl N-methyl taurate, oleth-X, oleth-X phosphate, oleth-X carboxylate where X is 3-20, oleth sulfate, oleamidopropyl betaine, ricinoleamidopropyl betaine, oylethmethylbis(hydroxyethyl)ammonium chloride and mixtures thereof.

20 26. A two-part disinfecting system comprising a first part and a second part adapted to be mixed to yield an aqueous disinfecting composition, wherein the first part comprises a chlorite and the second part comprises an acid, and wherein the second part contains an olefin compound having the formula:



wherein the olefin compound is present in the second part so that, when the first part and second part are combined, the olefin compound is present within the disinfecting composition at a concentration ranging from  
5 about 0.005% to about 50% by weight.

27. The system of claim 26, further comprising an oxidizable colorant present in the first part, the second part, or both the first and second part.

28. The system of claim 26, wherein the olefin compound is present in the second part in an amount so that, when the first part and second part are  
10 combined, it is present within the disinfecting composition at a concentration ranging from 0.05% to 10% by weight.

29. The system of claim 26, wherein the chlorite is a metal chlorite.

30. The system of claim 29, wherein the metal chlorite is an alkali or alkaline earth metal chlorite.

15 31. The system of claim 30, wherein the metal chlorite is sodium chlorite or potassium chlorite.

32. The system of claim 26, wherein the chlorite is present in the first part in an amount so that, when combined with the second part, it is present within the disinfecting composition at a concentration ranging from about  
20 0.005% to about 3% by weight.

33. The system of claim 26, wherein the acid is lactic acid.

34. The system of claim 26, wherein the acid is present in the second part in an amount so that, when combined with the first part, it is present

within the disinfecting composition at a concentration ranging from about 0.05% to about 10% by weight.

35. The system of claim 27, wherein the oxidizable colorant is a food, drug and cosmetic colorant.

5