

(19)



(11) Publication number:

SG 182137 A1

(43) Publication date:

30.07.2012

(51) Int. Cl:

;

(12)

Patent Application

(21) Application number: **2012038915**

(71) Applicant:

**IRON THERAPEUTICS HOLDINGS AG
SIHLEGGSTRASSE 23 8832 WOLLERAU
CH**

(22) Date of filing: **14.05.2009**

(30) Priority: **GB 0808835.3 15.05.2008**

(72) Inventor:

**STOCKHAM, MICHAEL ARTHUR
"BALTROY" STICKLING GREEN
CLAVERING SAFFRON WALDEN ESSEX
CB11 4QX GB**

(54) Title:

**MONO (IRON HYDROXYPYRONE) AND COMBINATION
(IRON HYDROXYPYRONE AND GI INFLAMMATION
INHIBITING AGENTS) COMPOSITIONS FOR ANAEMIA OR
H. PYLORI INFECTIONS**

(57) Abstract:

28 ABSTRACT MONO (IRON HYDROXYPYRONE) AND COMBINATION (IRON HYDROXYPYRONE AND GI INFLAMMATION INHIBITING AGENTS) COMPOSITIONS FOR ANAEMIA OR H. PYLORI INFECTIONS There is provided a composition or kit of parts comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract: and an iron hydroxypyrrone, for increasing the level of iron in a patient's bloodstream and/or treating and/or preventing anaemia such as iron deficiency anaemia. A composition comprising iron hydroxypyrrone is also provided for administration to a subject: having or at risk of having achlorhydria; wherein the gastric pH of the subject is equal to or greater than about 4; or wherein the subject has an inflammatory disease of the gastrointestinal tract. Fig. None

ABSTRACT

MONO (IRON HYDROXYPYRONE) AND COMBINATION (IRON HYDROXYPYRONE AND GI INFLAMMATION INHIBITING AGENTS) COMPOSITIONS FOR ANAEMIA OR H. PYLORI INFECTIONS

There is provided a composition or kit of parts comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyronone, for increasing the level of iron in a patient's bloodstream and/or treating and/or preventing anaemia such as iron deficiency anaemia. A composition comprising iron hydroxypyronone is also provided for administration to a subject: having or at risk of having achlorhydria; wherein the gastric pH of the subject is equal to or greater than about 4; or wherein the subject has an inflammatory disease of the gastrointestinal tract.

Fig. None

MONO (IRON HYDROXYPYRONE) AND COMBINATION (IRON
HYDROXYPYRONE AND GI INFLAMMATION INHIBITING AGENTS)
COMPOSITIONS FOR ANAEMIA OR H. PYLORI INFECTIONS

This invention relates to therapeutic compositions which comprise iron ions and to their medical applications.

5

An adequate supply of iron to the body is an essential requirement for tissue growth and the maintenance of good health in both man and animals. Moreover, in certain pathological conditions where there is an insidious blood loss, or where there is a mal-distribution of iron in the body, there may be a state of low iron stores in the body and a concomitant chronic anaemia. This is seen in inflammatory diseases of the gastrointestinal tract, such as gastric and peptic ulcers, reflux oesophagitis, ulcerative colitis and Crohn's disease.

10

Anaemia can also follow operations which result in serious blood loss and can be associated with gastrointestinal infections, such as those caused by *Helicobacter pylori*.

15

Achlorhydria (and hypochlorhydria) refer to states where the production of gastric acid in the stomach is absent or low, respectively. Achlorhydria can be due to many diverse causes including: pernicious anaemia, an autoimmune gastritis, other autoimmune conditions, such as autoimmune thyroid disease, any cause of severe chronic gastritis (*H. pylori* is the most common agent that may lead to the destruction of parietal cells (the cells that make hydrochloric acid) in the stomach, resulting in achlorhydria), mucopolidosis type IV (an autosomal recessive lysosomal storage disease), the use of antacids or drugs that decrease gastric acid production or transport, or atrophic gastritis.

20

25

The listing or discussion of an apparently prior-published document in this specification should not necessarily be taken as an acknowledgement that the document is part of the state of the art or is common general knowledge.

30

WO 96/41627 describes ferric iron complexes of hydroxypyrones, which comprise a carboxylic acid such as citric acid. The compositions are useful for the treatment of iron deficiency anaemia.

35

The use of certain metal ion complexes for treating gastrointestinal infection with *Helicobacter pylori* is disclosed in WO 98/16218. The complexes include

complexes of iron in the ferric state. The patients were not, however, treated with acid reducing regimens.

- 5 WO 01/89534 describes the use of cobalt salts to selectively treat gastrointestinal infections caused by *H. pylori*.

WO 02/24196 discloses compositions in solid form, such as powders, comprising a mixture of a ferrous salt and a hydroxypyrrone which may be used to increase
10 the level of iron in a patient's bloodstream or to treat and/or prevent gastrointestinal infection.

WO 2005/048912 describes methods and compositions for the treatment of helicobacter pylori-associated disorders using endoperoxide bridge-containing
15 compounds.

WO 03/097627 discloses a method of forming an iron hydroxypyrrone.

US 2007/0161600 describes the treatment of iron related conditions with iron-
20 carbohydrate complexes.

It is known that the absorption of iron is aided by the acid secretion of the stomach and that absorption is more readily affected when iron is in the ferrous state. It is also known that antacids may impair the absorption of iron by the
25 formation of insoluble complexes (Martindale, page 1347, 32nd Edition, 1999). Research suggests that taking acid-reducing drugs can cause iron deficiency due to reduced absorption (Aymard *et al*, Med. Toxicol. Adverse Drug Exp. 1988, vol. 3, pp 430-448). Hence, a person skilled in the art would not have expected iron to be available for absorption in the stomach in the presence of acid reducing
30 agents (www.numarkpharmacists.com/hn/Drug/Famotidine.html).

There are many medical conditions where the gastric environment is unfavourable for iron absorption yet iron absorption is required in order to address anaemia. For example, where gastritis and a high pH (such as above
35 6.5) prevails the absorption of iron may be too low to effectively treat anaemia.

Furthermore, the treatment regimen for conditions, such as Helicobacter infection, typically involves the use of acid reducing agents. The present invention can be considered, at least in part, to relate to the unexpected finding that acid-reducing regimen, such as are required for the treatment of, for example, ulcers, gastritis and acid reflux, can be compatible with certain iron hydroxypyrones and that the iron may be absorbed effectively from the iron hydroxypyrones despite the use of an acid-reducing regimen.

The present invention aims to alleviate some of the problems of existing anaemia treatments.

According to the invention in a first aspect there is provided a composition comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrene, for increasing the level of iron in a patient's bloodstream.

In another aspect there is provided a composition comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrene, for preventing and/or treating anaemia such as iron-deficiency anaemia.

In a further aspect there is provided a composition or kit of parts for increasing the level of iron in a patient's bloodstream comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrene, preferably wherein the iron hydroxypyrene comprises iron in the ferrous state.

In a still further aspect there is provided a composition or kit of parts for increasing the level of iron in a patient's bloodstream comprising one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract and an iron hydroxypyrene preferably with the proviso that when iron is in the ferric state, the compound is not a cobalt salt. Preferably, the composition does not comprise an antibiotic.

In another aspect, there is provided a composition comprising an iron hydroxypyrrone, for administration to a subject having, or at risk of having, achlorhydria.

5 In one aspect of the invention there is provided a composition comprising an iron hydroxypyrrone, for administration to a subject, wherein the gastric pH of the subject is equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.

10 In another aspect of the invention, there is provided a composition comprising an iron hydroxypyrrone for administration to a subject, wherein the subject has, or is suffering from, an inflammatory disease of the gastrointestinal tract and preferably the gastric pH of the subject is equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.

15

In one embodiment of the invention, the inflammatory disease of the gastrointestinal tract requires a stomach acid reducing treatment regimen as defined herein. However, the inflammatory disease may, in one embodiment, not require an acid reducing treatment regimen as defined herein.

20

The iron hydroxypyrrone, the inflammatory disease and the gastric pH, and combinations thereof, in any of the above aspects is preferably as defined herein.

25 In one embodiment of the invention, the iron hydroxypyrrone is administered alone, such as in the absence of other pharmaceutically active compounds, for example in the absence of one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract.

30 In one embodiment of the invention, the composition and/or iron hydroxypyrrone is for the absorption of iron, preferably following oral administration, for example into the blood and/or body of a subject.

Examples of an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen include gastric or

peptic ulcers, reflux, oesophagitis, ulcerative colitis, Crohn's disease, gastritis, Zollinger-Ellison syndrome or acid reflux.

In all aspects or embodiments of the invention, the one or more compounds
5 capable of treating and/or preventing an inflammatory disease of the
gastrointestinal tract which preferably requires a stomach acid reducing treatment
regimen preferably comprise an acid reducing regimen. Acid reducing regimens
are well known to a person skilled in the art. For example, the acid reducing
regimen may comprise one or more compounds capable of treating and/or
10 preventing inflammatory diseases of the gastrointestinal tract, such as, for
example, gastric or peptic ulcers, reflux oesophagitis, ulcerative colitis, Crohn's
disease, gastritis, Zollinger-Ellison syndrome or acid reflux.

In all aspects and embodiments of the invention, the peptic ulcer is preferably a
15 gastric ulcer and the anaemia is preferably iron-deficiency anaemia. The anaemia
is preferably associated with an inflammatory disease of the gastrointestinal tract
as defined herein.

In one embodiment of the invention, the anaemia is caused by blood loss
20 following an operation and/or is associated with a gastrointestinal infection, such
as those caused by *Helicobacter pylori*.

The composition and/or kit of parts according to the invention is preferably for
administration to a human.

25

The term "subject" as used herein preferably refers to a mammal, such as a
human. The term "mammal" also includes veterinary animals such as for
example, horses, cows, sheep and pigs, as well as pets such as, for example,
dogs, cats and hamsters.

30

In one embodiment of the invention, the subject has been, will be, or is being
administered an acid reducing regimen or one or more compounds capable of
treating and/or preventing an inflammatory disease of the gastrointestinal tract, as
defined herein.

35

In one embodiment of the invention, the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract comprise an acid reducing regimen. Surprisingly, it has been found that an iron hydroxypyrrone according to the invention can be administered to patients in combination with acid reducing therapies for treating gastrointestinal conditions in which there is produced excess stomach acid. The iron hydroxypyrrone can also be administered to patients generally having a high gastric pH, such as equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above, for example, from greater than 4 to 8, 4.5 to 7, 5 to 6.5 or 5.5 to 6.0, and be absorbed effectively. The high gastric pH may be, for example, as a result of achlorhydria, administration of an acid reducing regimen, such as defined herein, or an inflammatory disease of the gastrointestinal tract as defined herein, as well as combinations thereof. The administration of the iron hydroxypyrrone may be for the absorption of iron, for example, following oral administration and/or may be used to maintain levels of iron in the body of a subject i.e. to provide or maintain normal, healthy or nutritional levels of iron.

According to the present invention, the iron hydroxypyrrone may be administered at the same time, before or after the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract, or the acid reducing regimen. In one embodiment, the iron hydroxypyrrone is administered at the same time or subsequent to the one or more compounds or the acid reducing regimen. For example, the iron hydroxypyrrone may be administered from 1 minute to 2 hours after the compound, more preferably from 10 minutes to 1 hour after the compound, most preferably about 30 minutes after the administration of the compound.

In one embodiment of the invention, the composition, as defined in any of the aspects or embodiments herein, is for administration to a human or animal having or suffering from an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen. For example, the human or animal may be suffering from a gastric or peptic ulcer, reflux oesophagitis, ulcerative colitis, Crohn's disease, gastritis, Zollinger-Ellison syndrome, acid reflux and/or a peptic ulcer and is in need of treatment therefor.

In a preferred embodiment of the invention, as defined herein, the composition is for administration to a human or animal having a gastrointestinal infection.

Although the gastrointestinal infection may be any infection, the gastrointestinal
5 infection is preferably caused by *Helicobacter pylori*.

Suitable molar ratios of the one or more compounds, or acid reducing regimen, to the iron hydroxypyrrone are from 100:1 to 1:100, preferably from 50:1 to 1:50, more preferably from 10:1 to 1:10, such as from 5:1 to 1:5, for example 2:1 to 1:2.
10 Preferably the ratio of one or more compounds to iron hydroxypyrrone is from 10:1 to 1:10 by weight.

The acid reducing regimen may be administered in accordance with the normal dosage ranges for human patients for therapeutic or preventative treatment. In
15 one embodiment of the invention, the composition may be adapted for maintenance therapy, or the prevention of an inflammatory disease of the gastrointestinal tract, wherein the disease state is as defined in any of the embodiments herein. Thus, the composition may comprise lower doses of the compound and iron hydroxypyrrone than are used for the initial treatment of an
20 established disease.

In one embodiment of the invention, the composition comprises a maintenance dose of one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract, or the acid reducing regimen
25 and a nutritional dose of iron hydroxypyrrone, as iron. For example, the daily dose of the compound may be less than 400mg, for example, less than 300mg, 200mg or 100mg. Typical daily maintenance doses of the compound (such as omeprazole, pantoprazole or ranitidine) may be from 5 to 140 mg daily, such as from 10 to 100 mg, preferably 20 to 50 mg, as one or more administrations.

30

Examples of suitable nutritional doses, preferably daily, of the iron hydroxypyrrone include less than 40 or 60 mg as iron, such as from 5 to 20 mg, as iron, for example about 15 mg (as iron). The dosages may be administered for 1 to 4 weeks typically.

35

In a further embodiment of the invention, as defined herein, the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract comprise one or more acid reducing agents. The acid reducing agent may reduce the acid level directly i.e., by neutralisation of the acid, or indirectly, i.e., by inhibiting receptors which cause the secretion of excess acid or by treating the underlying disease state, for example.

Compositions for treating gastrointestinal conditions containing a metal salt, an antibiotic and a proton pump inhibitor, in so-called "triple therapy" methods are known. Antibiotics and proton pump inhibitors are suitable for use in the compositions of the present invention, as defined herein. Suitable antibiotics include, for example, amoxicillin, metronidazole, clarithromycin and mixtures thereof. Suitable proton pump inhibitors include lansoprazole, omeprazole, pantoprazole and rabeprazole for example. H₂ receptor antagonists, such as ranitidine, famotidine, nizatidine, cimetidine, for example, may also be used in the present invention. Omeprazole may be in the form of omeprazole magnesium or sodium. Ranitidine may be in the form of ranitidine bismuth citrate or ranitidine hydrochloride.

In any aspect or embodiment of the invention, the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract, or acid reducing agent, may be selected from a proton pump inhibitor, an H₂ receptor antagonist, or a substance capable of neutralising acids, such as, for example, an alkaline agent, or a combination thereof, as defined herein or a cobalt compound as defined herein. The term "substances capable of neutralising acids" is intended to include pharmaceutically acceptable bases. Suitable examples of bases include hydrogen carbonates (bicarbonates), hydroxides, oxides, or carbonates, which are commonly used in the treatment of acid reflux. Preferably, the bicarbonates, hydroxides, oxides or carbonates are in combination with a pharmaceutically acceptable cation, such as an alkali metal, for example, sodium or potassium, or alkaline earth metal, for example, magnesium, calcium or a transition metal, such as, for example, cobalt or a metal such as aluminium.

In one embodiment of the invention, as defined herein, the acid reducing agent is selected from: omeprazole, lansoprazole, pantoprazole, ranitidine, famotidine, nizatidine, cimetidine, sodium bicarbonate, potassium bicarbonate, calcium carbonate, aluminium hydroxide, magnesium carbonate, cobalt carbonate or a
5 combination thereof.

In any aspects or embodiments of the invention, the composition may further comprise an antibiotic such as, for example, amoxicillin, metronidazole, clarithromycin and mixtures thereof.

10

In the embodiments or aspects of the invention, the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract preferably comprises or consists of a cobalt compound.

15 The cobalt compound may comprise a cobalt salt. Suitable examples of cobalt salts are defined in WO 01/89534, the teachings of which are incorporated herein by reference. The cobalt salt is preferably a cobalt (II) salt, since it is well-known that cobalt (II) salts are readily available unlike cobalt (III) salts, and because there is concern over the long term safety of cobalt (III) salts.

20

The cobalt salt must be capable of providing cobalt ions (in hydrated form) at the site of a gastrointestinal infection. It is preferred that the cobalt salt, such as cobalt (II), comprises an anion which does not complex too strongly with the cobalt ion. Preferably the anion is the anion of a strong acid (ie, having a pKa of
25 less than about 3, 4, 5, or 6, for example less than 2) and/or is monovalent.

Suitable anions include chloride, gluconate, nitrate, sulphate, phosphate, carbonate, hydroxide, acetate and mixtures thereof (including salts containing two or more anions). Of these, chloride, carbonate, gluconate, hydroxide and acetate
30 are more preferred.

The cobalt salts preferably contain cobalt ions in uncomplexed form (ie, such that $\text{Co}(\text{H}_2\text{O})_6^{2+}$ ions are formed when the salt is in aqueous solution, for example) or in the form of complexes in which the cobalt ion is relatively weakly bound such

that the complex at least partly dissociates to provide cobalt ions in a form suitable for uptake into *H. pylori* at the site of infection, for example.

Alternatively, the cobalt compound may comprise a cobalt complex. Suitable
5 examples of ligands to form cobalt complexes are defined in WO 98/16218, the teachings of which are incorporated herein by reference. Preferred cobalt complexes include, for example, cobalt, preferably in the (II) or (III) state, complexed with ascorbate, aspartate, citrate, gluconate, histidine, malate, maltol, glutamate, glutamine, succinate, gluconate, or tartrate or combinations thereof.

10

In one embodiment of the invention, as defined herein, the cobalt compound comprises cobalt (II) gluconate or cobalt (II) chloride, preferably in their hydrated forms (i.e., the dihydrate for cobalt (II) gluconate or the hexahydrate for cobalt (II) chloride), or a combination thereof.

15

In one embodiment of the invention, the amount of cobalt compound is selected such that it is selective for the treatment of gastrointestinal infections caused by *Helicobacter pylori* i.e., it is present in an amount lower than the MIC for other microorganisms, particularly beneficial bacteria, in the intestinal tract.

20

The daily amount of cobalt compound, such as cobalt (II) gluconate or cobalt (II) chloride, is preferably less than 200 mg (as cobalt), such as less than 100 mg (as cobalt), for example from 2 to 50 mg, such as from 5 to 20 mg (as cobalt).

25 In one embodiment, the amount of compound (for example cobalt), such as a cobalt salt, which is required for the treatment of gastrointestinal conditions according to the invention varies from 1 to 50 mg of compound per day, more preferably 1 to 30 mg per day, such as 5 mg or 20 mg per day (preferably as cobalt), for example. A suitable dosage form comprises 10 mg of compound
30 (preferably as cobalt) and such a dosage will typically be administered twice daily for several weeks (eg 1 to 4 weeks) in order to treat the infection. It will be appreciated by those skilled in the art that other dosage regimens may be equally applicable in the method of the invention. The cobalt compound may be in a slow release form as described in WO 01/89534, for example.

35

The term "iron hydroxypyronone" as used herein is intended to include compositions which comprise a hydroxypyronone and iron. The term includes, for example, complexes of iron with a hydroxypyronone, such as, for example, ferric trimaltol, as well as mixtures comprising an iron compound, such as a salt or a complex of iron, and a hydroxypyronone in a substantially non-complexed form (such as less than 10%, 5%, 2%, or 1% hydroxypyronone complexed), for example ferrous or ferric gluconate and maltol, preferably in the solid state.

The iron hydroxypyronone may suitably comprise iron in the ferrous (Fe^{2+}) or ferric (Fe^{3+}) oxidation states. Alternatively, the iron hydroxypyronone may comprise a mixture of iron in the ferrous and ferric oxidation states.

In one embodiment of the invention, the iron hydroxypyronone comprises iron in the ferric state. In an alternative embodiment, the iron hydroxypyronone comprises iron in the ferrous state, particularly where the iron hydroxypyronone is in the form of a solid, such as a powder or tablet.

The hydroxypyronone is preferably a 3-hydroxy-4-pyrone. Suitable pyrones include 3-hydroxy-4-pyrone itself or a 3-hydroxy-4-pyrone in which one or more of the hydrogen atoms attached to ring carbon atoms are replaced by an aliphatic hydrocarbon group of 1 to 6 carbon atoms, or a 5-hydroxypyronone, such as Kojic acid. The most preferred pyrones are maltol (3-hydroxy-2-methyl-4-pyrone) and ethylmaltol. The hydroxypyronone may, alternatively, be a natural product (such as meconic acid or iso-maltol) which may be converted to another hydroxypyronone *in vivo*.

Certain hydroxypyrones, such as maltol, are available commercially. With others, a convenient starting material in many instances consists of 3-hydroxy-4-pyrone which is readily obtainable by the decarboxylation of 2,6-dicarboxy-3-hydroxy-4-pyrone (meconic acid). For example, 3-hydroxy-4-pyrone may be reacted with an aldehyde to insert a 1-hydroxyalkyl group at the 2-position, which group may then be reduced to produce a 2-alkyl-3-hydroxy-4-pyrone. The preparation of 2-ethyl-3-hydroxy-4-pyrone, etc, by this route is described in the published US application serial number 310,141 (series of 1960). Other preparation methods

are described by Spielman, Freifelder, J. Am. Chem. Soc. Vol 69 Page 2908 (1947).

It will be appreciated by a person skilled in the art that these are not the only
5 routes available to these compounds and their iron complexes and that various alternatives may be used.

In one embodiment of the invention, the iron hydroxypyrrone is a ferric complex with hydroxypyrrone. It is preferred that the ferric complex is a 1:3 molar complex
10 of iron to the hydroxypyrrone.

Preferably, the iron hydroxypyrrone is a ferric maltol, a ferric ethylmaltol or mixtures thereof. It is particularly preferred that the ferric maltol is ferric trimaltol. The ferric trimaltol may be prepared according to any method known in the art.
15 Preferably, the ferric trimaltol is prepared as set out in WO 03/097627, the teachings of which are incorporated herein by reference.

In another embodiment of the invention, the iron hydroxypyrrone is in the form of a solid, a liquid or a suspension in a liquid and comprises a mixture of a ferrous or a
20 ferric salt and a hydroxypyrrone as defined herein. The solid mixture may also be dry (i.e., substantially free of liquid such as water i.e., the mixture comprises less than 10 wt.%, 5 wt.% or 2 wt.%, such as less than 1 wt.% liquid). Solid mixtures of ferrous salts and hydroxypyrrones are stable on storage for a moderate period of time under substantially dry conditions. However, on dissolution in water, the
25 compositions undergo a reaction in which some or all of the ferrous iron is oxidised to ferric iron.

The ferrous or ferric salt can be an iron (II) or iron (III) salt with any pharmaceutically acceptable anion. Preferably, the iron (II) salt is iron (II)
30 carbonate or an iron (II) carboxylate.

The ferrous or ferric salt is preferably an iron (II) or iron (III) carboxylate. The iron (II) carboxylate may be selected from, for example, iron (II) gluconate, iron (II) succinate or iron (II) fumarate and mixtures thereof. The iron (III) carboxylate is

preferably selected from ferric citrate, ferric ammonium citrate or ferric tartrate and mixtures thereof.

In any aspect or embodiment of the invention, the compound (or acid reducing regimen) preferably comprises a cobalt compound, such as, for example, cobalt
5 (II) gluconate or cobalt (II) chloride, preferably in a daily dosage of, for example, less than 100 mg (as cobalt), and the iron hydroxypyrrone comprises, in a solid form, a mixture of a ferrous or ferric salt of a carboxylic acid and a hydroxypyrrone, such as maltol or ethylmaltol, preferably in a daily dosage in terms of iron content
10 of, for example, 15 to 50 mg (as iron). In any embodiment of the invention, the daily dosage of the compound is preferably a maintenance dosage and the daily dosage of the iron hydroxypyrrone is preferably nutritional (i.e., it can maintain the levels of iron stored in the body by, for example, replacing any iron "leakage").

15 The composition or kit of parts for increasing the level of iron in a patient's bloodstream and/or treating and/or preventing anaemia according to the invention may comprise any of the compounds or iron hydroxypyrrones set out above, or combinations thereof.

20 In particular, the kit of parts may comprise the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract, or acid reducing agent, such as one or more compounds capable of treating gastritis, acid reflux and/or a peptic ulcer, and an iron hydroxypyrrone, wherein the compounds and the iron hydroxypyrrone are packaged separately and optionally
25 instructions are provided for the administration of the two separate parts. This may be appropriate where there is otherwise a storage incompatibility between the compound and iron hydroxypyrrone.

In one embodiment, the composition comprises a co-formulation (for example, in
30 the same solid or liquid) of the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract, or acid reducing agent such as, for example, one or more compounds capable of treating gastritis, acid reflux and/or a peptic ulcer, as defined herein, and an iron hydroxypyrrone, as defined herein. This may be suitable where there is no storage incompatibility
35 between the compound and the hydroxypyrrone.

In a preferred composition or kit of parts of the invention, the iron hydroxypyrrone comprises a solid or dry (i.e., substantially free of liquid such as water i.e., the liquid comprises less than 10 wt.%, 5 wt.% or 2 wt.%, such as less than 1 wt.%)
5 mixture of a ferrous carboxylate, such as gluconate, and maltol or ethyl maltol and the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract or acid reducing agent comprises a cobalt compound, such as cobalt gluconate (preferably cobalt(II) and hydrated). The molar ratio of ferrous salt to maltol or ethylmaltol is preferably
10 about 1:3.

In an alternative preferred composition or kit of parts of the invention, the iron hydroxypyrrone comprises ferric trimaltol and the one or more compounds capable of treating and/or preventing an inflammatory disease of the
15 gastrointestinal tract or acid reducing agent, such as one or more compounds capable of treating gastritis, acid reflux and/or a peptic ulcer, comprises potassium bicarbonate. The potassium bicarbonate may be present in the solid or liquid state, such as an aqueous solution.

20 Conveniently, the composition of the invention is in the form of a powder (which term covers fine powders and granulates) comprising a mixture of the powdered iron hydroxypyrrone and the one or more compounds. The iron hydroxypyrrone may be in crystalline form, in amorphous form or in other solid forms but is preferably crystalline. The ferrous or ferric salt may contain water of
25 crystallisation i.e., it may be in the form of a hydrate.

In a further aspect, the compositions of the invention are used in medicine and, in another aspect, pharmaceutical compositions are provided which comprise the composition of the invention together with a pharmaceutically acceptable diluent
30 or carrier.

A method of forming a composition according to the invention is also provided which comprises mixing the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract, or acid reducing
35 agent, and an iron hydroxypyrrone, all as defined herein. The mixing may take

place in the solid or liquid state. For example, both components may be in the form of substantially dry solid such as powders, or tablets, preferably having a moisture or water content less than 10 wt.%, 5 wt.%, 2 wt.% or 1wt.%.

5 In the above aspects and embodiments of the invention, the pharmaceutical composition or kit of parts of the invention may be adapted for oral administration. Suitable forms for oral administration include powders, tablets and capsules (such as gelatin capsules) as well as solutions or suspensions in aqueous or non-aqueous liquids, oil-in-water liquid emulsions, or water-in-oil liquid emulsions.

10

Suitable pharmaceutically acceptable diluents and carriers include, for example, lubricants such as magnesium stearate, stabilising and suspending agents such as methylcellulose and povidone and other tableting agents and dose bulking agents such as lactose and flow aids such as Aerosil 2000™. Particularly useful
15 diluents and carriers are wetting agents or surfactants, preferably nonionic or ionic surfactants. Examples of suitable nonionic surfactants include polyoxyl 10 oleyl ether and polysorbates. An example of a suitable ionic surfactant is sodium lauryl sulphate.

20 Alternatively, the pharmaceutical composition may be provided as a liquid or a suspension in liquid form, as a powder for reconstitution prior to oral or parental administration or it may be formulated for use as a suppository. Liquids carriers are preferably sterile and pyrogen-free: examples are saline and water. Liquid formulations are particularly suitable for oral and parenteral administration.

25

More than one iron hydroxypyrrone or one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract or acid reducing agent of the invention may be contained in the pharmaceutical composition of the invention, and other active compounds may also be included.

30 Typical possible additives include, for example, flavourings and colourings known to a person skilled in the art, as well as those set out herein.

The pharmaceutical compositions of the invention may be formulated in unit dosage form ie, in the form of discrete portions containing a unit dose, or a

multiple or sub-unit dose. Preferably, the compositions of the invention are formulated to give a rapid release of the iron for optimal absorption in the body.

5 Whilst the dosage of the composition given in each particular case will depend upon various factors, including the particular components of the composition, it may be stated by way of guidance that maintenance at a satisfactory level of the amount of iron present in the human (or animal) body will often be achieved using a daily dosage, in terms of the iron content of the compound, which lies in a range from about 1 to 150 mg, such as from 10 to 120 mg (preferably as iron).
10 However, it may be appropriate in certain cases to give daily dosages either below or above these levels. Compositions containing 15 to 50 mg iron, to be taken once daily, twice daily or three times daily (depending on the severity of the anaemia) are, for example, suitable for the treatment of anaemia.

15 The compositions of the invention suitably contain from 0.1% to 20% by weight iron, such as 0.1% to 10% by weight, for example, preferably 2 to 10% by weight.

In one embodiment of the invention, there is provided a composition comprising: an iron hydroxypyrrone; and optionally one or more compounds capable of
20 treating and/or preventing an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen, for administration to a subject having, or at risk of having, achlorhydria.

The composition is preferably for increasing the level of iron in a patient's
25 bloodstream and/or for preventing and/or treating anaemia such as iron deficiency anaemia as described above.

The achlorhydria may be a result of any of the causes set out above or known to a person skilled in the art, such as, for example, pernicious anaemia, an
30 autoimmune gastritis, other autoimmune conditions, such as autoimmune thyroid disease, any cause of severe chronic gastritis (*H. pylori* is the most common agent that may lead to the destruction of parietal cells (the cells that make hydrochloric acid) in the stomach, resulting in achlorhydria), mucopolipidosis type IV (an autosomal recessive lysosomal storage disease), the use of antacids or drugs
35 that decrease gastric acid production or transport, for example one or more

compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract which requires a stomach acid reducing treatment regimen as defined herein, or atrophic gastritis.

- 5 In one embodiment of the invention the achlorhydria is associated with, or caused by, an inflammatory disease of the gastrointestinal tract such as, for example, a gastric or peptic ulcer, reflux oesophagitis, ulcerative colitis, Crohn's disease, gastritis, Zollinger-Ellison syndrome, acid reflux and/or a peptic ulcer.
- 10 In another embodiment of the invention, the achlorhydria is associated with, or caused by, atrophic gastritis or a gastrointestinal infection, such as caused by *Helicobacter pylori*.

The composition for administration to a subject having, or at risk of having, achlorhydria, may be as defined in any of the aspects or embodiments herein, in particular, the compound and the iron hydroxypyronone may be as defined in any of the above embodiments.

20 In one embodiment of the invention, the subject may have detectable antibodies against the H^+/K^+ ATP-ase proton pump and/or the gastric pH may remain high (such as greater than 4.0, 5.0, 6.0 or 6.5), despite stimulus, such as maximum pentagastrin stimulation and/or high gastrin levels may be detected and/or a maximal acid output of less than 6.9 m/mole/h, such as less than 5.0 m/mole/h.

25 In any of the above aspects or embodiments of the invention, the pH of the environment where iron is absorbed, such as the gastric environment or gastrointestinal tract, for example, the gastric pH, is preferably equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above, for iron absorption, for example, from 4 to 8, 4.5 to 7, 5 to 6.5 or 5.5 to 6.0. Thus, the one
30 or more compounds or acid reducing regimen in any aspect or embodiment may also be defined as adapted to increase, or capable of increasing, the pH of the environment for iron absorption, such as the gastric environment or gastrointestinal tract, for example, the gastric pH, to greater than about 4, such as

greater than 5 or 6, for example about 6.5 or above (for example from for example, from 4 to 8, 4.5 to 7, 5 to 6.5 or 5.5 to 6.0, 5 to 8, preferably 6 to 7), for or before iron absorption.

- 5 The following non-limiting examples illustrate the invention and do not limit its scope in any way. In the examples and throughout this specification, all percentages, parts and ratios are by weight unless indicated otherwise.

EXAMPLES:Example 1

5

Patient 1

The patient had been taking Gaviscon™ which contains potassium bicarbonate as an acid reducer to reduce acid reflux. The patient had a gall bladder rupture requiring surgery and became very anaemic due to blood loss. He continued to take Gaviscon™ but was also given ferrous sulphate tablets for his anaemia. He had serious side effects of gastric intolerance and severe headaches and the ferrous sulphate failed to correct his anaemia. He commenced treatment using maltol and ferrous gluconate, whilst continuing Gaviscon™ and his anaemia was corrected (haemoglobin rose from 8.8g% to 11.8g%) after a month. At a dose of 45mg/day his iron was well tolerated.

Example 2

20 Patient 2

On a long term basis this patient takes Losec™, omeprazole, a proton pump inhibitor of gastric acid secretion. He had a hip replacement operation where a significant amount of blood was lost causing iron deficiency anaemia. His anaemia was not corrected (Hb > 11.0g%) until a course of maltol and ferrous gluconate at 15mg daily was undertaken. He continues to use maltol and ferrous gluconate as a "boost" when feeling tired as his haemoglobin levels are always on the low side of normal.

30 Example 3

The following is an example of a pharmaceutical composition according to the invention, which is suitable for formulation into gelatin capsules:

35

20

<u>Component</u>	<u>Amount (mg)</u>
Ferrous gluconate	240
Maltol	200
5 Cobalt gluconate	100
Carrier (lactose)	balance to fill capsule

Example 4

10 Method

Thirteen normal female volunteers were treated such that their gastric pH was adjusted to > 6.5 with 0.5M sodium bicarbonate solution (acid reducing regimen). Using a Heidelberg capsule the maltol iron formulation was given orally and the % of the dose absorbed into the blood and the whole body determined.

15

Results

Absorption of iron from the 10 mg dose of the maltol iron formulation was 10.72%. For the 20mg dose the maltol iron formulation gave 8.00%.

20 This is a level of absorption that would be expected with normal volunteers with a standard iron preparation, such as ferrous sulphate, at normal gastric pH.

Conclusions of studies

25 In the study involving 13 female subjects the maltol iron formulation was given along with sodium bicarbonate (acid reducing regimen) and the gastric pH was in excess of 6.5. Surprisingly, there was no inhibition of absorption at this higher pH. The presence of the maltol makes the formulation suitable for special medical conditions with gastritis and higher pH values associated with iron deficiency anaemia.

30

Discussion

The work shows that iron hydroxypyrones according to the invention are well absorbed at a high pH (greater than 6.5). Surprisingly, the iron hydroxypyrones can therefore deliver iron in a high pH environment, for example the presence of
35 an acid reducing regimen such as used for the treatment of Helicobacter

infection. Thus, there is an unexpected advantage to use an iron preparation containing a hydroxypyrrone such as maltol since a high pH, such as caused by the presence of the acid reducing regimen does not alter the solubility or bioavailability of the iron, which would have been expected. Other work has

5 shown that cobalt at low doses can have beneficial activity in the treatment of Helicobacter infection. In the light of these results a co-formulation of cobalt with iron products containing maltol would be particularly useful for treating the anaemia arising with the Helicobacter infection.

CLAIMS:

1. A composition comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrrone, for increasing the level of iron in a patient's bloodstream.
5
2. A composition comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrrone, for preventing and/or treating anaemia, such as iron deficiency anaemia.
10
3. The composition according to Claim 1 or Claim 2, wherein the inflammatory disease of the gastrointestinal tract requires a stomach acid reducing treatment regimen and/or wherein the one or more compounds or regimen optionally provides a gastric pH of equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.
15
4. The composition according to any one of Claims 1 to 3, wherein the composition is for administration to a human or animal, preferably having an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen.
20
5. The composition according to any one of Claims 1 to 4, wherein the composition is for administration to a human or animal, preferably having a gastrointestinal infection.
25
6. The composition according to Claim 5 wherein the gastrointestinal infection is caused by *Helicobacter pylori*.
- 30 7. The composition according to any one of Claims 1 to 6 wherein the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen comprise an acid reducing agent.

8. The composition according to Claim 7, wherein the acid reducing agent is selected from a proton pump inhibitor, an H₂ receptor antagonist, a substance capable of neutralising acids, or a combination thereof.

5 9. The composition according to Claim 7, wherein the acid reducing agent is selected from omeprazole, lansoprazole, pantoprazole, ranitidine, famotidine, nizatidine, cimetidine, aluminium hydroxide, magnesium carbonate, sodium or potassium bicarbonate, or calcium carbonate, or a combination thereof.

10 10. The composition according to any one of Claims 1 to 9, wherein the one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen comprises a cobalt compound.

15 11. The composition according to Claim 10, wherein the cobalt compound comprises a cobalt salt or a cobalt complex.

12. The composition according to Claim 10, wherein the cobalt compound comprises cobalt (II) gluconate or cobalt (II) chloride, or a combination thereof.

20

13. The composition according to any one of Claims 1 to 12, wherein the iron hydroxypyrrone comprises ferric trimaltol.

14. The composition according to any one of Claims 1 to 12, wherein the iron hydroxypyrrone comprises a mixture of a ferrous or a ferric salt and a hydroxypyrrone, preferably in the solid form.

25

15. The composition according to Claim 14, wherein the ferrous salt is an iron (II) carboxylate.

30

16. The composition according to Claim 15, wherein the iron (II) carboxylate is selected from iron (II) gluconate, iron (II) succinate or iron (II) fumarate.

17. A composition or kit of parts for increasing the level of iron in a patient's bloodstream comprising: one or more compounds capable of treating and/or

35

preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrrone, wherein the iron hydroxypyrrone comprises iron in the ferrous state.

5 18. The composition according to Claim 17, wherein the compound is a cobalt salt comprising cobalt gluconate.

10 19. A composition or kit of parts for increasing the level of iron in a patient's bloodstream comprising: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrrone with the proviso that when iron is in the ferric state, the compound is not a cobalt salt.

15 20. The composition according to Claim 18, wherein the compound is a substance capable of neutralising an acid.

20 21. The composition according to any one of Claims 17 to 20, wherein the inflammatory disease of the gastrointestinal tract requires a stomach acid reducing treatment regimen.

22. The composition according to any one of Claims 1 to 21, which is a pharmaceutical composition comprising a pharmaceutically acceptable diluent or carrier and wherein the composition is adapted for oral administration.

25 23. A composition comprising an iron hydroxypyrrone, for administration to a subject having, or at risk of having, achlorhydria.

30 24. A composition comprising an iron hydroxypyrrone for administration to a subject, wherein the gastric pH of the subject is equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.

35 25. A composition comprising an iron hydroxypyrrone for administration to a subject, wherein the subject has an inflammatory disease of the gastrointestinal tract and preferably the gastric pH of the subject is equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.

26. The composition according to Claim 24 or Claim 25 wherein the subject has or is at risk of having achlorhydria.
- 5 27. The composition according to any one of Claims 23 to 26 wherein the subject has been, will be, or is being administered an acid reducing regimen or one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract.
- 10 28. The composition according to any one of Claims 23 to 26, further comprising one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract or an acid reducing regimen.
- 15 29. The composition of Claim 27 or Claim 28, wherein the inflammatory disease of the gastrointestinal tract requires a stomach acid reducing treatment regimen.
- 20 30. The composition according to any one of Claims 23 to 29, wherein the composition is for increasing the level of iron in a patient's bloodstream and/or for preventing and/or treating anaemia such as iron deficiency anaemia.
- 25 31. The composition according to Claim 23 or Claim 26, wherein the achlorhydria is associated with an inflammatory disease of the gastrointestinal tract.
- 30 32. The composition according to Claim 23 or Claim 26, wherein the achlorhydria is associated with atrophic gastritis or a gastrointestinal infection, such as caused by *Helicobacter pylori*.
- 35 33. The composition according to any one of Claims 23 to 32 as defined in any one of Claims 3 to 22.
34. The composition according to any one of Claims 1 to 33, wherein the iron hydroxypyrrone is administered at the same time, before or after one or more compounds capable of treating and/or preventing an inflammatory disease of the

gastrointestinal tract, or the acid reducing regimen, preferably the iron hydroxypyrrone is administered at the same time or subsequent to the one or more compounds or the acid reducing regimen.

5 35. The composition according to any one of Claims 1 to 34, wherein the composition is administered by oral administration.

36. The composition according to any one of Claims 1 to 35, wherein the composition is for the absorption of iron.

10

37. A method for increasing the level of iron in a patient's bloodstream and/or for preventing and/or treating anaemia, such as iron-deficiency anaemia, which comprises the step of administering to a subject: one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract; and an iron hydroxypyrrone.

15

38. Use of one or more compounds capable of treating and/or preventing an inflammatory disease of the gastrointestinal tract which preferably requires a stomach acid reducing treatment regimen, and an iron hydroxypyrrone, in the manufacture of a medicament for increasing the level of iron in a patient's bloodstream and/or for preventing and/or treating anaemia such as iron deficiency anaemia.

20

39. Use of an iron hydroxypyrrone in the manufacture of a medicament for administration to a subject having, or at risk of having, achlorhydria.

25

40. Use of an iron hydroxypyrrone in the manufacture of a medicament for administration to a subject, wherein the gastric pH of the subject is equal to or greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.

30

41. Use of an iron hydroxypyrrone in the manufacture of a medicament for administration to a subject, wherein the subject has an inflammatory disease of the gastrointestinal tract and preferably the gastric pH of the subject is equal to or

greater than about 4, such as greater than 5 or 6, for example about 6.5 or above.