

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : G01N 33/53, A01N 37/18, A61K 38/00, C07K 1/00, 14/00, 17/00		A1	(11) International Publication Number: WO 96/18103
			(43) International Publication Date: 13 June 1996 (13.06.96)
(21) International Application Number: PCT/US95/15991			(81) Designated States: AU, BR, CA, CN, FI, JP, MX, SG, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).
(22) International Filing Date: 11 December 1995 (11.12.95)			
(30) Priority Data: 353,658 9 December 1994 (09.12.94) US 440,725 8 May 1995 (08.05.95) US			
(71) Applicants: THE SCRIPPS RESEARCH INSTITUTE [US/US]; Mail Drop TPC-8, 10666 North Torrey Pines Road, La Jolla, CA 92037 (US). NEUROSCIENCES RESEARCH FOUNDATION [US/US]; 10640 John Jay Hopkins Drive, San Diego, CA 92121 (US).			
(72) Inventors: EDELMAN, Gerald, M.; 7428 High Avenue, La Jolla, CA 92037 (US). CROSSIN, Kathryn, L.; 12533 Maestro Court, San Diego, CA 92130 (US). SPORNS, Olaf; 1453 Wilbur Avenue, San Diego, CA 92109 (US). KRUSHEL, Leslie; 4473 Donald, San Diego, CA 92117 (US).			
(74) Agents: CEPURITIS, Talivaldis et al.; Olson & Hierl, Ltd., Suite 3000, 20 North Wacker Drive, Chicago, IL 60606 (US).			Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: INHIBITION OF GLIAL CELL PROLIFERATION			
(57) Abstract Methods and compositions are disclosed that are useful for the inhibition of glial cell proliferation by means of binding inhibiting agents to neural cell adhesion (N-CAM) molecules present on the cell surface. Such inhibiting agents can be peptides, monoclonal antibodies, polyclonal antibodies, Fab' fragments, and the like.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

INHIBITION OF GLIAL CELL PROLIFERATION

Specification

Cross-Reference to Related Application

5 This application is a continuation-in-part of co-pending U.S. patent application Serial No. 08/353,658, filed December 9, 1994.

Governmental Rights

10 This invention was made with governmental support from the United States Government, National Institutes of Health, Grant HD09635; the United States Government has certain rights in the invention.

Field of Invention:

15 The invention relates to methods and compositions suitable for inhibiting glial cell proliferation. The invention relates to the use of peptides, antibodies and oligonucleotides for inhibiting glial cell proliferation. More particularly, the invention relates to peptides comprising at least part of the amino acid sequence of
20 neural cell adhesion molecule (N-CAM) and to compositions that utilize these peptides to inhibit DNA synthesis and cell division of glial cells.

Background:

25 During embryogenesis, the proliferation of glioblasts and their differentiation into several glial cell types appear to be closely coordinated with the development of neurons. During early postnatal development of the brain, glial cell proliferation continues at
30 significant levels after neuronal proliferation has mostly ceased. In general, glial cell proliferation in the brain remains strongly inhibited throughout adult life, except in tumorigenesis and after injury.

35 The control of glial cell proliferation is an important factor in the regulation of neural development,

- 2 -

regeneration, and tumorigenesis. The experiments described herein provide evidence for a new functional role of N-CAM in the control of glial cell proliferation in vitro and in vivo.

5 After physical injury to the central nervous system (CNS), astrocytes respond by changes in cell morphology, increased expression of glial fibrillary acidic protein (GFAP), hypertrophy, and rapid proliferation (Ludwin, S. K. (1984) *Nature* **308**, 274-275). Such reactive
10 astrocytes form glial scars in a process of gliosis that appears to interfere with neuronal regeneration in the adult CNS (Barrett, C. P. et al. (1984) *Exp. Neurol.* **84**, 374-385; Reier, P. J. & Houle, J. D. (1988) *In Advances in Neurology*, (Waxman, S. G., ed.). Raven Press, New York, pp.
15 87-138).

 Glial cell proliferation can be influenced by a variety of factors. Stimulatory effects are exerted by various growth factors (Kniss, D. A. & Burry, R. W. (1988) *Brain Res.* **439**, 281-288), interleukin-6 (IL-6) (Selmaj, K.
20 W. et al. (1990) *J. Immunol.* **144**, 129-135), interleukin-1 β (IL-1 β) (Giulian, D. & Lachman, L. B. (1985) *Science* **228**, 497-499), tumor necrosis factor (Barna, B. P. et al. (1990) *J. Neuroimmunol.* **30**, 239-243) and γ -interferon (Yong, V. et al. (1991) *Proc. Natl. Acad. Sci. USA* **88**, 7016-7020). The
25 mitogenic effects of several of these factors can be inhibited by transforming growth factor- β (Hunter, K. E., Sporn, M. B. & Davies, A. M. (1993) *Glia* **7**, 203-211) which is produced by neurons (Flanders, K. C. et al. (1991) *Development* **113**, 183-191).

30 Further evidence points to a direct involvement of neurons in the control of glial cell proliferation. The addition of purified cerebellar granule neurons to cerebellar astrocytes (Hatten, M. E. (1987) *J. Cell Biol.* **104**, 1353-1360) or of hippocampal neurons to glia (Gasser,
35 U. E. & Hatten, M. E. (1990) *J. Neurosci.* **10**, 1276-1285)

- 3 -

resulted in a decrease in glial cell proliferation. This inhibition depends on a proper ratio of neurons to glia of about 4 to 1, ensuring that astrocytes make surface contact with the added neurons. A cell membrane preparation of cerebellar granule neurons also inhibited proliferation, while medium conditioned by such neurons did not, indicating that membrane-associated molecules might be involved in the neuronal control of glial cell proliferation.

10 N-CAM (Edelman, G. M. (1986) *Ann. Rev. Cell Biol.* 2, 81-116; Cunningham, B. A. et al. (1987) *Science* 236, 799-806) has an important regulatory role in the developing nervous system and is present in adult nervous tissue. While the traditional view centers in the role of N-CAM in providing adhesion to the substrate and other cells, there is evidence that N-CAM can mediate signalling across the cell membrane in N-CAM mediated neurite outgrowth (Doherty, P. & Walsh, F. S. (1994) *Current Opinion in Neurobiology* 4, 49-55).

20 Astrocytes express a variety of adhesion molecules, among them the cell adhesion molecules N-CAM, and L1, and the extracellular matrix protein laminin. N-CAM has been found in astrocytes in vivo (Bartsch, U. et al. (1989) *J. Comp. Neurol.* 284, 451-462) as well as in primary astrocyte cultures (Noble, M. et al. (1985) *Nature (London)* 316, 725-728) although normal levels of N-CAM in glia are low in comparison to those in neurons (Nybroe, O. et al. (1985) *J. Cell. Biol.* 101, 2310-2315). The amount of N-CAM expressed on the surface of astrocytes (Smith, G. M. et al. (1993) *J. Neurochem.* 60, 1453-1466) as well as in the nervous system as a whole (Chuong, C.-M. & Edelman, G. M. (1984) *J. Neurosci.* 4, 2354-2368; Linnemann, D. et al. (1993) *Int. J. Devl. Neurosci.* 11, 71-81) decreases significantly as the CNS matures as compared to the amounts seen in prenatal and early postnatal development.

35

- 4 -

Levels of N-CAM have been shown to change after injury to nervous tissue. For example, after neurotoxin-induced brain damage, glial cells re-express high levels of the highly sialylated embryonic form of N-CAM (Le Gal La Salle, G. et al. (1992) *J. Neurosci.* **12**, 872-882). Peripheral nerve injury leads to the expression of high levels of sialic-acid rich N-CAM in both neural and glial tissue (Daniloff, J. K. et al. (1986) *J. Cell Biol.* **103**, 929-945). The increased expression of N-CAM after injury and the presence of N-CAM on reactive astrocytes indicates that cell adhesion molecules may play a functional role in regeneration and potential healing after such insults.

Renewed glial cell proliferation and gliosis are important in the response of neural tissue to injury. A reduction in the amount of gliosis after CNS injury may be a significant factor in permitting axonal regeneration across a lesion site. Neuronal regeneration in the adult CNS can occur if regenerating axons are presented with an appropriate environment (Aguayo, A. J. (1985) *In Synaptic plasticity*, (Cotman, C. W., ed.). Guilford, New York, pp. 457-484). Injury to the CNS of neonatal rats results in less extensive gliosis as compared to the adult (Barrett et al., 1984) possibly facilitating neuronal regeneration. Moreover, gliosis in the adult can be reduced by transplantation of immature astrocytes into the lesion site (Smith, D. B. & Johnson, K. S. (1988) *Gene* **67**, 31-40). Our experiments raise the possibility that this might be due to the high levels of N-CAM expressed by immature astrocytes. Further elucidation of these responses and definition of the signalling pathways that mediate the inhibition of glial cell proliferation facilitate the design of rational therapies to enhance regeneration by CNS neurons.

- 5 -

Summary of the Invention:

The present invention provides a method as well as a composition suitable for inhibiting glial cell proliferation. To this end, a mixed population of neural cells and glial cells is exposed to peptides that interact with N-CAM. The present invention also provides a method for identifying populations of glial cells susceptible to proliferation inhibition as well as a method for enriching a mixed population of glial and neural cells in glial cells susceptible to proliferation inhibition. The present invention contemplates the inhibition of the proliferation of cells: any cell having N-CAM on the cell surface is suitable for the practice of the invention.

The present invention provides, in general, a method of inhibiting proliferation of cells having neural cell adhesion molecule (N-CAM) on the cell surface. Proliferation of such cells can be inhibited by contacting the cells with an proliferation inhibiting amount of a N-CAM homophilic peptide having at least 7 amino acid residues, preferably 7 to 90 amino acid residues. Such N-CAM homophilic peptides include a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues.

Brief Description of the Drawings:

FIGURE 1: Inhibition of proliferation as a function of the amount of N-CAM present in the medium for different culture conditions: (A) SFM (n = 10), (B) SFM plus 1% FBS (n = 5), (C) SFM plus 20 ng/ml bFGF (n = 5). The number in the upper left hand corner gives the level of [³H]-thymidine incorporation (percent of SFM control) with no N-CAM added. Throughout the paper, [³H]-thymidine incorporation is

- 6 -

normalized relative to [³⁵S]-methionine cell labeling, calculated as the ratio [³H]dpm/[³⁵S]dpm and expressed as a percentage (mean ± S.E.M.) of the respective control (no N-CAM added) for each of the three conditions. Cell
5 density is 7 x 10⁴ cells/cm² throughout.

FIGURE 2: Dependence of baseline proliferation levels (A) and inhibition of proliferation by 2 µg/ml N-CAM (B) on cell density. In (A), results are expressed as percent
10 [³H]-thymidine incorporation relative to a density of 7 x 10⁴ cells/cm² (the density used in all other experiments). (B) shows the [³H]-thymidine incorporation expressed as percent of the baseline (control) proliferation level (see A) for the corresponding cell density. n = 5 throughout.

15

FIGURE 3: Inhibitory effect of synthetic peptides with sequences containing a putative N-CAM homophilic binding site. Peptides are added at 0.1 mg/ml to SFM, with astrocytes at a density of 7 x 10⁴ cells/cm². From left to
20 right: chick N-CAM peptide and corresponding random peptide, rat N-CAM peptide and corresponding random peptide.

FIGURE 4: Effect of N-CAM and random antisense
25 oligonucleotides on the inhibition of glial cell proliferation by N-CAM. Left side (no N-CAM added): addition of 200 nM N-CAM antisense or random antisense oligonucleotides alone has no effect on proliferation. Right side (N-CAM added): SFM (Control) plus 2 µg/ml N-CAM
30 produces inhibition, which is partially reversed by addition (48 hours prior to N-CAM) of 200 nM N-CAM antisense, but not by 200 nM random antisense oligonucleotides. * P < 0.02 vs. Control + N-CAM (paired sample Student's t-test).

- 7 -

FIGURE 5: The labeling index of dividing astrocytes infused with rat N-CAM, the third Ig domain of chicken N-CAM, or the peptide derived from the third Ig domain of rat N-CAM in the cerebral cortex, hippocampus, or striatum 3 days after the lesion (n = 3 for each experimental condition; data are presented as mean $\pm$ S.E.M.). The labeling index is defined as the percentage of labeled astrocytes (S100 or GFAP positive) that were also BrdU positive. For each brain region, the labeling index for the control condition (infusion of the random peptide) was set to 100%. In all cases, there were significantly fewer (p<0.05; paired sample Student's t-test) dividing astrocytes after infusion of N-CAM or related molecules as compared to the control condition.

FIGURE 6: Inhibition of glial cell proliferation by N-CAM peptides injected in vivo. Rats were lesioned with the needle of a Hamilton syringe and concomitantly injected with a N-CAM peptide into either the cerebral cortex (A), the hippocampus (B), or the striatum (C). The control peptide was injected into the corresponding site on the contralateral side of the brain. The rats then received five bromodeoxyuridine injections (intraperitoneal) over the following 72 hours. The percentage difference in labeling index is a reduction in astrocytes labeled with bromodeoxyuridine at the N-CAM peptide injection site compared to the contralateral control. Mean values $\pm$ S.E.M for three animals.

FIGURE 7: The labeling index of dividing astrocytes 3 days after a cortical lesion infused with N-CAM, the control random peptide, bFGF plus N-CAM, or the random peptide plus bFGF (N=3 for each condition).

FIGURE 8: An autoradiogram of an immunoprecipitation experiment showing that at least one protein in the molecular weight range of 60-70 kDa and one protein at

- 8 -

approximately 120 kDa showed diminished tyrosine phosphorylation after N-CAM treatment.

Detailed Description

5 Abbreviations:

	bFGF	basic fibroblast growth factor
	FBS	fetal bovine serum
	FITC	fluorescein isothiocyanate
	GFAP	glial fibrillary acidic protein
10	N-CAM	neural cell adhesion molecule
	PBS	phosphate buffered saline
	SFM	serum free medium

Cell proliferation, comprising growth and
15 reproduction of cells, is a key primary process not only during neural development but also in the regenerative response of neural tissue to injury. Proliferation of the glial cells of neural tissue has been reported to be controlled, at least in part, by a neuronal signal that may
20 involve cell surface molecules.

N-CAM, as well as peptides comprising a portion of the amino acid sequence of N-CAM and including the homophilic binding region (herein termed N-CAM homophilic peptides), are capable of inhibiting glial cell
25 proliferation. The addition of soluble N-CAM portions to primary cultures of astrocytes inhibits their proliferation. This inhibitory effect is believed to be mediated by homophilic N-CAM binding. This inhibitory effect can be elicited by addition of short synthetic
30 peptides with sequences corresponding to parts of the third immunoglobulin-like domain of N-CAM.

It has recently been found that N-CAM inhibits glial cell proliferation of both astrocytes *in vitro* and brain glial cells *in vivo*. This inhibitory effect has been
35 elicited in primary cultures of rat forebrain astrocytes

- 9 -

grown in chemically defined serum free medium or in a medium that had been supplemented with growth factors. The present invention provides methods of using N-CAM and related molecules to control glial cell proliferation. The present invention is suitable for controlling the formation of glial scars after central nervous system injury, which may in turn aid neuronal regeneration. The inhibitory effect on proliferation of short synthetic peptides with sequences located in the third immunoglobulin-like domain of N-CAM lends additional support to the hypothesis that N-CAM homophilic binding can lead to decreased proliferation. This result suggests that binding to this region of N-CAM is sufficient to elicit a signal leading to an intracellular response. Although any influence of such peptides on cell proliferation has not been reported previously, it has been shown that the decapeptide sequence KYSFNYDGSE is directly involved in chick N-CAM homophilic binding (Rao et al., 1992). In solution, this peptide disrupted N-CAM-homophilic binding and inhibited N-CAM-dependent neurite outgrowth from retinal cells (Sandig, M., et al. (1994) *J. Biol. Chem.* **269**, 14841-14848).

Such decapeptides inhibit glial cell proliferation and can mimic the effect of whole purified N-CAM. These peptides comprise a portion of the amino acid sequence of N-CAM and include the homophilic binding region as N-CAM homophilic peptides. Binding of such peptides to N-CAM is sufficient to elicit a signaling event across the cell membrane.

The magnitude of the inhibitory effect of N-CAM was dependent on cell density: it was maximal at low cell densities and weakened progressively as cells approached confluency. Synthetic N-CAM homophilic peptides mimicked the effect of purified N-CAM, while peptides of the same length and amino acid composition but with a randomized sequence did not. The addition of N-CAM antisense

- 10 -

oligonucleotides to astrocyte primary cultures for 48 hours resulted in reduced levels of N-CAM expression. After N-CAM levels on astrocytes were diminished by this treatment, the anti-proliferative effect of N-CAM added to
5 the medium was significantly reduced. These results indicate that N-CAM homophilic peptides may be used to control proliferation of glial cells.


The experimental results indicate that the proliferation of rat forebrain astrocytes in primary
10 culture can be strongly inhibited by addition of purified N-CAM to the culture medium. The proliferation of other cell types expressing N-CAM also seem to be susceptible to N-CAM mediated inhibition. It has recently been found that C6 glioma cells which express N-CAM on their cell surface
15 also decrease their rate of proliferation in response to externally added N-CAM when grown in SFM at low cell densities. An assay for the presence of N-CAM on the cell surface helps to predict the ability of N-CAM homophilic peptides to inhibit the proliferation of a cell type.

20 The effect on astrocytes is mimicked by polyclonal N-CAM antibodies and N-CAM Fab' fragments. These findings indicate that the inhibition of proliferation does not result from crosslinking of N-CAM at the cell surface, but is the result of a signal mediated by
25 homophilic binding to N-CAM at the glial surface. This conclusion is also supported by the finding that N-CAM peptides can have the same effect. The results of antisense experiments indicate that a threshold amount of N-CAM at the glial surface is required.


30 N-CAM homophilic peptides or N-CAM homophilic antibodies are useful agents for inhibiting the proliferation of glial cells in a mixed culture of glial and neural cells when the desired goal is culturing neural cells free of glial cell contamination.

- 11 -


A suitable N-CAM homophilic decapeptide contemplated by the present invention is described by the formula:



where Xaa^{+} is an amino acid residue with a positively charged side chain, Xaa^{j} is an amino acid residue with a nonpositive side chain, Xaa^{o} is an amino acid residue with a nonpolar side chain, Xaa^{u} is an amino acid residue with an uncharged side chain, Xaa^{x} is an amino acid residue with no specific constraints on the nature of the side chain, and Xaa^{-} is an amino acid residue with a negatively charged side chain. A suitable N-CAM homophilic heptapeptide is described by the formula:



A preferred N-CAM homophilic decapeptide contemplated by the present invention is described by the formula:



25 A preferred N-CAM homophilic heptapeptide contemplated by the present invention is described by the formula:



30 The term "antibody" in its various grammatical forms is used herein to refer to immunoglobulin molecules and immunologically active portions of immunoglobulin molecules, i.e., molecules that contain an antibody combining site or paratope. Exemplary antibody molecules are intact immunoglobulin molecules, substantially intact

35

- 12 -

immunoglobulin molecules and those portions of an immunoglobulin molecule that contain the paratope, including those portions known in the art as Fab, Fab', F(ab')₂ and F(v).

5 An antibody composition of the present invention is characterized as containing antibody molecules that immunoreact with N-CAM peptides or portions thereof.

 An antibody composition of the present invention is typically produced by immunizing a mammal with a
10 inoculum of the present invention and thereby induce in the mammal antibody molecules having the appropriate peptide immunospecificity. The antibody molecules are then collected from the mammal and isolated to the extent desired by well known techniques such as, for example,
15 immunoaffinity chromatography. The antibody composition so produced can be used, inter alia, in the diagnostic methods and systems of the present invention to detect cells expressing N-CAM in a sample of tissue.

 Monoclonal antibody compositions are also
20 contemplated by the present invention. A monoclonal antibody composition contains, within detectable limits, only one species of antibody combining site capable of effectively binding a N-CAM homophilic peptide. Thus, a monoclonal antibody composition of the present invention
25 typically displays a single binding affinity for a N-CAM homophilic peptide even though it may contain antibodies capable of binding proteins other than a N-CAM homophilic peptide.

 Vectors which comprise the isolated nucleic acid
30 molecule coding for a portion of N-CAM may be prepared by means well known to one skilled in the art. Suitable vectors comprise, but are not limited to, a plasmid or a virus. These vectors may be transformed into a suitable host cell to form a host cell vector system for the
35 production of a N-CAM homophilic peptide. This invention

- 13 -

further provides an isolated DNA or cDNA molecule described hereinabove wherein the host cell is selected from the group consisting of bacterial cells (such as E. coli), yeast cells, fungal cells, insect cells and animal cells. Suitable animal cells include, but are not limited to Vero cells, HeLa cells, Cos cells, CVI cells and various primary mammalian cells. Cos cells are transfected by the DEAE-Dextran method as follows: 80% confluent overnight cultures are transfected with 5 μ g DNA, per 100 mm dish, in 250 μ g/ml DEAE Dextran (Pharmacia), 100 mM Tris pH 7.3, in DMEM. After 6 h cells are washed and incubated in DMEM 10% calf serum, 0.1 mM chloroquine (Sigma) for 2.5 h, followed by incubation in DMEM 10% calf serum overnight. For isolation of peptide the medium is changed to OPTI-MEM (BRL), and the cells are incubated for 48 h.

Methods for facilitating the transport of drugs across the blood-brain barrier are known (see, for example Bickel, U. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90 2618-2622). Suitable brain transport vectors include proteins such as cationized albumin or the OX26 monoclonal antibody to the transferrin receptor.

- 14 -

EXAMPLE 1

Purified N-CAM inhibits astrocyte proliferation.

5 Addition of N-CAM purified from rat forebrain
to primary cultures of astrocytes (at a density of 7×10^4 cells/cm²) in chemically defined medium (SFM)
inhibited their proliferation in a concentration
dependent manner (FIGURE 1A).

10 Astrocyte Cultures. Primary cultures of
astrocytes were prepared from the forebrains of 4-5 day
old neonatal rats. After decapitation and removal of the
forebrains from the skull, the meninges were carefully
peeled off, and forebrain tissue was triturated in
15 sterile phosphate buffered saline (PBS) using
firepolished pipettes. The resulting suspension of
tissue fragments was incubated in Hank's balanced salt
solution (Gibco BRL) with 2.5 mg/ml trypsin, 1 mM EDTA
and 0.1 mg/ml DNase (Boehringer) at 37°C for 15 minutes.
20 Undissociated tissue was removed by centrifugation and
the remaining cell suspension was centrifuged through a
cushion of 3.5% bovine serum albumin to remove cell
debris. The pellet was resuspended in serum free medium
(SFM) consisting of Dulbecco's Modified Eagle Medium (D-
25 MEM), 2 mM L-glutamine, 50 U/ml penicillin G sodium and
50 µg/ml streptomycin sulfate (all from Gibco BRL)
supplemented by 10% (vol/vol) fetal bovine serum (FBS).
The cells were transferred to a standard 100 mm tissue
culture dish (Falcon) and incubated at 37°C for 1-2
30 hours. After mild shaking, the supernatant containing
unattached cells was transferred to a 100 mm collagen-
coated dish. Cultures were maintained in SFM
supplemented with 10% FBS in 5% CO₂/95% air for about 7
days before the start of an experiment. Cell samples
35 were checked regularly for the expression of GFAP, a

- 15 -

marker for astrocytes. Only those cultures containing more than 95% astrocytes were used.

Purification of N-CAM. N-CAM was purified from early postnatal rat brains by affinity chromatography using a cross-reactive monoclonal anti-human N-CAM antibody (a generous gift of Dr. John Hemperly, Becton Dickinson Research Center, Research Triangle Park, NC). Purity of the preparations was tested by SDS gel electrophoresis using standard Coomassie blue staining and Western blotting techniques. Protein concentrations were estimated using the Bio Rad Micro Bradford assay. In some experiments, purified N-CAM was treated with 1 U/ml neuraminidase (from *Vibrio cholerae*, Calbiochem) in 100 mM sodium acetate, 2 mM CaCl_2 , 2 mM EDTA, for 24 hours at 35°C and pH = 5.5 (Hoffman, S. et al. (1984) *Proc. Natl. Acad. Sci. USA* 81, 6881-6885) before being added to cell cultures. SDS gel electrophoresis revealed that the purified N-CAM consisted mainly of the polysialylated embryonic form of N-CAM (MW 200-250 kD) with other molecular forms including an approximately 65 kD hydrolyzed N-CAM fragment present in lower amounts.

Proliferation Assay. Astrocytes were labeled by addition of 10 μl 8.72 μM [^{35}S]-L-methionine (New England Nuclear, approximately 1200 Ci/mmol) to 10 ml culture medium and transferred into 96-well plates (Falcon) usually at a density of 2×10^5 cells/ml (equivalent to 7×10^4 cells/cm² or 2×10^4 cells/well). After cell attachment, the culture medium was exchanged for SFM. In antisense experiments, oligonucleotides were added at the same time (T = 0). Growth factors, purified N-CAM, or synthetic peptides were added after the cells had remained in SFM for 48 hours (T = 48); at this time most of the cells are synchronized with respect to their position in the cell cycle at G₁/G₀ (Kniss and Burry, 1988). After another 12 hours (at T = 60), [^3H]-

- 16 -

thymidine (New England Nuclear, 20 Ci/mmol) was added (0.38 μ M, 10 μ Ci/ml). Following an additional 12 hours (T = 72), the wells were drained, counts precipitated by ice-cold 10% trichloroacetic acid, and the plates left to dry. Counts were solubilized overnight by adding 100 μ l 1 N KOH to each well, neutralized with 100 μ l 1 N HCl, transferred to liquid scintillation vials and counted in 3 ml of Beckman Ready Safe liquid scintillation cocktail on a Packard 1600 TR counter. To determine levels of [3 H]-thymidine incorporation relative to the number of cells, a dual dpm counting protocol was used (Crossin, K. L. (1991) *Proc. Natl. Acad. Sci. USA* 88, 11403-11407), which minimizes the overlap between the 3 H and 35 S energy spectra and takes into account detection efficiencies for both isotopes. In this paper, proliferation is expressed as [3 H]-thymidine incorporation normalized to [35 S]-methionine incorporation and calculated as [3 H]dpm/[35 S]dpm.

As shown by testing three separate preparations of rat N-CAM, maximal (> 90%) suppression of proliferation could be effected at concentrations of 10 μ g/ml (FIGURE 1A). N-CAM in a concentration of about 0.2 μ g/ml to about 2 μ g/ml was sufficient for half-maximal (50%) suppression. Similar effects were observed after the addition of N-CAM purified from embryonic chick brain, which has been shown to bind to rodent N-CAM (Hoffman *et al.*, 1984) also inhibited proliferation. At an N-CAM concentration of 10 μ g/ml the cells tended to assume a somewhat rounded appearance; at concentrations $\leq$ 2 μ g/ml (a range that still exhibited substantial inhibitory activity) no gross changes in cell morphology could be observed.

The addition of 1% FBS (containing various growth factors) and 20 ng/ml basic fibroblast growth factor (bFGF, Upstate Biotechnology Incorporated) to

- 17 -

astrocyte cultures led to a marked increase in proliferation as compared to cultures in SFM ($285 \pm 24\%$ for FBS, $n = 14$, and $347 \pm 56\%$ for bFGF, $n = 12$). Addition of N-CAM together with either 1% FBS or 20 ng/ml bFGF elicited also a suppressive effect similar in magnitude to that observed in SFM (FIGURE 1B,C). Addition of 2 μ g/ml N-CAM treated with neuraminidase to remove sialic acid produced an inhibitory effect comparable to untreated N-CAM ($24 \pm 4\%$ [3 H]-thymidine incorporation vs. SFM control, $n = 2$).

EXAMPLE 2

Contact inhibition and dependence of N-CAM effect on cell density

Astrocytes growing in SFM showed different levels of proliferative activity depending on their cell density. Levels of [3 H]-thymidine incorporation per cell were high at low cell densities, while confluent or nearly confluent cultures proliferated more slowly (presumably due to the phenomenon of "contact inhibition", FIGURE 2A). When N-CAM was added to these cultures, the inhibitory effect of N-CAM on proliferation was critically dependent on cell density and thus on baseline proliferative activity. Inhibition was maximal at lower cell density (1.8×10^4 cells/cm 2) when cells are actively proliferating, while N-CAM was virtually ineffective at higher cell densities (28×10^4 cells/cm 2) when cultures are confluent.

It is unlikely that N-CAM added in solution acts by disrupting cell-cell adhesion itself given the observation that the effect is maximal at low cell density. The finding that N-CAM becomes progressively less effective in inhibiting proliferation as cell density rises may be because baseline proliferation levels of cells in confluent culture are already low and

- 18 -

therefore cannot be suppressed further. The quiescence of confluent cultures could also be due in part to an autocrine effect via secretion of an inhibitory substance into the culture medium (e.g. Nieto-Sampedro, M. & Broderick, J. T. (1989) *J. Neurosci. Res.* 22, 28-35).

EXAMPLE 3

Anti-proliferative activity of peptides

Several synthetic N-CAM homophilic peptides were added to astrocyte cultures.

10 Peptide Synthesis. Peptides were synthesized on a Gilson AMS-422 automated peptide synthesizer. The lyophilized peptides were resuspended in water, desalted and re-lyophilized at least once before use. Purity and concentrations of synthetic peptides were confirmed using
15 mass spectrometry, analytical high pressure liquid chromatography (HPLC) and amino acid analysis.

At a concentration of 0.1 mg/ml, the peptide KYSFNYDGSE, identical to an N-CAM homophilic binding region located in the third immunoglobulin-like domain of
20 chick N-CAM, inhibited proliferation, while a peptide of the same length and amino acid composition but with randomized sequence (GFNSSYEDYK) had no effect (FIGURE 3). The direct homologue to the chick peptide for rat N-CAM, KHIFSDDSSE, had a similar inhibitory effect, while
25 the corresponding random peptide SFSISDEDHK did not affect proliferation.

EXAMPLE 4

Partial reversal of the inhibition of proliferation
30 after adding antisense oligonucleotides

The amounts of N-CAM present in astrocytes were quantified by Western blotting. Cultures were grown as described and cellular material was lysed in buffer (phosphate buffered saline, pH 7.4, containing 0.5% NP-
35 40, 350 µg/ml PMSF, 2 µg/ml aprotinin, 0.5 µg/ml

- 19 -

leupeptin, and 0.7 μ g/ml pepstatin A) at 4°C. Standard Western blotting techniques were employed either using cell lysate directly or after N-CAM immunoprecipitation. [125I]-Immunolabeled N-CAM was estimated by densitometry on a Molecular Dynamics PhosphorImager using ImageQuant software.

Phosphorothioate oligoribonucleotides were synthesized on an Applied Biosystems Model 380B DNA synthesizer using phosphoramidite chemistry. Two sequences of length 24 were used: one sequence 5'-AGAUCUUAGUUCGAGCAUUGUA-3' was complementary to position 205 to 228 of rat N-CAM, starting 4 bases before the translation initiation codon (Small, S. J., Shull, G. E., Santoni, M. J. & Akesson, R. (1987) *J. Cell Biol.* 105, 2335-2345) ("N-CAM antisense"), another sequence 5'-GUCCAUGUACAUCUAGUCAGUUGA-3' consisted of the same nucleotides arranged in random order ("random antisense"). After deprotection, the oligonucleotides were purified using HPLC, ethanol precipitated, solubilized in distilled water and added to the culture medium at the indicated concentrations.

Table 1

Levels of N-CAM expression (sum of all molecular forms, mostly 140 kD and 120 kD, in % Control) in antisense experiments as determined by Western blotting. Data are expressed as mean $\pm$ S.E.M. (n) with n = number of experiments. * $P < 0.02$ vs. control (paired sample Student's t-test).

Condition	N-CAM level
Control	100 $\pm$ 0 (6)
+ N-CAM antisense RNA	65* $\pm$ 9 (5)
+ random antisense RNA	87 $\pm$ 18 (6)

The addition of N-CAM antisense oligonucleotides for 48 hours resulted in significantly reduced N-CAM expression ($P < 0.02$, Table 1). A random antisense oligonucleotide had, on average, no effect on N-CAM expression. Neither oligonucleotide led to changes

- 20 -

in the gross morphology of astrocytes or had an influence on the baseline level of proliferation (FIGURE 4, left). Addition of 2 μ g/ml N-CAM to astrocyte cultures that had been preincubated with 200 nM N-CAM antisense
5 oligonucleotides for 48 hours revealed a significantly ($P < 0.02$) weaker inhibitory effect of N-CAM on proliferation (FIGURE 4, right). Addition of N-CAM to astrocytes preincubated with the random antisense oligonucleotide revealed no such effect.

10 When the level of N-CAM expression by astrocytes is diminished by adding specific antisense oligonucleotides, the effect on proliferation elicited by N-CAM added to the medium is reduced. Because of the specificity of antisense probes, this finding suggests
15 that expression of N-CAM on the glial cell surface is required for the inhibitory effect of soluble N-CAM and supports the notion that homophilic N-CAM binding mediates the decrease of astrocyte proliferation. Furthermore, addition of antisense oligonucleotides also
20 reduces the effectiveness of the ongoing endogenous inhibition of glial proliferation caused by the neurons in the culture, having the net effect of enhancing the proliferation of glia and enriching the proportion of glial cells. The ability to obtain a culture enriched in
25 glial cells is useful for experimental and testing purposes.

 The antisense oligonucleotides were added to the culture medium at relatively low concentrations in order to reduce possible non-specific effects. For
30 example, we found that oligonucleotide concentrations in the micromolar range reduced baseline proliferation levels irrespective of their sequence. Previous studies using antisense probes against bFGF in astrocyte cultures (Gerdes, W. et al. (1992) *NeuroReport* 3, 43-46) have
35 shown that phosphorothioate oligonucleotides can be

- 21 -

effective within a few hours. In the present experiments, however, long incubation periods with N-CAM antisense oligonucleotides were required before significant reductions in N-CAM levels could be observed (see Perides, G., *et al.* (1992) *Proc. Natl. Acad. Sci. USA* **89**, 10326-10330); this may be due to slow N-CAM turnover rates (Friedlander, D. R., *et al.* (1985) *J. Cell Biol.* **101**, 412-419). Even after 48 hours, only a gradual reduction in N-CAM levels was achieved. This finding would account for the significant but only partial reversal of the N-CAM effect on proliferation after N-CAM antisense oligonucleotide treatment.

EXAMPLE 5

In Vivo Effects on Glial Cell Proliferation

Experiments have shown the effectiveness in vivo of N-CAM and shorter peptides in reducing glial proliferation induced by lesions and brain trauma. Lesions were made by a hypodermic needle placed into the brain and N-CAM peptides or shorter fragments thereof were applied through the needle to the affected area.

Purification of Proteins and Peptides.

The purification of N-CAM from early postnatal rat brains was done as described in Example 1 and the synthesis of peptides was done as described in Example 3. The recombinant protein Ig3, corresponding to the chicken N-CAM third Ig domain, was affinity purified on a column of monoclonal antibody CAM 1-Sepharose (Hoffman, S. *et al.*, (1982) *J. Biol. Chem.* **257**, 7720-7729. N-CAMs from different species bind with each other and the chicken clone was readily available. Hoffman, S. & Edelman, G. M. (1984) The mechanism of binding of neural cell adhesion molecules. In (Lauder, J. M. & Nelson, P., eds.). Plenum Press, New York, pp. 147-160.

- 22 -

Construction of N-CAM Expression Vectors

A peptide comprising the third Ig domain of chick N-CAM was produced recombinantly using an expression vector. The peptide, comprising Val₆₇ through Lys₁₈₀ of chick N-CAM, was produced by an expression vector constructed from a segment including nucleotides 259 to 540 of clone pEC 208 (Hemperly, et al. (1986) *Proc. Natl. Acad. Sci. USA* 83, 3037-3041). Clone pEC 208 was generated by PCR amplification, its DNA sequence was confirmed and it was inserted into the Nco I/Bam I sites of pET 3d (Novagen). PCR primers were chosen to amplify the region coding for the third Ig domain and add a Nco I 5' and a BamH I site 3' to the coding region. The sequence coding for the peptide is given by SEQ ID NO:4, the 3' BamH I primer by SEQ ID NO:5, and the 5' Nco I primer by SEQ ID NO:6. The PCR conditions were 60 mM Tris HCl, 15 mM NH₄SO₄, 2 mM MgCl₂, pH 9.5 (PCR Optimizer Buffer G, Invitrogen), 1 minute cycles at 95, 60 and 72 °C.

The PCR product (SEQ ID NO:7) was digested with BamH I and Nco I, and ligated into a vector (pET-3d) that had been digested separately with BamH I and Nco I to form the expression vector. There are several suitable prokaryotic expression vectors, of these pET-3d is preferred. Suitable eukaryotic expression vectors include pNUT (Searle, P.F. et al. (1985) *Molec. & Cell. Biol.* 5, 1480-1489) and pcDNA1 neo (Invitrogen).

E. coli cells were transfected with the expression vector and cultured. Protein was extracted from the cultured cells, and the recombinant peptide was isolated by affinity purification using a CAM 6 (Hoffman, S. et al. (1982) *J. Biol. Chem.* 257, 7720-7729) monoclonal antibody column.

- 23 -

Stereotactic placement of Lesions

Samples were injected into the brains of rats (Wistar Albino, Male, 200-250 g, Charles River, anesthetized with xylazine (12 mg/kg) and ketamine (90 mg/kg) through a 26 gauge needle. The needle point was placed at the stereotactic coordinates of either the hippocampus, cerebral cortex or striatum (Table 2). Bilateral injections of 3-4 μ l were made using an Hamilton syringe. Each animal received an infusion of either the purified rat N-CAM (250 μ g/ml), the third Ig domain of chicken N-CAM (5 mg/ml), or the rat N-CAM peptide KHIFSDDSSSE (1 mg/ml) in PBS into one hemisphere; the other hemisphere was infused with the random peptide (SFSISDEDHK, 1 mg/ml in PBS) as a control. The experimental and control sides varied between animals and choice of side had no significant effect on the results. Three μ l of each protein or peptide was infused at a rate of 0.2 μ l per minute after which the needle was left in place for 5 minutes and then removed. In one experiment, a group of animals received basic fibroblast growth factor (bFGF, 450 ng) in the solutions which contained either the random peptide or the purified N-CAM.

The samples were either purified N-CAM in phosphate buffered saline (PBS), N-CAM peptide KHIFSDDSSSE (1mg/ml or 10mg/ml) in PBS, or the corresponding random peptide SFSISDEDHK (1mg/ml) (see Example 3 above). The injection rate was about 0.2 μ l/min; the needle was left in place for 5 minute after the injection. Each animal received either the purified N-CAM or the N-CAM peptide injected into one side of the brain and the corresponding structure on the other side was injected with the random peptide as a control.

Table 2

Stereotactic Coordinates of Lesions
(using bregma as a zero point)

	Anterior- Posterior (AP)	Lateral (L)	Ventral (V, from dura)	
5				
	Hippocampus:	-3.30 mm	+ 1.60 mm	-3.20 mm
	Cerebral Cortex:	÷ 0.70 mm	+ 2.70 mm	-1.60 mm
	(The needle entered the cerebral cortex at a 20 degree angle)			
10	Striatum:	÷ 0.70 mm	+ 2.60 mm	-4.70 mm

Three animals were examined for each treatment and brain region described. Animals were maintained for a postinjection survival time of about 70 - 72 hours. During that time they received five injections of bromodeoxyuridine (Brdu) dissolved in 0.007 N NaOH (120 mg/kg, intraperitoneal). Two of the three animals that had been injected in the hippocampus received Brdu approximately 18, 28, 42, 52, and 70 hours after the stereotactic surgery. All of the animals that had been injected in the cerebral cortex or the striatum and one animal injected in the hippocampus received Brdu approximately 2, 18, 28, 42, and 52 hours after the stereotactic surgery.

Animals were perfused under anesthesia with about 30 ml of ice-cold PBS followed by about 400ml of ice-cold 4% paraformaldehyde in PBS. The brains were then removed and postfixed for 2 - 16 hours in 4% paraformaldehyde in PBS then placed in succeeding solutions of ascending concentrations of sucrose (12%, 18%, and 24%). The brains were then sectioned on a cryostat at a thickness of 10 μ m.

- 25 -

ImmunocytochemistryHippocampal tissue.

Non-specific binding on the sections was blocked by incubation in 8% fetal bovine serum (FBS) in PBS. Sections were then incubated in polyclonal antibody to glial fibrillary acidic protein (GFAP; 1:100 dilution, Dako) in 0.2% Triton X-100, 8% FBS/PBS for 24 - 48 hr at 4°C. The sections were processed using a biotinylated secondary antibody and the Vectastain ABC kit (Vector), and stained with a diaminobenzidine peroxidase substrate kit from Vector or Pierce. The sections were then incubated for 6 minutes in a solution of Proteinase K (Dako), rinsed and incubated in 2 N HCL for 20 minutes at 37°C. The sections were rinsed and blocked in 8% FBS/PBS and incubated in a monoclonal antibody to Brdu (1:40 dilution, Dako), 0.01% Triton X-100, 8% fetal bovine serum/PBS for 24-48 hr at 4°C. The sections were processed using a biotinylated secondary antibody and the Vectastain ABC kit (Vector), and stained with a VIP peroxidase substrate kit from Vector giving the Brdu positive cells a red color. The sections were then rinsed in PBS and coverslipped in glycerol/PBS.

GFAP positive cells and cells positive for both GFAP and BRDU were counted under a Zeiss Axiophot light microscope. Three sections which contained the center of the lesion were examined for each animal. An area on either side of the lesion having a radius of about 600 μ m was examined.

Cerebral cortical and Striatal tissue.

The sections were incubated for 3 minutes in Proteinase K (Dako), rinsed and then incubated in 2 N HCL for 20 min at 37°C. The sections were rinsed and blocked in 8% FBS/PBS and incubated in a monoclonal antibody to Brdu (1:40 dilution, Dako), a polyclonal antibody to S100 (1:100 dilution, Dako or Sigma) in 0.3% Triton X-100 and 8%

- 26 -

FBS/PBS for 3 - 4 hours at room temperature. A solution containing a FITC conjugated anti-mouse secondary antibody and a Texas red conjugated anti-rabbit secondary antibody (Vector) were then applied for 1 hour. The
 5 tissue was rinsed in PBS and coverslipped with SlowFade (Molecular Probes).

S100 immunoreactive cells and cells immunoreactive for both S100 and Brdu were counted under a Zeiss Axiophot fluorescent microscope using a filter
 10 whose bandwidth contained the excitation and emission wavelengths for FITC and Texas Red (Omega Optical). Three sections which contained the center of the lesion per animal were examined. Approximately 300 μ m on either side of the lesion was examined.

15 Table 3. Labeling index after different treatments following a stab injury.

Brain Region	Infusion (Experimental)	Labeling Index (Control)	Labeling Index (Experimental)	Percent Difference
Cerebral Cortex	N-CAM	21.3 $\pm$ 1.2	10.2 $\pm$ 1.5	-52.1 $\pm$ 4.5
	3rd Ig Domain	24.9 $\pm$ 2.9	13.9 $\pm$ 1.9	-44.3 $\pm$ 9.4
	N-CAM Peptide	18.9 $\pm$ 4.8	13.2 $\pm$ 3.8	-30.4 $\pm$ 7.8
Hippocampus	N-CAM	19.6 $\pm$ 3.3	14.6 $\pm$ 2.3	-25.2 $\pm$ 1.6
	3rd Ig Domain	18.4 $\pm$ 2.6	15.6 $\pm$ 2.2	-15.0 $\pm$ 3.1
	N-CAM Peptide	19.9 $\pm$ 2.1	14.5 $\pm$ 2.0	-27.1 $\pm$ 2.9
Striatum	N-CAM	26.1 $\pm$ 0.7	20.7 $\pm$ 0.7	-20.8 $\pm$ 2.1
	3rd Ig Domain	24.3 $\pm$ 5.2	19.6 $\pm$ 3.9	-19.5 $\pm$ 5.1
	N-CAM Peptide	19.1 $\pm$ 0.9	16.3 $\pm$ 0.9	-14.5 $\pm$ 2.3

20 The data are presented as mean $\pm$ S.E.M., n = 3 for each condition. All comparisons of the labeling indices between the N-CAM or N-CAM related molecules and the random peptide within each brain region were found to be significant ($p < 0.05$; paired sample Student's t-test).
 30

Analysis of the data showed that there was less glial proliferation at the injection sites for N-CAM, 250 μ g/ml, the recombinant third Ig domain of chicken N-CAM
 35 (5mg/ml) or the N-CAM KHIFSDDSSSE peptide (1mg/ml) than

- 27 -

for the control random peptide SFSISDEDHK. The results are summarized in Table 3 and FIGURES 5, 6 and 7.

Infusion of the whole N-CAM into the cerebral cortex at the lower dose of 100 μ g/ml inhibited astrocyte proliferation by 0-10%; at 150 μ g/ml, the inhibition ranged between about 4 to about 20%.

These results indicate that N-CAM partially inhibited astrocyte cell division in the three forebrain regions examined. Nevertheless, there was regional variability in the extent to which N-CAM or its related peptides can inhibit glial proliferation. The strongest effects (~50% inhibition) were seen in the cerebral cortex with smaller effects in the hippocampus and striatum. The variability between regions may be due to several factors. Astrocytes from different regions of the brain have been shown to have variable responses to injury. In addition, differences in the ability of the injected molecules to diffuse within a structure may contribute to this variation. Overall, the entire N-CAM molecule was the most efficacious in its ability to inhibit astrocyte proliferation even though it was injected at a lower concentration than the N-CAM peptide and the recombinant third Ig domain. The entire N-CAM molecule was also able to oppose the mitogenic effects of bFGF (FIGURE 7). This is consistent with our previous studies on the inhibition of astrocyte proliferation by N-CAM *in vitro*.

Manipulation of the timing of N-CAM applications and of other reagents, such as glial cell proliferation inhibitors, may enhance the inhibition of proliferation. One useful approach would include insertion of a confined flow passageway, such as a hypodermic needle, catheter, permanent canula or the like, for supplying N-CAM, N-CAM peptides or antibodies intracranially, intraventricularly, into the subarachnoid

- 28 -

space or directly to the site of the lesion. Such an approach could provide the N-CAM, N-CAM peptides or antibodies continuously or in multiple applications.

5

EXAMPLE 6

Antibodies Immunoreactive with N-CAM Are Capable of Suppressing Glial Cell Proliferation

Polyclonal antibodies to N-CAM or their Fab' fragments also inhibited glial cell proliferation. Experimental conditions were the same as described in Example 1. Fab' fragments were prepared from anti-mouse N-CAM polyclonal antibodies as previously described (Brackenbury, R. et al. (1977) *J. Biol. Chem.* 252, 6835-6840).

Anti-mouse N-CAM polyclonal antibody (0.1 mg/ml) and Fab' fragments (0.03 mg/ml) prepared from this antibody exerted an inhibitory effect on astrocyte proliferation ($53 \pm 15\%$, $n = 2$, and $56 \pm 13\%$, $n = 4$, respectively, vs. SFM).

EXAMPLE 7

Modulation of the Inhibition of Glial Cell Proliferation by Drugs

The effects of drugs that affect particular signalling pathways were studied in the presence or absence of N-CAM to determine which pathways may have synergistic effects with N-CAM, or which agents may reverse the effects of N-CAM, are shown in Figure 8 and Table 4. We focussed on three areas: (1) activation of tyrosine kinases and protein kinases A and C, (2) activation of calcium channels, and (3) activation of G proteins.

Immunoprecipitation

Astrocyte cells were seeded at a density of $7 \times 10^4/\text{cm}^2$ in a 100 mm tissue culture dish in Dulbecco's modified essential medium (DMEM) containing 10% fetal calf

- 29 -

serum and gentamycin (5 μ g/ml) at 37 degrees C with 7% CO₂. After overnight culture, the medium was replaced with serum-free medium and the cells are cultured for an additional 48 hr. The serum-starved cells were treated for 30 minutes in the presence of N-CAM (2 μ g/ml) or buffer alone and then stimulated for 10 minutes with bFGF (100 ng/ml) to stimulate tyrosine phosphorylation through the FGF receptor.

Cells were solubilized in a lysis buffer composed of 50 mM tris-HCl, pH 7.4, 150 mM NaCl, 1.5 mM MgCl₂, 1.5 mM EGTA, 1% Triton X-100, phosphatase inhibitors (100 mM sodium fluoride, 30 mM p-nitrophenylphosphate, 10 mM sodium pyrophosphate, 500 μ M sodium orthovanadate, 5 μ M ZnCl₂), and protease inhibitors (10 μ l/ml leupeptin, 10 μ g/ml aprotinin, 1 mM PMSF). Cell lysates were immunoprecipitated either with 5 μ g/ml anti-phosphotyrosine antibodies or 10 μ g/ml of anti FGF receptor antibodies for 2 hrs. at 4°C followed by a 30 minute addition of 5 mg protein A-Sepharose beads.

Immunoprecipitates were washed in tris-buffered saline (50 mM tris-HCl, pH 7.4, 150 mM NaCl) containing 0.2% Triton X-100, boiled in Laemmli sample buffer, and the proteins resolved on a 7% SDS-polyacrylamide gel and electrophoretically transferred to Immobilon membranes (Millipore, Bedford, MA). Duplicate samples for each immunoprecipitate were blotted with antibodies to phosphotyrosine and with the particular immunoprecipitating antibody.

Immunoblotting was performed by incubation with the indicated antibodies followed by ¹²⁵I-labeled protein A; the resulting autoradiogram was exposed for 14 hr on XAR film at -70°C or scanned on a Molecular Dynamics PhosphorImager (Sunnyvale, CA). The amount of reactivity with antibodies to phosphotyrosine relative to the amount of reactivity with antibodies to each specific protein can

- 30 -

be calculated from imaging these gel blots, in extracts from N-CAM treated cells vs. non treated cells. This determines whether the FGF receptor or other proteins analyzed in this same paradigm are changed in their levels of tyrosine phosphorylation upon N-CAM treatment.

We found that several tyrosine kinase inhibitors completely blocked astrocyte proliferation, suggesting that the activity of at least one protein tyrosine kinase is required to sustain astrocyte proliferation. We examined differences in the patterns of tyrosine phosphorylated proteins between proliferating astrocytes and astrocytes treated with N-CAM. Astrocytes were treated with bFGF to stimulate tyrosine phosphorylation and cell proliferation with and without prior exposure to N-CAM. Cell extracts were then immunoprecipitated with anti-phosphotyrosine antibodies followed by immunoblots of the immunoprecipitated material with the same antibodies. These experiments indicated that at least one protein in the molecular weight range of 60-70 kDa and one protein at approximately 120 kDa (postulated to be the FGF receptor itself) showed diminished tyrosine phosphorylation after N-CAM treatment (Figure 8). The strong band at 55 kDa is the heavy chain of the immunoprecipitating Ig revealed with the second antibody and iodinated protein A.

Agents that stimulate or inhibit particular signalling pathways in the presence or absence of N-CAM can affect cell proliferation measured by thymidine incorporation (Table 4). We chose these agents based on their ability to activate or inhibit signalling pathways that have been implicated in the ability of N-CAM to support neurite outgrowth. They are grouped into agents that affect the activity of protein kinase A and protein kinase C, and agents that affect G proteins and calcium channels (Table 4).

- 31 -

The activation of protein kinase C (PKC) by phorbol (PMA) treatment led to a strong mitogenic response in astrocyte cultures. Conversely, the inhibition of protein kinase C with calphostin C, which is a highly specific inhibitor of protein kinase C, inhibits proliferation. This strongly suggests that astrocyte proliferation *in vitro* requires the activity of PKC. One effect of N-CAM might therefore be to decrease protein kinase C activity.

Activation of adenylate cyclase and protein kinase A by forskolin blocks proliferation of astrocytes in culture. The presence of K252b, a more general inhibitor of protein kinases A and C, allowed the astrocytes cells to grow normally in the presence of concentrations of N-CAM, which would otherwise inhibit cell proliferation. These results indicate that kinases other than PKC, e.g. PKA, may also be important in decreasing astrocyte proliferation. It is therefore possible that N-CAM inhibits proliferation by activating PKA and that K252b reverses this effect by its inhibition of PKA.

In contrast to these results and to those reported from experiments on neurons, we found that calcium channel blockers and arachidonic acid had no effect on the ability of N-CAM to inhibit astrocyte proliferation in culture. Activation of G proteins with mastoparan stimulated astrocyte proliferation and this was not inhibited by the concentration of N-CAM used, a concentration that inhibited glial proliferation to 54% of control in the absence of mastoparan (Table 4).

Inhibition of G proteins with pertussis toxin slightly inhibited astrocyte proliferation but N-CAM was still able to inhibit proliferation in the presence of pertussis toxin. These results are of interest since these agents that affect G proteins and calcium have been shown to inhibit the ability of N-CAM to support neurite

- 32 -

outgrowth, but they showed minimal effects on N-CAM inhibition of astrocyte proliferation. Astrocyte proliferation and neurite outgrowth are two very different biological measurements, and these results indicate that N-CAM may activate different signalling pathways in these different cell types.

Table 4:
Components of N-CAM signalling pathways.

Numbers are mean $\pm$ S.E.M. for each condition. Additions of N-CAM were 2 μ g/ml, cell density was 2×10^5 cells/ml. The control condition is serum-free medium (SFM) with no additions.

15

Reagent	Mode of Action	Concentration	[³ H] -thymidine incorporation (% SFM Ctrl)			
			Drug alone	Drug + N-CAM	SFM Ctrl	SFM + N-CAM
PKC/PKA Activators/Inhibitors						
PMA	PKC +	10 nM	138 ± 16	-	-	-
		100 nM	268 ± 34	228	100	50
Calphostin	PKC-	0.1 μM	111 ± 35	-	-	-
		0.5 μM	54 ± 35	-	-	-
Forskolin	PKA +	1 μM	73 ± 2	-	-	-
		10 μM	32 ± 6	-	-	-
K252b	PKA-/	100 nM	98 ± 17	85 ± 5	100	53 ± 8
	PKC-	200 nM	78 ± 15	64 ± 1	100	43 ± 6

20

G-Proteins, Calcium Channel Blockers						
Mastoparan	Gprot +	50 μM	213 ± 47	205 ± 30	100	54 ± 6
Pertussis Toxin	Gprot-	.5 μg/ml	74 ± 20	53 ± 9	100	50 ± 10
		1 μg/ml	67 ± 13	57 ± 13	100	49 ± 9
ω-Conotoxin/ diltiazem	Ca ⁺⁺ channel blocker	0.25 μM/ 10 μM	94 ± 17	57 ± 9	100	64 ± 2

25

Arachidonic Acid		10 μM	107 ± 10	60 ± 12	100	52 ± 4
------------------	--	-------	----------	---------	-----	--------

The effects of two drugs, mastoparan and methyl 2,5-dihydroxycinnamate, on glial cell proliferation inhibition of Ig3 were also examined. 2,5-dihydroxycinnamate (Calbiochem, San Diego, CA) is an erbstatin analogue that specifically decreases receptor-associated tyrosine kinase activity. The results (Table 5)

- 33 -

indicate that Ig3 and N-CAM effects are altered in a similar way by these drugs.

Table 5:
Drug Effects on Ig3 and N-CAM Glial Cell Proliferation Inhibition

Values are means of triplicate experiments. Additions of N-CAM and Ig3 were 2 μ g/ml, cell density was 2×10^5 cells/ml. The control condition is serum-free medium (SFM) with no additions.

	% of SFM Medium Control	Drug Alone	Drug + N-CAM	Drug + Ig3	N-CAM Alone	Ig3 Alone
10	Mastoparan (50 μ M)	210	202.5	192.5	32.5	30
	Methyl 2,5-dihydroxycinnamate (0.25 μ g/ml)	65	60	59	34	31

EXAMPLE 8

Suppression of Glial Cell Proliferation by Monoclonal Antibodies Immunoreactive with N-CAM

Glial cells and culture conditions were the same as used in Example 1. A mouse monoclonal antibody derived from clone N-CAM 16.2 was used. The antibody is an immunoglobulin type IgG_{2b} directed against the third Ig domain of human brain N-CAM, and is commercially available as Anti-N-CAM 16 from Becton-Dickinson (San Jose, CA).

The monoclonal antibody was added to cultures as diluted ascites fluid at a final concentration of about 15 μ g/ml. The antibody inhibited 60% of astrocyte proliferation.

The foregoing discussion and the accompanying examples are presented as illustrative, and are not to be taken as limiting. Still other variations within the spirit and scope of this invention are possible and will readily present themselves to those skilled in the art.

- 34 -

SEQUENCE LISTING

(1) GENERAL INFORMATION:

5

(i) APPLICANT: Edelman, Gerald M.
Crossin, Kathryn L.
Sporns, Olaf
Krushel, Leslie

10

(ii) TITLE OF INVENTION: INHIBITION OF GLIAL CELL
PROLIFERATION

(iii) NUMBER OF SEQUENCES: 7

15

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Olson & Hierl, Ltd.
(B) STREET: 20 North Wacker Drive Suite 3000
(C) CITY: Chicago
(D) STATE: IL
(E) COUNTRY: USA
(F) ZIP: 60606

20

(v) COMPUTER READABLE FORM:

25

(A) MEDIUM TYPE: Floppy disk
(B) COMPUTER: IBM PC compatible
(C) OPERATING SYSTEM: PC-DOS/MS-DOS
(D) SOFTWARE: PatentIn Release #1.0, Version

#1.30

30

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE: 11-DEC-1995
(C) CLASSIFICATION:

35

- 35 -

(vii) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/353,658

(B) FILING DATE: 09-DEC-1994

(A) APPLICATION NUMBER: US 08/440,725

5 (B) FILING DATE: 08-MAY-1995

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Cepuritis, Talivaldis

(B) REGISTRATION NUMBER: 20,818

10 (C) REFERENCE/DOCKET NUMBER: TSRI 457.1

(ix) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: 312-580-1180

(B) TELEFAX: 312-580-1189

15

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 10 amino acids

(B) TYPE: amino acid

20 (C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

25 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Gallus gallus

(F) TISSUE TYPE: Brain

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

30 Lys Tyr Ser Phe Asn Tyr Asp Gly Ser Glu
1 5 10

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

35 (A) LENGTH: 10 amino acids

- 36 -

(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear
(ii) MOLECULE TYPE: peptide
5 (iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: NO
(vi) ORIGINAL SOURCE:
(A) ORGANISM: Rattus norvegicus
(F) TISSUE TYPE: Brain
10 (xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Lys His Ile Phe Ser Asp Asp Ser Ser Glu
1 5 10

15 (2) INFORMATION FOR SEQ ID NO:3:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 24 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
20 (D) TOPOLOGY: linear
(ii) MOLECULE TYPE: mRNA
(iii) HYPOTHETICAL: NO
(iv) ANTI-SENSE: YES
(vi) ORIGINAL SOURCE:
25 (A) ORGANISM: Rattus norvegicus
(F) TISSUE TYPE: Brain
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:
AGAUCUUAG UUCGCAGCAU UGUA
24

30

(2) INFORMATION FOR SEQ ID NO:4:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 282 base pairs
(B) TYPE: nucleic acid
35 (C) STRANDEDNESS: double

- 37 -

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

5 (vi) ORIGINAL SOURCE:

(A) ORGANISM: Gallus gallus

(F) TISSUE TYPE: Brain

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

10 GTGCGTGCCA GGCAGAGCAC TATGAACGCC ACTGCCAACC TCAGCCAGTC TGTCACCTTA 60

GCATGTGATG CTGATGGCTT TCCTGAGCCA ACCATGACGT GGACAAAGGA TGGAGAGCCA 120

ATAGAGCAGG AGGATAACGA AGAGAAATAC AGTTTTAACT ACGATGGGTC CGAGCTGATC 180

15 ATCAAGAAGG TGGATAAGAG TGACGAAGCA GAGTACATCT GCATCGCTGA GAACAAGGCT 240

GGCGAGCAGG ATGCCACCAT TCATCTCAA GTCTTTGCAA AA 282

(2) INFORMATION FOR SEQ ID NO:5:

20 (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 27 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

25 (ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

30 ATATGTGGAT CCCTATTTTG CAAAGAC 27

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

35 (B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

- 38 -

(ii) MOLECULE TYPE: cDNA

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

ATGTACCCAT CGGCGTGCGT GCCAGG

26

5

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 334 base pairs

(B) TYPE: nucleic acid

10 (C) STRANDEDNESS: double

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

15

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

ATGTACCCAT GGGCGTGCGT GCCAGGCAGA GCACTATGAA CGCCACTGCC AACCTCAGCC

60

20 AGTCTGTCAC CTTAGCATGT GATGCTGATG GCTTTCCTGA GCCAACCATG ACGTGGACAA

120

AGGATGGAGA GCCAATAGAG CAGGAGGATA ACGAAGAGAA ATACAGTTTT AACTACGATG

180

GGTCCGAGCT GATCATCAAG AAGGTGGATA AGAGTGACGA AGCAGAGTAC ATCTGCATCG

240

25 CTGAGAACAA GGCTGGCGAG CAGGATGCCA CCATTCATCT CAAAGTCTTT GCAAAACCCA

300

AAATCACATA TGTGGATCCC TATTTTGCAA AGAC

334

- 39 -

WE CLAIM:

1. A method of inhibiting proliferation of cells having neural cell adhesion molecule (N-CAM) on the cell surface comprising: contacting a population of the
5 cells with a proliferation inhibiting amount of a N-CAM homophilic peptide comprising at least 7 amino acid residues, said peptide including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being
10 separated from the negatively charged amino acid residue by five intervening amino acid residues.
2. The method of claim 1 wherein the cells are glial cells.
3. The method of claim 1 wherein the positively
15 charged amino acid residue is lysine.
4. The method of claim 1 wherein the negatively charged amino acid residue is aspartate.
5. The method of claim 1 wherein an intervening amino acid residue is phenylalanine.
- 20 6. The method of claim 3 wherein the negatively charged amino acid residue is aspartate.
7. The method of claim 6 wherein an intervening amino acid residue is phenylalanine.
8. The method of claim 6 wherein the
25 intervening amino acid residue adjacent to the negatively charged amino acid residue is a negatively charged amino acid residue.
9. The method of claim 6 wherein an intervening amino acid residue is tyrosine.
- 30 10. The method of claim 7 wherein the peptide contains 7 to 90 amino acid residues.
11. The method of claim 7 wherein the peptide contains 10 amino acid residues.
12. The method of activating an intracellular
35 messenger system which comprises exposing neural cell

- 40 -

adhesion molecules on the glial cell membrane to a N-CAM homophilic peptide comprising at least 7 amino acid residues, including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues in an amount sufficient to bind the peptide to neural cell adhesion molecules.

13. A composition suitable for inhibiting proliferation of glial cells and consisting essentially of at least 0.2 $\mu\text{g}/\text{gram}$ to about 1,000 $\mu\text{g}/\text{gram}$ of a N-CAM homophilic peptide of about 7 to about 90 amino acid residues, including a positively charged amino acid residue and a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues in a physiologically acceptable vehicle adapted for transport across the blood-brain barrier.

14. A method for reducing glial cell proliferation in a mixed neuron - glial cell culture which comprises adding to the culture medium a proliferation inhibiting amount of a N-CAM homophilic peptide comprising at least 7 but no more than 90 amino acid residues, and including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues.

15. An isolated antibody reactive against a N-CAM homophilic peptide of at least 7 but no more than 90 amino acid residues, and including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues.

- 41 -

16. A method of inhibiting proliferation of glial cells which comprises the step of contacting a population of glial cells with an effective amount of the antibody of claim 15.

5 17. A method of screening for tumors that express N-CAM which comprises the steps of exposing the suspect tissue to the antibody of claim 15 and thereafter detecting the presence of bound antibody.

10 18. A method for producing a glial cell population enriched in glial cells susceptible to N-CAM homophilic peptide mediated proliferation suppression which comprises exposing the glial cell population to an oligonucleotide capable of hybridizing to messenger RNA coding for at least a portion of the third IgG domain of
15 N-CAM for a time period of about 24 to about 72 hours.

19. The method of claim 18 wherein the oligonucleotide is deoxyribonucleic acid.

20. The method of claim 18 wherein the oligonucleotide is ribonucleic acid.

20 21. The method of claim 18 wherein the oligonucleotide comprises peptide nucleic acids.

22. The method of claim 18 wherein the oligonucleotide is 24 base pairs long.

25 23. The method of claim 18 wherein the oligonucleotide is 5'-AGA UCCUAGUUCG CAGCAUUGUA-3'.

24. The method of claim 18 wherein the time period is about 48 hours.

25. An isolated nucleic acid molecule coding for a N-CAM homophilic peptide comprising at least 7 but no
30 more than 90 amino acid residues, including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues
35 operatively linked to a promoter of RNA transcription.

- 42 -

26. A vector which comprises an isolated nucleic acid molecule coding for the N-CAM homophilic peptide having at least 7 but no more than 90 amino acid residues, and including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues.

27. The vector of claim 26, wherein the isolated nucleic acid molecule is linked to a plasmid.

28. A host vector system for the production of a peptide having substantially the biological activity of the peptide of a N-CAM homophilic peptide, having at least 7 but no more than 90 amino acid residues, and including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues which system comprises the vector of claim 26 in a host which is a member of the group consisting of a bacterial cell, an insect cell, and an animal cell.

29. A method of inhibiting proliferation of cells having neural cell adhesion molecule (N-CAM) on the cell surface comprising: providing a confined flow passageway in communication with fluid surrounding the population of said cells, and administering through said confined flow passageway a proliferation inhibiting amount of a N-CAM homophilic peptide comprising at least 7 amino acid residues, said peptide including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues.

- 43 -

30. The method of claim 29 wherein the cells are glial cells.

31. The method of claim 29 wherein the positively charged amino acid residue is lysine.

5 32. The method of claim 29 wherein the negatively charged amino acid residue is aspartate.

33. The method of claim 29 wherein an intervening amino acid residue is phenylalanine.

10 34. The method of claim 31 wherein the negatively charged amino acid residue is aspartate.

35. The method of claim 34 wherein an intervening amino acid residue is phenylalanine.

15 36. The method of claim 34 wherein the intervening amino acid residue adjacent to the negatively charged amino acid residue is a negatively charged amino acid residue.

37. The method of claim 34 wherein an intervening amino acid residue is tyrosine.

20 38. The method of claim 35 wherein the peptide contains 7 to 90 amino acid residues.

39. The method of claim 35 wherein the peptide contains 10 amino acid residues.

25 40. A composition suitable for inhibiting proliferation of glial cells and consisting essentially of at least 0.2 μ g/gram to about 1,000 μ g/gram of a N-CAM homophilic peptide of about 7 to about 90 amino acid residues, including a positively charged amino acid residue and a negatively charged amino acid residue, the positively charged amino acid residue being separated from the
30 negatively charged amino acid residue by five intervening amino acid residues, and a glial cell proliferation inhibitor in a physiologically acceptable vehicle adapted for transport across the blood-brain barrier.

35 41. The composition of claim 40 wherein the glial cell proliferation inhibitor is calphostin.

- 44 -

42. The composition of claim 40 wherein the glial cell proliferation inhibitor is forskolin.

43. The composition of claim 40 wherein the glial cell proliferation inhibitor is pertussis toxin.

5 44. A method for reducing glial cell proliferation in a mixed neuron - glial cell culture which comprises adding to the culture medium a proliferation inhibiting amount of a N-CAM homophilic peptide comprising at least 7 but no more than 90 amino acid residues, and
10 including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five
15 intervening amino acid residues, and a glial cell proliferation inhibitor.

45. The composition of claim 44 wherein the glial cell proliferation inhibitor is calphostin.

46. The composition of claim 44 wherein the glial cell proliferation inhibitor is forskolin.

20 47. The composition of claim 44 wherein the glial cell proliferation inhibitor is pertussis toxin.

48. A method of inhibiting proliferation of glial cells which comprises the step of contacting a population of glial cells with an effective amount of a
25 monoclonal antibody immunoreactive with at least a portion of the third Ig domain of N-CAM.

49. The method of claim 48 where in the monoclonal antibody is anti-N-CAM 16.


30 50. A method of screening for tumors that express N-CAM which comprises the steps of exposing the suspect tissue to the antibody of claim 48 and thereafter detecting the presence of bound antibody.

51. A method of inhibiting proliferation of cells having neural cell adhesion molecule (N-CAM) on the
35 cell surface comprising: providing a confined flow

- 45 -

passageway that is in communication with the fluid surrounding the population of cells, administering through the confined flow passageway a proliferation inhibiting amount of a N-CAM homophilic peptide comprising at least 7
5 amino acid residues, said peptide including a positively charged amino acid residue spaced from a negatively charged amino acid residue, the positively charged amino acid residue being separated from the negatively charged amino acid residue by five intervening amino acid residues; and
10 repeating the administration sufficiently to optimize the inhibition of glial proliferation.

52. A method of inhibiting proliferation of cells having neural cell adhesion molecule (N-CAM) on the cell surface comprising: providing a confined flow
15 passageway that is in communication with the fluid surrounding the population of cells, and administering through the confined flow passageway an amount of a monoclonal antibody immunoreactive with at least a portion of the third Ig domain of N-CAM sufficient to inhibit glial
20 cell proliferation.



2 / 8




FIG. 2A





FIG. 2B

3 / 8

**FIG. 3**

4 / 8




FIG. 4








FIG. 5

6 / 8



7/8

**FIG. 7**




FIG. 8

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US95/15991

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) : G01N 33/53; A01N 37/18, A61K 38/00; C07K 1/00, 14/00, 17/00

US CL : 435/7.1; 514/2; 530/300, 350

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 435/7.1; 514/2; 530/300, 350

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X, P ----- Y, P	Proc. Natl. Acad. Sci. USA, Volume 92, issued May 1995, Krushel et al, "Neural cell adhesion molecule (N-CAM) inhibits astrocyte proliferation after injury to different regions of the adult rat brain", pages 4323-4327, see entire document.	13, 29-39 and 51 ----- 1-11, 14 and 40-47
X, P ----- Y, P	Proc. Natl. Acad. Sci. USA, Volume 92, issued January 1995, Sporns et al, "The neural cell adhesion molecule (N-CAM) inhibits proliferation in primary cultures of rat astrocytes", pages 542-546, see entire document.	1-11, 13 and 14 ----- 29-47 and 51



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be part of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

07 MARCH 1996

Date of mailing of the international search report

09 APR 1996

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PATRICIA A. DUFFY

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15991

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Proc. Natl. Acad. Sci. USA, Volume 83, issued December 1986, Hemperly et al, "cDNA clones of the neural cell adhesion molecule (N-CAM) lacking a membrane-spanning region consistent with evidence for membrane attachment via a phosphatidylinositol intermediate", pages 9822-9826, see entire document.	25-28
X	Proc. Natl. Acad. Sci. USA, Volume 84, issued December 1987, Edelman et al, "Cellular expression of liver and neural cell adhesion molecules after transfection of their cDNAs results in specific cell-cell binding", pages 8502-8506, see entire document.	25-28
Y	Int. J. Cancer, Volume 45, issued 1990, Fantini et al, "Suramin inhibits proliferation of rat glioma cells and alters N-CAM cell surface expression", pages 554-561, see entire document.	1-11, 14, and 40-47

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US95/15991

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:

2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:

3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☒ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
1-11, 13, 14, 25-47 and 51
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, MEDLINE, BIOSYS, EMBASE, DERWENT WPI, CAB ABSTRACTS

search terms: sequences, N-CAM, neural cell adhesion molecule, proliferation, growth, inhibits, inhibition, reduction, astrocytes, neural cells, glial cells, in vivo, in vitro, peptides, receptor, growth inhibitor, TGF-B, transforming growth factor-beta, neurons.

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

Group I, claim(s) 1-11, 13, 14, 29-39 and 51, drawn to N-CAM homophilic peptide and method of inhibiting the proliferation of cells having N-CAM adhesion molecule on their surface using only the homophilic peptide.
Group II, claim(s) 12, drawn to method of activating an intracellular messenger system by exposing neural cell adhesion molecules to a N-CAM homophilic peptide.
Group III, claim(s) 15, drawn to an antibody which is reactive against the N-CAM homophilic peptide.
Group IV, claim(s) 16, 48, 49, and 52 drawn to methods of reducing cell proliferation using the antibody.
Group V, claim(s) 17 and 50 drawn to methods of screening for tumors of administration of the antibody.
Group VI, claim(s) 18-24, drawn to methods for producing a glial cell population enriched in glial cells.
Group VII, claim(s) 25-28, drawn to nucleic acid, vector and transformed host.
Group VIII, claim(s) 40-47, drawn to a composition containing a glial cell proliferation inhibitor and N-CAM homophilic peptide and methods of use therefore.

The inventions listed as Groups I-VIII do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons: The special technical features are defined below: the special technical feature of Group I is the N-CAM homophilic peptide, and the methods drawn to inhibition of proliferation of cells having neural cell adhesion molecule using the homophilic peptide; Group II is drawn to a different method of using the homophilic peptide which requires different reagents, different method steps and has a distinct final outcome; the special technical feature of Group III is an antibody product which specifically reacts with the N-CAM homophilic peptide; the special technical feature of Group IV are methods for using the antibody product to inhibit proliferation; the special technical feature of Group V is methods of screening for tumors using the antibody product; the special technical feature of Group VI is a method of producing a glial cell population using a oligonucleotide which hybridizes to messenger RNA encoding N-CAM; the special technical feature of Group VII is an isolated nucleic acid encoding the N-CAM homophilic peptide; the special technical feature of Group VIII is the composition of a proliferation inhibitor and homophilic peptide and methods of use to inhibit proliferation of cells the methods and compositions are distinct from Group I because they require different reagents and the methods use a distinct special technical feature as defined by the claimed composition.

The inventions listed as Groups III, IV, V, VI, and VII are distinct from Group I because the latter groups do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical feature as defined above. These Groups (III, IV, V, VI, VII and VIII) have distinct special technical features as defined above. The invention listed as Group II is distinct from Group I because the former group is drawn to distinct methods of using the special technical features which do not require the inhibition of proliferation. The invention listed as Group VIII is distinct from Group I because the former group is drawn distinct compositions and distinct methods of use of these compositions. Because the methods require additional compositions of reagents distinct from the homophilic peptide alone, and corresponding methods present different method steps, and different final outcomes the invention of Group VIII is deemed distinct from Group I. The inventions listed as Groups II and VIII are distinct from Group I because the latter groups do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical feature as defined above.