




(86) **Date de dépôt PCT/PCT Filing Date:** 2013/06/21
(87) **Date publication PCT/PCT Publication Date:** 2013/12/27
(45) **Date de délivrance/Issue Date:** 2017/02/28
(85) **Entrée phase nationale/National Entry:** 2014/12/09
(86) **N° demande PCT/PCT Application No.:** NZ 2013/000108
(87) **N° publication PCT/PCT Publication No.:** 2013/191567
(30) **Priorité/Priority:** 2012/06/21 (US61/662,467)

(51) **Cl.Int./Int.Cl.** **C12N 1/21** (2006.01),
C12N 15/52 (2006.01), **C12P 7/64** (2006.01)
(72) **Inventeurs/Inventors:**
LIEW, FUNGMIN, NZ;
KOEPEKE, MICHAEL, NZ
(73) **Propriétaire/Owner:**
LANZATECH NEW ZEALAND LIMITED, NZ
(74) **Agent:** BERESKIN & PARR LLP/S.E.N.C.R.L.,S.R.L.

(54) **Titre : MICRO-ORGANISMES RECOMBINANTS EXPRIMANT UN BIODIESEL FAIT D'ACYLTRANSFERASE NON SPECIFIQUE**
(54) **Title: RECOMBINANT MICROORGANISMS EXPRESSING A NON-SPECIFIC ACYLTRANSFERASE MAKE BIODIESEL**



(57) **Abrégé/Abstract:**

A carboxydutrophic acetogenic recombinant microorganism is modified so that it produces biodiesel and optionally one or more other products by fermentation of a substrate comprising CO. Biodiesel is produced by microbial fermentation of a substrate comprising CO. The recombinant microorganism is modified to express one or more exogenous enzymes in the biodiesel biosynthesis pathway not present in a parental microorganism from which the recombinant microorganism is derived. The one or more enzymes comprise a nonspecific acyltransferase.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau(43) International Publication Date
27 December 2013 (27.12.2013)

WIPO | PCT

(10) International Publication Number
WO 2013/191567 A1

(51) International Patent Classification:

C12N 1/21 (2006.01) C12P 7/64 (2006.01)
C12N 15/52 (2006.01)

(21) International Application Number:

PCT/NZ2013/000108

(22) International Filing Date:

21 June 2013 (21.06.2013)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

61/662,467 21 June 2012 (21.06.2012) US

(71) Applicant: LANZATECH NEW ZEALAND LIMITED
[NZ/NZ]; 24 Balfour Road, Parnell, Auckland, 1052 (NZ).(72) Inventors: LIEW, FungMin; 24 Balfour Road, Parnell,
Auckland, 1052 (NZ). KOEPKE, Michael; 24 Balfour
Road, Parnell, Auckland, 1052 (NZ).(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,
MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ,
OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ,
UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ,
TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK,
EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV,
MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM,
TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW,
KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- before the expiration of the time limit for amending the
claims and to be republished in the event of receipt of
amendments (Rule 48.2(h))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: RECOMBINANT MICROORGANISMS MAKE BIODIESEL

(57) Abstract: A carboxydutrophic acetogenic recombinant microorganism is modified so that it produces biodiesel and optionally one or more other products by fermentation of a substrate comprising CO. Biodiesel is produced by microbial fermentation of a substrate comprising CO. The recombinant microorganism is modified to express one or more exogenous enzymes in the biodiesel biosynthesis pathway not present in a parental microorganism from which the recombinant microorganism is derived. The one or more enzymes comprise a nonspecific acyltransferase.




FIG. 1

**RECOMBINANT MICROORGANISMS EXPRESSING A NON-SPECIFIC
ACYLTRANSFERASE MAKE BIODIESEL**

TECHNICAL FIELD OF THE INVENTION

[0001] The present invention relates to recombinant microorganisms and methods for the production of biodiesel by microbial fermentation of a substrate comprising CO.

BACKGROUND OF THE INVENTION

[0002] Bacteria including carboxydophilic acetogens *Clostridium autoethanogenum* or *C. ljungdahlii* produce fatty acids in biosynthesis of lipids and cell membranes.

[0003] In wild-type Clostridia strains, flux down the fatty-acid pathway is significant with lipids accounting typically for 5-6 % (w/w) of the dry cell mass (respectively 1-1.5 (w/w)% of the wet cell mass) (Lepage et al., 1987, Microbiology 133: 103-110). Typically more than 95% of the lipids are in very defined C16-C18 chain length range (Lepage et al., 1987, Microbiology 133: 103-110), with 12:0, 14:0, 14:1, 16:0, 16:1, 17 Δ , 18:0, 18:1, 19 Δ fatty acids present.

[0004] Fatty acids (FAs) and their derivatives are energy dense and therefore have potential as biofuels for use as a “drop-in” transportation/jet fuel and/or for the production of other industrial chemical compounds. Examples of fatty acid derivatives include biodiesel, free fatty acids, alkenes and alkanes.

[0005] Biodiesel is a mono-alkyl ester and can be used alone in standard diesel engines, or can be blended with petrodiesel. It can also be used as a low carbon alternative to heating oil. In 2009, worldwide more than 3.5 billion gallons of biodiesel were used. Biodiesel is normally derived chemically from vegetable or animal fat by transesterification of lipids in the presence of alcohol to yield glycerine and a mono-alkyl ester. Biodiesel produced by this process can however lead to damage of diesel engines due to variations in the oils from various animal and vegetable sources which are not very defined with a wide range of carbon chain length (Fukuda et al., 2001, *Biosci Bioeng* 92: 405-416). Critical points are dilution of motor oil, coking of piston rings, corrosion of hydraulic components, and depositions in the injection system, resulting from the production process and fuel aging, resulting in some automotive manufacturers to refuse the use of animal or vegetable derived biodiesel in some of their models (Köpke et al., 2011, The Past, Present, and Future of Biofuels – Biobutanol as

Promising Alternative, In: dos Santos Bernades (Ed.) Biofuel Production-Recent Developments and Prospects, InTech, 451-486).

[0006] The current generation of biofuels that use either food or non-food crops to produce sugar or cellulose-based feedstocks may have drawbacks relating to land-use, food-security, volatility of supply and environmental issues.

[0007] It is an object of the invention to overcome these issues and provide a method of production of biodiesel, or at least to provide the public with a useful choice.

SUMMARY OF INVENTION

[0008] The invention generally provides, *inter alia*, methods for the production of biodiesel by microbial fermentation of a substrate comprising CO, and recombinant microorganisms of use in such methods.

[0009] In a first aspect, the invention provides a carboxydutrophic acetogenic recombinant microorganism capable of producing biodiesel and optionally one or more other products by fermentation of a substrate comprising CO.

[0010] In one particular embodiment, the microorganism is adapted to express one or more exogenous enzymes in the biodiesel biosynthesis pathway not present in a parental microorganism from which the recombinant microorganism is derived (may be referred to herein as an exogenous enzyme). In another embodiment, the microorganism is adapted to over-express one or more endogenous enzymes in the biodiesel synthesis pathway which are present in a parental microorganism from which the recombinant microorganism is derived (may be referred to herein as an endogenous enzyme).

[0011] In one embodiment, the recombinant microorganism is adapted to produce a greater amount of biodiesel than would be produced by a parental microorganism from which the recombinant microorganism is derived.

[0012] In one embodiment, the one or more enzyme that the microorganism is adapted to express or overexpress is an acyltransferase.

[0013] In one embodiment, the enzyme is an acyltransferase enzyme as defined in SEQ ID NO: 1, or a functionally equivalent variant thereof.

[0014] In one embodiment, the parental microorganism is capable of fermenting a substrate comprising CO to produce an alcohol but not of converting the alcohol to biodiesel and the recombinant microorganism is adapted to express one or more enzymes involved in the conversion of ethanol to biodiesel.

[0015] In one embodiment, the microorganism comprises one or more exogenous nucleic acids adapted to increase expression of one or more endogenous nucleic acids and which one or more endogenous nucleic acids encode one or more of the enzymes referred to herein before.

[0016] In one embodiment, the one or more exogenous nucleic acids adapted to increase expression is a regulatory element. In one embodiment, the regulatory element is a promoter. In one embodiment, the promoter is a constitutive promoter. In one embodiment, the promoter is selected from the group comprising Wood-Ljungdahl gene cluster, a pyruvate:ferredoxin oxidoreductase promoter, an Rnf complex operon promoter, ATP synthase operon promoter and Phosphotransacetylase/Acetate kinase operon promoters.

[0017] In one embodiment, the acetogenic carboxydophilic recombinant microorganism is further adapted to express one or more exogenous enzymes in the fatty acid biosynthesis pathway. In a further aspect, the microorganism is adapted to over-express one or more endogenous enzymes in the fatty acid biosynthesis pathway.

[0018] In one embodiment, the microorganism comprises one or more exogenous nucleic acids encoding and adapted to express one or more of the enzymes referred to hereinbefore. In one embodiment, the microorganisms comprise one or more exogenous nucleic acids encoding and adapted to express at least two of the enzymes. In other embodiments, the microorganism comprises one or more exogenous nucleic acid encoding and adapted to express five or more of the enzymes.

[0019] In one embodiment, the one or more exogenous nucleic acid is a nucleic acid construct or vector, in one particular embodiment a plasmid, encoding one or more of the enzymes referred to hereinbefore in any combination.

[0020] In one embodiment, the exogenous nucleic acid is an expression plasmid.

[0021] In one particular embodiment, the parental microorganism is selected from the group of carboxydophilic acetogenic bacteria comprising *Clostridium autoethanogenum*, *Clostridium ljungdahlii*, *Clostridium ragsdalei*, *Clostridium carboxidivorans*, *Clostridium drakei*, *Clostridium scatologenes*, *Clostridium aceticum*, *Clostridium formicoaceticum*, *Clostridium magnum*, *Butyribacterium methylotrophicum*, *Acetobacterium woodii*, *Alkalibaculum bacchii*, *Blautia producta*, *Eubacterium limosum*, *Moorella thermoacetica*, *Moorella thermautotrophica*, *Sporomusa ovata*, *Sporomusa silvacetica*, *Sporomusa sphaeroides*, *Oxobacter pfennigii*, and *Thermoanaerobacter kiuvi*.

[0022] In one embodiment the parental microorganism is *Clostridium autoethanogenum* or *Clostridium ljungdahlii*. In one particular embodiment, the microorganism is *Clostridium autoethanogenum* DSM23693 a derivative of strain DSM10061. In another particular embodiment, the microorganism is *Clostridium ljungdahlii* DSM13528 (or ATCC55383).

[0023] In a second aspect, the invention provides a nucleic acid encoding one or more enzymes which when expressed in a microorganism allows the microorganism to produce biodiesel by fermentation of a substrate comprising CO.

[0024] In one embodiment, the nucleic acid encodes two or more enzymes which when expressed in a microorganism allow the microorganism to produce biodiesel by fermentation of a substrate comprising CO.

[0025] In one embodiment, the nucleic acids of the invention encode five or more such enzymes.

[0026] In one embodiment, the enzymes are chosen from the group consisting of acyl transferase and a functionally equivalent variant thereof.

[0027] In one embodiment, the nucleic acid encoding acyl transferase is SEQ ID NO: 1 or is a functionally equivalent variant thereof.

[0028] In one embodiment, the nucleic acids of the invention further comprise a promoter. In one embodiment, the promoter allows for constitutive expression of the genes under its control. In a particular embodiment a Wood-Ljungdahl cluster promoter is used. In other

particular embodiments a pyruvate:ferredoxin oxidoreductase promoter, an Rnf complex operon promoter, ATP synthase operon promoter or a Phosphotransacetylase/Acetate kinase operon promoter is used. In one particular embodiment, the promoter is from *C. autoethanogenum*.

[0029] In a third aspect, the invention provides a nucleic acid construct or vector comprising one or more nucleic acid of the second aspect.

[0030] In one particular embodiment, the nucleic acid construct or vector is an expression construct or vector. In one particular embodiment, the expression construct or vector is a plasmid.

[0031] In a fourth aspect, the invention provides a host organism comprising any one or more of the nucleic acids of the second aspect or vectors or constructs of the third aspect.

[0032] In a fifth aspect, the invention provides a composition comprising an expression construct or vector as referred to in the third aspect of the invention and a methylation construct or vector.

[0033] Preferably, the composition is able to produce a recombinant microorganism according to the first aspect of the invention.

[0034] In one particular embodiment, the expression construct/vector and/or the methylation construct/vector is a plasmid.

[0035] In a sixth aspect, the invention provides a method for the production of biodiesel and optionally one or more other products by microbial fermentation comprising fermenting a substrate comprising CO using a recombinant microorganism of the first aspect of the invention.

[0036] In one embodiment the method comprises the steps of:

- a. providing a substrate comprising CO to a bioreactor containing a culture of one or more microorganisms of the first aspect of the invention; and
- b. anaerobically fermenting the culture in the bioreactor to produce biodiesel.

[0037] In one embodiment the method comprises the steps of:

- a. capturing CO-containing gas produced as a result of an industrial process
- b. anaerobic fermentation of the CO-containing gas to produce biodiesel by a culture containing one or more microorganisms of the first aspect of the invention.

[0038] In particular embodiments of the method aspects, the fermentation occurs in an aqueous culture medium.

[0039] In particular embodiments of the method aspects, the fermentation of the substrate takes place in a bioreactor.

[0040] Preferably, the substrate comprising CO is a gaseous substrate comprising CO. In one embodiment, the substrate comprises an industrial waste gas. In certain embodiments, the gas is steel mill waste gas or syngas.

[0041] In one embodiment, the substrate will typically contain a major proportion of CO, such as at least about 20% to about 100% CO by volume, from 20% to 70% CO by volume, from 30% to 60% CO by volume, and from 40% to 55% CO by volume. In particular embodiments, the substrate comprises about 25%, or about 30%, or about 35%, or about 40%, or about 45%, or about 50% CO, or about 55% CO, or about 60% CO by volume.

[0042] In certain embodiments the methods further comprise the step of recovering the biodiesel and optionally one or more other products from the fermentation broth.

[0043] In a seventh aspect, the invention provides biodiesel when produced by the method of the sixth aspect.

[0044] In another aspect, the invention provides a method for the production of a microorganism of the first aspect of the invention comprising transforming a carboxydophilic acetogenic parental microorganism by introduction of one or more nucleic acids such that the microorganism is capable of producing biodiesel, or producing an increased amount of biodiesel compared to the parental microorganism, and optionally one or more other products by fermentation of a substrate comprising CO, wherein the parental microorganism is not capable of producing biodiesel, or produces biodeiesel at a lower level than the recombinant microorganism, by fermentation of a substrate comprising CO.

[0045] In one particular embodiment, a parental microorganism is transformed by introducing one or more exogenous nucleic acids adapted to express one or more enzymes in the biodiesel biosynthesis pathway. In a further embodiment, a parental microorganism is further transformed by introducing one or more exogenous nucleic acids adapted to express one or more enzyme in the fatty acid biosynthesis pathway. In a further embodiment, a parental microorganism is further transformed by expressing or overexpressing one or more endogenous nucleic acids adapted to express one or more enzyme in the fatty acid biosynthesis pathway. In one embodiment, a parental microorganism is transformed with one or more nucleic acids adapted to over-express one or more endogenous enzymes in the biodiesel pathway which are naturally present in the parental microorganism.

[0046] In certain embodiments, the one or more enzymes are as herein before described.

[0047] In one embodiment a genetically engineered carboxydophilic acetogenic bacterium comprises an exogenous nucleic acid encoding a nonspecific acetyltransferase (wax ester synthase/acyl Coenzyme A:diacylglycerol acyltransferase). The bacterium may be a Clostridium, including but not limited to *Clostridium autoethanogenum*, *Clostridium ljungdahlii*, *Clostridium ragsdalei*, *Clostridium carboxidivorans*, *Clostridium drakei*, *Clostridium scatologenes*, *Clostridium aceticum*, *Clostridium formicoaceticum*, and *Clostridium magnum*. Other Clostridia species which may be used, albeit not acetogenic, include, *Clostridium acetobutylicum*, *Clostridium beijerinckii*, *C. saccharobutylicum*, *C. saccharoperbutylacetonicum*, *C. thermocellum*, *C. cellulolyticum*, *C. phytofermentans*, *C. kluyveri*, and *C. pasterianum*.

[0048] . The bacterium may also be, for example, *Butyribacterium methylotrophicum*, *Acetobacterium woodii*, *Alkalibaculum bacchii*, *Blautia producta*, *Eubacterium limosum*, *Moorella thermoacetica*, *Moorella thermautotrophica*, *Sporomusa ovata*, *Sporomusa silvacetica*, *Sporomusa sphaeroides*, *Oxobacter pfennigii*, or *Thermoanaerobacter kiuvi*. The exogenous nonspecific acetyl transferase may be *Acinetobacter baylyi* nonspecific acetyl transferase. The nucleic acid may be on a plasmid. The nucleic acid encoding the nonspecific acetyltransferase may be codon optimized for *C. autoethanogenum* or for another host bacterium.

[0049] Another embodiment is a process for converting CO and/or CO₂ into biodiesel. A gaseous CO-containing and/or CO₂-containing substrate is passed to a bioreactor that contains a culture of carboxydophilic, acetogenic bacteria in a culture medium. The bacteria comprise an exogenous nucleic acid encoding a nonspecific acetyltransferase (wax ester synthase/acyl Coenzyme A:diacylglycerol acyltransferase). The bacteria convert the CO and/or CO₂ directly to biodiesel, without the need to supply alcohols (*e.g.*, ethanol or butanol) or fatty acids. The biodiesel is recovered from the bioreactor. The substrate may comprise an industrial waste gas. The culture may be grown and maintained strictly as anaerobically. The biodiesel may comprise fatty acid ethyl esters and/or fatty acid butyl esters.

[0050] Another embodiment is a plasmid which replicates in a carboxydophilic acetogenic bacterium. The plasmid comprises an exogenous nucleic acid encoding a nonspecific acetyltransferase (wax ester synthase/acyl Coenzyme A:diacylglycerol acyltransferase). The nucleic acid encoding the nonspecific acetyltransferase may be codon optimized for *C. autoethanogenum*. Optionally the plasmid may be methylated, for example by passage through a bacterium that contains a desired methylase.

BRIEF DESCRIPTION OF THE FIGURES

[0052] These and other aspects of the present invention, which should be considered in all its novel aspects, will become apparent from the following description, which is given by way of example only, with reference to the accompanying figures, in which:

[0053] Figure 1: Conversion of carbon monoxide and/or hydrogen to an alcohol such as ethanol or butanol, then subsequent conversion of the alcohol and a fatty acid-CoA ester to a fatty acid acylester (biodiesel) by an unspecific acyltransferase

[0054] Figure 2: Genetic map of expression plasmid pMTL85245-atf

[0055] Figure 3: GC-MS result confirming biodiesel production from CO.

DETAILED DESCRIPTION OF THE INVENTION

[0056] The following is a description of the present invention, including preferred embodiments thereof, given in general terms. The invention is further elucidated from the disclosure given under the heading “Examples” herein below, which provides experimental data supporting the invention, specific examples of various aspects of the invention, and means of performing the invention.

[0057] As referred to herein, a “fermentation broth” is a culture medium comprising at least a nutrient media and bacterial cells.

[0058] As referred to herein, a “shuttle microorganism” is a microorganism in which a methyltransferase enzyme is expressed and is distinct from the destination microorganism.

[0059] As referred to herein, a “destination microorganism” is a microorganism in which the genes included on an expression construct/vector are expressed and is distinct from the shuttle microorganism.

[0060] The term “main fermentation product” is intended to mean the one fermentation product which is produced in the highest concentration and/or yield.

[0061] The terms “increasing the efficiency,” “increased efficiency” and the like, when used in relation to a fermentation process, include, but are not limited to, increasing one or more of the rate of growth of microorganisms catalysing the fermentation, the growth and/or product production rate at elevated product concentrations, the volume of desired product produced per volume of substrate consumed, the rate of production or level of production of the desired product, and the relative proportion of the desired product produced compared with other by-products of the fermentation.

[0062] The phrase “substrate comprising carbon monoxide” and like terms should be understood to include any substrate in which carbon monoxide is available to one or more strains of bacteria for growth and/or fermentation, for example.

[0063] The phrase “gaseous substrate comprising carbon monoxide” and like phrases and terms includes any gas which contains a level of carbon monoxide. In certain embodiments

the substrate contains at least about 20% to about 100% CO by volume, from 20% to 70% CO by volume, from 30% to 60% CO by volume, and from 40% to 55% CO by volume. In particular embodiments, the substrate comprises about 25%, or about 30%, or about 35%, or about 40%, or about 45%, or about 50% CO, or about 55% CO, or about 60% CO by volume.

[0064] While it is not necessary for the substrate to contain any hydrogen, the presence of H₂ should not be detrimental to product formation in accordance with methods of the invention. In particular embodiments, the presence of hydrogen results in an improved overall efficiency of alcohol production. For example, in particular embodiments, the substrate may comprise an approx 2:1, or 1:1, or 1:2 ratio of H₂:CO. In one embodiment the substrate comprises about 30% or less H₂ by volume, 20% or less H₂ by volume, about 15% or less H₂ by volume or about 10% or less H₂ by volume. In other embodiments, the substrate stream comprises low concentrations of H₂, for example, less than 5%, or less than 4%, or less than 3%, or less than 2%, or less than 1%, or is substantially hydrogen free. The substrate may also contain some CO₂ for example, such as about 1% to about 80% CO₂ by volume, or 1% to about 30% CO₂ by volume. In one embodiment the substrate comprises less than or equal to about 20% CO₂ by volume. In particular embodiments the substrate comprises less than or equal to about 15% CO₂ by volume, less than or equal to about 10% CO₂ by volume, less than or equal to about 5% CO₂ by volume or substantially no CO₂.

[0065] In the description which follows, embodiments of the invention are described in terms of delivering and fermenting a "gaseous substrate containing CO." However, it should be appreciated that the gaseous substrate may be provided in alternative forms. For example, the gaseous substrate containing CO may be provided dissolved in a liquid. Essentially, a liquid is saturated with a carbon monoxide containing gas and then that liquid is added to the bioreactor. This may be achieved using standard methodology. By way of example, a microbubble dispersion generator (Hensirisak et. al. Scale-up of microbubble dispersion generator for aerobic fermentation; Applied Biochemistry and Biotechnology Volume 101, Number 3 / October, 2002) could be used. By way of further example, the gaseous substrate containing CO may be adsorbed onto a solid support. Such alternative methods are encompassed by use of the term "substrate containing CO" and the like.

[0066] In particular embodiments of the invention, the CO-containing gaseous substrate is an industrial off or waste gas. "Industrial waste or off gases" should be taken broadly to include

any gases comprising CO produced by an industrial process and include gases produced as a result of ferrous metal products manufacturing, non-ferrous products manufacturing, petroleum refining processes, gasification of coal, gasification of biomass, electric power production, carbon black production, and coke manufacturing. Further examples may be provided elsewhere herein.

[0067] Unless the context requires otherwise, the phrases “fermenting,” “fermentation process” or “fermentation reaction” and the like, as used herein, are intended to encompass both the growth phase and product biosynthesis phase of the process. As will be described further herein, in some embodiments the bioreactor may comprise a first growth reactor and a second fermentation reactor. As such, the addition of metals or compositions to a fermentation reaction should be understood to include addition to either or both of these reactors.

[0068] The term “bioreactor” includes a fermentation device consisting of one or more vessels and/or towers or piping arrangement, which includes the Continuous Stirred Tank Reactor (CSTR), Immobilized Cell Reactor (ICR), Trickle Bed Reactor (TBR), Bubble Column, Gas Lift Fermenter, Static Mixer, or other vessel or other device suitable for gas-liquid contact. In some embodiments the bioreactor may comprise a first growth reactor and a second fermentation reactor. As such, when referring to the addition of substrate to the bioreactor or fermentation reaction it should be understood to include addition to either or both of these reactors where appropriate.

[0069] “Exogenous nucleic acids” are nucleic acids which originate outside of the microorganism to which they are introduced. Exogenous nucleic acids may be derived from any appropriate source, including, but not limited to, the microorganism to which they are to be introduced (for example in a parental microorganism from which the recombinant microorganism is derived), strains or species of microorganisms which differ from the organism to which they are to be introduced, or they may be artificially or recombinantly created. In one embodiment, the exogenous nucleic acids represent nucleic acid sequences naturally present within the microorganism to which they are to be introduced, and they are introduced to increase expression of or over-express a particular gene (for example, by increasing the copy number of the sequence (for example a gene), or introducing a strong or constitutive promoter to increase expression). In another embodiment, the exogenous nucleic

acids represent nucleic acid sequences not naturally present within the microorganism to which they are to be introduced and allow for the expression of a product not naturally present within the microorganism or increased expression of a gene native to the microorganism (for example in the case of introduction of a regulatory element such as a promoter). The exogenous nucleic acid may be adapted to integrate into the genome of the microorganism to which it is to be introduced or to remain in an extra-chromosomal state.

[0070] “Exogenous” may also be used to refer to proteins. This refers to a protein that is not present in the parental microorganism from which the recombinant microorganism is derived.

[0071] The term “endogenous” as used herein in relation to a recombinant microorganism and a nucleic acid or protein refers to any nucleic acid or protein that is present in a parental microorganism from which the recombinant microorganism is derived.

[0072] “Biodiesel” as referred to herein refers to a fatty acid alkyl ester for example comprising either fatty acid ethyl ester (FAEE) and/or fatty acid butyl ester (FABE). The biodiesel produced may a mixture of fatty acid alkyl esters.

[0073] The “biodiesel biosynthesis pathway” as referred to herein refers to the pathway from fatty acyl CoA to biodiesel. Exemplary enzymes in this pathway include but are not limited to acyl transferase [EC:2.3.-.-] and acyl-CoA synthetase/long-chain-fatty-acid--CoA ligase [EC:6.2.3.1].

[0074] The “fatty acid biosynthesis pathway” refers to the pathway from acetyl CoA to the production of a fatty acyl CoA. Exemplary enzymes in this pathway include but are not limited to acetyl-CoA carboxylase/biotin carboxylase [EC:6.3.4.14/EC:6.4.1.2/EC:6.4.1.3], malonyltransferase/malonnate decarboxylase [EC:2.3.1.39], fatty acid synthase [EC:2.3.1.85/EC:2.3.1.86/EC:2.3.1.-], 3-oxoacyl-[acyl-carrier-protein] synthase [EC:2.3.1.41/EC:2.3.1.179/EC:2.3.1.180], 3-oxoacyl-[acyl-carrier protein] reductase [EC:1.1.1.100], 3-hydroxymyristoyl ACP dehydrase [EC:4.2.1.-], 3-hydroxydecanoyl-[acyl-carrier-protein] dehydratase [EC:4.2.1.60], enoyl-[acyl-carrier protein] reductase [EC:1.3.1.9, EC:1.3.1.-, EC:1.3.1.-], fatty acyl-ACP thioesterase [EC:3.1.2.- 3.1.2.14], oleoyl-[acyl-carrier-protein] hydrolase [EC:3.1.2.14], acyl-[acyl-carrier-protein] desaturase [EC:1.14.19.2], acetyl-CoA acyltransferase [EC:2.3.1.16], 3-hydroxyacyl-CoA dehydrogenase [EC:1.1.1.35], enoyl-CoA hydratase / long-chain 3-hydroxyacyl-CoA

dehydrogenase [EC:1.1.1.211, EC:4.2.1.17], enoyl-CoA hydratase [EC:4.2.1.17], trans-2-enoyl-CoA reductase [EC:1.3.1.38], palmitoyl-protein thioesterase [EC:3.1.2.22], fatty acid elongation protein [EC:2.3.1.-], 3-ketoacyl-CoA synthase [EC:2.3.1.-], beta-keto reductase [EC:1.1.1.-], 3-hydroxy acyl-CoA dehydratase [EC:4.2.1.-], enoyl reductase [EC:1.3.1.-], palmitoyl-CoA hydrolase [EC:3.1.2.2].

[0075] It should be appreciated that the invention may be practised using nucleic acids whose sequence varies from the sequences specifically exemplified herein provided they perform substantially the same function. For nucleic acid sequences that encode a protein or peptide this means that the encoded protein or peptide has substantially the same function. For nucleic acid sequences that represent promoter sequences, the variant sequence will have the ability to promote expression of one or more genes. Such nucleic acids may be referred to herein as “functionally equivalent variants.” By way of example, functionally equivalent variants of a nucleic acid include allelic variants, fragments of a gene, genes which include mutations (deletion, insertion, nucleotide substitutions and the like) and/or polymorphisms and the like. Homologous genes from other microorganisms may also be considered as examples of functionally equivalent variants of the sequences specifically exemplified herein. These include homologous genes in species such as *Clostridium acetobutylicum*, *Clostridium beijerinckii*, *C. ljungdahlii*, *Acinetobacter baylyi* details of which are publicly available on websites such as Genbank or NCBI. The phrase “functionally equivalent variants” should also be taken to include nucleic acids whose sequence varies as a result of codon optimisation for a particular organism. “Functionally equivalent variants” of a nucleic acid herein will preferably have at least approximately 70%, preferably approximately 80%, more preferably approximately 85%, preferably approximately 90%, preferably approximately 95% or greater nucleic acid sequence identity with the nucleic acid identified.

[0076] It should also be appreciated that the invention may be practised using polypeptides whose sequence varies from the amino acid sequences specifically exemplified herein. These variants may be referred to herein as “functionally equivalent variants.” A functionally equivalent variant of a protein or a peptide includes those proteins or peptides that share at least 40%, preferably 50%, preferably 60%, preferably 70%, preferably 75%, preferably 80%, preferably 85%, preferably 90%, preferably 95% or greater amino acid identity with the protein or peptide identified and has substantially the same function as the peptide or protein

of interest. Such variants include within their scope fragments of a protein or peptide wherein the fragment comprises a truncated form of the polypeptide wherein deletions may be from 1 to 5, to 10, to 15, to 20, to 25 amino acids, and may extend from residue 1 through 25 at either terminus of the polypeptide, and wherein deletions may be of any length within the region; or may be at an internal location. Functionally equivalent variants of the specific polypeptides herein should also be taken to include polypeptides expressed by homologous genes in other species of bacteria, for example as exemplified in the previous paragraph.

[0077] “Substantially the same function” as used herein is intended to mean that the nucleic acid or polypeptide is able to perform the function of the nucleic acid or polypeptide of which it is a variant. For example, a variant of an enzyme of the invention will be able to catalyse the same reaction as that enzyme. However, it should not be taken to mean that the variant has the same level of activity as the polypeptide or nucleic acid of which it is a variant.

[0078] One may assess whether a functionally equivalent variant has substantially the same function as the nucleic acid or polypeptide of which it is a variant using methods known to one of skill in the art. However, by way of example, assays to test for acyltransferase activity are described in Kalscheuer et al., 2004, *Appl. Environ. Microbiol.*, 70: 7119-25; Stöveken et al., 2005, *J. Bacteriol.*, 187: 1369-76.

[0079] “Over-express,” “over expression” and like terms and phrases when used in relation to the invention should be taken broadly to include any increase in expression of one or more proteins (including expression of one or more nucleic acids encoding same) as compared to the expression level of the protein (including nucleic acids) of a parental microorganism under the same conditions. It should not be taken to mean that the protein (or nucleic acid) is expressed at any particular level.

[0080] A “parental microorganism” is a microorganism used to generate a recombinant microorganism of the invention. The parental microorganism may be one that occurs in nature (ie a wild type microorganism) or one that has been previously modified but which does not express or over-express one or more of the enzymes the subject of the present invention. Accordingly, the recombinant microorganisms of the invention may have been modified to express or over-express one or more enzymes that were not expressed or over-expressed in the parental microorganism.

[0081] The terms nucleic acid “constructs” or “vectors” and like terms should be taken broadly to include any nucleic acid (including DNA and RNA) suitable for use as a vehicle to transfer genetic material into a cell. The terms should be taken to include plasmids, viruses (including bacteriophage), cosmids and artificial chromosomes. Constructs or vectors may include one or more regulatory elements, an origin of replication, a multicloning site and/or a selectable marker. In one particular embodiment, the constructs or vectors are adapted to allow expression of one or more genes encoded by the construct or vector. Nucleic acid constructs or vectors include naked nucleic acids as well as nucleic acids formulated with one or more agents to facilitate delivery to a cell (for example, liposome-conjugated nucleic acid, an organism in which the nucleic acid is contained).

[0082] The inventors have surprisingly shown that a recombinant microorganism can be engineered to produce a biodiesel from a CO-containing substrate. The inventors have engineered recombinant organisms and invented methods of use thereof for the production of the fatty acid derivative biodiesel. The inventors also contemplate that other fatty acid derivatives including free fatty acids, alkanes and alkenes could be produced as part of the invention. All these products can be derived from fatty acid key intermediates fatty acid acyl-CoA (thioesters with CoA) or fatty acid ACPs (Acyl carrier proteins).

[0083] The biodiesel produced by the invention is a long-chain, energy dense compound, and its synthesis requires the cell to invest energy in the form of nucleoside triphosphates such as ATP. In an aerobic process and/or using sugar as a substrate requires sufficient energy to be supplied from glycolysis to yield several molecules of ATP. The production of biodiesel via the fatty acid biosynthesis pathway in an aerobic process and/or using sugar as a substrate proceeds in a relatively straightforward manner due to the C5 pentose and C6 hexose molecules which are converted into longer chain fatty acids driven by the high ATP availability although a large number of reactions are required. The present invention may have advantages over producing biofuels from sugar based substrates and provides an alternative means for the production of biodiesel utilising waste gases including carbon monoxide from industrial processes.

[0084] For anaerobic acetogens using a C1 substrate like CO or CO₂, it is more difficult to build up long molecules such as fatty acids as unlike in glycolysis, no net energy is gained from substrate-level phosphorylation in the carbon fixating Wood-Ljungdahl pathway, in fact

activation of CO₂ to formate even requires one molecule of ATP and a membrane gradient is required. To date the product with most carbon atoms reported in acetogens (both native and recombinant organisms) are C₄ compounds butanol and 2,3-butanediol.

[0085] The inventors have shown that it is possible to produce these longer chain fatty acid molecules such as biodiesel using the C₁ feedstock CO via the acetyl CoA intermediate. As the substrate CO or CO₂ is in the Wood-Ljungdahl pathway directly channelled into acetyl-CoA (the starting point of the fatty acid biosynthesis), fewer reactions and enzymes are needed as from sugar via glycolysis, making the process faster and more efficient even though less ATP is available. Though less ATP is available in carboxydophilic acetogens, the inventors consider that this more direct pathway may sustain a higher metabolic flux (owing to higher chemical motive force of intermediate reactions).

[0086] In a particular embodiment of the invention, the inventors have found that the production of biodiesel (a fatty acid alkyl ester) by a recombinant microorganism of the invention is enabled by introduction to the microorganism of an exogenous acyl transferase. Traditional methods of production of biodiesel involve the transesterification of a lipid (triglyceride) in the presence of alcohol to yield glycerine and biodiesel. However, the present invention provides a recombinant microorganism that is able to co-produce both alcohol (including ethanol and/or butanol) as well as fatty acid. The inventors believe that this co-production provides the requisite substrates to provide a driving force for the *in vivo* production of biodiesel comprising either FAEE (Fatty acid ethyl esters) and/or FABE (Fatty acid butyl esters).

[0087] To achieve an embodiment of the invention, an unspecific acyltransferase (wax ester synthase/acyl Coenzyme A:diacylglycerol acyltransferase) from *Acinetobacter baylyi* was introduced to the acetogenic microorganism. The microorganism produces an alcohol (for example ethanol or butanol) and a fatty acyl-CoA which are converted to to a fatty acid alkyl ester (i.e. biodiesel) (Fig. 1) (Kalscheuer et al., 2004, *Appl. Environ. Microbiol.*, 70: 7119-25; Stöveken et al., 2005, *J. Bacteriol.*, 187: 1369-76). Fatty acid acyl-CoAs are a direct product of fatty acids and are produced by the action of for example acyl-CoA synthetase (long-chain-fatty-acid--CoA ligase) which may be present in carboxydophilic acetogens. *In vivo* production of FAEE using this enzyme has not been shown except with supplemental fatty acids or alcohol being supplied to the reaction externally (Kalscheuer et al., 2006,

Microbiology, 152, 2529-36). While all organisms produce fatty acid precursors, for organisms that don't produce (high amounts of) alcohols like *E. coli*, additional genetic modifications become necessary. Bacterial production of FABE has not been demonstrated previously at all. The present invention does not require such external supply of alcohol therefore may provide a number of advantages. Included in these advantages is the reduction in the cost of feedstock, a significant reduction in complexity of the equipment and parameter control and limited handling and separation steps required by the process.

[0088] While the inventors have demonstrated the efficacy of the invention in *Clostridium autoethanogenum*, they contemplate that the invention is applicable to the wider group of carboxydutrophic acetogenic microorganisms and discussed further herein.

Microorganisms

[0089] As discussed hereinbefore, the invention provides a recombinant microorganism capable of producing biodiesel, and optionally one or more other products, by fermentation of a substrate comprising CO.

[0090] In one particular embodiment, the microorganism is adapted to express one or more exogenous enzymes in the biodiesel biosynthesis pathway. In another embodiment, the microorganism is adapted to over-express one or more endogenous enzymes in the biodiesel biosynthesis pathway.

[0091] In one embodiment, the recombinant microorganism is adapted to produce a greater amount of biodiesel than would be produced by a parental microorganism from which the recombinant microorganism is derived.

[0092] In one embodiment, the parental microorganism from which the recombinant microorganism is derived is capable of fermenting a substrate comprising CO to produce an alcohol but not of converting the alcohol to a biodiesel, and the recombinant microorganism is adapted to express one or more enzymes involved in the conversion of ethanol to biodiesel.

[0093] In one embodiment, the acetogenic carboxydutrophic recombinant microorganism is further adapted to express one or more exogenous enzymes in the fatty acid biosynthesis pathway. In a further aspect, the microorganism is further adapted to over-express one or more endogenous enzymes in the fatty acid biosynthesis pathway.

[0094] The microorganism may be adapted to express or over-express the one or more enzymes by any number of recombinant methods including, for example, increasing expression of endogenous genes (for example, by introducing a stronger or constitutive promoter to drive expression of a gene), increasing the copy number of a gene encoding a particular enzyme by introducing exogenous nucleic acids encoding and adapted to express the enzyme, or introducing an exogenous nucleic acid encoding and adapted to express an enzyme not naturally present within the parental microorganism.

[0095] In certain embodiments, the parental microorganism may be transformed to provide a combination of a) increased or over-expression of one or more endogenous genes and b) introduction of one or more exogenous genes. For example, one or more genes encoding one or more enzymes in the biodiesel and optionally the fatty acid biosynthesis pathway may be native to the parental microorganism but it may not include one or more other genes encoding one or more other enzymes in the pathway.

[0096] In one embodiment the one or more enzymes in the biodiesel biosynthesis pathway are chosen from the group consisting of acyl transferase and a functionally equivalent variant thereof. By way of example only, sequence information for acyl transferase is provided.

[0097] The enzymes and functional variants of use in the microorganisms of the invention may be derived from any appropriate source, including different genera and species of bacteria, or other organisms. However, in one embodiment, the acyl transferase is that derived from *Acinetobacter baylyi* as described in SEQ ID NO: 1, or a functionally equivalent variant thereof. In a particular embodiment, the acyl transferase has the identifying characteristics of the unspecific acyltransferase YP_045555.1; Gene ID: 2879218 of *Acinetobacter baylyi*. An acyl-CoA synthetase/long-chain-fatty-acid--CoA ligase is for example given under accession numbers P69451 or GeneID: 946327.

[0098] In one embodiment, the microorganism comprises one or more exogenous nucleic acids adapted to increase expression of one or more nucleic acids native to the parental microorganism and which one or more nucleic acids encode one or more of the enzymes referred to herein before. In one embodiment, the one or more exogenous nucleic acid adapted to increase expression is a regulatory element. In one embodiment, the regulatory element is a promoter. In one embodiment, the promoter is a constitutive promoter that is

preferably highly active under appropriate fermentation conditions. Inducible promoters could also be used. In preferred embodiments, the promoter is selected from the group comprising Wood-Ljungdahl gene cluster, a pyruvate:ferredoxin oxidoreductase promoter, an Rnf complex operon promoter, ATP synthase operon promoter or Phosphotransacetylase/Acetate kinase operon promoters. It will be appreciated by those of skill in the art that other promoters which can direct expression, preferably a high level of expression under appropriate fermentation conditions, would be effective as alternatives to the exemplified embodiments.

[0099] In one embodiment, the microorganism comprises one or more exogenous nucleic acids encoding and adapted to express one or more of the enzymes referred to herein before. In one embodiment, the microorganisms comprise one or more exogenous nucleic acids encoding and adapted to express at least two of the enzymes. In other embodiments, the microorganism comprises one or more exogenous nucleic acid encoding and adapted to express three of the enzymes. In other embodiments, the microorganism comprises one or more exogenous nucleic acid encoding and adapted to express five of the enzymes.

[0100] In one particular embodiment, the microorganism comprises one or more exogenous nucleic acids encoding an acyl transferase or a functionally equivalent variant thereof.

[0101] In one embodiment, the acyl transferase is encoded by the nucleic acid sequence exemplified in SEQ ID NO: 1, or a functionally equivalent variant thereof.

[0102] The microorganism may comprise one or more exogenous nucleic acids. Where it is desirable to transform the parental microorganism with two or more genetic elements (such as genes or regulatory elements (for example a promoter)) they may be contained on one or more exogenous nucleic acids.

[0103] In one embodiment, the one or more exogenous nucleic acid is a nucleic acid construct or vector, in one particular embodiment a plasmid, encoding one or more of the enzymes referred to hereinbefore in any combination.

[0104] The exogenous nucleic acids may remain extra-chromosomal upon transformation of the parental microorganism or may intergrade into the genome of the parental microorganism. Accordingly, they may include additional nucleotide sequences adapted to assist integration

(for example, a region which allows for homologous recombination and targeted integration into the host genome) or expression and replication of an extrachromosomal construct (for example, origin of replication, promoter and other regulatory elements or sequences).

[0105] In one embodiment, the exogenous nucleic acids encoding one or enzymes as mentioned herein before will further comprise a promoter adapted to promote expression of the one or more enzymes encoded by the exogenous nucleic acids. In one embodiment, the promoter is a constitutive promoter that is preferably highly active under appropriate fermentation conditions. Inducible promoters could also be used. In preferred embodiments, the promoter is selected from the group comprising Wood-Ljungdahl gene cluster, a pyruvate:ferredoxin oxidoreductase promoter, an Rnf complex operon promoter, ATP synthase operon promoter and Phosphotransacetylase/Acetate kinase promoters. It will be appreciated by those of skill in the art that other promoters which can direct expression, preferably a high level of expression under appropriate fermentation conditions, would be effective as alternatives to the exemplified embodiments.

[0106] In one embodiment, the exogenous nucleic acid is an expression plasmid.

[0107] In one embodiment, the parental carboxydutrophic acetogenic microorganism is selected from the group consisting of *Clostridium autoethanogenum*, *Clostridium ljungdahlii*, *Clostridium ragsdalei*, *Clostridium carboxidivorans*, *Clostridium drakei*, *Clostridium scatologenes*, *Butyribacterium limosum*, *Butyribacterium methylotrophicum*, *Acetobacterium woodii*, *Alkalibaculum bacchii*, *Blautia producta*, *Eubacterium limosum*, *Moorella thermoacetica*, *Moorella thermautotrophica*, *Oxobacter pfennigii*, and *Thermoanaerobacter kiuvi*.

[0108] In one particular embodiment of the first or second aspects, the parental microorganism is selected from the group of carboxydutrophic Clostridia comprising *Clostridium autoethanogenum*, *Clostridium ljungdahlii*, *Clostridium ragsdalei*, *Clostridium carboxidivorans*, *Clostridium drakei*, *Clostridium scatologenes*, *Clostridium aceticum*, *Clostridium formicoaceticum*, *Clostridium magnum*.

[0109] In a one embodiment, the microorganism is selected from a cluster of carboxydutrophic Clostridia comprising the species *C. autoethanogenum*, *C. ljungdahlii*, and “*C. ragsdalei*” and related isolates. These include but are not limited to strains *C.*

autoethanogenum JAI-1^T (DSM10061) (Abrini, Naveau, & Nyns, 1994), *C. autoethanogenum* LBS1560 (DSM19630) (WO/2009/064200), *C. autoethanogenum* LBS1561 (DSM23693), *C. ljungdahlii* PETC^T (DSM13528 = ATCC 55383) (Tanner, Miller, & Yang, 1993), *C. ljungdahlii* ERI-2 (ATCC 55380) (US patent 5,593,886), *C. ljungdahlii* C-01 (ATCC 55988) (US patent 6,368,819), *C. ljungdahlii* O-52 (ATCC 55989) (US patent 6,368,819), or “*C. ragsdalei* P11^T” (ATCC BAA-622) (WO 2008/028055), and related isolates such as “*C. coskatii*” (US patent 2011/0229947), “*Clostridium* sp. MT351” (Michael Tyurin & Kiriukhin, 2012) and mutant strains thereof such as *C. ljungdahlii* OTA-1 (Tirado-Acevedo O. Production of Bioethanol from Synthesis Gas Using *Clostridium ljungdahlii*. PhD thesis, North Carolina State University, 2010).

[0110] These strains form a subcluster within the Clostridial rRNA cluster I (Collins et al., 1994), having at least 99% identity on 16S rRNA gene level, although being distinct species as determined by DNA-DNA reassociation and DNA fingerprinting experiments (WO 2008/028055, US patent 2011/0229947).

[0111] The strains of this cluster are defined by common characteristics, having both a similar genotype and phenotype, and they all share the same mode of energy conservation and fermentative metabolism. The strains of this cluster lack cytochromes and conserve energy via an Rnf complex.

[0112] All strains of this cluster have a genome size of around 4.2 MBp (Köpke et al., 2010) and a GC composition of around 32 %mol (Abrini et al., 1994; Köpke et al., 2010; Tanner et al., 1993) (WO 2008/028055; US patent 2011/0229947), and conserved essential key gene operons encoding for enzymes of Wood-Ljungdahl pathway (Carbon monoxide dehydrogenase, Formyl-tetrahydrofolate synthetase, Methylene-tetrahydrofolate dehydrogenase, Formyl-tetrahydrofolate cyclohydrolase, Methylene-tetrahydrofolate reductase, and Carbon monoxide dehydrogenase/Acetyl-CoA synthase), hydrogenase, formate dehydrogenase, Rnf complex (*rnfCDGEAB*), pyruvate:ferredoxin oxidoreductase, aldehyde:ferredoxin oxidoreductase (Köpke et al., 2010, 2011). The organization and number of Wood-Ljungdahl pathway genes, responsible for gas uptake, has been found to be the same in all species, despite differences in nucleic and amino acid sequences (Köpke et al., 2011).

[0113] The strains all have a similar morphology and size (logarithmic growing cells are between 0.5-0.7 x 3-5 μm), are mesophilic (optimal growth temperature between 30-37 °C) and strictly anaerobe (Abrini et al., 1994; Tanner et al., 1993)(WO 2008/028055). Moreover, they all share the same major phylogenetic traits, such as same pH range (pH 4-7.5, with an optimal initial pH of 5.5-6), strong autotrophic growth on CO containing gases with similar growth rates, and a metabolic profile with ethanol and acetic acid as main fermentation end product, with small amounts of 2,3-butanediol and lactic acid formed under certain conditions (Abrini et al., 1994; Köpke et al., 2011; Tanner et al., 1993) However, the species differentiate in substrate utilization of various sugars (e.g. rhamnose, arabinose), acids (e.g. gluconate, citrate), amino acids (e.g. arginine, histidine), or other substrates (e.g. betaine, butanol). Some of the species were found to be auxotroph to certain vitamins (e.g. thiamine, biotin) while others were not. Reduction of carboxylic acids into their corresponding alcohols has been shown in a range of these organisms (Perez, Richter, Loftus, & Angenent, 2012).

[0114] The traits described are therefore not specific to one organism like *C. autoethanogenum* or *C. ljungdahlii*, but rather general traits for carboxydutrophic, ethanol-synthesizing Clostridia. Thus, the invention can be anticipated to work across these strains, although there may be differences in performance.

[0115] The recombinant carboxydutrophic acetogenic microorganisms of the invention may be prepared from a parental carboxydutrophic acetogenic microorganism and one or more exogenous nucleic acids using any number of techniques known in the art for producing recombinant microorganisms. By way of example only, transformation (including transduction or transfection) may be achieved by electroporation, electrofusion, ultrasonication, polyethylene glycol-mediated transformation, conjugation, or chemical and natural competence. Suitable transformation techniques are described for example in Sambrook J, Fritsch EF, Maniatis T: Molecular Cloning: A laboratory Manual, Cold Spring Harbour Labrotary Press, Cold Spring Harbour, 1989.

[0116] Electroporation has been described for several carboxydutrophic acetogens as *C. ljungdahlii* (Köpke et al., 2010; Leang, Ueki, Nevin, & Lovley, 2012) (PCT/NZ2011/000203; WO2012/053905), *C. autoethanogenum* (PCT/NZ2011/000203; WO2012/053905), *Acetobacterium woodii* (Strätz, Sauer, Kuhn, & Dürre, 1994) or *Moorella thermoacetica* (Kita et al., 2012) and is a standard method used in many *Clostridia* such as *C.*

acetobutylicum (Mermelstein, Welker, Bennett, & Papoutsakis, 1992), *C. cellulolyticum* (Jennert, Tardif, Young, & Young, 2000) or *C. thermocellum* (MV Tyurin, Desai, & Lynd, 2004).

[0117] Electrofusion has been described for acetogenic *Clostridium sp.* MT351 (Tyurin and Kiriukhin, 2012).

[0118] Prophage induction has been described for carboxydophilic acetogen as well in case of *C. scatologenes* (Prasanna Tamarapu Parthasarathy, 2010, Development of a Genetic Modification System in *Clostridium scatologenes* ATCC 25775 for Generation of Mutants, Masters Project Western Kentucky University).

[0119] Conjugation has been described as method of choice for acetogen *Clostridium difficile* (Herbert, O'Keeffe, Purdy, Elmore, & Minton, 2003) and many other Clostridia including *C. acetobutylicum* (Williams, Young, & Young, 1990).

[0120] In one embodiment, the parental strain uses CO as its sole carbon and energy source.

[0121] In one embodiment the parental microorganism is *Clostridium autoethanogenum* or *Clostridium ljungdahlii*. In one particular embodiment, the microorganism is *Clostridium autoethanogenum* DSM23693. In another particular embodiment, the microorganism is *Clostridium ljungdahlii* DSM13528 (or ATCC55383).

Nucleic acids

[0122] The invention also provides one or more nucleic acids or nucleic acid constructs of use in generating a recombinant microorganism of the invention.

[0123] In one embodiment, the nucleic acids comprises sequences encoding one or more of the enzymes in the biodiesel biosynthesis pathway which when expressed in a microorganism allows the microorganism to produce biodiesel by fermentation of a substrate comprising CO. In one particular embodiment, the invention provides a nucleic acid encoding two or more enzymes which when expressed in a microorganism allows the microorganism to produce biodiesel by fermentation of a substrate comprising CO. In one embodiment, the nucleic acids of the invention encode three such enzymes, or five such enzymes.

[0124] In one particular embodiment, the enzymes are chosen from the group consisting of acyl transferase and a functionally equivalent variant thereof.

[0125] Exemplary amino acid sequences and nucleic acid sequences encoding enzymes described herein are provided herein or can be obtained from GenBank as mentioned hereinbefore. However, skilled persons will readily appreciate alternative nucleic acids sequences encoding the enzymes or functionally equivalent variants thereof, having regard to the information contained herein, in GenBank and other databases, and the genetic code.

[0126] In one embodiment, the acyl transferase is encoded by the sequence of SEQ ID NO: 1 or a functionally equivalent variant thereof.

[0127] In one embodiment, the nucleic acid further encodes one or more exogenous enzymes in the fatty acid biosynthesis pathway. In a further aspect, the nucleic acid further encodes one or more endogenous enzymes in the fatty acid biosynthesis pathway.

[0128] In one embodiment, the nucleic acids of the invention will further comprise a promoter. In one embodiment, the promoter allows for constitutive expression of the genes under its control. However, inducible promoters may also be employed. Persons of skill in the art will readily appreciate promoters of use in the invention. Preferably, the promoter can direct a high level of expression under appropriate fermentation conditions. In a particular embodiment a Wood-Ljungdahl cluster promoter is used. In another embodiment, a Phosphotransacetylase/Acetate kinase promoter is used. In another embodiment a pyruvate:ferredoxin oxidoreductase promoter, an Rnf complex operon promoter or an ATP synthase operon promoter. In one particular embodiment, the promoter is from *C. autoethanogenum*.

[0129] The nucleic acids of the invention may remain extra-chromosomal upon transformation of a parental microorganism or may be adapted for integration into the genome of the microorganism. Accordingly, nucleic acids of the invention may include additional nucleotide sequences adapted to assist integration (for example, a region which allows for homologous recombination and targeted integration into the host genome) or stable expression and replication of an extrachromosomal construct (for example, origin of replication, promoter and other regulatory sequences).

[0130] In one embodiment, the nucleic acid is nucleic acid construct or vector. In one particular embodiment, the nucleic acid construct or vector is an expression construct or vector, however other constructs and vectors, such as those used for cloning are encompassed by the invention. In one particular embodiment, the expression construct or vector is a plasmid.

[0131] It will be appreciated that an expression construct/vector of the present invention may contain any number of regulatory elements in addition to the promoter as well as additional genes suitable for expression of further proteins if desired. In one embodiment the expression construct/vector includes one promoter. In another embodiment, the expression construct/vector includes two or more promoters. In one particular embodiment, the expression construct/vector includes one promoter for each gene to be expressed. In one embodiment, the expression construct/vector includes one or more ribosomal binding sites, preferably a ribosomal binding site for each gene to be expressed.

[0132] It will be appreciated by those of skill in the art that the nucleic acid sequences and construct/vector sequences described herein may contain standard linker nucleotides such as those required for ribosome binding sites and/or restriction sites. Such linker sequences should not be interpreted as being required and do not provide a limitation on the sequences defined.

[0133] Nucleic acids and nucleic acid constructs, including expression constructs/vectors of the invention may be constructed using any number of techniques standard in the art. For example, chemical synthesis or recombinant techniques may be used. Such techniques are described, for example, in Sambrook et al (Molecular Cloning: A laboratory manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989). Further exemplary techniques are described in the Examples section herein after. Essentially, the individual genes and regulatory elements will be operably linked to one another such that the genes can be expressed to form the desired proteins. Suitable vectors for use in the invention will be appreciated by those of ordinary skill in the art. However, by way of example, the following vectors may be suitable: pMTL80000 vectors, pIMP1, pJIR750, and the plasmids exemplified in the Examples section herein after.

[0134] It should be appreciated that nucleic acids of the invention may be in any appropriate form, including RNA, DNA, or cDNA.

[0135] The invention also provides host organisms, particularly microorganisms, and including viruses, bacteria, and yeast, comprising any one or more of the nucleic acids described herein.

Method of producing microorganisms

[0136] The one or more exogenous nucleic acids may be delivered to a parental microorganism as naked nucleic acids or may be formulated with one or more agents to facilitate the transformation process (for example, liposome-conjugated nucleic acid, an organism in which the nucleic acid is contained). The one or more nucleic acids may be DNA, RNA, or combinations thereof, as is appropriate. Restriction inhibitors may be used in certain embodiments; see, for example Murray, N.E. *et al.* (2000) *Microbial. Molec. Biol. Rev.* 64, 412.)

[0137] The microorganisms of the invention may be prepared from a parental microorganism and one or more exogenous nucleic acids using any number of techniques known in the art for producing recombinant microorganisms. By way of example only, transformation (including transduction or transfection) may be achieved by electroporation, ultrasonication, polyethylene glycol-mediated transformation, chemical or natural competence, protoplast transformation, prophage induction or conjugation. Suitable transformation techniques are described for example in, Sambrook J, Fritsch EF, Maniatis T: *Molecular Cloning: A laboratory Manual*, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, 1989.

[0138] Electroporation has been described for several carboxydophilic acetogens as *C. ljungdahlii* (Köpke *et al.* 2010, *Proc. Nat. Acad. Sci. U.S.A.* 107: 13087-92; PCT/NZ2011/000203; WO2012/053905), *C. autoethanogenum* (PCT/NZ2011/000203; WO2012/053905), or *Acetobacterium woodii* (Straetz *et al.*, 1994, *Appl. Environ. Microbiol.* 60:1033-37) and is a standard method used in many Clostridia such as *C. acetobutylicum* (Mermelstein *et al.*, 1992, *Biotechnology*, 10, 190-195), *C. cellulolyticum* (Jennert *et al.*, 2000, *Microbiology*, 146: 3071-3080) or *C. thermocellum* (Tyurin *et al.*, 2004, *Appl. Environ. Microbiol.* 70: 883-890). Prophage induction has been demonstrated for carboxydophilic acetogen as well in case of *C. scatologenes* (Prasanna Tamarapu Parthasarathy, 2010,

Development of a Genetic Modification System in *Clostridium scatologenes* ATCC 25775 for Generation of Mutants, Masters Project Western Kentucky University), while conjugation has been described as method of choice for many Clostridia including *Clostridium difficile* (Herbert et al., 2003, *FEMS Microbiol. Lett.* 229: 103-110) or *C. acetobutylicum* (Williams et al., 1990, *J. Gen. Microbiol.* 136: 819-826) and could be used in a similar fashion for carboxydutrophic acetogens.

[0139] In certain embodiments, due to the restriction systems which are active in the microorganism to be transformed, it is necessary to methylate the nucleic acid to be introduced into the microorganism. This can be done using a variety of techniques, including those described below, and further exemplified in the Examples section herein after.

[0140] By way of example, in one embodiment, a recombinant microorganism of the invention is produced by a method comprises the following steps:

introduction into a shuttle microorganism of (i) of an expression construct/vector as described herein and (ii) a methylation construct/vector comprising a methyltransferase gene;
expression of the methyltransferase gene;
isolation of one or more constructs/vectors from the shuttle microorganism; and,
introduction of the one or more construct/vector into a destination microorganism.

[0141] In one embodiment, the methyltransferase gene is expressed constitutively. In another embodiment, expression of the methyltransferase gene of is induced.

[0142] The shuttle microorganism is a microorganism, preferably a restriction negative microorganism, that facilitates the methylation of the nucleic acid sequences that make up the expression construct/vector. In a particular embodiment, the shuttle microorganism is a restriction negative *E. coli*, *Bacillus subtilis*, or *Lactococcus lactis*.

[0143] The methylation construct/vector comprises a nucleic acid sequence encoding a methyltransferase.

[0144] Once the expression construct/vector and the methylation construct/vector are introduced into the shuttle microorganism, the methyltransferase gene present on the methylation construct/vector is induced. Induction may be by any suitable promoter system although in one particular embodiment of the invention, the methylation construct/vector

comprises an inducible lac promoter and is induced by addition of lactose or an analogue thereof, more preferably isopropyl- β -D-thio-galactoside (IPTG). Other suitable promoters include the ara, tet, or T7 system. In a further embodiment of the invention, the methylation construct/vector promoter is a constitutive promoter.

[0145] In a particular embodiment, the methylation construct/vector has an origin of replication specific to the identity of the shuttle microorganism so that any genes present on the methylation construct/vector are expressed in the shuttle microorganism. Preferably, the expression construct/vector has an origin of replication specific to the identity of the destination microorganism so that any genes present on the expression construct/vector are expressed in the destination microorganism.

[0146] Expression of the methyltransferase enzyme results in methylation of the genes present on the expression construct/vector. The expression construct/vector may then be isolated from the shuttle microorganism according to any one of a number of known methods. By way of example only, the methodology described in the Examples section described hereinafter may be used to isolate the expression construct/vector.

[0147] In one particular embodiment, both construct/vector are concurrently isolated.

[0148] The expression construct/vector may be introduced into the destination microorganism using any number of known methods. However, by way of example, the methodology described in the Examples section hereinafter may be used. Since the expression construct/vector is methylated, the nucleic acid sequences present on the expression construct/vector are able to be incorporated into the destination microorganism and successfully expressed.

[0149] It is envisaged that a methyltransferase gene may be introduced into a shuttle microorganism and over-expressed. Thus, in one embodiment, the resulting methyltransferase enzyme may be collected using known methods and used *in vitro* to methylate an expression plasmid. The expression construct/vector may then be introduced into the destination microorganism for expression. In another embodiment, the methyltransferase gene is introduced into the genome of the shuttle microorganism followed by introduction of the expression construct/vector into the shuttle microorganism, isolation of

one or more constructs/vectors from the shuttle microorganism and then introduction of the expression construct/vector into the destination microorganism.

[0150] It is envisaged that the expression construct/vector and the methylation construct/vector as defined above may be combined to provide a composition of matter. Such a composition has particular utility in circumventing restriction barrier mechanisms to produce the recombinant microorganisms of the invention.

[0151] In one particular embodiment, the expression construct/vector and/or the methylation construct/vector are plasmids.

[0152] Persons of ordinary skill in the art will appreciate a number of suitable methyltransferases of use in producing the microorganisms of the invention. However, by way of example the *Bacillus subtilis* phage Φ T1 methyltransferase and the methyltransferase described in the Examples herein after may be used. In one embodiment, the methyltransferase has the amino acid sequence of SEQ ID NO: 12, or is a functionally equivalent variant thereof. Nucleic acids encoding suitable methyltransferases will be readily appreciated having regard to the sequence of the desired methyltransferase and the genetic code. In one embodiment, the nucleic acid encoding a methyltransferase is as described in the Examples herein after (for example the nucleic acid of SEQ ID NO: 17, or it is a functionally equivalent variant thereof).

[0153] Any number of constructs/vectors adapted to allow expression of a methyltransferase gene may be used to generate the methylation construct/vector. However, by way of example, the plasmid described in the Examples section hereinafter may be used (for example, SEQ ID NO: 14).

Methods of production

[0154] The invention provides a method for the production of biodiesel, and optionally one or more other products, by microbial fermentation comprising fermenting a substrate comprising CO using a recombinant microorganism of the invention. The methods of the invention may be used to reduce the total atmospheric carbon emissions from an industrial process.

[0155] Preferably, the fermentation comprises the steps of anaerobically fermenting a substrate in a bioreactor to produce at least biodiesel using a recombinant microorganism of the invention.

[0156] In one embodiment the method comprises the steps of:

- a. providing a substrate comprising CO to a bioreactor containing a culture of one or more microorganisms of the invention; and
- b. anaerobically fermenting the culture in the bioreactor to produce at least biodiesel.

[0157] In one embodiment the method comprises the steps of:

- a. capturing CO-containing gas produced as a result of an industrial process;
- b. anaerobic fermentation of the CO-containing gas to produce biodiesel by a culture containing one or more microorganisms of the invention.

[0158] In an embodiment of the invention, the gaseous substrate fermented by the microorganism is a gaseous substrate containing CO. The gaseous substrate may be a CO-containing waste gas obtained as a by-product of an industrial process, or from some other source such as from automobile exhaust fumes. In certain embodiments, the industrial process is selected from the group consisting of ferrous metal products manufacturing, such as a steel mill, non-ferrous products manufacturing, petroleum refining processes, gasification of coal, electric power production, carbon black production, ammonia production, methanol production and coke manufacturing. In these embodiments, the CO-containing gas may be captured from the industrial process before it is emitted into the atmosphere, using any convenient method. The CO may be a component of syngas (gas comprising carbon monoxide and hydrogen). The CO produced from industrial processes is normally flared off to produce CO₂ and therefore the invention has particular utility in reducing CO₂ greenhouse gas emissions and producing biodiesel for use as a biofuel. Depending on the composition of the gaseous CO-containing substrate, it may also be desirable to treat it to remove any undesired impurities, such as dust particles before introducing it to the fermentation. For example, the gaseous substrate may be filtered or scrubbed using known methods.

[0159] It will be appreciated that for growth of the bacteria and the production of biodiesel to occur, in addition to the CO-containing substrate gas, a suitable liquid nutrient medium will need to be fed to the bioreactor.

[0160] In particular embodiments of the method aspects, the fermentation occurs in an aqueous culture medium. In particular embodiments of the method aspects, the fermentation of the substrate takes place in a bioreactor.

[0161] The substrate and media may be fed to the bioreactor in a continuous, batch or batch fed fashion. A nutrient medium will contain vitamins and minerals sufficient to permit growth of the micro-organism used. Anaerobic media suitable for fermentation using CO are known in the art. For example, suitable media are described Biebel (2001). In one embodiment of the invention the media is as described in the Examples section herein after.

[0162] The fermentation should desirably be carried out under appropriate fermentation conditions for the production of biodiesel to occur. Reaction conditions that should be considered include pressure, temperature, gas flow rate, liquid flow rate, media pH, media redox potential, agitation rate (if using a continuous stirred tank reactor), inoculum level, maximum gas substrate concentrations to ensure that CO in the liquid phase does not become limiting, and maximum product concentrations to avoid product inhibition.

[0163] In addition, it is often desirable to increase the CO concentration of a substrate stream (or CO partial pressure in a gaseous substrate) and thus increase the efficiency of fermentation reactions where CO is a substrate. Operating at increased pressures allows a significant increase in the rate of CO transfer from the gas phase to the liquid phase where it can be taken up by the micro-organism as a carbon source for the production of fermentation. This in turn means that the retention time (defined as the liquid volume in the bioreactor divided by the input gas flow rate) can be reduced when bioreactors are maintained at elevated pressure rather than atmospheric pressure. The optimum reaction conditions will depend partly on the particular micro-organism of the invention used. However, in general, it is preferred that the fermentation be performed at pressure higher than ambient pressure. Also, since a given CO-to-biodiesel conversion rate is in part a function of the substrate retention time, and achieving a desired retention time in turn dictates the required volume of a bioreactor, the use of pressurized systems can greatly reduce the volume of the bioreactor

required, and consequently the capital cost of the fermentation equipment. According to examples given in US Patent No. 5,593,886, reactor volume can be reduced in linear proportion to increases in reactor operating pressure, i.e. bioreactors operated at 10 atmospheres of pressure need only be one tenth the volume of those operated at 1 atmosphere of pressure.

[0164] By way of example, the benefits of conducting a gas-to-ethanol fermentation at elevated pressures has been described. For example, WO 02/08438 describes gas-to-ethanol fermentations performed under pressures of 30 psig and 75 psig, giving ethanol productivities of 150 g/l/day and 369 g/l/day respectively. However, example fermentations performed using similar media and input gas compositions at atmospheric pressure were found to produce between 10 and 20 times less ethanol per litre per day.

[0165] It is also desirable that the rate of introduction of the CO-containing gaseous substrate is such as to ensure that the concentration of CO in the liquid phase does not become limiting. This is because a consequence of CO-limited conditions may be that one or more product is consumed by the culture.

[0166] The composition of gas streams used to feed a fermentation reaction can have a significant impact on the efficiency and/or costs of that reaction. For example, O₂ may reduce the efficiency of an anaerobic fermentation process. Processing of unwanted or unnecessary gases in stages of a fermentation process before or after fermentation can increase the burden on such stages (e.g. where the gas stream is compressed before entering a bioreactor, unnecessary energy may be used to compress gases that are not needed in the fermentation). Accordingly, it may be desirable to treat substrate streams, particularly substrate streams derived from industrial sources, to remove unwanted components and increase the concentration of desirable components.

[0167] In certain embodiments a culture of a bacterium of the invention is maintained in an aqueous culture medium. Preferably the aqueous culture medium is a minimal anaerobic microbial growth medium. Suitable media are known in the art and described for example in U.S. Patent Nos. 5,173,429 and 5,593,886 and WO 02/08438, and as described in the Examples section herein after.

[0168] Biodiesel, or a mixed stream containing biodiesel and/or one or more other products, may be recovered from the fermentation broth by methods known in the art, such as fractional distillation or evaporation, pervaporation, gas stripping and extractive fermentation, including for example, liquid-liquid extraction. Products may also diffuse or secrete into media, from which they can be extracted by phase separation.

[0169] In certain preferred embodiments of the invention, biodiesel and one or more products are recovered from the fermentation broth by continuously removing a portion of the broth from the bioreactor, separating microbial cells from the broth (conveniently by filtration), and recovering one or more products from the broth. Alcohols may conveniently be recovered for example by distillation. Acetone may be recovered for example by distillation. Any acids produced may be recovered for example by adsorption on activated charcoal. The separated microbial cells are preferably returned to the fermentation bioreactor. The cell free permeate remaining after any alcohol(s) and acid(s) have been removed is also preferably returned to the fermentation bioreactor. Additional nutrients (such as B vitamins) may be added to the cell free permeate to replenish the nutrient medium before it is returned to the bioreactor.

[0170] Also, if the pH of the broth was adjusted as described above to enhance adsorption of acetic acid to the activated charcoal, the pH should be re-adjusted to a similar pH to that of the broth in the fermentation bioreactor, before being returned to the bioreactor.

EXAMPLES

[0171] The invention will now be described in more detail with reference to the following non-limiting examples.

Example 1 – Production of biodiesel from CO

[0172] An acetogenic carboxydrotroph *Clostridium autoethanogenum* was engineered with the unspecific acyltransferase of *Acinetobacter baylyi* for production of a biodiesel fatty acid acyl ester, butanoic acid butyl ester (FABE). Production of butanol was demonstrated earlier using a genetically modified strain of *Clostridium autoethanogenum* (WO 2012/053905).

Strains and growth conditions:

[0173] All subcloning steps were performed in *E. coli* using standard strains and growth conditions as described earlier (Sambrook et al, Molecular Cloning: A laboratory Manual, Cold Spring Harbour Laboratory Press, Cold Spring Harbour, 1989; Ausubel et al, Current protocols in molecular biology, John Wiley & Sons, Ltd., Hoboken, 1987).

[0174] *C. autoethanogenum* DSM10061 and DSM23693 (a derivative of DSM10061) were obtained from DSMZ (The German Collection of Microorganisms and Cell Cultures, Inhoffenstraße 7 B, 38124 Braunschweig, Germany). Growth was carried out at 37° C using strictly anaerobic conditions and techniques (Hungate, 1969, Methods in Microbiology, vol. 3B. Academic Press, New York: 117-132; Wolfe, 1971, Adv. Microb. Physiol., 6: 107-146). Chemically defined PETC media without yeast extract (Tab. 1) and 30 psi carbon monoxide containing steel mill waste gas (collected from New Zealand Steel site in Glenbrook, NZ; composition: 44% CO, 32% N₂, 22% CO₂, 2% H₂) as sole carbon and energy source was used.

Table 1: PETC medium

Media component	Concentration per 1.0L of media
NH ₄ Cl	1 g
KCl	0.1 g
MgSO ₄ ·7H ₂ O	0.2 g
NaCl	0.8 g
KH ₂ PO ₄	0.1 g
CaCl ₂	0.02 g
Trace metal solution	10 ml
Wolfe's vitamin solution	10 ml

Resazurin (2 g/L stock)	0.5 ml
NaHCO ₃	2 g
Reducing agent	0.006-0.008 % (v/v)
Distilled water	Up to 1 L, pH 5.5 (adjusted with HCl)

Wolfe's vitamin solution	per L of Stock
Biotin	2 mg
Folic acid	2 mg
Pyridoxine hydrochloride	10 mg
Riboflavin	5 mg
Nicotinic acid	5 mg
Calcium D-(+)-pantothenate	5 mg
Vitamin B12	0.1 mg
p-Aminobenzoic acid	5 mg
Lipoic acid	5 mg
Thiamine	5 mg
Distilled water	To 1 L

Trace metal solution	per L of stock
Nitrilotriacetic Acid	2 g
MnSO ₄ .H ₂ O	1 g
Fe (SO ₄) ₂ (NH ₄) ₂ .6H ₂ O	0.8 g
CoCl ₂ .6H ₂ O	0.2 g
ZnSO ₄ .7H ₂ O	0.2 mg
CuCl ₂ .2H ₂ O	0.02 g
NaMoO ₄ .2H ₂ O	0.02 g
Na ₂ SeO ₃	0.02 g

NiCl ₂ .6H ₂ O	0.02 g
Na ₂ WO ₄ .2H ₂ O	0.02 g
Distilled water	To 1 L

Reducing agent stock	per 100 mL of stock
NaOH	0.9 g
Cystein.HCl	4 g
Na ₂ S	4 g
Distilled water	To 100 mL

Construction of expression plasmid:

[0175] Standard Recombinant DNA and molecular cloning techniques were used in this invention and are described by Sambrook et al, 1989 and Ausubel et al, 1987. The unspecific acyltransferase (YP_045555.1; Gene ID: 2879218) of *Acinetobacter baylyi* was codon optimized and synthesized (SEQ ID NO: 1).

[0176] Genomic DNA from *Clostridium autoethanogenum* DSM 10061 was isolated using a modified method by Bertram and Dürre (1989). A 100-ml overnight culture was harvested (6,000 x g, 15 min, 4 °C), washed with potassium phosphate buffer (10 mM, pH 7.5) and suspended in 1.9 ml STE buffer (50 mM Tris-HCl, 1 mM EDTA, 200 mM sucrose; pH 8.0). 300 µl lysozyme (~100,000 U) were added and the mixture was incubated at 37 °C for 30 min, followed by addition of 280 µl of a 10 % (w/v) SDS solution and another incubation for 10 min. RNA was digested at room temperature by addition of 240 µl of an EDTA solution (0.5 M, pH 8), 20 µl Tris-HCl (1 M, pH 7.5), and 10 µl RNase A (Fermentas). Then, 100 µl Proteinase K (0.5 U) were added and proteolysis took place for 1-3 h at 37 °C. Finally, 600 µl of sodium perchlorate (5 M) were added, followed by a phenol-chloroform extraction and an isopropanol precipitation. DNA quantity and quality was inspected spectrophotometrically.

[0177] The phosphotransacetylase/acetate kinase promoter region of *C. autoethanogenum* (SEQ ID NO: 4) was amplified by PCR from genomic DNA with oligonucleotides Ppta-ack-NotI-F (SEQ ID NO: 2: GAGCGGCCGCAATATGATATTTATGTCC) and Ppta-ack-NdeI-R (SEQ ID NO: 3: TTCCATATGTTTCATGTTCATTTCTCC) and iProof High Fidelity

DNA Polymerase (Bio-Rad Laboratories) applying the following program: initial denaturation at 98 °C for 30 seconds, followed by 35 cycles of denaturation (98 °C for 10 seconds), annealing (55 °C for 30 seconds) and elongation (72 °C for 30 seconds), before a final extension step (72 °C for 10 minutes).

[0178] The amplified 498 bp promoter region of the phosphotransacetylase/acetate kinase operon (Ppta-ack) was cloned into the *E. coli*-*Clostridium* shuttle vector pMTL 85241 (FJ797651.1; Nigel Minton, University of Nottingham; Heap et al., 2009) using NotI and NdeI restriction sites and strain DH5 α -T1R (Invitrogen). Subsequently the synthesized acyltransferase gene (SEQ ID NO: 1) was cloned in using NdeI and EcoRI to form plasmid pMTL85245-atf (SEQ ID NO: 5; Fig. 2). The insert was completely sequenced using oligonucleotides given in Table 2 and results confirmed that the atf gene was free of mutations.

Table 2: Oligonucleotides for sequencing

Oligonucleotide Name	DNA Sequence (5' to 3')	SEQ ID NO:
Atf-F1	AGACAACAACCTATGCATGTTGGAGGA	6
Atf-R1	GGGGATGTGCTGCAAGGCGA	7
Atf-F2	CATCATCAAGAAGGTTTGCAGCACAAT	8
Atf-R2	AGAGGTTCTCTTGGACCTGGAACAT	9
Atf-F3	TCGGTACCCGGGGATCCTCTA	10
Atf-R3	CATTCCTGCTACTCCATCTACCATTGC	11

Methylation of DNA:

[0179] Methylation of the FAEE expression plasmid pMTL85245-atf was performed *in vivo* in *E. coli* using a synthesized hybrid Type II methyltransferase (SEQ ID NO: 12) designed from methyltransferase genes from *C. autoethanogenum*, *C. ragsdalei* and *C. ljungdahlii*. The methyltransferase is fused to an inducible lac promoter (SEQ ID NO: 13) in vector pGS20 (SEQ ID NO: 14).

[0180] Both expression plasmid and methylation plasmid were transformed into same cells of restriction negative *E. coli* XL1-Blue MRF' Kan (Stratagene), which is possible due to their compatible Gram-(-) origins of replication (high copy ColE1 in expression plasmid and low

copy p15A in methylation plasmid). In vivo methylation was induced by addition of 1 mM IPTG, and methylated plasmids were isolated using QIAGEN Plasmid Midi Kit (QIAGEN). The resulting mix was used for transformation experiments with *C. autoethanogenum* DSM23693, but only the abundant (high-copy) expression plasmid has a Gram-(+) replication origin (repL) allowing it to replicate in Clostridia.

Transformation into C. autoethanogenum:

[0181] During the complete transformation experiment, *C. autoethanogenum* DSM23693 was grown in PETC media (Tab. 1) supplemented with 1 g/L yeast extract and 10 g/l fructose as well as 30 psi steel mill waste gas (collected from New Zealand Steel site in Glenbrook, NZ; composition: 44% CO, 32% N₂, 22% CO₂, 2% H₂) as carbon source.

[0182] To make competent cells, a 50 ml culture of *C. autoethanogenum* DSM23693 was subcultured to fresh media for 3 consecutive days. These cells were used to inoculate 50 ml PETC media containing 40 mM DL-threonine at an OD_{600nm} of 0.05. When the culture reached an OD_{600nm} of 0.4, the cells were transferred into an anaerobic chamber and harvested at 4,700 x g and 4 °C. The culture was twice washed with ice-cold electroporation buffer (270 mM sucrose, 1 mM MgCl₂, 7 mM sodium phosphate, pH 7.4) and finally suspended in a volume of 600 µl fresh electroporation buffer. This mixture was transferred into a pre-cooled electroporation cuvette with a 0.4 cm electrode gap containing 1 µg of the methylated plasmid mix and immediately pulsed using the Gene pulser Xcell electroporation system (Bio-Rad) with the following settings: 2.5 kV, 600 Ω, and 25 µF. Time constants of 3.7-4.0 ms were achieved. The culture was transferred into 5 ml fresh media. Regeneration of the cells was monitored at a wavelength of 600 nm using a Spectronic Helios Epsilon Spectrophotometer (Thermo) equipped with a tube holder. After an initial drop in biomass, the cells started growing again. Once the biomass has doubled from that point, the cells were harvested, suspended in 200 µl fresh media and plated on selective PETC plates (containing 1.2 % Bacto™ Agar (BD)) with 4 µg/ml Clarithromycin. After 4-5 days of inoculation with 30 psi steel mill gas at 37 °C, colonies were visible.

[0183] The colonies were used to inoculate 2 ml PETC media containing 4 µg/µl Clarithromycin. When growth occurred, the culture was upscaled into 5 ml and later 50 ml

PETC media containing 4 µg/ml Clarithromycin and 30 psi steel mill gas as sole carbon source.

Confirmation of the successful transformation:

[0184] To verify the DNA transfer, a plasmid mini prep was performed from 10 ml culture volume using Zyppy plasmid miniprep kit (Zymo). Since the quality of the isolated plasmid was not sufficient for a restriction digest due to Clostridial exonuclease activity [Burchhardt and Dürre, 1990], a PCR was performed with the isolated plasmid and oligonucleotides given in Table 2 to confirm the presence of the plasmid. PCR was carried out using iNtRON Maximise Premix PCR kit (Intron Bio Technologies) with the following conditions: initial denaturation at 94 °C for 2 minutes, followed by 35 cycles of denaturation (94 °C for 20 seconds), annealing (55 °C for 20 seconds) and elongation (72 °C for 60 seconds), before a final extension step (72 °C for 5 minutes).

[0185] To confirm the identity of the clones, genomic DNA was isolated (see above) from 50 ml cultures of *C. autoethanogenum* DSM23693. A PCR was performed against the 16s rRNA gene using oligonucleotides fd1 (SEQ ID NO: 15: ccgaattcgtcgacaacAGAGTTTGATCCTGGCTCAG) and rP2 (SEQ ID NO: 16: cccgggatccaagcttACGGCTACCTTGTTACGACTT) [Weisberg et al., 1991] and iNtRON Maximise Premix PCR kit (Intron Bio Technologies) with the following conditions: initial denaturation at 94 °C for 2 minutes, followed by 35 cycles of denaturation (94 °C for 20 seconds), annealing (55 °C for 20 seconds) and elongation (72 °C for 60 seconds), before a final extension step (72 °C for 5 minutes). Sequencing results were at least 99.9 % identity against the 16s rRNA gene (rrsA) of *C. autoethanogenum* (Y18178, GI:7271109).

Growth experiments to confirm biodiesel production from CO:

[0186] To demonstrate FAEE production, PETC media were prepared and inoculated with *C. autoethanogenum* strain harboring expression plasmid pMTL85245-atf. Serum bottles with 50 mL PETC medium (Table 1) were pressurized with 30 psi of a CO containing gas stream from steel mill waste gas (collected from New Zealand Steel site in Glenbrook, NZ; composition: 44% CO, 32% N₂, 22% CO₂, 2% H₂) and cultivated for 5 days. The same experiment was also carried out with the wild-type *C. autoethanogenum* strain without plasmid.

[0187] The cultures were analyzed by GC-MS using headspace sampling. 2 mL sample in 20 mL vial were exposed for 10 min at 40°C to a fibre (Supelco PDMS 100 fibre) and then analyzed using an Agilent 6890 GC with 5973 MSD equipped with a 30m x 0.25mm x 0.25µm ZB-Wax column at following conditions: Injector temperature: 250 °C; Splitless injection; desorb for 10 min at 250 °C; 1mL/min constant flow; Oven: 40°C hold for 5 min, raise at 10°C/min to 190°C, hold for 5 min, raise at 3°C/min to 208°C, raise at 10°C/min to 220°C, hold 10 min, back to 40°C at 60°C/min; MSD: Scan mode, mass range 38-650 AMU at 1.47 scans per second. Two peaks which matches to biodiesel substance butanoic acid butyl ester against the national Institute of Standards and Technology (NIST) standard reference database were found in the strain carrying the expression plasmid but not in the wild-type strain without plasmid, as well as some fatty acid products in C14-C18 range like 1-Octadecanol (C18) or Tetradecanal (C14), Heptadecane (C17), 9-Octadecanal (C18) and 11-Hexadecanal (C16)(Tab. 3; Fig. 3) . Alcohols like ethanol and butanol were detected by HPLC performed using an Agilent 1100 Series HPLC system equipped with a RID operated at 35 °C (Refractive Index Detector) and an Aminex HPX-87H column (300 x 7.8 mm, particle size 9 µm) kept at 35 °C. The RID was operated at 35 °C (Refractive Index Detector) and an Alltech IOA-2000 Organic acid column (150 x 6.5 mm, particle size 8 µm) kept at 60 °C. Slightly acidified water was used (0.005 M H₂SO₄) as mobile phase with a flow rate of 0.25 ml/min. To remove proteins and other cell residues, 400 µl samples were mixed with 100 µl of a 2 % (w/v) 5-Sulfosalicylic acid and centrifuged at 14,000 x g for 3 min to separate precipitated residues. 10 µl of the supernatant were then injected into the HPLC for analyses.

Table 3: Results from GC-MS analysis of strain *C. autoethanogenum* harboring expression plasmid pMTL85245-atf:

Retention Time		% NIST match
1.5-2.8	CO ₂	90
3.84	Disulfide	90
4-4.5	Acetic acid	96

WO 2013/191567

PCT/NZ2013/000108

14.37	1-Octadecanol	<50
16.35	Butanoic acid butylester	50
16.64	Butanoic acid butylester	78
18.84	Tetradecanal	95
21	Heptadecane	>90
21.7	9-Octadecanal(Z) / 11-Hexadecanal(Z)	93 / 87

[0188] The invention has been described herein, with reference to certain preferred embodiments, in order to enable the reader to practice the invention without undue experimentation. However, a person having ordinary skill in the art will readily recognise that many of the components and parameters may be varied or modified to a certain extent or substituted for known equivalents without departing from the scope of the invention. It should be appreciated that such modifications and equivalents are herein incorporated as if individually set forth. Titles, headings, or the like are provided to enhance the reader's comprehension of this document, and should not be read as limiting the scope of the present invention.

[0189]

However, the reference to any applications, patents and publications in this specification is not, and should not be taken as an acknowledgment or any form of suggestion that they constitute valid prior art or form part of the common general knowledge in any country in the world.

[0190] Throughout this specification and any claims which follow, unless the context requires otherwise, the words "comprise," "comprising" and the like, are to be construed in an inclusive sense as opposed to an exclusive sense, that is to say, in the sense of "including, but not limited to."

References

- Abrini, J., Naveau, H., & Nyns, E. J. (1994). *Clostridium autoethanogenum*, sp. nov., an anaerobic bacterium that produces ethanol from carbon monoxide. *Archives of microbiology*, 161(4), 345–351.
- Collins, M. D., Lawson, P. A., Willems, A., Cordoba, J. J., Fernandez-Garayzabal, J., Garcia, P., Cai, J., et al. (1994). The phylogeny of the genus *Clostridium*: proposal of five new genera and eleven new species combinations. *International journal of systematic bacteriology*, 44(4), 812–26.
- Herbert, M., O’Keeffe, T. a., Purdy, D., Elmore, M., & Minton, N. P. (2003). Gene transfer into *Clostridium difficile* CD630 and characterisation of its methylase genes. *FEMS Microbiology Letters*, 229(1), 103–110.
- Jennert, K. C., Tardif, C., Young, D. I., & Young, M. (2000). Gene transfer to *Clostridium cellulolyticum* ATCC 35319. *Microbiology (Reading, England)*, 146 Pt 12, 3071–80.
- Kita, A., Iwasaki, Y., Sakai, S., Okuto, S., Takaoka, K., Suzuki, T., Yano, S., et al. (2012). Development of genetic transformation and heterologous expression system in carboxydotrophic thermophilic acetogen *Moorella thermoacetica*. *Journal of Bioscience and Bioengineering*, xx(xx).
- Köpke, M., Held, C., Hujer, S., Liesegang, H., Wiezer, A., Wollherr, A., Ehrenreich, A., et al. (2010). *Clostridium ljungdahlii* represents a microbial production platform based on syngas. *Proceedings of the National Academy of Sciences of the United States of America*, 107(29).
- Köpke, M., Mihalcea, C., Liew, F., Tizard, J. H., Ali, M. S., Conolly, J. J., Al-Sinawi, B., et al. (2011). 2,3-Butanediol Production By Acetogenic Bacteria, an Alternative Route To Chemical Synthesis, Using Industrial Waste Gas. *Applied and environmental microbiology*, 77(15), 5467–75.
- Leang, C., Ueki, T., Nevin, K. P., & Lovley, D. R. (2012). A Genetic System for *Clostridium ljungdahlii*: A Chassis for Autotrophic Production of Biocommodities and a Model Homoacetogen. *Applied and environmental microbiology*, (November).

Mermelstein, L. D., Welker, N. E., Bennett, G. N., & Papoutsakis, E. T. (1992). Expression of cloned homologous fermentative genes in *Clostridium acetobutylicum* ATCC 824. *Bio/technology (Nature Publishing Company)*, 10(2), 190–195.

Perez, J. M., Richter, H., Loftus, S. E., & Angenent, L. T. (2012). Biocatalytic reduction of short-chain carboxylic acids into their corresponding alcohols with syngas fermentation. *Biotechnology and bioengineering*, 1–30.

Strätz, M., Sauer, U., Kuhn, a, & Dürre, P. (1994). Plasmid Transfer into the Homoacetogen *Acetobacterium woodii* by Electroporation and Conjugation. *Applied and environmental microbiology*, 60(3), 1033–7.

Tanner, R. S., Miller, L. M., & Yang, D. (1993). *Clostridium ljungdahlii* sp. nov., an acetogenic species in clostridial rRNA homology group I. *International journal of systematic bacteriology*, 43(2), 232.

Tyurin, Michael, & Kiriukhin, M. (2012). Electrofusion of cells of Acetogen *Clostridium* sp. MT 351 with *erm* (B) or *cat* in the chromosome. *Journal of Biotech*, 1–12.

Tyurin, MV, Desai, S., & Lynd, L. (2004). Electrotransformation of *Clostridium thermocellum*. *Applied and environmental mictrobiology* 70(2), 883–890.

Williams, D. R., Young, D. I., & Young, M. (1990). Conjugative plasmid transfer from *Escherichia coli* to *Clostridium acetobutylicum*. *Journal of general microbiology*, 136(5), 819–26.

WHAT IS CLAIMED IS:

1. A genetically engineered *Clostridium autoethanogenum* or *Clostridium ljungdahlii* bacterium comprising an exogenous nucleic acid encoding a nonspecific acyltransferase comprising SEQ ID NO: 18.
2. The bacterium of claim 1 wherein the nucleic acid is a plasmid.
3. The bacterium of claim 1 wherein the nucleic acid encoding the nonspecific acyltransferase is codon optimized for *Clostridium autoethanogenum*.
4. A process for converting at least one of CO and CO₂ into biodiesel, the process comprising:

passing a gaseous substrate containing at least one of CO and CO₂ to a bioreactor containing a culture of the bacterium of claim 1 in a culture medium such that the bacterium converts at least one of the CO and CO₂ to biodiesel and

recovering the biodiesel from the bioreactor.
5. The process of claim 4 wherein the substrate comprises an industrial waste gas.
6. The process of claim 4 wherein culture is anaerobic.
7. The process of claim 4 wherein the biodiesel comprises fatty acid ethyl esters.
8. The process of claim 4 wherein the biodiesel comprises fatty acid butyl esters.
9. The process of claim 4 wherein the substrate comprises syngas.




FIG. 1




FIG. 2




FIG. 3

