



US00PP15242P2

(12) **United States Plant Patent**  
**Arimitsu**

(10) **Patent No.:** **US PP15,242 P2**

(45) **Date of Patent:** **Oct. 19, 2004**

(54) *VERBENA* PLANT NAMED 'BODCOMPIN'

(50) Latin Name: *Verbena hybrida*  
Varietal Denomination: **Bodcompin**

(75) Inventor: **Yoshiro Arimitsu**, Lompoc, CA (US)

(73) Assignee: **John Bodger + Sons Co.**, South El Monte, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **10/684,564**

(22) Filed: **Oct. 13, 2003**

(51) **Int. Cl.**<sup>7</sup> ..... **A01H 5/00**

(52) **U.S. Cl.** ..... **Plt./308**

(58) **Field of Search** ..... **Plt./308**

(56) **References Cited**

PUBLICATIONS

UPOV-hit on 'Bodcompin', Plant Variety Database, Feb. 12, 2002.\*

\* cited by examiner

*Primary Examiner*—Anne Marie Grunberg  
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(57) **ABSTRACT**

A new and distinct cultivar of *Verbena* plant named 'Bodcompin', characterized by its compact, upright and mounded plant habit; freely branching habit; dense and bushy growth habit; dark green-colored leaves; and light pink-colored flowers with flowers held above and beyond the foliage.

**1 Drawing Sheet**

**1**

Botanical classification/cultivar designation: *Verbena hybrida* cultivar Bodcompin.

**BACKGROUND OF THE INVENTION**

The present Invention relates to a new and distinct cultivar of *Verbena* plant, botanically known as *Verbena hybrida*, and hereinafter referred to by the name 'Bodcompin'.

The new *Verbena* is a product of a planned breeding program conducted by the Inventor in Lompoc, Calif. The objective of the breeding program is to develop new *Verbena* cultivars with a dense and bushy growth habit, numerous flowers, and interesting flower and foliage colors.

The new *Verbena* originated from a cross-pollination made by the Inventor in July, 1991 of a proprietary *Verbena hybrida* selection identified as 1V236, not patented, as the female, or seed, parent with the *Verbena hybrida* cultivar *Amour Light Pink*, not patented, as the male, or pollen, parent. The cultivar *Bodcompin* was discovered and selected by the Inventor as a flowering plant within the progeny from this cross-pollination in a controlled environment in Lompoc, Calif.

Asexual reproduction of the new cultivar by terminal cuttings in a controlled environment in Lompoc, Calif. has shown that the unique features of this new *Verbena* are stable and reproduced true to type in successive generations of asexual reproduction.

**SUMMARY OF THE INVENTION**

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Bodcompin'. These characteristics in combination distinguish 'Bodcompin' as a new and distinct cultivar:

1. Compact, upright and mounded plant habit.
2. Freely branching habit; dense and bushy growth habit.
3. Dark green-colored leaves.
4. Light pink-colored flowers with flowers held above and beyond the foliage.

**2**

Plants of the new *Verbena* differ primarily from plants of the parents in plant habit and flower coloration.

Plants of the new *Verbena* can be compared to plants of the cultivar *Obsession Pink*, not patented. In side-by-side comparisons conducted in Lompoc, Calif., plants of the new *Verbena* differed primarily from plants of the cultivar *Obsession Pink* in plant and flower form as plants of the cultivar *Obsession Pink* are more mounded and have narrower flower petals than plants of the new *Verbena*.

Plants of the new *Verbena* can be compared to plants of the cultivar *Sandy Rose*, not patented. In side-by-side comparisons conducted in Lompoc, Calif., plants of the new *Verbena* differed primarily from plants of the cultivar *Sandy Rose* in plant form as plants of the cultivar *Sandy Rose* are more upright and taller than plants of the new *Verbena*.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Verbena*.

The photograph at the top of the sheet comprises a side perspective view of a typical flowering plant of 'Bodcompin' grown in a container.

The photograph at the bottom of the sheet comprises a close-up view of typical inflorescences, flowers and leaves of 'Bodcompin'.

**DETAILED BOTANICAL DESCRIPTION**

The cultivar *Bodcompin* has not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype. The aforementioned photographs and following observations and measurements describe plants grown in

Lompoc, Calif., under commercial practice during the winter and spring in a polycarbonate-covered greenhouse with day temperatures about 18 to 24° C., night temperatures about 16 to 18° C., and light levels about 4,000 to 8,000 foot-candles. Cuttings were planted in 10-cm containers, pinched one time, and grown for about nine weeks. In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Verbena hybrida* cultivar Bodcompin.

Parentage:

*Female, or seed, parent.*—Proprietary *Verbena hybrida* selection identified as 1V236, not patented.

*Male, or pollen, parent.*—*Verbena hybrida* cultivar Amour Light Pink, not patented.

Propagation:

*Type cutting.*—Terminal cuttings.

*Time to initiate roots, summer.*—About 7 to 10 days at 26° C.

*Time to initiate roots, winter.*—About 21 days at 26° C.

*Time to produce a rooted cutting or liner, summer.*—About 21 days at 26° C.

*Time to produce a rooted cutting or liner, winter.*—About 28 days at 26° C.

*Root description.*—Fine, fibrous; white in color.

*Rooting habit.*—Freely branching; dense.

Plant description:

*Form.*—Compact, upright and mounded plant habit.

*Growth and branching habit.*—Vigorous and freely branching with about 12 to 14 lateral branches developing after the pinch, dense and bushy growth habit.

*Plant height.*—About 16 cm.

*Plant diameter or spread.*—About 23.5 cm.

*Lateral branches.*—Length: About 18 cm. Diameter: About 2 mm. Internode length: About 4.5 cm. Texture: Pubescent. Color: 144C.

*Foliage description.*—Arrangement: Opposite, simple. Length: About 4 cm. Width: About 1.5 cm. Shape: Oblong to deltoid. Apex: Acute. Base: Acute to attenuate. Margin: Irregularly crenate. Texture, upper and lower surfaces: Coarse, pubescent. Venation pattern: Pinnate. Color: Developing foliage, upper and lower surfaces: 146B. Fully expanded, upper surface: 146A. Fully expanded, lower surface: 146B. Venation, upper surface: 146B. Venation, lower surface: 144D. Petiole: Length: About 8 mm. Diameter: About 2 mm. Color: 146C.

Flower description:

*Flower type and habit.*—Single upright salverform flowers arranged on compact terminal racemes; flowers sessile. Freely flowering with about 18 flowers and flower buds per raceme; about two to three

racemes per lateral branch. Inflorescences positioned above and beyond the foliage. Flowers last about two to four days under greenhouse conditions. Flowers not persistent.

*Fragrance.*—None detected.

*Flowering season.*—In the garden, flowering is continuous from spring until fall.

*Inflorescence height.*—About 2.8 cm.

*Inflorescence diameter.*—About 6 cm.

*Flower size.*—Diameter: About 2.5 cm. Tube length: About 2.5 cm. Throat diameter: About 2.5 mm. Tube diameter, at base: About 2 mm.

*Flower buds.*—Rate of opening, from showing color to fully open flower: About 1 to 2 days. Length: About 1.2 cm. Diameter, apex: About 4 mm. Diameter, base: About 2.5 mm. Shape: Tubular, oblong. Color: 50D.

*Petals.*—Quantity/arrangement: Five per flower fused at base. Lobe length: About 1.2 cm. Lobe width: About 1 cm. Shape: Roughly cordate. Apex: Emarginate. Margin: Entire. Texture, upper and lower surfaces: Velvety, smooth. Color: When opening, upper surface: Towards margins, 55B; center, 55C; towards base, 55D. When opening, lower surface: 56C. Fully opened, upper surface: Towards margins, 55C; towards base, 55D; color becoming closer to 56C with development. Fully opened, lower surface: 56C to 56D. Throat: 55D. Tube: 56C to 56D.

*Sepals.*—Quantity/arrangement: Five, fused into a tube. Length: About 1.3 cm. Diameter: About 2 mm. Shape: Ligulate. Apex: Acute. Margin: Entire. Texture, upper and lower surfaces: Coarse, pubescent. Color, upper surface: 144C. Color, lower surface: 144A.

*Peduncles.*—Length: About 4.75 cm. Diameter: About 1.5 mm. Angle: Upright to about 45° from vertical. Strength: Strong. Color: 144A.

*Reproductive organs.*—Stamens: Quantity per flower: Four, adnate to pistil. Anther shape: Oval. Anther length: Less than 1 mm. Anther color: 144D. Pollen amount: Scarce. Pollen color: 2D. Pistils: Quantity per flower: One. Pistil length: About 1.7 cm. Stigma shape: Bi-parted. Stigma color: 144D. Style length: About 1.5 cm. Style color: 144D. Ovary color: 144C.

*Fruit/seed.*—Fruit and seed production has not been observed.

*Disease/pest resistance:* Plants of the new *Verbena* have not been observed to be resistant to pathogens and pests common to *Verbena*.

*Temperature tolerance:* Plants of the new *Verbena* have been observed to be tolerant to temperatures ranging from 2 to 40° C.

It is claimed:

1. A new and distinct cultivar of *Verbena* plant named 'Bodcompin', as illustrated and described.

\* \* \* \* \*

