

**(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE**

(11) Application No. AU 2017319702 B2

(54) Title
Methods and compositions involving interleukin-6 receptor alpha-binding single chain variable fragments

(51) International Patent Classification(s)
A61K 48/00 (2006.01) **A61K 38/20** (2006.01)

(21) Application No: **2017319702** (22) Date of Filing: **2017.09.01**

(87) WIPO No: **WO18/042385**

(30) Priority Data

(31) Number	(32) Date	(33) Country
62/440,991	2016.12.30	US
62/443,509	2017.01.06	US
62/383,237	2016.09.02	US

(43) Publication Date: **2018.03.08**

(44) Accepted Journal Date: **2024.11.14**

(71) Applicant(s)
The Regents of the University of California

(72) Inventor(s)
Chen, Yvonne Yu-Hsuan;Lin, Meng-Yin

(74) Agent / Attorney
FB Rice Pty Ltd, L 33 477 Collins Street, Melbourne, VIC, 3000, AU

(56) Related Art
EP 1810980 A1
MAUDE SL ET AL.: "Managing Cytokine Release Syndrome Associated With Novel T Cell -Engaging Therapies", CANCER J., vol. 20, no. 2, March 2014
(2014-03-01), pages 119 - 122, XP055188104

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number

WO 2018/042385 A3

(43) International Publication Date
08 March 2018 (08.03.2018)

(51) International Patent Classification:
A61K 39/395 (2006.01) *C07K 16/28* (2006.01)
A61K 51/10 (2006.01) *C07K 14/705* (2006.01)

(21) International Application Number:
PCT/IB2017/055281

(22) International Filing Date:
01 September 2017 (01.09.2017)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
62/383,237 02 September 2016 (02.09.2016) US
62/440,991 30 December 2016 (30.12.2016) US
62/443,509 06 January 2017 (06.01.2017) US

(71) Applicant: THE REGENTS OF THE UNIVERSITY OF CALIFORNIA [US/US]; 1111 Franklin Street, 12th Floor, Oakland, California 94607 (US).

(72) Inventors: CHEN, Yvonne Yu-Hsuan; c/o The University of California, Los Angeles, 11000 Kinross Ave., Suite 200, Los Angeles, California 90095 (US). LIN, Meng-Yin; c/o The University of California, Los Angeles, 11000 Kinross Ave., Suite 200, Los Angeles, California 90095 (US).

(74) Agent: STELLMAN, Laurie et al.; Norton Rose Fulbright US LLP, 98 San Jacinto Blvd., Suite 1100, Austin, Texas 78701 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DJ, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JO, JP, KE, KG, KH, KN, KP, KR, KW, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

(54) Title: METHODS AND COMPOSITIONS INVOLVING INTERLEUKIN-6 RECEPTOR ALPHA-BINDING SINGLE CHAIN VARIABLE FRAGMENTS

scFv condition	VL-VH scFv				VH-VL scFv				No scFv	
	1ug	1ug	2ug	5ug	1ug	1ug	2ug	5ug	-	-
IL-6 (4 ng/ml)	-	+	+	+	-	+	+	+	-	+

pSTAT3
88 kDa



GAPDH
36 kDa




FIG. 1

(57) Abstract: Disclosed are compositions comprising an isolated chimeric interleukin 6 receptor alpha (IL-6Ra) binding protein or cells expressing an isolated chimeric IL-6Ra binding protein. The isolated IL-6Ra chimeric binding protein and cells expressing the protein may be used in methods of treating cancer and reducing the risk of cytokine release syndrome.



Published:

- *with international search report (Art. 21(3))*
- *before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))*

(88) Date of publication of the international search report:

19 April 2018 (19.04.2018)

DESCRIPTION

**METHODS AND COMPOSITIONS INVOLVING INTERLEUKIN-6 RECEPTOR
ALPHA-BINDING SINGLE CHAIN VARIABLE FRAGMENTS**

CROSS-REFERENCE TO RELATED APPLICATIONS

5 [0001] This application claims the benefit of U.S. Provisional Patent Application No. 62/383,237, filed September 2, 2016, U.S. Provisional Patent Application No. 62/440,991, filed December 30, 2016, and U.S. Provisional Patent Application No. 62/443,509, filed January 6, 2017. The entire contents of which applications are incorporated herein by reference.

10

BACKGROUND OF THE INVENTION

15 [0002] The prospects of using antibodies that interact with T-cells for employing new and effective therapies for the treatment of a variety of diseases and conditions, including cancer, microbial infections, and autoimmune diseases, continue to develop. However, there are side effects and toxicities that can result, including cytokine release syndrome (CRS). CRS can cause a patient to experience hypotension, pyrexia, and rigors (a sudden feeling of cold with shivering accompanied by a rise in temperature) similar to the onset of a high fever. Treatments are needed that will prevent or alleviate the side effects and toxicities from T-cell engaging therapies.

20 [0002A] Any discussion of documents, acts, materials, devices, articles or the like which has been included in the present specification is not to be taken as an admission that any or all of these matters form part of the prior art base or were common general knowledge in the field relevant to the present disclosure as it existed before the priority date of each of the appended claims.

SUMMARY OF THE INVENTION

25 [0003] Embodiments are provided in the form of methods, compositions, medical devices, and kits involving an isolated and engineered chimeric interleukin 6 receptor alpha (IL-6Ra) binding protein.

[0004] Compositions, medical devices, and kits include therapeutic compositions or a kit that has at least one therapeutic composition that comprises one or more of the following: an

isolated chimeric interleukin 6 receptor alpha (IL-6R α) binding protein, a nucleic acid encoding a chimeric IL-6R α binding protein, one or more cells expressing a chimeric IL-6R α binding protein, or one or more cells that have a nucleic acid encoding a chimeric IL-6R α binding protein.

[0005] In some embodiments, there is a chimeric interleukin 6 receptor alpha (IL-6R α) binding protein comprising a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and

5 CDR6 (SEQ ID NO:10). In some embodiments, a chimeric IL-6R α binding protein has at least or at most 1, 2, 3, 4, 5, or 6 amino acid substitutions, deletions, or additions in one or more of CDR1, CDR2, CDR3, CDR4, CDR5, and/or CDR6. In other embodiments, a chimeric IL-6R α binding protein has 60, 65, 70, 75, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 10 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% (or any range derivable therein) sequence identity with the amino acid sequence of CDR1, CDR2, CDR3, CDR4, CDR5, and/or CDR6 or with the heavy chain variable domain (SEQ ID NO:3) or with light chain variable domain (SEQ ID NO:4) or with the V_L-V_H single chain variable fragment (scFv) (SEQ ID NO:1) or with the V_H-V_L scFv (SEQ ID NO:2). In certain embodiments the chimeric IL-6R α binding protein is isolated and/or purified. In some embodiments, the IL6R α binding protein 15 comprises SEQ ID NO:1 or SEQ ID NO:2. Further embodiments concern an scFv that is essentially or comprises SEQ ID NO:1 or SEQ ID NO:2.

[0006] In some embodiments, the chimeric IL-6R α binding protein is an scFv, (scFv)₂, scFvFc, Fab, Fab', or F(ab)₂. In some cases, the scFv portion comprises all or part of SEQ ID NOs:1-10. In certain embodiments, the protein is 1, 2, 3, 4 or more polypeptides. In specific 20 embodiments, the IL-6R α binding protein is one polypeptide. In other specific embodiments, the chimeric chimeric IL-6R α binding protein is a fusion protein. In some embodiments, the fusion protein is one polypeptide.

[0007] Certain embodiments concern a chimeric IL-6R α binding protein that is a single chain variable fragment (scFv). In embodiments involving a single polypeptide containing 25 both a heavy chain variable region and a light chain variable region, both orientations of these variable regions are contemplated. In some cases, the heavy chain variable region is on the N-terminal side of the light chain variable region, which means the heavy chain variable region is closer to the N-terminus of the polypeptide. SEQ ID NO:2 is an example of this configuration. In other cases, the light chain variable region is on the N-terminal side of the 30 heavy chain variable region, which means the light chain variable region is closer to the

N-terminus of the polypeptide than the heavy chain variable region. SEQ ID NO:1 is an example of this structure.

[0008] In some embodiments, a chimeric IL-6R α binding protein comprises a linker. In some cases, the linker is between the heavy chain variable region and the light chain variable region (or vice versa). The linker comprises the amino acid sequence GSTSGSGKPGSSEGSTKG (SEQ ID NO:11) in some embodiments or has 10, 11, 12, 13, 14, 15, 16, 17, or 18 contiguous amino acids from the sequence of SEQ ID NO:11. In other embodiments, the linker is 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100% identical in amino acid sequence to SEQ ID NO:11.

[0009] Some chimeric IL-6R α binding proteins also have a leader or signal peptide at or toward the N-terminus of a polypeptide. The leader or signal peptide refers to a short (5-30 amino acids long) peptide present at the N-terminus of a polypeptide to promote or allow for secretion of that polypeptide.

[0010] In particular embodiments, a chimeric IL-6R α binding protein may include an isolation tag, which refers to a peptide tag that is used to isolate or purify the protein or polypeptide. The tag may or may not be removable, such as by cleavage. The tag may be a chitin binding protein (CBP), a maltose binding protein (MBP), glutathione-S-transferase (GST), a His tag (such as His₆). In some instances, the isolation tag is a chromatography tag that is used to alter chromatographic properties of the protein to afford different resolution across a particular separation technique. An example of this is a FLAG tag such as DYKDDDDK, which is a series of polyanionic amino acids. Another isolation tag may be an epitope tag, which refers to a short peptide sequence can be used in conjunction with an antibody. Epitope tags include, but are not limited to, a V5 tag, a c-myc tag, an HA tag, or an NE tag.

[0011] Some embodiments involve an isolated chimeric interleukin 6 receptor alpha (IL-6R α) binding protein comprising a leader peptide, heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and CDR6 (SEQ ID NO:10), wherein the heterologous linker comprises the amino acid sequence GSTSGSGKPGSSEGSTKG (SEQ ID NO:11) and the IL-6R α binding protein is a single-chain variable fragment (scFv). Embodiments do not

involve tocilizumab (also known as atlizumab), which is sold under the trade names of Actemra and RoActemra and is a humanized monoclonal antibody unless specifically included. In specific embodiments, the IL-6R α binding protein reduces the expression of one or more cytokines from a T cell. In some embodiments, the T cell expresses a chimeric antigen receptor or another protein that activates T cells. In certain embodiments, the IL-6R α binding protein reduces the expression of any of the following cytokines: TNF- α , IL-6, IL-2, IL-4, and GM-CSF (or any combination thereof). In some instances, an IL-6R α binding protein reduces the expression of one or more of those cytokines by, by at least, or by at most 10%, 20%, 30%, 40%, or 50% (or any range derivable therein) as compared to a cell not exposed to the IL-6R α binding protein. In some embodiments, a IL-6R α scFv reduces the expression of any of the following cytokines: TNF- α , IL-6, IL-2, IL-4, and GM-CSF (or any combination thereof). In some instances, an IL-6R α scFv reduces the expression of one or more of those cytokines by, by at least, or by at most 10%, 20%, 30%, 40%, or 50% (or any range derivable therein) as compared to a cell exposed to tocilizumab. In other embodiments, an IL-6R α scFv reduces the expression of one or more cytokines greater than tocilizumab. Methods similarly may involve an IL-6R α scFv that reduces the expression of one or more cytokines greater than tocilizumab. In certain embodiments, methods involve an IL-6R α scFv that reduces the expression of TNF- α , IL-6, IL-2, IL-4, and/or GM-CSF greater than tocilizumab. In some embodiments, the IL-6 receptor alpha single chain variable fragment that reduces expression of one or more cytokines comprises a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and CDR6 (SEQ ID NO:10). Other IL-6R α scFvs discussed in other paragraphs can reduce the expression of a cytokine as described herein.

[0012] A “chimeric interleukin 6 receptor alpha (IL-6R α) binding protein” refers to a protein that specifically binds to IL-6 receptor alpha and that contains at least two amino acid regions (each region is at least 10 contiguous amino acids), each region being from a different gene, and wherein the chimeric protein is not found in nature. In some embodiments, the chimeric IL-6R α protein is a single polypeptide. A chimeric IL-6R α binding protein does not refer to an IL-6R α antibody that is found in nature. “Isolated” refers to the separation or purification of the protein from either other proteins or one or more other components of a cell.

[0013] Other aspects concern a cell or cells that express a chimeric IL-6R α binding protein. The cells may express the chimeric IL-6R α binding protein on its surface or they may express and secrete the chimeric IL-6R α binding protein. In certain embodiments, T cells express the chimeric IL-6R α binding protein. The T cells may be among other blood cells or they may be in a pool of enriched T cells. The T cells may be conventional T cells of naïve, central memory, effector memory, or effector phenotypes; or they may be regulatory T cells of natural or induced phenotypes. In some cases, the T cells express one or more of the following markers: CD4, CD8, CD25, CD27, CD28, CD45RA, CD45RO, CD57, CD62L, CD69, CD95, CD127, CCR7, IL-7R α , or Foxp3. In some embodiments the T cell is a CD4+ T cell. In some embodiments, the T cell is a CD8+ T cell.

[0014] In some embodiments, there is a nucleic acid molecule encoding the chimeric IL-6R α binding protein. Certain embodiments involve a nucleic acid molecule that also comprises a promoter controlling expression of the chimeric IL-6R binding protein. In some embodiments, the promoter is constitutive. In other embodiments, the promoter responds positively to at least one cytokine or is responsive to one or more transcription factors that are upregulated in expression or activity when cytokines are upregulated. Promoters may respond positively to one or more of the following cytokines: IL-6, TNF- α , IFN- γ , IL-1 β , IL-2, IL-8, or IL-10. In other embodiments, a promoter responsive to a transcription factor that is increased when T cells are activated; NFAT-1 is such a transcription factor, as is NF- κ B.

[0015] In further embodiments, there is an expression construct comprising the nucleic acid molecule encoding an IL-6R α binding protein. Some embodiments concern an expression construct that is a viral vector while others involve an expression construct that is a plasmid.

[0016] Certain embodiments concern one or more cells that contain a nucleic acid construct encoding an IL-6R α binding protein. In some embodiments, the cell is a T cell. In specific embodiments, the cells have been transfected or infected with the expression construct, which may or may not integrate into the genome of one or more host cells. In some cases, there is a T cell comprising a heterologous expression construct encoding an IL-6R α scFv comprising a leader peptide, a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9),

and CDR6 (SEQ ID NO:10), wherein the heterologous linker comprises the amino acid sequence GSTSGSGKPGSGEGSTKG (SEQ ID NO:11). In some embodiments, the IL-6R α binding protein is a single chain antibody fragment (scFv). In some embodiments, the expression construct comprises a cytokine-responsive promoter, meaning the promoter has increased expression when cytokines are increased. In some embodiments, cytokines are increased when T cells are activated. In some embodiments, cytokines are increased when T-cell activation triggers a systemic immune response. In some embodiments, cytokines are increased when tumor-cell lysis triggers a systemic immune response.

[0017] Other embodiments concern methods of making or using the IL-6R α binding proteins, nucleic acid molecules encoding such proteins, or cells expressing such proteins. In certain embodiments, there are methods for reducing the risk of cytokine release syndrome (CRS), methods for inhibiting IL-6R α , methods for treating cancer, methods for treating an autoimmune disease, methods for reducing the toxicity of one or more cytokines, methods for reducing toxicity of T-cell activation, and methods for manufacturing an IL-6R α therapeutic composition. Embodiments also concern use of a chimeric IL-6R α binding protein for the treatment of the cytokine release syndrome, for the treatment of cancer, and/or for the treatment of an autoimmune disease.

[0017A] Some embodiments concern a method for reducing the risk of cytokine release syndrome comprising administering to a patient at risk for cytokine release syndrome a composition comprising the T cells of the invention.

[0017B] Some embodiments concern a method for reducing the toxicity of adoptive cell therapy in a cancer patient comprising administering to the patient the T cells of the invention, wherein the patient has been or will be treated with adoptive cell therapy.

[0017C] Some embodiments concern a method for treating a cancer patient comprising administering to the patient a composition comprising the T cells of the invention.

[0017D] Some embodiments concern the use of T cells of the invention in the manufacture of a medicament for reducing the risk of cytokine release syndrome.

[0017E] Use of T cells of the invention in the manufacture of a medicament for reducing the toxicity of adoptive cell therapy in a cancer patient, wherein the patient has been or will be treated with adoptive cell therapy.

[0017F] Some embodiments concern the use of T cells of the invention in the manufacture of a medicament for treating a cancer patient.

[0018] In some embodiments, methods comprise administering to a patient at risk for cytokine release syndrome a composition comprising an isolated IL-6 receptor alpha (IL-6R α) binding protein or cells comprising a heterologous nucleic acid molecule encoding the IL-6R α binding protein. In some cases, the patient has cancer and/or an autoimmune disease. In some circumstances, the patient has or will receive adoptive T-cell therapy. In some instances, the patient is administered the composition after adoptive T-cell transfer. In some instances, the patient is administered the composition along with adoptive T-cell transfer. In some cases, the patient has or will receive lymphodepletion. The patient may receive or may have received adoptive T cell therapy within 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 10 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 hours or 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 days (or any range derivable within) of receiving a therapeutic composition involving an IL-6R α binding protein. In other embodiments, the patient has an autoimmune disease.

[0019] In some embodiments, the patient is administered T cells comprising a heterologous nucleic acid molecule encoding an IL-6R α binding protein. In further embodiments, the patient is administered T cells that are autologous. 15

[0019A] Some embodiments concern T cells comprising a heterologous nucleic acid molecule encoding a chimeric interleukin 6 receptor alpha (IL-6R α) binding protein comprising a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and CDR6 (SEQ ID NO:10). 20

[0019B] Some embodiments concern T cells comprising a heterologous nucleic acid molecule encoding an anti- IL-6R α scFv comprising a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and CDR6 (SEQ ID NO:10), wherein the heterologous linker comprises the amino acid sequence GSTSGSGKPGSSEGSTKG (SEQ ID NO:11).

[0020] In further embodiments, methods include also administering to the patient an antihistamine, a corticosteroid, a steroid, acetaminophen, furosemide, and/or intravenous fluids or any other treatment used for cytokine release syndrome or other toxicity or side effects associated with T-cell engaging therapies. In some cases, the patient has one or more symptoms of cytokine release syndrome.

[0021] In some embodiments, there are methods for reducing the toxicity of adoptive cell therapy in a cancer patient comprising administering to the patient a composition comprising an isolated IL-6 receptor alpha (IL-6R α) binding protein or cells comprising a heterologous nucleic acid molecule encoding the IL-6R α binding protein. Additional embodiments include methods for treating an autoimmune disease in a patient comprising administering to the patient a composition comprising a composition comprising an isolated IL-6 receptor alpha (IL-6R α) binding protein or cells comprising a heterologous nucleic acid molecule encoding the IL-6R α binding protein. Other embodiments concern methods for treating a cancer patient comprising administering to the patient a composition comprising an isolated IL-6 receptor alpha (IL-6R α) binding protein or cells comprising a heterologous nucleic acid molecule encoding the IL-6R α binding protein, wherein the patient has been or will be treated with immunotherapy. In some embodiments, the patient has been or will be treated with an anti-T cell antibody. Non-limiting examples of anti-T cell antibodies include anti-thymocyte globulin, Muromonab-CD3, TGN1412, anti-CD20, anti-CD19, rituximab, and tisagenlecleucel. In some embodiments, the immunotherapy is adoptive cell therapy. In some embodiments, the methods of the disclosure comprise administering to the patient a composition comprising cells comprising a heterologous nucleic acid molecule encoding the IL-6R α binding protein and a tumor-targeting receptor. Other embodiments concern methods for reducing the toxicity of one or more cytokines comprising administering to a patient with an autoimmune disease or condition or a patient being treated with immunotherapy an

effective amount of a composition comprising a single chain variable fragment comprising the isolated IL-6R α binding protein comprises a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and CDR6 (SEQ ID NO:10).

5 [0022] The polypeptides described herein may include 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 10 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, 100 or more variant amino acids within at least, or at most 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 15 101, 102, 103, 104, 105, 106, 107, 108, 109, 110, 111, 112, 113, 114, 115, 116, 117, 118, 119, 120, 121, 122, 123, 124, 125, 126, 127, 128, 129, 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150, 151, 152, 153, 154, 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, 185, 186, 187, 188, 189, 190, 20 191, 192, 193, 194, 195, 196, 197, 198, 199, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 215, 216, 217, 218, 219, 220, 221, 222, 223, 224, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 300, 400, 500, 550, 1000 or more contiguous amino acids, or any range derivable therein, of any of SEQ ID NOs: 1-11.

25 [0023] In one embodiment of the methods described herein, the subject is a human subject. The terms "individual," "subject," "host," and "patient," used interchangeably herein, refer to a mammal, including, but not limited to, murines (e.g., rats, mice), lagomorphs (e.g., rabbits), non-human primates, humans, canines, felines, ungulates (e.g., equines, bovines, ovines, porcines, caprines), etc.

30 [0024] Before the present invention is further described, it is to be understood that this invention is not limited to particular embodiments described, as such may, of course, vary. It

is also to be understood that the terminology used herein is for the purpose of describing particular embodiments only, and is not intended to be limiting, since the scope of the present invention will be limited only by the appended claims.

[0025] Where a range of values is provided, it is understood that each intervening value, 5 to the tenth of the unit of the lower limit unless the context clearly dictates otherwise, between the upper and lower limit of that range and any other stated or intervening value in that stated range, is encompassed within the invention. The upper and lower limits of these smaller ranges may independently be included in the smaller ranges, and are also encompassed within the invention, subject to any specifically excluded limit in the stated 10 range. Where the stated range includes one or both of the limits, ranges excluding either or both of those included limits are also included in the invention.

[0026] It is specifically contemplated that embodiments described herein may be excluded. It is further contemplated that, when a range is described, certain ranges may be excluded.

[0027] As used herein the specification, “a” or “an” may mean one or more. As used herein 15 in the claim(s), when used in conjunction with the word “comprising”, the words “a” or “an” may mean one or more than one.

[0028] The use of the term “or” in the claims is used to mean “and/or” unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and “and/or.” As used herein 20 “another” may mean at least a second or more.

[0029] Throughout this application, the term “about” is used to indicate that a value 25 includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects. Other features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

[0029A] Throughout this specification the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element, integer or step, or group of elements, integers or steps, but not the exclusion of any other element, integer or step, or group of elements, integers or steps.

BRIEF DESCRIPTION OF THE DRAWINGS

[0030] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better

understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0031] FIG. 1. Anti-IL-6R α scFvs inhibit IL-6 signaling in HepG2 cells. HepG2 human hepatocarcinoma cells were seeded in a 12-well plate and incubated with indicated amounts of scFv for 3 hours. The indicated wells were subsequently treated with 4 ng/ml of IL-6 and incubated for another 30 minutes before cell harvest. The presence of phosphorylated STAT3 (pSTAT3), a major signaling molecule in the IL-6 signaling pathway, was probed by western blot, along with GAPDH (loading control).

[0032] FIG. 2. Anti-IL-6R α scFv inhibits IL-6 signaling in primary human T cells. Primary human CD4+ and CD8+ T cells were seeded in a 12-well plate and treated as described for FIG. 1. The V_L-V_H scFv (SEQ ID NO:1) was added in the indicated amounts, and IL-6 was added to the indicated wells. pSTAT3 and GAPDH (loading control) were detected by western blot.

[0033] FIG. 3. Constitutive expression of anti-IL-6R α scFv significantly reduces the production of inflammatory cytokines by primary human T cells. Primary human CD4+ T cells expressing an anti-CD19 chimeric antigen receptor (CAR) with (“anti-IL-6R α scFv,” middle bars) or without (“No scFv or mAb”; left bars) constitutive expression of the anti-IL-6R α scFv were co-incubated with peripheral blood mononuclear cells (PBMCs) from the same donor plus wild-type (CD19+) Raji lymphoma cells at a 1:1:1 ratio for 12 hours. Non-scFv-expressing CD19 CAR-T cells treated with 90 μ g/ml of the monoclonal antibody tocilizumab were included as control samples (“Tocilizumab”; right bars). Cytokine levels in culture media were quantified by a cytometric bead-based assay. Values shown are the means of three technical replicates with error bars indicating \pm 1 standard deviation. Statistical significance was evaluated by two-tailed, equal-variance Student’s *t*-test; * *p* < 0.05; ** *p* < 0.01; *** *p* < 0.001. The results shown are representative of two independent experiments using cells from two different healthy blood donors.

[0034] FIG. 4. Constitutive expression of anti-IL-6R α scFv does not inhibit T-cell proliferation or cytotoxicity in CD4+ T cells. Primary human CD4+ T cells constitutively expressing the anti-CD19 CAR with or without the anti-IL-6R α scFv were co-incubated with wild-type (WT; CD19+) or CD19- mutant Raji lymphoma cells at a 2:1 effector-to-target

(E:T) ratio. Viable T-cell and Raji-cell counts were obtained by flow cytometry every two days. Both Raji cell lines stably express EGFP to enable separation from T cells during flow-cytometry analysis. After cell counting on day 8, T cells co-incubated with WT Raji were harvested and re-challenged with fresh WT Raji cells at a 2:1 E:T ratio. Viable cell counts 5 were monitored for another 12 days during the second challenge. Values shown are the means of three technical replicates with error bars indicating ± 1 standard deviation. The results shown are representative of four independent experiments using cells from two different healthy blood donors.

[0035] **FIG. 5. Constitutive expression of anti-IL-6R α scFv does not alter T-cell 10 subtype differentiation upon antigen stimulation in CD4+ T cells.** In the same experiment shown in **FIG. 4**, CD4+ primary human T cells were surface-stained with antibody for CCR7, CD45RA, and CD57 to determine the T-cell subtype. T_{E/EM}: CCR7-/CD45RA-/CD57-; T_{E/Exh}: CCR7-/CD45RA-/CD57+; T_{EMRA}: CCR7-/CD45RA+; T_{cm}: CCR7+/CD45RA-; T_{naive}: CCR7+/CD45RA+. Values shown are the means of three technical replicates with 15 error bars indicating ± 1 standard deviation. The results shown are representative of four independent experiments using cells from two different healthy blood donors.

[0036] **FIG. 6. Constitutive expression of anti-IL-6R α scFv does not inhibit T-cell 20 proliferation or cytotoxicity in CD8+ T cells.** Primary human CD8+ T cells constitutively expressing the anti-CD19 CAR with or without the anti-IL-6R α scFv were co-incubated with wild-type (WT; CD19+) or CD19- mutant Raji lymphoma cells at a 2:1 effector-to-target 25 (E:T) ratio. Viable T-cell and Raji-cell counts were obtained by flow cytometry every two days. Both Raji cell lines stably express EGFP to enable separation from T cells during flow-cytometry analysis. Values shown are the means of three technical replicates with error bars indicating ± 1 standard deviation. The results shown are representative of three independent experiments using cells from two different healthy blood donors.

[0037] **FIG. 7. Constitutive expression of anti-IL-6R α scFv does not alter T-cell 30 subtype differentiation upon antigen stimulation in CD8+ T cells.** In the same experiment as described in Figure 1, primary CD8+ human T cells were surface-stained with antibody for CCR7, CD45RA, and CD57 to determine the T-cell subtype. T_{E/EM}: CCR7-/CD45RA-/CD57-; T_{E/Exh}: CCR7-/CD45RA-/CD57+; T_{EMRA}: CCR7-/CD45RA+; T_{cm}: CCR7+/CD45RA-; T_{naive}: CCR7+/CD45RA+. Values shown are the means of three technical replicates with

error bars indicating ± 1 standard deviation. The results shown are representative of four independent experiments using cells from three different healthy blood donors.

[0038] FIG. 8. Constitutive expression of anti-IL-6R α scFv does not alter exhaustion marker expression in primary human T cells upon antigen stimulation.

Primary human T cells constitutively expressing the anti-CD19 CAR with or without anti-IL-6R α scFv were co-incubated with WT Raji cells at a 2:1 E:T ratio. PD-1, TIM-3, and LAG-3 expression levels were quantified by surface antibody staining and flow cytometry every two days. Three trials were performed with CD8+ T cells and four trials were performed with CD4+ T cells for a total of seven independent experiments. Each trial included three technical replicates per condition. Each data point in the plots represents the mean value of triplicates from one trial. The average value of seven trials for each time point is indicated by the black horizontal bar.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0039] Embodiments disclosed herein include polypeptides that bind to IL-6R α , which modulates the inflammatory effects of IL-6. Further embodiments include cells that express IL-6R α -binding polypeptides, as well as methods of using the cells and polypeptides in cancer therapy, autoimmune therapy, and anti-inflammatory therapy, among others. These and other non-limiting aspects of the present invention are discussed in further detail in the following sections.

I. Definitions

[0040] Embodiments disclosed herein may include chimeric antigen receptors (CARs), which may be expressed by T cells or other immune cell types. CARs are engineered receptors, which graft an arbitrary specificity onto an immune effector cell. Typically, these receptors are used to graft the specificity of a monoclonal antibody onto a T cell. The receptors are called chimeric because they are composed of parts from different sources.

[0041] The terms "protein," "polypeptide," and "peptide" are used interchangeably herein when referring to a gene product.

[0042] "Homology," "identity," or "similarity" refers to sequence similarity between two peptides or between two nucleic acid molecules. Identity can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position

in the compared sequence is occupied by the same base or amino acid, then the molecules share sequence identity at that position. A degree of identity between sequences is a function of the number of matching or homologous positions shared by the sequences. An "unrelated" or "non-homologous" sequence shares less than 40% identity, or less than 25% identity, with 5 one of the sequences of the current disclosure.

[0043] The terms "amino portion," "N-terminus," "amino terminus," and the like as used herein are used to refer to order of the regions of the polypeptide. Furthermore, when something is N-terminal to a region it is not necessarily at the terminus (or end) of the entire polypeptide, but just at the terminus of the region or domain. Similarly, the terms "carboxy portion," "C-terminus," "carboxy terminus," and the like as used herein is used to refer to order of the regions of the polypeptide, and when something is C-terminal to a region it is not necessarily at the terminus (or end) of the entire polypeptide, but just at the terminus of the 10 region or domain.

[0044] The terms "subject," "individual," or "patient" are used interchangeably herein and 15 refer to a vertebrate, for example a primate, a mammal or preferably a human. Mammals include, but are not limited to equines, canines, bovines, ovines, murines, rats, simians, humans, farm animals, sport animals and pets.

[0045] The term "xeno-free (XF)" or "animal component-free (ACF)" or "animal free," when used in relation to a medium, an extracellular matrix, or a culture condition, refers to a 20 medium, an extracellular matrix, or a culture condition which is essentially free from heterogeneous animal-derived components. For culturing human cells, any proteins of a non-human animal, such as mouse, would be xeno components. In certain aspects, the xeno-free matrix may be essentially free of any non-human animal-derived components, therefore excluding mouse feeder cells or Matrigel™. Matrigel™ is a solubilized basement 25 membrane preparation extracted from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma, a tumor rich in extracellular matrix proteins to include laminin (a major component), collagen IV, heparin sulfate proteoglycans, and entactin/nidogen. In some embodiments, the compositions described herein or cells of the disclosure are cultured in and/or prepared in/with xeno-free or animal component-free or animal free medium.

30 **[0046]** Cells are "substantially free" of certain reagents or elements, such as serum, signaling inhibitors, animal components or feeder cells, exogenous genetic elements or vector

elements, as used herein, when they have less than 10% of the element(s), and are “essentially free” of certain reagents or elements when they have less than 1% of the element(s). However, even more desirable are cell populations wherein less than 0.5% or less than 0.1% of the total cell population comprise exogenous genetic elements or vector elements.

5 [0047] A culture, matrix or medium are “essentially free” of certain reagents or elements, such as serum, signaling inhibitors, animal components or feeder cells, when the culture, matrix or medium respectively have a level of these reagents lower than a detectable level using conventional detection methods known to a person of ordinary skill in the art or these agents have not been extrinsically added to the culture, matrix or medium. The serum-free medium may be essentially free of serum.

10 [0048] A “gene,” “polynucleotide,” “coding region,” “sequence,” “segment,” “fragment,” or “transgene” which “encodes” a particular protein, is a nucleic acid molecule which is transcribed and optionally also translated into a gene product, *e.g.*, a polypeptide, *in vitro* or 15 *in vivo* when placed under the control of appropriate regulatory sequences. The coding region may be present in either a cDNA, genomic DNA, or RNA form. When present in a DNA form, the nucleic acid molecule may be single-stranded (i.e., the sense strand) or double-stranded. The boundaries of a coding region are determined by a start codon at the 5' (amino) terminus and a translation stop codon at the 3' (carboxy) terminus. A gene can 20 include, but is not limited to, cDNA from prokaryotic or eukaryotic mRNA, genomic DNA sequences from prokaryotic or eukaryotic DNA, and synthetic DNA sequences. A transcription termination sequence will usually be located 3' to the gene sequence.

25 [0049] The term “cell” is herein used in its broadest sense in the art and refers to a living body which is a structural unit of tissue of a multicellular organism, is surrounded by a membrane structure which isolates it from the outside, has the capability of self-replicating, and has genetic information and a mechanism for expressing it. Cells used herein may be naturally-occurring cells or artificially modified cells (*e.g.*, fusion cells, genetically modified cells, etc.).

30 [0050] As used herein, the term “stem cell” refers to a cell capable of self-replication and pluripotency or multipotency. Typically, stem cells can regenerate an injured tissue. Stem

cells herein may be, but are not limited to, embryonic stem (ES) cells, induced pluripotent stem cells or tissue stem cells (also called tissue-specific stem cell, or somatic stem cell).

5 [0051] “Embryonic stem (ES) cells” are pluripotent stem cells derived from early embryos. An ES cell was first established in 1981, which has also been applied to production of knockout mice since 1989. In 1998, a human ES cell was established, which is currently becoming available for regenerative medicine.

10 [0052] Unlike ES cells, tissue stem cells have a limited differentiation potential. Tissue stem cells are present at particular locations in tissues and have an undifferentiated intracellular structure. Therefore, the pluripotency of tissue stem cells is typically low. Tissue stem cells have a higher nucleus/cytoplasm ratio and have few intracellular organelles. Most tissue stem cells have low pluripotency, a long cell cycle, and proliferative ability beyond the life of the individual. Tissue stem cells are separated into categories, based on the sites from which the cells are derived, such as the dermal system, the digestive system, the bone marrow system, the nervous system, and the like. Tissue stem cells in the dermal system include 15 epidermal stem cells, hair follicle stem cells, and the like. Tissue stem cells in the digestive system include pancreatic (common) stem cells, liver stem cells, and the like. Tissue stem cells in the bone marrow system include hematopoietic stem cells, mesenchymal stem cells, and the like. Tissue stem cells in the nervous system include neural stem cells, retinal stem cells, and the like.

20 [0053] “Induced pluripotent stem cells,” commonly abbreviated as iPS cells or iPSCs, refer to a type of pluripotent stem cell artificially prepared from a non-pluripotent cell, typically an adult somatic cell, or terminally differentiated cell, such as fibroblast, a hematopoietic cell, a myocyte, a neuron, an epidermal cell, or the like, by introducing certain factors, referred to as reprogramming factors.

25 [0054] “Pluripotency” refers to a stem cell that has the potential to differentiate into all cells constituting one or more tissues or organs, or particularly, any of the three germ layers: endoderm (interior stomach lining, gastrointestinal tract, the lungs), mesoderm (muscle, bone, blood, urogenital), or ectoderm (epidermal tissues and nervous system). “Pluripotent stem cells” used herein refer to cells that can differentiate into cells derived from any of the three 30 germ layers, for example, direct descendants of totipotent cells or induced pluripotent cells.

[0055] As used herein, the terms "treatment," "treating," and the like, refer to obtaining a desired pharmacologic and/or physiologic effect. The effect may be prophylactic in terms of completely or partially preventing a disease or symptom thereof and/or may be therapeutic in terms of a partial or complete cure for a disease and/or adverse effect attributable to the disease. "Treatment," as used herein, covers any treatment of a disease in a mammal, e.g., in a human, and includes: (a) preventing the disease from occurring in a subject which may be predisposed to the disease but has not yet been diagnosed as having it; (b) inhibiting the disease, i.e., arresting its development; and (c) relieving the disease, i.e., causing regression of the disease.

10 **[0056]** In some embodiments, the methods are useful for reducing the size and/or cell number of a solid tumor. In some embodiments, the method of the disclosure are useful for inhibiting the growth of tumors, such as solid tumors, in a subject.

15 **[0057]** The term "antigen" refers to any substance that causes an immune system to produce antibodies against it, or to which a T cell responds. In some embodiments, an antigen is a peptide that is 5-50 amino acids in length or is at least, at most, or exactly 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 125, 150, 175, 200, 250, or 300 amino acids, or any derivable range therein.

20 **[0058]** The term "antibody" includes monoclonal antibodies, polyclonal antibodies, dimers, multimers, multispecific antibodies and antibody fragments that may be human, mouse, humanized, chimeric, or derived from another species. A "monoclonal antibody" is an antibody obtained from a population of substantially homogeneous antibodies that is being directed against a specific antigenic site.

25 **[0059]** "Antibody or functional fragment thereof" means an immunoglobulin molecule that specifically binds to, or is immunologically reactive with a particular antigen or epitope, and includes both polyclonal and monoclonal antibodies. The term antibody includes genetically engineered or otherwise modified forms of immunoglobulins, such as intrabodies, peptibodies, chimeric antibodies, fully human antibodies, humanized antibodies, and heteroconjugate antibodies (e.g., bispecific antibodies, diabodies, triabodies, and tetrabodies). The term functional antibody fragment includes antigen binding fragments of antibodies, 30 including e.g., Fab', F(ab')₂, Fab, Fv, rlgG, and scFv fragments. The term scFv refers to a

single chain Fv antibody in which the variable domains of the heavy chain and of the light chain of a traditional two chain antibody have been joined to form one chain.

[0060] The use of a single chain variable fragment (scFv) is of particular interest. scFvs are recombinant molecules in which the variable regions of light and heavy immunoglobulin chains encoding antigen-binding domains are engineered into a single polypeptide. Generally, the V_H and V_L sequences are joined by a linker sequence. See, for example, Ahmad (2012) Clinical and Developmental Immunology Article ID 980250, herein specifically incorporated by reference.

[0061] A "therapeutically effective amount" or "efficacious amount" refers to the amount of an agent, or combined amounts of two agents, that, when administered to a mammal or other subject for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the agent(s), the disease and its severity and the age, weight, etc., of the subject to be treated.

[0062] As used herein the specification, "a" or "an" may mean one or more. As used herein in the claim(s), when used in conjunction with the word "comprising", the words "a" or "an" may mean one or more than one.

[0063] The use of the term "or" in the claims is used to mean "and/or" unless explicitly indicated to refer to alternatives only or the alternatives are mutually exclusive, although the disclosure supports a definition that refers to only alternatives and "and/or." As used herein "another" may mean at least a second or more.

[0064] Throughout this application, the term "about" is used to indicate that a value includes the inherent variation of error for the device, the method being employed to determine the value, or the variation that exists among the study subjects.

II. Polypeptides

25 A. IL-6R α -binding Polypeptides

[0065] Some embodiments disclosed herein include polypeptides that bind to IL-6R α , including human IL-6R α . The IL-6R α -binding proteins can be administered in a pharmaceutical formulation or can be expressed from cells that are administered to a subject. In either case, the IL-6R α -binding proteins affect immune processes involving IL-6. When an

IL-6R α -binding protein is expressed by a cell, such as a T cell, the IL-6R α may modulate the immune function of the cell itself and/or other cells in the vicinity. In some embodiments, the IL-6R α -binding protein reduces the secretion of pro-inflammatory cytokines by a cell that expresses the IL-6R α binding protein or by other cells in the vicinity. Such pro-inflammatory cytokines may include TNF- α , IL-6, IL-2, IL-4, GM-CSF, and IL-1 β . The reduction in cytokine secretion by a cell expressing an IL-6R α -binding protein can make the cell more effective and safe for use in therapies, such as adoptive T cell therapy. Cells that express an IL-6R α -binding protein can self-modulate IL-6 signaling levels to prevent and/or ameliorate adverse inflammatory events such as cytokine release syndrome.

[0066] IL-6R α -binding proteins can also be used in treating autoimmune conditions. In some embodiments, this may be accomplished by expressing an IL-6R α -binding protein in a regulatory T cell, which will suppress undesirable inflammatory responses and ameliorate the severity of autoimmune attacks. This may also be accomplished by administering an IL-6R α -binding protein itself to modulate signaling through IL-6R α .

[0067] In some embodiments, the IL-6R α -binding polypeptide is a single chain Fv (scFv), which comprise the V_H and V_L domains of an antibody, wherein these domains are present in a single polypeptide chain. In some embodiments, the antigen-binding domain further comprises a peptide linker between the V_H and V_L domains, which can enable the scFv to form the desired structure for antigen binding. Two different scFv constructs are used in the Examples section below: (1) the V_L-V_H scFv and (2) the V_H-V_L scFv. These two constructs differ only in the order of the sequences. The V_L-V_H scFv configuration is N-terminus—V_L—linker—V_H—C-terminus, and the V_H-V_L scFv configuration is N-terminus—V_H—linker—V_L—C-terminus, where V_L refers to the light chain variable fragment and V_H refers to the heavy chain variable fragment. The amino acid sequences for these scFv constructs are set forth in the table below:

Single Chain Variable Fragment	Amino Acid Sequence (Single Letter Abbreviations)
anti-IL-6R α V _L -V _H scFv	METDTLLWVLLWVPGSTGDGGSDYKDDDDKGGSDIQMTQSP SSLSASVGDRVTITCRASQDISSYLNWYQQKPGKAPKLLIYYTSR LHSGVPSRFSGSGSGTDFFTISSLQPEDIATYYCQQGNTLPYTFG QGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAK VQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTTLSKADYEKH

	KVYACEVTHQGLSSPVTKSFNRGECKSTSGSGKPGSGEGSTKGE VQLQESGPLVRPSQTLSTCTVSGYSITSDDHAWSWVRQPPGRGL EWIGYISYSGITTYNPSLKSRTMLRDTSKNQFSLRLSSVTAADTA VYYCARSLARTTAMDYWGQGSLTVSSASTKGPSVFPLAPSSKS TSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSG LYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKT HTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSH DPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQD WLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDE LTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP G (SEQ ID NO:1)
anti-IL-6R α V _H -V _L scFv	METDTLLLWVLLWVPGSTGDGGSDYKDDDDKGGSEVQLQESG PGLVRPSQTLSTCTVSGYSITSDDHAWSWVRQPPGRGLEWIGYIS YSGITTYNPSLKSRTMLRDTSKNQFSLRLSSVTAADTAVYYCAR SLARTTAMDYWGQGSLTVSSASTKGPSVFPLAPSSKSTSGGTA ALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSSGLYSLSSV VTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDKTHTCPCP APELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSHEDPEVKF NWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDWLNGK EYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQ VSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSDGSFFLY SKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPGGSTSG SGKPGSGEGSTKGDIQMTQSPSSLASVGDRVITCRASQDISSYL NWYQQKPGKAPKLLIYYTSRLHSGVPSRFSGSGSGTDFTFTISSLQ PEDIATYYCQQGNTLPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQL KSGTASVVCLNNFYPREAKVQWKVDNALQSGNSQESVTEQDS KDSTYSLSSTLTLKADYEKHKVYACEVTHQGLSSPVTKSFNRGE C (SEQ ID NO:2)

[0068] The V_H and V_L sequences of the anti-IL-6R α scFvs of SEQ ID NOs:1 and 2, along with the sequences for the CDRs within these sequences and the linker sequence between the V_H and V_L sequences in the scFvs are set forth in the following table:

Sequence name	Amino Acid Sequence (Single Letter Abbreviations)
V _H of the anti-IL-6R α scFvs	EVQLQESGPLVRPSQTLSTCTVSGYSITSDDHAWSWVRQPPGRG LEWIGYISYSGITTYNPSLKSRTMLRDTSKNQFSLRLSSVTAADT AVYYCARSLARTTAMDYWGQGSLTVSSASTKGPSVFPLAPSSK STSGGTAALGCLVKDYFPEPVTVSWNSGALTSGVHTFPAVLQSS GLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKKVEPKSCDK THTCPCPAPELLGGPSVFLFPPKPKDTLMISRTPEVTCVVVDVSH EDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQ DWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRD ELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTPPVLDSD GSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP

	G (SEQ ID NO:3)
V _L of the anti-IL-6R α scFvs	DIQMTQSPSSLSASVGDRVTITCRASQDISSYLNWYQQKPGKAPK LLIYYTSRLHSGVPSRFGSGSGTDFTFTISSLQPEDIAYYCQQGN TLPYTFGQGTKVEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNN FYPREAKVQWKVDNALQSGNSQESVTEQDSKDSTYSLSSTTLS KADYEKHKVYACEVTHQGLSSPVTKSFNRGEC (SEQ ID NO:4)
CDR1 (V _H)	SDHAWS (SEQ ID NO:5)
CDR2 (V _H)	YISYSGITTYNPSLKS (SEQ ID NO:6)
CDR3 (V _H)	SLARTTAMDY (SEQ ID NO:7)
CDR4 (V _L)	RASQDISSYLN (SEQ ID NO:8)
CDR5 (V _L)	YTSRLHS (SEQ ID NO:9)
CDR6 (V _L)	QQGNTLPYT (SEQ ID NO:10)
Linker	GSTSGSGKPGSGEGSTKG (SEQ ID NO:11)

[0069] The scFvs of SEQ ID NOs:1 and 2 are constructed from tocilizumab (WO 1992019759 A1, US 5670373, US 7479543, US 8580264), and previously developed anti-IL-6R α monoclonal antibody, by connecting the V_H domain to the V_L domain with the linker of SEQ ID NO:11.

5 **[0070]** Besides scFvs, the IL-6R α -bindng protein can take a variety of forms that may incorporate one or more of the V_H (SEQ ID NO:3), V_L, or CDR (SEQ ID NOs:5-11) sequences disclosed herein or derivatives or variations thereof. For example, the binding protein can be an antibody-like molecule that has an antigen binding region, including antibody fragments such as Fab', Fab, F(ab')2, F(ab)2, (scFv)2, single domain antibodies (DABs), Fv, and polypeptides with antibody CDRs, scaffolding domains that display the CDRs (e.g., anticalins) or a nanobody. For example, the nanobody can be antigen-specific VHH (e.g., a recombinant VHH) from a camelid IgG2 or IgG3, or a CDR-displaying frame from such camelid Ig. The techniques for preparing and using various antibody-based constructs and fragments are well known in the art. Means for preparing and characterizing antibodies are also well known in the art (See, e.g., Antibodies: A Laboratory Manual, Cold Spring Harbor Laboratory, 1988; incorporated herein by reference).

10 **[0071]** “Mini-antibodies” or “minibodies” are also contemplated for use with embodiments. Minibodies are sFv polypeptide chains which include oligomerization domains at their C-termini, separated from the sFv by a hinge region. Pack et al. (1992). The

oligomerization domain comprises self-associating α -helices, e.g., leucine zippers, that can be further stabilized by additional disulfide bonds. The oligomerization domain is designed to be compatible with vectorial folding across a membrane, a process thought to facilitate in vivo folding of the polypeptide into a functional binding protein. Generally, minibodies are 5 produced using recombinant methods well known in the art. See, e.g., Pack et al. (1992); Cumber et al. (1992).

[0072] Antibody-like binding peptidomimetics are also contemplated in embodiments. Liu et al. (2003) describe “antibody like binding peptidomimetics” (ABiPs), which are peptides that act as pared-down antibodies and have certain advantages of longer serum 10 half-life as well as less cumbersome synthesis methods.

[0073] Alternative scaffolds for antigen binding peptides, such as CDRs are also available and can be used to generate IL-6R α -binding molecules in accordance with the embodiments. Generally, a person skilled in the art knows how to determine the type of protein scaffold on which to graft at least one of the CDRs arising from the original antibody. 15 More particularly, it is known that to be selected such scaffolds must meet the greatest number of criteria as follows (Skerra, 2000): good phylogenetic conservation; known three-dimensional structure (as, for example, by crystallography, NMR spectroscopy or any other technique known to a person skilled in the art); small size; few or no post-transcriptional modifications; and/or easy to produce, express and purify.

[0074] The origin of such protein scaffolds can be, but is not limited to, the structures selected among: fibronectin and preferentially fibronectin type III domain 10, lipocalin, anticalin (Skerra, 2001), protein Z arising from domain B of protein A of *Staphylococcus aureus*, thioredoxin A or proteins with a repeated motif such as the “ankyrin repeat” (Kohl et al., 2003), the “armadillo repeat”, the “leucine-rich repeat” and the “tetrastricopeptide repeat”. For example, anticalins or lipocalin derivatives are a type of binding proteins that have affinities and specificities for various target molecules and can be used as target binding molecules. Such proteins are described in US Patent Publication Nos. 20100285564, 20060058510, 20060088908, 20050106660, and PCT Publication No. WO2006/056464, 25 incorporated herein by reference.

[0075] Scaffolds derived from toxins such as, for example, toxins from scorpions, insects, plants, mollusks, etc., and the protein inhibitors of neuronal NO synthase (PIN) may also be used in certain aspects.

[0076] Monoclonal antibodies (MAbs) are recognized to have certain advantages, 5 e.g., reproducibility and large-scale production. Embodiments include monoclonal antibodies of the human, murine, monkey, rat, hamster, rabbit and chicken origin.

[0077] “Humanized” antibodies are also contemplated, as are chimeric antibodies from mouse, rat, or other species, bearing human constant and/or variable region domains, bispecific antibodies, recombinant and engineered antibodies and fragments thereof. As used 10 herein, the term “humanized” immunoglobulin refers to an immunoglobulin comprising a human framework region and one or more CDR's from a non-human (usually a mouse or rat) immunoglobulin. The non-human immunoglobulin providing the CDR's is called the “donor” and the human immunoglobulin providing the framework is called the “acceptor”. A “humanized antibody” is an antibody comprising a humanized light chain and a humanized 15 heavy chain immunoglobulin. In order to describe antibodies of some embodiments, the strength with which an antibody molecule binds an epitope, known as affinity, can be measured. The affinity of an antibody may be determined by measuring an association constant (K_a) or dissociation constant (K_d). Antibodies deemed useful in certain embodiments may have an association constant of about, at least about, or at most about 10^6 , 10^7 , 10^8 , 10^9 or 20 10^{10} M or any range derivable therein. Similarly, in some embodiments antibodies may have a dissoaciation constant of about, at least about or at most about 10^{-6} , 10^{-7} , 10^{-8} , 10^{-9} or 10^{-10} M or any range derivable therein.

[0078] In certain embodiments, a polypeptide that specifically binds to IL-6R α is able to neutralize IL-6R α and/or prevent binding of IL-6 to IL-6R α or prevent signaling through 25 IL-6R α .

B. Modifications to IL-6R α -binding Polypeptides

1. Signal Peptides

[0079] Any IL-6R α -binding protein can be tagged with the leader (i.e. signal) peptide. A “signal peptide” refers to a peptide sequence that directs the transport and localization of the 30 protein within a cell, e.g. to a certain cell organelle (such as the endoplasmic reticulum)

and/or the cell surface. A signal peptide directs the nascent protein into the endoplasmic reticulum. Generally, the signal peptide natively attached to the amino-terminal most component is used (e.g. in a scFv with orientation light chain - linker - heavy chain, the native signal of the light-chain is used). In some embodiments the signal peptide from the 5 murine kappa light chain is used so that the proteins are secreted from the expressing cells. This also allows the expressed protein to be directly collected from the media in which the producer cells are cultured. Other leader/signal peptides known to those in the art can also be used.

[0080] In some embodiments, the signal peptide is cleaved after passage of the 10 endoplasmic reticulum (ER), i.e., is a cleavable signal peptide. In some embodiments, a restriction site is at the carboxy end of the signal peptide to facilitate cleavage.

2. Markers and Epitope Tags

[0081] The IL-6R α -binding proteins can also be tagged with markers such as the DYKDDDDK (FLAG) epitope, HA tag, or cMyc tag or other epitope tags, markers, and 15 affinity tags known in the art. This can facilitate isolation, purification, localization, and targeted binding of the IL-6R α -binding protein. In some embodiments, the marker is flanked by GGS linkers.

3. Linkers

[0082] In some embodiments, the polypeptides of the disclosure include peptide linkers 20 (sometimes referred to as a linker). A peptide linker may be separating any of the peptide domain/regions described herein. As an example, a linker may be between the signal peptide and the antigen binding domain, between a V_H and V_L of an antigen binding domain, between an antigen binding domain and a peptide spacer, between a peptide spacer and a transmembrane domain. A peptide linker may have any of a variety of amino acid sequences. 25 Domains and regions can be joined by a peptide linker that is generally of a flexible nature, although other chemical linkages are not excluded. A linker can be a peptide of between about 6 and about 40 amino acids in length, or between about 6 and about 25 amino acids in length. These linkers can be produced by using synthetic, linker-encoding oligonucleotides to couple the proteins.

[0083] Peptide linkers with a degree of flexibility can be used. The peptide linkers may have virtually any amino acid sequence, bearing in mind that suitable peptide linkers will have a sequence that results in a generally flexible peptide. The use of small amino acids, such as glycine and alanine, are of use in creating a flexible peptide. The creation of such sequences is routine to those of skill in the art.

[0084] Suitable linkers can be readily selected and can be of any of a suitable of different lengths, such as from 1 amino acid (e.g., Gly) to 20 amino acids, from 2 amino acids to 15 amino acids, from 3 amino acids to 12 amino acids, including 4 amino acids to 10 amino acids, 5 amino acids to 9 amino acids, 6 amino acids to 8 amino acids, or 7 amino acids to 8 amino acids, and may be 1, 2, 3, 4, 5, 6, or 7 amino acids.

[0085] Exemplary flexible linkers include glycine polymers $(G)_n$, glycine- serine polymers (including, for example, $(GS)_n$, $(GSGS)_n$ and $(GGGS)_n$, where n is an integer of at least one, glycine-alanine polymers, alanine-serine polymers, and other flexible linkers known in the art. Glycine and glycine-serine polymers can be used; both Gly and Ser are relatively unstructured, and therefore can serve as a neutral tether between components. Glycine polymers can be used; glycine accesses significantly more phi-psi space than even alanine, and is much less restricted than residues with longer side chains. Exemplary spacers can comprise amino acid sequences including, but not limited to, GGSG, GGSGG, GSGSG, GSAGGG, GGGSG, GSSSG, and the like.

20 4. Modifications to Antigen-binding Domains

[0086] The variable regions of antigen-binding domains of IL-6R α -binding proteins of the disclosure can be modified by mutating amino acid residues within V_H and/or V_L CDR regions to improve one or more binding properties (e.g., affinity) of the protein. The term “CDR” refers to a complementarity-determining region that is based on a part of the variable chains in immunoglobulins (antibodies) and T-cell receptors, generated by B cells and T cells respectively, where these molecules bind to their specific antigen. Since most sequence variation associated with immunoglobulins and T-cell receptors are found in the CDRs, these regions are sometimes referred to as hypervariable regions. Mutations may be introduced by site-directed mutagenesis or PCR-mediated mutagenesis and the effect on antibody binding, or other functional property of interest, can be evaluated in appropriate *in vitro* or *in vivo* assays. Preferably conservative modifications are introduced and typically no more than one,

two, three, four or five residues within a CDR region are altered. The mutations may be amino acid substitutions, additions or deletions.

5 [0087] Framework modifications can be made to IL-6R α -binding proteins to decrease immunogenicity, for example, by "backmutating" one or more framework residues to a corresponding germline sequence.

[0088] It is also contemplated that the antigen binding domain may be multi-specific or multivalent by multimerizing the antigen binding domain with V_H and V_L region pairs that bind either the same antigen (multi-valent) or a different antigen (multi-specific).

10 [0089] As used herein, the term "affinity" refers to the equilibrium constant for the reversible binding of two agents and is expressed as a dissociation constant (K_d). Affinity can be at least 1-fold greater, at least 2-fold greater, at least 3-fold greater, at least 4-fold greater, at least 5-fold greater, at least 6-fold greater, at least 7-fold greater, at least 8-fold greater, at least 9-fold greater, at least 10-fold greater, at least 20-fold greater, at least 30-fold greater, at least 40-fold greater, at least 50-fold greater, at least 60-fold greater, at least 70-fold greater, 15 at least 80-fold greater, at least 90-fold greater, at least 100-fold greater, or at least 1000-fold greater, or more (or any derivable range therein), than the affinity of an antibody for unrelated amino acid sequences. As used herein, the term "avidity" refers to the resistance of a complex of two or more agents to dissociation after dilution. The terms "immunoreactive" and "preferentially binds" are used interchangeably herein with respect to antibodies and/or 20 antigen-binding fragments.

[0090] The term "binding" refers to a direct association between two molecules, due to, for example, covalent, electrostatic, hydrophobic, and ionic and/or hydrogen-bond interactions, including interactions such as salt bridges and water bridges.

5. Chemical Modifications

25 [0091] Additionally, the polypeptides of the disclosure may be chemically modified. Glycosylation of the polypeptides can be altered, for example, by modifying one or more sites of glycosylation within the polypeptide sequence to increase the affinity of the polypeptide for antigen (U.S. Pat. Nos. 5,714,350 and 6,350,861).

[0092] The polypeptides of the invention can be pegylated to increase biological half-life by reacting the polypeptide with polyethylene glycol (PEG) or a reactive ester or aldehyde derivative of PEG, under conditions in which one or more PEG groups become attached to the polypeptide. Polypeptide pegylation may be carried out by an acylation reaction or an 5 alkylation reaction with a reactive PEG molecule (or an analogous reactive watersoluble polymer). As used herein, the term "polyethylene glycol" is intended to encompass any of the forms of PEG that have been used to derivatize other proteins, such as mono (C1-C10) alkoxy- or aryloxy-polyethylene glycol or polyethylene glycol-maleimide. Methods for pegylating proteins are known in the art and can be applied to the polypeptides of the 10 invention (EP 0 154 316 and EP 0 401 384).

[0093] Additionally, polypeptides may be chemically modified by conjugating or fusing the polypeptide to serum protein, such as human serum albumin, to increase half-life of the resulting molecule. Such approach is for example described in EP 0322094 and EP 0 486 525.

[0094] The polypeptides of the disclosure may be conjugated to a diagnostic or therapeutic agent and used diagnostically, for example, to monitor the development or progression of a disease and determine the efficacy of a given treatment regimen. The polypeptides may also be conjugated to a therapeutic agent to provide a therapy in combination with the therapeutic effect of the polypeptide. Examples of diagnostic agents 15 include enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, radioactive materials, positron emitting metals using various positron emission tomographies, and nonradioactive paramagnetic metal ions. The detectable substance may be coupled or conjugated either directly to the polypeptide, or indirectly, through a linker using techniques known in the art. Examples of suitable enzymes include 20 horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase. Examples of suitable prosthetic group complexes include streptavidin/ biotin and avidin/biotin. Examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate, rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin. An example of a luminescent material includes luminol. Examples of 25 suitable bioluminescent materials include luciferase, luciferin, and aequorin. Examples of suitable radioactive material include ¹²⁵I, ¹³¹I, Indium-111, Lutetium-171, Bismuth-212, 30

Bismuth-213, Astatine-211, Copper-62, Copper-64, Copper-67, Yttrium-90, Iodine-125, Iodine-131, Phosphorus-32, Phosphorus-33, Scandium-47, Silver-111, Gallium-67, Praseodymium-142, Samarium-153, Terbium-161, Dysprosium-166, Holmium-166, Rhenium-186, Ithenium-188, Rhenium-189, Lead-212, Radium-223, Actinium-225, Iron-59, 5 Selenium-75, Arsenic-77, Strontium-89, Molybdenum-99, Rhodium-1105, Palladium-109, Praseodymium-143, Promethium-149, Erbium-169, Iridium-194, Gold-198, Gold-199, and Lead-211. Chelating agents may be attached through amities (Meares et al., 1984 Anal. Biochem. 142: 68-78); sulphydral groups (Koyama 1994 Chem. Abstr. 120: 217262t) of amino acid residues and carbohydrate groups (Rodwell et al. 1986 PNAS USA 83: 10 2632-2636; Quadri et al. 1993 Nucl. Med. Biol. 20: 559-570).

[0095] The polypeptides may also be conjugated to a therapeutic agent to provide a therapy in combination with the therapeutic effect of the polypeptide.

[0096] Additional suitable conjugated molecules include ribonuclease (RNase), DNase I, an antisense nucleic acid, an inhibitory RNA molecule such as a siRNA molecule, an 15 immunostimulatory nucleic acid, aptamers, ribozymes, triplex forming molecules, and external guide sequences. Aptamers are small nucleic acids ranging from 15-50 bases in length that fold into defined secondary and tertiary structures, such as stern-loops or G-quartets, and can bind small molecules, such as ATP (U.S. Pat. No. 5,631,146) and theophiline (U.S. Pat. No. 5,580,737), as well as large molecules, such as reverse 20 transcriptase (U.S. Pat. No. 5,786,462) and thrombin (U.S. Pat. No. 5,543,293). Ribozymes are nucleic acid molecules that are capable of catalyzing a chemical reaction, either intramolecularly or intermolecularly. Ribozymes typically cleave nucleic acid substrates through recognition and binding of the target substrate with subsequent cleavage. Triplex forming function nucleic acid molecules can interact with double-stranded or single-stranded 25 nucleic acid by forming a triplex, in which three strands of DNA form a complex dependant on both Watson-Crick and Hoogsteen base-pairing. Triplex molecules can bind target regions with high affinity and specificity.

[0097] The functional nucleic acid molecules may act as effectors, inhibitors, modulators, and stimulators of a specific activity possessed by a target molecule, or the functional nucleic 30 acid molecules may possess a de novo activity independent of any other molecules.

C. Expression of IL-6Ra-binding Proteins

[0098] IL-6Ra-binding proteins can be expressed in a variety of cell types. An expression construct encoding an IL-6Ra-binding protein can be transfected into cells according to a variety of methods known in the art. In some embodiments, the IL-6Ra-binding protein expression construct can be under control of a constitutive promoter. In some embodiments, it can be under control of a regulatable promoter that drives expression only under certain conditions. This can provide for expression of a IL-6Ra-binding protein only when necessary to respond to a potentially harmful situation or a situation in which it is desirable to reduce signaling through IL-6Ra, such as, for example, when there are increased levels of cytokines present. To accomplish this, the IL-6Ra-binding protein expression construct may be placed under control of a promoter that is responsive to cytokines such as IL-6, TNF- α , IFN- γ , IL-1 β , IL-2, IL-8, and IL-10. The IL-6Ra-binding protein expression construct can also be placed under control of a promoter that is linked to T-cell activation, such as one that is controlled by NFAT-1 or NF- κ B, both of which are transcription factors that can be activated upon T-cell activation. Control of IL-6Ra-binding protein expression allows T cells, such as tumor-targeting T cells, to sense their surroundings and perform real-time modulation of cytokine signaling, both in the T cells themselves and in surrounding endogenous immune cells.

III. Nucleic Acids

[0099] In certain embodiments, there are recombinant polynucleotides encoding the proteins, polypeptides, or peptides described herein. Polynucleotide sequences contemplated include those encoding IL-6Ra-binding proteins and CARs.

[0100] As used in this application, the term “polynucleotide” refers to a nucleic acid molecule that either is recombinant or has been isolated free of total genomic nucleic acid. Included within the term “polynucleotide” are oligonucleotides (nucleic acids 100 residues or less in length), recombinant vectors, including, for example, plasmids, cosmids, phage, viruses, and the like. Polynucleotides include, in certain aspects, regulatory sequences, isolated substantially away from their naturally occurring genes or protein encoding sequences. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be RNA, DNA (genomic, cDNA or synthetic), analogs thereof, or a combination

thereof. Additional coding or non-coding sequences may, but need not, be present within a polynucleotide.

[0101] In this respect, the term “gene,” “polynucleotide,” or “nucleic acid” is used to refer to a nucleic acid that encodes a protein, polypeptide, or peptide (including any

5 sequences required for proper transcription, post-translational modification, or localization).

As will be understood by those in the art, this term encompasses genomic sequences, expression cassettes, cDNA sequences, and smaller engineered nucleic acid segments that express, or may be adapted to express, proteins, polypeptides, domains, peptides, fusion proteins, and mutants. A nucleic acid encoding all or part of a polypeptide may contain a

10 contiguous nucleic acid sequence encoding all or a portion of such a polypeptide. It also is contemplated that a particular polypeptide may be encoded by nucleic acids containing

variations having slightly different nucleic acid sequences but, nonetheless, encode the same or substantially similar protein.

[0102] In particular embodiments, there are isolated nucleic acid segments and

15 recombinant vectors incorporating nucleic acid sequences that encode a polypeptide (e.g., an

antibody or fragment thereof) that binds to IL-6Ra. The term “recombinant” may be used in conjunction with a polypeptide or the name of a specific polypeptide, and this generally refers to a polypeptide produced from a nucleic acid molecule that has been manipulated in vitro or that is a replication product of such a molecule.

20 **[0103]** The nucleic acid segments, regardless of the length of the coding sequence itself,

may be combined with other nucleic acid sequences, such as promoters, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that

25 a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant

nucleic acid protocol. In some cases, a nucleic acid sequence may encode a polypeptide sequence with additional heterologous coding sequences, for example to allow for purification of the polypeptide, transport, secretion, post-translational modification, or for therapeutic benefits such as targeting or efficacy. As discussed above, a tag or other

30 heterologous polypeptide may be added to the modified polypeptide-encoding sequence,

wherein “heterologous” refers to a polypeptide that is not the same as the modified polypeptide.

[0104] In certain embodiments, there are polynucleotide variants having substantial identity to the sequences disclosed herein; those comprising at least 70%, 75%, 80%, 85%, 5 90%, 95%, 96%, 97%, 98%, or 99% or higher sequence identity, including all values and ranges there between, compared to a polynucleotide sequence provided herein using the methods described herein (e.g., BLAST analysis using standard parameters). In certain aspects, the isolated polynucleotide will comprise a nucleotide sequence encoding a polypeptide that has at least 90%, preferably 95% and above, identity to an amino acid 10 sequence described herein, over the entire length of the sequence; or a nucleotide sequence complementary to said isolated polynucleotide.

A. Vectors

[0105] Polypeptides may be encoded by a nucleic acid molecule. The nucleic acid molecule can be in the form of a nucleic acid vector. The term “vector” is used to refer to a 15 carrier nucleic acid molecule into which a heterologous nucleic acid sequence can be inserted for introduction into a cell where it can be replicated and expressed. A nucleic acid sequence can be “heterologous,” which means that it is in a context foreign to the cell in which the vector is being introduced or to the nucleic acid in which it is incorporated, which includes a sequence homologous to a sequence in the cell or nucleic acid but in a position within the 20 host cell or nucleic acid where it is ordinarily not found. Vectors include DNAs, RNAs, plasmids, cosmids, viruses (bacteriophage, animal viruses, and plant viruses), and artificial chromosomes (e.g., YACs). One of skill in the art would be well equipped to construct a vector through standard recombinant techniques (for example Sambrook et al., 2001; Ausubel et al., 1996, both incorporated herein by reference). Vectors may be used in a host cell to 25 produce an antibody that binds IL-6Ra.

[0106] The term “expression vector” refers to a vector containing a nucleic acid sequence coding for at least part of a gene product capable of being transcribed. In some cases, RNA molecules are then translated into a protein, polypeptide, or peptide. Expression vectors can contain a variety of “control sequences,” which refer to nucleic acid sequences necessary for 30 the transcription and possibly translation of an operably linked coding sequence in a

particular host organism. In addition to control sequences that govern transcription and translation, vectors and expression vectors may contain nucleic acid sequences that serve other functions as well and are described herein.

[0107] A “promoter” is a control sequence. The promoter is typically a region of a nucleic acid sequence at which initiation and rate of transcription are controlled. It may contain genetic elements at which regulatory proteins and molecules may bind such as RNA polymerase and other transcription factors. The phrases “operatively positioned,” “operatively linked,” “under control,” and “under transcriptional control” mean that a promoter is in a correct functional location and/or orientation in relation to a nucleic acid sequence to control transcriptional initiation and expression of that sequence. A promoter may or may not be used in conjunction with an “enhancer,” which refers to a *cis*-acting regulatory sequence involved in the transcriptional activation of a nucleic acid sequence.

[0108] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon or adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals.

[0109] Vectors can include a multiple cloning site (MCS), which is a nucleic acid region that contains multiple restriction enzyme sites, any of which can be used in conjunction with standard recombinant technology to digest the vector. (See Carbonelli et al., 1999, Levenson et al., 1998, and Cocea, 1997, incorporated herein by reference.)

[0110] Most transcribed eukaryotic RNA molecules will undergo RNA splicing to remove introns from the primary transcripts. Vectors containing genomic eukaryotic sequences may require donor and/or acceptor splicing sites to ensure proper processing of the transcript for protein expression. (See Chandler et al., 1997, incorporated herein by reference.)

[0111] The vectors or constructs will generally comprise at least one termination signal. A “termination signal” or “terminator” is comprised of the DNA sequences involved in specific termination of an RNA transcript by an RNA polymerase. Thus, in certain embodiments a termination signal that ends the production of an RNA transcript is

contemplated. A terminator may be necessary *in vivo* to achieve desirable message levels. In eukaryotic systems, the terminator region may also comprise specific DNA sequences that permit site-specific cleavage of the new transcript so as to expose a polyadenylation site. This signals a specialized endogenous polymerase to add a stretch of about 200 A residues (polyA)

5 to the 3' end of the transcript. RNA molecules modified with this polyA tail appear to more stable and are translated more efficiently. Thus, in other embodiments involving eukaryotes, it is preferred that that terminator comprises a signal for the cleavage of the RNA, and it is more preferred that the terminator signal promotes polyadenylation of the message.

10 [0112] In expression, particularly eukaryotic expression, one will typically include a polyadenylation signal to effect proper polyadenylation of the transcript.

[0113] In order to propagate a vector in a host cell, it may contain one or more origins of replication sites (often termed “ori”), which is a specific nucleic acid sequence at which replication is initiated. Alternatively, an autonomously replicating sequence (ARS) can be employed if the host cell is yeast.

15 **B. Host Cells**

[0114] As used herein, the terms “cell,” “cell line,” and “cell culture” may be used interchangeably. All of these terms also include their progeny, which is any and all subsequent generations. It is understood that all progeny may not be identical due to deliberate or inadvertent mutations. In the context of expressing a heterologous nucleic acid 20 sequence, “host cell” refers to a prokaryotic or eukaryotic cell, and it includes any transformable organism that is capable of replicating a vector or expressing a heterologous gene encoded by a vector. A host cell can, and has been, used as a recipient for vectors or viruses. A host cell may be “transfected” or “transformed,” which refers to a process by which exogenous nucleic acid, such as a recombinant protein-encoding sequence, is 25 transferred or introduced into the host cell. A transformed cell includes the primary subject cell and its progeny.

[0115] Some vectors may employ control sequences that allow it to be replicated and/or expressed in both prokaryotic and eukaryotic cells. One of skill in the art would further understand the conditions under which to incubate all of the above described host cells to 30 maintain them and to permit replication of a vector. Also understood and known are

techniques and conditions that would allow large-scale production of vectors, as well as production of the nucleic acids encoded by vectors and their cognate polypeptides, proteins, or peptides.

C. Expression Systems

5 [0116] Numerous expression systems exist that comprise at least a part or all of the compositions discussed above. Prokaryote- and/or eukaryote-based systems can be employed for use with an embodiment to produce nucleic acid sequences, or their cognate polypeptides, proteins and peptides. Many such systems are commercially and widely available.

10 [0117] The insect cell/baculovirus system can produce a high level of protein expression of a heterologous nucleic acid segment, such as described in U.S. Patents 5,871,986, 4,879,236, both herein incorporated by reference, and which can be bought, for example, under the name MAXBAC® 2.0 from INVITROGEN® and BACPACK™ BACULOVIRUS EXPRESSION SYSTEM FROM CLONTECH®.

15 [0118] In addition to the disclosed expression systems, other examples of expression systems include STRATAGENE®'s COMPLETE CONTROL™ Inducible Mammalian Expression System, which involves a synthetic ecdysone-inducible receptor, or its pET Expression System, an E. coli expression system. Another example of an inducible expression system is available from INVITROGEN®, which carries the T-REX™ (tetracycline-regulated expression) System, an inducible mammalian expression system that 20 uses the full-length CMV promoter. INVITROGEN® also provides a yeast expression system called the Pichia methanolica Expression System, which is designed for high-level production of recombinant proteins in the methylotrophic yeast Pichia methanolica. One of skill in the art would know how to express a vector, such as an expression construct, to produce a nucleic acid sequence or its cognate polypeptide, protein, or peptide.

25 D. Methods of Gene Transfer

30 [0119] Suitable methods for nucleic acid delivery to effect expression of compositions are believed to include virtually any method by which a nucleic acid (e.g., DNA, including viral and nonviral vectors) can be introduced into a cell, a tissue or an organism, as described herein or as would be known to one of ordinary skill in the art. Such methods include, but are not limited to, direct delivery of DNA such as by injection (U.S. Patents 5,994,624,

5,981,274, 5,945,100, 5,780,448, 5,736,524, 5,702,932, 5,656,610, 5,589,466 and 5,580,859, each incorporated herein by reference), including microinjection (Harland and Weintraub, 1985; U.S. Patent 5,789,215, incorporated herein by reference); by electroporation (U.S. Patent No. 5,384,253, incorporated herein by reference); by calcium phosphate precipitation (Graham and Van Der Eb, 1973; Chen and Okayama, 1987; Rippe et al., 1990); by using DEAE dextran followed by polyethylene glycol (Gopal, 1985); by direct sonic loading (Fechheimer et al., 1987); by liposome mediated transfection (Nicolau and Sene, 1982; Fraley et al., 1979; Nicolau et al., 1987; Wong et al., 1980; Kaneda et al., 1989; Kato et al., 1991); by microprojectile bombardment (PCT Application Nos. WO 94/09699 and 10 95/06128; U.S. Patents 5,610,042; 5,322,783, 5,563,055, 5,550,318, 5,538,877 and 5,538,880, and each incorporated herein by reference); by agitation with silicon carbide fibers (Kaeppeler et al., 1990; U.S. Patents 5,302,523 and 5,464,765, each incorporated herein by reference); by Agrobacterium mediated transformation (U.S. Patents 5,591,616 and 5,563,055, each incorporated herein by reference); or by PEG mediated transformation of 15 protoplasts (Omirulleh et al., 1993; U.S. Patents 4,684,611 and 4,952,500, each incorporated herein by reference); by desiccation/inhibition mediated DNA uptake (Potrykus et al., 1985). Through the application of techniques such as these, organelle(s), cell(s), tissue(s) or organism(s) may be stably or transiently transformed. Other methods include viral transduction, such as gene transfer by lentiviral or retroviral transduction.

20 E. IL-6Ra-binding Protein Expression Construct

[0120] In some embodiments, an IL-6Ra-binding protein is encoded by an expression construct that has the following structure: murine 5' – IgG kappa leader sequence – FLAG tag – V_L – scFv linker – V_H – 3'. The following DNA sequence encodes such a construct (underlined sequences are the murine IgG kappa leader and the scFv linker sequences):

25 ATGGAGACAGACACACTCCTGCTATGGGTGCTGCTCTGGGTTCCAGGTTCCA
CAGGTGACGGAGGCAGCGACTATAAAGATGACGATGACAAAGGAGGTAGCGAT
ATACAAATGACTCAATCACCTCCTCACTTCCGCCAGCGTGGGTGATCGAGTGA
CCATCACATGCCCGCTTCACAGGACATCTCCTCTTACCTGAACCTGGTACCAAGCA
GAAACCCGGAAAGCCCCAAAGCTGTTGATCTACTATACATCAAGACTGCACTCT
30 GGCGTCCCCTCTAGGTTCAGTGGATCTGGCTCAGGTACTGACTTCACGTTCACTA
TTTCCAGCCTGCAGCCGGAAGATATTGCTACCTACTATTGTCAGCAGGGAAATAC

CCTCCCATATACCTTGGGCAGGGACAAAGGTGGAAATCAAGCGGACGGTAGC
TGCTCCGTCAGTGTTCATCTTCCACCTCAGATGAGCAGCTGAAGTCCGGAACC
GCCAGTGTGGTGTGCCTCCTGAATAACTTTATCCACGAGAAGCCAAGGTGCAGT
GGAAAGTGGATAACGCTTGCAAAGTGGCAATTCTCAAGAGAGCGTAAGTGAGC
5 AAGATAGCAAGGACAGTACTTACAGCTTGAGCAGCACGCTGACCCCTGAGCAAGG
CCGATTACGAGAACACAAGGTGTATGCCTGCGAGGTGACCCACCAGGGCCTTA
GCTCACCTGTGACCAAGTCCTCAACAGAGGCGAGTGTGGCAGCACCTCCGGCA
GGCGCAAGCCTGGCAGCGCGAGGGCAGCACCAAGGGCGAGGTCCAAGTGCAG
GAAAGCGGTCCAGGCTTGGTGAGACCATCCCAGACCCTGAGCCTCACCTGTACC
10 GTGTCCGGGTACAGTATCACCTCCGACCATGCATGGTCCTGGGTGCGCCAGCCTC
CCGGAAGAGGTCTGGAATGGATCGGGTATATCTCCTACTCAGGAATCACCACCTA
CAATCCCTCCCTTAAGTCAAGGGTGAATGCTCCGCGATACATCCAAAATCAG
TTCTCCCTCGGTTGTCAAGTGTACAGTGTACAGCCGCCGACACCGCAGTCTACTACTGTGC
AAGGAGCCTGCCAGGACGACTGCAATGGATTATTGGGGCCAGGGCTCCCTGGT
15 GACTGTCAGCAGCGCCTCAACAAAGGGCCATCTGTTTCCACTGCCCAAGC
TCTAAAAGTACCTCAGGAGGAACCGCTGCCCTGGGTGCCTCGTCAAAGATTACT
TTCCAGAACCCGTTACAGTGTCCCTGGAACCTCCGGAGCTCTTACCTCCGGGTGCA
TACATTCCCGGCCGTTCTCAAAGCTCTGGACTCTACTCCCTGAGCTCCGTTGTGA
CAGTTCTAGCAGCAGCCTCGGCACCCAGACATACATCTGCAACGTAAACCACA
20 AGCCTCAAACACTAAAGTGACAAAAAAAGTGGAACCGAACGTCTGTGACAAAA
CTCACACATGCCCTCCGTGCCCTGCCAGAGCTGCTGGCGGGCCATCCGTATT
TCTCTCCCTCCAAAGCAAAGGATACCCCTCATGATCAGCCGGACGCCGAAGTG
ACCTGCGTCGTAGTGGATGTGAGTCACGAGGACCCAGAGGTAAAGTTCAACTGG
TACGTAGACGGCGTGGAGGTCCATAATGCGAAGACGAAACCCAGAGAGGGAGCA
25 GTACAACCTCAACTTACAGAGTGGTCAGCGTGCTCACTGTCCTCCACCAGGACTGG
TTGAACGGCAAAGAATATAAGTGTAAAGGTGAGCAATAAGGCCCTCCCCGCCCT
ATCGAGAAAATTTCCAAGGCTAAAGGCCAGCCTCGAGAACCTCAGGTTAC
ACACTGCCCTCTAGCCGCGATGAACCTGACAAAAAATCAGGTGAGTCTCACCTGC
CTCGTTAAAGGTTCTACCCTCAGATATCGCAGTCGAGTGGAGAGCAACGGGC
30 AGCCGGAAAATAACTATAAGACCACCCCCCCCAGTCCTGACAGTGACGGCTTT
CTTCCTGTACTCCAAACTGACCGTGGACAAATCTAGGTGGCAGCAAGGAAATGTT

TTTAGTTCTGTAATGCACGAGGCTCTGCATAATCATTATAACCCAAAAGTCTTT
GTCCCTCTCCCCGGC (SEQ ID NO:89)

IV. Cells

A. Cell Types

5 [0121] Certain embodiments relate to cells comprising polypeptides or nucleic acids of the disclosure. In some embodiments the cell is an immune cell or a T cell. "T cell" includes all types of immune cells expressing CD3 including T-helper cells (CD4⁺ cells), cytotoxic T-cells (CD8⁺ cells), T-regulatory cells (T_{reg}), suppressor T cells, and gamma-delta T cells. A "cytotoxic cell" includes CD8⁺ T cells, natural-killer (NK) cells, and neutrophils, which cells 10 are capable of mediating cytotoxicity responses. In some embodiments, the cell is a progenitor cell or stem cell. In some embodiments, the progenitor or stem cell is *in vitro* differentiated into an immune cell. In some embodiments, the cell is *ex vivo*. The term immune cells includes cells of the immune system that are involved in defending the body against both infectious disease and foreign materials. Immune cells may include, for example, 15 neutrophils, eosinophils, basophils, natural killer cells, lymphocytes such as B cells and T cells, and monocytes.

[0122] Suitable mammalian cells include primary cells and immortalized cell lines. Suitable mammalian cell lines include human cell lines, non-human primate cell lines, rodent (e.g., mouse, rat) cell lines, and the like. Suitable mammalian cell lines include, but are not 20 limited to, HeLa cells (e.g., American Type Culture Collection (ATCC) No. CCL-2), CHO cells (e.g., ATCC Nos. CRL9618, CCL61, CRL9096), human embryonic kidney (HEK) 293 cells (e.g., ATCC No. CRL-1573), Vero cells, NIH 3T3 cells (e.g., ATCC No. CRL-1658), Huh-7 cells, BHK cells (e.g., ATCC No. CCL10), PC12 cells (ATCC No. CRL1721), COS cells, COS-7 cells (ATCC No. CRL1651), RATI cells, mouse L cells (ATCC No. CCL1.3), 25 HLHepG2 cells, Hut-78, Jurkat, HL-60, NK cell lines (e.g., NKL, NK92, and YTS), and the like.

[0123] In some instances, the cell is not an immortalized cell line, but is instead a cell (e.g., a primary cell) obtained from an individual. For example, in some cases, the cell is an 30 immune cell obtained from an individual. As an example, the cell is a T lymphocyte obtained from an individual. As another example, the cell is a cytotoxic cell obtained from an

individual. As another example, the cell is a stem cell or progenitor cell obtained from an individual.

B. Adoptive Cell Therapy

[0124] In some embodiments, cells disclosed herein are used in adoptive cell therapy or 5 adoptive cell transfer, which involves administration of therapeutic cells to subjects. This therapy can involve administration of tumor-infiltrating lymphocytes, cytotoxic T lymphocytes (CTLs), helper T (Th) cells, and Treg cells, and other immune cells and other types of cells. The therapeutic cells may originate from a patient or from a donor. In some embodiments, cells are extracted from a patient, genetically modified and cultured *in vitro* 10 and returned to the same patient.

[0125] Cells disclosed herein can be administered to a subject to treat, for example, cancer, autoimmune diseases, infections, and inflammatory disorders. In some embodiments, expression of IL-6R α binding proteins by cells used in adoptive T cell therapy can reduce or 15 eliminate side effects that may occur as a result of adoptive T cell therapy, such as cytokine release syndrome and other inflammatory or autoimmune conditions that involve IL-6 signaling through IL-6R α . In addition, administration of IL-6R α -binding proteins in conjunction with therapeutic T cells can eliminate, reduce, or mitigate the side effects of 20 adoptive T cell therapy.

1. Chimeric Antigen Receptors

20 a. Use in Adoptive Cell Therapies

[0126] In some embodiments, therapeutic cells used in adoptive cell therapies express 25 chimeric antigen receptors (CARs). CARs are fusion proteins that are commonly composed of an extracellular antigen-binding domain (which may be an scFv), an extracellular spacer, a transmembrane domain, costimulatory signaling regions (the number of which varies depending on the specific CAR design), and a CD3-zeta signaling domain/endodomain. Immune cells, including T cells and natural killer (NK) cells, can be engineered to express 30 CARs by a variety of methods known in the art, including viral transduction, DNA nucleofection, and RNA nucleofection. CAR binding to the antigen target can activate human T cells expressing the CAR, which may result in killing of the cell bearing the antigen or some other immunological response.

[0127] In some embodiments, the cells comprise a cancer-specific CAR. The term “cancer-specific” in the context of CARs refers to CARs that have an antigen binding specificity for a cancer-specific molecule, such as a cancer-specific antigen. In some embodiments, the cancer specific CAR is in a cell with a CAR specific for another target, 5 such as a TGF- β CAR. In some embodiments, the cancer-specific CAR and another CAR are on separate polypeptides. In some embodiments, the CAR is a bi-specific CAR that has antigen binding for a cancer-specific molecule and for another antigen, such as TGF- β . For example, a bi-specific CAR may have a signaling peptide, a cancer molecule-specific scFv, 10 optionally a peptide linker/spacer, followed by an scFv that binds another antigen, followed by a spacer, a transmembrane domain, and a costimulatory domain. In some embodiments, the bi-specific CAR comprises one or more additional peptide segments described herein.

[0128] In some embodiments, CARs of the disclosure may comprise a CD20 scFv. An exemplary CD20 scFv comprises the following: DIVLTQSPAILSASPGEKVTMTCRASSSVNYMDWYQKKPGSSPKPWYATSNLASGV 15 PARFSGSGSGTSYSLTISRVEAEDAATYYCQQWSFNPPTFGGGTKLEIKGSTSGGGSG GGSGGGGSSEVQLQQSGAELVKPGASVKMSCKASGYTFTSYNMHVWKQTPGQGLE WIGAIYPNGDTSYNQKFKGKATLTADKSSSTAYMQLSSLTSEDSADYYCARSNYY 20 GSSYWFFDVWGAGTTVTVSS (SEQ ID NO:12).

[0129] In some embodiments, CARs of the disclosure may comprise a CD19 scFv. An exemplary CD19 scFv comprises the following: DIQMTQTSSLSASLGDRVТИSCRASQDISKYLNWYQQKPDGTVKLLIYHTSRLHSGV 25 PSRFGSGSGTDYSLTISNLEQEDIATYFCQQGNTLPYTFGGGTKLEITGSTSGSGKPG SGEGSTKGEVKLQESGPGLVAPSQSLSVTCTVSGVSLPDYGVSWIRQPPRKGLEWLG VIWGSETYYNSALKSRLTIKDNSKSQVFLKMNSLQTDDTAIYYCAKHYYYGGSYA 30 MD YWGQGTSVTVSS (SEQ ID NO:13). In some embodiments, an anti-CD19 CAR is encoded by a DNA construct with the following structure: GM-CSF signal peptide – HA tag – CD19 scFv – IgG4 hinge – CD28 tm – 4-1BB co-stimulatory domain – CD3 zeta. This construct has the following sequence: ATGCTGCTGCTGGTGACCAGCCTGCTGCTGTGCGAGCTGCCACCCCGCCTTC TGCTGATCCCCGGCGGAAGTTACCCATATGACGTTCCGACTACGCTGGCGACAT 35 CCAGATGACCCAGACCACCTCCAGCCTGAGCGCCAGCCTGGCGACCGGGTGAC

CATCAGCTGCCGGGCCAGCCAGGACATCAGCAAGTACCTGAACCTGGTATCAGCA
GAAGCCCGACGGCACCGTCAAGCTGCTGATCTACCACACCAGCCGGCTGCACAG
CGGCGTGCCTAGCCGGTTAGCGGCAGCGGCTCCGGACCGACTACAGCCTGAC
CATCTCCAACCTGGAACAGGAAGATATGCCACCTACTTTGCCAGCAGGGCAA
5 CACACTGCCCTACACCTTGGCGGCGGAACAAAGCTGGAAATCACCAGCAC
CTCCGGCAGCGCAAGCCTGGCAGCGCGAGGGCAGCACCAAGGGCGAGGTGA
AGCTGCAGGAAAGCGGCCCTGGCCTGGTGGCCCCAGCCAGAGCCTGAGCGTGA
CCTGCACCGTGAGCGCGTGAGCCTGCCGACTACGGCGTGAGCTGGATCCGGC
AGCCCCCAGGAAGGGCTGGAATGGCTGGCGTGATCTGGGCAGCGAGACCA
10 CCTACTACAACAGCGCCCTGAAGAGCCGGCTGACCATCATCAAGGACAACAGCA
AGAGCCAGGTGTTCCCTGAAGATGAACAGCCTGCAGACCGACGACACCGCCATCT
ACTACTGCGCCAAGCACTACTACTACGGCGGCAGCTACGCCATGGACTACTGGG
GCCAGGGCACCAGCGTGACCGTGAGCAGCGAATCTAAGTACGGACCGCCCTGCC
CCCCTGCCCTATGTTCTGGGTGCTGGTGGTGGCGAGGCGTGCTGCCCTGCTA
15 CAGCCTGCTGGTCACCGTGGCCTCATCATCTTTGGGTGAAACGGGCAGAAAG
AAACTCCTGTATATATTCAAACAAACCATTATGAGACCAGTACAAACTACTCAAG
AGGAAGATGGCTGTAGCTGCCGATTCCAGAAGAAGAAGAAGGAGGATGTGAAC
TGCGGGTGAAGTCAGCAGAACGCGCCAGCCTGCCTACCAGCAGGGCCAGA
ATCAGCTGTACAACGAGCTGAACCTGGGCAGAAGGGAAAGAGTACGACGTCCTGG
20 ATAAGCGGAGAGGCCGGGACCTGAGATGGCGGCAAGCCTCGCGGAAGAAC
CCCCAGGAAGGCCTGTATAACGAACTGCAGAAAGACAAGATGGCCGAGGCCTAC
AGCGAGATCGGCATGAAGGGCGAGCGGAGGCAGGGCAAGGCCACGACGGCCT
GTATCAGGGCCTGTCCACCGCCACCAAGGATAACCTACGACGCCCTGCACATGCA
GGCCCTGCCCTCAAGG (SEQ ID NO:88).

25 **[0130]** Other cancer-specific molecules (in addition to CD19 and CD20) that can be targeted by CARs can include CAIX, CD33, CD44v7/8, CEA, EGP-2, EGP-40, erb-B2, erb-B3, erb-B4, FBP, fetal acetylcholine receptor, GD2, GD3, Her2/neu, IL-13R-a2, KDR, k-light chain, LeY, L1 cell adhesion molecule, MAGE-A1, mesothelin, MUC1, NKG2D ligands, oncofetal antigen (h5T4), PSCA, PSMA, TAA targeted by mAb IgE, TAG-72, and
30 VEGF-R2. In some embodiments, the cancer-specific molecule comprises Her2.

b. Signal Peptides

[0131] In some embodiments, CARs include a signal peptide directs the transport and localization of the protein within a cell, e.g. to a certain cell organelle (such as the endoplasmic reticulum) and/or the cell surface. A signal peptide directs the nascent protein 5 into the endoplasmic reticulum. This is essential if the receptor is to be glycosylated and anchored in the cell membrane. Generally, the signal peptide natively attached to the amino-terminal most component is used (e.g. in a scFv with orientation light chain - linker - heavy chain, the native signal of the light-chain is used). In some embodiments the signal peptide has the sequence METDTLLWVLLLWVPGSTG (SEQ ID NO:14),
10 MLLVTSLLLCELPHPAFLLIPDT (SEQ ID NO:15), or MGTSLLCWMALCLLGADHADG (SEQ ID NO:16). In some embodiments, the signal peptide is cleaved after passage of the endoplasmic reticulum (ER), i.e. is a cleavable signal peptide. In some embodiments, a restriction site is at the carboxy end of the signal peptide to facilitate cleavage.

15 c. Antigen-binding Domain

[0132] The antigen-binding domain of a CAR is an antigen-binding protein, such as a single-chain variable fragment (scFv) based on antibodies against the target antigen. Framework modifications can be made to the antigen-binding domain to decrease 20 immunogenicity, for example, by "backmutating" one or more framework residues to the corresponding germline sequence. It is also contemplated that the antigen binding domain may be multi-specific or multivalent by multimerizing the antigen binding domain with V_H and V_L region pairs that bind either the same antigen (multi-valent) or a different antigen (multi-specific).

d. Peptide Spacer

25 [0133] A spacer region links the antigen-binding domain to the transmembrane domain of a CAR. It should be flexible enough to allow the antigen-binding domain to orient in different directions to facilitate antigen recognition. The simplest form is the hinge region from IgG. Alternatives include the CH₂CH₃ region of immunoglobulin and portions of CD3. In some embodiments, the CH₂CH₃ region may have L235E/N297Q or L235D/N297Q modifications, 30 or at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%

amino acid sequence identity of the CH₂CH₃ region. For most scFv-based constructs, the IgG hinge suffices. However the best spacer often has to be determined empirically. In some embodiments, the spacer is from IgG4.

[0134] As used herein, the term "hinge" refers to a flexible polypeptide connector region 5 (also referred to herein as "hinge region" or "spacer") providing structural flexibility and spacing to flanking polypeptide regions and can consist of natural or synthetic polypeptides. A "hinge" derived from an immunoglobulin (e.g., IgG1) is generally defined as stretching from Glu216 to Pro230 of human IgG1 (Burton (1985) Molec. Immunol., 22: 161- 206). Hinge regions of other IgG isotypes may be aligned with the IgG1 sequence by placing the 10 first and last cysteine residues forming inter-heavy chain disulfide (S-S) bonds in the same positions. The hinge region may be of natural occurrence or non-natural occurrence, including but not limited to an altered hinge region as described in U.S. Pat. No. 5,677,425. The hinge region can include complete hinge region derived from an antibody of a different 15 class or subclass from that of the CH₁ domain. The term "hinge" can also include regions derived from CD8 and other receptors that provide a similar function in providing flexibility and spacing to flanking regions.

[0135] The peptide spacer can have a length of at least, at most, or exactly 4, 5, 6, 7, 8, 9, 10, 15, 16, 17, 18, 19, 20, 20, 25, 30, 35, 40, 45, 50, 75, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214, 20 215, 216, 217, 218, 219, 220, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 260, 270, 280, 290, 300, 325, 350, or 400 amino acids (or any derivable range therein). In some embodiments, the peptide spacer consists of or comprises a hinge region from an immunoglobulin. Immunoglobulin hinge region amino acid sequences are known in the art; see, e.g., Tan et al. 25 (1990) Proc. Natl. Acad. Sci. USA 87: 162; and Huck et al. (1986) Nucl. Acids Res.

[0136] The length of a peptide spacer may have effects on the response to the target antigen and/or expansion properties. In some embodiments, a shorter spacer such as less than 50, 45, 40, 30, 35, 30, 25, 20, 15, or 10 amino acids may have the advantage of a decrease in the concentration of target antigen required for an effective activation response. In some 30 embodiments, a longer spacer, such as one that is at least 50, 60, 70, 80, 90, 100, 110, 120, 130, 140, 150, 200, 201, 202, 203, 204, 205, 206, 207, 208, 209, 210, 211, 212, 213, 214,

215, 216, 217, 218, 219, 220, 225, 226, 227, 228, 229, 230, 231, 232, 233, 234, 235, 236, 237, 238, 239, 240, 241, 242, 243, 244, 245, 246, 247, 248, 249, 250, 260, 270, 280, or 290 amino acids may have the advantage of increased expansion in vivo or in vitro.

[0137] As non-limiting examples, an immunoglobulin hinge region can include one of the following amino acid sequences: DKTHT (SEQ ID NO:17); CPPC (SEQ ID NO:18); CPEPKSCDTPPPCPR (SEQ ID NO:19); ELKTPLGDTTHT (SEQ ID NO:20); KSCDKTHTCP (SEQ ID NO:21); KCCVDCP (SEQ ID NO:22); KYGPPCP (SEQ ID NO:23); EPKSCDKTHTCPPCP (SEQ ID NO:24) (human IgG1 hinge); ERKCCVECPCP (SEQ ID NO:25) (human IgG2 hinge); ELKTPLGDTTHTCPRCP (SEQ ID NO:26) (human IgG3 hinge); SPNMVPHAHHAQ (SEQ ID NO:27); ESKYGPPCP (SEQ ID NO:28) or ESKYGPPCPSCP (SEQ ID NO:29) (human IgG4 hinge-based) and the like.

[0138] The hinge region can comprise an amino acid sequence of a human IgG1, IgG2, IgG3, or IgG4, hinge region. The hinge region can include one or more amino acid substitutions and/or insertions and/or deletions compared to a wild-type (naturally-occurring) hinge region. For example, His229 of human IgG1 hinge can be substituted with Tyr, so that the hinge region comprises the sequence EPKSCDKTYTCPPCP (SEQ ID NO:30).

[0139] The hinge region can comprise an amino acid sequence derived from human CD8; e.g., the hinge region can comprise the amino acid sequence: TTTPAPRPPPTPAPTIASQPLSLRPEACRPAAGGA VHTRGLDFACD (SEQ ID NO:31), or a variant thereof.

e. Transmembrane Domain

[0140] The transmembrane domain of the CAR is a hydrophobic alpha helix that spans the membrane. Generally, the transmembrane domain from the most membrane proximal component of the endodomain is used. Different transmembrane domains result in different receptor stability.

[0141] The transmembrane domain is interposed between the peptide spacer and the endodomain. In some embodiments, the transmembrane domain is interposed between the peptide spacer and a co-stimulatory region. In some embodiments, a linker is between the transmembrane domain and a co-stimulatory region or endodomain.

[0142] Any transmembrane domain that provides for insertion of a polypeptide into the cell membrane of a eukaryotic (e.g., mammalian) cell is suitable for use. As one non-limiting example, the transmembrane sequence IYIWAPLAGTCGVLLSLVITLYC (SEQ ID NO:32) can be used. In some embodiments, the transmembrane domain is CD8 beta 5 derived: LGLLVAGVLVLLVSLGVAIHLCC (SEQ ID NO:33); CD4 derived: ALIVLGGVAGLLLFIGLGIFFCVRC (SEQ ID NO:34); CD3 zeta derived: LCYLLDGILFIYGVILTALFLRV (SEQ ID NO:35); CD28 derived: WVLVVVGGVLACYSLLVTVAIFIIFWV (SEQ ID NO:36); CD134 (OX40) derived: VAAILGLGLVLGLLGPLAILALYLL (SEQ ID NO:37); or CD7 derived: 10 ALPAALAVISFLLGLGLGVACVLA (SEQ ID NO:38).

f. Endodomain

[0143] After antigen recognition, receptors cluster and a signal is transmitted to the cell through the endodomain and/or co-stimulatory domain. In some embodiments, the co-stimulatory domains described herein are part of the endodomain. The most commonly 15 used endodomain component is CD3-zeta, which contains 3 ITAMs. This transmits an activation signal to the T cell after antigen is bound. CD3-zeta may not provide a fully competent activation signal and additional co-stimulatory signaling is needed. For example, chimeric CD28 and OX40 can be used with CD3-Zeta to transmit a proliferative/survival signal, or all three can be used together.

[0144] Further endodomains suitable for use in CARs include any desired signaling 20 domain that provides a distinct and detectable signal (e.g., increased production of one or more cytokines by the cell; change in transcription of a target gene; change in activity of a protein; change in cell behavior, e.g., cell death; cellular proliferation; cellular differentiation; cell survival; modulation of cellular signaling responses; etc.) in response to activation by 25 way of binding of the antigen to the antigen binding domain. In some embodiments, the endodomain includes at least one (e.g., one, two, three, four, five, six, etc.) ITAM motif as described herein. In some embodiments, the endodomain includes DAP10/CD28 type signaling chains.

[0145] Endodomains suitable for use in CARs include immunoreceptor tyrosine-based 30 activation motif (ITAM)-containing intracellular signaling polypeptides. An ITAM motif is YX₁X₂(L/I), where X₁ and X₂ are independently any amino acid (SEQ ID NO:39). In some

cases, the endodomain comprises 1, 2, 3, 4, or 5 ITAM motifs. In some cases, an ITAM motif is repeated twice in an endodomain, where the first and second instances of the ITAM motif are separated from one another by 6 to 8 amino acids, e.g., (YX₁X₂(L/I))(X₃)_n(YX₁X₂(L/I)), where n is an integer from 6 to 8, and each of the 6-8 X₃ can be any amino acid (SEQ ID

5 NO:40).

[0146] A suitable endodomain may be an ITAM motif-containing portion that is derived from a polypeptide that contains an ITAM motif. For example, a suitable endodomain can be an ITAM motif-containing domain from any ITAM motif-containing protein. Thus, a suitable endodomain need not contain the entire sequence of the entire protein from which it is derived. Examples of suitable ITAM motif-containing polypeptides include, but are not limited to: DAP12; FCER1G (Fc epsilon receptor I gamma chain); CD3D (CD3 delta); CD3E (CD3 epsilon); CD3G (CD3 gamma); CD3Z (CD3 zeta); and CD79A (antigen receptor complex-associated protein alpha chain).

[0147] In some cases, the endodomain is derived from DAP12 (also known as TYROBP;

15 TYRO protein tyrosine kinase binding protein; KARAP; PLOSL; DN AX- activation protein

12; KAR-associated protein; TYRO protein tyrosine kinase- binding protein; killer activating receptor associated protein; killer- activating receptor- associated protein; etc.). For example, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid

20 sequence identity to

MGGLEPCSRLLLLPLLLAVSGLRPVQAQAAQSDCSCSTVSPGVLAGIVMGDLVLTVAL
ALAVYFLGRLVPRGRGAAEAATRKORITETESEPYOEOLOGORSVYSDLNTQRPYYK
(SEQ ID NO:41);

MGGLEPCSRLLLLPLLLAVSGLRPVQAQAAQSDCSCSTVSPGVLAGIVMGDLVLTVAL

25 ALAVYFLGRLVPRGRGAAEAATRKORITETESEPYOEOLOGORSVYSDLNTQRPYYK
(SEQ ID NO:42);

MGGLEPCSRLLLLPLLLAVSDCSCSTVSPGVLAGIVMGDLVLTVALALAVYFLGRLVP

RGRGAAEAATRKORITETESEPYOEOLOGORSVYSDLNTQRPYYK (SEQ ID NO:43); or

MGGLEPCSRLLLLPLLLAVSDCSCSTVSPGVLAGIVMGDLVLTVALALAVYFLGRLVP

30 RGRGAAEAATRKORITETESEPYOEOLOGORSVYSDLNTQRPYYK (SEQ ID NO:44).

[0148] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM motif-containing portion of the full length DAP12 amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to ESPYOEOLOGORSVDVYSDLNTO (SEQ ID NO:45).

[0149] In some embodiments, the endodomain is derived from FCER1G (also known as FCRG; Fc epsilon receptor I gamma chain; Fc receptor gamma-chain; fc-epsilon R1-gamma; fcRgamma; fceRI gamma; high affinity immunoglobulin epsilon receptor subunit gamma; immunoglobulin E receptor, high affinity, gamma chain; etc.). For example, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to MIPAVVLLLLLVEQAAALGEPQLCYILDAILFLYGIVLTLLYCRLKIQVRKAAITSYE KSDGVYTGLSTRNQETYETLKHEKPPQ (SEQ ID NO:46).

[0150] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM motif-containing portion of the full length FCER1G amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to DGVYTGLSTRNOETYETLKHE (SEQ ID NO:47).

[0151] In some embodiments, the endodomain is derived from T-cell surface glycoprotein CD3 delta chain (also known as CD3D; CD3-DELTA; T3D; CD3 antigen, delta subunit; CD3 delta; CD3d antigen, delta polypeptide (TiT3 complex); OKT3, delta chain; T-cell receptor T3 delta chain; T-cell surface glycoprotein CD3 delta chain; etc.). For example, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to a contiguous stretch of from about 100 amino acids to about 110 amino acids (aa), from about 110 aa to about 115 aa, from about 115 aa to about 120 aa, from about 120 aa to about 130 aa, from about 130 aa to about 140 aa, from about 140 aa to about 150 aa, or from about 150 aa to about 170 aa, of either of the following amino acid sequences (2 isoforms):

MEHSTFLSGLVLATLLSQVSPFKIPIEELEDRVFVNCNTSITWVEGTVGTLSDITRLD

LGKRILDPRGIYRCNGTDIYKDKESTVQVHYRMCQSCVELDPATVAGIIVTDVIATLL
LALGVFCFAGHETGRLSGAADTOALLRNDOVYOPLRDRDDAOYSHLGGNWARNK
(SEQ ID NO:48) or

MEHSTFLSGLVLATLLSQVSPFKIPIEELDRVFVNCNTSITWVEGTVGTLSDITRLD
5 LGKRILDPRGIYRCNGTDIYKDKESTVOVHYRTADTOALLRNDOVYOPLRDRDDAQ
YSHLGGNWARNK (SEQ ID NO:49).

[0152] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM motif-containing portion of the full length CD3 delta amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 10 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to DOVYOPLRDRDDAOYSHLGGN (SEQ ID NO:50).

[0153] In some embodiments, the endodomain is derived from T-cell surface glycoprotein CD3 epsilon chain (also known as CD3e, T-cell surface antigen T3/Leu-4 epsilon chain, T-cell surface glycoprotein CD3 epsilon chain, AI504783, CD3, CD3epsilon, 15 T3e, etc.). For example, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to a contiguous stretch of from about 100 amino acids to about 110 amino acids (aa), from about 110 aa to about 115 aa, from about 115 aa to about 120 aa, from about 120 aa to about 130 aa, from about 130 aa to about 140 aa, from 20 about 140 aa to about 150 aa, or from about 150 aa to about 205 aa, of the following amino acid

sequence:

MQSGTHWRVLGLCLLSVGVWGQDGNEEMGGITQTPYKVSISGTTVILTCPQYPGSEI
LWQHNDKNIGGDEDDKNIGSDEDHSLKEFSELEQSGYYVCYPRGSKPEDANFYLY
LRARVCENCMDVMSVATIVDICITGGLLLVYYWSKNRKA
25 KAKPVTRGAGAG GRQRGQNKERPPPVPNPDYEPIRGQQRDLYSGLNQRRI (SEQ ID NO:51).

[0154] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM motif-containing portion of the full length CD3 epsilon amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence 30 identity to NPDYEPIRGQQRDLYSGLNQR (SEQ ID NO:52).

[0155] In some embodiments, the endodomain is derived from T-cell surface glycoprotein CD3 gamma chain (also known as CD3G, T-cell receptor T3 gamma chain, CD3-GAMMA, T3G, gamma polypeptide (TiT3 complex), etc.). For example, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least

5 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to a contiguous stretch of from about 100 amino acids to about 110 amino acids (aa), from about 110 aa to about 115 aa, from about 115 aa to about 120 aa, from about 120 aa to about 130 aa, from about 130 aa to about 140 aa, from about 140 aa to about 150 aa, or from about 150 aa to about 180 aa, of the following amino acid sequence:

10 MEQGKGLAVLILAIILLQGTLAQSIKGNHLVKVYDYQEDGSVLLTCDAEAKNITWFK
DGKMIGFLTEDKKWNLGSNAKDPRGMYQCKGSQNKSPLQVYYRMCQNCIELNA
ATISGFLFAEIVSIFVLAGVYFIAGODGVROSRAKOTLLPNDOLYOPLKDREDDQ
YSHLQGNQLRRN (SEQ ID NO:53).

[0156] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM motif-containing portion of the full length CD3 gamma amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to DOLYOPLKDREDDOYSHLOGN (SEQ ID NO:54).

[0157] In some embodiments, the endodomain is derived from T-cell surface glycoprotein CD3 zeta chain (also known as CD3Z, T-cell receptor T3 zeta chain, CD247, CD3-ZETA, CD3H, CD3Q, T3Z, TCRZ, etc.). For example, a suitable intracellular signaling domain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to a contiguous stretch of from about 100 amino acids to about 110 amino acids (aa), from about 110 aa to about 115 aa, from about 115 aa to about 120 aa, from about 120 aa to about 130 aa, from about 130 aa to about 140 aa, from about 140 aa to about 150 aa, or from about 150 aa to about 160 aa, of either of the following amino acid sequences (2 isoforms):
MKWKALFTAAILQAQLPITEAQSFGLDPKLCYLLDGILFIYGVILTAFLRVKFSRSA
DAPAYQQGQNQLYNELNLGRREEYDVLDKRRGRDPEMGGKPRRKNPQEGLYNELO
30 KDKMAEAYSEIGMKGERRRGKGHDGLYOGLSTATKDTYDALHMQALPPR (SEQ ID NO:

MWKALFTAAILQAQLPITEAQSFGLDPKLCYLLDGILFIYGVILTAFLRVKFSRSA
DAPAYQQGQNQLYNELNLGRREYDVLKDERRGRDPEMGGKPQRRKNPQEGLYNEL
OKDKMAEAYSEIGMKGERRRGKGHDGLYOGLSTATKDTYDALHMQALPPR
(SEQ ID NO: 56).

5 [0158] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM motif-containing portion of the full length CD3 zeta amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to any of the following amino acid sequences:

10 RVKFSRSADAPAYOQGONOLYNELNLGRREYDVLKDERRGRDPEMGGKPQRRKNPO
EGLYNEOKDKMAEAYSEIGMKGERRRGKGHDGLYOGLSTATKDTYDALHMQALPPR
(SEQ ID NO: 82); NOLYNELNLGRREYDVLDKR (SEQ ID NO:57);
EGLYNELQKDKMAEAYSEIGMK (SEQ ID NO:58); or
DGLYOGLSTATKDTYDALHMO (SEQ ID NO:59).

15 [0159] In some embodiments, the endodomain is derived from CD79A (also known as B-cell antigen receptor complex-associated protein alpha chain; CD79a antigen (immunoglobulin-associated alpha); MB-1 membrane glycoprotein; ig-alpha; membrane- bound immunoglobulin-associated protein; surface IgM-associated protein; etc.). For example, a suitable endodomain polypeptide can comprise an amino acid sequence 20 having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to a contiguous stretch of from about 100 amino acids to about 110 amino acids (aa), from about 110 aa to about 115 aa, from about 115 aa to about 120 aa, from about 120 aa to about 130 aa, from about 130 aa to about 150 aa, from about 150 aa to about 200 aa, or from about 200 aa to about 220 aa, of either of the following amino 25 acid sequences (2 isoforms):
MPGGPGVLQALPATIFLLFLSAVYLGPGCQALWMHKVPASLMVSLGEDAHFQCPH
NSSNNANVTWWRVLHGNYTWPPEFLGPAGEDPNTLIIQNVNKSHEGIYVCRVQEGN
ESYQQSCGTYLVRVRQPPPRPFLDMGEGTKNRIITAEGIILLFCAVVPGTLLLFRKRWO
NEKLGLDAGDEYEDEDENLYEGLNLDDCSMYEDISRGLOGTYQDVGSLNIGDVQLEKP
30 (SEQ ID NO:60); or
MPGGPGVLQALPATIFLLFLSAVYLGPGCQALWMHKVPASLMVSLGEDAHFQCPH

NSSNNANVTWWRVLHGNYTWPPEFLGPGEDPNEPPPRPFLDMGE GTKNRIITAEGIIL
LFCAVVPGTLLLFRKRWQNEKLGLDAGDEYEDENLYEGLNLDDCSMYE
DISRGLQGTYQDVGSNLIGDVQLEKP (SEQ ID NO:61).

[0160] In some embodiments, a suitable endodomain polypeptide can comprise an ITAM

5 motif-containing portion of the full length CD79A amino acid sequence. Thus, a suitable endodomain polypeptide can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100%, amino acid sequence identity to the following amino acid sequence: ENLYEGLNLDDCSMYEDISRG (SEQ ID NO:62).

10 **[0161]** In some embodiments, suitable endodomains can comprise a DAP10/CD28 type signaling chain. An example of a DAP 10 signaling chain is the amino acid sequence: RP RR SPAQDGK VYINMPGRG (SEQ ID NO:63). In some embodiments, a suitable endodomain comprises an amino acid sequence having at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to the entire length of the 15 amino acid sequence RP RR SPAQDGK VYINMPGRG (SEQ ID NO:64).

[0162] An example of a CD28 signaling chain is the amino acid sequence FWVLVVVGVLACYSLVTVAIFI FWVRSKRSRLLHSDYMNMTPRRPGPTRKHYQP YAPPRDFAAYRS (SEQ ID NO:65). In some embodiments, a suitable endodomain comprises an amino acid sequence having at least 85%, at least 90%, at least 95%, at least 98%, or at least 99%, amino acid sequence identity to the entire length of the amino acid sequence

FWVLVVVGVLACYSLVTVAIFI FWVRSKRSRLLHSDYMNMTPRRPGPTRKHYQP YAPPRDFAAYRS (SEQ ID NO:66).

[0163] Further endodomains suitable for use in the polypeptides of the disclosure include

25 a ZAP70 polypeptide, e.g., a polypeptide comprising an amino acid sequence having at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, or 100%, amino acid sequence identity to a contiguous stretch of from about 300 amino acids to about 400 amino acids, from about 400 amino acids to about 500 amino acids, or from about 500 amino acids to 619 amino acids, of the following amino acid sequence: MPDPA AHLPFFYGSIS
30 RAEAEELKLAGMADGLFLLRQCLRS LGGYVLSLVHDVRFHHFPIERQLNGTYAIAG GKAHCGPAELCEFYSRDPDGLPCNLRKPCNRPSGLEPQPGVFDCLR DAMVRDYVRQ

TWKLEGEALEQAIISQAPQVEKLIATTAKERMPWYHSSLTREEAERKLYSGAQTDGK
FLLRPRKEQGTYALSLIYGKTVYHYLISQDKAGKYCIPEGTKFDTLWQLVEYKLKA
DGLIYCLKEACPNSNASGAAAPTLPAHPSTLTHPQRIDTLNSDGYTPEPARITSP
DKPRPMPMDTSVYESPYSDPEELDKKFLKRDNLLIADIELGCGNFGSVRQGVYRM
5 RKKQIDVAIKVLKQGTEKADTEEMMREAQIMHQLDNPYIVRLIGVCQAEALMLVME
MAGGGPLHKFLVGKREEIPVSNAELLHQVSMGMKYLEEKNFVHRDLAARNVLLV
NRHYAKISDFGLSKALGADDSYYTARSAGKWPLKYAPECINFRKFSSRSDVWSYG
VTMWEALSYGQKPYKKMKGPEVMAFIEQGKRMECPPPELYALMSDCWIYKWE
DRPDFLTVEQRMRACYSLASKVEGPPGSTQKAEEACA (SEQ ID NO:67).

10 g. Detection Peptides

[0164] CARs can include detection peptides. Suitable detection peptides include hemagglutinin (HA; e.g., YPYDVPDYA (SEQ ID NO:68); FLAG (e.g., DYKDDDDK (SEQ ID NO:69); c-myc (e.g., EQKLISEEDL; SEQ ID NO:70), and the like. Other suitable detection peptides are known in the art.

15 h. Peptide Linkers

[0165] In some embodiments, CARs may also include peptide linkers (sometimes referred to as a linker). A peptide linker may be separating any of the peptide domain/regions described herein. As an example, a linker may be between the signal peptide and the antigen binding domain, between the V_H and V_L of the antigen binding domain, between the antigen binding domain and the peptide spacer, between the peptide spacer and the transmembrane domain, flanking the co-stimulatory region or on the N- or C- region of the co-stimulatory region, and/or between the transmembrane domain and the endodomain. The peptide linker may have any of a variety of amino acid sequences. Domains and regions can be joined by a peptide linker that is generally of a flexible nature, although other chemical linkages are not excluded. A linker can be a peptide of between about 6 and about 40 amino acids in length, or between about 6 and about 25 amino acids in length. These linkers can be produced by using synthetic, linker-encoding oligonucleotides to couple the proteins.

[0166] Peptide linkers with a degree of flexibility can be used. The peptide linkers may have virtually any amino acid sequence, bearing in mind that suitable peptide linkers will have a sequence that results in a generally flexible peptide. The use of small amino acids,

such as glycine and alanine, are of use in creating a flexible peptide. The creation of such sequences is routine to those of skill in the art.

[0167] Suitable linkers can be readily selected and can be of any of a suitable of different lengths, such as from 1 amino acid (e.g., Gly) to 20 amino acids, from 2 amino acids to 5 15 amino acids, from 3 amino acids to 12 amino acids, including 4 amino acids to 10 amino acids, 5 amino acids to 9 amino acids, 6 amino acids to 8 amino acids, or 7 amino acids to 8 amino acids, and may be 1, 2, 3, 4, 5, 6, or 7 amino acids.

[0168] Exemplary flexible linkers include glycine polymers (G)_n, glycine- serine polymers (including, for example, (GS)_n, (GSGGS)_n (SEQ ID NO:71) and (GGGS)_n 10 (SEQ ID NO:72), where n is an integer of at least one, glycine-alanine polymers, alanine-serine polymers, and other flexible linkers known in the art. Glycine and glycine- serine polymers can be used; both Gly and Ser are relatively unstructured, and therefore can serve as a neutral tether between components. Glycine polymers can be used; glycine accesses significantly more phi-psi space than even alanine, and is much less restricted than 15 residues with longer side chains. Exemplary spacers can comprise amino acid sequences including, but not limited to, GGSG (SEQ ID NO:73), GGS GG (SEQ ID NO:74), GSGSG (SEQ ID NO:75), GSGGG (SEQ ID NO:76), GGGSG (SEQ ID NO:77), GSSSG (SEQ ID NO:78), and the like.

i. Co-stimulatory Region

[0169] Non-limiting examples of suitable co-stimulatory regions in CARs include, but 20 are not limited to, polypeptides from 4-1BB (CD137), CD28, ICOS, OX-40, BTLA, CD27, CD30, GITR, and HVEM.

[0170] A co-stimulatory region may have a length of at least, at most, or exactly 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, or 300 amino acids or any range derivable 25 therein. In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein 4-1BB (also known as TNFRSF9; CD137; 4-1BB; CDw137; ILA; etc.). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to 30 KRGRKKLLYIFKQPFMRPVQTTQEEDGCSCRFPEEEEGGCEL (SEQ ID NO:79).

[0171] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein CD28 (also known as Tp44). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to FWVRSKRSRLLHSDYMNMTPRRPGPTRKHYQPYAPPRDFAAYRS (SEQ ID NO:80).

[0172] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein ICOS (also known as AILIM, CD278, and CVID1). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to TKKKYSSSVHDPNGEYMFMRNAVNTAKKSRLTDVTL (SEQ ID NO:81).

[0173] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein OX-40 (also known as TNFRSF4, RP5-902P8.3, ACT35, CD134, OX40, TXGP1L). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to RRDQRLPPDAHKPPGGGSFRTPIQEEQADAHSTLAKI (SEQ ID NO:82).

[0174] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein BTLA (also known as BTLA1 and CD272). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to CCLRRHQGKQNELSDTAGREINLVDAHLKSEQTEASTRQNSQVLLSETGIYDNDPDL CFRMQEGSEVYSNPCLEENKPGIVYASLNHSVIGPNSRLARNVKEAPTEYASICVRS (SEQ ID NO:83).

[0175] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein CD27 (also known as S 152, T14, TNFRSF7, and Tp55). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to

HQRRKYRSNKGESPVEPAEPCRYSCPREEGSTIPIQEDYRKPEPACSP (SEQ ID NO:84).

[0176] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein CD30 (also known as TNFRSF8, D1S166E, and Ki-1).

5 For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to
RRACRKIRQKLHLCYPVQTSQLPKLELVDSRPRRSSTQLRSGASVTEPVAEERGLMS
QPLMETCHSVGAAYLESLPLQDASPAGGPSSPRDLPEPRVSTEHTNNKIEKIYIMKAD
10 TVIVGTVKAELPEGRGLAGPAEPELEEELEADHTPHYPEQETEPPLGSCSDVMLSVEE
EGKEDPLPTAASGK (SEQ ID NO:85).

[0177] In some embodiments, the co-stimulatory region is derived from an intracellular portion of the transmembrane protein GITR (also known as TNFRSF18, RP5-902P8.2, AITR, CD357, and GITR-D). For example, a suitable co- stimulatory region can comprise an

15 amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to
HIWQLRSQCMWPRETQLLLEVPPSTEDARSCQFPEEERGERSAEEKGRLGDLWV
(SEQ ID NO:86).

[0178] In some embodiments, the co-stimulatory region derived from an intracellular portion of the transmembrane protein HVEM (also known as TNFRSF14, RP3-395M20.6,

20 ATAR, CD270, HVEA, HVEM, LIGHTR, and TR2). For example, a suitable co-stimulatory region can comprise an amino acid sequence having at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, or 100% amino acid sequence identity to
CVKRRKPRGDVVVKVIVSVQRKRQEAEGEATVIEALQAPPDVTTVAVEETIPSFTGRSP
25 NH (SEQ ID NO:87).

V. METHODS

[0179] Aspects of the current disclosure relate to methods of treating a condition, for stimulating an immune response, for regulating an immune response, for suppressing an immune response, and for reducing the risk of an adverse immune response, such as cytokine release syndrome. These actions may be done *in vitro*, *in vivo*, or *ex vivo*. In some

embodiments, the methods relate to cells capable of stimulating an immune response in the presence of a cancer cell. The method generally involves genetically modifying a mammalian cell with an expression vector, or an RNA (e.g., in vitro transcribed RNA), comprising nucleotide sequences encoding a polypeptide of the disclosure or directly transferring the 5 polypeptide to the cell. The cell can be an immune cell (e.g., a T lymphocyte or NK cell), a stem cell, a progenitor cell, etc. In some embodiments, the cell is a cell described herein.

[0180] In some embodiments, the genetic modification is carried out *ex vivo*. For example, a T lymphocyte, a stem cell, or an NK cell (or cell described herein) is obtained from an individual; and the cell obtained from the individual is genetically modified to 10 express a polypeptide of the disclosure. In some cases, the genetically modified cell is activated *ex vivo* (i.e., a target antigen is contacted with the cells *ex vivo*). In other cases, the genetically modified cell is introduced into an individual (e.g., the individual from whom the cell was obtained); and the genetically modified cell is activated *in vivo* (i.e., by endogenously produced target antigen).

15 **[0181]** In some embodiments, the methods further comprise the administration of additional therapeutic agents, such as bi-specific T cell engagers (BITE). Such therapeutic agents may be administered in peptide form to the patient or expressed in cells of the disclosure, such as those that comprise the IL-6Ra binding protein. The BITE may have antigen specificity for a cancer antigen/cancer molecule known in the art and/or described 20 herein and may also have antigen specificity for a T cell molecule such as CD3.

[0182] In some embodiments, the methods relate to administration of the cells or peptides described herein for the treatment of a cancer or administration to a person with a cancer. In some embodiments the cancer is adrenal cancer, anal cancer, bile duct cancer, bladder cancer, bone cancer, brain/CNS tumors in children or adults, breast cancer, cervical cancer, 25 colon/rectum cancer, endometrial cancer, esophagus cancer, Ewing family of tumors, eye cancer, gallbladder cancer, gastrointestinal carcinoid tumors, gastrointestinal stromal tumor (GIST), gestation trophoblastic disease, hodgkin disease, kaposi sarcoma, kidney cancer, laryngeal and hypopharyngeal cancer, leukemia, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, chronic myeloid leukemia, chronic 30 myelomonocytic leukemia, liver cancer, lung cancer, non-small cell lung cancer, small cell lung cancer, lung carcinoid tumor, lymphoma, malignant mesothelioma, multiple myeloma,

myelodysplastic syndrome, nasal cavity and paranasal sinum cancer, nasopharyngeal cancer, neuroblastoma, non-hodgkin lymphoma, oral cavity or oropharyngeal cancer, osteosarcoa, ovarian cancer, pancreatic cancer, penile cancer, pituitary tumors, prostate cancer, retinoblastoma, rhabdomyosarcoma, salivary gland cancer, skin cancer, sarcoma, basal skin 5 cancer, squamous cell skin cancer, melanoma, merkel cell skin cancer, small intestine cancer, stomach cancer, testicular cancer, thymus cancer, thyroid cancer, uterine sarcoma, vaginal cancer, vulvar cancer, waldenstrom macroglobulinemia, or wilms tumor.

[0183] Embodiments can be used to treat or ameliorate a number of immune-mediated, inflammatory, or autoimmune-inflammatory diseases, e.g., allergies, asthma, diabetes 10 (e.g. type 1 diabetes), graft rejection, etc. Examples of such diseases or disorders also include, but are not limited to arthritis (rheumatoid arthritis such as acute arthritis, chronic rheumatoid arthritis, gout or gouty arthritis, acute gouty arthritis, acute immunological arthritis, chronic inflammatory arthritis, degenerative arthritis, type II collagen-induced arthritis, infectious arthritis, Lyme arthritis, proliferative arthritis, psoriatic arthritis, Still's disease, vertebral 15 arthritis, and systemic juvenile-onset rheumatoid arthritis, osteoarthritis, arthritis chronica progradient, arthritis deformans, polyarthritis chronica primaria, reactive arthritis, and ankylosing spondylitis), inflammatory hyperproliferative skin diseases, psoriasis such as plaque psoriasis, gutatte psoriasis, pustular psoriasis, and psoriasis of the nails, atopy including atopic diseases such as hay fever and Job's syndrome, dermatitis including contact 20 dermatitis, chronic contact dermatitis, exfoliative dermatitis, allergic dermatitis, allergic contact dermatitis, dermatitis herpetiformis, nummular dermatitis, seborrheic dermatitis, non-specific dermatitis, primary irritant contact dermatitis, and atopic dermatitis, x-linked hyper IgM syndrome, allergic intraocular inflammatory diseases, urticaria such as chronic allergic urticaria and chronic idiopathic urticaria, including chronic autoimmune urticaria, 25 myositis, polymyositis/dermatomyositis, juvenile dermatomyositis, toxic epidermal necrolysis, scleroderma (including systemic scleroderma), sclerosis such as systemic sclerosis, multiple sclerosis (MS) such as spino-optical MS, primary progressive MS (PPMS), and relapsing remitting MS (RRMS), progressive systemic sclerosis, atherosclerosis, arteriosclerosis, sclerosis disseminata, ataxic sclerosis, neuromyelitis optica (NMO), 30 inflammatory bowel disease (IBD) (for example, Crohn's disease, autoimmune-mediated gastrointestinal diseases, colitis such as ulcerative colitis, colitis ulcerosa, microscopic colitis, collagenous colitis, colitis polyposa, necrotizing enterocolitis, and transmural colitis, and

autoimmune inflammatory bowel disease), bowel inflammation, pyoderma gangrenosum, erythema nodosum, primary sclerosing cholangitis, respiratory distress syndrome, including adult or acute respiratory distress syndrome (ARDS), meningitis, inflammation of all or part of the uvea, iritis, choroiditis, an autoimmune hematological disorder, rheumatoid 5 spondylitis, rheumatoid synovitis, hereditary angioedema, cranial nerve damage as in meningitis, herpes gestationis, pemphigoid gestationis, pruritis scroti, autoimmune premature ovarian failure, sudden hearing loss due to an autoimmune condition, IgE-mediated diseases such as anaphylaxis and allergic and atopic rhinitis, encephalitis such as Rasmussen's 10 encephalitis and limbic and/or brainstem encephalitis, uveitis, such as anterior uveitis, acute anterior uveitis, granulomatous uveitis, nongranulomatous uveitis, phacoantigenic uveitis, posterior uveitis, or autoimmune uveitis, glomerulonephritis (GN) with and without nephrotic 15 syndrome such as chronic or acute glomerulonephritis such as primary GN, immune-mediated GN, membranous GN (membranous nephropathy), idiopathic membranous GN or idiopathic membranous nephropathy, membrano- or membranous proliferative GN (MPGN), including Type I and Type II, and rapidly progressive GN, proliferative nephritis, autoimmune polyglandular endocrine failure, balanitis including balanitis circumscripta plasmacellularis, balanoposthitis, erythema annulare centrifugum, erythema dyschromicum 20 perstans, erythema multiform, granuloma annulare, lichen nitidus, lichen sclerosus et atrophicus, lichen simplex chronicus, lichen spinulosus, lichen planus, lamellar ichthyosis, epidermolytic hyperkeratosis, premalignant keratosis, pyoderma gangrenosum, allergic 25 conditions and responses, allergic reaction, eczema including allergic or atopic eczema, asteatotic eczema, dyshidrotic eczema, and vesicular palmoplantar eczema, asthma such as asthma bronchiale, bronchial asthma, and auto-immune asthma, conditions involving infiltration of T cells and chronic inflammatory responses, immune reactions against foreign antigens such as fetal A-B-O blood groups during pregnancy, chronic pulmonary 30 inflammatory disease, autoimmune myocarditis, leukocyte adhesion deficiency, lupus, including lupus nephritis, lupus cerebritis, pediatric lupus, non-renal lupus, extra-renal lupus, discoid lupus and discoid lupus erythematosus, alopecia lupus, systemic lupus erythematosus (SLE) such as cutaneous SLE or subacute cutaneous SLE, neonatal lupus syndrome (NLE), and lupus erythematosus disseminatus, juvenile onset (Type I) diabetes mellitus, including pediatric insulin-dependent diabetes mellitus (IDDM), and adult onset diabetes mellitus (Type II diabetes) and autoimmune diabetes. Also contemplated are immune responses

associated with acute and delayed hypersensitivity mediated by cytokines and T-lymphocytes, sarcoidosis, granulomatosis including lymphomatoid granulomatosis, Wegener's granulomatosis, agranulocytosis, vasculitides, including vasculitis, large-vessel vasculitis (including polymyalgia rheumatica and giant cell (Takayasu's) arteritis),

5 medium-vessel vasculitis (including Kawasaki's disease and polyarteritis nodosa/periarteritis nodosa), microscopic polyarteritis, immunovasculitis, CNS vasculitis, cutaneous vasculitis, hypersensitivity vasculitis, necrotizing vasculitis such as systemic necrotizing vasculitis, and ANCA-associated vasculitis, such as Churg-Strauss vasculitis or syndrome (CSS) and ANCA-associated small-vessel vasculitis, temporal arteritis, aplastic anemia, autoimmune

10 aplastic anemia, Coombs positive anemia, Diamond Blackfan anemia, hemolytic anemia or immune hemolytic anemia including autoimmune hemolytic anemia (AIHA), Addison's disease, autoimmune neutropenia, pancytopenia, leukopenia, diseases involving leukocyte diapedesis, CNS inflammatory disorders, Alzheimer's disease, Parkinson's disease, multiple organ injury syndrome such as those secondary to septicemia, trauma or hemorrhage,

15 antigen-antibody complex-mediated diseases, anti-glomerular basement membrane disease, anti-phospholipid antibody syndrome, allergic neuritis, Behcet's disease/syndrome, Castleman's syndrome, Goodpasture's syndrome, Reynaud's syndrome, Sjogren's syndrome, Stevens-Johnson syndrome, pemphigoid such as pemphigoid bullous and skin pemphigoid, pemphigus (including pemphigus vulgaris, pemphigus foliaceus, pemphigus

20 mucus-membrane pemphigoid, and pemphigus erythematosus), autoimmune polyendocrinopathies, Reiter's disease or syndrome, thermal injury, preeclampsia, an immune complex disorder such as immune complex nephritis, antibody-mediated nephritis, polyneuropathies, chronic neuropathy such as IgM polyneuropathies or IgM-mediated neuropathy, autoimmune or immune-mediated thrombocytopenia such as idiopathic

25 thrombocytopenic purpura (ITP) including chronic or acute ITP, scleritis such as idiopathic cerato-scleritis, episcleritis, autoimmune disease of the testis and ovary including autoimmune orchitis and oophoritis, primary hypothyroidism, hypoparathyroidism, autoimmune endocrine diseases including thyroiditis such as autoimmune thyroiditis, Hashimoto's disease, chronic thyroiditis (Hashimoto's thyroiditis), or subacute thyroiditis,

30 autoimmune thyroid disease, idiopathic hypothyroidism, Grave's disease, polyglandular syndromes such as autoimmune polyglandular syndromes (or polyglandular endocrinopathy syndromes), paraneoplastic syndromes, including neurologic paraneoplastic syndromes such

as Lambert-Eaton myasthenic syndrome or Eaton-Lambert syndrome, stiff-man or stiff-person syndrome, encephalomyelitis such as allergic encephalomyelitis or encephalomyelitis allergica and experimental allergic encephalomyelitis (EAE), experimental autoimmune encephalomyelitis, myasthenia gravis such as thymoma-associated myasthenia gravis, cerebellar degeneration, neuromyotonia, opsoclonus or opsoclonus myoclonus syndrome (OMS), and sensory neuropathy, multifocal motor neuropathy, Sheehan's syndrome, autoimmune hepatitis, chronic hepatitis, lupoid hepatitis, giant cell hepatitis, chronic active hepatitis or autoimmune chronic active hepatitis, lymphoid interstitial pneumonitis (LIP), bronchiolitis obliterans (non-transplant) vs NSIP, Guillain-Barre syndrome, Berger's disease (IgA nephropathy), idiopathic IgA nephropathy, linear IgA dermatosis, acute febrile neutrophilic dermatosis, subcorneal pustular dermatosis, transient acantholytic dermatosis, cirrhosis such as primary biliary cirrhosis and pneumonocirrhosis, autoimmune enteropathy syndrome, Celiac or Coeliac disease, celiac sprue (gluten enteropathy), refractory sprue, idiopathic sprue, cryoglobulinemia, amyotrophic lateral sclerosis (ALS; Lou Gehrig's disease), coronary artery disease, autoimmune ear disease such as autoimmune inner ear disease (AIED), autoimmune hearing loss, polychondritis such as refractory or relapsed or relapsing polychondritis, pulmonary alveolar proteinosis, Cogan's syndrome/nonsyphilitic interstitial keratitis, Bell's palsy, Sweet's disease/syndrome, rosacea autoimmune, zoster-associated pain, amyloidosis, a non-cancerous lymphocytosis, a primary lymphocytosis, which includes monoclonal B cell lymphocytosis (e.g., benign monoclonal gammopathy and monoclonal gammopathy of undetermined significance, MGUS), peripheral neuropathy, paraneoplastic syndrome, channelopathies such as epilepsy, migraine, arrhythmia, muscular disorders, deafness, blindness, periodic paralysis, and channelopathies of the CNS, autism, inflammatory myopathy, focal or segmental or focal segmental glomerulosclerosis (FSGS), endocrine ophthalmopathy, uveoretinitis, chorioretinitis, autoimmune hepatological disorder, fibromyalgia, multiple endocrine failure, Schmidt's syndrome, adrenalitis, gastric atrophy, presenile dementia, demyelinating diseases such as autoimmune demyelinating diseases and chronic inflammatory demyelinating polyneuropathy, Dressler's syndrome, alopecia areata, alopecia totalis, CREST syndrome (calcinosis, Raynaud's phenomenon, esophageal dysmotility, sclerodactyl), and telangiectasia), male and female autoimmune infertility, e.g., due to anti-spermatozoan antibodies, mixed connective tissue disease, Chagas' disease, rheumatic fever, recurrent

abortion, farmer's lung, erythema multiforme, post-cardiotomy syndrome, Cushing's syndrome, bird-fancier's lung, allergic granulomatous angiitis, benign lymphocytic angiitis, Alport's syndrome, alveolitis such as allergic alveolitis and fibrosing alveolitis, interstitial lung disease, transfusion reaction, leprosy, malaria, parasitic diseases such as leishmaniasis,

5 kypanosomiasis, schistosomiasis, ascariasis, aspergillosis, Sampter's syndrome, Caplan's syndrome, dengue, endocarditis, endomyocardial fibrosis, diffuse interstitial pulmonary fibrosis, interstitial lung fibrosis, pulmonary fibrosis, idiopathic pulmonary fibrosis, cystic fibrosis, endophthalmitis, erythema elevatum et diutinum, erythroblastosis fetalis, eosinophilic faciitis, Shulman's syndrome, Felty's syndrome, flariasis, cyclitis such as chronic

10 cyclitis, heterochronic cyclitis, iridocyclitis (acute or chronic), or Fuch's cyclitis, Henoch-Schonlein purpura, human immunodeficiency virus (HIV) infection, SCID, acquired immune deficiency syndrome (AIDS), echovirus infection, sepsis, endotoxemia, pancreatitis, thyroxicosis, parvovirus infection, rubella virus infection, post-vaccination syndromes, congenital rubella infection, Epstein-Barr virus infection, mumps, Evan's syndrome,

15 autoimmune gonadal failure, Sydenham's chorea, post-streptococcal nephritis, thromboangitis obliterans, thyrotoxicosis, tabes dorsalis, choroiditis, giant cell polymyopathy, chronic hypersensitivity pneumonitis, keratoconjunctivitis sicca, epidemic keratoconjunctivitis, idiopathic nephritic syndrome, minimal change nephropathy, benign familial and ischemia-reperfusion injury, transplant organ reperfusion, retinal autoimmunity, joint inflammation,

20 bronchitis, chronic obstructive airway/pulmonary disease, silicosis, aphthae, aphthous stomatitis, arteriosclerotic disorders, asperniogenesis, autoimmune hemolysis, Boeck's disease, cryoglobulinemia, Dupuytren's contracture, endophthalmia phacoanaphylactica, enteritis allergica, erythema nodosum leprosum, idiopathic facial paralysis, chronic fatigue syndrome, febris rheumatica, Hamman-Rich's disease, sensoneural hearing loss,

25 haemoglobinuria paroxysmata, hypogonadism, ileitis regionalis, leucopenia, mononucleosis infectiosa, traverse myelitis, primary idiopathic myxedema, nephrosis, ophthalmia sympathica, orchitis granulomatosa, pancreatitis, polyradiculitis acuta, pyoderma gangrenosum, Quervain's thyroiditis, acquired splenic atrophy, non-malignant thymoma, vitiligo, toxic-shock syndrome, food poisoning, conditions involving infiltration of T cells,

30 leukocyte-adhesion deficiency, immune responses associated with acute and delayed hypersensitivity mediated by cytokines and T-lymphocytes, diseases involving leukocyte diapedesis, multiple organ injury syndrome, antigen-antibody complex-mediated diseases,

antiglomerular basement membrane disease, allergic neuritis, autoimmune polyendocrinopathies, oophoritis, primary myxedema, autoimmune atrophic gastritis, sympathetic ophthalmia, rheumatic diseases, mixed connective tissue disease, nephrotic syndrome, insulitis, polyendocrine failure, autoimmune polyglandular syndrome type I, 5 adult-onset idiopathic hypoparathyroidism (AOIH), cardiomyopathy such as dilated cardiomyopathy, epidermolysis bullosa acquisita (EBA), hemochromatosis, myocarditis, nephrotic syndrome, primary sclerosing cholangitis, purulent or nonpurulent sinusitis, acute or chronic sinusitis, ethmoid, frontal, maxillary, or sphenoid sinusitis, an eosinophil-related disorder such as eosinophilia, pulmonary infiltration eosinophilia, eosinophilia-myalgia 10 syndrome, Loffler's syndrome, chronic eosinophilic pneumonia, tropical pulmonary eosinophilia, bronchopneumonic aspergillosis, aspergilloma, or granulomas containing eosinophils, anaphylaxis, seronegative spondyloarthritides, polyendocrine autoimmune disease, sclerosing cholangitis, sclera, episclera, chronic mucocutaneous candidiasis, Bruton's syndrome, transient hypogammaglobulinemia of infancy, Wiskott-Aldrich syndrome, ataxia 15 telangiectasia syndrome, angiectasis, autoimmune disorders associated with collagen disease, rheumatism, neurological disease, lymphadenitis, reduction in blood pressure response, vascular dysfunction, tissue injury, cardiovascular ischemia, hyperalgesia, renal ischemia, cerebral ischemia, and disease accompanying vascularization, allergic hypersensitivity disorders, glomerulonephritides, reperfusion injury, ischemic re-perfusion disorder, 20 reperfusion injury of myocardial or other tissues, lymphomatous tracheobronchitis, inflammatory dermatoses, dermatoses with acute inflammatory components, multiple organ failure, bullous diseases, renal cortical necrosis, acute purulent meningitis or other central nervous system inflammatory disorders, ocular and orbital inflammatory disorders, granulocyte transfusion-associated syndromes, cytokine-induced toxicity, narcolepsy, acute 25 serious inflammation, chronic intractable inflammation, pyelitis, endarterial hyperplasia, peptic ulcer, valvulitis, graft versus host disease, contact hypersensitivity, asthmatic airway hyperreaction, and endometriosis.

[0184] In some embodiments, the patient is one that does not have a particular indication described above. For example, in some embodiments, the patient does not have and/or has 30 not been diagnosed with rheumatoid arthritis. In some embodiments, the patient does not

have a chronic autoimmune condition. The term chronic refers to an illness that is persisting for a long time or constantly recurring.

VI. Pharmaceutical Compositions

[0185] The present disclosure includes methods for modulating immune responses in a subject in need thereof. The disclosure includes cells and proteins that may be in the form of a pharmaceutical composition that can be used to induce or modify an immune response.

[0186] Administration of the compositions according to the current disclosure will typically be via any common route. This includes, but is not limited to parenteral, orthotopic, intradermal, subcutaneous, intramuscular, intraperitoneal, intranasal, or intravenous injection.

[0187] Typically, compositions of the invention are administered in a manner compatible with the dosage formulation, and in such amount as will be therapeutically effective and immune modifying. The quantity to be administered depends on the subject to be treated. Precise amounts of active ingredient required to be administered depend on the judgment of the practitioner.

[0188] The manner of application may be varied widely. Any of the conventional methods for administration of pharmaceutical compositions comprising cellular components are applicable. The dosage of the pharmaceutical composition will depend on the route of administration and will vary according to the size and health of the subject.

[0189] In many instances, it will be desirable to have multiple administrations of at most about or at least about 3, 4, 5, 6, 7, 8, 9, 10 or more. The administrations may range from 2-day to 12-week intervals, more usually from one to two week intervals. The course of the administrations may be followed by assays for alloreactive immune responses and T cell activity.

[0190] The phrases "pharmaceutically acceptable" or "pharmacologically acceptable" refer to molecular entities and compositions that do not produce an adverse, allergic, or other untoward reaction when administered to an animal, or human. As used herein, "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutical active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active

ingredients, its use in immunogenic and therapeutic compositions is contemplated. The pharmaceutical compositions of the current disclosure are pharmaceutically acceptable compositions.

[0191] The compositions of the disclosure can be formulated for parenteral administration, e.g., formulated for injection via the intravenous, intramuscular, sub-cutaneous, or even intraperitoneal routes. Typically, such compositions can be prepared as injectables, either as liquid solutions or suspensions; solid forms suitable for use to prepare solutions or suspensions upon the addition of a liquid prior to injection can also be prepared; and, the preparations can also be emulsified.

5 **[0192]** The pharmaceutical forms suitable for injectable use include sterile aqueous solutions or dispersions; formulations including sesame oil, peanut oil, or aqueous propylene glycol. It also should be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi.

10 **[0193]** Sterile injectable solutions are prepared by incorporating the active ingredients (i.e. cells of the disclosure) in the required amount in the appropriate solvent with various of the other ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the various sterilized active ingredients into a sterile vehicle which contains the basic dispersion medium and the required other ingredients from those enumerated above.

15 **[0194]** An effective amount of a composition is determined based on the intended goal. The term "unit dose" or "dosage" refers to physically discrete units suitable for use in a subject, each unit containing a predetermined quantity of the composition calculated to produce the desired responses discussed herein in association with its administration, i.e., the appropriate route and regimen. The quantity to be administered, both according to number of treatments and unit dose, depends on the result and/or protection desired. Precise amounts of the composition also depend on the judgment of the practitioner and are peculiar to each individual. Factors affecting dose include physical and clinical state of the subject, route of administration, intended goal of treatment (alleviation of symptoms versus cure), and potency, stability, and toxicity of the particular composition. Upon formulation, solutions will 20 be administered in a manner compatible with the dosage formulation and in such amount as is

therapeutically or prophylactically effective. The formulations are easily administered in a variety of dosage forms, such as the type of injectable solutions described above.

VII. Examples

[0195] The following examples are included to demonstrate embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Anti-IL-6R α scFv constructs inhibit IL-6 signaling

[0196] This disclosure describes single-chain variable fragments (scFvs) that bind to IL-6R α . In this Example, two different anti-IL-6R α scFv constructs are used: a V_HV_L construct (SEQ ID NO:2) and a V_LV_H construct (SEQ ID NO:1). These constructs differ only in that the positions of the V_H and V_L sequences are switched.

[0197] IL-6 is a cytokine that has pro-inflammatory effects. IL-6 signals through IL-6R α , which is the ligand-binding component of IL-6's cell surface receptor. IL-6 signaling through IL-6R α plays a role in a pro-inflammatory cytokine signaling processes, including in cytokine release syndrome. IL-6 also stimulates the inflammatory and auto-immune processes in many diseases, including diabetes, atherosclerosis, depression, Alzheimer's Disease, systemic lupus erythematosus, multiple myeloma, prostate cancer, Behcet's disease, and rheumatoid arthritis.

[0198] Both of the anti-IL-6R α scFvs are effective at inhibiting IL-6 signaling in HepG2 cells. HepG2 human hepatocarcinoma cells were seeded in a 12-well plate and incubated with 1, 2, or 5 μ g of scFv for 3 hours. Some of the wells, as indicated in **FIG. 1**, were subsequently treated with 4 ng/ml of IL-6 and incubated for another 30 minutes before cell harvest. The presence of phosphorylated STAT3 (pSTAT3), a major signaling molecule in the IL-6 signaling pathway, was probed by western blot. The results in **FIG. 1** indicate that

scFvs with either V_L-V_H (SEQ ID NO:1) or V_H-V_L (SEQ ID NO:2) orientations are able to inhibit IL-6 signaling in a dose-dependent manner, as higher concentrations of the scFvs resulted in lower levels of pSTAT3 expression in response to IL-6. Subsequent experiments were performed using the V_L-V_H scFv (SEQ ID NO:1).

5 [0199] To determine whether anti-IL-6R α scFv inhibits IL-6 signaling in primary human T cells, primary human CD4+ and CD8+ T cells were seeded in a 12-well plate and treated as described for the HepG2 cells in **FIG. 1**. The V_L-V_H scFv (SEQ ID NO:1) was added to the wells in the concentrations indicated in **FIG. 2**, along with IL-6 in the indicated wells. The presence of pSTAT3 was once again probed by western blot. The results in **FIG. 2** indicate
10 that the anti-IL-6R α scFv inhibits IL-6 signaling in a dose-dependent manner in both CD4+ and CD8+ primary human T cells.

[0200] These results indicate that administering effective amounts of anti-IL-6R α scFv to a subject would treat inflammatory conditions such as CRS and autoimmune and other diseases in which IL-6 signaling is involved.

15

Example 2

Expression of anti-IL-6R α scFv constructs by T cells

[0201] Preparation of primary human CD4+ and CD8+ cells expressing an anti-CD19 chimeric antigen receptor (CAR) and anti-IL-6R α scFv: Human primary T cells were isolated and stimulated with CD3/CD28 dynabeads on the day of isolation. Two days after isolation, 20 T cells were transduced with lentivirus or retrovirus vectors encoding anti-CD19 CAR (SEQ ID NO:88) and IL-6R α scFv constructs encoding SEQ ID NO:1 or SEQ ID NO:2 (or no IL-6R α scFv for controls). Dynabeads were removed on day 9 post isolation. Cells were sorted by magnetic cell separation based on the HA-tag on anti-CD19 CAR. On day 21 of CD4 T cells and day 14 of CD8 T cells, primary T cells were expanded by a Rapid 25 Expansion Protocol, in which 0.5 x 10⁶ of T cells are mixed with 3.5 x 10⁶ irradiated TM-LCL, 50 U/ml rhIL-2 and 1 ng/ml rhIL-15 in 25 ml of complete RPMI media in a T25 flask. Cytokines are replenished every 2 days. Cells are used for experiments after day 9 of the REP.

[0202] The inventors tested the effect of expressing the anti-IL-6R α scFv in T cells. Primary human CD4+ T cells expressing the anti-CD19 chimeric antigen receptor (CAR) with or without constitutive expression of the anti-IL-6R α scFv were co-incubated with peripheral blood mononuclear cells (PBMCs) from the same donor as the CD4+ T cells plus 5 wild-type (CD19+) Raji lymphoma cells at a 1:1:1 ratio for 12 hours. Non-scFv-expressing CD19 CAR-T cells with 90 μ g/ml of the monoclonal antibody tocilizumab were included as control samples. Levels of the cytokines in culture media as indicated in **FIG. 3** were quantified by a cytometric bead-based assay. Values shown in **FIG. 3** are the means of three technical replicates with error bars indicating \pm 1 standard deviation. Statistical significance 10 was evaluated by two-tailed, equal-variance Student's *t*-test; * p < 0.05; ** p < 0.01; *** p < 0.001. The results shown are representative of two independent experiments using cells from two different healthy blood donors. The results in **FIG. 3** show that constitutive expression 15 by primary human T cells of anti-IL-6R α scFv significantly reduces the production of inflammatory cytokines by the T cells. This indicates that the incidence of CRS in adoptive T cell therapy could be reduced by using T cells expressing an anti-IL-6R α scFv.

[0203] The inventors next tested whether expression of anti-IL-6R α scFv by T cells would inhibit their proliferation or cytotoxicity. Primary human CD4+ and CD8+ T cells constitutively expressing the anti-CD19 CAR with or without the anti-IL-6R α scFv were co-incubated with wild-type (WT; CD19+) or CD19- mutant Raji lymphoma cells at a 2:1 20 effector-to-target (E:T) ratio. Viable T-cell and Raji-cell counts were obtained by flow cytometry every two days. Both Raji cell lines stably express EGFP to enable separation from T cells during flow-cytometry analysis. After cell counting on day 8, CD4+ T cells co-incubated with WT Raji were harvested and re-challenged with fresh WT Raji cells at a 2:1 E:T ratio. Viable cell counts were monitored for another 12 days during the second challenge. 25 Values shown in **FIGS. 4 and 6** are the means of three technical replicates with error bars indicating \pm 1 standard deviation. The results shown in **FIG. 4** are representative of four independent experiments using cells from two different healthy blood donors, and the results in **FIG. 6** are representative of three independent experiments using cells from two different healthy blood donors. The results in **FIGS. 4 and 6** demonstrate that constitutive expression 30 of anti-IL-6R α scFv by T cells does not inhibit T-cell proliferation or cytotoxicity, which

indicates that the efficacy of adoptive T cell therapy would not be diminished by using T cells that express an anti-IL-6R α scFv.

[0204] To test whether expression an anti-IL-6R α scFv alters the T-cell subtype differentiation upon antigen stimulation, the T cells in the experiment described in the

5 preceding paragraph were surface-stained with antibody for CCR7, CD45RA, and CD57 to determine the T-cell subtype. Subtype T_{E/EM}: CCR7-/CD45RA-/CD57-; subtype T_{E/Exh}: CCR7-/CD45RA-/CD57+; subtype T_{EMRA}: CCR7-/CD45RA+; subtype T_{cm}: CCR7+/CD45RA-; subtype T_{naive}: CCR7+/CD45RA+. Values shown in **FIGS. 5 and 7** are the means of three technical replicates with error bars indicating ± 1 standard deviation. The 10 results shown in **FIG. 5** are representative of four independent experiments using cells from two different healthy blood donors, and the results in **FIG. 7** are representative of four independent experiments using cells from three different healthy blood donors. As can be seen from **FIGS. 5 and 7**, there were very few T_{EMRA} or T_{naive} cells in any of the samples. The results in **FIGS. 5-7** show that constitutive expression of anti-IL-6R α scFv does not alter 15 T cell subtype differentiation upon antigen stimulation, further confirming that the efficacy of adoptive T cell therapy would not be diminished by using T cells that express an anti-IL-6R α scFv.

[0205] T cell exhaustion is a state of T cell dysfunction characterized by poor effector function, expression of inhibitory receptors, and a distinct transcriptional state. Exhaustion

20 can limit the efficacy of T cells used in adoptive T cell therapy. Markers of exhaustion include PD-1, TIM-3, and LAG-3. To determine whether expression of an anti-IL-6R α scFv would affect T cell exhaustion, primary human T cells constitutively expressing the anti-CD19 CAR with or without anti-IL-6R α scFv were co-incubated with WT Raji cells at a 2:1 E:T ratio. PD-1, TIM-3, and LAG-3 expression levels were quantified by surface 25 antibody staining and flow cytometry every two days. Three trials were performed with CD8+ T cells and four trials were performed with CD4+ T cells for a total of seven independent experiments. Each trial included three technical replicates per condition. Each data point in the plots in **FIG. 8** represents the mean value of triplicates from one trial. The average value of seven trials for each time point is indicated by the black horizontal bars in 30 **FIG. 8**. The results in **FIG. 8** show that constitutive expression of anti-IL-6R α scFv does not alter exhaustion marker expression in primary human T cells upon antigen stimulation,

further confirming that expression of an anti-IL-6R α scFv would not diminish the efficacy of adoptive T cell therapy.

* * * * *

[0206] All of the methods disclosed and claimed herein can be made and executed
5 without undue experimentation in light of the present disclosure. While the compositions and
methods of this invention have been described in terms of preferred embodiments, it will be
apparent to those of skill in the art that variations may be applied to the methods and in the
steps or in the sequence of steps of the method described herein without departing from the
concept, spirit and scope of the invention. More specifically, it will be apparent that certain
10 agents which are both chemically and physiologically related may be substituted for the
agents described herein while the same or similar results would be achieved. All such similar
substitutes and modifications apparent to those skilled in the art are deemed to be within the
spirit, scope and concept of the invention as defined by the appended claims. All references,
cited literature articles, patent publications, and sequences associated with any recited
15 GenBank accession numbers are specifically incorporated herein by reference in their entirety
for all purposes.

REFERENCES

The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

5

- U.S. Patent 4,684,611
- U.S. Patent 4,879,236
- U.S. Patent 4,952,500
- U.S. Patent 5,302,523
- 10 U.S. Patent 5,322,783
- U.S. Patent 5,384,253
- U.S. Patent 5,464,765
- U.S. Patent 5,538,877
- U.S. Patent 5,538,880
- 15 U.S. Patent 5,543,293
- U.S. Patent 5,550,318
- U.S. Patent 5,563,055
- U.S. Patent 5,580,737
- U.S. Patent 5,580,859
- 20 U.S. Patent 5,589,466
- U.S. Patent 5,591,616
- U.S. Patent 5,610,042
- U.S. Patent 5,631,146
- U.S. Patent 5,656,610
- 25 U.S. Patent 5,670,373
- U.S. Patent 5,677,425
- U.S. Patent 5,702,932
- U.S. Patent 5,714,350
- U.S. Patent 5,736,524
- 30 U.S. Patent 5,780,448
- U.S. Patent 5,786,462
- U.S. Patent 5,789,215

U.S. Patent 5,871,986
U.S. Patent 5,945,100
U.S. Patent 5,981,274
U.S. Patent 5,994,624
5 U.S. Patent 6,350,861
U.S. Patent 7,479,543
U.S. Patent 8,580,264
U.S. Patent Publn 2005/0106660
U.S. Patent Publn 2006/0058510
10 U.S. Patent Publn 2006/0088908
U.S. Patent Publn 2010/0285564

Ahmad (2012) Clinical and Developmental Immunology Article ID 980250
Burton (1985) *Molec. Immunol.*, 22: 161- 206
15 Carbonelli *et al.*, *FEMS Microbiol. Lett.*, 177(1):75-82, 1999.
Chandler *et al.*, *Proc. Natl. Acad. Sci. USA*, 94(8):3596-601, 1997.
Chen and Okayama, *Mol. Cell Biol.*, 7(8):2745-2752, 1987.
Cocea, *Biotechniques*, 23(5):814-816, 1997.
Cumber *et al.*, *J. Immunology*, 149B:120-126, 1992.
20 European Patent 0 154 316
European Patent 0 322 094
European Patent 0 401 384
European Patent 0 486 525
Fechheimer, *et al.*, *Proc Natl. Acad. Sci. USA*, 84:8463-8467, 1987.
25 Fraley *et al.*, *Proc. Natl. Acad. Sci. USA*, 76:3348-3352, 1979.
Gopal, *Mol. Cell Biol.*, 5:1188-1190, 1985.
Graham and Van Der Eb, *Virology*, 52:456-467, 1973.
Harland and Weintraub, *J. Cell Biol.*, 101(3):1094-1099, 1985.
30 Harlow *et al.*, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold
Spring Harbor, N.Y., Chapter 8, 1988. Ausubel *et al.*, 1996
Huck *et al.* (1986) *Nucl. Acids Res.*
Kaepller *et al.*, *Plant Cell Rep.*, 8:415-418, 1990.

Kaneda *et al.*, *Science*, 243:375-378, 1989.

Kato *et al.*, *J. Biol. Chem.*, 266:3361-3364, 1991.

Kohl *et al.*, *Proc. Natl. Acad. Sci., USA*, 100(4):1700-1705, 2003.

Koyama 1994 *Chem. Abstr.* 120: 217262t

5 Levenson *et al.*, *Hum. Gene Ther.*, 9(8):1233-1236, 1998.

Liu *et al.* *Cell Mol. Biol.*, 49(2):209-216, 2003.

Meares *et al.*, 1984 *Anal. Biochem.* 142: 68-78

Nicolau and Sene, *Biochim. Biophys. Acta*, 721:185-190, 1982.

Nicolau *et al.*, *Methods Enzymol.*, 149:157-176, 1987.

10 Omirulleh *et al.*, *Plant Mol. Biol.*, 21(3):415-28, 1993.

Pack *et al.*, *Biochem.*, 31:1579-1584, 1992.

PCT Appln WO 94/09699

PCT Appln WO 95/06128;

PCT Publn. WO 1992019759

15 PCT Publn. WO 2006/056464

Potrykus *et al.*, *Mol. Gen. Genet.*, 199(2):169-177, 1985.

Quadri *et al.* 1993 *Nucl. Med. Biol.* 20: 559-570

Rippe, *et al.*, *Mol. Cell Biol.*, 10:689-695, 1990.

Rodwell *et al.* 1986 *PNAS USA* 83: 2632-2636;

20 Sambrook *et al.*, In: *Molecular cloning*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 2001. Skerra, *J. Mol. Recogn.*, 13:167-187, 2000.

Skerra, *J. Biotechnol.*, 74(4):257-75, 2001.

Tan *et al.* (1990) *Proc. Natl. Acad. Sci. USA* 87: 162;

Wong *et al.*, *Gene*, 10:87-94, 1980.


CLAIMS

1. T cells comprising a heterologous nucleic acid molecule encoding a chimeric interleukin 6 receptor alpha (IL-6R α) binding protein comprising a heavy chain variable region comprising CDR1 (SEQ ID NO:5), CDR2 (SEQ ID NO:6), and CDR3 (SEQ ID NO:7) attached by a heterologous linker to a light chain variable region comprising CDR4 (SEQ ID NO:8), CDR5 (SEQ ID NO:9), and CDR6 (SEQ ID NO:10), wherein the T cell expresses a chimeric antigen receptor.
2. The T cells of claim 1, wherein the binding protein is an scFv, (scFv)2, scFvFc, Fab, Fab', or F(ab)2.
3. The T cells of claim 2, wherein the binding protein is a single chain variable fragment (scFv).
4. The T cells of claim 1, wherein the heavy chain variable region is on the N-terminal side of the light chain variable region.
5. The T cells of claim 1, wherein the light chain variable region is on the N-terminal side of the heavy chain variable region.
6. The T cells of any one of claims 1-5, wherein the linker comprises the amino acid sequence GSTSGSGKPGSSEGSTKG (SEQ ID NO:11).
7. The T cells of any one of claims 1-6, further comprising a leader peptide and/or isolation tag.
8. The T cells of any one of claims 1-7, wherein the heterologous nucleic acid encodes for a leader peptide.
9. The T cells of any one of claims 1-8, wherein the heterologous nucleic acid further comprises a promoter controlling expression of the chimeric IL-6R binding protein.
10. The T cells of claim 9, wherein the promoter is constitutive.

11. The T cells of claim 9, wherein the promoter responds positively to at least one cytokine or to T-cell activation and wherein the at least one cytokine is IL-6, TNF- α , IFN- γ , IL-1 β , IL-2, IL-8, or IL-10 or the promoter responds positively to NFAT-1 or NF- κ B.
12. A method for reducing the risk of cytokine release syndrome comprising administering to a patient at risk for cytokine release syndrome a composition comprising the T cells of any one of claims 1-11.
13. Use of T cells of any one of claims 1-12 in the manufacture of a medicament for reducing the risk of cytokine release syndrome.
14. The method of claim 12 or the use of claim 13, wherein the patient has cancer.
15. The method of claim 12 or 14 or the use of claim 13, wherein the patient has or will receive adoptive T-cell therapy.
16. The method of any one of claims 12 or 14-15, or the use of claim 13, wherein the patient has or will receive lymphodepletion.
17. The method of claim 12 or the use of claim 13, wherein the patient has an autoimmune disease.
18. The method of any one of claims 12 or 14-17 or the use of claim 13, wherein the T cells are autologous.
19. The method or the use of any one of claims 12 or 14-18, further comprising administering to the patient an antihistamine, a corticosteroid, a steroid, acetaminophen, furosemide, and/or intravenous fluids.
20. The method or the use of any one of claims 12 or 14-19, or the use of claim 13, wherein the patient has one or more symptoms of cytokine release syndrome.
21. A method for reducing the toxicity of adoptive cell therapy in a cancer patient comprising administering to the patient the T cells of any one of claims 1-11, wherein the patient has been or will be treated with adoptive cell therapy.

22. Use of T cells of any one of claims 1-11 in the manufacture of a medicament for reducing the toxicity of adoptive cell therapy in a cancer patient, wherein the patient has been or will be treated with adoptive cell therapy.
23. A method for treating a cancer patient comprising administering to the patient a composition comprising the T cells of any one of claims 1-11.
24. Use of T cells of any one of claims 1-11 in the manufacture of a medicament for treating a cancer patient.

scFv condition	VL-VH scFv				VH-VL scFv				No scFv
	1ug	1ug	2ug	5ug	1ug	1ug	2ug	5ug	
IL-6 (4 ng/ml)	-	+	+	+	-	+	+	+	-



pSTAT3
88 kDa





FIG. 1

scFv concentration	CD4+ T cell						CD8+ T cell						
	0 μ g	0 μ g	1 μ g	1 μ g	2 μ g	5 μ g	90 μ g	0 μ g	0 μ g	1 μ g	2 μ g	5 μ g	90 μ g
IL-6 (4 ng/ml)	-	+	-	+	+	+	+	-	+	-	+	+	+



Relative
pSTAT3

FIG. 2




FIG. 3




FIG. 4




FIG. 5

6/8




FIG. 6

7/8




FIG. 7




FIG. 8