




(86) Date de dépôt PCT/PCT Filing Date: 2015/04/16
(87) Date publication PCT/PCT Publication Date: 2015/10/22
(45) Date de délivrance/Issue Date: 2023/09/26
(85) Entrée phase nationale/National Entry: 2016/10/13
(86) N° demande PCT/PCT Application No.: US 2015/026239
(87) N° publication PCT/PCT Publication No.: 2015/161117
(30) Priorités/Priorities: 2014/04/16 (US61/980,415);
2014/04/16 (US61/980,393)

(51) Cl.Int./Int.Cl. *C12N 15/31* (2006.01),
C07K 14/35 (2006.01), *C12M 1/34* (2006.01),
C12Q 1/02 (2006.01), *G01N 33/483* (2006.01)
(72) Inventeurs/Inventors:
NIEDERWEIS, MICHAEL, US;
PAVLENOK, MIKHAIL, US
(73) Propriétaire/Owner:
THE UAB RESEARCH FOUNDATION, US
(74) Agent: MARKS & CLERK

(54) Titre : NANOPORES MSP ET UTILISATIONS DE CEUX-CI
(54) Title: MSP NANOPORES AND USES THEREOF



(57) Abrégé/Abstract:

Provided herein are mutant single-chain Mycobacterium smegmatis porin (Msp) and uses thereof.



(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(10) International Publication Number
WO 2015/161117 A2

(43) International Publication Date
22 October 2015 (22.10.2015)

- (51) International Patent Classification: Not classified
- (21) International Application Number: PCT/US2015/026239
- (22) International Filing Date: 16 April 2015 (16.04.2015)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/980,415 16 April 2014 (16.04.2014) US
61/980,393 16 April 2014 (16.04.2014) US
- (71) Applicant: THE UAB RESEARCH FOUNDATION [US/US]; 1720 2nd Avenue South, Administration Building 770, Birmingham, Alabama 35294 (US).
- (72) Inventors: NIEDERWEIS, Michael; 1703 Roseland Drive, Homewood, Alabama 35209 (US). PAVLENOK, Mikhail; 2928 Pawnee Avenue South, Apt. No. 6, Birmingham, Alabama 35205 (US).
- (74) Agents: MCKEON, Tina Williams et al.; Kilpatrick Townsend & Stockton LLP, 1100 Peachtree Street, Suite 2800, Atlanta, Georgia 30309 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

— as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))

[Continued on next page]

(54) Title: MSP NANOPORES AND USES THEREOF




FIGURE 21

(57) Abstract: Provided herein are mutant single-chain *Mycobacterium smegmatis* porin (Msp) and uses thereof.

WO 2015/161117 A2

WO 2015/161117 A2



-
- *as to the applicant's entitlement to claim the priority of the earlier application (Rule 4.17(iii))*
- *of inventorship (Rule 4.17(iv))*
- Published:**
- *without international search report and to be republished upon receipt of that report (Rule 48.2(g))*
- *with sequence listing part of description (Rule 5.2(a))*

MSP NANOPORES AND USES THEREOF

STATEMENT OF GOVERNMENT RIGHTS

This invention was made with government support under Grant No. R01HG005115 awarded by the National Institutes of Health. The government has certain rights in the invention.

BACKGROUND

Identification and characterization of analytes often involve expensive and time-consuming methodology. For example, current technologies require expensive custom-made reagents and/or detection methods to determine nucleic acid sequences. Similarly, methods for determining protein sequences, for example, peptide fingerprinting by mass spectrometry technologies, can be laborious and costly. Therefore, efficient methods of detecting and analyzing nucleic acids, proteins and other analytes are necessary.

SUMMARY

Provided herein are nucleic acid sequences encoding a mutant single-chain *Mycobacterium smegmatis* porin (Msp). The nucleic acid sequence optionally comprises a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence; and a third nucleotide sequence encoding an amino acid linker sequence. In some of the mutant single-chain Msps provided herein, at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence that comprises a mutation at position 97. In other mutant single-chain Msps provided herein, at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence comprising one or more mutations at any of the following amino acid positions: I68, S73, S116, P123 or V128.

Optionally, the nucleic acid sequence encoding the mutant single-chain Msp comprises a first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence or any subset thereof, and the first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequences encode a first, second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, respectively. The nucleic acid sequence further comprises a ninth nucleotide sequence encoding an amino acid linker sequence which is optionally present at

multiple sites (e.g., between the Msp monomer sequences). The first Msp monomer sequence can be a mutant Msp monomer sequence that comprises one or more mutations at any of the following amino acid positions: I68, S73, S116, P123 or V128 or a mutant Msp monomer sequence that comprises a mutation at position 97.

Further provided herein is a nucleic acid encoding a mutant Msp monomer. The Msp monomer comprises a mutation at one or more of the following positions: I68, S73, P97, S116, P123 or V128.

Further provided are polypeptides comprising the mutant Msps described herein and polypeptides encoded by the nucleic acids described herein. Also provided herein is a system comprising a mutant Msp described herein, wherein the Msp has a vestibule and a constriction zone that define a tunnel, wherein the tunnel is positioned between a first conductive liquid medium and a second conductive liquid medium, wherein at least one conductive liquid medium comprises an analyte, and wherein the system is operative to detect the analyte, when the system is subjected to an electric field sufficient to translocate the analyte from one conductive liquid medium to the other.

Further provided are methods for detecting the presence of an analyte. The methods include applying an electric field sufficient to translocate an analyte from a first conductive medium to a second conductive medium in liquid communication through a mutant Msp described herein and measuring an ion current, wherein a reduction in the ion current indicates the presence of the analyte in the first medium.

DESCRIPTION OF THE DRAWINGS

Figure 1 shows an alignment of the MspA, MspB, MspC and MspD monomers of *Mycobacterium smegmatis*. The numbering of each protein starts with the first amino acid of the mature portion of the sequence. The MspA, MspB, MspC and MspD monomer sequences without a signal/leader sequence are provided as SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 and SEQ ID NO: 4, respectively.

Figure 2 shows schematic representations of the plasmids used for construction of ML712 *M. smegmatis*. pML1611 – mspB is a deletion vector: *mspBup*, *mspBdown* – regions homologous upstream and downstream of to mspB chromosomal gene; *loxP* – recombination site for Cre recombinase; *hyg* – hygromycin resistance; *sacB* – levansucrase; *xylE* - catechol-2,3-dioxygenase; *gfp2+* - green fluorescent protein; *tsPAL5000*: temperature-sensitive origin of replication for mycobacteria; *ColE1* – *E. coli*

origin of replication. pML2622 – integrative plasmid for expression of the N-terminal channel-forming domain (d1) of *cpnT*. *hyg*: hygromycin resistance gene; *pBR322-Ori*: *E. coli* origin of replication; *pNIT1* and *pNIT2* are required for nitrile-inducible expression of *cpnT*, *attP*: chromosome attachment site of the phage L5; *int*: L5 integrase; *FRT*: Flp recombines site.

Figure 3 is a Southern blot of *msp* loci in *M. smegmatis* strains. Chromosomal DNA was isolated from SMR5 (1), ML16 (2), and ML712 (3) *M. smegmatis* strains. DNA fragments were separated on 1% agarose gel, blotted onto a nitrocellulose membrane, and detected using specific probes labeled with digoxigenin. The sizes of the DNA fragments that hybridized with the probe were consistent with predicted values. Expected fragment sizes: *mshA* locus: 1 – 1500bp, 2,3 – 1250bp; *mshB* locus: 1,2 – 1140bp, 3 – 1850bp; *mshC* locus: 1 – 1200bp, 2,3 – 2100bp; *mshD* locus: 1 – 1730bp, 2,3 – 1090bp.

Figure 4 is a Western blot of Msp porin expression in *M. smegmatis* porin gene deletion mutants. Msp porins were extracted with 0.5% octylpolyethyleneoxide and 10 µl were loaded in each lane. Proteins were separated on 8% SDS-PAGE and detected by Western blot using a polyclonal antibody against MspA. Lanes: 1, SMR5 (wt; expression of *mshA*); 2, ML16 (triple porin deletion mutant; *mshB* expression); 3, ML712/pMS2 (quadruple porin deletion mutant; empty vector); 4, ML712/pMN016 (+ *mshA* expression plasmid); 5, ML712/pML904 (+ M1 *mshA* expression plasmid).

Figure 5 is a schematic representation of the mycobacterial expression plasmid pML3213 encoding single-chain M1₈-MspA. The plasmid comprises the following: *hyg*, hygromycin resistance gene; *ory myc*, a mycobacterial origin of replication; *COLE1 ORI*, an *E. coli* origin of replication; *psmyc*, constitutive mycobacterial promoter; *m1-1*, *m1-2*, etc., *m1 mshA* genes with an index number corresponding to position in the single-chain gene construct. Tetrad A is flanked by PacI at the beginning of the first *m1-mshA* and by MluI at the end of the fourth *m1-mshA*; tetrad B is flanked by EcoRV at the beginning of the fifth *m1-mshA* and by HindII at the end of the eighth *m1-mshA*. Individual *m1-mshA* genes within the tetrads are flanked by KpnI, NsiI, NdeI, PstI, ScaI, NheI. Connecting (GGGGS)₃ (SEQ ID NO: 5) linkers are located between restriction sites, but are not shown.

Figure 6 is a Western blot showing the analysis of single-chain MspA expression in *M. smegmatis* ML714. Msp porins were extracted with 0.5% octylpolyethyleneoxide and 10 µl were loaded in each lane. Proteins were separated in 10% SDS-PAGE followed by transfer onto PVDF membrane, and probed with αMspA monoclonal antibodies. The lanes are

marked as follows: M, protein ladder; 1, expression wt MspA (SMR5); 2, M1₂-MspA; 3, M1₄-MspA; 4, M1₈-MspA (single-chain M1-MspA) The amount of MspA was determined by quantitative image analysis.

Figure 7 shows the thermal stability of single-chain M1₈-MspA. Single-chain M1 MspA porins were extracted with 0.5% n-octylpolyethyleneoxide. 15 μ l of the sample were incubated in a buffer containing 2% SDS for 15 min at temperatures indicated above each lane. Proteins were separated in 8% SDS PAGE followed by transfer onto PVDF membrane and probed with α MspA polyclonal antibodies. The lanes are marked as follows: M, protein ladder; RT, room temperature; 40, 50, etc. indicate incubation temperature. Equal amounts of the protein samples were loaded onto each lane.

Figure 8A shows the results of lipid bilayer experiments with M1₈-MspA. Approximately 70 ng of protein were added to the bilayer chamber. Current traces of M1₈-MspA in a diphtanoylphosphatidylcholine lipid bilayer were recorded in 1M KCl at a potential of -10 mV. This resulted in the step-wise increase in the current across the lipid bilayer indicating the insertion of M1₈-MspA channels into the membrane.

Figure 8B is a histogram of the conductance of M1₈-MspA. 269 pores in four different membranes were analyzed. Single-chain M1₈-MspA showed a predominant conductance peak at 1.1nS.

Figure 9 shows the results of voltage gating experiments. M1₈-MspA was added to the *cis*-side of a diphtanoylphosphatidylcholine membrane. Increasingly positive (*upper traces*) and negative (*lower traces*) voltages were applied to the membrane when ~220 channels were reconstituted into the membrane. The membrane current was recorded at each applied voltage. The critical voltage at which the channels began to close (V_c) was determined as the voltage when the conductance of the lipid bilayer decreased after its initial maximal value. The critical voltage V_c of M1₈-MspA was +90 mV. The M1₈-MspA channel was stable at all applied negative voltages.

Figure 10A shows an example of a positive ramp created in a single-chain Msp comprising a first mutant Msp monomer that comprises mutations at positions D56, I68, S73, D118, D134 and E139 and a seventh mutant Msp monomer that comprises a mutation at positions L88 and I105.

Figure 10B shows a top view of a positive ramp created in a single-chain Msp comprising a first mutant Msp monomer that comprises mutations at positions D56, I68, S73, D118, D134 and E139 and a seventh mutant Msp monomer that comprises a mutation at positions L88 and I105.

Figure 10C is a schematic of a single-chain Msp. Numbers under subunits #1 and #7 represent locations of the positive ramp.

Figure 11 is a schematic representation of plasmid pML3216, for the expression of single-chain *m2-mspA* in *E. coli*. *bla*, ampicillin resistance gene; *pBR322-Ori*, *E. coli* origin of replication; *lacI*, lac repressor protein; *T7 P*, T7 promoter; *SD*, Shine-Dalgarno sequence; *m2-1 – m2-8*, *m2 mspA* codon.

Figure 12 shows that *scm2-mspA* can be produced in *E. coli*. *E. coli* Omp8 cells were induced with 1.5 mM IPTG at OD600 of 0.5. At different time points, cells were collected and lysed. Equal amounts of protein sample were loaded onto 10% polyacrylamide gel. After electrophoresis, the gel was stained with Coomassie and scanned for quantification by LabWorks software (Waltham, Massachusetts). Lanes: *M*, molecular weight marker with masses indicated on the left (kDa); *Un*, non-induced cells; 2, 4 and 6 hours after induction with IPTG.

Figure 13 shows a Western blot of scMspA M2 refolding. Samples before and after refolding were loaded onto 8% polyacrylamide gel followed by overnight transfer onto PVDF membrane. The membrane was stained with MspA-specific rabbit antibodies. Lanes: *M*, molecular weight marker with masses indicated on the left (kDa); *Un*, unfolded sample after anion exchange chromatography; *F*, folded scMspA M2.

Figures 14(A-B) show the channel-forming properties of scMspA M2. (A) Current trace recording of scMspA M2 in planar lipid bilayer. After addition of refolded scMspA M2 protein as shown in Fig. 3 a step-wise current increase is observed indicative of channel insertions. (B) Histogram of single-channel conductance distribution. A total of 392 channels from 8 membranes were analyzed. The dotted line represents a Gaussian fit of the data distribution. The major conductance of scMspA M2 is 2.3 nS.

Figure 15 shows voltage-dependent channel closure of scMspA M2. Protein was added to the *cis*-side of the cuvette. After insertion of approximately 200 pores, positive or negative voltage was increased in 10 mV increments and current was recorded for 2 minutes. A flat line represents open pores, whereas a sloping line represents closing of the pores. Critical voltages for scMspA M2 were determined to be about +80mV/-70mV.

Figure 16 shows a schematic representation of plasmid pML3222, for the expression of single-chain *scmspA PNI* (*scmspA PNI*) in *E. coli*. *bla*, ampicillin resistance gene; *pBR322-Ori*, *E. coli* origin of replication; *lacI*, lac repressor protein; *T7 P*, T7 promoter; *SD*, Shine-Dalgarno sequence; *m2-97-1 – m2-97-8*, codon optimized genes of *m2 mspA* with a P97F mutation.

Figure 17 shows that *scmspA* PN1 can be produced in *E. coli*. *E. coli* Omp8 cells were induced with 1.5 mM IPTG at OD₆₀₀ of 0.5. At 2 hours post induction cells were collected and lysed. Equal amounts of protein sample were loaded onto 8% polyacrylamide gel. After electrophoresis, the gel was stained with Coomassie and scanned for quantification by LabWorks software (Waltham, Massachusetts). Lanes: *M*, molecular weight marker with masses indicated on the left (kDa); *Un*, non-induced cells; 2 hours after induction with IPTG.

Figure 18 shows a Western blot of scMspA PN1 refolding. Samples were loaded onto 8% polyacrylamide gel followed by staining with Coomassie and scanned for quantification by LabWorks software. Lanes: *M*, molecular weight marker with masses indicated on the left (kDa); IB, inclusion bodies purified from Omp8 *E. coli*; AE, sample after anion exchange chromatography; D, sample after dialysis; R, folded scMspA PN1 protein.

Figure 19 shows that scMspA PN1 is inserted into lipid membranes. DPhPC liposomes were loaded with 30mM carboxyfluorescein dye. Dye release was monitored at 517 nm emission (492 nm excitation) using Varian Cary fluorimeter (Palo Alto, CA). Symbols: open squares - Triton X-100 (0.1% v/v); closed circles – LDAO (0.1% v/v); open circles - OPOE (0.5% v/v); closed triangles – wtMspA (60 ng/ml); closed squares - scMspA PN1 (60 ng/ml), open triangles - scMspA M2 (120 ng/ml). No significant dye release was observed when liposomes were only in PBS buffer.

Figure 20 shows the distribution of pores' reconstitution time into artificial DPhPC membrane. A cuvette was filled with KCl in the indicated concentrations. Protein samples of 100 ng/ml final were added to the both sides. -10mV potential was applied and the data were recorded with TestPoint software. Each dot represents time of the first pore insertion. Median insertion time for scMspA PN1 in 1M KCl was 399 seconds (9 membranes with 89% successful insertion events analyzed). Median insertion time for scMspA PN1 in 0.3MKCl was 859 seconds (8 membranes with 50% successful insertion events analyzed). Median insertion time for scMspA M2 in 1M KCl was 695 seconds (10 membranes with 40% successful insertion events analyzed). In 0.3M KCl scMspA M2 had insertion time of 1270 seconds (8 membranes, 12% successful insertion). * - $P = 0.028$ as determined by Mann-Whitney Rank Sum test.

Figures 21(A-D) show single-channel conductances of scMspA PN1 and scMspA M2 in 1.0M KCl. (A) Current trace recording of scMspA PN1 in planar lipid bilayer. After addition of refolded scMspA PN1 protein a step-wise current increase is observed indicative of channel insertions. (B) Histogram of single-channel conductance distribution. A total of 137 channels from 4 membranes were analyzed. The dotted line represents a Gaussian fit of

the data distribution. The major conductance of scMspA PN1 is 2.0 nS. (C) Current trace recording of scMspA M2 in planar lipid bilayer. (D) Histogram of single-channel conductance distribution. A total of 238 channels from 6 membranes were analyzed. The dotted line represents a Gaussian fit of the data distribution. The major conductance of scMspA M2 is 1.3 nS.

Figure 22 shows single channel conductance of single-chain MspA PN1 at 0.3M KCl and 1.0M KCl and the increase of channel insertions of scMspA PN1 into membranes at 0.3M KCl after it was in contact with membranes.

DETAILED DESCRIPTION

Provided herein are mutant *Mycobacterium smegmatis* porins (Msp). A mutant Msp can be a multimer complex comprised of two or more Msp monomers, wherein at least one of the monomers is a mutant Msp monomer. An Msp monomer is encoded by a gene in *Mycobacterium smegmatis*. *Mycobacterium smegmatis* has four identified Msp genes, denoted MspA, MspB, MspC, and MspD. An alignment of the wild-type polypeptide sequences for the MspA, MspB, MspC and MspD monomers of *Mycobacterium smegmatis* is shown in Figure 1. The numbering of each protein starts with the first amino acid of the mature portion of the sequence, as indicated by the number “1” above the first amino acid of the mature amino acid sequence. The amino acid sequences for a MspA, MspB, MspC and a MspD monomer without a signal sequence, i.e., the mature portion of the sequence, are provided as SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 and SEQ ID NO: 4, respectively. The amino acid sequences for a MspA, MspB, MspC and a MspD monomer with a signal/leader sequence are provided as SEQ ID NO: 6, SEQ ID NO: 7, SEQ ID NO: 8 and SEQ ID NO: 9, respectively.

Further, sequences of wild-type Msp monomers that can be modified are disclosed in GenBank. For example, the nucleotide and amino acid sequences of a wild-type MspA monomer can be found at GenBank Accession Nos. AJ001442 and CAB56052, respectively. The nucleotide and amino acid sequences of a wild-type MspB monomer can be found, for example, at GenBank Accession Nos. NC_008596.1 (from nucleotide 600086 to 600730) and YP_884932.1, respectively. The nucleotide and amino acid sequences of a wild-type MspC monomer can be found, for example, at GenBank Accession Nos. AJ299735 and CAC82509, respectively. The nucleotide and amino acid sequences of a wild-type

MspD monomer can be found, for example, at GenBank Accession Nos. AJ300774 and CAC83628, respectively.

A mutant Msp monomer can be a full-length monomer or a functional fragment thereof encoded by a MspA, MspB, MspC or MspD-encoding nucleic acid, for example, an mRNA or a genomic sequence encoding MspA, MspB, MspC or MspD, wherein the monomer comprises one or more modifications.

Optionally, a mutant Msp is a mutant single-chain Msp or is a multimer of several single-chain Msps, wherein the multimer comprises at least one mutant single-chain Msp. A mutant Msp can also be a multimer of several Msp monomers wherein at least one Msp monomer is a mutant Msp monomer.

A single-chain Msp can, for example, comprise a multimer formed by two or more Msp monomers (e.g., eight monomers) connected by one or more amino acid linker peptides. A partial single-chain Msp refers to a single-chain multimer complex that dimerizes, trimerizes, or the like to form a porin. A full single-chain Msp porin refers to a single-chain multimer complex that forms a porin without the need to dimerize, trimerize or the like to form a porin. Stated differently, the single-chain folds to form a porin, but all components are in one amino acid chain, as compared to a porin that must associate with other partial single-chain Msp(s) or monomeric Msp monomers to form a porin.

Mutant single-chain Msps and the Nucleic Acids Encoding Them

Provided herein are nucleic acid sequences encoding mutant single-chain Msps. For example, the nucleic acid sequence encoding a mutant single-chain Msp comprises: (a) a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence; and (b) a third nucleotide sequence encoding an amino acid linker sequence, wherein at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence has a mutation at position P97. Optionally, the mutant Msp monomer sequence can comprise a mutation at P97, wherein the mutation is not a P97S mutation or a P97C mutation. Optionally, the mutant Msp monomer sequence can comprise a P97F mutation. As shown in the Examples, additional hydrophobic residues, for example, phenylalanine, located in loop 6 of scMspA (amino acids 91-103) promote faster and more efficient insertion of the pores into lipid bilayers. For a description of loop 6 of MspA and residues contained therein, see Hufte et al., *J. Biol. Chem.* 284: 10223-10231 (2009).

Therefore, provided herein is a single chain Msp comprising one or more hydrophobic substitutions in loop 6 (amino acids 91-103) of Msp.

For example, provided herein is a nucleic acid sequence encoding a mutant single-chain *Mycobacterium smegmatis* porin (Msp), wherein the nucleic acid sequence comprises (a) a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence and (b) a third nucleotide sequence encoding an amino acid linker sequence, wherein at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence comprising one or more mutations at any of amino acid positions D91, G92, D93, I94, T95, A96, P97, P98, F99, G100, L101, N102 or S103, wherein one or more of D91, G92, D93, I94, T95, A96, P97, P98, F99, G100, L101, N102 or S103 is substituted with a hydrophobic amino acid. For example, hydrophobic amino acids can be selected from the group consisting of alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tyrosine, tryptophan, proline and glycine. By way of example, and not to be limiting, a mutant Msp monomer sequence can comprise (i) a D90N, a D91N and a D93N mutation; and one or more of (ii) a G92F, T95F, A96F, P97F, P98F, G100F, L101F, N102F or S103F mutation. As set forth above, substitutions at position G92, T95, A96, P97, P98, G100, L101, N102 or S103 are not limited to phenylalanine, as one or more of these amino acids can be replaced with another hydrophobic residue, for example, alanine, valine, leucine, isoleucine, proline, methionine, tyrosine, tryptophan, proline and glycine. One or more of G92, T95, A96, P97, P98, G100, L101, N102 or S103 can be substituted with the same hydrophobic amino acid or different hydrophobic amino acids.

As used throughout, a mutation at a specific amino acid is indicated by the single letter code for the amino acid at a position, followed by the number of the amino acid position in an Msp polypeptide sequence (for example, an amino acid position in SEQ ID NO: 1), and the single letter code for the amino acid substitution at this position. Therefore, it is understood that a P97 mutation is a proline to phenylalanine substitution at amino acid 97 of SEQ ID NO: 1. Similarly, a D90N mutation is an aspartic acid to arginine substitution at amino acid 90 of SEQ ID NO: 1, a D91N mutation is an aspartic to arginine substitution at amino acid 91 of SEQ ID NO: 1, etc. It is also understood that amino acids corresponding to positions in SEQ ID NO: 1 are also provided herein (See Figure 1). For example, and not to be limiting, one of skill in the art would understand that, the corresponding amino acid for

E139 of SEQ ID NO: 1 in MspB (SEQ ID NO:2), MspC (SEQ ID NO: 3) and MspD (SEQ ID NO: 4) is A139, A139 and K138, respectively.

Optionally, any mutant Msp monomer sequence described herein can further comprise a mutation at amino acid position D118, a mutation at position D134 or a mutation at position E139. Optionally, a mutation at position E139 can be an E to R (arginine) or an E to K (lysine) substitution. Optionally, a mutation at position D118 can be a D to R substitution or a D to K substitution. Optionally, a mutation at position D134 can be a D to R substitution or a D to K substitution. For example, any mutant Msp monomer sequence described herein can comprise one or more mutations selected from the group consisting of: a D118R mutation, a D134R mutation and a E139K mutation. Optionally, any mutant Msp monomer sequence described herein can further comprise at least one of (i) a mutation at position 93 and (ii) a mutation at position D90, position D91 or both positions D90 and D91. Optionally, the amino acid at position 90, 91 or 93 is substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine or tryptophan. Optionally, any mutant Msp monomer sequence described herein can further comprise a D90N, a D91N and a D93N mutation.

For example, a mutant Msp monomer sequence comprising a mutation at position 97 can further comprise (i) a mutation at amino acid position D118, D134 and/or E139 (ii) a mutation at position D93, and/or (iii) a mutation at position D90, position D91 or both positions D90 and D91. For example, a mutant MspA monomer sequence can comprise a D90N mutation, a D91N mutation, a D93N mutation, a P97F mutation, a D118R mutation, a D134R mutation and a E139K mutation. The mutant MspA monomer sequence can also comprise a D90N mutation, a D91N mutation, a D93N mutation, a P97F mutation, a D118R mutation, a D134R mutation and a E139K mutation.

Also provided herein is a nucleic acid sequence encoding a mutant single-chain Msp which comprises (a) a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence, and (b) a third nucleotide sequence encoding an amino acid linker sequence, wherein at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence comprising one or more mutations at any of the following amino acid positions: I68, S73, S116, P123 or V128. Provided herein is a nucleic acid encoding a mutant single-chain Msp, wherein the mutant Msp monomer sequence comprises one or more mutations at any of I68, S73, S116, P123 or V128, wherein the mutation is not a I68V mutation or a S73C mutation.

Provided herein is a mutant Msp monomer sequence comprising one or more mutations at I68, S73, S116, P123 and/or V128 further comprising (i) a mutation at amino acid position D118, (ii) a mutation at position D93, and/or (iii) a mutation at position D90, position D91 or both positions D90 and D91. For example, a mutant MspA monomer sequence can comprise one or more mutations at amino acid positions I68, S73, S116, P123 or V128, a mutation at D93, a mutation at D118, a mutation at D134 and a mutation at E139 or any subset thereof. In another example, a mutant MspA monomer sequence can comprise one or more mutations at amino acid positions I68, S73, S116, P123 or V128, a mutation at amino acid position D118, a mutation at D134, a D90N mutation and/or a D91N mutation. In yet another example, a mutant MspA monomer sequence can comprise one or more mutations at amino acid positions I68, S73, S116, P123 or V128, a mutation at amino acid position D118, a mutation at D134 and a mutation at E139, a D90N mutation, a D91N mutation and a D93N mutation. Provided herein is a nucleic acid encoding a mutant single-chain Msp, wherein the mutant Msp monomer sequence comprises one or more mutations at any of I68, S73, S116, P123 or V128, wherein the mutation is not a I68V mutation or a S73C mutation. In any of the mutant single-chain Msps provided herein, the mutant Msp monomer sequence can comprise one or more mutations at any of I68, S73, S116, P123 or V128, wherein the mutation is not a I68V mutation or a S73C mutation.

Optionally, any of the mutant Msp monomer sequences described herein can further comprise one or more mutations at any of the following amino acid positions: D13, A55, D56, E57, F58, E63, S136, G137 or D172. Optionally, one or more of D13, A55, D56, E57, F58, E63, S136, G137 or D172 in a mutant Msp monomer sequence provided herein can be substituted with lysine or arginine. Optionally, any mutant Msp monomer sequence described herein comprising one or more mutations at D13, A55, D56, E57, F58, E63, S136, G137 or D172 can further comprise one or more mutations at the following positions: D118, D134 or E139. Optionally, any mutant Msp monomer sequence described herein comprising one or more mutations at D13, A55, D56, E57, F58, E63, S136, G137 or D172 can further comprise a mutation at position 93, and/or a mutation at position 90, position 91 or both positions 90 and 91.

Therefore, provided herein is a nucleic acid sequence encoding a mutant single-chain *Mycobacterium smegmatis* porin (Msp), wherein the nucleic acid sequence comprises (a) a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence, and (b) a third nucleotide sequence encoding an amino acid linker sequence,

wherein at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence comprising one or more mutations at any of the following amino acid positions: D13, A55, D56, E57, F58, E63, S136, G137 or D172. Optionally, the third nucleotide sequence encoding the linker is located between the first and second nucleotide sequence.

Also provided is a nucleic acid sequence encoding a mutant single-chain *Mycobacterium smegmatis* porin (Msp), wherein the nucleic acid sequence comprises: (a) a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence, and (b) a third nucleotide sequence encoding an amino acid linker sequence, wherein at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence comprising (i) a mutation at position 93, and/or (ii) a mutation at position 90, position 91 or both positions 90 and 91 and (iii) one or more mutations at any of the following amino acid positions: D13, A55, D56, E57, F58, E63, S136, G137 or D172.

Further provided is a nucleic acid encoding a mutant Msp monomer, wherein the Msp monomer comprises a mutation at one or more of the following positions: D13, A55, D56, E57, F58, E63, S136, G137 or D172. Optionally, a mutant Msp monomer sequence comprising one or more mutations at D13, A55, D56, E57, F58, E63, S136, G137 or D172 can further comprise a mutation at position 93 and/or a mutation at position 90, position 91 or both positions 90 and 91. Optionally, a mutant Msp monomer sequence comprising one or more mutations at positions D13, A55, D56, E57, F58, E63, S136, G137 or D172 can further comprise a D90N, a D91N and a D93N mutation.

In the mutant single-chain mutant Msps provided herein, the first monomer sequence can be any mutant monomer sequence described herein. For example, the mutant monomer sequence can be a mutant MspA sequence. The second monomer can be selected from the group consisting of a wildtype Msp monomer, a second mutant Msp monomer, a wild-type Msp paralog or homolog monomer, and a mutant Msp paralog or homolog monomer. It is understood that the second mutant Msp monomer can be the same or different than the first mutant Msp monomer. These include, but are not limited to, MspA/Msmeg0965, MspB/Msmeg0520, MspC/Msmeg5483, MspD/Msmeg6057, MppA, PorM1, PorM2, PorM1, Mmcs4296, Mmcs4297, Mmcs3857, Mmcs4382, Mmcs4383, Mjls3843, Mjls3857, Mjls3931, Mjls4674, Mjls4675, Mjls4677, Map3123c, Mav3943, Mvan1836, Mvan4117, Mvan4839, Mvan4840, Mvan5016, Mvan5017, Mvan5768, MUL_2391, Mflv1734, Mflv1735, Mflv2295, Mflv1891, MCH4691c, MCH4689c, MCH4690c, MAB1080, MAB1081,

MAB2800, RHA1 ro08561, RHA1 ro04074, and RHA1 ro03127. A wild-type MspA paralog or homolog monomer may be a wild-type MspB monomer. Wild-type MspA paralog and homolog monomers are well-known in the art. Table 1 provides a non-limiting list of such paralogs and homologs.

Table 1. Wild-type MspA and Wild-type MspA paralogs and homolog monomers

Protein#	Organism	Identity/ Similarity to MspA (%)	Length (aa)	Reference
MspA/Msmeg0965	<i>M. smegmatis</i>	100/100	211	gb ABK74363.1 , (Stahl <i>et al.</i> , 2001)*
MspB/Msmeg0520	<i>M. smegmatis</i>	94/95	215	gb ABK73437.1 , (Stahl <i>et al.</i> , 2001)*
MspC/Msmeg5483	<i>M. smegmatis</i>	93/95	215	gb ABK74976.1 , (Stahl <i>et al.</i> , 2001)*
MspD/Msmeg6057	<i>M. smegmatis</i>	82/89	207	gb ABK72453.1 , (Stahl <i>et al.</i> , 2001)*
MppA	<i>M. phlei</i>	100/100	211	AJ812030, (Dorner <i>et al.</i> , 2004)**
PorM1	<i>M. fortuitum</i>	95/96	211	emb CAI54228.1
PorM2	<i>M. fortuitum</i>	91/93	215	emb CAL29811.1
PorM1	<i>M. peregrinum</i>	94/96	211	emb CAI54230.1
Mmcs4296	<i>Mycobacterium sp. MCS</i>	85/91	216	gb ABG10401.1
Mmcs4297	<i>Mycobacterium sp. MCS</i>	85/91	216	gb ABG10402.1
Mmcs3857	<i>Mycobacterium sp. MCS</i>	30/44	235	gb ABG09962.1
Mmcs4382	<i>Mycobacterium sp. MCS</i>	85/91	216	gb ABL93573.1
Mmcs4383	<i>Mycobacterium sp. MCS</i>	85/91	216	gb ABL93574.1
Mjls3843	<i>Mycobacterium sp. JLS</i>	26/40	235	gb ABN99619.1
Mjls3857	<i>Mycobacterium sp. JLS</i>	26/40	235	gb ABG09962.1
Mjls3931	<i>Mycobacterium sp. JLS</i>	26/40	235	gb ABL93123.1
Mjls4674	<i>Mycobacterium sp. JLS</i>	85/89	216	gb ABO00440.1
Mjls4675	<i>Mycobacterium sp. JLS</i>	83/89	216	gb ABO00441.1
Mjls4677	<i>Mycobacterium sp. JLS</i>	84/89	216	gb ABO00443.1

Map3123c	<i>M. avium paratuberculosis</i>	24/39	220	gb AAS05671.1
Mav3943	<i>M. avium</i>	24/39	227	gb ABK66660.1
Mvan1836	<i>M. vanbaalenii</i> PYR-1	82/88	209	gb ABM12657.1
Mvan4117	<i>M. vanbaalenii</i> PYR-1	32/43	239	gb ABM14894.1
Mvan4839	<i>M. vanbaalenii</i> PYR-1	83/88	209	gb ABM15612.1
Mvan4840	<i>M. vanbaalenii</i> PYR-1	83/89	209	gb ABM15613.1
Mvan5016	<i>M. vanbaalenii</i> PYR-1	30/41	238	gb ABM15788.1
Mvan5017	<i>M. vanbaalenii</i> PYR-1	25/35	227	gb ABM15789.1
Mvan5768	<i>M. vanbaalenii</i> PYR-1	21/32	216	gb ABM16533.1
MUL_2391	<i>M. ulcerans</i> Agy99	21/34	233	gb ABL04749.1
Mflv1734	<i>M. gilvum</i> PYR-GCK	21/32	225	gb ABP44214.1
Mflv1735	<i>M. gilvum</i> PYR-GCK	32/41	226	gb ABP44215.1
Mflv2295	<i>M. gilvum</i> PYR-GCK	25/40	250	gb ABP44773.1
Mflv1891	<i>M. gilvum</i> PYR-GCK	84/90	217	gb ABP44371.1
MCH4691c	<i>M. chelonae</i>	70/80	223	gb ACV04474.1
MCH4689c	<i>M. chelonae</i>	66/78	223	gb ACV04472.1
MCH4690c	<i>M. chelonae</i>	72/81	217	gb ACV04473.1
MAB1080	<i>M. abscessus</i>	69/79	223	emb CAM61170.1
MAB1081	<i>M. abscessus</i>	68/78	222	emb CAM61171.1
MAB2800	<i>M. abscessus</i>	27/44	246	emb CAM62879.1
RHA1 ro08561	<i>Rhodococcus jostii</i> RHA1	34/51	233	gb ABG99605.1
n.d.	<i>Rhodococcus opacus</i> B4	34/51	233	gbj BAH52196.1
RHA1 ro04074	<i>Rhodococcus</i> sp. RHA1	34/50	233	gb ABG95871.1
RHA1 ro03127	<i>Rhodococcus</i> sp. RHA1	34/50	233	gb ABG94930.1
n.d.	<i>Rhodococcus erythropolis</i> PR4	35/50	229	gbj BAH30938.1

Only proteins with significant amino acid similarities over the full length of the protein were included. Data were obtained by PSI-Blast algorithm (BLOSUM62 matrix) using the NIH GenBank database on the world wide web at ncbi.nlm.nih.gov/blast/Blast.cgi.

n. d.: "not determined"

*Stahl et al., Mol. Microbial. 40:451 (2001)

**Domer et al., Biochim. Biophys. Acta. 1667:47-55 (2004)

As used herein, a mutant single-chain Msp is a polypeptide comprising at least two Msp monomers, or functional fragments thereof, connected by one or more amino acid linker

peptides wherein at least one of the Msp monomers is a mutant Msp monomer. For example, the mutant single-chain Msp can comprise two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, or more Msp monomers connected by one or more amino acid linker peptides, wherein at least one of the Msp monomers is a mutant Msp monomer. As set forth above, a single-chain mutant Msp can form a porin, for example, by folding, without the need to dimerize, trimerize or the like to form a porin. Alternatively, a mutant single-chain Msp can be a partial single-chain mutant Msp comprising at least two Msp monomers or fragments thereof connected by one or more amino acid linked peptides, that can dimerize, trimerize or the like to form a porin.

Optionally, a Msp porin comprising a mutant single-chain Msp can, for example, comprise two or more single-chain Msp porin dimers, two or more single-chain Msp porin trimers, two or more single-chain Msp porin tetramers, two or more single-chain Msp porin pentamers, one or more single-chain Msp porin hexamers, one or more single-chain Msp porin septamers, one or more single-chain Msp porin octamers, or combinations thereof. For example, a Msp porin can comprise a single-chain Msp porin dimer and two single-chain Msp porin trimers. By way of another example, a Msp porin can comprise a single-chain Msp porin tetramer and two single-chain Msp porin dimers.

Amino acid linker sequences are described herein. In any single-chain Msp described herein, a linker sequence can, for example, comprise 10 to 20 amino acids. For example, an amino acid linker sequence comprises 15 amino acids. Optionally, the amino acid linker sequence comprises a (GGGGS)₃ (SEQ ID NO: 5) peptide sequence. The same or different nucleic acid encoding linker sequence can be provided between nucleic acid sequences encoding more than two Msp monomers. Optionally, a linker sequence can be provided between all or some of the nucleic acid sequences encoding Msp monomers in the single chain Msps provided herein.

Further provided is a nucleic acid sequence encoding a mutant single-chain Msp, wherein the nucleic acid sequence comprises (a) a first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence or any subset thereof, wherein the first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequences encode a first, second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, respectively; and (b) a ninth nucleotide sequence encoding an amino acid linker sequence, wherein the first Msp monomer sequence is a mutant Msp monomer sequence that comprises a mutation at position P97.

The mutant Msp monomer sequence can comprise a mutation at P97, wherein the mutation is not a P97S mutation or a P97C mutation. The mutant Msp monomer sequence can comprise a P97F mutation. As set forth above, any mutant Msp monomer sequence described herein can further comprise a mutation at amino acid position D118, a mutation at position D134 or a mutation at position E139. For example, any mutant Msp monomer sequence described herein can comprise a D118R mutation, a D134R mutation and/or a E139K mutation. Any mutant Msp monomer sequence described herein can further comprise (i) a mutation at position 93 and/or (ii) a mutation at position D90, position D91 or both positions D90 and D91. Optionally, the amino acid at position 90, 91 or 93 is substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine or tryptophan. Any mutant Msp monomer sequence described herein can further comprise a D90N, a D91N and a D93N mutation. For example, provided herein is a nucleic acid sequence encoding a mutant single-chain Msp, wherein the nucleic acid sequence comprises (a) a first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence or any subset thereof, wherein the first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequences encode a first, second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, respectively; and (b) a ninth nucleotide sequence encoding an amino acid linker sequence, wherein the first Msp monomer sequence is a mutant Msp monomer sequence that comprises a mutation at position P97 can further comprise (i) a mutation at amino acid position D118, D134 and/or E139 (ii) a mutation at position D93, and/or (iii) a mutation at position D90, position D91 or both positions D90 and D91. For example, the first Msp monomer sequence can be a mutant Msp monomer sequence that comprises a D90N mutation, a D91N mutation, a D93N mutation, a P97F mutation, a D118R mutation, a D134R mutation and a E139K mutation.

Further provided is a nucleic acid sequence encoding a mutant single-chain Msp, wherein the nucleic acid sequence comprises (a) a first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence or any subset thereof, wherein the first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequences encode a first, second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, respectively, and (b) a ninth nucleotide sequence encoding an amino acid linker sequence, wherein the first Msp monomer sequence is a mutant Msp monomer sequence that comprises one or more mutations at any of the following amino acid positions: I68, S73, S116, P123 or V128.

The first Msp monomer sequence can also be a mutant Msp monomer sequence that comprises one or more mutations at any of the following amino acid positions: I68, S73,

S116, P123 or V128 and further comprises a mutation at amino acid position D118, optionally with (i) a mutation at position 93, and/or (ii) a mutation at position D90, position D91 or both positions D90 and D91. In any of the mutant Msp monomer sequences described herein, the amino acid at position 91 or the amino acid at position 90 can be substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine or tryptophan. The mutant Msp monomer sequence can further comprise a D90N, a D91N and a D93N mutation. The mutant Msp monomer comprising one or more mutations at amino acids I68, S73, S116, P123 or V128 can further comprise a mutation in one or more of the amino acids at positions D13, A55, D56, E57, F58, E63, S136, D134, G137, E139 or D172. In the mutant Msp monomer sequences described herein, D13, A55, D56, E57, F58, E63, S136, D134, G137, E139 or D172 can be substituted with lysine or arginine. Therefore, a mutant Msp monomer comprising one or more mutations at amino acids I68, S73, S116, P123 or V128, for example, can further comprise (i) a mutation at amino acid position D118, (ii) a mutation at position D93, (iii) a mutation at position D90, position D91 or both positions D90 and D91, (iv) a D90N, a D91N and a D93N mutation and/or (v) a mutation in one or more of the amino acids at positions D13, A55, D56, E57, F58, E63, S136, D134, G137, E139 or D172.

For example, and not to be limiting, a first mutant Msp monomer can be a mutant Msp monomer comprising a mutation at positions D56, I68, S73, D118, D134 and E139. Optionally, the mutant Msp monomer can further comprise a D90N, a D91N and D93N mutation. Optionally, one or more of the amino acids selected from the group consisting of D56, I68, S73, D118, D134 and E139 can be substituted with lysine or arginine.

Further provided is a nucleic acid sequence encoding a mutant single-chain Msp, wherein the nucleic acid sequence comprises: (a) a first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence or any subset thereof, wherein the first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequences encode a first, second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, respectively; and (b) a ninth nucleotide sequence encoding an amino acid linker sequence, wherein the first Msp monomer sequence is a mutant Msp monomer sequence that comprises one or more mutations at any of the following amino acid positions: I68, S73, S116, P123 or V128; and wherein one or more of the first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence encodes a mutant Msp monomer sequence comprising a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. For example, and not to be limiting, the seventh nucleotide can encode a mutant Msp monomer

sequence comprising a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96.

It is understood that the first nucleotide sequence and the seventh nucleotide sequence can be arranged, but are not necessarily arranged as the first nucleotide sequence and the seventh nucleotide sequence in the nucleic acid sequence that comprises a first, second, third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence in that order. In this context, the first nucleotide sequence is a nucleotide sequence encoding the first or starting monomer of a single-chain Msp, and can be the first, second, third, fourth, fifth, sixth, seventh, or eighth nucleotide sequence of the single-chain Msp. The starting nucleotide sequence is referred to as the first nucleotide sequence no matter where it occurs in the single-chain Msp. For example, if the starting subunit of the single-chain Msp is the first Msp monomer (first nucleotide sequence), then the seventh Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the second Msp monomer (first nucleotide sequence), then the eighth Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the third Msp monomer (first nucleotide sequence), then the first Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the fourth Msp monomer (first nucleotide sequence), then the second Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the fifth Msp monomer (first nucleotide sequence), then the third Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the sixth Msp monomer (first nucleotide sequence), then the fourth Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the seventh Msp monomer (first nucleotide sequence), then the fifth Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, if the starting subunit of the single-chain Msp is the eighth Msp

monomer (first nucleotide sequence), then the sixth Msp monomer (seventh nucleotide sequence) comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96.

For example, and not to be limiting, the first Msp monomer can be a mutant Msp monomer that comprises a mutation at positions D56, I68, S73, D118, D134 and E139 and the seventh monomer can be a mutant Msp monomer that comprises a mutation at positions L88 and I105. Optionally, each of the amino acid positions at positions D56, I68, S73, D118, D134 and E139 of the first mutant Msp monomer can be substituted with lysine or arginine. Optionally, each of the amino acid positions at positions D56, I68, S73, D118, D134 and E139 of the first mutant Msp monomer can be substituted with phenylalanine, tryptophan, histidine or tyrosine. Optionally, each of the amino acid positions at positions L88 and I105 of the seventh mutant Msp monomer can be substituted with lysine or arginine. Optionally, each of the amino acid positions at positions L88 and I105 of the seventh mutant Msp monomer can be substituted with phenylalanine, tryptophan, histidine or tyrosine. Substitution of D56, I68, S73, D118, D134, E139, L88 and/or I105 with aromatic amino acids, such as, phenylalanine, tryptophan, histidine or tyrosine can promote π -stacking interactions with an analyte, for example, nucleotides, to decrease translocation velocity. Optionally, the first, second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, or a subset thereof can comprise a D90N, a D91N and a D93N mutation.

Figures 10A and 10B show a non-limiting example of a positive ramp created in a single-chain Msp comprising a first mutant Msp monomer that comprises a mutation at positions D56, I68, S73, D118, D134 and E139 and a seventh mutant Msp monomer that comprises a mutation at positions L88 and I105. This positively charged ramp inside the vestibule of the MspA guides single-stranded nucleic acids, for example DNA, through the Msp. The electrostatic interactions between the nucleic acid and the ramp enable controlled translocation of DNA through the pore. This reduces Brownian motion of the nucleic acid and the translocation rate. This also increases the precision and the interaction between the nucleic acid bases and the amino acids in the constriction zone. Figure 10C is a schematic of a single-chain Msp. Numbers under subunits #1 and #7 represent locations of the positive ramp.

In any of the mutant single-chain Msps set forth herein, the constriction zone can be modified to increase the nucleobase, protein or analyte recognition properties of MspA. Modifications to the constriction zone can create a reading head that increases, for example, base-specific interactions. A reading head can be created by introducing an amino acid with

a longer side chain that protrudes into the path of DNA or another analyte. For example, and not to be limiting, in order to create one or more reading heads, the amino acid at position 90 and/or 91 in any of the mutant Msp monomers of the single-chain Msps described herein can be substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine, tryptophan or an unnatural amino acid. Positioning heads can also be created to increase the efficiency of one or more reading heads. For example, amino acids with longer side chains, preferably hydrophobic or negatively charged, can be introduced, opposite to the reading head, in order to reduce escape motions of DNA or another analyte in the constriction zone. Amino acids that are suitable, include but are not limited to, aspartate, glutamate, valine, leucine, isoleucine, phenylalanine, tyrosine, tryptophan and unnatural amino acids. In order to further slow down translocation rates, a stacking slide can be created by mutating one or more of the amino acids at positions 83, 86, 88 and 105. For example, and not to be limiting, one or more of the amino acids at positions 83, 86, 88 and 105 can be substituted with tryptophan, tyrosine or phenylalanine. Optionally, the stacking slide is positioned such that it is located in proximity to a positive ramp.

One or more of the second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence or any subset thereof, can be independently selected from the group consisting of a wildtype MspA monomer, a mutant MspA monomer, a wild-type MspA paralog or homolog monomer, and a mutant MspA paralog or homolog monomer. It is understood that, when the second, third, fourth, fifth, sixth, seventh and/or eighth Msp monomer sequence is a mutant MspA monomer sequence, the mutant MspA monomer sequence can be the same or different than the first mutant MspA monomer sequence. Optionally, the second, third, fourth, fifth, sixth, seventh, and eighth Msp monomer sequence, or any subset thereof, is a wild-type MspA paralog or homolog monomer. These include, but are not limited to, MspA/Msmeg0965, MspB/Msmeg0520, MspC/Msmeg5483, MspD/Msmeg6057, MppA, PorM1, PorM2, PorM1, Mmcs4296, Mmcs4297, Mmcs3857, Mmcs4382, Mmcs4383, Mjls3843, Mjls3857, Mjls3931 Mjls4674, Mjls4675, Mjls4677, Map3123c, Mav3943, Mvan1836, Mvan4117, Mvan4839, Mvan4840, Mvan5016, Mvan5017, Mvan5768, MUL_2391, Mflv1734, Mflv1735, Mflv2295, Mflv1891, MCH4691c, MCH4689c, MCH4690c, MAB1080, MAB1081, MAB2800, RHA1 ro08561, RHA1 ro04074, and RHA1 ro03127. A wild-type MspA paralog or homolog monomer may be a wild-type MspB monomer.

Mutant Msp Monomers and the Nucleic Acids Encoding Them

Further provided is a nucleic acid encoding a mutant Msp monomer, wherein the Msp monomer comprises a mutation at position 97. Optionally, the mutant Msp monomer can comprise a mutation at P97, wherein the mutation is not a P97S mutation or a P97C mutation. Optionally, the mutant Msp monomer can comprise a P97F mutation. Optionally the mutant Msp monomer can further comprise a mutation at amino acid position D118, a mutation at position D134 or a mutation at position E139. For example, the mutant Msp monomer comprising a mutation at position 97 can further comprise a D118R mutation, a D134R mutation and/or a E139K mutation. Optionally, the mutant Msp monomer comprising a mutation at position 97 can further comprise (i) a mutation at position 93 and/or (ii) a mutation at position D90, position D91 or both positions D90 and D91. Optionally, the amino acid at position 90 or 91 is substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine or tryptophan. Optionally, the mutant Msp monomer can further comprise a D90N, a D91N and a D93N mutation. For example, and not to be limiting, a mutant MspA monomer sequence can comprise a D90N mutation, a D91N mutation, a D93N mutation, a P97F mutation, a D118R mutation, a D134R mutation and a E139K mutation.

Further provided is a nucleic acid encoding a mutant Msp monomer, wherein the Msp monomer comprises a mutation at one or more of the following positions: I68, S73, S116, P123 or V128. Optionally, the mutant monomer further comprises a mutation at amino acid position D118. Optionally, the Msp monomer further comprises a mutation at position D90, position D91 or both positions D90 and D91. Optionally, the amino acid at position 91 or the amino acid at position 90 can be substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine or tryptophan. Optionally, the mutant Msp monomer sequence can further comprise a D90N, a D91N and a D93N mutation. Optionally, the mutant Msp monomer sequence can further comprise a mutation in one or more of the amino acids at positions D13, A55, D56, E57, F58, E63, S136, D134, G137, E139 or D172. Optionally, one or more of D13, A55, D56, E57, F58, E63, S136, D134, G137, E139 or D172 can be substituted with lysine or arginine.

Further provided is a nucleic acid encoding a mutant Msp monomer, wherein the Msp monomer comprises a mutation at one or more of the following positions: T83, N86, G92 or A96. Optionally, the mutant Msp monomer sequence further comprises a mutation at position L88 or I105. Optionally, the mutant monomer further comprises a mutation at amino

acid positions D118. Optionally, the Msp monomer further comprises a mutation at position D90, position D91 or both positions D90 and D91. Optionally, the amino acid at position 91 or the amino acid at position 90 can be substituted with arginine, lysine, histidine, glutamine, methionine, threonine, phenylalanine, tyrosine or tryptophan. Optionally, the mutant Msp monomer sequence can further comprise a D90N, a D91N and a D93N mutation.

As used herein, a mutant Msp monomer refers to an Msp monomer that has at least or at most 70, 75, 80, 85, 90, 95, 98, or 99 percent or more identity, or any range derivable therein, but less than 100% identity, as compared to a wild-type Msp monomer, and retains tunnel-forming capability when associated with one or more other Msp monomers (wild-type or mutant). Therefore, in addition to the mutations described herein, any mutant Msp provided herein can further comprise additional modifications such as substitutions, insertions, deletions, and/or additions, as long as the mutant Msp monomer has at least or at most 70, 75, 80, 85, 90, 95, 98, or 99 percent or more identity, or any range derivable therein, but less than 100%, to a wild-type Msp monomer, and retains tunnel-forming capability when associated with one or more other Msp monomers.

Any mutant Msp described herein can comprise 2-15 Msp monomers that are the same or different, wherein at least one of the Msp monomers is a mutant Msp monomer. Optionally, a mutant Msp comprises 7-9 Msp monomers that are the same or different. Optionally, at least a second monomer is selected from the group consisting of a wildtype Msp monomer, a second mutant Msp monomer, a wild-type Msp paralog or homolog monomer, and a mutant Msp paralog or homolog monomer, wherein the second mutant Msp monomer may be the same or different than the first mutant Msp monomer. For example, any mutant Msp described herein can comprise 2-15 Msp monomers wherein at least one of the Msp monomers is a mutant MspA monomer. Optionally, at least a second monomer is selected from the group consisting of a wildtype MspA monomer, a second mutant MspA monomer, a wild-type MspA paralog or homolog monomer, and a mutant MspA paralog or homolog monomer, wherein the second mutant MspA monomer can be the same or different than the first mutant MspA monomer. Optionally, the second monomer is a wild-type MspA paralog or homolog monomer.

For example, a mutant Msp can comprise one or more Msp monomers comprising a mutation at position 97. In another example, a mutant Msp can comprise one or more Msp monomers comprising a mutation at one or more of I68, S73, S116, P123 or V128 and one or more Msp monomers comprising a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96. In another example, a mutant Msp can comprise one

or more Msp monomers with mutations at positions D56, I68, S73, D118, D134 and E139 and one or more Msp monomers with mutations at positions L88 and I105.

Modifications in amino acid sequence may arise as allelic variations (e.g., due to genetic polymorphism), may arise due to environmental influence (e.g., due to exposure to ultraviolet radiation), or other human intervention (e.g., by mutagenesis of cloned DNA sequences), such as induced point, deletion, insertion, and substitution mutants. These modifications can result in changes in the amino acid sequence, provide silent mutations, modify a restriction site, or provide other specific mutations. Amino acid sequence modifications typically fall into one or more of three classes: substitutional, insertional, or deletional modifications. Insertions include amino and/or terminal fusions as well as intrasequence insertions of single or multiple amino acid residues. Insertions ordinarily will be smaller insertions than those of amino or carboxyl terminal fusions, for example, on the order of one to four residues. Deletions are characterized by the removal of one or more amino acid residues from the protein sequence. Typically, no more than about from 2 to about 6 residues are deleted at any one site within the protein molecule. Amino acid substitutions are typically of single residues, but can occur at a number of different locations at once; insertions usually will be on the order of about from 1 to about 10 amino acid residues; and deletions will range from about 1 to about 30 residues. Deletions or insertions preferably are made in adjacent pairs, i.e., a deletion of 2 residues or insertion of 2 residues. Substitutions, deletions, insertions or any combination thereof can be combined to arrive at a final construct. The mutations may or may not place the sequence out of reading frame and may or may not create complementary regions that could produce secondary mRNA structure. Substitutional modifications are those in which at least one residue has been removed and a different residue inserted in its place.

Modifications, including the specific amino acid substitutions disclosed herein, are made by known methods. By way of example, modifications are made by site specific mutagenesis of nucleotides in the DNA encoding the protein, thereby producing a DNA encoding the modification, and thereafter expressing the DNA in recombinant cell culture to produce the Msp monomers or single chain multimers. Techniques for making substitution mutations at predetermined sites in DNA having a known sequence are well known, for example M13 primer mutagenesis and PCR mutagenesis.

The amino acids in the Msp proteins described herein can be any of the 20 naturally occurring amino acids, D-stereoisomers of the naturally occurring amino acids, unnatural amino acids and chemically modified amino acids. Unnatural amino acids (that is, those that

are not naturally found in proteins) are also known in the art, as set forth in, for example, Williams et al., *Mol. Cell. Biol.* 9:2574 (1989); Evans et al., *J. Amer. Chem. Soc.* 112:4011-4030 (1990); Pu et al., *J. Amer. Chem. Soc.* 56:1280-1283 (1991); Williams et al., *J. Amer. Chem. Soc.* 113:9276-9286 (1991); and all references cited therein. B and γ amino acids are known in the art and are also contemplated herein as unnatural amino acids.

As used herein, a chemically modified amino acid refers to an amino acid whose side chain has been chemically modified. For example, a side chain can be modified to comprise a signaling moiety, such as a fluorophore or a radiolabel. A side chain can also be modified to comprise a new functional group, such as a thiol, carboxylic acid, or amino group. Post-translationally modified amino acids are also included in the definition of chemically modified amino acids.

Also contemplated are conservative amino acid substitutions. By way of example, conservative amino acid substitutions can be made in one or more of the amino acid residues of any Msp monomer provided herein. One of skill in the art would know that a conservative substitution is the replacement of one amino acid residue with another that is biologically and/or chemically similar. The following eight groups each contain amino acids that are conservative substitutions for one another:

- 1) Alanine (A), Glycine (G);
- 2) Aspartic acid (D), Glutamic acid (E);
- 3) Asparagine (N), Glutamine (Q);
- 4) Arginine (R), Lysine (K);
- 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);
- 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);
- 7) Serine (S), Threonine (T); and
- 8) Cysteine (C), Methionine (M)

Nonconservative substitutions, for example, substituting a proline with glycine are also contemplated.

Those of skill in the art readily understand how to determine the identity of two polypeptides or nucleic acids. For example, the identity can be calculated after aligning the two sequences so that the identity is at its highest level. Another way of calculating identity can be performed by published algorithms. Optimal alignment of sequences for comparison can be conducted using the algorithm of Smith and Waterman, *Adv. Appl. Math.* 2: 482 (1981); by the alignment algorithm of Needleman and Wunsch, *J. Mol. Biol.* 48: 443 (1970); by the search for similarity method of Pearson and Lipman, *Proc. Natl. Acad. Sci. U.S.A.* 85:

2444 (1988); by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, WI; the BLAST algorithm of Tatusova and Madden FEMS Microbiol. Lett. 174: 247-250 (1999) available from the National Center for Biotechnology Information; or by inspection.

The same types of identity can be obtained for nucleic acids by, for example, the algorithms disclosed in Zuker, *Science* 244:48-52, 1989; Jaeger et al. *Proc. Natl. Acad. Sci. USA* 86:7706-7710, 1989; Jaeger et al. *Methods Enzymol.* 183:281-306, 1989 including at least material related to nucleic acid alignment. It is understood that any of the methods typically can be used and that, in certain instances, the results of these various methods may differ, but the skilled artisan understands if identity is found with at least one of these methods, the sequences would be said to have the stated identity.

For example, as used herein, a sequence recited as having a particular percent identity to another sequence refers to sequences that have the recited identity as calculated by any one or more of the calculation methods described above. For example, a first sequence has 80 percent identity, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent identity to the second sequence using the Zuker calculation method even if the first sequence does not have 80 percent identity to the second sequence as calculated by any of the other calculation methods. As yet another example, a first sequence has 80 percent identity, as defined herein, to a second sequence if the first sequence is calculated to have 80 percent identity to the second sequence using each of calculation methods (although, in practice, the different calculation methods will often result in different calculated identity percentages).

Further, any Msp or Msp monomer can also be chemically or biologically modified. For example, one can modify an Msp or Msp monomer with chemicals to produce disulfide bridges, as is known by those of skill in the art.

An Msp can comprise a nucleotide binding site. As used herein, a nucleotide binding site refers to a site in an Msp where a nucleotide stays in contact with, or resides at, an amino acid for a period of time that is longer than attributable to diffusion movement, such as greater than one picosecond or one nanosecond. Molecular dynamics calculations can be employed to assess these temporary resting times.

Polypeptides encoded by nucleic acids described herein are also provided. Therefore polypeptides comprising a mutant Msp monomer or functional fragment thereof, are provided. Non-limiting examples of mutant Msp monomers include but are not limited to, polypeptides comprising SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 and SEQ ID NO: 4 comprising any of the mutations described herein. Further provided is a Msp monomer comprising an amino acid sequence that has least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 99% identity or any percentage in between to a polypeptide comprising SEQ ID NO: 1, SEQ ID NO: 2, SEQ ID NO: 3 and SEQ ID NO: 4, wherein the polypeptide comprises any of the mutations described herein. Also provided are polypeptides comprising a mutant single-chain Msp or functional fragment thereof. Also provided are polypeptides comprising a mutant single-chain Msp comprising any of the mutant Msp monomers described herein, or a fragment thereof.

Non-limiting examples of mutant Msp monomers comprising mutations set forth herein are provided in Table 2. Each exemplary mutant Msp monomer comprises all of the mutations listed for each monomer. For example, D90N/D91N/D93N/P97F indicates that all four mutations are present. Single chain Msps comprising any of the exemplary mutant Msp monomer sequences provided in Table 2 are also provided herein. It is understood that amino acids listed in parentheses are listed as alternatives for substitutions at that position. For example, P97 (A/V/L/I/F/M/Y/W or G) means that P97 can be substituted with A, V, L, I, F, M, Y W or G.

Table 2

D90N/D91N/D93N/P97(A/V/L/I/F/M/Y/W or G)
D90N/ D91N/D93N/P97F/D118R/D134R/E139K
D90N/D91N/D93N/P97F
D90N/D91N/D93N/G92(A/V/L/I/P/F/M/Y or W)
D90N/D91N/D93N/I94(A/V/L/P/F/M/Y/W or G)
D90N/D91N/D93N/T95(A/V/L/P/F/M/Y/W or G)
D90N/D91N/D93N/A96(V/L/P/F/I/M/Y/W or G)
D90N/D91N/D93N/P98(A/V/L/F/I/M/Y/W or G)
D90N/D91N/D93N/F99(A/V/L/P/I/M/Y/W or G)
D90N/D91N/D93N/G100(A/V/L/P/F/I/M/Y or W)
D90N/ D91N/D93N/L101(A/V/I/F/M/Y/W or G)
D90N/ D91N/D93N/N102((A/V/L/I/P/F/M/Y/W or G)
D90N/ D91N/D93N/S103((A/V/L/I/P/F/M/Y/W or G)
D90N/G92F/ D91N /D93N/P97F
D90N/T95F/ D91N /D93N/P97F
D90N/A96F/ D91N /D93N/P97F
D90N/A96F/ D91N /D93N/P97F/P98F
D90N/G92F/ D91N /D93N/P97F/D118R/D134R/E139K
I68(R/K/F/W/Y or H)/D90N/ D91N /D93N
I68(R/K/F/W/Y or H)/D90N/ D91N /D93N/D118R/D134R/E139K
I68(K/R/F/W/Y or H)/D90N/D91N/D93N
I68(K/R/F/W/Y or H)/D90N/D91N/D93N/D118R/D134R/E139K
S73(K/R/F/W/Y or H)/D90N/D91N/D93N
S73(K/R/F/W/Y or H)/D90N/D91N/D93N/D118R/D134R/E139K
S73(K/R/F/W/Y or H)/D90N/D91N/D93N
S73(K/R/F/W/Y or H)/D90N/D91N/D93N/D118R/D134R/E139K

D90N/D91N/D93N/S116R
D90N/D91N/D93N/S116R/D118R/D134R/E139K
D90N/D91N/D93N/S116K
D90N/D91N/D93N/S116K/D118R/D134R/E139K
D90N/D91N/D93N/P123R
D90N/D91N/D93N/P123R/D118R/D134R/E139K
D90N/D91N/D93N/P123K
D90N/D91N/D93N/P123K/D118R/D134R/E139K
D90N/D91N/D93N/L88(K/R/F/W/H or Y)
D90N/D91N/D93N/I105(K/R/F/W/H or Y)
D90N/D91N/D93N/L88(K/R/F/W/H or Y)/D118R/D134R/E139K
D90N/D91N/D93N/I105(K/R/F/W/H or Y)/D118R/D134R/E139K
D90N/D91N/D93N/L88(K/R/F/W/H or Y)/I105(K/R/F/W/H or Y)
D90N/D91N/D93N/L88(K/R/F/W/H or Y)/I105(K/R/F/W/H or Y)/D118R/D134R/E139K
D90N/D91N/D93N/L88(K/R/F/W/H or Y)
D90N/D91N/D93N/T83(K/R/F/W/H or Y)/D118R/D134R/E139K
D90N/D91N/D93N/T83(K/R/F/W/H or Y)
D90N/D91N/D93N/N86(K/R/F/W/H or Y)
D90N/D91N/D93N/N86(K/R/F/W/H or Y)/D118R/D134R/E139K

Tunnel-forming Proteins

Methods of determining whether a protein is a tunnel-forming protein are well known in the art. One can determine if an Msp forms a tunnel by determining whether the protein inserts into a bilayer, such as described in Example 2 of U.S. Patent Publication No. 20120055792. All of the methods of making and using porins described in U.S. Patent Publication No. 20120055792 can be employed to make and use the Msp porins described herein. If the protein inserts into the bilayer, then the porin is a tunnel-forming protein. Typically, tunnel formation is detected by observing a discrete change in conductivity. See, U.S. Patent Publication No. 20120055792, and Niederweis et al., *Mol. Microbiol.* 33:933 (1999).

Bilayers are described herein. An Msp will typically be able to be inserted in a lipid bilayer or other thin film, which are each well-known in the art. An example of inserting a mutant MspA into a lipid bilayer is provided in U.S. Patent Publication No. 20120055792; this technique can be applied to other Msp proteins as well. In addition, U.S. Patent No. 6,746,594 describes a variety of lipid bilayers and thin films, including inorganic materials, that can be employed with respect to the Msps discussed herein. Methods, apparatuses, and techniques described in U.S. Patent No. 6,267,872 are also employable with respect to Msps discussed herein. Moreover, more than one Msp can be comprised in a lipid bilayer. For example, 2, 3, 4, 5, 10, 20, 200, 2000, or more can be comprised in a lipid bilayer. Optionally, anywhere from 2 to 10¹⁰ Msps can be employed in methods described herein. Such a plurality of Msps can be in the form of clusters of Msps.

Clusters can be randomly assembled or can adopt a pattern. As used herein, a cluster refers to molecules that are grouped together and move as a unit, but are not covalently bound to one another.

Optionally, Msps do not gate spontaneously. As used herein, to gate or gating refers to the spontaneous change of electrical conductance through the tunnel of the protein that is usually temporary (e.g., lasting for as few as 1-10 milliseconds to up to a second). Long lasting gating events can often be reversed by changing the polarity. Under most circumstances, the probability of gating increases with the application of higher voltages. Gating and the degree of conductance through the tunnel change are highly variable among Msps, depending on, for example, the make-up of the vestibule and constriction zone as well as the properties of the liquid medium in which the protein is submerged. Typically, the protein becomes less conductive during gating, and conductance can permanently stop (i.e., the tunnel may permanently shut) as a result, such that the process is irreversible. Optionally, gating refers to the conductance through the tunnel of a protein spontaneously changing to less than 75% of its open state current.

Various conditions such as light and liquid medium, including its pH, buffer composition, detergent composition, and temperature, can affect the behavior of an Msp, particularly with respect to its conductance through the tunnel as well as the movement of an analyte with respect to the tunnel, either temporarily or permanently.

As used throughout, a tunnel refers to the central, empty portion of an Msp that is defined by the vestibule and the constriction zone, through which a gas, liquid, ion, or analyte can pass. As used herein, "cis" refers to the side of an Msp tunnel through which an analyte enters the tunnel or across the face of which the analyte moves. As used herein, "trans" refers to the side of an Msp tunnel through which an analyte (or fragments thereof) exits the tunnel or across the face of which the analyte does not move.

Any mutant Msp described herein, for example a mutant MspA, can comprise a vestibule and a constriction zone that define a tunnel. Further, the diameter of a mutant Msp or mutant Msp paralog or homolog can be less than the diameter of the constriction zone of a corresponding wild-type Msp or wild-type Msp paralog or homolog. A mutant Msp or mutant Msp paralog or homolog can have a mutation in the vestibule or the constriction zone that permits an analyte to translocate, electrophoretically or otherwise, through the tunnel of the mutant Msp or mutant Msp paralog or homolog with a translocation velocity or an average translocation velocity that is less than the translocation velocity or average translocation velocity at which the analyte translocates through the tunnel of a wild-type Msp or wild-type Msp paralog or homolog. Also, any mutant Msp described herein can comprise a vestibule having a length from about 2 to about 6 nm and a diameter from about 2 to about 6 nm, and a constriction zone 5 having a length from about 0.3 to

about 3 nm and a diameter from about 0.3 to about 3 nm, wherein the vestibule and constriction zone together define a tunnel. It is understood that, one or more mutations, can be made in the vestibule or the constriction zone of any of the Msp described herein in order to increase or decrease conductance through the tunnel of an Msp. For example, any of the mutant Msps described herein can further comprise a deletion, substitution or insertion of an amino acid in the vestibule and/or the constriction zone in order to modify conductance.

As used throughout, a vestibule refers to the cone-shaped portion of the interior of an Msp whose diameter generally decreases from one end to the other along a central axis, where the narrowest portion of the vestibule is connected to the constriction zone. A vestibule can also be referred to as a goblet. The vestibule and the constriction zone together define the tunnel of an Msp. When referring to a diameter of the vestibule, it is understood that because the vestibule is cone-like in shape, the diameter changes along the path of a central axis, where the diameter is larger at one end than the opposite end. The diameter can range from about 2 nm to about 6 nm. Optionally, the diameter is about, at least about, or at most about 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, or 6.0 nm, or any range derivable therein. The length of the central axis can range from about 2 nm to about 6 nm. Optionally, the length is about, at least about, or at most about 2, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4, 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, or 6.0 nm, or any range derivable therein. When referring to diameter herein, one can determine a diameter by measuring center-to-center distances or atomic surface-to-surface distances.

As used throughout, a constriction zone refers to the narrowest portion of the tunnel of an Msp, in terms of diameter, that is connected to the vestibule. The length of the constriction zone can range from about 0.3 nm to about 2 nm. Optionally, the length is about, at most about, or at least about 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, or 3 nm, or any range derivable therein. The diameter of the constriction zone can range from about 0.3 nm to about 2 nm. Optionally, the diameter is about, at most about, or at least about 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2, or 3 nm, or any range derivable therein.

Any Msp discussed herein can be comprised in a lipid bilayer. Optionally, an analyte electrophoretically or otherwise translocates from the cis side through a tunnel to the trans side. Optionally, an analyte electrophoretically or otherwise translocates from the trans side through a tunnel to the cis side. Optionally, an analyte is electrophoretically or otherwise driven from the cis side or the trans side into a tunnel and stays in the tunnel or then retracts

to the cis side or the trans side, respectively. It is understood that analytes can translocate through the tunnel in the presence or absence of an electric field.

Single-chain Msps function at a wide range of electrolyte concentration, for example from about 0.3-1M KCl (see Figure 22). To optimize channel activity, lipid association can be performed prior to insertion of Msp in a membrane or lipid bilayer. In a non-limiting example, Figure 22 shows that no channel activity was observed in a buffer containing only 0.3 M KCl at pH 8.0. However, breaking the membrane and subsequent repainting of the membrane leads to increased channel activity of scMspA PN1 in the electrolyte containing 0.3 M KCl at pH 8.0. Therefore, in any of the methods set forth herein, an Msp can be contacted or preincubated with one or more lipids to optimize channel activity.

Vectors and Cells

A vector comprising a nucleic acid encoding a polypeptide described herein is also provided. The vector can further comprise a promoter sequence, for example, a constitutive promoter or an inducible promoter. Examples of constitutive promoter include, but are not limited to, the p_{smc} promoter and Phsp60. Examples of inducible promoters include, but are not limited to, an acetamide-inducible promoter and a tetracycline inducible promoter.

Cultured cells transfected with any vector described herein, or progeny thereof, wherein the cell is capable of expressing a Msp (either as a single-chain Msp, an Msp comprising Msp monomers or an Msp monomer, are also provided). A *Mycobacterium smegmatis* strain comprising any vector described herein is also provided. A *Mycobacterium smegmatis* strain free of endogenous porins is also contemplated and can further comprise any vector described herein. By "free" is meant that an endogenous porin cannot be detected in an immunoblot when using an appropriate Msp-specific antiserum, or comprising less than 1% endogenous porins.

Any of the Msp monomers or single-chain Msps disclosed herein can be produced by transforming a mutant bacterial strain comprising a deletion of a wild-type MspA, a wild-type MspB, a wild-type MspC, a wildtype MspD, with a vector comprising an inducible promoter operably linked to a nucleic acid sequence encoding the Msp monomer or single-chain Msp porin; and purifying the Msp monomer or single-chain Msp porin (See, for example, U.S. Patent No. 6,746,594). Optionally, the mutant bacterial strain comprises a deletion of a *recA* gene. Optionally, the vector comprises any of the nucleic acids encoding an Msp monomer or single-chain Msp described herein. The bacterial strain can further comprise *M. smegmatis* strain ML16, ML714 or ML712.

Systems and Methods of Use

Also provided is a system comprising a mutant Msp described herein having a vestibule and a constriction zone that define a tunnel, wherein the tunnel is positioned between a first liquid medium and a second liquid medium, wherein at least one liquid medium comprises an analyte, and wherein the system is operative to detect a property of the analyte. A system can be operative to detect a property of any analyte comprising subjecting an Msp to an electric field such that the analyte interacts with the Msp. A system can be operative to detect a property of the analyte comprising subjecting the Msp to an electric field such that the analyte electrophoretically translocates through the tunnel of the Msp. Also provided is a system comprising an Msp having a vestibule and a constriction zone that define a tunnel, wherein the tunnel is positioned in a lipid bilayer between a first liquid medium and a second liquid medium, and wherein the only point of liquid communication between the first and second liquid media occurs in the tunnel. Moreover, any system described herein can comprise any mutant Msp described herein.

The first and second liquid media can be the same or different, and either one or both can comprise one or more salts, detergents, or buffers. In fact, any liquid media described herein can comprise one or more of a salt, a detergent, or a buffer. Optionally, at least one liquid medium is conductive. Optionally, at least one liquid medium is not conductive. Any liquid medium described herein can comprise a viscosity-altering substance or a velocity-altering substance. The liquid medium can comprise any analyte described herein.

A property of an analyte can be an electrical, chemical, or physical property. An Msp can be comprised in a lipid bilayer in a system or any other embodiment described herein. A system can comprise a plurality of Msps. A system can comprise any Msp described herein, such as a single-chain mutant MspA or a mutant Msp comprising at least 2-15 monomers, wherein at least one of the monomers is a mutant MspA monomer. A mutant Msp comprised in a system can comprise a vestibule having a length from about 2 to about 6 nm and a diameter from about 2 to about 6 nm, and a constriction zone having a length from about 0.3 to about 3 nm and a diameter from about 0.3 to about 3 nm, wherein the vestibule and constriction zone together define a tunnel.

Any Msp described herein, including an Msp comprised in a system, can further comprise a molecular motor. The molecular motor in a system is capable of moving an analyte into or through a tunnel with a translocation velocity or an average translocation velocity that is less than the translocation velocity or average translocation velocity at which the analyte translocates into or through the tunnel in the absence of the molecular motor. The molecular motor can be, for example, an enzyme, such as a polymerase, an exonuclease, or a helicase such as DnaB or a phage nucleic acid packing motors (see, for example, Serwer *Viruses* 3(7): 1249-80 (2011)). Any

system described herein can further comprise a patch-clamp amplifier or a data acquisition device. A system can further comprise one or more temperature regulating devices in communication with the first liquid medium, the second liquid medium, or both. Any system described herein can be operative to translocate an analyte through an Msp tunnel either electrophoretically or otherwise.

The mutant MspA can have a mutation in the vestibule or the constriction zone that permits an analyte to translocate, e.g., electrophoretically, through the tunnel with an average translocation velocity of less than 0.5 nm/ μ s or less than 0.05 nm/ μ s. The analyte can be selected from the group consisting of a nucleotide(s), a nucleic acid, amino acid(s), a peptide, a protein, a polymer, a drug, an ion, a biological warfare agent, a pollutant, a nanoscopic object, or a combination or cluster thereof. Optionally, the analyte is further defined as a nucleic acid. The nucleic acid can translocate, electrophoretically or otherwise, through the tunnel with an average translocation velocity of less than 1 nucleotide/ μ s, or less than 0.1 nucleotide/ μ s. A nucleic acid can be further defined as ssDNA, dsDNA, RNA, or a combination thereof.

As used herein, electrophoretically translocating an analyte, refers to applying an electric field to an Msp porin that is in contact with one or more solutions (e.g., immersed in a solution), such that current flows through the Msp tunnel. The electric field moves an analyte such that it interacts with the tunnel. As used herein, "interacts" means that the analyte moves into and, optionally, through the tunnel, where "through the Msp tunnel" (or "translocates") means to enter one side of the tunnel and move to and out of the other side of the tunnel. It is specifically contemplated that any analyte discussed herein can translocate through an Msp tunnel, either electrophoretically or otherwise, in any embodiment discussed herein. In this regard, it is specifically contemplated that any embodiment herein comprising translocation can refer to electrophoretic translocation or nonelectrophoretic translocation, unless specifically noted. Optionally, methods that do not employ electrophoretic translocation are contemplated.

As used throughout, a liquid medium includes aqueous, organic-aqueous, and organic-only liquid media. Organic media include, e.g., methanol, ethanol, dimethylsulfoxide, and mixtures thereof. Liquids employable in methods described herein are well-known in the art. Descriptions and examples of such media, including conductive liquid media, are provided in U.S. Patent No. 7,189,503. Salts, detergents, or buffers may be added to such media. Such agents can be employed to alter pH or ionic strength of the liquid medium. Viscosity-altering substances, such as glycerol or various polymers (e.g., polyvinylpyrrolidone, polyethylene glycol, polyvinyl alcohol, cellulose polymers), and mixtures thereof, can be included in liquid media. Methods of measuring viscosity are well-known in the art.

Any agent that can be added to a liquid medium can also alter the velocity of an analyte that is being studied. As such, a velocity-altering agent may be a salt, a detergent, a buffer, a viscosity-altering substance, or any other agent added to a liquid medium that increases or decreases the velocity of an analyte. Typically, an analyte employed herein is soluble or partially soluble in at least one liquid medium that is in contact with an Msp described herein.

As used herein, nucleic acid refers to a deoxyribonucleotide or ribonucleotide polymer in either single- or double-stranded form, and unless otherwise limited, encompasses known analogs of natural nucleotides that hybridize to nucleic acids in a manner similar to naturally occurring nucleotides, such as peptide nucleic acids (PNAs) and phosphorothioate DNA. Unless otherwise indicated, a particular nucleic acid sequence includes the complementary sequence thereof. Nucleotides include, but are not limited to, ATP, dATP, CTP, dCTP, GTP, dGTP, UTP, TTP, dUTP, 5-methyl-CTP, 5-methyl dCTP, ITP, diTP, 2-amino-adenosine-TP, 2-amino-deoxyadenosine-TP, 2-thiothymidine triphosphate, pyrrolo-pyrimidine triphosphate, and 2-thiocytidine, as well as the alphathiotriphosphates for all of the above, and 2'-O-methyl-ribonucleotide triphosphates for all the above bases. Modified bases include, but are not limited to, 5-Br-UTP, 5-BrdUTP, 5-F-UTP, 5-F-dUTP, 5-propynyl dCTP, and 5-propynyl-dUTP.

As used herein, a drug refers to any substance that may alter a biological process of a subject. Drugs can be designed or used for or in the diagnosis, treatment, or prevention of a disease, disorder, syndrome, or other health affliction of a subject. Drugs can be recreational in nature, that is, used simply to alter a biological process and not used for or in the diagnosis, treatment, or prevention of a disease, disorder, syndrome, or other health affliction of a subject. Biologics, which refer to substances produced by biological mechanisms involving recombinant DNA technology, are also encompassed by the term drug. Drugs include, for example, antibacterials, anti-inflammatories, anticoagulants, antivirals, antihypertensives, antidepressants, antimicrobials, analgesics, anesthetics, beta-blockers, bisphosphonates, chemotherapeutics, contrast agents, fertility medications, hallucinogens, hormones, narcotics, opiates, sedatives, statins, steroids, and vasodilators. Non-limiting examples of drugs can also be found in the *Merck Index: an Encyclopedia of Chemicals, Drugs, and Biologicals*, 15th ed. New Jersey: Merck, 2013. Antibacterial drugs used in the treatment of tuberculosis, for example, include isoniazid, rifampicin, pyrazinamide, and ethambutol. Methods employing a drug as an analyte can further comprise drug screening. For example, uptake of a drug into a cell or an organism can be investigated using an Msp by observing ion current blockades. Specific Msp porin constriction zones and/or vestibules with various sizes, electrostatic properties, and chemical properties can be constructed to closely emulate the desired pathway for drugs to enter or exit a cell or organism.

These methods could greatly accelerate screening for drugs as well as drug design (see, for example, Pagel et al., *J. Bacteriology* 189:8593 (2007)).

As used herein, a biological warfare agent refers to any organism or any naturally occurring, bioengineered, or synthesized component of any such microorganism capable of causing death or disease in plants or animals (including humans) or degradation of food or water supplies, or degradation of the environment. Non-limiting examples include Ebola viruses, Marburg virus, *Bacillus anthracis* and *Clostridium botulinum*, *Variola major*, *Variola minor*, anthrax, and ricin.

As used herein, a pollutant refers to a material that pollutes air, water, or soil. Non-limiting examples of pollutants include fertilizers, pesticides, insecticides, detergents, petroleum hydrocarbons, smoke, and heavy metal-containing substances, such as those containing zinc, copper, or mercury (e.g., methylmercury).

Any analyte can be used herein, including, for example, a nucleotide(s), a nucleic acid, an amino acid(s), a peptide, a protein, a polymer, a drug, an ion, a biological warfare agent, a pollutant, a nanoscopic object, or any other molecule comprising one of these analytes or a combination of thereof. An analyte can be a cluster of molecules (e.g. 2-10 nucleotides or amino acids), in that the cluster as a whole is considered an analyte. Typically, an analyte's size will not be so great such that it cannot enter a tunnel of an Msp. In other words, a typical analyte will be smaller in size than the opening of a tunnel of an Msp. However, an analyte having a size larger than the opening of a tunnel can be employed, and it can be determined that the analyte's size is too large to enter the tunnel. Optionally, the molecular weight of the analyte is less than one million Da. Optionally, the molecular weight of the analyte is about, at most about, or at least about 1,000,000, 950,000, 900,000, 850,000, 800,000, 750,000, 700,000, 650,000, 600,000, 550,000, 500,000, 450,000, 400,000, 350,000, 300,000, 250,000, 200,000, 150,000, 100,000, 75,000, 50,000, 25,000, 20,000, 15,000, 10,000, 7,500, 5,000, 2,500, 2,000, 1,500, 1,000, or 500 Da or less, or any range derivable therein.

An analyte can also be a nanoscopic object, which is an object that is smaller than 100 nm in two of its dimensions. As used herein, an analyte can further comprise a magnetic bead. A magnetic bead can be further defined as a streptavidin-coated magnetic bead. An analyte can further comprise an optical bead. Any analyte described herein can be an ion or can be neutral. An analyte can comprise biotin.

Beads that can be employed include magnetic beads and optical beads. For example, one can use streptavidin-coated magnetic beads to apply an opposing force to the electrostatic forces that pull DNA through the tunnel of an Msp. In this latter technique a magnetic bead is attached to biotinylated DNA, and a force comparable to the electrostatic driving force (-10 pN) would be

applied using a strong magnetic field gradient. See Gosse and Croquette, *Biophys. J.* 82:3314 (2002). In this way, the blockade-current readout would be unaffected, but the forces on the DNA could be independently controlled. Tens or hundreds of complete, independent reads of each DNA could then be correlated and assembled to reconstruct an accurate DNA sequence.

Optical beads manipulated by "optical tweezers" are also known in the art, and such methods can be applied to the Msp's described herein. Optical tweezers are a common tool used to exert a force on a nanoscopic object. An analyte is attached on one end of the bead, while the other end can be inserted into the tunnel of the porin. The position and force of the bead is controlled and measured with the optical tweezers. Such methods control the passage of the analyte into the tunnel and allow for more control of the reading of the analyte, such as the reading of the units of a polymer. See, e.g., Trepagnier et al., *Nano Lett.* 7:2824 (2007) for a description of such methods in the context of artificial nanopores. U.S. Patent No. 5,795,782 also discusses the use of optical tweezers.

Fluorescence resonance energy transfer (FRET), a well-known technique, can be employed in analytical methods described herein. For example, a fluorescent FRET acceptor or FRET-donor molecule can be incorporated into an Msp. The analyte is then labeled with a matching FRET -donor or FRET -acceptor. When the matching FRET donor is within the Forster distance to the FRET acceptor, energy transfer will likely occur. The resulting signal could be used for analytical purposes instead of or in addition to methods using ion current as described herein. Accordingly, methods of detection, identification, or sequencing can comprise FRET technology. Other optical methods that can be employed include introducing optically active molecules into the interior of an Msp (such as the vestibule or the constriction zone). External light would be applied to affect the interior of the protein. Such methods could be used to affect the translocation velocity of an analyte or could allow the analyte's entry or exit from the tunnel, offering controlled passage of the analyte. Alternatively, optical pulses focused onto the pore could be used to heat the pore to affect how it interacts with the analyte. Such control could be very fast as the heat from a small volume of a focal point would dissipate rapidly. Methods of controlling the translocation velocity of an analyte can therefore employ such optically active molecules or optical pulses. Manipulation of translocation velocity can also be accomplished by attaching an object to one end of an analyte, and the other end of the analyte then interacts with the Msp. The object can be a bead (e.g., a polystyrene bead), a cell, a large molecule such as streptavidin, neutravidin, DNA, etc., or a nanoscopic object. The object could then be subjected to a fluid flow or could be subject to passive viscous drag.

Molecular motors are well-known in the art and refer to a molecule (e.g., an enzyme) that physically interacts with an analyte, such as a polymer (e.g., a 15 polynucleotide), and is capable

of physically moving the analyte with respect to a fixed location, such as the vestibule, constriction zone, or tunnel of an Msp. Although not intending to be bound by theory, molecular motors utilize chemical energy to generate mechanical force. A molecular motor can interact with each unit (or "mer") of a polymer in a sequential manner. Non-limiting examples of molecular motors include DNA polymerases, RNA polymerases, helicases, ribosomes, and exonucleases. Nonenzymatic motors are also known, such as virus motors that pack DNA. See Smith et al., Nature 413:748 (2001). A variety of molecular motors and desirable properties of such motors are described in U.S. Patent No. 7,238,485.

A molecular motor can be disposed on the cis side or the trans side of an Msp porin and can optionally be immobilized, such as described by the '485 patent. Methods of incorporating a molecular motor into an Msp can be performed using methods described in the '485 patent. Systems and apparatuses described in the '485 patent can be employed with respect to an Msp described herein as well. Indeed, any embodiment discussed in the '485 patent can be employed using an Msp, as described herein. Molecular motors are also discussed in, e.g., Cockroft et al., J. Amer. Chem. Soc. 130:818 (2008); Benner et al., Nature Nanotech. 2:718 (2007); and Gyarfás et al., ACS Nano 3:1457 (2009).

A molecular motor is typically employed to regulate the rate or translocation velocity at which an analyte interacts with an Msp. Any Msp described herein can comprise a molecular motor. Optionally, a molecular motor is employed to decrease the rate at which an analyte enters an Msp porin tunnel or to decrease the translocation velocity at which an analyte translocates through an Msp tunnel. Optionally, the translocation velocity or average translocation velocity is less than 0.5 nm/ μ s. Optionally, the translocation velocity or average translocation velocity is less than 0.05 nm/ μ s. Optionally, the translocation velocity or average translocation velocity is less than 1 nucleotide/ μ s. Optionally, the translocation velocity or average translocation velocity is less than 0.1 nucleotide/ μ s.

Optionally, the rate of movement of an analyte ranges from greater than 0 Hz to 2000Hz. Here, rate refers to the number of subunits (or "mers") of a regular polymer advancing in one second (Hz). Optionally, the range is between about 50-1500Hz, 100-1500 Hz, or 350-1500 Hz. Optionally, the rate of movement is about, at most about, or at least about 25, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, 1450, 1500, 1550, 1600, 1650, 1700, 1750, 1800, 1850, 1900, 1950, or 2000Hz, or any range derivable therein. The rate can be controlled by the use of a molecular motor that moves an analyte at a substantially constant rate, at least for a portion of time during a characterization. In addition, the range of rate of movement can depend on the molecular motor. For example, for an RNA polymerase, a range can be 350-1500 Hz; for a DNA

polymerase, a range can be 75-1500 Hz; and for ribosomes, helicases, and exonucleases, a range can be 50-1500 Hz.

Recording and detection techniques can be employed in the methods described herein. In addition, U.S. Patent Nos. 5,795,782 and 7,189,503 also describes recording methods and instrumentation that can be employed with respect to Msps, as well as methods for optimizing conductance readings. U.S. Patent No. 6,746,594 describes a support for thin films containing nanopores and methods for using such supports that can be employed with respect to the Msps described herein.

Method of Making a Single Chain Msp

The Msp pore, for example, MspA, is currently the best available channel protein for nanopore sequencing of nucleic acids. However, its composition of eight subunits makes it impossible to introduce asymmetric changes in the pore that could optimize the properties of the Msp for nucleic acid sequencing. In order to overcome this difficulty, methods of making a single chain Msp are provided herein. These methods can be used to produce a full or partial single-chain Msp. Generally, the method comprises transforming a mutant bacterial strain. The mutant strain comprises a deletion of a wild-type MspA, a wild-type MspB, a wild-type MspC, a wild-type MspD, and optionally a deletion of the *recA* gene. The mutant strain is transformed with a vector comprising a nucleic acid sequence encoding a single-chain Msp porin. The single-chain Msp porin is then purified from the bacteria. Optionally, the single-chain Msp porin comprises a single-chain MspA porin. Optionally, the vector comprises any of the nucleic acids described herein.

As described in the Examples, in order to combine the superior sequencing capabilities of MspA with an increased ability to adapt vestibule and constriction properties to DNA sequencing, a single-chain MspA porin octamer was constructed that allows for the optimal properties of the vestibule and the constriction zone for DNA sequencing. The MspA chain termini are close together in the MspA porin and are connected by a short peptide linker. The (GGGS)₃ (SEQ ID NO:3) peptide, for example, is used to connect the carboxy-terminus of the preceding MspA monomer (or multimer) to the amino-terminus of the following MspA monomer (or multimer), which lacks signal peptide. To create a vector comprising the MspA porin sequence, each MspA monomer sequence is flanked by a unique restriction site, which allows the capability to mutate any individual monomer. To create an

MspA porin sequence, each MspA sequence can be assembled stepwise to form a dimeric, tetrameric, and octameric single-chain MspA utilizing the unique restriction sites. To avoid problems of recombination in creating the single-chain MspA multimer, seven MspA genes are synthesized with different codon usages i.e., the genes encode the exact same amino acid sequence, however, the DNA sequence has been altered from the native MspA nucleotide sequence (SEQ ID NO: 10). To create the MspA porin sequence, the nucleotide sequence encoding the first Msp monomer can optionally contain a nucleic acid sequence encoding a leader sequence (e.g., amino acids 1 to 27 of SEQ ID NO: 6). Each of the seven Msp monomer sequences following the first Msp monomer sequence can comprise SEQ ID NO: 1 or SEQ ID NO: 1 with one or more mutations described herein. The vector comprising the MspA porin sequence is transformed into the quadruple porin mutant bacterial strain, as described in the Examples. Optionally, single chain Msps can be purified and subjected to a refolding procedure. For example, anion exchange chromatography in the presence of 8M urea can be used to obtain a pure fraction of a single chain Msp which is dialyzed against a buffer to remove urea. After dialysis, a refolding buffer comprising a refolding agent, for example, L-arginine and detergent, are added to the sample and purified, refolded single chain Msp is obtained. Refolding agents are known to those of skill in the art. These include, but are not limited to, arginine, arginine hydrochloride, arginineamide, glycineamide, proline, glycerol, and cyclodextrins (see, for example, Yamaguchi et al. *Biomolecules* 4: 235-251 (2014); and expression levels and oligomeric status of the MspA porin can be checked by Western blot or other immunohistochemical techniques known to those of skill in the art. The tunnel activity of the MspA porin can be determined by lipid bilayer experiments, as described in the Examples and as known to those of skill in the art.

Single chain M1₈-MspA pores insert much more frequently into lipid bilayers than a similar amount of octameric M1-MspA. Insertion of octameric MspA is a tedious procedure. Thus, single-chain Msps, such as those described herein facilitate setup of systems and methods of using Msp for detecting and identifying analytes, for example, for nucleic acid sequencing.

Method of Increasing Msp Insertion in a Lipid Bilayer

Provided herein is a method of increasing the number of Msp insertions in a lipid bilayer, comprising contacting any Msp described herein with a lipid to form a lipid-associated Msp and inserting the lipid-associated Msp of step into a lipid bilayer. Optionally,

the contacting step comprises inserting the Msp in a lipid bilayer and disrupting the lipid bilayer to form a lipid-associated Msp. For example, an Msp can be inserted in a lipid bilayer that is subsequently disrupted. The disrupted lipid bilayer comprises Msp(s). Therefore, the Msp(s) are lipid associated. The lipid-associated Msp can then be contacted with other lipids to form another lipid bilayer that comprises the lipid-associated Msps. As used herein, a lipid bilayer is a thin membrane comprising lipid molecules, for example, phospholipids, that can be used to insert any Msp provided herein. Therefore, in the methods provided herein, the Msp can be contacted with phospholipids, either as part of a lipid bilayer or not, in order to form lipid-associated Msp.

As set forth above, one of skill in the art can determine if an Msp inserts into a bilayer, by using techniques such as those described in Example 2 of U.S. Patent Publication No. 20120055792. All of the methods of making and using porins described in U.S. Patent Publication No. 20120055792 can be employed to make and use the Msp porins described herein. If the protein inserts into the bilayer, then the porin is a tunnel-forming protein. Typically, tunnel formation is detected by observing a discrete change in conductivity. See, U.S. Patent Publication No. 20120055792, and Niederweis et al., *Mol. Microbiol.* 33:933 (1999). The increase in Msp insertions can be an increase of about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, 100%, 200%, 300%, 400% or greater.

Detection Methods

Further provided is a method for detecting the presence of an analyte, comprising: (a) applying an electric field sufficient to translocate an analyte from a first conductive medium to a second conductive medium in liquid communication through any mutant Msp described herein; and (b) measuring an ion current, wherein a reduction in the ion current indicates the presence of the analyte in the first medium. Optionally, the first and second liquid conductive media are the same. Optionally, the first and second liquid conductive media are different. The mutant Msp porin can be any mutant Msp porin discussed herein. For example, the Msp porin can be a mutant single-chain Msp, a mutant Msp comprising 2-15 monomers or combinations thereof. As set forth above, a plurality of Msps can also be used in the methods described herein.

In the methods disclosed herein, an Msp can further comprise a molecular motor. The molecular motor can be capable of moving an analyte into or through a tunnel with a translocation velocity or an average translocation velocity that is less than the translocation velocity or average translocation velocity at which the analyte electrophoretically translocates into or through the tunnel in the absence of the molecular motor. Accordingly, in any

embodiment herein comprising application of an electric field, the electric field can be sufficient to cause the analyte to electrophoretically translocate through the tunnel. Any liquid medium discussed herein, such as a conductive liquid medium, can comprise an analyte. In the methods comprising measuring an ion current, the analyte interacts with an Msp porin tunnel to provide a current pattern, wherein the appearance of a blockade in the current pattern indicates the presence of the analyte.

The methods disclosed herein can further comprise identifying the analyte. For example, such methods can comprise comparing the current pattern obtained with respect to an unknown analyte to that of a known current pattern obtained using a known analyte under the same conditions. In another example, and not to be limiting, identifying the analyte can comprise (a) measuring the ion current to provide a current pattern, wherein a reduction in the current defines a blockade in the current pattern, and (b) comparing one or more blockades in the current pattern to (i) one or more blockades in the current pattern, or (ii) one or more blockades in a known current pattern obtained using a known analyte.

The analyte can be any analyte described herein. For example, the analyte can be a nucleotide(s), a nucleic acid, an amino acid(s), a peptide, a protein, a polymer, a drug, an ion, a pollutant, a nanoscopic object, or a biological warfare agent. In the methods provided herein, optionally, at least one of the first or second conductive liquid media comprises a plurality of different analytes.

In methods where the analyte is a polymer, for example, a protein, a peptide or a nucleic acid, the method can further comprise identifying one or more units of the polymer. For example, identifying one or more units of the polymer can comprise measuring the ion current to provide a current pattern comprising a blockade for each polymer unit, and comparing one or more blockades in the current pattern to (i) one or more other blockades in the current pattern or (ii) one or more blockades in a current pattern obtained using a polymer having known units. These methods can comprise identifying sequential units of the polymer, for example, and not to be limiting, sequential or consecutive nucleotides in a nucleic acid. In another example, sequential or consecutive amino acids in a polypeptide can be identified using the methods described herein.

The methods provided herein can comprise distinguishing at least a first unit within a polymer from at least a second unit within the polymer. Distinguishing can comprise measuring the ion current produced as the first and second units separately translocate through a tunnel to produce a first and a second current pattern, respectively, where the first and second current patterns differ from each other.

The methods provided herein can further comprise sequencing a polymer. Sequencing can comprise measuring the ion current or optical signals as each unit of the polymer is separately translocated through the tunnel to provide a current pattern that is associated with each unit, and comparing each current pattern to the current pattern of a known unit obtained under the same conditions, such that the polymer is sequenced.

Further provided is a method of sequencing nucleic acids or polypeptides using any of the mutant Msps provided herein. The method comprises creating a lipid bilayer comprising a first and second side, adding a purified Msp to the first side of the lipid bilayer, applying positive voltage to the second side of the lipid bilayer, translocating an experimental nucleic acid or polypeptide sequence through the Msp porin, comparing the experimental blockade current with a blockade current standard, and determining the experimental sequence.

Any of the detection methods provided herein can further comprise determining the concentration, size, molecular weight, shape, or orientation of the analyte, or any combination thereof.

As used herein, a polymer refers to a molecule that comprises two or more linear units (also known as a "mers"), where each unit may be the same or different. Non-limiting examples of polymers include nucleic acids, peptides, and proteins, as well as a variety of hydrocarbon polymers (e.g., polyethylene, polystyrene) and functionalized hydrocarbon polymers, wherein the backbone of the polymer comprises a carbon chain (e.g., polyvinyl chloride, polymethacrylates). Polymers include copolymers, block copolymers, and branched polymers such as star polymers and dendrimers.

Methods of sequencing polymers using Msp are described herein. In addition, sequencing methods can be performed in methods analogous to those described in U.S. Patent No. 7,189,503. See also U.S. Patent No. 6,015,714. More than one read can be performed in such sequencing methods to improve accuracy. Methods of analyzing characteristics of polymers (e.g., size, length, concentration, identity) and identifying discrete units (or "mers") of polymers are discussed in the '503 patent as well, and can be employed with respect to the present Msps. Indeed, an Msp can be employed with respect to any method discussed in the '503 patent.

At present, several types of observable signals can be used as readout mechanisms in nanopore sequencing and analyte detection. An exemplary readout method relies on an ionic blockade current or copassing current, uniquely determined by the identity of a nucleotide or other analyte occupying the narrowest constriction in the pore. This method is referred to as blockade current nanopore sequencing or BCNS. Blockade current detection and characterization

of nucleic acids has been demonstrated in both the protein pore ahemolysin (aHL) and solid-state nanopores.

Blockade current detection and characterization has been shown to provide a host of information about the structure of DNA passing through, or held in, a nanopore in various contexts. In general, a blockade is evidenced by a change in ion current that is clearly distinguishable from noise fluctuations and is usually associated with the presence of an analyte molecule at the pore's central opening. The strength of the blockade will depend on the type of analyte that is present. More particularly, a blockade refers to an interval where the ionic current drops below a threshold of about 5-100% of the unblocked current level, remains there for at least 1.0 μ s, and returns spontaneously to the unblocked level. For example, the ionic current may drop below a threshold of about, at least about, or at most about 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, or 100%, or any range derivable therein. Blockades are rejected if the unblocked signal directly preceding or following it has an average current that deviates from the typical unblocked level by more than twice the rms noise of the unblocked signal. Deep blockades are identified as intervals where the ionic current drops <50% of the unblocked level. Intervals where the current remains between 80% and 50% of the unblocked level are identified as partial blockades.

Disclosed are materials, compositions, and components that can be used for, can be used in conjunction with, can be used in preparation for, or are products of the disclosed methods and compositions. These and other materials are disclosed herein, and it is understood that when combinations, subsets, interactions, groups, etc. of these materials are disclosed that while specific reference of each various individual and collective combinations and permutations may not be explicitly disclosed, each is specifically contemplated and described herein. For example, if a method is disclosed and discussed and a number of modifications that can be made to a number of compositions included in the method are discussed, each and every combination and permutation of the method, and the modifications that are possible are specifically contemplated unless specifically indicated to the contrary. Likewise, any subset or combination of these is also specifically contemplated and disclosed. This concept applies to all aspects of this disclosure including, but not limited to, steps in methods. Thus, if there are a variety of additional steps that can be performed, it is understood that each of these additional steps can be performed with any specific method steps or combination of method steps of the disclosed methods, and that each such combination or subset of combinations is specifically contemplated and should be considered disclosed. It is also contemplated that any embodiment discussed in this specification can be

implemented with respect to any method, compound, protein, porin, peptide, polypeptide, multimer, monomer, nucleic acid, vector, strain, cultured cell, system, or composition, etc., described herein, and *vice versa*.

A number of embodiments have been described. Nevertheless, it will be understood that various modifications can be made. Accordingly, other embodiments are within the scope of the following claims.

EXAMPLES

A single-chain MspA was constructed consisting of eight covalently connected monomers. As shown herein, an expression plasmid encoding single-chain M1-MspA is not stable in wild-type *M. smegmatis* but can be propagated in an *M. smegmatis* lacking the *recA* gene, which is required for homologous recombination. The *recA* gene was deleted in the quadruple porin mutant *M. smegmatis* ML712, which lacks the four known Msp porins. This strain enabled expression of single-chain M1-MspA. Tunnels made by single-chain M1-MspA had a similar conductance as octameric M1-MspA tunnels, but had drastically improved resistance to voltage gating. This unforeseen advantage of single-chain MspA is of great importance for nanopore sequencing of nucleic acids, for example, DNA.

Construction of an Msp quadruple deletion mutant of *M. smegmatis*

For isolation of mutant MspA porins a triple porin deletion mutant *Mycobacterium smegmatis* ML16 strain ($\Delta mspA::FRT$, $\Delta mspC::FRT$, $\Delta mspD::FRT$) was used (see Stephan et al., Mol. Microbiol. 58: 714-730 (2005)). However, low levels of MspB could still be detected in this strain in immunoblots with MspA-specific rabbit antiserum. The presence of MspB can contribute to the heterogeneity observed in single-channel experiments and complicates data analysis. In order to overcome this problem and improve MspA preparations an *M. smegmatis* strain lacking all four *msp* genes was constructed.

Since the deletion of all four *msp* genes of *M. smegmatis* is lethal, the first step was to integrate an expression cassette for the N-terminal channel-forming domain of CpnT (1) into the *M. smegmatis* porin triple mutant ML16. To this end, the plasmid pML2622 was constructed, which carries the N-terminal domain of CpnT tagged with His6 and HA under the control of a nitrile-inducible promoter (Fig. 2). The N-terminal domain of CpnT formed channels in planar bilayer experiments and complemented the porin mutant *M. smegmatis*

ML16 strain in glycerol uptake experiments. The rescue plasmid pML2622 was integrated into the mycobacteriophage L5-site in the chromosome of the porin triple mutant *M. smegmatis* ML16. Integration was confirmed by PCR with the sets of primers complementary to L5 genomic region. This strain was named *M. smegmatis* ML709. After integration of pML2622, the plasmid backbone was excised from the chromosome by Flp recombinase as described in Stephan et al. (*Gene* 343: 181-190 (2004)) to remove the genes encoding hygromycin phosphotransferase and L5 integrase. This strain was named *M. smegmatis* ML709-234.

To delete the remaining *mshB* gene in *M. smegmatis* ML711 the *mshB* deletion vector pML1611 containing the two reporter genes *gfp* and *xylE* as markers for integration and allelic replacement was constructed (Fig. 2). The *mshB* deletion vector pML1611 carries 863 bp and 946 bp-long upstream and downstream regions of *mshB* and was used to delete the *mshB* gene in *M. smegmatis* ML709-234 by allelic exchange. This Msp porin quadruple mutant was named *M. smegmatis* ML711. The plasmid pCreSacB containing Cre recombinase was used to excise *gfp* and *hyg* genes flanked by *loxP* sites from the chromosome. The deletion of all four *msh* genes was confirmed by PCR with chromosomal DNA using specific sets of primers and by Southern blot with chromosomal DNA using specific probe (Fig. 3). This unmarked Msp porin quadruple mutant was named *M. smegmatis* ML712 (relevant genotype: $\Delta mshA::FRT$, $\Delta mshB::loxP$, $\Delta mshC::FRT$, $\Delta mshD::FRT$, *attB* L5::FRT-pNIT-cpnT_{d1}-FRT). The growth of the strain ML712 on Middlebrook 7H10 agar plates was impaired in comparison to wt and ML16 strains. Next, the expression of *msh* genes in the ML712 strain was assessed by extraction of *M. smegmatis* cells using the detergent octylpolyethyleneoxide as described in Heinz et al. (*Anal. Biochem.* 285: 113-120 (2000)). The Msp quadruple porin deletion mutant *M. smegmatis* ML712 grown in Middlebrook 7H9 medium does not produce any Msp protein in contrast to the porin triple mutant ML16 (Fig. 4). This demonstrated that deletion of all *msh* genes was achieved in *M. smegmatis* ML712. The expression levels of the MspA were similar to that of wt *M. smegmatis*, when wt *mshA* or *M1 mshA* were expressed in ML712 using the plasmids pMN016 and pML904, respectively (Fig. 4).

Construction of single-chain M1-MspA

Previously a M1₂-MspA subunit dimer was constructed (Pavlenok et al. *PLoS One* 7(6): e38726). As the next step towards single-chain MspA, four *mshA-M1* genes were fused to encode a M1₄-MspA subunit tetramer. The resulting plasmid pML2647 was transformed

into the quadruple porin deletion strain of *M. smegmatis* ML712 for protein production and purification. However, the tetrameric *mspA* plasmid was unstable. In order to avoid recombination, the *recA* gene was deleted in the quadruple porin mutant ML712 and the strain *M. smegmatis* ML714 was created.

Then, a gene encoding single-chain MspA, in which eight M1-MspA subunits are linked (M1₈-MspA), was cloned into *E. coli*. Each of the subunits has a D90N mutation, a D91N mutation and a D93N mutation. This was achieved by fusing two genes encoding tetrameric M1 MspA together (M1₄-MspA) using pML2647 as a template. The individual MspA subunits are separated by (GGGS)₃ linkers. In the resulting plasmid pML3213, the two tetrameric *M1-mspA* constructs are flanked by unique restriction sites (tetrad A: PacI, MluI; tetrad B: EcoRV, HindIII) (Fig. 5). Genes within the tetramers are flanked by the same restriction sites with the exception of the first and last genes of the tetrads.

The resulting *mI₈-mspA* gene was placed under the control of the constitutive p_{smyc} promoter (Fig. 5). The plasmid pML3213 was transformed into *M. smegmatis* ML714 (quadruple porin mutant with *recA* deletion) for protein production and purification. Western blot experiments showed that the expression level of single-chain M1₈-MspA in *M. smegmatis* ML714 is lower compared to M1₂-MspA and is reduced to approximately 7% of wt MspA levels (Fig. 6).

Stability of single-chain M1-MspA

The MspA pore is very resistant against thermal and chemical denaturation (Heinz et al. *J. Biol. Chem.* 278: 8678-8685 (2003)). To test the thermal stability of single-chain MspA, the M1₈-MspA protein was subjected to increasing temperatures for 15 min in the presence of 2% SDS. (Fig. 7) A significant amount of M1₈-MspA is stable even after heating the protein sample to 100 °C (Fig. 7). This result shows that M1₈-MspA is at least as stable against thermal denaturation as the wt MspA protein.

Channel properties of single-chain M1-MspA

To examine whether M1₈-MspA forms functional channels, *in vitro* lipid bilayer experiments were performed. No insertions were recorded when only n-octyl-POE buffer was added to the lipid bilayer. Addition of approximately 70 ng of M1₈-MspA protein resulted in the step-wise increase in the current across the lipid bilayer indicating the insertion of M1₈-MspA channels into the membrane (Fig. 8A). Analysis of the current recordings of M1₈-MspA showed a major peak of 1.1 nS (Fig. 8B). This channel conductance is similar to those

of the pores made from the M1-MspA dimer (1.3 nS) and M1 MspA monomers (1.4 nS) (see Pavlenok et al.).

Voltage gating

Voltage gating is defined as a spontaneous channel closure at a certain voltage threshold and is an intrinsic property of bacterial β -barrel channel proteins (Bainbridge et al. *FEBS Lett* 431(3): 305-308 (1998)). Resistance to voltage gating is very important for nanopore sequencing experiments since voltages as high as +180mV are used to translocate ssDNA through MspA pore (Manrao et al. *Nat. Biotechnol.* 30(4): 349-353 (2012); Derrington et al. *Proc. Natl. Acad. Sci. USA* 107(37): 16060-16065 (2010); Butler et al. *Proc. Natl. Acad. Sci. USA* 105(52): 20647-20652 (2008)). Therefore, the voltage gating of M1₈-MspA in lipid bilayer experiments was analyzed. After insertion of approximately 220 M1₈-MspA pores, the voltage across the lipid bilayer was sequentially increased in 10 mV increments, and the ion current passing through the pores was measured for three minutes. The critical voltage V_c is defined as the voltage at which pores start to close, and is measured in these experiments as decrease of ion current.

The M1₈-MspA channels started to close at +90 mV and were completely stable at all applied negative voltages (Fig. 9). In a second experiment with gel-purified M1₈-MspA protein no voltage gating, up to voltages of ± 100 mV, was observed. Thus, the critical voltage V_c of M1₈-MspA is two-fold higher than that of M1-MspA or M1₂-MspA (V_{crit} +40mV, -50mV for both proteins). These results show that linking all eight subunits into a single polypeptide drastically increased the resistance of single-chain MspA to voltage gating. This unforeseen advantage of single-chain MspA is of great importance for nanopore sequencing of nucleic acids, for example, DNA.

Construction of a mutant single chain MspA (scMspA M2)

As described herein, mutations in MspA are useful for improving its interactions with DNA, its base recognition properties and its interactions with membranes and accessory proteins such, for example, Phi29 DNA polymerase. Using the approach described above for single-chain M1 MspA, a mutant single-chain MspA (MspA M2), in which eight mutant MspA monomers are linked together was constructed. Expression of both single-chain M1 MspA and single-chain MspA M2 constructs in *M. smegmatis* ML712 was shown by Western blots using an MspA antibody demonstrating that production of scMspA in *M. smegmatis* is feasible. As shown herein, single chain Msps can be expressed in *E. coli*. The single chain

M2 MspA (scMspA M2) protein is made in mg amounts, but is not folded. A folding protocol has been developed that allows isolation of active scMspA M2.

A single-chain *m2-mspA* (*scm2-mspA*) where eight *m2-mspA* genes (containing the mutations D90N/D91N/D93N/D118R/D134R/E139K as described in Butler et al. (*PNAS* 105: 20647-20652 (2008))) were connected by DNA fragments encoding (GGGGS)₃ polypeptide linkers. In addition, each gene was flanked by unique restriction sites to enable specific modifications of each MspA subunit. The genes in the sequence are named *m2-1* through *m2-8* beginning from the ATG start codon (Fig. 11 and Table 3). For protein production and purification of the single-chain MspA M2 protein in *E. coli* cells the signal peptide of MspA was removed. The *scm2-mspA* sequence was codon optimized for optimal expression in *E. coli* and was synthesized by GenScript. The resulting *scm2-mspA* gene was flanked by EcoRI and HindIII and was obtained in a pUC57 plasmid from GenScript. Next, the whole *sc m2-mspA* was excised and cloned into the pET-21(a)+ vector. The *scm2-mspA* gene is under the control of the T7 promoter in the resulting plasmid pML3216 (Fig. 11).

For scMspA M2 protein production and purification, the plasmid pML3216 was transformed into *E. coli* BL21(DE3)Omp8 strain which lacks 3 major porins (See Prilipov et al. *FEMS Microbiol. Lett* 163: 65-72 (1998)). The BL21(DE3) Omp8 strain was chosen to avoid contamination of scMspA M2 with endogenous porins of *E. coli*. After induction of *scm2-mspA* expression with 1.5 mM IPTG cells were grown at 37°C in LB medium supplemented with ampicillin. Maximal expression of the target protein was observed two hours after induction accounting for approximately 4% of the total protein in the cell lysate (Fig. 12). A protein band corresponding to scMspA M2 had an apparent mass of 170 kDa which is consistent with its predicted molecular mass of 165.6 kDa (Fig. 12). Next, scMspA M2 from inclusion bodies was isolated and purified as described in Sambrook et al. (*CSH Protocols* 2006) Inclusion bodies containing predominantly scMspA M2 protein were solubilized in 8 M urea. This sample was later a subject to anion exchange chromatography using HiTrap QFF column (GE HealthCare, United Kingdom) in the presence of 8 M urea. The elution profile of scMspA M2 protein was very similar to that of wt MspA published previously (Heinz *et al.*, 2003). This protein is probably not folded and has no channel activity.

Then, scMspA M2 was purified and subjected to a refolding procedure. After anion exchange chromatography a pure fraction of scMspA M2 with a concentration of 50 µg/ml was diluted by a factor of 10 in a buffer containing 10 mM NaCl, 25 mM HEPES, 0.6 M L-Arginine, 0.1% (v/v) LDAO, pH 8.0 to give final volume of 1 ml. The mixture was incubated

overnight at room temperature (approximately 21°C) on a rotating mixer. Then, the sample was transferred into a dialysis tube with 3.5 kDa MWCO and dialyzed against 2L of a buffer containing 10 mM NaCl, 25 mM HEPES (pH 8.0), 0.023% (v/v) LDAO overnight at room temperature. The dialyzed protein was transferred into a microtube and incubated at a room temperature for an additional day. Next, the refolding efficiency was assessed by Western blot analysis using MspA-specific rabbit antiserum. After the refolding procedure, the band which reacts with MspA polyclonal antibodies migrated from 170 kDa to approximately 130 kDa indicating that folding of scMspA M2 to a more compact form with an increased electrophoretic mobility had occurred (Fig. 13). Such an electrophoretic mobility shift upon folding has been observed for outer membrane proteins of *E. coli* previously. However, it was not clear whether MspA would show a similar phenomenon.

In order to examine if scMspA M2 forms functional channels *in vitro* after the refolding procedure lipid bilayer experiments were performed. No channel activity was observed when only 0.023% LDAO-buffer was added to the planar bilayer. In contrast, addition of scMspA M2 protein after the refolding step resulted in a step-wise current increase indicative of channel insertions into lipid bilayer (Fig. 14). Analysis of the current traces showed an average conductance of 2.3 nS (Fig. 14). Of interest, analysis of MspA M2 made from monomers showed two peaks at 1.2 nS and 2.4 nS suggesting two different protein conformations. In addition, a multi-channel experiment with scMspA M2 showed improved voltage-gating resistance with a critical voltage of +80mV/-70mV (Fig. 15). The increased voltage resistance is beneficial for example, for ssDNA experiments performed at relatively high voltages.

Table 3. Restriction sites of scMspA M2

# of RS	RS	Sequence	Ends	Amino acids	Gene flanked
1	EcoRI	GAATTC	cohesive	EF	m2-1
2	KpnI	GGTACC	cohesive	GT	
3	NsiI	ATGCAT	cohesive	MH	m2-2
4	ScaI	AGTACT	blunt	ST	
5	NheI	GCTAGC	cohesive	AS	m2-3
6	HpaI	GTTAAC	blunt	VN	
7	XbaI	TCTAGA	cohesive	SR	m2-4
8	NdeI	CATATG	cohesive	HM	
9	EcoRV	GATATC	blunt	DI	m2-5
10	PstI	CTGCAG	cohesive	LQ	
11	BstBI	TTCGAA	cohesive	FE	m2-6
12	BamHI	GGATCC	cohesive	GS	
13	MluI	ACGCGT	cohesive	TR	m2-7

14	PvuII	CAGCTG	blunt	QL	
15	Afl II	CTTAAG	cohesive	LK	
16	HindIII	AAGCTT	cohesive	KL	m2-8

Construction of mutant single chain MspA (MspA PN1)

A single-chain *mspA pn1* (*scmspA PN1*) gene where eight *mspA* genes (containing a P97F mutation and mutations D90N/D91N/D93N/D118R/D134R/E139K as described in Butler et al. (*PNAS* 105: 20647-20652 (2008))) were connected by DNA fragments encoding (GGGGS)₃ polypeptide linkers. In addition, each gene was flanked by unique restriction sites to enable specific modifications of each MspA subunit. The genes in the sequence are named *m2-97-1* through *m2-97-8* beginning from the ATG start codon (Fig. 16 and Table 3). For protein production and purification of the single-chain MspA PN1 protein in *E. coli* cells the signal peptide of MspA was removed. The *scmspA PN1* sequence was codon optimized for optimal expression in *E. coli* and was synthesized by GenScript. The resulting *scmspA PN1* gene was flanked by EcoRI and HindIII and was obtained in a pUC57 plasmid from GenScript. Next, the entire *scmspA PN1* was excised and cloned into the pET-21(a)+ vector. The *scmspA PN1* gene is under the control of the T7 promoter in the resulting plasmid pML3216 (Fig. 16).

For scMspA PN1 protein production and purification the plasmid pML3216 was transformed into *E. coli* BL21(DE3)Omp8 strain which lacks 3 major porins (See Prilipov et al. *FEMS Microbiol. Lett* 163: 65-72 (1998)). The BL21(DE3) Omp8 strain was chosen to avoid contamination of scMspA PN1 with endogenous porins of *E. coli*. After induction of *scmspA PN1* expression with 1.5 mM IPTG, cells were grown at 37°C in LB medium supplemented with ampicillin. Maximal expression of the target protein was observed two hours after induction accounting for approximately 5% of the total protein in the cell lysate (Fig. 17). A protein band corresponding to scMspA PN1 had an apparent mass of 170 kDa which is consistent with its predicted molecular mass of 165.6 kDa (Fig. 17). Next, scMspA PN1 from inclusion bodies was isolated and purified as described in Sambrook et al. (*CSH Protocols* 2006) Inclusion bodies containing predominantly scMspA PN1 protein were solubilized in 8 M urea. This sample was later a subject to anion exchange chromatography using HiTrap QFF column (GE HealthCare, United Kingdom) in the presence of 8 M urea. The elution profile of scMspA PN1 protein was very similar to that of wt MspA published previously (Heinz et al., 2003). This protein is probably not folded and has no channel activity.

Then, scMspA PN1 was purified and subjected to a refolding procedure. After anion exchange chromatography a pure fraction of scMspA PN1 was dialyzed against 2L of buffer containing 140 mM NaCl, 10mM K₂HPO₄/KH₂PO₄, 2mM KCl (pH 7.5) to remove urea. The mixture was incubated overnight at room temperature (approximately 21°C). After dialysis, L-arginine and LDAO were added to the sample to give a final concentration of 0.6M and 0.1% (v/v), respectively. The protein sample in the refolding buffer (140 mM NaCl, 10 mM K₂HPO₄/KH₂PO₄, 2mM KCl, 0.6 M L-Arginine, 0.1% (v/v) LDAO, pH 7.5) was incubated overnight on an orbital shaker (Fig. 18). The concentration of the purified sample was calculated to be 1.37 mg/ml as determined by absorbance at 280 nm. The protein yield was 0.45 mg per 1 liter of bacterial culture.

To test the effect of phenylalanine at position 97 on incorporation of the single-chain MspA into artificial lipid membranes the insertion activity of different MspA constructs was measured by monitoring the release of fluorescent carboxyfluorescein dye from the liposomes as described (See Schwarz et al., *Biophys. J.* 58(3):577-83 (1990); Schwarz et al., *Biochim. Biophys. Acta* 1239(1): 51-7 (1995)). Briefly, DPhPC liposomes were prepared by extrusion in the presence of 30 mM carboxyfluorescein. Carboxyfluorescein is self-quenched when it is enclosed into lipid vesicles. After insertion of MspA pore into the dye-loaded liposome, diffusion-mediated efflux of the dye results in the increase of fluorescence in the reaction mixture. Figure 19 shows the results of these carboxyfluorescein release experiments. Addition of buffers containing either LDAO (0.1% v/v) or OPOE (0.5% v/v) resulted in only minimal dye release from the liposomes, in contrast to Triton X-100 (1% v/v) buffer that was used as a positive control. Importantly, addition of scMspA PN1 (60 ng/ml, final) lead to faster and larger release of carboxyfluorscein than addition of scMspA M2(120 ng/ml, final). Interestingly, wt MspA (60 ng/ml, final) resulted in slower dye diffusion from the liposomes than scMspA PN1 (Fig. 19). These data indicate that additional phenylalanines located in the loop 6 of scMspA promote faster and more efficient insertion of the pores into lipid bilayers.

Next, the time of the first pore insertion into DPhPC membrane was measured in a bilayer set up. It was hypothesized that pores with enhanced insertion abilities would require less time to insert into lipid membrane. To examine the effect of phenylalanines in loop 6 on the time of membrane insertion of scMspA, scMspA PN1 was compared with scMspA M2. Briefly, the bilayer cuvette was filled with electrolyte, -10 mV potential was applied, and the data were acquired and recorded using TestPoint software. The same cuvette was always used in these experiments. The protein was added to both sides at a final concentration of 100

ng/ml. Importantly, successful insertion events were observed in 89% of the experiments for scMspA PN1, but only in 40% of the experiments for scMspA M2. This is consistent with the results of the carboxyfluorescein release experiments. Although the median insertion time for scMspA PN1 was 399 seconds as opposed to 695 seconds for scMspA M2, this difference was not significant. Surprisingly, the rate of insertion decreased when scMspA PN1 was analyzed in 0.3M KCl solution (median time: 859 seconds, 50%). However, half of the experiments resulted in successful insertions with scMspA PN1, while only one successful insertion was observed with scMspA M2 in 0.3M KCl with a time of 1270 seconds (8 membranes analyzed, 12% successful insertions) (Figure 20). This result shows the beneficial effect of phenylalanines in loop 6 for membrane insertion by single-chain MspA.

In order to examine whether scMspA PN1 forms functional channels *in vitro* after the refolding procedure, lipid bilayer experiments were performed. No channel activity was observed when only 0.1% LDAO-buffer was added to the planar bilayer. In contrast, addition of scMspA PN1 protein after the refolding step resulted in a step-wise current increase indicative of channel insertions into the lipid bilayer (Fig. 21). Analysis of the current traces showed an average conductance of 2.0 nS (Fig. 21). This could translate into larger residual currents for each nucleotide and better signals in DNA sequencing experiments.

Effect of lipids on channel activity of single-chain MspA PN1

scMspA PN1 was stored for more than a month at room temperature in 1 µg/mg and 0.2 µg/ml amounts. The scMspA PN1 was diluted in 0.1% LDAO, 140mM NaCl, 10mM K₂HPO₄/KH₂PO₄ (pH 7.5), 2mM KCL. Methods for making horizontal bilayers for channel experiments are known in the art. See, for example, Butler et al. (2008) and Akeson et al. Biophysical Journal; 77: 3227-3233 (1999). For the channel experiments, 2% diphtanoyl-phosphatidylcholine (DiphPC) in chloroform was used to form membrane bilayers for insertion of MspA essentially as described in Butler et al. and Akeson et al. After insertion of the MspA into the bilayers, the membrane was broken and the membrane was reapplied using 1% DiphPC in n-decane. The electrolyte used in these experiments was 0.3 or 1M KCl, 10mM Hepes, pH 8.0 or pH 7.0, respectively.

As shown in Figure 22, single-chain MspAs function at a wide range of electrolyte concentration, for example from about 0.3-1M KCl. To optimize channel activity, lipid association can be performed prior to insertion of MspA in a membrane or lipid bilayer.

Therefore, in any of the methods set forth herein, an MspA can be contacted or preincubated with one or more lipids to optimize channel activity. In a non-limiting example, Figure 22 shows that no channel activity was observed in a buffer containing only 0.3 M KCl at pH 8.0. However, breaking the membrane and subsequent repainting of the membrane leads to increased channel activity of scMspA PN1 in the electrolyte containing 0.3 M KCl at pH 8.0.

The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows:

1. A nucleic acid encoding a mutant single-chain *Mycobacterium smegmatis* porin (Msp), wherein the nucleic acid sequence comprises:

(a) a first and second nucleotide sequence, wherein the first nucleotide sequence encodes a first Msp monomer sequence and the second nucleotide sequence encodes a second Msp monomer sequence; and

(b) a third nucleotide sequence encoding an amino acid linker sequence, wherein at least one of the first and second Msp monomer sequences is a mutant Msp monomer sequence, wherein the mutant Msp monomer sequence has at least 95% sequence identity to the full-length of SEQ ID NO: 1 and wherein the mutant Msp monomer sequence comprises a P97F mutation.

2. The nucleic acid of claim 1, wherein the mutant Msp monomer sequence further comprises a mutation at one or more amino acid positions D118, D134 or E139.

3. The nucleic acid of claim 1 or 2, wherein the mutant Msp monomer sequence further comprises (i) a mutation at position 93, and/or (ii) a mutation at position 90, position 91 or both positions 90 and 91.

4. The nucleic acid of claim 3, wherein the mutant Msp monomer sequence comprises a D90N, a D91N and a D93N mutation.

5. The nucleic acid of claim 1, wherein the mutant Msp monomer sequence comprises a D90N mutation, a D91N mutation, a D93N mutation, the P97F mutation, a D118 mutation, a D134 mutation and a E139 mutation.

6. The nucleic acid of any one of claims 1-5, wherein the second Msp monomer sequence is a wild-type MspA monomer SEQ ID NO: 1 or a sequence having at least 95% sequence identity to the full-length of SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4.

7. A nucleic acid encoding a mutant single-chain *Mycobacterium smegmatis* porin (Msp), wherein the nucleic acid sequence comprises:

(a) a first and second nucleotide sequence wherein the first nucleotide sequence encodes a first Msp monomer and the second nucleotide sequence encodes a second Msp monomer;

(b) a third, fourth, fifth, sixth, seventh, and eighth nucleotide sequence or any subset thereof, wherein the third, fourth, fifth, sixth, seventh, and eighth nucleotide sequences encode a third, fourth, fifth, sixth, seventh, and eighth Msp monomer, respectively, wherein the first, second, third, fourth, fifth, sixth, seventh and eighth nucleotide sequence or any subset thereof are arranged consecutively in the nucleic acid; and

(c) at least one ninth nucleotide sequence encoding an amino acid linker sequence, wherein the ninth nucleotide sequence is positioned between any two consecutive nucleotide sequences encoding Msp monomers;

wherein at least one of the first and second Msp monomers is a mutant Msp monomer, wherein the mutant Msp monomer has at least 95% sequence identity to the full-length of SEQ ID NO: 1 and wherein the mutant Msp monomer comprises a P97F mutation.

8. The nucleic acid of claim 7, wherein the mutant Msp monomer sequence further comprises a mutation at one or more of amino acid position D118, D134 or E139.

9. The nucleic acid of claim 7 or 8, wherein the mutant Msp monomer further comprises (i) a mutation at position 93, and/or (ii) a mutation at position 90, position 91 or both positions 90 and 91.

10. The nucleic acid of claim 9, wherein the mutant Msp monomer sequence comprises a D90N, a D91N and a D93N mutation.
11. The nucleic acid of claim 7, wherein the mutant Msp monomer sequence comprises a D90N mutation, a D91N mutation, a D93N mutation, the P97F mutation, a D118 mutation, a D134 mutation and a E139 mutation.
12. The nucleic acid of any one of claims 7-11, wherein one or more of the second, third, fourth, fifth, sixth, seventh or eighth Msp monomer sequence is a wild-type MspA monomer SEQ ID NO: 1 or a sequence having at least 95% sequence identity to the full-length of SEQ ID NO: 2, SEQ ID NO: 3 or SEQ ID NO: 4.
13. A nucleic acid encoding a mutant *Mycobacterium smegmatis* porin (Msp) monomer, wherein the mutant Msp monomer sequence has at least 95% sequence identity to the full-length of SEQ ID NO: 1, and wherein the Msp monomer comprises a P97F mutation.
14. The nucleic acid of claim 13, wherein the mutant Msp monomer further comprises a mutation at amino acid position D118, D134 or E139.
15. The nucleic acid of claim 13 or 14, wherein the Msp monomer further comprises (i) a mutation at position 93, and/or (ii) a mutation at position 90, position 91 or both positions 90 and 91.
16. The nucleic acid of claim 15, wherein the mutant Msp monomer comprises a D90N, a D91N and a D93N mutation.
17. The nucleic acid of claim 13, wherein the mutant Msp monomer sequence comprises a D90N mutation, a D91N mutation, a D93N mutation, the P97F mutation, a D118 mutation, a D134 mutation and a E139 mutation.

18. A mutant single-chain Msp encoded by any one of the nucleic acids of any one of claims 1-12.
19. The mutant Msp monomer encoded by any one of the nucleic acids of any one of claims 13-17.
20. A vector comprising the nucleic acid of any one of claims 1-17.
21. A mutant *Mycobacterium smegmatis* porin (Msp) comprising at least 2-15 Msp monomers, wherein the mutant Msp comprises a first Msp monomer encoded by any of the nucleic acids set forth in any one of claims 13-17.
22. The mutant Msp of claim 21, further comprising at least a second mutant Msp monomer wherein the second mutant Msp monomer sequence has at least 95% sequence identity to the full-length of SEQ ID NO: 1, and wherein the second mutant Msp monomer comprises a mutation at one or more of the following positions: T83, N86, L88, I105, D90, D91, G92, D93 or A96.
23. A system comprising the mutant single-chain Msp encoded by any one of the nucleic acids of any one of claims 1-12, wherein the Msp has a vestibule and a constriction zone that define a tunnel, wherein the tunnel is positioned between a first conductive liquid medium and a second conductive liquid medium, wherein at least one conductive liquid medium comprises an analyte, and wherein the system is operative to detect the analyte, when the system is subjected to an electric field sufficient to translocate the analyte from one conductive liquid medium to the other.
24. A system comprising the mutant single-chain Msp of any one of claims 18, 21 and 22, wherein the Msp has a vestibule and a constriction zone that define a tunnel, wherein the

tunnel is positioned between a first conductive liquid medium and a second conductive liquid medium, wherein at least one conductive liquid medium comprises an analyte, and wherein the system is operative to detect the analyte, when the system is subjected to an electric field sufficient to translocate the analyte from one conductive liquid medium to the other.

25. The system of claim 23 or 24, wherein the system comprises a plurality of mutant single-chain Msp.

26. The system of any one of claims 23-25, wherein the Msp is further defined as an Msp comprising the vestibule having a length from about 2 to about 6 nm and a diameter from about 2 to about 6 nm, and the constriction zone having a length from about 0.3 to about 3 nm and a diameter from about 0.3 to about 3 nm, wherein the vestibule and constriction zone together define the tunnel.

27. The system of any one of claims 23-26, wherein the Msp further comprises a molecular motor, wherein the molecular motor moves the analyte into or through the tunnel with an average translocation velocity that is less than the average translocation velocity at which the analyte translocates into or through the tunnel in the absence of the molecular motor.

28. The system of any one of claims 23-27, further comprising a patch-clamp amplifier.

29. The system of any one of claims 23-28, further comprising a data acquisition device.

30. The system of any one of claims 23-29, further comprising one or more temperature regulating devices in communication with the first conductive liquid medium, the second conductive liquid medium, or both.

31. A method for detecting the presence of an analyte, comprising:
- a) applying an electric field sufficient to translocate an analyte from a first conductive medium to a second conductive medium in liquid communication through the mutant Msp of claim 18 or 21; and
 - b) measuring an ion current, wherein a 5% or more reduction in the ion current compared to an ion current level for the mutant single-chain Msp without an analyte present indicates the presence of the analyte in the first conductive medium.
32. The method of claim 31, wherein the mutant single-chain Msp further comprises a molecular motor, wherein the molecular motor moves the analyte into or through a vestibule and a constriction zone that define a tunnel with an average translocation velocity that is less than the average translocation velocity at which the analyte electrophoretically translocates into or through the tunnel in the absence of the molecular motor.
33. The method of claim 31 or 32, further comprising a step of identifying the analyte.
34. The method of claim 33, wherein identifying the analyte comprises measuring the ion current to provide a current pattern, wherein a reduction in the current defines a blockade in the current pattern, and comparing one or more blockades in the current pattern to one or more blockades in a reference current pattern obtained using a known analyte.
35. The method of any one of claims 31-34, wherein the analyte is a nucleotide, a nucleic acid, an amino acid, a peptide, a protein, a polymer, a drug, an ion, a pollutant, a nanoscopic object, or a biological warfare agent.
36. The method of any one of claims 31-34, wherein the analyte is a polymer.
37. The method of claim 36, wherein the polymer is a protein, a peptide or a nucleic acid.

38. The method of claim 36, wherein the polymer is a nucleic acid.
39. The method of claim 38, wherein the nucleic acid is ssDNA, dsDNA, RNA, or a combination thereof.
40. The method of any one of claims 36-39, further comprising identifying one or more units of the polymer.
41. The method of claim 40, wherein identifying one or more units of the polymer comprises measuring the ion current to provide a current pattern comprising a blockade for each polymer unit, and comparing one or more blockades in the current pattern to one or more blockades in a reference current pattern obtained using a polymer having known units.
42. The method of any one of claims 31-41, further comprising determining the concentration, size, molecular weight, shape, or orientation, or any combination thereof, of the analyte.
43. The method of any one of claims 31-42, wherein at least one of the first or second conductive liquid media comprises a plurality of different analytes.
44. The system of any one of claims 23-30, wherein the Msp is positioned in a lipid bilayer between the first conductive liquid medium and the second conductive liquid medium.
45. The system of claim 44, wherein the Msp is contacted with a lipid prior to being positioned in the lipid bilayer.
46. The method of any one of claims 31-43, wherein the Msp is positioned in a lipid bilayer between a first liquid medium and a second liquid medium.

47. The method of claim 46, wherein the Msp is contacted with a lipid prior to being positioned in the lipid bilayer.

48. The system of claim 44, wherein the Msp is refolded in a buffer comprising a refolding agent and a detergent prior to being positioned in the lipid bilayer.

49. The method of claim 46, wherein the Msp is refolded in a buffer comprising a refolding agent and a detergent prior to being positioned in the lipid bilayer.

50. A method of increasing the number of mutant single-chain Msp insertions in a lipid bilayer, comprising:

(a) contacting the mutant single-chain Msp of claim 18 or 21 with a lipid to form a lipid-associated mutant single-chain Msp; and

(b) inserting the lipid-associated mutant single-chain Msp of step (a) into the lipid bilayer.

	1	*	*	*	*	*	
MspA	MKAISRVLIA	MTA	MS	MS	MS	MS	40
MspB	MKAISRVLIA	MTA	MS	MS	MS	MS	40
MspC	MKAISRVLIA	MTA	MS	MS	MS	MS	40
MspD	-----	MTA	MS	MS	MS	MS	39
MspA	FHSGRAKYI	VAGEG	AD	EG	FE	GT	110
MspB	FHSGRAKYI	VAGEG	AD	EG	FE	GT	110
MspC	FHSGRAKYI	VAGEG	AD	EG	FE	GT	110
MspD	FHSGRAKYI	VAGEG	AD	EG	FE	GT	109
MspA	PGVSI	SAD	LGN	GP	GI	QEV	184
MspB	PGVSI	SAD	LGN	GP	GI	QEV	184
MspC	PGVSI	SAD	LGN	GP	GI	QEV	184
MspD	PGVSI	SAD	LGN	GP	GI	QEV	183

1/21

FIGURE 1




FIGURE 2




FIGURE 3

1 2 3 4 5




FIGURE 4

5/21




FIGURE 5

6/21




FIGURE 6

7/21




FIGURE 7

8/21




FIGURE 8A

9/21




FIGURE 8B

10/21




FIGURE 9

11/21




FIGURE 10

12/21




FIGURE 11

13/21




FIGURE 12

14/21




FIGURE 13




FIGURE 14

16/21




FIGURE 15

17/21




FIGURE 16




FIGURE 17

18/21




FIGURE 18




FIGURE 19

19/21




FIGURE 20




FIGURE 21

Channel activity of single-chain MspA PN1

Current trace recordings:




FIGURE 22

