



US00PP18601P2

(12) **United States Plant Patent**  
**Spencer**

(10) **Patent No.:** **US PP18,601 P2**

(45) **Date of Patent:** **Mar. 18, 2008**

(54) **PHYGELIUS PLANT NAMED ‘CROCORPRI’**

(52) **U.S. Cl.** ..... **Plt./263**

(50) Latin Name: *Phygelius aequalis*  
Varietal Denomination: **CROCORPRI**

(58) **Field of Classification Search** ..... Plt./263  
See application file for complete search history.

(76) Inventor: **Malcolm Spencer**, Croftway Hall,  
Yapton Road, Barnham, Bognor Regis,  
West Sussex (GB), PO22 OBQ

*Primary Examiner*—Kent Bell  
*Assistant Examiner*—S. B. McCormick-Ewoldt

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 106 days.

(57) **ABSTRACT**

A new cultivar of *Phygelius* named ‘CROCORPRI’ that is  
characterized by dense compact habit, purple stems, dark-  
green foliage, and coral flowers that bloom late summer and  
fall. In combination these traits set ‘CROCORPRI’ apart  
from all other existing varieties of *Phygelius* known to the  
inventor.

(21) Appl. No.: **11/494,379**

(22) Filed: **Jul. 26, 2006**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

**2 Drawing Sheets**

**1**

**2**

Genus: *Phygelius*. Species: *aequalis*.  
Denomination ‘CROCORPRI’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of *Phygelius*, commonly known as Cape Fuchsia which was  
bred by the inventor at the inventor’s nursery in West  
Sussex, United Kingdom. The new cultivar is known botani-  
cally as *Phygelius aequalis* and is referred to hereinafter by  
the cultivar name ‘CROCORPRI’. ‘CROCORPRI’ is one of  
four co-pending applications by the inventor relating to new  
cultivars of *Phygelius*. The other three co-pending applica-  
tions are titled ‘CROSNOQUE’ (U.S. patent application Ser.  
No. 11/494,374), ‘CROPURPRI’ (U.S. patent application  
Ser. No. 11/493,957) and ‘CROYELSOV’ (U.S. patent  
application Ser. No. 11/494,350). Taken together, these four  
cultivars constitute the Croftway Series of *Phygelius* and  
individual plants are grown as annual container plants in  
most regions of the United States or as perennial plants in  
regions in U.S.D.A. hardiness zones of 8 or higher.

The new *Phygelius* variety named ‘CROCORPRI’ is a  
hybrid plant that resulted from a formal breeding program  
developed by the inventor in 1997. The breeding program  
was developed with the goal of producing new varieties of  
*Phygelius* that exhibit new flower color. In August 1999 the  
inventor chose the parents as breeding candidates. The  
inventor conducted controlled cross-pollination using an  
individual unnamed seedling of *Phygelius aequalis* as the  
female parent, and an individual *Phygelius aequalis* ‘Yellow  
Trumpet’ (unpatented) as the male parent. Cross-pollination  
was conducted at the inventor’s nursery in West Sussex,  
United Kingdom in 1999. The seed was collected and sown  
by the inventor the following February 2000. First flowering  
occurred, and initial assessment was made in July 2000.  
Back-crosses were carried out in September, from which  
new seed was collected and sown in February 2001. The  
first-flowering and final assessment, were made by the inven-  
tor in September 2001.

‘CROCORPRI’ was selected by the inventor at his nurs-  
ery in West Sussex, United Kingdom in 2001. Selection was

based on the criteria of flower color, habit, and free flow-  
ering. ‘CROCORPRI’ is distinguishable from the female  
parent by flower color, free flowering, density, and pro-  
nounced compact habit. The female parent exhibits deep-  
pink flowers, whereas ‘CROCORPRI’ exhibits flowers that  
are coral in color. ‘CROCORPRI’ is distinguishable from the  
male parent by flower color, free flowering, density, and  
pronounced compact habit. The closest comparison plant is  
*Phygelius* ‘Trewidden Pink’ (unpatched). The traits that  
distinguish ‘CROCORPRI’ from ‘Trewidden Pink’ are  
darker flower color, larger flowers, and pronounced free  
flowering.

The new *Phygelius* variety named ‘CROCORPRI’ exhib-  
its dense compact habit, free flowering, purple stems, dark-  
green foliage, and flowers that are coral in color. The plant  
dimensions at maturity are 75 cm. in height and 45 cm. in  
width. ‘CROCORPRI’ blooms late summer and fall, and is  
hardy to USDA Zone 8. Cultural conditions include open  
well-draining soil with good humus content, full sun, and  
moderate to regular water.

‘CROCORPRI’ was first asexually propagated by the  
inventor in 2001 in West Sussex, United Kingdom. The  
method of asexual propagation used was softwood cuttings.  
Since that time, under careful observation, the distinguishing  
characteristics have been determined stable, uniform, and  
reproduce true to type in successive generations of asexual  
propagation.

**SUMMARY OF THE INVENTION**

The following represent the distinguishing characteristics  
of the new *Phygelius* cultivar named ‘CROCORPRI’. In  
combination these traits set ‘CROCORPRI’ apart from all  
other existing varieties of *Phygelius* known to the inventor.  
‘CROCORPRI’ has not been tested under all possible con-  
ditions and phenotypic differences may be observed with  
variations in environmental, climatic, and cultural  
conditions, however, without any variance in genotype.

1. ‘CROCORPRI’ exhibits dense compact habit.
2. ‘CROCORPRI’ exhibits coral flowers late summer and  
fall.

3. 'CROCORPRI' is free flowering.
4. 'CROCORPRI' exhibits dark-green foliage.
5. 'CROCORPRI' exhibits purple stems.
6. 'CROCORPRI' reaches 75 cm. in height and 45 cm. in width at maturity.
7. Cultural conditions of 'CROCORPRI' include full sun, open well-draining soil with good humus content, and moderate to regular water.
8. 'CROCORPRI' is hardy to USDA Zone 8.

## BRIEF DESCRIPTION OF THE DRAWINGS

The accompanying color drawings illustrate the overall appearance of the new *Phygелиus* cultivar named 'CROCORPRI' showing the color as true as it is reasonably possible to obtain in color reproductions of this type. Color in the drawings may differ from the color values cited in the detailed botanical description, which accurately describes the actual color of the new variety 'CROCORPRI'.

The drawing labeled FIG. 1 depicts a side view of the plant illustrating habit.

The drawing labeled FIG. 2 depicts a close-up view of the flower.

All photographs were taken using conventional techniques and although color may appear different from actual color due to light reflectance, they are as accurate as possible by conventional photography.

## BOTANICAL DESCRIPTION OF THE PLANT

The following is a detailed botanical description of the *Phygелиus* cultivar named 'CROCORPRI'. Data was collected in Arroyo Grande, Calif. from plants grown out-of-doors in 1-liter containers. Color determinations are in accordance with the 2001 Royal Horticultural Society Colour Chart of London, England except where general color terms of ordinary dictionary significance are used. The growing requirements are similar to the species. The age of the plant described is 4 months.

Botanical classification: *Phygелиus aequalis* 'CROCORPRI'.

Family: Schrophulariaceae.

Genus: *PHYGELIUS*.

Species: *aequalis*.

Denomination: 'CROCORPRI'.

Common name: Cape fuchsia.

Plant use: Ornamental for use in the landscape.

Parentage: *Phygелиus aequalis* 'CROCORPRI' is a hybrid plant seedling resulting from controlled cross-pollination of the following parents:

*Female parent*.—An individual unnamed *Phygелиus aequalis*.

*Male parent*.—An individual *Phygелиus aequalis* 'Yellow Trumpet'.

Type: Shrub.

Vigor: Vigorous.

Habit: Dense compact.

Dimensions (at one year): 45 cm. in height and 30 cm. in width.

Dimensions (at maturity): 75 cm. in height and 45 cm. in width.

Hardiness: USDA Zone 8.

Asexual propagation: Softwood cuttings.

Root system: Fine and fibrous.

Cultural conditions: Full sun, open well-draining soil with good humus content, and moderate to regular water.

Time to initiate roots: 4–6 weeks.

Crop time: 4–8 months to produce a finished 1-liter container.

Disease susceptibility: Powdery mildew when grown under cover.

Pest susceptibility: Whitefly and greenfly.

Seasonal interest: Flowers late summer and fall.

Stem:

*Stem shape*.—Angular.

*Internode length (average)*.—4 cm.

*Stem width*.—0.40 cm. in width.

*Stem length*.—Ranges from 23–33 cm.

*Stem surface*.—Glabrous.

*Stem color*.—N77A.

Foliage:

*Foliage type*.—Evergreen.

*Leaf arrangement*.—Opposite.

*Leaf division*.—Simple.

*Leaf shape*.—Broadly lanceolate.

*Leaf base*.—Rounded.

*Leaf apex*.—Acute.

*Leaf venation*.—Pinnate.

*Vein color (adaxial and abaxial surfaces)*.—147D.

*Leaf attachment*.—Petiolate.

*Petiole dimensions*.—3.5 cm. in length and 0.25 cm. in width.

*Petiole shape*.—Sulcate.

*Petiole color*.—N77A.

*Petiole surface*.—Glabrous.

*Leaf margin*.—Crenate.

*Leaf dimensions (average)*.—5.50 cm. in length and 3.25 cm. in width.

*Leaf color (adaxial surface)*.—139A.

*Leaf color (abaxial surface)*.—N138B.

*Leaf surfaces (adaxial and abaxial)*.—Glabrous.

*Leaf appearance (adaxial surface)*.—Semi-glossy.

*Leaf appearance (abaxial surface)*.—Matte.

*Stipules*.—Present.

*Stipule color*.—139A.

*Stipule apex*.—Acute.

*Stipule margin*.—Crenate.

*Stipule dimensions*.—6 mm. in length and 6 mm. in width.

*Foliar fragrance*.—None observed.

Flower:

*Inflorescence*.—Raceme.

*Inflorescence dimensions*.—9 cm. in length and 6 cm. in width.

*Inflorescence form*.—One sided.

*Persistent or self-cleaning*.—Self-cleaning.

*Peduncle dimensions*.—1.25 cm. in length and 2 mm. in diameter.

*Peduncle shape*.—Cylindroid.

*Peduncle surface*.—Glabrous.

*Peduncle color*.—N77B.

*Pedicel dimensions*.—0.75 cm. in length and 1.50 mm. in width.

*Pedicel shape*.—Cylindroid.

*Pedicel surface*.—Glabrous.

*Pedicel color*.—N77B.

*Bud shape*.—Claviform.

*Bud surface*.—Glabrous.

*Bud dimensions*.—2.50 cm. in length and 0.50 cm. in width.

*Bud apex.*—Truncate.  
*Bud color.*—51A.  
*Flower shape.*—Tubular.  
*Flower quantity.*—25–30 flowers per inflorescence.  
*Flower diameter.*—1.75 cm.  
*Flower depth.*—4.50 cm.  
*Flower aspect.*—Pendulous.  
*Corolla tube diameter.*—1 cm.  
*Corolla tube surface.*—Glabrous.  
*Corolla tube color.*—51A.  
*Petals.*—5 in number.  
*Petal dimensions.*—5 mm. in length and 5 mm. in width.  
*Petal color (abaxial surface).*—N57A.  
*Petal color (adaxial surface).*—51A.  
*Petals fused or unfused.*—Fused.  
*Petal margin.*—Entire.  
*Petal apex.*—Subacute.  
*Petal surface (abaxial).*—Glabrous.  
*Petal surface (adaxial).*—Stipitate glandular.  
*Calyx dimensions.*—0.50 cm. in diameter and 1 cm in length.  
*Calyx surface.*—Glabrous.  
*Calyx color.*—N77C and 191A.  
*Sepals.*—Five in number.  
*Sepal margin.*—Entire.  
*Sepal surface.*—Glabrous.  
*Sepal dimensions.*—6 mm. in length and 3 mm. in width.  
*Fused or unfused.*—Sepals are overlapping.

*Sepal apex.*—Acute.  
*Blooming period.*—Late summer and fall.  
*Fragrance.*—None observed.

Reproduction organs:

*Stamens.*—4 in number.  
*Stamen attachment.*—Filament adnate to corolla tube.  
*Stamen color.*—58A.  
*Stamen length.*—5 cm.  
*Anther shape.*—Rotund.  
*Anther dimensions.*—2 mm. in width and 3 mm. in length.  
*Anther color.*—161D.  
*Pollen color.*—161D.  
*Pollen quantity.*—Moderate.  
*Pistil.*—One.  
*Pistil color.*—58A.  
*Pistil length.*—5.25 cm.  
*Stigma.*—Penicillate.  
*Stigma color.*—58A.  
*Ovary position.*—Superior.  
*Ovary color.*—N144C.  
*Ovary dimensions.*—0.60 cm. in height and 0.40 cm. in width.  
*Ovary shape.*—Ovoid.

Seed: None observed to date.

It is claimed:

1. A new and distinct variety of *Phygelius* plant named ‘CROCORPRI’ as described and illustrated.

\* \* \* \* \*



Fig. 1



FIG. 2