



US00PP22253P2

(12) **United States Plant Patent**  
**Hurkman**

(10) **Patent No.:** **US PP22,253 P2**

(45) **Date of Patent:** **Nov. 15, 2011**

(54) **IMPATIENS PLANT NAMED ‘BALFIESUNRE’**

(50) Latin Name: *Impatiens walleriana*  
Varietal Denomination: **Balfiesunre**

(75) Inventor: **Margaret M. Hurkman**, Santa Maria,  
CA (US)

(73) Assignee: **Ball Horticultural Company**, West  
Chicago, IL (US)

(\* ) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/803,160**

(22) Filed: **Jun. 21, 2010**

(51) **Int. Cl.**  
**A01H 5/00** (2006.01)

(52) **U.S. Cl.** ..... **Plt./317**

(58) **Field of Classification Search** ..... **Plt./317**  
See application file for complete search history.

*Primary Examiner* — Susan McCormick Ewoldt

(74) *Attorney, Agent, or Firm* — Audrey Charles

(57) **ABSTRACT**

A new and distinct cultivar of *Impatiens* plant named  
‘Balfiesunre’, characterized by its double-type, red-orange  
colored flowers, dark green-colored foliage, and moderately  
vigorous, compact-mounded growth habit, is disclosed.

**1 Drawing Sheet**

**1**

Latin name of genus and species of plant claimed: *Impatiens walleriana*.

Variety denomination: ‘Balfiesunre’.

**BACKGROUND OF THE INVENTION**

The present invention relates to a new and distinct cultivar  
of *Impatiens* plant botanically known as *Impatiens walleri-*  
*ana* and hereinafter referred to by the cultivar name  
‘Balfiesunre’.

The new cultivar originated in a controlled breeding pro-  
gram in Arroyo Grande, Calif. during April 2005. The objec-  
tive of the breeding program was the development of *Impa-*  
*tiens* cultivars with fully double-type flowers, attractive  
flower and foliage coloration, and a freely branching, com-  
pact growth habit.

The new *Impatiens* cultivar is the result of cross-pollina-  
tion. The female (seed) parent of the new cultivar is the  
proprietary *Impatiens walleriana* breeding selection desig-  
nated m3792c-2, not patented, characterized by its semi-  
double type, dark cherry-red colored flowers, dark green-  
colored foliage, and vigorous, upright growth habit. The male  
(pollen) parent of the new cultivar is the proprietary *Impatiens*  
*walleriana* breeding selection designated m3420-1-1-2-1-6,  
not patented, characterized by its double-type, medium red-  
colored flowers, dark green-colored foliage, and vigorous,  
mounded growth habit. The new cultivar was discovered and  
selected as a single flowering plant within the progeny of the  
above stated cross-pollination during June 2006 in a con-  
trolled environment at Arroyo Grande, Calif.

Asexual reproduction of the new cultivar by terminal stem  
cuttings since June 2006 at Arroyo Grande, Calif. and West  
Chicago, Ill. has demonstrated that the new cultivar repro-  
duces true to type with all of the characteristics, as herein  
described, firmly fixed and retained through successive gen-  
erations of such asexual propagation.

**SUMMARY OF THE INVENTION**

The following characteristics of the new cultivar have been  
repeatedly observed and can be used to distinguish  
‘Balfiesunre’ as a new and distinct cultivar of *Impatiens* plant:

**2**

1. Double-type, red-orange colored flowers;
2. Dark green-colored foliage; and
3. Moderately vigorous, compact-mounded growth habit.

Plants of the new cultivar differ from plants of the female  
parent primarily in flower form, flower color, and growth  
vigor and habit. Plants of the new cultivar differ from plants of  
the male parent primarily in flower color and growth vigor.  
The new cultivar has a flower color with more orange coloration  
and is less vigorous than either parent.

Of the many commercially available *Impatiens* cultivars,  
the most similar in comparison to the new cultivar is Musica  
Red Esprit, not patented. However, in side by side compari-  
sons, plants of the new cultivar differ from plants of Musica  
Red Esprit in at least the following characteristics:

1. Plants of the new cultivar have a more iridescent petal  
color than plants of Musica Red Esprit; and
2. Plants of the new cultivar have more flowers than plants  
of Musica Red Esprit. At six weeks after transplant,  
plants of ‘Balfiesunre’ had approximately 8 fully-  
opened flowers per plant, while those of Musica Red  
Esprit had approximately 1 fully-opened flower per  
plant.

**BRIEF DESCRIPTION OF THE PHOTOGRAPHS**

The accompanying photographs show, as nearly true as it is  
reasonably possible to make the same in color illustrations of  
this type, typical flower and foliage characteristics of the new  
cultivar. Colors in the photographs differ slightly from the  
color values cited in the detailed description, which accu-  
rately describes the colors of ‘Balfiesunre’. The plants were  
grown in 4-inch pots for 6 weeks in a greenhouse at West  
Chicago, Ill. Plants were given one pinch at transplant.

FIG. 1 illustrates a side view of the overall growth and  
flowering habit of ‘Balfiesunre’.

FIG. 2 illustrates a close-up view of an individual flower of  
‘Balfiesunre’.

**DETAILED BOTANICAL DESCRIPTION**

The new cultivar has not been observed under all possible  
environmental conditions to date. Accordingly, it is possible

that the phenotype may vary somewhat with variations in the environment, such as temperature, light intensity, and day length, without, however, any variance in genotype.

The chart used in the identification of colors described herein is The R.H.S. Colour Chart of The Royal Horticultural Society, London, England, 2001 edition, except where general color terms of ordinary significance are used. The color values were determined in April 2010 under natural light conditions in West Chicago, Ill.

The following descriptions and measurements describe plants produced from cuttings from stock plants and grown in a glass-covered greenhouse under conditions comparable to those used in commercial practice. The plants were grown in West Chicago, Ill. in 4-inch pots for 6 weeks utilizing a soilless growth medium. Plants were given one pinch at transplant. Greenhouse temperatures were maintained at approximately 70° F. to 77° F. (21° C. to 25° C.) during the day and approximately 65° F. to 68° F. (18° C. to 20° C.) during the night. Greenhouse light levels of 2,500 footcandles to 6,000 footcandles were maintained during the day. Measurements and numerical values represent averages of typical plants.

Botanical classification: *Impatiens walleriana* cultivar Balfiesunre.

Parentage:

*Female parent*.—Proprietary *Impatiens walleriana* breeding selection designated m3792c-2, not patented.

*Male parent*.—Proprietary *Impatiens walleriana* breeding selection designated m3420-1-1-2-1-6, not patented.

Propagation:

*Type cutting*.—Terminal stem.

*Time to initiate roots*.—Approximately 7 to 8 days.

*Time to produce a rooted cutting*.—Approximately 21 to 24 days.

*Root description*.—Fibrous.

*Rooting habit*.—Freely branching.

Plant description:

*Commercial crop time*.—Approximately 5 to 7 weeks from a rooted cutting to finish in a 10 cm pot.

*Growth habit and general appearance*.—Moderately vigorous, compact-mounded growth habit.

*Size*.—Height from soil level to top of plant plane: Approximately 8.0 cm. Width: Approximately 18.8 cm.

*Branching habit*.—Freely branching, pinching improves basal branching. Quantity of main branches per plant: Approximately 4.

*Branch*.—Strength: Moderately strong. Length: Approximately 7.5 cm. Diameter at central internode: Approximately 5.0 mm. Length of central internode: Approximately 1.2 cm. Texture: Glabrous. Color of young stem: 145C with streaks of 187B at the node regions. Color of mature stem: 145C with streaks of 187B at the node regions.

Foliage description:

*General description*.—Quantity of leaves per main branch: Approximately 8. Fragrance: None. Form: Simple. Arrangement: Alternate.

*Leaves*.—Aspect: Acute angle to stem, becoming perpendicular with age. Shape: Ovate. Margin: Crenate, ciliate. Apex: Acute. Base: Attenuate. Venation pattern: Pinnate. Length of mature leaf at center of stem: Approximately 3.1 cm. Width of mature leaf at center of stem: Approximately 2.8 cm. Texture of upper surface: Sparsely pubescent. Texture of lower surface: Glabrous. Color of upper surface of young foliage: 137A with venation of 137C. Color of lower surface

of young and mature foliage: 191B with venation of 138A. Color of upper surface of mature foliage: Slightly darker than 137A with venation of 137C.

*Petiole*.—Length: Approximately 1.2 cm. Diameter: Approximately 2.0 mm. Texture: Glabrous. Color: 145D.

Flowering description:

*Flowering habit*.—‘Balfiesunre’ is freely flowering under outdoor growing conditions with substantially continuous blooming from spring through autumn and year-round in greenhouse environment.

*Lastingness of individual flower on the plant*.—Approximately 6 to 7 days.

Flower description:

*General description*.—Type: Double. Quantity per plant: Approximately 8. Fragrance: None.

*Bud*.—Rate of opening: Generally takes 2 to 3 days for bud to progress from first color to fully open flower. Quantity showing color per plant: Approximately 7.

*Bud just before opening*.—Shape: Ovoid. Length: Approximately 1.3 cm. Diameter: Approximately 1.1 cm. Color: 39B.

*Corolla*.—Shape: Hemispherical. Width: Approximately 4.5 cm. Depth: Approximately 2.3 cm. Borne: Above and below foliage.

*Petals*.—Quantity: 36. Shape: Obovate; innermost are irregular. Aspect: Flat. Appearance: Iridescent. Margin: Entire. Apex: Obtuse; some are emarginate. Base: Attenuate. Length of outermost petals: Approximately 2.3 cm. Width of outermost petals: Approximately 2.3 cm. Length of innermost petals: Approximately 5.0 mm. Width of innermost petals: Approximately 2.0 mm. Texture of upper and lower surfaces: Glabrous. Color of upper surface when first open: 46A. Color of lower surface when first and fully open: 39B with centers of 39C. Color of upper surface when fully open: Between 45A and 45B.

*Spur*.—Quantity: 1 per flower. Length: Approximately 1.3 cm. Diameter at proximal end: Approximately 1.0 mm. Diameter at distal end: Less than 1.0 mm. Texture: Glabrous. Color: 145C, transparent.

*Peduncle*.—Strength: Moderately strong. Aspect: Acute angle to stem. Length: Approximately 2.9 cm. Diameter: Approximately 2.0 mm. Texture: Glabrous. Color: 144B.

*Sepals*.—Quantity per flower: 4, with one fused to base of upper petal. Shape of lateral sepals: Lanceolate. Shape of lower sepal: Ovate. Apex of all sepals: Acute. Length of lateral sepals: Approximately 3.0 mm. Width of lateral sepals: Approximately 1.0 mm. Length of lower sepal: Approximately 1.0 cm. Width of lower sepal: Approximately 8.0 mm. Texture of upper surface of all sepals: Glabrous. Texture of lower surface of all sepals: Glabrous. Color of upper and lower surfaces of lateral sepals: 144A. Color of upper and lower surfaces of lower sepal: 145D with 145B at tip.

*Reproductive organs*.—None observed.

Seed and fruit production: Neither seed nor fruit production has been observed.

Disease and pest resistance: Resistance to pathogens and pests common to *Impatiens* has not been observed.

What is claimed is:

1. A new and distinct cultivar of *Impatiens* plant named ‘Balfiesunre’, substantially as herein shown and described.



FIG. 1



FIG. 2