



US008450091B2

(12) **United States Patent**
Dicosimo et al.

(10) **Patent No.:** **US 8,450,091 B2**
(45) **Date of Patent:** **May 28, 2013**

(54) **PERHYDROLASE PROVIDING IMPROVED SPECIFIC ACTIVITY**

(75) Inventors: **Robert Dicosimo**, Chadds Ford, PA (US); **John Edward Gavagan**, Wilmington, DE (US); **Mark Scott Payne**, Wilmington, DE (US)

(73) Assignee: **E. I. du Pont de Nemours and Company**, Wilmington, DE (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 222 days.

(21) Appl. No.: **13/070,144**

(22) Filed: **Mar. 23, 2011**

(65) **Prior Publication Data**

US 2011/0236338 A1 Sep. 29, 2011

Related U.S. Application Data

(60) Provisional application No. 61/318,050, filed on Mar. 26, 2010.

(51) **Int. Cl.**

C12P 7/24 (2006.01)

C12N 9/00 (2006.01)

C12N 9/16 (2006.01)

C12N 1/20 (2006.01)

C12N 15/00 (2006.01)

C07H 21/04 (2006.01)

(52) **U.S. Cl.**

USPC **435/147**; 435/183; 435/196; 435/252.3; 435/320.1; 536/23.2

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,974,082 A 8/1976 Weyn
4,444,886 A 4/1984 Esders et al.
4,585,150 A 4/1986 Beacham et al.
5,108,457 A 4/1992 Poulouse et al.
5,116,575 A 5/1992 Badertscher et al.
5,281,525 A 1/1994 Mitsushima et al.
5,296,161 A 3/1994 Wiersema et al.
5,338,676 A 8/1994 Mitsushima et al.
5,364,554 A 11/1994 Stanislawski et al.
5,398,846 A 3/1995 Corba et al.
5,528,152 A 6/1996 Hinoshita et al.
5,552,018 A 9/1996 Devenyns
5,624,634 A 4/1997 Brougham et al.
5,683,724 A 11/1997 Hei et al.
5,932,532 A 8/1999 Ghosh et al.
6,183,807 B1 2/2001 Gutzmann et al.
6,210,639 B1 4/2001 Vlass et al.
6,319,888 B2 11/2001 Wei et al.
6,391,840 B1 5/2002 Thompson et al.
6,465,233 B1 10/2002 Knauseder et al.
6,518,307 B2 2/2003 McKenzie et al.
6,545,047 B2 4/2003 Gutzmann et al.
6,645,233 B1 11/2003 Ayers et al.
6,995,125 B2 2/2006 Dasque et al.

7,384,787 B2 6/2008 Kazlauskas et al.
7,550,420 B2 6/2009 DiCosimo et al.
7,723,083 B2 5/2010 DiCosimo et al.
7,807,425 B2 10/2010 DiCosimo et al.
7,829,315 B2 11/2010 DiCosimo et al.
7,910,347 B1 3/2011 DiCosimo et al.
7,923,233 B1 4/2011 DiCosimo et al.
7,927,854 B1 4/2011 DiCosimo et al.
7,932,072 B1 4/2011 DiCosimo et al.
7,951,566 B2 5/2011 DiCosimo et al.
7,951,567 B2 5/2011 DiCosimo et al.
7,960,151 B1 6/2011 DiCosimo et al.
7,960,528 B1 6/2011 DiCosimo et al.
7,964,378 B2 6/2011 DiCosimo et al.
7,964,383 B1 6/2011 DiCosimo et al.
7,981,643 B2 7/2011 DiCosimo et al.
7,981,644 B2 7/2011 DiCosimo et al.
2003/0026846 A1 2/2003 Hei et al.
2005/0008526 A1 1/2005 Bianchetti et al.
2005/0139608 A1 6/2005 Muehlhausen et al.
2008/0176299 A1 7/2008 DiCosimo et al.
2008/0176783 A1 7/2008 DiCosimo et al.
2009/0005590 A1 1/2009 DiCosimo et al.
2009/0311763 A1 12/2009 DiCosimo et al.
2009/0312420 A1 12/2009 DiCosimo et al.
2009/0325266 A1 12/2009 DiCosimo et al.
2010/0041752 A1 2/2010 DiCosimo et al.
2010/0086510 A1 4/2010 DiCosimo et al.
2010/0086621 A1 4/2010 DiCosimo et al.
2010/0087529 A1 4/2010 DiCosimo et al.
2010/0168234 A1 7/2010 DiCosimo et al.
2010/0168235 A1 7/2010 DiCosimo et al.
2010/0168236 A1 7/2010 DiCosimo et al.
2010/0168237 A1 7/2010 DiCosimo et al.
2011/0150857 A1 6/2011 DiCosimo et al.
2011/0152368 A1 6/2011 DiCosimo et al.

FOREIGN PATENT DOCUMENTS

EP 0807156 B1 11/1997
EP 1040222 B1 10/2000

(Continued)

OTHER PUBLICATIONS

Abbott et al., "Physical Properties and Kinetic Behavior of a Cephalosporin Acetylesterase Produced by *Bacillus* . . .", Appl. Microbiol., vol. 30, pp. 413-419 (1975).

(Continued)

Primary Examiner — Christian Fronda

(57) **ABSTRACT**

An acetyl xylan esterase variant having perhydrolytic activity is provided for producing peroxycarboxylic acids from carboxylic acid esters and a source of peroxxygen. More specifically, a *Thermotoga maritima* acetyl xylan esterase gene was modified using error-prone PCR and site-directed mutagenesis to create an enzyme catalyst characterized by an increase in specific activity. The variant acetyl xylan esterase may be used to produce peroxycarboxylic acids suitable for use in a variety of applications such as cleaning, disinfecting, sanitizing, bleaching, wood pulp processing, and paper pulp processing applications.

17 Claims, No Drawings

FOREIGN PATENT DOCUMENTS

WO	WO 97/30160	A1	8/1997
WO	WO 00/11713	A1	2/2000
WO	WO 00/61713	A1	10/2000
WO	WO 2004/058961	A1	7/2004
WO	WO 2007/070609	A1	6/2007
WO	WO 2008/073139	A1	6/2008
WO	WO 2010/019544	A1	2/2010
WO	WO 2010/039953	A1	4/2010
WO	WO 2010/039958	A1	4/2010
WO	WO 2010/039960	A1	4/2010

OTHER PUBLICATIONS

Altschul et al., "Basic Local Alignment Search Tool", J. Mol. Biol., vol. 215, pp. 403-410 (1990).

Bernhardt et al., "Molecular basis of perhydrolase activity in Serine hydrolases", Enzyme Catalysis, vol. 44, pp. 2742-2746 (2005).

Chica et al., "Semi-rational approaches to engineering enzyme activity . . .", Curr. Opin. Biotechnol., vol. 16, pp. 378-384 (2005).

Cardoza et al., "A Cephalosporin C Acetylhydrolase is Present in the Culture of *Nocardia lactamdurans*", Appl. Microbiol. Biotechnol., vol. 54, pp. 406-412 (2000).

Copeland et al., "Thermatogal lettingae acetyl xylan transferase", A8F440_THELT, XP002501372, Nov. 13, 2007.

Degrassi et al., "The Acetyl Xylan Esterase of *Bacillus Pumilus* Belongs to a Family of Esterases with Broad Substrate Specificity", Microbiology, vol. 146, pp. 1585-1591 (2000).

Deshpande et al., "Ethanol Production from Cellulose by Coupled Saccharification/Fermentation . . .", Appl. Biochem. Biotechnol., vol. 36, pp. 227-234 (1992).

Gabrielson et al., "Evaluation of Redox Indicators and the Use of Digital Scanners and . . . Growth in Microplates", J. Microbiol. Methods, vol. 50, pp. 63-73 (2002).

Genbank Accession No. NP_227893.1., (2012).

Guo et al., "Protein tolerance to random amino acid change" PNAS, vol. 101, pp. 9205-9210 (2004).

Higgins et al., "Fast and Sensitive Multiple Sequence Alignments on a Microcomputer", CABIOS, vol. 5, pp. 151-153 (1989).

Pinkernell et al., "Simultaneous HPLC Determination of Peroxyacetic Acid and Hydrogen Peroxide" Anal. Chem., 69(17):3623-3627 (1997).

Kirk et al. Enzyme Catalyzed Degradation and Formation of Peroxycarboxylic Acids, Biocatalysis, vol. 11, pp. 65-77 (1994).

Lennon et al., "The I.M.A.G.E. Consortium: An Integrated Molecular Analysis of Genomes and Their Expression", Genomics, vol. 33, pp. 151-152 (1996).

Lorenz et al., "Isolation, Analysis and Expression of two Genes from Thermoanaerobacterium Sp. Strain . . .", Bacteriol., vol. 179, pp. 5436-5441 (1997).

Minning et al., "Determination of Peracid and Putative Enzymatic Peracid Formation by an Easy Colorimetric Assay", Analytica Chimica Acta, vol. 378, pp. 293-298 (1999).

Mitsushima et al., "Gene cloning, nucleotide sequence, and expression of a cephalosporin-C deacetylase . . ." Appl. Environ. Microbiol., vol. 61, pp. 2224-2229 (1995).

Mitsushima et al, NCBI Gen. Id. No. 550075, Gene Cloning, Nucleotide Sequence & Expression Accession No. BAA01729.1, pp. 1-2 (1999).

Payne et al., "Use of Alkaline Phosphatase Fusions to Studt Protein Secretion in *Bacillus subtilis*", J. Bacteriol., vol. 173, pp. 2278-2282 (1991).

Pearson, "Searching Protein Sequence Databases is Optimal Best?" Comput. Methods Genome Res., [Proc. Int. Symp.] (1994), Meeting Date 1992, pp. 111-120.

Politino et al., "Purification and Characterization of a Cephalosporin Esterase from *Rhodospiridium Toruloides*", Appl. Environ. Microbiol., vol. 63, 4807-4811 (1997).

Rey et al., "Complete Genome Sequence of the Industrial Bacterium *Bacillus licheniformis* and Comparisons with . . .", Genome Biol., vol. 5, pp. 1-13 (2004).

Sakai, et al., "Purification and Properties of Cephalosporin-C Deacetylase from the Yeast . . .", J. Ferment Bioeng., vol. 85, pp. 53-57 (1998).

Seffernick et al., "Melamine deaminase and atrozine chlorohydrolase . . .", J. Bacteriol., vol. 183, pp. 2405-2410 (2001).

Sulter et al., "Proliferation and Metabolic Significance of Peroxisomes in Canadian . . .", Arch. Microbiol., vol. 153, pp. 485-489 (1990).

Takami et al., "Complete Genome Sequence of the . . . Bacterium *Bacillus halodurans* & Genomic Sequence Comparison with . . .", NAR, vol. 28, pp. 4317-4331 (2000).

Vincent et al., "Multifunctional xylooligosaccharide/cephalosporin C deacetylase revealed", J. Mol. Biol., vol. 330, pp. 593-606 (2003).

Witkowski et al., "Conversion of a beta-ketoacyl synthase to a malonyl decarboxylase . . .", Biochemistry, vol. 38, pp. 11643-11650 (1999).

Dicosimo et al., "Thermophilic perhydrolases for peracetic acid production", Internet Citation, XP0002557717, Jul. 30, 3009.

International Search Report in PCT/US2006/047632, mailed Jul. 16, 2007.

International Search Report in PCT/US2007/010644, mailed Jan. 18, 2008.

International Search Report in PCT/US2008/067712, mailed Jan. 16, 2009.

International Search Report and Written Opinion in PCT/US2011/029594, mailed May 30, 2011.

U.S. Appl. No. 12/538,525, filed Aug. 10, 2009, Robert DiCosimo et al.

U.S. Appl. No. 13/070,123, filed Mar. 23, 2011, Robert DiCosimo et al.

U.S. Appl. No. 13/070,130, filed Mar. 23, 2011, Robert DiCosimo et al.

U.S. Appl. No. 13/070,135, filed Mar. 23, 2011, Robert DiCosimo et al.

U.S. Appl. No. 13/070,149, filed Mar. 23, 2011, Robert DiCosimo et al.

U.S. Appl. No. 13/087,800, filed Apr. 15, 2011, Robert DiCosimo et al.

U.S. Appl. No. 13/070,118, filed Mar. 23, 2011, Robert DiCosimo et al.

1

PERHYDROLASE PROVIDING IMPROVED SPECIFIC ACTIVITY

CROSS-REFERENCE TO RELATED APPLICATION

This application claims benefit of U.S. Provisional Patent Application No. 61/318,050, filed Mar. 26, 2010, which is incorporated by reference herein in its entirety.

TECHNICAL FIELD

The invention relates to the field of peroxycarboxylic acid biosynthesis and enzyme catalysis. More specifically, an enzyme catalyst comprising a variant enzyme having perhydrolytic activity is provided having an increase in specific activity. Methods of using the present enzyme catalyst to produce peroxycarboxylic acids are also provided.

BACKGROUND

Peroxycarboxylic acid compositions can be effective antimicrobial agents. Methods of using peroxycarboxylic acids to clean, disinfect, and/or sanitize hard surfaces, textiles, meat products, living plant tissues, and medical devices against undesirable microbial growth have been described (U.S. Pat. Nos. 6,545,047; 6,183,807; 6,518,307; U.S. Patent Application Publication No. 2003-0026846; and U.S. Pat. No. 5,683,724). Peroxycarboxylic acids have also been used in a various bleaching applications including, but not limited to, wood pulp bleaching/delignification and laundry care applications (European Patent 1040222B1; U.S. Pat. Nos. 5,552,018; 3,974,082; 5,296,161; and U.S. Pat. No. 5,364,554). The desired efficacious concentration of peroxycarboxylic acid may vary according to the product application (for example, ca. 500 ppm to 1000 ppm for medical instrument disinfection, ca. 30 ppm to 80 ppm for laundry bleaching or disinfection applications) in 1 min to 5 min reaction time at neutral pH.

Enzymes structurally classified as members of family 7 of the carbohydrate esterases (CE-7) have been employed as perhydrolases to catalyze the reaction of hydrogen peroxide (or alternative peroxide reagent) with alkyl esters of carboxylic acids in water at a basic to acidic pH range (from ca. pH 10 to ca. pH 5) to produce an efficacious concentration of a peroxycarboxylic acid for such applications as disinfection (such as medical instruments, hard surfaces, textiles), bleaching (such as wood pulp or paper pulp processing/delignification, textile bleaching and laundry care applications), and other laundry care applications such as destaining, deodorizing, and sanitization (Published U.S. Patent Application Nos. 2008-0176783, 2008-0176299, 2009-0005590, and 2010-0041752 to DiCosimo et al.). The CE-7 enzymes have been found to have high specific activity for perhydrolysis of esters, particularly acetyl esters of alcohols, diols and glycerols. Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al. describes several variant CE-7 perhydrolases derived from several *Thermotoga* sp. having higher perhydrolytic specific activity and/or improved selectivity for perhydrolysis when used to prepare peroxycarboxylic acid from carboxylic acid esters. One of the variants described in Published U.S. Patent Application No. 2010-0087529, *Thermotoga maritima* C277S, exhibited a significant improvement in specific activity relative to the *T. maritima* wild-type enzyme. However, there remains an ongoing need to identify additional variants having even higher perhydrolytic specific

2

activity as a relative increase in specific activity reduces the amount of enzyme used to achieve a desired peroxycarboxylic acid concentration.

The problem to be solved is to provide an enzyme catalyst comprising a CE-7 carbohydrate esterase having perhydrolytic activity wherein the perhydrolytic enzyme is characterized by a higher specific activity for perhydrolysis of esters, particularly acetyl esters of alcohols, diols and glycerols, when compared to the *Thermotoga maritima* C277S variant.

SUMMARY

A nucleic acid molecule encoding the *Thermotoga maritima* acetyl xylan esterase variant C277S was mutated to create a library of variant enzymes having perhydrolytic activity. Several perhydrolase variants were identified exhibiting an increase in specific activity when compared to the *Thermotoga maritima* C277S perhydrolase under the same assay conditions.

In one embodiment, an isolated nucleic acid molecule encoding a polypeptide having perhydrolytic activity is provided selected from the group consisting of:

- (a) a polynucleotide encoding a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 22;
- (b) a polynucleotide comprising the nucleic acid sequence of SEQ ID NO: 21; and
- (c) a polynucleotide fully complementary to the polynucleotide of (a) or (b).

In other embodiments, a vector, a recombinant DNA construct, and a recombinant host cell comprising the present polynucleotide are also provided.

In another embodiment, a method for transforming a cell is provided comprising transforming a cell with the above nucleic acid molecule.

In another embodiment, an isolated polypeptide having perhydrolysis activity is provided comprising the amino acid sequence of SEQ ID NO: 22.

In one embodiment, the variant polypeptide having perhydrolytic activity is characterized by an increase in specific activity (as determined by an increased amount of peroxycarboxylic acid produced) when compared to the specific activity of the *Thermotoga maritima* C277S variant (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.) under identical reaction conditions.

In another embodiment, a process for producing a peroxycarboxylic acid is also provided comprising:

- (a) providing a set of reaction components comprising:
 - (1) at least one substrate selected from the group consisting of:
 - (i) one or more esters having the structure

$[X]_mR_5$

wherein


X=an ester group of the formula $R_6-C(O)O$;

R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;

R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;


3

- m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and wherein said esters have solubility in water of at least 5 ppm at 25° C.;
- (ii) one or more glycerides having the structure



wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

- (iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

- (iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
- (v) any combination of (i) through (iv);
- (2) a source of peroxygen; and
- (3) an enzyme catalyst comprising a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 22;
- (b) combining the set of reaction components under suitable reaction conditions whereby peroxycarboxylic acid is produced; and
- (c) optionally diluting the peroxycarboxylic acid produced in step (b).

In another embodiment, a process is provided further comprising a step (d) wherein the peroxycarboxylic acid produced in step (b) or step (c) is contacted with a hard surface, an article of clothing or an inanimate object whereby the hard surface, article of clothing or inanimate object is disinfected, sanitized, bleached, destained, deodorized or any combination thereof.

In another embodiment, a composition is provided comprising:

- (a) a set of reaction components comprising:
- (1) at least one substrate selected from the group consisting of:
- (i) one or more esters having the structure



wherein

X=an ester group of the formula $R_6-C(O)O$;

R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;

R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in


4

R_5 individually comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;

m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and


wherein said esters have solubility in water of at least 5 ppm at 25° C.;

- (ii) one or more glycerides having the structure



wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

- (iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

- (iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
- (v) any combination of (i) through (iv);
- (2) a source of peroxygen; and
- (3) an enzyme catalyst comprising a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 22; and
- (b) at least one peroxycarboxylic acid formed upon combining the set of reaction components of (a).

The present process produces the desired peroxycarboxylic acid upon combining the reaction components. The reaction components may remain separated until use.

In a further aspect, a peroxycarboxylic acid generation and delivery system is provided comprising:

- (a) a first compartment comprising
- (1) an enzyme catalyst comprising a polypeptide having perhydrolytic activity, said polypeptide comprising the amino acid sequence of SEQ ID NO: 22;
- (2) at least one substrate selected from the group consisting of:
- (i) one or more esters having the structure



wherein


X=an ester group of the formula $R_6-C(O)O$;

R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;


R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individu-

5

- ally comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;
 m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and
 wherein said esters have solubility in water of at least 5 ppm at 25° C.;
- (ii) one or more glycerides having the structure



- wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;
- (iii) one or more esters of the formula:



- wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;
- (iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
 (v) any combination of (i) through (iv); and
 (3) an optional buffer; and
 (b) a second compartment comprising
 (1) source of peroxygen;
 (2) a peroxide stabilizer; and
 (3) an optional buffer.

In a further embodiment, a laundry care composition is provided comprising a polypeptide comprising the amino acid sequence of SEQ ID NO: 22.

In a further embodiment, a personal care composition is provided comprising a polypeptide comprising the amino acid sequence of SEQ ID NO: 22.

BRIEF DESCRIPTION OF THE BIOLOGICAL SEQUENCES

The following sequences comply with 37 C.F.R. §§1.821-1.825 ("Requirements for Patent Applications Containing Nucleotide Sequences and/or Amino Acid Sequence Disclosures—the Sequence Rules") and are consistent with World Intellectual Property Organization (WIPO) Standard ST.25 (1998) and the sequence listing requirements of the European Patent Convention (EPC) and the Patent Cooperation Treaty (PCT) Rules 5.2 and 49.5(a-bis), and Section 208 and Annex C of the Administrative Instructions. The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. §1.822.

SEQ ID NO: 1 is the nucleic acid sequence of the codon-optimized coding region encoding the wild-type *Thermotoga maritima* acetyl xylan esterase having perhydrolytic activity.

6

SEQ ID NO: 2 is the amino acid sequence of the wild-type *Thermotoga maritima* acetyl xylan esterase having perhydrolytic activity.

SEQ ID NOs: 3 and 4 are the nucleic acid sequences of primers used to prepare the C277S variant acetyl xylan esterase.

SEQ ID NO: 5 is the amino acid sequence of the C277S variant acetyl xylan esterase having perhydrolytic activity (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.).

SEQ ID NO: 6 is the nucleic acid sequence of the plasmid pSW202/C277S.

SEQ ID NOs: 7 and 8 are the nucleic acid sequences of primers used for error-prone PCR.

SEQ ID NO: 9 is the nucleic acid sequence encoding the variant acetyl xylan esterase 843H9 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: (L8R/L125Q/Q176L/V183D/F2471/C277S/P292L).

SEQ ID NO: 10 is the amino acid sequence of the 843H9 variant acetyl xylan esterase.

SEQ ID NO: 11 is the nucleic acid sequence encoding the variant acetyl xylan esterase 843B12 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: A206E/C277S.

SEQ ID NO: 12 is the amino acid sequence of the 843B12 variant acetyl xylan esterase.

SEQ ID NO: 13 is the nucleic acid sequence encoding the variant acetyl xylan esterase 843D9 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: S35R/H50R/C277S.

SEQ ID NO: 14 is the amino acid sequence of the 843D9 variant acetyl xylan esterase.

SEQ ID NO: 15 is the nucleic acid sequence encoding the variant acetyl xylan esterase 843F12 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: K77E/A266E/C277S.

SEQ ID NO: 16 is the amino acid sequence of the 843F12 variant acetyl xylan esterase.

SEQ ID NO: 17 is the nucleic acid sequence encoding the variant acetyl xylan esterase 843C12 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: F27Y/I149V/A266V/C277S/I295T/N302S.

SEQ ID NO: 18 is the amino acid sequence of the 843C12 variant acetyl xylan esterase.

SEQ ID NO: 19 is the nucleic acid sequence encoding the variant acetyl xylan esterase 843H7 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: G119S/L207P/C277S.

SEQ ID NO: 20 is the amino acid sequence of the 843H7 variant acetyl xylan esterase.

SEQ ID NO: 21 is the nucleic acid sequence encoding the variant acetyl xylan esterase 842H3 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: L195Q/C277S.

SEQ ID NO: 22 is the amino acid sequence of the 842H3 variant acetyl xylan esterase.

SEQ ID NO: 23 is the nucleic acid sequence encoding the variant acetyl xylan esterase 841 D3 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: K52E/C277S/Y298F.

SEQ ID NO: 24 is the amino acid sequence of the 841D3 variant acetyl xylan esterase.

SEQ ID NO: 25 is the nucleic acid sequence encoding the variant acetyl xylan esterase 841 D1 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: L125Q/T249S/C277S/A311V.

SEQ ID NO: 26 is the amino acid sequence of the 841D1 variant acetyl xylan esterase.

SEQ ID NO: 27 is the nucleic acid sequence encoding the variant acetyl xylan esterase 841D4 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: F38S/V197I/N275T/C277S.

SEQ ID NO: 28 is the amino acid sequence of the 841D4 variant acetyl xylan esterase.

SEQ ID NO: 29 is the nucleic acid sequence encoding the variant acetyl xylan esterase 841F11 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: K77I/K126N/C277S/K293R.

SEQ ID NO: 30 is the amino acid sequence of the 841F11 variant acetyl xylan esterase.

SEQ ID NO: 31 is the nucleic acid sequence encoding the variant acetyl xylan esterase 841A7 having the following substitutions relative to the wild-type *Thermotoga maritima* acetyl xylan esterase amino acid sequence: Y110F/C277S.

SEQ ID NO: 32 is the amino acid sequence of the 841A7 variant acetyl xylan esterase.

DETAILED DESCRIPTION

A nucleic acid molecule encoding the *Thermotoga maritima* C277S acetyl xylan esterase (SEQ ID NO: 5) was mutated by error-prone PCR and/or site-directed mutagenesis to create a library of variant perhydrolases (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.). Several perhydrolase variants were identified exhibiting an increase in specific activity when compared to the specific activity of the *Thermotoga maritima* C277S perhydrolase having amino acid sequence SEQ ID NO: 5.

In this disclosure, a number of terms and abbreviations are used. The following definitions apply unless specifically stated otherwise.

As used herein, the articles “a”, “an”, and “the” preceding an element or component of the invention are intended to be nonrestrictive regarding the number of instances (i.e., occurrences) of the element or component. Therefore “a”, “an” and “the” should be read to include one or at least one, and the singular word form of the element or component also includes the plural unless the number is obviously meant to be singular.

The term “comprising” means the presence of the stated features, integers, steps, or components as referred to in the claims, but does not preclude the presence or addition of one or more other features, integers, steps, components or groups thereof. The term “comprising” is intended to include embodiments encompassed by the terms “consisting essentially of” and “consisting of”. Similarly, the term “consisting essentially of” is intended to include embodiments encompassed by the term “consisting of”.

As used herein, the term “about” modifying the quantity of an ingredient or reactant employed refers to variation in the numerical quantity that can occur, for example, through typical measuring and liquid handling procedures used for making concentrates or use solutions in the real world; through inadvertent error in these procedures; through differences in

the manufacture, source, or purity of the ingredients employed to make the compositions or carry out the methods; and the like. The term “about” also encompasses amounts that differ due to different equilibrium conditions for a composition resulting from a particular initial mixture. Whether or not modified by the term “about”, the claims include equivalents to the quantities.

Where present, all ranges are inclusive and combinable. For example, when a range of “1 to 5” is recited, the recited range should be construed as including ranges “1 to 4”, “1 to 3”, “1-2”, “1-2 & 4-5”, “1-3 & 5”, and the like.

As used herein, the term “multi-component system” will refer to a system of enzymatically generating peroxy-carboxylic acid wherein the components remain separated until use. As such, the multi-component system will include at least one first component that remains separated from at least one second component. The first and second components are separated in different compartments until use (i.e., using first and second compartments). The design of the multi-component systems will often depend on the physical form of the components to be combined and are described in more detail below.

As used herein, the term “peroxy-carboxylic acid” is synonymous with peracid, peroxyacid, peroxy acid, percarboxylic acid and peroxy acid.

As used herein, the term “peracetic acid” is abbreviated as “PAA” and is synonymous with peroxyacetic acid, ethaneperoxy acid and all other synonyms of CAS Registry Number 79-21-0.

As used herein, the term “monoacetin” is synonymous with glycerol monoacetate, glycerin monoacetate, and glyceryl monoacetate.

As used herein, the term “diacetin” is synonymous with glycerol diacetate; glycerin diacetate, glyceryl diacetate, and all other synonyms of CAS Registry Number 25395-31-7.

As used herein, the term “triacetin” is synonymous with glycerin triacetate; glycerol triacetate; glyceryl triacetate, 1,2,3-triacetoxypropane, 1,2,3-propanetriol triacetate and all other synonyms of CAS Registry Number 102-76-1.

As used herein, the term “monobutylin” is synonymous with glycerol monobutyrate, glycerin monobutyrate, and glyceryl monobutyrate.

As used herein, the term “dibutylin” is synonymous with glycerol dibutyrate and glyceryl dibutyrate.

As used herein, the term “tributylin” is synonymous with glycerol tributyrate, 1,2,3-tributyrylglycerol, and all other synonyms of CAS Registry Number 60-01-5.

As used herein, the term “monopropionin” is synonymous with glycerol monopropionate, glycerin monopropionate, and glyceryl monopropionate.

As used herein, the term “dipropionin” is synonymous with glycerol dipropionate and glyceryl dipropionate.

As used herein, the term “tripropionin” is synonymous with glyceryl tripropionate, glycerol tripropionate, 1,2,3-tripropionylglycerol, and all other synonyms of CAS Registry Number 139-45-7.

As used herein, the term “ethyl acetate” is synonymous with acetic acid ethyl ester, acetoxyethane, ethyl ethanoate, acetic acid ethyl ester, ethanoic acid ethyl ester, ethyl acetic ester and all other synonyms of CAS Registry Number 141-78-6.

As used herein, the term “ethyl lactate” is synonymous with lactic acid ethyl ester and all other synonyms of CAS Registry Number 97-64-3.

As used herein, the terms “acylated sugar” and “acylated saccharide” refer to mono-, di- and polysaccharides comprising at least one acyl group, where the acyl group is selected from the group consisting of straight chain aliphatic carboxy-

lates having a chain length from C2 to C8. Examples include, but are not limited to, glucose pentaacetate, xylose tetraacetate, acetylated xylan, acetylated xylan fragments, β -D-ribofuranose-1,2,3,5-tetraacetate, tri-O-acetyl-D-galactal, and tri-O-acetyl-glucal.

As used herein, the terms “hydrocarbyl”, “hydrocarbyl group”, and “hydrocarbyl moiety” mean a straight chain, branched or cyclic arrangement of carbon atoms connected by single, double, or triple carbon to carbon bonds and/or by ether linkages, and substituted accordingly with hydrogen atoms. Such hydrocarbyl groups may be aliphatic and/or aromatic. Examples of hydrocarbyl groups include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, t-butyl, cyclopropyl, cyclobutyl, pentyl, cyclopentyl, methylcyclopentyl, hexyl, cyclohexyl, benzyl, and phenyl. In one embodiment, the hydrocarbyl moiety is a straight chain, branched or cyclic arrangement of carbon atoms connected by single carbon to carbon bonds and/or by ether linkages, and substituted accordingly with hydrogen atoms.

As used herein, the terms “monoesters” and “diesters” of 1,2-ethanediol, 1,2-propanediol, 1,3-propanediol, 1,2-butanediol, 1,3-butanediol, 2,3-butanediol, 1,4-butanediol, 1,2-pentanediol, 2,5-pentanediol, 1,6-pentanediol, 1,2-hexanediol, 2,5-hexanediol, 1,6-hexanediol, refer to said compounds comprising at least one ester group of the formula RC(O)O, wherein R is a C1 to C7 linear hydrocarbyl moiety.

As used herein, the terms “suitable enzymatic reaction formulation”, “components suitable for generation of a peroxycarboxylic acid”, “suitable reaction components”, “reaction components”, “reaction formulation”, and “suitable aqueous reaction formulation” refer to the materials and water in which the reactants and the enzyme catalyst comprising the present variant polypeptide having perhydrolytic activity come into contact to form the desired peroxycarboxylic acid. The components of the reaction formulation are provided herein and those skilled in the art appreciate the range of component variations suitable for this process. In one embodiment, the enzymatic reaction formulation produces peroxycarboxylic acid in situ upon combining the reaction components. As such, the reaction components may be provided as a multi-component system wherein one or more of the reaction components remains separated until use. The design of systems and means for separating and combining multiple active components are known in the art and generally will depend upon the physical form of the individual reaction components. For example, multiple active fluids (liquid-liquid) systems typically use multi-chamber dispenser bottles or two-phase systems (U.S. Patent Application Publication No. 2005-0139608; U.S. Pat. Nos. 5,398,846; 5,624,634; 6,391,840; E.P. Patent 0807156B1; U.S. Patent Application Publication No. 2005-0008526; and PCT Publication No. WO 00/61713A1) such as found in some bleaching applications wherein the desired bleaching agent is produced upon mixing the reactive fluids. Multi-component formulations and multi-component generation systems to enzymatically produce peroxycarboxylic acids from carboxylic acid esters are described by DiCosimo et al. in Published U.S. Patent Application Nos. 2010-0086510 and 2010-0086621, respectively. Other forms of multi-component systems used to generate peroxycarboxylic acid may include, but are not limited to, those designed for one or more solid components or combinations of solid-liquid components, such as powders used in many commercially available bleaching compositions (e.g., U.S. Pat. No. 5,116,575), multi-layered tablets (e.g., U.S. Pat. No. 6,210,639), water dissolvable packets having multiple com-

partments (e.g., U.S. Pat. No. 6,995,125) and solid agglomerates that react upon the addition of water (e.g., U.S. Pat. No. 6,319,888).

As used herein, the term “substrate” or “carboxylic acid ester substrate” will refer to the reaction components enzymatically perhydrolyzed using the present enzyme catalyst in the presence of a suitable source of peroxygen, such as hydrogen peroxide. In one embodiment, the substrate comprises at least one ester group capable of being enzymatically perhydrolyzed using the enzyme catalyst, whereby a peroxycarboxylic acid is produced.

As used herein, the term “perhydrolysis” is defined as the reaction of a selected substrate with a source of hydrogen peroxide to form a peroxycarboxylic acid. Typically, inorganic peroxide is reacted with the selected substrate in the presence of a catalyst to produce the peroxycarboxylic acid. As used herein, the term “chemical perhydrolysis” includes perhydrolysis reactions in which a substrate (such as a peroxycarboxylic acid precursor) is combined with a source of hydrogen peroxide wherein peroxycarboxylic acid is formed in the absence of an enzyme catalyst. As used herein, the term “enzymatic perhydrolysis” refers to a reaction of a selected substrate with a source of hydrogen peroxide to form a peroxycarboxylic acid, wherein the reaction is catalyzed by an enzyme catalyst having perhydrolysis activity.

As used herein, the term “perhydrolyase activity” refers to the enzyme catalyst activity per unit mass (for example, milligram) of protein, dry cell weight, or immobilized catalyst weight.

As used herein, “one unit of enzyme activity” or “one unit of activity” or “U” is defined as the amount of perhydrolyase activity required for the production of 1 μ mol of peroxycarboxylic acid product (such as peracetic acid) per minute at a specified temperature. “One unit of enzyme activity” may also be used herein to refer to the amount of peroxycarboxylic acid hydrolysis activity required for the hydrolysis of 1 μ mol of peroxycarboxylic acid (e.g., peracetic acid) per minute at a specified temperature.

The present variant CE-7 carbohydrate esterase is characterized by an increase in specific activity when compared to the *Thermotoga maritima* C277S (Published U.S. Patent Application No. 2010-0087529) under the same reaction conditions. As used herein, the “fold increase” in specific activity is measured relative to the specific activity of the *Thermotoga maritima* C277S perhydrolyase (SEQ ID NO: 5) under the same reaction conditions. In one embodiment, the fold increase in specific activity of the variant polypeptide (i.e., variant perhydrolyase) relative to the *Thermotoga maritima* C277S perhydrolyase is at least 1.05, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 3.0, 4.0, 5.0, 6.0, 7.0, 8.0, 9.0, or 10-fold when compared under identical reaction/assay conditions.

As used herein, “identical assay conditions” or “same assay conditions” refer to the conditions used to measure the peracid formation (i.e., perhydrolysis of a carboxylic acid ester substrate) specific activity of the variant polypeptide relative to the respective specific activity of the polypeptide from which it was derived (i.e., *Thermotoga maritima* C277S acetyl xylan esterase of SEQ ID NO: 5). The assay conditions used to measure the respective specific activities should be as close to identical as possible such that only the structure of the polypeptide having perhydrolytic activity varies.

As used herein, the terms “enzyme catalyst” and “perhydrolyase catalyst” refer to a catalyst comprising an enzyme (i.e., a polypeptide) having perhydrolysis activity and may be in the form of a whole microbial cell, permeabilized microbial cell(s), one or more cell components of a microbial cell extract, partially purified enzyme, or purified enzyme. The

enzyme catalyst may also be chemically modified (for example, by pegylation or by reaction with cross-linking reagents). The perhydrolase catalyst may also be immobilized on a soluble or insoluble support using methods well-known to those skilled in the art; see for example, *Immobilization of Enzymes and Cells*; Gordon F. Bickerstaff, Editor; Humana Press, Totowa, N.J., USA; 1997.

The present enzyme catalyst comprises a variant polypeptide having perhydrolytic activity and is structurally classified as a member of the carbohydrase family 7 (CE-7 family) of enzymes (see Coutinho, P.M., Henrissat, B. "Carbohydrate-active enzymes: an integrated database approach" in *Recent Advances in Carbohydrate Bioengineering*, H. J. Gilbert, G. Davies, B. Henrissat and B. Svensson eds., (1999) The Royal Society of Chemistry, Cambridge, pp. 3-12.). The CE-7 family of enzymes has been demonstrated to be particularly effective for producing peroxycarboxylic acids from a variety of carboxylic acid ester substrates when combined with a source of peroxygen (See PCT publication No. WO2007/070609 and U.S. Patent Application Publication Nos. 2008-0176299, 2008-0176783, and 2009-0005590 to DiCosimo et al.; each herein incorporated by reference in their entireties). The CE-7 enzyme family includes cephalosporin C deacetylases (CAHs; E.C. 3.1.1.41) and acetyl xylan esterases (AXEs; E.C. 3.1.1.72). Members of the CE-7 enzyme family share a conserved signature motif (Vincent et al., *J. Mol. Biol.*, 330:593-606 (2003)).

As used herein, the terms "signature motif" and "CE-7 signature motif", refer to conserved structures shared among a family of enzymes having a perhydrolytic activity.

As used herein, "structurally classified as a CE-7 enzyme", "structurally classified as a carbohydrase family 7 enzyme", "structurally classified as a CE-7 carbohydrase esterase", and "CE-7 perhydrolase" will be used to refer to enzymes having perhydrolysis activity that are structurally classified as a CE-7 carbohydrase esterase based on the presence of the CE-7 signature motif (Vincent et al., *supra*). The "signature motif" for CE-7 esterases comprises three conserved motifs (residue position numbering relative to reference sequence SEQ ID NO: 2; the wild-type *Thermotoga*

- a) Arg118-Gly119-Gln120;
- b) Gly186-Xaa187-Ser188-Gln189-Gly190; and
- c) His303-Glu304.

Typically, the Xaa at amino acid residue position 187 is glycine, alanine, proline, tryptophan, or threonine. Two of the three amino acid residues belonging to the catalytic triad are in bold. In one embodiment, the Xaa at amino acid residue position 187 is selected from the group consisting of glycine, alanine, proline, tryptophan, and threonine.

Further analysis of the conserved motifs within the CE-7 carbohydrase esterase family indicates the presence of an additional conserved motif (LXD at amino acid positions 272-274 of SEQ ID NO: 2) that may be used to further define a member of the CE-7 carbohydrase esterase family. In a further embodiment, the signature motif defined above includes a fourth conserved motif defined as:

Leu272-Xaa273-Asp274.

The Xaa at amino acid residue position 273 is typically isoleucine, valine, or methionine. The fourth motif includes the aspartic acid residue (bold) belonging to the catalytic triad (Ser188-Asp274-His303).

As used herein, the terms "cephalosporin C deacetylase" and "cephalosporin C acetyl hydrolase" refer to an enzyme (E.C. 3.1.1.41) that catalyzes the deacetylation of cephalosporins such as cephalosporin C and 7-aminocephalospo-

ranic acid (Mitsushima et al., *Appl. Environ. Microbiol.*, 61(6): 2224-2229 (1995); U.S. Pat. Nos. 5,528,152; and 5,338,676). Enzymes classified as cephalosporin C deacetylases have been shown to often have significant perhydrolytic activity (U.S. Patent Application Publication Nos. 2008-0176783 and 2008-0176299 to DiCosimo et al.).

As used herein, "acetyl xylan esterase" refers to an enzyme (E.C. 3.1.1.72; AXEs) that catalyzes the deacetylation of acetylated xylans and other acetylated saccharides. Enzymes classified as acetyl xylan esterases have been shown to have significant perhydrolytic activity (U.S. Patent Application Publication Nos. 2008-0176783, 2008-0176299, and 2009-0005590, each to DiCosimo et al.).

As used herein, the term "*Thermotoga maritima*" refers to a bacterial cell reported to have acetyl xylan esterase activity (GENBANK® NP_227893.1). In one aspect, the *Thermotoga maritima* strain is *Thermotoga maritima* MSB8. The amino acid sequence of the wild-type enzyme having perhydrolase activity from *Thermotoga maritima* is provided as SEQ ID NO: 2.

As used herein, the terms "variant", "variant polypeptide", and "variant enzyme catalyst" refer to an enzyme catalyst comprising at least one polypeptide (i.e., a perhydrolase) having perhydrolytic activity wherein the polypeptide comprises at least one amino acid change relative to the enzyme/polypeptide from which it was derived (typically the wild-type perhydrolase). Several variant polypeptides are provided herein having perhydrolytic activity and are characterized by an increase in specific activity relative to the *Thermotoga maritima* C277S acetyl xylan esterase having amino acid sequence SEQ ID NO: 5 (see co-owned and copending Published U.S. Patent Application No. 2010-0087529).

For a particular variant perhydrolase, amino acid substitutions are specified with reference to the *Thermotoga maritima* amino acid sequence (SEQ ID NO: 2). The wild-type amino acid (denoted by the standard single letter abbreviation) is followed by the amino acid residue position of SEQ ID NO: 2 followed by the amino acid of the variant (also denoted by the standard single letter abbreviation). For example, "C277S" describes a change in SEQ ID NO: 2 at amino acid residue position 277 where cysteine was changed to serine. The variant polypeptide may be comprised of multiple point substitutions. For example, Y110F/C277S refers to a variant polypeptide having two point substitutions: 1) a change at amino acid residue position 110 where a tyrosine was changed to a phenylalanine, and 2) a change at residue position 277 wherein a cysteine was changed to a serine.

The term "amino acid" refers to the basic chemical structural unit of a protein or polypeptide. The following abbreviations are used herein to identify specific amino acids:

Amino Acid	Three-Letter Abbreviation	One-Letter Abbreviation
Alanine	Ala	A
Arginine	Arg	R
Asparagine	Asn	N
Aspartic acid	Asp	D
Cysteine	Cys	C
Glutamine	Gln	Q
Glutamic acid	Glu	E
Glycine	Gly	G
Histidine	His	H
Isoleucine	Ile	I
Leucine	Leu	L
Lysine	Lys	K
Methionine	Met	M

-continued

Amino Acid	Three-Letter Abbreviation	One-Letter Abbreviation
Phenylalanine	Phe	F
Proline	Pro	P
Serine	Ser	S
Threonine	Thr	T
Tryptophan	Trp	W
Tyrosine	Tyr	Y
Valine	Val	V
Any amino acid (or as defined herein)	Xaa	X

As used herein, the term “biological contaminants” refers to one or more unwanted and/or pathogenic biological entities including, but not limited to microorganisms, spores, viruses, prions, and mixtures thereof. The present enzyme can be used to produce an efficacious concentration of at least one peroxycarboxylic acid useful to reduce and/or eliminate the presence of the viable biological contaminants. In a preferred embodiment, the biological contaminant is a viable pathogenic microorganism.

As used herein, the term “disinfect” refers to the process of destruction of or prevention of the growth of biological contaminants. As used herein, the term “disinfectant” refers to an agent that disinfects by destroying, neutralizing, or inhibiting the growth of biological contaminants. Typically, disinfectants are used to treat inanimate objects or surfaces. As used herein, the term “antiseptic” refers to a chemical agent that inhibits the growth of disease-carrying microorganisms. In one aspect of the embodiment, the biological contaminants are pathogenic microorganisms.

As used herein, the term “sanitary” means of or relating to the restoration or preservation of health, typically by removing, preventing or controlling an agent that may be injurious to health. As used herein, the term “sanitize” means to make sanitary. As used herein, the term “sanitizer” refers to a sanitizing agent. As used herein the term “sanitization” refers to the act or process of sanitizing.

As used herein, the term “virucide” refers to an agent that inhibits or destroys viruses, and is synonymous with “viricide”. An agent that exhibits the ability to inhibit or destroy viruses is described as having “virucidal” activity. Peroxycarboxylic acids can have virucidal activity. Typical alternative virucides known in the art which may be suitable for use with the present invention include, for example, alcohols, ethers, chloroform, formaldehyde, phenols, beta propiolactone, iodine, chlorine, mercury salts, hydroxylamine, ethylene oxide, ethylene glycol, quaternary ammonium compounds, enzymes, and detergents.

As used herein, the term “biocide” refers to a chemical agent, typically broad spectrum, which inactivates or destroys microorganisms. A chemical agent that exhibits the ability to inactivate or destroy microorganisms is described as having “biocidal” activity. Peroxycarboxylic acids can have biocidal activity. Typical alternative biocides known in the art, which may be suitable for use in the present invention include, for example, chlorine, chlorine dioxide, chloroisocyanurates, hypochlorites, ozone, acrolein, amines, chlorinated phenolics, copper salts, organo-sulphur compounds, and quaternary ammonium salts.

As used herein, the phrase “minimum biocidal concentration” refers to the minimum concentration of a biocidal agent that, for a specific contact time, will produce a desired lethal, irreversible reduction in the viable population of the targeted microorganisms. The effectiveness can be measured by the log₁₀ reduction in viable microorganisms after treatment. In

one aspect, the targeted reduction in viable microorganisms after treatment is at least a 3-log reduction, more preferably at least a 4-log reduction, and most preferably at least a 5-log reduction. In another aspect, the minimum biocidal concentration is at least a 6-log reduction in viable microbial cells.

As used herein, the terms “peroxygen source” and “source of peroxygen” refer to compounds capable of providing hydrogen peroxide at a concentration of about 1 mM or more when in an aqueous solution including, but not limited to, hydrogen peroxide, hydrogen peroxide adducts (e.g., urea-hydrogen peroxide adduct (carbamide peroxide)), perborates, and percarbonates, such as sodium percarbonate. As described herein, the concentration of the hydrogen peroxide provided by the peroxygen compound in the aqueous reaction formulation is initially at least 1 mM or more upon combining the reaction components. In one embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 10 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 100 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is at least 200 mM. In another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is 500 mM or more. In yet another embodiment, the hydrogen peroxide concentration in the aqueous reaction formulation is 1000 mM or more. The molar ratio of the hydrogen peroxide to enzyme substrate, such as triglyceride, (H₂O₂:substrate) in the aqueous reaction formulation may be from about 0.002 to 20, preferably about 0.1 to 10, and most preferably about 0.5 to 5.

As used herein, the term “benefit agent” refers to a material that promotes or enhances a useful advantage, a favorable/desirable effect or benefit. In one embodiment, a process is provided whereby a benefit agent, such as a composition comprising a peroxycarboxylic acid, is applied to a textile or article of clothing to achieve a desired benefit, such as disinfecting, bleaching, destaining, deodorizing, and any combination thereof. In another embodiment, the present variant polypeptide having perhydrolytic activity may be used to produce a peracid-based benefit agent for use in personal care products (such as hair care products, skin care products, nail care products or oral care products). In one embodiment, a personal care product is provided comprising the variant perhydrolytic having amino acid sequence SEQ ID NO: 22. The personal care products are formulated to provide a safe and efficacious concentration of the desired peracid benefit agent. Variant Polypeptides Having an Increase in Specific Activity.

The present variant polypeptides were derived from the *Thermotoga maritima* C277S acetyl xylan esterase that has been previously demonstrated to have significant perhydrolytic activity for producing peroxycarboxylic acids from carboxylic acid esters and a source of peroxygen, such as hydrogen peroxide (U.S. Patent Application Publication Nos. 2008-0176299 and 2010-0087529, each to DiCosimo et al.).

A library of variant polypeptides was created from the C277S *Thermotoga maritima* perhydrolytic (SEQ ID NO: 5) and assayed for an increase in the specific activity for producing peroxycarboxylic acids from carboxylic acid ester substrates. The assay conditions used to measure the respective specific activities should be as close to identical as possible such that only the structure of the polypeptide having perhydrolytic activity varies. In one embodiment, reactions used to measure specific activity are run at ca. 25° C. in phosphate buffer (50 mM, pH 7.2) containing 25 mM triacetin, 25 mM hydrogen peroxide and approximately 2.0 µg/mL of heat-treated extract supernatant total protein from *E. coli* strain KLP18 expressing the C277S perhydrolytic or variant

15

perhydrolase (see Example 5). In another embodiment, the reactions to measure specific activity are conducted under simulated laundry care conditions at ca. 20° C. using 0 to 6 mg/mL liquid laundry detergent, 2 mM triacetin, 10 mM hydrogen peroxide (from sodium percarbonate), hard water plus percarbonate as a buffer (pH 10), and 2 µg/mL to 6 µg/mL of heat-treated extract supernatant total soluble protein from *E. coli* strain KLP18 expressing the *T. maritima* C277S perhydrolase or the present variant perhydrolase (see Example 6).

Suitable Reaction Conditions for the Enzyme-catalyzed Preparation of Peroxycarboxylic acids from Carboxylic Acid Esters and Hydrogen Peroxide

A process is provided to produce an aqueous formulation comprising at least one peroxycarboxylic acid by reacting carboxylic acid esters and an inorganic peroxide (such as hydrogen peroxide, sodium perborate or sodium percarbonate) in the presence of an enzyme catalyst having perhydrolysis activity, wherein the enzyme catalyst comprises, in one embodiment, a polypeptide having an amino acid sequence selected from the group consisting of SEQ ID NOs: 10, 16, 24, 28, and 32. In another embodiment, the polypeptide has an amino acid sequence selected from the group consisting of SEQ ID NOs: 10, 16, 18, 22, and 32. In a further embodiment, the polypeptide has the amino acid sequence of SEQ ID NO: 22.

In one embodiment, suitable substrates include one or more esters provided by the following formula:



wherein X=an ester group of the formula $R_6C(O)O$


R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;

R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;

m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and wherein said esters have solubility in water of at least 5 ppm at 25° C.


In another embodiment, R_6 is C1 to C7 linear hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, optionally comprising one or more ether linkages. In a further preferred embodiment, R_6 is C2 to C7 linear hydrocarbyl moiety, optionally substituted with hydroxyl groups, and/or optionally comprising one or more ether linkages.

In another embodiment, suitable substrates also include one or more glycerides of the formula:



wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$.

In another aspect, suitable substrates may also include one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_n$, H and n is 1 to 10.

Suitable substrates may also include one or more acylated saccharides selected from the group consisting of acylated mono-, di-, and polysaccharides. In another embodiment, the acylated saccharides are selected from the group consisting of acetylated xylan, fragments of acetylated xylan, acetylated xylose (such as xylose tetraacetate), acetylated glucose (such as glucose pentaacetate), β -D-ribofuranose-1,2,3,5-tetraacetate, tri-O-acetyl-D-galactal, tri-O-acetyl-D-glucal, and acetylated cellulose. In a preferred embodiment, the acetylated saccharide is selected from the group consisting of β -D-ribofuranose-1,2,3,5-tetraacetate, tri-O-acetyl-D-galactal, tri-O-acetyl-D-glucal, and acetylated cellulose.

In another embodiment, suitable substrates are selected from the group consisting of: monoacetin; diacetin; triacetin; monopropionin; dipropionin; tripropionin; monobutyryn; dibutyryn; tributyrin; glucose pentaacetate; xylose tetraacetate; acetylated xylan; acetylated xylan fragments; β -D-ribofuranose-1,2,3,5-tetraacetate; tri-O-acetyl-D-galactal; tri-O-acetyl-D-glucal; monoesters or diesters of 1,2-ethanediol, 1,2-propanediol, 1,3-propanediol, 1,2-butanediol, 1,3-butanediol, 2,3-butanediol, 1,4-butanediol, 1,2-pentanediol, 2,5-pentanediol, 1,6-pentanediol, 1,2-hexanediol, 2,5-hexanediol, 1,6-hexanediol; and mixtures thereof.

In another embodiment, the carboxylic acid ester is selected from the group consisting of monoacetin, diacetin, triacetin, and combinations thereof. In another embodiment, the substrate is a C1 to C6 polyol comprising one or more ester groups. In a preferred embodiment, one or more of the hydroxyl groups on the C1 to C6 polyol are substituted with one or more acetoxy groups (such as 1,3-propanediol diacetate, 1,4-butanediol diacetate, etc.). In a further embodiment, the substrate is propylene glycol diacetate (PGDA), ethylene glycol diacetate (EGDA), or a mixture thereof.

In another embodiment, suitable substrates are selected from the group consisting of ethyl acetate; methyl lactate; ethyl lactate; methyl glycolate; ethyl glycolate; methyl methoxyacetate; ethyl methoxyacetate; methyl 3-hydroxybutyrate; ethyl 3-hydroxybutyrate; triethyl 2-acetyl citrate; glucose pentaacetate; gluconolactone; glycerides (mono-, di-, and triglycerides) such as monoacetin, diacetin, triacetin, monopropionin, dipropionin (glyceryl dipropionate), tripropionin (1,2,3-tripropionylglycerol), monobutyryn, dibutyryn (glyceryl dibutyrate), tributyrin (1,2,3-tributyrylglycerol); acetylated saccharides; and mixtures thereof.

In a further embodiment, suitable substrates are selected from the group consisting of monoacetin, diacetin, triacetin, monopropionin, dipropionin, tripropionin, monobutyryn, dibutyryn, tributyrin, ethyl acetate, and ethyl lactate. In yet another aspect, the substrate is selected from the group consisting of diacetin, triacetin, ethyl acetate, and ethyl lactate. In a most preferred embodiment, the suitable substrate comprises triacetin.

The carboxylic acid ester is present in the aqueous reaction formulation at a concentration sufficient to produce the desired concentration of peroxycarboxylic acid upon enzyme-catalyzed perhydrolysis. The carboxylic acid ester

16

need not be completely soluble in the aqueous reaction formulation, but has sufficient solubility to permit conversion of the ester by the perhydrolase catalyst to the corresponding peroxycarboxylic acid. The carboxylic acid ester is present in the aqueous reaction formulation at a concentration of 0.0005 wt % to 40 wt % of the aqueous reaction formulation, preferably at a concentration of 0.01 wt % to 20 wt % of the aqueous reaction formulation, and more preferably at a concentration of 0.05 wt % to 10 wt % of the aqueous reaction formulation. The wt % of carboxylic acid ester may optionally be greater than the solubility limit of the carboxylic acid ester, such that the concentration of the carboxylic acid ester is at least 0.0005 wt % in the aqueous reaction formulation that is comprised of water, enzyme catalyst, and source of peroxide, where the remainder of the carboxylic acid ester remains as a second separate phase of a two-phase aqueous/organic reaction formulation. Not all of the added carboxylic acid ester must immediately dissolve in the aqueous reaction formulation, and after an initial mixing of all reaction components, additional continuous or discontinuous mixing is optional.

The peroxycarboxylic acids produced by the present reaction components may vary depending upon the selected substrates, so long as the present enzyme catalyst is used. In one embodiment, the peroxycarboxylic acid produced is peracetic acid, perpropionic acid, perbutyric acid, perlactic acid, perglycolic acid, permethoxyacetic acid, per- β -hydroxybutyric acid, or mixtures thereof.

The peroxygen source may include, but is not limited to, hydrogen peroxide, hydrogen peroxide adducts (e.g., urea-hydrogen peroxide adduct (carbamide peroxide)), perborate salts and percarbonate salts. The concentration of peroxygen compound in the aqueous reaction formulation may range from 0.0033 wt % to about 50 wt %, preferably from 0.033 wt % to about 40 wt %, more preferably from 0.33 wt % to about 30 wt %.

Many perhydrolase catalysts (such as whole cells, permeabilized whole cells, and partially purified whole cell extracts) have been reported to have catalase activity (EC 1.11.1.6). Catalases catalyze the conversion of hydrogen peroxide into oxygen and water. In one aspect, the enzyme catalyst having perhydrolase activity lacks catalase activity. In another aspect, a catalase inhibitor is added to the aqueous reaction formulation. Examples of catalase inhibitors include, but are not limited to, sodium azide and hydroxylamine sulfate. One of skill in the art can adjust the concentration of catalase inhibitor as needed. The concentration of the catalase inhibitor typically ranges from 0.1 mM to about 1 M; preferably about 1 mM to about 50 mM; more preferably from about 1 mM to about 20 mM. In one aspect, sodium azide concentration typically ranges from about 20 mM to about 60 mM while hydroxylamine sulfate is concentration is typically about 0.5 mM to about 30 mM, preferably about 10 mM.

The catalase activity in a host cell can be down-regulated or eliminated by disrupting expression of the gene(s) responsible for the catalase activity using well known techniques including, but not limited to, transposon mutagenesis, RNA antisense expression, targeted mutagenesis, and random mutagenesis. In a preferred embodiment, the gene(s) encoding the endogenous catalase activity are down-regulated or disrupted (i.e., "knocked-out"). As used herein, a "disrupted" gene is one where the activity and/or function of the protein encoded by the modified gene is no longer present. Means to disrupt a gene are well-known in the art and may include, but are not limited to, insertions, deletions, or mutations to the gene so long as the activity and/or function of the correspond-

ing protein is no longer present. In a further preferred embodiment, the production host is an *E. coli* production host comprising a disrupted catalase gene selected from the group consisting of *katG* and *katE* (see U.S. Patent Application Publication No. 2008-0176783 to DiCosimo et al.). In another embodiment, the production host is an *E. coli* strain comprising a down-regulation and/or disruption in both *katG* and *katE* catalase genes. An *E. coli* strain comprising a double-knockout of *katG* and *katE* has been prepared and is described as *E. coli* strain KLP18 (U.S. Patent Application Publication No. 2008-0176783 to DiCosimo et al.).

The concentration of the catalyst in the aqueous reaction formulation depends on the specific catalytic activity of the catalyst, and is chosen to obtain the desired rate of reaction. The weight of catalyst in perhydrolysis reactions typically ranges from 0.0001 mg to 50 mg per mL of total reaction volume, preferably from 0.0005 mg to 10 mg per mL, more preferably from 0.0010 mg to 2.0 mg per mL. The catalyst may also be immobilized on a soluble or insoluble support using methods well-known to those skilled in the art; see for example, *Immobilization of Enzymes and Cells*; Gordon F. Bickerstaff, Editor; Humana Press, Totowa, N.J., USA; 1997. The use of immobilized catalysts permits the recovery and reuse of the catalyst in subsequent reactions. The enzyme catalyst may be in the form of whole microbial cells, permeabilized microbial cells, microbial cell extracts, partially-purified or purified enzymes, and mixtures thereof.

In one aspect, the concentration of peroxycarboxylic acid generated by the combination of chemical perhydrolysis and enzymatic perhydrolysis of the carboxylic acid ester is sufficient to provide an effective concentration of peroxycarboxylic acid for disinfection, bleaching, sanitization, deodorizing or destaining at a desired pH. In another aspect, the peroxycarboxylic acid is generated at a safe and efficacious concentration suitable for use in a personal care product to be applied to the hair, skin, nails or tissues of the oral cavity, such as tooth enamel, tooth pellicle or the gums. In another aspect, the present methods provide combinations of enzymes and enzyme substrates to produce the desired effective concentration of peroxycarboxylic acid, where, in the absence of added enzyme, there is a significantly lower concentration of peroxycarboxylic acid produced. Although there may be some chemical perhydrolysis of the enzyme substrate by direct chemical reaction of inorganic peroxide with the enzyme substrate, there may not be a sufficient concentration of peroxycarboxylic acid generated to provide an effective concentration of peroxycarboxylic acid in the desired applications, and a significant increase in total peroxycarboxylic acid concentration is achieved by the addition of an appropriate perhydrolase catalyst to the aqueous reaction formulation.

In one aspect of the invention, the concentration of peroxycarboxylic acid generated (e.g. peracetic acid) by the enzymatic perhydrolysis is at least about 2 ppm, preferably at least 20 ppm, preferably at least 100 ppm, more preferably at least 200 ppm peroxycarboxylic acid, more preferably at least 300 ppm, more preferably at least 500 ppm, more preferably at least 700 ppm, more preferably at least about 1000 ppm peroxycarboxylic acid, most preferably at least 2000 ppm peroxycarboxylic acid within 5 minutes more preferably within 1 minute of initiating the enzymatic perhydrolysis reaction. In a second aspect of the invention, the concentration of peroxycarboxylic acid generated (e.g. peracetic acid) by the enzymatic perhydrolysis is at least about 2 ppm, preferably at least 20 ppm, preferably at least 30 ppm, more preferably at least about 40 ppm peroxycarboxylic acid, more preferably at least 50 ppm, more preferably at least 60 ppm, more preferably at least 70 ppm, more preferably at least

about 80 ppm peroxycarboxylic acid, most preferably at least 100 ppm peroxycarboxylic acid within 5 minutes, more preferably within 1 minute, of initiating the enzymatic perhydrolysis reaction (i.e., time measured from combining the reaction components to form the formulation).

The aqueous formulation comprising the peroxycarboxylic acid may be optionally diluted with diluent comprising water, or a solution predominantly comprised of water, to produce a formulation with the desired lower target concentration of peroxycarboxylic acid. In one aspect, the reaction time required to produce the desired concentration (or concentration range) of peroxycarboxylic acid is about 20 minutes or less, preferable about 5 minutes or less, most preferably about 1 minute or less.

In other aspects, the surface or inanimate object contaminated with a concentration of a biological contaminant(s) is contacted with the peroxycarboxylic acid formed in accordance with the processes described herein within about 1 minute to about 168 hours of combining said reaction components, or within about 1 minute to about 48 hours, or within about 1 minute to 2 hours of combining said reaction components, or any such time interval therein.

In another aspect, the peroxycarboxylic acid formed in accordance with the processes describe herein is used in a laundry care application wherein the peroxycarboxylic acid is contacted with clothing or a textile to provide a benefit, such as disinfecting, bleaching, destaining, deodorizing and/or a combination thereof. The peroxycarboxylic acid may be used in a variety of laundry care products including, but not limited to, laundry or textile pre-wash treatments, laundry detergents or additives, stain removers, bleaching compositions, deodorizing compositions, and rinsing agents. In one embodiment, the present process to produce a peroxycarboxylic acid for a target surface is conducted in situ.

In the context of laundry care applications, the term "contacting an article of clothing or textile" means that the article of clothing or textile is exposed to a formulation disclosed herein. To this end, there are a number of formats the formulation may be used to treat articles of clothing or textiles including, but not limited to, liquid, solids, gel, paste, bars, tablets, spray, foam, powder, or granules and can be delivered via hand dosing, unit dosing, dosing from a substrate, spraying and automatic dosing from a laundry washing or drying machine. Granular compositions can also be in compact form; liquid compositions can also be in a concentrated form.

When the formulations disclosed herein are used in a laundry washing machine, the formulation can further contain components typical to laundry detergents. For example, typical components included, but are not limited to, surfactants, bleaching agents, bleach activators, additional enzymes, suds suppressors, dispersants, lime-soap dispersants, soil suspension and anti-redeposition agents, softening agents, corrosion inhibitors, tarnish inhibitors, germicides, pH adjusting agents, non-builder alkalinity sources, chelating agents, organic and/or inorganic fillers, solvents, hydrotropes, optical brighteners, dyes, and perfumes. For example, a liquid laundry detergent was used in Example 6 comprising 15-30% anionic and non-ionic surfactants, 5-15% soap, and <5% polycarboxylates, perfume, phosphonates, optical brighteners.

The formulations disclosed herein can also be used as detergent additive products in solid or liquid form. Such additive products are intended to supplement or boost the performance of conventional detergent compositions and can be added at any stage of the cleaning process.

In connection with the present systems and methods for laundry care where the peracid is generated for one or more of

bleaching, stain removal, and odor reduction, the concentration of peracid generated (e.g., peracetic acid) by the perhydrolysis of at least one carboxylic acid ester may be at least about 2 ppm, preferably at least 20 ppm, preferably at least 100 ppm, and more preferably at least about 200 ppm peracid. In connection with the present systems and methods for laundry care where the peracid is generated for disinfection or sanitization, the concentration of peracid generated (e.g., peracetic acid) by the perhydrolysis of at least one carboxylic acid ester may be at least about 2 ppm, more preferably at least 20 ppm, more preferably at least 200 ppm, more preferably at least 500 ppm, more preferably at least 700 ppm, more preferably at least about 1000 ppm peracid, most preferably at least 2000 ppm peracid within 10 minutes, preferably within 5 minutes, and most preferably within 1 minute of initiating the perhydrolysis reaction. The product formulation comprising the peracid may be optionally diluted with water, or a solution predominantly comprised of water, to produce a formulation with the desired lower concentration of peracid. In one aspect of the present methods and systems, the reaction time required to produce the desired concentration of peracid is not greater than about two hours, preferably not greater than about 30 minutes, more preferably not greater than about 10 minutes, even more preferably not greater than about 5 minutes, and most preferably in about 1 minute or less.

The temperature of the reaction is chosen to control both the reaction rate and the stability of the enzyme catalyst activity. The temperature of the reaction may range from just above the freezing point of the aqueous reaction formulation (approximately 0° C.) to about 85° C., with a preferred range of reaction temperature of from about 5° C. to about 75° C.

The pH of the aqueous reaction formulation while enzymatically producing peroxycarboxylic acid is maintained at a pH ranging from about 5.0 to about 10.0, preferably about 6.5 to about 8.5, and yet even more preferably about 6.5 to about 7.5. In one embodiment, the pH of the aqueous reaction formulation ranges from about 6.5 to about 8.5 for at least 30 minutes after combining the reaction components. The pH of the aqueous reaction formulation may be adjusted or controlled by the addition or incorporation of a suitable buffer, including, but not limited to, phosphate, pyrophosphate, bicarbonate, acetate, or citrate. In one embodiment, the buffer is selected from a phosphate buffer, a bicarbonate buffer, or a buffer formed by the combination of hard ward (tap water to simulate laundry care applications) and percarbonate (from sodium percarbonate used to generate hydrogen peroxide). The concentration of buffer, when employed, is typically from 0.1 mM to 1.0 M, preferably from 1 mM to 300 mM, most preferably from 10 mM to 100 mM. In another aspect of the present invention, no buffer is added to the reaction mixture while enzymatically producing peroxycarboxylic acid.

In yet another aspect, the enzymatic perhydrolysis aqueous reaction formulation may contain an organic solvent that acts as a dispersant to enhance the rate of dissolution of the carboxylic acid ester in the aqueous reaction formulation. Such solvents include, but are not limited to, propylene glycol methyl ether, acetone, cyclohexanone, diethylene glycol butyl ether, tripropylene glycol methyl ether, diethylene glycol methyl ether, propylene glycol butyl ether, dipropylene glycol methyl ether, cyclohexanol, benzyl alcohol, isopropanol, ethanol, propylene glycol, and mixtures thereof.

In another aspect, the enzymatic perhydrolysis product may contain additional components that provide desirable functionality. These additional components include, but are not limited to, buffers, detergent builders, thickening agents, emulsifiers, surfactants, wetting agents, corrosion inhibitors (e.g., benzotriazole), enzyme stabilizers, and peroxide stabi-

lizers (e.g., metal ion chelating agents). Many of the additional components are well known in the detergent industry (see, for example, U.S. Pat. No. 5,932,532; hereby incorporated by reference). Examples of emulsifiers include, but are not limited to, polyvinyl alcohol or polyvinylpyrrolidone. Examples of thickening agents include, but are not limited to, LAPONITE® RD, corn starch, PVP, CARBOWAX®, CARBOPOL®, CABOSIL®, polysorbate 20, PVA, and lecithin. Examples of buffering systems include, but are not limited to, sodium phosphate monobasic/sodium phosphate dibasic; sulfamic acid/triethanolamine; citric acid/triethanolamine; tartaric acid/triethanolamine; succinic acid/triethanolamine; and acetic acid/triethanolamine. Examples of surfactants include, but are not limited to, a) non-ionic surfactants such as block copolymers of ethylene oxide or propylene oxide, ethoxylated or propoxylated linear and branched primary and secondary alcohols, and aliphatic phosphine oxides b) cationic surfactants such as quaternary ammonium compounds, particularly quaternary ammonium compounds having a C8-C20 alkyl group bound to a nitrogen atom additionally bound to three C1-C2 alkyl groups, c) anionic surfactants such as alkane carboxylic acids (e.g., C8-C20 fatty acids), alkyl phosphonates, alkane sulfonates (e.g., sodium dodecylsulphate "SDS") or linear or branched alkyl benzene sulfonates, alkene sulfonates and d) amphoteric and zwitterionic surfactants such as aminocarboxylic acids, aminodicarboxylic acids, alkybetaines, and mixtures thereof. Additional components may include fragrances, dyes, stabilizers of hydrogen peroxide (e.g., metal chelators such as 1-hydroxyethylidene-1,1-diphosphonic acid (DEQUEST® 2010, Solutia Inc., St. Louis, Mo.) and ethylenediaminetetraacetic acid (EDTA)), TURPINAL® SL, DEQUEST® 0520, DEQUEST® 0531, stabilizers of enzyme activity (e.g., polyethylene glycol (PEG)), and detergent builders.

In another aspect, the enzymatic perhydrolysis product may be pre-mixed to generate the desired concentration of peroxycarboxylic acid prior to contacting the surface or inanimate object to be disinfected.

In another aspect, the enzymatic perhydrolysis product is not pre-mixed to generate the desired concentration of peroxycarboxylic acid prior to contacting the surface or inanimate object to be disinfected, but instead, the components of the aqueous reaction formulation that generate the desired concentration of peroxycarboxylic acid are contacted with the surface or inanimate object to be disinfected and/or bleached or destained, generating the desired concentration of peroxycarboxylic acid. In some embodiments, the components of the aqueous reaction formulation combine or mix at the locus. In some embodiments, the reaction components are delivered or applied to the locus and subsequently mix or combine to generate the desired concentration of peroxycarboxylic acid.

Production of Peroxycarboxylic acids using a Perhydrolase Catalyst

The peroxycarboxylic acids, once produced, are quite reactive and may decrease in concentration over extended periods of time, depending on variables that include, but are not limited to, temperature and pH. As such, it may be desirable to keep the various reaction components separated, especially for liquid formulations. In one aspect, the hydrogen peroxide source is separate from either the substrate or the perhydrolase catalyst, preferably from both. This can be accomplished using a variety of techniques including, but not limited to, the use of multicompartment chambered dispensers (U.S. Pat. No. 4,585,150) and at the time of use physically combining the perhydrolase catalyst with a source of peroxygen (such as hydrogen peroxide) and the present substrates to initiate the

aqueous enzymatic perhydrolysis reaction. The perhydrolase catalyst may optionally be immobilized within the body of reaction chamber or separated (e.g., filtered, etc.) from the reaction product comprising the peroxycarboxylic acid prior to contacting the surface and/or object targeted for treatment. The perhydrolase catalyst may be in a liquid matrix or in a solid form (e.g., powder or tablet) or embedded within a solid matrix that is subsequently mixed with the substrates to initiate the enzymatic perhydrolysis reaction. In a further aspect, the perhydrolase catalyst may be contained within a dissolvable or porous pouch that may be added to the aqueous substrate matrix to initiate enzymatic perhydrolysis. In yet a further aspect, the perhydrolase catalyst may comprise the contents contained within a separate compartment of a dissolvable or porous pouch that has at least one additional compartment for the containment contents comprising the ester substrate and/or source of peroxide. In an additional further aspect, a powder comprising the enzyme catalyst is suspended in the substrate (e.g., triacetin), and at time of use is mixed with a source of peroxygen in water.

Method for Determining the Concentration of Peroxycarboxylic acid and Hydrogen Peroxide.

A variety of analytical methods can be used in the present method to analyze the reactants and products including, but not limited to, titration, high performance liquid chromatography (HPLC), gas chromatography (GC), mass spectrometry (MS), capillary electrophoresis (CE), the analytical procedure described by U. Karst et al. (*Anal. Chem.*, 69(17): 3623-3627 (1997)), and the 2,2'-azino-bis(3-ethylbenzothiazoline)-6-sulfonate (ABTS) assay (S. Minning, et al., *Analytica Chimica Acta* 378:293-298 (1999) and WO 2004/058961 A1) as described in U.S. Patent Application Publication No. 2008-0176783.

Determination of Minimum Biocidal Concentration of Peroxycarboxylic acids

The method described by J. Gabrielson et al. (*J. Microbiol. Methods* 50: 63-73 (2002)) can be employed for determination of the Minimum Biocidal Concentration (MBC) of peroxycarboxylic acids, or of hydrogen peroxide and enzyme substrates. The assay method is based on XTT reduction inhibition, where XTT (2,3-bis[2-methoxy-4-nitro-5-sulphophenyl]-5-[(phenylamino)carbonyl]-2H-tetrazolium, inner salt, monosodium salt) is a redox dye that indicates microbial respiratory activity by a change in optical density (OD) measured at 490 nm or 450 nm. However, there are a variety of other methods available for testing the activity of disinfectants and antiseptics including, but not limited to, viable plate counts, direct microscopic counts, dry weight, turbidity measurements, absorbance, and bioluminescence (see, for example Brock, Semour S., *Disinfection, Sterilization, and Preservation*, 5th edition, Lippincott Williams & Wilkins, Philadelphia, Pa., USA; 2001).

Uses of Enzymatically Prepared Peroxycarboxylic acid Compositions

The enzyme catalyst-generated peroxycarboxylic acid produced according to the present method can be used in a variety of hard surface/inanimate object applications for reduction of concentrations of biological contaminants, such as decontamination of medical instruments (e.g., endoscopes), textiles (such as garments and carpets), food preparation surfaces, food storage and food-packaging equipment, materials used for the packaging of food products, chicken hatcheries and grow-out facilities, animal enclosures, and spent process waters that have microbial and/or virucidal activity. The enzyme-generated peroxycarboxylic acids may be used in formulations designed to inactivate prions (e.g.,

certain proteases) to additionally provide biocidal activity (see U.S. Pat. No. 7,550,420 to DiCosimo et al.).

In one aspect, the peroxycarboxylic acid composition is useful as a disinfecting agent for non-autoclavable medical instruments and food packaging equipment. As the peroxycarboxylic acid-containing formulation may be prepared using GRAS or food-grade components (enzyme, enzyme substrate, hydrogen peroxide, and buffer), the enzyme-generated peroxycarboxylic acid may also be used for decontamination of animal carcasses, meat, fruits and vegetables, or for decontamination of prepared foods. The enzyme-generated peroxycarboxylic acid may be incorporated into a product whose final form is a powder, liquid, gel, film, solid or aerosol. The enzyme-generated peroxycarboxylic acid may be diluted to a concentration that still provides an efficacious decontamination.

The compositions comprising an efficacious concentration of peroxycarboxylic acid can be used to disinfect surfaces and/or objects contaminated (or suspected of being contaminated) with biological contaminants, such as pathogenic microbial contaminants, by contacting the surface or object with the products produced by the present processes. As used herein, "contacting" refers to placing a disinfecting composition comprising an effective concentration of peroxycarboxylic acid in contact with the surface or inanimate object suspected of contamination with a biological contaminant for a period of time sufficient to clean and disinfect. Contacting includes spraying, treating, immersing, flushing, pouring on or in, mixing, combining, painting, coating, applying, affixing to and otherwise communicating a peroxycarboxylic acid solution or composition comprising an efficacious concentration of peroxycarboxylic acid, or a solution or composition that forms an efficacious concentration of peroxycarboxylic acid, with the surface or inanimate object suspected of being contaminated with a concentration of a biological contaminant. The disinfectant compositions may be combined with a cleaning composition to provide both cleaning and disinfection. Alternatively, a cleaning agent (e.g., a surfactant or detergent) may be incorporated into the formulation to provide both cleaning and disinfection in a single composition.

The compositions comprising an efficacious concentration of peroxycarboxylic acid can also contain at least one additional antimicrobial agent, combinations of prion-degrading proteases, a virucide, a sporicide, or a biocide. Combinations of these agents with the peroxycarboxylic acid produced by the claimed processes can provide for increased and/or synergistic effects when used to clean and disinfect surfaces and/or objects contaminated (or suspected of being contaminated) with biological contaminants. Suitable antimicrobial agents include carboxylic esters (e.g., p-hydroxy alkyl benzoates and alkyl cinnamates), sulfonic acids (e.g., dodecylbenzene sulfonic acid), iodo-compounds or active halogen compounds (e.g., elemental halogens, halogen oxides (e.g., NaOCl, HOCl, HOBr, ClO₂), iodine, interhalides (e.g., iodine monochloride, iodine dichloride, iodine trichloride, iodine tetrachloride, bromine chloride, iodine monobromide, or iodine dibromide), polyhalides, hypochlorite salts, hypochlorous acid, hypobromite salts, hypobromous acid, chloro- and bromo-hydantoin, chlorine dioxide, and sodium chlorite), organic peroxides including benzoyl peroxide, alkyl benzoyl peroxides, ozone, singlet oxygen generators, and mixtures thereof, phenolic derivatives (e.g., o-phenyl phenol, o-benzyl-p-chlorophenol, tert-amyl phenol and C₁-C₆ alkyl hydroxy benzoates), quaternary ammonium compounds (e.g., alkyldimethylbenzyl ammonium chloride, dialkyldimethyl ammonium chloride and mixtures thereof), and mixtures of such antimicrobial agents, in an amount

sufficient to provide the desired degree of microbial protection. Effective amounts of antimicrobial agents include about 0.001 wt % to about 60 wt % antimicrobial agent, about 0.01 wt % to about 15 wt % antimicrobial agent, or about 0.08 wt % to about 2.5 wt % antimicrobial agent.

In one aspect, the peroxycarboxylic acids formed by the process can be used to reduce the concentration of viable biological contaminants (such as a microbial population) when applied on and/or at a locus. As used herein, a "locus" comprises part or all of a target surface suitable for disinfecting or bleaching. Target surfaces include all surfaces that can potentially be contaminated with biological contaminants. Non-limiting examples include equipment surfaces found in the food or beverage industry (such as tanks, conveyors, floors, drains, coolers, freezers, equipment surfaces, walls, valves, belts, pipes, drains, joints, crevasses, combinations thereof, and the like); building surfaces (such as walls, floors and windows); non-food-industry related pipes and drains, including water treatment facilities, pools and spas, and fermentation tanks; hospital or veterinary surfaces (such as walls, floors, beds, equipment (such as endoscopes), clothing worn in hospital/veterinary or other healthcare settings, including clothing, scrubs, shoes, and other hospital or veterinary surfaces); restaurant surfaces; bathroom surfaces; toilets; clothes and shoes; surfaces of barns or stables for livestock, such as poultry, cattle, dairy cows, goats, horses and pigs; hatcheries for poultry or for shrimp; and pharmaceutical or biopharmaceutical surfaces (e.g., pharmaceutical or biopharmaceutical manufacturing equipment, pharmaceutical or biopharmaceutical ingredients, pharmaceutical or biopharmaceutical excipients). Additional hard surfaces include food products, such as beef, poultry, pork, vegetables, fruits, seafood, combinations thereof, and the like. The locus can also include water absorbent materials such as infected linens or other textiles. The locus also includes harvested plants or plant products including seeds, corms, tubers, fruit, and vegetables, growing plants, and especially crop growing plants, including cereals, leaf vegetables and salad crops, root vegetables, legumes, berried fruits, citrus fruits and hard fruits.

Non-limiting examples of hard surface materials are metals (e.g., steel, stainless steel, chrome, titanium, iron, copper, brass, aluminum, and alloys thereof), minerals (e.g., concrete), polymers and plastics (e.g., polyolefins, such as polyethylene, polypropylene, polystyrene, poly(meth)acrylate, polyacrylonitrile, polybutadiene, poly(acrylonitrile, butadiene, styrene), poly(acrylonitrile, butadiene), acrylonitrile butadiene; polyesters such as polyethylene terephthalate; and polyamides such as nylon). Additional surfaces include brick, tile, ceramic, porcelain, wood, wood pulp, paper, vinyl, linoleum, and carpet.

The peroxycarboxylic acids formed by the present process may be used to provide a benefit to an article of clothing or a textile including, but not limited to disinfecting, sanitizing, bleaching, destaining, and deodorizing. The peroxycarboxylic acids formed by the present process may be used in any number of laundry care products including, but not limited to textile pre-wash treatments, laundry detergents, laundry detergents or additives, stain removers, bleaching compositions, deodorizing compositions, and rinsing agents, to name a few.

The peroxycarboxylic acids formed by the present process can be used in one or more steps of the wood pulp or paper pulp bleaching/delignification process, particularly where peracetic acid is used (for example, see EP1040222 B1 and U.S. Pat. No. 5,552,018 to Devenyns, J.)

Recombinant Microbial Expression

The genes and gene products of the instant sequences may be produced in heterologous host cells, particularly in the cells of microbial hosts. Preferred heterologous host cells for expression of the instant genes and nucleic acid molecules are microbial hosts that can be found within the fungal or bacterial families and which grow over a wide range of temperature, pH values, and solvent tolerances. For example, it is contemplated that any of bacteria, yeast, and filamentous fungi may suitably host the expression of the present nucleic acid molecules. The perhydrolase may be expressed intracellularly, extracellularly, or a combination of both intracellularly and extracellularly, where extracellular expression renders recovery of the desired protein from a fermentation product more facile than methods for recovery of protein produced by intracellular expression. Transcription, translation and the protein biosynthetic apparatus remain invariant relative to the cellular feedstock used to generate cellular biomass; functional genes will be expressed regardless. Examples of host strains include, but are not limited to, bacterial, fungal or yeast species such as *Aspergillus*, *Trichoderma*, *Saccharomyces*, *Pichia*, *Phaffia*, *Kluyveromyces*, *Candida*, *Hansenula*, *Yarrowia*, *Salmonella*, *Bacillus*, *Acinetobacter*, *Zymomonas*, *Agrobacterium*, *Erythrobacter*, *Chlorobium*, *Chromatium*, *Flavobacterium*, *Cytophaga*, *Rhodobacter*, *Rhodococcus*, *Streptomyces*, *Brevibacterium*, *Corynebacteria*, *Mycobacterium*, *Deinococcus*, *Escherichia*, *Erwinia*, *Pantoea*, *Pseudomonas*, *Sphingomonas*, *Methylobacter*, *Methylococcus*, *Methylosinus*, *Methylomicrobium*, *Methylocystis*, *Alcaligenes*, *Synechocystis*, *Synechococcus*, *Anabaena*, *Thiobacillus*, *Methanobacterium*, *Klebsiella*, and *Myxococcus*. In one embodiment, bacterial host strains include *Escherichia*, *Bacillus*, and *Pseudomonas*. In a preferred embodiment, the bacterial host cell is *Escherichia coli*.

Industrial Production

A variety of culture methodologies may be applied to produce the perhydrolase catalyst. Large-scale production of a specific gene product over expressed from a recombinant microbial host may be produced by batch, fed-batch or continuous culture methodologies. Batch and fed-batch culturing methods are common and well known in the art and examples may be found in Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition, Sinauer Associates, Inc., Sunderland, Mass. (1989) and Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36:227 (1992).

In one embodiment, commercial production of the desired perhydrolase catalyst is accomplished with a continuous culture. Continuous cultures are an open system where a defined culture media is added continuously to a bioreactor and an equal amount of conditioned media is removed simultaneously for processing. Continuous cultures generally maintain the cells at a constant high liquid phase density where cells are primarily in log phase growth. Alternatively, continuous culture may be practiced with immobilized cells where carbon and nutrients are continuously added and valuable products, by-products or waste products are continuously removed from the cell mass. Cell immobilization may be performed using a wide range of solid supports composed of natural and/or synthetic materials.

Recovery of the desired perhydrolase catalysts from a batch or fed-batch fermentation, or continuous culture may be accomplished by any of the methods that are known to those skilled in the art. For example, when the enzyme catalyst is produced intracellularly, the cell paste is separated from the culture medium by centrifugation or membrane filtration, optionally washed with water or an aqueous buffer at a

desired pH, then a suspension of the cell paste in an aqueous buffer at a desired pH is homogenized to produce a cell extract containing the desired enzyme catalyst. The cell extract may optionally be filtered through an appropriate filter aid such as celite or silica to remove cell debris prior to a heat-treatment step to precipitate undesired protein from the enzyme catalyst solution. The solution containing the desired enzyme catalyst may then be separated from the precipitated cell debris and protein produced during the heat-treatment step by membrane filtration or centrifugation, and the resulting partially-purified enzyme catalyst solution concentrated by additional membrane filtration, then optionally mixed with an appropriate excipient (for example, maltodextrin, trehalose, sucrose, lactose, sorbitol, mannitol, phosphate buffer, citrate buffer, or mixtures thereof) and spray-dried to produce a solid powder comprising the desired enzyme catalyst. Alternatively, the resulting partially-purified enzyme catalyst solution prepared as described above may be optionally concentrated by additional membrane filtration, and the partially-purified enzyme catalyst solution or resulting enzyme concentrate is then optionally mixed with one or more stabilizing agents (e.g., glycerol, sorbitol, propylene glycol, 1,3-propanediol, polyols, polymeric polyols, polyvinylalcohol), one or more salts (e.g., sodium chloride, sodium sulfate, potassium chloride, potassium sulfate, or mixtures thereof), and one or more biocides, and maintained as an aqueous solution until used.

When an amount, concentration, or other value or parameter is given either as a range, preferred range, or a list of upper preferable values and lower preferable values, this is to be understood as specifically disclosing all ranges formed from any pair of any upper range limit or preferred value and any lower range limit or preferred value, regardless of whether ranges are separately disclosed. Where a range of numerical values is recited herein, unless otherwise stated, the range is intended to include the endpoints thereof, and all integers and fractions within the range. It is not intended that the scope be limited to the specific values recited when defining a range.

General Methods

The following examples are provided to demonstrate preferred embodiments. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the methods disclosed herein, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the presently disclosed methods.

All reagents and materials were obtained from DIFCO Laboratories (Detroit, Mich.), GIBCO/BRL (Gaithersburg, Md.), TCI America (Portland, Oreg.), Roche Diagnostics Corporation (Indianapolis, Ind.) or Sigma-Aldrich Chemical Company (St. Louis, Mo.), unless otherwise specified.

The following abbreviations in the specification correspond to units of measure, techniques, properties, or compounds as follows: "sec" or "s" means second(s), "min" means minute(s), "h" or "hr" means hour(s), "4" means microliter(s), "mL" means milliliter(s), "L" means liter(s), "mM" means millimolar, "M" means molar, "mmol" means millimole(s), "ppm" means part(s) per million, "wt" means weight, "wt %" means weight percent, "g" means gram(s), "µg" means microgram(s), "ng" means nanogram(s), "g" means gravity, "HPLC" means high performance liquid chromatography, "dd H₂O" means distilled and deionized water,

27

“dew” means dry cell weight, “ATCC” or ATCC® means the American Type Culture Collection (Manassas, Va.), “U” means unit(s) of perhydrolase activity, “rpm” means revolution(s) per minute, “EDTA” means ethylenediaminetetraacetic acid, “IPTG” means isopropyl β-D-1-thiogalactopyranoside, “DTT” means dithiothreitol, “BCA” means bicinchoninic acid.

EXAMPLE 1

Construction of a Random Mutagenesis Library of
Thermotoga maritima Acetyl Xylan Esterase C277S
Variant

The coding sequence of a *Thermotoga maritima* acetyl xylan esterase (GENBANK® accession # NP_227893.1) was synthesized using codons optimized for expression in *E. coli* (DNA 2.0, Menlo Park, Calif.), and cloned into pUC19 between PstI and XbaI to create the plasmid known as pSW202 (U.S. Patent Application Publication No. 2008-0176299). The codon-optimized sequence is provided as SEQ ID NO: 1 encoding the wild-type *Thermotoga maritima* acetyl xylan esterase provided as SEQ ID NO: 2.

A codon change was made in the gene using primer pairs identified as SEQ ID NO: 3 and SEQ ID NO: 4, and the QUIKCHANGE® site-directed mutagenesis kit (Stratagene, La Jolla, Calif.), according to the manufacturer's instructions, resulting in the amino acid change C277S (SEQ ID NO: 5), to create the plasmid known as pSW202/C277S (SEQ ID NO: 6). Plasmid pSW202/C277S served as a template for error-prone PCR using primers identified as SEQ ID NO: 7 and SEQ ID NO: 8, and the GENEMORPH® II random mutagenesis kit (Stratagene), according to the manufacturer's recommendations. The resulting PCR product was digested with PstI and XbaI and ligated with pUC19 also digested with PstI and XbaI. *E. coli* KLP18 (see Published U.S. Patent Application No. 2008-0176299; herein incorporated by reference in its entirety) was transformed with the ligation mixture and plated onto LB plates supplemented with 0.1 mg ampicillin/mL. Nucleotide sequencing of a random sample indicated a mutation frequency of 2-8 changes per PCR product.

EXAMPLE 2

Screening of *Thermotoga maritima* Error-prone PCR
Library for Increased Enzyme Activity

Colonies were picked (automated) and placed into 96-well “master plates” containing 0.1 mL LB media supplemented with 0.1 mg ampicillin/mL and grown 16-18 h at 37° C. and 80% humidity. From each well of the master plates, 0.003 mL of culture was transferred to 96-well “induction plates” containing 0.3 mL LB media supplemented with 0.1 mg ampicillin/mL and 0.5 mM IPTG, which were incubated for 16-18 h with shaking at 37° C. and 80% humidity. Separately, 0.1 mL of 50% glycerol was added to each well of the master plates, which were stored at -80° C. as stocks. From each well of the induction plates, 0.01 mL of culture was transferred to 96-well “lysis plates” containing 0.09 mL of 56 mg/mL CEL-LYTIC™ Express (Sigma Aldrich, St. Louis, Mo.), which were incubated for 30 minutes at 30° C. From each well of the lysis plates, 0.01 mL of material was transferred to 96-well “assay plates” containing 0.045 mL “assay solution part 1” (50 mM triacetin, 50 mM potassium phosphate buffer, pH 7.0). To each well of the assay plates was then added 0.045 mL of “assay solution part 2” (50 mM hydrogen peroxide).

28

Plates were gently mixed and incubated for 10 minutes at 30° C. To each well of the assay plate was added 0.1 mL of “stop buffer” (100 mM o-phenylenediamine and 500 mM sodium dihydrogen phosphate, pH 2.0). The plates were incubated for 30 minutes at 30° C., after which absorbance at 458 nm was read. *T. maritima* WT (codon optimized gene encoding the wild type enzyme) and *T. maritima* C277S were incorporated into each plate as controls. Screening approximately 7000 colonies resulted in the identification of numerous “hits” demonstrating activity significantly greater than *T. maritima* C277S. Nucleotide sequencing was used to determine the amino acid changes in the perhydrolase enzyme of these hits (Table 1).

TABLE 1

Variant CE-7 carbohydrate esterases having perhydrolytic activity identified by preliminary screening.				
Variant ID	# of amino acid changes	Amino acid residue changes relative to the <i>Thermotoga maritima</i> wild-type sequence (SEQ ID NO: 2)	Nucleic Acid SEQ ID NO.	Amino Acid SEQ ID NO.
843H9	7	L8R/L125Q/Q176L/V183D/F247I/C277S/P292L	9	10
843B12	2	A206E/C277S	11	12
843D9	3	S35R/H50R/C277S	13	14
843F12	3	K77E/A266E/C277S	15	16
843C12	6	F27Y/I149V/A266V/C277S/I295T/N302S	17	18
843H7	3	G119S/L207P/C277S	19	20
842H3	2	L195Q/C277S	21	22
841D3	3	K52E/C277S/Y298F	23	24
841D1	4	L125Q/T249S/C277S/A311V	25	26
841D4	4	F38S/V197I/N275T/C277S	27	28
841F11	4	K77I/K126N/C277S/K293R	29	30
841A7	2	Y110F/C277S	31	32
control	1	C277S	—	5

EXAMPLE 3

Production of Variant *Thermotoga maritima*
Perhydrolases

Variant strains identified in the screen (KLP18/pSW202/L8R/L125Q/Q176L/V183D/F247I/C277S/P292L; KLP18/pSW202/A206E/C277S; KLP18/pSW202/S35R/H50R/C277S; KLP18/pSW202/K77E/A266E/C277S; KLP18/pSW202/F27Y/I149V/A266V/C277S/I295T/N302S; KLP18/pSW202/G119S/L207P/C277S; KLP18/pSW202/L195Q/C277S; KLP18/pSW202/K52E/C277S/Y298F; KLP18/pSW202/L125Q/T249S/C277S/A311V; KLP18/pSW202/F38S/V197I/N275T/C277S; KLP18/pSW202/K77I/K126N/C277S/K293R; and KLP18/pSW202/Y110F/C277S) were grown in LB media at 37° C. with shaking up to OD_{600nm}=0.4-0.5, at which time IPTG was added to a final concentration of 1 mM, and incubation continued for 2-3 h. Cells were harvested by centrifugation and SDS-PAGE was performed to confirm expression of the perhydrolase enzyme at 20-40% of total soluble protein.

EXAMPLE 4

Preparation of Heat-Treated Cell Extracts Containing
Semi-Purified Variant *Thermotoga maritima*
Perhydrolases

Cell cultures (prepared as described in Example 3) were harvested by centrifugation at 5,000×g for 15 minutes then

resuspended (20% w/v) in 50 mM phosphate buffer pH 7.0 supplemented with 1.0 mM DTT. Resuspended cells were passed through a French pressure cell twice to ensure >95% cell lysis. Lysed cells were centrifuged for 30 minutes at 12,000×g, and the supernatant was heated at 75° C. for 20 minutes, followed by quenching in an ice bath for 2 minutes. Precipitated protein was removed by centrifugation for 10 minutes at 11,000×g. The resulting heat-treated extract supernatants were analyzed for mg total soluble protein/mL using BCA assay, and stored frozen at -80° C. Analysis of a first set of heat-treated extract supernatants produced by this method using PAGE indicated that the perhydrolase variant comprised approximately 85-90% of the total soluble protein in the heat-treated extract supernatant. Analysis of a second set of heat-treated extract supernatants produced by this method using PAGE indicated that the perhydrolase variant comprised approximately 65-70% of the total soluble protein in the heat-treated extract supernatant. In the following Examples 6 and 7, the individual Tables contain data obtained using a single set of heat-treated extract supernatants that contained the same concentration of perhydrolase variant as a percentage of total soluble protein.

EXAMPLE 5

Comparison of Perhydrolase Variants vs.
Thermotoga maritima C277S Variant Perhydrolase
Using 25 mM Triacetin and 25 mM H₂O₂ in
Phosphate Buffer (pH 7.2) at 25° C.

Reactions (10 mL total volume) were run at 25° C. in phosphate buffer (50 mM, pH 7.2) containing triacetin (25 mM), hydrogen peroxide (25 mM) and 2.0 µg/mL of heat-treated extract total soluble protein containing an error-prone PCR (epPCR)-generated variant perhydrolase prepared as described in Example 4. Reactions were stirred for only the first 45 seconds of reaction to initially mix the reactants and enzyme.

A comparative reaction was also run under identical conditions to that described immediately above using 2.0 µg/mL of heat-treated extract total soluble protein isolated from *E. coli* KLP18/pSW228/C277S (expressing *Thermotoga maritima* C277S perhydrolase variant), where the heat-treated extract supernatant was prepared according to the procedure of Example 4, and where this heat-treated extract supernatant contained the same concentration of *Thermotoga maritima* C277S perhydrolase variant as a percentage of total soluble protein as was present in the heat-treated extract total soluble protein containing an error-prone PCR (epPCR)-generated variant perhydrolase.

Reaction samples were analyzed for the amount of peracetic acid (PAA) produced at 5 minutes and 20 minutes after combining the reaction components using a modification of the method described by Karst et al., supra. A sample (0.200

mL) of the reaction mixture was removed at a predetermined time and immediately mixed with 0.200 mL of 50 mM phosphoric acid in water to terminate the reaction by adjusting the pH of the diluted sample to less than pH 4. The resulting solution was filtered using an ULTRAFREE® MC-filter unit (30,000 Normal Molecular Weight Limit (NMWL), Millipore Corp., Billerica, Mass.; cat #UFC3LKT 00) by centrifugation for 2 min at 12,000 rpm. An aliquot (0.100 mL) of the resulting filtrate was transferred to a 1.5-mL screw cap HPLC vial (Agilent Technologies, Palo Alto, Calif.; #5182-0715) containing 0.300 mL of deionized water, then 0.100 mL of 20 mM MTS (methyl-p-tolyl sulfide) in acetonitrile was added, the vial capped, and the contents briefly mixed prior to a 10 min incubation at ca. 25° C. in the absence of light. To the vial was then added 0.400 mL of acetonitrile and 0.100 mL of a solution of triphenylphosphine (TPP, 120 mM) in acetonitrile, the vial re-capped, and the resulting solution mixed and incubated at ca. 25° C. for 30 min in the absence of light. To the vial was then added 0.100 mL of 2.5 mM N,N-diethyl-m-toluamide (DEET; HPLC external standard) and the resulting solution analyzed by HPLC for MTSSO (methyl-p-tolyl sulfoxide), the stoichiometric oxidation product produced by reaction of MTS with peracetic acid. A control reaction was run in the absence of added extract protein or triacetin to determine the rate of oxidation of MTS in the assay mixture by hydrogen peroxide, for correction of the rate of peracetic acid production for background MTS oxidation. HPLC method: Supelco Discovery C8 column (10-cm×4.0-mm, 5 µm) (catalog #569422-U) with Supelco Supelguard Discovery C8 precolumn (Sigma-Aldrich; catalog #59590-U); 10 microliter injection volume; gradient method with CH₃CN (Sigma-Aldrich; catalog #270717) and deionized water at 1.0 mL/min and ambient temperature (Table 4).

TABLE 4

HPLC Gradient for analysis of peracetic acid.	
Time (min:sec)	(% CH ₃ CN)
0:00	40
3:00	40
3:10	100
4:00	100
4:10	40
7:00 (stop)	40

Several variants were identified that produced a higher concentration of PAA at one or both time points when compared to the *T. maritima* C277S variant perhydrolase (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.). As shown in Tables 2, 3 and 4, variants 843H9, 843F12, 841D3, 841D4, and 841A7 were able to produce more peracetic acid when compared to the *T. maritima* C277S perhydrolase under the specified conditions.

TABLE 2

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 843H9 and 843F12 vs. <i>T. maritima</i> C277S in phosphate buffer (50 mM, pH 7.2) at 25° C., 2.0 µg/mL total soluble protein.						
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	PAA produced at 5 min. (ppm)	PAA produced at 20 min. (ppm)
	no perhydrolase	25	25	0	0	1.1
Control	C277S	25	25	2.0	184	367
Control	C277S	25	25	2.0	179	383

TABLE 2-continued

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 843H9 and 843F12 vs. <i>T. maritima</i> C277S in phosphate buffer (50 mM, pH 7.2) at 25° C., 2.0 µg/mL total soluble protein.						
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	PAA produced at 5 min. (ppm)	PAA produced at 20 min. (ppm)
843H9	L8R/L125Q/Q176L/V183D/F247I/C277S/P292L	25	25	2.0	320	512
843H9	L8R/L125Q/Q176L/V183D/F247I/C277S/P292L	25	25	2.0	344	604
843F12	K77E/A266E/C277S	25	25	2.0	223	239
843F12	K77E/A266E/C277S	25	25	2.0	227	383

TABLE 3

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 841D3 and 841D4 vs. <i>T. maritima</i> C277S in phosphate buffer (50 mM, pH 7.2) at 25° C., 2.0 µg/mL total soluble protein.						
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	PAA produced at 5 min. (ppm)	PAA produced at 20 min. (ppm)
Control	no perhydrolase	25	25	0	1.7	0.5
	C277S	25	25	2.0	163	399
Control	C277S	25	25	2.0	174	373
841D3	K52E/277S/Y238F	25	25	2.0	163	421
841D3	K52E/277S/Y238F	25	25	2.0	194	408
841D4	F38S/V197I/N275T/C277S	25	25	2.0	215	430
841D4	F38S/V197I/N275T/C277S	25	25	2.0	232	439

TABLE 4

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variant 841A7 vs. <i>T. maritima</i> C277S in phosphate buffer (50 mM, pH 7.2) at 25° C., 2.0 µg/mL total soluble protein.						
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ (mM)	Total soluble protein (µg/mL)	PAA produced at 5 min. (ppm)	PAA produced at 20 min. (ppm)
Control	no perhydrolase	25	25	0	1.7	0.5
	C277S	25	25	2.0	227	491
Control	C277S	25	25	2.0	222	496
841A7	Y110F/C277S	25	25	2.0	290	545
841A7	Y110F/C277S	25	25	2.0	299	561

EXAMPLE 6

Comparison of *Thermotoga maritima* Perhydrolase
Variants vs. *Thermotoga maritima* C277S
Perhydrolase Under Simulated Laundry Care
Application Conditions

Five error-prone PCR (epPCR)-generated variant perhydrolases were assayed for peracetic acid (PAA) production under laundry application conditions (2 mM triacetin (TA), 10 mM H₂O₂ from sodium percarbonate in hard water (HW, 400 ppm Ca ion), 2.0 µg/mL of heat-treated extract total soluble protein containing an error-prone PCR (epPCR)-generated variant perhydrolase prepared as described in Example 4) with and without added liquid laundry detergent (comprising 15-30% anionic and non-ionic surfactants, 5-15% soap, and <5% polycarboxylates, perfume, phosphonates, optical

brighteners) and compared to PAA production using the same concentration (see Example 5) of C277S variant perhydrolase (Published U.S. Patent Application No. 2010-0087529 to DiCosimo et al.). Under the reaction conditions, several epPCR-generated variants showed a measurable improvement in specific activity for peracetic acid production when compared to the *T. maritima* C277S variant under these reaction conditions. These reactions were performed using 0, 4 and 6 g/L of liquid laundry detergent. The results of the various comparisons under simulated laundry care conditions are provided in Tables 5, 6, 7, 8, 9, 10, and 11. Under these reaction conditions, variants 842H3 (L195/C277S), 841A7 (Y110F/C277S), 843H9 (L8R/L125Q/Q176L/V183D/F247I/C277S/P292L), 843F 12 (K77E/A266E/C277S), and 843C12 (F27Y/I149V/A266V/C277S/1295T/N₃₀₂S) show a measurable improvement specific activity for peracetic acid production when compared to the C277S perhydrolase from which the variants were prepared.

TABLE 5

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 842H3 and 841A7 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 2.0 µg/mL total soluble protein and 0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (µg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
	no perhydrolase	2	10	0	0	10	19	28
Control	C277S	2	10	2.0	0	10	40	65
Control	C277S	2	10	2.0	0	10	44	66
Control	C277S	2	10	2.0	0	10	44	86
842H3	L195Q/C277S	2	10	2.0	0	10	51	99
842H3	L195Q/C277S	2	10	2.0	0	10	51	102
842H3	L195Q/C277S	2	10	2.0	0	10	50	101
841A7	Y110F/C277S	2	10	2.0	0	10	49	99
841A7	Y110F/C277S	2	10	2.0	0	10	51	101
841A7	Y110F/C277S	2	10	2.0	0	10	50	102

¹= Hydrogen peroxide generated from sodium percarbonate.

TABLE 6

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 842H3 and 841A7 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 6.0 µg/mL total soluble protein and 4.0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (µg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
	no perhydrolase	2	10	0	4.0	10	6.3	9.3
Control	C277S	2	10	6.0	4.0	10	42	58
Control	C277S	2	10	6.0	4.0	10	41	55
842H3	L195Q/C277S	2	10	6.0	4.0	10	47	66
842H3	L195Q/C277S	2	10	6.0	4.0	10	47	62
841A7	Y110F/C277S	2	10	6.0	4.0	10	46	63
841A7	Y110F/C277S	2	10	6.0	4.0	10	46	64

¹= Hydrogen peroxide generated from sodium percarbonate

TABLE 7

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 842H3 and 841A7 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 6.0 µg/mL total soluble protein and 6.0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (µg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
	no perhydrolase	2	10	0	6.0	10	7.6	6.4
Control	C277S	2	10	6.0	6.0	10	28	31
Control	C277S	2	10	6.0	6.0	10	35	32
842H3	L195Q/C277S	2	10	6.0	6.0	10	35	36
842H3	L195Q/C277S	2	10	6.0	6.0	10	36	35
841A7	Y110F/C277S	2	10	6.0	6.0	10	36	35
841A7	Y110F/C277S	2	10	6.0	6.0	10	33	32

¹= Hydrogen peroxide generated from sodium percarbonate

TABLE 8

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 843H9 and 843F12 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 2.0 µg/mL total soluble protein and 0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (µg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
Control	no perhydrolase	2	10	0	0	10	16	41
	C277S	2	10	2.0	0	10	26	73
Control	C277S	2	10	2.0	0	10	24	77
843H9	L8R/L125Q/Q176L/V183D/ F247I/C277S/P292L	2	10	2.0	0	10	58	104
843H9	L8R/L125Q/Q176L/V183D/ F247I/C277S/P292L	2	10	2.0	0	10	65	89
843F12	K77E/A266E/C277S	2	10	2.0	0	10	43	95
843F12	K77E/A266E/C277S	2	10	2.0	0	10	38	84

¹= Hydrogen peroxide generated from sodium percarbonate

TABLE 9

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variants 843H9 and 843F12 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 6.0 µg/mL total soluble protein and 4.0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (µg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
Control	no perhydrolase	2	10	0	4.0	10	3.6	4.1
	C277S	2	10	6.0	4.0	10	27	41
Control	C277S	2	10	6.0	4.0	10	31	45
843H9	L8R/L125Q/Q176L/V183D/ F247I/C277S/P292L	2	10	6.0	4.0	10	40	54
843H9	L8R/L125Q/Q176L/V183D/ F247I/C277S/P292L	2	10	6.0	4.0	10	40	57
843F12	K77E/A266E/C277S	2	10	6.0	4.0	10	36	44
843F12	K77E/A266E/C277S	2	10	6.0	4.0	10	33	43

¹= Hydrogen peroxide generated from sodium percarbonate

TABLE 10

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variant 843C12 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 2.0 µg/mL total soluble protein and 0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (µg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
Control	no perhydrolase	2	10	0	0	10	20	45
	C277S	2	10	2.0	0	10	30	79
Control	C277S	2	10	2.0	0	10	34	81
843C12	F27Y/I149V/A266V/C277S/ I295T/N302S	2	10	2.0	0	10	40	91
843C12	F27Y/I149V/A266V/C277S/ I295T/N302S	2	10	2.0	0	10	36	86

¹= Hydrogen peroxide generated from sodium percarbonate

TABLE 11

Peracetic acid (PAA) production at 5 minutes and 20 minutes from variant 843C12 vs. <i>T. maritima</i> C277S in hard water (HW) and carbonate buffer (pH 10) at 20° C., 6.0 mg/mL total soluble protein and 4.0 mg/mL liquid laundry detergent.								
Variant ID	Samples	Triacetin (mM)	H ₂ O ₂ ¹ (mM)	Total soluble protein (μg/mL)	Liquid detergent (mg/mL)	Initial pH	PAA @ 5 min. (ppm)	PAA @ 20 min. (ppm)
	no perhydrolase	2	10	0	0	10	4.7	7.9
Control	C277S	2	10	6.0	4.0	10	26	43
Control	C277S	2	10	6.0	4.0	10	27	41
843C12	F27Y/I149V/A266V/C277S/I295T/N302S	2	10	6.0	4.0	10	33	43
843C12	F27Y/I149V/A266V/C277S/I295T/N302S	2	10	6.0	4.0	10	27	42

¹ = Hydrogen peroxide generated from sodium percarbonate

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 32

<210> SEQ ID NO 1

<211> LENGTH: 978

<212> TYPE: DNA

<213> ORGANISM: Thermotoga maritima

<400> SEQUENCE: 1

```

atggccttct tcgatttacc actcgaagaa ctgaagaaat atcgtccaga gcggtacgaa      60
gagaaagact tcgatgagtt ctgggaagag acaactcgcag agagcgaaaa gtteccctta    120
gaccccgctc tcgagaggat ggagtctcac ctcaaaacag tcgaagcgta cgatgtcacc    180
ttctccggat acaggggaca gaggatcaaa ggggtggctcc ttgttccaaa actggaagaa    240
gaaaaacttc cctgcgttgt gcagtacata ggatacaacg gtggaagagg attccctcac    300
gactggctgt tctggccttc tatgggttac atatgtttcg tcatggatac tcgaggtcag    360
ggaagcggct ggctgaaagg agacacaccg gattaccctg aggggtcccg tgaccctcag    420
tatecaggat tcatgacaag aggaatactg gatcccagaa cttactacta cagacgagtc    480
ttcacggacg ctgtcagagc cgttgaagct gctgcttctt ttcctcaggt agatcaagaa    540
agaatcgtga tagctggagg cagtcagggt ggcggaatag cccttgcggt gagcgctctc    600
tcaaagaaag caaaggctct tctgtgcgat gtgccgtttc tgtgtcactt cagaagagca    660
gtacagcttg tggatacgca tccatacgcg gagatcacga actttctaaa gaccacaga     720
gacaaggaag aaatcgtgtt caggactctt tcctatttcg atggagttaa cttcgcagcc    780
agagcgaaga tccttcgcgt gttttctgtg ggtctcatgg acaacatttg tcctccttca    840
acgggttttc ctgcctacaa ttactacgct ggaccgaagg aaatcagaat ctatccgtac    900
aacaaccacg agggaggagg ctctttccaa gcggttgaac aggtgaaatt cttgaaaaaa    960
ctatttgaga aaggctaa                                     978

```

<210> SEQ ID NO 2

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: Thermotoga maritima

<400> SEQUENCE: 2

```

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1           5           10           15

```

-continued

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
 20 25 30
 Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
 35 40 45
 Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Cys Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 3
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 3

tcggacaaca tcacctcctt cta

23

<210> SEQ ID NO 4
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

-continued

<400> SEQUENCE: 4

tagaaggagg tgagatgttg tcc

23

<210> SEQ ID NO 5

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<400> SEQUENCE: 5

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1 5 10 15Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20 25 30Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35 40 45Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65 70 75 80Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100 105 110Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115 120 125Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130 135 140Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145 150 155 160Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165 170 175Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180 185 190Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195 200 205Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210 215 220Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225 230 235 240Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245 250 255Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260 265 270Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275 280 285Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
290 295 300Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305 310 315 320Leu Phe Glu Lys Gly
325

<210> SEQ ID NO 6

<211> LENGTH: 3674

-continued

<212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: plasmid

<400> SEQUENCE: 6

```

gacgaaaggg cctcgtgata cgcctatddd tataggttaa tgtcatgata ataatggddd    60
cttagacgtc aggtggcact ttccggggaa atgtgcgcgg aaccctatt tgtttatddd    120
tctaaataca ttcaaatatg tatccgtca tgagacaata accctgataa atgcttcaat    180
aatattgaaa aaggaagagt atgagtattc aacatttccg tgcgccctt attccctddd    240
ttgcggcatt ttgccttctc gtttttgctc acccagaaac gctggtgaaa gtaaaagatg    300
ctgaagatca gttgggtgca cgagtgggtt acatcgaaact ggatctcaac agcggtaaga    360
tccttgagag ttttcgcccc gaagaacgtt ttccaatgat gagcactddd aaagttctgc    420
tatgtggcgc ggtattatcc cgtattgacg ccgggcaaga gcaactcggg cgcgcgatac    480
actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg    540
gcatgacagt aagagaatta tgcagtgtcg ccataaccat gagtataac actgcggcca    600
acttacttct gacaacgacg ggaggaccga aggagctaac cgcttttttg cacaacatgg    660
gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacg    720
acgagcgtga caccacgatg cctgtagcaa tggcaacaac gttgcgcaaa ctattaactg    780
gcgaactact tactctagct tcccggcaac aattaataga ctggatggag gcggataaag    840
ttgcaggacc acttctgcgc tcggcccttc cggttggtg gtttattgct gataaatctg    900
gagccggtga gcgtgggtct cgcggtatca ttgcagcact ggggccagat ggtaagccct    960
cccgtatcgt agttatctac acgacgggga gtcaggcaac tatggatgaa cgaaaatagc   1020
agatcgtcga gataggtgcc tcaactgatta agcattggta actgtcagac caagtttact   1080
catatatact ttagattgat ttaaaacttc atttttaatt taaaaggatc taggtgaaga   1140
tcctttttga taatctcatg accaaaatcc cttaactgta gttttcgttc cactgagcgt   1200
cagaccccggt agaaaagatc aaaggatctt cttgagatcc tttttttctg cgcgtaatct   1260
gctgcttgca aacaaaaaaa ccaccgctac cagcgggtggg ttgtttgccg gatcaagagc   1320
taccaaactct ttttccgaag gtaactggct tcagcagagc gcagatacca aatactgtcc   1380
ttctagtgtg gccgtagtta ggccaccact tcaagaactc tgtagcaccg cctacatacc   1440
tcgctctgct aatcctgtta ccagtggctg ctgccagtgg cgataagtcg tgtcttaccg   1500
ggttggactc aagacgatag ttaccggata aggcgcagcg gtcgggctga acgggggggt   1560
cgtgcacaca gcccagcttg gagcgaacga cctacaccga actgagatac ctacagcgtg   1620
agctatgaga aagcgccacg cttccgaag ggagaaaggc ggacaggtat ccggtaaagc   1680
gcagggtcgg aacagagagag cgcacgaggg agcttccagg gggaaacgcc tggtatcttt   1740
atagtcctgt cgggtttcgc cacctctgac ttgagcgtcg atttttgtga tgcctgtcag   1800
gggggcggag cctatggaaa aacgcagca acgcggcctt ttacgggttc ctggcctttt   1860
gctggccttt tgctcacatg ttctttctcg cgttatcccc tgattctgtg gataaccgta   1920
ttaccgcctt tgagttagct gataccgctc gccgcagccg aacgaccgag cgcagcgagt   1980
cagttagcga ggaagcggaa gagcgcccaa tacgcaaacc gcctctcccc gcgcgttggc   2040
cgattcatta atgcagctgg cacgacaggt ttcccgaact gaaagcgggc agtgagcgca   2100
acgcaattaa tgtgagttag ctcaactcatt aggcacccca ggctttacac tttatgcttc   2160
cggtcgtgat gttgtgtgga attgtgagcg gataacaatt tcacacagga aacagctatg   2220

```


-continued

```

accatgatta cgccaagctt gcatgcctgc agtaaggagg aataggacat ggcgttcttc 2280
gacctgcctc tggaagaact gaagaaatac cgtccagagc gttacgaaga gaaggacttc 2340
gacgagttct gggaggaaac tctggcggag agcgaaaagt ttccgctgga cccagtgttc 2400
gagcgtatgg aatctcacct gaaaaccgtg gaggcataatg acgttacttt ttctggttac 2460
cgtggccagc gtatcaaagg ctggctgctg gttccgaaac tggaggaaga aaaactgccg 2520
tgcgtagttc agtacatcgg ttacaacggt ggccgtggct ttccgcacga ttggtgttc 2580
tggcgtctta tgggctacat ttgcttcgtc atggatactc gtggtcaggg ttccgctgg 2640
ctgaaaggcg atactccgga ttatccggag ggcccggtag acccgagta ccttggttc 2700
atgacgcgtg gtattctgga tccgcgtacc tattactatc gccgcgtttt taccgatgca 2760
gttcgtgccg tagaggccgc ggcttcttcc cctcaggttg accaggagcg tattgttacc 2820
gctggtggct cccagggtgg cggcatcgcc ctggcggtat ctgcgctgag caagaaagct 2880
aaggcactgc tgtgtgacgt cccgttctctg tgtcacttcc gtcgcgtgtg tcagctggta 2940
gatacccatc cgtacgcgga gattactaac ttctgaaaa ctcaccgcga caaagaagaa 3000
atcgttttcc gcacctgtgc ctatttcgac ggcgtaact tcgcggctcg tgcaaaaatt 3060
ccggcactgt tctctgttgg tctgatggac aacatcagcc ctcttctac cgttttcgcg 3120
gcatataact attatgcggg tccgaaagaa atccgtatct atccgtacaa caaccacgaa 3180
ggcgggtgga gctttcagcg tgttgaacaa gtgaaattcc tgaagaaact gtttgagaag 3240
ggctaatact gaggatcccc gggtaccgag ctcgaaattca ctggccgtcg ttttacaacg 3300
tcgtgactgg gaaaaccctg gcgttaccca acttaatcgc cttgcagcac atcccccttt 3360
cgccagctgg cgtaaatcgc aagaggcccg caccgatcgc ccttcccaac agttgcgcag 3420
cctgaatggc gaatggcgcc tgatgcggta ttttctcett acgcatctgt gcggtatttc 3480
acaccgcata tgggtgcactc tcagtacaat ctgctctgat gccgcatagt taagccagcc 3540
ccgacacccg ccaacacccg ctgacgcgcc ctgacgggct tgtctgctcc cggcatccgc 3600
ttacagacaa gctgtgacgg tctccgggag ctgcatgtgt cagaggtttt caccgtcatc 3660
accgaaacgc gcga 3674

```

```

<210> SEQ ID NO 7
<211> LENGTH: 49
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

```

```

<400> SEQUENCE: 7

```

```

taactgcagt aaggaggaat aggacatggc gttcttcgac ctgcctctg 49

```

```

<210> SEQ ID NO 8
<211> LENGTH: 38
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

```

```

<400> SEQUENCE: 8

```

```

tgatctagat tagcccttct caaacagttt ctttcagg 38

```

```

<210> SEQ ID NO 9
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: artificial sequence

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (978)

<400> SEQUENCE: 9

atg gcg ttc ttc gac ctg cct cgg gaa gaa ctg aag aaa tac cgt cca      48
Met Ala Phe Phe Asp Leu Pro Arg Glu Glu Leu Lys Lys Tyr Arg Pro
1          5          10          15

gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg      96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20          25          30

gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa      144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35          40          45

tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac      192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50          55          60

cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg aaa ctg gag gaa      240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65          70          75          80

gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt      288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85          90          95

ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc tac att tgc      336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100         105         110

ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg cag aaa gcc gat      384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Gln Lys Gly Asp
115         120         125

act ccg gat tat ccg gag gcc ccg gta gac ccg cag tac cct gcc ttc      432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130         135         140

atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt      480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145         150         155         160

ttt acc gat gca gtt cgt gcc gta gag gcc gcc gct tct ttc cct ctg      528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Leu
165         170         175

gtt gac cag gag cgt att gat atc gct ggt gcc tcc cag ggt gcc gcc      576
Val Asp Gln Glu Arg Ile Asp Ile Ala Gly Gly Ser Gln Gly Gly Gly
180         185         190

atc gcc ctg ccg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg      624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195         200         205

tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta      672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210         215         220

gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc      720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225         230         235         240

gac aaa gaa gaa atc gtt atc cgc acc ctg tcc tat ttc gac gcc gtt      768
Asp Lys Glu Glu Ile Val Ile Arg Thr Leu Ser Tyr Phe Asp Gly Val
245         250         255

aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg      816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260         265         270

atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat      864
Met Asp Asn Ile Ser Pro Pro Thr Val Phe Ala Ala Tyr Asn Tyr
275         280         285

```

-continued

tat gcg ggt ctg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa	912
Tyr Ala Gly Leu Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa	960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	
305 310 315 320	
ctg ttt gag aag ggc taa	978
Leu Phe Glu Lys Gly	
325	

<210> SEQ ID NO 10
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 10

Met Ala Phe Phe Asp Leu Pro Arg Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Gln Lys Gly Asp	
115 120 125	
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val	
145 150 155 160	
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Leu	
165 170 175	
Val Asp Gln Glu Arg Ile Asp Ile Ala Gly Gly Ser Gln Gly Gly Gly	
180 185 190	
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	
195 200 205	
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	
210 215 220	
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	
225 230 235 240	
Asp Lys Glu Glu Ile Val Ile Arg Thr Leu Ser Tyr Phe Asp Gly Val	
245 250 255	
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu	
260 265 270	
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	
275 280 285	
Tyr Ala Gly Leu Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	

-continued

305	310	315	320	
Leu Phe Glu Lys Gly				
325				
 <210> SEQ ID NO 11				
<211> LENGTH: 978				
<212> TYPE: DNA				
<213> ORGANISM: artificial sequence				
<220> FEATURE:				
<223> OTHER INFORMATION: synthetic construct				
<220> FEATURE:				
<221> NAME/KEY: CDS				
<222> LOCATION: (1)..(978)				
 <400> SEQUENCE: 11				
atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca				48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro				
1 5 10 15				
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg				96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu				
20 25 30				
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa				144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu				
35 40 45				
tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac				192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr				
50 55 60				
cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg aaa ctg gag gaa				240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu				
65 70 75 80				
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt				288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg				
85 90 95				
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc tac att tgc				336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys				
100 105 110				
ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg ctg aaa gcc gat				384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp				
115 120 125				
act ccg gat tat ccg gag gcc ccg gta gac ccg cag tac cct gcc ttc				432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe				
130 135 140				
atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt				480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val				
145 150 155 160				
ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag				528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln				
165 170 175				
gtt gac cag gag cgt att gtt atc gct ggt gcc tcc cag ggt gcc gcc				576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly				
180 185 190				
atc gcc ctg gcg gta tct gcg ctg agc aag aaa gct aag gaa ctg ctg				624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Glu Leu Leu				
195 200 205				
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta				672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val				
210 215 220				
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc				720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg				
225 230 235 240				
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac gcc gtt				768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val				

-continued

	245	250	255	
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg				816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu				
	260	265	270	
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat				864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr				
	275	280	285	
tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa				912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu				
	290	295	300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa				960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys				
	305	310	315	320
ctg ttt gag aag ggc taa				978
Leu Phe Glu Lys Gly				
	325			

<210> SEQ ID NO 12

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 12

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro			
1 5 10 15			
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu			
20 25 30			
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu			
35 40 45			
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr			
50 55 60			
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu			
65 70 75 80			
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg			
85 90 95			
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys			
100 105 110			
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp			
115 120 125			
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe			
130 135 140			
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val			
145 150 155 160			
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln			
165 170 175			
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly			
180 185 190			
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Glu Leu Leu			
195 200 205			
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val			
210 215 220			
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg			
225 230 235 240			
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val			
245 250 255			

[illegible]

```
<210> SEQ ID NO 13
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(978)
```

<400> SEQUENCE: 13

atg	gcg	ttc	ttc	gac	ctg	cct	ctg	gaa	gaa	ctg	aag	aaa	tac	cgt	cca	48
Met	Ala	Phe	Phe	Asp	Leu	Pro	Leu	Glu	Glu	Leu	Lys	Lys	Tyr	Arg	Pro	
1			5					10						15		
gag	cgt	tac	gaa	gag	aag	gac	ttc	gac	gag	ttc	tgg	gag	gaa	act	ctg	96
Glu	Arg	Tyr	20	Glu	Lys	Asp	Phe	Asp	Glu	Phe	Trp	Glu	Glu	Thr	Leu	
								25					30			
gcg	gag	aga	gaa	aag	ttt	ccg	ctg	gac	cca	gtg	ttc	gag	cgt	atg	gaa	144
Ala	Glu	Arg	Glu	Lys	Phe	Pro	Leu	Asp	Pro	Val	Phe	Glu	Arg	Met	Glu	
		35					40					45				
tct	cgc	ctg	aaa	acc	gtg	gag	gca	tat	gac	gtt	act	ttt	tct	ggg	tac	192
Ser	Arg	Leu	Lys	Thr	Val	Glu	Ala	Tyr	Asp	Val	Thr	Phe	Ser	Gly	Tyr	
						55					60					
cgt	ggc	cag	cgt	atc	aaa	ggc	tgg	ctg	ctg	gtt	ccg	aaa	ctg	gag	gaa	240
Arg	Gly	Gln	Arg	Ile	Lys	Gly	Trp	Leu	Leu	Val	Pro	Lys	Leu	Glu	Glu	
65					70					75					80	
gaa	aaa	ctg	ccg	tgc	gta	gtt	cag	tac	atc	ggg	tac	aac	ggg	ggc	cgt	288
Glu	Lys	Leu	Pro	Cys	Val	Val	Gln	Tyr	Ile	Gly	Tyr	Asn	Gly	Gly	Arg	
				85					90				95			
ggc	ttt	ccg	cac	gat	tgg	ctg	ttc	tgg	ccg	tct	atg	ggc	tac	att	tgc	336
Gly	Phe	Pro	His	Asp	Trp	Leu	Phe	Trp	Pro	Ser	Met	Gly	Tyr	Ile	Cys	
				100				105					110			
ttc	gtc	atg	gat	act	cgt	ggg	cag	ggg	tcc	ggc	tgg	ctg	aaa	ggc	gat	384
Phe	Val	Met	Asp	Thr	Arg	Gly	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp	
			115				120					125				
act	ccg	gat	tat	ccg	gag	ggc	ccg	gta	gac	ccg	cag	tac	cct	ggc	ttc	432
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe	
						135					140					
atg	acg	cgt	ggg	att	ctg	gat	ccg	cgt	acc	tat	tac	tat	cgc	cgc	gtt	480
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Tyr	Arg	Arg	Val	
145					150					155					160	
ttt	acc	gat	gca	gtt	cgt	gcc	gta	gag	gcc	gcg	gct	tct	ttc	cct	cag	528
Phe	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Phe	Pro	Gln	
				165					170				175			
gtt	gac	cag	gag	cgt	att	gtt	atc	gct	ggg	ggc	tcc	cag	ggg	ggc	ggc	576
Val	Asp	Gln	Glu	Arg	Ile	Val	Ile	Ala	Gly	Gly	Ser	Gln	Gly	Gly	Gly	
				180				185				190				
atc	gcc	ctg	gcg	gta	tct	gcg	ctg	agc	aag	aaa	gct	aag	gca	ctg	ctg	624
Ile	Ala	Leu	Ala	Val	Ser	Ala	Leu	Ser	Lys	Lys	Ala	Lys	Ala	Leu	Leu	
							200					205				

-continued

tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta	672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	
210 215 220	
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc	720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	
225 230 235 240	
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt	768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val	
245 250 255	
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg	816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu	
260 265 270	
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat	864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	
275 280 285	
tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa	912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa	960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	
305 310 315 320	
ctg ttt gag aag ggc taa	978
Leu Phe Glu Lys Gly	
325	

<210> SEQ ID NO 14
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 14

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
Ala Glu Arg Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
Ser Arg Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp	
115 120 125	
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val	
145 150 155 160	
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln	
165 170 175	
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly	
180 185 190	
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	
195 200 205	

-continued

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 15
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1) .. (978)

<400> SEQUENCE: 15

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca	48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg	96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa	144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac	192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg gaa ctg gag gaa	240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Glu Leu Glu Glu	
65 70 75 80	
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt	288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc tac att tgc	336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg ctg aaa gcc gat	384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp	
115 120 125	
act ccg gat tat ccg gag gcc ccg gta gac ccg cag tac cct gcc ttc	432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt	480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val	
145 150 155 160	
ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag	528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln	

-continued

	165	170	175	
ggt gac cag gag cgt att gtt atc gct ggt ggc tcc cag ggt ggc ggc				576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly				
	180	185	190	
atc gcc ctg gcg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg				624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu				
	195	200	205	
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta				672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val				
	210	215	220	
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc				720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg				
	225	230	235	240
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt				768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val				
	245	250	255	
aac ttc gcg gct cgt gca aaa att ccg gaa ctg ttc tct gtt ggt ctg				816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Glu Leu Phe Ser Val Gly Leu				
	260	265	270	
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat				864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr				
	275	280	285	
tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa				912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu				
	290	295	300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa				960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys				
	305	310	315	320
ctg ttt gag aag ggc taa				978
Leu Phe Glu Lys Gly				
	325			

<210> SEQ ID NO 16

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 16

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Glu Leu Glu Glu
65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val

-continued

145	150	155	160
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ser Phe Pro Gln	165	170	175
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly	180	185	190
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	195	200	205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	210	215	220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	225	230	235
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val	245	250	255
Asn Phe Ala Ala Arg Ala Lys Ile Pro Glu Leu Phe Ser Val Gly Leu	260	265	270
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	275	280	285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	290	295	300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	305	310	315
Leu Phe Glu Lys Gly	325		

<210> SEQ ID NO 17
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(978)

<400> SEQUENCE: 17

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca	48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
gag cgt tac gaa gag aag gac ttc gac gag tac tgg gag gaa act ctg	96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Tyr Trp Glu Glu Thr Leu	
20 25 30	
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa	144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac	192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg aaa ctg gag gaa	240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt	288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc tac att tgc	336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg ctg aaa gcc gat	384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp	
115 120 125	

-continued

act ccg gat tat ccg gag ggc ccg gta gac ccg cag tac cct ggc ttc	432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
atg acg cgt ggt gtt ctg gat ccg cgt acc tat tac tat cgc cgc gtt	480
Met Thr Arg Gly Val Leu Asp Pro Arg Thr Tyr Tyr Arg Arg Val	
145 150 155 160	
ttt acc gat gca gtt cgt gcc gta gag gcc cgc gct tct ttc cct cag	528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ser Phe Pro Gln	
165 170 175	
gtt gac cag gag cgt att gtt atc gct ggt ggc tcc cag ggt ggc ggc	576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly	
180 185 190	
atc gcc ctg cgc gta tct gcg ctg agc aag aaa gct aag gca ctg ctg	624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	
195 200 205	
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta	672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	
210 215 220	
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc	720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	
225 230 235 240	
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt	768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val	
245 250 255	
aac ttc cgc gct cgt gca aaa att ccg gta ctg ttc tct gtt ggt ctg	816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Val Leu Phe Ser Val Gly Leu	
260 265 270	
atg gac aac atc agc cct cct tct acc gtt ttc cgc gca tat aac tat	864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	
275 280 285	
tat cgc ggt ccg aaa gaa acc cgt atc tat ccg tac aac agc cac gaa	912
Tyr Ala Gly Pro Lys Glu Thr Arg Ile Tyr Pro Tyr Asn Ser His Glu	
290 295 300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa	960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	
305 310 315 320	
ctg ttt gag aag ggc taa	978
Leu Phe Glu Lys Gly	
325	

<210> SEQ ID NO 18

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 18

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Tyr Trp Glu Glu Thr Leu
20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95

-continued

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Val Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Val Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Thr Arg Ile Tyr Pro Tyr Asn Ser His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 19
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(978)

<400> SEQUENCE: 19

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca	48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg	96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa	144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac	192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
cgt ggc cag cgt atc aaa ggc tgg ctg ctg gtt ccg aaa ctg gag gaa	240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt ggc cgt	288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	

85										90										95										
ggc	ttt	ccg	cac	gat	tgg	ctg	ttc	tgg	ccg	tct	atg	ggc	tac	att	tgc	336														
Gly	Phe	Pro	His	Asp	Trp	Leu	Phe	Trp	Pro	Ser	Met	Gly	Tyr	Ile	Cys															
100										105										110										
ttc	gtc	atg	gat	act	cgt	agt	cag	ggg	tcc	ggc	tgg	ctg	aaa	ggc	gat	384														
Phe	Val	Met	Asp	Thr	Arg	Ser	Gln	Gly	Ser	Gly	Trp	Leu	Lys	Gly	Asp															
115										120										125										
act	ccg	gat	tat	ccg	gag	ggc	ccg	gta	gac	ccg	cag	tac	cct	ggc	ttc	432														
Thr	Pro	Asp	Tyr	Pro	Glu	Gly	Pro	Val	Asp	Pro	Gln	Tyr	Pro	Gly	Phe															
130										135										140										
atg	acg	cgt	ggg	att	ctg	gat	ccg	cgt	acc	tat	tac	tat	cgc	cgc	gtt	480														
Met	Thr	Arg	Gly	Ile	Leu	Asp	Pro	Arg	Thr	Tyr	Tyr	Arg	Arg	Val																
145										150										155										
ttt	acc	gat	gca	gtt	cgt	gcc	gta	gag	gcc	gcg	gct	tct	ttc	cct	cag	528														
Phe	Thr	Asp	Ala	Val	Arg	Ala	Val	Glu	Ala	Ala	Ala	Ser	Phe	Pro	Gln															
165										170										175										
gtt	gac	cag	gag	cgt	att	gtt	atc	gct	ggg	ggc	tcc	cag	ggg	ggc	ggc	576														
Val	Asp	Gln	Glu	Arg	Ile	Val	Ile	Ala	Gly	Gly	Ser	Gln	Gly	Gly	Gly															
180										185										190										
atc	gcc	ctg	gcg	gta	tct	gcg	ctg	agc	aag	aaa	gct	aag	gca	ccg	ctg	624														
Ile	Ala	Leu	Ala	Val	Ser	Ala	Leu	Ser	Lys	Lys	Ala	Lys	Ala	Pro	Leu															
195										200										205										
tgt	gac	gtc	ccg	ttc	ctg	tgt	cac	ttc	cgt	cgc	gct	gtt	cag	ctg	gta	672														
Cys	Asp	Val	Pro	Phe	Leu	Cys	His	Phe	Arg	Arg	Ala	Val	Gln	Leu	Val															
210										215										220										
gat	acc	cat	ccg	tac	gcg	gag	att	act	aac	ttc	ctg	aaa	act	cac	cgc	720														
Asp	Thr	His	Pro	Tyr	Ala	Glu	Ile	Thr	Asn	Phe	Leu	Lys	Thr	His	Arg															
225										230										235										
gac	aaa	gaa	gaa	atc	gtt	ttc	ccg	acc	ctg	tcc	tat	ttc	gac	ggc	gtt	768														
Asp	Lys	Glu	Glu	Ile	Val	Phe	Arg	Thr	Leu	Ser	Tyr	Phe	Asp	Gly	Val															
245										250										255										
aac	ttc	gcg	gct	cgt	gca	aaa	att	ccg	gca	ctg	ttc	tct	gtt	ggg	ctg	816														
Asn	Phe	Ala	Ala	Arg	Ala	Lys	Ile	Pro	Ala	Leu	Phe	Ser	Val	Gly	Leu															
260										265										270										
atg	gac	aac	atc	agc	cct	cct	tct	acc	gtt	ttc	gcg	gca	tat	aac	tat	864														
Met	Asp	Asn	Ile	Ser	Pro	Pro	Ser	Thr	Val	Phe	Ala	Ala	Tyr	Asn	Tyr															
275										280										285										
tat	gcg	ggg	ccg	aaa	gaa	atc	cgt	atc	tat	ccg	tac	aac	aac	cac	gaa	912														
Tyr	Ala	Gly	Pro	Lys	Glu	Ile	Arg	Ile	Tyr	Pro	Tyr	Asn	Asn	His	Glu															
290										295										300										
ggc	ggg	ggg	agc	ttt	cag	gct	gtt	gaa	caa	gtg	aaa	ttc	ctg	aag	aaa	960														
Gly	Gly	Gly	Ser	Phe	Gln	Ala	Val	Glu	Gln	Val	Lys	Phe	Leu	Lys	Lys															
305										310										315										
ctg	ttt	gag	aag	ggc	taa																									

-continued

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
 50 55 60
 Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
 65 70 75 80
 Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
 85 90 95
 Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
 100 105 110
 Phe Val Met Asp Thr Arg Ser Gln Gly Ser Gly Trp Leu Lys Gly Asp
 115 120 125
 Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
 130 135 140
 Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
 145 150 155 160
 Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
 165 170 175
 Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
 180 185 190
 Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Pro Leu
 195 200 205
 Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 21
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1) .. (978)

<400> SEQUENCE: 21

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca	48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg	96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa	144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	

-continued

tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac	192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
cggtt ggc cag cgt atc aaa ggc tgg ctg ctg gtt ccg aaa ctg gag gaa	240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt ggc cgt	288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg ggc tac att tgc	336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
ttc gtc atg gat act cgt ggt cag ggt tcc ggc tgg ctg aaa ggc gat	384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp	
115 120 125	
act ccg gat tat ccg gag ggc ccg gta gac ccg cag tac cct ggc ttc	432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt	480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val	
145 150 155 160	
ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag	528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln	
165 170 175	
gtt gac cag gag cgt att gtt atc gct ggt ggc tcc cag ggt ggc ggc	576
Val Asp Gln Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly	
180 185 190	
atc gcc cag gcg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg	624
Ile Ala Gln Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	
195 200 205	
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta	672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	
210 215 220	
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc	720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	
225 230 235 240	
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt	768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val	
245 250 255	
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg	816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu	
260 265 270	
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat	864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	
275 280 285	
tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa	912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa	960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	
305 310 315 320	
ctg ttt gag aag ggc taa	978
Leu Phe Glu Lys Gly	
325	

<210> SEQ ID NO 22

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 22

```

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1      5      10      15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20     25     30
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35     40     45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50     55     60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65     70     75     80
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85     90     95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100    105    110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115    120    125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130    135    140
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145    150    155    160
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165    170    175
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180    185    190
Ile Ala Gln Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195    200    205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210    215    220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225    230    235    240
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245    250    255
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260    265    270
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275    280    285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
290    295    300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305    310    315    320
Leu Phe Glu Lys Gly
325

```

<210> SEQ ID NO 23

<211> LENGTH: 978

<212> TYPE: DNA

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: synthetic construct

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (1)..(978)

<400> SEQUENCE: 23

```

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro

```


-continued

1	5	10	15	
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg				96
Glu Arg Tyr 20	Glu Lys Asp 25	Phe Asp Glu Phe Trp 30	Glu Glu Thr Leu	
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa				144
Ala Glu Ser 35	Glu Lys Phe Pro 40	Val Phe Glu Arg Met 45	Glu	
tct cac ctg gaa acc gtg gag gca tat gac gtt act ttt tct ggt tac				192
Ser His Leu Glu Thr 50	Val Glu Ala Tyr 55	Val Thr Phe Ser 60	Gly Tyr	
cgt ggc cag cgt atc aaa ggc tgg ctg ctg gtt ccg aaa ctg gag gaa				240
Arg Gly Gln Arg Ile 65	Lys Gly Trp Leu 70	Val Pro Lys Leu 75	Glu Glu	
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt ggc cgt				288
Glu Lys Leu Pro 85	Cys Val Val Gln Tyr 90	Ile Gly Tyr Asn Gly 95	Gly Arg	
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg ggc tac att tgc				336
Gly Phe Pro 100	His Asp Trp Leu Phe 105	Trp Pro Ser Met 110	Gly Tyr Ile Cys	
ttc gtc atg gat act cgt ggt cag ggt tcc ggc tgg ctg aaa ggc gat				384
Phe Val Met Asp Thr 115	Arg Gly Gln Gly Ser 120	Gly Trp Leu Lys 125	Gly Asp	
act ccg gat tat ccg gag ggc ccg gta gac ccg cag tac cct ggc ttc				432
Thr Pro Asp Tyr Pro 130	Glu Gly Pro Val Asp 135	Pro Gln Tyr Pro 140	Gly Phe	
atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt				480
Met Thr Arg Gly Ile 145	Leu Asp Pro Arg Thr 150	Tyr Tyr Tyr Arg Arg 155	Val 160	
ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag				528
Phe Thr Asp Ala Val 165	Arg Ala Val Glu Ala Ala 170	Ser Phe Pro Gln 175		
gtt gac cag gag cgt att gtt atc gct ggt ggc tcc cag ggt ggc ggc				576
Val Asp Gln Glu Arg Ile Val 180	Ile Ala Gly Gly Ser 185	Gln Gly Gly Gly 190		
atc gcc ctg gcg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg				624
Ile Ala Leu Ala Val 195	Ser Ala Leu Ser Lys 200	Lys Ala Lys Ala Leu Leu 205		
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta				672
Cys Asp Val Pro Phe 210	Leu Cys His Phe Arg Arg 215	Ala Val Gln Leu Val 220		
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc				720
Asp Thr His Pro Tyr 225	Ala Glu Ile Thr Asn Phe 230	Leu Lys Thr His Arg 235	240	
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt				768
Asp Lys Glu Glu Ile Val 245	Phe Arg Thr Leu Ser Tyr 250	Phe Asp Gly Val 255		
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg				816
Asn Phe Ala Ala Arg Ala 260	Lys Ile Pro Ala Leu Phe 265	Ser Val Gly Leu 270		
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat				864
Met Asp Asn Ile Ser Pro 275	Pro Ser Thr Val Phe Ala 280	Ala Ala Tyr Asn Tyr 285		
tat gcg ggt ccg aaa gaa atc cgt atc ttt ccg tac aac aac cac gaa				912
Tyr Ala Gly Pro Lys Glu 290	Ile Arg Ile Phe Pro 295	Tyr Asn Asn His Glu 300		
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa				960
Gly Gly Gly Ser Phe 305	Gln Ala Val Glu Gln 310	Val Lys Phe Leu Lys 315	Lys 320	
ctg ttt gag aag ggc taa				978
Leu Phe Glu Lys Gly				

-continued

325

```

<210> SEQ ID NO 24
<211> LENGTH: 325
<212> TYPE: PRT
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 24

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1          5          10          15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20          25          30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35          40          45

Ser His Leu Glu Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50          55          60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65          70          75          80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85          90          95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100         105         110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115         120         125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130         135         140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145         150         155         160

Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165         170         175

Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180         185         190

Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195         200         205

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210         215         220

Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225         230         235         240

Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
245         250         255

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260         265         270

Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
275         280         285

Tyr Ala Gly Pro Lys Glu Ile Arg Ile Phe Pro Tyr Asn Asn His Glu
290         295         300

Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305         310         315         320

Leu Phe Glu Lys Gly
325

```

```

<210> SEQ ID NO 25
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: artificial sequence

```

-continued

```

<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1) .. (978)

<400> SEQUENCE: 25

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca      48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1          5          10          15

gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg      96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20          25          30

gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa      144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35          40          45

tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac      192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50          55          60

cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg aaa ctg gag gaa      240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65          70          75          80

gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt      288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85          90          95

ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc tac att tgc      336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
100         105         110

ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg cag aaa gcc gat      384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Gln Lys Gly Asp
115         120         125

act ccg gat tat ccg gag gcc ccg gta gac ccg cag tac cct gcc ttc      432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130         135         140

atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt      480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val
145         150         155         160

ttt acc gat gca gtt cgt gcc gta gag gcc gcc gct tct ttc cct cag      528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
165         170         175

gtt gac cag gag cgt att gtt atc gct ggt gcc tcc cag ggt gcc gcc      576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
180         185         190

atc gcc ctg ccg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg      624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195         200         205

tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta      672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
210         215         220

gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc      720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
225         230         235         240

gac aaa gaa gaa atc gtt ttc cgc tcc ctg tcc tat ttc gac gcc gtt      768
Asp Lys Glu Glu Ile Val Phe Arg Ser Leu Ser Tyr Phe Asp Gly Val
245         250         255

aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg      816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
260         265         270

atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat      864
Met Asp Asn Ile Ser Pro Pro Thr Val Phe Ala Ala Tyr Asn Tyr
275         280         285

```

-continued

tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa	912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
ggc ggt ggt agc ttt cag gtt gtt gaa caa gtg aaa ttc ctg aag aaa	960
Gly Gly Gly Ser Phe Gln Val Val Glu Gln Val Lys Phe Leu Lys Lys	
305 310 315 320	
ctg ttt gag aag ggc taa	978
Leu Phe Glu Lys Gly	
325	

<210> SEQ ID NO 26
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 26

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Gln Lys Gly Asp	
115 120 125	
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val	
145 150 155 160	
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln	
165 170 175	
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly	
180 185 190	
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	
195 200 205	
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	
210 215 220	
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	
225 230 235 240	
Asp Lys Glu Glu Ile Val Phe Arg Ser Leu Ser Tyr Phe Asp Gly Val	
245 250 255	
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu	
260 265 270	
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	
275 280 285	
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
Gly Gly Gly Ser Phe Gln Val Val Glu Gln Val Lys Phe Leu Lys Lys	

-continued

305	310	315	320	
Leu Phe Glu Lys Gly				
325				
 <210> SEQ ID NO 27				
<211> LENGTH: 978				
<212> TYPE: DNA				
<213> ORGANISM: artificial sequence				
<220> FEATURE:				
<223> OTHER INFORMATION: synthetic construct				
<220> FEATURE:				
<221> NAME/KEY: CDS				
<222> LOCATION: (1)..(978)				
 <400> SEQUENCE: 27				
atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca				48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro				
1 5 10 15				
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg				96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu				
20 25 30				
gcg gag agc gaa aag tct ccg ctg gac cca gtg ttc gag cgt atg gaa				144
Ala Glu Ser Glu Lys Ser Pro Leu Asp Pro Val Phe Glu Arg Met Glu				
35 40 45				
tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac				192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr				
50 55 60				
cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg aaa ctg gag gaa				240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu				
65 70 75 80				
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt				288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg				
85 90 95				
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc tac att tgc				336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys				
100 105 110				
ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg ctg aaa gcc gat				384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp				
115 120 125				
act ccg gat tat ccg gag gcc ccg gta gac ccg cag tac cct gcc ttc				432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe				
130 135 140				
atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt				480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val				
145 150 155 160				
ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag				528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln				
165 170 175				
gtt gac cag gag cgt att gtt atc gct ggt gcc tcc cag ggt gcc gcc				576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly				
180 185 190				
atc gcc ctg gcg ata tct gcg ctg agc aag aaa gct aag gca ctg ctg				624
Ile Ala Leu Ala Ile Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu				
195 200 205				
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta				672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val				
210 215 220				
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc				720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg				
225 230 235 240				
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac gcc gtt				768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val				

-continued

	245	250	255	
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg				816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu				
	260	265	270	
atg gac acc atc agc cct cct tct acc gtt ttc gcg gca tat aac tat				864
Met Asp Thr Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr				
	275	280	285	
tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa				912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu				
	290	295	300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa				960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys				
	305	310	315	320
ctg ttt gag aag ggc taa				978
Leu Phe Glu Lys Gly				
	325			

<210> SEQ ID NO 28

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 28

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro			
1	5	10	15
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu			
	20	25	30
Ala Glu Ser Glu Lys Ser Pro Leu Asp Pro Val Phe Glu Arg Met Glu			
	35	40	45
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr			
	50	55	60
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu			
	65	70	75
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg			
	85	90	95
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys			
	100	105	110
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp			
	115	120	125
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe			
	130	135	140
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val			
	145	150	155
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln			
	165	170	175
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly			
	180	185	190
Ile Ala Leu Ala Ile Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu			
	195	200	205
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val			
	210	215	220
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg			
	225	230	235
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val			
	245	250	255

-continued

```

Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
      260                      265                      270
Met Asp Thr Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
      275                      280                      285
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
      290                      295                      300
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
305                      310                      315                      320
Leu Phe Glu Lys Gly
      325

```

```

<210> SEQ ID NO 29
<211> LENGTH: 978
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: synthetic construct
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1)..(978)

```

```

<400> SEQUENCE: 29

```

```

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca      48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1                      5                      10                      15

gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg      96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
      20                      25                      30

gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa      144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
      35                      40                      45

tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac      192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
      50                      55                      60

cgt ggc cag cgt atc aaa ggc tgg ctg ctg gtt ccg ata ctg gag gaa      240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Ile Leu Glu Glu
65                      70                      75                      80

gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt ggc cgt      288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
      85                      90                      95

ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg ggc tac att tgc      336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys
      100                      105                      110

ttc gtc atg gat act cgt ggt cag ggt tcc ggc tgg ctg aat ggc gat      384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Asn Gly Asp
      115                      120                      125

act ccg gat tat ccg gag ggc ccg gta gac ccg cag tac cct ggc ttc      432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
      130                      135                      140

atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt      480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Arg Arg Val
145                      150                      155                      160

ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag      528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln
      165                      170                      175

gtt gac cag gag cgt att gtt atc gct ggt ggc tcc cag ggt ggc ggc      576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly
      180                      185                      190

atc gcc ctg gcg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg      624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu
195                      200                      205

```

-continued

tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta	672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val	
210 215 220	
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc	720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg	
225 230 235 240	
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt	768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val	
245 250 255	
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg	816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu	
260 265 270	
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat	864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr	
275 280 285	
tat gcg ggt ccg aga gaa atc cgt atc tat ccg tac aac aac cac gaa	912
Tyr Ala Gly Pro Arg Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu	
290 295 300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa	960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys	
305 310 315 320	
ctg ttt gag aag ggc taa	978
Leu Phe Glu Lys Gly	
325	

<210> SEQ ID NO 30
 <211> LENGTH: 325
 <212> TYPE: PRT
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 30

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Ile Leu Glu Glu	
65 70 75 80	
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Tyr Ile Cys	
100 105 110	
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Asn Gly Asp	
115 120 125	
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val	
145 150 155 160	
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln	
165 170 175	
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly	
180 185 190	
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu	
195 200 205	

-continued

Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val
 210 215 220
 Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg
 225 230 235 240
 Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val
 245 250 255
 Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu
 260 265 270
 Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr
 275 280 285
 Tyr Ala Gly Pro Arg Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu
 290 295 300
 Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys
 305 310 315 320
 Leu Phe Glu Lys Gly
 325

<210> SEQ ID NO 31
 <211> LENGTH: 978
 <212> TYPE: DNA
 <213> ORGANISM: artificial sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: synthetic construct
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1) .. (978)

<400> SEQUENCE: 31

atg gcg ttc ttc gac ctg cct ctg gaa gaa ctg aag aaa tac cgt cca	48
Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro	
1 5 10 15	
gag cgt tac gaa gag aag gac ttc gac gag ttc tgg gag gaa act ctg	96
Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu	
20 25 30	
gcg gag agc gaa aag ttt ccg ctg gac cca gtg ttc gag cgt atg gaa	144
Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu	
35 40 45	
tct cac ctg aaa acc gtg gag gca tat gac gtt act ttt tct ggt tac	192
Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr	
50 55 60	
cgt gcc cag cgt atc aaa gcc tgg ctg ctg gtt ccg aaa ctg gag gaa	240
Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu	
65 70 75 80	
gaa aaa ctg ccg tgc gta gtt cag tac atc ggt tac aac ggt gcc cgt	288
Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg	
85 90 95	
ggc ttt ccg cac gat tgg ctg ttc tgg ccg tct atg gcc ttc att tgc	336
Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Phe Ile Cys	
100 105 110	
ttc gtc atg gat act cgt ggt cag ggt tcc gcc tgg ctg aaa gcc gat	384
Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp	
115 120 125	
act ccg gat tat ccg gag gcc ccg gta gac ccg cag tac cct gcc ttc	432
Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe	
130 135 140	
atg acg cgt ggt att ctg gat ccg cgt acc tat tac tat cgc cgc gtt	480
Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Arg Arg Val	
145 150 155 160	
ttt acc gat gca gtt cgt gcc gta gag gcc gcg gct tct ttc cct cag	528
Phe Thr Asp Ala Val Arg Ala Val Glu Ala Ala Ala Ser Phe Pro Gln	

-continued

	165	170	175	
ggt gac cag gag cgt att gtt atc gct ggt ggc tcc cag ggt ggc ggc				576
Val Asp Gln Glu Arg Ile Val Ile Ala Gly Gly Ser Gln Gly Gly Gly				
	180	185	190	
atc gcc ctg gcg gta tct gcg ctg agc aag aaa gct aag gca ctg ctg				624
Ile Ala Leu Ala Val Ser Ala Leu Ser Lys Lys Ala Lys Ala Leu Leu				
	195	200	205	
tgt gac gtc ccg ttc ctg tgt cac ttc cgt cgc gct gtt cag ctg gta				672
Cys Asp Val Pro Phe Leu Cys His Phe Arg Arg Ala Val Gln Leu Val				
	210	215	220	
gat acc cat ccg tac gcg gag att act aac ttc ctg aaa act cac cgc				720
Asp Thr His Pro Tyr Ala Glu Ile Thr Asn Phe Leu Lys Thr His Arg				
	225	230	235	240
gac aaa gaa gaa atc gtt ttc cgc acc ctg tcc tat ttc gac ggc gtt				768
Asp Lys Glu Glu Ile Val Phe Arg Thr Leu Ser Tyr Phe Asp Gly Val				
	245	250	255	
aac ttc gcg gct cgt gca aaa att ccg gca ctg ttc tct gtt ggt ctg				816
Asn Phe Ala Ala Arg Ala Lys Ile Pro Ala Leu Phe Ser Val Gly Leu				
	260	265	270	
atg gac aac atc agc cct cct tct acc gtt ttc gcg gca tat aac tat				864
Met Asp Asn Ile Ser Pro Pro Ser Thr Val Phe Ala Ala Tyr Asn Tyr				
	275	280	285	
tat gcg ggt ccg aaa gaa atc cgt atc tat ccg tac aac aac cac gaa				912
Tyr Ala Gly Pro Lys Glu Ile Arg Ile Tyr Pro Tyr Asn Asn His Glu				
	290	295	300	
ggc ggt ggt agc ttt cag gct gtt gaa caa gtg aaa ttc ctg aag aaa				960
Gly Gly Gly Ser Phe Gln Ala Val Glu Gln Val Lys Phe Leu Lys Lys				
	305	310	315	320
ctg ttt gag aag ggc taa				978
Leu Phe Glu Lys Gly				
	325			

<210> SEQ ID NO 32

<211> LENGTH: 325

<212> TYPE: PRT

<213> ORGANISM: artificial sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 32

Met Ala Phe Phe Asp Leu Pro Leu Glu Glu Leu Lys Lys Tyr Arg Pro
1 5 10 15

Glu Arg Tyr Glu Glu Lys Asp Phe Asp Glu Phe Trp Glu Glu Thr Leu
20 25 30

Ala Glu Ser Glu Lys Phe Pro Leu Asp Pro Val Phe Glu Arg Met Glu
35 40 45

Ser His Leu Lys Thr Val Glu Ala Tyr Asp Val Thr Phe Ser Gly Tyr
50 55 60

Arg Gly Gln Arg Ile Lys Gly Trp Leu Leu Val Pro Lys Leu Glu Glu
65 70 75 80

Glu Lys Leu Pro Cys Val Val Gln Tyr Ile Gly Tyr Asn Gly Gly Arg
85 90 95

Gly Phe Pro His Asp Trp Leu Phe Trp Pro Ser Met Gly Phe Ile Cys
100 105 110

Phe Val Met Asp Thr Arg Gly Gln Gly Ser Gly Trp Leu Lys Gly Asp
115 120 125

Thr Pro Asp Tyr Pro Glu Gly Pro Val Asp Pro Gln Tyr Pro Gly Phe
130 135 140

Met Thr Arg Gly Ile Leu Asp Pro Arg Thr Tyr Tyr Tyr Arg Arg Val

-continued

145		150		155		160
Phe Thr Asp	Ala Val Arg	Ala Val Glu	Ala Ala Ser	Phe Pro Gln		
	165		170		175	
Val Asp Gln	Glu Arg Ile Val	Ile Ala Gly	Gly Ser Gln	Gly Gly Gly		
	180		185		190	
Ile Ala Leu	Ala Val Ser	Ala Leu Ser	Lys Lys Ala	Lys Ala Leu	Leu	
	195		200		205	
Cys Asp Val	Pro Phe Leu	Cys His Phe	Arg Arg Ala	Val Gln Leu	Val	
	210		215		220	
Asp Thr His	Pro Tyr Ala	Glu Ile Thr	Asn Phe Leu	Lys Thr His	Arg	
	225		230		235	
Asp Lys Glu	Glu Ile Val	Phe Arg Thr	Leu Ser Tyr	Phe Asp Gly	Val	
	245		250		255	
Asn Phe Ala	Ala Arg Ala	Lys Ile Pro	Ala Leu Phe	Ser Val Gly	Leu	
	260		265		270	
Met Asp Asn	Ile Ser Pro	Pro Ser Thr	Val Phe Ala	Ala Tyr Asn	Tyr	
	275		280		285	
Tyr Ala Gly	Pro Lys Glu	Ile Arg Ile	Tyr Pro Tyr	Asn Asn His	Glu	
	290		295		300	
Gly Gly Gly	Ser Phe Gln	Ala Val Glu	Gln Val Lys	Phe Leu Lys	Lys	
	305		310		315	
Leu Phe Glu	Lys Gly					
	325					

What is claimed is:

1. An isolated polypeptide having perhydrolytic activity comprising the amino acid sequence of SEQ ID NO: 22.

2. The polypeptide of claim 1; wherein said polypeptide is characterized by a peracetic acid formation specific activity that is at least 1.05-fold higher than the peracetic acid formation specific activity of the *Thermotoga maritime* C277S acetyl xylan esterase having amino acid sequence SEQ ID NO: 5.

3. A process for producing a peroxycarboxylic acid comprising:

(a) providing a set of reaction components comprising:

(1) at least one substrate selected from the group consisting of:

(i) one or more esters having the structure



wherein


X=an ester group of the formula $R_6-C(O)O$;

R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;

R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;


m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and wherein said esters have solubility in water of at least 5 ppm at 25° C.;

(ii) one or more glycerides having the structure



wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

(iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_n$, H and n is 1 to 10;

(iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
(v) any combination of (i) through (iv);

(2) a source of peroxygen; and

(3) an enzyme catalyst comprising the polypeptide of claim 1;

(b) combining the set of reaction components under suitable reaction conditions whereby peroxycarboxylic acid is produced; and

99

(c) optionally diluting the peroxycarboxylic acid produced in step (b).

4. The process of claim 3 further comprising the step of: d) contacting a hard surface or inanimate object with the peroxycarboxylic acid produced in step (b) or step (c); whereby said hard surface or said inanimate object is disinfected, bleached, destained or a combination thereof.

5. The process of claim 3 wherein the inanimate object is a medical instrument.

6. The process of claim 3 further comprising the step of: d) contacting an article of clothing or a textile with peroxycarboxylic acid produced in step (b) or step (c); whereby the article of clothing or textile receives a benefit.

7. The process of claim 6 wherein the benefit is selected from the group consisting of a disinfecting, sanitizing, bleaching, destaining, deodorizing, and combinations thereof.

8. The process of claim 3 further comprising the step of: d) contacting wood pulp or paper pulp with peroxycarboxylic acid produced in step (b) or step (c); whereby the wood pulp or paper pulp is bleached.

9. The process of claim 3 wherein the substrate is selected from the group consisting of: monoacetin; diacetin; triacetin; monopropionin; dipropionin; tripropionin; monobutylin; dibutylin; tributyrin; glucose pentaacetate; xylose tetraacetate; acetylated xylan; acetylated xylan fragments; β -D-ribofuranose-1,2,3,5-tetraacetate; tri-O-acetyl-D-galactal; tri-O-acetyl-D-glucal; monoesters or diesters of 1,2-ethanediol, 1,2-propanediol, 1,3-propanediol, 1,2-butanediol, 1,3-butanediol, 2,3-butanediol, 1,4-butanediol, 1,2-pentanediol, 2,5-pentanediol, 1,6-pentanediol, 1,2-hexanediol, 2,5-hexanediol, 1,6-hexanediol; and mixtures thereof.

10. The process of claim 9 wherein the substrate is triacetin.

11. The process of claim 3 wherein the peroxycarboxylic acid produced is peracetic acid, perpropionic acid, perbutyric acid, perlactic acid, perglycolic acid, permethoxyacetic acid, per- β -hydroxybutyric acid, or mixtures thereof.

12. The process of claim 3 wherein the enzyme catalyst is in the form of a microbial cell, a permeabilized microbial cell, a microbial cell extract, a partially purified enzyme, or a purified enzyme.

13. A composition comprising:

(a) a set of reaction components comprising:

(1) at least one substrate selected from the group consisting of:

(i) one or more esters having the structure



wherein

X=an ester group of the formula $R_6-C(O)O$;

R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;


R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;

m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and

wherein said esters have solubility in water of at least 5 ppm at 25° C.;


100

(ii) one or more glycerides having the structure



wherein R_1 =C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $R_1C(O)$;

(iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(CH_2CH_2O)_n$, or $(CH_2CH(CH_3)-O)_nH$ and n is 1 to 10;

(iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and

(v) any combination of (i) through (iv);

(2) a source of peroxygen; and

(3) an enzyme catalyst comprising the polypeptide of claim 1; and

(b) at least one peroxycarboxylic acid formed upon combining the set of reaction components of (a).

14. A peracid generation and delivery system comprising:

(a) a first compartment comprising

(1) an enzyme catalyst comprising the polypeptide of claim 1;

(2) at least one substrate selected from the group consisting of:

(i) one or more esters having the structure



wherein

X=an ester group of the formula $R_6-C(O)O$;

R_6 =C1 to C7 linear, branched or cyclic hydrocarbyl moiety, optionally substituted with hydroxyl groups or C1 to C4 alkoxy groups, wherein R_6 optionally comprises one or more ether linkages for R_6 =C2 to C7;


R_5 =a C1 to C6 linear, branched, or cyclic hydrocarbyl moiety optionally substituted with hydroxyl groups; wherein each carbon atom in R_5 individually comprises no more than one hydroxyl group or no more than one ester group; wherein R_5 optionally comprises one or more ether linkages;

m is an integer ranging from 1 to the number of carbon atoms in R_5 ; and

wherein said esters have solubility in water of at least 5 ppm at 25° C.;


101

(ii) one or more glycerides having the structure



wherein R_1 = C1 to C21 straight chain or branched chain alkyl optionally substituted with an hydroxyl or a C1 to C4 alkoxy group and R_3 and R_4 are individually H or $\text{R}_1\text{C}(\text{O})$;

(iii) one or more esters of the formula:



wherein R_1 is a C1 to C7 straight chain or branched chain alkyl optionally substituted with an

102

hydroxyl or a C1 to C4 alkoxy group and R_2 is a C1 to C10 straight chain or branched chain alkyl, alkenyl, alkynyl, aryl, alkylaryl, alkylheteroaryl, heteroaryl, $(\text{CH}_2\text{CH}_2\text{O})_n$, or $(\text{CH}_2\text{CH}(\text{CH}_3)-\text{O})_n\text{H}$ and n is 1 to 10;

(iv) one or more acylated monosaccharides, acylated disaccharides, or acylated polysaccharides; and
(v) any combination of (i) through (iv); and

(3) an optional buffer; and

(b) a second compartment comprising

(1) source of peroxygen;

(2) a peroxide stabilizer; and

(3) an optional buffer.

15 **15.** The peracid generation and delivery system of claim **14** wherein the substrate comprises triacetin.

16. A laundry care product comprising the polypeptide of claim **1**.

17. A personal care product comprising the polypeptide of claim **1**.

* * * * *