



(11) **EP 2 081 925 B1**

(12) **EUROPEAN PATENT SPECIFICATION**

- (45) Date of publication and mention of the grant of the patent: **15.02.2012 Bulletin 2012/07**
- (51) Int Cl.: **C07D 403/12** <sup>(2006.01)</sup> **A01N 43/56** <sup>(2006.01)</sup>  
**A01N 43/713** <sup>(2006.01)</sup>
- (21) Application number: **07818631.9**
- (86) International application number: **PCT/EP2007/008551**
- (22) Date of filing: **02.10.2007**
- (87) International publication number: **WO 2008/043456 (17.04.2008 Gazette 2008/16)**

(54) **Benzoylpyrazoles, their use as herbicides, and intermediates**

Benzoylpyrazole, ihre Verwendung als Herbizide, und Zwischenprodukte

Benzoylpyrazoles, utilisation de ceux-ci en tant qu'herbicides et intermédiaires pour la fabrication de ceux-ci

- (84) Designated Contracting States:  
**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IS IT LI LT LU LV MC MT NL PL PT RO SE SI SK TR**
- (30) Priority: **13.10.2006 JP 2006279533**
- (43) Date of publication of application:  
**29.07.2009 Bulletin 2009/31**
- (73) Proprietor: **Bayer CropScience AG**  
**40789 Monheim (DE)**
- (72) Inventors:  
• **NARABU, Shinichi**  
**Yuki-shi, Ibaraki 307-0001 (JP)**  
• **YONETA, Yasushi**  
**Kazo-shi, Saitama 347-0056 (JP)**  
• **YANAGI, Akihiko**  
**Tochigi 323-0825 (JP)**  
• **SHIRAKURA, Shinichi**  
**Oyama-shi, Tochigi 323-0807 (JP)**  
• **UKAWA, Seiji**  
**Oyama -shi, Tochigi 323-0820 (JP)**
- **ICHIHARA, Teruyuki**  
**Tochigi 323-0822 (JP)**
  - **NAKAMURA, Shin**  
**Oyama-shi, Tochigi 323-0807 (JP)**
  - **HILLS, Martin**  
**65510 Idstein (DE)**
  - **KEHNE, Heinz**  
**65719 Hofheim (DE)**
  - **DITTGEN, Jan**  
**60316 Frankfurt a. M. (DE)**
  - **FEUCHT, Dieter**  
**65760 Eschborn (DE)**
  - **ROSINGER, Christopher**  
**65719 Hofheim (DE)**
- (56) References cited:  
**EP-A- 0 629 623 WO-A-01/10850**  
**WO-A-03/066607 DE-A1- 10 039 723**
- Remarks:  
The file contains technical information submitted after the application was filed and not included in this specification

**EP 2 081 925 B1**

Note: Within nine months of the publication of the mention of the grant of the European patent in the European Patent Bulletin, any person may give notice to the European Patent Office of opposition to that patent, in accordance with the Implementing Regulations. Notice of opposition shall not be deemed to have been filed until the opposition fee has been paid. (Art. 99(1) European Patent Convention).

## Description

Technical Field

5 **[0001]** The present invention relates to novel benzoylpyrazoles, their use as herbicides, to process for their preparation and to their novel intermediates.

Background Art


10 **[0002]** It has already been known that a certain kind of benzoylpyrazoles exhibits an action as herbicides (e.g., PCT International Publications WO03/66607, WO01/10850 and WO01/53275).

Disclosure of the invention

15 **[0003]** However, compounds disclosed in the above publications are not satisfactory enough from the viewpoints of efficacy as herbicides and/or their safety.

**[0004]** The present inventors have intensively studied to create novel compounds having higher efficacy as herbicides and being safer. As a result, the present inventors have found novel benzoylpyrazoles represented by the formula set forth below which exhibit superior herbicidal activity and safety against crops.

20 **[0005]** Benzoylpyrazoles represented by the formula (I):



30 wherein

R<sup>1</sup> represents alkyl;

R<sup>2</sup> represents alkyl or cycloalkyl;

35 R<sup>3</sup> represents hydrogen, alkyl, alkenyl, alkynyl, aralkyl, alkylsulfonyl, optionally substituted phenylsulfonyl, acyl or acylalkyl;

X represents halogen or alkyl;

Y represents halogen;

n represents 2 or 3, and

40 T represents a group T1:




or a group T2:




55 wherein R<sup>4</sup> represents hydrogen, alkyl, alkoxy or alkylthio.

**[0006]** The compounds of the above formula (I) of the present disclosure can be synthesized, for example, by either of the preparation process (a) or (b) set forth below. Preparation process (a): compounds of formula (I) wherein R<sup>3</sup> represents hydrogen:

**[0007]** A process of rearranging compounds represented by the formula (II):




10 wherein X, Y, n and T are as defined above and Z represents a group:



20 wherein R<sup>1</sup> and R<sup>2</sup> are as defined above.

**[0008]** Preparation process (b): compounds of formula (I) wherein R<sup>3</sup> represents alkyl, alkenyl, alkynyl, aralkyl, alkyl-sulfonyl, optionally substituted phenylsulfonyl, acyl or acylalkyl:

**[0009]** A process of reacting compounds represented by the formula (Ia):



35 wherein R<sup>1</sup>, R<sup>2</sup>, X, Y, n and T are as defined above, with compounds represented by the formula (III):



wherein

40 R<sup>3</sup> is as defined above and  
L represents halogen.

**[0010]** The benzoylpyrazoles of the formula (I) provided by the present disclosure have a more potent herbicidal activity than the compounds described in the above prior art documents while exhibiting extremely superior efficacy as selective herbicides without substantially no phytotoxicity to crops. Particularly, the benzoylpyrazoles of the formula (I) exhibit extremely superior efficacy as herbicides for Leguminous dry field crops such as soy beans and Poaceae dry field crops such as wheat, barley, oats and corn.

**[0011]** Accordingly, the benzoylpyrazoles of the formula (I) of the present disclosure are particularly useful as herbicides for dry field.

50 Best Mode for Carrying out the Invention

**[0012]** In the present specification, the term "halogen" means fluorine, chlorine, bromine or iodine, and preferably chlorine and bromine.

55 **[0013]** The term "alkyl" may be those of a linear or branched chain, and means for example C<sub>1-6</sub> alkyl. Specifically, examples of "alkyl" include methyl, ethyl, n- or iso-propyl, n-, iso-, sec- or tert-butyl, n-, iso-, neo- or tert-pentyl, n- or iso-hexyl.

**[0014]** Examples of "cycloalkyl" include cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl.

[0015] Examples of "alkenyl" include allyl, 1-methylallyl, 1,1-dimethylallyl, 2-butenyl, 2-pentenyl, 2-hexenyl, 2-heptenyl.

[0016] Examples of "alkynyl" include ethynyl, 2-propynyl, 1-methyl-2-propynyl, 1,1-dimethyl-2-propynyl, 2-butyne, 2-pentyne, 2-hexynyl, 2-heptynyl.

5 [0017] The term "aralkyl" means, for example, an arylalkyl group the alkyl moiety of which has the above meaning, and the "aryl" means, for example, phenyl, naphthyl, and specifically, benzyl.

[0018] The term "alkoxy" means, for example, an alkyl-O- group the alkyl moiety of which has the above meaning, and can be, for example, C<sub>1-6</sub> alkoxy. Specifically, examples of "alkoxy" include methoxy, ethoxy, n- or iso-propoxy, n-, iso-, sec- or tert-butoxy, n-pentyloxy, n-hexyloxy.

10 [0019] The term "alkylthio" means an alkyl-S- group the alkyl moiety of which has the above meaning, and can be, for example, C<sub>1-6</sub> alkylthio. Specifically, examples of "alkylthio" include methylthio, ethylthio, n- or iso-propylthio, n-, iso-, sec- or tert-butylthio, n-pentylthio, n-hexylthio.

[0020] The term "alkylsulfonyl" means an alkyl-SO<sub>2</sub>- group the alkyl moiety of which has the above meaning, and can be, for example, C<sub>1-6</sub> alkylsulfonyl. Specifically, examples of "alkylsulfonyl" include methylsulfonyl, ethylsulfonyl, n- or iso-propylsulfonyl, n-, iso-, sec- or tert-butylsulfonyl, n-pentylsulfonyl, n-hexylsulfonyl.

15 [0021] "Acyl" is an acyl group derived from aliphatic saturated monocarboxylic acid, or aliphatic or aromatic hydrocarbon ring carboxylic acid, and means for example, a C<sub>1-6</sub> alkyl-CO- group the alkyl moiety of which has the above meaning, or an aryl-CO- group the aryl moiety of which is as defined above. Specifically, examples of "acyl" include acetyl, propionyl, butyryl, isobutyryl, valeryl, isovaleryl, pivaloyl, n-pentylcarbonyl, n-hexylcarbonyl, benzoyl.

20 [0022] "Acylalkyl" is an acylalkyl group the acyl and alkyl moieties of which have the above meaning, and means for example, a benzoyl-C<sub>1-6</sub> alkyl group, and specifically, includes phenacyl.

[0023] The compounds representing the invention are those of the formula (I), wherein

R<sup>1</sup> represents C<sub>1-6</sub> alkyl;

R<sup>2</sup> represents C<sub>1-6</sub> alkyl or C<sub>3-7</sub> cycloalkyl;


25 R<sup>3</sup> represents hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> alkenyl, C<sub>3-7</sub> alkynyl, C<sub>7-12</sub> aralkyl, C<sub>1-6</sub> alkylsulfonyl, optionally C<sub>1-6</sub> alkyl-substituted phenylsulfonyl, C<sub>1-6</sub> alkyl-carbonyl, benzoyl or benzoyl-C<sub>1-6</sub> alkyl;

X represents halogen or C<sub>1-6</sub> alkyl;

Y represents halogen;

n represents 2 or 3; and

30 T represents a group represented by



wherein R<sup>4</sup> represents hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> alkylthio.

40 [0024] A preferable group of the compounds of the invention includes those of the formula (I), wherein

R<sup>1</sup> represents methyl or ethyl;

R<sup>2</sup> represents methyl, ethyl or cyclopropyl;


45 R<sup>3</sup> represents hydrogen, methyl, allyl, propargyl, benzyl, methanesulfonyl, n-propanesulfonyl, optionally methyl-substituted phenylsulfonyl, acetyl, benzoyl or phenacyl;

X represents chlorine, bromine or methyl;

Y represents chlorine or bromine;


n represents 2 or 3; and

50 T represents a group represented by




wherein R<sup>4</sup> represents hydrogen, methyl, ethyl, methoxy or methylthio.



**[0025]** Using, for example, 5-{2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoyloxy}-1,3-dimethylpyrazole, acetone cyanohydrin as a cyanide, and triethylamine as a base, the preparation process (a) mentioned above is represented by the following reaction scheme:




**[0026]** Using, for example, 4-{2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoyl}-1,3-dimethyl-5-hydroxypyrazole, benzyl bromide and potassium carbonate as an acid coupling agent, the preparation process (b) mentioned above is represented by the following reaction scheme:



**[0027]** The compounds of the formula (I) of the present invention can exist as tautomers when R<sup>3</sup> represents hydrogen, as shown by the following formulae (Ia), (Ib) and (Ic).




or



**[0028]** The compounds of the formula (II) used as raw materials in the above preparation process (a) are novel compounds not described in prior literatures. They may be prepared in accordance with the processes described in, for example, Japanese Patent KOKAI Publication No. 2-173 (hereinafter "Japanese Patent KOKAI Publication No." is referred to as "JP-A-") and WO93/18031. Namely, they can be prepared by reacting compounds represented by the formula (IV):




wherein Z is as defined above with compounds represented by the formula (V):



wherein X, Y, n and T are as defined above and Hal represents halogen, preferably chlorine or bromine, in the presence of a suitable acid coupling agent, for example, triethylamine in a suitable diluent, for example, dichloromethane.


**[0029]** The compounds represented by the above formula (IV) per se are known and can be prepared according to the processes described in literatures, for example, JP-A-61-257974 and J-A-10-109976.

**[0030]** A part of the compounds represented by the above formula (V) are novel compounds not described in prior literatures and can be prepared in accordance with the processes described in literatures, for example, JP-A-2-173, WO93/18031, and WO03/66607. Namely, they can be prepared by reacting compounds represented by the formula (VI):




wherein X, Y, n and T are as defined above with halogenating agents such as, for example, phosphorus oxychloride, phosphorus oxybromide, phosphorus trichloride, phosphorus tribromide, phosgene, oxalyl dichloride, thionyl chloride and thionyl bromide.

**[0031]** A part of the compounds represented by the above formula (VI) are novel compounds not described in prior literatures and can be prepared in accordance with the processes described in literatures, for example, JP-A-2-1-73 and WO03/66607. Namely, they can be prepared by hydrolyzing compounds represented by the formula (VII):



wherein X, Y, n and T are as defined above and  $\text{R}^5$  represents  $\text{C}_{1-4}$  alkyl, preferably methyl or ethyl in the presence of a suitable base, for example, sodium hydroxide in a diluent, for example, aqueous dioxane.

**[0032]** A part of the compounds represented by the above formula (VII) are novel compounds not described in prior literatures and can be prepared in accordance with the processes described in literatures, for example, German Patent No.10039723 and WO03/66607. Namely, they can be prepared by reacting compounds represented by the formula (VIII):



wherein X, Y and n and are as defined above and R<sup>5</sup> represents C<sub>1-4</sub> alkyl, preferably methyl or ethyl, and Hal represents halogen, preferably chlorine or bromine with the compounds represented by the formula (IX)



wherein T is as defined above, in the presence of a suitable condensing agent, for example, potassium carbonate in a suitable diluent, for example, N,N-dimethylformamide.

**[0033]** The compounds represented by the above formula (VIII) per se are known and can be easily prepared by the processes described in, for example, JP-A-6-247891, JP-A-7-206808 and European Patent No.369803

**[0034]** The compounds represented by the above formula (IX) per se are known and can be easily prepared by the processes described in, for example, Journal of the American Chemical Society, Vol. 80, p. 3908-3911 (1958), Journal of Organic Chemistry, Vol. 22, p. 1142-1145 (1957), Journal of Organic Chemistry, Vol. 56, p. 2395-2400 (1991) and JP-A-9-309883.

**[0035]** Representative examples of the compounds of the formula (II) used as raw materials in the above preparation process (a) include those described below.

5- $\{2,4\}$ -Dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]-benzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2,4\}$ -dichloro-3-[2-(2H-tetrazol-2-yl)ethyloxy]-benzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2,4\}$ -dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]-benzoyloxy}-1-methyl-3-cyclopropylpyrazole,  
 5- $\{2\}$ -methyl-3-[2-(5-methyl-1H-tetrazol-1-yl)-ethyloxy]-4-bromobenzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2\}$ -methyl-3-[2-(5-methyl-2H-tetrazol-2-yl)-ethyloxy]-4-bromobenzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2\}$ -methyl-3-[2-(5-methyl-1H-tetrazol-1-yl)-ethyloxy]-4-chlorobenzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2\}$ -methyl-3-[2-(5-ethyl-1H-tetrazol-1-yl)ethyloxy]-4-bromobenzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2,4\}$ -dichloro-3-[2-(5-methylthio-1H-tetrazol-1-yl)ethyloxy]benzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2\}$ -methyl-3-[2-(5-methoxy-1H-tetrazol-1-yl)-ethyloxy]-4-bromobenzoyloxy}-1,3-dimethylpyrazole,  
 5- $\{2\}$ -methyl-3-[2-(2H-tetrazol-2-yl)ethyloxy]-4-bromobenzoyloxy}-1-methyl-3-ethylpyrazole,  
 5- $\{2,4\}$ -dichloro-3-[3-(1H-tetrazol-1-yl)propyloxy]-benzoyloxy}-1,3-dimethylpyrazole,  
 1- $\{2,4\}$ -dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]-benzoyl]-2,5-dimethyl-1,2-dihydro-3H-pyrazol-3-one, and etc.

**[0036]** Representative examples of the compounds of the formula (IV) used as raw materials in the preparation of the compounds of the above formula (II) include those described below.

1,3-Dimethyl-5-hydroxypyrazole,  
 1-methyl-3-ethyl-5-hydroxypyrazole,  
 1-methyl-3-cyclopropyl-5-hydroxypyrazole,  
 1-ethyl-3-methyl-5-hydroxypyrazole,  
 1-ethyl-3-cyclopropyl-5-hydroxypyrazole, and etc.

**[0037]** Representative examples of the compounds of the formula (V) used as raw materials in the preparation of the compounds of the above formula (II) include those described below.

2,4-Dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]benzoyl chloride,  
 2,4-dichloro-3-[2-(2H-tetrazol-2-yl)ethyloxy]benzoyl chloride,  
 2-methyl-3-[2-(5-methyl-1H-tetrazol-1-yl)ethyloxy]-4-bromobenzoyl chloride,  
 2-methyl-3-[2-(5-methyl-2H-tetrazol-2-yl)ethyloxy]-4-bromobenzoyl chloride,  
 2-methyl-3-[2-(1H-tetrazol-1-yl)ethyloxy]-4-chlorobenzoyl chloride,  
 2,4-dichloro-3-[2-(5-thiomethyl-1H-tetrazol-1-yl)ethyloxy]benzoyl chloride,  
 2-methyl-3-[2-(5-methoxy-1H-tetrazol-1-yl)ethyloxy]-4-bromobenzoyl chloride,  
 2,4-dichloro-3-[3-(1H-tetrazol-1-yl)propyloxy]benzoyl chloride, and etc.

**[0038]** Representative examples of the compounds of the formula (VI) used as raw materials in the preparation of the compounds of the above formula (V) include those described below.

2,4-Dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]benzoic acid,  
 2,4-dichloro-3-[2-(2H-tetrazol-2-yl)ethyloxy]benzoic acid,  
 2-methyl-3-[2-(5-methyl-1H-tetrazol-1-yl)ethyloxy]-4-bromobenzoic acid,  
 2-methyl-3-[2-(5-methyl-2H-tetrazol-2-yl)ethyloxy]-4-bromobenzoic acid,  
 2-methyl-3-[2-(1H-tetrazol-1-yl)ethyloxy]-4-chlorobenzoic acid,  
 2,4-dichloro-3-[2-(5-thiomethyl-1H-tetrazol-1-yl)-ethyloxy] benzoic acid,  
 2-methyl-3-[2-(5-methoxy-1H-tetrazol-1-yl)ethyloxy]-4-bromobenzoic acid,  
 2,4-dichloro-3-[3-(1H-tetrazol-1-yl)propyloxy]benzoic acid, and etc.

**[0039]** Representative examples of the compounds of the formula (VII) used as raw materials in the preparation of the compounds of the above formula (VI) include those described below.

Ethyl 2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]-benzoate,  
 ethyl 2,4-dichloro-3-[2-(2H-tetrazol-2-yl)ethyloxy]-benzoate,

ethyl 2-methyl-3-[2-(5-methyl-1H-tetrazol-1-yl)-ethyloxy]-4-bromobenzoate,  
 ethyl 2-methyl-3-[2-(5-methyl-2H-tetrazol-2-yl)-ethyloxy]-4-bromobenzoate,  
 methyl 2-methyl-3-[2-(1H-tetrazol-1-yl)ethyloxy]-4-chlorobenzoate,  
 ethyl 2,4-dichloro-3-[2-(5-thiomethyl-1H-tetrazol-1-yl)ethyloxy]benzoate,  
 ethyl 2-methyl-3-[2-(5-methoxy-1H-tetrazol-1-yl)-ethyloxy]-4-bromobenzoate,  
 ethyl 2,4-dichloro-3-[3-(1H-tetrazol-1-yl)-propyloxy]benzoate, and etc.

**[0040]** Representative examples of the compounds of the formula (VIII) used as raw materials in the preparation of the compounds of the above formula (VII) include those described below.

Ethyl 2,4-dichloro-3-(2-bromoethoxy)benzoate,  
 ethyl 2-methyl-3-(2-bromoethoxy)-4-bromobenzoate,  
 methyl 2-methyl-3-(2-bromoethoxy)-4-chlorobenzoate,  
 ethyl 2,4-dichloro-3-(2-chloroethoxy)benzoate,  
 ethyl 2,4-dichloro-3-(2-iodoethoxy)benzoate,  
 ethyl 2,4-dichloro-3-(3-bromopropoxy)benzoate, and etc.

**[0041]** Representative examples of the compounds of the formula (IX) used as raw materials in the preparation of the compounds of the above formula (VII) include those described below.

1H-tetrazole,  
 5-methyl-1H-tetrazole,  
 5-ethyl-1H-tetrazole,  
 5-methylthio-1H-tetrazole,  
 5-methoxy-1H-tetrazole, and etc.

**[0042]** The compounds of the formula (Ia), which are raw materials in the above preparation process (b), constitute a part of the compounds represented by the formula (I) of the present invention and can be easily prepared in accordance with the above preparation process (a).

**[0043]** Representative examples of the compounds of the formula (Ia) used as raw materials in the preparation process (b) include those described below which are encompassed by the formula (I).

4-{2,4-Dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]-benzoyl}-1,3-dimethyl-5-hydroxypyrazole,  
 4-{2,4-dichloro-3-[2-(2H-tetrazol-2-yl)ethyloxy]-benzoyl}-1,3-dimethyl-5-hydroxypyrazole,  
 4-{2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethyloxy]-benzoyl}-1-methyl-3-cyclopropyl-5-hydroxypyrazole,  
 4-{2-methyl-3-[2-(1H-tetrazol-1-yl)ethyloxy]-4-bromobenzoyl}-1,3-dimethyl-5-hydroxypyrazole,  
 4-{2-methyl-3-[2-(2H-tetrazol-2-yl)ethyloxy]-4-bromobenzoyl}-1,3-dimethyl-5-hydroxypyrazole,  
 4-{2-methyl-3-[2-(1H-tetrazol-1-yl)ethyloxy]-4-chlorobenzoyl}-1,3-dimethyl-5-hydroxypyrazole, and etc.

**[0044]** The compounds of the formula (III), which are raw materials in the above preparation process (b), are halogenated alkyl, halogenated alkenyl, halogenated alkynyl, halogenated aralkyl, halogenated acyl, halogenated sulfonyl and acylalkyl halide that are well known themselves in the field of organic chemistry, and those described below can be exemplified.

Methyl iodide,  
 propargyl bromide,  
 benzyl bromide,  
 n-propane sulfonyl chloride,  
 4-methylbenzene sulfonyl chloride,  
 acetyl chloride,  
 phenacyl bromide and etc.

**[0045]** The reaction of the above preparation process (a) can be carried out in a suitable diluent and examples thereof that may be used during the process include aliphatic, alicyclic and aromatic hydrocarbons (which may be optionally chlorinated), for example, pentane, hexane, cyclohexane, petroleum ether, ligroin, toluene, xylene, dichloromethane, 1,2-dichloroethane;

ethers such as for example, ethyl ether, methyl ethyl ether, isopropyl ether, butyl ether, dioxane, dimethoxyethane (DME), tetrahydrofuran (THF) and diethylene glycol dimethyl ether (DGM);

ketones such as for example, acetone, methyl ethyl ketone (MEK), methyl isopropyl ketone and methyl isobutyl ketone (MIBK);

nitriles such as for example, acetonitrile and propionitrile;

alcohols such as for example, methanol, ethanol and isopropanol;

esters such as for example, ethyl acetate, amyl acetate;

acid amides such as for example, dimethylformamide (DMF), dimethylacetamide (DMA), N-methylpyrrolidone and 1,3-dimethyl-2-imidazolidinone;

sulfones and sulfoxides such as dimethylsulfoxide (DMSO) and sulfolane; and bases such as for example, pyridine.

**[0046]** The preparation process (a) can be carried out in the presence of a base. Examples of the base includes, from

inorganic bases, carbonates and bicarbonates of alkali metal and alkali earth metal such as for example, sodium bicarbonate, potassium bicarbonate, sodium carbonate and potassium carbonate; and from organic bases, tertiary amines, dialkylaminoanilines and pyridines such as for example, triethylamine, 1,1,4,4-tetramethylethylenediamine (TMEDA), N,N-dimethylaniline, N,N-diethylaniline, pyridine, 4-dimethylaminopyridine (DMAP), 1,4-diazabicyclo[2,2,2]octane (DABCO) and 1,8-diazabicyclo[5,4,0]undec-7-ene (DBU).

**[0047]** The preparation process (a) can be carried out in the presence of cyanide. Examples of the catalysts include sodium cyanide, potassium cyanide, acetone cyanohydrin and hydrogen cyanide.

**[0048]** When the above preparation process (a) is carried out in the presence of cyanide, it can also be carried out by a method using phase-transfer catalysts. Examples of a diluent that may be used during the preparation include aliphatic, alicyclic and aromatic hydrocarbons (which may be optionally chlorinated), for example, pentane, hexane, cyclohexane, petroleum ether, ligroin, toluene, xylene and dichloromethane.

ethers such as for example, ethyl ether, methyl ethyl ether, isopropyl ether, butyl ether, dioxane, dimethoxyethane (DME), tetrahydrofuran (THF) and diethylene glycol dimethyl ether (DGM); nitriles such as for example, acetonitrile and propionitrile.

**[0049]** Examples of the phase-transfer catalysts include crown ethers such as for example, dibenzo-18-crown-6, dicyclohexyl-18-crown-6, 18-crown-6 and 15-crown-5.

**[0050]** The preparation process (a) can be carried out within a substantially wide range of temperatures.

**[0051]** In general, the suitable range of temperatures is between 5°C to 200°C and preferably 25°C to 130°C. Further, the reaction is conducted desirably under normal pressure, but it can also be operated under elevated pressure or reduced pressure.

**[0052]** In carrying out the preparation process (a), 0.5 to 2 mol of potassium carbonate are reacted per mole of the compound of the formula (II) in a diluent, for example, dioxane to give the objective compound of the formula (I).

**[0053]** When the preparation process (a) is carried out in the presence of cyanide, it can be carried out within a substantially wide range of temperatures.

**[0054]** In general, the suitable range of temperatures is between -10°C to 80°C and preferably 5°C to 40°C. Further, the reaction is conducted desirably under normal pressure, but it can also be operated under elevated pressure or reduced pressure.

**[0055]** In carrying out the preparation process (a) in the presence of cyanide, 1 to 4 mol of triethylamine is reacted per mole of the compound of the formula (II) in the presence of 0.01 to 0.5 mol of acetone cyanohydrin in a diluent, for example, acetonitrile to give the objective compound of the formula (I).

**[0056]** In carrying out the preparation process (a), the compounds of the formula (I) can be obtained by starting from the compounds of the formula (VI) and continuously carrying out one pot reaction without isolating the compounds of the formula (V) and those of the formula (II).

**[0057]** The reaction of the above preparation process (b) can be carried out in a suitable diluent and examples thereof that may be used during the process include water;

aliphatic, alicyclic and aromatic hydrocarbons (which may be optionally chlorinated), for example, pentane, hexane, cyclohexane, petroleum ether, ligroin, benzene, toluene, xylene,

dichloromethane, chloroform, carbon tetrachloride, 1,2-dichloroethane, chlorobenzene, dichlorobenzene and the like;

ethers such as for example, ethyl ether, methyl ethyl ether, isopropyl ether, butyl ether, dioxane, dimethoxyethane (DME), tetrahydrofuran (THF) and diethylene glycol dimethyl ether (DGM);

ketones such as for example, acetone, methyl ethyl ketone (MEK), methyl isopropyl ketone and methyl isobutyl ketone (MIBK);

nitriles such as for example, acetonitrile, propionitrile and acrylonitrile;

esters such as for example, ethyl acetate and amyl acetate;

acid amides such as for example, dimethylformamide (DMF), dimethylacetamide (DMA), N-methylpyrrolidone and 1,3-dimethyl-2-imidazolidinone, hexamethylphosphoric triamide (HMPA);

sulfones and sulfoxides such as for example, dimethylsulfoxide (DMSO) and sulfolane; and

bases such as for example, pyridine.

**[0058]** The preparation process (b) can be carried out in the presence of acid coupling agents. The acid coupling agents include, from inorganic bases, hydrides, hydroxides, carbonates and bicarbonates of alkali metal and alkali earth metal such as for example,

sodium hydride, lithium hydride, sodium bicarbonate, potassium bicarbonate, sodium carbonate, potassium carbonate, lithium hydroxide, sodium hydroxide, potassium hydroxide and calcium hydroxide; and

from organic bases, alkalates, tertiary amines, dialkylaminoanilines and pyridines such as for example, triethylamine, 1,1,4,4-tetramethylethylenediamine (TMEDA), N,N-dimethylaniline, N,N-diethylaniline, pyridine, 4-dimethylaminopyridine (DMAP),

1,4-diazabicyclo[2,2,2]octane (DABCO) and 1,8-diazabicyclo[5,4,0]undec-7-ene (DBU).

**[0059]** The preparation process (b) can also be carried out by a process of using phase-transfer catalysts. Examples

of a diluent that may be used during the preparation include water;

aliphatic, alicyclic and aromatic hydrocarbons (which may be optionally chlorinated), for example, pentane, hexane, cyclohexane, petroleum ether, ligroin, benzene, toluene, xylene, dichloromethane, chlorobenzene, dichlorobenzene and the like; ethers such as for example, ethyl ether, methyl ethyl ether, isopropyl ether, butyl ether, dioxane, dimethoxyethane (DME), tetrahydrofuran (THF) and diethylene glycol dimethyl ether (DGM); nitriles such as for example, acetonitrile, propionitrile and acrylonitrile.

**[0060]** The examples of the phase-transfer catalysts include quaternary ions such as for example, tetramethylammonium bromide, tetrapropylammonium bromide, tetrabutylammonium bromide, tetrabutylammonium bisulfate, tetrabutylammonium iodide,

trioctylmethylammonium chloride, benzyltriethylammonium bromide, butylpyridinium bromide, heptylpyridinium bromide and benzyltriethylammonium chloride;

crown ethers such as for example, dibenzo-18-crown-6, dicyclohexyl-18-crown-6 and 18-crown-6; cryptands such as for example, [2.2.2]-cryptate, [2.1.1]-cryptate, [2.2.1]-cryptate, [2.2.B]-cryptate, [20202S]-cryptate and [3.2.2]-cryptate.

**[0061]** The preparation process (b) can be carried out within a substantially wide range of temperatures.

**[0062]** In general, the suitable range of temperatures is between -20°C to 140°C and preferably 0°C to 100°C. Further, the reaction is conducted desirably under normal pressure, but it can also be operated under elevated pressure or reduced pressure.

**[0063]** In carrying out the preparation process (b), 1 mol to 5 mol of the compound of the formula (III) is reacted per mole of the compound of the formula (Ia) in the presence of potassium carbonate in a diluent, for example, DMF to give the objective compound.

**[0064]** The active compounds of the formula (I) according to the present invention exhibit superior herbicidal activity for various weeds as shown in Biological Test Examples to be described later and can be used as herbicides. In the present specification weeds mean, in the broadest sense, all plants that grow in locations where they are undesired. The compounds of the present invention act as selective herbicides depending on the application concentration. The active compounds according to the present invention can be used, for example, between the following weeds and cultures.

**[0065]** From the genus of dicotyledonous weeds: *Sinapis*, *Capsella*, *Lepidium*, *Galium*, *Stellaria*, *Chenopodium*, *Kochia*, *Urtica*, *Senecio*, *Amaranthus*, *Portulaca*, *Xanthium*, *Ipomoea*, *Polygonum*, *Ambrosia*, *Cirsium*, *Sonchus*, *Solanum*, *Rorippa*, *Lamium*, *Veronica*, *Datura*, *Viola*, *Galeopsis*, *Papaver*, *Centaurea*, *Galinsoga*, *Rotala*, *Lindernia*, *Sesbania*, *Trifolium*, *Abutilon*, *Lamium*, *Matricaria*, *Artemisia*, *Sesbania*, *Pharbitis*, *Amaranthus*, and the like.

**[0066]** From the genus of dicotyledonous cultures: *Gossypium*, *Glycine*, *Beta*, *Daucus*, *Phaseolus*, *Pisum*, *Solanum*, *Linum*, *Ipomoea*, *Vicia*, *Nicotiana*, *Lycopersicon*, *Arachis*, *Brassica*, *Lactuca*, *Cucumis*, *Cucurbita*, and the like.

**[0067]** From the genus of monocotyledonous weeds: *Echinochloa*, *Setaria*, *Panicum*, *Digitaria*, *Phleum*, *Poa*, *Festuca*, *Eleusine*, *Lolium*, *Bromus*, *Avena*, *Cyperus*, *Sorghum*, *Agropyron*, *Monochoria*, *Fimbristylis*, *Sagittaria*, *Eleocharis*, *Scirpus*, *Paspalum*, *Ischaemum*, *Agrostis*, *Alopecurus*, *Cynodon*, *Commelina*, *Brachiaria*, *Leptochloa*, and the like.

**[0068]** From the genus of monocotyledonous cultures: *Oryza*, *Zea*, *Triticum*, *Hordeum*, *Avena*, *Secale*, *Sorghum*, *Panicum*, *Saccharum*, *Ananas*, *Asparagus*, *Allium*, and the like.

**[0069]** However, the use of the active compounds of the formula (I) of the present invention is not limited to the above species of weeds but can also be applied to other species of weeds similarly.

**[0070]** The use of the active compounds of the present invention is not limited to the above plants but can also be applied to other plants similarly. Further, the active compounds of the present invention can non-selectively control weeds and can be used, for example, in industrial areas such as factories, railway tracks, roads, plantations, non plantation and the like, depending on the application concentration. Similarly, the compounds can also be used for controlling weeds in perennial plant cultivation such as, for example, afforestations, decorative tree plantings, orchards, vineyards, citrus groves, nut orchards, banana plantations, coffee plantations, tee plantations, rubber plantations, oil palm plantations, cocoa plantations, soft fruit plantings and hopfields. The compounds of the present invention can also be applied for selectively controlling weeds in annual plant cultivation.

**[0071]** The active compounds of the present invention can be made into the customary formulations for actual use. Examples of the formulation forms include solutions, wettable powders, emulsions, suspensions, dusts, water-dispersible granules, tablets, granules, suspended emulsifiable concentrates, microcapsules in a polymer substance and etc.

**[0072]** These formulations can be produced by known methods per se. For example, they can be prepared by mixing the active compounds with extenders, namely, liquid or solid diluents or carriers and, optionally, with surfactants, namely, emulsifiers and/or dispersants and/or foam formers.

**[0073]** Examples of the liquid diluents or carriers include aromatic hydrocarbons (for example, xylene, toluene, alkyl-naphthalene and the like), chlorinated aromatic hydrocarbons or chlorinated aliphatic hydrocarbons (for example, chlorobenzenes, ethylene chlorides, methylene chloride and the like), aliphatic hydrocarbons [for example, cyclohexane and the like, paraffins (mineral oil fractions and the like)], alcohols (for example, butanol, glycol and the like), and ethers and esters thereof, ketones (for example, acetone, methyl ethyl ketone, methyl isobutyl ketone, cyclohexanone and the like), strongly polar solvents (for example, dimethylformamide, dimethylsulfoxide and the like), water, etc. In case of using water as an extender, for example, organic solvents can be used as auxiliary solvents.

**[0074]** Examples of the solid diluents may include ground natural minerals (for example, kaolins, clays, talc, chalk, quartz, attapulgite, montmorillonite or diatomaceous earth) and ground synthetic minerals (for example, highly dispersed silicic acid, alumina and silicate).

**[0075]** Examples of the solid carriers for granules may include crushed and fractionated rocks (for example, calcite, marble, pumice, sepiolite and dolomite), synthetic granules of inorganic or organic powders, and organic materials (for example, sawdust, coconut shells, maize cobs and tobacco stalks).

**[0076]** Examples of the emulsifiers and/or foam formers may include nonionic and anionic emulsifiers [for example, polyoxyethylene fatty acid esters, polyoxyethylene fatty acid alcohol ethers (for example, alkylaryl polyglycol ether), alkyl sulfonates, alkyl sulfates and aryl sulfonates] and albumin hydrolysates.

**[0077]** The dispersants include lignin sulfite waste liquor and methylcellulose.

**[0078]** Binders may also be used in the formulations (powders, granules and emulsifiable concentrates). Examples of the binders may include carboxymethyl cellulose, natural or synthetic polymers (for example, gum arabic, polyvinyl alcohol and polyvinyl acetate).

**[0079]** Colorants may also be used. Examples of the colorants may include inorganic pigments (for example, iron oxide, titanium oxide and Prussian blue), organic colorants such as Alizarin colorants, azo colorants or metal phthalocyanine colorants, and further, trace nutrients such as salts of iron, manganese, boron, copper, cobalt, molybdenum or zinc.

**[0080]** The formulations can contain the active compounds of the formula (I) in a range of generally 0.1 to 95% by weight and preferably 0.5 to 90% by weight.

**[0081]** Further, the active compounds of the formula (I) of the present invention can be used for controlling weeds as such or in their formulation forms. The active compounds of the formula (I) of the present invention can be used also in combination with known herbicides. A mixed herbicide composition with known herbicides may be previously prepared as a final formulation form, or can be prepared by tank-mixing on occasion of application.

**[0082]** For example, the following herbicides shown in common names can be exemplified as specific examples of herbicides that can be used in combination with the active compounds of the formula (I) of the present invention.

**[0083]** Sulfonylurea type herbicides: for example, chloresulfuron, sulfometuron methyl, chlorimuron ethyl, triasulfuron, amidosulfuron, oxasulfuron, tribenuron ethyl, prosulfuron, ethametsulfuron methyl, triflurosulfuron methyl, thifensulfuron methyl, flazasulfuron, rimsulfuron, nicosulfuron, flupyrsulfuron, bensulfuron methyl, pyrazosulfuron ethyl, foramsulfuron, sulfosulfuron, cinosulfuron, azimsulfuron, metsulfuronmethyl, halosulfuron methyl, ethoxysulfuron, cyclosulfamuron and iodiosulfuron;

carbamate type herbicides: for example, phenmedipham, chloroprotham, asulam, bentiocarb, molinate, esprocarb, pyributicarb, dimepiperate and swep;

chloroacetanilide type herbicides: for example, propachlor, metazachlor, alachlor, acetochlor, metolachlor, butachlor, pretilachlor and thenylchlor;

diphenylether type herbicides: for example, acifluorfen, oxyfluorfen, lactofen, fomesafen, aclonifen, chlomethoxynil, bifenox and CNP;

triazine type herbicides: for example, simazine, atrazine, propazine, cyanazine, ametryn, simetryn, dimethametryn and prometryn;

phenoxy acid or benzoic acid type herbicides: for example, 2,3,6-TBA, dicamba, quinclorac, quinmerac, clopyralid, picloram, triclopyr, fluroxypyr, fenoxaprop, diclofop methyl, fluzifop butyl, haloxyfop methyl, quizalofop ethyl, cyhalofop butyl, 2,4-PA, MCP, MCPB and phenothiol;

acid amide or urea type herbicides: for example, isoxaben, diflufenican, daiuron, linuron, fluometuron, difenoxuron, methylglyphosate, isoproturon, isouron, tebuthiuron, methabenzothiazuron, propanil, mefenacet, chlomeprop, naproanilide, bromobutide, daimuron, cumyluron, etobenzanid and oxadichlomefon;

organic phosphorous type herbicides: for example, glyphosate, bialaphos, glufosinate, amiprofos methyl, anilophos, bensulide, piperophos and butamiphos;

dinitroaniline type herbicides: for example, trifluralin and prodiamine;

phenolic type herbicides: for example, bromoxynil, ioxynil and dinoseb;

cyclohexanedione type herbicides: for example, alloxydim, sethoxydim, cloproxydim, clethodim, cycloxydim and tralkoxydim;

imidazolinone type herbicides: for example, imazametabenz, imazapyr, imazamethapyr, imazethapyr, imazamox and imazaquin;

5 bipyridium type herbicides: for example, paraquat and diquat;

carbamoyltetrazolinone type herbicides: for example, fentrazamide;

nitrile type herbicides: for example, dichlobenil; and

other herbicides: for example, bentazone, tridiphane, indanofan, amitrol, carfentrazone ethyl, sulfentrazone, fenchlorazole ethyl, isoxaflutole, clomazone, maleic acid hydrazide, pyridate, chloridazon, norflurazon, pyriothiac, bromacil, terbacil, metribuzin, oxaziclomefone, cinmethylin, flumiclorac pentyl, flumioxadin, fluthiacet methyl, azafenidin, benfuresate, oxadiazon, oxadiargyl, pentoxazone, cafenstrole, pyriminobac, bispiribac sodium, pyribenzoxim, pyrifthalid, pyraflufen ethyl, benzobicyclon, dithiopyr, dalapon, and chlorthiamid.

[0084] The active compounds listed above are known herbicides described in "Pesticide Manual," 2000, published by British Crop Protect Council.

15 [0085] Further, the active compounds of the formula (I) of the present invention can be also mixed with safeners to reduce phytotoxicity and to provide wider weed-controlling spectrum by the mixing, thereby their application as selective herbicides can be broadened.

[0086] As examples of the safeners, the following compounds shown in common names or development codes can be mentioned;

20 [0087] AD-67, BAS-145138, benoxacor, cloquintocet-methyl, cyometrinil, 2,4-D, DKA-24, dichlormid, dymron, fenclorin, fenchlorazole-ethyl, flurazole, fluxofenim, furilazole, isoxadifen-ethyl), mefenpyr-diethyl), MG-191, naphthalic anhydride, oxabetrinil, PPG-1292, R-29148 and etc.

[0088] The above safeners are also described in "Pesticide Manual" 2000, published by The British Crop Protect Council.

25 [0089] Furthermore, the mixed herbicide compositions comprising the active compounds of the formula (I) of the present invention and the above herbicides can be further mixed with the above safeners to reduce phytotoxicity and to provide wider weed-control spectrum by the mixing, thereby their application as selective herbicides can be broadened.

[0090] Surprisingly, some of the mixed herbicide compositions of the compounds according to the present invention with known herbicides and/or safeners show synergetic effects.

30 [0091] In case of using the active compounds of the formula (I) of the present invention, they can be directly used as such or used in formulation forms such as formulated solutions for spraying, emulsifiable concentrates, tablets, suspensions, powders, granules or used in the use forms prepared by further dilution. The active compounds of the present invention can be applied, for example, watering, spraying, atomizing, scattering granules, and etc.

[0092] The active compounds of the formula (I) of the present invention can be used at any stage before and after germination of plants. Further, they can be also taken into the soil before sowing.


35 [0093] The application amount of the active compounds of the formula (I) can vary and fundamentally differs depending on the nature of the desired effect. In case of using the active compounds as herbicides, examples of the application amount, in terms of the active compounds, can be in the range of 0.005 to 4 kg per hectare and preferably 0.01 to 2 kg per hectare.

40 [0094] The preparation and application of the compounds of the present invention will be described more specifically by way of the following Examples.

<Compound Examples>

45 Synthesis Example 1

[0095]



5-{2,4-Dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]-benzoyloxy}-1,3-dimethylpyrazole (0.65 g) was dissolved in acetonitrile (10 ml), triethylamine (0.33 g) and acetone cyanohydrin (40 mg) were added thereto and the mixture was stirred at room


## EP 2 081 925 B1

temperature for 12 hours. The solvent was distilled off, acidified by the addition of diluted hydrochloric acid and extracted with ethyl acetate (150 ml). The organic layer was washed with saturated brine and dried with anhydrous magnesium sulfate. Ethyl acetate was distilled off to obtain objective 4-{2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoyl}-1,3-dimethyl-5-hydroxy-pyrazole (0.59 g).

mp: 63-68°C.

### Synthesis Example 2

#### [0096]




4-{2-Methyl-3-[2-(1H-tetrazol-1-yl)ethoxy]-4-bromobenzoyl}-1,3-dimethyl-5-hydroxypyrazole (0.40 g) and potassium carbonate (0.15 g) were suspended in DMF (3 ml), benzyl bromide (0.17 g) was added thereto and the mixture was stirred at room temperature for 5 hours. After the reaction, cool water was added, the mixture was extracted with ethyl acetate (100 ml) and dried over anhydrous magnesium sulfate. The residue obtained by distilling off ethyl acetate was purified with silica gel column chromatography (elution solvent; ethyl acetate : hexane = 1 : 1) to obtain the desired 5-{2-methyl-3-[2-(1H-tetrazol-1-yl)ethoxy]-4-bromobenzoyloxy}-1,3-dimethyl-5-benzyloxy-pyrazole (0.37 g).


mp: 122-125°C.

[0097] The compounds obtained by operation according to the preparation processes of the compounds of the present invention exemplified in the above Synthesis Examples 1 and 2 are shown in the following Table 1 together with the compounds synthesized in Synthesis Examples 1 and 2.

[0098] In Table 1, T1 represents a group:




and T2 represents a group:



[0099] Further, Me represents methyl, Et represents ethyl, Pr-n represents n-propyl, cyclo-Pro represents cyclopropyl, OMe represents methoxy, SMe represents methylthio, SO<sub>2</sub>Me represents methanesulfonyl (methylsulfonyl), Ph represents phenyl and Ph-4-Me represents 4-methylphenyl.

Table 1



Compound No.	X	Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	n	T	R <sup>4</sup>	mp (°C) or n <sub>D</sub> (°C)
1	Cl	Cl	Me	Me	H	2	T1	H	63-68
2	Cl	Cl	Me	Me	Me	2	T1	H	
3	Cl	Cl	Me	Me	allyl	2	T1	H	
4	Cl	Cl	Me	Me	propargyl	2	T1	H	
5	Cl	Cl	Me	Me	benzyl	2	T1	H	
6	Cl	Cl	Me	Me	SO <sub>2</sub> -Me	2	T1	H	
7	Cl	Cl	Me	Me	SO <sub>2</sub> -Pr-n	2	T1	H	
8	Cl	Cl	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T1	H	
9	Cl	Cl	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T1	H	
10	Cl	Cl	Me	Me	C(=O)Me	2	T1	H	
11	Cl	Cl	Me	Me	C(=O)Ph	2	T1	H	
12	Cl	Cl	Me	Me	H	3	T1	H	53-63
13	Cl	Cl	Me	Et	H	2	T1	H	
14	Cl	Cl	Me	cyclo-Pr	H	2	T1	H	179-181
15	Cl	Cl	Et	Me	H	2	T1	H	1.569(20)
16	Cl	Cl	Et	Et	H	2	T1	H	
17	Cl	Cl	Et	cyclo-Pr	H	2	T1	H	
18	Cl	Cl	Me	Me	H	2	T1	Me	142-151
19	Cl	Cl	Me	cyclo-Pr	H	2	T1	Me	136-143
20	Cl	Cl	Me	Me	H	2	T1	Et	
21	Cl	Cl	Me	Me	H	2	T1	OMe	
22	Cl	Cl	Me	Me	H	2	T1	SMe	
23	Br	Br	Me	Me	H	2	T1	H	62-70
24	Br	Br	Me	cyclo-Pr	H	2	T1	H	146-154
25	Br	Br	Me	Me	H	2	T1	Me	55-64
26	Br	Br	Me	cyclo-Pr	H	2	T1	Me	150-154
27	Me	Cl	Me	Me	H	2	T1	H	135-140
28	Me	Cl	Me	Me	Me	2	T1	H	
29	Me	Cl	Me	Me	allyl	2	T1	H	
30	Me	Cl	Me	Me	propargyl	2	T1	H	
31	Me	Cl	Me	Me	benzyl	2	T1	H	
32	Me	Cl	Me	Me	SO <sub>2</sub> -Me	2	T1	H	
33	Me	Cl	Me	Me	SO <sub>2</sub> -Pr-n	2	T1	H	
34	Me	Cl	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T1	H	
35	Me	Cl	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T1	H	
36	Me	Cl	Me	Me	C(=O)Me	2	T1	H	
37	Me	Cl	Me	Me	C(=O)Ph	2	T1	H	
38	Me	Cl	Me	Me	H	3	T1	H	
39	Me	Cl	Me	Et	H	2	T1	H	
40	Me	Cl	Me	cyclo-Pr	H	2	T1	H	
41	Me	Cl	Et	Me	H	2	T1	H	
42	Me	Cl	Et	Et	H	2	T1	H	

EP 2 081 925 B1

(continued)

	Compound No.	X	Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	n	T	R <sup>4</sup>	mp (°C) or n <sub>D</sub> (°C)
	43	Me	Cl	Et	cyclo-Pr	H	2	T1	H	
5	44	Me	Cl	Me	Me	H	2	T1	Me	183-188
	45	Me	Cl	Me	Me	SO <sub>2</sub> -Pr-n	2	T1	Me	
	46	Me	Cl	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T1	Me	
	47	Me	Cl	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T1	Me	
10	48	Me	Cl	Me	Me	H	3	T1	Me	
	49	Me	Cl	Me	Et	H	2	T1	Me	
	50	Me	Cl	Me	cyclo-Pr	H	2	T1	Me	
	51	Me	Cl	Et	Me	H	2	T1	Me	
	52	Me	Cl	Et	Et	H	2	T1	Me	
15	53	Me	Cl	Et	cyclo-Pr	H	2	T1	Me	
	54	Me	Cl	Me	Me	H	2	T1	Et	
	55	Me	Cl	Me	Me	H	2	T1	OMe	
	56	Me	Cl	Me	Me	H	2	T1	SMe	
20	57	Me	Br	Me	Me	H	2	T1	H	153-166
	58	Me	Br	Me	Me	SO <sub>2</sub> -Pr-n	2	T1	H	
	59	Me	Br	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T1	H	
	60	Me	Br	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T1	H	
	61	Me	Br	Me	Me	H	3	T1	H	
25	62	Me	Br	Me	Et	H	2	T1	H	
	63	Me	Br	Me	cyclo-Pr	H	2	T1	H	151-156
	64	Me	Br	Et	Me	H	2	T1	H	108-115
	65	Me	Br	Et	Et	H	2	T1	H	
	66	Me	Br	Et	cyclo-Pr	H	2	T1	H	
30	67	Me	Br	Me	Me	H	2	T1	Me	183-184
	68	Me	Br	Me	Me	Me	2	T1	Me	
	69	Me	Br	Me	Me	allyl	2	T1	Me	
	70	Me	Br	Me	Me	propargyl	2	T1	Me	
35	71	Me	Br	Me	Me	benzyl	2	T1	Me	122-125
	72	Me	Br	Me	Me	SO <sub>2</sub> -Me	2	T1	Me	
	73	Me	Br	Me	Me	SO <sub>2</sub> -Pr-n	2	T1	Me	1.5475(24)
	74	Me	Br	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T1	Me	1.5668(24)
	75	Me	Br	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T1	Me	1.7445(24)
40	76	Me	Br	Me	Me	C(=O)Me	2	T1	Me	
	77	Me	Br	Me	Me	C(=O)Ph	2	T1	Me	
	78	Me	Br	Me	Me	H	3	T1	Me	
	79	Me	Br	Me	Et	H	2	T1	Me	
45	80	Me	Br	Me	cyclo-Pr	H	2	T1	Me	1.5506(20)
	81	Me	Br	Et	Me	H	2	T1	Me	166-171
	82	Me	Br	Et	Et	H	2	T1	Me	
	83	Me	Br	Et	cyclo-Pr	H	2	T1	Me	
	84	Me	Br	Me	Me	H	2	T1	Et	136-144
50	85	Me	Br	Me	Me	H	2	T1	OMe	
	86	Me	Br	Me	Me	H	2	T1	SMe	
	87	Cl	Cl	Me	Me	H	2	T2	H	
	88	Cl	Cl	Me	Me	SO <sub>2</sub> -Pr-n	2	T2	H	
55	89	Cl	Cl	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T2	H	
	90	Cl	Cl	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T2	H	
	91	Cl	Cl	Me	Et	H	2	T2	H	
	92	Cl	Cl	Me	cyclo-Pr	H	2	T2	H	

## EP 2 081 925 B1

(continued)

	Compound No.	X	Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	n	T	R <sup>4</sup>	mp (°C) or n <sub>D</sub> (°C)
	93	Cl	Cl	Et	Me	H	2	T2	H	
5	94	Cl	Cl	Et	Et	H	2	T2	H	
	95	Cl	Cl	Et	cyclo-Pr	H	2	T2	H	
	96	Cl	Cl	Me	Me	H	2	T2	Me	1.5807(20)
	97	Cl	Cl	Me	cyclo-Pr	H	2	T2	Me	1.5852(20)
10	98	Cl	Cl	Me	Me	H	2	T2	Et	
	99	Cl	Cl	Me	Me	H	2	T2	OMe	
	100	Cl	Cl	Me	Me	H	2	T2	SMe	
	101	Br	Br	Me	Me	H	2	T2	H	1.6034(24)
	102	Br	Br	Me	cyclo-Pr	H	2	T2	H	1.5986(24)
15	103	Br	Br	Me	Me	H	2	T2	Me	1.5905(24)
	104	Br	Br	Me	cyclo-Pr	H	2	T2	Me	47-55
	105	Me	Cl	Me	Me	H	2	T2	H	
	106	Me	Cl	Me	Me	Me	2	T2	H	
20	107	Me	Cl	Me	Me	allyl	2	T2	H	
	108	Me	Cl	Me	Me	propargyl	2	T2	H	
	109	Me	Cl	Me	Me	benzyl	2	T2	H	
	110	Me	Cl	Me	Me	SO <sub>2</sub> -Me	2	T2	H	
	111	Me	Cl	Me	Me	SO <sub>2</sub> -Pr-n	2	T2	H	
25	112	Me	Cl	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T2	H	
	113	Me	Cl	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T2	H	
	114	Me	Cl	Me	Me	C(=O)Me	2	T2	H	
	115	Me	Cl	Me	Me	C(=O)Ph	2	T2	H	
30	116	Me	Cl	Me	Et	H	2	T2	H	
	117	Me	Cl	Me	cyclo-Pr	H	2	T2	H	
	118	Me	Cl	Et	Me	H	2	T2	H	
	119	Me	Cl	Et	Et	H	2	T2	H	
	120	Me	Cl	Et	cyclo-Pr	H	2	T2	H	
35	121	Me	Cl	Me	Me	H	2	T2	Me	1.5524(24)
	122	Me	Cl	Me	Me	SO <sub>2</sub> -Pr-n	2	T2	Me	
	123	Me	Cl	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T2	Me	
	124	Me	Cl	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T2	Me	
	125	Me	Cl	Me	Et	H	2	T2	Me	
40	126	Me	Cl	Me	cyclo-Pr	H	2	T2	Me	
	127	Me	Cl	Et	Me	H	2	T2	Me	
	128	Me	Cl	Et	Et	H	2	T2	Me	
	129	Me	Cl	Et	cyclo-Pr	H	2	T2	Me	
45	130	Me	Cl	Me	Me	H	2	T2	Et	
	131	Me	Cl	Me	Me	H	2	T2	OMe	
	132	Me	Cl	Me	Me	H	2	T2	SMe	
	133	Me	Br	Me	Me	H	2	T2	H	
	134	Me	Br	Me	Me	SO <sub>2</sub> -Pr-n	2	T2	H	124-142
50	135	Me	Br	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T2	H	
	136	Me	Br	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T2	H	
	137	Me	Br	Me	Et	H	2	T2	H	
	138	Me	Br	Me	cyclo-Pr	H	2	T2	H	1.5818(24)
55	139	Me	Br	Et	Me	H	2	T2	H	119-122
	140	Me	Br	Et	Et	H	2	T2	H	
	141	Me	Br	Et	cyclo-Pr	H	2	T2	H	
	142	Me	Br	Me	Me	H	2	T2	Me	143-147


EP 2 081 925 B1

(continued)

Compound No.	X	Y	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	n	T	R <sup>4</sup>	mp (°C) or n <sub>D</sub> (°C)
143	Me	Br	Me	Me	Me	2	T2	Me	
144	Me	Br	Me	Me	allyl	2	T2	Me	
145	Me	Br	Me	Me	propargyl	2	T2	Me	
146	Me	Br	Me	Me	benzyl	2	T2	Me	
147	Me	Br	Me	Me	SO <sub>2</sub> -Me	2	T2	Me	
148	Me	Br	Me	Me	SO <sub>2</sub> -Pr-n	2	T2	Me	
149	Me	Br	Me	Me	SO <sub>2</sub> -Ph-4-Me	2	T2	Me	
150	Me	Br	Me	Me	CH <sub>2</sub> C(=O)Ph	2	T2	Me	
151	Me	Br	Me	Me	C(=O)Me	2	T2	Me	
152	Me	Br	Me	Me	C(=O)Ph	2	T2	Me	
153	Me	Br	Me	Et	H	2	T2	Me	
154	Me	Br	Me	cyclo-Pr	H	2	T2	Me	1.5772(24)
155	Me	Br	Et	Me	H	2	T2	Me	1.5705(24)
156	Me	Br	Et	Et	H	2	T2	Me	
157	Me	Br	Et	cyclo-Pr	H	2	T2	Me	
158	Me	Br	Me	Me	H	2	T2	Et	147-156
159	Me	Br	Me	Me	H	2	T2	OMe	
160	Me	Br	Me	Me	H	2	T2	SMe	
161	Me	Br	Me	cyclo-Pr	H	2	T2	Et	1.5602(24)

Reference Example 1


[0100]



2,4-Dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoyl chloride (0.60 g) was dissolved in tetrahydrofuran (5 ml), this solution was added dropwise to a tetrahydrofuran (5 ml) solution of 1,3-dimethyl-5-hydroxypyrazole (0.22 g) and triethylamine (0.23 g) at 5°C and the mixture was stirred at room temperature for 6 hours. After the reaction, the mixture was extracted with ethyl acetate (100 ml) and then washed with diluted hydrochloric acid and an aqueous sodium bicarbonate solution and dried over anhydrous magnesium sulfate. Ethyl acetate was distilled off to obtain the desired 5-{2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]-benzoyloxy}-1,3-dimethylpyrazole (0.65 g).  
mp: 138-140°C.

Reference Example 2

[0101]




2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoic acid (0.46 g) and thionyl chloride (0.36 g) were added in 1,2-dichlo-

roethane (10 ml), two drops of DMF were added and the mixture was refluxed by heating for 3 hours. After cooling, the solvent was distilled off to obtain the desired 2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoyl chloride (0.49 g). mp: 69-73°C.

5 Reference Example 3

[0102]



10  
15  
20  
25  
[0103] To a methanol (15 ml) solution of ethyl 2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]benzoate (0.88 g) were added sodium hydroxide (0.16 g) and water (5 ml), and the mixture was stirred at room temperature for 7 hours. After the addition of water (20 ml) and concentration at reduced pressure, an aqueous sodium hydroxide solution was added and the mixture was washed with ethyl acetate (50 ml). The aqueous layer was acidified with hydrochloric acid and extracted with ethyl acetate. The organic layer was washed with saturated brine and dried over anhydrous magnesium sulfate. Ethyl acetate was distilled off to obtain the desired 2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]-benzoic acid (0.69 g). mp: 145-146°C.

Reference Example 4

25

[0104]



30  
35  
40  
1H-tetrazol (0.45 g) and ethyl 3-(2-bromoethoxy)-2,4-dichlorobenzoate (2.00 g) were suspended in DMF (15 ml), potassium carbonate (1.21 g) and sodium iodide (0.04 g) were added thereto and the mixture was stirred at 80°C for 3 hours. After the reaction, cool water was added, extracted with ethyl acetate (100 ml) and dried over anhydrous magnesium sulfate. The residue obtained by distilling off ethyl acetate was purified with silica gel column chromatography (elution solvent; ethyl acetate : hexane = 1 : 1) to obtain the desired ethyl 2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy] benzoate (0.88 g) and ethyl 2,4-dichloro-3-[2-(1H-tetrazol-2-yl)ethoxy]benzoate (0.99 g).

[0105] Ethyl 2,4-dichloro-3-[2-(1H-tetrazol-1-yl)ethoxy]-benzoate;  $n_D^{20} = 1.5535$ . Ethyl 2,4-dichloro-3-[2-(1H-tetrazol-2-yl)ethoxy]-benzoate;  $n_D^{20} = 1.5483$ .


<Biological Test Examples>

45

Comparative compound

[0106]

50



Sulcotrione (common name)

55

## EP 2 081 925 B1

### Test Example 1

**[0107]** Test for herbicidal efficacy against field weeds (Pre-emergence soil spray treatment) Preparation of formulation of the active compound

Carrier: DMF 5 parts by weight

Emulsifier: Benzyloxy polyglycol ether 1 part by weight

**[0108]** A suitable formulation of an active compound is obtained as an emulsifiable concentrate by mixing 1 part by weight of the active compound with the stated amounts of the carrier and emulsifier. A prescribed amount of the formulation is diluted with water.

**[0109]** In a green house, the seeds of field weeds (*Amaranthus lividus* and *Setaria viridis*) were inoculated by one species per pot on the surface layer in 16 cm<sup>2</sup> pots filled with field soil (sand loam) and covered with soil. Immediately after the inoculation, the prescribed and diluted solutions of the formulations of respective active compounds prepared according to the above preparation method, were sprayed to the soil. After 2 weeks from the treatment, the herbicidal efficacy of respective active compounds was examined. The evaluation of the herbicidal efficacy was carried out by rating complete death as 100% and as 0% in case of no herbicidal efficacy. In case of showing herbicidal efficacy of 80% or more, such active compounds are evaluated to be practical as herbicides. As typical examples, the study results with Compound Nos. 19 and 96 and those with Comparative compound C-1 are shown in Table 2 below.

Table 2: Pre-emergence soil spray treatment

Compound No.	Amount of drug (g ai/ha)	<i>Setaria viridis</i>	<i>Amaranthus lividus</i>
19	500	80	90
96	500	90	80
C-1	500	30	70
Note: ai = active ingredient			

### Test Example 2

Test for herbicidal efficacy against field weeds (After-emerge stalk and leave spray treatment)

**[0110]** In a green house, young plants (in 2- to 3-leafstages) of field weeds (*Setaria viridis*) were transplanted to 16 cm<sup>2</sup> pots filled with field soil (sand loam) and the prescribed and diluted solutions of the formulations of respective active compounds, prepared according to the Test Example 1 above were sprayed from the upside of the plants. After 2 weeks from the treatment, the herbicidal efficacy of respective active compounds was examined. The evaluation of the herbicidal efficacy was carried out in the same manner as in Test Example 1 mentioned above. As typical examples, the study results with Compound Nos. 19 and 96 and Comparative compound C-1 are shown in Table 3 below.

Table 3: After-emergence stalk and leaf spray treatment

Compound No.	Amount of drug (g ai/ha)	<i>Setaria viridis</i>
19	500	90
96	500	90
C-1	500	40

### Test Example 3

Test for selective herbicidal efficacy against field crops (Pre-emergence soil spray treatment)

**[0111]** In a green house, seeds of field crops [*Zea mays* (maize) and *Glycine max* (soy beans)] and field weeds (*Echinochloa crus-galli*, *Digitaria ciliaris*, *Amaranthus lividus* and *Chenopodium album*) were inoculated on the surface layer in 16 cm<sup>2</sup> pots filled with field soil (sand loam) and covered with soil. After one day, the prescribed and diluted solutions of the formulations of the active compound, prepared according to the Test Example 1 above were sprayed to the soil. After 2 weeks from the treatment, phytotoxicity to the crops and herbicidal efficacy of the active compound were examined. The evaluation of the herbicidal efficacy and crop phytotoxicity was carried out by rating complete death as 100% and as 0% in case of no herbicidal efficacy and no crop phytotoxicity. In case of showing herbicidal efficacy

of 80% or more, such active compounds are evaluated to be practical as herbicides and in case of showing phytotoxicity of 20% or less, such active compounds are evaluated as being superior in safety as herbicides. As typical examples, the study results with Compound No. 12 are shown in Table 4 below.

Table 4: Pre-emerge soil spray treatment

Compound No.	Amount of drug (g ai/ha)	Maize	Soy beans
12	320	0	10

(continued)

<i>Echinochloa crus-galli</i>	<i>Digitaria ciliaris</i>	<i>Amaranthus lividus</i>	<i>Chenopodium album</i>
80	80	85	95

<Formulation Examples>

Formulation Example 1 (granules)

**[0112]** To a mixture of Compound No. 14 (10 parts) of the present invention, bentonite (montmorillonite) (30 parts), talc (58 parts) and ligninsulfonate (2 parts) is added water (25 parts), and the mixture is well kneaded, made in granules of 10-40 mesh by an extrusion granulator, which are dried at 40-50°C to obtain granules.

Formulation Example 2 (granules)

**[0113]** Clay mineral particles (95 parts) having particle size distribution of 0.2-2 mm are charged in a rotary mixer, Compound No.14 (5 parts) of the present invention was sprayed together with a liquid diluent under rotation, wetted uniformly and then dried at 40-50°C to obtain granules.

Formulation Example 3 (emulsifiable concentrate)

**[0114]** Compound No.103 (30 parts) of the present invention, xylene (55 parts), polyoxyethylene alkylphenyl ether (8 parts) and calcium alkylbenzene sulfonate (7 parts) were mixed by stirring to obtain emulsions.

Formulation Example 4 (wetable powder)


**[0115]** Compound No.154 (15 parts) of the present invention, a mixture of white carbon (hydrous amorphous silicon oxide fine powder) and powder clay (1 : 5) (80 parts), sodium alkylbenzene sulfonate (2 parts) and a polymer (3 parts) of sodium alkyl naphthalene sulfonate formalin polymer were mixed in powder form and made into to wettable powders.

Formulation Example 5 (water-dispersible granules)

**[0116]** Compound No.104 (20 parts) of the present invention, sodium ligninsulfonate (30 parts), bentonite (15 parts) and calcined diatomaceous earth powder (35 parts) are well mixed, water was added thereto and the mixture was extruded with a screen of 0.3 mm and dried to obtain water-dispersible granules.

## Claims

1. Benzoylpyrazoles represented by the formula (I):



10 wherein

R<sup>1</sup> represents C<sub>1-6</sub> alkyl;

R<sup>2</sup> represents C<sub>1-6</sub> alkyl or C<sub>3-7</sub>-cycloalkyl;


R<sup>3</sup> represents hydrogen, C<sub>1-6</sub> alkyl, C<sub>3-7</sub> alkenyl, C<sub>3-7</sub> alkynyl, C<sub>7-12</sub> aralkyl, C<sub>1-6</sub> alkylsulfonyl, optionally C<sub>1-6</sub> alkyl-substituted phenylsulfonyl, C<sub>1-6</sub> alkylcarbonyl, benzoyl or benzoyl-C<sub>1-6</sub> alkyl;

X represents halogen or C<sub>1-6</sub> alkyl;


Y represents halogen;

n represents 2 or 3; and

T represents a group T1 or a group T2:



(T1)



(T2)

25 wherein R<sup>4</sup> represents hydrogen, C<sub>1-6</sub> alkyl, C<sub>1-6</sub> alkoxy or C<sub>1-6</sub> alkylthio.

30 2. The compound according to Claim 1, wherein

R<sup>1</sup> represents methyl or ethyl;

R<sup>2</sup> represents methyl, ethyl or cyclopropyl;


R<sup>3</sup> represents hydrogen, methyl, allyl, propargyl, benzyl, methanesulfonyl, n-propanesulfonyl, optionally methyl-substituted phenylsulfonyl, acyl, benzoyl or phenacyl;

X represents chlorine, bromine or methyl;


Y represents chlorine or bromine;

n represents 2 or 3; and

T represents a group T1 or a group T2:



(T1)




(T2)


45 wherein R<sup>4</sup> represents hydrogen, methyl, ethyl, methoxy or methylthio.

50 3. A herbicide comprising, as an effective ingredient, the compound according to Claim 1 or 2.

55 4. The compound represented by the formula (II):



wherein X, Y, n and T have the same meaning as defined in Claim 1 or 2, and Z represents a group:




wherein R<sup>1</sup> and R<sup>2</sup> have the same meaning as defined in Claim 1 or 2.

20

### Patentansprüche

1. Benzoylpyrazole gemäß Formel (I):



in der

R<sup>1</sup> C<sub>1-6</sub>-Alkyl bedeutet;

R<sup>2</sup> C<sub>1-6</sub>-Alkyl oder C<sub>3-7</sub>-Cycloalkyl bedeutet;



R<sup>3</sup> Wasserstoff, C<sub>1-6</sub>-Alkyl, C<sub>3-7</sub>-Alkenyl, C<sub>3-7</sub>-Alkynyl, C<sub>7-12</sub>-Aralkyl, C<sub>1-6</sub>-Alkylsulfonyl, gegebenenfalls C<sub>1-6</sub>-alkylsubstituiertes Phenylsulfonyl, C<sub>1-6</sub>-Alkylcarbonyl, Benzoyl oder Benzoyl-C<sub>1-6</sub>-alkyl bedeutet;

X Halogen oder C<sub>1-6</sub>-Alkyl bedeutet;

Y Halogen bedeutet;

n 2 oder 3 bedeutet; und

T eine Gruppe T1 oder eine Gruppe T2 bedeutet:



in der R<sup>4</sup> Wasserstoff, C<sub>1-6</sub>-Alkyl, C<sub>1-6</sub>-Alkoxy oder C<sub>1-6</sub>-Alkylthio bedeutet.

2. Verbindung nach Anspruch 1, in der

R<sup>1</sup> Methyl oder Ethyl bedeutet;

R<sup>2</sup> Methyl, Ethyl oder Cyclopropyl bedeutet;

R<sup>3</sup> Wasserstoff, Methyl, Allyl, Propargyl, Benzyl, Methansulfonyl, n-Propansulfonyl, gegebenenfalls durch Methyl substituiertes Phenylsulfonyl, Acyl, Benzoyl oder Phenacyl bedeutet;


X Chlor, Brom oder Methyl bedeutet;  
 Y Chlor oder Brom bedeutet;  
 n 2 oder 3 bedeutet; und  
 T eine Gruppe T1 oder eine Gruppe T2 bedeutet:




in der R<sup>4</sup> Wasserstoff, Methyl, Ethyl, Methoxy oder Methylthio bedeutet.

3. Herbizid, das als Wirkstoff die Verbindung nach Anspruch 1 oder 2 umfasst.

4. Verbindung gemäß Formel (II):




in der X, Y, n und T dieselbe Bedeutung wie in Anspruch 1 oder 2 definiert aufweisen und Z eine Gruppe



bedeutet, in der R<sup>1</sup> und R<sup>2</sup> dieselbe Bedeutung wie in Anspruch 1 oder 2 definiert aufweisen.

### Revendications

1. Benzoylpyrazoles représentés par la formule (I) :



dans laquelle


R<sup>1</sup> représente C<sub>1-6</sub>alkyle ;

R<sup>2</sup> représente C<sub>1-6</sub>alkyle ou C<sub>3-7</sub>cycloalkyle ;


R<sup>3</sup> représente hydrogène, C<sub>1-6</sub>alkyle, C<sub>3-7</sub>alcényle, C<sub>3-7</sub>alcynyle, C<sub>7-12</sub>aralkyle, C<sub>1-6</sub>alkylsulfonyle, phénylsulfonyle éventuellement substitué par C<sub>1-6</sub>alkyle, C<sub>1-6</sub>alkylcarbonyle, benzoyle ou benzoyl-C<sub>1-6</sub>alkyle ;

X représente halogène ou C<sub>1-6</sub>alkyle ;  
 Y représente halogène ;  
 n représente 2 ou 3 ; et  
 T représente un groupement T1 ou un groupement T2 :

5



(T1)



(T2)

10

dans lesquels R<sup>4</sup> représente hydrogène, C<sub>1-6</sub>alkyle, C<sub>1-6</sub>alcoxy ou C<sub>1-6</sub>alkylthio.


15

2. Composé selon la revendication 1, caractérisé en ce que


R<sup>1</sup> représente méthyle ou éthyle ;  
 R<sup>2</sup> représente méthyle, éthyle ou cyclopropyle ;  
 R<sup>3</sup> représente hydrogène, méthyle, allyle, propargyle, benzyle, méthanesulfonyle, n-propanesulfonyle, phénylsulfonyle éventuellement substitué par méthyle, acyle, benzoyle ou phénacyle ;  
 X représente chlore, brome ou méthyle ;  
 Y représente chlore ou brome ;  
 n représente 2 ou 3 ; et  
 T représente un groupement T1 ou un groupement T2 :

20

25



(T1)



(T2)

30


dans lesquels R<sup>4</sup> représente hydrogène, méthyle, éthyle, méthoxy ou méthylthio.

35

3. Herbicide comprenant, à titre d'ingrédient efficace, le composé selon la revendication 1 ou 2.

4. Composé représenté par la formule (II) :


40




45

dans laquelle X, Y, n et T ont les mêmes significations que définies dans la revendication 1 ou 2, et Z représente un groupement :

50



ou



55

**EP 2 081 925 B1**

dans lesquels  $R^1$  et  $R^2$  sont tels que définis dans la revendication 1 ou 2.

5

10

15

20

25

30

35

40

45

50

55

**REFERENCES CITED IN THE DESCRIPTION**

*This list of references cited by the applicant is for the reader's convenience only. It does not form part of the European patent document. Even though great care has been taken in compiling the references, errors or omissions cannot be excluded and the EPO disclaims all liability in this regard.*

**Patent documents cited in the description**

- WO 0366607 A [0002] [0030] [0031] [0032]
- WO 0110850 A [0002]
- WO 0153275 A [0002]
- JP 2000173 A [0028] [0030]
- WO 9318031 A [0028] [0030]
- JP 61257974 A [0029]
- JP 10109976 A [0029]
- JP 2173 A [0031]
- DE 10039723 [0032]
- JP 6247891 A [0033]
- JP 7206808 A [0033]
- EP 369803 A [0033]
- JP 9309883 A [0034]

**Non-patent literature cited in the description**

- *Journal of the American Chemical Society*, 1958, vol. 80, 3908-3911 [0034]
- *Journal of Organic Chemistry*, 1957, vol. 22, 1142-1145 [0034]
- *Journal of Organic Chemistry*, 1991, vol. 56, 2395-2400 [0034]
- Pesticide Manual. British Crop Protect Council, 2000 [0084]