



US00PP17778P2

(12) **United States Plant Patent**
Roberson

(10) **Patent No.:** **US PP17,778 P2**

(45) **Date of Patent:** **Jun. 5, 2007**

(54) **LANTANA PLANT NAMED 'ROBPATLOL'**

(52) **U.S. Cl.** **Plt./227**

(50) Latin Name: *Lantana compara*
Varietal Denomination: **Robpatlol**

(58) **Field of Classification Search** **Plt./227**
See application file for complete search history.

(76) Inventor: **Robert J. Roberson**, P.O. Box 210,
Grain Valley, MO (US) 64029

Primary Examiner—Kent Bell
Assistant Examiner—Annette H Para
(74) *Attorney, Agent, or Firm*—C. A. Whealy

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 32 days.

(57) **ABSTRACT**

A new and distinct cultivar of *Lantana* plant named
'Robpatlol', characterized by its upright and outwardly
spreading plant habit; uniformly mounded plant form; semi-
compact growth habit; short internodes; dense and bushy
habit; freely and continuously flowering habit; and flowers
that are initially yellow and become red purple in color.

(21) Appl. No.: **11/185,457**

(22) Filed: **Jul. 20, 2005**

(51) **Int. Cl.**
A01H 5/00 (2006.01)

2 Drawing Sheets

1

2

Botanical designation: *Lantana camara*.
Cultivar denomination: 'Robpatlol'.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct culti-
var of *Lantana* plant, botanically known as *Lantana camara*,
and hereinafter referred to by the cultivar name Robpatlol.

The new *Lantana* is a product of a planned breeding
program conducted by the Inventor in Grain Valley, Mo. The
objective of the breeding program is to create new semi-
compact and freely-flowering *Lantanas* with attractive
flower coloration.

The new *Lantana* originated from a cross-pollination
made by the Inventor in August, 2001 of the *Lantana*
cultivar Robpatpas, disclosed in U.S. Plant Pat. No. 13,424,
as the female, or seed, parent with the *Lantana* cultivar
Robcomplan, disclosed in U.S. Plant Pat. No. 9,837, as the
male, or pollen, parent. The new *Lantana* was selected as a
single plant from the resulting progeny of the cross-
pollination by the Inventor in a controlled environment in
Grain Valley, Mo., on the basis of its semi-compact growth
habit and attractive flower coloration.

Asexual reproduction of the new cultivar by terminal
cuttings in a controlled environment in Grain Valley, Mo.,
since September, 2002, has shown that the unique features
of this new *Lantana* are stable and reproduced true to type
in successive generations.

SUMMARY OF THE INVENTION

Plants of the cultivar Robpatlol have not been observed
under all possible environmental conditions. The phenotype
may vary somewhat with variations in environment and
culture such as temperature and light intensity, however, any
variance in genotype.

The following traits have been repeatedly observed and
are determined to be the unique characteristics of 'Rob-
patlol'. These characteristics in combination distinguish
'Robpatlol' as a new and distinct *Lantana* cultivar:

1. Upright and outwardly spreading plant habit; uniformly
mounded plant form.

2. Semi-compact growth habit.
3. Short internodes, dense and bushy habit.
4. Freely and continuously flowering habit.
5. Flowers that are initially yellow and become red purple
in color.

Plants of the new *Lantana* can be compared to plants of
the parents. In side-by-side comparisons conducted in Grain
Valley, Mo., plants of the new *Lantana* differed from plants
of the female parent, the cultivar Robpatpas, in the following
characteristics:

1. Plants of the new *Lantana* were more compact and had
shorter internodes than plants of the cultivar Robpat-
pas.
2. Plants of the new *Lantana* were more freely branching
and denser than plants of the cultivar Robpatpas.
3. Plants of the new *Lantana* had smaller leaves than
plants of the cultivar Robpatpas.

In side-by-side comparisons conducted in Grain Valley,
Mo., plants of the new *Lantana* differed from plants of the
male parent, the cultivar Robcomplan, in the following
characteristics:

1. Plants of the new *Lantana* were not as compact as and
had longer internodes than plants of the cultivar Rob-
complan.
2. Plants of the new *Lantana* and the cultivar Robcomplan
differed in flower coloration as plants of the cultivar
Robcomplan had yellow flowers that turned orange in
color before becoming red purple.

Plants of the new *Lantana* can also be compared to plants
of the *Lantana* cultivar Robpathot, disclosed in U.S. Plant
Pat. No. 12,266. In side-by-side comparisons conducted in
Grain Valley, Mo., plants of the new *Lantana* differed from
plants of the cultivar Robpathot in the following character-
istics:

1. Plants of the new *Lantana* were more compact, denser
and had shorter internodes than plants of the cultivar
Robpathot.
2. Plants of the new *Lantana* had smaller leaves than
plants of the cultivar Robpathot.
3. Plants of the new *Lantana* and the cultivar Robpathot
differed in flower color.

BRIEF DESCRIPTION OF THE PHOTOGRAPHS

The accompanying colored photographs illustrate the overall appearance of the new cultivar, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photographs may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new *Lantana*. Plants used for the photographs were grown in an outdoor nursery in Grain Valley, Mo. for about three months.

The photograph on the first sheet comprises a side perspective view of a typical flowering plant of 'Robpatlol' grown in the landscape for about three months.

The photograph on the second sheet comprises a close-up view of typical inflorescences and leaves of 'Robpatlol'.

DETAILED BOTANICAL DESCRIPTION

In the following description, color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used. Plants used for the description were planted in containers after rooting and grown for about 3 months during the spring in an outdoor nursery in Tolar, Tex. During the production of the plants in the greenhouse, temperatures ranged from about 18° C. to about 35° C.

Botanical classification: *Lantana camara* cultivar Robpatlol.
Parentage:

Female, or seed, parent.—*Lantana camara* cultivar Robpatpas, disclosed in U.S. Plant Pat. No. 13,424.

Male, or pollen, parent.—*Lantana camara* cultivar Robcomplan, disclosed in U.S. Plant Pat. No. 9,837.

Propagation:

Type cutting.—Terminal cuttings.

Time to initiate roots, summer.—About 14 days at 27° C.

Time to initiate roots, winter.—About 16 days at 27° C.

Time to develop roots, summer.—About 28 days at 24° C.

Time to develop roots, winter.—About 38 days at 24° C.
Root description.—Fine, fibrous; initially glaucous white in color then becoming closer to 161D with development.

Rooting habit.—Freely branching.

Plant description:

Form.—Flowering subshrub; upright and outwardly spreading plant habit; uniformly mounded plant form; compact growth habit. Freely branching; two lateral branches potentially forming at every node; pinching enhances lateral branch development.

Plant height.—About 20 cm.

Plant diameter.—About 52 cm.

Lateral branches.—Length: About 23 cm. Diameter: About 4 mm. Internode length: About 2.3 cm. Strength: Strong, but flexible. Texture: Rough, pubescent. Color: Young: Close to 144A. Woody: Close to 199A.

Foliage description.—Arrangement: Opposite; simple. Length: About 6.7 cm. Width: About 4.4 cm. Shape: Ovate. Apex: Acute. Base: Obtuse with truncate tendencies. Margin: Serrate. Texture, upper and lower surfaces: Leathery, rough, coarse; pubescent. Luster: Upper surface: Slightly glossy. Lower surface: Dull. Venation pattern: Pinnate, arcuate. Color: Developing and fully expanded foliage, upper surface: Close to 147A. Developing and fully expanded

foliage, lower surface: Darker green than 147B. Venation, upper surface: Close to 146A. Venation, lower surface: Close to 147C to 147D. Petiole length: About 1.6 cm. Petiole diameter: About 2 mm. Petiole texture, both surfaces: Slightly pubescent. Petiole color, upper surface: Close to 146A. Petiole color, lower surface: Close to 147C to 147D.

Flower description:

Flower type and habit.—Small salverform flowers arranged in axillary umbels; flowers face mostly upward or outward. Flowers self-cleaning. Very freely flowering with potentially two inflorescences per node; typically about 24 flowers per umbel.

Natural flowering season.—Spring until frost in the autumn; flowering continuous and consistent.

Flower longevity on the plant.—About one week.

Fragrance.—Faint, pleasant.

Inflorescence diameter.—About 5.1 cm.

Inflorescence height.—About 2.75 cm.

Flowers.—Appearance: Flared trumpet, corolla fused, four-parted; flowers roughly rectangular in shape. Diameter: About 1.4 cm by 1.3 cm. Corolla tube length: About 1.2 cm.

Flower buds (before showing color).—Length: About 8 mm. Diameter: About 7 mm. Shape: Roughly spherical to ovoid. Color: Close to 144A.

Corolla.—Arrangement/appearance: Single whorl of four petals, fused into flared trumpet. Petal length from throat: Upper and lower petals: About 6 mm. Lateral petals: About 5 mm. Petal width: Upper and lower petals: About 8 mm. Lateral petals: About 5 mm. Petal shape: Spatulate to somewhat orbicular. Petal apex: Rounded. Petal margin: Entire. Petal lobe texture, upper and lower surfaces: Smooth, velvety. Corolla throat and tube texture: Pubescent. Color: Petal lobes, when opening, upper surface: Close to 12A. Petal lobes, when opening, lower surface: Close to 12B. Petal lobes, fully opened, upper surface: Close to 57A; color fading to 69A overlain with 57A with development. Petal lobes, fully opened, lower surface: Close to 58B to 58C. Throat: Close to 9A and 12A. Tube: Close to 58B to 58C.

Calyx.—Quantity: One sepal per flower. Length: About 7.5 mm. Width: About 2 mm. Shape: Lanceolate. Apex: Acute. Base: Truncate. Texture, upper and lower surface: Pubescent. Color, upper and lower surfaces: Close to 144A.

Peduncles.—Length: About 3.5 cm. Diameter: About 1 mm. Angle: About 45° from the stem. Strength: Flexible, but strong. Texture: Pubescent. Color: Close to 144A.

Pedicels.—Not observed, flowers not stalked.

Reproductive organs.—Stamens: Quantity/arrangement: Four per flower, adnate to floral tube. Anther shape: Oblong. Anther length: Less than 1 mm. Anther color: Close to 9A. Pollen amount: None observed. Pistils: Quantity: One per flower. Pistil length: About 2 mm. Stigma shape: Rounded. Stigma color: Close to 144B to 144C. Ovary color: Close to 144A.

Fruit.—Quantity: One per flower. Diameter: About 6 mm. Shape: Spherical. Texture: Smooth. Color: Close to 202A.

Disease/pest resistance: Plants of the new *Lantana* grown in the garden have not been noted to be resistant to pathogens and pests common to *Lantana*.

5

Weather tolerance: Plants of the new *Lantana* have been observed to be very tolerant to rain and wind.
Temperature tolerance: Plants of the new *Lantana* have been observed to be tolerant to temperatures ranging from 0° C. to 38° C.

6

It is claimed:

1. A new and distinct cultivar of *Lantana* plant named 'Robpatlol', as illustrated and described.

* * * * *



