PCT WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C11D 3/50, C11B 9/00, A61K 7/46	A1	(11) International Publication Number: **WO 97/06234**
		(43) International Publication Date: 20 February 1997 (20.02.97)

(21) International Application Number: PCT/JP96/02176

(22) International Filing Date: 2 August 1996 (02.08.96)

(30) Priority Data:
7/199637 4 August 1995 (04.08.95) JP

(71) Applicant *(for all designated States except US)*: KAO CORPORATION [JP/JP]; 14-10, Nihonbashikayabacho 1-chome, Chuo-ku, Tokyo 103 (JP).

(72) Inventors; and
(75) Inventors/Applicants *(for US only)*: KOHAMA, Makoto [JP/JP]; Kao Corporation, Research Laboratories, 1334, Minato, Wakayama-shi, Wakayama 640 (JP). KOSHINO, Junji [JP/JP]; Kao Corporation, Research Laboratories, 1334, Minato, Wakayama-shi, Wakayama 640 (JP). FUKUDA, Kazuyuki [JP/JP]; Kao Corporation, Research Laboratories, 1-3, Bunka 2-chome, Sumida-ku, Tokyo 131 (JP). TOI, Nao [JP/JP]; Kao Corporation, Research Laboratories, 1-3, Bunka 2-chome, Sumida-ku, Tokyo 131 (JP).

(74) Agents: ARUGA, Mitsuyuki et al.; Kyodo Building, 3-6, Nihonbashiningyocho 1-chome, Chuo-ku, Tokyo 103 (JP).


(81) Designated States: CN, US, VN, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).

Published
With international search report.

(54) Title: PERFUME BASE COMPOSITION

(57) Abstract

The present invention relates to a perfume base composition comprising a carrier and a benzyl-substituted cyclohexanol having formula (1), wherein the dashed line denotes an optional double bond and the benzyl group is in the 2-, 3- or 4-position of the cyclohexanol ring, with the proviso that the benzyl group is in the 4-position when a double bond is present in the cyclohexanol ring. The composition has not only a well-balanced fragrance but also excellent retentivity of fragrance.



(1)

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	**GB**	United Kingdom	**MW**	Malawi
AT	Austria	**GE**	Georgia	**MX**	Mexico
AU	Australia	**GN**	Guinea	**NE**	Niger
BB	Barbados	**GR**	Greece	**NL**	Netherlands
BE	Belgium	**HU**	Hungary	**NO**	Norway
BF	Burkina Faso	**IE**	Ireland	**NZ**	New Zealand
BG	Bulgaria	**IT**	Italy	**PL**	Poland
BJ	Benin	**JP**	Japan	**PT**	Portugal
BR	Brazil	**KE**	Kenya	**RO**	Romania
BY	Belarus	**KG**	Kyrgystan	**RU**	Russian Federation
CA	Canada	**KP**	Democratic People's Republic of Korea	**SD**	Sudan
CF	Central African Republic			**SE**	Sweden
CG	Congo	**KR**	Republic of Korea	**SG**	Singapore
CH	Switzerland	**KZ**	Kazakhstan	**SI**	Slovenia
CI	Côte d'Ivoire	**LI**	Liechtenstein	**SK**	Slovakia
CM	Cameroon	**LK**	Sri Lanka	**SN**	Senegal
CN	China	**LR**	Liberia	**SZ**	Swaziland
CS	Czechoslovakia	**LT**	Lithuania	**TD**	Chad
CZ	Czech Republic	**LU**	Luxembourg	**TG**	Togo
DE	Germany	**LV**	Latvia	**TJ**	Tajikistan
DK	Denmark	**MC**	Monaco	**TT**	Trinidad and Tobago
EE	Estonia	**MD**	Republic of Moldova	**UA**	Ukraine
ES	Spain	**MG**	Madagascar	**UG**	Uganda
FI	Finland	**ML**	Mali	**US**	United States of America
FR	France	**MN**	Mongolia	**UZ**	Uzbekistan
GA	Gabon	**MR**	Mauritania	**VN**	Viet Nam

DESCRIPTION

PERFUME BASE COMPOSITION

TECHNICAL FIELD

The present invention relates to a perfume base composition which excellently retains its fragrance. This composition can be used for the production of perfume and other products to be scented such as detergents, cosmetics and sprays.

BACKGROUND ART


A great number of perfume base materials having a floral fragrance have heretofore been known. Among others, aldehyde type perfume bases are the most important perfume base materials. However, since aldehydes are not always stable in fragrant products having different pHs and product forms, these products often do not smell fresh and natural. Moreover, these products do not retain their fragrance for a very long time, which is disadvantageous.

It is accordingly the object of the present invention to provide a perfume base composition which has a floral fragrance and high stability and can impart excellent retentivity of fragrance to blending systems.

DISCLOSURE OF THE INVENTION

This object is achieved by the use of specific benzyl-substituted cyclohexanols, which compounds can impart freshness to the perfume base composition and the products made therefrom. Due to the use of these benzyl-substituted cyclohexanols the articles made from the perfume base composition additionally have a natural, soft and voluminous feeling.

According to the present invention, there is thus provided a perfume base composition comprising a benzyl-substituted cyclohexanol having the following formula (1):



(1)

wherein the dashed line denotes an optional double bond and the benzyl group is in the 2-, 3- or 4-position of the cyclohexanol ring, with the proviso that the benzyl group is in the 4-position when a double bond is present in the cyclohexanol ring.

BEST MODE FOR CARRYING OUT THE INVENTION

The benzyl-substituted cyclohexanols (1) are known compounds. For example, 2-benzylcyclohexanol is described


in Tetrahedron, 48, 2059 (1992), Tetrahedron Lett., 36, 123 (1994) and J. Chem. Soc., 1809 (1956); 3-benzylcyclohexanol is described in J. Chem. Soc., 1809 (1956); and 4-benzylcyclohexanol and 4-benzyl-2-cyclohexen-1-ol are described in Tetrahedron, 48, 2059 (1992). These documents, however, do not disclose the use of benzylcyclohexanols as a fragrance. Moreover, the prior art does not suggest to use these compounds to retain the fragrance of perfume base compositions and of the products made therefrom.

Examples of the benzyl-substituted cyclohexanols (1) useful in the practice of the present invention include 2-benzylcyclohexanol, 3-benzylcyclohexanol, 4-benzylcyclohexanol and 4-benzyl-2-cyclohexen-1-ol. Of these, 2-benzylcyclohexanol, 3-benzylcyclohexanol and 4-benzylcyclohexanol are preferred with 3-benzylcyclohexanol being particularly preferred.

Cis-trans isomerism exists in the benzyl-substituted cyclohexanols (1) according to the substitution state between the benzyl group and the hydroxyl group on the cyclohexane ring (or cyclohexene ring). In the present invention both cis- and trans-benzylcyclohexanols and mixtures thereof may be used. Among these isomers, the cis-form is particularly preferred.


The benzyl-substituted cyclohexanols (1) can be prepared in accordance with the processes described in the above literature. For example, 2-benzylcyclohexanol (1a) can be prepared by subjecting benzaldehyde (2) and

cyclohexanone (3) to aldol condensation and hydrogenating the resulting enone (4) (see the following reaction scheme).




(2) (3) (4) (1a)

3-Benzylcyclohexanol (1b) can be prepared by subjecting a Grignard reagent (5) prepared from a benzyl halide, and 2-cyclohexen-1-one (6) to 1,4-addition and hydrogenating the resulting ketone (7) (see the following reaction scheme).



(5) (6) (7) (1b)

4-Benzylcyclohexanol (1c) and 4-benzyl-2-cyclohexen-1-ol (1d) can be prepared by using an enamine (9) obtained by the dehydration-condensation of 3-phenylpropionaldehyde (8) and morpholine to form 4-benzyl-2-cyclohexen-1-ol (10) by Robinson annellation and selectively hydrogenating only the carbonyl of the compound (10) to form 4-benzyl-2-cyclohexen-1-ol (1d) or hydrogenating both olefin and carbonyl of the compound (10) to form 4-benzylcyclohexanol (1c) (see the following reaction scheme).



(8) (9) (10) (1d) (1c)

Since most of the benzyl-substituted cyclohexanols (1) obtained in such a manner are mixtures of cis- and trans-forms, their isomers have to be separated from each other by column chromatography, recrystallization from a hydrocarbon solvent, superfractionation and/or the like if necessary.

The benzyl-substituted cyclohexanols (1) may be used as a deodorant component either singly or in combination with a carrier. Any carrier may be used without any limitation so far as it does not impair the fragrance of the benzyl-substituted cyclohexanols (1). Examples thereof include gaseous, liquid and solid carriers, which may contain other perfume compounds. The examples of the preferable gaseous carriers include a gas for propelland agents. The preferable liquid carriers include water, various organic solvents, and volatile oily substances. The preferable solid carriers include solid oily substances such as various waxes, and polymers. The amount of the benzyl-substituted cyclohexanol (1) to be incorporated in the perfume base composition according to the present invention varies according to the kind of the formulated perfume base used in combination, the kind and intesity of the intended fragrance, and the like, and no particular limitation is imposed on the amount. However, it is generally preferable to use it in a proportion of 0.1-90 wt.%, particularly preferably 0.5-70 wt.% based on the perfume base composition.

In addition to the benzyl-substituted cyclohexanol (1) and the carrier, optional components routinely incorporated in perfume base compositions may be incorporated into the perfume base composition according to the present invention within limits not impeding the effect of the present invention.

The perfume base composition according to the present invention can be obtained by mixing and stirring the benzyl-substituted cyclohexanol (1), carrier and optional components in accordance with a method known per se in the art.

The perfume base composition according to the present invention may be suitably applied to products required to be scented, such as perfumes, detergents, cosmetics, various sprays and fragrances, in particular, toiletry products such as soap, shampoos and rinses.

The perfume base composition according to the present invention has a well-balanced fragrance and is also excellent in retentivity of fragrance.

EXAMPLES

The present invention will hereinafter be described in more detail by the following examples.

Preparation Example 1:

A 1-liter four-necked flask equipped with a Dean-Stark trap and a thermometer was charged with 71 g of morpholine

and 400 ml of toluene, and 100 g of 3-phenylpropionaldehyde were added dropwise thereto while chilling with ice water, followed by azeotropic dehydration for 1 hour. After cooling the reaction mixture, excess morpholine was distilled off with toluene. The residue was dissolved in 400 ml of toluene. In a 1-liter four-necked flask equipped with a thermometer and a condenser, 70 g of methyl vinyl ketone were added dropwise under reflux to the solution over 1 hour. After aging under reflux for 1 hour, the reaction mixture was cooled, and 10% sulfuric acid was added dropwise to the reaction mixture until the pH of the reaction mixture reached 3. The thus-acidified reaction mixture was then stirred at room temperature for 1.5 hours. After separating the resulting lower water layer, the residual organic layer was neutralized, washed with water, dried and concentrated, thereby obtaining 137 g of a crude product. Dissolved in 500 ml of methanol were 137 g of the crude product, and 4 ml of 4.5 mol/liter aqueous KOH were added dropwise to the solution at room temperature, followed by stirring for 3 hours at the same temperature. After neutralizing the resultant mixture with acetic acid, methanol was distilled off. After the residue was charged into a 1-liter four-necked flask equipped with a Dean-Stark trap and a thermometer to dissolve it in 500 ml of toluene, 2 g of p-toluenesulfonic acid were added to the solution, followed by azeotropic dehydration for 3 hours. After cooling, the residue was neutralized and washed with water, and the

resultant organic layer was dried and concentrated, thereby obtaining 119 g of a crude product. This product was purified by column chromatography to obtain 41 g (yield: 30%) of 4-benzyl-2-cyclohexen-1-one.

To a suspension of 3.2 g of lithium aluminum hydride in 500 ml of ether, 30 g of 4-benzyl-2-cyclohexen-1-one were added dropwise at 0°C. After stirring the mixture for 2 hours at the same temperature, it was washed with diluted hydrochloric acid and water, and the resultant organic layer was dried and concentrated, thereby obtaining 31 g of a crude product. This product was purified by column chromatography to obtain 30 g (yield: 98%) of 4-benzyl-2-cyclohexen-1-ol. A ratio of a cis-form to a trans-form was found to be 32:68 by ^{1}H-NMR.

4-Benzyl-2-cyclohexen-1-ol:

^{1}H-NMR δ: 1.17-2.77(8H,m), 4.10-4.30(1H,m), 5.1-5.35(2H,m)
7.1-7.35(5H,m).

Besides, 30 g of the above-obtained 4-benzyl-2-cyclohexen-1-one were hydrogenated at room temperature for 6 hours using 5 g of Raney nickel in methanol under hydrogen pressure (3 kg/cm^2) until the absorption of hydrogen was stopped. After an organic layer was collected by decantation and concentrated, the residue was purified by column chromatography, thereby obtaining 13 g of cis-4-benzylcyclo-hexanol and 14 g of trans-4-benzylcyclohexanol (yield: 90% in total).

Cis-4-benzylcyclohexanol:

^{1}H-NMR δ: 1.30-1.80(10H,m), 2.54(2H,d,J=6.97Hz),
3.90-4.01(1H,br), 7.08-7.32(5H,m).

Trans-4-benzylcyclohexanol:

^{1}H-NMR δ: 0.87-1.30(4H,m), 1.35-1.62(1H,m), 1.63-1.85(2H,m),
1.85-2.03(3H,m), 2.47(2H,d,J=7.1Hz),
3.42-3.62(1H,m), 7.08-7.35(5H,m).

Preparation Example 2:

A 500-ml four-necked flask equipped with a thermometer and a condenser was charged with 2.4 g of a piece of magnesium and 120 ml of absolute ether, to which a part of 17 g of benzyl bromide was added with heating. After a reaction started, the remaining amount of benzyl bromide was added dropwise to such a degree that reflux continued. After completion of the drop addition, the resultant mixture was stirred further for 1 hour at room temperature to obtain a liquid reaction mixture. A 500-ml four-necked flask equipped with a thermometer was charged with a suspension of 1 g of copper iodide in 60 ml of absolute ether, to which the liquid reaction mixture was added dropwise at -5°C. After stirring the resultant mixture for 1 hour at the same temperature, it was cooled to -20°C. A solution of 8.6 g of 2-cyclohexen-1-one in 10 ml of absolute ether was added dropwise to the mixture, and the temperature of the resulting liquid reaction mixture was then raised to room temperature while stirring for 24 hours. The liquid reaction mixture was washed with water, dried, filtered and concentrated, and 21 g of the resulting crude product were

purified by column chromatography, thereby obtaining 9.6 g (yield: 57%) of 3-benzylcyclohexenone.

Then, 9.6 g of 3-benzylcyclohexenone were hydrogenated at room temperature using 1 g of Raney nickel in methanol under hydrogen pressure (3 kg/cm^2) until the absorption of hydrogen was stopped. After an organic layer was collected by decantation and concentrated, the residue was purified by column chromatography, thereby obtaining 4.5 g of cis-3-benzylcyclohexanol and 4.5 g of trans-3-benzylcyclohexanol (yield: 92% in total).

Cis-3-benzylcyclohexanol:

^{1}H-NMR δ: 0.9-1.15(1H,m), 1.17-1.82(8H,m), 1.87-2.15(1H,m), 2.55(2H,d), 7.1-7.4(1H,m), 7.12-7.35(5H,m).

Trans-3-benzylcyclohexanol:

^{1}H-NMR δ: 0.75-1.35(4H,m), 1.45-1.85(4H,m), 1.85-2.03(2H,m), 2.24-2.62(2H,m), 3.42-3.62(1H,m), 7.08-7.35(5H,m).

Preparation Example 3:

A 500-ml four-necked flask equipped with a thermometer was charged with 53 g of benzaldehyde, 25 ml of water and 1 g of sodium hydroxide, to which 68 g of cyclohexanone were added dropwise at 30°C over 30 minutes. The resultant mixture was stirred for 1 hour at the same temperature. After neutralizing the reaction mixture to separate liquid layers from each other, the resultant reaction product was dissolved in 500 ml of toluene. To the solution, 0.5 g of PTS (p-toluenesulfonic acid) was added to conduct azeotropic dehydration for 3 hours in a 1-liter four-necked flask

equipped with a Dean-Stark trap and a thermometer. After cooling, the residue was neutralized and washed with water, and the resultant organic layer was dried and concentrated, thereby obtaining 119 g of a crude product. This product was purified by column chromatography to obtain 24 g (yield: 28%) of 2-benzylidenecyclohexanone.

Then, 10 g of 2-benzylidenecyclohexanone were hydrogenated at room temperature using 1 g of Raney nickel in methanol under hydrogen pressure (3 kg/cm^2) until the absorption of hydrogen was stopped. After an organic layer was collected by decantation and concentrated, the residue was purified by column chromatography, thereby obtaining 4.6 g of cis-2-benzylcyclohexanol and 4.4 g of trans-2-benzyl-cyclohexanol (yield: 88% in total).

Cis-2-benzylcyclohexanol:

^{1}H-NMR δ: 1.1-1.35(1H,m), 1.35-1.85(9H,m),
2.53(1H,dd,J=7.7,13.4Hz),
2.70(1H,dd,J=9.2,13.3Hz), 3.75-3.82(1H,bs),
7.12-7.35(5H,m).

Trans-2-benzylcyclohexanol:

^{1}H-NMR δ: 0.8-1.85(9H,m), 1.91-2.08(1H,m),
2.34(1H,dd,J=9.2,13.3Hz),
3.17(1H,dd,J=4.0,13.3Hz), 3.28-3.4(1H,m),
7.12-7.35(5H,m).

The fragrance of the benzyl-substituted cyclohexanols (1) obtained in the above-described preparation examples are shown in Table 1.

4-benzyl-2-cyclohexen-1-ol, 2-benzyl-cyclohexanol, 3-benzylcyclohexanol and 4-benzylcyclohexanol will hereinafter be referred to as "1d", "1a", "1b" and "1c", respectively. Numerals in (:) following each of the symbols indicate the porportions of the cis-form and the trans-form. For example, 1c (0:100) means that the ratio of the cis-form to the trans-form in 4-benzylcyclohexanol is 0:100.

Table 1

Compound	Odor
1d (32:68)	Floral-Green-Muguet Rose-Geranium
1c (100:0)	Floral-Rose-Geranium
1c (0:100)	Floral-Green-Sweet
1b (100:0)	Floral-Grapefruit-Woody-Vetiver
1b (0:100)	Floral-Grapefruit-Vetiver
1a (100:0)	Floral-Grapefruit
1a (0:100)	Floral-Grapefruit

Example 1:

1c (100:0) or the like were added to a rose base having a composition shown in Table 2 and the effect brought about by the addition was evaluated in accordance with an organoleptic test by 5 expert panelists.

Organoleptic evaluation standard:

1: The effect of the addition was recognized to a very desirable extent;

2: The effect of the addition was recognized to a

desirable extent;

3: The effect of the addition was scarcely recognized;

4: No effect of the addition was recognized, and the balance of fragrance was poor.

Table 2

	Invention product 1	Comparative product 1	Comparative product 2
Rose base:			
Geraniol	400	400	400
Phenylethyl alcohol	80	80	80
Geranyl acetate	80	80	80
Geranium oil	50	50	50
Rhodinol	50	50	50
α-Ionone	50	50	50
Diphenyl oxide	40	40	40
Linalol	30	30	30
1c (100:0)	220	-	-
ℓ-Citronellol	-	220	-
Dipropylene glycol	-	-	220
Total (parts by weight)	1000	1000	1000
Organoleptic evaluation	1	2	4

The invention product 1 had a voluminous, well-balanced, rose-like fragrance. The comparative product 1 had a rose-like fragrance, but was lacking in a voluminous feeling. The comparative product 2 had an ill-balanced fragrance not associated with roses.

Example 2:

A composition having a gorgeous, soft, voluminous, lily of the valley-like fragrance was obtained in accordance

with the following formulation.

(Components)	(parts by weight)
Phenylethyl alcohol	250
Hydroxycitronellal	250
Bergamot oil	100
Jasmine oil	100
Heliotropin	100
Linalyl benzoate	50
Phenylethyl acetate	10
Cinnamic alcohol	10
1c (100:0)	130
Total	1000

Example 3:

A floral perfume base for soap, which had a soft, sweet fragrance, was obtained in accordance with the following formulation. In particular, voluminous sweetness was recognized during its use.

(Components)	(parts by weight)
Bois de Rose oil	250
Terpineol	150
Lavender oil	100
Cederwood oil	100
Citronella oil	150
Eugenol	50
Linalyl acetate	50
Diphenyl oxide	30
Pearlide BB (product of Kao Corporation)	70

Styrax resinoid	20
1c (0:100)	80
Total	1000

Example 4:

Added to a grapefruit base having a composition shown in Table 3 were 1a (50:50) or 1b (50:50), thereby obtaining grapefruit type perfume base compositions. These compositions were organoleptically evaluated.

Table 3

	Invention product 2	Invention product 3
Grapefruit base:		
Orange oil	718	718
Orange terpeneless oil	20	20
Fruitate*1	20	20
Pollenal II*2	20	20
Linalol	10	10
Ethyl octanoate	10	10
2-Methyl-4-propyl-1,3-oxathiane	2	2
1a (50:50)	200	-
1b (50:50)	-	200
Total (parts by weight)	1000	1000

*1: Kao's specialty chemical; ethyl-bicyclo[5.2.1.$0^{2,6}$]-decane-2-carboxylate.
*2: Kao's specialty chemical; 2-cyclohexylpropanal.

As a result, the invention product 2 was found to have a grapefruit type natural fragrance with better softness. On the other hand, the invention product 3 was found to have a grapefruit type, natural, fresh,

voluminous, and long-lasting fragrance. Of these, the formulated fragrance of the invention product 3, in which 1b (50:50) had been incorporated, was very excellent from the viewpoint of freshness.

Example 5:

1a (50:50) or 1b (50:50) were added to a citron base having the composition shown in Table 4 to obtain citron type perfume base compositions. These compositions were organoleptically evaluated.

Table 4


	Invention product 4	Invention product 5
Citron base:		
Orange oil	298	298
Mandarin oil	150	150
γ-Terpinene	100	100
Linalol	200	200
Dihydromyrcenol	50	50
Dimethyloctenone	50	50
Dihydrojasmonate	50	50
2-Methyl-4-propyl-1,3-oxathiane	2	2
1a (50:50)	100	-
1b (50:50)	-	100
Total (parts by weight)	1000	1000

As a result, the invention product 4 was found to be a citron type natural perfume base composition having better softness. On the other hand, the invention product 5 was found to be a natural, fresh citron type perfume base

composition feeling like the rind of a citron. Of these, the formulated fragrance of the invention product 5, in which 1b (50:50) had been incorporated, was very excellent from the viewpoint of freshness.

CLAIMS

1. A perfume base composition comprising a carrier and a benzyl-substituted cyclohexanol having the following formula (1):



(1)

wherein the dashed line denotes an optional double bond and the benzyl group is in the 2-, 3- or 4-position of the cyclohexanol ring, with the proviso that the benzyl group is in the 4-position when a double bond is present in the cyclohexanol ring.

2. The perfume base composition according to Claim 1, wherein the benzyl-substituted cyclohexanol is a cis- or trans-benzylcyclohexanol or a mixture thereof.


3. The perfume base composition according to Claim 1 or 2, wherein the benzyl-substituted cyclohexanol is a cis-benzylcyclohexanol.

4. The perfume base composition according to Claim 3, wherein the benzyl-substituted cyclohexanol is 2-benzylcyclohexanol, 3-benzylcyclohexanol or 4-benzyl-2-cyclohexen-1-ol.

5. The perfume base composition according to Claim 3, wherein the benzyl-sustituted cyclohexanol is 3-benzylcyclohexanol.

6. The perfume base composition according to any one of Claim 1, wherein the content of the benzyl-substituted cyclohexanol is 0.1-90 wt.% based on the perfume base composition.

7. Use of a benzyl-substituted cyclohexanol having the following formula (1):



(1)

wherein the dashed line denotes an optional double bond and the benzyl group is in the 2-, 3- or 4-position of the cyclohexanol ring, with the proviso that the benzyl group is in the 4-position when a double bond is present in the cyclohexanol ring as a fragrance.

8. The use according to Claim 7 for the production of toiletry, perfumes, detergents, cosmetics and sprays.

INTERNATIONAL SEARCH REPORT

Intern al Application No
PCT/JP 96/02176

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C11D3/50 C11B9/00 A61K7/46

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C11D C11B A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,A	EP,A,0 717 100 (INT FLAVORS & FRAGRANCES INC) 19 June 1996 see page 2, line 48 - line 50; claim 1	1-8
A	U.S. PUBLIC HEALTH REPORTS, vol. 63, 1948, pages 339-346, XP002011893 J.M. BRENNAN: "Field tests with tick repellents" see page 346, paragraph 5; tables 1,2 see page 344, line 4 see page 344, paragraph 3	1

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search	Date of mailing of the international search report
21 October 1996	14. 11. 96

Name and mailing address of the ISA	Authorized officer
European Patent Office, P.B. 5818 Patentlaan 2 NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax (+31-70) 340-3016	Loiselet-Taisne, S

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

..ormation on patent family members

Interna al Application No
PCT/JP 96/02176

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0717100	19-06-96	US-A- 5527768	18-06-96