



(43) International Publication Date
7 February 2013 (07.02.2013)

(10) International Publication Number
WO 2013/017522 A1

- (51) International Patent Classification:
A61K 38/00 (2006.01) *C07K 14/435* (2006.01)
- (21) International Application Number:
PCT/EP2012/064690
- (22) International Filing Date:
26 July 2012 (26.07.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
11176531.9 4 August 2011 (04.08.2011) EP
61/574,596 4 August 2011 (04.08.2011) US
- (71) Applicant (for all designated States except US): **MERZ PHARMA GMBH & CO. KGAA** [DE/DE]; Eckenheimer Landstraße 100, 60318 Frankfurt am Main (DE).
- (72) Inventor; and
- (75) Inventor/Applicant (for US only): **FREVERT, Jürgen** [DE/DE]; Weimarer Stasse 6, c/o Vorderhaus / Etage 2, 10625 Berlin (DE).
- (74) Agent: **DICK, Alexander**; Herzog Fiesser & Partner, Dudenstrasse 46, 68167 Mannheim (DE).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))



WO 2013/017522 A1

(54) Title: ALTERATION OF PROTEOLYTIC CLEAVAGE OF BOTULINUM NEUROTOXINS

(57) Abstract: The present invention pertains to a polynucleotide encoding a modified neurotoxin polypeptide comprising a modified neurotoxin light chain and a heavy chain, said modified light chain having at least one modification conferring altered cleavage by calpain proteases. Further encompassed by the present invention are vectors and host cells comprising the polynucleotide of the invention as well as polypeptides encoded by the said polynucleotide. In addition, the invention relates to compositions comprising the polynucleotide, vector, host cell or polypeptide of the invention as a medicament.

[0004] Despite its toxic effects, BoNTs have been used as therapeutic agents in a large number of diseases. BoNT serotype A (BoNT/A) was approved for human use in the United States in 1989 for the treatment of strabism, blepharospasm, and other disorders. It is commercially available as a protein preparation, for example, under the tradename BOTOX (Allergan Inc.) or under the tradename DYSPORT (Ipsen Ltd.). In these
5 preparations, the neurotoxin is integrated in a protein complex with so called complexing proteins. For therapeutic application, the complex is injected directly into the muscle to be treated. At physiological pH, the toxin is released from the protein complex to exert the desired pharmacological effect. An improved BoNT/A preparation being free of
10 complexing proteins is available under the tradename XEOMIN (Merz Pharmaceuticals GmbH).

[0005] BoNTs, in principle, weaken voluntary muscle strength and are, therefore, effective therapeutic agents for the therapy of diseases such as strabism, focal dystonia,
15 including cervical dystonia, and benign essential blepharospasm. They have been further shown to relief hemifacial spasm, and focal spasticity, and moreover, to be effective in a wide range of other indications, such as gastrointestinal disorders, hyperhidrosis, and cosmetic wrinkle correction, see Jost 2007, Drugs 67, 669.

[0006] However, the effect of BoNTs is only temporary, which is the reason why repeated administration of BoNTs may be required to maintain a therapeutic effect. Moreover, the more frequently these drugs are applied the higher will be the risk for an adverse immune response against the neurotoxin applied. Further, some patients develop anti-neurotoxin antibodies and, thereby, become non-responders to therapies by
25 conventional BoNTs. BoNTs are in some indications applied only locally. However, their diffusion potential makes a controlled local application difficult. In general, the production of these highly toxic polypeptides is cumbersome and needs special care with respect to safety issues and thus is expensive. In light of these drawbacks of conventional neurotoxin polypeptides as drugs, means for controlling and/or improving the biological activity of
30 neurotoxins would be highly appreciated for the neurotoxin therapy.

[0007] Thus, the technical problem underlying the present invention could be seen as the provision of means and methods which comply with the aforementioned needs. This technical problem has been solved by the embodiments characterized in the claims and
35 herein below.

[0008] The present invention relates to a polynucleotide encoding a modified neurotoxin polypeptide comprising a modified neurotoxin light chain and a heavy chain, said modified

light chain having at least one modification conferring altered cleavage by calpain proteases.

[0009] The term “neurotoxin” as used herein means a Clostridial molecule which is capable of interfering with the functions of a cell, including a neuron. Preferably, the neurotoxin is a polynucleotide encoding the neurotoxin polypeptide or a neurotoxin polypeptide. The interfered cell function can be exocytosis. The neurotoxin can be naturally occurring or recombinant. Active di-chain neurotoxin polypeptide consists of two chains, an N-terminal light chain of approx. 50 kDa and a heavy chain of approx. 100 kDa linked by a disulfide bond. Neurotoxins structurally consist of three domains, i.e. the catalytic light chain, the heavy chain encompassing the translocation domain (N-terminal half) and the receptor binding domain (C-terminal half).

[0010] The term “modified neurotoxin” as used herein means a Clostridial neurotoxin which includes a modification. Preferably, the modification is within the neurotoxin light chain.

[0011] The term “light chain” as used herein means the light chain of a Clostridial neurotoxin. It has a molecular weight of about 50kDa, and can be referred to as light chain or as the proteolytic domain of a Clostridial neurotoxin. The light chain is believed to be effective as an inhibitor of exocytosis, including as an inhibitor of neurotransmitter, e.g. acetylcholine, release when the light chain is present in the cytoplasm of a target cell, such as a neuron.

[0012] The term “modified light chain” as used herein denotes the light chain of a Clostridial neurotoxin which includes a modification. Preferably, the modification is a structural modification. The modified neurotoxin light chain is structurally different from a naturally occurring neurotoxin light chain, i.e. a non-modified neurotoxin light chain. This structural modification in the light chain of the neurotoxin changes (i) the half-life (time), (ii) the biological activity, (iii) the biological persistence, and/or (iv) the immunogenicity of the neurotoxin in an organism, relative to the neurotoxin from which the modified light chain is derived, i.e. a non-modified neurotoxin.

[0013] The term “heavy chain” as used herein refers to the heavy chain of a Clostridial neurotoxin. It has a molecular weight of about 100kDa.

[0014] The term “polynucleotide” as used herein refers to single- or double-stranded DNA molecules as well as to RNA molecules. Encompassed by the said term is genomic

DNA, cDNA, hnRNA, mRNA as well as all naturally occurring or artificially modified derivatives of such molecular species. The polynucleotide may be in an aspect a linear or circular molecule. Moreover, in addition to the nucleic acid sequences encoding the aforementioned modified neurotoxin polypeptide, a polynucleotide of the present invention may comprise additional sequences required for proper transcription and/or translation such as 5'- or 3'-UTR sequences. The polynucleotide of the present invention encodes a modified neurotoxin polypeptide as described in more detail herein. The modified neurotoxin polypeptide and, in particular, its modified light chain and heavy chain are derivable from one of the antigenically different serotypes of Botulinum Neurotoxins (BoNT), i.e. BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G, or Tetanus Neurotoxin (TeNT). Neurotoxin polypeptides comprise an N-terminal light chain of approximately 50kDa, and a C-terminal heavy chain of approximately 100kDa, linked by a disulfide bond. The neurotoxins are translated as single chain precursor molecules and become proteolytically cleaved into a mature, biologically active di-chain form during processing. The neurotoxin polypeptide (prior to the modification of the invention) comprises the light and heavy chain of neurotoxin BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G or TeNT. Said light and heavy chain of the neurotoxin polypeptide (prior to the modification) comprise an amino acid sequence as shown in any one of SEQ ID NO: 1 (BoNT/A), SEQ ID NO: 2 (BoNT/C1), or SEQ ID NO: 3 (BoNT/E) or in Swiss-Prot: B1INP5.1 (BoNT/B). The amino acid sequence of BoNT/A (comprising the heavy chain and light chain) is, for example, shown in GenBank accession number YP_001253342.1. The amino acid sequence of BoNT/C1 (comprising the heavy chain and light chain) is, for example, shown in Swiss-Prot. accession number P18640.2. The amino acid sequence of BoNT/E (comprising the heavy chain and light chain) is, for example, shown in GenBank accession number CAA44558.1. In one aspect, the structural modification in the BoNT/A light chain comprises one or more mutations selected from the group consisting of a (i) E126A mutation (glutamic acid at position 126 of the light chain is replaced by an alanine), a (ii) L127A mutation (leucine at position 127 is replaced by an alanine), a (iii) F213I mutation (phenylalanine at position 213 of the light chain is replaced by an isoleucine) or an (iv) A214I mutation (alanine at position 214 of the light chain is replaced by an isoleucine), with the amino acid sequence numbering as in GenBank accession number YP_001253342.1 or in SEQ ID NO: 1. In another aspect, the structural modification in the BoNT/A light chain comprises one or more mutations selected from the group consisting of a (i) E126D mutation (glutamic acid at position 126 of the light chain is replaced by an aspartic acid), a (ii) L127V mutation (leucine at position 127 is replaced by a valine), a (iii) F213Y mutation (phenylalanine at position 213 of the light chain is replaced by a tyrosine) or an (iv) A214G mutation (alanine at position 214 of the light chain is replaced by a glycine), with the amino acid sequence numbering as in GenBank

accession number YP_001253342.1 or in SEQ ID NO: 1. In yet another aspect, the structural modification in the BoNT/A light chain comprises one or more mutations selected from the group consisting of a (i) E126A or D mutation, a (ii) L127A or V mutation, a (iii) F213I or Y mutation or an (iv) A214I or G mutation, with the amino acid sequence numbering as in GenBank accession number YP_001253342.1 or in SEQ ID NO: 1. Preferably, said one or more mutations is/are within the BoNT/A light chain. More preferably, the BoNT/A light chain comprises an E126A mutation, a L127A mutation, a F213I mutation and an A214I mutation, with the amino acid sequence numbering as in GenBank accession number YP_001253342.1 or in SEQ ID NO: 1. In another aspect, the BoNT/A light chain comprises an E126D mutation, L127V mutation, F213Y mutation, and A214G mutation, with the amino acid sequence numbering as in GenBank accession number YP_001253342.1 or in SEQ ID NO 1. However, it is encompassed by the scope of the present invention that other and/or further mutations can be introduced into the light chain, either in addition to one or more of the mutations mentioned above or instead of these mutations. Preferably, said mutation(s) is/are within (a) calpain protease cleavage site(s).

[0015] The term “calpain proteases” or “calpains” as used herein refers to proteins belonging to the family of calcium-dependent, non-lysosomal cysteine proteases, i.e. proteolytic enzymes expressed ubiquitously in mammals and many other organisms. Calpains constitute the C2 family of protease clan CA in the MEROPS database. The calpain proteolytic system includes the calpain proteases, the small regulatory subunit CAPNS1, and the endogenous calpain-specific inhibitor, calpastatin.

The calpain protease family contains 14 members with μ -calpain (calpain-1) and m-calpain (calpain-2) being the most well-characterized. The accession number of μ -calpain (calpain-1, catalytic subunit, isoform a) is shown in NP001185798.1, whereas the accession number of m-calpain (calpain-2, catalytic subunit, isoform 1) is depicted in NP001739.2. Structurally, the calpains contain two subunits; an 80 kDa catalytic subunit and a 28 kDa regulatory subunit that functions as a chaperone to stabilize the 80 kDa structure. Calpains are regulated by Ca^{2+} concentration, phosphorylation, calpastatin and probably by altering their subcellular localization, e.g. by limiting access to the substrate. These endopeptidases have numerous biological functions including, but not limited to, remodeling of cytoskeletal attachments to the plasma membrane during cell fusion and cell motility, proteolytic modification of molecules in signal transduction pathways, degradation of enzymes controlling progression through the cell cycle, regulation of gene expression, substrate degradation in some apoptotic pathways, and an involvement in long-term potentiation (Chowdhury et al. 2008, *Comp Biochem Physiol B Biochem Mol Biol.* 151, 10). μ -calpain (calpain-1) and m-calpain (calpain-2) have also been found to be involved in

the degradation of botulinum neurotoxins in neurons, which means that botulinum neurotoxins are recognized and cleaved by calpains. Amongst protein substrates, primary amino acid sequences and tertiary structure elements are likely responsible for directing cleavage to a specific substrate; see e.g. Tompa 2004, J Biol Chem 279, 20775 or Cuerrier 2005, J Biol Chem 280, 40632.

[0016] The term “modified light chain having at least one modification conferring altered cleavage by calpain proteases” as used herein denotes a modification within the neurotoxin light chain which results in (i) increased, (ii) decreased, or (iii) no cleavage, by calpains of the modified neurotoxin light chain.

The term “modification” means any change to a neurotoxin light chain which makes it physically or chemically different from a neurotoxin light chain without the structural modification, e.g. a naturally occurring neurotoxin light chain. Preferably, the modification is a structural modification. The structural modification can comprise e.g. the introduction of one or more exogenous or endogenous calpain recognition and/or cleavage site(s) into the neurotoxin light chain. Proteolytic cleavage by the calpains of a neurotoxin light chain includes recognition and cleavage of the neurotoxin light chain by the calpains. Recognition of the neurotoxin light chain occurs at the calpain recognition site within the neurotoxin light chain. Proteolytic cleavage occurs at the calpain cleavage site within the neurotoxin light chain. Recognition and proteolysis by calpains of the neurotoxin light chain results in the degradation of the neurotoxin light chain, thereby inactivating the proteolytic activity. For example, it has been found that μ -calpain (calpain-1) and m-calpain (calpain-2) are involved in the degradation processes of neurotoxins.

In light of the above, the term “calpain recognition site” as used herein refers to a site, e.g. a primary amino acid sequence or tertiary structure elements, on the neurotoxin light chain recognized by the calpain proteases. The term “calpain cleavage site” as used herein refers to the cleavage site within the neurotoxin light chain cleaved by a calpain protease.

The recognition site and cleavage site of the calpain proteases in the neurotoxin light chain can be identical or they can differ from each other; see, e.g. Tompa 2004, J Biol Chem 279, 20775.

The term “endogenous calpain recognition and/or cleavage site” as used herein means a calpain recognition and/or cleavage site naturally occurring in the neurotoxin light chain. The term “exogenous calpain recognition and/or cleavage site” as used herein means a calpain recognition and/or cleavage site which does not naturally occur in the neurotoxin light chain, e.g. a heterologous calpain recognition and/or cleavage site derived from a different organism or a recombinant calpain recognition and/or cleavage site. The structural modification can also be a mutation within an exogenous or endogenous calpain recognition and/or cleavage site in the neurotoxin light chain. A mutation in a nucleic acid

sequence as used herein can be a deletion, addition or substitution of one or more nucleotides in a DNA sequence coding for a calpain recognition and/or cleavage site. A mutation in a protein sequence can be a deletion, addition or substitution of one or more amino acid residues in a protein sequence of the calpain recognition and/or cleavage site.

5 Such a mutation in a protein sequence of the calpain recognition and/or cleavage site can be, for example, a substitution of one or more amino acid residue(s) at positions P1, P2, P3, P4, P5, P1', P2', and/or P3' position of the calpain recognition and/or cleavage site in the neurotoxin light chain. Preferably, the protein sequence of the calpain recognition and/or cleavage site within the light chain of BoNT/A comprises, for example, Gly-Lys-

10 Phe-Ala-Thr-Asp-Pro (GKFATDP) (SEQ ID NO: 4), with the Glycine corresponding to amino acid residue 211 of the BoNT/A light chain sequence as shown, e.g. in GenBank number YP_001253342.1 or SEQ ID NO: 1. More specifically, Gly corresponds to the P3 position, Lys to the P2 position, Phe to the P1 position, Ala to the P1' position, Thr to the P2' position, Asp to the P3' position and Pro to the P4' position. In another aspect, the

15 protein sequence of the calpain recognition and/or cleavage site within the light chain of BoNT/A comprises Glu-Leu-Lys-Val-Ile-Asp (ELKVID) (SEQ ID NO: 5), with the Glu corresponding to amino acid residue 126 of the BoNT/A light chain sequence as shown, e.g. in GenBank number YP_001253342.1 or SEQ ID NO: 1. More specifically, Glu corresponds to the P3 position, Leu to the P2 position, Lys to the P1 position, Val to the

20 P1' position, Ile to the P2' position and Asp to the P3' position. In other aspects, the calpain recognition and/or cleavage site for calpain-1 within the light chain of BoNT/A comprises Glu-Asp-Thr-Ser-Gly-Lys (SEQ ID NO: 6), Gly-Leu-Glu-Val-Ser-Phe (SEQ ID NO: 7), Leu-Asn-Lys-Ala-Lys-Ser (SEQ ID NO: 8), Val-Asp-Lys-Leu-Lys-Phe (SEQ ID NO: 9), or Val-Leu-Asn-Arg-Lys-Thr (SEQ ID NO: 10), wherein the BoNT/A light chain

25 sequence is as shown, e.g. in GenBank number YP_001253342.1 or SEQ ID NO: 1. In other aspects, the calpain recognition and/or cleavage site for calpain-2 within the light chain of BoNT/A comprises Ile-Val-Gly-Thr-Thr-Ala (SEQ ID NO: 11), or Gly-Thr-Thr-Ala-Ser-Leu (SEQ ID NO: 12), wherein the BoNT/A light chain sequence is as shown, e.g. in GenBank number YP_001253342.1 or SEQ ID NO: 1. One, two, three, four, five, or

30 even all of the mentioned amino acid residues can be replaced by another amino acid residue. A preferred structural modification within the calpain recognition and/or cleavage site Gly-Lys-Phe-Ala-Thr-Asp-Pro (GKFATDP) (SEQ ID NO: 4) of the light chain of BoNT/A comprises an F213I or F213Y mutation (phenylalanine at position 213 of the light chain is replaced by an isoleucine or tyrosine) and/or an A214I or A214G mutation

35 (alanine at position 214 of the light chain is replaced by and isoleucine or glycine), with the BoNT/A light chain sequence as indicated above. A preferred structural modification within the calpain recognition and/or cleavage site Glu-Leu-Lys-Val-Ile-Asp (ELKVID) (SEQ ID NO: 5) comprises an E126A or E126D mutation (glutamic acid at position 126

of the light chain or the P3 position is replaced by an alanine or aspartic acid) and/or a L127A or L127I mutation (leucine at position 127 or P2 position is replaced by an alanine or isoleucine), with the BoNT/A light chain sequence as indicated above. Mutations to a protein sequence can be the result of mutations to DNA sequences that when transcribed and the resulting mRNA translated produce the mutated protein sequence. The term
5 “having at least one modification” as used herein means that the modified neurotoxin light chain has one, two, three or even more modifications conferring altered cleavage by calpain proteases. “Altered cleavage” as used herein means increased cleavage or decreased cleavage or no cleavage of the modified neurotoxin light chain by calpains, in
10 comparison to a naturally occurring neurotoxin light chain, i.e. a non-modified light chain. Said modification in the neurotoxin light chain alters advantageously the biological persistence, biological half life, biological activity and/or immunogenicity of a thus modified neurotoxin, preferably the duration of the biological activity of the modified neurotoxin.

15

[0017] The term “biological activity” or “activity” of a non-modified or modified neurotoxin as used herein denotes the amount of cellular exocytosis inhibited from a cell per unit of time, such as exocytosis of a neurotransmitter, e.g. acetylcholine, from a target cell, such as a neuron. More specifically, it refers to the biological activity of a mature
20 (non-modified or modified) di-chain neurotoxin polypeptide exhibiting a) receptor binding, b) internalization, c) translocation across the endosomal membrane into the cytosol, and/or d) endoproteolytic cleavage of proteins involved in synaptic vesicle fusion. The term “duration of biological activity” as used herein means the time period of the biological activity of a neurotoxin which can be influenced, i.e. altered, by the modification of the
25 neurotoxin light chain, as described herein.

30

[0018] The term “biological persistence” or “persistence” as used herein means the time of duration of interference or influence caused by a non-modified neurotoxin or a modified neurotoxin with a cellular, such as a neuronal, function, including the temporal duration of
30 an inhibition of exocytosis, such as exocytosis of neurotransmitter (e.g. acetylcholine) from a cell, such as a neuron.

35

[0019] The term “biological half life (time)” or “half life (time)” as used herein means the time that the concentration of a non-modified neurotoxin or a modified neurotoxin is reduced to half of the original concentration in a mammalian cell, such as in a mammalian
35 neuron. Preferably, the mammalian neuron is a human neuron.

[0020] The term “immunogenicity” as used herein means the ability of a particular substance, such as an antigen (, e.g. a neurotoxin,) or epitope, to provoke an immune response in the body of a human or animal.

5 [0021] It is to be understood that the definitions and explanations of the terms made above apply mutatis mutandis for all aspects described in this specification in the following except as otherwise indicated.

[0022] The present invention is based on the finding that the biological persistence,
10 biological half life, biological activity and/or immunogenicity of a neurotoxin can be altered by structurally modifying the neurotoxin light chain. In other words, a modified neurotoxin polypeptide comprising a modified neurotoxin light chain with an altered biological persistence, biological half life, biological activity and/or immunogenicity can be formed from a neurotoxin containing or including a structural modification. Preferably,
15 the duration of biological activity of a neurotoxin can be altered by structurally modifying the neurotoxin light chain as described herein.

[0023] In one aspect of the polynucleotide of the invention, said modification confers an
20 increased cleavage by calpain proteases compared to a non-modified light chain.

The term “increased cleavage” as used herein denotes that the cleavage of the modified neurotoxin light chain by calpains is at least 1.5, 2, 3, 4, 5, 10, 50, 100 fold or even higher, in comparison to the cleavage by calpains of a non-modified light chain. The cleavage can be tested by assays well described in the art, for example by ELISA assays, SDS-PAGE,
25 Western blot analysis, and/or HPLC (, e.g. size exclusion). An increased cleavage by calpain proteases of the modified neurotoxin polypeptide comprising a modified neurotoxin light chain described herein results in an increased degradation of the modified neurotoxin in the target cell, e.g. a neuron.

30 [0024] Advantageously, it has been found in accordance with the present invention that the biological persistence, biological half life, biological activity and/or immunogenicity of a neurotoxin can be altered by structurally modifying the light chain of the neurotoxin. The biological persistence and biological activity of BoNT are mainly dependent on the presence of the proteolytically active neurotoxin light chain in the cytoplasm of the target
35 cell, i.e. the neuron. SNARE proteins are cleaved and thereby inactivated by the neurotoxin light chain in sufficient amount only if enough BoNT light chain is present within the neuron. The inactivation of the SNARE proteins, e.g. SNAP-25, within the neuron by the neurotoxin light chain inhibits the exocytosis of neurotransmitter such as acetylcholine. By

inhibiting acetylcholine release, the toxin interferes with nerve impulses and causes flaccid (sagging) paralysis of muscles. Thus, the degree of paralysis depends on the concentration of the neurotoxin light chain in the neuron. Over time, the concentration of the neurotoxin light chain is decreased by cellular degradation processes thereby abolishing the blocking effect of the neurotoxin. For example, for BoNT/A the biological persistence is about three months, whereas for BoNT/E it is about four to six weeks. The cellular degradation of neurotoxins is performed by the ubiquitin proteasome system and the calpains. Whereas the light chain of BoNT/E is ubiquitinated and degraded by the proteasomes the light chain of BoNT/A is resistant against a stable ubiquitination and ensuing degradation by the proteasomes. This finding explains as to why BoNT/E is degraded relatively fast in the neuron, resulting in shorter biological persistence and/or duration of biological activity, whereas BoNT/A exhibits a longer biological persistence and/or duration of biological activity. In the recovery process the light chain seems to be continuously degraded by the proteolytic activity of the calpains. Based on these observations, structural modifications have been introduced into the neurotoxin light chains which alter the biological persistence, biological half life, biological activity and/or immunogenicity of the thus modified neurotoxins. Thereby, the neurotoxin can be optimized for the respective disorders to be treated.

[0025] In a further aspect of the polynucleotide of the invention, said modification is at least one calpain cleavage site which has been introduced into the light chain.

In this aspect, one, two, three, four, five or even more calpain recognition and/or cleavage site(s) have been introduced into the light chain of neurotoxins in order to increase the degradation of said neurotoxins by calpains, thereby reducing the biological persistence, biological half life, biological activity and/or immunogenicity of the neurotoxin. Advantageously, a thus modified neurotoxin can be used as a medicament for the treatment or prevention of diseases defined herein below. In these indications it is particularly beneficial to use a neurotoxin with a reduced biological persistence, biological half life, biological activity and/or immunogenicity.

According to the literature, no specific amino acid sequence is uniquely recognized by calpains. Amongst protein substrates, primary amino acid sequences and tertiary structure elements seem to be responsible for directing cleavage to a specific substrate. Amongst peptide and small-molecule substrates, the most consistently reported specificity is for small, hydrophobic amino acids (e.g. leucine, valine and isoleucine) at the P2 position, and large hydrophobic amino acids (e.g. phenylalanine and tyrosine) at the P1 position; see e.g. Tompa 2004, J Biol Chem 279, 20775 or Cuerrier 2005, J Biol Chem 280, 40632.

[0026] In an aspect, the calpain cleavage site to be introduced into the neurotoxin light chain can be an endogenous calpain recognition and/or cleavage site, i.e. a calpain recognition and/or cleavage site naturally occurring in the neurotoxin light chain. In another aspect, it can be an exogenous calpain recognition and/or cleavage site, i.e. a calpain recognition and/or cleavage site which does not naturally occur in the neurotoxin light chain. Said exogenous calpain recognition and/or cleavage site can also be understood as a heterologous calpain recognition and/or cleavage site or a recombinant calpain recognition and/or cleavage site well understood in the art and as defined herein. Preferably, the calpain recognition and/or cleavage site is a μ -calpain (calpain-1) or m-calpain (calpain-2) recognition and/or cleavage site. More preferably, it is a calpain cleavage site as described in Tompa 2004, J Biol Chem 279, 20775. It is also encompassed by the present invention that the calpain recognition and/or cleavage site comprises the amino acid sequence Pro-Leu-Lys-Ser-Pro-Pro [SEQ ID NO. 13]. In one aspect, said calpain recognition and/or cleavage site replaces the amino acid sequence Ile-Lys-Phe-Ser-Asn-Gly (IKFSNG) [SEQ ID NO. 14] in the BoNT/E light chain, with the isoleucine corresponding to amino acid residue 134, of the BoNT/E light chain as shown, e.g. in GenBank accession number CAA44558.1 or SEQ ID NO: 3.

The introduction of the at least one calpain recognition and/or cleavage site can be carried out by methods described in the art and include mutagenesis techniques as well as standard cloning and PCR based techniques. Recognition and cleavage of the neurotoxin light chain by calpains results in the degradation of the modified neurotoxin light chain. Thereby, the biological persistence, biological half life, biological activity and/or immunogenicity of the neurotoxin is being altered, i.e. decreased, by the incorporation of one or more calpain recognition and/or cleavage site(s) into the neurotoxin light chain, in comparison to a non-modified neurotoxin light chain. This aspect of the invention has been exemplified for BoNT/E.

[0027] In still another aspect of the polynucleotide of the invention, said modified light chain and the heavy chain are derived from BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/F, or BoNT/G.

In this aspect, the light chain (prior to the modification) and the heavy chain come from BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/F, or BoNT/G. The heavy chain is preferably a non-modified heavy chain, i.e. a naturally-occurring heavy chain. However, it is envisaged that the heavy chain comprises for instance tags which allow for the purification of the modified neurotoxin polypeptide, such as His tag or the like. It is preferred that a protease cleavage site is being introduced between the heavy chain and light chain which allows for proteolytic activation of the modified light chain upon cleavage, e.g. by thrombin or an *E. coli* protease known in the art.

[0028] In an aspect of the polynucleotide of the invention, the modified neurotoxin polypeptide comprising the modified light chain exhibits at least one of the following properties, compared to a non-modified neurotoxin polypeptide: (i) altered, i.e. increased or decreased, half-life time in a cellular system, (ii) altered, i.e. increased or decreased biological persistence, and/or (iii) reduced immunogenicity in an organism, preferably a mammal, more preferably a human. It is preferred that the duration of biological activity is altered, i.e. shortened or prolonged. It is envisaged that in some aspects, the biological activity is also being altered, i.e. increased or decreased.

[0029] In a further aspect of the polynucleotide of the invention, said modified neurotoxin polypeptide exhibits a shortened duration of biological activity.

The term “shortened duration of biological activity” as used herein refers to a reduced time period in which the modified neurotoxin light chain exerts its biological activity, in comparison to a non-modified neurotoxin light chain.

Assays for testing the biological activity and the duration of biological activity of neurotoxins are described in the art and comprise, e.g. the Digit Abduction Scoring (DAS) (Aoki 2001, Toxicon 39, 1815) or the voluntary running assay (Keller 2006, Neuroscience 139, 629) or assays described in the following examples. The incorporation of at least one calpain recognition and/or cleavage site into the neurotoxin light chain, results in increased degradation of the neurotoxin light chain, thereby shortening the duration of biological activity. It will be understood that the modified neurotoxin polypeptides according to the present invention will, in an aspect, have reduced side effects when applied to an organism.

[0030] In still another aspect of the polynucleotide of the invention, said modification confers a decreased cleavage by calpain proteases compared to a non-modified light chain.

The term “said modification confers a decreased cleavage by calpain proteases compared to a non-modified light chain” as used herein means a modification within the neurotoxin light chain which results in decreased cleavage or no cleavage by calpains of the modified light chain, in comparison to a non-modified light chain. Thereby, the degradation of the modified neurotoxin by calpains occurs slower compared to a non-modified neurotoxin, or not at all.

More specifically, the term “decreased cleavage” as used herein denotes that the cleavage of the modified neurotoxin light chain by calpains is at least 1.5, 2, 3, 4, 5 fold or even lower, in comparison to the cleavage by calpains of a non-modified light chain, or no cleavage at all. Assays for testing decreased cleavage by calpain proteases have been mentioned elsewhere herein.

[0031] Advantageously, it has been found by the present inventors, that the calpain cleavage site(s) within the light chain of neurotoxins can be mutated, in order to decrease the degradation of said neurotoxins, thereby extending the biological persistence and increasing the biological activity and/or half life. The mutation in an aspect is preferably within an endogenous calpain recognition and/or cleavage site in the neurotoxin light chain. However, it is also envisaged that the mutation is within an exogenous (or heterologous or recombinant) calpain recognition and/or cleavage site in the neurotoxin light chain. A mutation in a nucleic acid sequence as used herein can be a deletion, addition or substitution of one or more nucleotides in the DNA sequence of a calpain recognition and/or cleavage site of a neurotoxin light chain. A mutation in a protein sequence of a calpain recognition and/or cleavage site can be a deletion, addition or substitution of one or more amino acid residues in said protein sequence. As a result of such a mutation, the neurotoxin light chain is no longer recognized, cleaved and thereby degraded by calpains. Mutations to a protein sequence can be the result of mutations to DNA sequences that when transcribed and the resulting mRNA translated produce the mutated protein sequence. In another aspect, the mutation is a deletion of one or more of the endogenous calpain recognition and/or cleavage site(s) within the neurotoxin light chain. In this aspect, the complete endogenous calpain recognition and/or cleavage site(s) within the neurotoxin light chain is/are deleted which results in a neurotoxin light chain no longer being degraded by calpains. One of the reasons, that BoNT/A has originally been selected over the other serotypes, i.e. serotypes BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F and BoNT/G, for clinical use is that BoNT/A has a substantially longer lasting therapeutic effect, i.e. the inhibitory effect of BoNT/A is more persistent. However, for effective therapy of some medical indications such as, e.g., the treatment of dystonia or for cosmetic purposes it is beneficial to use a BoNT/A with a prolonged persistence, increased biological half life and/or biological activity so that the therapeutic effect can be improved and/or extended. Advantageously, a thus modified neurotoxin can be used as a medicament for the treatment or prevention of diseases specified herein below.

[0032] In an aspect of the polynucleotide of the invention, said modification is at least one substitution within a calpain cleavage site in the light chain.

The term "substitution" as used herein means a mutation in which one or more nucleotides within the encoded calpain recognition and/or cleavage site(s) in the neurotoxin light chain is/are replaced by other nucleotides. A substitution in a protein sequence can be a replacement of one or more amino acid residues in the amino acid sequence of the calpain cleavage site. Such a mutation in a protein sequence can be, for example, a substitution of one or more amino acid residues at positions P1, P2, P3, P4, P5, P1', P2', and/or P3' position of the calpain recognition and/or cleavage site in the neurotoxin light chain. One,

two, three, four, five, six, seven or even all of the eight amino acid residues can be replaced by another amino acid residue. Preferably, the calpain recognition and/or cleavage site is a μ -calpain (calpain-1) or m-calpain (calpain-2) recognition and/or cleavage site. Mutations to a protein sequence can be the result of mutations to DNA sequences that when transcribed and the resulting mRNA translated produce the mutated protein sequence. Suitable techniques for carrying out such modifications are well known in the art and include standard cloning, mutagenesis techniques as well as PCR based techniques.

[0033] In still another aspect of the polynucleotide of the invention, said substitution is a substitution at the P1 or P1' position of the calpain cleavage site.

In this aspect, position P1 of the calpain recognition and/or cleavage site in the neurotoxin light chain can be substituted e.g. with Valine or Isoleucine.

The P2 position of the calpain recognition and/or cleavage site in the neurotoxin light chain can be replaced with e.g. Tryptophan or Alanine.

The P3 position of the calpain recognition and/or cleavage site in the neurotoxin light chain can be replaced with e.g. Tyrosine or Alanine.

The P4 position of the calpain recognition and/or cleavage site in the neurotoxin light chain can be replaced with e.g. Tryptophan or Phenylalanine.

The P1' position of the calpain recognition and/or cleavage site in the neurotoxin light chain can be replaced with e.g. Tyrosine or Tryptophan.

The P2' position of the calpain recognition and/or cleavage site in the neurotoxin light chain can be replaced with e.g. Leucine or Isoleucine.

The P3' position of the calpain recognition and/or cleavage site in the neurotoxin light chain can be replaced with e.g. Glutamine or Asparagine.

[0034] This approach has been exemplified by a modified light chain of BoNT/A in which, inter alia, the P1 position Phenylalanine (corresponding to amino acid residue 213) has been mutated to an Isoleucine residue and the P1' position Alanine (corresponding to amino acid residue 214) has been mutated to a Tyrosine residue. This modification results in a reduced degradation of the modified neurotoxin light chain by calpains which leads to an extended inhibition of the exocytosis of neurotransmitter, e.g. acetylcholine, thereby increasing the biological persistence, half life and/or biological activity of the neurotoxin.

[0035] In an aspect of the polynucleotide of the invention, said modified light chain and the heavy chain are derived from BoNT/A, BoNT/B or BoNT/C.

[0036] In a further aspect of the polynucleotide of the invention, said modified neurotoxin polypeptide exhibits a prolonged duration of biological activity.

The term “prolonged duration of biological activity” as used herein refers to a prolonged time period in which the modified neurotoxin light chain exerts its biological activity, in comparison to a non-modified neurotoxin light chain. This can be tested as set forth elsewhere herein.

5

[0037] The present invention also relates to a vector comprising the polynucleotide of the invention. In an aspect, the said vector is an expression vector. The term “vector”, preferably, encompasses phage, plasmid, viral or retroviral vectors as well as artificial
10 chromosomes, such as bacterial or yeast artificial chromosomes. Moreover, the term also relates to targeting constructs which allow for random or site- directed integration of the targeting construct into genomic DNA. Such target constructs, preferably, comprise DNA of sufficient length for either homologous or heterologous recombination as described in detail below. The vector encompassing the polynucleotides of the present invention, in an
15 aspect, further comprises selectable markers for propagation and/or selection in a host. The vector may be incorporated into a host cell by various techniques well known in the art. For example, a plasmid vector can be introduced in a precipitate such as a calcium phosphate precipitate or rubidium chloride precipitate, or in a complex with a charged lipid or in carbon-based clusters, such as fullerenes. Alternatively, a plasmid vector may be
20 introduced by heat shock or electroporation techniques. Should the vector be a virus, it may be packaged in vitro using an appropriate packaging cell line prior to application to host cells. Retroviral vectors may be replication competent or replication defective. In the latter case, viral propagation generally will occur only in complementing host/cells. Moreover, in an aspect of the invention, the polynucleotide is operatively linked to
25 expression control sequences allowing expression in prokaryotic or eukaryotic host cells or isolated fractions thereof in the said vector. Expression of the polynucleotide comprises transcription of the polynucleotide into a translatable mRNA. Regulatory elements ensuring expression in host cells are well known in the art. In an aspect, they comprise regulatory sequences ensuring initiation of transcription and/or poly-A signals ensuring
30 termination of transcription and stabilization of the transcript. Additional regulatory elements may include transcriptional as well as translational enhancers. Possible regulatory elements permitting expression in prokaryotic host cells comprise, e.g., the lac-, trp- or tac-promoter in *E. coli*, and examples for regulatory elements permitting expression in eukaryotic host cells are the AOX1- or the GAL1- promoter in yeast or the CMV-, SV40-,
35 RSV-promoter (Rous sarcoma virus), CMV-enhancer, SV40-enhancer or a globin intron in mammalian and other animal cells. Other expression systems envisaged by the invention shall permit expression in insect cells, such as polyhedrin promoter based systems.

[0038] Moreover, inducible expression control sequences may be used in an expression vector encompassed by the present invention. Such inducible vectors may comprise tet or lac operator sequences or sequences inducible by heat shock or other environmental factors. Suitable expression control sequences are well known in the art. Beside elements which are responsible for the initiation of transcription such regulatory elements may also comprise transcription termination signals, such as the SV40-poly-A site or the tk-poly-A site, downstream of the polynucleotide. In this context, suitable expression vectors are known in the art such as Okayama-Berg cDNA expression vector pcDV1 (Pharmacia), pBluescript (Stratagene), pCDM8, pRc/CMV, pcDNA1, pcDNA3 (Invitrogen) or pSPORT1 (Invitrogen) or baculovirus-derived vectors. Preferably, said vector is an expression vector and a gene transfer or targeting vector. Expression vectors derived from viruses such as retroviruses, vaccinia virus, adeno-associated virus, herpes viruses, or bovine papilloma virus, may be used for delivery of the polynucleotides or vector of the invention into targeted cell population. Methods which are well known to those skilled in the art can be used to construct recombinant viral vectors; see, for example, the techniques described in Sambrook, *Molecular Cloning A Laboratory Manual*, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y. (1994).

[0039] The present invention further relates to a host cell comprising the polynucleotide or the vector of the invention.

The term "host cell" as used herein encompasses prokaryotic and eukaryotic host cells. In an aspect the host cell is a bacterial cell and, in another aspect, a Firmicutes bacterial cell. In one aspect, the said bacterial host cell is an E.coli host cell. In another aspect, it is a Clostridium host cell. In a further aspect, the said Clostridium host cell is a Clostridium botulinum host cell, in even a further aspect, a cell of one of the aforementioned seven different serotypes of Clostridium botulinum. In yet another aspect, the bacterial host cell is a Clostridium tetani host cell. In a further aspect, the host cell is a Bacillus host cell and in a particular aspect a Bacillus megaterium host cell. A eukaryotic host cell, in an aspect, is a cell of an animal cell line suitable for production of toxic proteins or a fungal host cell such as a yeast host cell. A host cell as referred to herein, thus, encompasses in an aspect yeast, mammalian, plant or insect cells either as primary cells or as cell lines.

[0040] The present invention also relates to a polypeptide encoded by the polynucleotide of the invention.

The term "polypeptide" as used herein denotes a modified neurotoxin polypeptide comprising a modified neurotoxin light chain and a heavy chain, said modified light chain having at least one modification conferring altered cleavage by calpain proteases, as defined elsewhere herein.

5

[0041] In another aspect, the said light and heavy chain of the neurotoxin polypeptide (prior to the modification of the invention) is a variant having one or more amino acid substitutions, deletions and/or additions. Moreover, such a variant polypeptide, in an aspect, is an amino acid sequence being at least 40%, at least 50%, at least 60%, at least 10 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the amino acid sequence as shown in any one of SEQ ID NOs: 1, 2, or 3, or in Swiss-Prot: B1INP5.1, or encoded by a polynucleotide being at least 40%, at least 50%, at least 60%, at least 70%, at least 75%, at least 80%, at least 85%, at least 90%, at least 95%, at least 98% or at least 99% identical to the polynucleotide encoding the 15 amino acid sequence as shown in any one of SEQ ID NOs: 1, 2, or 3, or in Swiss-Prot: B1INP5.1. The term "identical" as used herein refers to sequence identity characterized by determining the number of identical amino acids between two nucleic acid sequences or amino acid sequences wherein the sequences are aligned so that the highest order match is obtained. It can be calculated using published techniques or methods codified in computer 20 programs such as, for example, BLASTP, BLASTN or FASTA (Altschul 1990, J Mol Biol 215, 403). The percent identity values are, in one aspect, calculated over the entire amino acid sequence. A series of programs based on a variety of algorithms is available to the skilled worker for comparing different sequences. In this context, the algorithms of Needleman and Wunsch or Smith and Waterman give particularly reliable results. To carry 25 out the sequence alignments, the program PileUp (Higgins 1989, CABIOS 5, 151) or the programs Gap and BestFit (Needleman 1970, J Mol Biol 48; 443; Smith 1981, Adv Appl Math 2, 482), which are part of the GCG software packet (Genetics Computer Group 1991, 575 Science Drive, Madison, Wisconsin, USA 53711), may be used. The sequence identity values recited above in percent (%) are to be determined, in another aspect of the 30 invention, using the program GAP over the entire sequence region with the following settings: Gap Weight: 50, Length Weight: 3, Average Match: 10.000 and Average Mismatch: 0.000, which, unless otherwise specified, shall always be used as standard settings for sequence alignments. In an aspect, each of the aforementioned variant polypeptides (prior to the modification) retains one or more and, in another aspect, all of 35 the biological properties of the respective original neurotoxin polypeptide, i.e. the BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/E, BoNT/F, BoNT/G or Tetanus Neurotoxin (TeNT). Those of skill in the art will appreciate that full biological activity is maintained only after proteolytic activation, even though it is conceivable that the

unprocessed precursor can exert some biological functions or be partially active. "Biological properties" as used herein refers to (a) receptor binding, (b) internalization, (c) translocation across the endosomal membrane into the cytosol, and/or (d) endoproteolytic cleavage of proteins involved in synaptic vesicle membrane fusion. In vivo assays for assessing biological activity include the mouse LD50 assay and the ex vivo mouse hemidiaphragm assay as described by Pearce et al. (Pearce 1994, Toxicol Appl Pharmacol 128: 69-77) and Dressler et al. (Dressler 2005, Mov Disord 20:1617-1619, Keller 2006, Neuroscience 139: 629-637). The biological activity is commonly expressed in Mouse Units (MU). As used herein, 1 MU is the amount of neurotoxic component, which kills 50% of a specified mouse population after intraperitoneal injection, i.e. the mouse i.p. LD50. In an aspect, the variant polynucleotides can be modified according to the invention in that the encoded light chain has at least one modification conferring altered cleavage by calpain proteases as defined herein. In this way, the duration of biological activity of the variant polynucleotides can be changed.

15

[0042] The modified neurotoxin polypeptide of the invention, in an aspect, can be manufactured entirely or in part by chemical synthesis or recombinant molecular biology techniques well known for the skilled artisan. In an aspect, such a method of manufacturing the modified neurotoxin polypeptide of the invention comprises (a) culturing the host cell of the present invention described elsewhere herein and (b) obtaining from the said host cell the polypeptide of the present invention. In an aspect of this method, the polypeptide can be obtained by conventional purification techniques from a host cell lysate including affinity chromatography, ion exchange chromatography, size exclusion chromatography and/or preparative gel electrophoresis. The modified neurotoxin polypeptide of the invention can be analyzed quantitatively or qualitatively by methods described in the art, such as ELISA assays, Western blot analysis, SDS-PAGE, and/or HPLC (Reverse phase, Size exclusion).

20

25

[0043] The present invention also relates to a composition comprising the polynucleotide, the vector or the polypeptide of the invention, as a medicament.

30

The term "medicament" as used herein refers, in one aspect, to a pharmaceutical composition containing the polypeptide, polynucleotide or vector of the present invention as pharmaceutical active compound, wherein the pharmaceutical composition may be used for human or non-human therapy of various diseases or disorders in a therapeutically effective dose.

35

[0044] In an aspect, the polypeptide, polynucleotide or vector of the present invention can be present in liquid or lyophilized form. In an aspect, said compound can be present

together with glycerol, protein stabilizers (e.g., human serum albumin (HSA)) or non-protein stabilizers.

5 [0045] The medicament is, in one aspect, administered topically. Conventionally used drug administration is administered intra-muscular, subcutaneous (near glands). However, depending on the nature and the mode of action of a compound (i.e. the polypeptide, polynucleotide or vector of the present invention), the medicament may be administered by other routes as well.

10 [0046] The compound is the active ingredient of the composition, and is in one aspect, administered in conventional dosage forms prepared by combining the drug with standard pharmaceutical carriers according to conventional procedures. These procedures may involve mixing, granulating, and compression, or dissolving the ingredients as appropriate to the desired preparation. It will be appreciated that the form and character of the
15 pharmaceutical acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration, and other well-known variables.

[0047] The carrier(s) must be acceptable in the sense of being compatible with the other
20 ingredients of the formulation and being not deleterious to the recipient thereof. The pharmaceutical carrier employed may include a solid, a gel, or a liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are phosphate buffered saline solution, syrup, oil, water, emulsions, various types of wetting agents, and the like.
25 Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl mono-stearate or glyceryl distearate alone or with a wax. Said suitable carriers comprise those mentioned above and others well known in the art, see, e.g., Remington's Pharmaceutical Sciences, Mack Publishing Company, Easton, Pennsylvania.

30 [0048] The diluent(s) is/are selected so as not to affect the biological activity of the combination. Examples of such diluents are distilled water, physiological saline, Ringer's solutions, dextrose solution, and Hank's solution. In addition, the pharmaceutical composition or formulation may also include other carriers, adjuvants, or non-toxic, non-therapeutic, non-immunogenic stabilizers and the like.

35

[0049] A therapeutically effective dose refers to an amount of the compound to be used in medicament of the present invention which prevents, ameliorates or treats the symptoms accompanying a disease or condition referred to in this specification. Therapeutic efficacy

and toxicity of the compound can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., ED50 (the dose therapeutically effective in 50% of the population) and LD50 (the dose lethal to 50% of the population). The dose ratio between therapeutic and toxic effects is the therapeutic index, and it can be expressed as the ratio, LD50/ED50.

[0050] The dosage regimen will be determined by the attending physician and other clinical factors. As is well known in the medical arts, dosages for any one patient depends upon many factors, including the patient's size, body surface area, age, the particular compound to be administered, sex, time and route of administration, general health, and other drugs being administered concurrently. Progress can be monitored by periodic assessment.

[0051] The medicament referred to herein is administered at least once in order to treat or ameliorate or prevent a disease or condition recited in this specification. However, the said medicament may be administered more than one time.

[0052] Specific medicaments are prepared in a manner well known in the pharmaceutical art and comprise at least one active compound referred to herein above in admixture or otherwise associated with a pharmaceutically acceptable carrier or diluent. For making those specific pharmaceutical compositions, the active compound(s) will usually be mixed with a carrier or the diluent. The resulting formulations are to be adapted to the mode of administration. Dosage recommendations shall be indicated in the prescribers or users instructions in order to anticipate dose adjustments depending on the considered recipient.

[0053] The medicament according to the present invention may in a further aspect of the invention comprise drugs in addition to the polypeptide, polynucleotide or vector of the present invention which are added to the medicament during its formulation. Finally, it is to be understood that the formulation of a medicament takes place under GMP standardized conditions or the like in order to ensure quality, pharmaceutical security, and effectiveness of the medicament.

[0054] In addition, the present invention relates to a composition comprising the polynucleotide of the invention or a polypeptide encoded thereby for use as a medicament for or treating and/or preventing a disease selected from the group consisting of: wound healing, immobilisation for bone and tendon fracture treatment, post surgery immobilization, specifically in connection with haemorrhoidectomy, introduction of dental implants, or hip joint replacement (endoprosthesis), knee arthroplasty, ophthalmological

surgery, acne, irritable bowel disease or prostate hyperplasia. In an aspect of the composition of the invention, the polynucleotide encodes a modified neurotoxin polypeptide comprising a modified neurotoxin light chain and a heavy chain, said modified light chain having at least one modification conferring altered cleavage by calpain proteases. In one aspect of the composition of the invention, the polynucleotide of the invention or polypeptide encoded thereby comprise a modification conferring an increased cleavage by calpain proteases, compared to a non-modified neurotoxin light chain. In another aspect of the composition of the invention, the modification is at least one calpain cleavage site which has been introduced into the light chain. In a further aspect of the composition of the invention, the modified light chain and the heavy chain are derived from BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/F, or BoNT/G. In still another aspect of the composition of the invention, the modified neurotoxin polypeptide exhibits a shortened duration of biological activity.

[0055] The present invention furthermore relates to a composition comprising the polynucleotide of the invention or a polypeptide encoded thereby for use as a medicament for or treating and/or preventing a disease selected from the group consisting of: voluntary muscle strength, focal dystonia, including cervical, cranial dystonia, and benign essential blepharospasm, hemifacial spasm, and focal spasticity, gastrointestinal disorders, hyperhidrosis, and cosmetic wrinkle correction, Blepharospasm, oromandibular dystonia, jaw opening type, jaw closing type, bruxism, Meige syndrome, lingual dystonia, apraxia of eyelid, opening cervical dystonia, antecollis, retrocollis, laterocollis, torticollis, pharyngeal dystonia, laryngeal dystonia, spasmodic dysphonia/adductor type, spasmodic dysphonia/abductor type, spasmodic dyspnea, limb dystonia, arm dystonia, task specific dystonia, writer's cramp, musician's cramps, golfer's cramp, leg dystonia, thigh adduction, thigh abduction knee flexion, knee extension, ankle flexion, ankle extension, equinovarus, deformity foot dystonia, striatal toe, toe flexion, toe extension, axial dystonia, pisa syndrome, belly dancer dystonia, segmental dystonia, hemidystonia, generalised dystonia, dystonia in lubag, dystonia in corticobasal degeneration, dystonia in lubag, tardive dystonia, dystonia in spinocerebellar ataxia, dystonia in Parkinson's disease, dystonia in Huntington's disease, dystonia in Hallervorden-Spatz disease, dopa-induced dyskinesias/dopa-induced dystonia, tardive dyskinesias/tardive dystonia, paroxysmal dyskinesias/dystonias, kinesiogenic non-kinesiogenic action-induced palatal myoclonus, myoclonus myokymia, rigidity, benign muscle cramps, hereditary chin trembling, paradoxical jaw muscle activity, hemimasticatory spasms, hypertrophic branchial myopathy, masseteric hypertrophy, tibialis anterior hypertrophy, nystagmus, oscillopsia supranuclear gaze palsy, epilepsy, partialis continua, planning of spasmodic torticollis operation, abductor vocal cord paralysis, recalcitrant mutational dysphonia, upper oesophageal

sphincter dysfunction, vocal fold granuloma, stuttering Gilles de la Tourette syndrome, middle ear myoclonus, protective larynx closure, postlaryngectomy, speech failure, protective ptosis, entropion sphincter Odii dysfunction, pseudoachalasia, nonachalsia, oesophageal motor disorders, vaginismus, postoperative immobilisation tremor, bladder
5 dysfunction, detrusor sphincter dyssynergia, bladder sphincter spasm, hemifacial spasm, reinnervation dyskinesias, cosmetic use crow's feet, frowning facial asymmetries, mentalis dimples, stiff person syndrome, tetanus prostate hyperplasia, adipositas, treatment infantile cerebral palsy strabismus, mixed paralytic concomitant, after retinal detachment surgery, after cataract surgery, in aphakia myositic strabismus, myopathic strabismus, dissociated
10 vertical deviation, as an adjunct to strabismus surgery, esotropia, exotropia, achalasia, anal fissures, exocrine gland hyperactivity, Frey syndrome, Crocodile Tears syndrome, hyperhidrosis, axillar palmar plantar rhinorrhea, relative hypersalivation in stroke, in Parkinson's, in amyotrophic lateral sclerosis, spastic conditions, in encephalitis and myelitis autoimmune processes, multiple sclerosis, transverse myelitis, Devic syndrome,
15 viral infections, bacterial infections, parasitic infections, fungal infections, in hereditary spastic paraparesis postapoplectic syndrome hemispheric infarction, brainstem infarction, myelon infarction, in central nervous system trauma, hemispheric lesions, brainstem lesions, myelon lesion, in central nervous system hemorrhage, intracerebral hemorrhage, subarachnoidal hemorrhage, subdural hemorrhage, intraspinal hemorrhage, in neoplasias,
20 hemispheric tumors, brainstem tumors, myelon tumor and vaginism.

In an aspect of the composition of the invention, the polynucleotide encodes a modified neurotoxin polypeptide comprising a modified neurotoxin light chain and a heavy chain, said modified light chain having at least one modification conferring altered cleavage by calpain proteases. In one aspect of the composition of the invention, the polynucleotide of
25 the invention or polypeptide encoded thereby comprise a modification which confers a decreased cleavage by calpain proteases compared to a non-modified light chain. In another aspect of the composition of the invention, the modification is at least one substitution within a calpain cleavage in the light chain. In still another aspect of the composition of the invention, substitution is a substitution at the P1, P2, P3, P4, P5, P1',
30 P2', and/or P3' position of the calpain cleavage site. In a further aspect of the composition of the invention, modified light chain and the heavy chain are derived from BoNT/A, BoNT/B or BoNT/C1. In a still further aspect of the composition of the invention, the modified neurotoxin polypeptide exhibits a prolonged duration of biological activity.

35

[0056] Another aspect of the invention, the composition relates to a cosmetic composition which can be formulated as described for a medicament above. For a cosmetic composition, likewise, it is envisaged that the modified neurotoxin of the present invention

is in an aspect used in substantially pure form. Cosmetic compositions are, in a further aspect, to be applied intramuscular. In an even further aspect, cosmetic compositions comprising the neurotoxin can be formulated as anti-wrinkle solution.

- 5 [0057] All references cited in this specification are herewith incorporated by reference with respect to their entire disclosure content and the disclosure content specifically mentioned in this specification.

EXAMPLES:

10

[0058] The invention will now be illustrated by examples which shall, however, not be construed as limiting the scope of the invention.

15

[0059] Example 1: Different susceptibility of BoNT/E and BoNT/A for m-calpain

20

30 μ g BoNT/E (SEQ ID NO: 3) in 162 mM Na Acetate buffer were supplemented with 14,7 μ L 119 mM dithiothreitol to cleave the disulphide bond between heavy and light chain. After incubation at 30°C for 30 min 90 μ L 125 mM HEPES, 375 mM NaCl and 10mM CaCl₂, pH 7.5 were added followed by 2 μ g calpain (Calbiochem) in 3 μ L. After 6 ,
24 and 46 hours incubation 23 μ L sample were supplemented with 2 μ L EGTA and then analysed by SDS-PAGE. 30 μ g of BoNT/A (SEQ ID NO: 1) were analysed in parallel in the same buffer and under the same conditions and analysed by SDS-PAGE. The electropherogram shows that the light chain of BoNT/A was resistant against proteolysis for 46 hours whereas the heavy chain is slightly degraded, a band with a molecular weight of about 80 kD appeared and the intensity of the heavy chain (about 100 kD) was reduced. In contrast the heavy chain of BoNT/E was completely degraded after 360 min. The light chain of BoNT/E was still visible after 48 hours but showed a loss of intensity. After 48 hours the light chain had disappeared i.e. was completely degraded.

30

[0060] Example 2: Preparation of a mutated BoNT/A

35

The light chain of BoNT/A comprised by SEQ ID NO: 1 contains a putative cleavage site for m-calpain, as shown in SEQ ID NO: 5. The following mutants were synthesised: E126A / L127A and F213I / A214I, with the mentioned mutations at the indicated positions within the BoNT/A light chain comprised by SEQ ID NO: 1. The double mutations were introduced into the expressin plasmid pET29c-mod Strep-BoNTA-Strep-His, containing the BoNT/A gene and the genetic information for the c-terminal Strep- and His-tag and a thrombin cleavage site between heavy and light chain. The introduction was

accomplished by site directed mutagenesis according to the Gentailor protocol (Invitrogen).

Plasmids with the verified DNA were transformed into *E. coli* BL21DE3. Colonies were grown in YT medium + 50µg/mL kanamycin over night at 37°C. 3 L 2YT medium were inoculated with this pre-culture. After reaching an OD 0.6 the expression was started by adding 50 µM IPTG until a final concentration of 0.2 M IPTG was attained. After further growth for 14 hours the cells were harvested. The cell pellets were lysed by ultrasonication in 50 mM Tris, 150 mM NaCl, 1,5 mM imidazole in the presence of complete protease inhibitor cocktail without EDTA (Roche). After centrifugation the mutants were purified over 2 mL Talon, Matrix (Clontech), bound proteins were eluted with 250 mM imidazole. Mutant containing fractions were further purified by cation chromatography HiTrap SP FF in 50 mM Tris /HCl pH 8.0. The mutants were eluted with a NaCl gradient 0-1.0 M NaCl. The purified single chain mutated BoNT/A was cleaved into heavy and light chain with 0.01 U Thrombin per µg protein at ambient temperature overnight. Thrombin was removed by size exclusion chromatography over a HiLoad Superdex 200 16/60 column in phosphate buffered saline at pH 7.4. The biological activity was analysed in the mouse hemidiaphragm assay.

[0061] Example 3: Analysis of the duration of effect in the mouse running assay

Mice were trained for about 14 days in a cage equipped with a running wheel according to Keller (loc. cit.). The running distance per night was monitored electronically. 0.8 Units of unmodified BoNT/A in 20 µl were injected into the gastrocnemius muscle of 8 mice. The running distance was reduced to zero after 3-4 days. Over a period of 28 days the running distance continuously increased to reach the initial value. In a further experiment a group of mice is treated with 0.8 U of the mutant BoNT/A generated in Example 2 in parallel with unmodified BoNT/A. Again after 3-4 days the running distance is reduced to zero and continuously increases for both groups. The recovery time for mice treated with the mutant BoNT/A is markedly longer.

Claims

- 5 1. A polynucleotide encoding a modified neurotoxin polypeptide comprising a modified neurotoxin light chain and a heavy chain, said modified light chain having at least one modification conferring altered cleavage by calpain proteases.
- 10 2. The polynucleotide of claim 1, wherein said modification confers an increased cleavage by calpain proteases compared to a non-modified light chain.
- 15 3. The polynucleotide of claim 2, wherein said modification is at least one calpain cleavage site which has been introduced into the light chain.
- 20 4. The polynucleotide of claim 3, wherein said modified light chain and the heavy chain are derived from BoNT/A, BoNT/B, BoNT/C1, BoNT/D, BoNT/F, or BoNT/G.
- 25 5. The polynucleotide of any one of claims 1 to 4, wherein said modified neurotoxin polypeptide exhibits a shortened duration of biological activity.
- 30 6. The polynucleotide of claim 1, wherein said modification confers a decreased cleavage by calpain proteases compared to a non-modified light chain.
- 35 7. The polynucleotide of claim 6, wherein said modification is at least one substitution within a calpain cleavage in the light chain.
8. The polynucleotide of claim 7, wherein said substitution is a substitution at the P1, P2, P3, P4, P5, P1', P2', and/or P3' position of the calpain cleavage site.
9. The polynucleotide of claim 7 or 8, wherein said modified light chain and the heavy chain are derived from BoNT/A, BoNT/B or BoNT/C1.
10. The polynucleotide of any one of claims 6 to 9, wherein said modified neurotoxin polypeptide exhibits a prolonged duration of biological activity.
11. A vector comprising the polynucleotide of any one of claims 1 to 10.

12. A host cell comprising the polynucleotide of any one of claims 1 to 10 or the vector of claim 11.
13. A polypeptide encoded by the polynucleotide of any one of claims 1 to 10.
- 5 14. A composition comprising the polynucleotide of any one of claims 1 to 10, the vector of claim 11, or the polypeptide of claim 13 as a medicament.
- 10 15. A composition comprising the polynucleotide of any one of claims 1 to 5 or a polypeptide encoded thereby for use as a medicament for or treating and/or preventing a disease selected from the group consisting of: wound healing, immobilisation for bone and tendon fracture treatment, post surgery immobilization, specifically in connection with haemorrhoidectomy, introduction of dental implants, or hip joint replacement (endoprosthesis), knee arthroplasty, 15 ophthalmological surgery, acne, irritable bowel disease or prostate hyperplasia.
16. A composition comprising the polynucleotide of any one of claims 6 to 10 or a polypeptide encoded thereby for use as a medicament for or treating and/or preventing a disease selected from the group consisting of: voluntary muscle 20 strength, focal dystonia, including cervical, cranial dystonia, and benign essential blepharospasm, hemifacial spasm, and focal spasticity, gastrointestinal disorders, hyperhidrosis, and cosmetic wrinkle correction, Blepharospasm, oromandibular dystonia, jaw opening type, jaw closing type, bruxism, Meige syndrome, lingual dystonia, apraxia of eyelid, opening cervical dystonia, antecollis, retrocollis, 25 laterocollis, torticollis, pharyngeal dystonia, laryngeal dystonia, spasmodic dysphonia/adductor type, spasmodic dysphonia/abductor type, spasmodic dyspnea, limb dystonia, arm dystonia, task specific dystonia, writer's cramp, musician's cramps, golfer's cramp, leg dystonia, thigh adduction, thigh abduction knee flexion, knee extension, ankle flexion, ankle extension, equinovarus, deformity foot 30 dystonia, striatal toe, toe flexion, toe extension, axial dystonia, pisa syndrome, belly dancer dystonia, segmental dystonia, hemidystonia, generalised dystonia, dystonia in lubag, dystonia in corticobasal degeneration, dystonia in lubag, tardive dystonia, dystonia in spinocerebellar ataxia, dystonia in Parkinson's disease, dystonia in Huntington's disease, dystonia in Hallervorden-Spatz disease, dopa-induced 35 dyskinesias/dopa-induced dystonia, tardive dyskinesias/tardive dystonia, paroxysmal dyskinesias/dystonias, kinesiogenic non-kinesiogenic action-induced palatal myoclonus, myoclonus myokymia, rigidity, benign muscle cramps,

hereditary chin trembling, paradoxical jaw muscle activity, hemimasticatory spasms, hypertrophic branchial myopathy, masseteric hypertrophy, tibialis anterior hypertrophy, nystagmus, oscillopsia supranuclear gaze palsy, epilepsy, partialis continua, planning of spasmodic torticollis operation, abductor vocal cord paralysis, 5 recalitant mutational dysphonia, upper oesophageal sphincter dysfunction, vocal fold granuloma, stuttering Gilles de la Tourette syndrome, middle ear myoclonus, protective larynx closure, postlaryngectomy, speech failure, protective ptosis, entropion sphincter Oculi dysfunction, pseudoachalasia, nonachalasia, oesophageal motor disorders, vaginismus, postoperative immobilisation tremor, bladder 10 dysfunction, detrusor sphincter dyssynergia, bladder sphincter spasm, hemifacial spasm, reinnervation dyskinesias, cosmetic use of feet, frowning facial asymmetries, mentalis dimples, stiff person syndrome, tetanus prostate hyperplasia, adipositas, treatment of infantile cerebral palsy strabismus, mixed paralytic concomitant, after retinal detachment surgery, after cataract surgery, in aphakia 15 myositic strabismus, myopathic strabismus, dissociated vertical deviation, as an adjunct to strabismus surgery, esotropia, exotropia, achalasia, anal fissures, exocrine gland hyperactivity, Frey syndrome, Crocodile Tears syndrome, hyperhidrosis, axillary palmar plantar rhinorrhea, relative hypersalivation in stroke, in Parkinson's, in amyotrophic lateral sclerosis, spastic conditions, in encephalitis 20 and myelitis autoimmune processes, multiple sclerosis, transverse myelitis, Devic syndrome, viral infections, bacterial infections, parasitic infections, fungal infections, in hereditary spastic paraparesis postapoplectic syndrome hemispheric infarction, brainstem infarction, myelom infarction, in central nervous system trauma, hemispheric lesions, brainstem lesions, myelom lesion, in central nervous 25 system hemorrhage, intracerebral hemorrhage, subarachnoidal hemorrhage, subdural hemorrhage, intraspinal hemorrhage, in neoplasias, hemispheric tumors, brainstem tumors, myelom tumor and vaginismus.

eolf-seq1
SEQUENCE LISTING

<110> Merz Pharma GmbH & Co. KGaA
<120> Alteration of proteolytic cleavage of botulinum neurotoxins
<130> MP10484PC
<160> 14
<170> PatentIn version 3.3
<210> 1
<211> 1296
<212> PRT
<213> Clostridium botulinum

<400> 1

Met Pro Phe Val Asn Lys Gln Phe Asn Tyr Lys Asp Pro Val Asn Gly
1 5 10 15

Val Asp Ile Ala Tyr Ile Lys Ile Pro Asn Ala Gly Gln Met Gln Pro
20 25 30

Val Lys Ala Phe Lys Ile His Asn Lys Ile Trp Val Ile Pro Glu Arg
35 40 45

Asp Thr Phe Thr Asn Pro Glu Glu Gly Asp Leu Asn Pro Pro Pro Glu
50 55 60

Ala Lys Gln Val Pro Val Ser Tyr Tyr Asp Ser Thr Tyr Leu Ser Thr
65 70 75 80

Asp Asn Glu Lys Asp Asn Tyr Leu Lys Gly Val Thr Lys Leu Phe Glu
85 90 95

Arg Ile Tyr Ser Thr Asp Leu Gly Arg Met Leu Leu Thr Ser Ile Val
100 105 110

Arg Gly Ile Pro Phe Trp Gly Gly Ser Thr Ile Asp Thr Glu Leu Lys
115 120 125

Val Ile Asp Thr Asn Cys Ile Asn Val Ile Gln Pro Asp Gly Ser Tyr
130 135 140

Arg Ser Glu Glu Leu Asn Leu Val Ile Ile Gly Pro Ser Ala Asp Ile
145 150 155 160

Ile Gln Phe Glu Cys Lys Ser Phe Gly His Glu Val Leu Asn Leu Thr
165 170 175

Arg Asn Gly Tyr Gly Ser Thr Gln Tyr Ile Arg Phe Ser Pro Asp Phe
180 185 190

Thr Phe Gly Phe Glu Glu Ser Leu Glu Val Asp Thr Asn Pro Leu Leu

eof-seq1

740

745

750

Gln Tyr Thr Glu Glu Glu Lys Asn Asn Ile Asn Phe Asn Ile Asp Asp
755 760 765

Leu Ser Ser Lys Leu Asn Glu Ser Ile Asn Lys Ala Met Ile Asn Ile
770 775 780

Asn Lys Phe Leu Asn Gln Cys Ser Val Ser Tyr Leu Met Asn Ser Met
785 790 795 800

Ile Pro Tyr Gly Val Lys Arg Leu Glu Asp Phe Asp Ala Ser Leu Lys
805 810 815

Asp Ala Leu Leu Lys Tyr Ile Tyr Asp Asn Arg Gly Thr Leu Ile Gly
820 825 830

Gln Val Asp Arg Leu Lys Asp Lys Val Asn Asn Thr Leu Ser Thr Asp
835 840 845

Ile Pro Phe Gln Leu Ser Lys Tyr Val Asp Asn Gln Arg Leu Leu Ser
850 855 860

Thr Phe Thr Glu Tyr Ile Lys Asn Ile Ile Asn Thr Ser Ile Leu Asn
865 870 875 880

Leu Arg Tyr Glu Ser Asn His Leu Ile Asp Leu Ser Arg Tyr Ala Ser
885 890 895

Lys Ile Asn Ile Gly Ser Lys Val Asn Phe Asp Pro Ile Asp Lys Asn
900 905 910

Gln Ile Gln Leu Phe Asn Leu Glu Ser Ser Lys Ile Glu Val Ile Leu
915 920 925

Lys Asn Ala Ile Val Tyr Asn Ser Met Tyr Glu Asn Phe Ser Thr Ser
930 935 940

Phe Trp Ile Arg Ile Pro Lys Tyr Phe Asn Ser Ile Ser Leu Asn Asn
945 950 955 960

Glu Tyr Thr Ile Ile Asn Cys Met Glu Asn Asn Ser Gly Trp Lys Val
965 970 975

Ser Leu Asn Tyr Gly Glu Ile Ile Trp Thr Leu Gln Asp Thr Gln Glu
980 985 990

Ile Lys Gln Arg Val Val Phe Lys Tyr Ser Gln Met Ile Asn Ile Ser
995 1000 1005

Asp Tyr Ile Asn Arg Trp Ile Phe Val Thr Ile Thr Asn Asn Arg

eof-seq1

1010						1015					1020			
Leu	Asn	Asn	Ser	Lys	Ile	Tyr	Ile	Asn	Gly	Arg	Leu	Ile	Asp	Gln
	1025					1030					1035			
Lys	Pro	Ile	Ser	Asn	Leu	Gly	Asn	Ile	His	Ala	Ser	Asn	Asn	Ile
	1040					1045					1050			
Met	Phe	Lys	Leu	Asp	Gly	Cys	Arg	Asp	Thr	His	Arg	Tyr	Ile	Trp
	1055					1060					1065			
Ile	Lys	Tyr	Phe	Asn	Leu	Phe	Asp	Lys	Glu	Leu	Asn	Glu	Lys	Glu
	1070					1075					1080			
Ile	Lys	Asp	Leu	Tyr	Asp	Asn	Gln	Ser	Asn	Ser	Gly	Ile	Leu	Lys
	1085					1090					1095			
Asp	Phe	Trp	Gly	Asp	Tyr	Leu	Gln	Tyr	Asp	Lys	Pro	Tyr	Tyr	Met
	1100					1105					1110			
Leu	Asn	Leu	Tyr	Asp	Pro	Asn	Lys	Tyr	Val	Asp	Val	Asn	Asn	Val
	1115					1120					1125			
Gly	Ile	Arg	Gly	Tyr	Met	Tyr	Leu	Lys	Gly	Pro	Arg	Gly	Ser	Val
	1130					1135					1140			
Met	Thr	Thr	Asn	Ile	Tyr	Leu	Asn	Ser	Ser	Leu	Tyr	Arg	Gly	Thr
	1145					1150					1155			
Lys	Phe	Ile	Ile	Lys	Lys	Tyr	Ala	Ser	Gly	Asn	Lys	Asp	Asn	Ile
	1160					1165					1170			
Val	Arg	Asn	Asn	Asp	Arg	Val	Tyr	Ile	Asn	Val	Val	Val	Lys	Asn
	1175					1180					1185			
Lys	Glu	Tyr	Arg	Leu	Ala	Thr	Asn	Ala	Ser	Gln	Ala	Gly	Val	Glu
	1190					1195					1200			
Lys	Ile	Leu	Ser	Ala	Leu	Glu	Ile	Pro	Asp	Val	Gly	Asn	Leu	Ser
	1205					1210					1215			
Gln	Val	Val	Val	Met	Lys	Ser	Lys	Asn	Asp	Gln	Gly	Ile	Thr	Asn
	1220					1225					1230			
Lys	Cys	Lys	Met	Asn	Leu	Gln	Asp	Asn	Asn	Gly	Asn	Asp	Ile	Gly
	1235					1240					1245			
Phe	Ile	Gly	Phe	His	Gln	Phe	Asn	Asn	Ile	Ala	Lys	Leu	Val	Ala
	1250					1255					1260			
Ser	Asn	Trp	Tyr	Asn	Arg	Gln	Ile	Glu	Arg	Ser	Ser	Arg	Thr	Leu

eof-seq1

1265

1270

1275

Gly Cys Ser Trp Glu Phe Ile Pro Val Asp Asp Gly Trp Gly Glu
 1280 1285 1290

Arg Pro Leu
 1295

<210> 2
 <211> 1291
 <212> PRT
 <213> Clostridium botulinum

<400> 2

Met Pro Ile Thr Ile Asn Asn Phe Asn Tyr Ser Asp Pro Val Asp Asn
 1 5 10 15

Lys Asn Ile Leu Tyr Leu Asp Thr His Leu Asn Thr Leu Ala Asn Glu
 20 25 30

Pro Glu Lys Ala Phe Arg Ile Thr Gly Asn Ile Trp Val Ile Pro Asp
 35 40 45

Arg Phe Ser Arg Asn Ser Asn Pro Asn Leu Asn Lys Pro Pro Arg Val
 50 55 60

Thr Ser Pro Lys Ser Gly Tyr Tyr Asp Pro Asn Tyr Leu Ser Thr Asp
 65 70 75 80

Ser Asp Lys Asp Pro Phe Leu Lys Glu Ile Ile Lys Leu Phe Lys Arg
 85 90 95

Ile Asn Ser Arg Glu Ile Gly Glu Glu Leu Ile Tyr Arg Leu Ser Thr
 100 105 110

Asp Ile Pro Phe Pro Gly Asn Asn Asn Thr Pro Ile Asn Thr Phe Asp
 115 120 125

Phe Asp Val Asp Phe Asn Ser Val Asp Val Lys Thr Arg Gln Gly Asn
 130 135 140

Asn Trp Val Lys Thr Gly Ser Ile Asn Pro Ser Val Ile Ile Thr Gly
 145 150 155 160

Pro Arg Glu Asn Ile Ile Asp Pro Glu Thr Ser Thr Phe Lys Leu Thr
 165 170 175

Asn Asn Thr Phe Ala Ala Gln Glu Gly Phe Gly Ala Leu Ser Ile Ile
 180 185 190

Ser Ile Ser Pro Arg Phe Met Leu Thr Tyr Ser Asn Ala Thr Asn Asp
 195 200 205

eof-seq1

Val Gly Glu Gly Arg Phe Ser Lys Ser Glu Phe Cys Met Asp Pro Ile
 210 215 220
 Leu Ile Leu Met His Glu Leu Asn His Ala Met His Asn Leu Tyr Gly
 225 230 235 240
 Ile Ala Ile Pro Asn Asp Gln Thr Ile Ser Ser Val Thr Ser Asn Ile
 245 250 255
 Phe Tyr Ser Gln Tyr Asn Val Lys Leu Glu Tyr Ala Glu Ile Tyr Ala
 260 265 270
 Phe Gly Gly Pro Thr Ile Asp Leu Ile Pro Lys Ser Ala Arg Lys Tyr
 275 280 285
 Phe Glu Glu Lys Ala Leu Asp Tyr Tyr Arg Ser Ile Ala Lys Arg Leu
 290 295 300
 Asn Ser Ile Thr Thr Ala Asn Pro Ser Ser Phe Asn Lys Tyr Ile Gly
 305 310 315 320
 Glu Tyr Lys Gln Lys Leu Ile Arg Lys Tyr Arg Phe Val Val Glu Ser
 325 330 335
 Ser Gly Glu Val Thr Val Asn Arg Asn Lys Phe Val Glu Leu Tyr Asn
 340 345 350
 Glu Leu Thr Gln Ile Phe Thr Glu Phe Asn Tyr Ala Lys Ile Tyr Asn
 355 360 365
 Val Gln Asn Arg Lys Ile Tyr Leu Ser Asn Val Tyr Thr Pro Val Thr
 370 375 380
 Ala Asn Ile Leu Asp Asp Asn Val Tyr Asp Ile Gln Asn Gly Phe Asn
 385 390 395 400
 Ile Pro Lys Ser Asn Leu Asn Val Leu Phe Met Gly Gln Asn Leu Ser
 405 410 415
 Arg Asn Pro Ala Leu Arg Lys Val Asn Pro Glu Asn Met Leu Tyr Leu
 420 425 430
 Phe Thr Lys Phe Cys His Lys Ala Ile Asp Gly Arg Ser Leu Tyr Asn
 435 440 445
 Lys Thr Leu Asp Cys Arg Glu Leu Leu Val Lys Asn Thr Asp Leu Pro
 450 455 460
 Phe Ile Gly Asp Ile Ser Asp Val Lys Thr Asp Ile Phe Leu Arg Lys
 465 470 475 480

e0lf-seq1

Asp Ile Asn Glu Glu Thr Glu Val Ile Tyr Tyr Pro Asp Asn Val Ser
485 490 495

Val Asp Gln Val Ile Leu Ser Lys Asn Thr Ser Glu His Gly Gln Leu
500 505 510

Asp Leu Leu Tyr Pro Ser Ile Asp Ser Glu Ser Glu Ile Leu Pro Gly
515 520 525

Glu Asn Gln Val Phe Tyr Asp Asn Arg Thr Gln Asn Val Asp Tyr Leu
530 535 540

Asn Ser Tyr Tyr Tyr Leu Glu Ser Gln Lys Leu Ser Asp Asn Val Glu
545 550 555 560

Asp Phe Thr Phe Thr Arg Ser Ile Glu Glu Ala Leu Asp Asn Ser Ala
565 570 575

Lys Val Tyr Thr Tyr Phe Pro Thr Leu Ala Asn Lys Val Asn Ala Gly
580 585 590

Val Gln Gly Gly Leu Phe Leu Met Trp Ala Asn Asp Val Val Glu Asp
595 600 605

Phe Thr Thr Asn Ile Leu Arg Lys Asp Thr Leu Asp Lys Ile Ser Asp
610 615 620

Val Ser Ala Ile Ile Pro Tyr Ile Gly Pro Ala Leu Asn Ile Ser Asn
625 630 635 640

Ser Val Arg Arg Gly Asn Phe Thr Glu Ala Phe Ala Val Thr Gly Val
645 650 655

Thr Ile Leu Leu Glu Ala Phe Pro Glu Phe Thr Ile Pro Ala Leu Gly
660 665 670

Ala Phe Val Ile Tyr Ser Lys Val Gln Glu Arg Asn Glu Ile Ile Lys
675 680 685

Thr Ile Asp Asn Cys Leu Glu Gln Arg Ile Lys Arg Trp Lys Asp Ser
690 695 700

Tyr Glu Trp Met Met Gly Thr Trp Leu Ser Arg Ile Ile Thr Gln Phe
705 710 715 720

Asn Asn Ile Ser Tyr Gln Met Tyr Asp Ser Leu Asn Tyr Gln Ala Gly
725 730 735

Ala Ile Lys Ala Lys Ile Asp Leu Glu Tyr Lys Lys Tyr Ser Gly Ser
740 745 750

e0lf-seq1

Asp Lys Glu Asn Ile Lys Ser Gln Val Glu Asn Leu Lys Asn Ser Leu
755 760 765

Asp Val Lys Ile Ser Glu Ala Met Asn Asn Ile Asn Lys Phe Ile Arg
770 775 780

Glu Cys Ser Val Thr Tyr Leu Phe Lys Asn Met Leu Pro Lys Val Ile
785 790 795 800

Asp Glu Leu Asn Glu Phe Asp Arg Asn Thr Lys Ala Lys Leu Ile Asn
805 810 815

Leu Ile Asp Ser His Asn Ile Ile Leu Val Gly Glu Val Asp Lys Leu
820 825 830

Lys Ala Lys Val Asn Asn Ser Phe Gln Asn Thr Ile Pro Phe Asn Ile
835 840 845

Phe Ser Tyr Thr Asn Asn Ser Leu Leu Lys Asp Ile Ile Asn Glu Tyr
850 855 860

Phe Asn Asn Ile Asn Asp Ser Lys Ile Leu Ser Leu Gln Asn Arg Lys
865 870 875 880

Asn Thr Leu Val Asp Thr Ser Gly Tyr Asn Ala Glu Val Ser Glu Glu
885 890 895

Gly Asp Val Gln Leu Asn Pro Ile Phe Pro Phe Asp Phe Lys Leu Gly
900 905 910

Ser Ser Gly Glu Asp Arg Gly Lys Val Ile Val Thr Gln Asn Glu Asn
915 920 925

Ile Val Tyr Asn Ser Met Tyr Glu Ser Phe Ser Ile Ser Phe Trp Ile
930 935 940

Arg Ile Asn Lys Trp Val Ser Asn Leu Pro Gly Tyr Thr Ile Ile Asp
945 950 955 960

Ser Val Lys Asn Asn Ser Gly Trp Ser Ile Gly Ile Ile Ser Asn Phe
965 970 975

Leu Val Phe Thr Leu Lys Gln Asn Glu Asp Ser Glu Gln Ser Ile Asn
980 985 990

Phe Ser Tyr Asp Ile Ser Asn Asn Ala Pro Gly Tyr Asn Lys Trp Phe
995 1000 1005

Phe Val Thr Val Thr Asn Asn Met Met Gly Asn Met Lys Ile Tyr
1010 1015 1020

e0lf-seq1

Ile Asn Gly Lys Leu Ile Asp Thr Ile Lys Val Lys Glu Leu Thr
1025 1030 1035

Gly Ile Asn Phe Ser Lys Thr Ile Thr Phe Glu Ile Asn Lys Ile
1040 1045 1050

Pro Asp Thr Gly Leu Ile Thr Ser Asp Ser Asp Asn Ile Asn Met
1055 1060 1065

Trp Ile Arg Asp Phe Tyr Ile Phe Ala Lys Glu Leu Asp Gly Lys
1070 1075 1080

Asp Ile Asn Ile Leu Phe Asn Ser Leu Gln Tyr Thr Asn Val Val
1085 1090 1095

Lys Asp Tyr Trp Gly Asn Asp Leu Arg Tyr Asn Lys Glu Tyr Tyr
1100 1105 1110

Met Val Asn Ile Asp Tyr Leu Asn Arg Tyr Met Tyr Ala Asn Ser
1115 1120 1125

Arg Gln Ile Val Phe Asn Thr Arg Arg Asn Asn Asn Asp Phe Asn
1130 1135 1140

Glu Gly Tyr Lys Ile Ile Ile Lys Arg Ile Arg Gly Asn Thr Asn
1145 1150 1155

Asp Thr Arg Val Arg Gly Gly Asp Ile Leu Tyr Phe Asp Met Thr
1160 1165 1170

Ile Asn Asn Lys Ala Tyr Asn Leu Phe Met Lys Asn Glu Thr Met
1175 1180 1185

Tyr Ala Asp Asn His Ser Thr Glu Asp Ile Tyr Ala Ile Gly Leu
1190 1195 1200

Arg Glu Gln Thr Lys Asp Ile Asn Asp Asn Ile Ile Phe Gln Ile
1205 1210 1215

Gln Pro Met Asn Asn Thr Tyr Tyr Tyr Ala Ser Gln Ile Phe Lys
1220 1225 1230

Ser Asn Phe Asn Gly Glu Asn Ile Ser Gly Ile Cys Ser Ile Gly
1235 1240 1245

Thr Tyr Arg Phe Arg Leu Gly Gly Asp Trp Tyr Arg His Asn Tyr
1250 1255 1260

Leu Val Pro Thr Val Lys Gln Gly Asn Tyr Ala Ser Leu Leu Glu
1265 1270 1275

e0lf-seq1

Ser Thr Ser Thr His Trp Gly Phe Val Pro Val Ser Glu
 1280 1285 1290

<210> 3
 <211> 1252
 <212> PRT
 <213> Clostridium botulinum

<400> 3

Met Pro Lys Ile Asn Ser Phe Asn Tyr Asn Asp Pro Val Asn Asp Arg
 1 5 10 15

Thr Ile Leu Tyr Ile Lys Pro Gly Gly Cys Gln Glu Phe Tyr Lys Ser
 20 25 30

Phe Asn Ile Met Lys Asn Ile Trp Ile Ile Pro Glu Arg Asn Val Ile
 35 40 45

Gly Thr Thr Pro Gln Asp Phe His Pro Pro Thr Ser Leu Lys Asn Gly
 50 55 60

Asp Ser Ser Tyr Tyr Asp Pro Asn Tyr Leu Gln Ser Asp Glu Glu Lys
 65 70 75 80

Asp Arg Phe Leu Lys Ile Val Thr Lys Ile Phe Asn Arg Ile Asn Asn
 85 90 95

Asn Leu Ser Gly Gly Ile Leu Leu Glu Glu Leu Ser Lys Ala Asn Pro
 100 105 110

Tyr Leu Gly Asn Asp Asn Thr Pro Asp Asn Gln Phe His Ile Gly Asp
 115 120 125

Ala Ser Ala Val Glu Ile Lys Phe Ser Asn Gly Ser Gln Asp Ile Leu
 130 135 140

Leu Pro Asn Val Ile Ile Met Gly Ala Glu Pro Asp Leu Phe Glu Thr
 145 150 155 160

Asn Ser Ser Asn Ile Ser Leu Arg Asn Asn Tyr Met Pro Ser Asn His
 165 170 175

Gly Phe Gly Ser Ile Ala Ile Val Thr Phe Ser Pro Glu Tyr Ser Phe
 180 185 190

Arg Phe Asn Asp Asn Ser Met Asn Glu Phe Ile Gln Asp Pro Ala Leu
 195 200 205

Thr Leu Met His Glu Leu Ile His Ser Leu His Gly Leu Tyr Gly Ala
 210 215 220

eof-seq1

Lys Gly Ile Thr Thr Lys Tyr Thr Ile Thr Gln Lys Gln Asn Pro Leu
 225 230 235 240

Ile Thr Asn Ile Arg Gly Thr Asn Ile Glu Glu Phe Leu Thr Phe Gly
 245 250 255

Gly Thr Asp Leu Asn Ile Ile Thr Ser Ala Gln Ser Asn Asp Ile Tyr
 260 265 270

Thr Asn Leu Leu Ala Asp Tyr Lys Lys Ile Ala Ser Lys Leu Ser Lys
 275 280 285

Val Gln Val Ser Asn Pro Leu Leu Asn Pro Tyr Lys Asp Val Phe Glu
 290 295 300

Ala Lys Tyr Gly Leu Asp Lys Asp Ala Ser Gly Ile Tyr Ser Val Asn
 305 310 315 320

Ile Asn Lys Phe Asn Asp Ile Phe Lys Lys Leu Tyr Ser Phe Thr Glu
 325 330 335

Phe Asp Leu Ala Thr Lys Phe Gln Val Lys Cys Arg Gln Thr Tyr Ile
 340 345 350

Gly Gln Tyr Lys Tyr Phe Lys Leu Ser Asn Leu Leu Asn Asp Ser Ile
 355 360 365

Tyr Asn Ile Ser Glu Gly Tyr Asn Ile Asn Asn Leu Lys Val Asn Phe
 370 375 380

Arg Gly Gln Asn Ala Asn Leu Asn Pro Arg Ile Ile Thr Pro Ile Thr
 385 390 395 400

Gly Arg Gly Leu Val Lys Lys Ile Ile Arg Phe Cys Lys Asn Ile Val
 405 410 415

Ser Val Lys Gly Ile Arg Lys Ser Ile Cys Ile Glu Ile Asn Asn Gly
 420 425 430

Glu Leu Phe Phe Val Ala Ser Glu Asn Ser Tyr Asn Asp Asp Asn Ile
 435 440 445

Asn Thr Pro Lys Glu Ile Asp Asp Thr Val Thr Ser Asn Asn Asn Tyr
 450 455 460

Glu Asn Asp Leu Asp Gln Val Ile Leu Asn Phe Asn Ser Glu Ser Ala
 465 470 475 480

Pro Gly Leu Ser Asp Glu Lys Leu Asn Leu Thr Ile Gln Asn Asp Ala
 485 490 495

e0lf-seq1

Tyr Ile Pro Lys Tyr Asp Ser Asn Gly Thr Ser Asp Ile Glu Gln His
 500 505 510

Asp Val Asn Glu Leu Asn Val Phe Phe Tyr Leu Asp Ala Gln Lys Val
 515 520 525

Pro Glu Gly Glu Asn Asn Val Asn Leu Thr Ser Ser Ile Asp Thr Ala
 530 535 540

Leu Leu Glu Gln Pro Lys Ile Tyr Thr Phe Phe Ser Ser Glu Phe Ile
 545 550 555

Asn Asn Val Asn Lys Pro Val Gln Ala Ala Leu Phe Val Ser Trp Ile
 565 570 575

Gln Gln Val Leu Val Asp Phe Thr Thr Glu Ala Asn Gln Lys Ser Thr
 580 585 590

Val Asp Lys Ile Ala Asp Ile Ser Ile Val Val Pro Tyr Ile Gly Leu
 595 600 605

Ala Leu Asn Ile Gly Asn Glu Ala Gln Lys Gly Asn Phe Lys Asp Ala
 610 615 620

Leu Glu Leu Leu Gly Ala Gly Ile Leu Leu Glu Phe Glu Pro Glu Leu
 625 630 635

Leu Ile Pro Thr Ile Leu Val Phe Thr Ile Lys Ser Phe Leu Gly Ser
 645 650 655

Ser Asp Asn Lys Asn Lys Val Ile Lys Ala Ile Asn Asn Ala Leu Lys
 660 665 670

Glu Arg Asp Glu Lys Trp Lys Glu Val Tyr Ser Phe Ile Val Ser Asn
 675 680 685

Trp Met Thr Lys Ile Asn Thr Gln Phe Asn Lys Arg Lys Glu Gln Met
 690 695 700

Tyr Gln Ala Leu Gln Asn Gln Val Asn Ala Ile Lys Thr Ile Ile Glu
 705 710 715 720

Ser Lys Tyr Asn Ser Tyr Thr Leu Glu Glu Lys Asn Glu Leu Thr Asn
 725 730 735

Lys Tyr Asp Ile Lys Gln Ile Glu Asn Glu Leu Asn Gln Lys Val Ser
 740 745 750

Ile Ala Met Asn Asn Ile Asp Arg Phe Leu Thr Glu Ser Ser Ile Ser
 755 760 765

eof-seq1

Tyr Leu Met Lys Leu Ile Asn Glu Val Lys Ile Asn Lys Leu Arg Glu
 770 775 780

Tyr Asp Glu Asn Val Lys Thr Tyr Leu Leu Asn Tyr Ile Ile Gln His
 785 790 795 800

Gly Ser Ile Leu Gly Glu Ser Gln Gln Glu Leu Asn Ser Met Val Thr
 805 810 815

Asp Thr Leu Asn Asn Ser Ile Pro Phe Lys Leu Ser Ser Tyr Thr Asp
 820 825 830

Asp Lys Ile Leu Ile Ser Tyr Phe Asn Lys Phe Phe Lys Arg Ile Lys
 835 840 845

Ser Ser Ser Val Leu Asn Met Arg Tyr Lys Asn Asp Lys Tyr Val Asp
 850 855 860

Thr Ser Gly Tyr Asp Ser Asn Ile Asn Ile Asn Gly Asp Val Tyr Lys
 865 870 875 880

Tyr Pro Thr Asn Lys Asn Gln Phe Gly Ile Tyr Asn Asp Lys Leu Ser
 885 890 895

Glu Val Asn Ile Ser Gln Asn Asp Tyr Ile Ile Tyr Asp Asn Lys Tyr
 900 905 910

Lys Asn Phe Ser Ile Ser Phe Trp Val Arg Ile Pro Asn Tyr Asp Asn
 915 920 925

Lys Ile Val Asn Val Asn Asn Glu Tyr Thr Ile Ile Asn Cys Met Arg
 930 935 940

Asp Asn Asn Ser Gly Trp Lys Val Ser Leu Asn His Asn Glu Ile Ile
 945 950 955 960

Trp Thr Leu Gln Asp Asn Ala Gly Ile Asn Gln Lys Leu Ala Phe Asn
 965 970 975

Tyr Gly Asn Ala Asn Gly Ile Ser Asp Tyr Ile Asn Lys Trp Ile Phe
 980 985 990

Val Thr Ile Thr Asn Asp Arg Leu Gly Asp Ser Lys Leu Tyr Ile Asn
 995 1000 1005

Gly Asn Leu Ile Asp Gln Lys Ser Ile Leu Asn Leu Gly Asn Ile
 1010 1015 1020

His Val Ser Asp Asn Ile Leu Phe Lys Ile Val Asn Cys Ser Tyr
 1025 1030 1035

e0lf-seq1

Thr Arg Tyr Ile Gly Ile Arg Tyr Phe Asn Ile Phe Asp Lys Glu
 1040 1045 1050

Leu Asp Glu Thr Glu Ile Gln Thr Leu Tyr Ser Asn Glu Pro Asn
 1055 1060 1065

Thr Asn Ile Leu Lys Asp Phe Trp Gly Asn Tyr Leu Leu Tyr Asp
 1070 1075 1080

Lys Glu Tyr Tyr Leu Leu Asn Val Leu Lys Pro Asn Asn Phe Ile
 1085 1090 1095

Asp Arg Arg Lys Asp Ser Thr Leu Ser Ile Asn Asn Ile Arg Ser
 1100 1105 1110

Thr Ile Leu Leu Ala Asn Arg Leu Tyr Ser Gly Ile Lys Val Lys
 1115 1120 1125

Ile Gln Arg Val Asn Asn Ser Ser Thr Asn Asp Asn Leu Val Arg
 1130 1135 1140

Lys Asn Asp Gln Val Tyr Ile Asn Phe Val Ala Ser Lys Thr His
 1145 1150 1155

Leu Phe Pro Leu Tyr Ala Asp Thr Ala Thr Thr Asn Lys Glu Lys
 1160 1165 1170

Thr Ile Lys Ile Ser Ser Ser Gly Asn Arg Phe Asn Gln Val Val
 1175 1180 1185

Val Met Asn Ser Val Gly Asn Asn Cys Thr Met Asn Phe Lys Asn
 1190 1195 1200

Asn Asn Gly Asn Asn Ile Gly Leu Leu Gly Phe Lys Ala Asp Thr
 1205 1210 1215

Val Val Ala Ser Thr Trp Tyr Tyr Thr His Met Arg Asp His Thr
 1220 1225 1230

Asn Ser Asn Gly Cys Phe Trp Asn Phe Ile Ser Glu Glu His Gly
 1235 1240 1245

Trp Gln Glu Lys
 1250

<210> 4
 <211> 7
 <212> PRT
 <213> Artificial

<220>
 <223> Calpain recognition and/or cleavage site within BoNT/A

eolf-seq1

<400> 4

Gly Lys Phe Ala Thr Asp Pro
1 5

<210> 5

<211> 6

<212> PRT

<213> Artificial

<220>

<223> Calpain recognition and/or cleavage site within BoNT/A

<400> 5

Glu Leu Lys Val Ile Asp
1 5

<210> 6

<211> 6

<212> PRT

<213> Artificial

<220>

<223> Calpain recognition and/or cleavage site within BoNT/A

<400> 6

Glu Asp Thr Ser Gly Lys
1 5

<210> 7

<211> 6

<212> PRT

<213> Artificial

<220>

<223> Calpain recognition and/or cleavage site within BoNT/A

<400> 7

Gly Leu Glu Val Ser Phe
1 5

<210> 8

<211> 6

<212> PRT

<213> Artificial

<220>

<223> Calpain recognition and/or cleavage site within BoNT/A

<400> 8

Leu Asn Lys Ala Lys Ser
1 5

<210> 9

<211> 6

<212> PRT

<213> Artificial

eolf-seq1

<220>
<223> Calpain recognition and/or cleavage site within BoNT/A
<400> 9

Val Asp Lys Leu Lys Phe
1 5

<210> 10
<211> 6
<212> PRT
<213> Artificial

<220>
<223> Calpain recognition and/or cleavage site within BoNT/A
<400> 10

Val Leu Asn Arg Lys Thr
1 5

<210> 11
<211> 6
<212> PRT
<213> Artificial

<220>
<223> Calpain recognition and/or cleavage site within BoNT/A
<400> 11

Ile Val Gly Thr Thr Ala
1 5

<210> 12
<211> 6
<212> PRT
<213> Artificial

<220>
<223> Calpain recognition and/or cleavage site within BoNT/A
<400> 12

Gly Thr Thr Ala Ser Leu
1 5

<210> 13
<211> 6
<212> PRT
<213> Artificial

<220>
<223> Calpain recognition and/or cleavage site
<400> 13

Pro Leu Lys Ser Pro Pro
1 5

<210> 14
<211> 6
<212> PRT

eolf-seq1

<213> Artificial

<220>

<223> Calpain recognition and/or cleavage site within BoNT/E

<400> 14

Ile Lys Phe Ser Asn Gly
1 5