



(51) International Patent Classification:
C12Q 1/68 (2006.01)

(21) International Application Number:
PCT/EP2014/076044

(22) International Filing Date:
1 December 2014 (01.12.2014)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
13195330.9 2 December 2013 (02.12.2013) EP

(71) Applicant: **ALBERT-LUDWIGS-UNIVERSITÄT
FREIBURG** [DE/DE]; Fahnenbergplatz, 79085 Freiburg
(DE).

(72) Inventors: **NAZARENKO, Irina**; Buchsweilerstrasse 7,
79331 Teningen Nimburg (DE). **CATO, Andrew**; Lauter-
str. 4A, 76344 Eggenstein-Leopoldshafen (DE). **NEEB,
Antje**; Rheinstraße 37, 76479 Steinmauern (DE). **BOR-
MANN, Stefanie**; Volkhovener Weg 195, 50765 Köln
(DE). **SCHÖNTHALER, Martin**; Harriet-Straub-Str. 17,
79100 Freiburg (DE). **MIERNIK, Arkadiusz**; Starkenstr.
2, 79104 Freiburg (DE). **KRÖNIG, Malte**; Hirschen-
hofweg 6, 79117 Freiburg (DE). **WILHELM, Konrad**;
Bienenweg 6, 79110 Freiburg (DE). **HÄNSELMANN,
Benjamin**; Johanna-Kohlund-Straße 20, 79111 Freiburg
im Breisgau (DE). **HEFELE, Simon**; Günterstalstrasse 78,
79100 Freiburg (DE).

(74) Agents: **KELLER, Günter** et al.; Unsöldstr. 2, 80538
München (DE).

(81) Designated States (unless otherwise indicated, for every
kind of national protection available): AE, AG, AL, AM,
AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY,
BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM,
DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT,
HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR,
KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG,
MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM,
PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC,
SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN,
TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every
kind of regional protection available): ARIPO (BW, GH,
GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,
TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU,
TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE,
DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU,
LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK,
SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ,
GW, KM, ML, MR, NE, SN, TD, TG).

Published:

- with international search report (Art. 21(3))
- with sequence listing part of description (Rule 5.2(a))

(54) Title: PROCESS FOR THE DIAGNOSIS OF CANCER BY USING EXOSOMES

(57) Abstract: The present invention relates to an in vitro process for the diagnosis of prostate cancer and other tumor types in exo-
somes obtained from a body fluid which comprises a) concentrating the exosomes and other extracellular vesicles from a body fluid
of a patient, b) extraction of total RNA from the exosomes obtained in step a), c) conversion of the RNA obtained in step b) to
cDNA, d) amplification of the cDNA obtained in step c) with a polymerase chain reaction whereby primers derived from the AGR2
nucleotide sequence or the complement thereof are used, and e) determining whether in the amplification product variants of the
AGR2 gene can be identified.



Process for the Diagnosis of Cancer by Using Exosomes

Prostate cancer has become a major public health problem. In many developed countries it is not only the most commonly diagnosed malignancy but it is the second leading cause of cancer related deaths in males.

Since its discovery more than 20 years ago, prostate specific antigen (PSA) has been the most valuable tool in the detection, staging and monitoring of prostate cancer. Although widely accepted as a prostate tumour marker, PSA is known to be prostate tissue- but not prostate cancer-specific. PSA levels have been reported to increase in men with benign prostatic hyperplasia (BPH) and prostatitis. This substantial overlap in serum PSA values between men with non-malignant prostatic diseases and prostate cancer is the limitation of PSA as a prostate tumour marker. Moreover, a single reading of PSA cannot be used to differentiate the aggressive tumours from the indolent ones. Thus, (non-invasive) molecular tests that can accurately identify those men who have early stage, clinically localized prostate cancer, and who would gain prolonged survival and quality of life from early radical intervention are urgently needed.

Looking for new diagnostic, prognostic markers as well as understanding the molecular mechanisms underlying the disease progress and progression process and identify new treatment targets is the main focus for current prostate cancer research. For the identification of new candidate markers for prostate cancer it is necessary to study expression patterns in malignant as well as non-malignant prostate tissues. One of the genes identified to be overexpressed in prostate cancer is the anterior gradient 2 (AGR2) gene (Bu et al., The Prostate (2011), 575-587). The authors describe long and short transcripts of AGR2 (anterior gradient-2) being up-regulated in prostate cancer cells and tissues. RNAs were isolated from microdissected frozen prostate tissue samples obtained from radical prostatectomy specimens. Total RNA was amplified, reverse transcribed into cDNA and subjected to quantitative real-time PCR. Two transcript variants of the AGR2 gene were detected. Expression of short AGR2 transcript was up-regulated in tumors. Although AGR2 short transcript mRNA level showed slightly higher sensitivity and specificity as compared to the PSA serum level, no significant differences could be

observed. Therefore, mRNA level of the AGR2 short transcript in tissue or in the urine sediment are considered as not applicable for a reliable detection of prostate carcinoma.

Nilsson et al., British Journal of Cancer (2009) **100**, 1603-1607 analyze prostate cancer-derived urine exosomes as novel approach to biomarkers for prostate cancer. The authors describe the application of urine exosomes as a source for prostate cancer biomarkers. Thereby, mRNA levels of two known prostate cancer biomarkers designated as PCA-3 and TMPRSS2:ERG were tested in several patients. For the detection of corresponding mRNA a prostate massage was applied prior to urine harvesting which is not applicable in routine daily praxis. The article does, however, not provide data about the status of the tumor markers in tumor tissues. Consequently it is not possible to evaluate the diagnostic and/or prognostic values of the suggested biomarkers. A potential enrichment of the two biomarkers PCA-3 and TMPRSS2:ERG in urine exosomes seems to be at least questionable. Since the article shows only slightly higher specificity and sensitivity for prostate cancer detection as compared to the frequently used PSA serum level the test described in this article does not seem promising for the implementation into daily clinical routine.

The present invention as defined in more detail in the claims shows significantly better sensitivity and specificity of AGR2 splice variants as compared to the serum PSA. This makes the AGR2 splice variants as described herein highly specific and promising diagnostic markers.

AGR2 was first described in *Xenopus laevis* embryos where it induces the formation of the forebrain and the cement gland. In humans, it is predominantly found in mucus-secreting tissues or in endocrine cells. It is a member of the protein disulfide isomerase family of endoplasmic reticulum-resident proteins that has been implicated in the folding of proteins. It became of interest in cancer research, when it was found to be differential expressed in estrogen receptor (ER)-positive breast cancer cell lines. Since then, elevated AGR2 has been reported in subsets of breast, prostate, non-small cell lung, pancreatic, and hepatic adenocarcinomas. In some cases its increased expression has been shown to be of prognostic value. Furthermore elevated AGR2 RNA levels have been shown to correlate with decreased efficacy of tamoxifen treatment. AGR2 transcript level is therefore postulated to serve as a potential predictive biomarker for selection of optimal algorithm for adjuvant hormonal therapy in postmenopausal ER-positive breast cancer patients.

The AGR2 gene has been deposited at the Genebank and is available under accession number AF038451. The nucleotide sequence (SEQ ID NO:1) can be downloaded from the gene bank as well as the protein sequence (SEQ ID NO:2).

With the use of transcript-specific primers the expression of AGR2 transcript variants in microdissected prostate tissues was investigated. A real-time reverse transcription polymerase chain reaction (PCR) method to detect AGR2 in exosomes was established in order to test its potential as a predictive marker for prostate cancer and potentially in other tumors. AGR2 is highly expressed in prostate cancer especially in low-grade tumors and PIN lesions compared to benign tissue. AGR2 transcripts were also detected in urine sediments of patients undergoing prostate biopsy with significantly higher levels in tumor patients. The urine AGR2/PSA transcript ratio allowed much better discrimination between cancer and benign patients than serum total PSA or % free PSA suggesting that urine AGR2 transcripts could be a better marker for the detection of prostate cancer.

In the course of the present invention several new splice variants were cloned and sequenced. The AGR2 transcript variants are shown schematically in Figure 5. The AGR2 gene comprises 8 exons.

Variants of the AGR2 gene differ from the wild-type AGR2 gene insofar as either complete exons or parts of some exons are deleted. The new splice transcript variants were termed AGR2 SV-C, AGR2 SV-E, AGR2 SV-F, AGR2 SV-G and AGR2 SV-H. The variants are shown schematically in Figure 5 wherein as wild-type AGR2 the variant B containing shorter part of the untranslated region of the first exon, is designated. In variants C exon 1 is shortened. Variant E comprises exons 1, 5, 7 and 8. Variant F comprises exons 1, 7 and 8. Variant G comprises part of exon 1 and part of exon 7 and exon 8. Variant H comprises exon 2, part of exon 3 and part of exon 8.

As shown in the examples below, exosomes isolated from the urine or from the blood of patients can be used as a source for the detection of the biomarkers (ARG2 transcript variants), which allows the application of a non-invasive or only minimally invasive diagnostic routine method. When exosomes are isolated from the urine of patients, the diagnostic method can be performed without any treatment performed by a medical doctor.

Surprisingly, it has been found that distinct ARG2 splice variants recovered from the exosomes can be used as specific diagnostic and/or prognostic markers for different tumor entities. Such tumors comprise not only the preferred prostate tumor but also breast cancer or cancer of colon, lung, pancreas or small intestine to name only a few.

Splice variants E and G (compare with Figure 5) are enriched in the exosomes isolated from the blood of patients with prostate cancer but not in the cells isolated from the blood. This can be seen from Figure 2A.

Splice variants E, F, G and H (compare with Figure 5) are enriched in the exosomes isolated from urine or patients with prostate cancer compared to the urine sediment. This confirms the superior results obtainable by the method of the present invention.

A pilot clinical study confirmed that splice variants G and H are enriched in the exosomes isolated from urine of prostate cancer patients and are significantly better diagnostic markers as compared to the serum PSA. This can be seen from Figures 3A and 3C of the present application.

In the present invention the following primers were preferred:

SV C: Forward: 5' CAC AAG GCA GAG TTG CCA TGG 3' (SEQ ID NO:3)
SV E: Forward: 5' ATC TGG TCA CCC ATC TCT GA 3' (SEQ ID NO:4)
SV F: Forward: 5' GGA AAT CCA GAC CCA TCT CTG 3' (SEQ ID NO:5)
SV G: Forward: 5' AAG GCA GGT ACA GCT CTG 3' (SEQ ID NO:6)
Reverse SV C, E, F G: 5' TCC ACA CTA GCC AGT CTT CTC A 3' (SEQ ID NO:7)

SV H Forward: 5' ATG GAG AAA ATT CCA GTG TCA GCA 3' (SEQ ID NO:8)
Reverse: 5'- ACT TGA GAG CTT TCT TCA TAT GTC TG-3' (SEQ ID NO:9)

AGR2 long transcript: A (was not detected in the exosomes)
AGR2Lt Forward GCCAACAGACAACCCAAAGT (SEQ ID NO:10)
AGR2Lt Reverse GCAAGAATGCTGACACTGGA (SEQ ID NO:11)

AGR2 short transcript: B (also designated as a wt AGR2)
AGR2St Forward -CGACTCACACAAGGCAGGT- (SEQ ID NO:12)
AGR2St Reverse -GCAAGAATGCTGACACTGGA (SEQ ID NO:13)

PSA:
PSA Forward ACCAGAGGAGTTCTTGACCCCA (SEQ ID
NO:14)
PSA Reverse CCCCAGAATCACCCGAGCAG
(SEQ ID NO:15)

By selection of appropriate forward and reverse primers specific variants could be amplified. The person skilled in the art can, however, easily design alternative primers since the splice variants are described herein and the complete sequence of the gene to be analyzed is known.

Exosomes (extracellular vesicles, EV) are specialized nanovesicles (30–100 nm) that are actively secreted by a variety of normal and tumor cells. Elevated exosome amount has been found in malignancy effusions, serum and urine from cancer patients. Certain RNA transcripts are enriched several 100-fold in the exosomes compared with the donor cells and in particular exosomes are enriched in unique transcripts specific to tumor cells that may be below detection limit even in the tumor cells themselves. Therefore, the mRNA levels of *svB*, *svC*, *svE*, *svF*, *svG* and *svH* (shown in Figure 5) variant of *AGR2* in urine exosomes was analyzed, in order to define the most reliable source of the *AGR2* variant as biomarkers for prostate cancer.

The present invention provides an *in vitro* process for the diagnosis of prostate cancer in exosomes. The term "exosome" is used as a term for "extracellular vesicles". The term "diagnosis" means to obtain more insight in the specific forms of cancer and whether the patient suffers from an aggressive form of prostate cancer or whether possibly other diagnostic markers speak for a benign form which does not require surgery or treatment with strong chemostatic agents. It is an advantage of the present invention that the diagnosis can be performed in a sample obtained from the patient without the action of a doctor. The sample which is used for the diagnostic process is preferably urine. The exosomes can, however, also be obtained from other sources like from bloods serum or plasma. In general the diagnostic method can be performed with fluids obtained from the human body. Body fluids comprise urine, semen, blood (serum or plasma), saliva and any other fluid which can be obtained from the human body.

The diagnostic method of the present invention is performed on exosomes. Therefore, it is essential that the exosomes are enriched and the other components or the fluid containing

the exosomes are at least partially removed. Since the detection step is based on the amplification of nucleic acids it is essential that nucleic acids which might interfere with the test results are efficiently removed. Protein components may strongly interfere with the test results when such components inhibit the enrichment of exosomes. Such proteins may for example entrap the exosomes and the nucleic acids contained therein cannot be made accessible to the amplification reaction.

In the following a preferred process for the diagnosis of prostate cancer is described when the body fluid is urine. One of the advantages of using as starting material for diagnosis urine is that the sample can be obtained from the patient without any invasive interaction of a doctor. Moreover, in some cases it may be difficult to obtain a sufficient amount of serum or plasma from a patient. To obtain a sample via biopsy from patients may be even more difficult and may increase the risk to cause metastases by circulating tumor cells or activation of dormant tumor cells.

In the first step of the diagnostic process the exosomes are concentrated from the body fluid, in particular urine of the patient. In a preferred embodiment the urine is first centrifuged at a low speed (preferably up to 4,000 G) at a low temperature (2-15°C) for 5 to 30 minutes in order to remove remaining cells and debris.

Preferably the supernatant obtained from the first centrifugation step is filtered through suitable filtering means in order to remove remaining large particles from the supernatant. The filter pores preferably have a diameter of about 1 μm to about 150 μm , more preferred about 5 μm to 100 μm .

The urine which has been treated by a low speed centrifugation and possibly an additional filtration step is then preferably ultracentrifuged. In a particularly preferred embodiment the method of the present invention comprises the step of ultracentrifugation. This purification step results in superior test results. The ultracentrifugation can take place either in the presence of a commonly used buffering agent. The ultracentrifugation is usually performed at +4°C and at $G \geq 100,000$ for 20 minutes, preferably from one hour to two hours. As an alternative the samples can be purified by chromatography- or precipitation-based approaches. It is important to enrich the extracellular vesicles with suitable purification steps from the body fluids.

The sediment obtained in the ultracentrifugation step is then further treated to extract the RNA from the exosomes. It is preferred to treat the sediment with proteases and DNAses or other suitable chemicals in order to extract the RNA from the sediment as efficiently as possible.

For the analysis of the RNA fractions obtained from this step the RNA is first transcribed into cDNA with the help of a reverse transcription.

The cDNA obtained in this manner is amplified with a polymerase chain reaction whereby specific primers developed in the course of the present invention are preferably used. From the nucleotide sequence of *AGR2* (SEQ ID NO:1) and the complement thereof suitable parts are selected. Then it has to be checked whether said primers do potentially amplify other sequences which may result in false-positive reactions. In the course of the present invention the following primers are especially preferred: SEQ ID NO:3 to SEQ ID NO:13 which allow the amplification of variants of distinct splice variants of *AGR2*. To avoid artifacts, for a PCR reaction usage of 2 different primer pairs or preferably 3 primer pairs is more preferred

After having performed the PCR reaction the amplification products are analyzed. It is well-known in the art, how such analysis can be performed. In a preferred embodiment the nucleic acid is amplified in a quantitative PCR.

Depending on the results the diagnosis can be performed. The presence or absence of specific variants of the *AGR2* gene allows a diagnosis whether the patient suffers from a malignant prostatic cancer or whether the clinical symptoms are benign.

The experimental results show that mRNAs coding for isoforms B, E, G and H are significantly elevated in tumor patients compared with samples obtained from patients with benign forms according to the student $-t$ test. The isoforms G and H demonstrated significantly higher specificity and sensitivity compared to the currently applied standard PSA test. Therefore, it is preferred to detect the presence and/or absence of the isoforms G, H, B and E either alone or in combination.

Since the diagnostic process of the present invention is also based on the observation whether certain variants are overly expressed or expressed only to a very low level it is

preferred to standardize the test method. This can be done for example by using always the same concentration of RNA in the test. It is preferred to standardize the RNA level to 10-50 ng, preferably to 30 ng per sample to be amplified.

Another alternative to improve the reliability of the diagnostic result is to amplify together with the AGR2 variants also a housekeeping gene. A so-called "housekeeping gene" may serve as internal standard which allows a comparison of results obtained in different experiments. Such as housekeeping gene is preferably the gene GAPDH which is present in all samples. Of course other housekeeping genes, recruited to the exosomes can also be used. In case PSA is amplified, suitable primers are SEQ ID NO:14 and SEQ ID NO:15.

It is one object of the present invention to determine whether splice variants of the AGR2 gene can be amplified and thereafter be assigned to certain types of diseases, in particular different tumors, and furthermore whether nucleic acid coding for certain splice variants can be detected in exosomes obtained from the urine of patients.

AGR2 has also been shown to be expressed in several human tissues rich in epithelial cells, like prostate, breast, small intestine, colon, lung, and pancreas. Its protein level was found increased in prostate tumor tissue and its expression level was associated with poor survival of prostate cancer patients. Therefore, the method of the present invention may serve as means for a prediction of the further development of the disease and as means for differentiating one type of tumor from another.

In breast and prostate adenocarcinoma, AGR2 seems to be under the control of ER or androgen receptor expression. Up-regulation of AGR2 enhances classical hallmarks of cancer, such as metastasis, invasion, colony formation, and proliferation.

The present invention is further illustrated in the following examples whereby the results are shown in the enclosed Figures. Important aspects of the present invention can be summarized as follows:

Splice variants E and G are enriched in exosomes isolated from the blood of patients suffering from prostate cancer. Such splice variants were, however, not detected in cells isolated from blood (compare with Figure 2A).

Splice variants E, F, G and H are enriched in exosomes isolated from urine of patients suffering from prostate cancer as compared to the urine sediment. The result that the exosomes allow superior diagnostic results could not have been expected. This can be seen from Figure 2B.

Clinical pilot studies confirmed that splice variants G and H are enriched in the exosomes isolated from urine of patients suffering from prostate cancer patients. Those splice variants are significantly better diagnostic markers as compared to serum PSA. This can be seen from Figure 3A or 3C, respectively.

The analysis of ARG2 splice variants from the exosomes obtained from blood and/or urine is tumor type specific. Figure 6 for example shows the enrichment of AGR2 splice variants in breast cancer exosomes which may allow the distinction between breast and prostate cancer.

Splice variant C is present in breast cancer exosomes but absent in prostate cancer whereas variant A is present in the exosomes derived from patients suffering from prostate cancer but absent in exosomes obtained from patients suffering from breast cancer. Even if prostate cancer and breast cancer are highly specific to gender such distinction may also be useful in the molecular diagnostic of other cancer types.

The present invention is summarized in the Figures and further illustrated in the following examples:

Figure 1 *Characterization of extracellular vesicles (EV) isolated from the plasma and urine of patients with colorectal carcinoma*

EVs isolated from the blood plasma (A) and urine (B) by differential centrifugation were characterized by transmission electron microscopy (upper panels) and dynamic light scattering analysis (Zetasizer, Malvern Instruments, Germany). To detect different population of vesicles, three main parameter were applied: distribution by intensity (left panel), allowing to visualize vesicles according to the intensity of their fluorescence independent on number; distribution by mass (right panel) according to which the amount of vesicles with certain mass can be measured, and distribution by number (table below), showing the relative amount of vesicles of certain size. (A) Three populations of vesicles

were detected in the blood plasma (left panel). According to the distribution by intensity first population (15% of total amount) consists of EVs with the radius 4,5 nm; second population (43%) consists of EVs with the radius of 23,6 nm which be the size corresponding to exosomes; 41% of total EVs comprise a third population of EVs with the mean radius of about. 254,2 nm and might contain bigger vesicles or agglomerates. The analysis of the distribution of EVs by mass clearly shows that the majority of EVs (95%) is represented by the population of vesicles with the radius of 4.7 nm which might correspond to liposomes. Only a minor part of EVs (4,6%) consists of EVs with the radius of 43 nm, which might correspond to exosomes. Analysis of the distribution by number show that the vesicles with the radius of 4,7 nm represent 99,9%, and the vesicles with the radius of 23,6 nm, which correspond the size of the exosomes, represent 0,09%, and the vesicles with the radius of 254,2 nm comprise about 0,01% of the whole amount.

(Figure 1B) Vesicles isolated from the urine represented a homogenous population with the average radius of about 74 nm, which as contain EVs with the size, corresponding exosomes, and bigger vesicles as it is shown by electron microscopy (upper panel).

Figure 2 *Comparative analysis of the AGR2wt and the splice variants svC, svE, svF, svG, svH mRNA levels in different components of the blood plasma and urine of patients with prostate cancer.*

For this analysis 5 patients with prostate adenocarcinoma were chosen. RNA was isolated either from the total blood plasma or from the EVs isolated from the blood plasma (A).

As a second source of AGR2 and its splice variants, urine sediment and urine EVs were examined (Figure 2B).

RNA was isolated from these four sources and subjected to the quantitative RT-PCR. GAPDH was used as a reference gene for the normalization of signal intensity.

The expression of AGR2 and the splice variants differ between total plasma and plasma vesicles. It is likely that the splice variant AGR2 SV-E might be specifically enriched in the plasma EVs of patients with the prostate cancer. However the analysis of urine EVs delivers more consistent results, supporting application of urine EVs for further analysis.

(Figure 2C) First, it was verified if the population of urine EVs isolated by differential centrifugation contains exosomes. Urine from the patients 1, 2, and 3 was used for the isolation of proteins from the cell sediment, designated as Se; and from the vesicles, designated as EV.

As exosome markers Tsg101, CD9 and Hsp70 were used. GAPDH served as a loading control. Additionally, the presence of prostate specific proteins PSMA was analyzed. In all three patients analyzed PSMA was present on the vesicles isolated from the urine, supporting their origin from the prostate.

Figure 3 *mRNA levels of AGR2 wt and the splice variants in the urine EVs can serve as an independent diagnostic marker for prostate cancer.*

39 patients (24 tumors; 15 BPH) with 15 benign and 24 malign adenocarcinoma were included into analysis. EVs were isolated from the urine and were directly used for RNA preparation. 30 ng of RNA were used for the followed quantitative RT-PCR analysis.

(Figure 3A) AGR2 and all splice variants can serve as potential diagnostic markers allowing to differentiate between benign (BPH) and adenocarcinomas. Additionally, potential predictive value of AGR2 and its splice variants was controlled.

(Figure 3B) Additionally, The AGR2 svF was demonstrated to show a difference between high and low Gleason scores. Further analysis of a bigger cohort of patients will verify this correlation.

(Figure 3C) ROC (receiver operator characteristic) plots of the AGR2 variants on the exosomes and PSA on the exosome and protein level in serum (analysis regularly used in the diagnostics) was performed with 27 patients (12 tumor and 9 benign) and revealed significant differences between AGR2 svG and PSA level in serum (p 0,044); AGR2 svG and PSA mRNA in the exosomes (p 0,0195), AGR2svH and PSA in serum (p 0,0269) and AGR2svH and PSA level in the vesicles (0,0274) confirming that the exosome AGR2 can serve as an additional independent diagnostic marker for the prostate cancer allowing to differ between benign disease and prostate cancer with the higher specificity as common markers (e.g. PSA).

Figure 4 *mRNA of AGR2 wt and the splice variants are most probably enclosed in the EVs and are resistant to the RNase treatment.*

To verify if detected mRNAs of AGR2 and its splice variants are indeed enclosed into vesicles, the vesicles from 2 independent samples were treated with RNase. After treatment the RNase was inactivated and the samples were subjected to the RNA isolation followed by the Q RT-PCR. The results showed that the AGR2 and splice variants mRNAs are resistant to the RNase treatment, which support their localization within the EVs.

Figure 5 *shows schematically the structure of different splice variants of the AGR2 gene.*

The AGR2 gene comprises 8 exons. The exons are shown schematically and the different variants are designated as B-H. A (upper line) corresponds to wild type. It is shown which exon or which part of an exon is deleted in splice variants B to H.

Example 1: Clinical samples

The use of clinical samples for the study was approved by the ethics committee of the Innsbruck Medical University. Frozen prostate tissue samples from previously untreated patients who had undergone radical prostatectomy after tumor diagnosis in a PSA-based prostate cancer early detection program were obtained from the prostate center of the Department of Urology of the Innsbruck Medical University. Frozen tissue samples were processed and microdissected and RNA was isolated. Tumor samples were obtained from a cohort of Gleason score (GSC) 6 tumors (Gleason pattern 3) and a group of GSC 8–10 tumors (Gleason patterns 4 and 5). Benign epithelial cell samples were isolated from the same specimens apart from tumor loci.

Example 2: Patients with clinical diagnosis

This example included 30 patients, with prostate cancer and with benign prostate hyperplasia. Analysed were in each case a urine sample and a blood sample (serum).

The cancer cases were all detected by biopsy. Before prostatic specific antigen (PSA) was measured and patients with a high PSA-level were advised to undergo a transrectal guided

biopsy. Their age was in between 28 and 78. As regards the histologic classification only patients with a gleason score from 6 to 9 were chosen.

All patients with prostate cancer were treated by radical prostatectomy and pelvic lymphadenectomy.

The patients with benign prostate hyperplasia with a low PSA-level (<4ng/ml) were treated by transurethral resection. If the PSA-level of these patients was high prostate cancer was obviated by transrectal guided biopsy following by transurethral resection.

Example 3: Splice variants of AGR2

AGR2 has a fairly broad expression pattern in human tissues. However splice variants of AGR2 (e.g. the long form, $\Delta 6$ and $\Delta 4-6$) are reported to provide some selectivity by being predominantly expressed in certain tissues or in distinct hepatocellular neoplasms. It was therefore investigated whether further splice variants of AGR2 exist and whether they would be a better predictor of prostate cancer development.

In PCR amplification reactions which were carried out with primers in exon I and 8, and 6 new splice transcripts were identified which were termed sv C, E, F, G, H in addition to transcripts A and B that have already been reported as long and short transcripts of AGR2 (compare with Figure 5). The transcripts C-H most likely arose from alternative splicing as they lack partial or complete exons. In contrast to transcripts A and B that are known to encode the full length AGR2 protein, no protein products for transcripts C, E, F, G and H have been detected so far.

Example 4

As a further characterization of the splice variants, their expression pattern was examined in different cell lines to find out whether they would differentiate between cells of prostate from cells of non-prostate origin. For prostate cells, VCaP, LNCaP, PNT2 and 22Rv.1 cell lines were used whereas the non-prostate cells which were analyzed consisted of breast tumor (T47D, MCF7), endometrial tumor (Ishikawa), cervical tumor (HeLa), kidney tumor (Hek 293), liver tumor (HepG2) and choriocarcinoma cells (JEG-3). AGR2 (short form transcript; svB) was expressed in several cell lines in agreement with reports on its

widespread expression. In contrast, it was found that the AGR2 long isoform (sv A) which was previously reported to be expressed in predominantly prostate tissue, was not only expressed in the prostate tumor cell lines but also in mammary (MCF7 and T47D) and endometrial cell lines (Ishikawa). The sv C, E, F, G and H showed varying levels of expression among the different cell lines with sv C being mainly expressed in prostate tumor cell lines. This differential expression pattern of the splice variants and their levels can be used to differentiate in the progression of prostate cancer from the benign to the more advanced stages of the tumor.

Example 5: AGR2 splice variants in prostate cancer progression

A real-time qPCR was performed for the different AGR2 transcripts on prostate biopsies which consisted of 32 benign biopsies and 32 tumor samples (16 from Gleason pattern 3 and 16 from Gleason pattern 4). In the quantitative RT-PCR the splice variant sv B (short transcript) again showed a significant difference ($p = 0.0406$) between benign and prostate tumor. In addition sv H was also identified as a predictive discriminator of benign and prostate tumor ($p = 0.0476$). All the other splice variants did not show any significant difference in the two sample pools. Since non-invasive diagnostics (using urine or saliva samples) are superior alternatives to traditional needle or excision biopsies due to the reduced patient pain and inconvenience, and greater speed and lower cost of analysis, it was decided to examine the expression of the splice variants in urine sediments.

Example 6: Urine exosomes for studying AGR2 expression

AGR2 (svB) splice variant is present in urine sediments of patients with prostate cancer and their levels correlate with the tumor aggressiveness allowing discrimination between benign and malignant neoplasm outperforming markers currently used for the diagnostic of prostate cancer. It was tested whether newly identified splice variants are also present in urine sediments and these studies were extended to determine whether they are also present in urine exosomes.

Urine extracellular membrane vesicles were isolated by differential centrifugation and characterized by dynamic light scattering. This showed that the urine exosomes are of sizes 74.0 ± 85.9 nm. The expression of the AGR2 splice variants in urine sediments and urine exosomes in 5 patients diagnosed with prostate carcinoma was compared to find out

whether the spliced variants are indeed enriched in exosomes. In all 5 samples, the level of expression of the splice variants was higher in the urine exosomes than in the sediments suggesting that these transcripts are enriched in the exosomes and are most likely a more appropriate source for AGR2 as a non-invasive diagnostic marker.

As exosomes are somewhat unique in their protein composition, the level of a few markers was analyzed as additional traits for the identity of the exosomes in the urine sediments and exosome preparation of 3 of the isolated samples. It could be shown that TSG191, CD9, Hsp70 are all overexpressed in the exosome preparation compared to the urine sediments of the 3 samples. Furthermore we could show that PSMA is enriched in the exosomal vesicle fraction indicating the prostate origin of these samples.

For that purpose *AGR2* wt and sv mRNA levels were examined in urine sediments and in exosomes isolated from 5 patients diagnosed with prostate carcinoma. The urine extracellular membrane vesicles were isolated by differential centrifugation and characterized by dynamic light scattering using Zetasizer instrument (Malvern Instruments GmbH).

Example 7: The *AGR2* wt and splice variants mRNA is enriched in the urine exosomes

The previous work showed that *AGR2* wt mRNA and the transcripts of splice variants (sv) were present in urine sediments of patients with prostate cancer. Statistical analysis revealed that the *AGR2* mRNA levels correlate with the tumor aggressiveness and allows discriminating between malign and benign tumors, outperforming markers currently used for the diagnostic of prostate cancer. Based on these findings the mRNA levels of *AGR2* wt and svC, svE, svF, svG, svH were compared in different physiological fluids and tested if these mRNA is available as a circulating cell-free component or as an entire part of the extracellular membrane vesicles, e.g. exosomes, in order to define the most reliable source of the *AGR2* wt or sv mRNA as biomarkers for prostate cancer.

For that purpose *AGR2* wt and sv mRNA levels were examined in blood plasma and in the extracellular membrane vesicles isolated from the plasma in 4 patients diagnosed with prostate carcinoma. Exosome-depleted plasma did not contain detectable levels of RNA and therefore was not included into analysis. Additionally, urine sediments and urine

exosomes from the same patients were examined. Prostate specific gene *PSA* and *GAPDH* mRNA were analyzed as controls.

The plasma and urine extracellular membrane vesicles were isolated by differential centrifugation and characterized by dynamic light scattering using Zetasizer instrument (Malvern Instruments GmbH).

The analysis shows three distinct vesicles population in the blood: population1 with radius 4.7 nm, population 2 with radius 23.6 nm, which might correspond exosomes, and the population 3 with radius 254.2 nm, which might contain either bigger vesicles or the agglomerates of vesicles.

Example 8: AGR2 splice variants obtained from exosomes are tumor type specific

By using the methods as described in the previous examples the presence or absence of ARG2 splice variants was analyzed in samples obtained from patients suffering from other tumor types or in the exosomes derived from cancer cell lines. The results are shown in Figures 6-9B.

In the experiments leading to the results shown in Figure 6, ARG2 splice variants were analyzed in breast cancer cell lines and in the exosomes derived from these cell lines.

The experiments leading to the results shown in Figure 6 allow to speculate that the recruitment of AGR2 splice variants is tumor type -specific and will for instance allow a discrimination between breast cancer and prostate cancer. Splice variant C is present in breast cancer exosome but absent in prostate cancer. On the other side splice variant A is present in the exosomes derived from a patient suffering from prostate cancer but absent in samples obtained from breast cancer cells. Consequently the combinatorial application of translated and non-translated AGR2 splice variants offers a unique possibility for the diagnosis and prognosis of different tumor entities.

Description of the Figure 6-9

Figure 6 *Comparative analysis of the AGR2 SVs mRNA levels in the breast cancer cells and in exosomes*

mRNA levels of AGR2 wild type and of splice variants were analyzed in three breast cancer cell lines: MCF7, MDA-MB-361 and MDA-MB-231. The cells were routinely cultured at 37°C by 95% humidity and 5% CO₂. The cellular RNA was isolated using RNeasy Qiagen kit according to the manufacture of the supplier. Exosomes were isolated from the supernatants of the cells grown in serum-free medium for at least 48 hours. Serum-free medium was used to avoid contaminations with the bovine exosomes. Three hundred fifty ml of cell culture supernatant was used from each cell line for the exosome isolation. Exosomes were isolated by differential ultracentrifugation as described elsewhere (Nazarenko et al., 2010 Cancer Research). After sedimentation by ultracentrifugation, exosomes were directly re-suspended in RLT lysis buffer, supplied with the RNeasy Qiagen Kit. Then RNA was isolated according to the manufacture of the supplied. For the production of cDNA, 500 ng of cellular RNA and 30 ng of exosomal RNA were used. Quantitative PCR, using *GAPDH* and *AGR2*- specific primers was performed as described above.

Splice variants B, C, G and H were detected in both, cells and exosomes, whereas only splice variant G was strongly enriched on the exosomes. Splice variants A, E and F were detectable only in the cells but not in the exosomes, which differed from the data obtained by the analysis of exosomes derived from the prostate cancer cells. These data suggest a difference in the recruitment of *AGR2* splice variants mRNAs to exosomes in different tumor types, which may potentially be used for the diagnosis.

Figure 7 *AGR2 SVs mRNA levels in the exosomes isolated from the urine of 4 patients with urothelial carcinoma.*

EVs were isolated from the urine of 4 patients diagnosed with urothelial carcinoma using protocol applied for the isolation of EVs from the urine of the patient with prostate carcinoma. Following differences were observed as compared to the prostate and breast cancer EVs: SV-A is present in EVs isolated from the urine of the patients with urothelial and prostate cancer, but is absent in breast cancer EVs (Fig. 6); The SVs-E, -F, -G shown

to be enriched in the EVs of prostate cancer patients, were detectable at low levels only in one out of four patients with urothelial carcinoma.

Figure 8 *AGR2 SVs mRNA levels in the exosomes isolated from the urine of 4 patients with renal cell carcinoma.*

EVs were isolated from the urine of 4 patients diagnosed with renal cell carcinoma using protocol applied for the isolation of EVs from the urine of the patient with prostate carcinoma. Following differences were observed as compared to the prostate, breast and urothelial cancer: only SV-A, B, C were detectable in 2 from 4 samples. SV-E, F, G seems to be prostate cancer specific and could not be detected in the EVs derived from the urine of patients with renal cell carcinoma.

Figure 9 *AGR2 SVs mRNA levels in the exosomes isolated from the blood (serum or plasma) of patients with different tumor types.*

To assess if blood can be used as a source of EVs for the *AGR2*-based diagnosis of other different tumor types, blood samples from patients with urothelial and renal cell carcinoma, melanoma, breast and colorectal cancer were used for the isolation of EVs. The *GAPDH* mRNA was analyzed as a control; additionally, breast cancer cell lines T47D was used as a positive control for the *AGR2* wt and splice variants. No *AGR2*-specific signal could be detected in the EVs isolated from the blood samples of these tumor entities, suggesting that only a very low proportion of EVs, derived from the tumor cells were present in the blood and supporting that urine may be a better source for EV for the diagnosis of prostate, urothelial, renal carcinomas.

Claims

1. In vitro process for the diagnosis of prostate cancer and other tumor types in exosomes obtained from a body fluid which comprises

- a) concentrating the exosomes from a body fluid of a patient,
- b) extraction of total RNA from the exosomes and other types of extracellular vesicles obtained in step a),
- c) conversion of the RNA obtained in step b) to cDNA,
- d) amplification of the cDNA obtained in step c) with a polymerase chain reaction whereby primers derived from the AGR2 nucleotide sequence or the complement thereof are used, and
- e) determining whether in the amplification product variants of the AGR2 gene can be identified.

2. Process according to claim 1 wherein the concentration of the exosomes from the body fluid of a patient according to step a) is performed an ultracentrifugation of the body fluid in a buffer at $G \geq 100,000$ for at least 30 minutes.



3. Process according to claim 1 whereby for the concentration of the exosomes from the body fluid of a patient according to step a) is

- a1) centrifuged at low speed in order to remove cells and debris,
- a2) the supernatant is filtered through filtering means, and
- a3) the supernatant is ultracentrifuged, and
- a4) the fraction containing the exosomes is harvested.

4. Process according to any of the preceding claims wherein the RNA is extracted from the exosomes by treating the sediment with at least one protease and DNase.
5. Process according to any of the preceding claims wherein additionally the concentration of RNA is determined and the sample is adjusted to a concentration of 10 to 50, in particular 30 ng RNA.
6. Process according to any of the preceding claims wherein the conversion of the RNA to cDNA is performed by reverse transcription.
7. Process according to any of the preceding claims whereby in addition to the primers derived from the ARG2 gene or complement therefrom a housekeeping gene is amplified.
8. Process according to claim 7 wherein the housekeeping gene is GapDH and/or PSA.
9. Process according to claims 1 to 8 wherein at least two primers are selected from the group consisting of SEQ ID NO:3 to SEQ ID NO:15.
10. Process according to any of claims 1 to 9 wherein the PCR is a quantitative PCR.
11. Process according to any of the preceding claims wherein the extracellular vesicles are further purified from the body fluid by a chromatography step.
12. Process according to any of the preceding claims wherein the extracellular vesicles are further purified from the body fluid by a precipitation step.
13. Process according to any of the preceding claims wherein the body fluid is urine.
14. Process according to any of claims 1 to 12 wherein the body fluid is plasma.
15. Process according to any of claims 1 to 14 wherein the other tumor type is selected from tumors of breast, kidney, bladder, ovarian, small intestines, colon, lung and pancreas.

1/15



Fig. 1A



blood plasma EV	radius (nm)	StdDev (nm)	distribution by signal intensity (%)	distribution by mass (%)	distribution by number (%)
population 1	4.7	0.7	15	95	99.9
population 2	23.6	3.2	43	4.6	0.09
population 3	254.2	29.5	41	0.5	0.01

2/15


Fig. 1B



urine EV	radius (nm)	StdDev (nm)	distribution by signal intensity (%)	distribution by mass (%)	distribution by number (%)
population 1	74	85.9	100	52.90	100

3/15

Fig. 2A



4/15

Fig. 2B






Fig. 2C




6/15

Fig. 3A




7/15

Fig. 3B




8/15

Fig. 3C



9/15

Fig. 4



10/15

Fig. 5




Fig. 6 Comparative analysis of *AGR2* SVs mRNA levels in breast cancer cells and in exosomes





Fig. 7 *AGR2* SVs mRNA levels in exosomes isolated from the urine of 4 patients with urothelial carcinoma



13/15

Fig. 8 *AGR2* SVs mRNA levels in exosomes isolated from the urine of 4 patients with renal cell carcinoma




Fig. 9A *AGR2* SVs mRNA levels in exosomes isolated from blood (serum or plasma) of patients with different tumor types





Fig. 9B *AGR2* SVs mRNA levels in exosomes isolated from blood (serum or plasma) of patients with different tumor types



INTERNATIONAL SEARCH REPORT

International application No
PCT/EP2014/076044

A. CLASSIFICATION OF SUBJECT MATTER
INV. C12Q1/68
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EPO-Internal, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HUAIJIE BU ET AL: "The anterior gradient 2 (AGR2) gene is overexpressed in prostate cancer and may be useful as a urine sediment marker for prostate cancer detection", THE PROSTATE, vol. 71, no. 6, 13 October 2010 (2010-10-13), pages 575-587, XP055107581, ISSN: 0270-4137, DOI: 10.1002/pros.21273 cited in the application	1-14
Y	the whole document Para. "AGR2 Expression Analysis"; Figs. 1 and 2; Pages 579, 581 ----- -/--	15



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

19 February 2015

Date of mailing of the international search report

02/03/2015

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Costa Roldán, Nuria

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2014/076044

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	KIAN KANI ET AL: "Anterior gradient 2 (AGR2): Blood-based biomarker elevated in metastatic prostate cancer associated with the neuroendocrine phenotype", THE PROSTATE, vol. 73, no. 3, 21 August 2012 (2012-08-21), pages 306-315, XP055107578, ISSN: 0270-4137, DOI: 10.1002/pros.22569 the whole document	1-15
Y	----- WO 2012/115885 A1 (CARIS LIFE SCIENCES LUXEMBOURG HOLDINGS S A R L [LU]; SPETZLER DAVID []) 30 August 2012 (2012-08-30) abstract; claims 5-12 paragraph [0195] paragraph [0199] paragraph [0429] paragraph [1285] -----	1-15
A	NILSSON J ET AL: "Prostate cancer-derived urine exosomes: a novel approach to biomarkers for prostate cancer", BRITISH JOURNAL OF CANCER, NATURE PUBLISHING GROUP, GB, vol. 100, no. 10, 1 May 2009 (2009-05-01), pages 1603-1607, XP008149487, ISSN: 0007-0920, DOI: 10.1038/SJ.BJC.6605058 [retrieved on 2009-04-28] the whole document	1-15
A	----- DIEDERICK DUIJVESZ ET AL: "Exosomes as Biomarker Treasure Chests for Prostate Cancer", EUROPEAN UROLOGY, ELSEVIER BV, NL, vol. 59, no. 5, 20 December 2010 (2010-12-20), pages 823-831, XP028160348, ISSN: 0302-2838, DOI: 10.1016/J.EURURO.2010.12.031 [retrieved on 2010-12-22] the whole document	1-15
A	----- WO 2004/031239 A2 (UNIV LIVERPOOL [GB]; CLATTERBRIDGE CANCER RES TRUST [GB]; RUDLAND PHIL) 15 April 2004 (2004-04-15) the whole document ----- -/--	1-15

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2014/076044

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	VERONIKA BRYCHTOVA ET AL: "Anterior gradient 2: A novel player in tumor cell biology", CANCER LETTERS, NEW YORK, NY, US, vol. 304, no. 1, 30 December 2010 (2010-12-30), pages 1-7, XP028173826, ISSN: 0304-3835, DOI: 10.1016/J.CANLET.2010.12.023 [retrieved on 2011-01-19] the whole document -----	1-15
A	VIVEKANANDAN P ET AL: "Anterior gradient-2 is overexpressed by fibrolamellar carcinomas", HUMAN PATHOLOGY, SAUNDERS, PHILADELPHIA, PA, US, vol. 40, no. 3, 1 March 2009 (2009-03-01), pages 293-299, XP025892877, ISSN: 0046-8177, DOI: 10.1016/J.HUMPATH.2008.08.003 [retrieved on 2008-10-29] abstract; figure 2; table 1 -----	1-15
A	PIZZI MARCO ET AL: "Anterior gradient 2 overexpression in lung adenocarcinoma.", APPLIED IMMUNOHISTOCHEMISTRY & MOLECULAR MORPHOLOGY : AIMM / OFFICIAL PUBLICATION OF THE SOCIETY FOR APPLIED IMMUNOHISTOCHEMISTRY JAN 2012, vol. 20, no. 1, January 2012 (2012-01), pages 31-36, XP009182635, ISSN: 1533-4058 abstract -----	1-15
A	WO 2013/134786 A2 (CARIS LIFE SCIENCES LUXEMBOURG HOLDINGS S A R L [LU]; PAWLOWSKI TRACI) 12 September 2013 (2013-09-12) the whole document [00384]; Tab. 2; Tab. 5, page 89, last two rows -----	1-15
A	WO 2013/090620 A1 (GENOMEDX BIOSCIENCES INC [US]) 20 June 2013 (2013-06-20) the whole document [00500]-[00501]; [00257] ----- -/--	1-14

INTERNATIONAL SEARCH REPORT

International application No

PCT/EP2014/076044

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
T	Antje Neeb ET AL: "Splice variant transcripts of the anterior gradient 2 gene as a marker of prostate cancer", Oncotarget, 30 September 2014 (2014-09-30), page 8681, XP055169911, United States Retrieved from the Internet: URL:http://www.ncbi.nlm.nih.gov/pubmed/25237833 the whole document -----	

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/EP2014/076044

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 2012115885 A1	30-08-2012	AU 2012220872 A1 CA 2827894 A1 CN 103492590 A EP 2678448 A1 JP 2014507160 A US 2014141986 A1 WO 2012115885 A1	12-09-2013 30-08-2012 01-01-2014 01-01-2014 27-03-2014 22-05-2014 30-08-2012
WO 2004031239 A2	15-04-2004	AU 2003273502 A1 WO 2004031239 A2	23-04-2004 15-04-2004
WO 2013134786 A2	12-09-2013	AU 2013229762 A1 CA 2865335 A1 EP 2823306 A2 WO 2013134786 A2	25-09-2014 12-09-2013 14-01-2015 12-09-2013
WO 2013090620 A1	20-06-2013	AU 2012352153 A1 CA 2858581 A1 EP 2791359 A1 US 2015011401 A1 WO 2013090620 A1	26-06-2014 20-06-2013 22-10-2014 08-01-2015 20-06-2013