




(86) Date de dépôt PCT/PCT Filing Date: 2008/11/19
(87) Date publication PCT/PCT Publication Date: 2009/05/28
(85) Entrée phase nationale/National Entry: 2010/05/19
(86) N° demande PCT/PCT Application No.: JP 2008/003376
(87) N° publication PCT/PCT Publication No.: 2009/066447
(30) Priorité/Priority: 2007/11/19 (JP2007-299914)

(51) Cl.Int./Int.Cl. *C07H 19/04* (2006.01),
C07D 209/56 (2006.01)
(71) Demandeur/Applicant:
JAPAN ADVANCED INSTITUTE OF SCIENCE AND
TECHNOLOGY, JP
(72) Inventeurs/Inventors:
FUJIMOTO, KENZO, JP;
YOSHIMURA, YOSHINAGA, JP;
TOBA, SHINYA, JP;
NITTA, YUKARI, JP
(74) Agent: RIDOUT & MAYBEE LLP

(54) Titre : NUCLEOTIDE ARTIFICIEL SENSIBLE A LA LUMIERE AYANT UNE APTITUDE DE PHOTO-RETICULATION
(54) Title: LIGHT-RESPONSIVE ARTIFICIAL NUCLEOTIDE HAVING PHOTO-CROSSLINKING ABILITY

[FIG. 4]



(57) Abrégé/Abstract:

The present invention provides a photoreactive crosslinking agent that is capable of crosslinking a sequence which cannot be photo-crosslinked by psoralen, and is capable of photo-crosslinking using a light having a longer wavelength, as compared with psoralen. The present invention also provides a compound having a group represented by formula (I) coupled with a group represented by formula (II).



ABSTRACT

The present invention provides a photoreactive crosslinking agent that is capable of crosslinking a sequence which cannot be photo-crosslinked by psoralen, and is capable of photo-crosslinking using a light having a longer wavelength, as compared with psoralen. The present invention also provides a compound having a group represented by formula (I) coupled with a group represented by formula (II).

DESCRIPTION

LIGHT-RESPONSIVE ARTIFICIAL NUCLEOTIDE HAVING
PHOTO-CROSSLINKING ABILITYTECHNICAL FIELD

[0001]

The present invention relates to a photoreactive crosslinking agent having a photoreactive nucleobase-like structure, which can be crosslinked with a nucleic acid compound, and a light-responsive artificial nucleotide having a photo-crosslinking ability, which has the nucleobase-like structure as a base moiety.

BACKGROUND ART

[0002]

A fundamental technology employed in the field of molecular biology is coupling of nucleic acids. The coupling of nucleic acids is used, for example, in combination with hybridization, for the introduction of genes or detection of base sequences. Therefore, coupling of nucleic acids is an extremely important technology used not only in the basic research in molecular biology, but also in, for example, the diagnosis or treatment in the field of medicine, the development or production of therapeutic drugs, diagnostic drugs or the like, or the development or production of enzymes, microorganisms or the like in industrial and agricultural fields.

[0003]

Coupling of nucleic acids has conventionally been carried out using, for example, DNA ligases and the like. However, reactions carried out by taking out such in vivo enzymatic reactions must be carried out under specially set conditions, and the reactions have disadvantages such as relatively high prices of the enzymes used and insufficient stability. In order to overcome such disadvantages, research has been conducted to find technologies for the coupling of nucleic acids which do not make use of enzymes.

[0004]

As one such technology for the coupling of nucleic acids that does not use enzymes, there is an available method of using an organic compound which is capable of reacting with a nucleic acid. In recent years, technologies for the coupling of nucleic acids utilizing photoreaction are increasingly attracting attention because of their advantages such as that the temporal and spatial control of the reaction is freely achieved, and the reaction can be carried out under milder conditions as compared with general organic chemical reactions.

Among photocoupling technologies as such, those photocoupling technologies making use of 5-cyanovinyldeoxyuridine (Patent Document 1: Japanese Patent No. 3753938 and Patent Document 2: Japanese Patent No. 3753942) are known.

[0005]

Another important technology similar to the coupling of

nucleic acids is crosslinking of nucleic acids. For example, a crosslinking reaction of DNA or RNA blocks the intracellular flow of genetic information, and thus is used in the inhibition of gene expression. As a crosslinking agent for DNA, a compound called psoralen has been traditionally used as a photo-crosslinking agent (photo-crosslinking agent) that induces crosslinking through a photoreaction. A therapeutic method of internally taking psoralen, which is a photo-crosslinking agent, as a medicine, and carrying out photoirradiation, is in widespread use as one of standard therapeutic methods for psoriasis, which is a skin disease. [0006]

However, since the photo-crosslinking reaction of psoralen occurs preferentially with a 5'-TA-3' sequence between the two strands of nucleic acid, there is a problem that the target of the reaction that can be used is restricted. Furthermore, since the photocoupling wavelength for psoralen is 350 nm while its photocleavage wavelength is 250 nm, it is needed to use a short wavelength light source, in the case of using psoralen as a photo-crosslinking agent. Therefore, there is a problem that photoirradiation is likely to cause damages to DNA or cells.

Patent Document 1: Japanese Patent No. 3753938

Patent Document 2: Japanese Patent No. 3753942

DISCLOSURE OF THE INVENTION

PROBLEM TO BE SOLVED BY THE INVENTION

[0007]

Because of the problems as described above, there has been a demand for a photoreactive crosslinking agent that can be used for those sequences which cannot be photo-crosslinked by psoralen. There has also been a demand for a photoreactive crosslinking agent that is capable of photo-crosslinking using a light having a longer wavelength, as compared with psoralen.

[0008]

Therefore, it is an object of the present invention to provide a photoreactive crosslinking agent (a photoreactive crosslinker) that is capable of crosslinking a sequence which cannot be photo-crosslinked by psoralen. It is another object of the present invention to provide a photoreactive crosslinking agent capable of photo-crosslinking using a light having a longer wavelength, as compared with psoralen.

MEANS FOR SOLVING THE PROBLEM

[0009]

The inventors of the present invention devotedly conducted search and investigation on photoreactive crosslinking agents, and as a result, the inventors found a novel photoreactive crosslinking agent (photoreactive crosslinker) compound that is capable of crosslinking a sequence which cannot be photo-crosslinked by psoralen. Thus, the inventors found that the objects described above can be achieved by using this photoreactive crosslinking agent compound.

[0010]

When this photoreactive crosslinking agent compound is used, crosslinks induced by photoirradiation can be formed for those sequences which cannot be photo-crosslinked by psoralen, and this crosslink-forming reaction can be carried out using a light having a longer wavelength, as compared with psoralen.

[0011]

The compound according to the present invention has a characteristic structure in which a vinyl group has been added to a carbazole structure, and exhibits photo-crosslinkability through this relatively small structure. Thus, the compound can be modified by various ways and can be used for various purposes, as in the case of psoralen. Furthermore, this characteristic structure of the compound according to the present invention has a structure similar to the base of nucleic acids, and therefore can be used as an artificial base (artificial nucleobase). That is, an artificial nucleoside and an artificial nucleotide can be produced by introducing the characteristic structure of the compound according to the present invention as an artificial base, and an artificial nucleic acid compound containing such an artificial nucleotide in the sequence can be produced. When such an artificial nucleic acid compound forms a crosslink by a photoreaction, this is led to photo-crosslinks formed (photo-crosslinking) from one of the strands of a double helix to the other strand. Thus, a photoreactive nucleic acid compound can be used as a photo-crosslinking agent for double helix, which is capable of


specifically reacting with a desired sequence.

[0012]

Therefore, the present invention provides the following items [1] to [9].

[1] A compound having a group represented by the following formula (I):

[Chemical Formula 1]



(wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; and

R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen),



coupled with a group represented by the following formula (II):

Formula (II) Rb-

(wherein Rb represents hydrogen, a sugar (the sugar includes ribose and deoxyribose), a polysaccharide (the polysaccharide includes a polyribose chain and a polydeoxyribose chain of nucleic acids), a polyether, a polyol, a polypeptide chain (the

polypeptide chain includes a polypeptide chain of peptide nucleic acids), or a water-soluble synthetic polymer).

[2] The compound according to [1], wherein Rb represents a group represented by the following formula (III) or formula (IV):



or hydrogen.

[3] A photoreactive crosslinking agent containing the compound according to any one of [1] and [2].

[4] A nucleoside having a group represented by the formula (I) according to [1] (wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen) as a base moiety.

[5] A photoreactive crosslinking agent containing the nucleoside according to [4].

[6] A nucleotide having a group represented by the formula (I) according to [1] (wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen) as a base moiety.

[7] A photoreactive crosslinking agent containing the nucleotide according to [6].

[8] A nucleic acid compound (the nucleic acid compound includes a nucleic acid and a peptide nucleic acid) having a group represented by the formula (I) according to [1] (wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen) as a base moiety.

[9] A photoreactive crosslinking agent containing the nucleic acid compound according to [8].

[0013]

The present invention also provides the following items [10] to [13].

[10] A method for forming a photo-crosslink(s) (a photo-crosslinking(s)) using the compound according to any one of [1] and [2], between the compound and a nucleobase having a pyrimidine ring.

[11] A method for forming a photo-crosslink(s) (a photo-crosslinking(s)) using the nucleoside according to [4],

between the nucleoside and a nucleobase having a pyrimidine ring.

[12] A method for forming a photo-crosslink(s) (a photo-crosslinking(s)) using the nucleotide according to [6], between the nucleotide and a nucleobase having a pyrimidine ring.

[13] A method for forming a photo-crosslink(s) (a photo-crosslinking(s)), the method including the steps of:

hybridizing the nucleic acid compound according to [5] and another nucleic acid compound having a pyrimidine ring as a nucleobase to form a double helix; and

irradiating the formed double helix with light.

[0014]

Furthermore, the present invention also provides the following items [14] to [17].

[14] Use of the compound according to any one of [1] and [2], for forming a photo-crosslink(s) (a photo-crosslinking(s)) between the compound and a nucleobase having a pyrimidine ring.

[15] Use of the nucleoside according to [4], for forming a photo-crosslink(s) (a photo-crosslinking(s)) between the nucleoside and a nucleobase having a pyrimidine ring.

[16] Use of the nucleotide according to [6], for forming a photo-crosslink(s) (a photo-crosslinking(s)) between the nucleotide and a nucleobase having a pyrimidine ring.

[17] Use of the nucleic acid compound according to [5], for forming a photo-crosslink(s) (a photo-crosslinking(s))

between the nucleic acid compound and another nucleic acid compound having a pyrimidine ring as a nucleobase, in the double helix formed by hybridization.

EFFECT OF THE INVENTION

[0015]

According to the present invention, a crosslink resulting from photoirradiation can be formed for sequences which cannot be photo-crosslinked by psoralen. That is, according to the present invention, crosslinking can be extensively carried out for the bases in a sequence that cannot be used with psoralen, of which the target of crosslinking is limited to 5'-TA-3' sequences.

[0016]

Furthermore, the characteristic structure of the compound according to the present invention, in which a vinyl group has been added to a carbazole structure, can be used as an artificial nucleobase. When nucleic acid compounds which contain this artificial base and are included in a specific sequence, are hybridized with nucleic acid compounds in a sequence that is complementary to the foregoing sequence, and thereby form a double helix, a photo-crosslink can be formed (photo-crosslinking) for the base of a nucleic acid compound in the complementary sequence, which is located at a specific position in the sequence.

[0017]

Therefore, according to the present invention, not only

a photo-crosslink can be formed for the bases in various sequences which cannot be photo-crosslinked by psoralen, but also a photo-crosslink can be formed for the bases that are at specific positions in a certain specific sequence, by forming a complementary strand.

[0018]

The crosslink-forming reaction according to the present invention can be carried out using a light having a longer wavelength, as compared with psoralen. Accordingly, it can be avoided to use a light source of short wavelength, and thus there is less concern for any damage to DNA or cells due to photoirradiation.

[0019]

Furthermore, the crosslink-forming reaction according to the present invention can be carried out with photoirradiation of a very short duration, as compared with psoralen. That is, since crosslink formation is carried out rapidly, the reaction can be used also for the uses where rapidity of reaction is required. Moreover, even from the viewpoint of the time for photoirradiation being short, there is less concern for any damage to DNA or cells due to photoirradiation.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020]

FIG. 1 is a diagram showing the NOESY spectrum of 3-iodocarbazole-1'-deoxyriboside;

FIG. 2 is a set of chromatograms showing the results of

an HPLC analysis of a photo-crosslinking reaction experiment for ODN(XT) and ODN(A^{CNV}K);

FIG. 3 is a set of chromatograms showing the results of an HPLC analysis of a photo-crosslinking reaction experiment for ODN(XC) and ODN(G^{CNV}K);

FIG. 4 is a graph comparing the rates of a photo-crosslink (photo-crosslinking) reaction between ODN(A^{CNV}K) and ODN(G^{CNV}K);

FIG. 5 is a set of diagrams showing the experiment results for examining the sequence-specificity of ODNs containing ^{CNV}K;

FIG. 6 is a set of chromatograms showing the results of an HPLC analysis of a photo-crosslinking reaction experiment for ODN (A^{OMeV}K) and ODN(GT);

FIG. 7 is a set of chromatograms showing the results of an HPLC analysis of a photo-crosslinking reaction experiment for ODN(G^{OMeV}K) and ODN(GC);

FIG. 8 is a set of chromatograms showing the results of an HPLC analysis of a photo-crosslinking reaction experiment for ODN(A^{OHV}K) and ODN(GT);

FIG. 9 is a set of chromatograms showing the results of an HPLC analysis of a photo-crosslinking reaction experiment for ODN(G^{OHV}K) and ODN(GC);

FIG. 10 is a graph comparing the rates of a photo-crosslinking reaction of ODN(A^{CNV}K), ODN(A^{OMeV}K) and ODN(A^{OHV}K);

FIG. 11 is a set of chromatograms showing the results of photoirradiation and HPLC analysis of ODN(A^{CNV}K-GT); and

FIG. 12 is a set of chromatograms showing the results of photoirradiation and HPLC analysis of ODN(G^{CNV}K-GC).

BEST MODES FOR CARRYING OUT THE INVENTION


[0021]

The present invention will be explained below in detail. The present invention is not intended to be limited to the following specific embodiments.

[0022]

The present invention lies in a compound having a group represented by the following formula (I):

[Chemical Formula 4]



(wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; and

R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen,

coupled with a group represented by the following formula (II):


Formula II Rb-

(wherein Rb represents hydrogen, a sugar (the sugar includes ribose and deoxyribose), a polysaccharide (the polysaccharide includes a polyribose chain and a polydeoxyribose chain of nucleic acids), a polyether, a polyol, a polypeptide chain (the polypeptide chain includes a polypeptide chain of peptide nucleic acids), or a water-soluble synthetic polymer).

[0023]

That is, the present invention lies in a compound represented by the following formula (V), in which the group represented by the formula (I) is coupled with the group represented by the formula (II):

[Chemical Formula 5]



(wherein in the formula (V), Ra, R1, R2 and Rb have the same definitions as described for the formula (I) and formula (II)).

[0024]

The above compound according to the present invention can form crosslinks by photoirradiation for a sequence that cannot be photo-crosslinked by psoralen, and can also induce a photo-crosslinking reaction using a light having a longer wavelength, as compared with psoralen.

[0025]

Ra represents a cyano group, an amide group, a carboxyl group, an alkoxycarbonyl group, or hydrogen; preferably a cyano group, an amide group, a carboxyl group, an alkoxycarbonyl group, or hydrogen; and even more preferably a cyano group, an amide group, a carboxyl group, or an alkoxycarbonyl group. For the alkoxycarbonyl group, preferably a C2-C7 group, more preferably a C2-C6 group, even more preferably a C2-C5 group, even more preferably a C2-C4 group, even more preferably a C2-C3 group, and particularly preferably a C2 group can be used.

[0026]

R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, an alkoxycarbonyl group, or hydrogen; preferably a cyano group, an amide group, a carboxyl group, an alkoxycarbonyl group, or hydrogen; and even more preferably a cyano group, an amide group, a carboxyl group, or an alkoxycarbonyl group. For the alkoxycarbonyl group, preferably a C2-C7 group, more preferably a C2-C6 group, even more preferably a C2-C5 group, even more preferably a C2-C4 group, even more preferably a C2-C3 group, and particularly preferably a C2 group can be used.


[0027]

Rb represents hydrogen, a sugar (the sugar includes ribose and deoxyribose), a polysaccharide (the polysaccharide includes a polyribose chain and a polydeoxyribose chain of nucleic acids), a polyether, a polyol, a polypeptide chain (the polypeptide chain includes a polypeptide chain of peptide nucleic acids), or a water-soluble synthetic polymer.

[0028]

According to a preferred embodiment, hydrogen can be used as Rb, and the compound according to the present invention in this case is represented by the following formula (VI):

[Chemical Formula 6]




(wherein in the formula (VI), Ra, R1 and R2 have the same definitions as described for the formula (I)).

[0029]

According to another preferred embodiment, ribose can be used as Rb, and an example of the compound according to the present invention in this case is a nucleoside (ribonucleoside) represented by the following formula (VII):

[Chemical Formula 7]




(wherein in the formula (VII), R_a , R_1 and R_2 have the same definitions as described for the formula (I)).

[0030]

According to another preferred embodiment, deoxyribose can be used as R_b , and an example of the compound according to the present invention in this case is a nucleoside (deoxyribonucleoside) represented by the following formula (VIII):

[Chemical Formula 8]



(wherein in the formula (VIII), R_a , R_1 and R_2 have the same definitions as described for the formula (I)).

[0031]

The compound according to the present invention can be used as a nucleic acid compound (a photo-crosslinkable nucleic acid compound) having a characteristic structure in which a vinyl group has been added to a carbazole structure as a photoreactive artificial nucleobase. The nucleic acid compound can be hybridized with a sequence having a base sequence that is complementary to the nucleic acid compound,

and can thereby form a double helix with the complementary chain. When the formed double helix is subjected to photoirradiation, the photoreactive artificial nucleobase forms a photo-crosslink with a base which is present in the complementary chain of the photo-crosslinkable nucleic acid compound. The base to be crosslinked to the photoreactive artificial nucleobase is the next base on the 3'-terminal side to a base which is located to a position suitable for base-pairing with the photoreactive artificial nucleobase.

[0032]

The counterpart base with which the photoreactive artificial nucleobase according to the present invention can form a photo-crosslink, is a base having a pyrimidine ring. On the other hand, the photoreactive artificial nucleobase according to the present invention does not form a photo-crosslink with a base having a purine ring. That is, the photo-crosslinkable compound according to the present invention has a strong specificity such that the compound forms a photo-crosslink with cytosine, uracil and thymine among naturally occurring-nucleobases, but does not form a photo-crosslink with guanine and adenine.

[0033]

The photo-crosslinking agent (photo cross-linker) compound according to the present invention, when used as a nucleic acid compound having a photoreactive artificial nucleobase (i.e. photo-crosslinkable nucleic acid compound), can be hybridized with a sequence having a base sequence that

is complementary to the nucleic acid compound, and can thereby form a double helix. Therefore, the photo-crosslinking agent compound can cause a photo-crosslinking reaction to be carried out only for an intended specific sequence (a target sequence). That is, the photo-crosslinking agent compound according to the present invention can impart a very high sequence-selectivity that is impossible with conventional photo-crosslinking agents such as psoralen, through designing of a sequence as desired.

[0034]

Furthermore, when the photo-crosslinking agent compound according to the present invention is used as a nucleic acid compound having a photoreactive artificial nucleic acid base (a photo-crosslinkable nucleic acid compound), is hybridized with a sequence having a base sequence complementary to the nucleic acid compound to thereby form a double helix, the base which should form a base pair with the photoreactive artificial nucleobase according to the present invention in the complementary chain of photo-crosslinkable nucleic acid compounds, is not particularly limited, and can be freely selected.

[0035]

The nucleic acid compound according to the present invention includes a nucleic acid and a peptide nucleic acid (PNA), and also includes a mononucleotide. The nucleic acid includes DNA and RNA, which are natural nucleic acids, and also includes modified nucleic acids such as LNA (BNA), which are non-natural (artificial) nucleic acids.

[0036]

The light that is irradiated for photo-crosslinking is preferably a light having a wavelength in the range of generally 350 to 380 nm, more preferably in the range of 360 to 370 nm, and even more preferably 366 nm. The light is particularly preferably a laser light having a single wavelength of 366 nm.

[0037]

The photo-crosslinking agent compound according to the present invention can further undergo photocleavage by photoirradiation, after being photo-crosslinked with another nucleic acid (nucleic acid compound) by photoirradiation. That is, the photo-crosslinking agent compound according to the present invention enables reversible photo-crosslinking, and thus can be used as a reversible photo-crosslinking agent.

[0038]

Regarding the light that is irradiated for photocleavage, a light having a wavelength in the range of generally 330 to 370 nm, and preferably in the range of 330 to 360 nm, can be used. Furthermore, according to a preferred embodiment, a light having a wavelength of 366 nm, and particularly preferably a laser light having a single wavelength of 366 nm, can be used.

[0039]

According to a preferred embodiment, the photoreactions according to the present invention involving photo-crosslinking and photocleavage can be carried out by irradiation with a light having a wavelength in the range of 350 to 370 nm, and preferably, the photoreactions can be carried

out using a laser light having a single wavelength of 366 nm. When a light having a wavelength in this range is used, both of the photoreactions such as photo-crosslinking and photocleavage can be carried out by means of a same light source, and thus it is advantageous in that there is no need to provide two types of light sources. In the case of using a light having a wavelength in this range, control of which photoreaction between photo-crosslinking and photocleavage would proceed, can be achieved by the temperature conditions. In order to make a photo-crosslinking reaction to proceed, photoirradiation is carried out at a temperature in the range of generally 0 to 50°C, preferably 0 to 40°C, more preferably 0 to 30°C, even more preferably 0 to 20°C, even more preferably 0 to 10°C, and even more preferably 0 to 5°C, and particularly preferably at 0°C. In order to make a photocleavage reaction to proceed, photoirradiation is carried out at a temperature in the range of generally 60 to 100°C, preferably 60 to 90°C, and more preferably 70 to 90°C.

[0040]

Since the photo-crosslinking and photocleavage according to the present invention utilize photoreactions, there is no particular limitation on the pH, temperature, salt concentration or the like, and the reactions can be carried out by photoirradiation in a solution set at a pH, a temperature and a salt concentration, at which biopolymers such as nucleic acid compounds can stably exist.

[0041]

The photo-crosslinking and photocleavage according to the present invention proceed very rapidly, and for example, the photoreaction proceeds for only one second (under irradiation with a light at 366 nm) under the conditions at which psoralen would require several hours (under irradiation with a light at 350 nm). That is, when the photo-crosslinking agent according to the present invention is used, photo-crosslinks can be formed by carrying out the photoreaction by photoirradiation for a time period of several seconds, for example, for 1 to 9 seconds, for 1 to 7 seconds, for 1 to 5 seconds, or for 1 to 3 seconds.

EXAMPLES

[0042]


The present invention will be described below by way of Examples. The present invention is not intended to be limited to the following Examples.

[0043]

[1. Synthesis of ODN having vinylcarbazole at base site]

Synthesis was carried out according to the following Scheme 1. In the explanation given below, the compounds may be occasionally represented by the numbers attached to the compounds.

[Chemical Formula 9]



Scheme 1

[0044]

[3-Iodocarbazole (1)]

To an ethanol solution (500 mL) of carbazole (2.50 g, 15.0 mmol), NaIO₄ (0.80 g, 3.75 mmol) and I₂ (1.89 g, 7.45 mmol) were added in this order, and then an ethanol solution (100 mL) of H₂SO₄ (1.60 mL, 30.0 mmol) was added thereto. The reaction solution was heated to reflux for one hour at 65°C.

Disappearance of the raw materials was confirmed by TLC

(HexH:AcOEt = 4:1), and an ethanol solution (100 mL) of NaOH

(1.4 g) was added thereto to neutralize the reaction solution. Ethanol was removed, and then the reaction solution was extracted two times with chloroform and washed two times with water. The organic phase was dried over Na_2SO_4 , and the solvent was removed. The residue was purified by column chromatography (HexH:AcOEt = 4:1), and compound (1) (3.06 g, 70%) was obtained as a white powder. Thus, 3,6-diiodocarbazole (0.47 g, 7.5%) was obtained as a white powder.

1: ^1H NMR ($\text{DMSO}-d_6$) δ 11.4 (s, 1H), 8.49 (d, 1H, $J = 1.7$ Hz), 8.14 (d, 1H, $J = 8.0$ Hz), 7.62 (dd, 1H, $J = 8.4, 1.7$ Hz), 7.48 (d, 1H, $J = 8.0$ Hz), 7.40 (m, 1H), 7.33 (d, 2H, $J = 8.4$ Hz), 7.16 (m, 1H).

3,6-Diiodocarbazole: ^1H NMR ($\text{DMSO}-d_6$) δ 11.5 (s, 1H), 8.56 (d, 2H, $J = 1.7$ Hz), 7.65 (dd, 2H, $J = 8.5, 1.7$ Hz), 7.34 (d, 2H, $J = 8.5$ Hz).

[0045]

[3-Cyanovinylcarbazole (2)]

To a dioxane solution (10 mL) of triphenylphosphine (139 mg, 0.53 μmol), palladium acetate (40.0 mg, 0.18 μmol) and triethylamine (0.59 μL , 4.23 mmol) were added in this order. The mixture was stirred for 5 minutes at 75°C . A dioxane solution (15 mL) of the compound (1) (1.03 g, 3.52 mmol) and acrylonitrile (0.46 μL , 7.04 mmol) were added thereto, and the reaction solution was heated to reflux for 11.5 hours. Generation of a product was confirmed by TLC (HexH:AcOEt = 4:1), and then palladium powder was removed by cotton filtration. The residue was purified by column chromatography (HexH:AcOEt = 4:1), and compound (2) (0.14 g, 18%, trans:cis = 97:3) was

obtained as a white powder. Thus, compound (1) (0.37 g, recovery rate 37%) was recovered as a white powder.

2: ^1H NMR (DMSO-d_6) δ 11.6 (s, 1H), 8.44 (s, 1H), 8.11 (d, 1H, $J = 8.0$ Hz), 7.75 (d, 1H, $J = 16.7$ Hz), 7.69–7.72 (m, 1H), 7.40–7.52 (m, 3H), 7.19–7.24 (m, 1H), 6.36 (d, 1H, $J = 16.7$ Hz).

[0046]

[3-Cyanovinylcarbazole-1'- β -deoxyriboside-3',5'-di-(p-toluo yl) ester (3)]

To an acetonitrile solution (20 mL) of KOH (0.22 g, 3.87 mmol) and TDA-1 (11 mg, 34 μmol), compound (2) (0.26 g, 1.20 mmol) was added at room temperature. The mixture was stirred for 20 minutes. Chlorosugar (0.53 g, 1.24 mmol) was added to the reaction solution, and the resulting mixture was stirred for 20 minutes at room temperature. Disappearance of the raw materials was confirmed by TLC (HexH:AcOEt = 4:1).

Precipitates were removed, and then the reaction solution was purified by column chromatography (CHCl_3). Thus, compound (3) (0.23 g, 33%) was obtained as a yellow oil.

3: ^1H NMR (CDCl_3) δ 8.09 (s, 1H), 8.02 (d, 2H, $J = 8.4$ Hz), 7.98 (d, 2H, $J = 8.4$ Hz), 7.62–7.65 (m, 1H), 7.62 (d, 1H, $J = 8.8$ Hz), 7.49 (d, 1H, $J = 16.5$ Hz), 7.25–7.31 (m, 7H), 7.17–7.20 (m, 1H), 6.68 (dd, 1H, $J = 9.3, 5.8$ Hz), 5.78 (m, 1H), 5.76 (d, 1H, $J = 16.5$ Hz), 4.91 (dd, 1H, $J = 12.4, 2.7$ Hz), 4.78 (dd, 1H, $J = 12.4, 3.3$ Hz), 4.55–4.57 (m, 1H), 3.09–3.20 (m, 1H), 2.45–2.52 (m, 1H), 2.45 (s, 3H), 2.44 (s, 3H), HRMS (MALDI): calcd. for $\text{C}_{36}\text{H}_{30}\text{N}_2\text{O}_5\text{Na}$ [$(\text{M} + \text{Na})^+$] 593.2053, found 593.2018.

[0047]

[3-Cyanovinylcarbazole-1'- β -deoxyriboside (4)]

To a methanol solution (20 mL) of compound (3) (0.22 g, 0.39 mmol), 0.5 M methanolic NaOMe (2.3 mL, 1.2 mmol) and chloroform (5.0 mL) were added, and the reaction solution was stirred for 3.5 hours at room temperature. Disappearance of the raw materials was confirmed by TLC (CHCl₃:MeOH = 9:1). The solvent was removed, and then the residue was purified by column chromatography (CHCl₃:MeOH = 9:1). Thus, compound (4) (0.11 g, 81%) was obtained as a white powder.

4: ¹H NMR (CDCl₃) δ 8.12 (d, 1H, J = 1.7 Hz), 8.06 (d, 1H, J = 7.7 Hz), 7.59 (d, 1H, J = 9.1 Hz), 7.43-7.57 (m, 4H), 7.26-7.31 (m, 1H), 6.64 (dd, 1H, J = 8.2, 6.9 Hz), 5.87 (d, 1H, J = 16.5 Hz), 4.77-4.82 (m, 1H), 3.95-4.06 (m, 3H), 2.95 (dt, 1H, J = 14.0, 8.2 Hz), 2.30 (ddd, 1H, J = 14.0, 6.9, 3.3 Hz), HRMS (MALDI): calcd. for C₂₀H₁₈N₂O₃Na [(M+ Na)⁺] 357.1215, found 357.1265.

[0048]

[5'-O-(4,4'-dimethoxytrityl)-3-cyanovinylcarbazole-1'-β-deoxyriboside (5)]

Pyridine (0.5 mL) was added to compound (4) (97 mg, 0.29 mmol), which had been azeotropically boiled with pyridine (1.0 mL × 2). To the reaction solution, a pyridine solution (1.0 mL) of 4,4'-dimethoxytrityl chloride (118 mg, 0.35 mmol) and 4-(dimethylamino)pyridine (7.0 mg, 58 μmol) was added, and the reaction solution was stirred for 18 hours at room temperature. Generation of a product was confirmed by TLC (CHCl₃:MeOH = 95:5), and then pyridine was removed. The residue was purified by column chromatography (CHCl₃:MeOH = 98:2), and thus compound (5) (113 mg, 61%) was obtained as a yellow powder.

5: ^1H NMR (CDCl_3) δ 8.07 (d, 1H, $J = 1.7$ Hz), 8.02-8.05 (m, 1H), 7.71 (d, 1H, $J = 8.5$ Hz), 7.62-7.65 (m, 1H), 7.45-7.52 (m, 3H), 7.33-7.37 (m, 4H), 7.25-7.28 (m, 4H), 7.12 (dd, 1H, $J = 8.8, 1.7$ Hz), 6.81 (dd, 4H, $J = 8.8, 1.7$ Hz), 6.61 (dd, 1H, $J = 8.2, 6.3$ Hz), 5.77 (d, 1H, $J = 16.7$ Hz), 4.80-4.82 (m, 1H), 4.05-4.07 (m, 1H), 3.77 (s, 3H), 3.76 (s, 3H), 3.56-3.58 (m, 2H), 2.89 (dt, 1H, $J = 13.8, 8.2$ Hz), 2.23 (ddd, 1H, $J = 13.8, 6.3, 2.7$ Hz), 1.98 (d, 1H, $J = 3.6$ Hz), HRMS (MALDI): calcd. for $\text{C}_{41}\text{H}_{36}\text{N}_2\text{O}_5\text{Na}$ $[(\text{M} + \text{Na})^+]$ 659.2522, found 659.2485.

[0049]

[5'-O-(4,4'-dimethoxytrityl)-3-cyanovinylcarbazole-1'- β -deoxyriboside-3'-O-(cyanoethoxy-N,N-diisopropylamino)phosphoramidite (6)]

Acetonitrile (1.5 mL) was added to compound (5) (0.11 g, 0.17 mol), which had been azeotropically boiled with acetonitrile (1.5 mL). To the reaction solution, 2-cyanoethyl-N,N,N',N'-tetraisopropylphosphorodiamidite (52 μL , 0.17 mol) and an acetonitrile solution (0.37 mL, 0.17 mol) of 0.45 M tetrazole was added, and the reaction solution was stirred for 1.0 hour at room temperature. The reaction solution was extracted two times with deacetylation-treated ethyl acetate, and the residue was washed with a saturated aqueous solution of NaHCO_3 and H_2O . The organic phase was dried over MgSO_4 , and the solvent was removed. Compound (6) (0.12 g), which was a crude product in the form of yellow oil, was transferred in acetonitrile to a rubber-sealed bottle, and was azeotropically boiled three times. The resultant was used in DNA synthesis without further purification.

[0050]

[Synthesis of ODN containing

3-cyanovinylcarbazole-1'- β -deoxyriboside ($^{\text{CNV}}$ K)]

ODNs containing ^{CNV}K as shown below were synthesized.


ODN(A^{CNV}K): 5'-TGCA^{CNV}KCCGT-3'

ODN(G^{CNV}K): 5'-TGCG^{CNV}KCCGT-3'

ODN(C^{CNV}K): 5'-TGCC^{CNV}KCCGT-3'


ODN(T^{CNV}K): 5'-TGCT^{CNV}KCCGT-3'

[Chemical Formula 10]



Synthesis was carried out according to the following Scheme 2.

[Chemical Formula 11]



Scheme 2

[0051]

[3-Methoxycarbonylvinylcarbazole (7)]

To a DMF solution (0.25 mL) of palladium acetate (38.0 mg, 0.17 μmol), compound (1) (0.50 g, 1.71 mmol) as well as tributylamine (0.41 μL , 1.71 mmol), methyl acrylate (0.38 L, 4.27 mmol) and H_2O (1.0 mL) were added in this order. The reaction solution was allowed to react for 10 minutes at 160°C

using microwaves, and the reaction was traced by TLC to confirm the disappearance of compound (1). The palladium powder was removed by Kiriya filtration, and then the residue was purified by column chromatography (HexH:AcOEt = 3:1). Thus, compound (7) (0.26 g, 62%) was obtained as a white powder.

7: ^1H NMR (CDCl_3) δ 8.26 (s, 1H), 8.21 (s, 1H), 8.07 (d, 1H, $J = 8.0$ Hz), 7.89 (d, 1H, $J = 15.9$ Hz), 7.61 (dd, 1H, $J = 1.7, 8.5$ Hz), 7.39-7.44 (m, 3H), 7.23-7.29 (m, 1H), 6.47 (d, 1H, $J = 15.9$ Hz), 3.81 (s, 3H).

[0052]

[3-Methoxycarbonylcarbazole-1'- β -deoxyribose-3',5'-di-(p-toluoyl) ester (8)]

To an acetonitrile solution (49 mL) of compound (7) (0.55 g, 2.22 mmol), NaH (92.0 mg, 2.31 mmol) was added at room temperature, and the mixture was stirred for 10 minutes. Chlorosugar (1.14 g, 2.66 mmol) was added to the reaction solution, and the resulting mixture was stirred for 60 minutes at room temperature. Disappearance of the raw materials was confirmed by TLC (HexH:AcOEt = 4:1). Precipitates were removed, and then the residue was purified by column chromatography (HexH:AcOEt = 4:1). Thus, compound (8) (0.98 g, 71%) was obtained as a white powder.

[0053]

[3-Methoxycarbonylcarbazole-1'- β -deoxyribose (9)]

To a methanol solution (46 mL) of compound (8) (0.96 g, 1.59 mmol), 0.5 M methanolic NaOMe (9.6 mL, 4.8 mmol) and dichloromethane (12 mL) were added, and the reaction solution

was stirred for one hour at room temperature. Disappearance of the raw materials was confirmed by TLC (CHCl_3 :MeOH = 9:1). The solvent was removed, and then the residue was purified by column chromatography (CHCl_3 :MeOH = 9:1). Thus, compound (9) (0.28 g, 48%) was obtained as a white powder.

9: ^1H NMR (CDCl_3) δ 8.20 (s, 1H), 8.06 (d, 1H, $J = 7.7$ Hz), 7.86 (d, 1H, $J = 15.9$ Hz), 7.53–7.61 (m, 3H), 7.44 (t, 1H, $J = 7.1$ Hz), 7.24–7.27 (m, 1H), 6.63 (dd, 1H, $J = 8.2, 7.0$ Hz), 6.46 (d, 1H, $J = 15.9$ Hz), 4.75–4.80 (m, 1H), 3.95–4.04 (m, 3H), 3.81 (s, 1H), 2.95 (dt, 1H, $J = 14.0, 8.2$ Hz), 2.28 (ddd, 1H, $J = 14.0, 7.0, 3.6$ Hz).

[0054]

[5'-O-(4,4'-dimethoxytrityl)-3-methoxycarbonylvinylcarbazole-1'- β -deoxyriboside (10)]

Pyridine (0.5 mL) was added to compound (9) (0.23 g, 0.63 mmol), which had been azeotropically boiled with pyridine (1.0 mL $\times$ 2). To the reaction solution, a pyridine solution (2.2 mL) of 4,4'-dimethoxytrityl chloride (0.26 g, 0.75 mmol) and 4-(dimethylamino)pyridine (15.0 mg, 0.13 μmol) was added, and the reaction solution was stirred for 16 hours at room temperature. Generation of a product was confirmed by TLC (CHCl_3 :MeOH = 95:5), and then pyridine was removed. The residue was purified by column chromatography (CHCl_3 :MeOH = 99:1), and thus compound (10) (0.21 g, 51%) was obtained as a yellow powder.

10: ^1H NMR (CDCl_3) δ 8.17 (s, 1H), 8.02–8.05 (m, 1H), 7.83 (d, 1H, $J = 15.9$ Hz), 7.62–7.66 (m, 3H), 7.46–7.49 (m, 2H), 7.34–7.38 (m, 4H), 7.25–7.28 (m, 4H), 7.15 (d, 1H, $J = 8.8$ Hz), 6.81 (dd, 4H, $J = 8.8, 1.4$ Hz), 6.61 (dd, 1H, $J = 8.5, 6.3$ Hz), 6.40 (d, 1H, $J = 15.9$ Hz), 4.76–4.80 (m, 1H), 4.05–4.09 (m, 1H), 3.80 (s, 3H), 3.77 (s, 3H), 3.76 (s, 3H), 3.56–3.57 (m, 2H), 2.89 (dt, 1H, $J = 14.0, 8.5$ Hz), 2.18 (m, 1H), 2.17 (d, 1H, $J = 3.8$ Hz).

[0055]

[5'-O-(4,4'-dimethoxytrityl)-3-methoxycarbonylvinylcarbazole-1'- β -deoxyriboside-3'-O-(cyanoethoxy-N,N-diisopropylamino)phosphoramidite (11)]

Acetonitrile (1.3 mL) was added to compound (10) (0.20 g, 0.29 μ mol), which had been azeotropically boiled with acetonitrile (1.0 mL). To the reaction solution, 2-cyanoethyl-N,N,N',N'-tetraisopropylphosphorodiamidite (92 μ L, 0.29 μ mol) and an acetonitrile solution (0.65 mL, 0.29 μ mol) of 0.45 M tetrazole was added, and the reaction solution was stirred for 2 hours at room temperature. The reaction solution was extracted two times with deacetylation-treated ethyl acetate, and the residue was washed with a saturated aqueous solution of NaHCO₃ and H₂O. The organic phase was dried over MgSO₄, and the solvent was removed. Compound (11) (0.25 g), which was a crude product in the form of yellow oil, was transferred in acetonitrile to a rubber-sealed bottle, and was azeotropically boiled three times. The resultant was used in DNA synthesis without further purification.

[0056]

[Synthesis of modified ODN]

ODN (AX) (5'-TGCA_XCCGT-3', X = 9) and ODN (GX) (5'-TGCG_XCCGT-3', X = 9) were synthesized using an ABI 3400 DNA synthesizer. The CPG of each of the obtained reaction mixtures was divided into two portions, and one portion of the reaction mixture was deprotected by incubating the mixture at 37°C for

17 hours using 0.4 M NaOH in $\text{H}_2\text{O}:\text{CH}_3\text{OH} = 1:4$, and was subjected to neutralization with 2 M TEAA, followed by freeze-drying. The other portion of the reaction mixture was deprotected by incubating the mixture at room temperature for 17 hours using 0.05 M K_2CO_3 in CH_3OH , and was subjected to neutralization with 2 M TEAA, followed by freeze-drying. The DNAs of $\text{ODN}(\text{A}^{\text{OHV}}\text{K})$, $\text{ODN}(\text{G}^{\text{OHV}}\text{K})$, $\text{ODN}(\text{A}^{\text{OMeV}}\text{K})$ and $\text{ODN}(\text{G}^{\text{OMeV}}\text{K})$ were purified by reverse phase HPLC. Each of the DNAs was enzymatically degraded. The isolation yields were 5, 10, 11 and 13%, respectively. The molecular weights were measured by MALDI-TOF-MS.

calcd. for $\text{ODN}(\text{A}^{\text{OHV}}\text{K})$, 5'-TGCA^{OHV}KCCGT-3': $[(\text{M}+\text{H})^+]$ 2801.93, found 2802.12.

calcd. for $\text{ODN}(\text{G}^{\text{OHV}}\text{K})$, 5'-TGCG^{OHV}KCCGT-3': $[(\text{M}+\text{H})^+]$ 2817.93, found 2818.08.

calcd. for $\text{ODN}(\text{A}^{\text{OMeV}}\text{K})$, 5'-TGCA^{OMeV}KCCGT-3': $[(\text{M}+\text{H})^+]$ 2815.95, found 2816.07.

calcd. for $\text{ODN}(\text{G}^{\text{OMeV}}\text{K})$, 5'-TGCG^{OMeV}KCCGT-3': $[(\text{M}+\text{H})^+]$ 2831.95, found 2831.98.


[0057]

ODNs containing $\text{NH}_2\text{V}\text{K}$ as shown below were synthesized.

$\text{ODN}(\text{A}^{\text{NH}_2\text{V}}\text{K})$: 5'-TGCA^{NH₂V}KCCGT-3'

$\text{ODN}(\text{G}^{\text{NH}_2\text{V}}\text{K})$: 5'-TGCG^{NH₂V}KCCGT-3'

[Chemical Formula 12]



[0058]

[2. Measurement of NOESY spectrum of nucleoside having

vinylcarbazole at base site]

A NOESY spectrum of 3-iodocarbazole-1'-deoxyriboside was measured. In the NOESY spectrum, cross peaks were seen between H1' and H4' (FIG. 1). 3-Iodocarbazole-1'-deoxyriboside is believed to be a β -form.


[0059]

[3. Photo-crosslinking reaction using ODNs containing ^{CNV}K]

Photo-crosslinking reactions were carried out using ODNs containing ^{CNV}K (see the following Scheme 3).

Photo-crosslinking reactions of ODN(A^{CNV}K) (20 μ M) and ODN(XT) (20 μ M, X = A, G, C and T) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μ L). A light at 366 nm was irradiated using a UV-LED for one second at 0°C. The results of an HPLC analysis of the resulting photoreaction products are presented in FIG. 2 (elution with a solvent mixture of 50 mM ammonium formate, pH 7.0; linear gradient over 30 min from 3% to 20% acetonitrile). Photoirradiation of ODN(A^{CNV}K) was carried out in the presence of ODN(XT). FIG. 2(a) presents the results of an HPLC analysis before irradiation, and FIG. 2(b) presents the results of the HPLC analysis obtained after irradiation with a light at 366 nm for one second.

[Chemical Formula 13]




Scheme 3

[0060]

Similarly, a photo-crosslinking reaction was carried out using ODNs containing ^{CNV}K, in which C serves as a photo-crosslinking site (see the following Scheme 4). Photo-crosslinking reactions of ODN(^{CNV}K) (20 μM) and ODN(XC) (20 μM, X = A, G, C and T) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μL). A light at 366 nm was irradiated using a UV-LED for 25 seconds at 0°C. The results of an HPLC analysis of the resulting photoreaction products are presented in FIG. 3. Photoirradiation of ODN(^{CNV}K) was carried out in the presence of ODN(XC). FIG. 3(a) presents the results of an HPLC analysis before irradiation, and FIG. 3(b) presents the results of the HPLC analysis obtained after irradiation with a light at 366 nm for 25 seconds.

[Chemical Formula 14]



Scheme 4

[0061]

In the photo-crosslinking reaction using ODNs containing ^{CNV}K, the photo-crosslinking reaction proceeded sufficiently regardless of whether the photo-crosslinking site was T or C (FIG. 4). FIG. 4 is a graph comparing the rates of photo-crosslink (photo-crosslinking) reactions of ODN(^{CNV}K) (closed circles) and ODN(^{CNV}K) (open circles).

[0062]

[4. Sixteen kinds of photo-crosslinking reactions using ODNs containing ^{CNV}K]

Sixteen kinds of photo-crosslinking reactions were carried out using ODNs containing ^{CNV}K. Photo-crosslinking reactions of ODN(A^{CNV}K), ODN(G^{CNV}K), ODN(C^{CNV}K) and ODN(T^{CNV}K) (10 μM) as well as ODN(XT), ODN(XC), ODN(XG) and ODN(XA) (X = A, G, C and T, 10 μM) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 200 μL). A light at 366 nm was irradiated using a UV-LED for one second at 0°C, and an analysis was made using UPLC (elution with a solvent mixture of 50 mM ammonium formate, pH 7.0; linear gradient over 3.4 min from 2% to 15% acetonitrile, column temperature 30°C). The results are presented in FIG. 5.

[0063]

Sixteen kinds of photo-crosslinking reactions were carried out, and as a result, the following was discovered.

The photo-crosslinking reactions proceed efficiently when the target of crosslinking is T and C. Particularly, if the target of crosslinking is T, the reaction proceeds quantitatively in one second. On the other hand, if the target of crosslinking is A or G, the reaction does not proceed.


[0064]

[5. Photo-crosslinking reaction using ODNs containing ^{OMeV}K or ^{OHV}K]

Photo-crosslinking reactions were carried out using ODNs containing ^{OMeV}K. Photo-crosslinking reactions of ODN(A^{OMeV}K)

(20 μM) and ODN(GT) (20 μM) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 105 μL). A light at 366 nm was irradiated using a UV-LED for 0.5, 1, 2, 3, 5, 10 and 20 seconds, respectively, at 0°C (see the following Scheme 5). The results of an HPLC analysis of the photoreaction products are presented in FIG. 6 (elution with a solvent mixture of 50 mM ammonium formate, pH 7.0; linear gradient over 30 min from 3% to 20% acetonitrile).




[Chemical Formula 15]



[0065]

Similarly, photo-crosslinking reactions were carried out using ODNs containing ^{OMeV}K, in which C serves as a photo-crosslinking site. Photo-crosslinking reactions of ODN(G^{OMeV}K) (20 μM) and ODN(GC) (20 μM) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μL). A light at 366 nm was irradiated using a UV-LED for 180 seconds at 0°C (Scheme 6). The results of an HPLC analysis of the resulting photoreaction products are presented in FIG. 7.

[Chemical Formula 16]






Scheme 6

[0066]

Photo-crosslinking reactions were carried out using ODNs containing $^{\text{OHV}}\text{K}$. Photo-crosslinking reactions of ODN($\text{A}^{\text{OHV}}\text{K}$) (20 μM) and ODN(GT) (20 μM) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μL). A light at 366 nm was irradiated using a UV-LED for 1, 10, 25, 40, 60, 120 and 180 seconds, respectively, at 0°C (see the following Scheme 7). The results of an HPLC analysis of the photoreaction products are presented in FIG. 8.

[Chemical Formula 17]





Scheme 7

[0067]

Similarly, photo-crosslinking reactions were carried out using ODNs containing $^{\text{OHV}}\text{K}$, in which C serves as a photo-crosslinking site. Photo-crosslinking reactions of ODN($\text{G}^{\text{OHV}}\text{K}$) (20 μM) and ODN(GC) (20 μM) were carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μL). A light at 366 nm was irradiated using a UV-LED for 180 seconds at 0°C (Scheme 8). The results of an HPLC analysis of the resulting photoreaction products are presented in FIG. 9.

[Chemical Formula 18]



Scheme 8

[0068]

In the photo-crosslinking reactions using ODNs containing $^{\text{OMeV}}\text{K}$, when the photo-crosslinking site was T, the photo-crosslinking reaction was completed in 10 seconds. When the photo-crosslinking site was C, the photo-crosslinking reaction was completed in 180 seconds. On the other hand, in the photo-crosslinking reactions using ODN containing $^{\text{OHV}}\text{K}$, when the photo-crosslinking site was T, the photo-crosslinking reaction was completed in 180 seconds, with a yield of 75%. When the photo-crosslinking site was C, the photo-crosslinking reaction was completed in 180 seconds. A comparison of changes over time of $^{\text{CNV}}\text{K}$ and $^{\text{OMeV}}\text{K}$ shows significant differences (FIG. 10). FIG. 10 is a graph comparing the rates of photo-crosslinking reactions of $\text{ODN}(\text{A}^{\text{CNV}}\text{K})$ (closed circles), $\text{ODN}(\text{A}^{\text{OMeV}}\text{K})$ (open circles) and $\text{ODN}(\text{A}^{\text{OHV}}\text{K})$ (closed squares).


[0069]

[6. Photocleavage reaction of photo-crosslinked body]

A photo-crosslinked body, $\text{ODN}(\text{A}^{\text{CNV}}\text{K-GT})$ or $\text{ODN}(\text{G}^{\text{CNV}}\text{K-GC})$,

in which the photo-crosslinking site was T or C, was prepared by a photoreaction and was preparatively isolated by HPLC. Then, a photo-crosslinking reaction of ODN(A^{CNV}K-GT) (20 μM) was carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μL). A light at 366 nm was irradiated using a transilluminator for 5 minutes at 70°C (see the following Scheme 9). The results of an HPLC analysis of the photoreaction product are presented in FIG. 11. FIG. 11 shows the results of photoirradiating ODN(A^{CNV}K-GT) at 0°C and analyzing the resultant by HPLC, and FIG. 11(a) presents the results obtained before irradiation, while FIG. 11(b) presents the results obtained after irradiation with a light at 366 nm for 5 minutes.

[Chemical Formula 19]




Scheme 9

[0070]

Similarly, a photo-crosslinking reaction of ODN(G^{CNV}K-GC) was carried out (50 mM sodium cacodylate, 100 mM NaCl, total volume: 30 μL). A light at 366 nm was irradiated using a transilluminator for 5 minutes at 70°C (see the following Scheme 10). The results of an HPLC analysis of the photoreaction product are presented in FIG. 12. The photocleavage reaction proceeded with a yield of 29 and 28%, respectively, when the photo-crosslinking site was T or C. In a photo-crosslinking

reaction using ODNs containing ^{CNV}K, photo-reversible reactivity was observed by changing the reaction temperature. FIG. 12 shows the results of photoirradiating ODN(G^{CNV}K-GC) at 70°C and analyzing the resultant by HPLC, and FIG. 12(a) presents the results obtained before irradiation, while FIG. 12(b) presents the results obtained after irradiation with a light at 366 nm for 5 minutes.

[Chemical Formula 20]




Scheme 10

CLAIMS

1. A compound having a group represented by the following formula (I):

[Chemical Formula 21]



(wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; and

R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen),

coupled with a group represented by the following formula (II):


Formula II Rb-

(wherein Rb represents hydrogen, a sugar (the sugar includes ribose and deoxyribose), a polysaccharide (the polysaccharide includes a polyribose chain and a polydeoxyribose chain of nucleic acids), a polyether, a polyol, a polypeptide chain (the polypeptide chain includes a polypeptide chain of peptide nucleic acids), or a water-soluble synthetic polymer).


2. The compound according to claim 1, wherein

Rb represents a group represented by the following formula (III) or formula (IV):

[Chemical Formula 22]



[Chemical Formula 23]



or hydrogen.

3. A photoreactive crosslinking agent comprising the compound according to any one of claims 1 and 2.

4. A nucleoside having a group represented by the formula (I) according to claim 1 (wherein in the formula (I), Ra represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; and

R1 and R2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen), as a base moiety.

5. A photoreactive crosslinking agent comprising the nucleoside according to claim 4.

6. A nucleotide having a group represented by the formula (I) according to claim 1 (wherein in the formula (I), R_a represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen; and

R_1 and R_2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen), as a base moiety.



7. A photoreactive crosslinking agent comprising the nucleotide according to claim 6.

8. A nucleic acid compound (provided that the nucleic acid compound includes a nucleic acid and a peptide nucleic acid) having a group represented by the formula (I) according to claim 1 (wherein in the formula (I), R_a represents a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen;

and R_1 and R_2 each independently represent a cyano group, an amide group, a carboxyl group, a C2-C7 alkoxy carbonyl group, or hydrogen), as a base moiety.


9. A photoreactive crosslinking agent comprising the nucleic acid compound according to claim 8.

[FIG. 1]




[FIG. 2]


(a) 0 s




(b) 1 s




(a) 0 s




(b) 1 s




(a) 0 s




(b) 1 s







(a) 0 s




(b) 1 s




[FIG. 3]




[FIG. 4]




[FIG. 5]




[FIG. 6]




[FIG. 7]




[FIG. 8]




[FIG. 9]




[FIG. 10]



[FIG. 11]



[FIG. 12]



[FIG. 4]

