



(86) Date de dépôt PCT/PCT Filing Date: 2009/11/18
(87) Date publication PCT/PCT Publication Date: 2010/05/27
(85) Entrée phase nationale/National Entry: 2011/05/16
(86) N° demande PCT/PCT Application No.: IN 2009/000658
(87) N° publication PCT/PCT Publication No.: 2010/058423
(30) Priorité/Priority: 2008/11/20 (IN2623/DEL/2008)

(51) Cl.Int./Int.Cl. *C07D 413/10* (2006.01),
C07D 413/14 (2006.01)


(71) Demandeur/Applicant:
PANACEA BIOTEC LTD., IN

(72) Inventeurs/Inventors:
JAIN, RAJESH, IN;
TREHAN, SANJAY, IN;
DAS, JAGATTARAN, IN;
KANWAR, SANDEEP, IN;
MAGADI, SITARAM KUMAR, IN;
SHARMA, SUDHIR KUMAR, IN

(74) Agent: SIM & MCBURNEY

(54) Titre : NOUVEAUX AGENTS ANTIMICROBIENS

(54) Title: NOVEL ANTIMICROBIALS



Formula I

(57) **Abrégé/Abstract:**

The present invention relates to novel phenyl oxazolidinone compounds of formula I, their pharmaceutically acceptable analogs, tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof. The invention also relates to the processes for the synthesis of novel compounds of formula I or their pharmaceutically acceptable analogs, tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof. The present invention also provides pharmaceutical compositions comprising novel compounds of formula I and methods of using them. The compounds of the present invention are useful as antimicrobial agents, effective against a number of aerobic and/or anaerobic Gram positive and/or Gram negative pathogens such as multi drug resistant species of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacterioides*, *Clostridia*, *H. influenza*, *Moraxella*, acid-fast organisms such as *Mycobacterium tuberculosis* as well as Linezolid resistant species of *Staphylococcus* and *Enterococcus*.

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

REVISED VERSION

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
27 May 2010 (27.05.2010)

PCT

(10) International Publication Number
WO 2010/058423 A9

(51) International Patent Classification:

C07D 413/10 (2006.01) C07D 413/14 (2006.01)

(21) International Application Number:

PCT/IN2009/000658

(22) International Filing Date:

18 November 2009 (18.11.2009)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

2623/DEL/2008 20 November 2008 (20.11.2008) IN

(71) Applicant (for all designated States except US):

PANACEA BIOTEC LTD. [IN/IN]; B-1 Extn./ A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **JAIN, Rajesh** [IN/IN]; B-1 Extn. / A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN). **TREHAN, Sanjay** [IN/IN]; B-1 Extn./ A-27, Mohan Co-operative Industrial Estate, New Delhi 110 044 (IN). **DAS, Jagattaran** [IN/IN]; B-1 Extn. / A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN). **KANWAR, Sandeep** [IN/IN]; B-1 Extn. / A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN). **MAGADI, Sitaram, Kumar** [IN/IN]; B-1 Extn. / A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN). **SHARMA, Sudhir, Kumar** [IN/IN]; B-1 Extn. / A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN).

(74) Agent: **GUPTA, Bhartee**; B-1 Extn. / A-27, Mohan Co-operative Industrial Estate, Mathura Road, New Delhi 110 044 (IN).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM,

AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Declarations under Rule 4.17:

- as to the identity of the inventor (Rule 4.17(i))
- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii))
- of inventorship (Rule 4.17(iv))

Published:

- with international search report (Art. 21(3))

(88) Date of publication of the international search report:

9 December 2010


Date of publication of the revised international search report:

28 April 2011

(15) Information about Correction:

see Notice of 28 April 2011

(54) Title: NOVEL ANTIMICROBIALS



(I)

(57) Abstract: The present invention relates to novel phenyl oxazolidinone compounds of formula I, their pharmaceutically acceptable analogs, tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof. The invention also relates to the processes for the synthesis of novel compounds of formula I or their pharmaceutically acceptable analogs, tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof. The present invention also provides pharmaceutical compositions comprising novel compounds of formula I and methods of using them. The compounds of the present invention are useful as antimicrobial agents, effective against a number of aerobic and/or anaerobic Gram positive and/or Gram negative pathogens such as multi drug resistant species of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacterioides*, *Clostridia*, *H. influenza*, *Moraxella*, acid-fast organisms such as *Mycobacterium tuberculosis* as well as Linezolid resistant species of *Staphylococcus* and *Enterococcus*.

REPLACED SHEET

TITLE: NOVEL ANTIMICROBIALS

Field of the invention

The present invention relates to novel phenyl oxazolidinone compounds of Formula I, their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof. The invention also relates to the processes for the synthesis of novel compounds of Formula I or their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof. The present invention also provides pharmaceutical compositions comprising novel compounds of Formula I and methods of using them. The compounds of the present invention are useful as antimicrobial agents, effective against a number of aerobic and/or anaerobic Gram positive and/or Gram negative pathogens such as multi drug resistant species of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacterioides*, *Clostridia*, *H. influenza*, *Moraxella*, acid-fast organisms such as *Mycobacterium tuberculosis* as well as Linezolid resistant species of *Staphylococcus* and *Enterococcus*.

Background of the invention

Antibacterial resistance has increased alarmingly in the recent years resulting in bacterial strains against which currently available antimicrobial agents are ineffective. In particular, Gram positive bacteria are presenting a formidable treatment problem. Methicillin Resistant *Staphylococcus aureus* (MRSA), Vancomycin Resistant *Enterococci* (VRE) and Glycopeptide Resistant *Staphylococcus aureus* (GRSA) are no longer objects of scientific curiosity but a life threatening proposition that is confronting physicians all over the world. The 'super-bugs' are here to stay and in addition to the several measures to control the spread of drug resistance, a concerted effort is still needed to develop new antibiotics to control life threatening bacterial infections. This growing multidrug resistance has recently rekindled interest in the search for new structural class of antibiotics that kill or inhibit the growth of these bacteria. See, *Chemical Reviews*, "Antibiotic Resistance", 105 (2), February 2005.

Oxazolidinones are a class of antibacterial agents with a unique mechanism of inhibiting bacterial protein synthesis. They inhibit the formation of ribosomal initiation complex involving

REPLACED SHEET

30S and 50S ribosomes leading to prevention of initiation complex formation at the stage of protein synthesis. Due to their unique mechanism of action, these compounds are active against pathogens resistant to other clinically useful antibiotics.

5 Several patent publications disclose oxazolidinones as antimicrobial agents. For example, PCT publications bearing numbers WO 93/09103, WO 93/023384, WO 97/030995, WO 99/064417, WO 00/29396, WO 01/94342, WO 01/081350, WO 02/81469, WO 02/81470, WO 02/02095, WO 03/072553, WO 03/006447, WO 03/07870, WO 03/08389, WO 03/97059, WO 04/045616, WO 04/056817, WO 04/056818, WO 04/14392, WO 04/009587, WO 04/018439A1, 10 WO 05/058886, WO 05/082897, WO 05/116024, WO 05/116021, WO 05/082900, WO 05/003087, WO 06/043121 and WO 09/001192, US patents having numbers US 6,689,779, US 5,565,571, US 5,801,246, US 5,756,732, US 5,654,435 and US 5,654,428, EP patent application EP 1130016 and Chinese patent applications CN 1510032 and CN 1749256 disclose oxazolidinone compounds having antibacterial activity and useful as antimicrobial agents.

15 Some recent publications such as WO 07/114326, US 07/0155798, WO 07/040326, WO 07/095784, WO 07/000432, WO 07/004037 and WO 07/093904 disclose phenyl oxazolidinone derivatives as antibacterial agents. WO 06/109056, WO 06/035283, WO 03/072553, WO 03/064415 disclose heterobicyclic substituted phenyl oxazolidinones as antibacterial agents. WO 96/35691 and WO 00/073301 disclose bicyclic oxazolidinones as antibacterial agents. WO 20 02/064547 discloses pyridoarylphenyl oxazolidinones as antibacterial agents. WO 04/033451, WO 04/089943, WO 05/005422 and WO 05/005399 disclose bicyclo[3.1.0]hexyl-phenyl-oxazolidinone derivatives useful for treating bacterial infections. PCT publication WO 07/082910 discloses dicarbonyl compounds having antibacterial activity. A recent Chinese patent application CN 101434584 discloses phenyl oxazolidinones with glyciny l substitutions 25 having antibacterial activity.


 Linezolid (sold under the trade name Zyvox[®]), the first oxazolidinone to receive regulatory approval, has become an important clinical option in the treatment of serious Gram-positive bacterial infections, including those caused by multidrug resistant pathogens such as MRSA and VRE (see WO 95/07272). In spite of its high potential as an antibiotic and its unique 30 mode of action, no other molecule from oxazolidinone class, except for linezolid, could make it to the clinic. Moreover, development of resistance to an antibiotic is inevitable, and linezolid has been no exception. (See, Mutnick, A. H.; Enne, V.; Jones, R. N. *Ann. Pharmacother.*, 2003, 37,

REPLACED SHEET

769-774). Further, due to myelosuppression, linezolid is not suitable for long duration therapy, although there are cases where patients receiving linezolid for more than two years are without serious side effects. (See, Hutchinson, D. K. *Expert Opin. Ther. Patents* 2004, 14, 1309-1328). Linezolid and its analogs (first generation oxazolidinones) are generally limited in their antimicrobial spectrum to Gram-positive pathogens only. An expanded spectrum and enhanced potency of newer second generation oxazolidinones with activity against Gram-negative pathogens could expand the utility of this class beyond the hospital setting into the treatment of community acquired infections. Thus, there is an ongoing need to develop more effective and safe compounds. The compounds of the present invention are novel, none of them having been previously reported in the prior art. The novel compounds of Formula I according to the present invention possess improved efficacy, particularly enhanced activity against bacterial infections, appreciable bioavailability, reduced associated side effects, good solubility and can be made into formulations with ease.

15 **Summary of the invention**

The present invention relates to novel phenyl oxazolidinones of Formula I,



Formula I

their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof, wherein:

'---' is independently a single bond or absent;

when '---' is a single bond, 'A' represents carbon atom and when '---' is absent, 'A' is CH or N;

Y and Y' are same or different and independently represent O or S;

25 R^1 and R^2 are same or different and independently represent hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_1-C_{12} alkoxy, C_1-12 haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -

REPLACED SHEET

heteroaryl, $-(CH_2)_n C(=Y)NR^5R^6$, $-(CH_2)_n C(=Y)OR^5$, $-(CH_2)_n NR^5R^6$, $-(CH_2)_n OC(=Y)R^5$, $-(CH_2)_n OC(=Y)OR^5$, $-(CH_2)_n OC(=Y)NR^5R^6$, $-(CH_2)_n N(R^5)C(=Y)OR^6$, $-(CH_2)_n N(R^5)C(=Y)NR^5R^6$, $-(CH_2)_n NR^5C(=Y)R^6$, $-(CH_2)_n C(=Y)R^5$, $-(CH_2)_n YR^5$ (wherein each methylene group may be substituted by one or more halogen atoms), $-C(=Y)NR^5R^6$, $-OC(=Y)R^5$, $-OC(=Y)NR^5R^6$, $-C(=Y)OR^5$, $-OR^5$, $-OC(=Y)OR^5$, $-SR^5$, $-NO_2$, $-NR^5R^6$, $-N(R^5)C(=Y)R^6$, $-N(R^5)-C(=Y)OR^6$, or $-N(R^5)C(=Y)NR^5R^6$, each of which may be optionally substituted at any available position by one or more substituents R^a ; or

R^1 and R^2 can together with the carbon atom to which they are attached form a 3 to 10 membered monocyclic ring, partially unsaturated or saturated, which may contain from one to three heteroatoms independently selected from O, S or N; the ring thus formed may be fused with one or two rings independently selected from the group comprising an aryl ring, a cycloalkyl ring, a heterocyclyl ring or monocyclic heteroaryl ring; the ring thus formed can be optionally substituted at any available position by one or more substituents R^a ;

R^3 represents hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_n YR^5$, $-(CH_2)_n C(=Y)R^5$, $-(CH_2)_n NR^5R^6$, $-(CH_2)_n C(=Y)NR^5R^6$, $-(CH_2)_n C(=Y)OR^5$, $-(CH_2)_n OC(=Y)R^5$, $-(CH_2)_n OC(=Y)OR^5$, $-(CH_2)_n NR^5C(=Y)R^6$, $-(CH_2)_n N(R^5)C(=Y)OR^6$, $-(CH_2)_n N(R^5)C(=Y)NR^5R^6$, $-(CH_2)_n OC(=Y)NR^5R^6$, or $-(CH_2)_n N(R^5)C(=Y)NR^5R^6$, (wherein each methylene group may be substituted by one or more halogen atoms), each of which may be optionally substituted at any available position by one or more substituents R^a ;

R^4 represents aryl, heteroaryl or $C(=Y)R^5$ each of which may be optionally substituted at any available position by one or more substituents R^a ;

with the proviso that when m is equal to 2 or 3, then R^4 can not be phenyl substituted with substituents selected from $-OH$, $-OC_{1-4}alkyl$, $-NH_2$, aminoacyl, $-CH_2-NH_2$ and aminoacylalkyl;

Z represents C_{2-12} alkenyl, C_{2-12} alkynyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, $-C_{3-20}$ cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_n NCS$, $-C(=Y)R^5$, $-C(=Y)OR^6$, $-C(=Y)NR^5R^6$, $-OC(=Y)OR^5$, $-(CH_2)_n OP(=O)R^5R^6$, $-(CH_2)_n NHP(=O)R^5R^6$, $-(CH_2)_n OC(=Y)OR^5$, $-(CH_2)_n C(=Y)R^5$, $-(CH_2)_n C(=Y)NR^5R^6$ or $-(CH_2)_n C(=Y)OR^5$ (wherein

REPLACED SHEET

each methylene group may be substituted by one or more halogen atoms), each of which may be optionally substituted at any available position by one or more substituents R^a ;

T, U, V and W are same or different and independently represent hydrogen or halogen;

R^5 and R^6 are same or different and are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12}

5 alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, aryloxy, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted at any available position with halogen, hydroxyl, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} alkoxy, C_{1-12} alkylcarbonyl, C_{1-12} alkoxy carbonyl, C_{3-8} cycloalkyl, C_{1-12} haloalkyl, C_{1-12} haloalkoxy, C_{2-12} haloalkenyl, aryl, 10 heterocyclyl, heteroaryl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-CN$, $-OR^7$, $-NO_2$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-C(=Y)OR^7$, $-OC(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-NR^7SO_2R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, or $-P(=O)R^7R^8$; or R^5 and R^6 may be joined together along with the heteroatom to which they are 15 joined to form a heterocyclic or heteroaryl ring which may additionally contain from one to three heteroatoms independently selected from O, S or N, the ring formed may optionally be substituted with one or more substituents selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{1-12} alkoxy, C_{1-12} haloalkoxy, C_{1-6} alkoxy C_{1-6} alkyl, C_{1-6} alkoxy C_{1-6} alkoxy C_{1-3} alkyl, C_{3-8} cycloalkyl, 20 heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-OR^7$, $-CF_3$, $-OCF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-NO_2$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-OC(=Y)OR^7$, $-C(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-NR^7SO_2R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, or $-P(O)R^7R^8$; the ring thus formed 25 may further be fused with 3 to 7 membered unsaturated or saturated ring, which may contain from one to three heteroatoms independently selected from O, S or N, the fused ring may optionally be substituted at any available position by one or more substituents R^a ;

R^a is independently selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, oxo, C_{1-12} alkoxy, C_{1-6} alkoxy C_{1-6} alkyl, 30 C_{1-6} alkoxy C_{1-6} alkoxy C_{1-3} alkyl, C_{1-12} haloalkoxy, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$

REPLACED SHEET

alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-YR^7$, $-(CH_2)_n YR^7$, $-NO_2$, $=NOR^7$, $-NR^7 R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7 R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7 R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7 R^8$, $-C(=Y)OR^7$, $-OC(=Y)OR^7$, $-SR^7$, $-S(O)_d R^7$, $-SO_2 NR^7 R^8$, $-OP(=O)R^7 R^8$, $-NHP(=O)R^7 R^8$, $-P(O)R^7 R^8$, $-(CH_2)_n CN$, $-YR^7$, $-(CH_2)_n YR^7$, $-NO_2$, $=NOR^7$, $-(CH_2)_n NR^7 R^8$, $-(CH_2)_n N(R^7)C(=Y)R^8$, $-(CH_2)_n N(R^7)C(=Y)OR^8$, $-(CH_2)_n N(R^7)C(=Y)NR^7 R^8$, $-(CH_2)_n C(=Y)R^7$, $-(CH_2)_n C(=Y)NR^7 R^8$, $-(CH_2)_n OC(=Y)R^7$, $-(CH_2)_n OC(=Y)NR^7 R^8$, $-(CH_2)_n C(=Y)OR^7$, $-(CH_2)_n OC(=Y)OR^7$, $-(CH_2)_n SR^7$, $-(CH_2)_n S(O)_d R^7$, $-(CH_2)_n SO_2 NR^7 R^8$, $-(CH_2)_n OP(=O)R^7 R^8$, $-(CH_2)_n NHP(=O)R^7 R^8$, or $-(CH_2)_n P(O)R^7 R^8$; each of which may optionally be substituted at any available position by one or more substituents selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, oxo, C_{1-12} alkoxy, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-OR^9$, $-(CH_2)_n OR^9$, $-CF_3$, $-NO_2$, $-NR^9 R^{10}$, $-N(R^9)C(=Y)R^{10}$, $-N(R^9)C(=Y)OR^{10}$, $-N(R^9)C(=Y)NR^9 R^{10}$, $-C(=Y)R^9$, $-C(=Y)NR^9 R^{10}$, $-OC(=Y)R^9$, $-OC(=Y)NR^9 R^{10}$, $-OC(=Y)OR^9$, $-C(=Y)OR^9$, $-SR^9$, $-S(O)_d R^9$, $-SO_2 NR^9 R^{10}$, $-NR^9 SO_2 R^{10}$, $-OP(=O)R^9 R^{10}$, $-NHP(=O)R^9 R^{10}$, $-P(O)R^9 R^{10}$, $-(CH_2)_n CN$, $-OR^9$, $-(CH_2)_n OR^9$, $-CF_3$, $-NO_2$, $-(CH_2)_n NR^9 R^{10}$, $-(CH_2)_n N(R^9)C(=Y)R^{10}$, $-(CH_2)_n N(R^9)C(=Y)OR^{10}$, $-(CH_2)_n N(R^9)C(=Y)NR^9 R^{10}$, $-(CH_2)_n C(=Y)R^9$, $-(CH_2)_n C(=Y)NR^9 R^{10}$, $-(CH_2)_n OC(=Y)R^9$, $-(CH_2)_n OC(=Y)NR^9 R^{10}$, $-(CH_2)_n OC(=Y)OR^9$, $-(CH_2)_n C(=Y)OR^9$, $-(CH_2)_n SR^9$, $-(CH_2)_n S(O)_d R^9$, $-(CH_2)_n SO_2 NR^9 R^{10}$, $-(CH_2)_n NR^9 SO_2 R^{10}$, $-(CH_2)_n OP(=O)R^9 R^{10}$, $-(CH_2)_n NHP(=O)R^9 R^{10}$, or $-(CH_2)_n P(O)R^9 R^{10}$;

R^7 and R^8 are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, or $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted with halogen, hydroxyl or C_{1-6} alkoxy, or R^7 and R^8 may be joined together along with the heteroatom to which they are attached to form a heterocyclic or heteroaryl ring which may contain from one to three heteroatoms independently selected from O, S or N, each of which may be optionally substituted with halogen, hydroxyl, C_{1-6} alkyl or C_{1-6} alkoxy.

R^9 and R^{10} are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -

REPLACED SHEET

cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, or $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted with halogen, hydroxyl or C_{1-6} alkoxy, or R^9 and R^{10} may be joined together along with the heteroatom to which they are attached to form a heterocyclic or heteroaryl ring which may contain from one to three heteroatoms independently selected from O, S or N, each of which may be optionally substituted with halogen, hydroxyl, C_{1-6} alkyl or C_{1-6} alkoxy.

m is 1, 2, 3 or 4;

m' is 0, 1, 2, 3 or 4;

n is 1, 2, 3 or 4;

10 d is 1 or 2.

Another aspect of the invention provides the processes for the preparation of the novel compounds of Formula I or their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof.

15 A further aspect of the present invention provides pharmaceutical compositions, containing compounds of Formula I or their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof in combination with one or more pharmaceutically acceptable carrier(s).

20 Another aspect of the invention is to provide methods of using the compounds of Formula I of the present invention or compositions comprising the compounds of Formula I for the management such as prophylaxis, amelioration and/or treatment of disease(s)/ disorder(s) especially caused by microbial infections which comprises administering to a subject in need thereof the compounds of Formula I or compositions comprising a pharmaceutically effective amount of the compounds of Formula I.

25 Yet another aspect of the invention is the use of the compounds of Formula I as antimicrobial agents, effective against a number of aerobic and/or anaerobic Gram positive and/or Gram negative pathogens such as multi drug resistant species of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacterioides*, *Clostridia*, *H. influenza*, *Moraxella*, acid-fast organisms such as *Mycobacterium tuberculosis* as well as Linezolid resistant species of *Staphylococcus* and *Enterococcus*.

REPLACED SHEET


In another aspect, the present invention provides a method for treating Gram positive and/or Gram negative pathogens in a mammal by administering a therapeutically effective amount of a compound of Formula I or a pharmaceutically acceptable salt.

5 The present invention also encompasses prodrugs and active metabolites of the compounds of the Formula I.

Other aspects of the invention will be set forth in the description which follows, and in part will
10 be apparent from the description, or may be learnt by the practice of the invention.

Detailed description of the invention

The present invention relates to novel phenyl oxazolidinones of Formula I,



Formula I

15 their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof, wherein:

'---' is independently a single bond or absent;

when '---' is a single bond, 'A' represents carbon atom and when '---' is absent, 'A' is CH or
20 N;

Y and Y' are same or different and independently represent O or S;

R¹ and R² are same or different and independently represent hydrogen, halogen, C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, C₂₋₁₂ alkynyl, C₁₋₁₂ haloalkyl, C₂₋₁₂ haloalkenyl, C₂₋₁₂ haloalkynyl, C₁₋₁₂ alkoxy, C₁₋₁₂ haloalkoxy, C₁₋₆ alkoxyC₁₋₆alkyl, C₁₋₆ alkoxyC₁₋₆alkoxyC₁₋₃alkyl, C₃₋₂₀ cycloalkyl,
25 heterocyclyl, aryl, heteroaryl, -(CH₂)_n-cycloalkyl, -(CH₂)_n-heterocyclyl, -(CH₂)_n-aryl, -(CH₂)_n-heteroaryl, -(CH₂)_nC(=Y)NR⁵R⁶, -(CH₂)_nC(=Y)OR⁵, -(CH₂)_nNR⁵R⁶, -(CH₂)_nOC(=Y)R⁵, -(CH₂)_nOC(=Y)OR⁵, -(CH₂)_nOC(=Y)NR⁵R⁶, -(CH₂)_nOC(=Y)NR⁵R⁶, -(CH₂)_nN(R⁵)C(=Y)OR⁶, -(CH₂)_nN(R⁵)C(=Y)NR⁵R⁶, -(CH₂)_nNR⁵C(=Y)R⁶, -(CH₂)_nC(=Y)R⁵, -(CH₂)_nYR⁵ (wherein each methylene group may be substituted by one or more halogen atoms), -C(=Y)NR⁵R⁶, -OC(=Y)R⁵, -OC(=Y)NR⁵R⁶, -C(=Y)OR⁵, -OR⁵, -OC(=Y)OR⁵, -SR⁵, -NO₂, -NR⁵R⁶, -
30

REPLACED SHEET

$N(R^5)C(=Y)R^6$, $-N(R^5)-C(=Y)OR^6$, or $-N(R^5)C(=Y)NR^5R^6$, each of which may be optionally substituted at any available position by one or more substituents R^a ; or

R^1 and R^2 can together with the carbon atom to which they are attached form a 3 to 10 membered monocyclic ring, partially unsaturated or saturated, which may contain from one to three heteroatoms independently selected from O, S or N; the ring thus formed may be fused with one or two rings independently selected from the group comprising an aryl ring, a cycloalkyl ring, a heterocyclyl ring or monocyclic heteroaryl ring; the ring thus formed can be optionally substituted at any available position by one or more substituents R^a ;

R^3 represents hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_nYR^5$, $-(CH_2)_nC(=Y)R^5$, $-(CH_2)_nNR^5R^6$, $-(CH_2)_nC(=Y)NR^5R^6$, $-(CH_2)_nC(=Y)OR^5$, $-(CH_2)_nOC(=Y)R^5$, $-(CH_2)_nOC(=Y)OR^5$, $-(CH_2)_nNR^5C(=Y)R^6$, $-(CH_2)_nN(R^5)C(=Y)OR^6$, $-(CH_2)_nN(R^5)C(=Y)NR^5R^6$, $-(CH_2)_nOC(=Y)NR^5R^6$, or $-(CH_2)_nN(R^5)C(=Y)NR^5R^6$, (wherein each methylene group may be substituted by one or more halogen atoms), each of which may be optionally substituted at any available position by one or more substituents R^a ;

R^4 represents aryl, heteroaryl or $C(=Y)R^5$ each of which may be optionally substituted at any available position by one or more substituents R^a ;

with the proviso that when m is equal to 2 or 3, then R^4 can not be phenyl substituted with substituents selected from $-OH$, $-OC_{1-4}alkyl$, $-NH_2$, aminoacyl, $-CH_2-NH_2$ and aminoacylalkyl;

Z represents C_{2-12} alkenyl, C_{2-12} alkynyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_nNCS$, $-C(=Y)R^5$, $-C(=Y)OR^6$, $-C(=Y)NR^5R^6$, $-OC(=Y)OR^5$, $-(CH_2)_nOP(=O)R^5R^6$, $-(CH_2)_nNHP(=O)R^5R^6$, $-(CH_2)_nOC(=Y)OR^5$, $-(CH_2)_nC(=Y)R^5$, $-(CH_2)_nC(=Y)NR^5R^6$ or $-(CH_2)_nC(=Y)OR^5$ (wherein each methylene group may be substituted by one or more halogen atoms), each of which may be optionally substituted at any available position by one or more substituents R^a ;

T , U , V and W are same or different and independently represent hydrogen or halogen;

R^5 and R^6 are same or different and are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{3-8} cycloalkyl,

REPLACED SHEET

heterocyclyl, aryl, heteroaryl, aryloxy, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted at any available position with halogen, hydroxyl, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} alkoxy, C_{1-12} alkylcarbonyl, C_{1-12} alkoxycarbonyl, C_{3-8} cycloalkyl, C_{1-12} haloalkyl, C_{1-12} haloalkoxy, C_{2-12} haloalkenyl, aryl, heterocyclyl, heteroaryl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_n$ -cycloalkyl, -CN, $-OR^7$, $-NO_2$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-C(=Y)OR^7$, $-OC(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-NR^7SO_2R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, or $-P(=O)R^7R^8$; or R^5 and R^6 may be joined together along with the heteroatom to which they are joined to form a heterocyclic or heteroaryl ring which may additionally contain from one to three heteroatoms independently selected from O, S or N, the ring formed may optionally be substituted with one or more substituents selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{1-12} alkoxy, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxycarbonyl, -CN, $-OR^7$, $-CF_3$, $-OCF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-NO_2$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-OC(=Y)OR^7$, $-C(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-NR^7SO_2R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, or $-P(O)R^7R^8$; the ring thus formed may further be fused with 3 to 7 membered unsaturated or saturated ring, which may contain from one to three heteroatoms independently selected from O, S or N, the fused ring may optionally be substituted at any available position by one or more substituents R^8 ;

R^8 is independently selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, oxo, C_{1-12} alkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{1-12} haloalkoxy, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxycarbonyl, -CN, $-YR^7$, $-(CH_2)_nYR^7$, $-NO_2$, $=NOR^7$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-C(=Y)OR^7$, $-OC(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, $-P(O)R^7R^8$, $-(CH_2)_nCN$, $-YR^7$, $-(CH_2)_nYR^7$, $-NO_2$, $=NOR^7$, $-(CH_2)_nNR^7R^8$, $-(CH_2)_nN(R^7)C(=Y)R^8$, $-(CH_2)_nN(R^7)C(=Y)OR^8$, $-(CH_2)_nN(R^7)C(=Y)NR^7R^8$, $-(CH_2)_nC(=Y)R^7$,

REPLACED SHEET

- $-(CH_2)_n C(=Y)NR^7R^8$, $-(CH_2)_n OC(=Y)R^7$, $-(CH_2)_n OC(=Y)NR^7R^8$, $-(CH_2)_n C(=Y)OR^7$, $-(CH_2)_n OC(=Y)OR^7$, $-(CH_2)_n SR^7$, $-(CH_2)_n S(O)_dR^7$, $-(CH_2)_n SO_2NR^7R^8$, $-(CH_2)_n OP(=O)R^7R^8$, $-(CH_2)_n NHP(=O)R^7R^8$, or $-(CH_2)_n P(O)R^7R^8$; each of which may optionally be substituted at any available position by one or more substituents selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, oxo, C_{1-12} alkoxy, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-OR^9$, $-(CH_2)_n OR^9$, $-CF_3$, $-NO_2$, $-NR^9R^{10}$, $-N(R^9)C(=Y)R^{10}$, $-N(R^9)C(=Y)OR^{10}$, $-N(R^9)C(=Y)NR^9R^{10}$, $-C(=Y)R^9$, $-C(=Y)NR^9R^{10}$, $-OC(=Y)R^9$, $-OC(=Y)NR^9R^{10}$, $-OC(=Y)OR^9$, $-C(=Y)OR^9$, $-SR^9$, $-S(O)_dR^9$, $-SO_2NR^9R^{10}$, $-NR^9SO_2R^{10}$, $-OP(=O)R^9R^{10}$, $-NHP(=O)R^9R^{10}$, $-P(O)R^9R^{10}$, $-(CH_2)_n CN$, $-OR^9$, $-(CH_2)_n OR^9$, $-CF_3$, $-NO_2$, $-(CH_2)_n NR^9R^{10}$, $-(CH_2)_n N(R^9)C(=Y)R^{10}$, $-(CH_2)_n N(R^9)C(=Y)OR^{10}$, $-(CH_2)_n N(R^9)C(=Y)NR^9R^{10}$, $-(CH_2)_n C(=Y)R^9$, $-(CH_2)_n C(=Y)NR^9R^{10}$, $-(CH_2)_n OC(=Y)R^9$, $-(CH_2)_n OC(=Y)NR^9R^{10}$, $-(CH_2)_n OC(=Y)OR^9$, $-(CH_2)_n C(=Y)OR^9$, $-(CH_2)_n SR^9$, $-(CH_2)_n S(O)_dR^9$, $-(CH_2)_n SO_2NR^9R^{10}$, $-(CH_2)_n NR^9SO_2R^{10}$, $-(CH_2)_n OP(=O)R^9R^{10}$, $-(CH_2)_n NHP(=O)R^9R^{10}$, or $-(CH_2)_n P(O)R^9R^{10}$;
- R^7 and R^8 are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, or $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted with halogen, hydroxyl or C_{1-6} alkoxy, or R^7 and R^8 may be joined together along with the heteroatom to which they are attached to form a heterocyclic or heteroaryl ring which may contain from one to three heteroatoms independently selected from O, S or N, each of which may be optionally substituted with halogen, hydroxyl, C_{1-6} alkyl or C_{1-6} alkoxy.
- R^9 and R^{10} are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, or $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted with halogen, hydroxyl or C_{1-6} alkoxy, or R^9 and R^{10} may be joined together along with the heteroatom to which they are attached to form a heterocyclic or heteroaryl ring which may contain from one to three heteroatoms independently selected from

REPLACED SHEET

O, S or N, each of which may be optionally substituted with halogen, hydroxyl, C₁₋₆ alkyl or C₁₋₆ alkoxy.


m is 1, 2, 3 or 4;

m' is 0, 1, 2, 3 or 4;

5 n is 1, 2, 3 or 4;

d is 1 or 2.

One embodiment of the present invention provides compounds of Formula Ia, wherein




Formula Ia

10 R¹, R², R³, R⁴, Y, Y', A, T, U, V, W, Z, m and m' are as defined herein;

their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof.

Another embodiment of the present invention provides compounds of Formula Ib, wherein



Formula Ib

15 R¹, R², R³, R⁴, T, U, V, W, Z, m, m' are as defined herein; their pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof.


20 In another embodiment of the compounds of the present invention, R¹ and R² are independently selected from hydrogen, C₁₋₁₂ alkyl, C₂₋₁₂ alkenyl, C₂₋₁₂ alkynyl, C₃₋₈ cycloalkyl or aryl, each of which may optionally be substituted at any available position by one or more substituents R^a or R¹ and R² together with the carbon atom to which they are attached form a 3 to 10 membered monocyclic ring, partially unsaturated or saturated, which may contain from
25 one to three heteroatoms independently selected from O, S or N, the ring thus formed may be

REPLACED SHEET

fused with one or two rings independently selected from the group comprising an aryl ring, a cycloalkyl ring, a heterocyclyl ring or monocyclic heteroaryl ring; the ring thus formed can be optionally substituted at any available position by one or more substituents R^a.


In another embodiment of the compounds of the present invention R^3 is selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl or C_{2-12} alkynyl, each of which may be optionally substituted at any available position by one or more substituents R^a .

In a further embodiment of the compounds of the present invention, it is preferred that R^4 is selected from



wherein R¹¹ is selected from -H, -CH₃, -OH, -F, -Cl, or -N $\overline{\text{O}}$

10




In a further embodiment of the compounds of the present invention, it is preferred that T and W independently represent fluorine and U and V both represent hydrogen.

In another embodiment of the compounds of the present invention, m is selected from 1 or 2 and m' is 0.

REPLACED SHEET

In another embodiment of the compounds of the present invention, Z represents -CH₂-triazole, which may be optionally substituted at any available position by one or more substituents R^a.

In a further embodiment of the compounds of the present invention, it is preferred that Z
5 is selected from



† Definitions

Relative to the above description of the oxazolidinone compounds of the present invention, the following definitions apply.

10 The term "alkyl" as used herein refers to a straight or branched hydrocarbon chain, having from 1 to 12 carbon atoms. Examples of alkyl include, but are not limited to, for example, methyl, ethyl, *n*-propyl, isoprppyl, *n*-butyl, *n*-pentyl, *t*-butyl and the like. These groups may further be substituted with one or more substituents selected from but not limited to, for example,
15 halogen, hydroxyl, oxo, carboxyl, carboxyalkyl, azido, cyano, amino, nitro, alkenyl, alkynyl, alkoxy, cycloalkyl, acyl acyloxy, aryl, heterocyclyl and heteroaryl.

The term "alkenyl" as used herein refers to an aliphatic hydrocarbon group containing at least one carbon-carbon double bond and which may be straight or branched hydrocarbon chain
20 having from 1 to 12 carbon atoms. Examples of alkenyl include, but are not limited to, for example, ethenyl, 1-propenyl, 2-propenyl, iso-propenyl, 1-butenyl, 2-butenyl, and the like. These groups may further be substituted with one or more substituents selected from but not limited to, for example, halogen, hydroxyl, oxo, carboxyl, carboxyalkyl, azido, cyano, amino, nitro, alkenyl, alkynyl, alkoxy, cycloalkyl, acyl acyloxy, aryl, heterocyclyl and heteroaryl.

25 The term "alkynyl" as used herein refers to a straight or branched hydrocarbon group containing at least one carbon-carbon triple bond and which may be straight or branched chain having from 1 to 12 carbon atoms. . Examples of alkynyl include, but are not limited to, for example, ethynyl, propynyl, and butynyl. These groups may further be substituted with one or
30 more substituents selected from but not limited to, for example, halogen, hydroxyl, oxo,

REPLACED SHEET

carboxyl, carboxyalkyl, azido, cyano, amino, nitro, alkenyl, alkynyl, alkoxy, cycloalkyl, acyl acyloxy, aryl, heterocyclyl and heteroaryl.

5 The term "alkoxy" refers to an above defined alkyl group attached via an oxygen linkage to the rest of the molecule. Non-limiting examples of such groups include $-OCH_3$, $-OC_2H_5$ and the like.

"Halogen" refers to fluorine, chlorine, bromine or iodine.

10 The term "haloalkyl" refers to an above-defined "alkyl" group, which is substituted with one or more "halogen" groups, as defined herein, at any one or more of the 1 to 12 carbon atoms of the alkyl group. Representative examples of haloalkyl include, but are not limited to, chloromethyl, fluoromethyl, trifluoromethyl, trichloromethyl, difluoroethyl, trifluoroethyl, dichloroethyl, and the like.

15 The term "haloalkenyl" refers to an above-defined "alkenyl" group, which is substituted with one or more "halogen" groups, as defined herein, at any one or more of the carbon atoms of the alkenyl group.. Representative examples of haloalkenyl include, but are not limited to, chloroethenyl, 2-fluoroethenyl, trifluorobutenyl and dichloropropenyl.

20 The term "haloalkynyl" refers to an above-defined "alkynyl" group, which is substituted with one or more "halogen" groups, as defined herein, at any one or more of the carbon atoms of the alkynyl group. Representative examples of haloalkynyl include, but are not limited to, 2-fluoroethynyl, trifluorobutynyl and dichloropropynyl.

25 The term "haloalkoxy" refers to an above defined "haloalkyl" group, appended to the parent molecular moiety through an oxygen atom.

30 The term "cycloalkyl" refers to cyclic alkyl groups constituting of 3 to 20 carbon atoms having a single cyclic ring or multiple condensed rings, for example, fused or spiro systems, unless otherwise constrained by the definition. Such cycloalkyl groups include, by way of example, single ring structures, for example, cyclopropyl, cyclobutyl, cyclopentenyl, cyclohexyl, cyclooctyl, and the like, or multiple ring structures, for example, adamantyl, and bicyclo[2.2.1] heptane, or cyclic alkyl groups to which is fused an aryl group, for example, indane and the like.

REPLACED SHEET

Cycloalkyl groups may further be substituted with one or more substituents selected from but not limited to, for example, halogen, hydroxyl, oxo, carboxy, carboxyalkyl, azido, alkenyl, alkynyl, alkoxy, cycloalkyl, acyl acyloxy, aryl, heterocyclyl, heteroaryl.

- 5 The term "aryl" herein refers to a mono- or poly- carbocyclic aromatic group, for example phenyl or naphthyl ring and the like optionally substituted with one or more substituents selected from but not limited to, for example, halogen, hydroxyl, alkyl, alkenyl, alkynyl, cycloalkyl, alkoxy, acyl, amino, aryloxy, CF₃, COOR_d (wherein R_d can be hydrogen, alkyl, alkenyl, cycloalkyl, aralkyl, heterocyclylalkyl or heteroarylalkyl), cyano, nitro, carboxy, heterocyclyl, 10 heteroaryl, heterocyclylalkyl or heteroarylalkyl. The aryl group may optionally be fused with cycloalkyl group, heteroaryl group, heterocyclyl group or another aryl group. The fused group may be further substituted at any available position with one or more substituents selected from but not limited to, for example, halogen, hydroxyl, oxo, carboxy, amino, nitro, cyano, carboxyalkyl, azido, alkenyl, alkynyl, alkoxy, cycloalkyl, acyl, acyloxy, aryl, heterocyclyl, 15 heteroaryl.

The term "aryloxy" refers to an above defined aryl group attached via an oxygen linkage to the rest of the molecule, for example -OPh and the like.

- 20 The term "heteroaryl" unless and otherwise specified refers to an aromatic monocyclic or polycyclic ring structure, containing one to four heteroatoms independently selected from N, O or S. The nitrogen heteroatoms may optionally be oxidized. The nitrogen atoms may optionally be quaternized. "Heteroaryl" also includes, but is not limited to, bicyclic or tricyclic rings, wherein the heteroaryl ring is fused to one or two rings independently selected from the group consisting of 25 an aryl ring, a cycloalkyl ring, a heterocyclyl ring and another monocyclic heteroaryl ring. Examples of heteroaryl groups include, but not limited to, oxazolyl, imidazolyl, pyrrolyl, 1,2,3-triazolyl, 1,2,4-triazolyl, tetrazolyl, thiazolyl, oxadiazolyl, benzoimidazolyl, thiadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, thienyl, isoxazolyl, triazinyl, furanyl, benzofuranyl, indolyl, benzothiazolyl, benzoxazolyl, imidazo[1,2-*a*]pyrimidine, imidazo[1,2-*a*]pyrazine, and the like. The 30 bicyclic or tricyclic heteroaryl rings have to be attached through the heteroaryl group only. The heteroaryl group may be further substituted at any available position with one or more substituents selected from but not limited to, for example, halogen, hydroxyl, oxo, carboxy, amino, nitro,

REPLACED SHEET

cyano, carboxyalkyl, azido, alkenyl, alkynyl, alkoxy, cycloalkyl, cycloalkynyl, acyl acyloxy, aryl, heterocyclyl, heteroaryl.

5 The term "heterocyclyl" unless otherwise specified refers to a non-aromatic monocyclic or polycyclic cycloalkyl group, fully or partially unsaturated, containing one or more heteroatom(s) independently selected from N, O or S. The heterocyclyl ring may be fused with another cycloalkyl, aryl, heterocyclyl or heteroaryl ring and are optionally benzofused or fused heteroaryl of 5-6 ring members and/or are optionally substituted wherein the substituents are selected from but not limited to halogen, hydroxyl, alkyl, alkenyl, alkynyl, cycloalkyl, acyl, carboxy, aryl, 10 alkoxy, aralkyl, cyano, nitro, amino, heterocyclyl, or heteroaryl. Examples of heterocyclyl groups include but are not limited to, morpholinyl, oxazolidinyl, tetrahydrofuranyl, dihydrofuranyl, dihydropyridinyl, dihydroisooxazolyl, dihydrobenzofuryl, azabicyclohexyl, dihydroindonyl, piperidinyl or piperazinyl. The fused group may be further substituted at any available position with one or more substituents selected from but not limited to, for example, 15 halogen, hydroxyl, oxo, carboxy, amino, nitro, cyano, carboxyalkyl, azido, alkenyl, alkynyl, alkoxy, cycloalkyl, acyl acyloxy, aryl, heterocyclyl, heteroaryl. The nitrogen and sulphur heteroatoms may optionally be oxidized. The nitrogen atoms may optionally be quaternized.

"Hydroxy" or "hydroxyl" refers to the group -OH.

20 The term "Protecting Group" or "PG" refers to a group which is in a modified form to preclude undesired side reactions at the protected site. The term protecting group, unless otherwise specified, may be used with groups, for example, hydroxyl, amino, carboxyl and examples of such groups are found in T.W. Greene. et al. "*Protecting Groups in Organic Synthesis*," 3rd Ed, 25 Wiley, New York, which is incorporated herein by reference. The species of the carboxylic protecting groups, amino protecting groups or hydroxyl protecting groups employed are not critical, as long as the derivatised moieties/moiety is/are stable to conditions of subsequent reactions and can be removed without disrupting the remainder of the molecule. Examples of suitable hydroxyl and amino protecting groups include but are not limited to trimethylsilyl, 30 triethylsilyl, *o*-nitrobenzyloxycarbonyl, *p*-nitrobenzyloxycarbonyl, *t*-butyldiphenylsilyl, *t*-butyldimethylsilyl, acetyl, trifluoroacetyl, benzyloxycarbonyl (CBz), *t*-butoxycarbonyl (Boc), 9-fluorenylethoxycarbonyl (Fmoc), 2,2,2-trichloroethyloxycarbonyl, allyloxycarbonyl and the

REPLACED SHEET

like. Examples of suitable carboxyl protecting groups are benzhydryl, *o*-nitrobenzyl, *p*-nitrobenzyl, 2-naphthylmethyl, allyl, 2-chloroallyl, benzyl, 2,2,2-trichloroethyl, trimethylsilyl, *t*-butyldimethylsilyl, *t*-butyldiphenylsilyl, 2-(trimethylsilyl)ethyl, phenacyl, *p*-methoxybenzyl, acetyl, *p*-methoxyphenyl, 4-pyridylmethyl, *t*-butyl and the like.

5

"Subject" includes humans, non-human mammals (e.g., dogs, cats, rabbits, cattle, horses, sheep and the like) or non-mammals (e.g., birds and the like).

10 The term "therapeutically effective amount" means the amount of a compound that, when administered to a subject for treating a disease, is sufficient to effect such treatment for the disease. The "therapeutically effective amount" will vary depending on the compound, the disease and its severity, weight, physical condition and responsiveness of the subject to be treated, among other factors.

15 A "pharmaceutically acceptable salt" encompasses any compound according to the present invention that is utilized in the form of a salt thereof, especially where the salt confers on a compound improved pharmacokinetic properties as compared to the free form of compound or a different salt form of the compound.

20 Asymmetric centres may exist in the compounds of the present invention. The compounds of Formula I may have one or more stereogenic centres and so can exhibit optical isomerism. All such isomers including enantiomers, diastereomers, and epimers are included within the scope of this invention. Furthermore, the invention includes such compounds as single isomers (*R* and /or *S*) and as mixtures, including racemates. If desired, racemic mixtures of the
25 compounds may be separated so that the individual enantiomers are isolated. The separation may be carried out by methods well known in the art, such as the coupling of a racemic mixture of compounds to an enantiomerically pure compound to form a diastereomeric mixture, followed by separation of the individual diastereomers by standard methods, such as fractional crystallization or chromatography. Starting materials of particular stereochemistry may either be
30 commercially available or may be made by the methods described herein and resolved by techniques well known in the art. The independent syntheses of these diastereomers or their chromatographic separations may be achieved as known in the art by appropriate modifications.

REPLACED SHEET

Certain compounds according to Formula I, can also exist as tautomers, which have different points of attachment of hydrogen accompanied by one or more double bond shifts. These tautomers, either separately or as mixtures, are also considered to be within the scope of the invention.

5 Certain compounds according to Formula I, may also exist as polymorphs.

The present invention also encompasses geometrical isomers of compounds of Formula I and the mixtures thereof.

10 Particularly useful examples of the present invention include but are not limited to the compounds selected from Table 1:

Table 1

Compound No.	Structure	Compound No.	Structure
1		2	
3		4	
5		6	
7		8	
9		10	

REPLACED SHEET

11		12	
13		14	
15		16	
17		18	
19		20	
21		22	
23		24	
25		26	


REPLACED SHEET

27		28	
29		30	
31		32	
33		34	

The compounds of the present invention can be prepared in accordance with one or more of the Schemes discussed herein. All of the starting materials are either commercially available or can be prepared by procedures that would be well known to one of ordinary skill in organic chemistry.

“L” denotes an appropriate leaving group and as such may vary in nature depending on the exact reaction conditions employed. Some typical leaving groups may be fluorine, chlorine, bromine, iodine, tosyl, mesyl, trifluoromethanesulfonyl and the like, but these should not be construed as limiting as many other leaving groups are also well known to those skilled in the art.

REPLACED SHEET



The compounds of Formula I can be prepared from the compounds of Formula II by following

5 Scheme 1. Deprotection of amino protecting group in the compounds of Formula II is carried out using standard deprotecting reagents for example, trifluoroacetic acid, HCl(g) saturated solution of a solvent such as methanol, ethyl acetate, diethyl ether, dioxane and the like, hydrogenation using Pd/C in a suitable polar solvent or by using a basic amine such as piperidine resulting in compounds of Formula III. (The resulting compounds may be in the form of free


10 amine or salt depending upon the nature of protecting group and corresponding deprotecting agent used). Compounds of Formula III and Formula IV are then coupled using standard peptide coupling conditions, for example, using EDC [1-ethyl-3-(3-dimethylaminopropyl)carbodiimide]/HOBT (1-hydroxybenzotriazole) or DCC (dicyclohexyl carbodiimide), DMAP (4-dimethylaminopyridine) or HATU [O-(7-azabenzotriazole-yl)-

REPLACED SHEET

N,N,N',N'-tetramethyluronium hexafluorophosphate/HOAT (1-hydroxy-7-azabenzotriazole) or by mixed anhydride method using ethyl chloroformate or methyl chloroformate in a suitable solvent such as DMF, DCM (dichloromethane) or THF and the like or mixtures thereof and in the presence of a suitable base such as NMM (*N*-methylmorpholine), DIPEA (*N,N*-diisopropylethylamine) or triethylamine to form compounds of Formula V. The amino protecting group (PG) is then removed by using standard deprotecting reagents, for example, trifluoroacetic acid, HCl(g) saturated solution of a solvent such as methanol, ethyl acetate, diethyl ether, dioxane and the like, hydrogenation using Pd/C in a suitable polar solvent or by using a basic amine such as piperidine to give the compounds of Formula VI. Compounds of Formula VI are then reacted with compounds of Formula VII in the presence of EDC, HOBt and the like and in the presence of a suitable base such as triethyl amine, pyridine, NMM, DMAP, DIPEA and the like and in the presence of a suitable solvents such as DMF, toluene, THF, chloroform, dichloromethane and the like or mixtures thereof to give compounds of Formula I.

REPLACED SHEET

Scheme 2




- 5 The compounds of Formula I can also be obtained by following the Scheme 2. Compounds of Formula IX can be obtained by reacting compounds of Formula VII with compounds of Formula VIII in the presence of EDC, HOBT and the like. The reaction may be carried out in the presence of a suitable base such as triethyl amine, pyridine, NMM, DMAP, DIPEA and the like and in the presence of a suitable solvent such as DMF, toluene, THF, chloroform, dichloromethane and the like or mixtures thereof. The compounds of Formula IX are then converted to compound of Formula X using standard deprotecting reagents, familiar to those skilled in the art. Compounds of Formula X and Formula III are then coupled using standard peptide coupling conditions, for example, using EDC [1-ethyl-3-(3-dimethylaminopropyl)carbodiimide]/HOBT (1-hydroxybenzotriazole or DCC (dicyclohexyl carbodiimide), DMAP (4-dimethylaminopyridine) or HATU [O-(7-azabenzotriazole-yl)-N,N,N',N'-tetramethyluronium hexafluorophosphate/HOAT (1-hydroxy-7-azabenzotriazole) or by mixed anhydride method using ethyl chloroformate or methyl chloroformate in a suitable
- 10
- 15

REPLACED SHEET

solvent such as DMF, DCM (dichloromethane) or THF and the like or mixtures thereof and in the presence of a suitable base such as NMM (*N*-methylmorpholine), DIPEA (*N,N*-diisopropylethylamine) or triethylamine to form compounds of Formula I.

Compounds of Formula II can be easily prepared by those skilled in art. For example
 5 compound of Formula II (when A is Nitrogen and '---' is absent) can be prepared following Scheme 3. Compounds of Formula II (when A is carbon atom and '---' is a single bond or when A is CH and '---' is absent) can be prepared following procedure reported in patent number US 6051716 or WO 2003/097640 or WO 2004/113329.

Scheme 3



10

Compounds of Formula XIII can be obtained by reacting compounds of Formula XI with substituted nitrobenzene derivatives of Formula XII (wherein T,U,V,W are the same as defined earlier and L is an appropriate leaving group such as fluoro, chloro, bromo, iodo) in an
 15 appropriate solvent and base. Examples of appropriate solvents include acetonitrile,

REPLACED SHEET

tetrahydrofuran, methylene dichloride, dichloroethane, DMF, DMSO and the like or mixtures thereof. Examples of appropriate bases include triethylamine, potassium carbonate, diisopropylethyl amine, KOH and the like. The compounds of Formula XIII can then be reacted with a suitable standard amino protecting group (PG), familiar to those skilled in the art, to form compounds of Formula XIV in the presence of a suitable solvent such as methylene dichloride, chloroform, THF and the like or mixtures thereof and in presence of a suitable base such as triethylamine, sodium bicarbonate, diisopropylethyl amine and the like. The nitro derivatives of Formula XIV can then be reduced to the corresponding amino compounds of Formula XV by a variety of reducing agents familiar to those skilled in the art such as hydrogenation over an appropriate catalyst such as palladium, platinum, or ruthenium on activated charcoal or chemical methods such as reaction with Fe/HCl or SnCl₂/HCl or NiCl₂/NaBH₄ or Fe/NH₄Cl. The resulting amines XV can then be treated with benzyl or methyl chloroformate and sodium bicarbonate in presence of water and acetone to form the corresponding benzyl or methyl carbamate derivatives XVI which are then deprotonated in the next step using a lithium base such as *n*-butyllithium followed by the addition of Glycidyl butyrate in presence of a suitable solvent such as diethylether or tetrahydrofuran to afford the oxazolidinones XVII. The hydroxyl group (Formula XVII) can then be converted to Z (Formula II) (wherein Z is as defined earlier). The exact nature of the reagents used for this conversion is dependent on the exact nature of the Z desired. For example, if Z is desired to be -(CH₂)_n-1,2,3-triazolyl group, the hydroxyl group is first converted to azide group which is then reacted with bicyclo[2.2.1]hepta-2,5-diene in suitable solvent like dioxane to give 1,2,3-triazolyl group. If Z is desired to be -(CH₂)_n-4-hydroxymethyl-1,2,3-triazolyl group, the hydroxyl group is first converted to the azide and then reacted with propargyl alcohol in the presence of copper iodide and suitable base and solvent such as DIPEA and THF or the like. The appropriate conditions and reagents for any particular Z group can be readily selected by those having well known skill in the art.

It is understood that, as used herein, references to the compounds of structural Formula I are meant to also include the pharmaceutically acceptable salts, and also salts that are not pharmaceutically acceptable when they are used as precursors to the free compounds or their pharmaceutically acceptable salts or in other synthetic manipulations. The compounds of the present invention may be administered in the form of a pharmaceutically acceptable salt. The term "pharmaceutically acceptable salt" refers to salts prepared from pharmaceutically

REPLACED SHEET

acceptable non-toxic bases or acids including inorganic or organic bases and inorganic or organic acids. The salts may be prepared during the final isolation and purification of the compounds or separately by making basic or acidic addition salts. Representative salts of basic compounds of the present invention can be prepared by reacting free base form of the compound with a suitable acid, including, but not limited to acetate, trifluoroacetate, adipate, citrate, aspartate, benzoate, benzenesulphonate, bisulfate, besylate, butyrate, camphorsulphonate, difluconate, hemisulfate, heptanoate, formate, fumarate, lactate, maleate, methanesulfonate, naphthylsulfonate, nicotinate, oxalate, picrate, pivalate, succinate, tartrate, tirchloracetat, glutamate, *p*-toluenesulphonate, hydrochloric, hydrobromic, sulphuric, phosphoric and the like. Representative salts of acidic compounds of the present invention can be prepared by reacting free acid form of the compound with a suitable base, including, but not limited to ammonium, calcium, magnesium, potassium, sodium salts, salts of primary, secondary and tertiary amines, substituted amines including naturally occurring ones e.g., arginine, betaine, caffeine, choline, glucamine, glucosamine, histidine, lysine, morpholine, piperazine, piperidine, purine, triethylamine and the like. Compounds of the present invention that contain a carboxylic acid (-COOH) or alcohol group, their pharmaceutically acceptable esters of carboxylic acids such as methyl, ethyl and the like, or acyl derivatives of alcohols such as acetate and the like, can be employed. Compounds of the present invention that comprise basic nitrogen atom may be quaternized with alkyl halides, alkyl sulfates and the like. Such salts permit the preparation of both water soluble and oil soluble compounds of the present invention. It should be recognized that the free base or free acid forms will typically differ from their respective salt forms somewhat in physical properties such as solubility in polar solvents, but otherwise the salts are equivalent to their respective free forms for the purpose of the invention.

The "pharmaceutically acceptable solvates" refer to solvates with water (i.e., hydrates) or pharmaceutically acceptable solvents, for example, ethanol and the like.

The invention also encompasses "prodrugs" of the compounds of the present invention which upon *in-vivo* administration undergo cleavage by metabolic processes before becoming active pharmacological substances. In general such prodrugs are derivatives of functional group of a compound of the invention which are readily convertible *in vivo* into the compound of the invention. Conventional procedures for the selection and preparation of suitable prodrug

REPLACED SHEET

derivatives are described, for example, in "Targeted prodrug design to optimize drug delivery", *AAPS PharmaSci* (2000), 2(1), E6.

The invention also encompasses active "metabolites" of the compound of the present invention.

5 Various "polymorphs" of a compound of general Formula I forming part of this invention may be prepared by crystallization of a compound of Formula I under different conditions. For example, by using different solvents commonly used or their mixtures for recrystallization; crystallizations at different temperatures; various modes of cooling, ranging from very fast to very slow cooling during crystallizations, heating or melting the compound
10 followed by gradual or fast cooling may also obtain polymorphs. The presence of polymorphs may be determined by solid probe NMR spectroscopy, IR spectroscopy, differential scanning calorimetry, powder X-ray diffraction or such other techniques.

The present invention also provides pharmaceutical compositions, comprising compounds of the present invention or their pharmaceutically acceptable derivatives,
15 tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof optionally in combination with one or more pharmaceutically acceptable carriers comprising excipients and auxiliaries. The pharmaceutical compositions may be in any form known in the art, such as tablets, capsules, powders, syrups, solutions, suspensions and the like, may contain
20 media to form injectable solutions or suspensions. Such compositions typically contain active compound optionally in combination with pharmaceutically acceptable carriers, diluents or solvents.

The pharmaceutical compositions of the present invention can be manufactured by the processes well known in the art, for example, by means of conventional mixing, dissolving, dry
25 granulation, wet granulation, dragee-making, levigating, emulsifying, encapsulating, entrapping, lyophilizing processes or spray drying. The compounds or the pharmaceutical compositions comprising such compounds of the present invention may be administered in the form of any pharmaceutical Formulation. The pharmaceutical formulation will depend upon the nature of the active compound and its route of administration. Any route of administration
30 may be used, for example oral, buccal, pulmonary, topical, parenteral (including subcutaneous, intramuscular and intravenous), transdermal, ocular (ophthalmic), by inhalation, intranasal,

REPLACED SHEET

transmucosal, implant or rectal administration. Preferably the compounds of the present invention are administered orally, parenterally or topically.

5 In an embodiment, the amount of the novel compounds having the Formula I according to the present invention to be incorporated into the pharmaceutical compositions of the present invention can vary over a wide range depending on known factors such as, for example, the disorder to be treated, the severity of the disorder, the patient's body weight, the dosage form, the chosen route of administration and the number of administrations per day. Typically, the amount of the compound of Formula I in the pharmaceutical compositions of the present invention will range from approximately 0.01 mg to about 5000 mg. In an embodiment, the
10 daily dose of composition comprising the novel compounds having the Formula I is in the range of about 0.01 mg/kg to about 100 mg/kg based on the body weight of the subject in need thereof which may be administered as a single or multiple doses.

In an embodiment, the novel compounds having the Formula I according to the present invention are particularly useful for the treatment of disease(s) or disorder(s) which are
15 particularly acute in nature and which require a short term but mild to moderate treatment, or even some chronic conditions which favorably respond to or are alleviated by the novel compounds having the Formula I or compositions comprising them. The compositions comprising the novel compounds having the Formula I are useful prophylactically or therapeutically depending upon the pathological condition intended to be prevented or treated
20 respectively.

The compounds of the present invention are effective against a number of aerobic and/or anaerobic Gram positive and/or Gram negative pathogens such as multi drug resistant species of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacterioides*, *Clostridia*, *H. influenza*, *Moraxella*, acid-fast organisms such as *Mycobacterium tuberculosis* as well as
25 Linezolid resistant species of *Staphylococcus* and *Enterococcus*.

Thus, a further embodiment of the present invention is the use of a compound of Formula I for the manufacture of a medicament for the prophylaxis, amelioration and/or treatment of bacterial infections in a subject in need thereof preferably a mammal including a human. Another embodiment of the present invention provides methods for the management
30 such as prophylaxis, amelioration and/or treatment of bacterial infections in a subject in need thereof preferably a mammal including a human that comprises administering a therapeutically

REPLACED SHEET

effective amount of compound of Formula I. In still another embodiment of the present invention is provided use of the dosage form compositions comprising the novel compounds of Formula I for the treatment of disease(s)/disorder(s) which comprises administering to a subject in need thereof a pharmaceutically effective amount of the composition.

5

The compounds of the present invention may be used in combination with one or more other active ingredients such as quinolones, β -lactams e.g., cephalosporins, penicillins, penams, penems and the like in the prophylaxis, amelioration and/or treatment of bacterial infections, where the combination of the active ingredients together are safer or more effective than either active ingredient alone or where incorporation of another active ingredient might reduce the dose of the compound of Formula I.

10

***In-vitro* antibacterial activity :**

The *in-vitro* antibacterial activity of the compounds of the present invention (as described in Table 2) was determined by a broth microdilution following the guidelines prescribed by the Clinical and Laboratory Standards Institute (CLSI). This method is described in the CLSI Document M7-A7, Vol.26, No.2, "*Methods for Dilution Antimicrobial Susceptibility Test for Bacteria that Grow Aerobically; Approved Standard-Seven Edition*", which is incorporated herein by reference. Minimum Inhibitory Concentration (MIC) is defined as the minimum concentration of test compound which inhibits the growth of bacteria as visible or seen by the naked eye. This test can also be carried out by agar dilution method.

15

The compounds of the present invention were tested against a panel of standard microorganisms obtained from ATCC (American type culture collection), and a Linezolid resistant strain (LRSA) i.e. PTCC 100 (Panacea type culture collection). PTCC 100 is a repository that has been created by Panacea Biotec Ltd. at Mohali, India for storage and maintenance of clinical, bacterial and other isolates developed in-house which are used for testing the test compounds. Linezolid was used as comparator in all the tests.

20

Organism	Culture No.	Type
<i>Staphylococcus aureus</i>	ATCC 29213	MSSA (Methicillin sensitive)
<i>Staphylococcus aureus</i>	ATCC 33591	MRSA (Methicillin resistant)
<i>Enterococcus faecalis</i>	ATCC 29212	Vancomycin Sensitive

REPLACED SHEET

<i>Enterococcus faecium</i>	ATCC 700221	VRE (Vancomycin resistant <i>E.faecium</i>)
<i>Staphylococcus aureus</i>	PTCC 100	LRSA (Linezolid resistant <i>S.aureus</i>)

In the broth microdilution method, the compound was dissolved in dimethylsulfoxide and two fold serial dilutions were carried out in 96 well microtitre plates. The inoculum was prepared by adjusting the turbidity of actively growing broth culture and added to the wells to obtain a final bacterial count of $\sim 2.5 \times 10^4$ CFU/well. The microtitre plates were incubated at 35 ± 2 °C for 16-20 h and then read visually. MICs ($\mu\text{g/mL}$) values of some of the compounds of Formula 1 are presented in the Table 2 and Table 3.

Table 2: *In-vitro* antibacterial activity MICs ($\mu\text{g/mL}$)

Compound No.	MIC($\mu\text{g/mL}$)			
	<i>S. aureus</i> ATCC 29213	<i>S. aureus</i> ATCC 33591 (MRSA)	<i>E. faecalis</i> ATCC 29212	<i>E. faecium</i> ATCC 700221 (VRE)
1	0.5	0.25	0.5	0.5
2	0.25	0.25	0.25	0.25
3	0.5	0.5	0.5	0.5
4	0.25	0.125	0.25	0.125
5	2	0.5	0.5	0.5
6	16	8	16	4
7	0.5	0.25	0.25	0.25
8	1	0.5	0.5	0.5
9	0.5	0.5	0.25	0.25
10	1	0.5	0.5	0.25
11	2	2	1	1
12	0.5	0.5	0.25	0.25
13	0.5	0.25	0.25	0.25
14	0.5	0.25	0.25	0.25
15	0.5	0.5	0.25	0.25

REPLACED SHEET

16	>32	>32	>32	32
17	16	8	16	4
18	4	2	2	1
19	8	4	2	2
20	4	2	1	2
21	2	2	1	0.5
22	8	4	4	8
23	2	2	2	2
24	0.5	0.25	0.25	0.25
25	0.25	0.25	0.25	0.25
26	0.25	0.25	0.25	0.25
27	0.5	0.5	0.5	0.25
28	2	1	2	2
29	0.5	0.25	0.5	0.25
30	0.125	0.125	0.125	0.125
31	1	0.5	0.5	0.25
32	2	1	1	2
33	1	1	1	1
34	8	4	8	2
Linezolid	2	1	2	2

Development of in-house LRSA strain (PTCC 100):

- 5 PTCC 100 was developed by a procedure similar to the one cited in *Antimicrobial Agents and Chemotherapy*, 2008, 52, 1940. Female Swiss albino mice (18-22gm) bred in-house were inoculated with *S.aureus* ATCC 29213 strain and dosed orally with Linezolid at 5 mg/kg/p.o, next day mice were sacrificed after 20-22 h and intraperitoneal swab was taken and streaked onto Mueller Hinton Agar plates containing 4 and 8 µg/mL of Linezolid. Colonies obtained on 4 µg/mL and 8 µg/mL were selected from plates and further passaged into mice (SAM), dosed orally with Linezolid at 7.5 mg/kg/p.o. Mice were sacrificed and intraperitoneal swabs were streaked onto plates containing higher linezolid concentration i.e 16 and 32 µg/mL and the process was

REPLACED SHEET

repeated by incrementally increasing the concentration of Linezolid upto 10 mg/kg/p.o in mice, to finally obtain *S.aureus* strains resistant to Linezolid at 64 µg/mL. Minimum inhibitory concentration (MIC) of isolated colonies was determined by broth microdilution assay and MIC values of 64 µg/ mL for Linezolid confirmed the development of in house LRSA strain, PTCC 100.

Table 3: *In-vitro* antibacterial activity MICs (µg/mL) against *S. aureus* PTCC 100 (LRSA)

Compound No.	MIC(µg/mL) <i>S. aureus</i> PTCC 100 (LRSA)
1	4
4	1
7	4
12	2
13	1
14	1
15	2
18	8
19	8
20	8
21	16
22	8
23	16
25	2
27	2
28	4
Linezolid	32

REPLACED SHEET

In-vivo* efficacy studies:*Systemic Model of infection in Mice**

Female Swiss albino mice bred in-house were selected in weight range of 19-23 gm (n=6/group).

- 5 *S.aureus* ATCC 29213/ *S.aureus* PTCC 100 (LRSA) was grown overnight for 18-20 h, on Columbia Blood Agar (Difco;BD). Next day bacterial inoculum was prepared with optical density (O.D) corresponding to cell density of $\sim 2 \times 10^9$ CFU/ml and mixed with 10% of gastric mucin (Difco;BD) in ratio of 1:1 to obtain final mucin concentration of 5% w/v. 0.5 mL of bacterial inoculum was injected intraperitoneally (*i.p*) into all the mice. Compounds obtained in
- 10 the present invention, hereinafter referred as Test compounds were formulated in 0.25% Carboxymethylcellulose (C.M.C) and Tween 80, at different dose levels and then administered orally at 1h and 5h post infection. Linezolid was used as standard control. Saline was administered to the infection control group which received neither the test compound nor the standard drug. Mice were observed for 7 days post treatment. Numbers of survivors in each
- 15 group were noted and ED₅₀ of test compound on the basis of 50% survival was calculated through regression analysis.

Table 4: ED₅₀ values (mg/kg/p.o.) against *S.aureus* ATCC 29213

Compound No.	ED ₅₀ values (mg/kg/p.o.) <i>S.aureus</i> ATCC 29213
4	3.3
10	5.3
12	3.8
13	1.8
14	0.9
27	2.2
Linezolid	6.0

20

REPLACED SHEET

Table 5: ED₅₀ values (mg/kg/p.o.) against *S.aureus* PTCC 100 (LRSA)

Compound No.	ED ₅₀ values (mg/kg/p.o.) <i>S.aureus</i> PTCC 100 (LRSA)
13	< 10
14	< 10
Linezolid	45

EXAMPLES

- 5 The invention is explained in detail in the following examples which are given solely for the purpose of illustration only and therefore should not be construed to limit the scope of the invention. All of the starting materials are either commercially available or can be prepared by procedures that would be well known to one of ordinary skill in organic chemistry. Solvents
- 10 were dried prior to use wherever necessary by standard methods (Perrin, D.D.; Armarego, W.L.F. Purification of Laboratory Chemicals, Pergamon Press: Oxford, 1988). Mass spectra (MS) were obtained by electron spray ionization (ESI) eV using Applied biosystem 4000 Q TRAP. ¹H NMR were recorded on Bruker 400 MHz Avance II NMR spectrometer. Chemical shifts are reported as δ values in parts per million (ppm), relative to TMS as internal standard.
- 15 All coupling constants (*J*) values are given in Hz.

Abbreviations

The following abbreviations are employed in the examples and elsewhere herein:

¹ H NMR	proton nuclear magnetic resonance
C	centigrade
CDCl ₃	deuterated chloroform
CuI	copper(I) iodide
DCM	dichloromethane
d	doublet
DAST	(diethylamino)sulfur trifluoride
DCC	dicyclohexyl carbodiimide

REPLACED SHEET

dd	doublet of doublet
DIPEA	diisopropylethylamine
DMAP	4-dimethylaminopyridine
DMF	<i>N,N</i> -dimethylformamide
DMSO	dimethylsulfoxide
EDC	1-ethyl-3-(3-dimethylaminopropyl)carbodiimide
ESIMS	electron spray ionization mass Spectroscopy
Fe	iron
g	gram(s)
h	hour(s)
HATU	O-(7-azabenzotriazole-1-yl)- <i>N,N,N',N'</i> -tetramethyluronium hexafluorophosphate
HCl	hydrochloric acid
HOAt	1-hydroxy-7-azabenzotriazole
HOBt	1-hydroxybenzotriazole
Hz	hertz
<i>J</i>	coupling constant
KOH	potassium hydroxide
<i>M</i>	molar
m	multiplet
mg	milligram
min	minutes
mL	milliliter
mmol	millimoles
mol	moles
<i>n</i> -BuLi	<i>n</i> -butyl lithium
NaBH ₄	sodium borohydride
NH ₄ Cl	ammonium chloride
NMM	<i>N</i> -methylmorpholine
NMR	nuclear magnetic resonance

REPLACED SHEET

NiCl ₂	nickel(II) chloride
Pd/C	palladium on carbon
Pet. Ether	petroleum ether
q	quartet
r. t.	room temperature
s	singlet
SnCl ₂	tin(II) chloride
t	triplet
TFA	trifluoroacetic acid
THF	tetrahydrofuran
TLC	thin layer chromatography

Preparation of starting materials:**5 Intermediate I: 1-(2,6-difluoro-4-nitrophenyl)piperazine:**

A solution of piperazine (24 g, 0.28 mol) and 3,4,5-trifluoronitrobenzene (13 mL, 0.11 mol) in acetonitrile (200 mL) was stirred at 60 °C. The progress of reaction was monitored by TLC. On completion, acetonitrile was evaporated under reduced pressure. The residue was dissolved in ethyl acetate (300 mL) and resulting solution was washed with water (100 mL), brine (100 mL),
 10 dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, 1:9 methanol:chloroform) to provide title compound (25.8 g, 92%) as orange solid.

ESIMS (m/z): 244.1 (M+1)

Intermediate II: *tert*-butyl 4-(2,6-difluoro-4-nitrophenyl)piperazine-1-carboxylate:

15 To a solution of 1-(2,6-difluoro-4-nitrophenyl)piperazine (Intermediate I) (25 g, 0.1 mol), in THF (200 mL) was added Boc anhydride (26.2 g, 0.12 mol) at 0 °C. The solution was stirred at 0 °C and progress of reaction was monitored by TLC. On completion, THF was evaporated under reduced pressure and the solid obtained was washed with Pet. ether (3X100 mL). The yellow solid (34 g, 96%) obtained was subjected to next reaction without further purification.

20 ESIMS (m/z): 344.1 (M+1)

Intermediate III: *tert*-butyl 4-(4-amino-2,6-difluorophenyl)piperazine-1-carboxylate:

REPLACED SHEET

To a solution of *tert*-butyl 4-(2,6-difluoro-4-nitrophenyl)piperazine-1-carboxylate (Intermediate II) (30 g, 0.09 mol), in methanol (500 mL) under argon atmosphere was added 10% Pd/C (4.5 g, 15 mol% by weight). Flask was evacuated and hydrogen was introduced with the help of balloon. The reaction mixture was stirred under hydrogen and progress of the reaction was monitored by TLC. On completion, the reaction mixture was filtered through celite pad using methanol as solvent. The filtrate was evaporated to provide title compound (26 g, 95%) as pale yellow solid.

ESIMS (m/z): 336.7 (M+23), 314.8 (M+1)

Intermediate IV: *tert*-butyl 4-(4-(((benzyloxy)carbonyl)amino)-2,6-difluorophenyl)piperazine-1-carboxylate:

To a solution of *tert*-butyl 4-(4-amino-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate III) (25 g, 0.08 mol), in 1:1 acetone:water (300 mL) was added sodium bicarbonate (15.1 g, 0.18 mol). The resulting solution was cooled to 0 °C and benzyl chloroformate (40 mL, 0.24 mol, 50% solution in toluene) was added dropwise. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, solvent was evaporated under reduced pressure and the residue was dissolved in ethyl acetate (500 mL). The organic layer was washed with water (2X50 mL), brine (100 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, 2:3 ethyl acetate:Pet. ether) to provide title compound (32 g, 90%) as off white solid.

ESIMS (m/z): 448.0 (M+1)

Intermediate V: (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-(hydroxymethyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate:

To a solution of *tert*-butyl 4-(4-(((benzyloxy)carbonyl)amino)-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate IV) (30 g, 0.067 mol), in dry THF (300 mL) was added *n*-BuLi (75 mL, 0.12 mol, 1.6 M solution in hexane) dropwise under nitrogen atmosphere at -78 °C. The reaction mixture was stirred at the same temperature for one hour and then (*R*)-glycidyl butyrate (10.4 mL, 0.074 mol) was added dropwise over a period of 5 min. The reaction mixture was stirred at -78 °C for additional two hours and then warmed to r.t. The progress of reaction was monitored by TLC and on completion, the reaction mixture was quenched with saturated NH₄Cl solution (400 mL) and extracted with ethyl acetate (4X200 mL). The organic layer was washed with brine (100 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The

REPLACED SHEET

residue was purified by column chromatography (silica gel, 3:2 ethyl acetate:Pet. ether) to provide title compound (18 g, 65%) as off white solid.

ESIMS (m/z): 452.7 (M+39), 436.6 (M+23), 414.7 (M+1)

Intermediate VI: (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-(((methylsulfonyl)oxy)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate:

To a solution of (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-(hydroxymethyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate (Intermediate V) (10 g, 24.2 mmol), in DCM (100 mL) was added triethylamine (10.5 mL, 73 mmol). The reaction mixture was cooled to 0 °C and methanesulfonyl chloride (2.8 mL, 36 mmol) was added dropwise. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, the reaction mixture was diluted with DCM (100 mL). The organic layer was washed with water (25 mL), brine (25 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The crude product (11.3 g, 95%) was obtained as brown solid and subjected to further reaction without any purification.

ESIMS (m/z): 514.8 (M+23), 492.6 (M+1)

Intermediate VII: (*R*)-*tert*-butyl 4-(4-(5-(azidomethyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate:

To a solution of (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-(((methylsulfonyl)oxy)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate (Intermediate VI) (11 g, 22.4 mmol), in DMF (50 mL) was added sodium azide (4.37 g, 67.2 mmol). The reaction mixture was stirred at 80 °C and progress of the reaction was monitored by TLC. On completion, the reaction mixture was diluted with water (100 mL) and extracted with ethyl acetate (2X100 mL). The organic layer was washed with brine (50 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The crude product (8 g, 82%) was obtained as off white solid and subjected to further reaction without any purification.

ESIMS (m/z): 439.7 (M+1)

Intermediate VIII: (*S*)-*tert*-butyl 4-(4-(5-(aminomethyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate:

A mixture of (*R*)-*tert*-butyl 4-(4-(5-(azidomethyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate VII) (5 g, 11.4 mmol) and triphenylphosphine (3.3 g, 12.5 mmol) in THF (80 mL) was stirred at r.t. for 3 h. Water (3 mL)

REPLACED SHEET

was added and the reaction mixture was stirred at 40 °C for 16 h. The reaction mixture was then diluted with water (50 mL) and extracted with ethyl acetate (4X50 mL). The organic layer was washed with brine (50 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, 1:9 methanol:chloroform) to provide title compound (3.5 g, 74%) as off white solid.


ESIMS (*m/z*): 435.8 (*M*+23), 413.7 (*M*+1)

Intermediate IX: (*R*)-*tert*-butyl 4-(4-(5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate:

To a solution of (*R*)-*tert*-butyl 4-(4-(5-(azidomethyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate VII) (2 g, 4.6 mmol), in dioxane (25 mL) was added bicyclo[2.2.1]hepta-2,5-diene (1.9 mL, 18.5 mmol) and the resulting solution was stirred at 60 °C for 8 h. The solvent was evaporated and the residue was purified by column chromatography (silica gel, 1:20 methanol:chloroform) to provide title compound (1.47 g, 70%) as white solid.

ESIMS (*m/z*): 465.8 (*M*+1)

Example I: *N*-((*S*)-1-(4-(4-((*R*)-5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazin-1-yl)-1-oxopropan-2-yl)nicotinamide



Step 1: (*R*)-4-(4-(5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazin-1-ium 2,2,2-trifluoroacetate:

To a stirred solution of (*R*)-*tert*-butyl 4-(4-(5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate IX) (650 mg, 1.4 mmol), in DCM (5 mL) at 0 °C under nitrogen atmosphere was added TFA (4.5 mL) dropwise. The mixture was stirred at 0 °C and progress of the reaction was monitored by TLC. On completion, excess TFA and DCM were evaporated under reduced pressure to obtain the title compound (660 mg, 98%) as brown solid and subjected to further reaction without any purification.

ESIMS (*m/z*): 365.6 (*M*+1) of free amine.

REPLACED SHEET

Step 2: *tert*-butyl ((*S*)-1-(4-(4-((*R*)-5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazin-1-yl)-1-oxopropan-2-yl)carbamate:

To a solution of compound (650 mg, 1.36 mmol) obtained in step 1, in DCM (30 mL), were added (*S*)-*N*-(*tert*-butoxycarbonyl)alanine (309 mg, 1.63 mmol), EDC (339 mg, 1.77 mmol),
5 HOBt (239 mg, 1.77 mmol) and DIPEA (0.72 mL, 4.08 mmol) at 0° C. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, the reaction mixture was diluted with DCM (100 mL). The organic layer was washed with water (50 mL), brine (50 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, 3:5 ethyl acetate:Pet. ether) to provide title
10 compound (550 mg, 76%) as white solid.

ESIMS (*m/z*): 558.8 (*M*+23), 536.6 (*M*+1).

Step 3: (*S*)-1-(4-(4-((*R*)-5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazin-1-yl)-1-oxopropan-2-aminium 2,2,2-trifluoroacetate:

15 To a stirred solution of compound (535 mg, 1 mmol) obtained in Step 2, in DCM (2 mL) at 0 °C under nitrogen atmosphere was added TFA (3 mL) dropwise. The mixture was stirred at 0 °C and progress of the reaction was monitored by TLC. On completion, excess TFA and DCM were evaporated under reduced pressure to obtain the title compound (540 mg, 98%) as brown solid and subjected to further reaction without any purification.

20 ESIMS (*m/z*): 458.8 (*M*+23), 436.6 (*M*+1) of free amine.

Step 4: *N*-((*S*)-1-(4-(4-((*R*)-5-((1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazin-1-yl)-1-oxopropan-2-yl)nicotinamide:


To a solution of compound (300 mg, 0.57 mmol) obtained in step 3, in DMF (10 mL), were
25 added nicotinic acid (81 mg, 0.66 mmol), EDC (127 mg, 0.66 mmol), HOBt (89 mg, 0.66 mmol) and NMM (0.15 mL, 1.35 mmol) at 0 °C. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, DMF was evaporated *in vacuo* and residue was dissolved in chloroform (100 mL). The organic layer was washed with water (50 mL), brine (50 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was
30 purified by column chromatography (silica gel, 1:20 methanol:chloroform) to provide title compound (204 mg, 69%) as white solid.

REPLACED SHEET

¹H NMR (400 MHz, CDCl₃, δ): 1.49 (d, *J* = 6.8 Hz, 3H), 3.10-3.30 (m, 4H), 3.65-3.80 (m, 3H), 3.80-3.90 (m, 1H), 3.92 (dd, *J* = 9.3 and 6.0 Hz, 1H), 4.11 (t, *J* = 9.1 Hz, 1H), 5.00-5.20 (m, 2H), 6.99 (d, *J* = 10.7 Hz, 2H), 7.39 (dd, *J* = 7.9 and 4.9 Hz, 1H), 7.47 (d, *J* = 6.8 Hz, 1H), 7.76 (d, *J* = 8.5 Hz, 2H), 8.10-8.15 (m, 1H), 8.75 (dd, *J* = 4.7 and 1.2 Hz, 1H), 9.06 (d, *J* = 2.0 Hz, 1H).

ESIMS (*m/z*): 563.7 (M+23), 541.5 (M+1)

Example II: *N*-((*S*)-1-(4-(2,6-difluoro-4-((*R*)-5-((4-(fluoromethyl)-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazin-1-yl)-1-oxopropan-2-yl)nicotinamide



Step 1: (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-((4-(hydroxymethyl)-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate:

To a stirred solution of (*R*)-*tert*-butyl 4-(4-(5-(azidomethyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate VII) (1.3 g, 2.97 mmol) in THF (25 mL) was added propargyl alcohol (0.5 mL, 8.9 mmol), DIPEA (1.1 mL, 5.94 mmol) and CuI (0.28 g, 1.48 mmol) at 0 °C. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, reaction mixture was quenched with saturated solution of ammonium chloride in liquor ammonia (20 mL), diluted with water (50 mL) and extracted with ethyl acetate (4X50 mL). The organic layer was washed with brine (25 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, 1:10 methanol:chloroform) to provide title compound (1.24 g, 85%) as cream solid.

ESIMS (*m/z*): 495.5 (M+1)

Step 2: (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-((4-(fluoromethyl)-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate:

To a stirred solution of compound (500 mg, 1.01 mmol) obtained in Step 1, in DCM (15 mL) was added DAST (0.5 mL, 4.04 mmol) at -20 °C. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, reaction mixture was quenched

REPLACED SHEET

with saturated solution of sodium bicarbonate (25 mL) and extracted with DCM (4X50 mL). The organic layer was washed with water (50 mL), brine (25 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The residue was purified by column chromatography (silica gel, 1:10 methanol:chloroform) to provide title compound (300 mg, 60%) as cream solid.


5 ESIMS (m/z): 497.5 (M+1)

Compound obtained in Step 2 Example II, was converted to *N*-((*S*)-1-(4-(2,6-difluoro-4-((*R*)-5-((4-(fluoromethyl)-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazin-1-yl)-1-oxopropan-2-yl)nicotinamide following the procedure described in steps 1 to 4 in Example I.

10 ¹H NMR (400 MHz, CDCl₃, δ): 1.49 (d, *J* = 6.8 Hz, 3H), 3.10-3.25 (m, 4H), 3.60-3.75 (m, 3H), 3.80-3.90 (m, 1H), 3.91 (dd, *J* = 9.4 and 6.3 Hz, 1H), 4.13 (t, *J* = 9.1 Hz, 1H), 4.60-4.70 (m, 2H), 5.05-5.20 (m, 2H), 5.43 (s, 1H), 5.55 (s, 1H), 7.01 (d, *J* = 10.8 Hz, 2H), 7.39 (dd, *J* = 7.8 and 4.9 Hz, 1H), 7.48 (d, *J* = 6.8 Hz, 1H), 7.88 (d, *J* = 2.4 Hz, 1H), 8.10-8.15 (m, 1H), 8.74 (dd, *J* = 4.8 and 1.6 Hz, 1H), 9.06 (d, *J* = 2.0 Hz, 1H).

15 ESIMS (m/z): 595.9 (M+23), 573.6 (M+1)

Example III: *N*-((*S*)-1-(4-(2,6-difluoro-4-((*R*)-5-((4-methyl-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazin-1-yl)-1-oxopropan-2-yl)imidazo[1,2-*a*]pyrimidine-2-carboxamide



20 **Step 1:** (*R*)-*tert*-butyl 4-(2,6-difluoro-4-(5-((4-methyl-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazine-1-carboxylate:

To a stirred solution of (*S*)-*tert*-butyl 4-(4-(5-(aminomethyl)-2-oxooxazolidin-3-yl)-2,6-difluorophenyl)piperazine-1-carboxylate (Intermediate VIII) (780 mg, 1.89 mmol) in methanol
25 (15 mL) was added DIPEA (1.2 mL, 7.56 mmol) followed by *N*-(1,1-dichloropropan-2-ylidene)-4-methylbenzenesulfonohydrazide (680 mg, 2.2 mmol) at 0 °C. The reaction mixture was stirred at r.t. and progress of the reaction was monitored by TLC. On completion, the solvent was evaporated *in vacuo*. The residue was dissolved in chloroform (25 mL) and washed with water (20 mL), brine (20 mL), dried over anhydrous sodium sulphate and concentrated *in vacuo*. The

REPLACED SHEET


residue was purified by column chromatography (silica gel, 1:10 methanol:chloroform) to provide title compound (590 mg, 65%) as cream solid.

ESIMS (m/z): 479.5 (M+1)

- 5 Compound obtained in Step 1 Example III, was converted to *N*-((*S*)-1-(4-(2,6-difluoro-4-((*R*)-5-((4-methyl-1*H*-1,2,3-triazol-1-yl)methyl)-2-oxooxazolidin-3-yl)phenyl)piperazin-1-yl)-1-oxopropan-2-yl)imidazo[1,2-*a*]pyrimidine-2-carboxamide following the procedure described in steps 1 to 4 in Example I. Imidazo[1,2-*a*]pyrimidine-2-carboxylic acid was used in place of nicotinic acid in step 4.
- 10 ¹H NMR (400 MHz, CDCl₃, δ): 1.50 (d, *J* = 6.8 Hz, 3H), 2.35 (s, 3H), 3.10-3.25 (m, 4H), 3.65-3.80 (m, 3H), 3.80-3.90 (m, 1H), 3.91 (dd, *J* = 9.3 and 6.3 Hz, 1H), 4.09 (t, *J* = 9.1 Hz, 1H), 4.71 (d, *J* = 3.94 Hz, 2H), 4.95-5.05 (m, 1H), 5.05-5.20 (m, 1H), 7.01 (d, *J* = 10.8 Hz, 2H), 7.05 (dd, *J* = 6.9 and 4.1 Hz, 1H), 7.39 (d, *J* = 7.2 Hz, 1H), 7.49 (s, 1H), 8.30 (s, 1H), 8.68 (dd, *J* = 4.1 and 2.1 Hz, 1H), 9.75 (dd, *J* = 7.0 and 2.1 Hz, 1H).
- 15 ESIMS (m/z): 617.9 (M+23), 595.7 (M+1)

























The compounds listed in Table 5 were prepared essentially following the procedures described for Examples I to III.

Table 6

























S.No.	R ⁴	R ²	R ³	T	Z	ESIMS
1		-H	-H	-H		571.2 (M+23) 549.4 (M+1)
2		-H	-H	-F		605.3 (M+39) 589.3 (M+23) 567.4 (M+1)
3		-H	-H	-F		605.1 (M+39) 589.3 (M+23)



REPLACED SHEET

						567.4 (M+1)
4		-CH ₃	-H	-F		619.3 (M+39) 603.5 (M+23) 581.4 (M+1)
5		-CH(CH ₃) ₂	-H	-F		570.1 (M+1)
6		-CH ₃	-H	-F		594.0 (M+23) 572.1 (M+1)
7		-CH ₃	-H	-F		577.9 (M+23) 555.5 (M+1)
8		-CH ₃	-H	-F		636.1 (M+23) 613.9 (M+1)
9		-CH ₃	-H	-F		654.0 (M+1)
10		-CH ₃	-H	-F		569.8 (M+39) 553.9 (M+23) 531.8 (M+1)
11		-CH ₃	-H	-F		564.0 (M+23) 541.6 (M+1)
12		-CH ₃	-H	-F		597.7 (M+23) 575.4 (M+1)
13		-CH ₃	-H	-H		563.9 (M+23) 541.9 (M+1)
14		-CH ₃	-H	-H		579.9, 581.8 (M+23) 557.7, 559.7 (M+1)
15		-CH ₃	-H	-F		581.8 (M+23) 559.7 (M+1)

REPLACED SHEET

16		-CH ₃	-H	-F		581.7 (M+23) 559.7 (M+1)
17		-CH ₃	-H	-F		595.3 (M+23) 573.5 (M+1)
18		-CH ₃	-H	-F		571.5 (M-1)
19		-CH ₃	-H	-F		612.0 (M+23) 590.0 (M+1)
20		-CH ₃	-H	-F		613.5 (M+23) 591.5 (M+1)
21		-CH ₃	-H	-F		611.8 (M+23) 589.6 (M+1)
22		-CH ₃	-H	-F		613.9 (M+23) 591.5 (M+1)
23		-CH ₃	-H	-F		597.7, 599.7 (M+23) 575.4, 577.4 (M+1)
24		-H	-H	-F		567.8 (M+23) 545.8 (M+1)
25		-CH ₃	-H	-F		597.8, 599.6 (M+23) 575.5, 577.7 (M+1)
26		-CH ₃	-H	-F		579.8 (M+23) 557.5 (M+1)


REPLACED SHEET

27		-CH ₃	-H	-F		564.8 (M+23) 542.8 (M+1)
----	---	------------------	----	----	---	-----------------------------

REPLACED SHEET

We claim:

1. A compound of Formula I,



Formula I

its pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof, wherein:

'---' is independently a single bond or absent;

when '---' is a single bond, 'A' represents carbon atom and when '---' is absent, 'A' is CH or N;

Y and Y' are same or different and independently represent O or S;

R^1 and R^2 are same or different and independently represent hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_1-C_{12} alkoxy, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_nC(=Y)NR^5R^6$, $-(CH_2)_nC(=Y)OR^5$, $-(CH_2)_nNR^5R^6$, $-(CH_2)_nOC(=Y)R^5$, $-(CH_2)_nOC(=Y)OR^5$, $-(CH_2)_nOC(=Y)NR^5R^6$, $-(CH_2)_nN(R^5)C(=Y)OR^6$, $-(CH_2)_nN(R^5)C(=Y)NR^5R^6$, $-(CH_2)_nNR^5C(=Y)R^6$, $-(CH_2)_nC(=Y)R^5$, $-(CH_2)_nYR^5$ (wherein each methylene group may be substituted by one or more halogen atoms), $-C(=Y)NR^5R^6$, $-OC(=Y)R^5$, $-OC(=Y)NR^5R^6$, $-C(=Y)OR^5$, $-OR^5$, $-OC(=Y)OR^5$, $-SR^5$, $-NO_2$, $-NR^5R^6$, $-N(R^5)C(=Y)R^6$, $-N(R^5)-C(=Y)OR^6$, or $-N(R^5)C(=Y)NR^5R^6$, each of which may be optionally substituted at any available position by one or more substituents R^a ; or

R^1 and R^2 can together with the carbon atom to which they are attached form a 3 to 10 membered monocyclic ring, partially unsaturated or saturated, which may contain from one to three heteroatoms independently selected from O, S or N; the ring thus formed may be fused with one or two rings independently selected from the group comprising an aryl ring, a cycloalkyl ring, a heterocyclyl ring or monocyclic heteroaryl ring; the ring thus formed can be optionally substituted at any available position by one or more substituents R^a ;

REPLACED SHEET

R^3 represents hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_nYR^5$, $-(CH_2)_nC(=Y)R^5$, $-(CH_2)_nNR^5R^6$, $-(CH_2)_nC(=Y)NR^5R^6$, $-(CH_2)_nC(=Y)OR^5$, $-(CH_2)_nOC(=Y)R^5$, $-(CH_2)_nOC(=Y)OR^5$, $-(CH_2)_nNR^5C(=Y)R^6$, $-(CH_2)_nN(R^5)C(=Y)OR^6$, $-(CH_2)_nN(R^5)C(=Y)NR^5R^6$, $-(CH_2)_nOC(=Y)NR^5R^6$, or $-(CH_2)_nN(R^5)C(=Y)NR^5R^6$, (wherein each methylene group may be substituted by one or more halogen atoms), each of which may be optionally substituted at any available position by one or more substituents R^a ;

R^4 represents aryl, heteroaryl or $C(=Y)R^5$ each of which may be optionally substituted at any available position by one or more substituents R^a ;

with the proviso that when m is equal to 2 or 3, then R^4 can not be phenyl substituted with substituents selected from $-OH$, $-OC_{1-4}$ alkyl, $-NH_2$, aminoacyl, $-CH_2-NH_2$ and aminoacylalkyl;

Z represents C_{2-12} alkenyl, C_{2-12} alkynyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-20} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_n-NCS$, $-C(=Y)R^5$, $-C(=Y)OR^6$, $-C(=Y)NR^5R^6$, $-OC(=Y)OR^5$, $-(CH_2)_nOP(=O)R^5R^6$, $-(CH_2)_nNHP(=O)R^5R^6$, $-(CH_2)_nOC(=Y)OR^5$, $-(CH_2)_nC(=Y)R^5$, $-(CH_2)_nC(=Y)NR^5R^6$ or $-(CH_2)_nC(=Y)OR^5$ (wherein each methylene group may be substituted by one or more halogen atoms), each of which may be optionally substituted at any available position by one or more substituents R^a ;

T , U , V and W are same or different and independently represent hydrogen or halogen;

R^5 and R^6 are same or different and are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, aryloxy, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, each of which may be optionally substituted at any available position with halogen, hydroxyl, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} alkoxy, C_{1-12} alkylcarbonyl, C_{1-12} alkoxycarbonyl, C_{3-8} cycloalkyl, C_{1-12} haloalkyl, C_{1-12} haloalkoxy, C_{2-12} haloalkenyl, aryl, heterocyclyl, heteroaryl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-CN$, $-OR^7$, $-NO_2$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-C(=Y)OR^7$, $-OC(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-NR^7SO_2R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, or -

REPLACED SHEET

$P(=O)R^7R^8$; or R^5 and R^6 may be joined together along with the heteroatom to which they are joined to form a heterocyclic or heteroaryl ring which may additionally contain from one to three heteroatoms independently selected from O, S or N, the ring formed may optionally be substituted with one or more substituents selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, C_1-C_{12} alkoxy, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-OR^7$, $-CF_3$, $-OCF_3$, $-CH_2CF_3$, $-CF_2CF_3$, $-NO_2$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-OC(=Y)OR^7$, $-C(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-NR^7SO_2R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, or $-P(O)R^7R^8$; the ring thus formed may further be fused with 3 to 7 membered unsaturated or saturated ring, which may contain from one to three heteroatoms independently selected from O, S or N, the fused ring may optionally be substituted at any available position by one or more substituents R^8 ;

R^8 is independently selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, oxo, C_{1-12} alkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{1-12} haloalkoxy, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-YR^7$, $-(CH_2)_nYR^7$, $-NO_2$, $=NOR^7$, $-NR^7R^8$, $-N(R^7)C(=Y)R^8$, $-N(R^7)C(=Y)OR^8$, $-N(R^7)C(=Y)NR^7R^8$, $-C(=Y)R^7$, $-C(=Y)NR^7R^8$, $-OC(=Y)R^7$, $-OC(=Y)NR^7R^8$, $-C(=Y)OR^7$, $-OC(=Y)OR^7$, $-SR^7$, $-S(O)_dR^7$, $-SO_2NR^7R^8$, $-OP(=O)R^7R^8$, $-NHP(=O)R^7R^8$, $-P(O)R^7R^8$, $-(CH_2)_nCN$, $-YR^7$, $-(CH_2)_nYR^7$, $-NO_2$, $=NOR^7$, $-(CH_2)_nNR^7R^8$, $-(CH_2)_nN(R^7)C(=Y)R^8$, $-(CH_2)_nN(R^7)C(=Y)OR^8$, $-(CH_2)_nN(R^7)C(=Y)NR^7R^8$, $-(CH_2)_nC(=Y)R^7$, $-(CH_2)_nC(=Y)NR^7R^8$, $-(CH_2)_nOC(=Y)R^7$, $-(CH_2)_nOC(=Y)NR^7R^8$, $-(CH_2)_nC(=Y)OR^7$, $-(CH_2)_nOC(=Y)OR^7$, $-(CH_2)_nSR^7$, $-(CH_2)_nS(O)_dR^7$, $-(CH_2)_nSO_2NR^7R^8$, $-(CH_2)_nOP(=O)R^7R^8$, $-(CH_2)_nNHP(=O)R^7R^8$, or $-(CH_2)_nP(O)R^7R^8$; each of which may optionally be substituted at any available position by one or more substituents selected from hydrogen, halogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{2-12} haloalkynyl, oxo, C_{1-12} alkoxy, C_{1-12} haloalkoxy, C_1-C_6 alkoxy C_1-C_6 alkyl, C_1-C_6 alkoxy C_1-C_6 alkoxy C_1-C_3 alkyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(CH_2)_n$ -cycloalkyl, $-(CH_2)_n$ -heterocyclyl, $-(CH_2)_n$ -aryl, $-(CH_2)_n$ -heteroaryl, $-C_{1-12}$ alkylcarbonyl, $-C_{1-12}$ alkoxy carbonyl, $-CN$, $-OR^9$,

REPLACED SHEET

$(\text{CH}_2)_n\text{OR}^9$, $-\text{CF}_3$, $-\text{NO}_2$, $-\text{NR}^9\text{R}^{10}$, $-\text{N}(\text{R}^9)\text{C}(=\text{Y})\text{R}^{10}$, $-\text{N}(\text{R}^9)\text{C}(=\text{Y})\text{OR}^{10}$, $-\text{N}(\text{R}^9)\text{C}(=\text{Y})\text{NR}^9\text{R}^{10}$, $-\text{C}(=\text{Y})\text{R}^9$, $-\text{C}(=\text{Y})\text{NR}^9\text{R}^{10}$, $-\text{OC}(=\text{Y})\text{R}^9$, $-\text{OC}(=\text{Y})\text{NR}^9\text{R}^{10}$, $-\text{OC}(=\text{Y})\text{OR}^9$, $-\text{C}(=\text{Y})\text{OR}^9$, $-\text{SR}^9$, $-\text{S}(\text{O})_d\text{R}^9$, $-\text{SO}_2\text{NR}^9\text{R}^{10}$, $-\text{NR}^9\text{SO}_2\text{R}^{10}$, $-\text{OP}(=\text{O})\text{R}^9\text{R}^{10}$, $-\text{NHP}(=\text{O})\text{R}^9\text{R}^{10}$, $-\text{P}(\text{O})\text{R}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{CN}$, $-\text{OR}^9$, $-(\text{CH}_2)_n\text{OR}^9$, $-\text{CF}_3$, $-\text{NO}_2$, $-(\text{CH}_2)_n\text{NR}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{N}(\text{R}^9)\text{C}(=\text{Y})\text{R}^{10}$, $-(\text{CH}_2)_n\text{N}(\text{R}^9)\text{C}(=\text{Y})\text{OR}^{10}$, $-(\text{CH}_2)_n\text{N}(\text{R}^9)\text{C}(=\text{Y})\text{NR}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{C}(=\text{Y})\text{R}^9$, $-(\text{CH}_2)_n\text{C}(=\text{Y})\text{NR}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{OC}(=\text{Y})\text{R}^9$, $-(\text{CH}_2)_n\text{OC}(=\text{Y})\text{NR}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{OC}(=\text{Y})\text{OR}^9$, $-(\text{CH}_2)_n\text{C}(=\text{Y})\text{OR}^9$, $-(\text{CH}_2)_n\text{SR}^9$, $-(\text{CH}_2)_n\text{S}(\text{O})_d\text{R}^9$, $-(\text{CH}_2)_n\text{SO}_2\text{NR}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{NR}^9\text{SO}_2\text{R}^{10}$, $-(\text{CH}_2)_n\text{OP}(=\text{O})\text{R}^9\text{R}^{10}$, $-(\text{CH}_2)_n\text{NHP}(=\text{O})\text{R}^9\text{R}^{10}$, or $-(\text{CH}_2)_n\text{P}(\text{O})\text{R}^9\text{R}^{10}$;

R^7 and R^8 are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(\text{CH}_2)_n$ -cycloalkyl, $-(\text{CH}_2)_n$ -heterocyclyl, $-(\text{CH}_2)_n$ -aryl, or $-(\text{CH}_2)_n$ -heteroaryl, each of which may be optionally substituted with halogen, hydroxyl or C_{1-6} alkoxy, or R^7 and R^8 may be joined together along with the heteroatom to which they are attached to form a heterocyclic or heteroaryl ring which may contain from one to three heteroatoms independently selected from O, S or N, each of which may be optionally substituted with halogen, hydroxyl, C_{1-6} alkyl or C_{1-6} alkoxy.

R^9 and R^{10} are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{1-12} haloalkyl, C_{2-12} haloalkenyl, C_{3-8} cycloalkyl, heterocyclyl, aryl, heteroaryl, $-(\text{CH}_2)_n$ -cycloalkyl, $-(\text{CH}_2)_n$ -heterocyclyl, $-(\text{CH}_2)_n$ -aryl, or $-(\text{CH}_2)_n$ -heteroaryl, each of which may be optionally substituted with halogen, hydroxyl or C_{1-6} alkoxy, or R^9 and R^{10} may be joined together along with the heteroatom to which they are attached to form a heterocyclic or heteroaryl ring which may contain from one to three heteroatoms independently selected from O, S or N, each of which may be optionally substituted with halogen, hydroxyl, C_{1-6} alkyl or C_{1-6} alkoxy.

m is 1, 2, 3 or 4;


m' is 0, 1, 2, 3 or 4;

n is 1, 2, 3 or 4;

d is 1 or 2.

2. The compound according to claim 1 having the Formula Ia, wherein,


REPLACED SHEET



Formula Ia

R^1 , R^2 , R^3 , R^4 , Y , Y' , A , T , U , V , W , Z , m and m' are as defined in claim 1; its pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof.

3. The compound according to claim 1 having the Formula Ib, wherein,




Formula Ib


R^1 , R^2 , R^3 , R^4 , T , U , V , W , Z , m , m' are as defined in claim 1; its pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers including *R* and *S* isomers, polymorphs, prodrugs, metabolites, salts or solvates thereof.

4. The compound according to claim 1, wherein R^1 and R^2 are independently selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl, C_{2-12} alkynyl, C_{3-8} cycloalkyl or aryl, each of which may optionally be substituted at any available position by one or more substituents R^a as defined in claim 1 or R^1 and R^2 together with the carbon atom to which they are attached form a 3 to 10 membered monocyclic ring, partially unsaturated or saturated, which may contain from one to three heteroatoms independently selected from O, S or N, the ring thus formed may be fused with one or two rings independently selected from the group comprising an aryl ring, a cycloalkyl ring, a heterocyclyl ring or monocyclic heteroaryl ring; the ring thus formed is optionally substituted at any available position by one or more substituents R^a as defined in claim 1.
5. The compound according to claim 1, wherein R^3 is selected from hydrogen, C_{1-12} alkyl, C_{2-12} alkenyl or C_{2-12} alkynyl, each of which may be optionally substituted at any available position by one or more substituents R^a as defined in claim 1.
6. The compound according to claim 1, wherein R^4 is selected from

REPLACED SHEET




7. The compound according to claim 1, wherein T and W independently represent fluorine and U and V both represent hydrogen.
8. The compound according to claim 1, wherein m is selected from 1 or 2 and m' is 0.
9. The compound according to claim 1, wherein Z represents -CH₂-triazole, which may be optionally substituted at any available position by one or more substituents R^a as defined in claim 1.
10. The compound according to claim 9, wherein Z is selected from:




- 11. A compound which is selected from the group comprising of:**



REPLACED SHEET




REPLACED SHEET



REPLACED SHEET



12. A pharmaceutical composition comprising a therapeutically effective amount of a compound of Formula I according to claim 1 or its pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof.
13. A method for the prophylaxis, amelioration and/or treatment of bacterial infections in a subject in need thereof, that comprises administering a therapeutically effective amount of compound of Formula I according to claim 1.
14. A method according to claim 13, wherein the bacterial infection is caused by multi drug resistant species of *Staphylococcus*, *Streptococcus*, *Enterococcus*, *Bacterioides*, *Clostridia*, *H. influenza*, *Moraxella*, acid-fast organisms such as *Mycobacterium tuberculosis* as well as Linezolid resistant species of *Staphylococcus* and *Enterococcus*.
15. Use of a compound of Formula I according to claim 1 or its pharmaceutically acceptable derivatives, tautomeric forms, stereoisomers, polymorphs, prodrugs, metabolites, salts or solvates thereof, for the manufacture of a medicament for the prophylaxis, amelioration and/or treatment of bacterial infections in a subject in need thereof.
16. Use according to claim 15, wherein the compound is in combination with other therapeutic agents.
17. Use according to claim 15 and 16, wherein the medicament is administered via oral, buccal, pulmonary, topical, subcutaneous, intramuscular, intravenous, transdermal, ocular (ophthalmic), by inhalation, intranasal, transmucosal, implant or rectal route.
18. The compounds of Formula I, processes, methods and compositions as described and illustrated herein.



Formula I