

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
16 June 2005 (16.06.2005)

PCT

(10) International Publication Number
WO 2005/054286 A2

(51) International Patent Classification⁷: **C07K 14/54**,
14/765

(21) International Application Number:
PCT/GB2004/005091

(22) International Filing Date: 3 December 2004 (03.12.2004)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
03027770.1 3 December 2003 (03.12.2003) EP

(71) Applicant (for all designated States except US): **DELTA BIOTECHNOLOGY LIMITED** [GB/GB]; Castle Court, 59 Castle Boulevard, Nottingham NG7 1FD (GB).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **KRÖZ, M.** [DE/DE]; ZLB Behring GmbH, P.O. Box 1230, D-35002 Marburg (DE). **DICKNEITE, G.** [DE/DE]; ZLB Behring GmbH, P.O. Box 1230, D-35002 Marburg (DE). **HAUSER, Hans-Peter** [DE/DE]; Feuerdornweg 6, D-35041 Marburg

(DE). **WEIMER, T.** [DE/DE]; ZLB Behring GmbH, P.O. Box 1230, D-35002 Marburg (DE). **SLEEP, Darrell** [GB/GB]; 66 Ladybay Road, West Bridgford, Nottingham NG2 5DS (GB).


(74) Agent: **BASSETT, Richard**; Eric Potter Clarkson, Park View House, 58 The Ropewalk, Nottingham NG1 5DD (GB).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI,

[Continued on next page]

(54) Title: INTERLEUKIN-11 FUSION PROTEINS



(57) Abstract: The invention relates to proteins comprising Interleukin 11 (IL-11) (including, but not limited to, fragments and variants thereof), which exhibit thrombopoietic or antiinflammatory properties, fused to albumin (including, but not limited to fragments or variants of albumin). These fusion proteins are herein collectively referred to as "albumin fusion proteins of the invention". These fusion proteins exhibit extended shelf-life and/or pharmacokinetic properties and/or extended or therapeutic activity. The invention encompasses therapeutic albumin fusion proteins, compositions, pharmaceutical compositions, formulations and kits. The invention also encompasses nucleic acid molecules encoding the albumin fusion proteins of the invention, as well as vectors containing these nucleic acids, host cells transformed with these nucleic acids and vectors, and methods of making the albumin fusion proteins of the invention using these nucleic acids, vectors, and/or host cells. The invention also relates to compositions and methods for treatment or prophylaxis of thrombocytopenia or inflammatory diseases.

WO 2005/054286 A2



FR, GB, GR, HU, IE, IS, IT, LU, MC, NL, PL, PT, RO,
SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN,
GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished
upon receipt of that report*

Interleukin-11 fusion proteins

Field of the Invention

- 5 The invention relates to novel compositions for treatment or prophylaxis of thrombocytopenia, von Willebrand disease (vWD) or inflammatory diseases, such as inflammatory bowel disease (IBD).

Background of the Invention

Physiological function of IL-11

- 10 Interleukin eleven (IL-11) is a hematopoietic cytokine that promotes megakaryocytopoiesis and thrombocytopoiesis by stimulating the proliferation of primitive stem cells, multipotent and committed progenitor cells (synergistic with other hematopoietic growth factors like IL-3, IL-6, GM-CSF) resulting in megakaryocyte maturation and increased platelet production. Due to the activity
15 described above it stimulates erythropoiesis as a side effect but shows little effect on neutrophil proliferation. It also has trophic effects on intestinal mucosa cells. Furthermore IL-11 induces the secretion of acute phase proteins (ferritin, haptoglobin, CRP, fibrinogen) by hepatocytes. It also shows antiinflammatory properties by inhibiting macrophage and T cell effector function. It inhibits the
20 production of TNF α , IL-1 β , IL-12, IL-6 and NO from activated macrophages in-vitro. Physiologically, IL-11 is produced by a variety of stromal cells, like fibroblasts, epithelial cells, chondrocytes and osteoblasts. In normal individuals it is generally undetectable in plasma. Degradation and elimination of IL-11 is yet poorly understood.

25

Biochemical characteristics of IL-11

- IL-11 is a 19 kDa polypeptide consisting of 178 amino acids (AA), plus 21 AA secretory leader sequence, which does not contain potential glycosylation residues, disulphide bonds or other posttranslational modifications, and has a close
30 similarity to IL-6. It binds to a multimeric receptor complex which contains an IL-11 specific α -receptor subunit and a promiscuous β subunit (gp130).

Recombinant human IL-11 (Oprelvekin, Neumega[®] by Genetics Institute/Wyeth) is a 2-178-interleukin-11 produced in *E. coli* lacking the amino terminal proline. This can be necessary to achieve secretion from the host cell but reportedly does not affect the activity of the cytokine.

Chemotherapy-induced thrombocytopenia

Thrombocytopenia is a significant problem for patients receiving prolonged or aggressive chemotherapy for malignancies. For some chemotherapeutic agents, such as Carboplatin, it represents the predominant, dose-limiting toxicity and acts cumulatively.

Currently, platelet transfusion is the standard treatment for thrombocytopenia. However, platelet transfusions are expensive and associated with a significant risk of alloimmunisation and transmission of blood-borne diseases. Approximately 5 to 30% of platelet transfusions are associated with usually febrile, non-hemolytic reactions. Also, platelets are a limited resource with a shelf life of 5 days only. Thus, agents that promote platelet production are an attractive alternative to platelet transfusions for the prevention or treatment of thrombocytopenia. Recombinant human IL-11 (Oprelvekin, Neumega[®]) was approved in 1997 in the US for the secondary prophylaxis of chemotherapy-induced thrombocytopenia. Further potential indications include inflammatory bowel disease (IBD), Crohn's disease, colitis ulcerosa, psoriasis and von Willebrand's disease.

Expected advantages of fusing IL11 to albumin

Safe and effective treatment of thrombocytopenia remains an unmet medical need. Recombinant human IL-11 (Oprelvekin, Neumega[®]) has a very short half-life of 6.9 hrs in humans and at the same time a narrow therapeutic window. Prolongation of plasma-half-life and increased bioavailability through albumin-fusion are expected to result in an improved safety and efficacy profile. This means that therapeutic plasma levels can be achieved and maintained with lower doses and/or longer dose intervals, avoiding peak levels above the toxic threshold.

Currently, Neumega[®] is the only drug licensed for the treatment of chemotherapy-induced thrombocytopenia, representing a field with a high unmet medical need. Neumega[®] shows a high incidence of toxic effects, such as edema and cardiovascular irregularities, combined with low efficacy especially in severe cases of thrombocytopenia.

Summary of the Invention

The invention relates to proteins comprising IL-11 fused to albumin or fragments or variants thereof. These fusion proteins are herein collectively referred to as “albumin fusion proteins of the invention.” These fusion proteins of the invention exhibit extended *in vivo* half-life and/or extended therapeutic activity as compared to unfused IL11.

The invention encompasses therapeutic albumin fusion proteins, compositions, pharmaceutical compositions, formulations and kits. The invention also encompasses nucleic acid molecules encoding the albumin fusion proteins of the invention, as well as vectors containing these nucleic acids, host cells transformed with these nucleic acids and vectors, and methods of making the albumin fusion proteins of the invention using these nucleic acids, vectors, and/or host cells.

The invention also relates to compositions and methods for therapy and prevention of thrombocytopenia. The invention further relates to compositions and methods for antiinflammatory therapy and prevention. Also, the invention relates to compositions and methods for therapy and prevention of von Willebrand's disease.

Brief Description of the Drawings

Figure 1. IL-11 plasma concentration after intravenous administration of rhIL-11 or C-terminal IL-11-albumin fusion to rabbits

- Figure 2. IL-11 plasma concentration after subcutaneous administration of rhIL-11 or C-terminal IL-11-albumin fusion to rabbits
- Figure 3. IL-11 plasma concentration after intravenous administration of rhIL-11 or N-terminal IL-11-albumin fusion to rats
- 5 Figure 4. IL-11 plasma concentration after subcutaneous administration of rhIL-11 or N-terminal IL-11-albumin fusion to rats
- Figure 5. Course of platelet levels after treatment of naive rats with IL-11
- Figure 6. Course of platelet levels after treatment of rats under chemotherapy with IL-11
- 10 Figure 7. Development of body weight (in % of baseline) after IL-11 application in a mouse model for IBD
- Figure 8. Visual observation score (diarrhoea and gross rectal bleeding) after IL-11 application in a mouse model for IBD
- Figure 9. Colon length after IL-11 application in a mouse model for IBD
- 15 Figure 10. Histological disease score after IL-11 application in a mouse model for IBD
- Figure 11 SDS gel and Western blots of various compounds of the invention.

Detailed Description of the Invention

- 20 The present invention relates to fusion proteins comprising albumin coupled to IL-11. Such peptides include, but are not limited to, peptides binding to the gp130 receptor complex. These peptides include IL-11, or fragments or variants thereof, which have thrombopoietic or antiinflammatory properties.
- 25 The terms "protein" and "peptide" as used herein are non-limiting and include proteins and polypeptides as well as peptides.

Furthermore, chemical entities may be covalently attached to the fusion proteins of the invention or used in combinations to enhance a biological activity or to
 30 modulate a biological activity.

The albumin fusion proteins of the present invention are expected to prolong the half-life of IL-11 *in vivo*. The *in vitro* or *in vivo* half-life of said albumin-fused

peptide/protein is extended 2-fold, or 5-fold, or more, over the half-life of the peptide/protein lacking the linked albumin. Furthermore, the albumin fusion proteins of the present invention are expected to reduce the frequency of the dosing schedule of the therapeutic peptide. The dosing schedule frequency is
5 reduced by at least one-quarter, or by at least one-half, or more, as compared to the frequency of the dosing schedule of the therapeutic peptide lacking the linked albumin.

The albumin fusion proteins of the present invention are expected to prolong the shelf-life of the peptide, and/or stabilize the peptide and/or its activity in solution
10 (or in a pharmaceutical composition) *in vitro* and/or *in vivo*. These albumin-fusion proteins, which may be therapeutic agents, are expected to reduce the need to formulate protein solutions with large excesses of carrier proteins (such as albumin, unfused) to prevent loss of proteins due to factors such as non-specific
15 binding. An increased half-life is defined as a half-life that is at least 2 times higher (preferably at least 5, 10, 20 or 30 times higher) than that of the unfused IL-11 compound, when measured over the first 24 hours after sub-cutaneous injection according to Example 4 below, in male Wistar rats aged 6 months.

20 The present invention also encompasses nucleic acid molecules encoding the albumin fusion proteins as well as vectors containing these nucleic acids, host cells transformed with these nucleic acids vectors, and methods of making the albumin fusion proteins of the invention using these nucleic acids, vectors, and/or host cells. The present invention further includes transgenic organisms modified
25 to contain the nucleic acid molecules of the invention, optionally modified to express the albumin fusion proteins encoded by the nucleic acid molecules.

The present invention also encompasses pharmaceutical formulations comprising an albumin fusion protein of the invention and a pharmaceutically acceptable
30 diluent or carrier. Such formulations may be in a kit or container. Such kit or container may be packaged with instructions pertaining to the extended shelf-life of the protein. Such formulations may be used in methods of preventing, treating, ameliorating thrombocytopenia or inflammatory diseases, such as inflammatory

bowel disease, psoriasis, rheumatoid arthritis, von Willebrand's disease, etc., or a related disorder in a patient, such as a mammal, or a human, comprising the step of administering the pharmaceutical formulation to the patient.

- 5 The invention also encompasses a method for potentially minimizing side effects (e.g., injection site reaction, increase of plasma volume, arrhythmia, headache, nausea, fever, rash, asthenia, diarrhoea, dizziness, allergic reactions) associated with the treatment of a mammal with cytokines in moderately higher concentrations comprising administering an albumin-fused cytokine of the
10 invention to said mammal.

The present invention encompasses a method of preventing, treating or ameliorating thrombocytopenia or inflammatory diseases, such as inflammatory
15 bowel disease, psoriasis, rheumatoid arthritis, etc., comprising administering to a mammal, in which such prevention treatment or amelioration is desired, an albumin fusion protein of the invention that comprises a IL-11 peptide/protein (or fragment or variant thereof) in an amount effective to treat, prevent or ameliorate the disease or disorder. In the present invention, the IL-11 is also called the
"therapeutic protein".

20

The present invention encompasses albumin fusion proteins comprising IL-11 (including fragments and variants thereof) fused to albumin or multiple copies of albumin (including fragments and variants thereof).

- 25 The present invention also encompasses a method for extending the half-life of IL-11 in a mammal. The method entails linking IL-11 to an albumin to form albumin-fused IL-11 and administering the albumin-fused IL-11 to a mammal. Typically, the half-life of the albumin-fused IL-11 may be extended by at least 2-fold, 5-fold, 10-fold, 20-fold, 30-fold, 40-fold or at least 50-fold over the half-life
30 of IL-11 lacking the linked albumin.

Exemplified herein are fusion proteins comprising albumin fused to IL-11. The present invention also includes an improved method of manufacturing a

therapeutic moiety as compared to what is available in the art. For example, the present invention provides an enhanced means of manufacturing a protein with the active moiety IL-11. Various aspects of the present invention are discussed in further detail below

5

Albumin

The terms human serum albumin (HSA) and human albumin (HA) are used interchangeably herein. The terms, "albumin" and "serum albumin" are broader, and encompass human serum albumin (and fragments and variants thereof) as well as albumin from other species (and fragments and variants thereof).

As used herein, "albumin" refers collectively to albumin protein or amino acid sequence, or an albumin fragment or variant, having one or more functional activities (e.g., biological activities) of albumin. In particular, "albumin" refers to human albumin or fragments thereof (see EP 201 239, EP 322 094 and WO 97/24445) especially the mature form of human albumin as shown in Table 1 and SEQ ID NO:18 of WO 03/066824 and WO 01/79480, or residues 202 to 762 of SEQ ID No: 17 herein, or albumin from other vertebrates or fragments thereof, or analogs or variants of these molecules or fragments thereof (for example the modified albumins of WO95/23857).

The human serum albumin protein used in the albumin fusion proteins of the invention may contain one or both of the following sets of point mutations with reference to SEQ ID NO:18: Leu-407 to Ala, Leu-408 to Val, Val-409 to Ala, and Arg-410 to Ala; or Arg-410 to Ala, Lys-413 to Gln, and Lys-414 to Gln (see, e.g., International Publication No. WO95/23857, hereby incorporated in its entirety by reference herein). In other embodiments, albumin fusion proteins of the invention that contain one or both of above-described sets of point mutations have improved stability/resistance to yeast Yap3p proteolytic cleavage, allowing increased production of recombinant albumin fusion proteins expressed in yeast host cells.

As used herein, a portion of albumin sufficient to prolong or extend the *in vivo* half-life, therapeutic activity, or shelf-life of the Therapeutic protein refers to a

portion of albumin sufficient in length or structure to stabilize, prolong or extend the *in vivo* half-life, therapeutic activity or shelf-life of the Therapeutic protein portion of the albumin fusion protein compared to the *in vivo* half-life, therapeutic activity, or shelf-life of the Therapeutic protein in the non-fusion state. The

5 albumin portion of the albumin fusion proteins may comprise the full length of the HA sequence as described above, or may include one or more fragments thereof that are capable of stabilizing or prolonging the therapeutic activity. Such fragments may be of 10 or more amino acids in length or may include about 15, 20, 25, 30, 50, 100, 150 or more contiguous amino acids from the HA sequence or

10 may include part or all of specific domains of HA.

The albumin portion of the albumin fusion proteins of the invention may be a variant of normal HA. The Therapeutic protein portion of the albumin fusion proteins of the invention may also be variants of the Therapeutic proteins as described herein. The term "variants" includes insertions, deletions and

15 substitutions, either conservative or non conservative, where such changes do not substantially alter one or more of the oncotic, useful ligand-binding and non-immunogenic properties of albumin, or the active site, or active domain which confers the therapeutic activities of the Therapeutic proteins.

20 In particular, the albumin fusion proteins of the invention may include naturally occurring polymorphic variants of human albumin and fragments of human albumin, for example those fragments disclosed in EP 322 094 (namely HA (1-n), where n is 369 to 419). The albumin may be derived from any vertebrate, especially any mammal, for example human, cow, sheep or pig. Non-mammalian

25 albumins include, but are not limited to, hen and salmon. The albumin portion of the albumin fusion protein may be from a different animal than the Therapeutic protein portion.

Generally speaking, an HA fragment or variant will be at least 100 amino acids

30 long, optionally at least 150 amino acids long. The HA variant may consist of or alternatively comprise at least one whole domain of HA, for example domains 1 (amino acids 1-194 of SEQ ID NO:18), 2 (amino acids 195-387 of SEQ ID NO:18), 3 (amino acids 388-585 of SEQ ID NO:18), 1 + 2 (1-387 of SEQ ID

NO:18), 2 + 3 (195-585 of SEQ ID NO:18) or 1 + 3 (amino acids 1-194 of SEQ ID NO:18 + amino acids 388-585 of SEQ ID NO:18). Each domain is itself made up of two homologous subdomains namely 1-105, 120-194, 195-291, 316-387, 388-491 and 512-585, with flexible inter-subdomain linker regions comprising
5 residues Lys106 to Glu119, Glu292 to Val315 and Glu492 to Ala511.

The albumin portion of an albumin fusion protein of the invention may comprise at least one subdomain or domain of HA or conservative modifications thereof. If the fusion is based on subdomains, some or all of the adjacent linkers may
10 optionally be used to link to the Therapeutic protein moiety.

Albumin Fusion Proteins

The present invention relates generally to albumin fusion proteins and methods of treating, preventing or ameliorating diseases or disorders. As used herein,
15 “albumin fusion protein” refers to a protein formed by the fusion of at least one molecule of albumin (or a fragment or variant thereof) to at least one molecule of a Therapeutic protein (or fragment or variant thereof). An albumin fusion protein of the invention comprises at least a fragment or variant of a Therapeutic protein and at least a fragment or variant of human serum albumin, which are associated
20 with one another, such as by genetic fusion (i.e., the albumin fusion protein is generated by translation of a nucleic acid in which a polynucleotide encoding all or a portion of a Therapeutic protein is joined in-frame with a polynucleotide encoding all or a portion of albumin) to one another. The Therapeutic protein and albumin protein, once part of the albumin fusion protein, may be referred to as a
25 “portion”, “region” or “moiety” of the albumin fusion protein.

In one embodiment, the invention provides an albumin fusion protein comprising, or alternatively consisting of, a Therapeutic protein and a serum albumin protein. In other embodiments, the invention provides an albumin fusion protein
30 comprising, or alternatively consisting of, a biologically active and/or therapeutically active fragment of a Therapeutic protein and a serum albumin protein. In other embodiments, the invention provides an albumin fusion protein comprising, or alternatively consisting of, a biologically active and/or

therapeutically active variant of a Therapeutic protein and a serum albumin protein. In further embodiments, the serum albumin protein component of the albumin fusion protein is the mature portion of serum albumin.

- 5 In further embodiments, the invention provides an albumin fusion protein comprising, or alternatively consisting of, a Therapeutic protein, and a biologically active and/or therapeutically active fragment of serum albumin. In further embodiments, the invention provides an albumin fusion protein comprising, or alternatively consisting of, a Therapeutic protein and a biologically
10 active and/or therapeutically active variant of serum albumin. In some embodiments, the Therapeutic protein portion of the albumin fusion protein is the mature portion of the Therapeutic protein.

- In further embodiments, the invention provides an albumin fusion protein
15 comprising, or alternatively consisting of, a biologically active and/or therapeutically active fragment or variant of a Therapeutic protein and a biologically active and/or therapeutically active fragment or variant of serum albumin. In some embodiments, the invention provides an albumin fusion protein comprising, or alternatively consisting of, the mature portion of a Therapeutic
20 protein and the mature portion of serum albumin.

- In one embodiment, the albumin fusion protein comprises HA as the N-terminal portion, and a Therapeutic protein as the C-terminal portion. Alternatively, an albumin fusion protein comprising HA as the C-terminal portion, and a
25 Therapeutic protein as the N-terminal portion may also be used.

- In other embodiments, the albumin fusion protein has a Therapeutic protein fused to both the N-terminus and the C-terminus of albumin. In one embodiment, the Therapeutic proteins fused at the N- and C- termini are the same Therapeutic
30 proteins. In another embodiment, the Therapeutic proteins fused at the N- and C- termini are different Therapeutic proteins. In another embodiment, the Therapeutic proteins fused at the N- and C- termini are different Therapeutic proteins which may be used to treat or prevent the same disease, disorder, or

condition. In another embodiment, the Therapeutic proteins fused at the N- and C- termini are different Therapeutic proteins which may be used to treat or prevent diseases or disorders which are known in the art to commonly occur in patients simultaneously.

5

In addition to albumin fusion protein in which the albumin portion is fused N-terminal and/or C-terminal of the Therapeutic protein portion, albumin fusion proteins of the invention may also be produced by inserting the Therapeutic protein or peptide of interest into an internal region of HA. For instance, within
10 the protein sequence of the HA molecule a number of loops or turns exist between the end and beginning of α -helices, which are stabilized by disulphide bonds. The loops, as determined from the crystal structure of HA (PDB identifiers 1AO6, 1BJ5, 1BKE, 1BM0, 1E7E to 1E7I and 1UOR) for the most part extend away from the body of the molecule. These loops are useful for the insertion, or
15 internal fusion, of therapeutically active peptides, particularly those requiring a secondary structure to be functional, or Therapeutic proteins, to essentially generate an albumin molecule with specific biological activity.

Loops in human albumin structure into which peptides or polypeptides may be
20 inserted to generate albumin fusion proteins of the invention include: Val54-Asn61, Thr76-Asp89, Ala92-Glu100, Gln170-Ala176, His247-Glu252, Glu266-Glu277, Glu280-His288, Ala362-Glu368, Lys439-Pro447, Val462-Lys475, Thr478-Pro486, and Lys560-Thr566. In other embodiments, peptides or polypeptides are inserted into the Val54-Asn61, Gln170-Ala176, and/or Lys560-
25 Thr566 loops of mature human albumin (SEQ ID NO:18).

Peptides to be inserted may be derived from either phage display or synthetic peptide libraries screened for specific biological activity or from the active portions of a molecule with the desired function. Additionally, random peptide
30 libraries may be generated within particular loops or by insertions of randomized peptides into particular loops of the HA molecule and in which all possible combinations of amino acids are represented.

Such library(s) could be generated on HA or domain fragments of HA by one of the following methods:

- 5 (a) randomized mutation of amino acids within one or more peptide loops of HA or HA domain fragments. Either one, more or all the residues within a loop could be mutated in this manner;
 - (b) replacement of, or insertion into one or more loops of HA or HA domain fragments (*i.e.*, internal fusion) of a randomized peptide(s) of length X_n (where X
10 is an amino acid and n is the number of residues;
 - (c) N-, C- or N- and C- terminal peptide/protein fusions in addition to (a) and/or (b).
- 15 The HA or HA domain fragment may also be made multifunctional by grafting the peptides derived from different screens of different loops against different targets into the same HA or HA domain fragment.

Examples of peptides inserted into a loop of human serum albumin are
20 Therapeutic protein peptides or peptide fragments or peptide variants thereof. For example, peptides inserted into a loop of human serum albumin may include T-20 and/or T-1249 peptide or peptide fragments or peptide variants thereof. More particularly, the invention encompasses albumin fusion proteins which comprise peptide fragments or peptide variants of at least 7 at least 8, at least 9, at least 10,
25 at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 35, or at least 40 amino acids in length inserted into a loop of human serum albumin. The invention also encompasses albumin fusion proteins which comprise peptide fragments or peptide variants having at least 7 at least 8,
30 at least 9, at least 10, at least 11, at least 12, at least 13, at least 14, at least 15, at least 20, at least 25, at least 30, at least 35, or at least 40 amino acids fused to the N-terminus of human serum albumin. The invention also encompasses albumin fusion proteins which comprise peptide fragments or peptide variants having at least 7, at least 8, at least 9, at least 10, at least 11, at least 12, at least 13, at least

14, at least 15, at least 20, at least 25, at least 30, at least 35, or at least 40 amino acids fused to the C-terminus of human serum albumin.

Generally, the albumin fusion proteins of the invention may have one HA-derived
5 region and one Therapeutic protein-derived region. Multiple regions of each protein, however, may be used to make an albumin fusion protein of the invention. Similarly, more than one Therapeutic protein may be used to make an albumin fusion protein of the invention. For instance, a Therapeutic protein may be fused to both the N- and C-terminal ends of the HA. In such a configuration, the
10 Therapeutic protein portions may be the same or may be different Therapeutic protein molecules. The structure of bifunctional albumin fusion proteins may be represented as: X-HA-Y or Y-HA-X or X-Y-HA or HA-X-Y or HA-Y-X-HA or HA-X-X-HA or HA Y-Y-HA or HA-X-HA-Y or X-HA-Y-HA or multiple combinations and/or inserting X and/or Y within the HA sequence at any location.

15 Bi- or multi-functional albumin fusion proteins may be prepared in various ratios depending on function, half-life etc. Bi- or multi-functional albumin fusion proteins may also be prepared to target the Therapeutic protein portion of a fusion to a target organ or cell type via protein or peptide at the opposite terminus of HA.

20 As an alternative to the fusion of known therapeutic molecules, the peptides may be obtained by screening libraries constructed as fusions to the N-, C- or N- and C- termini of HA, or domain fragment of HA, of typically 6, 8, 12, 20 or 25 or X_n (where X is an amino acid (aa) and n equals the number of residues) randomized
25 amino acids, and in which all possible combinations of amino acids are represented. A particular advantage of this approach is that the peptides may be selected *in situ* on the HA molecule and the properties of the peptide would therefore be as selected for rather than, potentially, modified as might be the case for a peptide derived by any other method then being attached to HA.

30 Additionally, the albumin fusion proteins of the invention may include a linker peptide between the fused portions to provide greater physical separation between the moieties and thus maximize the accessibility of the Therapeutic protein

portion, for instance, for binding to its cognate receptor. The linker peptide may consist of amino acids such that it is flexible or more rigid.

Therefore, as described above, the albumin fusion proteins of the invention may have the following formula R2-R1; R1-R2; R2-R1-R2; R2-L-R1-L-R2; R1-L-R2; R2-L-R1; or R1-L-R2-L-R1, wherein R1 is at least one Therapeutic protein, peptide or polypeptide sequence (including fragments or variants thereof), and not necessarily the same Therapeutic protein, L is a linker and R2 is a serum albumin sequence (including fragments or variants thereof). Exemplary linkers include (GGGGS)_N (SEQ ID NO:8) or (GGGS)_N (SEQ ID NO:9) or (GGS)_N, wherein N is an integer greater than or equal to 1 and wherein G represents glycine and S represents serine. When R1 is two or more Therapeutic proteins, peptides or polypeptide sequence, these sequences may optionally be connected by a linker.

In further embodiments, albumin fusion proteins of the invention comprising a Therapeutic protein have extended shelf-life or *in vivo* half-life or therapeutic activity compared to the shelf-life or *in vivo* half-life or therapeutic activity of the same Therapeutic protein when not fused to albumin. Shelf-life typically refers to the time period over which the therapeutic activity of a Therapeutic protein in solution or in some other storage formulation, is stable without undue loss of therapeutic activity. Many of the Therapeutic proteins are highly labile in their unfused state. As described below, the typical shelf-life of these Therapeutic proteins is markedly prolonged upon incorporation into the albumin fusion protein of the invention.

Albumin fusion proteins of the invention with "prolonged" or "extended" shelf-life exhibit greater therapeutic activity relative to a standard that has been subjected to the same storage and handling conditions. The standard may be the unfused full-length Therapeutic protein. When the Therapeutic protein portion of the albumin fusion protein is an analogue, a variant, or is otherwise altered or does not include the complete sequence for that protein, the prolongation of therapeutic activity may alternatively be compared to the unfused equivalent of that analogue, variant, altered peptide or incomplete sequence. As an example, an albumin

fusion protein of the invention may retain greater than about 100% of the therapeutic activity, or greater than about 105%, 110%, 120%, 130%, 150% or 200% of the therapeutic activity of a standard when subjected to the same storage and handling conditions as the standard when compared at a given time point.

- 5 However, it is noted that the therapeutic activity depends on the Therapeutic protein's stability, and may be below 100%.

Shelf-life may also be assessed in terms of therapeutic activity remaining after storage, normalized to therapeutic activity when storage began. Albumin fusion
10 proteins of the invention with prolonged or extended shelf-life as exhibited by prolonged or extended therapeutic activity may retain greater than about 50% of the therapeutic activity, about 60%, 70%, 80%, or 90% or more of the therapeutic activity of the equivalent unfused Therapeutic protein when subjected to the same conditions.

15

Therapeutic proteins

As stated above, an albumin fusion protein of the invention comprises at least a fragment or variant of a therapeutic protein and at least a fragment or variant of human serum albumin, which are associated with one another by genetic fusion.

20

As used herein, "Therapeutic protein" refers to IL-11, or fragments or variants thereof, having one or more therapeutic and/or biological activities. Thus an albumin fusion protein of the invention may contain at least a fragment or variant of a Therapeutic protein. Additionally, the term "therapeutic protein" may refer
25 to the endogenous or naturally occurring correlate of a therapeutic protein. Variants include mutants, analogs, and mimetics, as well as homologues, including the endogenous or naturally occurring correlates of a therapeutic protein.

30 By a polypeptide displaying a "therapeutic activity" or a protein that is "therapeutically active" is meant a polypeptide that possesses one or more known biological and/or therapeutic activities associated with a therapeutic protein such as one or more of the therapeutic proteins described herein or otherwise known in

the art. As a non-limiting example, a “therapeutic protein” is a protein that is useful to treat, prevent or ameliorate a disease, condition or disorder.

As used herein, “therapeutic activity” or “activity” may refer to an activity whose effect is consistent with a desirable therapeutic outcome in humans, or to desired effects in non-human mammals or in other species or organisms. Therapeutic activity may be measured *in vivo* or *in vitro*. For example, a desirable effect may be assayed in cell culture. Such *in vitro* or cell culture assays are commonly available for many Therapeutic proteins as described in the art.

10

Therapeutic proteins corresponding to a therapeutic protein portion of an albumin fusion protein of the invention may be modified by the attachment of one or more oligosaccharide groups. The modification, referred to as glycosylation, can dramatically affect the physical properties of proteins and can be important in protein stability, secretion, and localization. Such modifications are described in detail in WO 03/066824 and WO 01/79480, which are incorporated herein by reference

Therapeutic proteins corresponding to a Therapeutic protein portion of an albumin fusion protein of the invention, as well as analogs and variants thereof, may be modified so that glycosylation at one or more sites is altered as a result of manipulation(s) of their nucleic acid sequence, by the host cell in which they are expressed, or due to other conditions of their expression. For example, glycosylation isomers may be produced by abolishing or introducing glycosylation sites, *e.g.*, by substitution or deletion of amino acid residues, such as substitution of glutamine for asparagine, or unglycosylated recombinant proteins may be produced by expressing the proteins in host cells that will not glycosylate them, *e.g.* in *E. coli* or glycosylation-deficient yeast. Examples of these approaches are described in more detail in WO 03/066824 and WO 01/79480, which are incorporated by reference, and are known in the art.

20
25
30

In various embodiments, the albumin fusion proteins of the invention are capable of a therapeutic activity and/or biologic activity corresponding to the therapeutic

activity and/or biologic activity of the Therapeutic protein corresponding to the Therapeutic protein portion of the albumin fusion. In further embodiments, the therapeutically active protein portions of the albumin fusion proteins of the invention are fragments or variants of the reference sequence and are capable of the therapeutic activity and/or biologic activity of the corresponding Therapeutic protein.

Polypeptide and Polynucleotide Fragments and Variants

Fragments

The present invention is further directed to fragments of the Therapeutic proteins, albumin proteins, and/or albumin fusion proteins of the invention. Even if deletion of one or more amino acids from the N-terminus of a protein results in modification or loss of one or more biological functions of the Therapeutic protein, albumin protein, and/or albumin fusion protein, other Therapeutic activities and/or functional activities (e.g., biological activities, ability to multimerize, ability to bind a ligand) may still be retained. For example, the ability of polypeptides with N-terminal deletions to induce and/or bind to antibodies which recognize the complete or mature forms of the polypeptides generally will be retained when less than the majority of the residues of the complete polypeptide are removed from the N-terminus. Whether a particular polypeptide lacking N-terminal residues of a complete polypeptide retains such immunologic activities can readily be determined by routine methods described herein and otherwise known in the art. It is not unlikely that a mutein with a large number of deleted N-terminal amino acid residues may retain some biological or immunogenic activities. In fact, peptides composed of as few as six amino acid residues may often evoke an immune response.

Accordingly, fragments of a Therapeutic protein corresponding to a Therapeutic protein portion of an albumin fusion protein of the invention, include the full length protein as well as polypeptides having one or more residues deleted from the amino terminus of the amino acid sequence of the reference polypeptide. Polynucleotides encoding these polypeptides are also encompassed by the invention.

In addition, fragments of serum albumin polypeptides corresponding to an albumin protein portion of an albumin fusion protein of the invention, include the full length protein as well as polypeptides having one or more residues deleted
5 from the amino terminus of the amino acid sequence of the reference polypeptide (i.e., serum albumin). Polynucleotides encoding these polypeptides are also encompassed by the invention.

Moreover, fragments of albumin fusion proteins of the invention include the full
10 length albumin fusion protein as well as polypeptides having one or more residues deleted from the amino terminus of the albumin fusion protein. Polynucleotides encoding these polypeptides are also encompassed by the invention.

The present invention further provides polypeptides having one or more residues
15 deleted from the carboxy terminus of the amino acid sequence of a Therapeutic protein corresponding to a Therapeutic protein portion of an albumin fusion protein of the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

20 In addition, the present invention provides polypeptides having one or more residues deleted from the carboxy terminus of the amino acid sequence of an albumin protein corresponding to an albumin protein portion of an albumin fusion protein of the invention (e.g., serum albumin). Polynucleotides encoding these polypeptides are also encompassed by the invention.

25 Moreover, the present invention provides polypeptides having one or more residues deleted from the carboxy terminus of an albumin fusion protein of the invention. Polynucleotides encoding these polypeptides are also encompassed by the invention.

30 In addition, any of the above described N- or C-terminal deletions can be combined to produce a N- and C-terminal deleted reference polypeptide (e.g., a Therapeutic protein referred to in Table 1, or serum albumin (e.g., SEQ ID

NO:18), or an albumin fusion protein of the invention). The invention also provides polypeptides having one or more amino acids deleted from both the amino and the carboxyl termini. Polynucleotides encoding these polypeptides are also encompassed by the invention.

5

The present application is also directed to proteins containing polypeptides at least 60%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to a reference polypeptide sequence (e.g., a Therapeutic protein, serum albumin protein or an albumin fusion protein of the invention) set forth herein, or fragments thereof. In

10 some embodiments, the application is directed to proteins comprising polypeptides at least 60%, 80%, 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to reference polypeptides having the amino acid sequence of N- and C-terminal deletions as described above. Polynucleotides encoding these polypeptides are also encompassed by the invention.

15

Other polypeptide fragments of the invention are fragments comprising, or alternatively, consisting of, an amino acid sequence that displays a Therapeutic activity and/or functional activity (e.g. biological activity) of the polypeptide sequence of the Therapeutic protein or serum albumin protein of which the amino

20 acid sequence is a fragment.

Other polypeptide fragments are biologically active fragments. Biologically active fragments are those exhibiting activity similar, but not necessarily identical, to an activity of the polypeptide of the present invention. The biological activity

25 of the fragments may include an improved desired activity, or a decreased undesirable activity.

Variants

“Variant” refers to a polynucleotide or nucleic acid differing from a reference nucleic acid or polypeptide, but retaining essential properties thereof. Generally, variants are overall closely similar, and, in many regions, identical to the reference
5 nucleic acid or polypeptide.

As used herein, “variant” refers to a Therapeutic protein portion of an albumin fusion protein of the invention, an albumin portion of an albumin fusion protein of the invention, or an albumin fusion protein differing in sequence from a
10 Therapeutic protein, albumin protein, and/or albumin fusion protein of the invention, respectively, but retaining at least one functional and/or therapeutic property thereof as described elsewhere herein or otherwise known in the art. Generally, variants are overall very similar, and, in many regions, identical to the amino acid sequence of the Therapeutic protein corresponding to a Therapeutic
15 protein portion of an albumin fusion protein of the invention, albumin protein corresponding to an albumin protein portion of an albumin fusion protein of the invention, and/or albumin fusion protein of the invention. Nucleic acids encoding these variants are also encompassed by the invention.

20 The present invention is also directed to proteins which comprise, or alternatively consist of, an amino acid sequence which is at least 60%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, 99% or 100%, identical to, for example, the amino acid sequence of a Therapeutic protein corresponding to a Therapeutic protein portion of an albumin fusion protein of the invention, albumin proteins (e.g., SEQ ID NO:18 or
25 fragments or variants thereof) corresponding to an albumin protein portion of an albumin fusion protein of the invention, and/or albumin fusion proteins of the invention. Fragments of these polypeptides are also provided (e.g., those fragments described herein). Further polypeptides encompassed by the invention are polypeptides encoded by polynucleotides which hybridize to the complement
30 of a nucleic acid molecule encoding an amino acid sequence of the invention under stringent hybridization conditions (e.g., hybridization to filter bound DNA in 6X Sodium chloride/Sodium citrate (SSC) at about 45 degrees Celsius, followed by one or more washes in 0.2X SSC, 0.1% SDS at about 50 - 65 degrees

- Celsius), under highly stringent conditions (e.g., hybridization to filter bound DNA in 6X sodium chloride/Sodium citrate (SSC) at about 45 degrees Celsius, followed by one or more washes in 0.1X SSC, 0.2% SDS at about 68 degrees Celsius), or under other stringent hybridization conditions which are known to those of skill in the art (see, for example, Ausubel, F.M. et al., eds., 1989 *Current protocol in Molecular Biology*, Green publishing associates, Inc., and John Wiley & Sons Inc., New York, at pages 6.3.1 - 6.3.6 and 2.10.3). Polynucleotides encoding these polypeptides are also encompassed by the invention.
- 10 By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a query amino acid sequence of the present invention, it is intended that the amino acid sequence of the subject polypeptide is identical to the query sequence except that the subject polypeptide sequence may include up to five amino acid alterations per each 100 amino acids of the query amino acid sequence.
- 15 In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a query amino acid sequence, up to 5% of the amino acid residues in the subject sequence may be inserted, deleted, or substituted with another amino acid. These alterations of the reference sequence may occur at the amino- or carboxy-terminal positions of the reference amino acid sequence or
- 20 anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.
- As a practical matter, whether any particular polypeptide is at least 60%, 80%, 25 85%, 90%, 95%, 96%, 97%, 98% or 99% identical to, for instance, the amino acid sequence of an albumin fusion protein of the invention or a fragment thereof (such as the Therapeutic protein portion of the albumin fusion protein or the albumin portion of the albumin fusion protein), can be determined conventionally using known computer programs. Such programs and methods of using them are
- 30 described, e.g., in WO 03/066824 and WO 01/79480 (pp. 41-43), which are incorporated by reference herein.

The polynucleotide variants of the invention may contain alterations in the coding regions, non-coding regions, or both. Polynucleotide variants include those containing alterations which produce silent substitutions, additions, or deletions, but do not alter the properties or activities of the encoded polypeptide. Such
5 nucleotide variants may be produced by silent substitutions due to the degeneracy of the genetic code. Polypeptide variants include those in which less than 50, less than 40, less than 30, less than 20, less than 10, or 5-50, 5-25, 5-10, 1-5, or 1-2 amino acids are substituted, deleted, or added in any combination. Polynucleotide variants can be produced for a variety of reasons, e.g., to optimize
10 codon expression for a particular host (change codons in the human mRNA to those preferred by a microbial host, such as, yeast or *E. coli*).

In another embodiment, a polynucleotide encoding an albumin portion of an albumin fusion protein of the invention is optimized for expression in yeast or
15 mammalian cells. In a further embodiment, a polynucleotide encoding a Therapeutic protein portion of an albumin fusion protein of the invention is optimized for expression in yeast or mammalian cells. In a still further embodiment, a polynucleotide encoding an albumin fusion protein of the invention is optimized for expression in yeast or mammalian cells.

20 In an alternative embodiment, a codon optimized polynucleotide encoding a Therapeutic protein portion of an albumin fusion protein of the invention does not hybridize to the wild type polynucleotide encoding the Therapeutic protein under stringent hybridization conditions as described herein. In a further embodiment, a
25 codon optimized polynucleotide encoding an albumin portion of an albumin fusion protein of the invention does not hybridize to the wild type polynucleotide encoding the albumin protein under stringent hybridization conditions as described herein. In another embodiment, a codon optimized polynucleotide encoding an albumin fusion protein of the invention does not hybridize to the wild
30 type polynucleotide encoding the Therapeutic protein portion or the albumin protein portion under stringent hybridization conditions as described herein.

In an additional embodiment, polynucleotides encoding a Therapeutic protein portion of an albumin fusion protein of the invention do not comprise, or alternatively consist of, the naturally occurring sequence of that Therapeutic protein. In a further embodiment, polynucleotides encoding an albumin protein
5 portion of an albumin fusion protein of the invention do not comprise, or alternatively consist of, the naturally occurring sequence of albumin protein. In an alternative embodiment, polynucleotides encoding an albumin fusion protein of the invention do not comprise, or alternatively consist of, the naturally occurring sequence of a Therapeutic protein portion or the albumin protein portion.

10

In an additional embodiment, the Therapeutic protein may be selected from a random peptide library by biopanning, as there will be no naturally occurring wild type polynucleotide.

15 Naturally occurring variants are called "allelic variants," and refer to one of several alternative forms of a gene occupying a given locus on a chromosome of an organism. (Genes II, Lewin, B., ed., John Wiley & Sons, New York (1985)). These allelic variants can vary at either the polynucleotide and/or polypeptide level and are included in the present invention. Alternatively, non-naturally
20 occurring variants may be produced by mutagenesis techniques or by direct synthesis.

Using known methods of protein engineering and recombinant DNA technology, variants may be generated to improve or alter the characteristics of the
25 polypeptides of the present invention. For instance, one or more amino acids may be deleted from the N-terminus or C-terminus of the polypeptide of the present invention without substantial loss of biological function. See, e.g., Ron *et al.*, *J. Biol. Chem.* 268: 2984-2988 (1993) (KGF variants) and Dobeli *et al.*, *J. Biotechnology* 7:199-216 (1988) (interferon gamma variants).

30

Moreover, ample evidence demonstrates that variants often retain a biological activity similar to that of the naturally occurring protein (e.g. Gayle and co-workers (*J. Biol. Chem.* 268:22105-22111 (1993) (IL-1a variants)).

Furthermore, even if deleting one or more amino acids from the N-terminus or C-terminus of a polypeptide results in modification or loss of one or more biological functions, other biological activities may still be retained. For example, the ability of a deletion variant to induce and/or to bind antibodies which recognize the secreted form will likely be retained when less than the majority of the residues of the secreted form are removed from the N-terminus or C-terminus. Whether a particular polypeptide lacking N- or C-terminal residues of a protein retains such immunogenic activities can readily be determined by routine methods described herein and otherwise known in the art.

Thus, the invention further includes polypeptide variants which have a functional activity (e.g., biological activity and/or therapeutic activity). In further embodiments the invention provides variants of albumin fusion proteins that have a functional activity (e.g., biological activity and/or therapeutic activity, such as that disclosed in the "Biological Activity" column in Table 1) that corresponds to one or more biological and/or therapeutic activities of the Therapeutic protein corresponding to the Therapeutic protein portion of the albumin fusion protein. Such variants include deletions, insertions, inversions, repeats, and substitutions selected according to general rules known in the art so as to have little effect on activity.

In other embodiments, the variants of the invention have conservative substitutions. By "conservative substitutions" is intended swaps within groups such as replacement of the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

Guidance concerning how to make phenotypically silent amino acid substitutions is provided, for example, in Bowie *et al.*, "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310

(1990), wherein the authors indicate that there are two main strategies for studying the tolerance of an amino acid sequence to change.

As the authors state, proteins are surprisingly tolerant of amino acid substitutions.

5 The authors further indicate which amino acid changes are likely to be permissive at certain amino acid positions in the protein. For example, most buried (within the tertiary structure of the protein) amino acid residues require nonpolar side chains, whereas few features of surface side chains are generally conserved. Moreover, tolerated conservative amino acid substitutions involve replacement of
10 the aliphatic or hydrophobic amino acids Ala, Val, Leu and Ile; replacement of the hydroxyl residues Ser and Thr; replacement of the acidic residues Asp and Glu; replacement of the amide residues Asn and Gln, replacement of the basic residues Lys, Arg, and His; replacement of the aromatic residues Phe, Tyr, and Trp, and replacement of the small-sized amino acids Ala, Ser, Thr, Met, and Gly.

15 Besides conservative amino acid substitution, variants of the present invention include (i) polypeptides containing substitutions of one or more of the non-conserved amino acid residues, where the substituted amino acid residues may or may not be one encoded by the genetic code, or (ii) polypeptides containing
20 substitutions of one or more of the amino acid residues having a substituent group, or (iii) polypeptides which have been fused with or chemically conjugated to another compound, such as a compound to increase the stability and/or solubility of the polypeptide (for example, polyethylene glycol), (iv) polypeptide containing additional amino acids, such as, for example, an IgG Fc fusion region peptide.
25 Such variant polypeptides are deemed to be within the scope of those skilled in the art from the teachings herein.

For example, polypeptide variants containing amino acid substitutions of charged amino acids with other charged or neutral amino acids may produce proteins with
30 improved characteristics, such as less aggregation. Aggregation of pharmaceutical formulations both reduces activity and increases clearance due to the aggregate's immunogenic activity. See Pinckard *et al.*, *Clin. Exp. Immunol.*

2:331-340 (1967); Robbins *et al.*, *Diabetes* 36: 838-845 (1987); Cleland *et al.*, *Crit. Rev. Therapeutic Drug Carrier Systems* 10:307-377 (1993).

In specific embodiments, the polypeptides of the invention comprise, or
5 alternatively, consist of, fragments or variants of the amino acid sequence of a
Therapeutic protein described herein and/or human serum albumin, and/or
albumin fusion protein of the invention, wherein the fragments or variants have 1-
5, 5-10, 5-25, 5-50, 10-50 or 50-150, amino acid residue additions, substitutions,
and/or deletions when compared to the reference amino acid sequence. In certain
10 embodiments, the amino acid substitutions are conservative. Nucleic acids
encoding these polypeptides are also encompassed by the invention.

The polypeptide of the present invention can be composed of amino acids joined
to each other by peptide bonds or modified peptide bonds, i.e., peptide isosteres,
15 and may contain amino acids other than the 20 gene-encoded amino acids. The
polypeptides may be modified by either natural processes, such as post-
translational processing, or by chemical modification techniques which are well
known in the art. Such modifications are well described in basic texts and in
more detailed monographs, as well as in a voluminous research literature.
20 Modifications can occur anywhere in a polypeptide, including the peptide
backbone, the amino acid side-chains and the amino or carboxyl termini. It will
be appreciated that the same type of modification may be present in the same or
varying degrees at several sites in a given polypeptide. Also, a given polypeptide
may contain many types of modifications. Polypeptides may be branched, for
25 example, as a result of ubiquitination, and they may be cyclic, with or without
branching. Cyclic, branched, and branched cyclic polypeptides may result from
posttranslational natural processes or may be made by synthetic methods.
Modifications include acetylation, acylation, ADP-ribosylation, amidation,
covalent attachment of flavin, covalent attachment of a heme moiety, covalent
30 attachment of a nucleotide or nucleotide derivative, covalent attachment of a lipid
or lipid derivative, covalent attachment of phosphatidylinositol, cross-linking,
cyclization, disulfide bond formation, demethylation, formation of covalent cross-
links, formation of cysteine, formation of pyroglutamate, formylation, gamma-

carboxylation, glycosylation, GPI anchor formation, hydroxylation, iodination, methylation, myristylation, oxidation, pegylation, proteolytic processing, phosphorylation, prenylation, racemization, selenoylation, sulfation, transfer-RNA mediated addition of amino acids to proteins such as arginylation, and
5 ubiquitination.

Furthermore, chemical entities may be covalently attached to the albumin fusion proteins to enhance or modulate a specific functional or biological activity such as by methods disclosed in *Current Opinions in Biotechnology*, 10:324 (1999).

10

Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains,
15 and addition or deletion of an N-terminal methionine residue as a result of prokaryotic host cell expression. The albumin fusion proteins may also be modified with, e.g., but not limited to, a chemotherapeutic agent, such as a drug, and/or a detectable label, such as an enzymatic, fluorescent, isotopic and/or affinity label to allow for detection and isolation of the protein. Examples of such
20 modifications are given, e.g., in WO 03/066824 and in WO 01/79480 (pp. 105-106), which are incorporated by reference herein.

Functional activity

"A polypeptide having functional activity" refers to a polypeptide capable of
25 displaying one or more known functional activities associated with the full-length, pro-protein, and/or mature form of a Therapeutic protein. Such functional activities include, but are not limited to, biological activity, antigenicity (ability to bind (or compete with a polypeptide for binding) to an anti-polypeptide antibody), immunogenicity (ability to generate antibody which binds to a specific
30 polypeptide of the invention), ability to form multimers with polypeptides of the invention, and ability to bind to a receptor or ligand for a polypeptide.

“A polypeptide having biological activity” refers to a polypeptide exhibiting activity similar to, but not necessarily identical to, an activity of a Therapeutic protein of the present invention, including mature forms, as measured in a particular biological assay, with or without dose dependency. In the case where
5 dose dependency does exist, it need not be identical to that of the polypeptide, but rather substantially similar to the dose-dependence in a given activity as compared to the polypeptide of the present invention.

In other embodiments, an albumin fusion protein of the invention has at least one
10 biological and/or therapeutic activity associated with the Therapeutic protein (or fragment or variant thereof) when it is not fused to albumin.

The albumin fusion proteins of the invention can be assayed for functional activity (e.g., biological activity) using or routinely modifying assays known in the art, as
15 well as assays described herein. Specifically, albumin fusion proteins may be assayed for functional activity. Additionally, one of skill in the art may routinely assay fragments of a Therapeutic protein corresponding to a Therapeutic protein portion of an albumin fusion protein of the invention, for activity using assays for IL-11 activity. Further, one of skill in the art may routinely assay fragments of an
20 albumin protein corresponding to an albumin protein portion of an albumin fusion protein of the invention, for activity using assays known in the art and/or as described in the Examples section in WO 03/066824 and WO 01/79480.

In addition, assays described herein (see Examples and Table 1) and otherwise
25 known in the art may routinely be applied to measure the ability of albumin fusion proteins of the present invention and fragments, variants and derivatives thereof to elicit biological activity and/or Therapeutic activity (either *in vitro* or *in vivo*) related to either the Therapeutic protein portion and/or albumin portion of the albumin fusion protein of the present invention. Other methods will be known to
30 the skilled artisan and are within the scope of the invention.

Expression of Fusion Proteins

The albumin fusion proteins of the invention may be produced as recombinant molecules by secretion from yeast, a microorganism such as a bacterium, or a human or animal cell line. Optionally, the polypeptide is secreted from the host
5 cells.

For expression of the albumin fusion proteins exemplified herein, yeast strains disrupted in the *HSP150* gene as exemplified in WO 95/33833, or yeast strains disrupted in the *PMT1* gene as exemplified in WO 00/44772 (serving to
10 reduce/eliminate O-linked glycosylation of the albumin fusions), or yeast strains disrupted in the *YAP3* gene as exemplified in WO 95/23857 were successfully used, in combination with the yeast *PRB1* promoter, the HSA/MF α -I fusion leader sequence exemplified in WO 90/01063, the yeast *ADHI* terminator, the *LEU2* selection marker and the disintegration vector pSAC35 exemplified in US
15 5,637,504.

Other yeast strains, promoters, leader sequences, terminators, markers and vectors which are expected to be useful in the invention are described in WO 03/066824 and in WO 01/74980 (pp. 94-99), which are incorporated herein by reference.
20

The present invention also includes a cell, optionally a yeast cell transformed to express an albumin fusion protein of the invention. In addition to the transformed host cells themselves, the present invention also contemplates a culture of those cells, optionally a monoclonal (clonally homogeneous) culture, or a culture
25 derived from a monoclonal culture, in a nutrient medium. If the polypeptide is secreted, the medium will contain the polypeptide, with the cells, or without the cells if they have been filtered or centrifuged away. Many expression systems are known and may be used, including bacteria (for example *E. coli* and *Bacillus subtilis*), yeasts (for example *Saccharomyces cerevisiae*, *Kluyveromyces lactis* and
30 *Pichia pastoris*), filamentous fungi (for example *Aspergillus*), plant cells, animal cells and insect cells.

The desired protein is produced in conventional ways, for example from a coding sequence inserted in the host chromosome or on a free plasmid. The yeasts are transformed with a coding sequence for the desired protein in any of the usual ways, for example electroporation. Methods for transformation of yeast by
5 electroporation are disclosed in Becker & Guarente (1990) *Methods Enzymol.* 194, 182.

Successfully transformed cells, *i.e.*, cells that contain a DNA construct of the present invention, can be identified by well known techniques. For example, cells
10 resulting from the introduction of an expression construct can be grown to produce the desired polypeptide. Cells can be harvested and lysed and their DNA content examined for the presence of the DNA using a method such as that described by Southern (1975) *J. Mol. Biol.* 98, 503 or Berent *et al.* (1985) *Biotech.* 3, 208. Alternatively, the presence of the protein in the supernatant can be detected using
15 antibodies.

Useful yeast plasmid vectors include pRS403-406 and pRS413-416 and are generally available from Stratagene Cloning Systems, La Jolla, CA 92037, USA. Plasmids pRS403, pRS404, pRS405 and pRS406 are Yeast Integrating plasmids
20 (YIps) and incorporate the yeast selectable markers *HIS3*, *TRP1*, *LEU2* and *URA3*. Plasmids pRS413-416 are Yeast Centromere plasmids (YCps).

Vectors for making albumin fusion proteins for expression in yeast include pPPC0005, pScCHSA, pScNHSA, and pC4:HSA which were deposited on April
25 11, 2001 at the American Type Culture Collection, 10801 University Boulevard, Manassas, Virginia 20110-2209 and which are described in WO 03/066824 and WO 01/79480, which are incorporated by reference herein.

Another vector which is expected to be useful for expressing an albumin fusion
30 protein in yeast is the pSAC35 vector which is described in Sleep *et al.*, *BioTechnology* 8:42 (1990), which is hereby incorporated by reference in its entirety. The plasmid pSAC35 is of the disintegration class of vector described in US 5,637 504.

A variety of methods have been developed to operably link DNA to vectors via complementary cohesive termini. For instance, complementary homopolymer tracts can be added to the DNA segment to be inserted to the vector DNA. The vector and DNA segment are then joined by hydrogen bonding between the complementary homopolymeric tails to form recombinant DNA molecules.

Synthetic linkers containing one or more restriction sites provide an alternative method of joining the DNA segment to vectors. The DNA segment, generated by endonuclease restriction digestion, is treated with bacteriophage T4 DNA polymerase or *E. coli* DNA polymerase I, which are enzymes that remove protruding, 5'-single-stranded termini with their 3' -5'-exonucleolytic activities and fill in recessed 3'-ends with their polymerizing activities. The combination of these activities therefore generates blunt-ended DNA segments. The blunt-ended segments are then incubated with a large molar excess of linker molecules in the presence of an enzyme that is able to catalyze the ligation of blunt-ended DNA molecules, such as bacteriophage T4 DNA ligase. Thus, the products of the reaction are DNA segments carrying polymeric linker sequences at their ends. These DNA segments are then cleaved with the appropriate restriction enzyme and ligated to an expression vector that has been cleaved with an enzyme that produces termini compatible with those of the DNA segment.

Synthetic linkers containing a variety of restriction endonuclease sites are commercially available from a number of commercial sources.

A desirable way to modify the DNA in accordance with the invention, if, for example, HA variants are to be prepared, is to use the polymerase chain reaction as disclosed by Saiki *et al.* (1988) *Science* 239, 487-491. In this method the DNA to be enzymatically amplified is flanked by two specific oligonucleotide primers which themselves become incorporated into the amplified DNA. The specific primers may contain restriction endonuclease recognition sites which can be used for cloning into expression vectors using methods known in the art.

Exemplary genera of yeast contemplated to be useful in the practice of the present invention as hosts for expressing the albumin fusion proteins are *Pichia* (formerly classified as *Hansenula*), *Saccharomyces*, *Kluyveromyces*, *Aspergillus*, *Candida*, *Torulopsis*, *Torulaspora*, *Schizosaccharomyces*, *Citeromyces*, *Pachysolen*,
5 *Zygosaccharomyces*, *Debaromyces*, *Trichoderma*, *Cephalosporium*, *Humicola*, *Mucor*, *Neurospora*, *Yarrowia*, *Metschnikowia*, *Rhodospiridium*, *Leucosporidium*, *Botryosacculus*, *Sporidiobolus*, *Endomycopsis*, and the like. Genera include those selected from the group consisting of *Saccharomyces*, *Schizosaccharomyces*, *Kluyveromyces*, *Pichia* and *Torulaspora*. Examples of
10 *Saccharomyces* spp. are *S. cerevisiae*, *S. italicus* and *S. rouxii*. Examples of other species, and methods of transforming them, are described in WO 03/066824 and WO 01/79480 (pp. 97-98), which are incorporated herein by reference.

Methods for the transformation of *S. cerevisiae* are taught generally in EP 251
15 744, EP 258 067 and WO 90/01063, all of which are incorporated herein by reference. Suitable promoters for *S. cerevisiae* include those associated with the *PGKI* gene, *GAL1* or *GAL10* genes, *CYCI*, *PHO5*, *TRPI*, *ADHI*, *ADH2*, the genes for glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, triose phosphate isomerase, phosphoglucose
20 isomerase, glucokinase, alpha-mating factor pheromone (a mating factor pheromone), the *PRBI* promoter, the *GUT2* promoter, the *GPDI* promoter, and hybrid promoters involving hybrids of parts of 5' regulatory regions with parts of 5' regulatory regions of other promoters or with upstream activation sites (e.g. the promoter of EP-A-258 067).

25

Convenient regulatable promoters for use in *Schizosaccharomyces pombe* are the thiamine-repressible promoter from the *nmt* gene as described by Maundrell (1990) *J. Biol. Chem.* 265, 10857-10864 and the glucose repressible *jbpl* gene promoter as described by Hoffman & Winston (1990) *Genetics* 124, 807-816.

30

Methods of transforming *Pichia* for expression of foreign genes are taught in, for example, Cregg *et al.* (1993), and various Phillips patents (e.g. US 4 857 467, incorporated herein by reference), and *Pichia* expression kits are commercially

available from Invitrogen BV, Leek, Netherlands, and Invitrogen Corp., San Diego, California. Suitable promoters include AOX1 and AOX2. Gleeson *et al.* (1986) *J. Gen. Microbiol.* 132, 3459-3465 include information on *Hansenula* vectors and transformation, suitable promoters being MOX1 and FMD1; whilst
5 EP 361 991, Fleer *et al.* (1991) and other publications from Rhone-Poulenc Rorer teach how to express foreign proteins in *Kluyveromyces* spp.

The transcription termination signal may be the 3' flanking sequence of a eukaryotic gene which contains proper signals for transcription termination and
10 polyadenylation. Suitable 3' flanking sequences may, for example, be those of the gene naturally linked to the expression control sequence used, *i.e.* may correspond to the promoter. Alternatively, they may be different in which case the termination signal of the *S. cerevisiae ADHI* gene is optionally used.

15 The desired albumin fusion protein may be initially expressed with a secretion leader sequence, which may be any leader effective in the yeast chosen. Leaders useful in *S. cerevisiae* include that from the mating factor polypeptide (MF α) and the hybrid leaders of EP-A-387 319. Such leaders (or signals) are cleaved by the yeast before the mature albumin is released into the surrounding medium. Further
20 such leaders include those of *S. cerevisiae* invertase (SUC2) disclosed in JP 62-096086 (granted as 911036516), acid phosphatase (PH05), the pre-sequence of MF α -1, β -glucanase (BGL2) and killer toxin; *S. diastaticus* glucoamylase II; *S. carlsbergensis* α -galactosidase (MEL1); *K. lactis* killer toxin; and *Candida glucoamylase*.

25

Additional Methods of Recombinant and Synthetic Production of Albumin Fusion Proteins

The present invention includes polynucleotides encoding albumin fusion proteins
30 of this invention, as well as vectors, host cells and organisms containing these polynucleotides. The present invention also includes methods of producing albumin fusion proteins of the invention by synthetic and recombinant techniques.

The polynucleotides, vectors, host cells, and organisms may be isolated and purified by methods known in the art

- 5 A vector useful in the invention may be, for example, a phage, plasmid, cosmid, mini-chromosome, viral or retroviral vector. The vectors which can be utilized to clone and/or express polynucleotides of the invention are vectors which are capable of replicating and/or expressing the polynucleotides in the host cell in which the polynucleotides are desired to be replicated and/or expressed. In general, the polynucleotides and/or vectors can be utilized in any cell, either
- 10 eukaryotic or prokaryotic, including mammalian cells (e.g., human (e.g., HeLa), monkey (e.g., Cos), rabbit (e.g., rabbit reticulocytes), rat, hamster (e.g., CHO, NSO and baby hamster kidney cells) or mouse cells (e.g., L cells), plant cells, yeast cells, insect cells or bacterial cells (e.g., *E. coli*). See, e.g., F. Ausubel *et al.*, *Current Protocols in Molecular Biology*, Greene Publishing Associates and
- 15 Wiley-Interscience (1992) and Sambrook *et al.* (1989) for examples of appropriate vectors for various types of host cells. Note, however, that when a retroviral vector that is replication defective is used, viral propagation generally will occur only in complementing host cells.
- 20 The host cells containing these polynucleotides can be used to express large amounts of the protein useful in, for example, pharmaceuticals, diagnostic reagents, vaccines and therapeutics. The protein may be isolated and purified by methods known in the art or described herein.
- 25 The polynucleotides encoding albumin fusion proteins of the invention may be joined to a vector containing a selectable marker for propagation in a host. Generally, a plasmid vector may be introduced in a precipitate, such as a calcium phosphate precipitate, or in a complex with a charged lipid. If the vector is a virus, it may be packaged in vitro using an appropriate packaging cell line and
- 30 then transduced into host cells.

The polynucleotide insert should be operatively linked to an appropriate promoter compatible with the host cell in which the polynucleotide is to be expressed. The

promoter may be a strong promoter and/or an inducible promoter. Examples of promoters include the phage lambda PL promoter, the *E. coli lac*, *trp*, *phoA* and *tac* promoters, the SV40 early and late promoters and promoters of retroviral LTRs, to name a few. Other suitable promoters will be known to the skilled artisan. The expression constructs will further contain sites for transcription initiation, termination, and, in the transcribed region, a ribosome binding site for translation. The coding portion of the transcripts expressed by the constructs may include a translation initiating codon at the beginning and a termination codon (UAA, UGA or UAG) appropriately positioned at the end of the polypeptide to be translated.

As indicated, the expression vectors may include at least one selectable marker. Such markers include dihydrofolate reductase, G418, glutamine synthase, or neomycin resistance for eukaryotic cell culture, and tetracycline, kanamycin or ampicillin resistance genes for culturing in *E. coli* and other bacteria. Representative examples of appropriate hosts include, but are not limited to, bacterial cells, such as *E. coli*, *Streptomyces* and *Salmonella typhimurium* cells; fungal cells, such as yeast cells (e.g., *Saccharomyces cerevisiae* or *Pichia pastoris* (ATCC Accession No. 201178)); insect cells such as *Drosophila* S2 and *Spodoptera Sf9* cells; animal cells such as CHO, COS, NSO, 293, and Bowes melanoma cells; and plant cells. Appropriate culture mediums and conditions for the above-described host cells are known in the art.

In one embodiment, polynucleotides encoding an albumin fusion protein of the invention may be fused to signal sequences which will direct the localization of a protein of the invention to particular compartments of a prokaryotic or eukaryotic cell and/or direct the secretion of a protein of the invention from a prokaryotic or eukaryotic cell. For example, in *E. coli*, one may wish to direct the expression of the protein to the periplasmic space. Examples of signal sequences or proteins (or fragments thereof) to which the albumin fusion proteins of the invention may be fused in order to direct the expression of the polypeptide to the periplasmic space of bacteria include, but are not limited to, the *pelB* signal sequence, the maltose binding protein (MBP) signal sequence, MBP, the *ompA* signal sequence, the

signal sequence of the periplasmic *E. coli* heat-labile enterotoxin B-subunit, and the signal sequence of alkaline phosphatase. Several vectors are commercially available for the construction of fusion proteins which will direct the localization of a protein, such as the pMAL series of vectors (particularly the pMAL-p series) available from New England Biolabs. In a specific embodiment, polynucleotides albumin fusion proteins of the invention may be fused to the *pelB* pectate lyase signal sequence to increase the efficiency of expression and purification of such polypeptides in Gram-negative bacteria. See, U.S. Patent Nos. 5,576,195 and 5,846,818, the contents of which are herein incorporated by reference in their entireties.

Examples of signal peptides that may be fused to an albumin fusion protein of the invention in order to direct its secretion in mammalian cells include, but are not limited to, the MPIF-1 signal sequence (e.g., amino acids 1-21 of GenBank Accession number AAB51134), the stanniocalcin signal sequence (MLQNSAVLLLLVISASA, SEQ ID NO:10) and a consensus signal sequence (MPTWAWWLFLVLLLALWAPARG, SEQ ID NO:11). A suitable signal sequence that may be used in conjunction with baculoviral expression systems is the gp67 signal sequence (e.g., amino acids 1-19 of GenBank Accession Number AAA72759).

Vectors which use glutamine synthase (GS) or DHFR as the selectable markers can be amplified in the presence of the drugs methionine sulfoximine or methotrexate, respectively. An advantage of glutamine synthase based vectors are the availability of cell lines (e.g., the murine myeloma cell line, NSO) which are glutamine synthase negative. Glutamine synthase expression systems can also function in glutamine synthase expressing cells (e.g., Chinese Hamster Ovary (CHO) cells) by providing additional inhibitor to prevent the functioning of the endogenous gene. A glutamine synthase expression system and components thereof are detailed in PCT publications: WO87/04462; WO86/05807; WO89/01036; WO89/10404; and WO91/06657, which are hereby incorporated in their entireties by reference herein. Additionally, glutamine synthase expression vectors can be obtained from Lonza Biologics, Inc. (Portsmouth, NH).

Expression and production of monoclonal antibodies using a GS expression system in murine myeloma cells is described in Bebbington *et al.*, *Bio/technology* 10:169(1992) and in Biblia & Robinson *Biotechnol. Prog.* 11:1 (1995) which are herein incorporated by reference.

5

The present invention also relates to host cells containing vector constructs, such as those described herein, and additionally encompasses host cells containing nucleotide sequences of the invention that are operably associated with one or more heterologous control regions (e.g., promoter and/or enhancer) using techniques known of in the art. The host cell can be a higher eukaryotic cell, such as a mammalian cell (e.g., a human derived cell), or a lower eukaryotic cell, such as a yeast cell, or the host cell can be a prokaryotic cell, such as a bacterial cell. A host strain may be chosen which modulates the expression of the inserted gene sequences, or modifies and processes the gene product in the specific fashion desired. Expression from certain promoters can be elevated in the presence of certain inducers; thus expression of the genetically engineered polypeptide may be controlled. Furthermore, different host cells have characteristics and specific mechanisms for the translational and post-translational processing and modification (e.g., phosphorylation, cleavage) of proteins. Appropriate cell lines can be chosen to ensure the desired modifications and processing of the foreign protein expressed.

Introduction of the nucleic acids and nucleic acid constructs of the invention into the host cell can be effected by calcium phosphate transfection, DEAE-dextran mediated transfection, cationic lipid-mediated transfection, electroporation, transduction, infection, or other methods. Such methods are described in many standard laboratory manuals, such as Davis *et al.*, *Basic Methods In Molecular Biology* (1986). It is specifically contemplated that the polypeptides of the present invention may in fact be expressed by a host cell lacking a recombinant vector.

In addition to encompassing host cells containing the vector constructs discussed herein, the invention also encompasses primary, secondary, and immortalized host

cells of vertebrate origin, particularly mammalian origin, that have been engineered to delete or replace endogenous genetic material (e.g., the coding sequence corresponding to a Therapeutic protein may be replaced with an albumin fusion protein corresponding to the Therapeutic protein), and/or to include genetic material (e.g., heterologous polynucleotide sequences such as for example, an albumin fusion protein of the invention corresponding to the Therapeutic protein may be included). The genetic material operably associated with the endogenous polynucleotide may activate, alter, and/or amplify endogenous polynucleotides.

10 In addition, techniques known in the art may be used to operably associate heterologous polynucleotides (e.g., polynucleotides encoding an albumin protein, or a fragment or variant thereof) and/or heterologous control regions (e.g., promoter and/or enhancer) with endogenous polynucleotide sequences encoding a Therapeutic protein via homologous recombination (see, e.g., US 5,641,670; WO 96/29411; WO 94/12650; Koller *et al.*, *Proc. Natl. Acad. Sci. USA* 86:8932-8935 (1989); and Zijlstra *et al.*, *Nature* 342:435-438 (1989), the disclosures of each of which are incorporated by reference in their entireties).

Advantageously, albumin fusion proteins of the invention can be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxyapatite chromatography, hydrophobic charge interaction chromatography and lectin chromatography. In some embodiments, high performance liquid chromatography ("HPLC") may be employed for purification.

In preferred some embodiments albumin fusion proteins of the invention are purified using one or more chromatography methods listed above. In other embodiments, albumin fusion proteins of the invention are purified using one or more of the following chromatography columns, Q Sepharose FF column, SP Sepharose FF column, Q Sepharose High Performance Column, Blue Sepharose

FF column, Blue Column, Phenyl Sepharose FF column, DEAE Sepharose FF, or Methyl Column. "Sepharose" is a trademark.

5 Additionally, albumin fusion proteins of the invention may be purified using the process described in WO 00/44772 which is herein incorporated by reference in its entirety. One of skill in the art could easily modify the process described therein for use in the purification of albumin fusion proteins of the invention.

10 Albumin fusion proteins of the present invention may be recovered from: products produced by recombinant techniques from a prokaryotic or eukaryotic host, including, for example, bacterial, yeast, higher plant, insect, and mammalian cells. Depending upon the host employed in a recombinant production procedure, the polypeptides of the present invention may be glycosylated or may be non-glycosylated. In addition, albumin fusion proteins of the invention may also
15 include an initial modified methionine residue, in some cases as a result of host-mediated processes. Thus, it is well known in the art that the N-terminal methionine encoded by the translation initiation codon generally is removed with high efficiency from any protein after translation in all eukaryotic cells. While the N-terminal methionine on most proteins also is efficiently removed in most
20 prokaryotes, for some proteins, this prokaryotic removal process is inefficient, depending on the nature of the amino acid to which the N-terminal methionine is covalently linked.

25 Albumin fusion proteins of the invention and antibodies that bind a Therapeutic protein or fragments or variants thereof can be fused to marker sequences, such as a peptide to facilitate purification. In one embodiment, the marker amino acid sequence is a hexa-histidine peptide, such as the tag provided in a pQE vector (QIAGEN, Inc., 9259 Eton Avenue, Chatsworth, CA, 91311), among others, many of which are commercially available. As described in Gentz *et al.*, *Proc. Natl. Acad. Sci. USA* 86:821-824 (1989), for instance, hexa-histidine provides for
30 convenient purification of the fusion protein. Other peptide tags useful for purification include, but are not limited to, the "HA" tag, which corresponds to an

epitope derived from the influenza hemagglutinin protein (Wilson *et al.*, *Cell* 37:767 (1984)) and the "FLAG" tag.

Further, an albumin fusion protein of the invention may be conjugated to a
5 therapeutic moiety such as a cytotoxin, e.g., a cytostatic or cytocidal agent, a therapeutic agent or a radioactive metal ion, e.g., alpha-emitters such as, for example, ²¹³Bi. Examples of such agents are given in WO 03/066824 and in WO 01/79480 (p. 107), which are incorporated herein by reference.

10 Albumin fusion proteins may also be attached to solid supports, which are particularly useful for immunoassays or purification of polypeptides that are bound by, that bind to, or associate with albumin fusion proteins of the invention. Such solid supports include, but are not limited to, glass, cellulose, polyacrylamide, nylon, polystyrene, polyvinyl chloride or polypropylene.

15 Also provided by the invention are chemically modified derivatives of the albumin fusion proteins of the invention which may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Patent No. 4,179,337). Examples involving the use of
20 polyethylene glycol are given in WO 01/79480 (pp. 109-111), which are incorporated by reference herein.

The presence and quantity of albumin fusion proteins of the invention may be determined using ELISA, a well known immunoassay known in the art.

25

Uses of the Polypeptides

Each of the polypeptides identified herein can be used in numerous ways. The following description should be considered exemplary and utilizes known techniques.

30

The albumin fusion proteins of the present invention are useful for treatment, prevention and/or prognosis of various disorders in mammals, preferably humans.

Such disorders include, but are not limited to thrombocytopenia, vWD and inflammatory diseases, such as IBD.

Moreover, albumin fusion proteins of the present invention can be used to treat or prevent diseases or conditions. In addition, the albumin fusion proteins of the invention may be used as a prophylactic measure. Albumin fusion proteins can be used to assay levels of polypeptides in a biological sample. Albumin fusion proteins of the invention can also be used to raise antibodies, which in turn may be used to measure protein expression of the Therapeutic protein, albumin protein, and/or the albumin fusion protein of the invention from a recombinant cell, as a way of assessing transformation of the host cell, or in a biological sample. Moreover, the albumin fusion proteins of the present invention can be used to test the biological activities described herein.

Transgenic Organisms

Transgenic organisms that express the albumin fusion proteins of the invention are also included in the invention. Transgenic organisms are genetically modified organisms into which recombinant, exogenous or cloned genetic material has been transferred. Such genetic material is often referred to as a transgene. The nucleic acid sequence of the transgene may include one or more transcriptional regulatory sequences and other nucleic acid sequences such as introns, that may be necessary for optimal expression and secretion of the encoded protein. The transgene may be designed to direct the expression of the encoded protein in a manner that facilitates its recovery from the organism or from a product produced by the organism, *e.g.* from the milk, blood, urine, eggs, hair or seeds of the organism. The transgene may consist of nucleic acid sequences derived from the genome of the same species or of a different species than the species of the target animal. The transgene may be integrated either at a locus of a genome where that particular nucleic acid sequence is not otherwise normally found or at the normal locus for the transgene.

The term "germ cell line transgenic organism" refers to a transgenic organism in which the genetic alteration or genetic information was introduced into a germ line cell, thereby conferring the ability of the transgenic organism to transfer the

genetic information to offspring. If such offspring in fact possess some or all of that alteration or genetic information, then they too are transgenic organisms. The alteration or genetic information may be foreign to the species of organism to which the recipient belongs, foreign only to the particular individual recipient, or
5 may be genetic information already possessed by the recipient. In the last case, the altered or introduced gene may be expressed differently than the native gene.

A transgenic organism may be a transgenic human, animal or plant. Transgenics can be produced by a variety of different methods including transfection,
10 electroporation, microinjection, gene targeting in embryonic stem cells and recombinant viral and retroviral infection (*see, e.g.*, U.S. Patent No. 4,736,866; U.S. Patent No. 5,602,307; Mullins *et al.* (1993) *Hypertension* 22(4):630-633; Brenin *et al.* (1997) *Surg. Oncol.* 6(2)99-110; Tuan (ed.), *Recombinant Gene Expression Protocols*, Methods in Molecular Biology No. 62, Humana Press
15 (1997)). The method of introduction of nucleic acid fragments into recombination competent mammalian cells can be by any method which favours co-transformation of multiple nucleic acid molecules. Detailed procedures for producing transgenic animals are readily available to one skilled in the art, including the disclosures in U.S. Patent No. 5,489,743 and U.S. Patent No.
20 5,602,307. Additional information is given in WO 03/066824 and WO 01/79480 (pp. 151-162), which are incorporated by reference herein.

Gene Therapy

Constructs encoding albumin fusion proteins of the invention can be used as a part
25 of a gene therapy protocol to deliver therapeutically effective doses of the albumin fusion protein. One approach for *in vivo* introduction of nucleic acid into a cell is by use of a viral vector containing nucleic acid, encoding an albumin fusion protein of the invention. Infection of cells with a viral vector has the advantage that a large proportion of the targeted cells can receive the nucleic acid.
30 Additionally, molecules encoded within the viral vector, *e.g.*, by a cDNA contained in the viral vector, are expressed efficiently in cells which have taken up viral vector nucleic acid. The extended plasma half-life of the described albumin fusion proteins might even compensate for a potentially low expression level.

Retrovirus vectors and adeno-associated virus vectors can be used as a recombinant gene delivery system for the transfer of exogenous nucleic acid molecules encoding albumin fusion proteins *in vivo*. These vectors provide efficient delivery of nucleic acids into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. Examples of such vectors, methods of using them, and their advantages, as well as non-viral delivery methods are described in detail in WO 03/066824 and WO 01/79480 (pp. 151-153), which are incorporated by reference herein.

10

Gene delivery systems for a gene encoding an albumin fusion protein of the invention can be introduced into a patient by any of a number of methods. For instance, a pharmaceutical preparation of the gene delivery system can be introduced systemically, *e.g.* by intravenous injection, and specific transduction of the protein in the target cells occurs predominantly from specificity of transfection provided by the gene delivery vehicle, cell-type or tissue-type expression due to the transcriptional regulatory sequences controlling expression of the receptor gene, or a combination thereof. In other embodiments, initial delivery of the recombinant gene is more limited with introduction into the animal being quite localized. For example, the gene delivery vehicle can be introduced by catheter (see U.S. Patent 5,328,470) or by stereotactic injection (*e.g.* Chen *et al.* (1994) *PNAS* 91: 3054-3057). The pharmaceutical preparation of the gene therapy construct can consist essentially of the gene delivery system in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Where the albumin fusion protein can be produced intact from recombinant cells, *e.g.* retroviral vectors, the pharmaceutical preparation can comprise one or more cells which produce the albumin fusion protein. Additional gene therapy methods are described in WO 03/066824 and in WO 01/79480 (pp. 153-162), which are incorporated herein by reference.

30 Pharmaceutical or Therapeutic Compositions

The albumin fusion proteins of the invention or formulations thereof may be administered by any conventional method including parenteral (*e.g.* subcutaneous

or intramuscular) injection or intravenous infusion. The treatment may consist of a single dose or a plurality of doses over a period of time. Furthermore, the dose, or plurality of doses, is administered less frequently than for the Therapeutic Protein which is not fused to albumin.

5

While it is possible for an albumin fusion protein of the invention to be administered alone, it is desirable to present it as a pharmaceutical formulation, together with one or more acceptable carriers. The carrier(s) must be “acceptable” in the sense of being compatible with the albumin fusion protein and not deleterious to the recipients thereof. Typically, the carriers will be water or saline which will be sterile and pyrogen free. Albumin fusion proteins of the invention are particularly well suited to formulation in aqueous carriers such as sterile pyrogen free water, saline or other isotonic solutions because of their extended shelf-life in solution. For instance, pharmaceutical compositions of the invention may be formulated well in advance in aqueous form, for instance, weeks or months or longer time periods before being dispensed.

Formulations containing the albumin fusion protein may be prepared taking into account the extended shelf-life of the albumin fusion protein in aqueous formulations. As discussed above, the shelf-life of many of these Therapeutic proteins are markedly increased or prolonged after fusion to HA.

In instances where aerosol administration is appropriate, the albumin fusion proteins of the invention can be formulated as aerosols using standard procedures. The term “aerosol” includes any gas-borne suspended phase of an albumin fusion protein of the instant invention which is capable of being inhaled into the bronchioles or nasal passages. Specifically, aerosol includes a gas-borne suspension of droplets of an albumin fusion protein of the instant invention, as may be produced in a metered dose inhaler or nebulizer, or in a mist sprayer. Aerosol also includes a dry powder composition of a compound of the instant invention suspended in air or other carrier gas, which may be delivered by insufflation from an inhaler device, for example.

The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. Such methods include the step of bringing into association the albumin fusion protein with the carrier that constitutes one or more accessory ingredients. In general the
5 formulations are prepared by uniformly and intimately bringing into association the active ingredient with liquid carriers or finely divided solid carriers or both, and then, if necessary, shaping the product.

Formulations suitable for parenteral administration include aqueous and
10 non-aqueous sterile injection solutions which may contain anti-oxidants, buffers, bacteriostats and solutes which render the formulation appropriate for the intended recipient; and aqueous and non-aqueous sterile suspensions which may include suspending agents and thickening agents. The formulations may be presented in unit-dose or multi-dose containers, for example sealed ampoules, vials or syringes,
15 and may be stored in a freeze-dried (lyophilised) condition requiring only the addition of the sterile liquid carrier, for example water for injections, immediately prior to use. Extemporaneous injection solutions and suspensions may be prepared from sterile powders. Dosage formulations may contain the Therapeutic protein portion at a lower molar concentration or lower dosage compared to the
20 non-fused standard formulation for the Therapeutic protein given the extended serum half-life exhibited by many of the albumin fusion proteins of the invention.

As an example, when an albumin fusion protein of the invention comprises one or more of the Therapeutic protein regions, the dosage form can be calculated on the
25 basis of the potency of the albumin fusion protein relative to the potency of the Therapeutic protein, while taking into account the prolonged serum half-life and shelf-life of the albumin fusion proteins compared to that of the native Therapeutic protein. For example, in an albumin fusion protein consisting of a full length HA fused to a full length Therapeutic protein, an equivalent dose in terms of units
30 would represent a greater weight of agent but the dosage frequency can be reduced.

Formulations or compositions of the invention may be packaged together with, or included in a kit with, instructions or a package insert referring to the extended shelf-life of the albumin fusion protein component. For instance, such instructions or package inserts may address recommended storage conditions, such as time, temperature and light, taking into account the extended or prolonged shelf-life of the albumin fusion proteins of the invention. Such instructions or package inserts may also address the particular advantages of the albumin fusion proteins of the inventions, such as the ease of storage for formulations that may require use in the field, outside of controlled hospital, clinic or office conditions.

As described above, formulations of the invention may be in aqueous form and may be stored under less than ideal circumstances without significant loss of therapeutic activity.

The invention also provides methods of treatment and/or prevention of diseases or disorders (such as, for example, any one or more of the diseases or disorders disclosed herein) by administration to a subject of an effective amount of an albumin fusion protein of the invention or a polynucleotide encoding an albumin fusion protein of the invention ("albumin fusion polynucleotide") in a pharmaceutically acceptable carrier.

Effective dosages of the albumin fusion protein and/or polynucleotide of the invention to be administered may be determined through procedures well known to those in the art which address such parameters as biological half-life, bioavailability, and toxicity, including using data from routine *in vitro* and *in vivo* studies, using methods well known to those skilled in the art.

The albumin fusion protein and/or polynucleotide will be formulated and dosed in a fashion consistent with good medical practice, taking into account the clinical condition of the individual patient (especially the side effects of treatment with the albumin fusion protein and/or polynucleotide alone), the site of delivery, the method of administration, the scheduling of administration, and other factors known to practitioners. The "effective amount" for purposes herein is thus determined by such considerations.

For example, determining an effective amount of substance to be delivered can depend upon a number of factors including, for example, the chemical structure and biological activity of the substance, the age and weight of the patient, the
5 precise condition requiring treatment and its severity, and the route of administration. The frequency of treatments depends upon a number of factors, such as the amount of albumin fusion protein or polynucleotide constructs administered per dose, as well as the health and history of the subject. The precise amount, number of doses, and timing of doses will be determined by the
10 attending physician or veterinarian.

Albumin fusion proteins and polynucleotides of the present invention can be administered to any animal, preferably to mammals and birds. Preferred mammals include humans, dogs, cats, mice, rats, rabbits sheep, cattle, horses and
15 pigs, with humans being particularly preferred.

As a general proposition, the albumin fusion protein of the invention will be dosed lower (on the molar basis of the unfused Therapeutic protein) or administered less frequently than the unfused Therapeutic protein. The albumin fusion proteins of
20 the invention are advantageous in that they can simulate continuous infusion of "classic drugs", i.e., less protein equivalent is needed for identical inhibitory activity.

Albumin fusion proteins and/or polynucleotides can be are administered orally,
25 rectally, parenterally, intracisternally, intravaginally, intraperitoneally, topically (as by powders, ointments, gels, drops or transdermal patch), buccally, or as an oral or nasal spray. "Pharmaceutically acceptable carrier" refers to a non-toxic solid, semisolid or liquid filler, diluent, encapsulating material or formulation auxiliary of any. The term "parenteral" as used herein refers to modes of administration
30 which include intravenous, intramuscular, intraperitoneal, intrasternal, subcutaneous and intra-articular injection and infusion.

Albumin fusion proteins and/or polynucleotides of the invention are also suitably administered by sustained-release systems such as those described in WO 03/066824 and WO 01/79480 (pp. 129-130), which are incorporated herein by reference.

5

For parenteral administration, in one embodiment, the albumin fusion protein and/or polynucleotide is formulated generally by mixing it at the desired degree of purity, in a unit dosage injectable form (solution, suspension, or emulsion), with a pharmaceutically acceptable carrier, i.e., one that is non-toxic to recipients at the dosages and concentrations employed and is compatible with other ingredients of the formulation. For example, the formulation optionally does not include oxidizing agents and other compounds that are known to be deleterious to the Therapeutic.

15 The albumin fusion proteins and/or polynucleotides of the invention may be administered alone or in combination with other therapeutic agents. Albumin fusion protein and/or polynucleotide agents that may be administered in combination with the albumin fusion proteins and/or polynucleotides of the invention include, but are not limited to, antiretroviral agents like protease, reverse transcriptase, integrase and assembly inhibitors, chemotherapeutic agents, antibiotics, steroidal and non-steroidal anti-inflammatories, conventional immunotherapeutic agents, and/or therapeutic treatments as described, e.g., in WO 03/066824 and WO 01/79480 (pp. 132-151) which are incorporated by reference herein. Combinations may be administered either concomitantly, e.g., as an admixture, separately but simultaneously or concurrently; or sequentially. This includes presentations in which the combined agents are administered together as a therapeutic mixture, and also procedures in which the combined agents are administered separately but simultaneously, e.g., as through separate intravenous lines into the same individual. Administration "in combination" further includes the separate administration of one of the compounds or agents given first, followed by the second.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. In certain embodiments, albumin fusion proteins and/or polynucleotides of the invention are administered in combination with
5 antiretroviral agents, nucleoside/nucleotide reverse transcriptase inhibitors (NRTIs), non-nucleoside reverse transcriptase inhibitors (NNRTIs), and/or protease inhibitors (PIs).

The invention also provides a pharmaceutical pack or kit comprising one or more
10 containers filled with one or more of the ingredients of the pharmaceutical compositions comprising albumin fusion proteins of the invention. Optionally associated with such container(s) can be a notice in the form prescribed by a governmental agency regulating the manufacture, use or sale of pharmaceuticals or biological products, which notice reflects approval by the agency of
15 manufacture, use or sale for human administration.

Having generally described the invention, the same will be more readily understood by reference to the following examples, which are provided by way of illustration and are not intended as limiting.

20 Without further description, it is believed that one of ordinary skill in the art can, using the preceding description and the following illustrative examples, make and utilize the alterations detected in the present invention and practice the claimed methods. The following working examples therefore, specifically point out
25 certain embodiments of the present invention, and are not to be construed as limiting in any way the remainder of the disclosure.

Example 1: Preparation of albumin-fused IL-11

The recombinant albumin expression vectors pAYE645 and pAYE646 have been
30 described previously in WO 2004/009819. Plasmid pAYE645 contained the HSA/MF α -1 fusion leader sequence, as well as the yeast *PRB1* promoter and the yeast *ADHI* terminator providing appropriate transcription promoter and

transcription terminator sequences, is described in WO 2004/009819. Plasmid pAYE645 was digested to completion with the restriction enzyme *Afl*III and partially digested with the restriction enzyme *Hind*III and the DNA fragment comprising the 3' end of the yeast *PRB1* promoter and the albumin coding sequence was isolated. Plasmid pDB2241, described in patent application WO 00/44772, was digested with *Afl*III/*Hind*III and the DNA fragment comprising the 5' end of the yeast *PRB1* promoter and the yeast *ADHI* terminator was isolated. The *Afl*III/*Hind*III DNA fragment from pAYE645 was then cloned into the *Afl*III/*Hind*III pDB2241 vector DNA fragment to create the plasmid pDB2302. Plasmid pDB2302 was digested to completion with *Pac*I/*Xho*I and the 6.19kb fragment isolated, the recessed ends were blunt ended with T4 DNA polymerase and dNTPs, and religated to generate plasmid pDB2465. Plasmid pDB2465 was linearised with *Cla*I, the recessed ends were blunt ended with T4 DNA polymerase and dNTPs, and religated to generate plasmid pDB2533. Plasmid pDB2533 was linearised with *Bln*I, the recessed ends were blunt ended with T4 DNA polymerase and dNTPs, and religated to generate plasmid pDB2534. Plasmid pDB2534 was digested to completion with *Bmg*BI/*Bg*II, the 6.96kb DNA fragment isolated and ligated to one of two double stranded oligonucleotide linkers, VC053/VC054 and VC057/VC058 to create plasmid pDB2540, or VC055/VC056 and VC057/VC058 to create plasmid pDB2541.

VC053: 5'-GATCTTTGGATAAGAGAGACGCTCACAAGTCCGAAGTCGCT-CACCGGT-3' (SEQ ID NO: 1)

VC054: 5'-pCCTTGAACCGGTGAGCGACTTCGGACTTGTGAGCGTCTCTCTTATCCAAA-3' (SEQ ID NO: 2)

VC055: 5'-GATCTTTGGATAAGAGAGACGCTCACAAGTCCGAAGTCGCTCATCGAT-3' (SEQ ID NO: 3)

VC056: 5'-pCCTTGAATCGATGAGCGACTTCGGACTTGTGAGCGTCTCTCTCTATCCAAA-3' (SEQ ID NO: 4)

VC057: 5'-pTCAAGGACCTAGGTGAGGAAAACCTTCAAGGCTTTGGTCTTGATCGCTTTCGCTCAATACTTGCAACAATGTCCATTCGAAGATCAC-3' (SEQ ID NO: 5)

5'-

5 A sample of a human IL11 cDNA was serially diluted from 100ng.mL⁻¹ to 10pg.mL⁻¹ (in 10 fold increments). PCR primers CF59 (SEQ ID No: 7) and CF60 (SEQ ID No: 12) were designed to allow the IL11 cDNA to be cloned as an N-terminal albumin fusion into pDB2541 linearised with *Bgl*III and *Cla*I, whilst at the same time deleting the codon encoding the N-terminal proline from the IL11 coding region. The DNA sequence of each primer were as follows:

	HSA/MF α -1	
<i>Bgl</i> III	fusion leader	IL11

Clal albumin IL11

51

The PCR DNA fragment was digested to completion with the restriction endonucleases *Bgl*II/*Cla*I and the 0.58kb fragment was ligated into the 6.15kb pDB2541 *Bgl*II/*Cla*I vector DNA fragment to create plasmid pDB2567.

5

Appropriate yeast vector sequences were provide by a “disintegration” plasmid pSAC35 generally disclosed in EP-A-286 424 and described by Sleep, D., *et al.* (1991) *Bio/Technology* 9, 183-187. The 3.52kb *Not*I N-terminal (des-pro)IL11-albumin expression cassette was isolated from pDB2567, purified and ligated into

10 *Not*I digested pSAC35 which had been treated with calf intestinal phosphatase, creating plasmid pDB2569 contained the *Not*I expression cassette in the same orientation to the *LEU2* selection marker, and plasmid pDB2570 contained the *Not*I expression cassette in the opposite orientation to the *LEU2* selection marker.

15 PCR primers CF61 and CF62 were designed to allow the IL11 cDNA to be cloned as a C-terminal albumin fusion into pDB2243 linearised with *Bsu*36I and partially digested with *Hind*III, whilst at the same time adding two “TAA” translation stop codons at the 3’ end of the IL11 open reading frame. Plasmid pDB2243, previously described in patent application WO 00/44772, which contained the

20 yeast *PRB1* promoter and the yeast *ADHI* terminator provided appropriate transcription promoter and transcription terminator sequences. The DNA sequence of the CF61 (SEQ ID No: 13) and CF62 (SEQ ID No: 14) primers was as follows:

25 CF61

*Bsu*36I albumin IL11

5’-CCGGCCTTAGGCTTACCTGGGCCACCACTGGCCCCCCTCG-
AGTTTCCCC-3’

30 CF62

*Hind*III IL11

5’-GGCCAAGCTTATTACAGCCGAGTCTTCAGCAGCAGTCCCCTC-
3’

STOP STOP

35

A master mix was prepared as follows: 2mM MgCl₂ PCR Buffer, 10μM PCR dNTP's, 0.2μM CF61, 0.2μM CF62, 2U FastStart *Taq*. DNA polymerase. 1μL

of the IL11 cDNA (10pg, 100pg, 1ng, 10ng, 100ng) was added to 49µL of reaction mix. The total reaction volume was 50µL. Perkin-Elmer Thermal Cycler 9600 was programmed as follows: Denature at 95°C for 4 mins [HOLD], then [CYCLE] denature at 95°C for 30s, anneal for 30s at 45°C, extend at 72°C
5 for 60s for 20 cycles, followed by a [HOLD] 72°C for 600 s and then [HOLD] 4°C. The products of the PCR amplification were analysed by gel electrophoresis and a band of expected size (0.55kb) was observed. The 0.59kb DNA fragment was isolated from the 1%(w/v) agarose TAE gel using Gene Clean III Kit (BIO101 Inc.).

10

The PCR DNA fragment was digested to completion with the restriction endonucleases *Bsu36I/HindIII* and the 0.55kb fragment was ligated into the 6.19kb pDB2243 *Bsu36I*, partially digested with *HindIII* vector DNA fragment to create plasmid pDB2568.

15

Appropriate yeast vector sequences were provide by a “disintegration” plasmid pSAC35 generally disclosed in EP-A-286 424 and described by Sleep, D., *et al.* (1991) Bio/Technology 9, 183-187. The 3.53kb *NotI* C-terminal albumin-IL11 expression cassette was isolated from pDB2568, purified and ligated into *NotI*
20 digested pSAC35 which had been treated with calf intestinal phosphatase, creating plasmid pDB2571 contained the *NotI* expression cassette in the same orientation to the *LEU2* selection marker, and plasmid pDB2572 contained the *NotI* expression cassette in the opposite orientation to the *LEU2* selection marker. Yeast strains disclosed in WO 95/23857, WO 95/33833 and WO 94/04687 were
25 transformed to leucine prototrophy as described in Sleep D., *et al.* (2001) *Yeast* 18, 403-421. The transformants were patched out onto Buffered Minimal Medium (BMM, described by Kerry-Williams, S.M. *et al.* (1998) *Yeast* 14, 161-169) and incubated at 30°C until grown sufficiently for further analysis.

30 DNA sequence of the N-terminal IL11-albumin fusion open reading frame forms SEQ ID No: 15.

Amino acid sequence of the N-terminal IL11-albumin fusion protein is presented as
SEQ ID No: 16.

5

Amino acid sequence of the mature N-terminal IL11-albumin fusion protein forms
SEQ ID No: 17.

GPPPGPPRVSPDPRAELDSTVLLTRSLLADTRQLAAQLRDKFPADGDHNLDS
10 LPTLAMSGALGALQLPGVLTRLRADLLSYLRHVQWLRRAGGSSSLKTLEPE
LGTLQARLDRLRLRLQLMSRLALPQPPDPPAPPLAPPSSAWGGIRAAHAI
LGGLHLTLDWAVRGLLLLKTRLDAHKSEVAHRFKDLGEENFKALVLIAFA
QYLQQCPFEDHVKL VNEVTEFAKTCVADESAENCDKSLHTLFGDKLCTVA
TLRETYGEMADCCAKQEPERNECFLQHKDDNPNLPRLVRPEVDVMCTAFH
15 DNEETFLKKYLYEIARRHPYFYAPELLFFAKRYKAAFTECCQAADKAACLL
PKLDEL RDEGKASSAKQRLKCASLQKFGERAFKAWAVARLSQRFPKAEFA
EVSKLVTDLT KVHTECCHGDLLECADDRADLAKYICENQDSISSKLKECCE
KPLLEKSHCIAEVENDEMPADLPSLAADFVESKDVCKNYAEAKDVFLGMF
LYEYARRHPDYSVVL LRLAKTYETTLEKCCAAADPHECYAKVFDEFKPL
20 VEPPQNLIKQNC ELFELGEYKFQNALLVRYTKKVPQVSTPTLVEVSRNLG
KVGSKCKHPEAKRMPCAEDYLSVVLNQLCVLHEKTPVSDRVTKCCTESL
VNRRPCFSALEVD ETYVPKEFNAETFTFHADICTLSEKERQIKKQTALVELV
KHKPKATKEQLKAVMDDFAAFVEKCCKADDKETCFAEEGKKLV AASQAA
LGL

25

DNA sequence of the C-terminal albumin-IL11 fusion open reading frame is SEQ ID
No: 18.

30 And the corresponding amino acid sequence of the C-terminal albumin-IL11 fusion
protein is SEQ ID No: 19.

The amino acid sequence of the mature C-terminal albumin-IL11 fusion protein is SEQ ID No: 20.

5 **Example 2: Purification**

C-Terminal IL11 Purification

The C-Terminal IL11 fusion contained high levels of clipped (i.e. not full length) material. It was purified using the standard rHA SP-FF conditions as described in
10 WO 00/44772 but in a negative mode whereby the fusion was in the flowthrough. The flowthrough was adjusted to pH 8 and $2.5\text{mS}\cdot\text{cm}^{-1}$ and loaded on a standard rHA DE-FF equilibrated in 15mM potassium tetraborate. This was operated in a negative mode. The conductivity of the DE-FF flowthrough was increased to $15\text{mS}\cdot\text{cm}^{-1}$ and the material purified using standard rHA DBA chromatography
15 with an extra elution of 50mM octanoate in the equilibration buffer. The material was then concentrated and diafiltered against 300mM glycine, 10mM phosphate pH7.

N-Terminal IL11 Purification (type A)

20 The N-Terminal IL11 contained some clipped material. It was purified using the standard rHA SP-FF conditions as described in WO 00/44772. The majority was in the flowthrough but sufficient bound for it to be necessary for it to be eluted using the standard elution buffer containing 200mM NaCl. The eluate was then adjusted to pH 8 and $2.5\text{mS}\cdot\text{cm}^{-1}$ and purified using standard rHA DE-FF
25 equilibrated in 15mM potassium tetraborate. The DE-FF was eluted using the standard rHA elution buffer. The purified material was then concentrated and diafiltered against 300mM glycine, 10mM phosphate pH7.

N-Terminal IL11 Purification (type B)

30 The N-terminal IL-11 fusion protein contained some clipped material. This material was separated using a S-Sepharose-FF column according to the standard rHA SP-FF conditions. Free rHA remained on the column. Then the fusion protein in the flowthrough was adsorbed to a monoclonal antibody Sepharose

specific for albumin. The column was washed with high salt and eluted at pH 2.5. The eluate was adjusted to neutral pH, concentrated and diafiltered against 300mM glycine, 10mM Na-phosphate pH 7.0.

5 Protein Characterisation after Purification

Table 1: IL-11 albumin fusion characterisation

	C-Terminal Fusion	N-Terminal Fusion (Type A)	N-Terminal Fusion (Type B)
% Purity by SDS-PAGE and colloidal blue staining	77	90	89
ESMS indication of post-translational modifications	Theoretical mass 85565. Measured mass 68000-70000.	Theoretical mass 85468. Measured mass 85471. Good evidence for correct primary structure.	Theoretical mass 85468. Measured mass 85471. Good evidence for correct primary structure.
N-Terminal Sequence	N/A	Correct NT sequence for IL 11.	
Endotoxin (EU.mL ⁻¹)	73	23	130
Fusion Concentration (mg.mL ⁻¹)	1.5	2.7	1

- 10 Images of the 12% Gradient SDS Non-Reducing Gel and Western Blots are shown in Figure 11.

Example 3: Pharmacokinetics of albumin-fused IL-11 versus recombinant human IL-11 after single intravenous or subcutaneous administration to rabbits

15

Three male and three female rabbits per group received Neumega® IL-11 (100 µg/kg) or C-terminal albumin-fused IL-11 (440 µg/kg) by a single i.v. or s.c. injection on day 0 (Table 2). Blood samples were drawn for the determination of the respective antigen levels at baseline and at 5 min, 10 min, 20 min, 30 min, 45 min, 1 h, 2 h, 4 h, 8 h, 24 h (1 d), 48 h (2 d), 72 h (3 d), 5 d, 7 d, 9 d, 11 d, and 14 d after i.v. administration of the respective test substance and at baseline, 30 min, 1 h, 2 h, 4 h, 8 h, 24 h (1 d), 48 h (2 d), 72 h (3 d), 5 d, 7 d, 9 d, 11 d and 14 d following s.c. injection. The doses of Neumega® IL-11 and C-terminal albumin-fused IL-11 were calculated on an equimolar basis.

10

Measurement of IL-11 plasma levels

Plasma levels of human IL-11 were measured by Quantikine® Human IL-11 Immunoassay (R&D Systems, Catalog No. D1100). This assay employs the quantitative sandwich enzyme immunoassay technique. A murine monoclonal antibody specific to IL-11 has been coated onto a microplate. Standards and samples are pipetted into the wells and any IL-11 present is bound by the immobilized antibody. After washing away any unbound substances, an enzyme-linked polyclonal antibody specific for IL-11 is added to the wells. Following a wash to remove any unbound antibody-enzyme reagent, a substrate solution is added to the wells and colour develops in proportion to the amount of IL-11 bound in the initial step. The colour development is stopped and the intensity of the colour is measured.

15
20

Plasma half life of fused vs. unfused protein in rabbits

The means and standard deviations of the IL-11 concentrations at every time point are shown in Figure 1 for the i.v.-treated groups and in Figure 2 for the s.c. treated groups.

In the animals treated intravenously, levels above 100 pg/mL can be found for 1 day after injection in the Neumega® IL-11 group and for 9 days after injection in the albumin-fused IL-11 group.

30

In the animals treated subcutaneously, the levels in the Neumega® IL-11 group reach their peak around 6 hours after injection and stay above 100 pg/mL for little

less than 2 days (Figure 2). The levels in the albumin-fused IL-11 group reach their peak 24 hours after injection and stay above 100 pg/mL for just under 7 days.

The pharmacokinetic results are presented in Table 2 for the i.v.-treated groups and in Table 3 for the s.c.-treated groups.

5

Table 2: Pharmacokinetic results following i.v. administration

		Neumega® IL-11	C-terminal albumin-fused IL-11
N		6	6
Initial half-life (hr)	Mean	0.12	3.61
	Std Dev	0.03	2.53
	Median	0.11	3.32
	Range	0.09 – 0.15	0.30 – 6.52
Terminal half-life (hr)	Mean	3.31	22.79
	Std Dev	2.30	8.07
	Median	2.12	22.08
	Range	1.59 – 7.01	14.73 – 35.93
AUC₀₋₁₄ (hr·pg/mL)	Mean	162,686	5,421,764
	Std Dev	60,920	513,997
	Median	156,014	5,552,217
	Range	99,915 – 232,051	4,719,147 – 5,946,916
	Geometric mean	152,862	5,397,976
	Scatter factor*	1.47	1.10
C_{max} (pg/mL)	Mean	547,425	395,308
	Std Dev	194,193	61,216
	Median	540,825	379,975
	Range	274,200 – 760,450	332,300 – 489,100
Total clearance (mL/hr/kg)	Mean	721	21.0
	Std Dev	335	3.1
	Median	662	19.5
	Range	382 – 1,123	18.5 – 19.5
Total volume of distribution (mL/kg)	Mean	771	537
	Std Dev	636	170
	Median	628	500
	Range	204 – 1,781	377 – 847

		Neumega® IL-11	C-terminal albumin-fused IL-11
Mean residence time (hr)	Mean	0.94	25.1
	Std Dev	0.49	4.3
	Median	0.81	25.2
	Range	0.44 – 1.63	20.4 – 31.9

* Scatter factor = $\exp[\text{standard deviation (log-transformed values)}]$

Table 3: Pharmacokinetic results following s.c. administration

5

		Neumega® IL-11	C-terminal albumin-fused IL-11
	N	6	6
Absorption half-life (hr)	Mean	1.14	16.5
	Std Dev	0.76	8.5
	Median	0.81	15.0
	Range	0.35 – 2.12	7.7 – 32.5
Terminal half-life (hr)	Mean	4.67	18.0
	Std Dev	1.43	3.9
	Median	4.40	19.2
	Range	3.07 – 6.67	11.7 – 22.3
AUC₀₋₁₄ (hr·pg/mL)	Mean	122,093	1,794,152
	Std Dev	28,159	753,505
	Median	118,860	1,665,227
	Range	95,376 – 172,688	1,113,203 – 3,000,848
	Geometric mean	119,635	1,670,725
	Scatter factor*	1.24	1.51
C_{max} (pg/mL)	Mean	8,098	30,590
	Std Dev	2,468	9,091
	Median	8,061	29,420
	Range	5,342 – 11,117	21,261 – 44,730
t_{max} (hr)	Mean	5.67	28.0
	Std Dev	2.66	9.80
	Median	6.00	24.0
	Range	2.00 – 8.00	24.0 – 48.0

		Neumega® IL-11	C-terminal albumin-fused IL-11
Relative clearance (mL/hr/kg)	total Mean	1,270	70.9
	Std Dev	407	27.5
	Median	1,286	69.9
	Range	639 – 1,842	38.3 – 102.3
Relative volume of distribution (mL/kg)	total Mean	8,880	1753
	Std Dev	4,969	532
	Median	6,910	1,599
	Range	4,512 – 15,836	1,041 – 2,495

* Scatter factor = $\exp[\text{standard deviation (log-transformed values)}]$

Subcutaneous versus intravenous administration in rabbits

- 5 C-terminal albumin-fused IL-11 showed an average elimination half-life that was 8 times longer than that of Neumega® IL-11 after i.v. application (Table 4). The area under the curve was 35 times larger. After s.c. injection, the average elimination half-life of C-terminal albumin-fused IL-11 was 4 times longer than that of Neumega® IL-11. The area under the curve was 14 times larger.

10

Table 4: Comparison of bioavailability between substances

Rou te	Parameter	Elimination half-life	AUC ₀₋₁₄	C _{max}
i.v.	Estimated ratio (C-terminal albumin-fused IL-11/ Neumega®)	7.85	35.3	0.76
	90% confidence limits	5.24 – 11.76	25.9 – 48.2	0.56 – 1.03
s.c.	Estimated ratio (C-terminal albumin-fused IL-11/ Neumega®)	3.93	14.0	3.80
	90% confidence limits	2.62 – 5.89	10.2 – 19.1	2.79 – 5.17

The bioavailability of Neumega® s.c. versus i.v. was 78 % (Table 5) while for C-terminal albumin fusion it was calculated as 31 %.

Table 5: Comparison of bioavailability between routes of application

5

Substance	Parameter		Elimination half-life	AUC ₀₋₁₄	C _{max}
Neumega® IL-11	Estimated	ratio	1.63	0.78	0.015
	(s.c. / i.v.)				
	90% confidence limits		1.09 – 2.44	0.57 – 1.07	0.011 – 0.021
C-term. albumin- fused IL-11	Estimated	ratio	0.81	0.31	0.075
	(s.c. / i.v.)				
	90% confidence limits		0.54 – 1.22	0.23 – 0.42	0.055 – 0.103

Example 4: Pharmacokinetics of albumin-fused IL-11 versus recombinant human IL-11 after intravenous or subcutaneous administration to rats

Three rats per group received Neumega® IL-11 (100 µg/kg) or N-terminal
 10 albumin-fused IL-11 (Type A) (440 µg/kg) by a single i.v. or s.c. injection on day
 0 (Table 6). Blood samples for the determination of the respective antigen levels
 were drawn at the following timepoints: pre-injection, 2 min, 5 min, 10 min, 15
 min, 30 min, 45 min, 1 h, 2 h, 8 h, 24 h, 48 h, 72 h (3d), 5 d, 7 d. The doses of
 Neumega® IL-11 and N-terminal albumin-fused IL-11 were calculated on an
 15 equimolar basis.

Study design

Measurement of IL-11 plasma levels

20

Plasma levels for groups 1 and 3 (Neumega® IL-11) were measured with an anti-
 human IL-11 ELISA by Quantikine® Human IL-11 Immunoassay (R&D Systems,
 Catalog No. D1100), as described in Example 3 above. Plasma levels for groups
 2 and 4 (IL-11-AFP) were determined with an anti-human albumin ELISA. The
 25 standard for this assay was IL-11-AFP.

Plasma half life of fused vs. unfused protein in rats

- In the animals treated intravenously (Figure 3), levels above 0.1 ng/mL were found for 6 to 12 hours after injection in the Neumega[®] group and for 3 days after injection in the albumin-fused IL-11 group. In the animals treated subcutaneously (Figure 4), the levels in the Neumega[®] IL-11 group reached their peak within 1 hour after injection and stayed above 0.1 ng/mL for about 9 hours. The levels in the albumin-fused IL-11 group reached their peak between 2 and 8 hours after injection and stayed above 0.1 ng/mL for at least 44 hours.
- 10 The pharmacokinetic results are presented in Table 6 for the i.v. treated groups and in Table 7 for the s.c. treated groups.

Table 6: Pharmacokinetic results following i.v. administration

		Neumega [®] IL-11	N-terminal albumin-fused IL-11
		N	3
Initial half-life (hr)	Mean	0.13	3.15
	Std Dev	0.09	2.45
	Median	0.09	3.09
	Range	0.07 – 0.24	0.73 – 5.62
Terminal half-life (hr)	Mean	1.39	28.08
	Std Dev	0.92	23.10
	Median	0.92	24.57
	Range	0.80 – 2.46	6.93 – 52.73
AUC (hr·pg/mL)	Mean	26,375	8,357,724
	Std Dev	4,595	1,214,981
	Median	25,108	8,164,280
	Range	22,548 – 31,471	7,251,070 – 9,657,822
	Geometric mean	26,118	8,299,777
		Scatter factor*	1.19
C_{max} (pg/mL)	Mean	33,510	1,065,644
	Std Dev	4,310	133,261

		Neumega® IL-11	N-terminal albumin-fused IL-11
	Median	31,565	1,134,432
	Range	30,515 – 38,450	912,045 – 1,150,455
Total clearance (mL/hr/kg)	Mean	4,324	14.1
	Std Dev	570	2.2
	Median	4,253	15.1
	Range	3,793 – 4,926	11.6 – 15.7
Total volume of distribution (mL/kg)	Mean	3,281	157
	Std Dev	538	27
	Median	2,971	150
	Range	2,970 – 3,902	134 – 187
Mean residence time (hr)	Mean	0.78	11.2
	Std Dev	0.22	2.1
	Median	0.70	11.9
	Range	0.60 – 1.03	8.9 – 12.9

* Scatter factor = $\exp[\text{standard deviation (log-transformed values)}]$

** measured at 10 minutes, the earliest post-injection measurement in study PSR

04/02

5

Table 7: Pharmacokinetic results following s.c. administration

		Neumega® IL-11	N-terminal albumin-fused IL-11
N		3	3
Absorption half-life (hr)	Mean	0.74	2.63
	Std Dev	0.19	1.97
	Median	0.85	3.24
	Range	0.52 – 0.85	0.42 – 4.22
Terminal half-life (hr)	Mean	0.87	6.70
	Std Dev	0.04	4.84
	Median	0.86	4.66
	Range	0.85 – 0.92	3.21 – 12.23
AUC (hr·pg/mL)	Mean	24,687	1,032,955

		Neumega® IL-11	N-terminal albumin- fused IL-11
	Std Dev	2,547	335,512
	Median	25,706	1,043,068
	Range	21,788 – 26,566	692,500 – 1,363,295
	Geometric mean	24,596	994,888
	Scatter factor*	1.11	1.41
C_{max} (pg/mL)	Mean	7,550	63,636
	Std Dev	135	14,205
	Median	7,612	62,159
	Range	7,395 – 7,642	50,227 – 78,523
t_{max} (hr)	Mean	1.0	6.0
	Std Dev	0.0	3.5
	Median	1.0	8.0
	Range	1.0 – 1.0	2.0 – 8.0
Relative total clearance (mL/hr/kg)	Mean	4,168	112.0
	Std Dev	480	47.5
	Median	3,962	87.9
	Range	3,826 – 4,718	81.3 – 166.6
Relative total volume of distribution (mL/kg)	Mean	5,275	956
	Std Dev	842	527
	Median	4,858	771
	Range	4,723 – 6,244	547 – 1,550

* Scatter factor = $\exp[\text{standard deviation (log-transformed values)}]$

The maximum IL-11 concentrations measured in Study PSR 01/03 are presented
5 in Table 8.

Table 8: Maximum values obtained within one hour post-injection

Neumega® IL-11	N-terminal albumin-fused IL-11
----------------	-----------------------------------

		Neumega® IL-11		N-terminal albumin-fused IL-11	
		N	3	3	
i.v.	C_{max} (pg/mL)	Mean	910,200	1,537,879	
		Std Dev	30,524	109,122	
		Median	911,600	1,581,818	
		Range	879,000 – 940,000	1,413,636 – 1,618,182	
s.c.	C_{max} (pg/mL)	Mean	7,086	0	
		Std Dev	618	0	
		Median	7,133	0	
		Range	6,445 – 7,679	0 – 0	
	t_{max} (hours)	Mean	0.42	–	
		Std Dev	0.29	–	
		Median	0.25	–	
		Range	0.25 – 0.75	0.25 – 1.00	

Subcutaneous versus intravenous administration in rats

Table 9 shows the results of the analyses of variance regarding the relative bioavailability. The differences between the two products were significant with respect to elimination half-life, AUC and C_{max} for both routes of application.

Table 9: Comparison of bioavailability between substances

Route	Parameter	Elimination half-life	AUC	C _{max}
i.v.	Estimated ratio (N-terminal albumin-fused IL-11/ Neumega®)	17.03	317.8	1.69
	90% confidence limits	5.98 – 48.49	230.8 – 437.5	1.41 – 2.02
s.c.	Estimated ratio (N-terminal albumin-fused IL-11/ Neumega®)	6.50	40.5	8.29
	90% confidence limits	2.28 – 18.50	29.4 – 55.7	6.92 – 9.93

Table 10 shows the results of the analyses of variance regarding the absolute bioavailability. For Neumega[®] IL-11, the differences between the two routes of application were not statistically significant with respect to elimination half-life and AUC. The difference in C_{max} was highly significant. For albumin-fused IL-11, the differences between the two routes of application were not significant with respect to elimination half-life. The differences regarding AUC and C_{max} were highly significant.

Table 10: Comparison of bioavailability between routes of application

Substance	Parameter		Elimination half-life	AUC	C _{max} ^a
Neumega [®] IL-11	Estimated (s.c. / i.v.)	ratio	0.72	0.942	0.008
	90% confidence limits		0.25 – 2.04	0.684 – 1.296	0.007 – 0.010
N-term. albumin-fused IL-11	Estimated (s.c. / i.v.)	ratio	0.27	0.120	0.041
	90% confidence limits		0.10 – 0.78	0.087 – 0.165	0.034 – 0.049

^a C_{max} values of i.v. route taken from Study PSR 01/03, measured 2 minutes post-injection.

In rats N-terminal albumin-fused IL-11 showed an average elimination half-life that was 17 times longer than that of Neumega[®] IL-11 after i.v. application. The area under the curve was 318 times larger. After s.c. injection, the average elimination half-life of N-terminal albumin-fused IL-11 was 6 times longer than that of Neumega[®] IL-11. The area under the curve was 40 times larger.

Example 5: Stimulation of thrombocytopoiesis in naive rats by IL-11-fusion protein

Naive CD-rats (10 per group) were treated with Neumega® IL-11, C-terminal- or N-terminal-fused IL-11 (type B), or placebo according to the following schedule.

5 **Table 11: Treatment schedule**

Group	Substance	Dose	Appl. Sch.	Vol.	N
1	Formulation solution	-	t= d 0 - d 9 s.c.	0.5 ml	10
2	Neumega® IL-11	100 µg/kg	t= d 0 - d 9 s.c.	0.5 ml	10
3	IL-11-AFP C-term	660 µg/kg	t= d 0, d 3, d 6, d 9 s.c.	0.5 ml	10
4	IL-11-AFP N-term	660 µg/kg	t= d 0, d 3, d 6, d 9 s.c.	0.5 ml	10
5	IL-11-AFP C-term	660 µg/kg	t= d 0 - d 9 s.c.	0.5 ml	10
6	IL-11-AFP N-term	660 µg/kg	t= d 0 - d 9 s.c.	0.5 ml	10

Blood samples were drawn at baseline, day 5, day 7, day 9, day 13 and day 16 and hematologic parameters (PLT, WBC, RBC, HCT, HGB) and body weight were measured. The dose of the fusion protein was calculated on an equimolar basis compared to Neumega® IL-11 and corrected by a factor of 1.5 due to SDS-PAGE results.

Results

Maximum platelet counts were achieved in all groups on day 7 (Figure 5). The highest mean levels of $1528 \times 103/\mu\text{L}$ were achieved with the N-terminal fusion administered daily (group 6) or every three days (group 4) with $1304 \times 103/\mu\text{L}$. The C-terminal fusion also showed a dose-interval related effect with maximum mean levels of $1238 \times 103/\mu\text{L}$ when administered daily compared to $977 \times 103/\mu\text{L}$ if given every three days. Neumega® IL-11 reached peak levels of $1032 \times 103/\mu\text{L}$ while the control animals remained at $864 \times 103/\mu\text{L}$.

Red blood parameters (RBC, HGB, HCT) remained constant for all groups. White blood cell counts showed a slight initial increase followed by a non-consistent fluctuation. This can probably be attributed to the microtrauma caused by frequent injections and blood sampling. All animals showed a normal, slight increase in body weight over the course of the observation period.

In summary, both the N- and the C-terminal fusion administered daily as well as the N-terminal fusion administered in 3-day-intervals were clearly superior to control and Neumega® IL-11 on days 5, 7 and 9. However, the N-terminal fusion seems to achieve higher levels than the C-terminal fusion.

The decrease of platelet counts after day 7 despite continuation of treatment until day 9 might occur due to the development of neutralizing antibodies to the heterologous human protein.

10 **Example 6: Treatment of chemotherapy-induced thrombocytopenia in rats by IL-11-fusion-protein**

Ten female CD-rats per group received Carboplatin at 35 mg/kg i.v. on day 0. Starting on day 5 they were treated with Neumega® IL-11, C-terminal- or N-terminal-fused IL-11 (type B), or placebo according to the following schedule.

15

Table 12: Treatment schedule

Group	Substance	Dose	Appl. Sch.	Vol.	N
1	Formulation solution	-	t= d 5 - d 14 s.c.	0.5 ml	10
2	Neumega®	50 µg /kg	t= d 5 - d 14 s.c.	0.5 ml	10
3	IL-11-AFP C-term	330 µg/kg	t= d 5, d 8, d 11, d 14 s.c.	0.5 ml	10
4	IL-11-AFP N-term	330 µg/kg	t= d 5, d 8, d 11, d 14 s.c.	0.5 ml	10
5	IL-11-AFP C-term	330 µg/kg	t= d 5 - d 14 s.c.	0.5 ml	10
6	IL-11-AFP N-term	330 µg/kg	t= d 5 - d 14 s.c.	0.5 ml	10

Blood samples were drawn at baseline, day 5, day 7, day 9, day 13 and day 16 and hematologic parameters (PLT, WBC, RBC, HCT, HGB) and body weight were measured. The dose of the fusion protein was calculated on an equimolar basis compared to Neumega® IL-11 and corrected by a factor of 1.5 due to SDS-PAGE results.

Results

In all groups platelet nadirs were observed between day 5 and day 9 (Table 13). Thereafter platelet levels recovered to above baseline on day 16. Mean platelet levels are shown in the following table.

5 *Table 13: Mean platelet levels ($\times 10^3/\mu\text{L}$)*

	Group 1	Group 2	Group 3	Group 4	Group 5	Group 6
day 0	782	783	788	703	813	810
day 5	477	281	368	458	344	365
day 7	388	302	303	330	348	374
day 9	365	323	365	375	328	432
day 13	603	702	681	832	689	1064
day 16	1003	1034	969	974	901	1287

For Neumega® IL-11, C- and N-terminal fusion injected daily, the platelet-nadir was observed on day 5, the first day of treatment (Figure 6). Thereafter in the group receiving the N-terminal fusion daily the platelet levels increased steadily and significantly, while the C-terminal fusion administered daily induced an increase comparable to Neumega® IL-11.

An analysis of covariance adjusted for the day 5 platelet value shows a significantly higher platelet count for N-terminal fusion injected daily as compared to control on days 9, 13 and 16, as well as significant superiority over Neumega® IL-11 on day 13. The same analysis demonstrates an overall treatment effect with $p \leq 0.004$ on days 7, 13 and 16.

Considering the duration of platelet levels below $500 \times 10^3/\mu\text{L}$ again the N-terminal fusion shows the best effects at a mean of 5.87 days when injected daily or 6.27 every 3 days as compared to 8.08 days for Neumega® IL-11 and 6.93 days for controls. The C-terminal fusion shows less efficacy in that respect with a duration of 7.16 days with daily administration and 6.69 every 3 days.

Red blood parameters (RBC, HGB, HCT) showed a slight decrease during the study period in all groups, which might reflect the blood loss caused by frequent blood sampling. White blood cell counts showed a slight increase which can

probably be attributed to the microtrauma caused by frequent injections and blood sampling. However this effect was especially prominent in the animals receiving N-terminal fusion of IL-11 daily, so that in this group a treatment related effect on WBC levels can not be excluded. All animals showed a normal, slight increase in
5 body weight over the course of the observation period.

In summary the most prominent effect can be observed with the N-terminal albumin fusion of IL-11, while the C-terminal fusion behaves comparable to Neumega® IL-11. Even though the absolute levels at nadir are not affected
10 significantly, the N-terminal fusion protein is able to reduce the duration of levels below $500 \times 10^3/\mu\text{L}$ and leads to a significantly steeper recovery as compared to Neumega® IL-11.

Example 7: IL-11-fusion protein in a mouse model of inflammatory bowel disease (IBD)

15 In this model colitis is induced in mice through oral administration of 3 % dextran sulfate disodium salt (DSS) in tap water. Ten mice per group were allocated to the following treatment groups.

Table 14: Treatment schedule

Group	Treatment	Dose	Appl scheme	Volume	n
1	Formulation solution, no DSS	n.a.	t = d 0-9 s.c.; 1x/day	0.2 mL	10
2	Formulation solution	n.a.	t = d 0-9 s.c.; 1x/day	0.2 mL	10
3	Sulfasalazine	100 mg/kg	t = d 0-10; p.o.	ad lib.	10
4	Neumega® IL-11	250 µg/kg	t = d 0-9 s.c.; 1x/day	0.2 mL	10
5	IL-11-AFP term	C- 1650 µg/kg	t = d 0, 3, 6, 9 s.c.; 1x/day	0.2 mL	10
6	IL-11-AFP term (type B)	N- 1650 µg/kg	t = d 0, 3, 6, 9 s.c.; 1x/day	0.2 mL	10

20 Body weight, rectal bleeding/diarrhoea and occult blood in faeces were assessed before induction and on days 3, 6 and 9. The experiment was terminated on day 10, the animals were necropsied, colon length was measured and samples of large intestine were retained for histological evaluation.

Results

Animals receiving placebo, Sulfasalazine or Neumega[®] IL-11 lost weight after the onset of colitis, while the animals treated with N- or C-terminal fusion even
5 gained weight (Figure 7). This was a surprising effect that could not have been predicted in advance.

The visual observation score evaluating diarrhoea and rectal bleeding also shows that there was a surprising, and significant, beneficial effect of the fusion proteins
10 compared to that of placebo, Sulfasalazine or Neumega[®] IL-11 (Figure 8). The same holds true for the length of the colon (Figure 9) and the histological score (Figure 10).

In summary these data show that both albumin fusions given in 3 day intervals are
15 able to significantly ameliorate the symptoms of DSS-induced colitis in mice, even superior to Neumega[®] IL-11 or the current standard treatment Sulfasalazine.

Example 8: Shelf-life

20 Shelf-life of the albumin-fused IL-11 products was determined according to the following method. A total of 25mg of the albumin-fused IL-11 products was formulated to a final volume of 5ml in a glass vial in sterile water-for-injection, with glycine, heptahydrate di-basic sodium phosphate, and mono-basic basic sodium phosphate as excipients. Vials of the products were incubated at 25⁰C for
25 1 month. Samples were assayed at the end of one month as described in Example 1 for post-translational modifications, by ESMS, N-terminal sequence, non-reducing SDS polyacrylamide gels and Western blots. Shelf-life is considered to be prolonged if the compound exhibits fewer changes, in any one of these tests, after this storage than the unfused IL-11 compound.

30

Example 9: Human Administration

At least in the case of the treatment of cancers, the albumin-fused IL-11 products may be administered by single s.c. injection once every four days, with a dose
5 regime of 25-1000 µg per kg body weight, preferably 25-50 µg per kg.

References

- 10 The present invention is not to be limited in scope by the specific embodiments described which are intended as single illustrations of individual aspects of the invention, and functionally equivalent methods and components are within the scope of the invention. Indeed, various modifications of the invention, in addition to those shown and described herein will become apparent to those skilled in the
15 art from the foregoing description and accompanying drawings. Such modifications are intended to fall within the scope of the appended claims. Every reference cited hereinabove is incorporated by reference in its entirety.

SEQUENCE LISTING

5 <110> Delta Biotechnology Limited
 <120> Interleukin-11 fusion protein
 <130> DELF P29719EP
 10 <140> 03027770.1
 <141> 13/12/03
 <160> 20
 15 <170> SeqWin99
 <210> 1
 <211> 48
 <212> DNA
 20 <213> Artificial Sequence
 <220>
 <223> Oligonucleotide Linker VC053
 25 <400> 1
 gatctttgga taagagagac gtcacaagt ccgaagtcgc tcaccggt 48
 <210> 2
 <211> 50
 30 <212> DNA
 <213> Artificial Sequence
 <220>
 <223> Oligonucleotide Linker VC054
 35 <400> 2
 ccttgaaccg gtgagcgact tcggacttgt gagcgtctct cttatccaaa 50
 <210> 3
 40 <211> 48
 <212> DNA
 <213> Artificial Sequence
 <220>
 45 <223> Oligonucleotide Linker VC055
 <400> 3
 gatctttgga taagagagac gtcacaagt ccgaagtcgc tcacgat 48
 50 <210> 4
 <211> 50
 <212> DNA
 <213> Artificial Sequence
 55 <220>
 <223> Oligonucleotide Linker VC056
 <400> 4
 ccttgaatcg atgagcgact tcggacttgt gagcgtctct cttatccaaa 50
 60 <210> 5

<211> 86
 <212> DNA
 <213> Artificial Sequence

5 <220>
 <223> Oligonucleotide Linker VC057

<400> 5
 tcaaggacct aggtgaggaa aacttcaagg ctttgggtctt gatcgctttc gctcaatact 60
 10 tgcaacaatg tccattcgaa gatcac 86

<210> 6
 <211> 80
 <212> DNA
 15 <213> Artificial Sequence

<220>
 <223> Oligonucleotide Linker VC058

20 <400> 6
 gtgatcttcg aatggacatt gttgcaagta ttgagcgaaa gcgatcaaga ccaaagcctt 60
 gaagttttcc tcacctaggt 80

<210> 7
 25 <211> 54
 <212> DNA
 <213> Oligonucleotide Primer CF59

<400> 7
 30 cgatagatct ttggataaga gagggccacc acctggcccc cctcgagttt cccc 54

<210> 8
 <211> 5
 <212> PRT
 35 <213> Artificial Sequence

<220>
 <223> Polypeptide linker SEQ ID 8

40 <400> 8
 Gly Gly Gly Gly Ser
 1 5

<210> 9
 45 <211> 4
 <212> PRT
 <213> Artificial Sequence

<220>
 50 <223> Polypeptide linker SEQ ID 9

<400> 9
 Gly Gly Gly Ser
 1

55 <210> 10
 <211> 17
 <212> PRT
 <213> Stanniocalcin signal peptide sequence

60

<400> 10
 Met Leu Gln Asn Ser Ala Val Leu Leu Leu Leu Val Ile Ser Ala Ser
 1 5 10 15
 5 Ala

 <210> 11
 <211> 22
 10 <212> PRT
 <213> Consensus signal sequence

 <400> 11
 Met Pro Thr Trp Ala Trp Trp Leu Phe Leu Val Leu Leu Leu Ala Leu
 15 1 5 10 15
 Trp Ala Pro Ala Arg Gly
 20
 20 <210> 12
 <211> 66
 <212> DNA
 <213> Artificial Sequence
 25 <220>
 <223> Oligonucleotide Primer CF60

 <400> 12
 30 ggccatcgat gagegacttc ggacttgtga gcgtccagcc gagtcttcag cagcagcagt 60
 cccctc 66

 <210> 13
 <211> 50
 <212> DNA
 35 <213> Artificial Sequence

 <220>
 <223> Oligonucleotide Primer CF61
 40 <400> 13
 ccggccttag gcttacctgg gccaccacct ggccccctc gagtttcccc 50

 <210> 14
 <211> 45
 45 <212> DNA
 <213> Artificial Sequence

 <220>
 <223> Oligonucleotide Primer CF62
 50 <400> 14
 ggccaagctt attacagccg agtcttcagc agcagcagtc ccctc 45

 <210> 15
 55 <211> 2358
 <212> DNA
 <213> Artificial Sequence

 <220>
 60 <223> N-terminal IL11-albumin fusion

```

<400> 15
5  atgaagtggg ttttcatcgt ctccattttt ttcttgttct cctctgctta ctctagatct 60
   ttggataaga gagggccacc acctggcccc cctcgagttt cccagagacc tcggggccgag 120
   ctggacagca ccgtgctcct gacctgctct ctcttgccgg acacgcggca gctggctgca 180
   cagctgaggg acaaattccc agctgacggg gaccacaacc tggattccct gccaccctg 240
   gccatgagtg cgggggacct gggagctcta cagctcccag gtgtgctgac aaggctgcga 300
   gcggacctac tgtcctacct gcggcacgtg cagtggctgc gccgggcagg tggctcttcc 360
   ctgaagaccc tggagcccga gctgggcacc ctgcaggccc gactggaccg gctgctgcgc 420
10  cggtctgcagc tcctgatgtc ccgcctggcc ctgccccagc ccccccgga ccgcccggcg 480
   ccccgctgg cgccccctc ctgagcctgg gggggcatca gggccgcca cgccatcctg 540
   ggggggctgc acctgacact tgactgggccc gtgaggggac tgctgctgct gaagactcgg 600
   ctggacgctc acaagtccga agtcgctcat cgattcaagg acctaggtga ggaaaacttc 660
   aaggcttttg tcttgatcgc ttctgctcaa tacttgcaac aatgtccatt cgaagatcac 720
15  gtcaagtttg tcaacgaagt taccgaattc gctaagactt gtgttctgta cgaatctgct 780
   gaaaactgtg acaagtcctt gcacaccttg ttccggtgata agttgtgtac tgttctgctacc 840
   ttgagagaaa cctacggtga aatggctgac tgttctgcta agcaagaacc agaaagaaac 900
   gaatgtttct tgcaacacaa ggacgacaac ccaaacttgc caagatttgt tagaccagaa 960
   gttgacgtca tgtgtactgc ttccacgac aacgaagaaa ccttcttgaa gaagtacttg 1020
20  tacgaaattg ctagaagaca ccatacttc tacgctccag aattgttgtt cttcgctaag 1080
   agatacaagg ctgctttcac cgaatgttgt caagctgctg ataaggctgc ttgtttgttg 1140
   ccaaagttgg atgaattgag agacgaaggt aaggcttctt ccgctaagca aagattgaag 1200
   tgtgcttctt tgcaaaagtt cggtgaaaga gctttcaagg cttgggctgt cgctagattg 1260
   totcaaagat tcccaaaggc tgaattcgct gaagtttcta agttggttac tgacttgact 1320
25  aaggttcaca ctgaatgttg tcacggtgac ttgttggaa gtgctgatga cagagctgac 1380
   ttggctaagt acatctgtga aaaccaagac tctatctctt ccaagttgaa ggaatgttgt 1440
   gaaaagccat tgttggaaaa gtctcactgt attgctgaag ttgaaaacga tgaatgccca 1500
   gctgacttgc catcttttggc tgctgacttc gttgaatcta aggacgtttg taagaactac 1560
   gctgaagcta aggacgtctt cttgggtatg ttcttctgac aatacgctag aagacacca 1620
30  gactactccg ttgtcttgtt gttgagattg gctaagacct acgaaactac cttggaaaag 1680
   tgttctgctg ctgctgacct acacgaatgt tacgctaagg tttctgatga attcaagcca 1740
   ttggtcgaag aaccacaaaa cttgatcaag caaaactgtg aattgttcga acaattgggt 1800
   gaatacaagt tccaaaacgc tttgttgggt agatacacta agaaggtccc acaagtctcc 1860
   accccaactt tgggtgaagt ctctagaaac ttgggtaagg tcggttctaa gtgttgtaag 1920
35  caccagaag ctaagagaat gccatgtgct gaagattact tgtccgtcgt tttgaaccaa 1980
   ttgtgtgttt tgcacgaaaa gacccagtc tctgatagag tcaccaagtg ttgtactgaa 2040
   tctttgggtt acagaagacc atgtttctct gctttggaag tcgacgaaac ttacgttcca 2100
   aagggaattca acgctgaaac ttacaccttc cacgctgata tctgtacctt gtccgaaaag 2160
   gaaagacaaa ttaagaagca aactgctttg gttgaattgg tcaagcacia gccaaaggct 2220
40  actaaggaac aattgaaggc tgtcatggat gatttcgctg ctttcgttga aaagtgttgt 2280
   aaggctgatg ataaggaac ttgtttcgct gaagaaggta agaagtttgt cgctgcttcc 2340
   caagctgctt tgggtttg

```

```

45  <210> 16
     <211> 786
     <212> PRT
     <213> Artificial Sequence

```

```

50  <220>
     <223> N-terminal IL11-albumin fusion

```

```

<400> 16
Met Lys Trp Val Phe Ile Val Ser Ile Leu Phe Leu Phe Ser Ser Ala
1          5          10          15
55  Tyr Ser Arg Ser Leu Asp Lys Arg Gly Pro Pro Pro Gly Pro Pro Arg
     20          25          30
60  Val Ser Pro Asp Pro Arg Ala Glu Leu Asp Ser Thr Val Leu Leu Thr
     35          40          45

```

Arg Ser Leu Leu Ala Asp Thr Arg Gln Leu Ala Ala Gln Leu Arg Asp
 50 55 60

5 Lys Phe Pro Ala Asp Gly Asp His Asn Leu Asp Ser Leu Pro Thr Leu
 65 70 75 80

Ala Met Ser Ala Gly Ala Leu Gly Ala Leu Gln Leu Pro Gly Val Leu
 85 90 95

10 Thr Arg Leu Arg Ala Asp Leu Leu Ser Tyr Leu Arg His Val Gln Trp
 100 105 110

15 Leu Arg Arg Ala Gly Gly Ser Ser Leu Lys Thr Leu Glu Pro Glu Leu
 115 120 125

Gly Thr Leu Gln Ala Arg Leu Asp Arg Leu Leu Arg Arg Leu Gln Leu
 130 135 140

20 Leu Met Ser Arg Leu Ala Leu Pro Gln Pro Pro Pro Asp Pro Pro Ala
 145 150 155 160

Pro Pro Leu Ala Pro Pro Ser Ser Ala Trp Gly Gly Ile Arg Ala Ala
 165 170 175

25 His Ala Ile Leu Gly Gly Leu His Leu Thr Leu Asp Trp Ala Val Arg
 180 185 190

30 Gly Leu Leu Leu Leu Lys Thr Arg Leu Asp Ala His Lys Ser Glu Val
 195 200 205

Ala His Arg Phe Lys Asp Leu Gly Glu Glu Asn Phe Lys Ala Leu Val
 210 215 220

35 Leu Ile Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His
 225 230 235 240

Val Lys Leu Val Asn Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala
 245 250 255

40 Asp Glu Ser Ala Glu Asn Cys Asp Lys Ser Leu His Thr Leu Phe Gly
 260 265 270

45 Asp Lys Leu Cys Thr Val Ala Thr Leu Arg Glu Thr Tyr Gly Glu Met
 275 280 285

Ala Asp Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn Glu Cys Phe Leu
 290 295 300

50 Gln His Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu Val Arg Pro Glu
 305 310 315 320

Val Asp Val Met Cys Thr Ala Phe His Asp Asn Glu Glu Thr Phe Leu
 325 330 335

55 Lys Lys Tyr Leu Tyr Glu Ile Ala Arg Arg His Pro Tyr Phe Tyr Ala
 340 345 350

60 Pro Glu Leu Leu Phe Phe Ala Lys Arg Tyr Lys Ala Ala Phe Thr Glu
 355 360 365

Cys Cys Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu Pro Lys Leu Asp
 370 375 380
 5 Glu Leu Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys Gln Arg Leu Lys
 385 390 395 400
 Cys Ala Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe Lys Ala Trp Ala
 405 410 415
 10 Val Ala Arg Leu Ser Gln Arg Phe Pro Lys Ala Glu Phe Ala Glu Val
 420 425 430
 Ser Lys Leu Val Thr Asp Leu Thr Lys Val His Thr Glu Cys Cys His
 435 440 445
 15 Gly Asp Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp Leu Ala Lys Tyr
 450 455 460
 Ile Cys Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu Lys Glu Cys Cys
 465 470 475 480
 Glu Lys Pro Leu Leu Glu Lys Ser His Cys Ile Ala Glu Val Glu Asn
 485 490 495
 25 Asp Glu Met Pro Ala Asp Leu Pro Ser Leu Ala Ala Asp Phe Val Glu
 500 505 510
 Ser Lys Asp Val Cys Lys Asn Tyr Ala Glu Ala Lys Asp Val Phe Leu
 515 520 525
 Gly Met Phe Leu Tyr Glu Tyr Ala Arg Arg His Pro Asp Tyr Ser Val
 530 535 540
 35 Val Leu Leu Leu Arg Leu Ala Lys Thr Tyr Glu Thr Thr Leu Glu Lys
 545 550 555 560
 Cys Cys Ala Ala Ala Asp Pro His Glu Cys Tyr Ala Lys Val Phe Asp
 565 570 575
 40 Glu Phe Lys Pro Leu Val Glu Glu Pro Gln Asn Leu Ile Lys Gln Asn
 580 585 590
 Cys Glu Leu Phe Glu Gln Leu Gly Glu Tyr Lys Phe Gln Asn Ala Leu
 595 600 605
 45 Leu Val Arg Tyr Thr Lys Lys Val Pro Gln Val Ser Thr Pro Thr Leu
 610 615 620
 Val Glu Val Ser Arg Asn Leu Gly Lys Val Gly Ser Lys Cys Cys Lys
 625 630 635 640
 His Pro Glu Ala Lys Arg Met Pro Cys Ala Glu Asp Tyr Leu Ser Val
 645 650 655
 55 Val Leu Asn Gln Leu Cys Val Leu His Glu Lys Thr Pro Val Ser Asp
 660 665 670
 Arg Val Thr Lys Cys Cys Thr Glu Ser Leu Val Asn Arg Arg Pro Cys
 675 680 685
 60

Phe Ser Ala Leu Glu Val Asp Glu Thr Tyr Val Pro Lys Glu Phe Asn
 690 695 700

5 Ala Glu Thr Phe Thr Phe His Ala Asp Ile Cys Thr Leu Ser Glu Lys
 705 710 715 720

Glu Arg Gln Ile Lys Lys Gln Thr Ala Leu Val Glu Leu Val Lys His
 725 730 735

10 Lys Pro Lys Ala Thr Lys Glu Gln Leu Lys Ala Val Met Asp Asp Phe
 740 745 750

15 Ala Ala Phe Val Glu Lys Cys Cys Lys Ala Asp Asp Lys Glu Thr Cys
 755 760 765

Phe Ala Glu Glu Gly Lys Lys Leu Val Ala Ala Ser Gln Ala Ala Leu
 770 775 780

20 Gly Leu
 785

<210> 17
 <211> 762
 25 <212> PRT
 <213> Artificial Sequence

<220>
 <223> Mature N-terminal IL11-albumin fusion

30 <400> 17
 Gly Pro Pro Pro Gly Pro Pro Arg Val Ser Pro Asp Pro Arg Ala Glu
 1 5 10 15

35 Leu Asp Ser Thr Val Leu Leu Thr Arg Ser Leu Leu Ala Asp Thr Arg
 20 25 30

Gln Leu Ala Ala Gln Leu Arg Asp Lys Phe Pro Ala Asp Gly Asp His
 35 40 45

40 Asn Leu Asp Ser Leu Pro Thr Leu Ala Met Ser Ala Gly Ala Leu Gly
 50 55 60

45 Ala Leu Gln Leu Pro Gly Val Leu Thr Arg Leu Arg Ala Asp Leu Leu
 65 70 75 80

Ser Tyr Leu Arg His Val Gln Trp Leu Arg Arg Ala Gly Gly Ser Ser
 85 90 95

50 Leu Lys Thr Leu Glu Pro Glu Leu Gly Thr Leu Gln Ala Arg Leu Asp
 100 105 110

Arg Leu Leu Arg Arg Leu Gln Leu Leu Met Ser Arg Leu Ala Leu Pro
 115 120 125

55 Gln Pro Pro Pro Asp Pro Pro Ala Pro Pro Leu Ala Pro Pro Ser Ser
 130 135 140

60 Ala Trp Gly Gly Ile Arg Ala Ala His Ala Ile Leu Gly Gly Leu His
 145 150 155 160

Leu Thr Leu Asp Trp Ala Val Arg Gly Leu Leu Leu Leu Lys Thr Arg
 165 170 175
 5 Leu Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly
 180 185 190
 Glu Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu
 195 200 205
 10 Gln Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr
 210 215 220
 Glu Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp
 15 225 230 235 240
 Lys Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr
 245 250 255
 20 Leu Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu
 260 265 270
 Pro Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn
 275 280 285
 25 Leu Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe
 290 295 300
 His Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala
 30 305 310 315 320
 Arg Arg His Pro Tyr Phe Tyr Ala Pro Glu Leu Leu Phe Phe Ala Lys
 325 330 335
 35 Arg Tyr Lys Ala Ala Phe Thr Glu Cys Cys Gln Ala Ala Asp Lys Ala
 340 345 350
 Ala Cys Leu Leu Pro Lys Leu Asp Glu Leu Arg Asp Glu Gly Lys Ala
 355 360 365
 40 Ser Ser Ala Lys Gln Arg Leu Lys Cys Ala Ser Leu Gln Lys Phe Gly
 370 375 380
 Glu Arg Ala Phe Lys Ala Trp Ala Val Ala Arg Leu Ser Gln Arg Phe
 45 385 390 395 400
 Pro Lys Ala Glu Phe Ala Glu Val Ser Lys Leu Val Thr Asp Leu Thr
 405 410 415
 50 Lys Val His Thr Glu Cys Cys His Gly Asp Leu Leu Glu Cys Ala Asp
 420 425 430
 Asp Arg Ala Asp Leu Ala Lys Tyr Ile Cys Glu Asn Gln Asp Ser Ile
 435 440 445
 55 Ser Ser Lys Leu Lys Glu Cys Cys Glu Lys Pro Leu Leu Glu Lys Ser
 450 455 460
 His Cys Ile Ala Glu Val Glu Asn Asp Glu Met Pro Ala Asp Leu Pro
 60 465 470 475 480

Ser Leu Ala Ala Asp Phe Val Glu Ser Lys Asp Val Cys Lys Asn Tyr
 485 490 495
 5 Ala Glu Ala Lys Asp Val Phe Leu Gly Met Phe Leu Tyr Glu Tyr Ala
 500 505 510
 Arg Arg His Pro Asp Tyr Ser Val Val Leu Leu Leu Arg Leu Ala Lys
 515 520 525
 10 Thr Tyr Glu Thr Thr Leu Glu Lys Cys Cys Ala Ala Ala Asp Pro His
 530 535 540
 Glu Cys Tyr Ala Lys Val Phe Asp Glu Phe Lys Pro Leu Val Glu Glu
 15 545 550 555 560
 Pro Gln Asn Leu Ile Lys Gln Asn Cys Glu Leu Phe Glu Gln Leu Gly
 565 570 575
 20 Glu Tyr Lys Phe Gln Asn Ala Leu Leu Val Arg Tyr Thr Lys Lys Val
 580 585 590
 Pro Gln Val Ser Thr Pro Thr Leu Val Glu Val Ser Arg Asn Leu Gly
 595 600 605
 25 Lys Val Gly Ser Lys Cys Cys Lys His Pro Glu Ala Lys Arg Met Pro
 610 615 620
 Cys Ala Glu Asp Tyr Leu Ser Val Val Leu Asn Gln Leu Cys Val Leu
 30 625 630 635 640
 His Glu Lys Thr Pro Val Ser Asp Arg Val Thr Lys Cys Cys Thr Glu
 645 650 655
 35 Ser Leu Val Asn Arg Arg Pro Cys Phe Ser Ala Leu Glu Val Asp Glu
 660 665 670
 Thr Tyr Val Pro Lys Glu Phe Asn Ala Glu Thr Phe Thr Phe His Ala
 675 680 685
 40 Asp Ile Cys Thr Leu Ser Glu Lys Glu Arg Gln Ile Lys Lys Gln Thr
 690 695 700
 Ala Leu Val Glu Leu Val Lys His Lys Pro Lys Ala Thr Lys Glu Gln
 45 705 710 715 720
 Leu Lys Ala Val Met Asp Asp Phe Ala Ala Phe Val Glu Lys Cys Cys
 725 730 735
 50 Lys Ala Asp Asp Lys Glu Thr Cys Phe Ala Glu Glu Gly Lys Lys Leu
 740 745 750
 Val Ala Ala Ser Gln Ala Ala Leu Gly Leu
 755 760
 55 <210> 18
 <211> 2361
 <212> DNA
 <213> Artificial Sequence
 60

<220>

<223> C-terminal albumin-IL11 fusion

<400> 18

```

5 atgaagtggg taagctttat ttcccttctt tttctcttta gctcggctta ttccaggagc 60
  ttggataaaa gagatgcaca caagagtgag gttgctcatc ggtttaaaga tttgggagaa 120
  gaaaatttca aagccttggg gttgattgcc tttgctcagt atcttcagca gtgtccattt 180
  gaagatcatg taaaattagt gaatgaagta actgaatttg caaaaacatg tgttgctgat 240
  gagtcagctg aaaattgtga caaatcactt catacccttt ttggagacaa attatgcaca 300
10 gttgcaactc ttcgtgaaac ctatggtgaa atggctgact gctgtgcaaa acaagaacct 360
  gagagaaatg aatgcttctt gcaacacaaa gatgacaacc caaacctccc ccgattgggtg 420
  agaccagagg ttgatgtgat gtgcactgct tttcatgaca atgaagagac atttttgaaa 480
  aaatacttat atgaaattgc cagaagacat ccttactttt atgccccgga actccttttc 540
  tttgctaaaaa ggtataaagc tgctttttaca gaatgttgcc aagctgctga taaagctgcc 600
15 tgctgtgtgc caaagctcga tgaacttcgg gatgaaggga aggcttcgtc tgccaaacag 660
  agactcaagt gtgccagtct ccaaaaattt ggagaaagag ctttcaaagc atgggcagta 720
  gctcgctga gccagagatt tcccaaagct gagtttgcag aagtttcaa gttagtga 780
  gatcttacca aagtccacac ggaatgctgc catggagatc tgcttgaatg tgctgatgac 840
  agggcggacc ttgccaagta tatctgtgaa aatcaagatt cgatctccag taaactgaag 900
20 gaatgctgtg aaaaacctct gttggaaaaa tcccactgca ttgccgaagt ggaaaatgat 960
  gagatgcctg ctgacttgcc ttcattagct gctgattttg ttgaaagtaa ggatgtttgc 1020
  aaaaactatg ctgaggcaaa ggatgtcttc ctgggcatgt ttttgtatga atatgcaaga 1080
  aggcatcctg attactctgt cgtgctgctg ctgagacttg ccaagacata tgaaaccact 1140
  ctagagaagt gctgtgccgc tgcagatcct catgaatgct atgccaaagt gttcgaatga 1200
25 tttaaacctc ttgtggaaga gcctcagaat ttaatcaaac aaaattgtga gctttttgag 1260
  cagcttggag agtacaaatt ccagaatgct ctattagttc gttacaccaa gaaagtacct 1320
  caagtgtcaa ctccaactct tgtagaggct tcaagaaacc taggaaaagt gggcagcaaa 1380
  tggtgtaaac atoctgaagc aaaaagaatg ccctgtgcag aagactatct atccgtggtc 1440
  ctgaaccagt tatgtgtgtt gcatgagaaa acgccagtaa gtgacagagt caccaaagtc 1500
30 tgcacagaat ccttggtgaa caggcgacca tgcttttcag ctctggaagt cgatgaaaca 1560
  tacgttccca aagagtttaa tgctgaaaca ttcaccttcc atgcagatat atgcacactt 1620
  tctgagaagg agagacaaat caagaaacaa actgcacttg ttgagctcgt gaaacacaag 1680
  cccaaggcaa caaaagagca actgaaagct gttatggatg atttcgcagc tttttagtag 1740
  aagtgtgca aggtgacga taaggagacc tgctttgccc aggagggtaa aaaacttggt 1800
35 gctgcaagtc aagctgcctt aggtttacct gggccaccac ctggccccc tgcagtttcc 1860
  ccagaccctc gggccgagct ggacagcacc gtgctcctga cccgctctct cctggcggac 1920
  acgcggcagc tggctgcaca gctgagggac aaattcccag ctgacgggga ccacaacctg 1980
  gattccctgc ccacctggc catgagtgcg ggggcactgg gagctctaca gctcccaggt 2040
  gtgctgacaa ggctgcgagc ggacctactg tccacctgc ggcacgtgca gtggctgcgc 2100
40 cgggcagggt gctcttccct gaagaccctg gagcccagc tgggcaccct gcagcccca 2160
  ctggaccggc tgctgcgcgc gctgcagctc ctcatgtccc gcctggccct gcccagcca 2220
  cccccggacc cgccggcgcc ccogctggcg cccccctcct cagcctgggg gggcatcagg 2280
  gccgccacg ccacctggg ggggctgcac ctgacacttg actgggcccgt gaggggactg 2340
  ctgctgctga agactcggct g 2361

```

<210> 19

<211> 787

<212> PRT

<213> Artificial Sequence

50

<220>

<223> C-terminal albumin-IL11 fusion

<400> 19

```

55 Met Lys Trp Val Ser Phe Ile Ser Leu Leu Phe Leu Phe Ser Ser Ala
   1             5             10             15

```

```

Tyr Ser Arg Ser Leu Asp Lys Arg Asp Ala His Lys Ser Glu Val Ala
   20             25             30

```

60

His Arg Phe Lys Asp Leu Gly Glu Glu Asn Phe Lys Ala Leu Val Leu
 35 40 45
 5 Ile Ala Phe Ala Gln Tyr Leu Gln Gln Cys Pro Phe Glu Asp His Val
 50 55 60
 Lys Leu Val Asn Glu Val Thr Glu Phe Ala Lys Thr Cys Val Ala Asp
 65 70 75 80
 10 Glu Ser Ala Glu Asn Cys Asp Lys Ser Leu His Thr Leu Phe Gly Asp
 85 90 95
 Lys Leu Cys Thr Val Ala Thr Leu Arg Glu Thr Tyr Gly Glu Met Ala
 100 105 110
 15 Asp Cys Cys Ala Lys Gln Glu Pro Glu Arg Asn Glu Cys Phe Leu Gln
 115 120 125
 His Lys Asp Asp Asn Pro Asn Leu Pro Arg Leu Val Arg Pro Glu Val
 130 135 140
 Asp Val Met Cys Thr Ala Phe His Asp Asn Glu Glu Thr Phe Leu Lys
 145 150 155 160
 25 Lys Tyr Leu Tyr Glu Ile Ala Arg Arg His Pro Tyr Phe Tyr Ala Pro
 165 170 175
 Glu Leu Leu Phe Phe Ala Lys Arg Tyr Lys Ala Ala Phe Thr Glu Cys
 180 185 190
 30 Cys Gln Ala Ala Asp Lys Ala Ala Cys Leu Leu Pro Lys Leu Asp Glu
 195 200 205
 Leu Arg Asp Glu Gly Lys Ala Ser Ser Ala Lys Gln Arg Leu Lys Cys
 210 215 220
 Ala Ser Leu Gln Lys Phe Gly Glu Arg Ala Phe Lys Ala Trp Ala Val
 225 230 235 240
 40 Ala Arg Leu Ser Gln Arg Phe Pro Lys Ala Glu Phe Ala Glu Val Ser
 245 250 255
 Lys Leu Val Thr Asp Leu Thr Lys Val His Thr Glu Cys Cys His Gly
 260 265 270
 45 Asp Leu Leu Glu Cys Ala Asp Asp Arg Ala Asp Leu Ala Lys Tyr Ile
 275 280 285
 Cys Glu Asn Gln Asp Ser Ile Ser Ser Lys Leu Lys Glu Cys Cys Glu
 290 295 300
 Lys Pro Leu Leu Glu Lys Ser His Cys Ile Ala Glu Val Glu Asn Asp
 305 310 315 320
 55 Glu Met Pro Ala Asp Leu Pro Ser Leu Ala Ala Asp Phe Val Glu Ser
 325 330 335
 Lys Asp Val Cys Lys Asn Tyr Ala Glu Ala Lys Asp Val Phe Leu Gly
 340 345 350
 60

	Met	Phe	Leu	Tyr	Glu	Tyr	Ala	Arg	Arg	His	Pro	Asp	Tyr	Ser	Val	Val	
			355					360					365				
5	Leu	Leu	Leu	Arg	Leu	Ala	Lys	Thr	Tyr	Glu	Thr	Thr	Leu	Glu	Lys	Cys	
			370				375					380					
	Cys	Ala	Ala	Ala	Asp	Pro	His	Glu	Cys	Tyr	Ala	Lys	Val	Phe	Asp	Glu	
			385			390					395					400	
10	Phe	Lys	Pro	Leu	Val	Glu	Glu	Pro	Gln	Asn	Leu	Ile	Lys	Gln	Asn	Cys	
				405						410					415		
	Glu	Leu	Phe	Glu	Gln	Leu	Gly	Glu	Tyr	Lys	Phe	Gln	Asn	Ala	Leu	Leu	
			420						425					430			
15	Val	Arg	Tyr	Thr	Lys	Lys	Val	Pro	Gln	Val	Ser	Thr	Pro	Thr	Leu	Val	
			435					440					445				
	Glu	Val	Ser	Arg	Asn	Leu	Gly	Lys	Val	Gly	Ser	Lys	Cys	Cys	Lys	His	
20			450				455					460					
	Pro	Glu	Ala	Lys	Arg	Met	Pro	Cys	Ala	Glu	Asp	Tyr	Leu	Ser	Val	Val	
						470					475					480	
25	Leu	Asn	Gln	Leu	Cys	Val	Leu	His	Glu	Lys	Thr	Pro	Val	Ser	Asp	Arg	
				485						490						495	
	Val	Thr	Lys	Cys	Cys	Thr	Glu	Ser	Leu	Val	Asn	Arg	Arg	Pro	Cys	Phe	
				500					505					510			
30	Ser	Ala	Leu	Glu	Val	Asp	Glu	Thr	Tyr	Val	Pro	Lys	Glu	Phe	Asn	Ala	
			515					520					525				
	Glu	Thr	Phe	Thr	Phe	His	Ala	Asp	Ile	Cys	Thr	Leu	Ser	Glu	Lys	Glu	
35			530				535					540					
	Arg	Gln	Ile	Lys	Lys	Gln	Thr	Ala	Leu	Val	Glu	Leu	Val	Lys	His	Lys	
						550					555					560	
40	Pro	Lys	Ala	Thr	Lys	Glu	Gln	Leu	Lys	Ala	Val	Met	Asp	Asp	Phe	Ala	
				565						570					575		
	Ala	Phe	Val	Glu	Lys	Cys	Cys	Lys	Ala	Asp	Asp	Lys	Glu	Thr	Cys	Phe	
				580					585					590			
45	Ala	Glu	Glu	Gly	Lys	Lys	Leu	Val	Ala	Ala	Ser	Gln	Ala	Ala	Leu	Gly	
			595					600					605				
	Leu	Pro	Gly	Pro	Pro	Pro	Gly	Pro	Pro	Arg	Val	Ser	Pro	Asp	Pro	Arg	
50			610				615					620					
	Ala	Glu	Leu	Asp	Ser	Thr	Val	Leu	Leu	Thr	Arg	Ser	Leu	Leu	Ala	Asp	
						630					635					640	
55	Thr	Arg	Gln	Leu	Ala	Ala	Gln	Leu	Arg	Asp	Lys	Phe	Pro	Ala	Asp	Gly	
				645						650					655		
	Asp	His	Asn	Leu	Asp	Ser	Leu	Pro	Thr	Leu	Ala	Met	Ser	Ala	Gly	Ala	
				660					665					670			
60																	

Leu Gly Ala Leu Gln Leu Pro Gly Val Leu Thr Arg Leu Arg Ala Asp
 675 680 685
 5 Leu Leu Ser Tyr Leu Arg His Val Gln Trp Leu Arg Arg Ala Gly Gly
 690 695 700
 Ser Ser Leu Lys Thr Leu Glu Pro Glu Leu Gly Thr Leu Gln Ala Arg
 705 710 715 720
 10 Leu Asp Arg Leu Leu Arg Arg Leu Gln Leu Leu Met Ser Arg Leu Ala
 725 730 735
 Leu Pro Gln Pro Pro Pro Asp Pro Pro Ala Pro Pro Leu Ala Pro Pro
 740 745 750
 15 Ser Ser Ala Trp Gly Gly Ile Arg Ala Ala His Ala Ile Leu Gly Gly
 755 760 765
 20 Leu His Leu Thr Leu Asp Trp Ala Val Arg Gly Leu Leu Leu Leu Lys
 770 775 780
 Thr Arg Leu
 785
 25 <210> 20
 <211> 763
 <212> PRT
 <213> Artificial Sequence
 30 <220>
 <223> Mature C-terminal IL11-albumin fusion
 <400> 20
 35 Asp Ala His Lys Ser Glu Val Ala His Arg Phe Lys Asp Leu Gly Glu
 1 5 10 15
 Glu Asn Phe Lys Ala Leu Val Leu Ile Ala Phe Ala Gln Tyr Leu Gln
 20 25 30
 40 Gln Cys Pro Phe Glu Asp His Val Lys Leu Val Asn Glu Val Thr Glu
 35 40 45
 Phe Ala Lys Thr Cys Val Ala Asp Glu Ser Ala Glu Asn Cys Asp Lys
 50 55 60
 45 Ser Leu His Thr Leu Phe Gly Asp Lys Leu Cys Thr Val Ala Thr Leu
 65 70 75 80
 Arg Glu Thr Tyr Gly Glu Met Ala Asp Cys Cys Ala Lys Gln Glu Pro
 85 90 95
 50 Glu Arg Asn Glu Cys Phe Leu Gln His Lys Asp Asp Asn Pro Asn Leu
 100 105 110
 55 Pro Arg Leu Val Arg Pro Glu Val Asp Val Met Cys Thr Ala Phe His
 115 120 125
 Asp Asn Glu Glu Thr Phe Leu Lys Lys Tyr Leu Tyr Glu Ile Ala Arg
 130 135 140
 60

	Arg	His	Pro	Tyr	Phe	Tyr	Ala	Pro	Glu	Leu	Leu	Phe	Phe	Ala	Lys	Arg	
	145					150					155					160	
5	Tyr	Lys	Ala	Ala	Phe	Thr	Glu	Cys	Cys	Gln	Ala	Ala	Asp	Lys	Ala	Ala	
					165					170					175		
	Cys	Leu	Leu	Pro	Lys	Leu	Asp	Glu	Leu	Arg	Asp	Glu	Gly	Lys	Ala	Ser	
				180					185					190			
10	Ser	Ala	Lys	Gln	Arg	Leu	Lys	Cys	Ala	Ser	Leu	Gln	Lys	Phe	Gly	Glu	
			195					200					205				
	Arg	Ala	Phe	Lys	Ala	Trp	Ala	Val	Ala	Arg	Leu	Ser	Gln	Arg	Phe	Pro	
	210						215					220					
15	Lys	Ala	Glu	Phe	Ala	Glu	Val	Ser	Lys	Leu	Val	Thr	Asp	Leu	Thr	Lys	
	225					230					235					240	
	Val	His	Thr	Glu	Cys	Cys	His	Gly	Asp	Leu	Leu	Glu	Cys	Ala	Asp	Asp	
20					245					250					255		
	Arg	Ala	Asp	Leu	Ala	Lys	Tyr	Ile	Cys	Glu	Asn	Gln	Asp	Ser	Ile	Ser	
				260					265					270			
25	Ser	Lys	Leu	Lys	Glu	Cys	Cys	Glu	Lys	Pro	Leu	Leu	Glu	Lys	Ser	His	
			275					280					285				
	Cys	Ile	Ala	Glu	Val	Glu	Asn	Asp	Glu	Met	Pro	Ala	Asp	Leu	Pro	Ser	
	290						295					300					
30	Leu	Ala	Ala	Asp	Phe	Val	Glu	Ser	Lys	Asp	Val	Cys	Lys	Asn	Tyr	Ala	
	305					310					315					320	
	Glu	Ala	Lys	Asp	Val	Phe	Leu	Gly	Met	Phe	Leu	Tyr	Glu	Tyr	Ala	Arg	
35					325					330					335		
	Arg	His	Pro	Asp	Tyr	Ser	Val	Val	Leu	Leu	Leu	Arg	Leu	Ala	Lys	Thr	
				340					345					350			
40	Tyr	Glu	Thr	Thr	Leu	Glu	Lys	Cys	Cys	Ala	Ala	Ala	Asp	Pro	His	Glu	
			355					360					365				
	Cys	Tyr	Ala	Lys	Val	Phe	Asp	Glu	Phe	Lys	Pro	Leu	Val	Glu	Glu	Pro	
	370						375					380					
45	Gln	Asn	Leu	Ile	Lys	Gln	Asn	Cys	Glu	Leu	Phe	Glu	Gln	Leu	Gly	Glu	
	385					390					395					400	
	Tyr	Lys	Phe	Gln	Asn	Ala	Leu	Leu	Val	Arg	Tyr	Thr	Lys	Lys	Val	Pro	
50					405					410					415		
	Gln	Val	Ser	Thr	Pro	Thr	Leu	Val	Glu	Val	Ser	Arg	Asn	Leu	Gly	Lys	
				420					425					430			
55	Val	Gly	Ser	Lys	Cys	Cys	Lys	His	Pro	Glu	Ala	Lys	Arg	Met	Pro	Cys	
			435					440					445				
	Ala	Glu	Asp	Tyr	Leu	Ser	Val	Val	Leu	Asn	Gln	Leu	Cys	Val	Leu	His	
	450						455					460					
60																	

	Glu	Lys	Thr	Pro	Val	Ser	Asp	Arg	Val	Thr	Lys	Cys	Cys	Thr	Glu	Ser
	465					470					475				480	
5	Leu	Val	Asn	Arg	Arg	Pro	Cys	Phe	Ser	Ala	Leu	Glu	Val	Asp	Glu	Thr
					485					490					495	
	Tyr	Val	Pro	Lys	Glu	Phe	Asn	Ala	Glu	Thr	Phe	Thr	Phe	His	Ala	Asp
				500					505					510		
10	Ile	Cys	Thr	Leu	Ser	Glu	Lys	Glu	Arg	Gln	Ile	Lys	Lys	Gln	Thr	Ala
			515					520					525			
	Leu	Val	Glu	Leu	Val	Lys	His	Lys	Pro	Lys	Ala	Thr	Lys	Glu	Gln	Leu
			530				535					540				
15	Lys	Ala	Val	Met	Asp	Asp	Phe	Ala	Ala	Phe	Val	Glu	Lys	Cys	Cys	Lys
	545					550					555					560
	Ala	Asp	Asp	Lys	Glu	Thr	Cys	Phe	Ala	Glu	Glu	Gly	Lys	Lys	Leu	Val
20					565					570					575	
	Ala	Ala	Ser	Gln	Ala	Ala	Leu	Gly	Leu	Pro	Gly	Pro	Pro	Pro	Gly	Pro
				580					585					590		
25	Pro	Arg	Val	Ser	Pro	Asp	Pro	Arg	Ala	Glu	Leu	Asp	Ser	Thr	Val	Leu
			595					600					605			
	Leu	Thr	Arg	Ser	Leu	Leu	Ala	Asp	Thr	Arg	Gln	Leu	Ala	Ala	Gln	Leu
		610					615					620				
30	Arg	Asp	Lys	Phe	Pro	Ala	Asp	Gly	Asp	His	Asn	Leu	Asp	Ser	Leu	Pro
	625					630					635					640
	Thr	Leu	Ala	Met	Ser	Ala	Gly	Ala	Leu	Gly	Ala	Leu	Gln	Leu	Pro	Gly
35					645					650					655	
	Val	Leu	Thr	Arg	Leu	Arg	Ala	Asp	Leu	Leu	Ser	Tyr	Leu	Arg	His	Val
				660					665					670		
40	Gln	Trp	Leu	Arg	Arg	Ala	Gly	Gly	Ser	Ser	Leu	Lys	Thr	Leu	Glu	Pro
			675					680					685			
	Glu	Leu	Gly	Thr	Leu	Gln	Ala	Arg	Leu	Asp	Arg	Leu	Leu	Arg	Arg	Leu
		690					695					700				
45	Gln	Leu	Leu	Met	Ser	Arg	Leu	Ala	Leu	Pro	Gln	Pro	Pro	Pro	Asp	Pro
	705					710					715					720
	Pro	Ala	Pro	Pro	Leu	Ala	Pro	Pro	Ser	Ser	Ala	Trp	Gly	Gly	Ile	Arg
50					725					730					735	
	Ala	Ala	His	Ala	Ile	Leu	Gly	Gly	Leu	His	Leu	Thr	Leu	Asp	Trp	Ala
				740					745					750		
55	Val	Arg	Gly	Leu	Leu	Leu	Leu	Lys	Thr	Arg	Leu					
			755					760								

Claims

1. An albumin fusion protein comprising IL-11, or a fragment or variant thereof, and albumin, or a fragment or variant thereof.
- 5 2. The albumin fusion protein of claim 1, wherein the IL-11 is human IL-11.
3. An albumin fusion protein according to Claim 1 or 2, comprising an albumin fused to IL-11.
- 10 4. The albumin fusion protein of any preceding claim, wherein the IL-11 is human IL-11.
5. The albumin fusion protein of claim 1 wherein the albumin has the ability to prolong the *in vivo* half-life of IL-11, or a fragment or variant thereof, compared
15 to the *in vivo* half-life of IL-11, or a fragment or variant thereof, in an unfused state.
6. The protein of claim 5, whereby the half-life of said albumin-fused IL-11 is extended at least 5-fold over the half-life of the IL-11 lacking the linked
20 albumin.
7. The protein of claim 6, whereby the half-life of said albumin-fused IL-11 is extended at least 10-fold over the half-life of the IL-11 lacking the linked
25 albumin.
8. The protein of claim 7 whereby the half-life of said albumin-fused IL-11 is extended at least 50-fold over the half-life of the IL-11 lacking the linked albumin.
9. The albumin fusion protein of claim 1 wherein IL-11, or a fragment or
30 variant thereof, is fused to the N-terminus of albumin, or the N-terminus of the fragment or variant of albumin.

10. The albumin fusion protein of claim 1 wherein IL-11, or a fragment or variant thereof, is fused to the C-terminus of albumin, or the C-terminus of the fragment or variant of albumin.
- 5 11. The albumin fusion protein of claim 1 wherein IL-11, or a fragment or variant thereof, is fused to an internal region of albumin, or an internal region of a fragment or variant of albumin.
12. The albumin fusion protein of claim 1 wherein IL-11, or a fragment or
10 variant thereof, is separated from the albumin or the fragment or variant of albumin by a linker.
13. The albumin fusion protein of any preceding claim wherein the *in vitro* biological activity of the IL-11, or fragment or variant thereof, fused to albumin,
15 or fragment or variant thereof, is greater than the *in vitro* biological activity IL-11, or fragment or variant thereof, in an unfused state.
14. The albumin fusion protein of any preceding claim wherein the *in vivo* biological activity of IL-11, or fragment or variant thereof, fused to albumin, or
20 fragment or variant thereof, is greater than the *in vivo* biological activity of IL-11, or fragment or variant thereof, in an unfused state.
15. A nucleic acid molecule comprising a polynucleotide sequence encoding the albumin fusion protein of any preceding claim.
- 25 16. A vector comprising the nucleic acid molecule of claim 15.
17. A host cell containing the nucleic acid molecule of claim 15.
- 30 18. A method for manufacturing an albumin fusion protein of any of claims 1-14, the method comprising (a) providing a nucleic acid comprising a nucleotide sequence encoding the albumin fusion protein expressible in a cell or organism;

(b) expressing the nucleic acid in the cell or organism to form an albumin fusion protein; and (c) purifying the albumin fusion protein.

19. The method of claim 18 wherein the albumin fusion protein is expressed in
5 a yeast.

20. The method of Claim 19 wherein the yeast is glycosylation deficient.

21. The method of claim 19 wherein the yeast is glycosylation competent.
10

22. The method of Claim 18 wherein the albumin fusion protein is expressed
in a mammalian cell in cell culture.

23. A composition comprising the albumin fusion protein of any one of claims
15 1-14 and a carrier.

24. A pharmaceutical composition comprising an effective amount of the
albumin fusion protein of any one of claims 1-14 and a pharmaceutically
acceptable carrier or excipient.
20

25. A method for minimizing a side effect associated with the treatment of a
mammal with IL-11 comprising administering an albumin-fused IL-11 to said
mammal.

25 26 A method according to Claim 25 wherein the mammal is suffering from a
bowel disorder and the side effect is weight loss, rectal bleeding or diarrhoea.

27 A method of increasing weight in a mammal suffering from a bowel disease
causing weight loss, the method comprising administering an albumin-fused IL-11
30 to said mammal.

28. A method of treating a disease or disorder in a patient, comprising the step of administering an effective amount of the albumin fusion protein of any of claims 1-14.
- 5 29. A method of treating a patient, comprising the step of administering an effective amount of the albumin fusion protein of any of claims 1-14.
30. A method of extending the *in vivo* half-life of IL-11, or a fragment or variant thereof, comprising the step of fusing IL-11, or fragment or variant
10 thereof, to albumin or a fragment or variant of albumin sufficient to extend the *in vivo* half-life of IL-11, or fragment or variant thereof, compared to the *in vivo* half-life of IL-11, or fragment or variant thereof, in an unfused state.
31. A method for extending the half-life of IL-11 in a mammal, the method
15 comprising linking said IL-11 to an albumin to form an albumin-fused IL-11 and administering said albumin-fused IL-11 to said mammal, whereby the half-life of said albumin-fused IL-11 is extended at least 2-fold over the half-life of IL-11 lacking the linked albumin.
- 20 32. A method for preventing or treating thrombocytopenia in a mammal, the method comprising administering an albumin-fused IL-11 to said mammal.
33. A method for minimizing a side effect associated with the treatment of a mammal with IL-11, the method comprising administering an albumin-fused IL-
25 11 to said mammal.

Figures


Figure 1: Mean IL-11 concentrations $\pm$ SD, following i.v. application

Figure 2: Mean IL-11 concentrations $\pm$ SD, following s.c. application




Figure 3: Mean IL-11 concentrations $\pm$ SD after intravenous administration of rhIL-11 or N-terminal IL-11-albumin fusion to rats




Figure 4: Mean IL-11 concentrations $\pm$ SD after subcutaneous administration of rhIL-11 or N-terminal IL-11-albumin fusion to rats




Figure 5: Course of platelet levels after treatment of naive rats with IL-11 and IL-11 fusions




Figure 6: Course of platelet levels after treating rats under chemotherapy with IL-11 and IL-11 fusions




Figure 7: Development of body weight (in % of baseline) after IL-11 and IL-11 fusions application in a mouse model for IBD




Figure 8: Visual observation score (diarrhoea and gross rectal bleeding) after IL-11 and IL-11 fusions application in a mouse model for IBD




Figure 9: Colon length after IL-11 and IL-11 fusions application in a mouse model for IBD




Figure 10: Histological disease score after IL-11 and IL-11 fusions application in a mouse model for IBD





Figure 11: 12% Gradient SDS Non-Reducing Gel and Western Blots

COLLOIDAL BLUE GEL


1 2 3 4 5 6 7 8 9 10



<u>Lane</u>	<u>Sample</u>	<u>Load</u>
1.	-	-
2.	Magic Marker	-
3.	-	-
4.	C Terminal IL11 (1721#11)	1µg
5.	N Terminal IL11 (Type A)	1µg
6.	N Terminal IL11 (Type B)	1µg
7.	HSA	1µg
8.	IL11 Standard (IL11 blot only)	50ng
9	-	-
10	SPT9901	100ng

ANTI-IL11 WESTERN BLOT

4 5 6 7 8 9 10



ANTI-HSA WESTERN BLOT

4 5 6 7 8 9 10

