




- (72) ANDERSON, DENISE, CH  
(72) FRATER, GEORG, CH  
(71) GIVAUDAN-ROURE (INTERNATIONAL) S.A., CH  
(51) Int.Cl.<sup>7</sup> C07C 69/675, C11D 3/50, A61K 7/46, C07C 69/007,  
C07C 311/39, A61K 7/32, C07C 235/06, C07C 235/02  
(30) 1997/06/21 (97110195.1) EP  
(54) **COMPOSES DE PRECURSEUR DE PARFUM**  
(54) **FRAGRANCE PRECURSOR COMPOUNDS**



(57) L'invention concerne des composés de la formule (I) dans laquelle n désigne 1, 2 ou 3 et R<sup>1</sup> à R<sup>6</sup> représentent indépendamment des radicaux alkyle, alcényle, alcynyle, cycloalkyle, cycloalcényle, ou aromatiques, ramifiés ou non ramifiés, substitués ou non substitués, ou de l'hydrogène, ces radicaux pouvant contenir en outre un ou plusieurs groupes - O - et/ou - (a), un ou deux noyaux pouvant être formés par combinaison de R<sup>1</sup> à R<sup>6</sup> respectivement et ce/ces noyaux pouvant être substitués par un groupe alkyle dans lequel soit X représente O et R<sup>7</sup> représente un radical d'un alcool ou d'un phénol

(57) Compounds having formula (I) in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or more - O - and/or (a) - groups, whereby one or two rings can be built by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group, in which X is either O and R<sup>7</sup> represents a radical of an alcohol or phenol R<sup>7</sup>OH, or X is N and R<sup>7</sup> represents the





(21) (A1) **2,293,252**  
(86) 1998/06/22  
(87) 1998/12/30

$R^7$ OH soit X représente N et  $R^7$  représente le radical d'une amine  $R^{7'}R^{7''}NH$ ,  $R^{7'}$  et  $R^{7''}$  représentant indépendamment des radicaux alkyle, alcényle, alcynyle, cycloalkyle, cycloalcényle ou aromatiques ou alors  $R^{7'}R^{7''}$  peuvent désigner un hydrogène, auquel cas l'amine est une amine parfumée ou l'amine renferme plus de 9 atomes de C, auquel cas  $R^7$  de l'alcool ou du phénol et  $R^{7'}$  et/ou  $R^{7''}$  de l'amine respectivement peuvent contenir également au moins un reste  $C(OH)R^1R^2-CR^3R^4-(CR^5R^6)_n-CO-$  de la formule (I); ces composés s'utilisent comme précurseurs dans la diffusion de composés odorants et/ou antibactériens dans des compositions cosmétiques, des produits cosmétiques, des désodorisants, des agents de nettoyage de surfaces dures ou des produits de lessive.

radical of an amine  $R^{7'}R^{7''}NH$ , whereby  $R^{7'}$  and  $R^{7''}$  represent independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either  $R^{7'}R^{7''}$  may be hydrogen, whereby the amine is a fragrant amine or the amine has more than 9 C atoms, whereby  $R^7$  of the alcohol or phenol and  $R^{7'}$  and/or  $R^{7''}$  of the amine, respectively, may further contain at least one remaining part  $C(OH)R^1R^2-CR^3R^4-(CR^5R^6)_n-CO-$  of formula (I), are useful as precursors for the delivery of odoriferous and/or antibacterial compounds in cosmetic compositions, cosmetic products, air fresheners, hard surface cleaners or laundry products.





PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)


<p>(51) International Patent Classification <sup>6</sup> : C07C 69/675, 69/007, C11D 3/50, A61K 7/32, 7/46, C07C 235/02, 235/06, 311/39</p>	A1	<p>(11) International Publication Number: <b>WO 98/58899</b> (43) International Publication Date: 30 December 1998 (30.12.98)</p>
<p>(21) International Application Number: PCT/EP98/03772 (22) International Filing Date: 22 June 1998 (22.06.98) (30) Priority Data: 97110195.1 21 June 1997 (21.06.97) EP (34) Countries for which the regional or international application was filed: AT et al. (71) Applicant (for all designated States except US): GIVAUDAN-ROURE (INTERNATIONAL) S.A. [CH/CH]; CH-1412 Vernier (CH). (72) Inventors; and (75) Inventors/Applicants (for US only): ANDERSON, Denise [US/CH]; 5 Hottingerstrasse, CH-8032 Zürich (CH). FRATER, Georg [CH/CH]; 61 Turmstrasse, CH-8400 Winterthur (CH). (74) Common Representative: GIVAUDAN-ROURE (INTERNATIONAL) S.A.; P.O. Box 3255, CH-4002 Basle (CH).</p>	<p>(81) Designated States: AU, BR, CA, JP, SG, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p>	

(54) Title: FRAGRANCE PRECURSOR COMPOUNDS

## (57) Abstract

Compounds having formula (I) in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or more - O - and

/or (a) - groups, whereby one or two rings can be built by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group, in which X is either O and R<sup>7</sup> represents a radical of an alcohol or phenol R<sup>7</sup>OH, or X is N and R<sup>7</sup> represents the radical of an amine R<sup>7</sup>R<sup>7'</sup>NH, whereby R<sup>7</sup> and R<sup>7'</sup> represent independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either R<sup>7</sup>R<sup>7'</sup> may be hydrogen, whereby the amine is a fragrant amine or the amine has more than 9 C atoms, whereby R<sup>7</sup> of the alcohol or phenol and R<sup>7</sup> and/or R<sup>7'</sup> of the amine, respectively, may further contain at least one remaining part C(OH)R<sup>1</sup>R<sup>2</sup>-CR<sup>3</sup>R<sup>4</sup>-(CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub>-CO- of formula (I), are useful as precursors for the delivery of odoriferous and/or antibacterial compounds in cosmetic compositions, cosmetic products, air fresheners, hard surface cleaners or laundry products.



## Fragrance Precursor Compounds

The invention relates to fragrance and/or antibacterial precursor compounds. In particular, the invention relates to the use of compounds which can act as fragrance and/or antibacterial precursors in cosmetic products such as deodorants and antiperspirants and in laundry products such as detergents and fabric softeners or in air fresheners or hard surface cleaners. These compounds are normally odourless or nearly so, but upon contacting the skin as for example, in skin care compositions or in personal care compositions, produce fragrances. The compounds also produce fragrances when used in the presence of enzymes such as lipases and proteases, e.g. used in (laundry) detergents and fabric softeners, thus providing a prolongation of the fabric scenting effect. The compounds can also produce fragrances when heated.


A principal strategy currently employed in imparting odours to consumer products is the admixing of the fragrance directly into the product. There are however, several drawbacks to this strategy. The fragrance material can be too volatile, resulting in fragrance loss during manufacturing, storage, and use. Many fragrance materials are also unstable over time. This again results in loss during storage.

In some cases, fragrances are microencapsulated or treated with cyclodextrins to form inclusion complexes to help decrease volatility and improve stability. However, these methods are for a number of reasons often not successful. In addition, cyclodextrins can be too expensive.

In many consumer products it is desirable for the fragrance to be released slowly over time. Microencapsulation and cyclodextrins have been used to provide slow-release properties, however, they are subject to the same limitations as above.

The present invention now provides compounds which show a low level of odour, or are even odourless, and can be cleaved under activating conditions, e.g. by heat or by an acid, a base, bacteria or enzymes, to give fragrant molecules. So, for example, the compounds of the invention are odourless prior to application to the skin, but release fragrant molecules after application to the skin, that is, they provide a delayed release of the fragrance, in particular to the skin in the axilla which is the result of the cleavage by bacteria. The compounds of the present invention also release fragrant molecules when used in the presence of enzyme-containing products and, this way, provide a prolongation of the fabric scenting effect. The compounds of the present invention also release fragrant molecules when heated.

The compounds under consideration are compounds of the formula



15


I

in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or more - O - and /or - C(=O) - groups, whereby one or two rings can be build by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group, in which X is either O and R<sup>7</sup> represents a radical of an alcohol or phenol R<sup>7</sup>OH or X is N and R<sup>7</sup> represents the radical of an amine R<sup>7'</sup>R<sup>7''</sup>NH, whereby R<sup>7'</sup> and R<sup>7''</sup> represent independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either R<sup>7'</sup> or R<sup>7''</sup> may be hydrogen, whereby the amine is a fragrant amine or the amine has more than 9 C atoms, whereby R<sup>7</sup> of the alcohol or phenol and R<sup>7'</sup> and/or R<sup>7''</sup> of the amine, respectively, may further contain at least one remaining part

C(OH)R<sup>1</sup>R<sup>2</sup>-CR<sup>3</sup>R<sup>4</sup>-(CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub>-CO- of formula I.

30

Thus, the precursor compounds are esters of formula Ia




5 in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or more - O - and /or - C(=O) - groups, R<sup>7</sup> represents a radical of an alcohol or phenol R<sup>7</sup>OH, whereby one or two rings can be build by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group, whereby R<sup>7</sup> may further contain the remaining part C(OH)R<sup>1</sup>R<sup>2</sup>-CR<sup>3</sup>R<sup>4</sup>-(CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub>-CO- of formula Ia,

10

or amides of formula Ib

15



in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or more - O - and /or - C(=O) - groups, whereby one or two rings can be build by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group, and R<sup>7</sup> represents a radical of an amine R<sup>7'</sup>R<sup>7''</sup>NH, whereby R<sup>7'</sup> and R<sup>7''</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either R<sup>7'</sup> or R<sup>7''</sup> may be hydrogen, whereby the amine is a fragrant amine or the amine has more than 9 C atoms, whereby R<sup>7'</sup> and/or R<sup>7''</sup> may further contain at least one remaining part C(OH)R<sup>1</sup>R<sup>2</sup>-CR<sup>3</sup>R<sup>4</sup>-(CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub>-CO- of formula Ib.

20

25

With respect to the precursor compounds of formula Ia specifically the invention is related to those compounds in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein

5 these radicals may in addition contain one or more - O - and /or -  $\overset{\text{O}}{\parallel}{\text{C}}$  - groups, R<sup>7</sup> represents a radical of a fragrant alcohol R<sup>7</sup>OH, whereby one or two rings can be build by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group.

10 Further characteristics and advantages of the invention are described by claims 4 to 29 and by the following description and examples.

The compounds of formula I are not limited to any particular stereoisomers, all possible stereoisomers (enantiomers, diastereomers) and all  
15 mixtures are thus included within the scope of formula I.

The compounds of formula I may preferably be used as sustained release odorants but also to mask or attenuate undesirable odours or to provide additional odours not initially present in consumer products, i.e.  
20 laundry detergents, fabric softeners, fabric softeners sheets, hard surface cleaners, automatic dishwasher detergents and (other) enzyme-containing consumer products. Additional applications include cosmetic products destined for application to human skin such as underarm deodorants or antiperspirants or other deodorants contacting the body, or in hand lotions, baby powders,  
25 baby lotions, ointments, foot products, facial cleansers, body wipes, facial make-up, colognes, after-shave lotions, shaving creams, etc. Further additional applications include ironing treatments and air fresheners dispensed via heat.

The compounds of formula I may be used individually in an amount  
30 effective to enhance the characteristic odour of a material. More commonly, however, the compounds are mixed with other fragrance components in an amount sufficient to provide the desired odour characteristics.

The amount required to produce the desired, overall effect varies  
35 depending upon the particular compounds of formula I chosen, the product in which it will be used, and the particular effect desired.

For example, depending upon the selection and concentration of the compound(s) of formula I chosen, when added either singly or as a mixture, e.g. to a deodorant, laundry product, hard surface cleaner or air freshener composition at levels ranging from about 0.1 to about 10 % by weight, or most preferred about 0.25 to about 4 % by weight, an odorant, i.e. an odoriferous alcohol and an odoriferous lactone in an organoleptically effective amount is released when the product is used. These newly formed odorants serve to enhance the odour of the fragrance.

The compounds of formula I can accordingly be used in the manufacture of odorant compositions used in the preparation of air fresheners, cosmetic and laundry products e.g. deodorants, antiperspirants, laundry detergents, fabric softeners, hard surface cleaners, and as is evident from the above compilation, a broad range of known odorants or odorant mixtures can be used. In the manufacture of such compositions the known odorants or odorant mixtures set forth above can be used according to methods known to a person skilled in the art, normally a perfumer, or described e.g. in W.A. Poucher, *Perfumes, Cosmetics, Soaps*, 2, 7th Edition, Chapman and Hall, London 1974.

Concerning the esters suitable examples of odoriferous alcohols  $R^7OH$  are alcohols or phenols such as listed in Tabel 1.

Table 1

amyl alcohol  
 hexyl alcohol\*  
 2-hexyl alcohol\*  
 heptyl alcohol\*  
 octyl alcohol\*  
 nonyl alcohol\*  
 decyl alcohol\*  
 undecyl alcohol\*  
 lauryl alcohol\*  
 myristic alcohol  
 3-methyl-but-2-en-1-ol\*  
 3-methyl-1-pentanol  
 cis-3-hexenol\*\*  
 cis-4-hexenol\*  
 3,5,5-trimethyl hexanol  
 3,4,5,6,6-pentamethylheptan-2-ol\*  
 citronellol\*\*  
 geraniol\*\*

- oct-1-en-3-ol
- 2,5,7-trimethyl octan-3-ol
- 2-cis-3,7-dimethyl-2,6-octadien-1-ol
- 6-ethyl-3-methyl-5-octen-1-ol\*
- 5 3,7-dimethyl-oct-3,6-dienol\*
- 3,7-dimethyloctanol\*\*
- 7-methoxy-3,7-dimethyl-octan-2-ol\*
- cis-6-nonenol\*
- 5-ethyl-2-nonanol
- 10 6,8-dimethyl-2-nonanol\*
- 2,2,8-trimethyl-7 (8)-nonene-3-ol
- nona-2,6-dien-1-ol
- 4-methyl-3-decen-5-ol\*\*
- dec-9-en-1-ol
- 15 benzylalcohol
- 2-methyl undecanol
- 10-undecen-1-ol
- 1-phenyl ethanol\*
- 2-phenyl ethanol\*\*
- 20 2-methyl-3-phenyl-3-propenol
- 2-phenyl propanol\*
- 3-phenyl propanol\*
- 4-phenyl-2-butanol
- 2-methyl-5-phenyl pentanol\*
- 25 2-methyl-4-phenyl-pentanol\*
- 3-methyl-5-phenyl-pentanol\*
- 2-(2-methylphenyl)-ethanol\*
- 4-(1-methylethyl)benzene methanol
- 4-(4-hydroxyphenyl)butan-2-one\*
- 30 2-phenoxy ethanol\*
- 4-(1-methylethyl)-2-hydroxy-1-methyl benzene
- 2-methoxy-4-methyl phenol
- 4-methyl phenol
- anisic alcohol\*
- 35 p-tolyl alcohol\*
- cinnamic alcohol\*\*
- vanillin\*
- ethyl vanillin\*
- eugenol\*\*
- 40 isoeugenol\*\*
- thymol
- anethol\*
- decahydro 2-naphthalenol
- borneol\*
- 45 cedrenol\*
- farnesol\*
- fenchyl alcohol\*
- menthol\*

- 3,7,11-trimethyl-2,6,10-dodecatrien-1-ol  
 alpha ionol\*  
 tetrahydro ionol\*  
 2-(1,1-dimethylethyl)cyclohexanol\*  
 5 3-(1,1-dimethylethyl)cyclohexanol\*  
 4-(1,1-dimethylethyl)cyclohexanol\*  
 4-isopropyl cyclohexanol  
 6,6-dimethyl-bicyclo [3.3.1]hept-2-ene-2-ethanol  
 6,6-dimethyl-bicyclo [3.1.1]hept-2-ene-methanol\*  
 10 p-menth-8-en-3-ol\*  
 3,3,5-trimethyl cyclohexanol  
 2,4,6-trimethyl-3-cyclohexenyl-methanol\*  
 4-(1-methylethyl)cyclohexyl-methanol\*  
 4-(1,1-dimethylethyl)cyclohexanol  
 15 2-(1,1-dimethylethyl)-cyclohexanol  
 2,2,6-trimethyl-alpha-propyl cyclohexane propanol\*  
 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol\*  
 3-methyl-5-(2,2,3-trimethylcyclopentyl-3-enyl)pent-4-en-2-ol\*\*  
 2-ethyl-4(2,2,3-trimethylcyclopentyl-3-enyl)but-2-en-1-ol\*\*  
 20 4-(5,5,6-trimethylbicyclo[2.2.1] hept-2-yl)-cyclohexanol\*  
 2-(2-methylpropyl)-4-hydroxy-4-methyl-tetrahydropyran\*  
 2-cyclohexyl propanol\*  
 2-(1,1-dimethylethyl)-4-methyl cyclohexanol\*  
 1-(2-tert-butyl-cyclohexyloxy)-2-butanol\*  
 25 1-(4-isopropyl-cyclohexyl)-ethanol\*  
 linalool \*\*  
 2,6-dimethyl-heptan-2-ol\*\*  
 2,6-dimethyl-oct-7-en-2-ol\*\*

30 whereby one asterisk indicates the preferred alcohols and phenols and two asterisks indicate the more preferred alcohols and phenols.

It is a matter of course, that it is not possible to give a complete list of the odoriferous alcohols and phenols  $R^7OH$  which are liberated as a result of  
 35 the desired cleavage of the compounds of formula I as mentioned above, especially by heat, enzymes or by bacteria, and which alcohols are then capable of imparting agreeable odours. The skilled artisan is, however, quite aware of those alcohols and phenols, which provide a positive contribution to the fragrance compositions.

40

But the alcohol or phenol constituting the radical  $R^7$  can also be a non-odoriferous one or a polyalcohol. Thus, the radical  $R^7$  can be derived from as well a simple alcohol like e.g. methanol or ethanol as also from a fatty alcohol or a compound such as 7-hydroxy-4-methyl coumarin.

Examples of polyalcohols constituting the radical R<sup>7</sup> in the compounds of formula I, or more specifically of formula Ia, are:

diols such as: diethylene glycol, propylene glycol, triethylene glycol, N-  
5 butyldiethanol amine, 1,3- bis -(4-hydroxy butyl)-1,1,3,3 tetramethyl-  
disiloxane, 4,4'-bicyclohexyldiol;  
triols such as: glycerol, 1,3,5-cyclohexanetriol;  
sugars such as: furanoside and pyranoside sugars such as glucose, fructose;  
polymers such as: hydroxyethylcellulose, hydroxypropylcellulose.

10

Compounds of formula I upon cleavage may also generate antimicrobial compounds. Examples of these compounds are e.g. presented by J.J. Kabara, Cosmet. Sci. Technol. Ser. (16) 1997, p 181-208, especially in Table 8.6.

15

Of course, the afore mentioned alcohols, phenols and antimicrobial compounds can serve mutually as fragrances and antimicrobial compounds, respectively. A person of skill in the art is well aware of these interrelationships and can make use thereof to solve a specific problem by using the precursors of the present invention.

20

The compounds of formula Ia are virtually odourless under room temperature and atmospheric conditions, i.e. about 10 to about 40 degrees Celsius and about 20 to 100 % relative humidity. However, when applied to the body or when used in an application in the presence of enzymes such as  
25 lipases and proteases, or when heated they undergo a transformation in which the alcohol or phenol and lactone are released.

The compounds of formula Ia, upon cleavage, provide alcohols or phenols and lactones having organoleptic properties and therefore permit the  
30 development of methods useful in enhancing the odour of consumer products.

Suitable examples of such lactones are listed in Table 2.

35

**Table 2**

6-methyl-pyran-2-one

5-heptyldihydro-2(3H)-furanone\*

- 5-pentyldihydro-2(3H)-furanone\*
- 5-(3-hexenyl)dihydro-5-methyl-(Z)-2(3H)-furanone
- 5-hexyldihydro-5-methyl-2(3H)-furanone
- 5-hexyldihydro-2(3H)-furanone\*
- 5 5-octyldihydro-2(3H)-furanone
- 8-(1-methylethyl)-1-oxaspiro[4.5]-decan-2-one\*
- 8-methyl-1-oxaspiro[4.5]-decan-2-one
- 8-ethyl-1-oxaspiro[4.5]-decan-2-one
- 5-(1,5-dimethyl-4-hexenyl)dihydro-2(3H)-furanone
- 10 2-oxo-5-butyl-tetrahydrofuran\*
- 4-methyl-5-pentyl-dihydro-2(3H)-furan-2-one
- 5-hexyldihydro-5-methyl-2(3H)-furanone
- dihydro-5-methyl-5-vinyl-2(3H)-furanone
- octahydro-2H-1-benzopyran-2-one
- 15 tetrahydro-6-pentyl-2H-pyran-2-one
- tetrahydro-6-hexyl-2H-pyran-2-one
- tetrahydro-6-heptyl-2H-pyran-2-one
- tetrahydro-6-(3-pentenyl)-(E)-2H-pyran-2-one
- tetrahydro-6-(2-pentenyl)-(Z)-2H-pyran-2-one

20

whereby the asterisks indicate the preferred lactones. Table 2 is not complete but shows only representative lactones with positive effect in connection with the present invention. A person skilled in the art is, however, quite aware to further lactones which provide a positive contribution to the enhancement of

25 the odour of consumer products.

The compounds of formula Ia can be prepared by using standard methods known to a person skilled in the art. For example, they can be prepared in two steps from keto acids. Esters of the keto acids may be

30 prepared by standard methods known to those skilled in the art, see Comprehensive Organic Chemistry, Derek Barton and W. David Ollis, eds., Vol. 2, 1979, pp.871-907. For example, esters are formed by the acid catalyzed reaction between a carboxylic acid and an alcohol. During the condensation water is usually removed.

35

Either protic or Lewis acids may be used. Some acids which may be used are p-toluenesulfonic acid, sulfuric acid, and pyridinium p-toluenesulfonate. A

variety of inert solvents may be used such as toluene, xylene, cyclohexane, and hexane.

In another method which may be used for the preparation of the  
5 compounds of formula Ia, an appropriate carboxylic acid and an appropriate alcohol react to form an ester when treated with N,N'-dicyclohexylcarbodiimide and 4-pyrrolidinopyridine, see e.g. the procedure of Hassner and Alexanian, *Tetrahedron Letters* 4475, (1978).

10 These esters can then be reduced to the compounds of formula I by using standard methods known to a person skilled in the art. Reagents for the transformation include sodium borohydride and lithium aluminum hydride.

15 Compounds of formula Ia can also be prepared directly from the corresponding alcohol and lactone, see e.g. *J. Org. Chem.* (1966), 485.

The most preferred precursor compounds according to the invention are esters of one alcohol out of the group citronellol, phenylethyl alcohol, geraniol and cis-3-hexanol, specifically compounds selected from the group consisting of  
20 4-hydroxy-decanoic acid 2-phenethyl ester, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-decanoic acid hex-3-enyl ester, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-6-enyl ester, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-  
25 hydroxy undecanoic acid 3,7-dimethyl-oct-2,6 dienyl ester, 4-hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hept-6-enyl ester, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-undecanoic acid phenethyl ester, 4-hydroxy-undecanoic acid hex-3-enyl ester,  
30 4-hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, 4-hydroxy-nonanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-nonanoic acid hex-3-enyl ester, 4-hydroxy-nonanoic acid phenethyl ester, 3-(1-hydroxy-4-isopropyl-cyclohexyl)-propionic acid 3,7-dimethyl-octa-2,6-dienyl ester and 4-hydroxy-undecanoic acid 2-(4-hydroxy-undecanoyloxy)-ethyl ester.

35

Concerning the second group of the precursor compounds according to the invention under activating conditions, the hydroxy amide will cleave to release a lactone and an amine. The lactone may have organoleptic properties.

There are basically three classes of amine products to build the amides according to the invention:

- non-odoriferous amines which can be primary or secondary amines and have more than nine carbon atoms. Preferable amines are those with an affinity to fiber and those used in cosmetic and laundry formulations,
- odoriferous amines,
- non-odoriferous amines substituted with an odoriferous group. For example, the amine may be substituted by a group such as  $-\text{CO}_2\text{R}$ ,  $-\text{OCO}_2\text{R}$  wherein R represents a radical of a fragrant alcohol or the enol form of a fragrant aldehyde or ketone, or any radical that can form another lactone.

Preferred examples of odoriferous amines are:

- 1-methyl-1-(4-methyl-3-cyclohexen-1-yl)ethyl anthranilic acid; benzopyrrole; 8,8-di(1H-indol-3-yl)-2,6-dimethyl-octane-2-ol; anthranilic acid allyl ester; anthranilic acid 1,5-dimethyl-1-vinyl-4-hexenyl ester; 2-amino-benzoic acid methyl ester\*; methyl anthranilic acid N-(2-methylpent-1-en-1-yl) ester; anthranilic acid phenylethyl ester\*; 2-methylamino-benzoic acid methyl ester\*;
- 6-methyltetrahydroquinoline; isobutyl N-methyl anthranilate; (Z)-3-hexenyl 2-aminobenzoate\*, whereby the asterisks indicate the more preferred odoriferous amides.

A wide variety of non-odoriferous amines can also be used for the preparation of the amides according to the invention. For example, a list of suitable primary and secondary cosmetic amines can be found in the 'Cosmetic Ingredient Handbook' edited by Joanne M. Nikitakis. Suitable surfactant amines can be found, for example, in 'Surfactants Europa' edited by Gordon L. Hollis. Amino acids such as glycine, leucine, tyrosine, serine, glutamic acid, aspartic acid, phenylalanine, alanine, lysine, arginine, histidine, cysteine, valine, proline, tryptophan, isoleucine, methionine, asparagine, glutamine and threonine may also be used.

The present invention is described in the following examples which are presented solely for the non-limiting purpose of further illustrating the invention.

**Example 1****(a) 4-Oxo-decanoic acid 3,7-dimethyl-oct-6-enyl ester**

5           According to Synthesis, 1987, 408, a solution of 20.0 g 4-oxo-1-decanoic acid 16.8 g citronellol extra and, 0.5 g p-toluenesulfonic acid in 150 ml of cyclohexane was refluxed in a flask equipped with a Dean-Stark trap for 3 hours. Then the reaction mixture was cooled, diluted with ether, washed with saturated NaHCO<sub>3</sub> and water. The organic phase was dried, filtered and  
10 evaporated to dryness. The resulting oil was purified by chromatography to yield 31.7 g of a yellow oil.

15           NMR (CDCl<sub>3</sub>)  $\delta$    5.04-5.19 (m, 1H), 4.11-4.17 (m, 2H), 2.68-2.79 (m, 2H),  
                          2.53-2.60 (m, 2H), 2.41-2.48 (t, 2H), 2.03-1.91 (q, 2H),  
                          1.75-1.44 (m, 6H), 1.43-1.08 (m, 12H), 0.92-0.84 (m, 6H)

20           (b) According to the same procedure, 5-oxo-dodecanoic acid 3,7-dimethyl-oct-6-enyl ester was prepared from 5-oxo-dodecanoic acid, citronellol and p-toluenesulfonic acid.

            (c) According to the same procedure, 4-oxo-decanoic acid hex-3-enyl ester was prepared from 4-oxo-decanoic acid, cis-3-hexenol and p-toluenesulfonic acid.

25           (d) According to the same procedure, 4-oxo-decanoic acid 2-phenethyl ester was prepared from 4-oxo-decanoic acid, 2-phenyl ethanol and p-toluenesulfonic acid.

30           (e) According to the same procedure, 4-oxo-nonanoic acid 3,7-dimethyl-oct-6-enyl ester was prepared from 4-oxo-nonanoic acid, citronellol and p-toluenesulfonic acid.

35           (f) According to the same procedure, 4-oxo-nonanoic acid phenethyl ester was prepared from 4-oxo-nonanoic acid, 2-phenyl ethanol and p-toluenesulfonic acid.

(g) According to the same procedure, 4-oxo-nonanoic acid hex-3-enyl ester was prepared from 4-oxo-nonanoic acid, cis-3-hexenol and p-toluenesulfonic acid.

5 (h) According to the same procedure, 4-oxo-undecanoic acid phenethyl ester was prepared from 4-oxo-undecanoic acid, 2-phenyl ethanol and p-toluenesulfonic acid.

10 (i) According to the same procedure, 4-oxo-undecanoic acid hex-3-enyl ester was prepared from 4-oxo-undecanoic acid, cis-3-hexenol and p-toluenesulfonic acid.

15 (j) According to the same procedure, 4-oxo-undecanoic acid 2-(4-oxo-undecanoyloxy)-ethyl ester was prepared from 4-oxo-undecanoic acid, ethylene glycol and p-toluenesulfonic acid.

### Example 2

20 (a) 4-Oxo-decanoic acid 3,7-dimethyl-oct-2,6-dienyl ester

A solution of 5.2 g 4-oxo-decanoic acid, 13.5 g geraniol, 17.5 g N,N'-dicyclohexyl-carbodiimide and 1.0 g 4-pyrrolidinopyridine in 250 ml of dichloromethane was stirred for 24 hours at room temperature. The precipitate  
25 was filtered off, the filtrate was diluted with ether, washed with aqueous hydrochloric acid, saturated NaHCO<sub>3</sub> and brine. The organic phase was dried, filtered and evaporated to dryness. The resulting oil-crystall mixture was purified by chromatography to yield 19.7 g of a colourless oil.

30 NMR (CDCl<sub>3</sub>)  $\delta$  5.13-5.04 (m, 1H), 4.15-4.02 (q, 2H), 3.60 (s, 1H), 2.49-2.37 (t, 2H), 2.06-1.92 (m, 2H), 1.60 (s, 6H), 1.56-1.17 (m, 12H), 0.93-0.85 (q, 3H)

35 (b) According to the same procedure, 5-oxo-decanoic acid 3,7-dimethyl-oct-2,6-dienyl ester was prepared from 5-oxo-decanoic acid and geraniol.

(c) According to the same procedure, 4-oxo-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 4-oxo-nonanoic acid and geraniol.

(d) According to the same procedure, 4-oxo-decanoic acid 1,1,5-trimethyl-hexyl ester was prepared from 4-oxo-decanoic acid and 2,6-dimethyl-heptan-2-ol.

5

(e) According to the same procedure, 4-oxo-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester was prepared from 4-oxo-decanoic acid and (+)-linalool.

(f) According to the same procedure, 4-oxo-decanoic acid 1,1,5-trimethyl-hept-6-enyl ester was prepared from 4-oxo-decanoic acid and 2,6-dimethyl-oct-7-en-2-ol.

(g) According to the same procedure, 4-oxo-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 4-oxo-undecanoic acid and geraniol.

(h) According to the same procedure, 4-oxo-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester was prepared from 4-oxoundecanoic acid and 2-benzyloxycarbonylamino-3-hydroxy-propionic acid benzyl ester.

### Example 3

#### (a) 4-Hydroxy-decanoic acid 3,7-dimethyl-oct-6-enyl ester

25

A solution of 2,0 g sodium borohydride in 30 ml of water was cooled to 5°C. A solution of 4-oxo-decanoic acid 3,7-dimethyl-oct-6-enyl ester in 75 ml of THF was added to the reaction during 12 minutes and the resulting reaction mixture was stirred at room temperature for 5 hours. Then the reaction mixture was diluted with ether, washed with saturated NaHCO<sub>3</sub>, brine and water. The organic phase was dried, filtered and evaporated to dryness. The resulting liquid was purified by chromatography to yield 7.6 g of a liquid.

(b) According to the same procedure, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-2,6-dienyl ester was prepared from 4-oxo-decanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, sodium borohydride and water.

35

(c) According to the same procedure, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-6-enyl ester was prepared from 5-oxo dodecanoic acid 3,7-dimethyl-oct-6-enyl ester, sodium borohydride and water.

5 (d) According to the same procedure, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester was prepared from 5-oxo dodecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, sodium borohydride and water.

10 (e) According to the same procedure, 4-hydroxy-decanoic acid hex-3-enyl ester was prepared from 4-oxo-decanoic acid hex-3-enyl ester, sodium borohydride and water.

15 (f) According to the same procedure, 4-hydroxy-decanoic acid 2-phenethyl ester was prepared from 4-oxo-decanoic acid 2-phenethyl ester, sodium borohydride and water.

20 (g) According to the same procedure, 4-hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester was prepared from 4-oxo-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester, sodium borohydride and water.

(h) According to the same procedure, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hept-6-enyl ester was prepared from 4-oxo-decanoic acid 1,1,5-trimethyl-hept-6-enyl ester, sodium borohydride and water.

25 (i) According to the same procedure, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester was prepared from 4-oxo-decanoic acid 1,1,5-trimethyl-hexyl ester, sodium borohydride and water.

30 (j) According to the same procedure, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester was prepared from 4-oxo-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, sodium borohydride and water.

35 (k) According to the same procedure, 4-hydroxy-undecanoic acid phenethyl ester was prepared from 4-oxo-undecanoic acid phenethyl ester, sodium borohydride and water.

(l) According to the same procedure, 4-hydroxy-undecanoic acid hex-3-enyl ester was prepared from 4-oxo-undecanoic acid hex-3-enyl ester, sodium borohydride and water.

40

(m) According to the same procedure, 4-hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 4-oxo-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, sodium borohydride and water.

5 (n) According to the same procedure, 4-hydroxy-nonanoic acid 3,7-dimethyl-oct-6-enyl ester was prepared from 4-oxo-nonanoic acid 3,7-dimethyl-oct-6-enyl ester, sodium borohydride and water.

10 (o) According to the same procedure, 4-hydroxy-nonanoic acid hex-3-enyl ester was prepared from 4-oxo-nonanoic acid hex-3-enyl ester, sodium borohydride and water.

15 (p) According to the same procedure, 4-hydroxy-nonanoic acid phenethyl ester was prepared from 4-oxo-nonanoic acid phenethyl ester, sodium borohydride and water.

#### Example 4

##### (a) 4-Hydroxy-undecanoic acid sodium salt

20

To a solution of 43.6 g sodium hydroxide in 150 ml of methanol heated to reflux, 200 g gamma-undecalactone were dropped in. After stirring 2 hours at reflux, the mixture was cooled to room temperature and evaporated to dryness. The resulting crystals were washed with hexane to yield 240 g white  
25 crystals.

NMR (CDCl<sub>3</sub>) δ 5.1-4.8 (br s, OH), 3.63-3.42 (m, 1H), 2.39-2.20 (t, 2H), 1.89-1.52 (m, 2H), 1.51-1.15 (m, 12H), 1.00-0.81 (t, 3H) ppm.

30

(b) According to the same procedure, 3-(1-hydroxy-4-isopropyl-cyclohexyl)-propionic acid sodium salt was prepared from 8-(1-methylethyl)-1-oxaspiro[4.5]-decan-2-one and sodium hydroxide.

#### Example 5

35

##### 1-Chloro-3,7-dimethyl-octa-2,6-diene

To a mixture of 170 g linalool and 20 mg bismuth(III)-oxide heated to 60 °C, 130.5 g trimethylchlorosilane were dropped in. Then the mixture was

cooled to room temperature and the organic layer was separated. The resulting oil was purified by distillation to yield 158.35 g of a colorless oil.

5 NMR (CDCl<sub>3</sub>) δ 5.56-5.35 (t, 1H), 5.18-4.99 (m, 1H), 4.16-4.02 (d, 2H), 2.26-1.91 (m, 4H), 1.89-1.45 (m, 9H) ppm.

### Example 6

#### 10 (a) 4-Hydroxy-undecanoic acid 3,7-dimethyl-octa-2,6-dienyl ester

A mixture of 155 g 1-chloro-3,7-dimethyl-octa-2,6-diene, 202 g 4-hydroxy-undecanoic acid sodium salt and 5 g tetrabutylammoniumbromide in 800 ml of dimethylformamide was heated to 50°C. After stirring for 24 hours, 15 the mixture was cooled to room temperature and filtered through Celite. The filtrate was diluted with ether, washed with water, 2N HCl, saturated sodium bicarbonate and brine. The organic phase was dried and evaporated to dryness. The resulting yellow oil was purified by wipe film distillation to yield 96.6 g of a yellow oil.

20 NMR (CDCl<sub>3</sub>) δ 5.42-5.37 (t, 1H), 5.16-5.01 (m, 1H), 4.65-4.53 (m, 2H), 3.69-3.52 (m, 1H), 2.60-2.22 (m, 2H), 2.20-1.95 (m, 4H), 1.89-1.12 (m, 24H), 1.02-0.78 (t, 3H) ppm.

25 (b) According to the same procedure, 3-(1-hydroxy-4-isopropyl-cyclohexyl)-propionic acid 3,7-dimethyl-octa-2,6-dienyl ester was prepared from 1-chloro-3,7-dimethyl-octa-2,6-diene, 3-(1-hydroxy-4-isopropyl-cyclohexyl)-propionic acid sodium salt and tetrabutylammoniumbromide.

### 30 Example 7

#### (a) 4-Hydroxy-undecanoic acid 2-(4-hydroxy-undecanoyloxy)-ethyl ester

To a solution of 21.54 g 4-oxo-undecanoic acid 2-(4-oxo-undecanoyloxy)-ethyl ester and a of bromocresol green in 150 ml of methanol, 6.34 g sodium cyanoborohydride was added. The resulting reaction mixture was stirred for 35 2.5 hours at room temperature, while dropping in 2N HCl to adjust the pH < 3.8. The solvent was then evaporated, the solid residue was dissolved in ether

and washed with water. The organic phase was dried and evaporated to dryness to yield the compound.

5 NMR (CDCl<sub>3</sub>) δ 4.35-4.20 (m, 4H), 3.87-3.53 (m, 2H), 2.54-2.39 (t, 4H),  
2.05-1.13 (m, 28H), 1.04-0.80 (t, 6H) ppm.

(b) According to the same procedure, 4-hydroxy-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester was prepared from 4-oxo-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester, sodium cyanoborohydride and methanol.  
10

### Example 8

Test cloth was washed with a lipase-containing detergent to which one or more delayed release fragrances had been added. Headspace analysis of the wet and dry laundry indicated the presence of the fragrant alcohols and lactones. The alcohol and lactone level was higher than when the test cloth was washed with a lipase-containing detergent to which one or more fragrant alcohols and lactones were added. The compounds in Examples 3, 6, 7, 12, 13 and 15 were tested in this way.  
15  
20

### Example 9

Test cloth was washed with a lipase-containing detergent and then a fabric softener, containing one or more delayed release fragrances, was added to the rinse cycle. Headspace analysis of the wet and dry laundry indicated the presence of the fragrant alcohols and lactones. The alcohol and lactone level was higher than when the test cloth was washed with a lipase-containing detergent and then a fabric softener, containing one or more fragrant alcohols and lactones, was added to the rinse cycle. The compounds in Examples 3, 6, 7, 12, 13 and 15 were tested in this way.  
25  
30

### Example 10

35 Axilla bacteria cultures containing 0.1 % precursor I were incubated for 20 hours at 30 °C. After filtration from the cells, the presence of the parent alcohol and lactone was in each case detected by headspace-GC techniques and/or the majority of an 18 member panel.

The same tests were carried out with inactivated cultures (85°/20 min). The odour of the parent alcohols and lactones could not be detected after incubation, excluding therefore a hydrolysis by the medium or the culture. The  
5 compounds in Examples 3, 6, 7, 12, 13 and 15 were tested in this way.

### Example 11

10 The following set forth examples for the use of the delayed release fragrances of the present invention in various products. The compounds in Examples 3, 6, 7, 12, 13, and 15 were tested in this way. The methods of forming the following compositions are well known to those skilled in the art. All formulations may contain additional ingredients known to those skilled in  
15 the art, e.g. colorants, opacifiers, buffers, antioxidants, vitamins, emulsifiers, UV absorbers, silicones and the like. All products can also be buffered to the desired pH. All values are % w/w.

20 <b>Deo-colognes</b>	<b>I</b>	<b>II</b>	<b>III</b>	<b>IV</b>
Delayed Release Fragrances	0.5	1.5	2.5	6.0
Fragrance	0.5	1.5	2.5	6.0
Triclosan (Ciba Geigy)	1.0	-	0.75	1.0
Alcohol to	100.0	100	100	100

25

### 30 **Deo-Sticks:**

#### **Antiperspirant**

Ethylene Glycol Monostearate	7.0
Shea butter	3.0
35 Neobee 1053 (PVO International)	12.0
Generol 122 (Henkel)	5.0
Kesscowax B (Akzo)	17.0
Dimethicone Dow Corning 345	35.0

	Aluminum Sesquichlorhydrate	20.0
	Delayed Release Fragrances	0.5
	Fragrance	0.5
<b>5</b>	<b>Antiperspirant</b>	
	Steary Alcohol	17.0
	Castor Wax	3.0
	Talc	5.0
	Aluminum Zirconium	
<b>10</b>	<b>Tetrachlorhydrate</b>	20.0
	Delayed Release Fragrances	1.0
	Fragrance	1.0
	Dimethicone Dow 245	to 100.0
<b>15</b>	<b>Clear Deodorant Stick</b>	
	Witconol APM	43.0
	Propylene Glycol	20.0
	Alcohol 39C	20.0
	Demin water	7.0
<b>20</b>	<b>Monamid 150ADD</b>	5.0
	Millithix 925	2.0
	Ottasept Extra	0.5
	Delayed Release Fragrances	0.75
	Fragrance	0.75
<b>25</b>	<b>Deodorant Stick</b>	
	Propylene Glycol	69.0
	Demin Water	21.8
	Triclosan	0.2
<b>30</b>	<b>Sodium Stearate</b>	8.0
	Delayed Release Fragrances	0.5
	Fragrance	0.5
	<b>Alcohol free Deodorant Stick</b>	
<b>35</b>	<b>PPG-3 Myristyl Ether (Witconol APM)</b>	36.0
	Propylene Glycol	36.0
	Demin Water	19.0
	Triclosan	0.25

	Sodium Stearate	7.75
	Delayed Release Fragrances	0.5
	Fragrance	0.5
<b>5</b>	<b>Antiperspirant Aerosol</b>	
	Absolute Ethanol	15.0
	Zirconium Aluminum	
	tetrachlorhydrate	5.0
	Bentone 38	1.5
<b>10</b>	Delayed Release Fragrances	0.75
	Fragrance	0.75
	S-31 Hydrocarbon propellant	to 100.0
	<b>Antiperspirant Pump</b>	
<b>15</b>	Demin water	57.5
	Aluminum Sesquichlorhydrate	20.0
	Triton X-102 (Union Carbide)	2.0
	Dimethyl Isosorbide (ICI)	20.0
	Delayed Release Fragrances	0.25
<b>20</b>	Fragrance	0.25
	<b>Roll-On</b>	
	Dimethicone DC 354 (Dow Corning)	69.0
	Bentone 38	10.0
<b>25</b>	Rezal 36 GP ( Reheis Chem. Co.)	20.0
	Delayed Release Fragrances	0.5
	Fragrance	0.5

In the above, the following components were used:

<b>30</b>	Triclosan-	5-chloro-2-(2,4-dichlorophenoxy)phenol
	Neobee 1053	glycerol tricaprinate/caprylate
	Generol 122	soya sterol
	Kesscowax B	cetyl alcohol and glycol polymer
<b>35</b>	Witconol APM	polypropylene glycol-3 myristyl ether
	Monamid 150 ADD	cocoamide diethanolamine
	Millithix 925	dibenzylidene sorbitol

	Ottasept Extra	quaternium 18 hectorite
	Bentone 38	quaternium 18 hectorite
	Triton X-102	octoxynol-13
5	Dimethicone DC 354	mixture of fully methylated linear siloxanepolymers end-blocked with trimethylsiloxy units
	Rezal 36 GP	Aluminium zirconium tetrachlorohydroxyglycine

10

**Example 12****4-Hydroxy-undecanoic acid dodecylamide**

To a suspension of 19.0 g aluminum trichloride in 50 ml of  
 15 dichloromethane, a solution of 46.5 g dodecylamine in 50 ml of  
 dichloromethane was dropped in at 15-20°C. Then 19.3 g gamma-  
 undecalactone were added quickly at room temperature and an additional 50  
 ml of dichloromethane was added. After 4 hours, the reaction mixture was  
 quenched with water and filtered through celite. The filtrate was diluted with  
 20 ether and washed with brine. The organic phase was dried, filtered and  
 evaporated to dryness. The resulting brown solid was purified by  
 recrystallisation to yield 2.65 g of colourless crystals.

25 NMR (CDCl<sub>3</sub>) δ 5.79-5.61 (m, 1H), 3.70-3.54 (m, 1H), 3.31-  
 3.15 (m, 2H), 2.42-2.28 (t, 2H), 1.96-1.10  
 (m, 34H), 0.97-0.78 (t, 6H).

**Example 13**30 **(a) 5-Hydroxy-decanoic acid dodecylamide**

A mixture of 9.34 g dodecylamine and 8.53 g d-decalactone was heated  
 to 50°C. After 10 minutes, the reaction mixture was cooled to room  
 temperature. The resulting solid was purified by recrystallisation to yield  
 35 16.70 g of colourless crystals.

NMR (CDCl<sub>3</sub>) δ 5.70-5.51 (m, 1H), 3.68-3.50 (m, 1H), 3.32-3.16 (m, 2H), 2.30-2.15 (t, 2H), 2.04-1.10 (m, 32H), 1.00-0.80 (t, 6H).

5

(b) According to the same procedure, 4-hydroxy-pentanoic acid phenethyl-amide was prepared from 6-methyl-pyran-2-one and phenethylamine.

10

(c) According to the same procedure, 4-hydroxy-undecanoic acid [12-(4-hydroxy-undecanoylamino)-dodecyl]-amide was prepared from 5-heptyldihydro-2(3H)-furanone and 1,12-diaminododecane.

#### Example 14

15

4-Oxo-pentanoic acid {4-[4-(4-oxo-pentanoylamino)-benzenesulfonyl]-phenyl}-amide

20

This compound was prepared by coupling N-{4-[4-(3-carboxy-propionylamino)-benzenesulfonyl]-phenyl}-succinamic acid (*J. Amer. Chem. Soc.*, 1945, 67, 1979) with N,O-dimethyl-hydroxyamine followed by reaction with methylmagnesium iodide to yield the solid product.

#### Example 15

25

4-Hydroxy-pentanoic acid {4-[4-(4-hydroxy-pentanoylamino)-benzenesulfonyl]-phenyl}-amide

30

A solution of sodium borohydride in methanol was cooled to 5°C. A solution of 4-oxo-pentanoic acid {4-[4-(4-oxo-pentanoylamino)-benzenesulfonyl]-phenyl}-amide in THF was added and the resulting reaction mixture was stirred at room temperature. After workup and purification, the product was obtained as a colourless solid.

35

**Example 16**

a) Fabric softener of the ester quat type (4 x concentrate):

5	INGREDIENTS	CHEMICAL NAME	%
	<u>PHASE A</u>		
10	DEIONISED WATER		to 100.0
	MgCl <sub>2</sub> (saturated sol.)	Magnesium chloride	1.0
15	<u>PHASE B</u>		
	REWOQUAT WE 18	Di-(tallow carboxyethyl)hydroxy ethyl methylammonium methosulfate	15.0
20	GENAPOL O 100	Ethoxylated fatty alcohol C16-C18 10EO	2.0
25	ANTIFOAM DB 31		0.5
	<u>PHASE C</u>		
30	ISOPROPYL ALCOHOL PRESERVATIVE		3.0 Qs
	PERFUME		Qs
35			

PROCESS:

While stirring and heating to 65° C phase A was mixed with phase B which has been preheated to 65° C. After cooling to room temperature phase C was  
5 added.

The PH value of the finished product is 2.60. It turned out that the recommended level of perfume is 1.0 %. Delayed release fragrances then could be any part of this 1.0 %.

10

b) Fabric softener of the ester quat type (1 x concentrate):

INGREDIENTS	CHEMICAL NAME	%
15	<u>PHASE A</u>	
	DEIONISED WATER	to 100.0
20	<u>PHASE B</u>	
	REWOQUAT WE 18	Di-(tallowcarboxyethyl)- hydroxy
25		6.0
		ethyl methylammonium methosulfate
	DOBANOL 25-9	Ethoxylated fatty alcohol C12-C15 9EO
30	ANTIFOAM DB 31	0.50
	<u>PHASE C</u>	0.10
	MYACIDE BT 30	2-bromo-2-nitropropane 1,3 diol
35		0.03
	PROXEL GXL	Benzisothiazolinone sodium salt
		0.02

## PERFUME

Qs

PROCESS:


5

While stirring and heating to 65° C phase A was mixed with phase B which has been preheated to 65° C. After cooling to room temperature, phase C was added.

10 The pH value of the finished product was 3.5. It turned out that the recommended level of perfume is 0.3 %. Delayed release fragrances then could be any part of this 0.3 %.

Claims

## 5 1. Precursor compounds having the formula




I

10 in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in

addition contain one or more - O - and /or -  $\overset{\text{O}}{\parallel}{\text{C}}$  - groups, whereby one or two rings can be build by the combination of the respective R<sup>1</sup> to R<sup>6</sup> and this/these ring(s) can be further substituted by an alkyl-group, in which X is either O and  
 15 R<sup>7</sup> represents a radical of an alcohol or phenol R<sup>7</sup>OH or X is N and R<sup>7</sup> represents the radical of an amine R<sup>7'</sup>R<sup>7''</sup>NH, whereby R<sup>7'</sup> and R<sup>7''</sup> represent independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either R<sup>7'</sup> R<sup>7''</sup> may be hydrogen, whereby the amine is a fragrant amine or the amine has  
 20 more than 9 C atoms, whereby R<sup>7</sup> of the alcohol or phenol and R<sup>7'</sup> and/or R<sup>7''</sup> of the amine, respectively, may further contain at least one remaining part C(OH)R<sup>1</sup>R<sup>2</sup>-CR<sup>3</sup>R<sup>4</sup>-(CR<sup>5</sup>R<sup>6</sup>)<sub>n</sub>-CO- of formula I.

## 25 2. Precursor compounds of claim 1 having the formula




Ia

30 in which n is 1, 2 or 3 and R<sup>1</sup> to R<sup>6</sup> represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in

addition contain one or more - O - and /or -  $\overset{\text{O}}{\parallel}{\text{C}}$  - groups,  $\text{R}^7$  represents a radical of an alcohol or phenol  $\text{R}^7\text{OH}$ , whereby one or two rings can be build by the combination of the respective  $\text{R}^1$  to  $\text{R}^6$  and this/these ring(s) can be further substituted by an alkyl-group, whereby  $\text{R}^7$  may further contain the remaining

5 part  $\text{C}(\text{OH})\text{R}^1\text{R}^2\text{-CR}^3\text{R}^4\text{-(CR}^5\text{R}^6)_n\text{-CO-}$  of formula Ia.

3. Precursor compounds of claim 2 having the formula



Ia

10


in which  $n$  is 1, 2 or 3 and  $\text{R}^1$  to  $\text{R}^6$  represent, independently, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or

more - O - and /or -  $\overset{\text{O}}{\parallel}{\text{C}}$  - groups,  $\text{R}^7$  represents a radical of a fragrant alcohol  $\text{R}^7\text{OH}$ , whereby one or two rings can be build by the combination of the respective  $\text{R}^1$  to  $\text{R}^6$  and this/these ring(s) can be further substituted by an alkyl-group.

15

4. Precursor compounds of claim 1 having the formula

20



Ib

in which  $n$  is 1, 2 or 3 and  $\text{R}^1$  to  $\text{R}^6$  represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in

25

addition contain one or more - O - and /or -  $\overset{\text{O}}{\parallel}{\text{C}}$  - groups, whereby one or two rings can be build by the combination of the respective  $\text{R}^1$  to  $\text{R}^6$  and this/these ring(s) can be further substituted by an alkyl-group, and  $\text{R}^7$  represents a

radical of an amine  $R^{7'}R^{7''}NH$ , whereby  $R^{7'}$  and  $R^{7''}$  represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkinyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either  $R^{7'}$  or  $R^{7''}$  may be hydrogen, whereby the amine is a fragrant amine or the amine has  
 5 more than 9 C atoms, whereby  $R^{7'}$  and/or  $R^{7''}$  may further contain at least one remaining part  $C(OH)R^1R^2-CR^3R^4-(CR^5R^6)_n-CO-$  of formula Ib.

5. Compounds of any one of the claims 1 to 4 wherein n is 1 or 2.

10 6. Compounds of any one of the claims 1 to 4 wherein  $R^1$  is an alkyl-radical and  $R^2$  to  $R^6$  are hydrogen atoms.

7. Compounds of any one of the claims 1 to 4 wherein  $R^1$ ,  $R^3$  are alkyl-radicals and  $R^2$ ,  $R^4$ ,  $R^5$ ,  $R^6$  are hydrogen atoms.

15

8. Compounds of any one of the claims 1 to 4 wherein  $R^1$  and  $R^2$  build a ring.

20

9. Compounds according to claim 8 wherein the ring is a 6-membered ring.

10. Compounds according to claim 9 wherein the ring is substituted by methyl-, ethyl- or isopropyl-.

25

11. Compounds of claim 1 or 2 wherein  $R^7$  is a radical derived from a fragrant alcohol or fragrant phenol.

12. Compounds of claim 3 or 11, wherein the fragrant alcohol or fragrant phenol is selected from the group listed in Table 1.

30

13. Compounds of claim 11 or 12 wherein the radical is derived from a fragrant alcohol or phenol selected from the group consisting of hexyl alcohol, 2-hexyl alcohol, heptyl alcohol, octyl alcohol, nonyl alcohol, decyl alcohol, undecyl alcohol, lauryl alcohol, 3-methyl-but-2-en-1-ol, cis-3-hexenol, cis-4-  
 35 hexenol, 3,4,5,6,6-pentamethylheptan-2-ol, citronellol, geraniol, 6-ethyl-3-methyl-5-octen-1-ol, 3,7-dimethyl-oct-3,6-dienol, 3,7-dimethyloctanol, 7-methoxy-3,7-dimethyl-octan-2-ol, cis-6-nonenol, 6,8-dimethyl-2-nonanol, 4-methyl-3-decen-5-ol, 1-phenyl ethanol, 2-phenyl ethanol, 2-phenyl propanol, 3-

phenyl propanol, 2-methyl-5-phenyl pentanol, 2-methyl-4-phenyl-pentanol, 3-methyl-5-phenyl-pentanol, 2-(2-methylphenyl)-ethanol, 4-(4-hydroxyphenyl)butan-2-one, 2-phenoxy ethanol, anisic alcohol, p-tolyl alcohol, cinnamic alcohol, vanillin, ethyl vanillin, eugenol, isoeugenol, anethol, borneol, cedrenol, farnesol, fenchyl alcohol, menthol, alpha ionol, tetrahydro ionol, 2-(1,1-dimethylethyl)cyclohexanol, 3-(1,1-dimethylethyl)cyclohexanol, 4-(1,1-dimethylethyl)cyclohexanol, 6,6-dimethyl-bicyclo [3.1.1]hept-2-ene-methanol, p-menth-8-en-3-ol, 2,4,6-trimethyl-3-cyclohexenyl-methanol, 4-(1-methylethyl)cyclohexyl-methanol, 2,2,6-trimethyl-alpha-propyl cyclohexane propanol, 5-(2,2,3-trimethyl-3-cyclopentenyl)-3-methylpentan-2-ol, 3-methyl-5-(2,2,3-trimethylcyclopentyl-3-enyl)pent-4-en-2-ol, 2-ethyl-4-(2,2,3-trimethylcyclopentyl-3-enyl)but-2-en-1-ol, 4-(5,5,6-trimethylbicyclo[2.2.1]hept-2-yl)-cyclohexanol, 2-(2-methylpropyl)-4-hydroxy-4-methyl-tetrahydropyran, 2-cyclohexyl propanol, 2-(1,1-dimethylethyl)-4-methyl cyclohexanol, 1-(2-tert-butyl-cyclohexyloxy)-2-butanol, 1-(4-isopropyl-cyclohexyl)-ethanol, 2,6-dimethyl-heptan-2-ol, 2,6-dimethyl-oct-7-en-2-ol.

14. Compounds of claim 13 wherein the radical is derived from a fragrant alcohol or phenol selected from the group consisting of citronellol, geraniol, linalool, cis-3-hexenol, 4-methyl-3-decen-5-ol, 2-phenyl ethanol, iso eugenol, 3-methyl-5-(2,2,3-trimethylcyclopentyl-3-enyl)pent-4-en-2-ol, 2-ethyl-4-(2,2,3-trimethylcyclopentyl-3-enyl)but-2-en-2-ol, 2,6-dimethyl-heptan-2-ol, 2,6-dimethyl-oct-7-en-2-ol, cinnamic alcohol and eugenol.

15. Compounds of claim 14 wherein the radical is derived from a fragrant alcohol selected from the group consisting of citronellol, 2-phenyl ethanol, linalool, geraniol and cis-3-hexanol.

16. Compounds according to any one of the claims 1 to 4 which upon cleavage provide lactones with organoleptic properties wherein the lactones are selected from the group consisting of 6-methyl-pyran-2-one, 5-heptyldihydro-2(3H)-furanone, 5-pentyldihydro-2(3H)-furanone, 5-(3-hexenyl)dihydro-5-methyl-(Z)-2(3H)-furanone, 5-hexyldihydro-5-methyl-2(3H)-furanone, 5-hexyldihydro-2(3H)-furanone, 5-octyldihydro-2(3H)-furanone, 8-(1-methylethyl)-1-oxaspiro[4.5]-decan-2-one, 8-methyl-1-oxaspiro[4.5]-decan-2-one, 8-ethyl-1-oxaspiro[4.5]-decan-2-one, 5-(1,5-dimethyl-4-hexenyl)dihydro-2(3H)-furanone, 2-oxo-5-butyl-tetrahydrofuran, 4-methyl-5-pentyl-dihydro-2(3H)-furan-2-one, 5-hexyldihydro-5-methyl-2(3H)-furanone, dihydro-5-

methyl-5-vinyl-2(3H)-furanone, octahydro-2H-1-benzopyran-2-one, tetrahydro-6-pentyl-2H-pyran-2-one, tetrahydro-6-hexyl-2H-pyran-2-one, tetrahydro-6-heptyl-2H-pyran-2-one, tetrahydro-6-(3-pentenyl)-(E)-2H-pyran-2-one and tetrahydro-6-(2-pentenyl)-(Z)-2H-pyran-2-one.

5

17. Compounds of claim 16, wherein the lactones are selected from the group consisting of 5-heptyldihydro-2(3H)-furanone, 5-pentyldihydro-2(3H)-furanone, 5-hexyldihydro-2(3H)-furanone, 8-(1-methylethyl)-1-oxaspiro[4.5]-decan-2-one and 2-oxo-5-butyl-tetrahydrofuran.

10

18. Compound according to claim 2 selected from the group consisting of 4-hydroxy-decanoic acid 2-phenethyl ester, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-decanoic acid hex-3-enyl ester, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-6-enyl ester, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hept-6-enyl ester, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-undecanoic acid phenethyl ester, 4-hydroxy-undecanoic acid hex-3-enyl ester, 4-hydroxy-nonanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-hydroxy-nonanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-nonanoic acid hex-3-enyl ester, 4-hydroxy-nonanoic acid phenethyl ester, 3-(1-hydroxy-4-isopropyl-cyclohexyl)-propionic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-hydroxy-undecanoic acid 2-(4-hydroxy-undecanoyloxy)-ethyl ester and 4-hydroxy-undecanoic acid 2-benzyloxycarbonyl-2-benzyloxycarbonylamino-ethyl ester.

15

20

25

30

35

40

45

50

55

60

65

70

75

80

85

90

95

100

105

110

115

120

125

130

135

140

145

150

155

160

165

170

175

180

185

190

195

200

205

210

215

220

225

230

235

240

245

250

255

260

265

270

275

280

285

290

295

300

305

310

315

320

325

330

335

340

345

350

355

360

365

370

375

380

385

390

395

400

405

410

415

420

425

430

435

440

445

450

455

460

465

470

475

480

485

490

495

500

505

510

515

520

525

530

535

540

545

550

555

560

565

570

575

580

585

590

595

600

605

610

615

620

625

630

635

640

645

650

655

660

665

670

675

680

685

690

695

700

705

710

715

720

725

730

735

740

745

750

755

760

765

770

775

780

785

790

795

800

805

810

815

820

825

830

835

840

845

850

855

860

865

870

875

880

885

890

895

900

905

910

915

920

925

930

935

940

945

950

955

960

965

970

975

980

985

990

995

1000

methylamino-benzoic acid methyl ester; 6-methyltetrahydro-quinoline; isobutyl N-methyl anthranilate; (Z)-3-hexenyl 2-aminobenzoate.

21. Compounds of claim 20 wherein the odoriferous amine is selected  
5 from the group consisting of 2-amino benzoic acid methyl ester; anthranilic acid phenylethyl ester; 2-methylamino-benzoic acid methyl ester and (Z)-3-hexenyl 2-aminobenzoate.

22. Compound according to claim 4 selected from the group consisting of  
10 4-hydroxy-undecanoic acid dodecylamide; 5-hydroxy-decanoic acid dodecylamide; 4-oxo-pentanoic acid {4-[4-(4-oxo-pentanoylamino)-benzenesulfonyl]-phenyl}-amide  
4-hydroxy-pentanoic acid {4-[4-(4-hydroxy-pentanoylamino)-benzenesulfonyl]-phenyl}-amide and 4-hydroxy-undecanoic acid [12-(4-hydroxy-undecanoylamino)-dodecyl]-amide.

15

23. Compound of any one of the claims 1 to 22 characterized in that the compound is cleavable by heat.

24. A composition for cosmetic application to the human skin, air  
20 fresheners, hard surface cleaners or laundry products, containing at least one of the compounds according to any one of the claims 1 to 23.

25. A process for prolonging the effect of diffusion of the characteristic odour of an odoriferous alcohol and/or amine and/or lactone on human skin or  
25 of air fresheners or in hard surface cleaners or in laundry products comprising applying at least one of the compounds of any one of the claims 1 to 23.

26. A method of suppressing human body malodour by means of applying a composition of claim 24 to the human skin.


30

27. The use of at least one of the compounds of any one of the claims 1 to 23 as a fragrance and/or antibacterial precursor in a cosmetic composition or cosmetic product, air freshener, hard surface cleaner or laundry product.

35

40

## 28. Compounds having the formula



I

- 5 in which  $n$  is 1, 2 or 3 and  $\text{R}^1$  to  $\text{R}^6$  represent, independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic-radicals or hydrogen wherein these radicals may in addition contain one or more - O - and /or -  $\overset{\text{O}}{\parallel}{\text{C}}$  - groups, whereby one or two rings can be build by the combination of the respective  $\text{R}^1$  to  $\text{R}^6$  and this/these
- 10 ring(s) can be further substituted by an alkyl-group, in which X is either O and  $\text{R}^7$  represents a radical of an alcohol or phenol  $\text{R}^7\text{OH}$  or X is N and  $\text{R}^7$  represents the radical of an amine  $\text{R}^7'\text{R}^7''\text{NH}$ , whereby  $\text{R}^7'$  and  $\text{R}^7''$  represent independently, branched or unbranched, substituted or unsubstituted alkyl-, alkenyl-, alkynyl-, cycloalkyl-, cycloalkenyl- or aromatic radicals or either  $\text{R}^7'$
- 15  $\text{R}^7''$  may be hydrogen, whereby the amine is a fragrant amine or the amine has more than 9 C atoms, whereby  $\text{R}^7$  of the alcohol or phenol and  $\text{R}^7'$  and/or  $\text{R}^7''$  of the amine, respectively, may further contain at least one remaining part  $\text{C}(\text{OH})\text{R}^1\text{R}^2\text{-CR}^3\text{R}^4\text{-(CR}^5\text{R}^6)_n\text{-CO-}$  of formula I.

- 20 29. Compound selected from the group consisting of 4-hydroxy-decanoic acid 2-phenethyl ester, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-decanoic acid hex-3-enyl ester, 4-hydroxy-decanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-6-enyl ester, 5-hydroxy-dodecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-
- 25 hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-2,6-dienyl ester, 4-hydroxy-decanoic acid 1,5-dimethyl-1-vinyl-hex-4-enyl ester, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hept-6-enyl ester, 4-hydroxy-decanoic acid 1,1,5-trimethyl-hexyl ester, 4-hydroxy-undecanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-undecanoic acid
- 30 phenethyl ester, 4-hydroxy-undecanoic acid hex-3-enyl ester, 4-hydroxy-nonanoic acid 3,7-dimethyl-octa-2,6-dienyl ester, 4-hydroxy-nonanoic acid 3,7-dimethyl-oct-6-enyl ester, 4-hydroxy-nonanoic acid hex-3-enyl ester, 4-hydroxy-nonanoic acid phenethyl ester, 3-(1-hydroxy-4-isopropyl-cyclohexyl)-propionic

acid 3,7-dimethyl-octa-2,6-dienyl ester and 4-hydroxy-undecanoic acid 2-(4-hydroxy-undecanoyloxy)-ethyl ester.

30. Compound selected from the group consisting of 4-hydroxy-undecanoic  
5 acid dodecylamide; 5-hydroxy-decanoic acid dodecylamide, 4-hydroxy-  
pentanoic acid {4-[4-(4-hydroxy-pentanoylamino)-benzenesulfonyl]-phenyl}-  
amide and 4-hydroxy-undecanoic acid [12-(4-hydroxy-undecanoylamino)-  
dodecyl]-amide.

10

\*\*\*