




(22) Date de dépôt/Filing Date: 2010/04/29
 (41) Mise à la disp. pub./Open to Public Insp.: 2010/11/04
 (45) Date de délivrance/Issue Date: 2022/07/12
 (62) Demande originale/Original Application: 2 760 629
 (30) Priorités/Priorities: 2009/04/29 (GB0907413.9);
 2009/05/13 (US61/177,811)

(51) Cl.Int./Int.Cl. *A61K 31/202* (2006.01),
A61P 17/00 (2006.01), *A61P 17/16* (2006.01)
 (72) Inventeurs/Inventors:
 KELLIHER, ADAM, GB;
 MORRISON, ANGUS, GB;
 KNOWLES, PHIL, GB
 (73) Propriétaire/Owner:
 DS BIOPHARMA LIMITED, IE
 (74) Agent: SMART & BIGGAR LLP

(54) Titre : UTILISATION DE PUFAS POUR TRAITER UNE INFLAMMATION CUTANEE
 (54) Title: USE OF PUFAS FOR TREATING SKIN INFLAMMATION




(I)

(57) **Abrégé/Abstract:**

The present invention provides a compound which is a polyunsaturated fatty acid (PUFA) derivative of formula (I), or a pharmaceutically acceptable salt, or solvate thereof, in the form of a racemate, a stereoisomer or a mixture of stereoisomers, for use in treating skin inflammation in a mammal, by topical administration. (see formula I)

Abstract

The present invention provides a compound which is a polyunsaturated fatty acid (PUFA) derivative of formula (I), or a pharmaceutically acceptable salt, or solvate thereof, in the form of a racemate, a stereoisomer or a mixture of stereoisomers, for use in treating skin inflammation in a mammal, by topical administration.



I

WO 2010/125340

PCT/GB2010/000844

USE OF PUFAS FOR TREATING SKIN INFLAMMATION

Field of the Invention

5 The present invention relates to novel methods of treating skin inflammation, in particular skin inflammation caused by atopic eczema, contact dermatitis, psoriasis or uremic pruritis.

Background of the Invention

10

Most common skin disorders and complaints comprise a number of different components. Thus, many skin disorders involve (a) hyperproliferation, (b) inflammation and/or (c) dehydration. Hyperproliferation involves a state of abnormally high cell division, which can lead to excess flaking skin. Inflammation
15 involves swelling and redness of the skin, as well as sensations of increased heat, and pain in the skin. Dehydration involves loss of water from the skin and may be due, for example, to damage to the normally waterproof top layer of the skin (epidermis).

Inflammation of the skin (dermatitis) in mammals can result from a number of
20 different etiologies. Dermatitis can be caused by eczema, in particular atopic eczema (atopic dermatitis), disseminated neurodermatitis, flexural eczema, infantile eczema, prurigo diathesique, contact dermatitis, (eg irritant contact dermatitis, allergic contact dermatitis and photocontact dermatitis), xerotic eczema, seborrheic eczema, dyshidrosis, discoid eczema, venous eczema, dermatitis herpetiformis,
25 neurodermatitis and autoeczematisation. Dermatitis can also be caused by skin inflammation resulting from exposure to radiation, in particular exposure to ultraviolet radiation. Other causes of dermatitis includes uremic pruritis and autoimmune diseases, in particular lupus and psoriasis.

30 Inflammation of the skin causes rashes, redness, skin edema (swelling), itching, blistering, sensations of pain and/or heat and can be unsightly. The itchiness caused by inflammation can lead to scratching. Scratching of skin that is already damaged in some way can easily lead to the barrier of the epidermis being broken, resulting in

WO 2010/125340

PCT/GB2010/000844


bleeding, and secondary infection with pathogens. Such secondary infection can require treatment with antibiotics.

It is well known that when treating a skin condition that has a number of different components, i.e. hyperproliferative, inflammatory and/or dehydrative components, a number of different treatments may be used. Thus, in the treatment of psoriasis, for example, an antihyperproliferative may be used to treat the hyperproliferative component of the disease, an anti-inflammatory may be used to treat the inflammatory component and an emollient may be used to treat the dehydrative component.

The most common form of treatment for inflammation of the skin is oral and/or topical steroids. There are, however, drawbacks associated with steroid treatments. Common side effects associated with steroids include stunting of growth, thinning of the skin, muscle loss and osteoporosis.

The present invention relates to new methods for treating skin inflammation, in particular skin inflammation caused by atopic eczema, contact dermatitis, psoriasis or uremic pruritis, in mammals.

Eicosa-8Z,11Z,14Z-trienoic acid (Dihomo- γ -linolenic acid or DGLA) is a commercially available polyunsaturated fatty acid (PUFA). DGLA has the structure shown below.




EP-A-0085579 describes the use of DGLA in combination with antipruritic lithium salts. EP-A-0173478 describes the use of DGLA in combination with anti-inflammatory glucocorticoids. In these applications, treatment with lithium salts and glucocorticoids is supplemented with DGLA, as both lithium salts and glucocorticoids are believed to block release of DGLA from endogenous stores in the body.

WO 2010/125340


PCT/GB2010/000844

Advantageously, it has now been discovered that DGLA can be used effectively as a monotherapy.


- 5 5-Hydroxy-eicosa-6E,8Z,11Z-trienoic acid (5-HETrE) is a commercially available PUFA derivative derived from mead acid. 5-HETrE has the structure shown below.



- 10 8-Hydroxy-eicosa-9E,11Z,14Z-trienoic acid (8-HETrE) is a commercially available PUFA derivative derived from eicosa-8Z,11Z,14Z-trienoic acid (Dihomo- γ -linolenic acid or DGLA). 8-HETrE has the structure shown below.



- 15 15-Hydroxy-eicosa-8Z,11Z,13E-trienoic acid (15-HETrE) is a commercially available PUFA derivative derived from eicosa-8Z,11Z,14Z-trienoic acid (Dihomo- γ -linolenic acid or DGLA). 15-HETrE has the structure shown below.



20


15-HETrE is known to have antiproliferative properties when applied directly to the skin (*Xi, et al*; Prostaglandins, Leukotrienes and Essential Fatty Acids (2000) 62(1), 13 to 19).

WO 2010/125340

PCT/GB2010/000844

13-Hydroxy-octadeca-6Z, 9Z, 11E-γ-trienoic acid (13-HOTrE(γ)) is a commercially available PUFA derivative derived from gamma-linolenic acid (GLA). 13-HOTrE(γ) has the structure shown below.

5




It has now been surprisingly found that DGLA, 5-HETrE, 8-HETrE, 15-HETrE, 13-HOTrE(γ) and their derivatives are clinically useful in treating skin inflammation, in particular skin inflammation caused by atopic eczema, contact dermatitis, psoriasis or uremic pruritis, by topical administration in mammals. A particular finding of the present invention is that these compounds reduce the level of COX-2 enzymes in the skin when applied topically. The COX-2 family of enzymes have been strongly linked to inflammation and have been found to be present in increased amounts in inflamed tissue.

Summary of the Invention

The present invention therefore provides a compound which is a polyunsaturated fatty acid (PUFA) derivative of formula (I),

20



I

25 or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is -CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-C₃H₆-,
 -(CH₂)₃-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-, -(CH₂)₃-
 [*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)- or -CH₂-
 [*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-;

WO 2010/125340

PCT/GB2010/000844

R₁ is a hydrogen atom; or

R₁ is a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl, 5- to 10-membered heteroaryl, C₃-C₇ carbocyclyl or 5- to 10-membered heterocyclyl group; or

5 R₁ is a group of formula -CH₂-CH(OR₃)-CH₂-(OR₄), wherein R₃ and R₄ are each independently hydrogen atoms or -(C=O)-R₆, wherein R₆ is an aliphatic group having from 3 to 29 carbon atoms; or

R₁ is a group of formula -(CH₂OCH₂)_mOH, wherein m is an integer of from 1 to 200;

10 R₂ is a hydrogen atom; or

R₂ is a group -(C=O)-R₅, wherein R₅ is a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl, 5- to 10-membered heteroaryl, C₃-C₇ carbocyclyl or 5- to 10-membered heterocyclyl group, or R₅ is an aliphatic group having from 3 to 29 carbon atoms; or

15 R₂ is a group of formula -(CH₂OCH₂)_nOH, wherein n is an integer of from 1 to 200;

and wherein

said alkyl, alkenyl, alkynyl and aliphatic groups are the same or different and are each unsubstituted or substituted with 1, 2 or 3 unsubstituted substituents which are the same or different and are selected from halogen atoms and C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₁-C₄ haloalkoxy, C₂-C₄ haloalkenyloxy, hydroxyl, -SR', and -NR'R'' groups where R' and R'' are the same or different and represent hydrogen or unsubstituted C₁-C₂ alkyl;

25 said aryl, heteroaryl, carbocyclyl and heterocyclyl groups are the same or different and are each unsubstituted or substituted by 1, 2, 3 or 4 unsubstituted substituents which are the same or different and are selected from halogen atoms, and cyano, nitro, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₂-C₄ alkenyl, C₂-C₄ alkenyloxy, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₁-C₄ haloalkoxy, C₂-C₄ haloalkenyloxy, hydroxyl, C₁-C₄ hydroxyalkyl, -SR' and -NR'R'' groups
30 wherein each R' and R'' is the same or different and represents hydrogen or unsubstituted C₁-C₄ alkyl;

and wherein the PUFA derivative is in the form of a racemate, a stereoisomer or a mixture of stereoisomers,


which compound is for use in treating skin inflammation in a mammal, by topical administration.

Various aspects of this disclosure relate to a pharmaceutical composition suitable for topical administration comprising a compound of the invention, and a pharmaceutically acceptable diluent or carrier, for use in treating skin inflammation in a mammal.

Various aspects of this disclosure relate to a use of a compound for the treatment of skin inflammation.

Various aspects of this disclosure relate to use of a compound in the manufacture of a medicament for the treatment of skin inflammation.


Various embodiments of the claimed invention relate to a topical monotherapy composition for the treatment of skin inflammation in a mammal, the topical monotherapy composition comprising: at least one excipient; and a polyunsaturated fatty acid (PUFA) derivative of formula (I):



or a salt thereof, wherein:

-Alk- is: $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; topical monotherapy composition for the treatment of skin inflammation and wherein the topical monotherapy composition is substantially free of glucocorticoids.


Various embodiments of the claimed invention relate to a use of a polyunsaturated fatty acid (PUFA) derivative of formula (I):



or a salt thereof, in the preparation of a medicament for the treatment of skin inflammation in a mammal as a topical monotherapy, wherein:

-Alk- is: $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient; and wherein the medicament is substantially free of glucocorticoids.

Various embodiments of the claimed invention relate to a use of a composition for the treatment of skin inflammation in a mammal as a topical monotherapy, the composition comprising: at least one excipient; and a polyunsaturated fatty acid (PUFA) derivative of formula (I):




or a salt thereof, wherein:

-Alk- is: $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; wherein the PUFA derivative is in a composition comprising at least one excipient,

wherein the composition contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient, and wherein the composition is substantially free of glucocorticoids.


Various embodiments of the claimed invention relate to a use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),



or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; and R_1 is a hydrogen atom, wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient, and wherein the amount of the PUFA derivative is 0.5 wt% to 25 wt% of the medicament.

Various embodiments of the claimed invention relate to a use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),




or a pharmaceutically acceptable salt thereof for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; and R_1 is a hydrogen atom;

wherein the PUFA derivative is in a medicament, wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient, and wherein the amount of the PUFA derivative is 0.5 wt% to 25 wt% of the medicament.

Various embodiments of the claimed invention relate to a use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),




(I)

or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; and R_1 is hydrogen atom;

wherein the medicament is free of lithium salts and glucocorticoids, and wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient.

Various embodiments of the claimed invention relate to a use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),




(I)

or a pharmaceutically acceptable salt thereof for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; and R_1 is hydrogen atom;

wherein the PUFA derivative is in a medicament, and wherein the medicament is free of lithium salts and glucocorticoids, and wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient.


Various aspects of the disclosure relate to a topical composition use in treating pruritus, the topical composition comprising: at least one excipient; and a polyunsaturated fatty acid (PUFA) derivative of formula (I):



or a salt thereof, wherein:

-Alk- is: $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; and wherein the topical composition is substantially free of glucocorticoids, and , wherein the composition is for co-administration with a corticosteroid selected from the group consisting of clobetasol dipropionate, betamethasone dipropionate, halbetasol dipropionate, diflorasone diacetate, fluocinonide, halcinonide, amcinonide, desoximetasone, triamcinolone acetonide, mometasone furoate, fluticasone dipropionate, flucinolone acetonide, hydrocortisone valerate, hydrocortisone butyrate, desonide, prednicarbate, prednisolone, methylprednisolone, dexamethasone, naflocort, deflazacort, halopredone acetate, budesonide, beclomethasone dipropionate, hydrocortisone, clocortolone pivalate, methylprednisolone aceponate, dexamethasone palmitoate, tipredane, hydrocortisone aceponate, alclometasone dipropionate, halometasone, methylprednisolone suleptanate, rimexolone, prednisolone farnesylate, ciclesonide, deprodone propionate, loteprednol etabonate, betamethasone butyrate propionate, flunisolide, prednisone, dexamethasone sodium phosphate, triamcinolone, betamethasone 17- valerate, betamethasone, hydrocortisone acetate, hydrocortisone sodium succinate, prednisolone sodium phosphate, hydrocortisone probutate, and combinations thereof.

Various embodiments of the claimed invention relate to a pharmaceutical composition for use in the treatment of skin inflammation in a mammal by topical administration, comprising a polyunsaturated fatty acid (PUFA) derivative, of formula (I),




or a pharmaceutically acceptable salt, or solvate thereof, wherein -Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; and wherein the PUFA derivative is the sole active ingredient.

Various embodiments of the claimed invention relate to a use of a composition comprising DGLA or an ester or pharmaceutically acceptable salt thereof, for treating uremic pruritus in a subject in need thereof, wherein the composition is formulated for administration to provide a therapeutically effective amount to skin of the subject, wherein the amount of the DGLA is 0.5 wt% to 25 wt% of the composition.

Various embodiments of the claimed invention relate to a use of DGLA or an ester or pharmaceutically acceptable salt thereof in the preparation of a composition for treating uremic pruritus in a subject in need thereof, wherein the composition formulated for administration to provide a therapeutically effective amount to skin of the subject, wherein the amount of the DGLA is 0.5 wt% to 25 wt% of the composition.

Various embodiments of the claimed invention relate to a pharmaceutical composition for use in the treatment of skin inflammation in a mammal by topical administration, comprising a polyunsaturated fatty acid (PUFA) derivative, of formula (I),




(I) or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom, wherein the PUFA is the sole active ingredient,

wherein the skin inflammation is caused by atopic dermatitis, and wherein the amount of the PUFA is 1 wt% to 10 wt% of the pharmaceutical composition.

Various aspects of this disclosure relate to a compound which is a polyunsaturated fatty acid (PUFA) derivative of formula (I)



(I)

6d


or is a pharmaceutically acceptable salt or solvate of the PUFA derivative, wherein: -Alk- is:

-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-(CH₂)₃-, -(CH₂)₃-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-, -(CH₂)₃-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)- or -CH₂-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-; R₁ is a hydrogen atom; or R₁ is a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl, 5- to 10-membered heteroaryl, C₃-C₇ carbocyclyl or 5- to 10-membered heterocyclyl group; or R₁ is a group of formula -CH₂-CH(OR₃)-CH₂-(OR₄), wherein R₃ and R₄ are each independently hydrogen atoms or -(C-O)-R₆, wherein R₆ is an aliphatic group having from 3 to 29 carbon atoms; or R₁ is a group of formula -(CH₂OCH₂)_mOH, wherein m is an integer of from 1 to 200; R₂ is a hydrogen atom; or R₂ is a group -(C=O)-R₅, wherein R₅ is a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, C₆-C₁₀ aryl, 5- to 10-membered heteroaryl, C₃-C₇ carbocyclyl or 5- to 10-membered heterocyclyl group, or R₅ is an aliphatic group having from 3 to 29 carbon atoms; or

R₂ is a group of formula -(CH₂OCH₂)_nOH, wherein n is an integer of from 1 to 200; and wherein said alkyl, alkenyl, alkynyl and aliphatic groups are the same or different and are each unsubstituted or substituted with 1, 2 or 3 unsubstituted substituents which are the same or different and are selected from halogen atoms, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₁-C₄ haloalkoxy, C₂-C₄ haloalkenyloxy, hydroxyl, -SR', and -NR'R" groups where R' and R" are the same or different and represent hydrogen or unsubstituted C₁-C₂ alkyl;

said aryl, heteroaryl, carbocyclyl and heterocyclyl groups are the same or different and are each unsubstituted or substituted by 1, 2, 3 or 4 unsubstituted substituents which are the same or different and are selected from halogen atoms, cyano, nitro, C₁-C₄ alkyl, C₁-C₄ alkoxy, C₂-C₄ alkenyl, C₂-C₄ alkenyloxy, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₁-C₄ haloalkoxy, C₂-C₄ haloalkenyloxy, hydroxyl, C₁-C₄ hydroxyalkyl, -SR' and -NR'R" groups wherein each R' and R" is the same or different and represents hydrogen or unsubstituted C₁-C₄ alkyl; and wherein the PUFA derivative is in the form of a racemate, a stereoisomer or a mixture of stereoisomers, which compound is for use in treating a skin inflammation in a mammal, by topical administration.

Various aspects of this disclosure also relate to use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),




(I)

or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is $-\text{CH}(\text{OR}_2)-[\textit{trans}]\text{CH}=\text{CH}-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-\text{C}_3\text{H}_6-$, $-(\text{CH}_2)_3-\text{CH}(\text{OR}_2)-[\textit{trans}]\text{CH}=\text{CH}-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-$, $-(\text{CH}_2)_3-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-[\textit{trans}]\text{CH}=\text{CH}-\text{CH}(\text{OR}_2)-$ or $-(\text{CH}_2)_3-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-$; R_1 is a hydrogen atom; or R_1 is a C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_6 - C_{10} aryl, 5- to 10-membered heteroaryl, C_3 - C_7 carbocyclyl or 5- to 10-membered heterocyclyl group; or R_1 is a group of formula $-\text{CH}_2-\text{CH}(\text{OR}_3)-\text{CH}_2-(\text{OR}_4)$, wherein R_3 and R_4 are each independently hydrogen atoms or $-(\text{C}=\text{O})-\text{R}_6$, wherein R_6 is an aliphatic group having from 3 to 29 carbon atoms; or R_1 is a group of formula $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is an integer of from 1 to 200; R_2 is a hydrogen atom; or R_2 is a group $-(\text{C}=\text{O})-\text{R}_5$, wherein R_5 is a C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_6 - C_{10} aryl, 5- to 10-membered heteroaryl, C_3 - C_7 carbocyclyl or 5- to 10-membered heterocyclyl group, or R_5 is an aliphatic group having from 3 to 29 carbon atoms; or R_2 is a group of formula $-(\text{CH}_2\text{OCH}_2)_n\text{OH}$, wherein n is an integer of from 1 to 200; and wherein said alkyl, alkenyl, alkynyl and aliphatic groups are the same or different and are each unsubstituted or substituted with 1, 2 or 3 unsubstituted substituents which are the same or different and are selected from halogen atoms and C_1 - C_4 alkoxy, C_2 - C_4 alkenyloxy, C_1 - C_4 haloalkyl, C_2 - C_4 haloalkenyl, C_1 - C_4 haloalkoxy, C_2 - C_4 haloalkenyloxy, hydroxyl, $-\text{SR}'$, and $-\text{NR}'\text{R}''$ groups where R' and R'' are the same or different and represent hydrogen or unsubstituted C_1 - C_2 alkyl; said aryl, heteroaryl, carbocyclyl and heterocyclyl groups are the same or different and are each unsubstituted or substituted by 1, 2, 3 or 4 unsubstituted substituents which are the same or different and are selected from halogen atoms, and cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4 alkoxy, C_2 - C_4 alkenyl, C_2 - C_4 alkenyloxy, C_1 - C_4 haloalkyl, C_2 - C_4 haloalkenyl, C_1 - C_4 haloalkoxy, C_2 - C_4 haloalkenyloxy, hydroxyl, C_1 - C_4 hydroxyalkyl, $-\text{SR}'$ and $-\text{NR}'\text{R}''$ groups wherein each R' and R'' is the same or different and represents hydrogen or unsubstituted C_1 - C_4 alkyl; and wherein the PUFA derivative is in the form of a racemate, a stereoisomer or a mixture of stereoisomers.

Various aspects of this disclosure also relate to a topical composition for use in treating skin inflammation in a mammal, comprising: at least one excipient; and a polyunsaturated fatty acid (PUFA) derivative of formula (I),




(I)

or a salt thereof, wherein

-Alk- is: -CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH- C₃H₆-, -(CH₂)₃-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-, -(CH₂)₃-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-or -CH₂-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-; R₁ is a group of formula -CH₂-CH(OR₃)-CH₂-(OR₄), wherein R₃ and R₄ each independently represent a hydrogen atom or -(C=O)-R₆, wherein R₆ is a saturated aliphatic group having from 3 to 29 carbon atoms, wherein at least one of R₃ or R₄ is -(C=O)-R₆; or R₁ is a group of formula -(CH₂OCH₂)_mOH, wherein m is an integer of from 1 to 200; and/or R₂ is a group -(C=O)-R₅, wherein R₅ is a saturated aliphatic group having from 3 to 29 carbon atoms; or R₂ is a group of formula -(CH₂OCH₂)_nOH, wherein n is an integer of from 1 to 200.

Various aspects of this disclosure also relate to a topical composition comprising: at least one excipient; and a polyunsaturated fatty acid (PUFA) derivative of formula (I):




(I)

or a salt thereof, wherein:

-Alk- is: -(CH₂)₃-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-; R₁ is a hydrogen atom; wherein the PUFA derivative is in the form of a racemate, a stereoisomer or a mixture of stereoisomers; and wherein the topical composition is substantially free of glucocorticoids.

Various aspects of this disclosure also relate to a pharmaceutical composition for use in the treatment of skin inflammation in a mammal by topical administration, comprising a polyunsaturated fatty acid (PUFA) derivative, of formula (I),



6g

(I)


or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; or R_1 is a $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_6\text{-C}_{10}$ aryl, 5- to 10-membered heteroaryl, $\text{C}_3\text{-C}_7$ carbocyclyl or 5- to 10-membered heterocyclyl group; or R_1 is a group of formula $-\text{CH}_2\text{-CH(OR}_3\text{)-CH}_2\text{-(OR}_4\text{)}$, wherein R_3 and R_4 are each independently hydrogen atoms or $-(\text{C=O})\text{-R}_6$, wherein R_6 is an aliphatic group having from 3 to 29 carbon atoms; or R_1 is a group of formula $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is an integer of from 1 to 200; and wherein said alkyl, alkenyl, alkynyl and aliphatic groups are the same or different and are each unsubstituted or substituted with 1, 2 or 3 unsubstituted substituents which are the same or different and are selected from halogen atoms and $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_2\text{-C}_4$ alkenyloxy, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_2\text{-C}_4$ haloalkenyl, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_2\text{-C}_4$ haloalkenyloxy, hydroxyl, $-\text{SR}'$, and $-\text{NR}'\text{R}''$ groups where R' and R'' are the same or different and represent hydrogen or unsubstituted $\text{C}_1\text{-C}_2$ alkyl; said aryl, heteroaryl, carbocyclyl and heterocyclyl groups are the same or different and are each unsubstituted or substituted by 1, 2, 3 or 4 unsubstituted substituents which are the same or different and are selected from halogen atoms, and cyano, nitro, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkenyloxy, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_2\text{-C}_4$ haloalkenyl, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_2\text{-C}_4$ haloalkenyloxy, hydroxyl, $\text{C}_1\text{-C}_4$ hydroxyalkyl, $-\text{SR}'$ and $-\text{NR}'\text{R}''$ groups wherein each R' and R'' is the same or different and represents hydrogen or unsubstituted $\text{C}_1\text{-C}_4$ alkyl; and wherein the PUFA derivative is in the form of a racemate, a stereoisomer or a mixture of stereoisomers, and wherein the PUFA derivative is the sole active ingredient.

Various aspects of this disclosure also relate to use of a composition comprising 15-HETrE for treating skin inflammation in a mammal caused by exposure to electromagnetic radiation, the wherein the composition is formulated for administering a therapeutically effective amount to the skin of the mammal.

Various aspects of this disclosure also relate to a topical composition comprising 15-HETrE or an ester thereof and an excipient.

Various aspects of this disclosure also relate to use of a composition comprising a therapeutically effective amount of a compound of formula:



or a pharmaceutically acceptable salt thereof for treating itching of the skin in a mammal.


Various aspects of this disclosure also relate to use of a composition comprising 5-HETrE, 8-HETrE or 15-HETrE, or an ester or pharmaceutically acceptable salt thereof for treating psoriasis in a subject in need thereof, wherein the composition is formulated to provide a therapeutically effective amount to skin of the subject.

Various aspects of this disclosure also relate to use of a composition comprising 5-HETrE or 8-HETrE, or an ester or pharmaceutically acceptable salt thereof for treating pruritus in a subject in need thereof, wherein the composition is formulated for topical administration to skin of the subject.

Various aspects of this disclosure also relate to use of DGLA or a C1-C2 alkyl ester or pharmaceutically acceptable salt thereof, for treating uremic pruritus in a subject in need thereof, wherein the DGLA is in a composition formulated for administration to provide a therapeutically effective amount to skin of the subject, wherein the amount of the DGLA is 0.5 wt% to 25 wt% of the composition.

Various aspects of this disclosure also relate to use of DGLA or a C1-C2 alkyl ester or pharmaceutically acceptable salt thereof in the preparation of a composition for treating uremic pruritus in a subject in need thereof, wherein the composition formulated for administration to provide a therapeutically effective amount to skin of the subject, wherein the amount of the DGLA is 0.5 wt% to 25 wt% of the composition.

Various aspects of this disclosure also relate to a pharmaceutical composition for use in the treatment of skin inflammation in a mammal by topical administration, comprising a polyunsaturated fatty acid (PUFA) derivative, of formula (I),



(I)

or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is $-(\text{CH}_2)_3\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-}$; R_1 is a hydrogen atom; or R_1 is a $\text{C}_1\text{-C}_6$ alkyl, $\text{C}_2\text{-C}_6$ alkenyl, $\text{C}_2\text{-C}_6$ alkynyl, $\text{C}_6\text{-C}_{10}$ aryl, 5- to 10-membered heteroaryl, $\text{C}_3\text{-C}_7$ carbocyclyl or 5- to 10-membered heterocyclyl group; or R_1 is a group of formula $-\text{CH}_2\text{-CH(OR}_3\text{)-CH}_2\text{-(OR}_4\text{)}$, wherein R_3 and R_4 are each independently hydrogen atoms or $-(\text{C=O})\text{-R}_6$, wherein R_6 is an aliphatic group having from 3 to 29 carbon atoms; or R_1 is a group of formula $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is an integer of from 1 to 200; and wherein said alkyl, alkenyl, alkynyl and aliphatic groups are the same or different and are each unsubstituted or substituted with 1, 2 or 3 unsubstituted substituents which are the same or different and are selected from halogen atoms and $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_2\text{-C}_4$ alkenyloxy, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_2\text{-C}_4$ haloalkenyl, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_2\text{-C}_4$ haloalkenyloxy, hydroxyl, $-\text{SR}'$, and $-\text{NR}'\text{R}''$ groups where R' and R'' are the same or different and represent hydrogen or unsubstituted $\text{C}_1\text{-C}_2$ alkyl; said aryl, heteroaryl, carbocyclyl and heterocyclyl groups are the same or different and are each unsubstituted or substituted by 1, 2, 3 or 4 unsubstituted substituents which are the same or different and are selected from halogen atoms, and cyano, nitro, $\text{C}_1\text{-C}_4$ alkyl, $\text{C}_1\text{-C}_4$ alkoxy, $\text{C}_2\text{-C}_4$ alkenyl, $\text{C}_2\text{-C}_4$ alkenyloxy, $\text{C}_1\text{-C}_4$ haloalkyl, $\text{C}_2\text{-C}_4$ haloalkenyl, $\text{C}_1\text{-C}_4$ haloalkoxy, $\text{C}_2\text{-C}_4$ haloalkenyloxy, hydroxyl, $\text{C}_1\text{-C}_4$ hydroxyalkyl, $-\text{SR}'$ and $-\text{NR}'\text{R}''$ groups wherein each R' and R'' is the same or different and represents hydrogen or unsubstituted $\text{C}_1\text{-C}_4$ alkyl; and wherein the PUFA derivative is in the form of a racemate, a stereoisomer or a mixture of stereoisomers, wherein the PUFA is the sole active ingredient, wherein the skin inflammation is caused by atopic dermatitis, and wherein the amount of the PUFA is 1 wt% to 10 wt% of the pharmaceutical composition.

Detailed description of the invention

Figure 1 shows the results of an immunohistochemistry assay to determine the amount of COX-2 enzymes present in *ex vivo* porcine ear skin 0 and 6 hours after staining with diaminobenzidine.

Figure 2 shows the results of an immunohistochemistry assay to determine the amount of COX-2 enzymes present in *ex vivo* porcine ear skin, which has been treated with ketoprofen in fish oil, 0 and 6 hours after staining with diaminobenzidine.

Figure 3 shows the results of an immunohistochemistry assay to determine the amount of COX-2 enzymes present in *ex vivo* porcine ear skin, which has been treated with a representative compound of the invention, DGLA, 0 and 6 hours after staining with diaminobenzidine.

Figure 4 shows the results of an immunohistochemistry assay to determine the amount of COX-2 enzymes present in *ex vivo* porcine ear skin, which has been treated with a representative compound of the invention, 15-HETrE, 0 and 6 hours after staining with diaminobenzidine.

Figure 5 shows the results of a Western Blotting analysis to determine the effect of DGLA (second bar) and 15-HETrE (third bar) on COX-2 expression in porcine skin relative to water (first) bar. Levels in control were assigned a value of 100%.

Preferably the alkyl, alkenyl, alkynyl and aliphatic groups are unsubstituted or substituted with 1, 2 or 3, preferably 1 or 2, more preferably 1, unsubstituted substituents which are the same or different and are selected from halogen atoms and C₁-C₄ alkoxy, hydroxyl, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₁-C₄ haloalkyloxy and -NR'R'' wherein R' and R'' are the same or different and represent hydrogen or C₁-C₂ alkyl. More preferred substituents are halogen, C₁-C₄ alkoxy, hydroxyl and -NR'R'' groups where R' and R'' are the same or different and represent hydrogen or

WO 2010/125340

PCT/GB2010/000844

unsubstituted C₁-C₂ alkyl. Particularly preferred substituents include hydroxyl and -NR'R'' groups where R' and R'' are the same and represent hydrogen.

When the alkyl, alkenyl, alkynyl and aliphatic groups above are substituted by two or
 5 three substituents, it is preferred that not more than two substituents are selected from hydroxyl. More preferably, not more than one substituent is selected from hydroxyl.

Most preferably, the alkyl, alkenyl and alkynyl groups above are unsubstituted.

10 As used herein, a C₁-C₆ alkyl group is a linear or branched alkyl group containing from 1 to 6 carbon atoms, for example a C₁-C₄ alkyl group containing from 1 to 4 carbon atoms, preferably a C₁-C₂ alkyl group containing from 1 to 2 carbon atoms. Examples of C₁-C₄ alkyl groups include methyl, ethyl, n-propyl, i-propyl, n-butyl, i-butyl and t-butyl. For the avoidance of doubt, where two alkyl groups are present in a
 15 compound of the present invention, the alkyl groups may be the same or different.

As used herein, a C₂-C₆ alkenyl group is a linear or branched alkenyl group having at least one double bond of either *cis* or *trans* configuration where applicable and containing from 2 to 6 carbon atoms, for example a C₂-C₄ alkenyl group containing
 20 from 2 to 4 carbon atoms, such as -CH=CH₂ or -CH₂-CH=CH₂, -CH₂-CH₂-CH=CH₂, -CH₂-CH=CH-CH₃, -CH=C(CH₃)-CH₃ and -CH₂-C(CH₃)=CH₂, preferably a C₂ alkenyl group having 2 carbon atoms. For the avoidance of doubt, where two alkenyl groups are present in a compound of the present invention, they may be the same or
 different.

25 As used herein, a C₂-C₆ alkynyl group is a linear or branched alkynyl group containing from 2 to 6 carbon atoms, for example a C₂-C₄ alkynyl group containing from 2 to 4 carbon atoms, preferably a C₂ alkynyl group containing 2 carbon atoms. Exemplary alkynyl groups include -C≡CH or -CH₂-C≡CH, as well as 1- and 2-butyne, 2-methyl-
 30 2-propyne, 2-pentyne, 3-pentyne, 4-pentyne, 2-hexyne, 3-hexyne, 4-hexyne and 5-hexyne. For the avoidance of doubt, where two alkynyl groups are present in a compound of the present invention, they may be the same or different.

WO 2010/125340

PCT/GB2010/000844

Preferably, said C₁-C₆ alkyl group is a C₁-C₂ alkyl group, said C₂-C₆ alkenyl group is a C₂ alkenyl group and said C₂-C₆ alkynyl group is a C₂ alkynyl group.

As used herein, a halogen atom is chlorine, fluorine, bromine or iodine.

5

As used herein, a C₁-C₆ alkoxy group or C₂-C₆ alkenyloxy group is typically a said C₁-C₆ alkyl (e.g. a C₁-C₄ alkyl) group or a said C₂-C₆ alkenyl (e.g. a C₂-C₄ alkenyl) group respectively which is attached to an oxygen atom.

10 A haloalkyl, haloalkenyl, haloalkoxy or haloalkenyloxy group is typically a said alkyl, alkenyl, alkoxy or alkenyloxy group respectively which is substituted by one or more said halogen atoms. Typically, it is substituted by 1, 2 or 3 said halogen atoms. Preferred haloalkyl and haloalkoxy groups include perhaloalkyl and perhaloalkoxy groups, such as -CX₃ and -OCX₃ wherein X is a said halogen atom, for example
15 chlorine and fluorine.

As used herein, a C₁-C₄ alkylthio or C₂-C₄ alkenylthio group is typically a said C₁-C₄ alkyl group or a C₂-C₄ alkenyl group respectively which is attached to a sulphur atom, for example -S-CH₃.

20

As used herein, a C₁-C₄ hydroxyalkyl group is a C₁-C₄ alkyl group substituted by one or more hydroxy groups. Typically, it is substituted by one, two or three hydroxy groups. Preferably, it is substituted by a single hydroxy group.

25 As used herein, a C₆-C₁₀ aryl group is a monocyclic or polycyclic, preferably monocyclic, aromatic ring containing from 6 to 10 carbon atoms, for example a C₆ aryl group containing 6 carbon atoms. Examples of such aryl groups include phenyl, naphthalene and azulene. Phenyl is preferred.

30 As used herein, a 5- to 10- membered heteroaryl group is a monocyclic or polycyclic, preferably monocyclic, 5- to 10- membered aromatic ring, such as a 5- or 6- membered ring, containing at least one heteroatom, for example 1, 2, 3 or 4 heteroatoms, selected from O, S and N. When the ring contains 4 heteroatoms these are preferably all nitrogen atoms. Examples include thienyl, furyl, pyrrolyl,

imidazolyl, thiazolyl, isothiazolyl, pyrazolyl, oxazolyl, isoxazolyl, triazolyl, thiadiazolyl, oxadiazolyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, triazinyl and tetrazolyl groups. Thienyl, pyrrolyl, imidazolyl, thiazolyl, isothiazolyl, pyrazolyl, oxazolyl, isoxazolyl, triazolyl, pyridinyl, pyridazinyl, pyrimidinyl and pyrazinyl groups are preferred, e.g. pyrrolyl, imidazolyl, thiazolyl, isothiazolyl, pyrazolyl, oxazolyl, isoxazolyl, triazolyl, pyridinyl, pyridazinyl, pyrimidinyl and pyrazinyl groups. More preferred groups are thienyl, pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl and triazinyl, e.g. pyridinyl, pyridazinyl, pyrimidinyl, pyrazinyl, pyrrolyl and triazinyl, most preferably pyridinyl.

10

As used herein, a 5- to 10- membered heterocyclyl group is a non-aromatic, saturated or unsaturated monocyclic or polycyclic, preferably monocyclic, C₅₋₁₀ carbocyclic ring in which one or more, for example 1, 2, 3 or 4, of the carbon atoms are replaced with a moiety selected from N, O, S, S(O) and S(O)₂, and wherein one or more of the remaining carbon atoms is optionally replaced by a group -C(O)- or -C(S)-. When one or more of the remaining carbon atoms is replaced by a group -C(O)- or -C(S)-, preferably only one or two (more preferably two) such carbon atoms are replaced. Typically, the 5- to 10- membered heterocyclyl ring is a 5- to 6- membered ring.

20

Suitable heterocyclyl groups include azetidiny, oxetanyl, thietanyl, pyrrolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, dithiolanyl, dioxolanyl, pyrazolidinyl, piperidinyl, piperazinyl, hexahydropyrimidinyl, methylenedioxyphenyl, ethylenedioxyphenyl, thiomorpholinyl, S-oxo-thiomorpholinyl, S,S-dioxo-thiomorpholinyl, morpholinyl, 1,3-dioxolanyl, 1,4-dioxolanyl, trioxolanyl, trithianyl, imidazoliny, pyranyl, pyrazoliny, thioxolanyl, thioxothiazolidinyl, 1H-pyrazol-5-(4H)-onyl, 1,3,4-thiadiazol-2(3H)-thionyl, oxopyrrolidinyl, oxothiazolidinyl, oxopyrazolidinyl, succinimido and maleimido groups and moieties. Preferred heterocyclyl groups are pyrrolidinyl, imidazolidinyl, oxazolidinyl, isoxazolidinyl, thiazolidinyl, isothiazolidinyl, tetrahydrofuranyl, tetrahydrothienyl, tetrahydropyranyl, tetrahydrothiopyranyl, dithiolanyl, dioxolanyl, pyrazolidinyl, piperidinyl, piperazinyl, hexahydropyrimidinyl, thiomorpholinyl and morpholinyl groups and moieties.

30

WO 2010/125340

PCT/GB2010/000844

For the avoidance of doubt, although the above definitions of heteroaryl and heterocyclyl groups refer to an "N" moiety which can be present in the ring, as will be evident to a skilled chemist the N atom will be protonated (or will carry a substituent as defined below) if it is attached to each of the adjacent ring atoms via a single bond.

5

As used herein, a C₃-C₇ carbocyclic group is a non-aromatic saturated or unsaturated hydrocarbon ring having from 3 to 7 carbon atoms. Preferably it is a saturated or mono-unsaturated hydrocarbon ring (i.e. a cycloalkyl moiety or a cycloalkenyl moiety) having from 3 to 7 carbon atoms, more preferably having from 3 to 6 carbon atoms. Examples include cyclopropyl, cyclobutyl, cyclopentyl and cyclohexyl and their mono-unsaturated variants, more particularly cyclopentyl and cyclohexyl. A C₃-C₇ carbocyclyl group also includes C₃-C₇ carbocyclyl groups described above but wherein one or more ring carbon atoms are replaced by a group -C(O)-. More preferably, 0, 1 or 2 ring carbon atoms (most preferably 0) are replaced by -C(O)-.

15 Most preferably, said C₃-C₇ carbocyclyl group is cyclohexyl

Typically the aryl, heteroaryl, heterocyclyl and carbocyclyl groups in R₁ and R₅ are unsubstituted or substituted by 1, 2, 3 or 4 unsubstituted substituents, for example by 1, 2 or 3 unsubstituted substituents. Preferred substituents include halogen atoms and C₁-C₄ alkyl, C₂-C₄ alkenyl, C₁-C₄ alkoxy, C₂-C₄ alkenyloxy, C₁-C₄ haloalkyl, C₂-C₄ haloalkenyl, C₁-C₄ haloalkoxy, C₂-C₄ haloalkenyloxy, hydroxyl, mercapto, cyano, nitro, C₁-C₄ hydroxyalkyl, C₂-C₄ hydroxyalkenyl, C₁-C₄ alkylthio, C₂-C₄ alkenylthio and -NR'R'' groups wherein each R' and R'' is the same or different and represents hydrogen or C₁-C₄ alkyl. More preferred substituents include halogen atoms and

20 unsubstituted C₁-C₄ alkyl, C₁-C₄ alkoxy, hydroxyl, C₁-C₄ haloalkyl, C₁-C₄ haloalkoxy, C₁-C₄ hydroxyalkyl, cyano, nitro, -SR' and -NR'R'' groups where R' and R'' are the same or different and represent hydrogen or unsubstituted C₁-C₂ alkyl. More preferred substituents include halogen atoms, hydroxyl groups and C₁-C₂ alkyl and C₁-C₂ alkoxy groups.

30

Most preferably, the aryl, heteroaryl, heterocyclyl and carbocyclyl groups above are unsubstituted.

WO 2010/125340

PCT/GB2010/000844

When the aryl, heteroaryl, heterocyclyl and carbocyclyl groups in R₁ and R₅ are substituted by two, three or four substituents, it is preferred that not more than two substituents are selected from hydroxyl, cyano and nitro. More preferably, not more than one substituent is selected from hydroxyl, cyano and nitro.

5

As used herein, a pharmaceutically acceptable salt is a salt with a pharmaceutically acceptable acid or base. Pharmaceutically acceptable acids include both inorganic acids such as hydrochloric, sulphuric, phosphoric, diphosphoric, hydrobromic or nitric acid and organic acids such as citric, fumaric, maleic, malic, ascorbic, succinic, tartaric, benzoic, acetic, methanesulphonic, ethanesulphonic, benzenesulphonic or *p*-toluenesulphonic acid. Pharmaceutically acceptable bases include alkali metal (e.g. sodium or potassium) and alkali earth metal (e.g. calcium or magnesium) hydroxides and organic bases such as alkyl amines, aralkyl amines and heterocyclic amines.

10

The term "solvate" refers to a complex or aggregate formed by one or more molecules of a solute, i.e. compounds of the invention or pharmaceutically-acceptable salts thereof, and one or more molecules of a solvent. Such solvates are typically crystalline solids having a substantially fixed molar ratio of solute and solvent. Representative solvents include by way of example, water, methanol, ethanol, isopropanol, acetic acid, and the like. When the solvent is water, the solvate formed is a hydrate.

15

20

The compounds of the invention may contain a chiral center. Accordingly, they can be used in the form of a racemic mixture, an enantiomer, or a mixture enriched in one or more stereoisomer. The scope of the invention as described and claimed encompasses the racemic forms of the compounds of the invention as well as the individual enantiomers, and stereoisomer-enriched mixtures.

25

It will be appreciated that the term "or a pharmaceutically acceptable salt or solvate thereof" is intended to include all permutations of salts and solvates, such as solvates of pharmaceutically-acceptable salts of compounds of the invention.

30

R₅ and R₆ may be an aliphatic group having 3 to 29 carbon atoms. Typically, the aliphatic group is not cyclic. The aliphatic group is typically linear or branched,

WO 2010/125340

PCT/GB2010/000844

preferably linear. Typically the aliphatic group has 7 to 25 carbon atoms, more preferably 11 to 25 carbon atoms. The aliphatic group is typically unsubstituted or substituted with one hydroxyl group. The aliphatic group is preferably unsubstituted.

- 5 Aliphatic groups may be saturated, monounsaturated or polyunsaturated. Saturated aliphatic groups are preferred.

Typically, saturated aliphatic groups have from 7 to 25 carbon atoms, preferably 11 to 17 carbon atoms.

10

Monounsaturated aliphatic groups typically contain a single C=C double bond. The double bond has *cis* or *trans* configuration. The single double bond may be present at any point in the aliphatic group, but is typically 7 or 9 carbon atoms from the end of the aliphatic group distal to the (C=O) group to which the aliphatic group is attached.

- 15 Typically, monounsaturated aliphatic groups have from 7 to 25 carbon atoms, preferably 15 to 23 carbon atoms.

- 20 Polyunsaturated aliphatic groups typically contain two or more C=C double bonds, for example 2, 3, 4, 5 or 6 C=C double bonds. Each double bond may have *cis* or *trans* configuration. The double bonds may be present at any point in the aliphatic chain, but typically, the C=C double bond furthest from the (C=O) group to which the aliphatic group is attached is 3, 6 or 9 carbon atoms from the end of the aliphatic group distal to the (C=O) group to which the aliphatic group is attached. Typically, polyunsaturated aliphatic groups have from 7 to 25 carbon atoms, preferably 15 to 23 carbon atoms.

- Typically, said aliphatic group is the group R, wherein R-CO₂H is a fatty acid. Preferably, said fatty acid is lauric acid, myristic acid, palmitic acid, stearic acid, palmitoleic acid, *cis*-vaccenic acid, oleic acid, eicosenoic acid, erucic acid, nervonic acid, alpha-linolenic acid, stearidonic acid, eicosatrienoic acid, eicosatetraenoic acid, eicosapentaenoic acid, docosapentaenoic acid, docosahexaenoic acid, tetracosapentaenoic acid, tetracosahexaenoic acid, linoleic acid, gamma-linolenic acid, eicosadienoic acid, dihomo-gamma-linolenic acid, arachidonic acid,

WO 2010/125340


PCT/GB2010/000844

docosadienoic acid, adrenic acid, docosapentaenoic acid, or mead acid. More preferably, said fatty acid is lauric acid, myristic acid, palmitic acid, or stearic acid.

In one embodiment, the aliphatic group having 3 to 29 carbon atoms is the aliphatic group of a PUFA derivative of formula (I) as defined herein, i.e. the aliphatic group is of formula $-(\text{CH}_2)_3\text{-Alk}-(\text{CH}_2)_4\text{CH}_3$, wherein -Alk- is as defined herein.

In a preferred embodiment, the aliphatic group having 3 to 29 carbon atoms is the aliphatic group of dihomo-gamma-linolenic acid or 15-hydroxyeicosatrienoic acid, i.e. the aliphatic group is $-(\text{CH}_2)_6\text{-CH=CH-CH}_2\text{-CH=CH-CH}_2\text{-CH=CH-}(\text{CH}_2)_4\text{CH}_3$, wherein all of the C=C double bonds have *cis* configuration, or $-(\text{CH}_2)_6\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-[trans]CH=CH-CH(OH)-}(\text{CH}_2)_4\text{CH}_3$. Preferably, the aliphatic group having 3 to 29 carbon atoms is the aliphatic group of 15-hydroxyeicosatrienoic acid, i.e. the aliphatic group is $-(\text{CH}_2)_6\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-[trans]CH=CH-CH(OH)-}(\text{CH}_2)_4\text{CH}_3$.

In a more preferred embodiment, the PUFA derivative of formula (I) is of formula $\text{R}'(\text{C}=\text{O})\text{O-CH}_2\text{-CH(O(C}=\text{O)R}')\text{-CH}_2\text{-O(C}=\text{O)R}'$, wherein each R' is the aliphatic group of 15-hydroxyeicosatrienoic acid, i.e. R' is $-(\text{CH}_2)_6\text{-[cis]CH=CH-CH}_2\text{-[cis]CH=CH-[trans]CH=CH-CH(OH)-}(\text{CH}_2)_4\text{CH}_3$. Thus, the PUFA derivative of formula (I) is preferably



It is to be understood that the left hand side of the -Alk- moiety is bonded to the unsaturated carbon chain bearing the -COOR_1 moiety and the right hand side of the -Alk- group is bonded to the saturated carbon chain.

WO 2010/125340

PCT/GB2010/000844

Typically, -Alk- is $-\text{CH}(\text{OR}_2)-[\textit{trans}]\text{CH}=\text{CH}-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-\text{C}_3\text{H}_6-$,
 $-(\text{CH}_2)_3-\text{CH}(\text{OR}_2)-[\textit{trans}]\text{CH}=\text{CH}-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-$, or $-(\text{CH}_2)_3-$
 $[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-[\textit{trans}]\text{CH}=\text{CH}-\text{CH}(\text{OR}_2)-$. Preferably, -Alk- is
 $(\text{CH}_2)_3-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-$.

5

Typically, R_1 is a hydrogen atom; or R_1 is a C_1 - C_4 alkyl, C_2 - C_4 alkenyl, C_2 - C_4 alkynyl,
 C_6 aryl, 5- to 6-membered heteroaryl, C_3 - C_6 carbocyclyl or 5- to 6-membered
heterocyclyl group; or R_1 is a group of formula $-\text{CH}_2-\text{CH}(\text{OR}_3)-\text{CH}_2-(\text{OR}_4)$, wherein
 R_3 and R_4 are as defined herein; or R_1 is a group of formula $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$,
10 wherein m is as defined herein, wherein said alkyl, alkenyl and alkynyl groups are the
same or different and are each unsubstituted or substituted with 1, or 2 unsubstituted
substituents which are the same or different and are selected from halogen atoms, C_1 -
 C_4 alkoxy, hydroxyl, and $-\text{NR}'\text{R}''$ groups where R' and R'' are the same or different
and represent hydrogen or unsubstituted C_1 - C_2 alkyl; and said aryl, heteroaryl,
15 carbocyclyl and heterocyclyl groups are the same or different and are each
unsubstituted or substituted by 1, 2 or 3 unsubstituted substituents which are the same
or different and are selected from halogen atoms, and cyano, nitro, C_1 - C_4 alkyl, C_1 - C_4
alkoxy, and $-\text{NR}'\text{R}''$ groups wherein each R' and R'' is the same or different and
represents hydrogen or unsubstituted C_1 - C_2 alkyl group.

20

Preferably, R_1 is a hydrogen atom; or R_1 is an unsubstituted C_1 - C_4 alkyl group; or R_1
is a group of formula $-\text{CH}_2-\text{CH}(\text{OR}_3)-\text{CH}_2-(\text{OR}_4)$, wherein R_3 and R_4 are as defined
herein; or R_1 is a group of formula $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is as defined herein.

25

More preferably, R_1 is a hydrogen atom; or R_1 is a group of formula $-\text{CH}_2-\text{CH}(\text{OR}_3)-$
 $\text{CH}_2-(\text{OR}_4)$, wherein R_3 and R_4 are as defined herein, and wherein at least one of R_3 or
 R_4 is $-(\text{C}=\text{O})-\text{R}_6$, wherein R_6 is as defined herein.

Most preferably, R_1 is a hydrogen atom.

30

m is typically an integer of from 5 to 150, preferably from 10 to 50.

R_3 is typically $-(\text{C}=\text{O})-\text{R}_6$, wherein R_6 is as defined herein.

WO 2010/125340

PCT/GB2010/000844

R₄ is typically -(C=O)-R₆, wherein R₆ is as defined herein.

Preferably, both R₃ and R₄ are -(C=O)-R₆, wherein each R₆ may be the same or different and is as defined herein.

5

Typically, when R₃ and R₄ are both -(C=O)-R₆, then R₅ is not an aliphatic group having 3 to 29 carbon atoms.

R₆ is an aliphatic group having from 3 to 29 carbon atoms, as defined herein.

10 Typically, said aliphatic group is saturated. Typically, R₆ is an aliphatic group having 7 to 25 carbon atoms, preferably 11 to 17 carbon atoms. Preferably, R₆ is a group R, wherein R-CO₂H is auric acid, myristic acid, palmitic acid, or stearic acid.

Typically, R₂ is a hydrogen atom; or R₂ is a group -(C=O)-R₅, wherein R₅ is a C₁-C₄
 15 alkyl, C₂-C₄ alkenyl, C₂-C₄ alkynyl, C₆ aryl, 5- to 6-membered heteroaryl, C₂-C₆ carbocyclyl or 5- to 6-membered heterocyclyl group, or R₅ is an aliphatic group having from 3 to 29 carbon atoms; or R₂ is a group of formula -(CH₂OCH₂)_nOH, wherein n is as defined herein, wherein said alkyl, alkenyl and alkynyl groups are the same or different and are each unsubstituted or substituted with 1, or 2 unsubstituted
 20 substituents which are the same or different and are selected from halogen atoms, C₁-C₄ alkoxy, hydroxyl, and -NR'R'' groups where R' and R'' are the same or different and represent hydrogen or unsubstituted C₁-C₂ alkyl; and said aryl, heteroaryl, carbocyclyl and heterocyclyl groups are the same or different and are each unsubstituted or substituted by 1, 2 or 3 unsubstituted substituents which are the same
 25 or different and are selected from halogen atoms, and cyano, nitro, C₁-C₄ alkyl, C₁-C₄ alkoxy, and -NR'R'' groups wherein each R' and R'' is the same or different and represents hydrogen or unsubstituted C₁-C₂ alkyl group.

Preferably, R₂ is a hydrogen atom; or R₂ is a group -(C=O)-R₅, wherein R₅ is
 30 unsubstituted C₁-C₄ alkyl; or R₂ is a group -(C=O)-R₅, wherein R₅ is an aliphatic group having from 3 to 29 carbon atoms; or R₂ is a group of formula -(CH₂OCH₂)_nOH, wherein n is as defined herein.

WO 2010/125340

PCT/GB2010/000844

More preferably, R_2 is a hydrogen atom; or R_2 is a group $-(C=O)-R_5$, wherein R_5 is an aliphatic group having from 3 to 29 carbon atoms; or R_2 is a group of formula $-(CH_2OCH_2)_nOH$, wherein n is as defined herein.

5 Most preferably, R_2 is a hydrogen atom.

n is typically an integer of from 5 to 150, preferably from 10 to 50.

10 When R_5 is an aliphatic group having 3 to 29 carbon atoms, said aliphatic group is as defined herein. Typically, said aliphatic group is saturated. Typically, R_5 is an aliphatic group having 7 to 25 carbon atoms, preferably 11 to 17 carbon atoms. Preferably, R_5 is a group R , wherein $R-CO_2H$ is auric acid, myristic acid, palmitic acid, or stearic acid.

15 In one embodiment, the PUFA derivative of formula (I) is present as a racemic mixture of the R and S enantiomers.

In another embodiment, the PUFA derivative of formula (I) is present as the R enantiomer.

20

In another embodiment, the PUFA derivative of formula (I) is present as the S enantiomer.

Typically, the mammal is a human.

25

In a preferred embodiment, -Alk- is $-(CH(OR_2)-[trans]CH=CH-[cis]CH=CH-CH_2-[cis]CH=CH-C_3H_6-$, $-(CH_2)_3-CH(OR_2)-[trans]CH=CH-[cis]CH=CH-CH_2-[cis]CH=CH-$, or $-(CH_2)_3-[cis]CH=CH-CH_2-[cis]CH=CH-[trans]CH=CH-CH(OR_2)-$; R_1 is a hydrogen atom, an unsubstituted C_1-C_4 alkyl group, or a group of formula $-CH_2-CH(OR_3)-CH_2-(OR_4)$, wherein R_3 and R_4 are each independently hydrogen atoms or $-(C=O)-R_6$, wherein R_6 is a linear aliphatic group having from 11 to 25 carbon atoms, which aliphatic group is unsubstituted or substituted with one hydroxyl group, or R_1 is a group of formula

30

WO 2010/125340

PCT/GB2010/000844

5 $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is an integer of from 5 to 150; and R_2 is a hydrogen atom, a group $-(\text{C}=\text{O})-\text{R}_5$, wherein R_5 is unsubstituted C_1 - C_4 alkyl, or a group $-(\text{C}=\text{O})-\text{R}_5$, wherein R_5 is a linear aliphatic group having from 11 to 25 carbon atoms, which aliphatic group is unsubstituted or substituted with one hydroxyl group; or R_2 is a group of formula $-(\text{CH}_2\text{OCH}_2)_n\text{OH}$, wherein n is an integer of from 5 to 150.

In a more preferred embodiment, $-\text{Alk}-$ is $-(\text{CH}_2)_3-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-[\textit{trans}]\text{CH}=\text{CH}-\text{CH}(\text{OR}_2)-$; R_1 is a hydrogen atom, an unsubstituted C_1 - C_4 alkyl group, or a group of formula $-\text{CH}_2-\text{CH}(\text{OR}_3)-\text{CH}_2-(\text{OR}_4)$, wherein R_3 and R_4 are each
 10 independently hydrogen atoms or $-(\text{C}=\text{O})-\text{R}_6$, wherein R_6 is a linear aliphatic group having from 11 to 25 carbon atoms, which aliphatic group is unsubstituted or substituted with one hydroxyl group, or R_1 is a group of formula
 $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is an integer of from 5 to 150; and R_2 is a hydrogen atom, a group $-(\text{C}=\text{O})-\text{R}_5$, wherein R_5 is unsubstituted C_1 - C_4 alkyl, or a group
 15 $-(\text{C}=\text{O})-\text{R}_5$, wherein R_5 is a linear aliphatic group having from 11 to 25 carbon atoms, which aliphatic group is unsubstituted or substituted with one hydroxyl group; or R_2 is a group of formula $-(\text{CH}_2\text{OCH}_2)_n\text{OH}$, wherein n is an integer of from 5 to 150.

In a most preferred embodiment, $-\text{Alk}-$ is $-(\text{CH}_2)_3-[\textit{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\textit{cis}]\text{CH}=\text{CH}-[\textit{trans}]\text{CH}=\text{CH}-\text{CH}(\text{OR}_2)-$; R_1 is a hydrogen atom, a group of formula
 $-\text{CH}_2-\text{CH}(\text{OR}_3)-\text{CH}_2-(\text{OR}_4)$, wherein R_3 and R_4 are each independently hydrogen atoms or $-(\text{C}=\text{O})-\text{R}_6$, wherein R_6 is an unsubstituted linear, saturated aliphatic group
 20 having from 11 to 17 carbon atoms, and wherein at least one of R_3 or R_4 is $-(\text{C}=\text{O})-\text{R}_6$; and R_2 is a hydrogen atom, or a group $-(\text{C}=\text{O})-\text{R}_5$, wherein R_5 is an unsubstituted
 25 linear, saturated aliphatic group having from 11 to 17 carbon atoms, or R_2 is a group of formula $-(\text{CH}_2\text{OCH}_2)_n\text{OH}$, wherein n is an integer of from 10 to 50.

Typically, in the compounds of the present invention, including the preferred embodiments set out above, (a) R_1 is a group of formula $-\text{CH}_2-\text{CH}(\text{OR}_3)-\text{CH}_2-(\text{OR}_4)$,
 30 wherein R_3 and R_4 each independently represent a hydrogen atom or $-(\text{C}=\text{O})-\text{R}_6$, wherein R_6 is a saturated aliphatic group having from 3 to 29 carbon atoms, wherein at least one of R_3 or R_4 is $-(\text{C}=\text{O})-\text{R}_6$; or
 (b) R_1 is a group of formula $-(\text{CH}_2\text{OCH}_2)_m\text{OH}$, wherein m is as defined herein; and/or

WO 2010/125340

PCT/GB2010/000844

(c) R_2 is a group $-(C=O)-R_5$; wherein R_5 is a saturated aliphatic group having from 3 to 29 carbon atoms; or

(d) R_2 is a group of formula $-(CH_2OCH_2)_nOH$, wherein n is as defined herein. Such compounds will be particularly lipophilic, which is advantageous in some instances.

5

In this preferred embodiment, preferably (a) R_1 is a group of formula $-CH_2-CH(OR_3)-CH_2-(OR_4)$, wherein R_3 and R_4 are each independently hydrogen atoms or $-(C=O)-R_6$, wherein R_6 is a saturated aliphatic group having 3 to 29 carbon atoms, wherein at least one of R_3 or R_4 is $-(C=O)-R_6$; and/or (c) R_2 is a group $-(C=O)-R_5$, wherein R_5 is a


10

In a more preferred embodiment, both R_1 and R_2 are hydrogen atoms.

In a particularly preferred embodiment, -Alk- is $-(CH_2)_3-[cis]CH=CH-CH_2-$

15

$[cis]CH=CH-[trans]CH=CH-CH(OR_2)-$, and R_1 and R_2 are both hydrogen atoms. In this embodiment, the PUFA derivative of formula (I) is 15-HETrE and is represented by the formula



20

Typically, compounds of the invention are administered as one or more treatments per day, preferably from 1 to 4 treatments per day, more preferably from 1 to 2 treatments

25

Typically, compounds of the invention are administered at a daily dosage of from 0.1 mg / m² / day to 1 kg / m² / day, preferably from 1 mg / m² / day to 100 g / m² / day, more preferably from 10 mg / m² / day to 10 g / m² / day, most preferably from 100

30

mg / m² / day to 1 g / m² / day.

WO 2010/125340

PCT/GB2010/000844

Skin inflammation includes rashes, hives, blisters and/or wheals and may be caused by eczema, exposure to radiation, autoimmune diseases, and/or uremic pruritis.

5 In a preferred embodiment, the skin inflammation is caused by atopic eczema, contact dermatitis, psoriasis or uremic pruritis.

In a further preferred embodiment, the skin inflammation is caused by exposure of the skin to electromagnetic radiation. This includes, for example, exposure to sunlight, heat, X-rays or radioactive materials. Thus, in this embodiment, the compound of the
10 present invention is typically used to treat sunburn.

The term eczema is applied to a wide range of skin conditions with a variety of aetiologies. In general, eczema is categorised by inflammation of the epidermis. Common symptoms associated with eczema include dryness, recurring skin rashes,
15 redness, skin edema (swelling), itching, dryness, crusting, flaking, blistering, cracking, oozing, and bleeding. Eczema includes atopic eczema (atopic dermatitis), contact dermatitis, xerotic eczema, seborrhoeic dermatitis, dyshydrosis, discoid eczema, venous eczema, dermatitis herpetiformis, neurodermatitis, and autoeczematisation. Eczema is typically atopic eczema or contact dermatitis.

20 Atopic eczema is primarily aggravated by contact with or intake of allergens, which include animal hair and dander, food allergens, for example nuts or shellfish, or drugs, for example penicillin.

25 Contact dermatitis includes allergic contact dermatitis, irritant contact dermatitis and photocontact dermatitis.

Electromagnetic radiation includes radio waves, microwaves, terahertz radiation, infrared radiation, visible light, ultraviolet radiation, X-rays and gamma rays.

30 Electromagnetic radiation is preferably infrared radiation, visible light, ultraviolet radiation, X-rays and gamma rays, more preferably ultraviolet radiation, X-rays and gamma rays.

Autoimmune diseases can involve an autoimmune response against the skin. Examples of such autoimmune diseases are lupus and psoriasis.

Uremic pruritis is a disorder of the skin associated with chronic renal failure. It also frequently affects patients undergoing dialysis treatment.

Typically, the compound of the present invention is co-administered with a corticosteroid.

Suitable corticosteroids to be used for co-administration with compounds of the invention are clobetasol dipropionate, betamethasone dipropionate, halbetasol propionate, diflorasone diacetate, fluocinonide, halcinonide, amcinonide, desoximetasone, triamcinolone acetonide, mometasone furoate, fluticasone propionate, fluocinolone acetonide, hydrocortisone valerate, hydrocortisone butyrate, triamcinalone acetonide, desonide, prednicarbate, prednisolone, methylprednisolone, dexamethasone, naflocort, deflazacort, halopredone acetate, budesonide, beclomethasone dipropionate, hydrocortisone, clocortolone pivalate, methylprednisolone aceponate, dexamethasone palmitoate, tipredane, hydrocortisone aceponate, alclometasone dipropionate, halometasone, methylprednisolone suleptanate, rimexolone, prednisolone farnesylate, ciclesonide, deprodone propionateloteprednol etabonate, betamethasone butyrate propionate, flunisolide, prednisone, dexamethasone sodium phosphate, triamcinolone, betamethasone 17-valerate, betamethasone, betamethasone dipropionate, hydrocortisone acetate, hydrocortisone sodium succinate, prednisolone sodium phosphate and hydrocortisone probutate.

Preferred corticosteroids corticosteroids to be used for co-administration with compounds of the invention are clobetasol dipropionate, betamethasone dipropionate, halbetasol propionate, diflorasone diacetate, fluocinonide, halcinonide, amcinonide, desoximetasone, triamcinolone acetonide, mometasone furoate, fluticasone propionate, fluocinolone acetonide, hydrocortisone valerate, hydrocortisone butyrate, triamcinalone acetonide, desonide, and prednicarbate.

Any reference to corticosteroids within the scope of the present invention includes a reference to salts or derivatives thereof which may be formed from the corticosteroids. Examples of possible salts or derivatives include: sodium salts, sulphobenzoates, phosphates, isonicotinates, acetates, propionates, dihydrogen phosphates, palmitates, pivalates, farnesylates, aceponates, suleptanates, prednicarbates, furoates or acetonides. In some cases the corticosteroids may also occur in the form of their hydrates.

It is a particular finding of the present invention that co-administration of a compound of the present invention with a corticosteroid enables the amount of corticosteroid administered to be reduced, without reducing the efficacy of the treatment. Thus, in a preferred embodiment, the compound of the present invention is co-administered with a corticosteroid, wherein the corticosteroid is administered at a daily dosage of 50% or less, preferably 25% or less, more preferably 10% or less, most preferably 5% or less, of the recommended daily dosage of said corticosteroid in said mammal.

The person skilled in the art is well aware of the recommended daily dosages of corticosteroids in mammals. For example, the recommended daily dosage of hydrocortisone in humans is approximately $0.35 \text{ g} / \text{m}^2 / \text{day}$. The recommended daily dosage of clobetasol dipropionate in humans is 0.009 to $0.018 \text{ g} / \text{m}^2 / \text{day}$.

Thus, compounds of the present invention are typically coadministered with hydrocortisone, wherein the daily dosage of hydrocortisone is $0.175 \text{ g} / \text{m}^2 / \text{day}$ or less, preferably $0.0875 \text{ g} / \text{m}^2 / \text{day}$ or less, more preferably $0.035 \text{ g} / \text{m}^2 / \text{day}$, most preferably $0.0175 \text{ g} / \text{m}^2 / \text{day}$ or less. Compounds of the present invention are typically coadministered with clobetasol dipropionate, wherein the daily dosage of clobetasol dipropionate is $0.009 \text{ g} / \text{m}^2 / \text{day}$ or less, preferably $0.0045 \text{ g} / \text{m}^2 / \text{day}$ or less, more preferably $0.0018 \text{ g} / \text{m}^2 / \text{day}$ or less, most preferably $0.0009 \text{ g} / \text{m}^2 / \text{day}$ or less.


The compounds of the present invention are typically coadministered with a further therapeutic agent, which is not a corticosteroid, which is effective in treating skin conditions/diseases. Such therapeutic agents are well known to the skilled person and

WO 2010/125340

PCT/GB2010/000844

include, but are not limited to, immunomodulators, antibiotics, immunosuppressants and anti-itch drugs.

- 5 Compounds of the invention are typically commercially available, or may be prepared by analogy with known methods. Thus, for example, both DGLA and 15-HETrE are commercially available. These available fatty acids can easily be derivatised to obtain PUFA derivatives of formula (I) by known methods.
- 10 For example, PUFA derivatives of formula (I) as defined herein, wherein R_1 is a C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_6 - C_{10} aryl, 5- to 10-membered heteroaryl, C_3 - C_7 carbocyclyl or 5- to 10-membered heterocyclyl group; or R_1 is a group of formula $-CH_2-CH(OR_3)-CH_2-(OR_4)$, wherein R_3 and R_4 are as defined herein; or R_1 is a group of formula $-(CH_2OCH_2)_mOH$, wherein m is as defined herein, can be prepared by
- 15 esterifying a compound of formula



- wherein -Alk- is as defined herein and X is a leaving group, for example a halogen
- 20 atom, a tosylate or mesylate group with an alcohol of formula $R_1'-OH$, wherein R_1' is a C_1 - C_6 alkyl, C_2 - C_6 alkenyl, C_2 - C_6 alkynyl, C_6 - C_{10} aryl, 5- to 10-membered heteroaryl, C_3 - C_7 carbocyclyl or 5- to 10-membered heterocyclyl group; or R_1' is a group of formula $-CH_2-CH(OR_3)-CH_2-(OR_4)$, wherein R_3 and R_4 are as defined herein; or R_1' is a group of formula $-(CH_2OCH_2)_mOH$, wherein m is as defined herein,
- 25 to obtain a PUFA derivative of formula (I) as defined herein. Alternatively, X may be a hydroxyl group. In that case, the reaction is preferably carried out under acidic conditions, or in the presence of a suitable catalyst, for example pyridine.
- Compounds of formula $R_1'-OH$ are typically commercially available or may be prepared by analogy with known methods.

30

When -Alk- is $-CH(OR_2)-[trans]CH=CH-[cis]CH=CH-CH_2-[cis]CH=CH-C_3H_6-$, $-(CH_2)_3-CH(OR_2)-[trans]CH=CH-[cis]CH=CH-CH_2-[cis]CH=CH-$, or $-(CH_2)_3-$

WO 2010/125340

PCT/GB2010/000844

[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-, PUFA derivatives of formula (I) as defined herein, wherein R₂ is a group -(C=O)-R₅, wherein R₅ is a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, a C₆-C₁₀ aryl, a 5- to 10-membered heteroaryl, a C₃-C₇ carbocyclyl or a 5- to 10-membered heterocyclyl group, or R₅ is an aliphatic group having from 3 to 29 carbon atoms, can be prepared by treating a compound of formula (I) as defined herein, wherein -Alk- is -CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-C₃H₆-, -(CH₂)₃-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-, or -(CH₂)₃-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-, and R₂ is a hydrogen atom, with a carboxylic acid derivative Y-(C=O)-R'₅, wherein R'₅ is a C₁-C₆ alkyl, C₂-C₆ alkenyl, C₂-C₆ alkynyl, a C₆-C₁₀ aryl, a 5- to 10-membered heteroaryl, a C₃-C₇ carbocyclyl or a 5- to 10-membered heterocyclyl group, or R'₅ is an aliphatic group having from 3 to 29 carbon atoms, and Y is a leaving group, for example a halogen atom, a tosylate or mesylate group. Compounds of formula Y-(C=O)-R'₅ are typically commercially available or may be prepared by analogy with known methods.

When -Alk- is -CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-C₃H₆-, -(CH₂)₃-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-, or -(CH₂)₃-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-, PUFA derivatives of formula (I) as defined herein, wherein R₂ is a group of formula -(CH₂OCH₂)_nOH, wherein n is as defined herein, can be prepared by treating a compound of formula (I) as defined herein, wherein -Alk- is -CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-C₃H₆-, -(CH₂)₃-CH(OR₂)-[*trans*]CH=CH-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-, or -(CH₂)₃-[*cis*]CH=CH-CH₂-[*cis*]CH=CH-[*trans*]CH=CH-CH(OR₂)-, and R₂ is a hydrogen atom, with a compound of formula Z-(CH₂OCH₂)_nOH, wherein n is as defined herein and Z is a good leaving group, for example a halogen atom, a tosylate or mesylate group. Compounds of formula Z-(CH₂OCH₂)_nOH are typically commercially available or may be prepared by analogy with known methods.

The present invention also provides a pharmaceutical composition suitable for topical administration comprising a PUFA derivative, as defined herein, or a pharmaceutically acceptable salt, or solvate thereof, and a pharmaceutically acceptable diluent or carrier, for use in treating skin inflammation, as defined herein, in a mammal, as defined herein.

WO 2010/125340

PCT/GB2010/000844

Preferred pharmaceutical compositions are sterile and pyrogen free.

The pharmaceutical composition is typically in the form of a gel, ointment, cream or
5 lotion.

When said pharmaceutical composition is a gel it typically comprises a hydrophilic
polymer such as cross-linked polyethylene glycol, cross-linked starch or polyvinyl
pyrrolidone.

10

An ointment, cream or lotion typically contains an aqueous phase and an oleaginous
phase in admixture.

The pharmaceutical composition may additionally contain one or more emollients,
15 emulsifiers, thickeners and/or preservatives, particularly when it is a cream or
ointment.

Emollients suitable for inclusion in creams or ointments of the present invention are
typically long chain alcohols, for example a C8-C22 alcohol such as cetyl alcohol,
20 stearyl alcohol and cetearyl alcohol, hydrocarbons such as petrolatum and light
mineral oil, or acetylated lanolin. The total amount of emollient in the pharmaceutical
composition is preferably about 5 wt% to about 30 wt%, and more preferably about 5
wt% to about 10 wt% based on the total weight of the pharmaceutical composition.

25 The emulsifier is typically a nonionic surface active agent, e.g., polysorbate 60
(available from ICI Americas), sorbitan monostearate, polyglyceryl-4 oleate and
polyoxyethylene(4)lauryl ether. Generally the total amount of emulsifier is about 2
wt% to about 14 wt%, and more preferably about 2 wt% to about 6 wt% by weight
based on the total weight of the pharmaceutical composition.

30

Pharmaceutically acceptable thickeners, such as Veegum.TM.K (available from R. T.
Vanderbilt Company, Inc.), and long chain alcohols (i.e. C8-C22 alcohols such as
cetyl alcohol, stearyl alcohol and cetearyl alcohol) can be used. The total amount of

WO 2010/125340

PCT/GB2010/000844

thickener present is preferably about 3 wt% to about 12 wt% based on the total weight of the pharmaceutical composition.

5 Preservatives such as methylparaben, propylparaben and benzyl alcohol can be present in the pharmaceutical composition. The appropriate amount of such preservative(s) is known to those skilled in the art. The pharmaceutical composition may also comprise a fat-soluble antioxidant such as ascorbyl palmitate, tocopherol and/or ascorbic acid in the presence of lecithin.

10 Optionally, an additional solubilizing agent such as benzyl alcohol, lactic acid, acetic acid, stearic acid or hydrochloric acid can be included in the pharmaceutical composition. If an additional solubilizing agent is used, the amount present is preferably about 1 wt% to about 12 wt% based on the total weight of the pharmaceutical composition.

15

Optionally, the pharmaceutical composition can contain a humectant such as glycerin and a skin penetration enhancer such as butyl stearate, urea and DMSO.

It is known to those skilled in the art that a single ingredient can perform more than one function in a cream, i.e., cetyl alcohol can serve both as an emollient and as a
20 thickener.

In one embodiment, the pharmaceutical composition is in the form of a cream. The cream typically consists of an oil phase and a water phase mixed together to form an emulsion. Preferably, the cream comprises an oil-in-water emulsion. Preferably, the
25 amount of water present in a cream of the invention is about 45 wt% to about 85 wt% based on the total weight of the cream.

Where the pharmaceutical composition is in the form of an ointment, it typically comprises a pharmaceutically acceptable ointment base such as petrolatum, or
30 polyethylene glycol 400 (available from Union Carbide) in combination with polyethylene glycol 3350 (available from Union Carbide). The amount of ointment base present in an ointment of the invention is preferably about 60 wt% to about 95 wt% based on the total weight of the ointment.

In the pharmaceutical composition of the present invention, the amount of the PUFA derivative of formula (I) is typically from 0.01 wt% to 50wt%, preferably from 0.5wt% to 25 wt%, more preferably from 1 wt% to 10 wt%, for example about 5 wt%, based on the total weight of the pharmaceutical composition.

The compound / composition of the present invention is formulated for topical administration and it may be administered to a patient in an amount such that from 0.00001 to 10 g, preferably from 0.0001 to 1 g active ingredient is delivered per m² of the area being treated.

Pharmaceutical compositions of the present invention may additionally comprise one or more corticosteroids as defined herein. The amount of the corticosteroid present in the pharmaceutical composition of the invention is typically 50% or less, preferably 25% or less, more preferably 10% or less, most preferably 5% or less, of the recommended amount of said corticosteroid in a commercially available formulation. The skilled person is well aware of the amount of corticosteroids present in various topical formulations. For example, the amount of hydrocortisone present in most commercially available formulations is typically 1 % w/w. The amount of clobetasol dipropionate present in most commercially available formulations is typically 0.0525 % w/w.

Thus, pharmaceutical compositions of the present invention typically comprise hydrocortisone in an amount of 0.5 % w/w or less, preferably 0.025 % w/w or less, more preferably 0.01 % w/w or less, most preferably 0.005 % w/w or less. Pharmaceutical compositions of the present invention typically comprise clobetasol dipropionate in an amount of 0.026 % w/w or less, preferably 0.013 % w/w or less, more preferably 0.0053 % w/w or less, most preferably 0.0026 % w/w or less.

The pharmaceutical compositions of the present invention typically comprise one or more further therapeutic agents, which are not corticosteroids, as defined herein. The amount of the one or more further therapeutic agents, which are not corticosteroids, present in the composition will be evident to the person skilled in the art.

WO 2010/125340

PCT/GB2010/000844


Pharmaceutical compositions of the present invention may be prepared simply by admixing the ingredients in a suitable manner.

The present invention also provides the use of a compound which is a PUFA derivative of formula (I) as defined herein or a pharmaceutically acceptable salt, or solvate thereof, in the manufacture of a medicament for use in treating skin inflammation, as defined herein, in a mammal, as defined herein, by topical administration.

The present invention also provides a method of treating skin inflammation, as defined herein, in a mammal, as defined herein, which method comprises administering to the skin of said mammal a therapeutically effective amount of a compound which is a PUFA derivative of formula (I) as defined herein or a pharmaceutically acceptable salt, or solvate thereof.

15

The present invention also provides a pharmaceutical composition comprising a polyunsaturated fatty acid (PUFA) derivative of formula (II),



20

II

or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is $-(\text{CH}_2)_3-[\text{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\text{cis}]\text{CH}=\text{CH}-\text{CH}_2-[\text{cis}]\text{CH}=\text{CH}-$; and
 R_1 is as defined herein;

25 which composition is for use in treating skin inflammation in a mammal, by topical administration, and is substantially free of lithium salts and glucocorticoids.

As the skilled person will appreciate, in this embodiment, the pharmaceutically acceptable salt of the PUFA derivative of formula (II) is not a lithium salt.

30


WO 2010/125340

PCT/GB2010/000844

Typically, the pharmaceutical composition comprising a polyunsaturated fatty acid (PUFA) derivative of formula (II) contains the PUFA derivative of formula (II) as sole active ingredient.


- 5 Preferably, the pharmaceutical composition comprising a polyunsaturated fatty acid (PUFA) derivative of formula (II) consists of the PUFA derivative of formula (II) and a pharmaceutically acceptable diluent or carrier, as defined herein.

Preferably, the PUFA derivative of formula (II) is of formula $R''(C=O)O-CH_2-$
 10 $CH(O(C=O)R'')-CH_2-O(C=O)R''$, wherein each R'' is the same and is the aliphatic group of dihomo- γ -linolenic acid, i.e. R'' is $-(CH_2)_6-CH=CH-CH_2-CH=CH-$
 $CH_2-CH=CH-(CH_2)_4CH_3$, wherein all of the C=C double bonds have *cis* configuration. Thus, the PUFA derivative of formula (II) is preferably



15

In a particularly preferred embodiment, R_1 is a hydrogen atom. In this embodiment, the PUFA derivative of formula (II) is DGLA and is represented by the formula



20

The present invention also provides the use of a pharmaceutical composition comprising a polyunsaturated fatty acid (PUFA) derivative of formula (II) as defined herein in the manufacture of a medicament for use in treating skin inflammation, as

WO 2010/125340

PCT/GB2010/000844

defined herein, in a mammal, as defined herein, by topical administration, wherein the composition is substantially free of lithium salts and glucocorticoids.

- 5 The present invention also provides a method of treating skin inflammation, as defined herein, in a mammal, as defined herein, which method comprises administering to the skin of said mammal a therapeutically effective amount of a pharmaceutical composition comprising a polyunsaturated fatty acid (PUFA) derivative of formula (II) as defined herein, wherein the composition is substantially free of lithium salts and glucocorticoids.

i0

WO 2010/125340

PCT/GB2010/000844

EXAMPLES

Immunohistochemistry analyses were carried out to measure the degree of expression of COX-2 enzymes in *ex vivo* porcine ear skin. The COX-2 family of enzymes have
5 been strongly linked to inflammation and have been found to be present in increased amounts in inflamed tissue. Thus, a decreased level of COX-2 in the skin corresponds with reduced inflammation of the skin.

Example 1

10

Freshly excised porcine ears were immersed in iced Hank's buffer for transport from the abattoir to the laboratory. Upon arrival, the porcine ears were first washed with running tap water, and the full thickness skin was liberated from underlying cartilage by blunt dissection using a scalpel and hairs were removed with an electrical razor.

15

The skin was used within 2 h of slaughter. The full thickness skin was cut into approximately 2cm×2cm and placed in Hanks balance salt to maintain skin viability.

20

A 2mm strip of skin was obtained using a surgical scalpel and the skin strip then fixed using a 40% formaldehyde solution, dehydrated and set in molten wax. The sections were cut using a Shandon Finesse microtome (Thermo Scientific, Waltham, MA, USA) to a thickness of 5µm and were placed onto 2.5 cm x 7.5 cm x 1mm pre-cleaned microslides. The section of skin was stained with diaminobenzidine, which binds to COX-2 enzymes and the slides visualised on a light microscope with image capture facility.

25

Skin membranes were mounted in Franz diffusion cells using Hanks buffer as receptor phase. Water was used as the donor phase. After 6 hours the skin was removed from the Franz cell apparatus, excess donor phase was removed and the skin wiped clean with a clean paper tissue. The diffusional area was then cut into
30 approximately 2mm strips using a surgical scalpel and the skin strips then using a 40% formaldehyde solution, dehydrated and set in molten wax. The sections were cut using a Shandon Finesse microtome (Thermo Scientific, Waltham, MA, USA) to a thickness of 5µm and were placed onto 2.5 cm x 7.5 cm x 1mm pre-cleaned microslides. The section of skin was stained with diaminobenzidine, which binds to

WO 2010/125340

PCT/GB2010/000844

COX-2 enzymes, and the slides visualised on a light microscope with image capture facility.

5 The results of this experiment are shown as Figure 1. A dark stain is present in the sample both after 0 hours and 6 hours, indicating the continuing presence of COX-2 in the skin. This indicates that water has no anti-inflammatory activity.

Example 2

10 An experiment was carried out as described in Example 1, except that the donor phase comprised ketoprofen, a known COX-2 inhibitor, in fish oil.

The results of this experiment are shown as Figure 2. After six hours, the amount of dark stain has reduced, indicating penetration of ketoprofen into the viable epidermis
15 and activity against COX-2 expression in the skin

Example 3

20 An experiment was carried out as described in Example 1, except that the donor phase comprised a representative compound of the invention, DGLA.

The results of this experiment are shown as Figure 3. After six hours, the amount of dark stain has reduced, indicating penetration of DGLA into the viable epidermis and activity against COX-2 expression in the skin.
25

Example 4

30 An experiment was carried out as described in Example 1, except that the donor phase comprised a representative compound of the invention, 15-HETrE.

The results of this experiment are shown as Figure 4. After six hours, the amount of dark stain has reduced, indicating penetration of 15-HETrE into the viable epidermis and activity against COX-2 expression in the skin.

WO 2010/125340

PCT/GB2010/000844

Example 5

A Western Blotting experiment was carried out to determine the effect of topically applied representative compounds of the invention DGLA and 15-HETrE on the expression of COX-2 in porcine skin. Water was used as a control.

The porcine skin membranes (as described in Example 1) were gently cleaned with de-ionised water before being homogenized (Silverion, Chesham, UK) in a RIPA lysis buffer [50 mM Tris-HCl (pH 7.4), 150 mM NaCl, 1 mM PMSF, 1 mM EDTA, 5 $\mu\text{g mL}^{-1}$ aprotinin, 5 $\mu\text{g mL}^{-1}$ leupeptin, 1% Triton X-100, 1% sodium deoxycholate, 0.1% SDS]. After 15 min incubation on ice, the lysates were clarified by centrifugation at 14000 x g for 2 x 15 min and the supernatant stored at -80 °C for subsequent protein analysis.

Aliquots of 30 μg total protein were separated by SDS PAGE, transferred to nitrocellulose membranes using the Trans-Blot electrophoretic Transfer Cell and briefly stained with Ponceau S to verify effective transfer. Immunoblots were incubated for 1 h in a blocking solution [tris-buffered saline (TBS)-Tween 20 containing 5% (w/v) commercial skimmed milk powder (Marvel) at room temperature. After washing, the membrane was probed overnight at 4 °C with COX-2 antibody at 1:1000, 5-LOX at 1:1000 and iNOS at 1:500 in (1:20 and 1:100 western blocking reagent and sodium azide solution respectively made up to volume with TBS tween). Membranes were then incubated for 1 h with HRP-conjugated anti-rabbit at 1:10000. For β -actin, membranes were probed with anti-actin and anti-mouse for 1h each at room temperature at 1:10000. After 3x10 min washes in TBS Tween, they were finally exposed to freshly prepared Dura Substrate for chemiluminescence for appropriate time before performing autoradiography.


The results of this experiment are shown as Figure 5. It can be seen that a significant reduction in COX-2 expression is seen following dosing with DGLA and 15-HETrE, relative to the control.

Claims:

1. A topical monotherapy composition for the treatment of skin inflammation in a mammal, the topical monotherapy composition comprising:

at least one excipient; and


a polyunsaturated fatty acid (PUFA) derivative of formula (I):



(I)

or a salt thereof, wherein:

-Alk- is:



R₁ is a hydrogen atom;


wherein the topical monotherapy composition contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient; and

wherein the topical monotherapy composition is substantially free of glucocorticoids.

2. The topical composition of claim 1, wherein the excipient comprises a preservative, wherein the preservative is methylparaben, propylparaben, benzyl alcohol, ascorbyl palmitate, ascorbic acid, or any combination thereof.

3. The topical composition of claim 1, wherein the excipient comprises an emollient, an emulsifier, a thickener, a preservative, or any combination thereof.


4. The topical composition of claim 1, wherein the excipient comprises a solubilizing agent, a humectant, or a combination thereof.
5. The topical composition of any one of claims 1 to 4, wherein the topical composition is in the form of a cream.
6. The topical composition of any one of claims 1 to 5, wherein the topical composition is substantially free of lithium salts.
7. The topical composition of any one of claims 1 to 6, wherein the PUFA derivative of Formula (I) is dihomo- γ -linolenic acid (DGLA), and wherein the composition comprises about 0.01 wt% to about 50 wt% DGLA.
8. The topical composition of any one of claims 1 to 7, wherein the mammal is a human.
9. Use of a polyunsaturated fatty acid (PUFA) derivative of formula (I):



(I)

or a salt thereof, in the preparation of a medicament for the treatment of skin inflammation in a mammal as a topical monotherapy, wherein:

-Alk- is:




R_1 is a hydrogen atom;

wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient; and

wherein the medicament is substantially free of glucocorticoids.

10. The use of claim 9, wherein the medicament is in the form of a cream.
11. The use of claim 9 or 10, wherein the medicament is substantially free of lithium salts.
12. The use of claim 9, 10, or 11, wherein the PUFA derivative of Formula (I) is dihomono- γ -linolenic acid (DGLA), and wherein the medicament comprises about 0.01 wt% to about 50 wt% DGLA.
13. The use of any one of claims 9 to 12, wherein the medicament further comprises at least one excipient.
14. Use of a polyunsaturated fatty acid (PUFA) derivative of formula (I):



(I)

or a salt thereof, for the treatment of skin inflammation in a mammal as a topical monotherapy, wherein:

-Alk- is:



R₁ is a hydrogen atom;


wherein the PUFA derivative is in a composition comprising at least one excipient,

wherein the composition contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient, and

wherein the composition is substantially free of glucocorticoids.

15. The use of claim 14, wherein the composition is in the form of a cream.

16. The use of claim 14 or 15, wherein the composition is substantially free of lithium salts.
17. The use of claim 14, 15, or 16, wherein the PUFA derivative of Formula (I) is dihomo- γ -linolenic acid (DGLA), and wherein the composition comprises about 0.01 wt% to about 50 wt% DGLA.
18. The use of any one of claims 13 to 17, wherein the excipient comprises a preservative, wherein the preservative is methylparaben, propylparaben, benzyl alcohol, ascorbyl palmitate, ascorbic acid, or any combination thereof.
19. The use of any one of claims 13 to 17, wherein the excipient comprises an emollient, an emulsifier, a thickener, a preservative, or any combination thereof.
20. The use of any one of claims 13 to 17, wherein the excipient comprises a solubilizing agent, a humectant, or a combination thereof.
21. The use of any one of claims 9 to 20, wherein the mammal is a human.
22. The use of any one of claims 9 to 21, wherein the skin inflammation is caused by atopic eczema, contact dermatitis, psoriasis or uremic pruritis.
23. The use of claim 22, wherein the skin inflammation is caused by exposure of the skin to electromagnetic radiation.
24. Use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),



(I)

or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is




R_1 is a hydrogen atom,

wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient, and

wherein the amount of the PUFA derivative is 0.5 wt% to 25 wt% of the medicament.

25. Use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),



(I)

or a pharmaceutically acceptable salt thereof for topical treatment of skin inflammation in a mammal, wherein:

-Alk- is



R_1 is a hydrogen atom;

wherein the PUFA derivative is in a medicament,


wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient,

and wherein the amount of the PUFA derivative is 0.5 wt% to 25 wt% of the medicament.

26. The use according to claim 24 or 25, wherein the mammal is a human.

27. The use according to claim 24, 25, or 26, wherein the skin inflammation is caused by atopic eczema, contact dermatitis, psoriasis or uremic pruritis.

28. The use according to any one of claims 24 to 27, which is for co-administration with a corticosteroid.
29. The use according to any one of claims 24 to 28, wherein the skin inflammation is caused by exposure of the skin to electromagnetic radiation.
30. The use according to any one of claims 24 to 29, wherein the medicament further comprises a pharmaceutically acceptable diluent or carrier.
31. The use according to claim 30, wherein the medicament is in the form of a gel, ointment, cream or lotion.
32. Use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),



(I)

or a pharmaceutically acceptable salt thereof, in the manufacture of a medicament for topical treatment of skin inflammation in a mammal, wherein:


-Alk- is:



R₁ is hydrogen atom;

wherein the medicament is free of lithium salts and glucocorticoids, and wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient.

33. Use of a polyunsaturated fatty acid (PUFA) derivative of formula (I),



(I)

or a pharmaceutically acceptable salt thereof for topical treatment of skin inflammation in a mammal, wherein:


-Alk- is:



R₁ is hydrogen atom;

wherein the PUFA derivative is in a medicament, and wherein the medicament is free of lithium salts and glucocorticoids, and wherein the medicament contains the polyunsaturated fatty acid (PUFA) derivative of formula (I) as sole active ingredient.


34. A pharmaceutical composition for use in the treatment of skin inflammation in a mammal by topical administration, comprising a polyunsaturated fatty acid (PUFA) derivative, of formula (I),



(I)

or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is



R₁ is a hydrogen atom; and wherein the PUFA derivative is the sole active ingredient.

35. A pharmaceutical composition according to claim 34, wherein the mammal is a human.

36. A pharmaceutical composition according to claim 34 or 35, wherein the skin inflammation is caused by atopic eczema, contact dermatitis, psoriasis, atopic dermatitis, or uremic pruritis.
37. A pharmaceutical composition according to claim 34, 35, or 36, which is for co-administration with a corticosteroid.
38. A pharmaceutical composition according to claim 34 or 35, wherein the skin inflammation is caused by exposure of the skin to electromagnetic radiation.
39. A pharmaceutical composition according to any one of the claims 34 to 38, which is in the form of a gel, ointment, cream or lotion.
40. Use of DGLA or a C1-C2 alkyl ester or pharmaceutically acceptable salt thereof, for treating uremic pruritus in a subject in need thereof, wherein the DGLA is in a composition formulated for administration to provide a therapeutically effective amount to skin of the subject, wherein the amount of the DGLA is 0.5 wt% to 25 wt% of the composition.
41. Use of DGLA or a C1-C2 alkyl ester or pharmaceutically acceptable salt thereof in the preparation of a composition for treating uremic pruritus in a subject in need thereof, wherein the composition formulated for administration to provide a therapeutically effective amount to skin of the subject, wherein the amount of the DGLA is 0.5 wt% to 25 wt% of the composition.
42. The use of claim 40 or 41, wherein the composition comprises about 1 mg to about 1000 mg of DGLA.
43. The use of claim 40, 41, or 42, wherein the composition further comprises a pharmaceutically acceptable diluent or carrier.
44. The use of any one of claims 40 to 43, wherein the composition is formulated for co-administration with a corticosteroid selected from the group consisting of clobetasol dipropionate, betamethasone dipropionate, halbetasol dipropionate, diflorasone diacetate, fluocinonide, halcinonide, amcinonide, desoximetasone, triamcinolone acetonide, mometasone furoate, fluticasone dipropionate, fluocinolone acetonide, hydrocortisone valerate,

hydrocortisone butyrate, desonide, prednicarbate, prednisolone, methylprednisolone, dexamethasone, naflocort, deflazacort, halopredone acetate, budesonide, beclomethasone dipropionate, hydrocortisone, clocortolone pivalate, methylprednisolone aceponate, dexamethasone palmitoate, tipredane, hydrocortisone aceponate, alclometasone dipropionate, halometasone, methylprednisolone suleptanate, rimexolone, prednisolone farnesylate, ciclesonide, deprodone propionate, loteprednol etabonate, betamethasone butyrate propionate, flunisolide, prednisone, dexamethasone sodium phosphate, triamcinolone, betamethasone 17-valerate, betamethasone, hydrocortisone acetate, hydrocortisone sodium succinate, prednisolone sodium phosphate, hydrocortisone probutate, and combinations thereof.


45. The use of claim 44, wherein the corticosteroid is for co-administration in an amount sufficient to provide 50% or less of the recommended daily dosage of said corticosteroid in said mammal.

46. The use of claim 45, wherein the corticosteroid is hydrocortisone.

47. The use of any one of claims 40 to 46, wherein the composition is further formulated for co-administration with a therapeutic agent selected from the group consisting of immunomodulators, antibiotics, immunosuppressants, and anti-itch drugs.

48. A pharmaceutical composition for use in the treatment of skin inflammation in a mammal by topical administration, comprising a polyunsaturated fatty acid (PUFA) derivative,


of formula (I),



(I)

or a pharmaceutically acceptable salt, or solvate thereof, wherein

-Alk- is



R₁ is a hydrogen atom, wherein the PUFA is the sole active ingredient,

wherein the skin inflammation is caused by atopic dermatitis,

and wherein the amount of the PUFA is 1 wt% to 10 wt% of the pharmaceutical composition.

49. A pharmaceutical composition for use according to claim 48, wherein the mammal is a human.
50. A pharmaceutical composition for use according to claim 48 or 49, which is for co-administration with a corticosteroid.
51. A pharmaceutical composition for use according to claim 48, 49, or 50, which is in the form of a gel, ointment, cream or lotion.

t=0



t=6hr



Figure 1

WO 2010/125340

PCT/GB2010/000844

2/5

t=0

t=6hr




Figure 2

WO 2010/125340

PCT/GB2010/000844

3/5

$t=0$

$t=6hr$





Figure 3

WO 2010/125340

4/5

PCT/GB2010/000844

t=0



t=6hr



Figure 4





Figure 5



(I)