

(19) World Intellectual Property Organization
International Bureau(43) International Publication Date
28 February 2002 (28.02.2002)

PCT

(10) International Publication Number
WO 02/16407 A2(51) International Patent Classification⁷: C07K 5/065,
5/087, C07F 9/40, C07C 211/30, A61K 38/05, 38/06,
G01N 33/68, A61P 35/00

(21) International Application Number: PCT/US01/26078

(22) International Filing Date: 22 August 2001 (22.08.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/226,671 22 August 2000 (22.08.2000) US(71) Applicant (for all designated States except US): THE
GOVERNMENT OF THE UNITED STATES, rep-
resented by SECRETARY, DEPARTMENT OF
HEALTH AND HUMAN SERVIC [US/US]; Suite 325,
6011 Executive Boulevard, Rockville, MD 20852-3804
(US).


(72) Inventors; and

(75) Inventors/Applicants (for US only): BURKE, Terrence,

R., Jr. [US/US]; 7400 Lakeview Drive # 410, Bethesda,
MD 20817 (US). WEI, Chang-Qing [CN/US]; 90 Waver-
ley Drive, #F302, Frederick, MD 21702 (US). VOIGT, Jo-
hannes, H. [DE/US]; 525 Quinton Avenue, Kenilworth,
NJ 07033 (US). GAO, Yang [CN/US]; 62 Montoya Cir-
cle, Branford, CT 06405 (US).(74) Agent: PILLAI, Xavier; Leydig, Voit & Mayer, Ltd.,
Suite 300, 700 13th Street, Washington, DC 20005 (US).(81) Designated States (national): AE, AG, AL, AM, AT, AU,
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU,
CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC,
LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW,
MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI,
SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU,
ZA, ZW.(84) Designated States (regional): ARIPO patent (GH, GM,
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,

[Continued on next page]

(54) Title: SH2 DOMAIN BINDING INHIBITORS



(57) **Abstract:** Disclosed are compounds for SH2 domain binding inhibition. For example, disclosed is a compound of formula (I) wherein R1 is a lipophile; R2, in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; R3 is hydrogen, azido, amino, carboxyalkyl, alkoxyalkyl, aminocarbonylalkyl, or alkylcarbonylamino, wherein the alkyl portion of R3 may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R6 is a linker; AA is an amino acid; and n is 1 to 6; or a salt thereof. The compounds of the present invention have the advantage that their conformation is constrained to provide enhanced binding affinity with SH2 domain protein. Also disclosed are a pharmaceutical composition comprising a pharmaceutically or pharmacologically acceptable carrier and a compound of the present invention, a method for inhibiting an SH2 domain from binding with a phosphoprotein comprising contacting an SH2 domain with a compound of the present invention, a method of preventing or treating a disease, state, or condition by the use of one or more of these compounds, and a method for preparing the compounds of the present invention. The present invention further provides intermediates useful in the preparation of the compounds.



CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

Published:

— *without international search report and to be republished upon receipt of that report*

SH2 DOMAIN BINDING INHIBITORS

This application claims the benefit of U.S. provisional patent application No. 60/226,671, filed August 22, 2000.

TECHNICAL FIELD

5 This invention relates to novel compounds, e.g., macrocyclic and other peptides, compositions comprising these compounds, and methods of using these compounds, e.g., in inhibiting SH2 domain binding with a phosphoprotein such as in the prevention or treatment of a disease, state, or condition in a mammal. The present invention also relates to a method for preparing these compounds and intermediates useful in their preparation.

10 BACKGROUND OF THE INVENTION

The pharmaceutical industry is in search for new classes of compounds for the therapy and prophylaxis of proliferative diseases such as cancer, autoimmune diseases, and hyperproliferative skin disorders such as psoriasis. These diseases or disorders affect a large portion of the population, leading to suffering and possibly death.

15 Some of these diseases or disorders may involve signal transduction. Signal transduction is critical to normal cellular homeostasis and is the process of relaying extracellular messages, e.g., chemical messages in the form of growth factors, hormones and neurotransmitters, via receptors, e.g., cell-surface receptors, to the interior of the cell. Protein-tyrosine kinases play a central role in this biological function. Among others, these
20 enzymes catalyze the phosphorylation of specific tyrosine residues to form tyrosine phosphorylated residues. Examples of this class of enzymes include the PDGF receptor, the FGF receptor, the HGF receptor, members of the EGF receptor family such as the EGF receptor, erb-B2, erb-B3 and erb-B4, the src kinase family, Fak kinase and the Jak kinase family. The tyrosine-phosphorylated proteins are involved in a range of metabolic
25 processes, from proliferation and growth to differentiation.

Protein-tyrosine phosphorylation is known to be involved in modulating the activity of some target enzymes as well as in generating specific complex networks involved in signal transduction via various proteins containing a specific amino acid sequence called a Src homology region or SH2 domain (see *Proc. Natl. Acad. Sci. USA*, 90, 5891 (1990)). A
30 malfunction in this protein-tyrosine phosphorylation through tyrosine kinase overexpression or deregulation is manifested by various oncogenic and (hyper-)proliferative disorders such

as cancer, inflammation, autoimmune disease, hyperproliferative skin disorders, such as psoriasis, and allergy/asthma.

SH2- and/or SH3- comprising proteins that play a role in cellular signaling and transformation include, but are not limited to, the following: Src, Lck, Eps, ras GTPase-activating protein (GAP), phospholipase C, phosphoinositol-3 (PI-3)kinase, Fyn, Lyk, Fgr, Fes, ZAP-70, Sem-5, p85, SHPTP1, SHPTP2, corkscrew, Syk, Lyn, Yes, Hck, Dsrc, Tec, Atk/Bpk, Itk/Tsk, Arg, Csk, tensin, Vav, Emt, Grb2, BCR-Abl, Shc, Nck, Crk, CrkL, Syp, Blk, 113TF, 91TF, Tyk2, especially Src, phospholipase c, phosphoinositol-3 (pi-3)kinase, Grb2, BCR-Abl, Shc, Nck, Crk, CrkL, Syp, Blk, 113TF, 91TF, and Tyk2. A direct link has been established between activated receptor kinases and Ras with the finding that the mammalian Grb2 protein, a 26 kilodalton (kD) protein comprising a single SH2 and two SH3 domains bind to proline-rich sequences present in the Sos exchange factor.

The significance of ras-regulatory proteins in human tumors is also highlighted by the critical role of Grb2 in BCR-Abl mediated oncogenesis (*J. Exp. Med.*, 179, 167-175 (1994)).

Central to the binding of SH2 domains with phosphotyrosine ("pTyr") containing ligands is the interaction of the doubly ionized ptyr phosphate with two invariant arginine residues in a well formed pocket. These arginine-phosphate interactions are particularly critical to the overall binding, such that high affinity binding is usually lost by removal of the phosphate group.

U.S. provisional patent application Nos. 60/160,899, filed October 22, 1999; and 60/221,525, filed July 28, 2000; and International Publication Nos. WO 00/73326 A2 and WO 00/56760 disclose certain SH2 domain binding inhibitors, the disclosures of which are incorporated herein in their entireties by reference.


There exists a need for molecules that have an ability to mimic the structure of the phosphotyrosine peptide binding site, as well as a need for compounds that have the ability to disrupt the interaction between SH2 domains of proteins (e.g., regulatory proteins) for example that of Grb2, and proteins with phosphorylated moieties. There also exists a need for suitable starting materials, intermediates, or precursors in the synthesis of the molecules that inhibit binding of SH2 domains. There further exists a need for compounds suitable for use in the therapy or prophylaxis of proliferative diseases or conditions, as well as in

diagnosis, assays, and testing.

These and other advantages of the present invention will be apparent from the description as set forth below.

SUMMARY OF THE INVENTION

5 The present invention provides a number of SH2 domain inhibiting compounds. e.g., peptides, preferably, macrocyclic peptides. Thus, the present invention provides a compound of formula (I)



(I)

10 wherein R_1 is a lipophile; R_2 in combination with the phenyl ring is a phenylphosphate mimic group or a protected phenylphosphate mimic group; R_3 is hydrogen, azido, amino, carboxyalkyl, alkoxy carbonylalkyl, aminocarbonylalkyl, or alkylcarbonylamino, wherein the alkyl portion of R_3 may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R_6 is a
15 linker; AA is an amino acid; and n is 1 to 6; or a salt thereof. The compounds of the present invention, in embodiments, have the advantage that their conformation is constrained to provide enhanced binding affinity with SH2 domain protein.

The present invention further provides a pharmaceutical composition comprising a pharmaceutically or pharmacologically acceptable carrier and a compound of the present
20 invention. The present invention also provides a method for inhibiting an SH2 domain from binding with a phosphoprotein comprising contacting an SH2 domain with a compound of the present invention. The present invention also provides a method of preventing or treating a disease, state, or condition by the use of one or more of these compounds. The present invention also provides a method for preparing the compounds of the present invention. The
25 present invention further provides intermediates useful in the preparation of the compounds.

While the invention has been described and disclosed below in connection with certain embodiments and procedures, it is not intended to limit the invention to those

specific embodiments. Rather it is intended to cover all such alternative embodiments and modifications as fall within the spirit and scope of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 depicts a method of preparing compound 9a in accordance with an
5 embodiment of the present invention. i. pivaloyl chloride, Et₃N, BuLi, -78°C~R.T., 81%; ii. t-butyl phosphite, BuLi, 0°C~R.T., 90%; iii. Et₃N, Pd(OAc)₂, tri-o-tolylphosphine, reflux, 85.5%.

Figure 2 depicts a method of preparing compound 6a in accordance with an
embodiment of the present invention. i. vinyl magnesium bromide, PhSCu, Et₂O-THF,
10 -40°C, 67%, 64% d.e.; ii H₂O₂, 2 eq. LiOH, THF-H₂O, 81%.

Figure 3 depicts a method of preparing compound 7a in accordance with an
embodiment of the present invention. i. Trimethylacetyl chloride, NMP, 70%, ii.
a) LiHMDS, THF; b) 1-bromomethyl-naphthalene, 88%; iii. LiAlH₄, THF, -78°C-R.T.,
100%; iv. DEAD, PPh₃, Phthalimide, THF, 73%; v. EtOH, H₂O, N₂H₄·H₂O, 91%.

15 Figure 4 depicts a method of preparing compound 5a in accordance with an
embodiment of the present invention. i. HOBT, DIPCDI, 95%; ii. a) TFA-CH₂Cl₂;
b) NHCO₃; c) Fmoc-1-amino-cyclohexane carboxylic acid, HOBT, DIPCDI; 70%; d) TFA-
CH₂Cl₂; e) NaHCO₃, 98%; iii. 6a, HOBT, DIPCDI, 67%.

Figure 5 depicts a method of preparing compound 4a in accordance with an
20 embodiment of the present invention. i. Grubbs Catalyst, CH₂Cl₂, reflux, 60 hr, 67%; ii.
TFA-H₂O-Trimethylsilane, 1hr; A: Grubbs Catalyst.

Figure 6 depicts a method of preparing compound 23a in accordance with an
embodiment of the present invention. i. 1) NaHMDS, -70°C, 2) Trisyl azide, 73%; ii. H₂O₂,
2 eq. LiOH, THF-H₂O, 92%.

25 Figure 7 depicts a method of preparing compound 3a in accordance with an
embodiment of the present invention. i. 23a HOBT, DIPCDI, 67%; ii. PPh₃, THF-H₂O, 77%;
iii. Ac₂O, Pyridine.

Figure 8 depicts a method of preparing compounds 34-35 in accordance with an
embodiment of the present invention. (a) (1) pivaloyl chloride, N-methylmorpholine,
30 -78°C; (2) lithium salt of 38 (49%); (b) Pd(OAc)₂, P(*o*-toluyl)₃, NEt₃, 85°C (79%); (c)
H₂/Pd⁰ (82%); (d) LiOH, H₂O₂, THF/H₂O, 0°C (81%); (e) KHMDS, trisyl azide (77%); (f)

LiOH, H₂O₂, THF/H₂O, 0°C (88%).

Figure 9 depicts a method of preparing compounds 44-45 in accordance with an embodiment of the present invention.

Figure 10 depicts a method of preparing compounds 64a-b in accordance with an embodiment of the present invention.

Figure 11 depicts the innate affinity data for inhibitors 1, 2, and 3a to the Grb2 SH2 domain protein measured by the ELISA techniques. The X-axis depicts the concentration of the inhibitor and the Y-axis depicts the decrease in binding relative to the control. IC₅₀ (μM) are 0.02 (■), 2 (▲) and 0.035 (•) (n=3).

Figure 12 depicts the Grb2 SH2 domain binding inhibition in whole cell preparations for compounds 3a and 51-52. Cells were treated with inhibitor at the concentrations indicated prior to stimulation with growth factor. Cells were washed, lysed and immunoprecipitated with anti-Grb2 antibody, then pTyr Western blots against the ErbB-2 protein were run.

Figure 13 depicts the effect of embodiments of the Grb2 SH2 domain inhibitors on the growth of MDA-453/m1 cells. Cells were treated with inhibitor at the concentrations indicated prior to stimulation with growth factor. Cells were washed, lysed and immunoprecipitated with anti-Grb2 antibody, then pTyr Western blots against the ErbB-2 protein were run.

Figures 14a-e depict the formulas of compounds Ii, 3a, 5b, Ih, and 126, respectively.

Figure 15 depicts the effect of compounds Ih and 126 on phosphorylation of MAP Kinase in MDA453 cells (4 hour treatment in 10% FBS).

Figure 16a depicts the dose response curves of compounds 126, Ih, Ii, 3a, and 5b, on the growth of MDA453 breast cancer cells after treatment with growth factor. Figure 16b depicts the dose response bar graphs of compounds 126, Ih, Ii, and 3a on the growth of MDA453 breast cancer cells after treatment with growth factor.

Figure 17 depicts the extracellular ELISA assay of Grb2 SH2 domain binding inhibitors. IC₅₀ of compounds 126, Ii, 3a, 5b, and Ih are respectively, 0.01, 0.25, 4.00, 2.00, and 0.004 μM.

Figures 18a-c depict the effect of inhibitors on the binding of Grb2 to erbB-2 tyrosine kinase in whole cell assays (IP: Grb2 c-23 WB: pTy (PY99)). Figure 18a depicts


the effect of compounds 1h and 126 on erbB-2 and Grb2 association in MDA453 cells; Figure 18b depicts the effect of compounds 3a and 5b on erbB-2 and Grb2 association in MDA453 cells; and Figure 18c depicts the effect of compound 1i on erbB-2 and Grb2 association in MDA453 cells.

5

SPECIFIC DESCRIPTION OF THE INVENTION

An aspect of the present invention is predicated on the concept that binding affinity for SH2 domain proteins can be envisioned to increase by a conformational constraint in a ligand. The conformational constraint is believed to lead to certain advantages, e.g., a reduction in binding entropy penalty. Binding of natural pTyr-containing ligands to Grb2 SH2 domains takes place in a β -bend fashion, with key interactions occurring in a pTyr binding pocket as well as in a proximal pocket which ligates the amino acid side chain of a pY+2 Asn residue. The present invention provides a novel platform which is expected provide enhanced binding outside the pTyr pocket.

In accordance with an embodiment, the present invention provides certain compounds, e.g., macrocyclic peptides. The present invention provides a compound of formula (I)



(I)

wherein R_1 is a lipophile; R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; R_3 is hydrogen, azido, amino, carboxyalkyl, alkoxyalkyl, aminocarbonylalkyl, or alkylcarbonylamino, wherein the alkyl portion of R_3 may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R_6 is a linker; AA is an amino acid; and n is 1 to 6; or a salt thereof.

R_1 can be any suitable lipophile, e.g., a lipophilic group that is capable of providing or enhancing cell membrane penetration. Examples of lipophiles include aralkyl, arylheterocyclalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminoacarbonyl,

alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl groups, wherein the aryl portion is optionally substituted a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, amido, aminoalkyl, alkyl, alkoxy, and keto. In embodiments, the aryl portion is phenyl or naphthyl and the alkyl portion is a C₁-C₆ alkyl, and the heterocyclyl is a 3-7

5 membered ring having one or more of N, O, or S. In a preferred embodiment, the heterocyclyl is a 5-membered ring, and in a further preferred embodiment, the heterocyclyl is a 5-membered ring containing N. In a preferred embodiment, the alkyl portion of the arylheterocyclylalkyl is a C₂-C₃ alkyl. Examples of suitable arylheterocyclylalkyls include indolylmethyl and naphthylmethyl, e.g., naphthyl-2-methyl.

10 R₂ can be any suitable phenylphosphate mimic group or a protected phenylphosphate mimic group. Examples of R₂ include hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkyloxy, dicarboxyhaloalkyl, dicarboxyhaloalkyloxy, phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl portion may be optionally
15 substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, amido, aminoalkyl, alkyl, alkoxy, and keto. Examples of the phenylphosphate mimic group include phosphonomethyl, phosphono-(α -fluoro)methyl, phosphono-(α,α -difluoro)methyl, phosphono-(α -hydroxy)methyl, O-sulfo, and dicarboxymethoxy. R₂ can be located at the *o*, *p*, or *meta* position, and preferably at the *para* position on the phenyl ring.
20 The phenylphosphate mimic group can be present as protected groups, e.g., protected as an ester, amide, or ether, of an alcohol such as C₁-C₆ alcohol, preferably t-butyl alcohol.

In accordance with embodiments, R₃ is hydrogen, azido, amino, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, oxalylamino, or alkylcarbonylamino; wherein the alkyl portion of R₃ is C₁-C₆ alkyl which may be optionally substituted with a substituent
25 selected from the group consisting of halo, hydroxy, carboxyl, amino, amido, aminoalkyl, alkyl, alkoxy, and keto. In a preferred embodiment, R₃ is carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, or alkylcarbonylamino, and wherein the alkyl is methyl.

Any suitable, natural or synthetic or modified amino acid can be employed.


30 Examples of amino acids include glycine, alanine, valine, norvaline, leucine, iso-leucine, norleucine, α -amino n-decanoic acid, serine, homoserine, threonine, methionine, cysteine, S-

acetylaminomethyl-cysteine, proline, trans-3- and trans-4-hydroxyproline, phenylalanine, tyrosine, 4-aminophenylalanine, 4- nitrophenylalanine, 4-chlorophenylalanine, 4-carboxyphenylalanine, β -phenylserine β -hydroxyphenylalanine, phenylglycine, α -naphthylalanine, cyclohexylalanine, cyclohexylglycine, tryptophan, indoline-2-carboxylic acid, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, aspartic acid, asparagine, aminomalonic acid, aminomalonic acid monoamide, glutamic acid, glutamine, histidine, arginine, lysine, N'-benzyl-N'-methyl-lysine, N',N'-dibenzyl-lysine, 6-hydroxylysine, ornithine, α -aminocyclopentane carboxylic acid, α -aminocyclohexane carboxylic acid, α -aminocycloheptane carboxylic acid, α -(2-amino-2-norbornane)-carboxylic acid, α,γ -diaminobutyric acid and α,β -diaminopropionic acid, homophenylalanine, and α -tert-butylglycine. Preferred examples include, α -aminocyclohexane carboxylic acid and asparagine.

The linker can be any suitable moiety that conformationally constrains the compound of the present invention, e.g., in the beta bend fashion. The linker is not limited by any chemical structure. The linker can be linear or branched, substituted or unsubstituted, group, e.g., a group having 1-6 carbon atoms, optionally with one or more N, O, or S. Examples of suitable linkers include alkylenyl, alkenylenyl, and alkynylenyl, preferably ethylenyl, ethenylenyl, or ethynylenyl. The points of attachment of the linker can have a suitable configuration, e.g., in R, S, or a mixture of R and S forms.

In preferred embodiments of the compounds of the present invention, n is 2-4, and in further preferred embodiments, n is 2 or 3.

In a preferred embodiment, the present invention provides a compound of formula (Ia)




(Ia)

wherein R₄ and R₅, independently, are hydrogen, alkyl, cycloalkyl, or heterocyclyl, or R₄ and

R₅ together form a cycloalkyl or heterocyclyl.

In another preferred embodiment, the present invention provides a compound of the formula (Ib):



(Ib)

In accordance with embodiments of the present invention, in the compounds described above, R₁ is aralkyl, arylheterocyclylalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminoacarbonyl, alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl, wherein the aryl portion is phenyl or naphthyl and the alkyl portion is a C₁-C₆ alkyl, and the heterocyclyl is a 3-7 membered ring having at least one of N, O, or S;

R₂ is hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkyloxy, dicarboxyhaloalkyl, dicarboxyhaloalkyloxy, phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl or alkoxy portion of R₂ is a C₁-C₆ alkyl or alkoxy and may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto;

R₃ is hydrogen, azido, amino, carboxyalkyl, alkoxycarbonylalkyl, aminocarbonylalkyl, or alkylcarbonylamino; wherein the alkyl portion of R₃ is C₁-C₆ alkyl which may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto;

R₄ and R₅, independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl, wherein the alkyl is a C₁-C₆ alkyl, the cycloalkyl is a C₃-C₇ cycloalkyl, and the heterocyclyl is a 3-7 membered ring with one or more of N, O, or S; and

R₆ is a C₂-C₄ alkylenyl or alkenylenyl group, which may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto.

In particular embodiments, R₁ is naphthylmethyl or indolyl and R₂ is carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkoxy, dicarboxyhaloalkyl, dicarboxyhaloalkoxy, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl or alkoxy portion of R₂ is a C₁-C₆ alkyl or alkoxy and may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto.

In certain embodiments, R₁ is naphthylmethyl or indolyl and R₂ is carboxyalkoxy, dicarboxyalkyl, dicarboxyalkoxy, dicarboxyhaloalkyl, dicarboxyhaloalkoxy, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl or alkoxy portion of R₂ is a C₁-C₆ alkyl or alkoxy and may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto. In certain preferred embodiments, R₂ is phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphonodihaloalkyl, or phosphoryl, e.g., phosphonomethyl, phosphonohalomethyl, or phosphonodihalomethyl.


In embodiments, the present invention provides compounds described above wherein R₃ is carboxy C₁-C₆ alkyl, e.g., carboxymethyl or dicarboxy C₁-C₆ alkyl, e.g., dicarboxymethyl.

In certain embodiments described above, R₃ is alkoxy carbonyl C₁-C₆ alkyl, aminocarbonyl C₁-C₆ alkyl, oxalylamino, or C₁-C₆ alkylcarbonylamino; wherein the alkyl portion of R₃ may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto. In certain preferred embodiments, R₃ is C₁-C₆ alkylcarbonylamino, e.g., acetylamino.

In certain embodiments described above, R₄ and R₅, independently, are hydrogen, alkyl, or together form cycloalkyl, wherein the alkyl is a C₁-C₆ alkyl, and the cycloalkyl is a C₃-C₇ cycloalkyl, e.g., a C₆ cycloalkyl.


In accordance with embodiments of the present invention, the present invention provides a compound of formula (Ic):

11




(Ic)

wherein R_4 and R_5 are independently C_1 - C_6 alkyl or hydrogen; a compound of formula (Id)



(Id)


wherein R_4 and R_5 are independently C_1 - C_6 alkyl or hydrogen, a compound of the formula (Ie)



(Ie)


wherein R_4 and R_5 together form cyclohexyl; a compound of the formula (If):

12



(If)

wherein R_7 is aryl or alkenyl, and R_4 and R_5 are independently C_1 - C_6 alkyl or hydrogen; and a compound of the formula (Ig)




(Ig)

wherein R_8 is aryl alkyl and R_4 and R_5 are independently C_1 - C_6 alkyl or hydrogen. In a preferred embodiment of the compound of formula (Ig), R_8 is benzyl or naphthylmethyl.

In a preferred embodiment, the present invention provides a compound of formula


10 (Ih):



(Ih)

wherein R₄ and R₅ are methyl.

In another embodiment, the present invention provides a compound of formula (Ii):



(Ii)

5 wherein R₄ and R₅ are methyl.

In another aspect, the present invention provides a compound of the formula:



wherein n is 0 to 15; Y' is a phenylalanyl radical having a phenyl ring, an amine end, and a carboxyl end, the phenyl ring having one or more substituents selected from the group
 10 consisting of hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkoxy, dicarboxyhaloalkyl, dicarboxyhaloalkoxy, and phosphonoalkyl, phosphonohaloalkyl, wherein the alkyl portion of the substituents may be unsubstituted or substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto and the amine end includes an azido group; W is
 15 a moiety attached to the nitrogen of Y' and is selected from the group consisting of alkylcarbonyl, oxalyl, alkylaminooxalyl, arylaminooxalyl, arylalkylaminooxalyl, alkoxyoxalyl, carboxyalkyl carbonyl, heterocyclyl carbonyl, heterocyclylalkyl carbonyl, arylalkyl heterocyclylalkyl carbonyl, aryloxy carbonyl, and arylalkoxy carbonyl, wherein the aryl and alkyl portions of the substituents may be unsubstituted or substituted with a
 20 substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; and the heterocyclyl portion of W contains at least 4 hetero atoms selected from the group consisting of O, N, and S; AA is an amino acid, the amine end of which is attached to the carboxyl end of Y'; and Z is an arylalkylamino or arylheterocyclyl alkylamino; or a salt thereof.

25 The alkyl portion of the various groups described for the compound of formula (II) can have any suitable number of carbon atoms, e.g., from 1 to about 12 carbon atoms,

preferably from 1 to 6 carbon atoms, and more preferably from 1 to 4 carbon atoms. The aryl portion of the various groups described can have any number of aromatic rings, e.g., from 1 to 3 rings, e.g., six membered rings, preferably 1 or 2 rings, and more preferably 1 ring. Thus, for example, the present invention provides a compound wherein Y' is a

5 phenylalanyl radical having a phenyl ring, an amine end, and a carboxyl end, the phenyl ring having one or more substituents selected from the group consisting of hydroxyl, carboxyl, formyl, carboxy C₁-C₆ alkyl, carboxy C₁-C₆ alkyloxy, dicarboxy C₁-C₆ alkyl, dicarboxy C₁-C₆ alkyloxy, dicarboxyhalo C₁-C₆ alkyl, dicarboxyhalo C₁-C₆ alkyloxy, and phosphono C₁-C₆ alkyl, phosphonohalo C₁-C₆ alkyl, wherein the alkyl portion of the substituents may be

10 unsubstituted or substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, amino C₁-C₆ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, and keto;

W is a moiety attached to the nitrogen of Y' and is selected from the group consisting of C₁-C₆ alkylcarbonyl, oxalyl, C₁-C₆ alkylaminooxalyl, arylaminooxalyl, aryl C₁-C₆ alkylaminooxalyl, C₁-C₆ alkoxyoxalyl, carboxy C₁-C₆ alkyl carbonyl, heterocyclyl

15 carbonyl, heterocyclyl C₁-C₆ alkyl carbonyl, aryl C₁-C₆ alkyl heterocyclyl C₁-C₆ alkyl carbonyl, aryloxy carbonyl, and aryl C₁-C₆ alkoxy carbonyl, wherein the aryl and alkyl portions of the substituents may be unsubstituted or substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, amino C₁-C₆ alkyl, C₁-C₆ alkyl, C₁-C₆ alkoxy, and keto; and the heterocyclyl portion of W contains at least 4 hetero

20 atoms selected from the group consisting of O, N, and S; AA is an amino acid, the amine end of which is attached to the carboxyl end of Y'; and Z is an aryl C₁-C₆ alkylamino or arylheterocyclyl C₁-C₆ alkylamino; or a salt thereof. The compounds can be in D, L, or a mixed form thereof.

The present invention further provides a composition comprising a pharmaceutically

25 acceptable carrier and an effective (e.g., therapeutically or prophylactically effective) amount of at least one of the compounds set forth above. The present invention further provides a method of inhibiting an SH2 domain from binding with a phosphoprotein comprising contacting a sample or substance containing an SH2 domain with a compound of the present invention.

30 The present invention discloses the use of above compounds in the manufacture of a medicament for the treatment of a condition that responds to the inhibition of

phosphoprotein binding to an SH2 domain of a mammal. The present invention further teaches the use of the above compounds in medicine. The compounds find use as a Grb2-SH2 domain inhibitor.

The pharmaceutically acceptable (e.g., pharmacologically acceptable) carriers
5 described herein, for example, vehicles, adjuvants, excipients, or diluents, are well-known to those who are skilled in the art and are readily available to the public. It is preferred that the pharmaceutically acceptable carrier be one which is chemically inert to the active compounds and one which has no detrimental side effects or toxicity under the conditions of use.

The choice of carrier will be determined in part by the particular active agent, as well
10 as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of the pharmaceutical composition of the present invention. The following formulations for oral, aerosol, parenteral, subcutaneous, intravenous, intraarterial, intramuscular, interperitoneal, intrathecal, rectal, and vaginal administration are merely exemplary and are in no way limiting.

15 Formulations suitable for oral administration can comprise (a) liquid solutions, such as an effective amount of the compound dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations can
20 include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant, suspending agent, or emulsifying agent. Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can
25 include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible
30 carriers. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base,

such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such carriers as are known in the art.

The compounds of the present invention, alone or in combination with other suitable components, can be made into aerosol formulations to be administered via inhalation. These
5 aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like. They also can be formulated as pharmaceuticals for non-pressured preparations, such as in a nebulizer or an atomizer.

Formulations suitable for parenteral administration include aqueous and non-aqueous, isotonic sterile injection solutions, which can contain anti-oxidants, buffers, bacteriostats, and
10 solutes that render the formulation isotonic with the blood of the intended recipient, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. The compound can be administered in a physiologically acceptable diluent in a pharmaceutical carrier, such as a sterile liquid or mixture of liquids, including water, saline, aqueous dextrose and related sugar solutions, an
15 alcohol, such as ethanol, isopropanol, or hexadecyl alcohol, glycols, such as propylene glycol or polyethylene glycol, glycerol ketals, such as 2,2-dimethyl-1,3-dioxolane-4-methanol, ethers, such as poly(ethyleneglycol) 400, an oil, a fatty acid, a fatty acid ester or glyceride, or an acetylated fatty acid glyceride with or without the addition of a pharmaceutically acceptable surfactant, such as a soap or a detergent, suspending agent, such as pectin, carbomers,
20 methylcellulose, hydroxypropylmethylcellulose, or carboxymethylcellulose, or emulsifying agents and other pharmaceutical adjuvants.

Oils, which can be used in parenteral formulations include petroleum, animal, vegetable, or synthetic oils. Specific examples of oils include peanut, soybean, sesame, cottonseed, corn, olive, petrolatum, and mineral. Suitable fatty acids for use in parenteral
25 formulations include oleic acid, stearic acid, and isostearic acid. Ethyl oleate and isopropyl myristate are examples of suitable fatty acid esters. Suitable soaps for use in parenteral formulations include fatty alkali metal, ammonium, and triethanolamine salts, and suitable detergents include (a) cationic detergents such as, for example, dimethyl dialkyl ammonium halides, and alkyl pyridinium halides, (b) anionic detergents such as, for example, alkyl, aryl,
30 and olefin sulfonates, alkyl, olefin, ether, and monoglyceride sulfates, and sulfosuccinates, (c) nonionic detergents such as, for example, fatty amine oxides, fatty acid alkanolamides, and

polyoxyethylenepolypropylene copolymers, (d) amphoteric detergents such as, for example, alkyl- β -aminopropionates, and 2-alkyl-imidazoline quaternary ammonium salts, and (e) mixtures thereof.

The parenteral formulations will typically contain from about 0.5 to about 25% by weight of the active ingredient in solution. Suitable preservatives and buffers can be used in such formulations. In order to minimize or eliminate irritation at the site of injection, such compositions may contain one or more nonionic surfactants. The quantity of surfactant in such formulations typically ranges from about 5 to about 15% by weight. Suitable surfactants include polyethylene sorbitan fatty acid esters, such as sorbitan monooleate and the high molecular weight adducts of ethylene oxide with a hydrophobic base, formed by the condensation of propylene oxide with propylene glycol. The parenteral formulations can be presented in unit-dose or multi-dose sealed containers, such as ampoules and vials, and can be stored in a freeze-dried (lyophilized) condition requiring only the addition of the sterile liquid carrier, for example, water, for injections, immediately prior to use. Extemporaneous injection solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described.

The compounds of the present invention may be made into injectable formulations. The requirements for effective pharmaceutical carriers for injectable compositions are well known to those of ordinary skill in the art. See *Pharmaceutics and Pharmacy Practice*, J.B. Lippincott Co., Philadelphia, PA, Banker and Chalmers, eds., pages 238-250 (1982), and *ASHP Handbook on Injectable Drugs*, Toissel, 4th ed., pages 622-630 (1986).

Additionally, the compounds of the present invention may be made into suppositories by mixing with a variety of bases, such as emulsifying bases or water-soluble bases. Formulations suitable for vaginal administration may be presented as pessaries, tampons, creams, gels, pastes, foams, or spray formulas containing, in addition to the active ingredient, such carriers as are known in the art to be appropriate. Suitable doses and dosage regimens can be determined by conventional range-finding techniques known to those of ordinary skill in the art. Generally, treatment is initiated with smaller dosages, which are less than the optimum dose of the compound. Thereafter, the dosage is increased by small increments until the optimum effect under the circumstances is reached. For convenience, the total daily dosage may be divided and administered in portions during the day if desired. In proper doses

and with suitable administration of certain compounds, the present invention provides for a wide range of responses. Typically the dosages range from about 0.001 to about 1000 mg/kg body weight of the animal being treated/day. Preferred dosages range from about 0.01 to about 10 mg/kg body weight/day, and further preferred dosages range from about 0.01 to about 1
5 mg/kg body weight/day.

Embodiments of the compounds have the advantage that they are stable to or in presence of enzymes encountered during *in vivo* use. Embodiments of the compounds can find use in *in vitro* and *in vivo* applications. For example, the compounds can find use as molecular probes as well as in assays to identify, isolate, and/or quantitate receptor or binding
10 sites in a cell or tissue. The compounds also can find use *in vivo* for studying the efficacy in the treatment of various diseases or conditions involving SH2 domains.

The present invention further provides a method of preventing or treating a disease, state, or condition in a mammal by the use of the compounds of the present invention. In an embodiment, the method involves preventing a disease, state, or condition. In another
15 embodiment, the method involves treating an existing disease, state, or condition.

Preferably, the method involves inhibition of SH2 domain binding with a phosphoprotein. The SH2 domain may involve one or more of the following proteins: Src, Lck, Eps, ras GTPase-activating protein (GAP), phospholipase C, phosphoinositol-3 (PI-3)kinase, Fyn, Lck, Fgr, Fes, ZAP-70, Sem-5, p85, SHPTP1, SHPTP2, corkscrew, Syk, Lyn,
20 Yes, Hck, Dsrc, Tec, Atk/Bpk, Itk/Tsk, Arg, Csk, tensin, Vav, Emt, Grb2, BCR-Abl, Shc, Nck, Crk, CrkL, Syp, Blk, 113TF, 91TF, Tyk2, especially Src, phospholipase c, phosphoinositol-3 (p1-3)kinase, Grb2, BCR-Abl, Shc, Nck, Crk, CrkL, Syp, Blk, 113TF, 91TF, and Tyk2.

The method comprises administering to the mammal one or more compounds of the
25 present invention. The disease, state, condition can be a cancer, e.g., a breast cancer or an ovarian cancer, or a tumor such as a solid tumor, e.g., a brain tumor, a prostate tumor, and the like, leukemia including chronic myelocytic leukemia, lymphoma, an autoimmune disease, an inflammatory disease, a metabolic disease, diabetes, obesity, or cardiovascular disease.

30 The present invention further provides a method of enhancing the therapeutic effect of a treatment rendered to a mammal comprising administering a compound in conjunction

with the treatment. By conjunction, it is meant that the inhibitor can be used in any suitable manner, for example, prior to, simultaneous with, or post- administration of the therapeutic agent. Synergistic effects are observed when the SH2 domain binding inhibitor is used in combination with other treatments known to those skilled in the art. The inhibitor enhances
5 the cytotoxicity of the chemotherapeutic treatments. Cancer treatment is particularly suitable for this combination treatment.

The cancer may involve any number of mechanisms. A majority of human breast cancer are dependent upon activation of the *Ras* signaling pathways through activation of growth factor receptor as the means to achieve continuous cellular proliferation. For
10 example, the cancer may involve overexpression of Her-2/neu. The cancer can be mediated through BCR-Abl or the expression of erbB-2 receptor. In cells transformed by p185 *erbB*-2 overexpression, therapeutic agents affecting Grb2 function at its SH2 domain may interrupt the flow of signal transduction to the *ras* pathway and thus result in reversal of the cancer phenotype.

15 The therapeutic treatment can include a chemotherapy, a radiation therapy, and/or a biological therapy. Examples of chemotherapy includes the use of cancer treatment agents such as alkylating agents, hormonal agents, antimetabolites, natural products, and miscellaneous agents. Particular examples of cancer treatment agents include paclitaxel, 5-fluoruracil, and doxorubicin. Examples of biological therapy includes the use of a protein
20 such as an antibody (monoclonal or polyclonal) or a recombinant protein. An example of an antibody is herceptin, which is targeted for inhibiting the erbB-2 receptor. In embodiments, the enhancement of the therapeutic effect comprises blocking of a cell survival factor in the mammal and/or triggering, e.g., enhancing or speeding up, of cell apoptosis. The treatment can be carried out *in vivo* and/or *in vitro*.

25 The present invention further provides a method of inhibiting the MAP kinase activity in a mammal. MAP kinases function in a protein kinase cascade that plays a critical role in the regulation of cell growth and differentiation. MAP kinases are activated by a variety of signals including growth factors, cytokines and hormones through Grb2 and other signaling proteins. For example, the state of threonine and tyrosine phosphorylation of
30 cellular MAP kinase is determined in MDA-453 cells treated with growth factor heregulin (HRG) using a polyclonal antibody specifically recognizing the phosphorylated threonine

and tyrosine residues of MAP kinase.

The Grb2 SH2 binding inhibitors are effective in inhibiting the association or binding of Grb2 with activated receptor PTKs. Interaction of native Grb2 protein with phosphotyrosinylated proteins including receptor PTKs can be monitored by

- 5 immunoprecipitating Grb2 and detecting the amount of phosphotyrosinylated proteins which are coprecipitated using anti-phosphotyrosine Western Blotting

The compounds of the present invention exert a cytostatic effect. In embodiments, the compounds of the present invention are free or substantially free of toxicity.

- The present invention provides a method for inhibiting cell motility. The present
10 invention also provides a method for inhibiting angiogenesis in an animal. The present invention further provides a method for preventing or treating a variety of diseases, disorders, states or conditions in a mammal, particularly in a human. The present invention provides a method of inhibiting cell motility in a mammal comprising administering to the mammal a peptide having cell signal inhibiting activity and cell motility inhibiting activity.
15 Advantageously, the peptide is free or substantially free of cytotoxicity.

- The present invention contemplates to retard or reduce the movement of cells. A number of factors, forces, and/or mechanisms are involved in the movement of cells from one location to another. The method of the present invention is not limited to inhibiting or interfering with one particular factor, force, or mechanism that is involved in the cell
20 movement.

- The process of cell movement begins with extension of the cell membrane, the push forward of cytosol (the inner material of the cell), and retraction of the rear of the cell. As the cell membrane initially is propelled forward, an attachment forms between the membrane and the substratum, thereby anchoring the "head" of the cell. Some believe that
25 the cytosol is pushed forward by restructuring of the cytoskeletal network within the cell, although the exact mechanism is unknown. The final step involves the detachment of the "tail" of the cell from the substratum.

- It is believed that growth factors activate a signal transduction pathway involving G-proteins, which promote cytoskeletal changes including actin polymerization. External
30 factors promote cell motility by binding to a cell surface receptor and activating a signal transduction pathway, e.g., one involving G-proteins. The signal transduction pathway, in

turn, promotes reorganization of the cytoskeleton. A variety of extracellular factors influence cell motility. The movement of a cell following soluble molecules along a concentration gradient is called chemotaxis. Intracellular calcium may play a role in the ability of a cell to recognize concentration gradients. Hormones such as insulin, cytokines,
5 and specific peptide fragments of the extracellular matrix have been identified which stimulate tumor cell motility and chemotaxis.

Aside from instigating cell motility, growth factors stimulate neovascularization, which involves, in part, cell movement. Angiogenesis begins with proteolytic enzyme-mediated breakdown of the basement membrane of a blood vessel. It is believed that
10 breakdown of the basement membrane is regulated by angiogenic factors, such as fibroblast growth factor. Endothelial cells migrate to the area of degradation and invade the surrounding extracellular matrix. Invading endothelial cells proliferate, forming an elongated column of cells. A lumen forms within the solid cell column, thereby forming a vessel, which eventually connects with an existing blood vessel forming a capillary loop
15 (Fotsis et al., *J. Nutr.*, 125: 790S-797S (1995)).

The present invention provides a method for inhibiting angiogenesis in an animal, e.g., a mammal. The method comprises administering to the animal, e.g., mammal, a peptide having cell signal inhibiting activity and cell motility inhibiting activity, wherein the peptide is substantially free of cytotoxicity. Preferably, the peptide affects multiple aspects
20 of the angiogenic process to effectively therapeutically or prophylactically treat angiogenesis. For example, in addition to inhibiting cell signaling and cell motility, the peptide preferably inhibits invasion of epithelial and/or endothelial cells into the extracellular matrix.

In one embodiment, the present invention provides a method of inhibiting cell
25 motility and angiogenesis induced by the hepatocyte growth factor (HGF), particularly the motility derived from a biological response mediated by its cell surface receptor, the c-Met proto-oncogene product, a transmembrane tyrosine kinase. Upon HGF binding, several tyrosine residues within the c-Met intracellular domain are phosphorylated. Some of the phosphorylated domains mediate binding with various signaling proteins, e.g., the Grb2
30 protein, the p85 subunit of phosphoinositide 3-kinase (PI3K), phospholipase C-gamma, Shc, and Gab1.

The compounds of the present invention inhibit Grb2 SH2 domain binding. In this regard, it is imperative to cellular function that a transducer protein accurately identify activated cellular receptors. Very often, recognition specificity stems from the ability of the transducer protein to recognize a phosphotyrosine surrounded by a specific amino acid
5 sequence. The recognition motif for Grb2 is pYXN wherein pY is phospho-Tyr, X is any amino acid, and N is Asn. The method of the present invention, in an embodiment, is directed to inhibiting cell motility induced or mediated by signaling due to one or more of the above HGF bindings, preferably the binding of HGF c-Met receptor with the Grb2 protein.

10 Compounds having cell signaling inhibitory activity and cell motility inhibiting activity, such as Grb2-SH2 domain mimetic compounds, are particularly useful in inhibiting neovascularization. Compounds having cell signaling inhibitory activity and cell motility inhibiting activity, such as the Grb2-SH2 domain mimetic compounds, e.g., macrocyclic peptides described herein, inhibit endothelial cell and epithelial cell invasion of matrices and
15 the formation of cell cords.

The compounds of the present invention interact with intracellular signal transducers, thus interfering in the pathways leading to cell proliferation and movement. These biological effects can be utilized to inhibit growth of neoplastic cells, inhibit angiogenesis, and to prevent metastatic spreading. The present invention provides a method for preventing or
20 treating a disease, condition, or state in a mammal that is mediated by the binding of an intracellular signal transducer to a receptor protein tyrosine kinase comprising administering to the mammal a peptide of the present invention.

The compounds of the present invention inhibit cell motility. The peptides prevent scattering of cells.

25 The cytotoxic effects of agents that disrupt the cytoskeleton, such as colchicine, taxol, cytochalasins, and phalloidin are well-characterized, and are fundamentally different from the anti-motility effects exerted by the compounds employed in the present invention. These compounds may be highly efficacious for the safe treatment of human diseases such as metastatic cancers, e.g., where the role of HGF plays a role in stimulating the invasion of
30 cells into tissue surrounding the tumors and the migration of metastases to distant sites.

The compounds of the present invention are of use in medicine, e.g., in the


manufacture of a medicament for the treatment of a condition that responds to the inhibition of phosphoprotein binding to an SH2 domain of a mammal.

The present invention further provides a method for determining the presence of an SH2 domain in a material comprising (a) exposing a sample of said material to a SH2
5 binding compound and obtaining a first binding result; (b) exposing another sample of said material to a compound described above and obtaining a second binding result; and (c) comparing the first and second binding results to determine whether an SH2 domain is present in the material.

The present invention further provides a method for inhibiting the binding of an
10 intracellular transducer to a receptor protein tyrosine kinase comprising contacting (a) a sample containing the receptor protein tyrosine kinase, (b) the intracellular transducer, and (c) a compound of the present invention under conditions wherein, in the absence of the compound, the receptor protein tyrosine kinase binds to the intracellular transducer; wherein the contacting results in the inhibition of binding of the intracellular transducer to the
15 receptor protein tyrosine kinase.


The present invention further provides for detecting the inhibition of binding of an intracellular transducer to a receptor protein tyrosine kinase comprising: (a) contacting a sample containing the receptor protein tyrosine kinase with the intracellular transducer, separately, in the presence and absence of the compound of the present invention under
20 conditions that allow for binding of the receptor protein tyrosine kinase to the intracellular transducer in the absence of the compound; (b) determining that binding has occurred between the receptor protein tyrosine kinase and the intracellular transducer; and (c) comparing the relative binding levels of the receptor protein tyrosine kinase to the intracellular transducer in the presence and absence of the compound.

25 The present invention provides (a) a compound of the formula




(b) a compound of the formula


24



(c) a compound of the formula:




(d) a compound of the formula:



5

and (e) a compound of formula 24:




24;

wherein R₁ is a lipophile, R₂, in combination with the phenyl ring, is a phenylphosphate
 10 mimic group or a protected phenylphosphate mimic group; and R₄ and R₅, independently,
 are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.

The present invention further provides a compound of formula 25:


25



25

wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate
mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently,
5 are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.


The present invention further provides a compound of formula 20



20


wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate
10 mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently,
are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.

The present invention further provides (a) a compound of the formulas




(b) a compound of the formula

26



and (c) a compound of the formula 31




31;

- 5 wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group, and R is aralkyl, aryl, or alkyl. In embodiments, R is benzyl.

In preferred embodiments of the intermediate compounds, the phenylphosphate mimic is phosphonomethyl or an ester thereof. In certain embodiments, the lipophile is
 10 aralkyl, arylheterocyclalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminoacarbonyl, alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl, wherein the aryl portion is phenyl or naphthyl and the alkyl portion is a C_1 - C_6 alkyl, and the heterocycl is a 3-7 membered ring having one or more of N, O, or S.


The present invention further provides a method for preparing a compound of
 15 formula 31 wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group, and R is aralkyl, aryl, or alkyl comprising: (a) treating a compound of the formula 26



27

26


with a compound of the formula 27



27


5

to obtain a compound of the formula 28




28

10 (b) treating the compound of formula 28 with a compound of formula 11



11


to obtain a compound of formula 29



29


15

(c) reducing the compound of formula 29 to obtain a compound of formula 30



(d) treating the compound of formula 30 with an alkali metal salt of bis(trimethylsilyl)amide and trisyl azide, to obtain the compound of formula 31. In embodiments, R is benzyl.

5 The present invention further provides a method for preparing a compound of formula 3; wherein R₁ is a lipophile; R₂, in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R₄ and R₅, independently, are hydrogen, alkyl, cycloalkyl, or heterocyclyl, or R₄ and R₅ together form a cycloalkyl or heterocyclyl, the method comprising:




10

(a) providing a compound of formula 24; (b) reducing the azido group in the compound of formula 24 to an amino group to obtain a compound of formula 25; (c) acetylating the amino group of the compound of formula 25 to obtain a compound of formula 20; and (d) carrying
15 out an olefin metathesis reaction on the compound of formula 20, thereby obtaining the compound of formula 3.

The azido group can be reduced by methods known to those skilled in the art, for example, by treating with triphenyl phosphine in a suitable medium, e.g., a mixture of THF and water. The amino group is acetylated to obtain the compound of formula 20.

20 Acetylation can be carried out by methods known to those skilled in the art, e.g., by the use of acetic anhydride and a base such as pyridine. An olefin metathesis reaction can be carried


out by the use of a suitable catalyst, e.g., catalysts that are functionally tolerant such as those developed by Schrock or Grubbs, preferably a ruthenium based catalyst such as the Grubbs catalyst A below:




A

The above catalyst A is not significantly affected by air or moisture and compounds containing nitrogen functional groups or bulky groups can be ring closed by the metathesis reaction. The metathesis reaction can be carried out in a suitable solvent, e.g., halogenated solvents such as dichloromethane. The reaction can be carried out under mild or moderate
 10 conditions, e.g., at the reflux temperature of dichloromethane.

The present invention further provides a method for preparing a compound of formula 24, wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or
 15 heterocyclyl, the method comprising treating a compound of formula 18 with a compound of formula 23:




18




The compound of formula 18 can be treated with the compound of formula 23 under a suitable base to promote the condensation of the free amine with the acid group.

- 5 Preferably, the acid group is pre-activated to an ester. For example, the compound of formula 23 can be pre-activated by reaction with an alcohol such t-butanol, a coupling agent such as DIPC DI in a dry solvent such as N,N-dimethylformamide. To the pre-activated solution of the compound of formula 23 is added the compound of formula 18, and the mixture is stirred, e.g., at room temperature ($20\pm 3^\circ\text{C}$). The reaction can be carried out for a
- 10 suitable length of time, e.g., by stirring for about 12 hours. The solvent is removed, e.g., by evaporation, and the residue is dissolved in ethyl acetate, washed with a saturated solution of sodium bicarbonate, and followed by water, and by brine washes. The resulting product is dried, e.g., over anhydrous sodium sulfate to obtain the compound of formula 24.

- The present invention further provides a method for preparing a compound of
- 15 formula 18 wherein R_1 is a lipophile; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl, the method comprising:
- (a) treating a compound of formula 7




- 20 with a compound of the formula B:



31


to obtain a compound of formula 16



16


5 wherein P is an amine protecting group;

(b) treating the compound of formula 16 with an amine protected amino acid of the formula C



C

10 to obtain a compound of formula 17




17

and (c) removing the amine protecting group to obtain the compound of formula 18.

15 The compound of formula 7 is combined with compound B, which is an amine protected asparagine. Any suitable amine protecting group, e.g., Boc or Fmoc, can be employed. The compound B is pre-activated, e.g., by reacting with t-butanol and DIPCDI in dry DMF. The compound of formula 7 is added to pre-activated B. The resulting mixture is stirred, e.g., at room temperature (20±3°C). The reaction can be carried out for a suitable


length of time, e.g., by stirring for about 12 hours. The solvent is removed, e.g., by evaporation, and the residue is washed with ethyl acetate (e.g., up to 5 times). The product can be purified, e.g., by chromatography.

The present invention further provides a method for preparing a compound of the
5 formula 7 comprising: (a) treating compound D




D

with compound E




E

to obtain a compound of formula 12




12

15 (b) replacing a hydrogen on the CH₂ adjacent to the carbonyl carbon (1) of the compound of formula 12 with a lipophile to obtain a compound of formula 13:




(c) reducing the compound of formula 13 to obtain the compound of formula 14




and (d) replacing the hydroxyl group of the compound of formula 14 with an amino group, thereby obtaining the compound of formula 7; wherein R_1 is a lipophile and M is a metal.

The compound of formula 7 can be prepared from the Evan's reagent, (R)-(-)-4-phenyl-2-oxazolidinone (compound of formula 12). The compound of formula 12 is prepared by coupling or condensing compounds D and E. The metal M can be an alkali metal such as lithium. The coupling is carried out by the use of trimethylacetyl chloride in a suitable solvent such as N-methyl pyrrolidone. The resulting compound of formula 12 is converted to the compound of formula 13 by treating the compound of formula 12 with a base and a haloalkylaryl compound, e.g., 1-bromomethyl naphthalene. A suitable base is LiHMDS. The compound of formula 14 can be obtained by treating the compound of formula 13 with a reducing agent, e.g., a metal hydride such as $LiAlH_4$, preferably at a low temperature, e.g., at -78°C to room temperature. The compound of formula 14 is contacted with phthalidimide and triphenyl phosphine. The compound of formula 7 can be obtained by treating with ethanol, water, and hydrazine.

The present invention further provides a method for preparing a compound of formula 23 comprising (a) providing a compound of formula 8




(b) treating the compound of formula 8 with a base and trisyl azide to obtain a compound of formula 22



and (c) treating the compound of formula 22 with an alkaline peroxide, thereby obtaining the compound of formula 23; wherein R₂, in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group. An example of a suitable base is NaHMDS. The treating (b) is carried out at a low temperature, e.g., at -78°C. In (c), an alkali such as lithium hydroxide can be employed in combination with hydrogen peroxide.

The present invention further provides a method for preparing a compound of formula 9 comprising: (a) providing a compound of formula 9




and (b) treating the compound of formula 9 with vinyl magnesium bromide and PhSCu; wherein R₂, in combination with the phenyl ring, is a phenylphosphate mimic group or a

35

protected phenylphosphate mimic group.

The present invention further provides a method for preparing a compound of formula 9 comprising treating a compound of formula 10



5


10

with a compound of formula 11, wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group, and X is a halogen. A preferred halogen is bromine. Compounds of formulas 10 and 11 are combined.

A base such as triethylamine, a palladium catalyst, e.g., $Pd(OAc)_2$, and a phosphine such as tri-o-tolylphosphine are employed. The reaction can be carried out at reflux.

10


The present invention further provides a method for preparing a compound of formula 10 comprising treating compound F



15

F

with compound G



G.


The treating (coupling or condensing) can be carried out in the presence of an acid chloride, e.g., pivalyl chloride, a trialkylamine such as triethylamine, and a base such as BuLi at a low temperature, e.g., $-78^\circ C$ to room temperature.

20

The present invention further provides a method for preparing a compound of


36

formula 4 comprising:



4


- 5 (a) treating the compound of formula 18 with a compound of formula 6



6


to obtain a compound of formula 5

10



5

- (b) treating the compound of formula 5 with a Grubbs catalyst to compound of formula 19



19

and (c) treating the compound of formula 19 with a mixture of trifluoroacetic acid, water, and trimethylsilane to obtain the compound of formula 4; wherein R₁ is a lipophile, R₂, in
 5 combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R₄ and R₅, independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl. The treating (coupling or condensing) of the compounds of formulas 18 and 6 can be carried out by pre-activating the acid to an ester, e.g., by contacting with t-butanol and DIPCDI, and the pre-activated
 10 compound is contacted with the compound of formula 18. The conversion of the compound of formula 5 to the compound of formula 19 can be carried out by the use of a Grubbs catalyst.

The following examples further illustrate the present invention but, of course, should not be construed as in any way limiting its scope.

15

EXAMPLE 1

This Example illustrates a method of preparing some of the compounds or intermediates of the present invention. The method is illustrated schematically in Figs. 1-9.
(S)-N-Acroyl-4-phenyl-2-oxazolidinone, 10. To a solution of acrylic acid (5.188g, 72 mmol) in 200 ml of anhydrous THF containing 7.283g of N-methylmorpholine was added
 20 pivaloyl chloride (8.68g, 72 mmol) at 0°C under argon atmosphere, then stirred at -78°C for 1 hr. Then a cooled solution of Lithium (s)-4-phenyl-2-oxazolidinone (prepared by addition of BuLi (1.6 M, 37.5 ml, 60 mmol) to a suspension of (s)-(+)-4-phenyl-2-oxazolidinone (9.791g, 60 mmol) in 200 ml of anhydrous THF at -78°C under argon, stirred for 30 min) was added by cannula. After being stirred at -78°C for 2 hr, the resulting solution was raised
 25 to room temperature, stirred overnight. 100 ml of saturated NH₄Cl solution was added to

quench the reaction, and the THF was evaporated under reduced pressure, the residue was extracted with ethyl acetate, washed with water and saturated brine, dried over Na₂SO₄.

Concentration and purification by silica gel chromatography (Hexanes and ethyl acetate, from 10:1 to 4:1) afforded 12.304 g of product 10 (78.7% of Yield). NMR (in CDCl₃):

5 7.584~7.475 (1H, dd, J = 10.5, 16.85 Hz), 7.443~7.302 (5H, m), 6.534~6.459 (1H, dd, J = 1.71, 17.09 Hz), 5.917~5.868 (1H, dd, J = 1.71, 10.26 Hz), 5.531~5.481 (1H, dd, J = 3.91, 8.55 Hz), 4.811 (1H, t, J = 8.79 Hz), 4.421~4.370 (1H, dd, J = 3.91, 8.79 Hz).

Di-tert-butyl p-Bromobenzylphosphite 11a. To the solution of di-tert-butylphosphite (9.715 g, 50 mmol) in 150 ml of anhydrous THF was added BuLi (2.5 M in hexanes, 20 ml, 10 50 mmol) dropwise at -78°C under Argon, the solution was stirred at -78°C for 30 min, then raised to 0°C, stirred for additional 1 hr. To the resulting solution was added a solution of 4-bromobenzyl bromide in 50 ml of anhydrous THF at 0°C, after stirring for additional 2 hr at the same temperature, the solution was raised to room temperature, stirred overnight. 100 ml of water was added to quench the reaction and the THF was evaporated under reduced 15 pressure. The residue was extracted with ethyl acetate, washed with water and saturated brine, dried over Na₂SO₄. Concentration and purification by silica gel chromatography (Hexanes and ethyl acetate, from 10:1 to 4:1) afforded 16.5 g of product as white solid 11a (91% of Yield). NMR (in CDCl₃): 7.415 (2H, d, J = 8.30 Hz), 7.157 (2H, dd, J = 2.44, 8.56 Hz), 2.987 (2H, d, J = 21.49 Hz), 1.431 (18H, s).

20 **Compound 9a.** To the mixture of N-acroyl-(s)-4-phenyl-2-oxazolidinone, 10 (6.402 g, 27.81 mmol), di-tert-butyl p-bromobenzylphosphite 11a (10.10g, 27.81 mmol), Palladium acetate (309 mg) and tri-*o*-tolylphosphine in 200 ml of round bottom flask was added, 100 ml of anhydrous triethylamine, the resulting solution was refluxed under Argon overnight, a large amount of solid precipitate appeared, and some silver mirror was generated on the 25 flask wall. The triethylamine solvent was evaporated and the residue was dissolved in 150 ml of dichloromethane, then the solid palladium compound was removed by filtration. The solution was washed with water and brine, dried over Na₂SO₄, and evaporated to dryness to give yellow solid product. The crude product was recrystallized through Hexanes-EtOAc-CH₂Cl₂ mixture to give 8.930 g of white solid product 9a, from the liquid solution, 2.95 g of 30 product was recovered by chromatography (combined yield 85.5%). NMR (in CDCl₃): 7.920 (1H, d, J = 14.89 Hz), 7.768 (1H, d, J = 16.36 Hz), 7.529 (2H, d, J = 7.57 Hz), 7.48~7.18 (7

H, m), 5.570 (1H, dd, $J = 3.66, 8.54$ Hz), 4.751 (1H, t, $J = 8.79$ Hz), 4.33 (1H, dd, $J = 3.91, 8.79$ Hz), 3.066 (2H, d, $J = 21.72$ Hz), 1.462 (18H, s); FAB-MS (+Ve): 500 (MH^+), 444 ($MH^+ - C_4H_8$), 388 ($MH^+ - 2C_4H_8$).

Compound 8a. To a slurry of cuprous thiophenoxide (1.7271g, 10 mmol) in 300 ml of dry ether under argon at -40°C was added vinyl magnesium bromide in THF (1.0 M, 30 ml, 30.0 mmol, 3 eq.) dropwise, the mixture was allowed to warm up to -25°C until a color change (from brown to black-green) indicating the formation of complex $PhSCu(RMgX)_n$ was observed, the mixture was allowed to stir at this temperature for additional 1 hr. Then to the mixture was added a pre-cooled (-40°C) solution of compound 9a in 100 ml of anhydrous THF dropwise, the reaction mixture was stirred at -40°C for 3 hr (the reaction was monitored by T.L.C., till the starting material disappeared). The reaction mixture was poured into ice-cooled NH_4Cl solution, the solid cuprous salt was removed by filtration. After evaporation of THF, the residue was extracted with ethyl acetate, washed with water, saturated brine, and dried over Na_2SO_4 . Concentration and purification by silica gel chromatography afforded 2.967g of product 8a as white solid (56% of yield). NMR (in $CDCl_3$): 7.410~7.100 (9H, m), 5.961 (1H, ddd, $J = 6.59$ Hz), 5.306 (1H, dd, $J = 3.41, 8.54$ Hz), 4.984 (1H, ddd, $J = 1.22, 1.22, 10.35$ Hz), 4.934 (1H, ddd, $J = 1.22, 1.22, 17.09$ Hz), 4.585 (1H, t, $J = 8.79$ Hz), 4.235 (1H, dd, $J = 3.32, 9.04$ Hz), 3.89 (1H, m), 3.49 (1H, dd, $J = 8.05, 16.35$ Hz), 3.334 (1H, dd, $J = 7.08, 16.36$ Hz), 3.006 (2H, d, $J = 21.49$ Hz), 1.426 (18H, s). FAB-MS (+VE): 472 ($MH^+ - C_4H_8$), 416 ($MH^+ - 2C_4H_8$).

Compound 6a. To the solution of compound 8a (527 mg, 1 mmol) in 16 ml of THF- H_2O mixture (3:1) at 0°C was added H_2O_2 (30%, 509 μl , 5.0 mmol) via a syringe over 1 min, this was followed by the addition of $LiOH$ (84 mg, 2 mmol in 2 ml of water). After stirring at 0°C for 1 hr, the reaction mixture was raised to room temperature, stirred overnight. 628 mg of Na_2SO_3 (5.0 mmol) in 4.0 ml of water was added to destroy the remained hydrogen peroxide, and THF was evaporated at 30°C . The remained mixture was extracted with dichloromethane to remove the Evan's reagent, then the aqueous solution was poured into 25 ml of ice-cooled 0.2 M HCl solution, extracted with ethyl acetate, washed with ice-cooled water and brine, dried over Na_2SO_4 , concentrated, and taken to dryness under high vacuum to give 204 mg (80% of yield) of product 6 as foam. NMR (in $CDCl_3$): 7.22~7.17 (2H, dd, $J = 2.20, 8.30$ Hz), 7.141 (2H, d, $J = 8.55$ Hz), 5.990 (1H, m), 5.12~5.00 (2H, m),

3.85 (1H, m), 2.955 (2H, d, $J = 21.48$ Hz), 2.798 (1H, dd, $J = 7.57, 15.13$ Hz), 2.696 (1H, dd, $J = 8.06, 15.38$ Hz), 1.409 (9H, s), 1.397 (9H, s). FAB-MS (+VE):

(R)-N-(4-pentenoic)-4-phenyl-2-oxazolidinone 12a. To a solution of 4-pentenoic acid (3.604g, 36mmol) in 100 ml of anhydrous THF containing 3.645g (36mmol) of N-methylmorpholine was added pivaloyl chloride (4.34g, 36mmol) at -78°C under argon atmosphere, the mixture was stirred at -78°C for 30 min, then raised to 0°C , stirred for additional 1 hr, then cooled to -78°C again, stirred for 15 min before adding the solution of lithium oxazolidinone [prepared from BuLi (1.6 M, 18.75 ml, 30mmol) and (R)-(-)-4-phenyl-2-oxazolidinone (4.895g, 30 mmol) in 200 ml of anhydrous THF at -78°C under argon, 30 min] through cannula. After stirred at -78°C for 2 hr, the resulting solution was raised to room temperature, stirred overnight. 100 ml of ice-cooled water was added to quench the reaction, and the THF was evaporated under reduced pressure, the residue was extracted with ethyl acetate, washed with water and saturated brine, dried over Na_2SO_4 . Concentration and purification by silica gel chromatography (Hexanes and ethyl acetate, from 10:1 to 4:1) afforded 6.82 g of product 12a (70% of Yield). NMR ((in CDCl_3): 7.435~7.270 (5H, m), 5.891~5.730 (1H, ddt, $J = 6.60, 10.26, 16.85$ Hz), 5.438 (1H, dd, $J = 3.66, 8.55$ Hz), 5.10~4.95 (2H, m), 4.704 (1H, t, $J = 8.79$ Hz), 4.297 (1H, dd, $J = 3.66, 9.03$ Hz), 3.061 (2H, t, $J = 7.08$ Hz), 2.370 (2H, m). FAB-MS (+VE): 246 (MH^+).

Compound 13a. To the solution of (R)-N-(4-pentenoic)-4-phenyl-2-oxazolidinone 12a (6.885 g, 27.3 mmol) in 150 ml of dry THF was added the solution of LiHMDS (1.0 M in THF, 27.3 ml, 27.3mmol) at -78°C under argon, the resulting solution was stirred at -78°C for 2 hr, then a pre-cooled solution of 1-(bromomethyl)-naphthalene (12.07g, 54.3mmol) in 50 ml of THF was added, the resulting solution was stirred at -78°C for 4hr, then raised to room temperature, stirred overnight. 100 ml of ice-cooled water was added to quench the reaction, and the THF was evaporated under reduced pressure, the residue was extracted with ethyl acetate, washed with water and saturated brine, dried over Na_2SO_4 . Concentration and purification by silica gel chromatography (Hexanes and ethyl acetate, from 20:1 to 6:1) afforded 9.256 g of product 13a (88% of Yield). NMR (in CDCl_3): 8.805 (1H, d, $J = 7.81$ Hz), 7.845 (1H, dd, $J = 2.19, 6.83$ Hz), 7.710 (1H, d, $J = 8.06$ Hz), 7.556~7.440 (2H, m), 7.304~7.201 (5H, m), 7.141 (1H, dd, $J = 0.73, 6.83$ Hz), 6.993 (2H, m), 5.926~5.761 (1H, ddt, $J = 7.32, 10.01, 16.84$ Hz), 5.398 (1H, dd, $J = 4.15, 8.79$ Hz), 5.142~5.045 (2H, m),

4.611 (1H, t, $J = 8.79$ Hz), 4.542 (1H, m), 4.144 (1H, dd, $J = 4.15, 8.79$ Hz), 3.424 (1H, dd, $J = 8.30, 13.67$ Hz), 3.165 (1H, dd, $J = 6.59, 13.91$ Hz), 2.519 (1H, m), 2.239 (1H, m). FAB-MS (+VE): 386 (MH^+)

Compound 14a. A solution of $LiAlH_4$ (0.90g, 95%, 22.47mmol) in 100 ml of dry THF was cooled to $-78^\circ C$, to the solution was added a pre-cooled solution of naphthalene compound 13a (8.654g, 22.47 mmol) in 50 ml of dry THF at $-78^\circ C$, the mixture was stirred at $-78^\circ C$ for 1 hr, then raised to $0^\circ C$ over 1 hr. After 0.5 hr at $0^\circ C$, the clear solution was cooled to $-78^\circ C$, and 15 ml of ethyl acetate was added to destroy the remained $LiAlH_4$. The reaction was quenched with aqueous NH_4Cl solution at $-78^\circ C$, and after warming up to room temperature, diluted with water, extracted with ether, washed with water and brine, dried over Na_2SO_4 . Concentration and purification by silica gel chromatography (Hexanes and ethyl acetate, from 20:1 to 10:1) afforded 3.630g (70% of yield) of product 14a as oil. NMR (in $CDCl_3$): 8.069 (1H, m), 7.871 (1H, m), 7.744 (1H, d, $J = 7.81$ Hz), 7.560~7.324 (1H, m), 5.976~5.810 (1H, m), 5.200~5.100 (2H, m), 3.610 (2H, d, $J = 5.13$ Hz), 3.156 (1H, dd, $J = 7.81, 13.92$ Hz), 3.060 (1H, dd, $J = 6.59, 13.91$ Hz), 2.30 ~2.20 (2H, m), 2.15~2.06 (1H, m); FAB-MS (+VE): 226 (M^+).

Compound 15a. To the solution of the alcohol 14a (3.492g, 15.43mmol) in 30 ml of THF was added phthalimide (2.282g, 15.43mmol) and triphenylphosphine (4.06g, 15.43mmol) at $0^\circ C$, this was followed by the addition of diisopropyl azodicarboxylate (3.131g, 15.43mmol). After stirring had been continued overnight at room temperature, THF was evaporated, the residue was purified by chromatography to give 4.01g (73% of yield) of product 15a as sticky oil. NMR (in $CDCl_3$): 7.962 (1H, m), 7.827~7.633 (6H, m), 7.518~7.309 (4H, m), 5.932~5.767 (1H, ddt, $J = 7.08, 10.25, 17.09$ Hz), 5.15~5.00 (2H, m), 3.785 (1H, dd, $J = 7.57, 13.67$ Hz), 3.688 (1H, dd, $J = 7.33, 13.67$ Hz), 3.094 (2H, d, $J = 7.32$ Hz), 2.696 (1H, m), 2.151 (2H, m). FAB-MS (+VE): 356 (MH^+).

Chiral amine 7a. To the solution of phthalimide 15a (3.945g, 11.1mmol) in 100 ml of ethanol containing 820 μ l of water was added 867 mg (26.7mmol) of hydrazine, the resulting solution was refluxed under argon for 4 hr, and a large amount of solid precipitate appeared. The solution was cooled to room temperature, the solid was filtered off, washed with small amount of ethanol. The combined liquid solution was evaporated to dryness, the residue oil was purified by silica gel chromatography (Hexanes and ethyl acetate, 2:1, plus

1% v/v of NH_3 -MeOH solution) to give 1.916g (77% of yield) of desired product 7a as oil. NMR (in CDCl_3): 8.072 (1H, m), 7.863 (1H, m), 7.733 (1H, d, $J=8.06$ Hz), 7.551~7.443 (2H, m), 7.398 (1H, t, $J=7.08$ Hz), 7.313 (1H, m), 5.950~5.785 (1H, ddt, $J=7.08, 10.36, 17.09$ Hz), 5.175~5.050 (2H, m), 3.065 (2H, d, $J=7.32$ Hz), 2.718 (2H, d, $J=5.61$ Hz),
5 2.30~2.10 (2H, m), 2.05~1.92 (1H, m); FAB-MS (+VE): 226 (MH^+).

Compound 16a. To the solution of amine 7a (1.9164g, 8.51mmol) was added the pre-activated Boc-Asn [Formed by the reaction of Boc-Asn (1.974g, 8.51mmol), HOBT (1.15g, 8.51mmol), and DIPCDI (1.078g, 8.51mmol) in 20 ml of dry DMF, 10 min]. The resulting mixture was stirred at room temperature for 12hr, then DMF was evaporated. The resulting
10 solid compound was washed with ethyl acetate (30 ml x 5) to give 3.44g of pure compound 16a, the combined liquid phase was concentrated, purified by silica gel flash chromatography (chloroform and methanol, from 20:1 to 9:1) to give 112mg of product 16a (95% of combined yield). NMR (in CDCl_3): 8.025~7.961 (1H, m), 7.870~7.831 (1H, m), 7.724 (1H, d, $J=7.82$ Hz), 7.551~7.297 (4H, m), 6.955 (1H, s, br), 6.30~6.10 (2H, m, br),
15 5.920~5.750 (1H, m), 5.568 (1H, s, br), 5.175~5.050 (2H, m), 4.50~4.350 (1H, m), 3.252 (2H, m), 3.108~2.838 (3H, m), 2.561~2.474 (1H, dd, $J=6.35, 15.63$ Hz), 2.121 (3H, m), 1.444 (9H, s); FAB-MS (+VE): 440 (MH^+), 384 ($\text{MH}^+ - \text{C}_4\text{H}_8$)

Compound 17a. To the suspension of compound 16a (3.56g, 8.08mmol) in 20ml of dichloromethane was added 15 ml of trifluoroacetic acid, the suspension turned to be a clear
20 solution. The solution was stirred at room temperature for 1 hr, then dichloromethane and TFA was evaporated. The residue was dissolved in 150 ml of ethyl acetate, neutralized with saturated NaHCO_3 solution, then washed with brine, dried over Na_2SO_4 , concentrated, taken to dryness under high vacuum to give quantity yield of free amine as an oil. NMR (in CDCl_3): 8.10~7.27 (7H, m), 5.87~5.65 (1H, m), 5.20~4.90 (2H, m), 3.95 (1H, m, br),
25 3.40~2.60 (5H, m), 2.25~2.00 (4H, m). The obtained amine was dissolved in 20 ml of dry DMF, to the solution was added the pre-activated Fmoc-1-amino-cyclohexane carboxylic acid [Formed by reaction of Fmoc-1-amino-cyclohexane carboxylic acid (3.107g, 8.51mmol), HOBT (1.15g, 8.51mmol), DIPCDI (1.078g, 8.51mmol) in 20 ml of DMF, 10 min], the resulting mixture was stirred at room temperature for 12 hr, the DMF was
30 evaporated, the residue was dissolved in 200 ml of dichloromethane, washed with saturated NaHCO_3 solution, water, and brine, dried over Na_2SO_4 . Concentration and purification by

silica gel chromatography (chloroform and methanol, from 20:1 to 9:1) afforded 4.088g of 17a (70% of yield). NMR (in CDCl₃): 8.105 (1H, d, J = 8.06 Hz), 7.997 (1H, d, J = 8.06 Hz), 7.806~7.645 (9H, m), 7.480~7.189 (11H, m), 6.061 (1H, s, br), 5.83~5.66 (1H, m), 5.267 (1H, s, br), 5.232 (1H, s, br), 5.07~4.90 (9H, m), 4.747 (1H, m), 4.306~4.006 (3H, m), 3.32~3.00 (4H, m), 2.890 (1H, dd, J = 5.13, 14.16 Hz), 2.426 (1H, dd, J = 4.39, 14.89 Hz), 2.158~1.530 (13H, m); FAB-MS (+VE): 687 (M⁺)

Compound 18a. Compound 17a (4.088g, 5.957mmol) was dissolved in 50 ml of acetonitrile, to the solution was added 4.0 ml of piperidine, the resulting solution was stirred at room temperature for 2 hr. The solvent was evaporated, the residue was purified by silica gel chromatography to give free amine 2.707g 18a (98% of yield). NMR (in CDCl₃): 8.893 (1H, d, J = 6.84 Hz), 7.982 (1H, m), 7.851 (1H, m), 7.724 (1H, d, J = 7.81 Hz), 7.55~7.258 (7H, m), 6.309 (1H, s, br), 5.905~5.754 (1H, m), 5.540 (1H, s, br), 5.134~5.070 (2H, m), 4.683 (1H, dd, J = 6.10, 10.74 Hz), 3.30~3.209 (2H, m), 3.076 (1H, dd, J = 6.59, 14.16 Hz), 2.966 (1H, dd, J = 6.35, 13.92 Hz), 2.825~2.611 (9H, dd, J = 4.39, 15.14 Hz), 2.561 (1H, dd, J = 6.53, 14.59 Hz), 2.166 (2H, d, J = 5.86 Hz), 2.054~1.234 (10H, m).

Compound 5a. To the solution of amine 18a (376g, 0.801 mmol) in 5 ml of dry DMF was added the pre-activated ester solution of 6a [Formed by the reaction of 6a (306mg, 0.801mmol), HOBt (109mg, 0.801mmol), and DIPCDI (101 mg, 0.801 mmol) in 5 ml of dry DMF, 10 min]. The resulting mixture was stirred at room temperature for 12hr, then DMF was evaporated. The residue was dissolved in 50 ml of ethyl acetate, washed with saturated NaHCO₃ solution, water, and brine, dried over Na₂SO₄. Concentration and purification by silica gel chromatography (chloroform and methanol, from 20:1 to 9:1) afforded 569mg (86% of yield) of product 5a as foam. NMR (in CDCl₃): 8.054 (1H, d, J = 8.05 Hz), 7.85~7.64 (4H, m), 7.550~7.384 (4H, m), 7.154 (2H, dd, J = 2.44, 7.81 Hz), 6.991 (2H, d, J = 8.05 Hz), 6.283 (1H, s, br), 5.959~5.772 (H, m), 5.414 (1H, s, br), 5.125~4.903 (4H, m), 4.719 (1H, m), 3.728 (1H, M), 3.367 (1H, m), 3.200~2.910 (3H, m), 2.650~2.450 (3H, m), 2.300~2.005 (5H, m), 1.950~1.090 (11H, m), 1.434 (9H, s), 1.414 (9H, s); FAB-MS (+VE): 828 (M⁺), 772 (M⁺-C₄H₈), 716 (M⁺-2C₄H₈).

Compound 19a. To the solution of compound 5a (429 mg, 0.518 mmol) in 120 ml of anhydrous dichloromethane (deoxygenated with Argon) was added via a syringe of Ruthenium catalyst (180 mg, 0.218 mmol, 0.4 eq.) in 37 ml of dichloromethane, the solution

was refluxed under Argon for 60hr, monitored by T.L.C.. Then, the solvent was evaporated, the residue was purified by silica gel chromatography (chloroform-EtOAc-MeOH, from 2:1:0 to 14:7:1) to give 279 mg (67% of yield) of product 19a. FAB-MS (+VE): 801 (MH^+), 745 ($MH^+ - C_4H_8$), 689 ($MH^+ - 2C_4H_8$).

- 5 **Product 4a.** The compound 19a (179 mg) was treated with a solution made of TFA-TES- H_2O (3.7 ml-0.1 ml-0.2 ml) at room temperature, after 1 hr, the solvents were evaporated. To the residue was added ether, a large amount of solid product precipitated, the product was separated by centrifuge, then washed with ether, separated by centrifuge again, the resulting solid product was dried under high vacuum to give 144 mg of crude product. The
- 10 crude compound was dissolved in 16 ml of acetonitrile-water solution (1:1), purified by HPLC to give 73 mg (60% of yield) of pure compound 4a. NMR (in DMSO): 8.395 (1H, s), 8.234 (1H, d, $J = 7.81$ Hz), 8.116 (1H, d, $J = 8.06$ Hz), 7.893 (1H, d, $J = 7.57$ Hz), 7.750 (1H, m), 7.561~7.375 (6H, m), 7.150~7.020 (5H, m), 5.550 (1H, dd, $J = 9.52, 14.16$ Hz), 5.43~5.30 (1H, m), 4.269 (1H, m), 3.861 (1H, m), 3.587 (1H, dd, $J = 5.61, 11.72$ Hz), 3.156
- 15 (1H, dd, $J = 5.66, 14.40$ Hz), 2.950~2.300 (9H, m), 2.250~1.100 (11H, m); FAB-MS (-VE): 687.6 ($M^+ - H$); HR-MS for $C_{37}H_{45}N_4O_7P$: Calcd.: 687.2948; Found: 687.2958.

- Compound 22a.** To the solution of sodium hexamethyl disilylamide (1.0 M in THF, 3.33 ml, 3.33 mmol) at $-78^\circ C$ was added via cannula a pre-cooled solution ($-78^\circ C$) of compound 8 (1.596g, 3.024mmol) in 10 ml of THF, the solution turned to be deep purple quickly. After
- 20 stirring at $-78^\circ C$ for 20 min, a pre-cooled ($-78^\circ C$) solution of trisyl azide (1.126g, 3.628mmol) was added via cannula, the resulting solution was stirred at $-78^\circ C$ for 5 min, then the reaction was quenched by the addition of glacial acetic acid (0.95g, 16.6mmol) followed by the addition of potassium acetate in 15 ml of THF, the mixture was then raised to $30^\circ C$, stirred for additional 1.5 hr. 20 ml of saturated $NaHCO_3$ solution was added to the
- 25 reaction mixture, and then THF was evaporated, the remained product was extracted with ethyl acetate, washed with brine, dried over Na_2SO_4 . Concentration and purification by silica gel chromatography (hexanes and ethyl acetate, from 4:1 to 2:1) afforded 1.26 g of product 22a (73.3% of yield). NMR (in $CDCl_3$): 7.40~7.14 (9H, m), 6.23~6.08 (1H, ddd, $J = 8.31, 10.01, 17.09$ Hz), 5.480 (1H, d, $J = 10.25$ Hz), 5.289 (1H, d, $J = 18.06$ Hz), 5.272 (1H, d, $J =$
- 30 9.27 Hz), 4.902 (1H, dd, $J = 2.44, 8.06$ Hz), 4.235 (1H, t, $J = 8.54$ Hz), 4.040 (1H, dd, $J = 2.69, 8.79$ Hz), 3.824 (1H, t, $J = 8.55$ Hz), 3.008 (2H, d, $J = 21.24$ Hz), 1.454 (9H, s). . FAB-

MS (+VE): 569 (MH^+), 513 ($M^+ - C_4H_8$), 457 ($M^+ - 2C_4H_8$).

Compound 23a. To the solution of compound 22a (1.2356 mg, 2.17 mmol) in 32 ml of THF-H₂O mixture (3:1) at 0°C was added H₂O₂ (30%, 1.107 ml, 10.85 mmol) via a syringe over 1 min, this was followed by the addition of LiOH (182.4 mg, 4.34 mmol in 4.4 ml of water). After stirring at 0°C for 1 hr, the reaction mixture was raised to 30°C, for additional 4 hr. 1.367g of Na₂SO₃ (10.85mmol) in 8.5 ml of water was added to destroy the remained hydrogen peroxide, and THF was evaporated at 30°C. The remained mixture was extracted with dichloromethane to remove the Evan's reagent, then the aqueous solution was poured into 54 ml of ice-cooled 0.2 M HCl solution, extracted with ethyl acetate, washed with ice-cooled water and brine, dried over Na₂SO₄, concentrated, and taken to dryness under high vacuum to give 841 mg (92% of yield) of product 23a as white foam. NMR (in CDCl₃): 7.218 (2H, d, J = 7.81 Hz), 7.143 (2H, dd, J = 2.44, 7.81 Hz), 6.208~6.060 (1H, m), 5.246 (1H, d, J = 13.67 Hz), 5.245 (1H, J = 16.60 Hz), 3.950 (1H, d, J = 9.28 Hz), 3.849 (1H, t, J = 9.27 Hz), 2.779 (2H, dd, J = 12.21, 21.24 Hz), 1.406 (9H, s), 1.346 (9H, s). FAB-MS (-VE): 422 (M-H).

Compound 24a. To the solution of amine 18a (290 mg, 0.626 mmol) in 5 ml of dry DMF was added the pre-activated ester solution of 23a [Formed by the reaction of 23a (265mg, 0.626mmol), HOBt (84.7 mg, 0.626mmol), and DIPCDI (79.1mg, 0.626mmol) in 5 ml of dry DMF, 10 min]. The resulting mixture was stirred at room temperature for 12hr, then DMF was evaporated. The residue was dissolved in 50 ml of ethyl acetate, washed with saturated NaHCO₃ solution, water, and brine, dried over Na₂SO₄. Concentration and purification by silica gel chromatography (chloroform and methanol, from 20:1 to 9:1) afforded 363mg (67% of yield) of product 24a as a foam.

Compound 25a. To the solution of compound 24a (363 mg, 0.412 mmol) in the mixture of THF-water (8 ml-0.5 ml) was added triphenyl phosphine (163 mg, 0.618mmol, 1.5eq), the resulting mixture was stirred at room temperature for 12 hr, then THF was evaporated and the residue was purified by chromatography (chloroform and methanol, from 100:0 to 9:1) to give compound 268 mg of 25a (77% of yield).

General Procedures employed in the following: Elemental analyses were obtained from Atlantic Microlab Inc., Norcross. Solvent was removed by rotary evaporation under reduced pressure and silica gel chromatography was performed using high performance silica gel (60

Å pore, 10 µ particle size). Anhydrous solvents were obtained commercially and used without further drying. Analytical HPLC were conducted using a Vydac C₁₈ column (10 mm dia. x 250 mm long; solvent A = 0.1% aqueous TFA; solvent B = 0.1% TFA in acetonitrile; flow rate = 2 mL/min.).

- 5 **(S)-4-Benzyl-3-(prop-2-enoyl)-1,3-oxazolidin-2-one (39).** To a solution of acrylic acid (7) (8.2 mL; 120 mmol) and N-methyl morpholine (NMM) (13.2 mL; 120 mmol) in anhydrous THF (100 mL) at -78° C under argon was added pivaloyl chloride (14.8 mL; 120 mmol) and the mixture was stirred at -78° C (1 h). To a separate round bottomed flask containing Evan's reagent, (S)-(-)-4-Benzyl-2-oxazolidinone (38) (17.7 g; 100 mmol) in anhydrous
- 10 THF (200 mL) at -78° C under argon, was added *n*-BuLi, 1.6 M in hexanes (62 mL; 100 mmol) and the solution was stirred at -78° C (30 min). To this was then added via cannula at -78° C, the suspension of acryloyl mixed anhydride, and the resulting mixture was stirred first at -78° C (1.5 h) then at room temperature (2 h). The mixture was partitioned between saturated NH₄Cl / EtOAc and the combined organics were washed with brine, dried
- 15 (MgSO₄) and taken to dryness to yield a syrup (27.6 g). Purification by silica gel chromatography (EtOAc in hexanes, from 0% to 50%) provided unreacted Evan's reagent (6.44 g) as well as product (39) as a white foam, which crystallized (7.16 g; 49% based on recovered starting material), mp 70 - 72° C. ¹H NMR (CDCl₃) δ 7.61 (dd, 1H, *J* = 10.7, 17.1 Hz), 7.49 ~ 7.26 (m, 5 Hz), 6.70 (dd, 1H, *J* = 1.7, 17.1 Hz), 6.03 (dd, 1H, *J* = 1.7, 10.2 Hz),
- 20 4.99 ~ 4.79 (m, 1H), 3.37 ~ 4.25 (m, 2 H), 3.44 (dd, 1H, 3.0, 13.2 Hz), 2.92 (dd, 1H, *J* = 9.8, 13.2 Hz). FABMS (⁺VE, NBA) *m/z* 232 (MH⁺). Anal. calcd for C₁₃H₁₃NO₃: C, 67.52; H, 5.67; N, 6.06. Found: C, 67.57; H, 5.82; N, 6.08.

- Bis-(*tert*-butyl) ((4-bromophenyl)methyl)phosphonate (40).** To a solution of di-*tert*-butyl phosphite (9.70 g; 50 mmol) in anhydrous THF (100 mL) at -78° C under argon, was added
- 25 *n*-BuLi, 1.6 M in hexanes (33.2 mL; 50 mmol) over 5 minutes and the solution was stirred first at -78° C (30 minutes), then at 0° C (30 minutes). To this was added a solution of 4-bromobenzyl bromide (12.5 g; 50 mmol) in anhydrous THF (20 mL) and the mixture was stirred at 0° C, then allowed to come to room temperature and stirred overnight. The mixture was partitioned between saturated aqueous NH₄Cl and EtOAc; the combined organic
- 30 extracts were washed with brine; dried (MgSO₄) and taken to dryness, yielding a light yellow crystalline solid (18.27 g). Purification by silica gel chromatography (EtOAc in

hexanes; from 0% to 100%), provided 40 as a cream colored crystalline solid (15.21 g; 82% yield), mp 45° (soften), 48° - 56° C. ¹H NMR (CDCl₃) δ 7.42 (d, 2H *J* = 8.3 Hz), 7.16 (dd, 2H *J* = 2.4, 8.3 Hz), 2.99 (d, 2H *J* = 21.5 Hz), 1.43 (s, 18 H). HR-FABMS (⁺VE) *m/z* calcd. for C₁₅H₂₅N₃O₃PBr (M⁺): 3363.0725. Found: 3963.0735 (δ *m* = 2.8 ppm). Anal. calcd for C₁₅H₂₄BrO₃P: C, 49.60; H, 6.66. Found: C, 50.20; H, 6.60.

3-(3-(4-((Bis-(*tert*-butyl)phosphono)methyl)phenyl)prop-2-enoyl)-4-(S)-benzyl-1,3-oxazolidin-2-one (41).

A mixture of acrylamide 39 (1.48 g; 6.40 mmol), di-*tert*-butyl (4-bromobenzyl)phosphonate (40) (2.36 g; 6.40 mmol), Pd(OAc)₂ (72 mg; 0.032 mmol) and tri-*o*-toluyl phosphine (390 mg; 1.28 mmol) in a round bottom flask sealed with a rubber septum, was alternately evacuated and flushed with argon (3 x), then to this was added NEt₃ (12 mL) and the mixture was heated with stirring at 85° C (overnight). The resulting suspension was partitioned between ice-cold saturated NH₄Cl/EtOAc and the combined organics were washed with brine, dried (MgSO₄) and taken to dryness to yield a foam (3.73 g). Purification by silica gel chromatography (EtOAc in hexanes, from 20% to 100%)

provided 41 as a crystalline solid (2.58 g, 79%). Recrystallization from ether: hexanes provided an analytical sample, mp 128 - 129° C. ¹H NMR (CDCl₃) δ 7.98 (s, 2H), 7.66 (d, sH, *J* = 8.1 Hz), 7.48 ~ 7.31 (m, 1H), 4.95 ~ 4.84 (m, 1H), 4.38 ~ 4.27 (m, 2H), 3.46 (dd, 1H, *J* = 3.0 13.2 Hz), 3.15 (d, 2H, *J* = 22.2 Hz), 2.94 (dd, 1H, *J* = 9.4, 13.2 Hz), 1.52 (s, 18 H). FABMS (⁺VE) *m/z* 514 (MH⁺). Anal. calcd for C₂₈H₃₆NO₆P: C, 65.48; H, 7.07; N, 2.73. Found: C, 65.74; H, 7.06; N, 2.81.

3-(3-(4-((Bis-(*tert*-butyl)phosphono)methyl)phenyl)propanoyl)-4-(S)-benzyl-1,3-oxazolidin-2-one (42).

A solution of 41 (6.48 g; 12.7 mmol) in absolute EtOH (50 mL) was hydrogenated over Pd black (200 mg) at 40 psi H₂ in a Parr apparatus (overnight).

Additional Pd black was added (200 mg) and hydrogenation continued (overnight). Catalyst

was removed by filtration and the filtrate was taken to dryness to yield a white crystalline solid, which was triturated with ether to provide a white solid. This was combined with additional product obtained by cooling the filtrate to -78° C, providing 42 as a combined total of 5.37 g (82% yield), mp 114 - 115° C. ¹H NMR (CDCl₃) δ 7.43 ~ 7.34 (m, 4H), 7.27 (s, 5H), 4.80 ~ 4.66 (m, 1H), 4.28 ~ 4.23 (m, 2H), 3.40 ~ 3.30 (m, 3H), 3.15 ~ 3.05 (m, 4H), 2.84 (dd, 1H, *J* = 9.4, 13.2 Hz), 1.53 (s, 18 H). FABMS (⁺VE, NBA) *m/z* 516 (MH⁺), 404 (MH⁺ - 2C₄H₈). Anal. calcd for C₂₈H₃₈NO₆P: C, 65.23; H, 7.43; N, 2.72. Found: C, 64.94;

H, 7.34; N, 2.79.

(3-(4-((Bis-(*tert*-butyl)phosphono)methyl)phenyl)propanoic acid (35). To a solution of 42 (1.0 g; 1.95 mmol) in THF (15 mL) with H₂O (5 mL) at 0° C, was added aqueous H₂O₂, 30% w/w (1.10 mL; 9.74 mmol) dropwise, followed by dropwise addition of an ice-cold solution of LiOH.H₂O (164 mg; 3.89 mmol) in H₂O (10 mL), and the mixture was then stirred at 0° C (4 h). The mixture was diluted with H₂O (100 mL), washed with CH₂Cl₂, then the aqueous was cooled to ~0° C and ice-cold 0.1 N HCl was added (~80 mL) and the mixture was then extracted with EtOAc. Combined organics were dried (MgSO₄) and taken to a syrup, which quickly crystallized to provide 35 as a white solid (563 mg; 81% yield), mp 101 (soften); 106 - 110° C. ¹H NMR (CDCl₃) δ 7.28 (dd, 2H, *J* = 2.1, 8.1 Hz), 7.21 (d, 2H, *J* = 8.1 Hz), 3.09 (d, 2H, *J* = 21.4 Hz), 3.02 (t, 2H, *J* = 7.7 Hz), 2.72 (t, 2H, *J* = 7.7 Hz), 1.49 (s, 18 Hz). FABMS (VE, NBA) *m/z* 355 (M-H). Anal. calcd for C₁₈H₂₉O₅P: C, 60.66; H, 8.20. Found: C, 60.90; H, 8.10.

3-(2-(S)-Azido-3-(4-((bis-(*tert*-butyl)phosphono)methyl)phenyl) propanoyl)-4-(S)-benzyl-1,3-oxazolidin-2-one (43). To anhydrous THF (50 mL) at -78° C under argon was added potassium bis(trimethylsilyl)amide, 0.5 M in toluene (29.0 mL; 14.5 mmol), followed via cannula, a pre-cooled (-78° C) solution of 42 (6.20 g; 12.1 mmol) in anhydrous THF (50 mL), and the resulting violet solution was stirred at -78° C (30 minutes). To this was added rapidly via cannula, a pre-cooled (-78° C) solution of (2,4,5-tri-isopropyl)phenylsulfonyl azide (4.50 g; 14.5 mmol). The resulting yellow solution was stirred at -78° C (2 minutes), quenched by addition of HOAc (3.8 mL; 66.6 mmol) followed by solid KOAc (4.87 g; 49.7 mmol). The mixture was stirred at room temperature (3.5 h), then partitioned between saturated NaHCO₃ in brine / EtOAc; washed with saturated NaHCO₃ in brine; dried (MgSO₄) and taken to dryness to yield a yellow resin (8.66 g). Purification by silica gel chromatography (50% EtOAc in hexanes) provided 43 as a white foam (5.13 g, 77%).

Crystallization from ether provided an analytical sample, mp 80° C (soften); 115 - 117° C. ¹H NMR (CDCl₃) δ 7.46 ~ 7.27 (m, 9H), 5.37 (dd, 1H, *J* = 6.0, 9.0 Hz), 4.70 ~ 4.58 (m, 1H), 4.30 ~ 4.14 (m, 2H), 3.40 (dd, 1H, *J* = 3.0, 13.2 Hz), 3.28 (dd, 1H, *J* = 6.0, 13.2 Hz), 3.10 (d, 2H, *J* = 21.4 Hz), 2.90 (dd, 1H, *J* = 9.4, 13.2 Hz), 1.50 (s, 9H), 1.49 (s, 9H). FABMS (⁺VE) *m/z* 557 (MH⁺), 501 (MH⁺ - C₄H₈), 445 (MH⁺ - 2C₄H₈). Anal.

3-(2-(S)-Azido-3-(4-((bis-(*tert*-butyl)phosphono)methyl)phenyl) propanoic acid (34).

To a solution of 43 (4.77 g; 8.60 mmol) in THF (40 mL) with H₂O (10 mL) at 0° C, was added aqueous H₂O₂, 30% w/w (4.88 mL; 43.0 mmol) dropwise, followed by dropwise addition of an ice-cold solution of LiOH.H₂O (722 mg; 17.2 mmol) in H₂O (40 mL), and the mixture was then stirred at 0° C (2 h). To the solution was added Na₂SO₃ (5.42 g; 43.0 mmol) in H₂O (20 mL), then the mixture was diluted with brine (300 mL), washed with CH₂Cl₂, then the aqueous was cooled to ~0° C and ice-cold 1.0 N HCl was added until the pH ≤ 3, and the mixture was extracted with EtOAc. Combined organics were dried (MgSO₄) and solvent removed to provide 36 as a highly crystalline white solid (3.00 g; 88% yield), mp 68 - 72° C. ¹H NMR (CDCl₃) δ 7.29 (brs, 4H), 4.24 ~ 4.14 (m, 1H), 3.25 ~ 3.00 (m, 4H), 1.49 (s, 9H), 1.44 (s, 9H). FABMS (VE) *m/z* 396 (M-H). HR-FABMS (VE, MCA, Gly) *m/z* calcd. for C₁₈H₂₇N₃O₃P (M-H): 396.1688. Found: 396.1667 (δ *m* = 2.1 mmu; 5.3 ppm). [α]_D = +14.1° (c 0.95, CHCl₃). Anal. calcd for C₁₈H₂₈N₃O₅P•³/₄H₂O: C, 52.61; H, 7.24; N, 10.23. Found: C, 52.63; H, 6.77; N, 10.36.

Determination of enantiomeric purity of analogue 34. Dipeptides 44 and 45 were prepared from protected azido acid 34 using Rink amide resin (0.4 meq/g, purchased from Bachem Corp., Torrance, CA) with Fmoc-protocols similar to those previously described. Fmoc-D,L-Leu and Fmoc-L-Leu-Rink amide resins were prepared by coupling the appropriate Fmoc-protected amino acids to Rink resin, with the resulting Fmoc-protected resins (12.5 mg) then being washed well with several 1 mL portions of N-methyl-2-pyrrolidinone (NMP). Fmoc amino protection was removed by treatment with 20% piperidine in NMP (0.5 mL, 2 minutes followed by 0.5 mL, 20 minutes). Resins were washed well with NMP (10 x 1 mL) then coupled overnight with a solution of active ester formed by reacting 12.5 μmol each of protected azido acid 34, 1-hydroxybenzotriazole (HOBt) and 1,3-diisopropylcarbodiimide (DIPCDI) in NMP (1.0 mL, 10 minutes). Resins were first washed with NMP (10 x 1 mL) and dichloromethane (10 x 1 mL), then dipeptides were cleaved from the resin using a mixture of trifluoroacetic acid (TFA, 1.80 mL) and H₂O (200 μl) (1h), taken to dryness and analyzed by HPLC (linear gradient from 10% B to 90% B over 20 minutes). Retention times of diastereomeric peaks as determined using dipeptide 44 prepared from racemic D,L-leucine, indicated diastereomers eluting at 18.6 minutes and 19.1 minutes. Enantiomeric contamination of azido acid 34 was then determined by similar analysis of dipeptide 45, where diastereomeric contamination accounted for an area less than

3% of that observed for the major diastereomer. These results indicated greater than 94% enantiomeric purity.

Compounds 61-64 are illustrated in Fig. 10

Compound 63a. To a solution of amine 61 (85 mg; 0.20 mmol) in anhydrous DMF (1 mL) was added a preactivated ester solution formed by reacting des-amino Pmp 62a (71 mg; 0.20 mmol), HOBt (27 mg; 0.20 mmol), diisopropylcarbodiimide (DIPCDI) (31 μ L; 0.20 mmol) in DMF (1 mL; 10 minutes). The mixture was stirred at room temperature (overnight) then taken to dryness under high vacuum and purified by silica gel chromatography (from 50% EtOAc in CHCl₃ to 10% MeOH in EtOAc) to provide 63a as a white foam (118 mg; 77% yield). ¹H NMR (CDCl₃) δ 8.20 ~ 8.10 (m, 2H), 7.94 ~ 7.84 (m, 2H), 7.80 ~ 7.72 (m, 1H), 7.58 ~ 7.48 (m, 2H), 7.45 ~ 7.38 (m, 2H), 7.23 (dd, 2H, J = 1.7, 7.7 Hz), 7.10 (dd, 2H, J = 7.7 Hz), 6.64 (brs, 1H), 6.28 (brs, 1H), 5.65 (brs, 1H), 4.86 ~ 4.75 (m, 1H), 3.54 ~ 3.35 (m, 2H), 3.25 ~ 3.15 (m, 4H), 3.06 (d, 2H, J = 21.4 Hz), 3.00 ~ 2.87 (m, 2H), 2.70 ~ 2.50 (m, 4H), 2.35 ~ 1.25 (m, 12H), 1.50 (s, 18H). FABMS (+VE) m/z 763 (MH⁺). HR-FABMS (+VE, MCA, Gly) m/z calcd. for C₄₂H₆₀N₄O₇P M+H: 763.4200. Found: 763.4244 (Dm = 4.4 mmu; 5.8 ppm).

Compound 64a. A solution of 63a (103 mg; 13.5 mmol) in TFA (1.9 mL): H₂O (100 μ L): triethylsilane (TES) (150 μ L) was stirred at room temperature (1 h) then taken to dryness under high vacuum and purified by HPLC to provide 64a as a white solid (60 mg; 68% yield). Analytical HPLC (Vydac C18 Peptide and Protein column; 10 mm dia x 250 mm long; flow rate = 2 mL/min; solvent A = 0.1% aqueous TFA, solvent B = 0.1% TFA in acetonitrile; linear gradient 10% B - 90% B over 20 minutes) retention time = 21.7 minutes (>99%). ¹H NMR (DMSO-d₆) δ 8.28 ~ 8.10 (m, 2H), 7.97 ~ 7.92 (m, 1H), 7.84 ~ 7.76 (m, 1H), 7.74 ~ 7.66 (m, 1H), 7.58 ~ 7.48 (m, 2H), 7.46 ~ 7.42 (m, 2H), 7.16 (dd, 2H, J = 1.7, 8.1 Hz), 7.03 (d, 2H, J = 8.1 Hz), 4.45 ~ 4.37 (m, 1H), 3.27 ~ 3.17 (m, 2H), 3.15 ~ 3.05 (m, 4H), 2.94 (d, 2H, J = 20.9 Hz), 2.85 ~ 2.68 (m, 4H), 2.13 ~ 1.10 (m, 12H). FABMS (-VE) m/z 649 (M-H). HR-FABMS (-VE, MCA, Gly) m/z calcd. for C₃₄H₄₂N₄O₇P (M-H): 649.279. Found: 649.2803 (Dm = 1.2 mmu; 1.8 ppm).

Compound 63b. To a solution of amine 61 (85 mg; 0.20 mmol) in anhydrous DMF (1 mL) was added a preactivated ester solution formed by reacting azido Pmp 2b (79 mg; 0.20 mmol), HOBt (27 mg; 0.20 mmol), diisopropylcarbodiimide (DIPCDI) (31 μ L; 0.20 mmol)

in DMF (1 mL; 10 minutes). The mixture was stirred at room temperature (overnight) then taken to dryness under high vacuum and purified by silica gel chromatography (from 50% EtOAc in CHCl₃ to 5% MeOH in EtOAc) to provide 63b as a white foam (94 mg; 58% yield). ¹H NMR (CDCl₃) δ 8.38 (d, 1H, J = 7.7 Hz), 8.15 (d, 1H, J = 8.1 Hz), 7.96 ~ 7.75 (m, 3H), 7.60 ~ 7.40 (m, 4H), 7.25 ~ 7.09 (m, 6H), 6.78 (brs, 1H), 5.65 (brs, 1H), 4.84 ~ 4.72 (m, 1H), 4.36 ~ 4.25 (m, 1H), 3.60 ~ 3.35 (m, 2H), 3.30 ~ 2.85 (m, 6H), 3.07 (d, 2H, J = 21 Hz), 2.57 (dd, 1H, J = 4.3, 9.0 Hz), 2.27 ~ 1.10 (m, 12H), 1.48 (s, 18H). FABMS (+VE, NBA) m/z 804 (MH⁺).

Compound 64b. A solution of 63b (47 mg; 58 μmol) in TFA (900 μL): H₂O (100 μL) was stirred at room temperature (1 h) then taken to dryness under high vacuum and purified by HPLC to provide 64b as a white solid (32 mg; 80% yield). Analytical HPLC (Vydac C18 Peptide and Protein column; 10 mm dia x 250 mm long; flow rate = 2 mL/min; solvent A = 0.1% aqueous TFA, solvent B = 0.1% TFA in acetonitrile; linear gradient 10% B - 90% B over 20 minutes) retention time = 22.8 minutes (>99%). ¹H NMR (DMSO-d₆) δ 8.62 (s, 1H), 8.24 (d, 1H, J = 7.7 Hz), 8.18 ~ 8.10 (m, 1H), 7.98 ~ 7.92 (m, 1H), 7.62 ~ 7.52 (m, 3H), 7.46 ~ 7.40 (m, 2H), 7.24 (dd, 2H, J = 1.7, 8.1 Hz), 7.15 (d, 2H, J = 8.1 Hz), 7.02 (s, 1H), 4.55 ~ 4.45 (m, 1H), 4.20 (dd, 1H, J 3.4, 10.7 Hz), 3.35 ~ 2.68 (m, 8H), 2.97 (d, 2H, J = 21.4 Hz), 2.15 ~ 1.15 (m, 12H). FABMS (-VE, Gly) m/z 690 (M-H). HR-FABMS (-VE, MCA, Gly) m/z calcd. for C₃₄H₄₁N₇O₇P (M-H): 690.2805. Found: 690.2780 (Δm = 2.5 mmu; 3.6 ppm).

EXAMPLE 2

This Example illustrates some of the properties of the compounds in accordance with embodiments of the present invention.

To characterize the time period and dose which the Grb2 SH2 domain inhibitor (antagonist) compound of formula 4 enters intact cells, the murine IL-3-dependent cell line 32D c-met was used. The effects of the inhibitor on c-Met-Grb2 interaction was examined by co-immunoprecipitation/ immunoblot analysis. Cells were cultured in RPMI 1640 + 15% FBS and 5% WEHI-3B conditioned medium. Intact 32Dc-met cells were treated with different amount (3, 30, 300 nM) of the compound for 1, 2, 4, and 16 h, then briefly stimulated with HGF/NK1 for 10 minutes before lysis, immunoprecipitated with anti-Grb2 antibodies (Santa Cruz), and immunoblotting with anti-c-Met (Santa Cruz), or anti-Grb2

(Transduction Lab). HGF/NK1 stimulated co-immunoprecipitation of the 145 kDa c-Met beta subunit with Grb2. When cells were pretreated for 16 h with 30 nM or 300 nM compound of formula 4 prior to HGF/NK1 stimulation, the amount of HGF receptor that was co-immunoprecipitated with Grb2 was reduced by approximately 50%.

- 5 **Inhibition of Grb2 SH2 Domain Binding using ELISA Techniques.** A biotinated phosphopeptide encompassing a Grb2 SH2 domain binding sequence derived from SHC protein, was bound at 20 ng / mL to 96-well plates overnight. Nonspecific interactions were inhibited by 5% bovine serum albumin containing TBS. Samples of recombinant purified Grb2 SH2-GST fusion protein were preincubated with serial of dilutions of inhibitors, then
- 10 added into each well. After extensive washing with 0.1% bovine serum albumin in TBS, bound Grb2 SH2 domain was detected using anti-GST antibodies and goat anti-mouse antibody conjugated to alkaline phosphatase. Quantitation of bound alkaline phosphatase was achieved by a colorimetric reaction employing para-nitrophenyl phosphate as substrate. The results obtained are set forth in Fig. 11. Compound 3a showed greater inhibition
- 15 compared to compound 2, a compound which is not constrained in the beta bend fashion.

- Inhibition of Grb2 SH2 Domain Binding in Whole Cells.** ErbB2 over expressing breast cancer cells, MDA-MB-453, were treated with inhibitors (25 μ M) for 3 h in serum-free IMEM medium (Gibco). Cells were washed twice with PBS to remove inhibitor, then cell lysates were prepared using 1% Triton X-100 in PBS containing 0.2 mM NaVO₄. Grb2 and
- 20 associated Grb2-binding proteins were immunoprecipitated from each lysate (500 ug) with anti-Grb2 antibodies and collected using protein A Sepharose. Immunoprecipitated proteins were separated by SDS-PAGE on 8-16% gradient gels (Novex) and pTyr-containing proteins were detected by Western blotting using anti-phosphotyrosine antibodies (Upstate Biochemicals Inc.). Previous experiments have shown that a major tyrosine phosphorylated
- 25 protein in these cells is the p185 erbB-2, which is over expressed as a consequence of gene amplification. Western blotting with Grb2 MAb was done as a control. The results obtained are depicted in Fig. 12.

- Assay of Cell Proliferation.** Cell lines were obtained from the American Type Culture Collection (Rockville, MD) and Lombardi Cancer Center, Georgetown University Medical
- 30 Center. Cells were routinely maintained in improved minimal essential medium (IMEM, Biofluids, Rockville, MD) with 10% fetal bovine serum. Cultures were maintained in a

humidified incubator at 37°C and 5% CO₂. The effect of Grb2 inhibitors on cell proliferation was determined by direct cell counting. Briefly, 25,000 cells were plated into 24-well plates and the Grb2 inhibitors at appropriate concentrations were added and cultured for 8 to 10 days. Cells were collected every other day and counted with a Coulter counter.


5 The results obtained are depicted in Fig. 13.

All of the references cited herein, including publications, patents, and patent applications, are hereby incorporated in their entireties by reference.

While the invention has been described in some detail by way of illustration and example, it should be understood that the invention is susceptible to various modifications
10 and alternative forms, and is not restricted to the specific embodiments set forth. It should be understood that these specific embodiments are not intended to limit the invention but, on the contrary, the intention is to cover all modifications, equivalents, and alternatives falling within the spirit and scope of the invention.


WHAT IS CLAIMED IS:

1. A compound of formula (I):



(I)


- 5 wherein R_1 is a lipophile; R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; R_3 is hydrogen, azido, amino, carboxyalkyl, alkoxy-carbonylalkyl, aminocarbonylalkyl, or alkylcarbonylamino, wherein the alkyl portion of R_3 may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R_6 is a
 10 linker; AA is an amino acid; and n is 1 to 6; or a salt thereof.
2. The compound of claim 1, wherein n is 2 or 3.
3. The compound of claim 1 or 2 having the formula



- wherein R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, or heterocyclyl, or R_4 and
 15 R_5 together form a cycloalkyl or heterocyclyl.
4. The compound of claim 3, wherein R_1 is aralkyl, arylheterocyclylalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminocarbonyl, alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl, wherein the aryl portion is substituted or unsubstituted; R_2 is hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkyloxy,
 20 dicarboxyhaloalkyl, dicarboxyhaloalkyloxy, phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl portion of the substituents may be optionally substituted with a substituent selected from the

group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; and R_6 is a substituted or unsubstituted group having 1-6 carbon atoms.

5. The compound of any one of claims 1-4 having the formula:



6. The compound of any one of claims 1-5, wherein R_1 is aralkyl, arylheterocyclalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminocarbonyl, alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl, wherein the aryl portion is phenyl or naphthyl and the alkyl portion is a C₁-C₆ alkyl, and the heterocycl is a 3-7 membered ring having at least one of N, O, and S; R_2 is hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkyloxy, dicarboxyhaloalkyl, dicarboxyhaloalkyloxy, phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl or alkoxy portion of R_2 is a C₁-C₆ alkyl or alkoxy and may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R_3 is hydrogen, azido, amino, oxalylamino, carboxyalkyl, alkoxy carbonylalkyl, aminocarbonylalkyl, or alkylcarbonylamino; wherein the alkyl portion of R_3 is C₁-C₆ alkyl which may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocycl, or together form cycloalkyl or heterocycl, wherein the alkyl is a C₁-C₆ alkyl, the cycloalkyl is a C₃-C₇ cycloalkyl, and the heterocycl is a 3-7 membered ring with at least one of N, O, and S; and R_6 is a C₂-C₄ alkylenyl or alkenylenyl group, which may optionally substituted.
7. The compound of any one of claims 1-6, wherein R_1 is naphthylmethyl or indolyl.
8. The compound of any one of claims 1-7, wherein R_2 is carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkoxy, dicarboxyhaloalkyl, dicarboxyhaloalkoxy, phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl or alkoxy portion of R_2 is a C₁-C₆ alkyl or alkoxy and may be optionally


substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto.

9. The compound of any one of claims 1-7, wherein R₂ is carboxyalkoxy, dicarboxyalkyl, dicarboxyalkoxy, dicarboxyhaloalkyl, dicarboxyhaloalkoxy, phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl or alkoxy portion of R₂ is a C₁-C₆ alkyl or alkoxy and may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto.
10. The compound of any one of claims 1-9, wherein R₂ is phosphono, phosphonoalkyl, phosphonohaloalkyl, phosphonodihaloalkyl, or phosphoryl.
11. The compound of any one of claims 1-9, wherein R₂ is phosphono, phosphonomethyl, phosphonohalomethyl, or phosphonodihalomethyl.
12. The compound of any one of claims 1-11, wherein R₃ is carboxy C₁-C₆ alkyl or dicarboxy C₁-C₆ alkyl.
13. The compound of any one of claims 1-12, wherein R₃ is carboxymethyl or dicarboxymethyl.
14. The compound of any one of claims 1-11, wherein R₃ is alkoxycarbonyl C₁-C₆ alkyl, aminocarbonyl C₁-C₆ alkyl, amino, oxalylamino, or C₁-C₆ alkylcarbonylamino; wherein the alkyl portion of R₃ may be optionally substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto.
15. The compound of any one of claims 1-11 and 14, wherein R₃ is amino or C₁-C₆ alkylcarbonylamino.
16. The compound of claim 15, wherein R₃ is acetylamino.
17. The compound of claim 14, wherein R₃ is oxalylamino.
18. The compound of any one of claims 3-17, wherein R₄ and R₅, independently, are hydrogen, alkyl, or together form cycloalkyl, wherein the alkyl is a C₁-C₆ alkyl, and the cycloalkyl is a C₃-C₇ cycloalkyl.
19. The compound of any one of claims 3-17, wherein R₄ and R₅, independently, are hydrogen, alkyl, or together form cycloalkyl, wherein the alkyl is a C₁-C₆ alkyl, and the cycloalkyl is a C₆ cycloalkyl.
20. The compound of any one of claims 1-19, wherein R₆ is a C₂-C₃ alkylene or alkenylene

57


group, which may be optionally substituted.

21. The compound of any one of claims 1-19, which has the formula:



wherein R₄ and R₅ are independently C₁-C₆ alkyl or hydrogen.


5 22. The compound of any one of claims 1-20 which has the formula:



wherein R₄ and R₅ are independently C₁-C₆ alkyl or hydrogen.

23. The compound of claim 22, wherein R₄ and R₅ are methyl.

24. The compound of any one of claims 1-20 which has the formula:




10

wherein R₄ and R₅ are methyl or R₄ and R₅ together form cyclohexyl.


25. The compound of any one of claims 1-20 which has the formula:

58



wherein R_7 is aryl or alkenyl, and R_4 and R_5 are independently C_1 - C_6 alkyl or hydrogen.

26. The compound of any one of claims 1-20 which has the formula:



5 wherein R_8 is aryl alkyl and R_4 and R_5 are independently C_1 - C_6 alkyl or hydrogen.

27. The compound of claim 26, wherein R_8 is a benzyl or naphthylmethyl.


28. The compound of claim 1 or 2, wherein said amino acid (AA) is selected from the group consisting of glycine, alanine, valine, norvaline, leucine, iso-leucine, norleucine, α -amino n-decanoic acid, serine, homoserine, threonine, methionine, cysteine, S-acetylaminoethyl-

- 10 cysteine, proline, trans-3- and trans-4-hydroxyproline, phenylalanine, tyrosine, 4-aminophenylalanine, 4-nitrophenylalanine, 4-chlorophenylalanine, 4-carboxyphenylalanine, β -phenylserine β -hydroxyphenylalanine, phenylglycine, α -naphthylalanine, cyclohexylalanine, cyclohexylglycine, tryptophan, indoline-2-carboxylic acid, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, aspartic acid, asparagine, aminomalonic acid, 15 aminomalonic acid monoamide, glutamic acid, glutamine, histidine, arginine, lysine, N' -benzyl- N' -methyl-lysine, N' , N' -dibenzyl-lysine, 6-hydroxylysine, ornithine, α -aminocyclopentane carboxylic acid, α -aminocyclohexane carboxylic acid, α -aminocycloheptane carboxylic acid, α -(2-amino-2-norbornane)-carboxylic acid, α,γ -

diaminobutyric acid and α,β -diaminopropionic acid, homophenylalanine, and α -tert-butylglycine.

29. The compound of claim 1 or 2, wherein the phenylphosphate mimic group is phosphonomethyl, phosphono-(α -fluoro)methyl, phosphono-(α,α -difluoro)methyl,
 5 phosphono-(α -hydroxy)methyl, O-sulfo, and dicarboxymethoxy.

30. A compound of the formula:



wherein n is 0 to 15; Y' is a phenylalanyl radical having a phenyl ring, an amine end, and a carboxyl end, the phenyl ring having one or more substituents selected from the group
 10 consisting of hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkyloxy, dicarboxyalkyl, dicarboxyalkyloxy, dicarboxyhaloalkyl, dicarboxyhaloalkyloxy, and phosphonoalkyl, phosphonohaloalkyl, wherein the alkyl portion of the substituents may be unsubstituted or substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto and the amine end includes an azido group; W is
 15 a moiety attached to the nitrogen of Y' and is selected from the group consisting of alkylcarbonyl, oxalyl, alkylaminooxalyl, arylaminooxalyl, arylalkylaminooxalyl, alkoxyoxalyl, carboxyalkyl carbonyl, heterocyclyl carbonyl, heterocyclylalkyl carbonyl, arylalkyl heterocyclylalkyl carbonyl, aryloxycarbonyl, and arylalkoxycarbonyl, wherein the aryl and alkyl portions of the substituents may be unsubstituted or substituted with a
 20 substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; and the heterocyclyl portion of W contains at least 4 hetero atoms selected from the group consisting of O, N, and S; AA is an amino acid, the amine end of which is attached to the carboxyl end of Y'; and Z is an arylalkylamino or arylheterocyclyl alkylamino; or a salt thereof.

25 31. A pharmaceutical composition comprising a pharmaceutically or pharmacologically acceptable carrier and a compound of any one of claims 1-30.

32. A method for inhibiting an SH2 domain from binding with a phosphoprotein comprising contacting an SH2 domain with a compound of any one of claims 1-30.

33. The method of claim 32, wherein said SH2 domain is in a mammal, and said compound
 30 is administered to said mammal.

34. The use of a compound of any one of claims 1-30 in the manufacture of a medicament

for the treatment of a condition that responds to the inhibition of phosphoprotein binding to an SH2 domain of a mammal.

35. The use of a compound of any one of claims 1-30 in medicine.

36. A compound of any one of claims 1-30 for use as a Grb2-SH2 domain binding inhibitor.

5 37. A method for inhibiting SH2 domain binding comprising exposing a material having an SH2 domain to a compound of any one of claims 1-30.

38. A method for determining the presence of an SH2 domain in a material comprising:

(a) exposing a sample of said material to a SH2 binding compound and obtaining a first binding result; (b) exposing another sample of said material to a compound of any one of
10 claims 1-30 and obtaining a second binding result; and (c) comparing the first and second binding results to determine whether an SH2 domain is present in the material.

39. A method of preventing or treating a disease, state, or condition in a mammal comprising administering a compound of any one of claims 1-30.

40. A method of inhibiting cell motility or angiogenesis in a mammal comprising
15 administering to said mammal a macrocyclic peptide having cell signal inhibiting activity, cell motility, or cell angiogenesis inhibiting activity.


41. The method of claim 40, wherein said macrocyclic peptide is substantially free of cytotoxicity.

42. The method of claim 40 or 41, wherein said macrocyclic peptide is a Grb2-SH2 domain
20 mimetic peptide.

43. The method of any one of claims 40-42, wherein said cell motility or angiogenesis is induced by the hepatocyte growth factor (HGF).

44. The method of any of claims 40-43, wherein said cell motility is induced by the binding of c-Met receptor with the Grb2 protein.

25 45. The method of any one of claims 40-44, wherein the compound is of the formula (I):




(I)

wherein R_1 is a lipophile; R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; R_3 is hydrogen, azido, carboxyalkyl, alkoxy-carbonylalkyl, aminocarbonylalkyl, or alkylcarbonylamino, wherein the alkyl portion of R_3 may be optionally substituted with a substituent selected from the group
5 consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto; R_6 is a linker; AA is an amino acid; and n is 1 to 6; or a salt thereof.

46. A method for inhibiting the binding of an intracellular signal transducer to a receptor protein tyrosine kinase comprising contacting (a) a sample containing the receptor protein tyrosine kinase, (b) the intracellular signal transducer, and (c) a compound of any one of
10 claims 1-30 under conditions wherein, in the absence of the compound, the receptor protein tyrosine kinase binds to the intracellular transducer; wherein the contacting results in the inhibition of binding of the intracellular signal transducer to the receptor protein tyrosine kinase.


47. A method for detecting the inhibition of binding of an intracellular signal transducer to a
15 receptor protein tyrosine kinase comprising: (a) contacting a sample containing the receptor protein tyrosine kinase with the intracellular signal transducer, separately, in the presence and absence of the compound of any one of claims 1-30 under conditions that allow for binding of the receptor protein tyrosine kinase to the intracellular signal transducer in the absence of the compound; (b) determining that binding has occurred between the receptor
20 protein tyrosine kinase and the intracellular signal transducer; and (c) comparing the relative binding levels of the receptor protein tyrosine kinase to the intracellular signal transducer in the presence and absence of the compound.

48. A compound of the formula




25 wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group.

49. A compound of the formula




wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group.

50. A compound of the formula




wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group.

51. A compound of the formula




wherein R_1 is a lipophile.

52. A compound of the formula:




wherein R_1 is a lipophile and R is hydrogen or amine protective group.

53. A compound of the formula:



5 wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group.


54. A compound of formula 24:



24

10 wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.

55. A compound of formula 25:



25


15

wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate

64

mimic group or a protected phenylphosphate mimic; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.

56. A compound of formula 20



20

wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.


57. The compound of any one of claims 48-50 and 53-56, wherein said phenylphosphate

10 mimic group is phosphono or phosphonomethyl.

58. The compound of any one of claims 48-50 and 53-56, wherein said protected phenylphosphate mimic group is phosphonomethyl ester.

59. The compound of any one of claims 51-52 and 54-56, wherein said lipophile is aralkyl, arylheterocyclylalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminoacarbonyl, 15 alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl, wherein the aryl portion is phenyl or naphthyl and the alkyl portion is a C_1 - C_6 alkyl, and the heterocyclyl is a 3-7 membered ring having one or more of N, O, and S.


60. A method for preparing a compound of formula 3 comprising:



65


3

(a) providing a compound of formula 24




24

- 5 (b) reducing the azido group in the compound of formula 24 to an amino group to obtain a compound of formula 25



25

- (c) acetylating the amino group of the compound of formula 25 to obtain a compound of
10 formula 20:




20

and (d) carrying out an olefin metathesis reaction on the compound of formula 20, thereby obtaining the compound of formula 3; wherein R₁ is a lipophile; R₂, in combination with the

66

phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, or heterocyclyl, or R_4 and R_5 together form a cycloalkyl or heterocyclyl.


61. A method for preparing a compound of formula 24:



5


24

comprising treating a compound of formula 18 with a compound of formula 23:



18

10




23

wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.


62. A method for preparing a compound of formula 18 comprising:

67




18

- 5 (a) treating a compound of formula 7




7

with a compound of the formula:




- 10 to obtain a compound of the formula 16




16

- (b) treating the compound of formula 16 with an amine protected amino acid of the formula

68



to obtain a compound of formula 17




5

17

and (c) removing the amine protecting group to obtain the compound of formula 18;
wherein R_1 is a lipophile; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl,
heterocyclyl, or together form cycloalkyl or heterocyclyl.


63. A method for preparing a compound of the formula 7 comprising:

10



7


(a) treating a compound of the formula




15

with a compound of the formula

69




to obtain a compound of formula 12



5


12

(b) replacing a hydrogen on the CH₂ adjacent to the carbonyl carbon (1) of the compound of formula 12 with a lipophile to obtain a compound of formula 13:



13

10 (c) reducing the compound of formula 13 to obtain the compound of formula 14




14

and (d) replacing the hydroxyl group of the compound of formula 14 with an amino group, thereby obtaining the compound of formula 7; wherein R₁ is a lipophile and M is a metal.


15 64. A method for preparing a compound of formula 23 comprising:

70



23


(a) providing a compound of formula 8



8

5


(b) treating the compound of formula 8 with a base and trisyl azide to obtain a compound of formula 22



22

10 and (c) treating the compound of formula 22 with an alkaline peroxide, thereby obtaining the compound of formula 23; wherein R₂, in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group.

65. A method for preparing a compound of formula 8 comprising:




8

15

71

(a) providing a compound of formula 9




9

and (b) treating the compound of formula 9 with vinyl magnesium bromide and PhSCu;


5 wherein R₂, in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group.

66. A method for preparing a compound of formula 9 comprising




9

10 treating a compound of formula 10



10

with a compound of formula 11




11

15 wherein R₂, in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group, and X is a halogen.


72

67. A method for preparing a compound of formula 10 comprising:




10

treating a compound of the formula




5

with a compound of the formula



68. A method for preparing a compound of formula 4 comprising:




4

10


(a) treating the compound of formula 18

73



18


with a compound of the formula 6



5


6

to obtain a compound of formula 5



5

(b) treating the compound of formula 5 with a Grubbs catalyst to compound of formula 19



10

74

19

and (c) treating the compound of formula 19 with a mixture of trifluoroacetic acid, water, and trimethylsilane to obtain the compound of formula 4; wherein R_1 is a lipophile, R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected

5 phenylphosphate mimic group; and R_4 and R_5 , independently, are hydrogen, alkyl, cycloalkyl, heterocyclyl, or together form cycloalkyl or heterocyclyl.

69. The method of any one of claims 60-61, 64-66, and 68, wherein said phenylphosphate mimic group is hydroxyl, carboxyl, formyl, carboxyalkyl, carboxyalkoxy, dicarboxyalkyl, dicarboxyalkyloxy, dicarboxyhaloalkyl, dicarboxyhaloalkyloxy, phosphono,

10 phosphonoalkyl, phosphonohaloalkyl, phosphoryl, phosphorylalkyl, or phosphorylalkoxy, wherein the alkyl portion of the substituents may be unsubstituted or substituted with a substituent selected from the group consisting of halo, hydroxy, carboxyl, amino, aminoalkyl, alkyl, alkoxy, and keto.

70. The method of any one of claims 60-61, 64-66, and 68, wherein said phenylphosphate 15 mimic group is phosphono or phosphonomethyl.

71. The method of any one of claims 60-61, 64-66, and 68, wherein said protected phenylphosphate mimic group is phosphonomethyl ester.

72. The method of any one of claims 60-63 and 68, wherein said lipophile is aralkyl, arylheterocyclylalkyl, alkylaminocarbonyl, alkenylaminocarbonyl, arylaminoacrabonyl, 20 alkoxyalkyl, aryloxyalkyl, or aralkoxyalkyl, wherein the aryl portion is substituted or unsubstituted.


73. The method of any one of claims 60-63 and 68, wherein said lipophile is naphthylmethyl.

74. The method of claim 46 or 47, wherein said method is carried out *in vitro*.

25 75. The method of claim 46 or 47, wherein said method is carried out *in vivo*.

76. The method of claim 39, wherein the disease, state, or condition is cancer.

77. A compound of the formula 30




75

30

wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group, and R is aralkyl, aryl, or alkyl.

78. The compound of claim 77, wherein R is benzyl.


5 79. A method for preparing a compound of the formula 31



31


wherein R_2 , in combination with the phenyl ring, is a phenylphosphate mimic group or a protected phenylphosphate mimic group and R is aralkyl, aryl, or alkyl comprising: (a)

10 treating a compound of the formula 26



26


with a compound of the formula 27



27

15


to obtain a compound of the formula 28



28


76

(b) treating the compound of formula 28 with a compound of formula 11




11

to obtain a compound of formula 29



29

(c) reducing the compound of formula 29 to obtain a compound of formula 30



30

(d) treating the compound of formula 30 with an alkali metal salt of bis(trimethylsilyl)amide and trisyl azide, to obtain the compound of formula 31.

80. The method of claim 79, wherein R is benzyl.

1/18




FIG. 1

2/18




FIG. 2

3/18




FIG. 3

4/18




FIG. 4

5/18




FIG. 5

6/18




FIG. 6

7/18




FIG. 7

8/18




FIG. 8

9/18




FIG. 9

10/18




FIG. 10

11/18





FIG. 11

12/18



13/18







FIG. 13

FIG. 14a


1i

FIG. 14b


3a

FIG. 14c

5b

FIG. 14d

1h

FIG. 14e

Compd 126

FIG. 15




FIG. 16a




FIG. 16b





FIG. 17






FIG. 18a

FIG. 18b

FIG. 18c