



(51) International Patent Classification:

C07C 319/18 (2006.01) C07D 277/04 (2006.01)

C07K 1/107 (2006.01) C07C 233/05 (2006.01)

C07K 1/113 (2006.01)

(21) International Application Number:

PCT/SG2012/000168

(22) International Filing Date:

14 May 2012 (14.05.2012)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

201103480-8

13 May 2011 (13.05.2011)

SG

(71) Applicant (for all designated States except US): NAN-

YANG TECHNOLOGICAL UNIVERSITY [SG/SG];

50 Nanyang Avenue, Singapore 639798 (SG).

(72) Inventors; and

(75) Inventors/Applicants (for US only): LIU, Chuan-Fa

[US/SG]; Nanyang Technological University, SBS 03S-85,

50 Nanyang Avenue, Singapore 639798 (SG). LI, Fupeng

[CN/SG]; Nanyang Technological University, SBS 03S-85,

50 Nanyang Avenue, Singapore 639798 (SG).

(74) Agent: Ella Cheong Spruson & Ferguson (Singapore)

Pte Ltd; Robinson Road Post Office, P.O. Box 1531,

Singapore 903031 (SG).

(81) Designated States (unless otherwise indicated, for every

kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every


kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:


— with international search report (Art. 21(3))

— before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments (Rule 48.2(h))


(54) Title: METHOD FOR MODIFICATION OF ORGANIC MOLECULES



(I)



(II)



(57) Abstract: The present invention is directed to a method of alkylating a thiol group (R-S-H) or seleno group (R-Se-H) in a target molecule wherein the method comprises: reacting a target molecule comprising at least one thiol group with a compound of formula (I) or (II); wherein R is an acetyl group or any other acyl group or is a group comprising any one of: or wherein R in formula (II) can also be an alkyl group; and wherein R' is selected from a group consisting of a hydrogen, a methyl group and an ethyl group.

METHOD FOR MODIFICATION OF ORGANIC MOLECULES**FIELD OF INVENTION**

5 The present invention relates to a method for alkylation of organic molecules.

BACKGROUND

10 Post-translational modification (PTM) is a fundamental mechanism for modulating protein function. One such PTM with increasingly recognized significance is protein lysine (Lys) acetylation. Like Tyrosine (Tyr) /Serine (Ser) / Threonine (Thr) phosphorylation, Lys acetylation is a
15 reversible biochemical process, where a lysine acetyltransferase adds an acetyl group onto the ϵ -amine of a lysine residue. A deacetylase acts in the opposite way to remove it. Initially discovered in histones, lysine acetylation has also been observed recently in a very large
20 number of other proteins, suggesting its diverse regulatory functions in the cell. There is mounting evidence that aberrant lysine acetylation is implicated in many disease conditions such as cancer and neurological disorders. Therefore, the study of lysine acetylation biology is of
25 great importance and will lead to continuous therapeutic innovations.

Although many years of intensive research has firmly established a broad role for lysine acetylation as a
30 histone epigenetic mark affecting chromatin structure and function, the effects of most individual acetylation events

- especially those identified more recently - remain to be elucidated. Recent research in genetics, cell biology and especially proteomics has identified lysine acetylation in a large number of non-histone proteins. The functions of most of these modifications are unknown. As a major limiting factor, the study of protein lysine acetylation is often hindered by a lack of homogeneous protein samples containing the acetylated lysine ("Lys-Ac") residue(s) of interest. Such homogenously acetylated proteins would be invaluable reagents for discerning the structural and functional effects of a particular Lys-Ac PTM via biophysical and biochemical means.

Several methods are useful to prepare site-specific modified proteins, such as unnatural amino-acid mutagenesis using the amber stop codon/suppressor tRNA pair, and protein chemical synthesis, but there remain significant technical barriers for the wide use of these methods by the bioscience research community at large. However, unnatural amino acid mutagenesis is not widely available because it is a proprietary technology; while protein chemical synthesis requires extensive expertise of a well trained chemist and is technically very challenging as well as labor-intensive.

Another known method is based on the combined use of unnatural amino acid mutagenesis and chemical modification to generate an acetyl-lysine analog but the optical purity is lost on the modified amino acid. Enzymatic Lys acetylation of recombinant proteins would appear to be an attractive approach. However, given the promiscuity of lysine acetyltransferases and the incomplete nature of

enzymatic reactions, it is difficult if not impossible to isolate or characterize the desired acetylation product for structural and functional investigations.


5 As for other enzymatic reactions that do offer the necessary specificity, they often must work in the context of large macromolecular complexes such as with the help of other adaptor molecules, which renders them of little practical value.

10

A chemical approach for selective installation of acetylated Lys residues in recombinant proteins is therefore highly desired. Evidently, with the presence of many possible lysine residues in a normal protein, direct
15 acetylation of a particular lysine residue is chemically not feasible.

The unique reactivity of the thiol group of cysteine (Cys) as a soft nucleophile has been exploited extensively for
20 selective protein modification. Previously, a chemical method was developed to install a close isosteric analog of N-methyl-lysine into recombinant proteins (See Scheme I below). The method is based on a traditional alkylation reaction of the thiol group of a Cys residue with
25 aminoethyl bromide or chloride, yielding aminoethylcysteine which is known to be a lysine equivalent.


When the alkylating agent is changed to N-methyaminoethyl halide, an N-methylated aminoethylcysteine residue or N-
30 methyl thiaLys ("thiaLys(Me)") is generated, which has been shown to be functionally similar to the natural Lys(Me).

Scheme I

5

As such, similar strategies have been proposed to prepare a close mimic of the native Lys(Ac) residue by making use of the unique reactivity of the cysteine thiol group. However, attempts to alkylate the Cys thiol with a similar N-acetyl-aminoethyl bromide and N-acetyl-aziridine failed to give the desired product (See Scheme IIa).

Further efforts in the prior art led to the development of methylthiocarbonyl-aziridine as the alkylating agent to afford methylthiocarbonyl-thiaLys as an N-acetyl-Lys mimic (See Scheme IIb). Although the alkylation reaction was successful and the resultant methylthiocarbonyl-thiaLys was shown to be recognized by a bromodomain-containing protein and by anti-acetyl-Lys antibodies, the presence of a large sulfur atom between the carbonyl and methyl in the thiocarbamate moiety makes it only a distant analog of the acetamide part in Lys(Ac). From both steric and electrochemical viewpoints, there are obvious and considerable differences between the acetyl and methylthiocarbonyl group (see structure in Scheme IIb), which may explain why the thiocarbamate modification is resistant to histone deacetylase cleavage.

Scheme II



Clearly, analogous to 2-aminoethyl-cysteine (i.e., thialysine) and N-methyl-thialysine being ideal mimics of lysine and N-methyl-lysine respectively, the best mimic of Lys(Ac) would still be N-acetyl-4-thialysine or sLys(Ac) in which the thioether linkage is a close isosteric replacement of the γ -methylene group in natural Lys(Ac). As the position of this substitution is rather far away - by 2 carbon atoms - from the acetamide nitrogen, little differences are expected between this Lys(Ac) mimic and its natural counterpart in their exhibited physicochemical and biochemical properties. Unfortunately, current attempts to use a similar alkylation reaction with N-acetyl-aminoethyl bromide or iodide and N-acetyl-aziridine are unsuccessful at producing acetyl-thialysine with acceptable yields and selectivity.

Furthermore, nucleophilic substitution with an alkyl halide or the equivalent arizidine compound (Scheme IIa) cannot be used to selectively alkylate the thiol under conditions that are acceptable for a protein, so a different
5 alkylation method must be provided.


Accordingly, there is a need to provide a method for alkylating the thiol group of Cys to obtain a Lys(Ac) mimic that overcomes or ameliorates one or more disadvantages
10 disclosed above. Additionally, such a method will also be useful for installing other modifications (e.g., pegylation and ubiquitination) onto a thiol-containing compound such as a peptide or a protein.

SUMMARY

In a first aspect, there is provided a method of alkylating a thiol group (R-S-H) or seleno group (R-Se-H) in a target
20 molecule wherein the method comprises: reacting a target molecule comprising at least one thiol group with a compound of formula (I) or (II):



wherein R is an acetyl group or any other acyl group or is a group comprising any one of:



wherein R in formula (II) can also be an alkyl group; and
 wherein R' is selected from a group consisting of a
 5 hydrogen, a methyl group and an ethyl group.

In one embodiment, the target molecule is an organic compound. In one embodiment, the organic compound is a peptide or a protein molecule. In still another embodiment,
 10 the peptide is selected from an oligopeptide, a polypeptide or a synthetic peptide. In yet another embodiment, the protein molecule is selected from the group consisting of a recombinant protein and a protein complex. One important
 15 aspect of the disclosed method resides in its ability to be performed on protein molecules under conditions which would not denature the protein or compromise its structural integrity.

In one embodiment, the disclosed method advantageously
 20 provides a simple, one-step process for modifying one or more amino acid residues present on a target molecule, such as biotin, histone or ubiquitin. In one embodiment, the disclosed method is capable of modifying specific amino acid residues to form ideal mimics of acetylated lysine. In
 25 one embodiment, the disclosed method selectively modifies one or more cysteine residues on the target molecule to form ideal mimics of acetylated lysine. More particularly,

the disclosed method is directed to the alkylation of one or more cysteine residues via reaction with an appropriate alkylation agent to form acetylated lysine analogs. The type of acetylated lysine analog formed depends on the
5 specific alkylating agent used in the reaction.

In one embodiment, the alkylation agent is a compound of formula (I) or (II) as described above. Advantageously, the inventors have found that the alkylation agent is
10 capable of reacting with the thiol group present on the cysteine residue to form N-acetyl-4-thialysine, in which the thioether linkage is a close isosteric replacement of the 4-methylene group in natural acetylated lysine. As the position of this substitution is sterically distant from
15 the acetamide nitrogen (by two carbon atoms), little differences are detected between the mimic and the natural acetylated lysine in physicochemical and biochemical properties.

Advantageously, the disclosed method overcomes technical
20 problems of feasibility, low yield and low selectivity which plague the prior art methods. In contrast, the disclosed method is very robust and provides near 100% yield of the Lys-Ac mimics (e.g., N-acetyl-4-thialysine) in
25 considerably short reaction times of from 30 minutes to 2 hours or lesser.

In another aspect, there is provided the use of a compound of Formula (I) as defined above as an alkylating agent for
30 alkylation of a target molecule.

In yet another aspect, there is provided the use of a compound of Formula (II) as defined above as an alkylating agent for alkylation of a target molecule comprising a thiol-group (R-S-H) or a seleno group (R-Se-H).

5

In still another aspect, there is provided the use of a compound of Formula (I) for installing an acetylated lysine residue analog in a target molecule.

- 10 In another aspect, there is provided a pharmaceutical composition comprising a target molecule modified according to the method as defined above.

Definitions

15

The following words and terms used herein shall have the meaning indicated:

- 20 The word "substantially" does not exclude "completely" e.g. a composition which is "substantially free" from Y may be completely free from Y. Where necessary, the word "substantially" may be omitted from the definition of the invention.

- 25 Unless specified otherwise, the terms "comprising" and "comprise", and grammatical variants thereof, are intended to represent "open" or "inclusive" language such that they include recited elements but also permit inclusion of additional, unrecited elements.

30

As used herein, the term "about", in the context of concentrations of components of the formulations, typically

means +/- 5% of the stated value, more typically +/- 4% of the stated value, more typically +/- 3% of the stated value, more typically, +/- 2% of the stated value, even more typically +/- 1% of the stated value, and even more typically +/- 0.5% of the stated value.

Throughout this disclosure, certain embodiments may be disclosed in a range format. It should be understood that the description in range format is merely for convenience and brevity and should not be construed as an inflexible limitation on the scope of the disclosed ranges. Accordingly, the description of a range should be considered to have specifically disclosed all the possible sub-ranges as well as individual numerical values within that range. For example, description of a range such as from 1 to 6 should be considered to have specifically disclosed sub-ranges such as from 1 to 3, from 1 to 4, from 1 to 5, from 2 to 4, from 2 to 6, from 3 to 6 etc., as well as individual numbers within that range, for example, 1, 2, 3, 4, 5, and 6. This applies regardless of the breadth of the range.

Disclosure on Optional Embodiments

Exemplary, non-limiting embodiments of the method according to the first aspect will now be disclosed.

In one embodiment, the disclosed method comprises alkylating a thiol group of a target molecule, wherein the thiol group is the thiol group of a cysteine and/ or a selenocysteine residue. In one embodiment, the thiol group is the thiol group of a cysteine residue.

In one embodiment, the R group of the compound of formula (I) or (II) may be an acyl group, wherein the acyl group is an aliphatic acyl group, or an alicyclic acyl group, or an aromatic acyl group, or an amino acyl or a peptidyl group
5 or a proteiny group.

In one embodiment, the R group of formula (I) or (II) is an aliphatic acyl group, and which is selected from the group consisting of a linear aliphatic acyl group, a branched aliphatic acyl group, a tert-butyloxycarbonyl group, a
10 derivative of tert-butyloxycarbonyl group, a benzyloxycarbonyl group and a derivative of benzyloxycarbonyl group.

In one embodiment, the acyl group is selected from the group consisting of a linear aliphatic acyl groups having 1
15 to 20 or 1 to 7 carbon atoms; a branched aliphatic acyl groups having 1 to 7 carbon atoms; and a poly(ethylene glycol) moiety.

In another embodiment, the acyl group is selected from the group consisting of formyl group, acetyl group, propionyl
20 group, 2-propionyl group, butyryl group, isobutyryl group, pentanoyl group, hexanoyl group, allylcarbonyl group, cyclohexylmethylcarbonyl group, and C₃-C₆ cycloalkylcarbonyl groups, such as cyclopropylcarbonyl group, cyclopentylcarbonyl group, cyclohexylcarbonyl group, and 1
25 cyclohexenylcarbonyl group.

In yet another embodiment, the R group of formula (I) or (II) is an aromatic acyl group, and which is selected from the group consisting of benzoyl group, 4-methylbenzoyl group, and 4 methoxybenzoyl group.

In another embodiment, the R group in Formula (II) is an alkyl group, wherein the alkyl group is selected from the group consisting of methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, sec-butyl, pentyl, hexyl, octyl, nonyl, decyl, undecyl, dodecyl, trideyl, tetradecyl, pentadecyl, hexadecyl, octadecyl, poly(ethylene.glycol)-ethyl and poly(ethylene.glycol)-propyl. Exemplary alkyl groups may include, but are not limited to, $\text{HO}(\text{CH}_2\text{CH}_2\text{O})_n\text{-CH}_2\text{CH}_2\text{-}$; $\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_n\text{-CH}_2\text{CH}_2\text{-}$; and $\text{CH}_3\text{O}(\text{CH}_2\text{CH}_2\text{O})_n\text{-CH}_2\text{CH}_2\text{CH}_2\text{-}$.

- 10 In one embodiment, the alkylating agent is a compound of formula (I), where R is an acetyl group and R' is hydrogen and n is 0 (i.e., N-vinyl acetamide or "NVA").

In yet another embodiment, the alkylating agent is a compound of formula (I), wherein R is propionyl ($\text{CH}_3\text{CH}_2\text{C}(\text{O})\text{-}$), R' is hydrogen and n is 0 (i.e., the alkylating agent is N-vinylpropionamide).

In still another embodiment, the alkylating agent is a compound of formula (I), wherein R is butyryl ($\text{CH}_3\text{CH}_2\text{CH}_2\text{C}(\text{O})\text{-}$), R' is hydrogen and n is 0 (i.e., the alkylating agent is N-vinylbutyramide)

In another embodiment, the alkylating agent is a compound of formula (II), where R is $\text{CH}_3\text{C}(\text{O})\text{-}$, i.e., vinyl acetate.


- 25 In another embodiment, the alkylating agent is a compound of formula (I), wherein R is $\text{CH}_3\text{C}(\text{O})\text{-}$, R' is H and n is 1, i.e., N-allylacetamide.

In still another embodiment, the alkylating agent is a compound of formula (I), wherein R is $\text{CH}_3\text{C}(\text{O})-$, R' is CH_3 and n is 0, i.e., N-methyl-N-vinylacetamide.

- 5 In yet another embodiment, the alkylating agent is a compound of formula (I), where the R is an acyl group selected from a peptidyl group or a proteiny group, R' is H and n is 0.
- 10 In one embodiment, the alkylating reaction between NVA and a thiol group may be represented by Scheme III as shown below:


Scheme III

15



- As can be seen from Scheme III, the thiol-ene coupling between the cysteine thiol and N-vinylacetamide directly generates the desired acetyl-thialysine in a single step reaction.
- 20

- In one embodiment, the reaction mechanism for the reaction in Scheme III may be described in three general steps as shown in Scheme IV below:
- 25

Scheme IV

5 The disclosed method may further comprise a step of irradiating the reaction mixture with ultra-violet (UV) radiation. The irradiation step may further comprise the addition of photoinitiators into the reaction mixture. The photoinitiators may be a water soluble photoinitiator. The

10 photoinitiator may be an azo-type initiator. In one embodiment, the photoinitiator is the compound 2,2'-Azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride ("VA-044"). In another embodiment, the photoinitiator is the compound lithium phenyl-2,4,6-trimethylbenzoylphosphinate

15 ("LAP").

The disclosed method may further comprise performing the alkylation reaction in the presence of an additional thiol compound. Advantageously, the additional thiol compound may

20 serve to suppress the undesirable side reaction of radical chain telomerization. In one embodiment, the additional thiol compound may be selected for its ability to react

with the radical intermediate formed by the addition of the thiyl radical to the NVA ethylene double bond to prevent the intermediate from reacting with another NVA molecule. In one embodiment, the additional thiol compound is
5 glutathione. In one embodiment, the glutathione is used in its reduced form.

In one embodiment, the alkylation reaction may be performed in acetate buffer at pH 4 to 7 and in the presence of VA-
10 044 as the initiator under UV irradiation at 365 nm.

In one embodiment, the molar ratio of the additional thiol compound to the NVA is about 3:10. In one embodiment, the molar ratio of the additional compound to the NVA to the
15 photoinitiator is about 3:10:1.

In another embodiment, the molar ratio of the additional thiol compound to the NVA may be selected from the group consisting of 1:10, 2:10, 3:10, 4:10, 5:10, 6:10, 7:10,
20 8:10, 9:10 and 1:1.

The disclosed molar ratios may also apply to embodiments of the present invention where the alkylating agent is not NVA.
25

Brief Description Of Drawings

The accompanying drawings illustrate a disclosed embodiment and serves to explain the principles of the disclosed
30 embodiment. It is to be understood, however, that the

drawings are designed for purposes of illustration only, and not as a definition of the limits of the invention.

Fig. 1A is a C18 analytical high pressure liquid chromatogram ("HPLC") monitoring of the thiol-ene coupling reaction between benzyl mercaptan (BM) and N-vinyl-acetamide ("NVA") at pH 4.0, wherein peak a corresponds to BM; Peak b corresponds to product; Peak c corresponds to side product 1, i.e. NVA dimer attached to BM; and Peak d corresponds to side product 2, NVA trimer attached to BM.

Fig. 1B shows an electrospray ionization - mass spectrometer ("ESI-MS") spectrum of peak b of Fig. 1A.

Fig. 1C shows an ESI-MS spectrum of peak c of Fig. 1A.

Fig. 1D shows an ESI-MS spectrum of peak d of Fig. 1A.

Fig. 2 shows a ^1H NMR spectra of the reaction product of the thiol-ene reaction of Fig. 1.

Fig. 3 shows a ^1H NMR spectra of the side product of the thiol-ene reaction of Fig. 1.

Fig. 4A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and NVA at pH 4.0. Peak a: peptide 1; Peak b: product.

Fig. 4B shows an ESI-MS spectrum of peak a.

Fig. 4C shows an ESI-MS spectrum of peak b.

Fig. 5A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and NVA at pH 6.0. Peak a: product; Peak b: disulfide-linked side product between peptide 1 and glutathione.

5

Fig. 5B shows an ESI-MS spectrum of peak a.

Fig. 5C shows an ESI-MS spectrum of peak b.

10 Fig. 6A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 2 and NVA at pH 4.0. Peak a: peptide 2; Peak b: product.

Fig. 6B shows an ESI-MS spectrum of peak a.

15

Fig. 6C shows an ESI-MS spectrum of peak b.

Fig. 7A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 3 and NVA at pH 4.0.

20 Peak a: peptide 3; Peak b: expected product; Peak c: oxidation product.

Fig. 7B shows an ESI-MS spectrum of peak a.

25 Fig. 7C shows an ESI-MS spectrum of the peak b.

Fig. 7D shows an ESI-MS spectrum of the peak c.

30 Fig. 8A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 4 and NVA at pH 4.0. Peak a: peptide 4; Peak b: product.

Fig. 8B shows a matrix assisted laser desorption/ionization - time of flight mass spectrometer ("MALDI-TOF MS") spectrum of peak a.

5 Fig. 8C shows a MALDI-TOF MS spectrum of peak b.

Fig. 9A shows a MALDI-TOF MS spectrum of the thiol-ene coupling reaction between peptide substrate 4 and NVA under non-standard conditions, i.e., when no glutathione was
10 added (other conditions were the same).

Fig. 9B shows a MALDI-TOF MS spectrum of the product if 15 mM tris(2-carboxyethyl)phosphine ("TCEP") was added to replace glutathione.

15

Fig 10A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide substrate 5 and NVA at pH4.0; Peak a: peptide 5; Peak b: expected product; Peak
c: oxidation product.

20

Fig. 10B shows a MALDI-TOF MS spectrum of peak a.

Fig. 10C shows a MALDI-TOF MS spectrum of peak b.

25 Fig. 10D shows a MALDI-TOF MS spectrum of peak c.

Fig. 11A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between ubiquitin K48C and NVA at pH 4.0. Peak a: ubiquitin K48C; Peak b: product.

30

Fig. 11B shows a MALDI-TOF MS spectrum of peak a.

Fig. 11C shows a MALDI-TOF MS spectrum of peak b.

FIG. 12A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between ubiquitin K48C and NVA at pH
5 7.0.

Fig. 12B shows the ESI-MS spectrum of the raw and deconvoluted mass of product.

10 FIG. 13A shows a C4 semi-prep HPLC profile of the thiol-ene coupling reaction between H4 K16C and NVA at pH 4.0 with or without dimethyl sulfide. Peak a: H4 K16C; Peak b: expected product; Peak c: oxidation product.

15 Fig. 13B is an ESI-MS spectrum of the raw and deconvoluted mass of the starting material H4 K16C (shown in peak a).

Fig. 13C is an ESI-MS spectrum of the raw and deconvoluted mass of peak b.

20

Fig. 13D is a MALDI-TOF MS spectrum of peak c.

FIG. 14A is a C4 semi-prep HPLC profile of the thiol-ene coupling reaction between H4 K16C and NVA at pH 7.0. Peak
25 a: expected product; Peak b: oxidation product.

Fig. 14B is an ESI-MS spectrum of the raw and deconvoluted mass of peak a.

30 Fig. 15A is C4 semi-prep HPLC profile of the thiol-ene coupling reaction between H3 K27C and NVA at pH 4.0. Peak a: H3 K27C; Peak b: product.

Fig. 15B is an ESI-MS spectrum of the raw and deconvoluted mass of peak a.

- 5 Fig. 15C is an ESI-MS spectrum of the raw and deconvoluted mass of peak b.

Fig. 16 shows the results of SIRT2 mediated deacetylation assay.

10

Fig. 16A is a C18 analytical HPLC profile of peptide 1 with the installed acetyl-lysine analog in SIRT2 assay. Peak a: deacetylated product; Peak b: Ac-FQPKKS(Ac)G-NH₂.

- 15 Fig. 16B is a C18 analytical HPLC profile of the native K(Ac)-peptide substrate in SIRT2 assay. Peak c: deacetylated product; Peak d: native peptide substrate.

Fig. 16C is an ESI-MS spectrum of peak a.

20

Fig. 16D is an ESI-MS spectrum of peak b.

Fig. 16E is an ESI-MS spectrum of peak c.

- 25 Fig. 16F is an ESI-MS spectrum of peak d.

Fig. 17 is an ESI-MS spectrum of the raw and deconvoluted mass of the alkylated product H4 K_s16.

- 30 Fig. 18A show Western blots with anti-H4 K16Ac Ab on the H4 proteins: H4 K16C and H4 K_s16Ac.

Fig. 18B shows a deacetylation assay of Ac-FQPKKs(Ac)G by SIRT2 after 16 h.

Fig. 18C shows the effects of H4 K16 acetylation on
5 nucleosome array folding as seen from sedimentation distributions of the nucleosome arrays before (no Mg^{2+} , open symbols) and after Mg^{2+} -induced folding (with 1.0 mM $MgCl_2$, solid symbols).

10 Fig 19A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-vinylpropionamide. Peak a: product; Peak b: peptide 1.

Fig. 19B shows an ESI-MS of peak a.

15

Fig. 20A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-vinylbutyramide. Peak a: product; Peak b: peptide 1; Peak *: unidentified peak.

20

Fig. 20B show an ESI-MS of peak a.

Fig. 21A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-methyl-N-
25 vinylacetamide. Peak a: product; Peak b: peptide 1.

Fig. 21B shows an ESI-MS of peak a.

Fig. 22A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and vinyl acetate.
30 Peak a: product; Peak b: peptide 1; Peak *: unidentified peak.

Fig. 22B shows an ESI-MS of peak a.

Fig. 23A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-allylacetamide. Peak a: product; Peak b: peptide 1.

Fig. 23B shows an ESI-MS of peak a.

Fig. 24 shows an ESI-MS of the isolated product formed from the thiol-ene coupling reaction between **RLYRAG**-vinyl amide and methyl mercaptoacetate.

Examples

Non-limiting examples of the invention will be further described in greater detail by reference to specific Examples, which should not be construed as in any way limiting the scope of the invention.

In the following methods, the amino acids and coupling agents used were purchased from GL Biochem (Shanghai, China) and Novabiochem (Germany). The human sirtuin 2 (SIRT2) enzyme, NAD⁺ and assay buffer were purchased from Enzo Life Sciences (New York, United States of America). The Histone H4 K16Ac antibody was from Active Motif (California, United States of America). All other chemical reagents were purchased from commercial suppliers.

Analytical and semi-prep HPLC were performed on a Shimadzu HPLC system equipped with a SPD-M20A prominence diode array detector. A C18 reverse-phase column (Jupiter 5 μ m 300 A,

250*4.6 mm) was used for analytical HPLC. The C18 reversed-phase column (Jupiter 5 μ m 300 A, 250*10 mm) and C4 reversed-phase column (Vydac 5 μ m 300 A, 250 * 10 mm) were used for semi-prep HPLC. The analytes were eluted using a gradient mixture of two solvents: solvent A was deionized water containing 0.05% trifluoroacetic acid (TFA) and solvent B was 90% acetonitrile (ACN) in deionized water containing 0.05% TFA. The mobile phase flow rate was 1 mL/min for analytical HPLC and 2.5 mL/min for semi-prep HPLC.

Peptide or protein masses were measured using a Thermo FINNIGAN LCQ Deca XP MAX equipped with electrospray ionization (ESI) ion source or a 4800 MALDI TOF/TOF Analyzer using α -cyano-4-hydroxycinnamic acid as the matrix.

Example 1

Model Study of the Thiol-ene Coupling Reaction Between NVA and Benzyl Mercaptan

To demonstrate the thiol-ene coupling mechanism, different concentrations of N-vinyl-acetamide (NVA) and benzyl mercaptan (BM) were mixed in 0.2 M acetate buffer (at pH 4.0, prepared from acetic acid and sodium acetate). Thereafter, 5 mM of initiator VA-044 (2,2'-Azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride) was added in a 0.5 ml thin wall, clear tube (Axygen, Inc., San Francisco, United States of America). The tube containing the reaction mixture was placed in a Cole Parmer 9818-series darkroom UV light box at about 10 cm under the lamp (365 nm) at room temperature for 30 min to obtain the reaction product, a small organic thiol compound benzyl mercaptan (BzSH).

The reaction was monitored by C18 analytical HPLC and the results are shown in Fig. 1A. The results were also confirmed by electrospray ionization mass spectrometry (ESI-MS) and shown in Figs. 1B, 1C and 1D. The HPLC conditions were 0% to 50% of buffer B in buffer A in 50 min.

As seen in Fig. 1A(i), when NVA and BM were at a concentration of 1 mM respectively, only a BM peak (denoted as peak a) was detected. No BzSH peak (denoted as peak b) was detected. In Figs. 1A(ii) and 1A(iii), when NVA and BM were at a concentration of 5 mM and 10 mM respectively, the conversion to BzSH did not improve with prolonged irradiation time, as evidenced by the presence of peak a. As seen in Figs. 1A(iv), 1A(v) and 1A(vi) where NVA was added in an excess amount as compared to BM, complete conversion to BzSH were obtained, as evidenced by the absence of the BM peak a.

However, in Figs 1A(iv), 1A(v) and 1A(vi), additional peaks c and d were observed. The presence of peak c is due to a side product of an NVA dimer attached to BM and the presence of peak d is due to a side product of an NVA trimer attached to BM.

Peak b was further characterized by ESI-MS. In Fig. 1B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of BzSH is shown. The $[M+H]^+$ found had a m/z value of 210.02 and molecular weight of 209.09.

Peak c was further characterized by ESI-MS. In Fig. 1C, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the NVA dimer side product is shown. The $[M+H]^+$ found had a m/z value of 295.08 and molecular weight of 294.14.

Peak d was further characterized by ESI-MS. In Fig. 1D, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the NVA trimer side product is shown. The $[M+H]^+$ found had a m/z value of 380.12 and molecular weight of 379.19.

The molecular structure and the 1H NMR spectra of the reaction product, BzSH, (in deuterated chloroform ($CDCl_3$) solvent) is shown in Fig. 2. The molecular structure and the 1H NMR spectra of the NVA dimer side product (in $CDCl_3$ solvent) is shown in Fig. 3.

This example demonstrated the mechanism of the classic radical thiol-ene addition reaction and helps to explain the formation of the side products. The carbon radical formed at step 2, Scheme IV, is usually much more likely to react with a thiol molecule in the critical rate-limiting step of radical chain transfer (step 3, Scheme IV) which generates another thiyl radical. However, in the presence of a large excess of the -ene compound (NVA), the carbon radical can also sometimes react with another NVA molecule in a phenomenon called telomerization (See Scheme IV above). Therefore, addition of glutathione is capable of suppressing this side reaction as it can act as a radical chain transferring agent and participate in step 3, Scheme IV.

Examples 2-6 and Comparative Examples***Synthesis of Peptide Substrates for Alkylation***

To obtain Cys-containing peptides, five model peptides were
5 synthesized as C-terminal carboxyamides using standard
fluorenylmethyloxycarbonyl (Fmoc) solid phase peptide
synthesis techniques. Rink-amide 4-methylbenzhydrylamine
(MBHA) resin was utilized for the synthesis. The Fmoc
protection group was removed by treatment with 20%
10 piperidine in dimethylformamide (DMF) twice (2 min for the
first time and 20 min for the second time).

All the amino acids were mixed with 4 molar equivalents
(eq.) of the resin, and the coupling was performed using 4
15 eq. of benzotriazol-1-yl-oxytripyrrolidinophosphonium
hexafluorophosphate (PyBOP) and 8 eq. of N,N-
diisopropylethylamine (DIEA) for 2 h.

In Examples 4 and 5, D-biotin was used at the last coupling
20 step to synthesize peptide substrates 3 and 4, as detailed
in Table 1 below. After sequence assembly, the final
deprotection and cleavage was performed by using a cocktail
of TFA : H₂O : triisopropylsilane : 2-mercaptoethanol in the
ratio of 94 : 2.5 : 2.5 : 1 for 3 h at room temperature.
25 The peptide was then precipitated with diethyl ether and
lyophilized. The crude peptides were purified by C18 semi-
prep HPLC.

The sequences of the Cys-containing peptides obtained in
30 Examples 2 to 6 are listed in Table 1 below.

Table 1

Ex.	No.	Substrate
2	1	Ac-Phe-Gln-Pro-Lys-Cys-Gly-NH ₂ (Ac-FQPKCG-NH ₂)
3	2	H-Val-Gly-Cys-Ala-Glu-Lys-Ser-Leu-NH ₂ (H-VGCAEKSL-NH ₂)
4	3	H-Trp-Ala-Cys-Tyr-Lys-Ser-Leu-NH ₂ (H-WACYKSL-NH ₂)
5	4	Biotin-Gly-Lys-Gly-Gly-Ala-Cys-Arg-His-Arg-Lys-Val-Leu-Arg-Asp-Asn-NH ₂ (Biotin-GKGGACRHRKVLIRD-NH ₂)
6	5	Biotin-Gly-Cys-Gly-Gly-Cys-Gly-Leu-Gly-Cys-Gly-Gly-Ala-Cys-Arg-NH ₂ (Biotin-GCGGCGLGCGGACR-NH ₂)

Furthermore, two control peptide substrates were synthesized without Cys residues: Ac-Phe-Gln-Pro-Lys-Ser-Gly-NH₂ and Ac-Phe-Gln-Pro-Lys-Lys(Ac)-Gly-NH₂.

Alkylation of Peptide Substrates

15 mM glutathione, 50 mM N-vinyl-acetamide (NVA) and 5 mM of initiator 2,2'-Azobis[2-(2-imidazolin-2-yl)propane]dihydrochloride (VA-044) in 0.2 M acetate buffer at pH 4 were mixed in a reaction tube to obtain a reaction mixture. The reaction mixture was added to each the synthesized peptide substrates in the amounts detailed in Table 2 below.

15

The reaction tube was irradiated under 365 nm ultraviolet (UV) for 30 min or 1 h to obtain reaction products. The reaction products were analyzed by C18 analytical HPLC and confirmed by ESI or matrix-assisted laser

desorption/ionization time-of-flight mass spectrometry (MALDI-TOF MS). The mass spectra are shown in Figs. 4 to 8. MS analysis was done on either desalted samples (using a C18 zip-tip) or on HPLC-purified fractions.

5

Table 2

Ex.	Substrate No.	Substrate	Amount (mM)	Reaction Time (h)	Yield (%)
2	1	Ac-Phe-Gln-Pro-Lys- <u>Cys</u> -Gly-NH ₂ (Ac-FQPK <u>C</u> G-NH ₂)	5	1	>95
3	2	H-Val-Gly- <u>Cys</u> -Ala-Glu-Lys-Ser-Leu-NH ₂ (H-VG <u>C</u> AEKSL-NH ₂)	5	1	>95
4	3	H-Trp-Ala- <u>Cys</u> -Tyr-Lys-Ser-Leu-NH ₂ (H-WA <u>C</u> YKSL-NH ₂)	5	0.5	95
5	4	Biotin-Gly-Lys-Gly-Gly-Ala- <u>Cys</u> -Arg-His-Arg-Lys-Val-Leu-Arg-Asp-Asn-NH ₂ (Biotin-GKGGAC <u>R</u> HRKVLRDN-NH ₂)	5	1	>95
6	5	Biotin-Gly- <u>Cys</u> -Gly-Gly- <u>Cys</u> -Gly-Leu-Gly- <u>Cys</u> -Gly-Gly-Ala- <u>Cys</u> -Arg-NH ₂ (Biotin-G <u>C</u> GG <u>C</u> GLG <u>C</u> GGAC <u>R</u> -NH ₂)	1.25	1	95

The C18 analytical HPLC profile of the reaction of Example 2 is shown in Fig. 4A. In Fig. 4A, peptide substrate 1 in

Example 2 is denoted by peak a, while the reaction product is denoted by peak b.

Peak a was further characterized by ESI-MS. In Fig. 4B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the peptide substrate is shown. The $[M+H]^+$ found had a m/z value of 720.29 and molecular weight of 719.34.

Peak b was further characterized by ESI-MS. In Fig. 4B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the reaction product is shown. The $[M+H]^+$ found had a m/z value of 805.28 and molecular weight of 804.39.

The HPLC conditions used in Example 2 were 0% to 40% of buffer B in buffer A in 40 min.

A comparative example was done for peptide substrate 1 at pH 6 instead, with all other conditions kept the same. The C18 analytical HPLC profile of the reaction of the comparative example is shown in Fig. 5A. In Fig. 5A, the reaction product is denoted by peak a, while the disulfide-linked side product between peptide substrate 1 and glutathione is denoted by peak b.

Peak a was further characterized by ESI-MS. In Fig. 5B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the reaction product is shown. The $[M+H]^+$ found had a m/z value of 805.37 and molecular weight of 804.39.

Peak b was further characterized by ESI-MS. In Fig. 5C, the mass spectrum of intensity versus the mass-to-charge ratio

(m/z) of the side product is shown. The $[M+H]^+$ found had a m/z value of 1025.36 and molecular weight of 1024.66.

In this comparative example, formation of a minute amount
5 (i.e. less than 3%) of disulfide cross-linked side product between the peptide substrate and glutathione was detected. This seems due to the presence in the glutathione sample of a small amount of an oxidized form of glutathione, which at the higher pH of pH 6 underwent a disulfide exchange with
10 the Cys thiol in the peptide substrate. At lower pH (e.g., pH 4), such an exchange reaction is inhibited. Accordingly, it can be concluded that a new and pure glutathione (reduced) sample should be used. Further, degassing of the buffer should be performed to prevent oxidation of the
15 glutathione.

The HPLC conditions used in the comparative example were 0% to 40% of buffer B in buffer A in 40 min.

20 The C18 analytical HPLC profile of the reaction of Example 3 is shown in Fig. 6A. In Fig. 6A, peptide substrate 2 in Example 3 is denoted by peak a, while the reaction product is denoted by peak b.

25 Peak a was further characterized by ESI-MS. In Fig. 6B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the peptide substrate is shown. The $[M+H]^+$ found had a m/z value of 805.26 and molecular weight of 804.42.

30 Peak b was further characterized by ESI-MS. In Fig. 4B, the mass spectrum of intensity versus the mass-to-charge ratio

(m/z) of the reaction product is shown. The $[M+H]^+$ found had a m/z value of 890.32 and molecular weight of 889.47.

5 The HPLC conditions used in Example 3 were 0 % to 30 % of buffer B in buffer A over 30 min.

The C18 analytical HPLC profile of the reaction of Example 4 is shown in Fig. 7A. In Fig. 7A, peptide substrate 3 in Example 4 is denoted by peak a, the expected reaction
10 product is denoted by peak b and the oxidation product is denoted by peak c.

Peak a was further characterized by ESI-MS. In Fig. 7B, the mass spectrum of intensity versus the mass-to-charge ratio
15 (m/z) of the peptide substrate is shown. The $[M+H]^+$ found had a m/z value of 869.34 and molecular weight of 869.04.

Peak b was further characterized by ESI-MS. In Fig. 7C, the mass spectrum of intensity versus the mass-to-charge ratio
20 (m/z) of the expected reaction product is shown. The $[M+H]^+$ found had a m/z value of 954.41 and molecular weight of 954.09.

Peak c was further characterized by ESI-MS. In Fig. 7D, the mass spectrum of intensity versus the mass-to-charge ratio
25 (m/z) of the oxidation reaction product is shown. The $[M+H]^+$ found had a m/z value of 970.39 and molecular weight of 970.08.

30 The HPLC conditions used in Example 4 were 0 % to 30 % of buffer B in buffer A over 30 min.

The C18 analytical HPLC profile of the reaction of Example 5 is shown in Fig. 8A. In Fig. 8A, peptide substrate 4 in Example 5 is denoted by peak a and the reaction product is denoted by peak b.

5

Peak a was further characterized by MALDI-TOF MS. In Fig. 8B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the peptide substrate is shown. The $[M+H]^+$ found had a m/z value of 1892.68 and molecular weight of 1890.99.

10

Peak b was further characterized by MALDI-TOF MS. In Fig. 8C, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the reaction product is shown. The $[M+H]^+$ found had a m/z value of 1977.93 and molecular weight of 1976.04.

15

The HPLC conditions used in Example 5 were 0 % to 30 % of buffer B in buffer A over 30 min.

20

Comparative examples were done for peptide substrate 4 under non-standard conditions. The MALDI-TOF MS of the reaction products when no glutathione added, while keeping all other conditions the same, is shown in Fig. 9A. As seen in Fig. 9A, additional side products were detected in addition to the desired thiol-ene coupling product. The side products were because the Cys residue was alkylated by di-, tri- and tetrameric NVAs. Further, the MALDI-TOF MS of the reaction products when glutathione was replaced with 15mM tris(2-carboxyethyl)phosphine (TCEP), keeping all other conditions the same, is shown in Fig. 9B. As seen in Fig. 9B, a desulfurization reaction took place to give a

25

30

product with a -32 MW. Thus, addition of the reducing agent TCEP was actually detrimental to the reaction as it led to desulfurization of the peptide substrate.

5 Accordingly, it can be concluded that glutathione, together with peptide substrate 4, participated in the critical rate-limiting chain transfer step to effectively intercept (or quench) the carbon free radical intermediate which was formed at the addition step of the thiyl radical to the NVA
10 ethylene double bond, thereby preventing it from reacting with another molecule of NVA. As expected, all the glutathione was also alkylated by NVA in the reaction.

The C18 analytical HPLC profile of the reaction of Example
15 6 is shown in Fig. 10A. In Fig. 10A, peptide substrate 5 in Example 6 is denoted by peak a, the expected reaction product is denoted by peak b and the oxidized thiol product is denoted by peak c.

20 Peak a was further characterized by MALDI-TOF MS. In Fig. 10B, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the peptide substrate is shown. The $[M+H]^+$ found had a m/z value of 1395.74 and molecular weight of 1394.52.

25 Peak b was further characterized by MALDI-TOF MS. In Fig. 10C, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the expected reaction product is shown. The $[M+H]^+$ found had a m/z value of 1735.89 and
30 molecular weight of 1734.72.

Peak c was further characterized by MALDI-TOF MS. In Fig. 10D, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the oxidized thiol product is shown. The $[M+H]^+$ found had a m/z value of 1751.96 and molecular weight of 1750.71.

The HPLC conditions used in Example 6 were 0 % to 30 % of buffer B in buffer A over 30 min.

It can thus be concluded from Example 6 that because peptide substrate 5 contains 4 Cys residues in its sequence, 1.25 mM of peptide 5 was sufficient to obtain the desired reaction product in a clean tetra-alkylating reaction with 95% yield, as shown in Table 2 above.

A comparative example was done with control peptide Ac-Phe-Gln-Pro-Lys-Ser-Gly-NH₂ which has no Cys residue. No reaction product was detected.

Further comparative examples were done without UV irradiation. No reaction products were detected.

Examples 7 to 9

Synthesis of Protein Substrates for Alkylation

Preparation of ubiquitin K48C

K48C point mutation was introduced to ubiquitin gene by QuikChangeTM site-directed mutagenesis kit (Stratagene) with the primers:

Ubi K48C F: 5'- GTCTGATATTTGCCGGCTGTCAGCTGGAGGATGGCCG-3';
and Ubi K48C R: 5'- CGGCCATCCTCCAGCTGACAGCCGGCAAATATCAGAC -
3'.

5 The plasmid containing ubiquitin K48C gene was transformed
into BL21 (DE3) cell. The cells were grown in 1 L LB media
containing ampicillin to OD600 of 0.6 and induced by a final
concentration of 0.5 mM IPTG for 4 h at 37°C. After
centrifugation at 6000 rpm for 10 min at 4°C, cells were
10 re-suspended in 50 ml lysis buffer (50 mM Tris-HCl, 100 mM
NaCl, 1 mM DTT, pH 7.5) and broken by microfluidization
(Microfluidics, Newton, USA). After centrifugation at
25,000 g for 30 min at 4°C, 70 % perchloric acid was added
to the supernatant at a ratio of 350 µl to 50 ml lysis
15 buffer. The mixture was stirred for 10 min and centrifuged
at 25,000 g for 30 min at 4°C. The supernatant was filtered
with 0.2 µm filter and dialyzed with 3.5 KDa cut-off
dialysis tubing against 50 mM ammonium acetate buffer (pH
4.5 with 1 mM DTT).

20

After filtration, the proteins were purified with a HiTrap™
SP FF 5 ml FPLC column (GE Healthcare Life Sciences). It
was eluted with a linear gradient from 0% to 100% FPLC
buffer B (50 mM ammonium acetate, pH 4.5, 1 mM DTT, 0.5 M
25 NaCl) in buffer A (50 mM ammonium acetate, pH 4.5, 1 mM
DTT) in 120 min at a flowrate 0.5 ml / min. Usually the
ubiquitin would be eluted at around 240 mM NaCl. After FPLC
purification, the proteins were dialyzed to ddH2O followed
by lyophilization.

30

Preparation of H4 K16C

K16C point mutation was introduced to H4 gene by the same mutagenesis kit with the primers:

H4 K16C F: 5'-GGTAAAGGTGGTGCTTGCCGTCACCGTAAAGTTC-3' and

5 H4 K16C R: 5'-GAACTTTACGGTGACGGCAAGCACCTTTACC-3'.

The plasmid containing H4 K16C gene was transformed into BL21 (DE3) pLysS cell. The cells were grown in 1 L LB media containing ampicilin and cholramphinicol to OD600 of 0.6 and induced by 0.4 mM IPTG for 3 h at 37°C. After centrifugation at 6000 rpm for 10 min, cells were re-suspended in 50 ml wash buffer (50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, pH 7.5) and broken by microfluidization.

15 The cell debris was removed by centrifugation at 20,000 g for 30 min at 4°C. The pellet was washed with wash buffer containing 1% Triton X-100 twice and one time without Triton X-100. Then 1 ml of DMSO was added to the pellet and the pellet was stirred for 30 min. After that 10 ml unfolding buffer (6 M Guanidinium HCl, 10 mM Tris-HCl, 10 mM DTT, pH 7.5) was added. After centrifugation, the supernatant was loaded to a 26/60 Sephacryl S-200 column and purified with gel filtration buffer (7 M de-ionized Urea, 20 mM sodium acetate, 1 M sodium chloride, 5 mM beta-mercaptoethanol, 0.5 M EDTA, pH 5.2). After FPLC purification, the protein was purified again by C4 semi-prep HPLC followed by lyophilization.

Preparation of H3 K27C

30 The C110A point mutation was introduced to H3 gene by mutagenesis kit with the primers:

H3 C110A F: 5'-GAGGACACCAACCTGGCCGCCATCCACGCCAAG-3'; and

H3 C110A R: 5'- CTTGGCGTGGATGGCGGCCAGGTTGGTGTCTCCTC-3'.

The K27C point mutation was introduced to H3 C110A gene by mutagenesis kit with the primers:

- 5 H3 K27C F: 5'- AAGGCAGCCAGGTGCTCCGCTCCTGCTACC -3'; and
H3 K27C R: 5'- AGCAGGAGCGGAGCACCTGGCTGCCTTGGTG-3'.

The expression of H3 K27C protein was the same as that of H4 K16C.

10

Alkylation of Protein Substrates

Introduction of sLys(Ac) into ubiquitin: Preparation of Ub_{K48Ac}

- 15 The freeze-dried ubiquitin K48C was dissolved in the 0.2 M acetate buffer (pH 4.0 or pH 7.0). The final concentrations of the reactants were as following:
Ubiquitin K48C: 0.5 mM.
NVA: 50 mM.
20 VA-044: 5 mM.
Glutathione: 15 mM.

- The reaction tube was irradiated under 365 nm UV for 2 h. The reaction product was analyzed by C18 analytical HPLC
25 and confirmed by ESI or MALDI-TOF MS (See Figs. 11 and 12). The protein remained soluble during the reaction period, indicating it stayed in its folded state.

- Figure 11A shows the C18 analytical HPLC profile of the
30 thiol-ene coupling reaction between ubiquitin K48C and NVA at pH 4.0, where Peak a corresponds to ubiquitin K48C; and peak b corresponds to the acetylated product.

The MALDI-TOF MS spectrum of peak a is show in Fig. 11B, where $[M+H]^+$ found is 8541.78, and the calculated MW is 8539.88.

5

The MALDI-TOF MS spectrum of peak b is shown in Fig. 11C wherein the $[M+H]^+$ found is 8626.58, and the calculated MW is 8624.93. The HPLC was carried out under the following conditions: 0 % to 30 % in 15 min, then to 50 % in 20 min of buffer B in buffer A.

10

Figure 12A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between ubiquitin K48C and NVA at pH 7.0.

15

The raw and deconvoluted mass of product determined by ESI-MS is shown in Fig. 12B, where the MW found is 8624.0, and the calculated MW is 8624.93).

20 The HPLC was carried out under the following conditions: 0 % to 30 % in 15 min, then to 50 % in 20 min of buffer B in buffer A.

This ubiquitin mutant contains a Cys residue at position 25 48. The protein (0.5 mM) was used in its native folded state for alkylation in the same reaction mixture at pH 4 or 7. The above MS analysis clearly showed an almost quantitative conversion, in 2 h, of the Cys residue to sLys(Ac) with the expected +85 Da MW for the alkylated 30 product. The protein remained soluble during the reaction, suggesting that no denaturing was occurring and that the presence of 50 mM NVA and 5 mM VA-044 did not affect the

structure of the folded protein. It is worth noticing that it would be difficult to use a semi-synthetic method to prepare such a modified protein since the modification site is in the middle of the sequence.

5

Two other proteins, histone H4 K16C and H3 K27C, were also modified with excellent results (see below). In these cases, 6 M Gdn.HCl was included in the alkylation reaction mixture. Interestingly, for the alkylation of H4 K16C at pH
10 4 or 7, in addition to the desired product, a side product was formed in significant amount (See below). The side product was more hydrophilic with a MW that was 16 Da higher than that of the expected acetyl-thialysine product. It appeared to result from oxidation of the thioether
15 linkage to sulfoxide, and inclusion of dimethylsulfide in the reaction mixture minimized the formation of this side product to about 5% (See Figs. 13 and 14).

No alkylation was detected when wild type H4 was subjected
20 to the same treatment. From these results, it can be seen that this free radical thiol reaction can tolerate various reaction conditions, e.g., native or denatured buffers, to modify a protein.

25 Introduction of sLys(Ac) into histone H4: Preparation of H4 K_S16Ac

The reaction was performed in the 0.2 M acetate buffer (pH 4.0 or pH 7.0) containing 6 M Guanidinium HCl. Dimethyl
30 sulfide was added to minimize oxidation of the thioester linkage in this product.

The final concentrations of the reactants were as following:

H4 K16C: 1 mM.

NVA: 50 mM.

5 VA-044: 5 mM.

Glutathione: 15 mM.

Dimethyl sulfide: 100 mM.

The reaction tube was irradiated under 365 nm UV for 2 h.

10 The reaction products were analyzed by C4 semi-prep HPLC and by ESI or MALDI-TOF MS (See Figs. 13 and 14).

Figure 13A is a C4 semi-prep HPLC profile of the thiol-ene coupling reaction between H4 K16C and NVA at pH 4.0 with or
15 without dimethyl sulfide. Peak a shown in Fig. 13A corresponds to H4 K16C; whereas peak b corresponds to expected product; and peak c corresponds to the oxidation product.

20 The raw and deconvoluted mass of the starting material H4 K16C in peak a determined by ESI-MS is shown in Fig. 13B, wherein the MW found is 11210.4, and the calculated MW is 11211.16).

25 The raw and deconvoluted mass of peak b determined by ESI-MS is shown in Fig. 13C, where the MW found is 11295.5, and the calculated MW is 11296.21.

The MALDI-TOF MS spectrum of peak c is shown in Fig. 13D,
30 where the $[M+H]^+$ found is 11311.56, and the calculated MW is 11312.2). The HPLC was performed under the following

conditions: 0 % to 40 % in 20 min, then to 60 % in 20 min of buffer B in buffer A.

5 Figure 14A is a C4 semi-prep HPLC profile of the thiol-ene coupling reaction between H4 K16C and NVA at pH 7.0, where peak a corresponds to the expected product; and peak b corresponds to the oxidation product.

10 The raw and deconvoluted mass of peak a as determined by ESI-MS is shown in Fig. 14B, where the MW found is 11296.5, and the calculated MW is 11296.21). The HPLC was performed under the following conditions: 0 % to 40 % in 20 min, then to 60 % in 20 min of buffer B in buffer A.

Introduction of sLys(Ac) into histone H3: Preparation of H3 K_s27Ac

The reaction was performed in the 0.2 M acetate buffer (pH 4.0) containing 6 M Guanidinium HCl. The final concentrations of the reactants were as following:

H3 K27C: 1 mM.

NVA: 50 mM.

VA-044: 5 mM.

10 Glutathione: 15 mM.

The reaction tube was irradiated under 365 nm for 2 h UV. The result was analyzed by C4 semi-prep HPLC and confirmed by ESI-MS (See Fig. 15).

15

The C4 semi-prep HPLC profile in Fig. 15A shows two peaks where peak a corresponds to the protein substrate H3 K27C and peak b corresponds to the acetylated product.

20 The ESI-MS spectrum of the raw and deconvoluted mass of peak a is provided in Fig. 15B, where the MW found is 15213.7, and the calculated MW is 15213.92.

25 The ESI-MS spectrum of the raw and deconvoluted mass of peak b is provided in Fig. 15C where the MW found is 15298.7, and the calculated MW is 15298.97. The HPLC were performed under the following conditions: 0 % to 40 % in 20 min, then to 60 % in 20 min of buffer B in buffer A.

30 Table 3 below summarizes the reaction yields of the above described protein alkylation reactions in Examples 7 to 9.

Table 3.

Ex.	Substrate No.	Substrate	Amount (mM)	Reaction Time (h)	Yield (%)
7	6	Ubiquitin K48C	0.5	2	>95
8	7	Histone H4 K16C	1	2	90
9	8	Histone H3 K27C	1	2	90

From the above, it can be seen that apart from being effective on peptide substrates, the thiol-ene coupling reaction is also highly effective on protein substrates.

Example 10

In this example, the generated sLys(Ac) was shown to be a good functional mimic of the natural Lys(Ac) using Western blots.

3 µg of the control protein (H4 K16C) and H4 K_s16Ac were dissolved respectively in loading buffer and run on 15 % SDS-PAGE, then transferred onto a polyvinylidene difluoride membrane. Thereafter, 20 ml of TBST (50 mM Tris.HCl, 150 mM NaCl, 0.1% Tween 20, pH 7.4) containing 5% non-fat milk was added for 1 h. The membrane was then incubated in 20 ml TBST with 5% non-fat milk containing Histone H4 K16Ac antibody (1:1000 dilution) overnight at 4°C. The membrane was then washed with TBST 4 times and incubated in 20 ml TBST with 5% non-fat milk containing the anti-rabbit IgG peroxidase conjugate (1:5000 dilution) for 1 h at room temperature. After washing with TBST for 4 times, the proteins were visualized by chemiluminescence.

As seen in Fig. 18A, the histone protein H4 K_s16Ac was recognized by a specific anti-H4 K16Ac antibody, as evidenced by the resultant darker stain on membrane. On the other hand, the unmodified H4 K16C was not recognized at all by the same antibody, as evidenced by the absence of a stain on the membrane.

Example 11

10 In this example, an enzymatic test was conducted to investigate whether the sLys(Ac) could be recognized by a histone deacetylase and used as substrate for deacetylation.

15 Sirtuin-2 (SIRT2), a class III NAD-dependent deacetylase, was used for the deacetylation reaction of the alkylated an acetyl-lysine analog of peptide 1 (Ac-FQPKK_s(Ac)G-NH₂) and the native peptide 1 (Ac-FQPKK(Ac)G-NH₂). The sequence of peptide 1 was based on residues 317-320 of protein p53
20 (i.e. Gln-Pro-Lys-Lys(Ac)) which is the best deacetylase substrate for the SIRT2 enzyme and the N-terminus was capped by an acetyl group to increase the hydrophobicity.

0.5 mM peptide and 0.1 mM NAD⁺ were mixed in SIRT2 assay
25 buffer (50 mM Tris, 137 mM NaCl, 2.7 mM KCl, 1 mM MgCl₂, 1 mg/ml BSA) and equilibrated at 37°C for 10 min. The reaction was initiated by adding 0.1 mM SIRT2 enzyme (0.1 U/μl) at 37°C and monitored by C18 analytical HPLC. It is to be noted that the deactylases were not very efficient in
30 deacetylating synthetic peptide substrates and required relatively large amount of the enzyme and a long reaction time for the deacetylation reaction. The results showed

that the deacetylation rate of the alkylated N-acetyl-thialysine peptide was about 1/3 of that on the control peptide containing the native N-acetyl-lysine.

5 The C18 analytical HPLC profile for Example 11 is shown in Figs. 16A and 16B (or Fig. 18B). Specifically in Fig. 16A, the C18 analytical HPLC profile for the deacetylated product of peptide 1 with the installed acetyl-lysine analog in the SIRT2 assay is shown. In Fig. 16A, peak a
10 denotes the deacetylated product, while peak b denotes the acetyl-lysine analog (Ac-FQPKK₅(Ac)G-NH₂). In Fig. 16B, the C18 analytical HPLC profile for the deacetylated product of the native K(Ac)-peptide substrate in the SIRT2 assay is shown. In Fig. 16B, peak c denotes the deacetylated
15 product, while peak d denotes the native peptide substrate.

Peak a was further characterized by ESI-MS in Fig. 16C. In Fig. 16C, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the deacetylated product of the
20 acetyl-lysine analog is shown. The [M+H]⁺ found had a m/z value of 763.30 and molecular weight of 762.38.

Peak b was further characterized by ESI-MS in Fig. 16D. In Fig. 16D, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the acetyl-lysine analog is shown.
25 The [M+H]⁺ found had a m/z value of 805.28 and molecular weight of 804.39.

Peak c was further characterized by ESI-MS in Fig. 16E. In
30 Fig. 16E, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the deacetylated product of the native native K(Ac)-peptide substrate is shown. The [M+H]⁺

found had a m/z value of 745.28 and molecular weight of 744.43.

Peak d was further characterized by ESI-MS in Fig. 16F. In Fig. 16F, the mass spectrum of intensity versus the mass-to-charge ratio (m/z) of the native K(Ac)-peptide substrate is shown. The $[M+H]^+$ found had a m/z value of 787.33 and molecular weight of 786.44.

The HPLC conditions used in Example 11 were 0% to 30% of buffer B in buffer A in 30 min.

It can thus be concluded that the sLys(Ac) residue in the alkylated peptide 1 was susceptible to enzymatic deacetylation, albeit to a lesser degree, compared to its native counterpart. This is evidenced by the smaller peak a as compared to peak c.

Example 12

Preparation of H4 K_s16

In this example, H4 K_s16 was synthesized using the classic aminoethylation reaction.

H4 K16C (0.5 mM) was dissolved in the 1 M HEPES [4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid] buffer (pH 7.8, 6 M Gdn.HCl, 5 mM D/L methionine, 20 mM dithiothreitol (DTT)) containing 2-bromoethylamine hydrobromide (160 mM).

After 11 h reaction at room temperature in the dark, the alkylated protein was dialyzed to ddH₂O followed by

lyophilization. The reaction product was analyzed by ESI-MS and is shown in Fig. 17.

In Fig. 17, the raw spectrum and deconvoluted mass of the alkylated product (H4 K_s16) as determined by ESI-MS had a molecular weight of 11253.4, while the molecular weight calculated was 11254.2.

EXAMPLE 13

Acetylation of Lys16 in histone H4 is known to inhibit the folding of nucleosome arrays and hence the formation of the compact 30-nm chromatin fiber. H4 K_s16Ac and three other control H4 proteins (H4 K16Ac, H4K_s16 and H4-WT, described below) were incorporated respectively, together with H3, H2A and H2B, into histone octamers.

H4 K16Ac was prepared using a semi-synthetic approach and H4 K_s16 was synthesized in Example 12 by alkylating H4 C16 with 2-bromoethylamine. The four different octamers were then individually combined with the 12-177-601 DNA to assemble into the 12-nucleosome arrays.

Using analytical ultracentrifugation, it was demonstrated that the K_s16Ac produced an identical effect as the native K16Ac in abolishing Mg²⁺-induced folding of the reconstituted nucleosome array, as seen in Fig. 18C.

The AUC data clearly showed that, in the presence of 1 mM MgCl₂, the nucleosome array containing wild-type H4 or its equivalent H4 K_s16 folded into a significantly more compact

state (sedimentation coefficient $S_{20^{\circ}\text{C}, w} = 52-53\text{S}$) than did the K16Ac and Ks16Ac arrays ($S_{20^{\circ}\text{C}, w} = 44-45\text{S}$). Remarkably, these results prove not only the functional equivalency between sLys(Ac) and Lys(Ac), but also the functional
5 equivalency between sLys and Lys.

Nucleosomal Array Reconstitution:

Plasmid containing the 12-177-601 DNA is transformed and
10 amplified in HB101 cell. The plasmid was extracted as described in Korolev et al, *Biophys. J.* 2010, 99, 1865-1905.

RNA and protein impurities were removed by gel filtration
15 on a Sepharose 6 column with the use of TES2000 buffer. After excision with EcoRV, the 12-177-601 DNA was separated from short plasmid fragment by polyethylene glycol (PEG 6000). Finally 12-177-601 DNA was purified on a Sephacryl SF1000 column with TES100 buffer.

20

Wild type *Xenopus laevis* histones H2A, H2B, H3, and H4 were individually over-expressed in BL21 (DE3) pLysS cell in presence of ampicillin and chloramphenicol. Each histone was purified by gel filtration on Sephacryl S-200 column
25 and subsequently on a Resource S cation exchange column. The H4 K16Ac was prepared by chemical semi-synthesis described in Allahverdi et al, *Nucleic Acids. Res.*, 2011, 39, 1680-1691.

30 The histone octamer was formed using a molar ratio of 1:1:1.2:1.2 for H2A, H2B, H3, and H4. The histone octamer was purified on a Sephacryl S-200 gel filtration column.

The nucleosome array was reconstituted by step-wise dialysis using the histone octamer and 12-177-601 DNA as described in Rorigo et al, *J. Mol. Biol.*, 2003, 327, 85-96 and T. Schalch, *The 30-nm chromatin fiber. In vitro reconstitution and structural analysis. PhD Thesis. 2004, Swiss Federal Institute of Technology, Zurich.* <http://e-collection.library.ethz.ch/eserv/eth:27516/eth-27516-02.pdf>.

To prevent excessive binding of the histone to the DNA, 0.5 molecule of competitor Core Length DNA (150-bp) per one array was added to the reconstitution mixture. The exact amount of histone octamer to 12-177-601 DNA to get the stoichiometry of 12:1 was determined empirically in small scale preparations using different ratios of histone octamer to the DNA template. The reconstituted arrays were purified as described above. Purified array material and also the respective digests by ScaI were checked on 5% polyacrylamide gel electrophoresis (PAGE) to verify the quality of the nucleosomal arrays.

Analytical ultracentrifugation

Sedimentation velocity experiment were carried out on a Beckman XL-I analytical ultracentrifuge equipment with AN-50T rotor and monochrome scanner. The stock solution of array were diluted in TEK buffer to get $A_{259} = 0.8 \text{ cm}^{-1}$ (DNA concentration $C_p = 121 \text{ } \mu\text{M}$).

The sample and reference (TEK buffer + salt at same concentration as in the sample) were loaded into the 12 mm double channel cells and equilibrated under vacuum for 30

min at 3000rpm and 20 °C. Data measurement and analyses were carried out following methods described in Korolev et al and Allahverdi et al described above.

- 5 Table 4 provides values of $S_{20^{\circ}\text{C},w} \pm \text{SD}$ of nucleosome arrays in the presence of different concentrations of Mg^{2+} .

Table 4.

Mg ²⁺ concentration	Array			
	H4 WT	H4 K _S 16	H4 K16Ac	H4 K _S 16Ac
0.0 mM	35.4±0.27	35.2±0.68	36.2±0.93	34.8±0.77
0.4 mM	42.5±0.3	40.7±0.42	40±1.32	41.6±0.76
0.8 mM	48.5±0.25	48.9±0.79	42.9±0.61	43.3±0.59
1.0 mM	52.7±0.4	52.3±1.03	44.4±0.6	44.7±0.56

10

Example 14

The reaction with vinyl acetate was also successful. The reaction with methyl vinyl ether and di(ethylene glycol) vinyl ether obtained similar results. The product of the
 15 reaction on the Cys residue of a target molecule with vinyl acetate is also a mimic of Lys(Ac). But more importantly, the reaction with poly(ethylene glycol) vinyl ether is a very useful method for PEGylation of peptides and proteins.

20

PEGylation is being increasingly used for peptide/protein modification as a way to prolong the half-life of therapeutic peptides and proteins. Several PEGylated peptides and proteins (e.g., PEG-interferon alpha, PEG-G-
 25 CSF) are already on the market.


An experiment was carried out on one exemplary peptide (Peptide 1 in Table 1 above: Ac-Phe-Gln-Pro-Lys-Cys-Gly-amide). The reaction was conducted in the acetate buffer (pH 4) containing 5 mM of peptide 1, 100 mM vinyl acetate, 5 mM VA-044 and 15 mM glutathione. After 2h UV (365 nm) irradiation at room temperature, >90% alkylation product was obtained ($[M+H]^+$ found 805.2).

EXAMPLE 15

Alkylation with N-vinylpropionamide or N-vinylbutyramide

Alkylation of a Cys residue with N-vinylpropionamide or N-butyrylamide generates a mimic of N^ε-propionyl-Lys or N^ε-butyryl-Lys residue, respectively. The mechanism of reaction is as provided in Scheme V below:

Scheme V



Peptide 1 from Table 1 was used as the substrate. All the reactants were dissolved in the 0.2 M acetate buffer (pH 4.0) as indicated above and the final concentrations were
5 as following:

Peptide 1: 5 mM

N-vinylpropionamide or N-vinylbutyramide: 50 mM

VA-044: 5 mM

10 Glutathione (reduced form): 15 mM

The reaction tube was irradiated under 365 nm UV for 1 h at room temperature. The reaction products were analyzed by C18 analytical HPLC and confirmed by ESI or MALDI-TOF MS.
15 MS analysis was done on either desalted samples (using a C18 zip-tip) or on HPLC-purified fractions.

Results

20 With reference to Figs. 19 and 20, it can be seen that the reaction of peptide 1 with N-vinylpropionamide or N-vinylbutyramide was similar to that with NVA. The reaction was clean and efficient with a yield of over 85% after 1 h. No telomerization sideproducts were detected.

25

Specifically, Fig 19A shows the C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-vinylpropionamide, where Peak a corresponds to the product; Peak b corresponds to peptide 1. Fig. 19B shows the ESI-MS
30 of peak a where the $[M+H]^+$ found was 820.26, and the calculated MW is 819.52. The HPLC was performed under the

following conditions: 0% to 40% of buffer B in buffer A in 40 min.


Fig 20A shows the C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-vinylbutyramide, where Peak a corresponds to the product; Peak b corresponds to peptide 1. Fig. 20B shows an ESI-MS of peak a where the $[M+H]^+$ found is 834.55, and the calculated MW is 833.67. The HPLC was performed under the following conditions: 0% to 40% of buffer B in buffer A in 40 min.

Example 16

15 ***Comparative study of the alkylation reaction with NVA, vinyl acetate, N-allylacetate or N-methyl-N-vinylacetate***

Several ene reagents (alkylating reagents), vinyl acetate, N-allylacetate and N-methyl-N-vinylacetate, were compared with NVA in the reaction with Cys to generate other N^ε-acetyl-lysine analogues by the thiol-ene coupling method. Their respective reaction mechanisms are provided in Scheme VI below.

Although these analogues are structurally similar, they differ from one another in electrosteric properties which may result in subtle differences in functions and therefore may be useful for better understanding the effects of protein modifications by the acetyl group.

Scheme VI5 Methods

For the site-specific installation of the acetyl-lysine analogs shown in Scheme VI, a model peptide (peptide 1) was modified by the thiol-ene coupling reaction. All the reactants were dissolved in the 0.2 M acetate buffer (pH 4.0) as indicated and the final concentrations are as follows:

10

Peptide 1: 5 mM

vinyl acetate, or N-allylacetamide, or N-methyl-N-vinylacetamide: 50 mM

VA-044: 5 mM

5 Glutathione (reduced form): 15 mM

The reaction tube was irradiated under 365 nm UV for 1 h at room temperature. The reaction products were analyzed by C18 analytical HPLC and confirmed by ESI or MALDI-TOF MS.

10 MS analysis was done on either desalted samples (using a C18 zip-tip) or on HPLC-purified fractions.

Peptide 1 was treated in a similar way with vinyl acetate, N-allylacetamide, and N-methyl-N-vinylacetamide

15 respectively to generate different acetyl-lysine analogues.

The experimental results are provided in Figs. 21 - 23.

A notable distinction from the reaction with NVA was that no telomerization side products were detected. Although

20 all these alkylating agents contain a C=C double bond, their reactivities appeared to be different. In particular, these alkylating agents appeared to be less reactive than NVA.

25 Table 5 lists the yields at 1 h of the reactions with the different alkylating agents used in this Example. As seen from Table 5, the reactivity order is: N-vinylacetamide > N-methyl-N-vinylacetamide > vinyl acetate > N-

30 allylacetamide. In particular, while a yield of >95% was obtained with NVA at 1 h, the respective yields obtained with N-methyl-N-vinylacetamide, vinyl acetate and N-allylacetamide were about 79%, 51% and 38%. This may have









resulted from the different stabilities of the radical intermediates or from the different steric effects of these alkylating agents. Notably, all the reactions were very clean (Figs. 21-23) and proceeded to completion with prolonged time (data not shown).

In particular, Figure 21A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-methyl-N-vinylacetamide, wherein Peak a corresponds to the product; and Peak b corresponds to peptide 1. Fig. 21B shows the ESI-MS of peak a, where the $[M+H]^+$ found is 819.36, whereas the calculated MW is 818.40. The HPLC was performed under the following conditions: 0% to 40% of buffer B in buffer A in 40 min.

Figure 22A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and vinyl acetate, wherein Peak a corresponds to the product; and Peak b corresponds to peptide 1. Fig. 22B shows the ESI-MS of peak a, where the $[M+H]^+$ found is 806.30, whereas the calculated MW is 805.37. The HPLC was performed under the following conditions: 0% to 40% of buffer B in buffer A in 40 min.


Figure 23A shows a C18 analytical HPLC profile of the thiol-ene coupling reaction between peptide 1 and N-allylacetamide, wherein Peak a corresponds to the product; and Peak b corresponds to peptide 1. Fig. 23B shows the ESI-MS of peak a, where the $[M+H]^+$ found is 819.30, whereas the calculated MW is 818.40. The HPLC was performed under the following conditions: 0% to 40% of buffer B in buffer A in 40 min.

Table 5.

Comparison of different ene reagents in thio-ene coupling reaction			
Peptide 1	Alkylating agent	Product	Yield (1h) (%)
Ac-FQPKCG	 <i>N</i> -vinylacetamide		>95
Ac-FQPKCG	 <i>N</i> -methyl- <i>N</i> -vinylacetamide		78.7
Ac-FQPKCG	 vinyl acetate		50.9
Ac-FQPKCG	 <i>N</i> -allylacetamide		38.3

Example 17**Alkylation with N-vinyl-peptidylamide or N-vinyl-proteinylamide**

5 In this Example, the same thiol-ene coupling reaction is carried out but with a peptidylamide and proteinylamide alkylating agent. In these two cases, the -ene compound has a large acyl group, i.e., the peptidyl or proteinyl group (See Scheme VII below). The thiol compound can be a
10 small organic thiol compound, a peptide/protein containing Cys residue(s) or a thiol-functionalized polymer (e.g., PEG).

Scheme VII


15 The initial step is to prepare a peptide or protein with a C-terminal vinyl (or allyl) amide. The present inventors have developed an indirect method for this synthesis.

20


A peptide or protein thioester is first prepared chemically or biosynthetically, and which is then reacted with N-vinyl-glycylamide (i.e., 2-amino-N-vinylacetamide) in the presence of silver ions (See Scheme VIII below). This
25 allows the introduction of the N-vinyl group onto the C-terminus of the peptide with the extension of one amino

acid (Gly in this case). A peptide thioester can be prepared by the solid phase peptide synthesis technique and a large peptide or protein thioester can be prepared by using the intein-catalyzed protein splicing technique.

5

Scheme VIII

- 10 Using this method, a small peptide (RLYRAG) and a protein (ubiquitin) were prepared, each containing the C-terminal N-vinyl amide (See Scheme IX below). Specifically, the peptide N-vinyl amide was prepared from RLYRA-COSR and the ubiquitinyl N-vinylamide was prepared from the Ub(1-75)-
- 15 COSR.

Scheme IX

- 20 A small thiol compound, methyl mercaptoacetate, was used to react with the peptide RLYRAG-vinyl amide in the thiol-ene coupling reaction. It was found that the vinyl group on

the C-terminal amide of the peptide was much less reactive than in NVA. When VA-044 was used as the radical initiator, the reaction could only take place at elevated temperature (at 60 °C or higher) with only a modest yield.

5

It was further found that a different photoinitiator lithium phenyl-2,4,6-trimethylbenzoylphosphinate (LAP) was better than VA-044 in promoting the thiol-ene radical reaction in this case. At 50°C, over 70% of reaction product was obtained after 2 h reaction with LAP as the photoinitiator.

10

Figure 24 shows the EMI-MS of the isolated product formed from the thiol-ene coupling reaction between RLYRAG-vinyl amide and methyl mercaptoacetate, wherein the $[M+H]^+$ found: was 972.82 (+113 TFA adduct, because of the highly basic nature of the peptide) and the calculated MW: is 894.88.

15

This initiator (LAP) was also used for the reaction of the ubiquitin-vinyl amide. A relative large thiol substrate Ubi-K48C was used, which is monoubiquitin in which Lys48 was mutated to Cys. After 1 h reaction at 50°C, the diubiquitin product was formed at about 35-40% as analyzed from SDS-PAGE gel.

20

25

Methods

Experimental procedures for peptide RLYRAG-vinyl amide

Preparation of peptide RLYRAG-vinyl amide

30

The thioester peptide RLYRA-COSR was reacted with 2-amino-N-vinylacetamide in the presence of silver ions at room

temperature. All the reagents were dissolved in DMSO and the reaction conditions are as follows:

RLYRA-COSR: 5 mM

5 2-amino-N-vinylacetamide: 100 mM

AgNO₃: 100 mM

After 16 h, the product was purified by C18 semi HPLC followed by lyophilization.

10

Thiol-ene coupling reaction between RLYRAG-vinyl amide and methyl mercaptoacetate

The purified RLYRAG-vinyl amide is thereafter reacted with
15 methyl mercaptoacetate by the thiol-ene coupling reaction. The reaction was performed in 0.2 M acetate buffer (pH 4 or 5).

The final concentrations of the reactants were as follows:

20

RLYRAG-vinyl amide: 10 mM

methyl mercaptoacetate: 40 mM

LAP: 5 mM

25 The reaction tube was irradiated under 365 nm UV at 50°C for 2 h. The reaction product was purified by HPLC and confirmed by ESI-MS (See Fig. 24).

30

2) Experimental procedures for N-vinyl-ubiquitinylamide (Ubi-vinyl amide)

Preparation of Ubi (1-75)-COSR

- 5 The human ubiquitin gene was amplified by PCR using the primers:

Ubi75_F: 5'-GGTGGTCATATGCAGATCTTTGTGAAG-3'; and

Ubi75_R: 5'-CTGGTCAGGTGGGATACCCTCCTTGTCTTGAATTTTG-3'.

- 10 The PCR product was purified and ligated into the T-easy vector (*Promega*). After digestion with *NdeI* and *SapI* restriction enzymes, the product was purified and ligated into the indentically digested pTYB1 vector (*New England Biolabs*). Then the correct insert was confirmed by
15 sequencing.

- The plasmid pUbi 75aa-TYB1 was transformed into *E. coli* BL21(DE3) cells. The cells were grown in LB medium (containing 100µg/ml Ampicillin) at 37°C to an OD600 of
20 0.6-0.8. The desired protein was induced by 50 µM IPTG at 15°C for 18 h. After centrifuge at 6000 rpm for 10 mins, cell pellets from 1 liter of cells were resuspended in 50ml lysis buffer (50 mM HEPES, 500 mM NaCl, 1 mM beta-mercaptoethanol, pH 7.5). Cells were then broken by
25 sonication and debris removed by centrifugation at 20,000g for 30 min. The supernatants are then mixed with pre-equilibrated 3 ml chitin beads (*New England Biolabs*) at 4°C for 2h. The beads were then poured into a column and washed with 40 ml lysis buffer. The fusion protein was
30 cleaved by adding 4 ml lysis buffer containing 100 mM 2-mercaptoethanesulfonic acid (MESNA) and incubating at 37°C overnight. The Ubi(1-75) thioester was eluted by 10 ml

lysis buffer. Affinity binding, cleavage and purification were monitored by SDS-PAGE and HPLC. The purified Ubi(1-75)-COSR was dried by lyophilization.

5 Preparation of Ubi-vinyl amide

The purified Ubi(1-75)-COSR was reacted with 2-amino-N-vinylacetamide to generate the Ubi-vinyl amide. All the reagents were dissolved in DMSO and the reaction conditions are as follows:

10

Ubi(1-75)-COSR: 2 mM

2-amino-N-vinylacetamide: 100 mM

AgNO₃: 100 mM

15 After 16 h, the product was purified by C4 semi HPLC followed by lyophilization.

Preparation of Ubi-K48C

K48C point mutation was introduced to ubiquitin gene by QuikChangeTM site-directed mutagenesis kit (Stratagene) with the primers:

Ubi K48C F: 5'- GTCTGATATTTGCCGGCTGTCAGCTGGAGGATGGCCG-3';
and

Ubi K48C R: 5'-CGGCCATCCTCCAGCTGACAGCCGGCAAATATCAGAC -3'.

25

The plasmid containing ubiquitin K48C gene was transformed into BL21 (DE3) cell. The cells were grown in 1 L LB media containing ampicillin to OD₆₀₀ of 0.6 and induced by a final concentration of 0.5 mM IPTG for 4 h at 37°C. After centrifugation at 6000 rpm for 10 min at 4°C, cells were resuspended in 50 ml lysis buffer (50 mM Tris-HCl, 100 mM NaCl, 1 mM DTT, pH 7.5) and broken by microfluidization

30

(Microfluidics, Newton, USA). After centrifugation at 25,000 g for 30 min at 4°C, 70 % perchloric acid was added to the supernatant at a ratio of 350 µl to 50 ml lysis buffer. The mixture was stirred for 10 min and centrifuged at 25,000 g for 30 min at 4°C. The supernatant was filtered with 0.2 µm filter and dialyzed with 3.5 KDa cut-off dialysis tubing against 50 mM ammonium acetate buffer (pH 4.5 with 1 mM DTT).

After filtration, the proteins were purified with a HiTrap™ SP FF 5 ml FPLC column (GE Healthcare Life Sciences). It was eluted with a linear gradient from 0% to 100% FPLC buffer B (50 mM ammonium acetate, pH 4.5, 1 mM DTT, 0.5 M NaCl) in buffer A (50 mM ammonium acetate, pH 4.5, 1 mM DTT) in 120 min at a flowrate 0.5 ml / min.

Usually the ubiquitin would be eluted at around 240 mM NaCl. After FPLC purification, the proteins were dialyzed to ddH₂O followed by lyophilization.

20

Synthesis of di-ubiquitin by thiol-ene coupling reaction

The purified Ubi-vinyl amide was reacted with Ubi-K48C to generate di-Ubi by thiol-ene coupling reaction. The reaction was performed in the 0.2 M acetate buffer (pH 5.0) containing 6 M Guanidinium HCl. The final concentrations of the reactants are as follows:

Ubi-vinyl amide: 2 mM

Ubi-K48C: 2 mM

30 LAP: 4 mM

The reaction tube was irradiated under 365 nm UV, at 50°C, for 2 h. The reaction products were analyzed by SDS-PAGE.

Applications

5

Site-specifically modified proteins are invaluable reagents for studying the biology of protein lysine acetylation - an intensive field in biomedical research at present. However, as discussed above, these proteins are difficult to come by and none of them is available commercially. The presently disclosed method will make such proteins easily available and an immediate application will be to produce these modified proteins as commercial products for the biomedical research community.

15

In the long term, since many proteins require modification on their structure to acquire desired activity and/or improved pharmacokinetic properties, the technology can be used to manufacture modified proteins with e.g. longer half-life for therapeutic applications.

20

Accordingly, in summary, the present disclosure provides a thiol-ene radical addition reaction involving, for instance, the commercially available NVA is well suited for the S-acetamidoethylation of cysteine residues in synthetic peptides and recombinant proteins. The resultant N-acetyl-thialysine differs with natural acetyl-lysine only isosterically at the γ -position of the amino acid structure and is functionally equivalent or similar to the latter.

25

30


The disclosed reaction and method has many potential applications, such as the histone epigenetic study. The disclosed reaction system is robust and gives near quantitative yields of site-specifically acetylated proteins which can be purified in a simple chromatography or dialysis step. The ease of implementation of this method also makes it easily adoptable by researchers from the bioscience research community. As such, this radical reaction approach provides a convenient enabling tool for the study of lysine acetylation biology and will help to advance research in this important field.

It will be apparent that various other modifications and adaptations of the invention will be apparent to the person skilled in the art after reading the foregoing disclosure without departing from the spirit and scope of the invention and it is intended that all such modifications and adaptations come within the scope of the appended claims.


Claims

1. A method of alkylating a thiol group (R-S-H) or seleno group (R-Se-H) in a target molecule wherein the method comprises:

reacting a target molecule comprising at least one thiol group with a compound of formula (I) or (II):




(I)



(II)

wherein R is an acetyl group or any other acyl group or is a group comprising any one of:



or

wherein R in formula (II) can also be an alkyl group;

and

wherein R' is selected from a group consisting of a hydrogen, a methyl group and an ethyl group.

2. The method of claim 1, wherein the target molecule is an organic molecule.

3. The method of claim 2, wherein the organic molecule is a peptide or protein.
4. The method of claim 3, wherein the peptide is selected from the group consisting of an oligopeptide, a polypeptide and a synthetic peptide.
5. The method of claim 3, wherein the protein is selected from the group consisting of a recombinant protein and a protein complex.
6. The method of claim 1, wherein the target molecule is histone or ubiquitin.
7. The method of any one of claims 1 to 6, wherein the thiol group is the thiol group of a cysteine and/ or a selenocysteine.
8. The method of claim 7, wherein the thiol group is the thiol group of a cysteine.
9. The method of any one of the preceding claims, wherein the acyl group is an aliphatic acyl group, or an alicyclic acyl group, or an aromatic acyl group, or an amino acyl or a peptidyl or a proteinyll group.
10. The method of claim 9, wherein the aliphatic acyl group is selected from the group consisting of a linear aliphatic acyl group, a branched aliphatic acyl group, a tert-butyloxycarbonyl group, a derivative of tert-butyloxycarbonyl group, a benzyloxycarbonyl group and a derivative of benzyloxycarbonyl group.

11. The method of claim 9, wherein the acyl group is selected from the group consisting of a linear aliphatic acyl groups having 1 to 20 or 1 to 7 carbon atoms; a branched aliphatic acyl groups having 1 to 7 carbon atoms; and a poly(ethylene glycol) moiety.
12. The method of claim 9, wherein the acyl group is selected from the group consisting of formyl group, acetyl group, propionyl group, 2-propionyl group, butyryl group, isobutyryl group, pentanoyl group, hexanoyl group, allylcarbonyl group, cyclohexylmethylcarbonyl group, and C₃-C₆ cycloalkylcarbonyl groups, such as cyclopropylcarbonyl group, cyclopentylcarbonyl group, cyclohexylcarbonyl group, and 1-cyclohexenylcarbonyl group.
13. The method of claim 9, wherein the aromatic acyl group is selected from the group consisting of benzoyl group, 4-methylbenzoyl group, and 4-methoxybenzoyl group.
14. The method of any one of the preceding claims, wherein the alkyl group is selected from the group consisting of methyl, ethyl, n-propyl, i-propyl, n-butyl, t-butyl, sec-butyl, pentyl, hexyl, octyl, nonyl, decyl, undecyl, dodecyl, trideyl, tetradecyl, pentadecyl, hexadecyl, octadecyl, poly(ethylene glycol)-ethyl and poly(ethylene glycol)-propyl [e.g., HO(CH₂CH₂O)_n-CH₂CH₂-; CH₃O(CH₂CH₂O)_n-CH₂CH₂-; CH₃O(CH₂CH₂O)_n-CH₂CH₂CH₂-].

15. Use of a compound of Formula (I) as defined in any one of claims 1 to 14 as alkylating agent for alkylation of a target molecule.
- 5 16. Use of a compound of Formula (II) as defined in any one of claims 1 to 14 as alkylating agent for alkylation of a target molecule comprising a thiol-group (R-S-H) or a seleno group (R-Se-H).
- 10 17. Use of a compound of Formula (I) for installing an acetylated lysine residue analog in a target molecule.
- 15 18. The use of any one of claims 15 to 17, wherein the target molecule is an organic molecule.
19. The use of claim 18, wherein the organic molecule is a peptide or protein.
- 20 20. The use of claim 19, wherein the peptide is selected from the group consisting of an oligopeptide, a polypeptide and a synthetic peptide.
- 25 21. The use of claim 19, wherein the protein is selected from the group consisting of a recombinant protein and a protein complex.
- 30 22. A pharmaceutical composition comprising a target molecule modified according to the method of any one of claims 1 to 14.

23. The method of any one of claims 1 to 14 wherein the compound of formula (I) is N-vinylacetamide.

FIGURES

1/21




Fig. 1

2/21




FIG. 2

3/21




FIG. 3

4/21




FIG. 4

5/21




FIG. 5

6/21




FIG. 6

7/21




FIG. 7

8/21




FIG. 8

9/21




FIG. 9

10/21




FIG. 10

11/21




FIG. 11

12/21




FIG. 12

13/21




FIG. 13




FIG. 14

15/21




FIG. 15

16/21




FIG. 16

17/21




FIG. 17

18/21




FIG. 18

19/21




FIG. 19




FIG. 20

20/21




FIG. 21




FIG. 22

21/21




FIG. 23




FIG. 24

INTERNATIONAL SEARCH REPORT

International application No.
PCT/SG2012/000168

A. CLASSIFICATION OF SUBJECT MATTER

C07C 319/18 (2006.01) C07K 1/107 (2006.01) C07K 1/113 (2006.01) C07D 277/04 (2006.01) C07C 233/05 (2006.01)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CAS Registry, CASREACT and CAPlus via STN Online: Sub-structure search in accordance with Formulae (I) and (II);
Keywords: CAS Registry Numbers [625-77-4], [3195-78-6], [108-05-4] and [692-33-1]; ALKYLAT?, PROTEIN, ?PEPTID?, ?CYSTEIN?, HISTONE, UBIQUITIN, DENDRIMER, ?GLYCON?, SUGAR

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
	Documents are listed in the continuation of Box C	

☒ Further documents are listed in the continuation of Box C

☒ See patent family annex

* Special categories of cited documents:	
"A" document defining the general state of the art which is not considered to be of particular relevance	"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"E" earlier application or patent but published on or after the international filing date	"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"O" document referring to an oral disclosure, use, exhibition or other means	"&" document member of the same patent family
"P" document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search
22 October 2012

Date of mailing of the international search report
22 October 2012

Name and mailing address of the ISA/AU

AUSTRALIAN PATENT OFFICE
PO BOX 200, WODEN ACT 2606, AUSTRALIA
Email address: pct@ipaaustralia.gov.au
Facsimile No.: +61 2 6283 7999

Authorised officer

Marie-Luise Huber
AUSTRALIAN PATENT OFFICE
(ISO 9001 Quality Certified Service)
Telephone No. 0262832788

INTERNATIONAL SEARCH REPORT		International application No.
C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		PCT/SG2012/000168
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	Li, F. et al., "A Direct Method for Site-Specific Protein Acetylation", Angew. Chem. Int. Ed., 2011, Vol. 50, pages 9611-9614 and Supporting Information (see http://dx.doi.org/10.1002/anie.201103754) & CASREACT Online Abstract Accession No. 156:94557 Page 9611, last paragraph to Page 9613, end of final paragraph; Supporting Information; and CASREACT Online Abstract Accession No. 156:94557, Reactions 1, 7, 8, 12, 13 and 14	1-23
X	Mustapa, M. et al., "Synthesis of Orthogonally Protected Lanthionines", Journal of Organic Chemistry, 2003, Vol. 68, No. 21, pages 8185-8192 & CASREACT Online Abstract Accession No. 139:350921 Page 8185 Abstract; page 8185, left column, first paragraph; page 8186 Scheme 1; page 8187 Scheme 2; Page 8189, left column, third paragraph - right column, second paragraph; and page 8189, Scheme 8; CASREACT Online Abstract Accession No. 139:350921: Reactions 32, 33 and 52	1-5, 7-9, 16 and 18-22
X	Yeung, C. L. et al., "Tuning Specific Biomolecular Interactions Using Electro-Switchable Surfaces", Advanced Functional Materials, 2010, Vol. 20, No. 16, pages 2657-2663. & CASREACT Online Abstract Accession No. 153:549060 Page 2657 Abstract, page 2658 left column paragraph 1, page 2658 Scheme 1, Supporting Information Scheme S1	1, 2, 14, 16 and 18
X	Lanza, T. et al., "Radical additions of thiols to alkenes and alkynes in ionic liquids", Current Organic Chemistry, 2009, Vol. 13, No. 17, pages 1726-1732 & CASREACT Online Abstract Accession No. 152:380733 Page 1726 Abstract, page 1727 left column paragraph 1 - right column paragraph 1, page 1727 Scheme 1, page 1729 Scheme 3, page 1730 left column paragraph 2	1-5, 7-14, 16 and 18-22
X	Merbouh, N. et al., "3-Mercaptopropanol as a Traceless Linker for Chemical and Enzymatic Synthesis of Oligosaccharides", Organic Letters, 2007, Vol. 9, No. 4, pages 651-653 & CASREACT Online Abstract Accession No. 146:338090 Page 651 Abstract, page 652 Scheme 1, page 652 Table 1, page 652 Figure 1, Page 652 right column paragraph 1.	1-2, 9-14, 16, 18 and 22
X	Gomez-Garcia, M. et al., "Probing Secondary Carbohydrate-Protein Interactions with Highly Dense Cyclodextrin-Centered Heteroglycoclusters: The Heterocluster Effect", Journal of the American Chemical Society, 2005, Vol. 127, No. 22, pages 7970-7971 & CASREACT Online Abstract Accession No. 143:128624 Page 7970 left column second paragraph, Supporting Information	1, 2, 14, 16 and 22
Form PCT/ISA/210 (fifth sheet) (July 2009)		

INTERNATIONAL SEARCH REPORT Information on patent family members		International application No. PCT/SG2012/000168	
This Annex lists known patent family members relating to the patent documents cited in the above-mentioned international search report. The Australian Patent Office is in no way liable for these particulars which are merely given for the purpose of information.			
Patent Document/s Cited in Search Report		Patent Family Member/s	
Publication Number	Publication Date	Publication Number	Publication Date
End of Annex			