

**(12) STANDARD PATENT  
(19) AUSTRALIAN PATENT OFFICE**

**(11) Application No. AU 2006236649 B2**

(54) Title  
**Methods and compositions for modulating and detecting WISP activity**

(51) International Patent Classification(s)  
**C07K 14/475 (2006.01) A61P 35/00 (2006.01)**  
**A61K 39/395 (2006.01) C07K 16/22 (2006.01)**

(21) Application No: **2006236649** (22) Date of Filing: **2006.04.13**

(87) WIPO No: **WO06/113402**

(30) Priority Data

(31) Number **11/105,876** (32) Date **2005.04.14** (33) Country **US**

(43) Publication Date: **2006.10.26**  
(44) Accepted Journal Date: **2012.03.29**

(71) Applicant(s)  
**Genentech, Inc.**

(72) Inventor(s)  
**Filvaroff, Ellen;Desnoyers, Luc**

(74) Agent / Attorney  
**Griffith Hack, GPO Box 1285, Melbourne, VIC, 3001**

(56) Related Art  
**WO 1999/021998 A1 (GENENTECH, INC.) 6 May 1999**  
**WO 2004/003158 A2 (GENENTECH, INC.) 8 January 2004**

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
26 October 2006 (26.10.2006)

PCT

(10) International Publication Number  
WO 2006/113402 A1

(51) International Patent Classification:

C07K 16/22 (2006.01) A61P 35/00 (2006.01)  
A61K 39/395 (2006.01) C07K 14/475 (2006.01)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(21) International Application Number:

PCT/US2006/014008

(22) International Filing Date: 13 April 2006 (13.04.2006)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
11/105,876 14 April 2005 (14.04.2005) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for all designated States except US): GENENTECH, INC. [US/US]; 1 Dna Way, Ms 49, South San Francisco, California 94080 (US).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

(72) Inventors; and

(75) Inventors/Applicants (for US only): DESNOYERS, Luc [CA/US]; 2050 Stockton Street, San Francisco, California 94133 (US). FILVAROFF, Ellen [US/US]; 538 18th Avenue, San Francisco, California 94121 (US).

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(74) Agent: HASAK, Janet, E. à; c/o GENENTECH, INC., 1 DNA Way, MS 49, South San Francisco, California 94080 (US).

WO 2006/113402 A1

(54) Title: METHODS AND COMPOSITIONS FOR MODULATING AND DETECTING WISP ACTIVITY

(57) Abstract: Methods and compositions for use in modulating the activity(s) of WISP-1 polypeptide are provided. WISP-1 antagonists include anti-WISP-1 antibodies, WISP-1 immunoadhesins and WISP-1 variants (and fusion proteins thereof) which inhibit or neutralize induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell. The invention also provides methods for *in vitro*, *in situ*, and/or *in vivo* diagnosis and/or treatment of mammalian cells or pathological conditions associated with native WISP-1 polypeptides.

## METHODS AND COMPOSITIONS FOR MODULATING AND DETECTING WISP ACTIVITY

Field of the Invention

The present invention relates generally to methods and compositions for use in modulating the activity(s) of WISP polypeptides, particularly, WISP-1 polypeptides. The invention also relates to methods and compositions for *in vitro*, *in situ*, and/or *in vivo* diagnosis and/or treatment of mammalian cells or pathological conditions associated with WISP polypeptides.

10

Background of the Invention

Connective tissue growth factor (CTGF) is a growth factor induced in fibroblasts by many factors, including TGF- $\beta$ , and is essential for the ability of TGF- $\beta$  to induce anchorage-independent growth (AIG), a property of transformed cells. CTGF also is mitogenic and chemotactic for cells, and hence growth factors in this family are believed to play a role in the normal development, growth, and repair of human tissue. Five proteins related to CTGF, including Cyr61, Nov, WISP-1, WISP2, and WISP-3 have been isolated, cloned, sequenced, and characterized as belonging to the CCN gene family. Oemar and Luescher, Arterioscler. Thromb. Vasc. Biol., 17: 1483-1489 (1997); Brigstock, Endocrine Rev., 20:189-206 (1999). The gene encoding Cyr61 has been found to promote angiogenesis, tumor growth, and vascularization. Babic et al., Proc. Natl. Acad. Sci. USA, 95: 6355-6360 (1998). The nov gene is expressed in the kidney essentially at the embryonic stage, and alterations of nov expression, relative to the normal kidney, have been detected in both avian nephroblastomas and human Wilms' tumors. Martinerie et al., Oncogene, 9: 2729-2732 (1994). Wt1 downregulates human nov expression, which downregulation might represent a key element in normal and tumoral nephrogenesis. Martinerie et al., Oncogene, 12: 1479-1492 (1996).

The different members of the CCN family interact with various soluble or matrix associated macromolecules in particular sulfated glycoconjugates (Bork, FEBS Letters, 327:125-130). This interaction was used to purify Cyr61 and CTGF by affinity chromatography on heparin-agarose (Frazier et al., J. Invest. Dermatol., 107:404-411 (1996); Kireeva et al., Mol. Cell. Biol., 16:1326-1334 (1996)). Cyr61 is secreted and associated with both the extracellular matrix and the cell surface due to its affinity for heparan sulfate (Yang et al., Cell. Growth Diff., 2:351-357 (1991)). Recently, WISP-1 was

shown to interact with decorin and biglycan, two secreted dermatan sulfate proteoglycans. (Desnoyers, et al., J. Biol. Chem., 276:47599-47607 (2001)).

The murine protein ELM1 was recently identified in low metastatic cells. Hashimoto et al., J. Exp. Med., 187:289-296 (1998). The *elm1* gene, a mouse orthologue of WISP-1 disclosed below, is another member of the CNN gene family. It suppresses *in vivo* tumor growth and metastasis of K-1735 murine melanoma cells. Another recent article on rCop-1, the rat orthologue of WISP-2 described below, describes the loss of expression of this gene after cell transformation. Zhang et al., Mol. Cell. Biol., 18:6131-6141 (1998).

Wnts are encoded by a large gene family whose members have been found in round worms, insects, cartilaginous fish, and vertebrates.

Holland et al., Dev. Suppl., 125-133 (1994). Wnts are thought to function in a variety of developmental and physiological processes since many diverse species have multiple conserved *Wnt* genes. McMahon, Trends Genet., 8: 236-242 (1992); Nusse and Varmus, Cell, 69: 1073-1087 (1992). *Wnt* genes encode secreted glycoproteins that are thought to function as paracrine or autocrine signals active in several primitive cell types. McMahon, *supra* (1992); Nusse and Varmus, *supra* (1992). The *Wnt* growth factor family includes more than ten genes identified in the mouse (*Wnt-1*, -2, -3A, -3B, -4, -5A, -5B, -6, -7A, -7B, -8A, -8B, -10B, -11, -12, and -13) (see, e.g., Gavin et al., Genes Dev., 4: 2319-2332 (1990); Lee et al., Proc. Natl. Acad. Sci. USA, 92: 2268-2272 (1995); Christiansen et al., Mech. Dev., 51: 341-350 (1995)) and at least nine genes identified in the human (*Wnt-1*, -2, -3, -5A, -7A, -7B, -8B, -10B, and -11) by cDNA cloning. See, e.g., Vant Veer et al., Mol. Cell. Biol., 4: 2532-2534 (1984).

The *Wnt-1* proto-oncogene (*int-1*) was originally identified from mammary tumors induced by mouse mammary tumor virus (MMTV) due to an insertion of viral DNA sequence. Nusse and Varmus, Cell, 31: 99-109 (1982). In adult mice, the expression level of *Wnt-1* mRNA is detected only in the testis during later stages of sperm development. *Wnt-1* protein is about 42 KDa and contains an amino-terminal hydrophobic region, which may function as a signal sequence for secretion (Nusse and Varmus, *supra*, 1992). The expression of *Wnt-2* is detected in mouse fetal and adult tissues and its distribution does not overlap with the expression pattern for *Wnt-1*. *Wnt-3* is associated with mouse mammary tumorigenesis. The expression of *Wnt-3*

in mouse embryos is detected in the neural tubes and in the limb buds. *Wnt-5a* transcripts are detected in the developing fore- and hind limbs at 9.5 through 14.5 days and highest levels are concentrated in apical ectoderm at the distal tip of limbs. Nusse and Varmus, *supra* (1992). Recently, a Wnt growth factor, termed *Wnt-x*, was described (WO95/17416) along with the detection of *Wnt-x* expression in bone tissues and in bone-derived cells. Also described was the role of *Wnt-x* in the maintenance of mature osteoblasts and the use of the *Wnt-x* growth factor as a therapeutic agent or in the development of other therapeutic agents to treat bone-related diseases.

Wnts may play a role in local cell signaling. Peifer and Polakis, *Science*, 287:1606-1609 (2000). Biochemical studies have shown that much of the secreted Wnt protein can be found associated with the cell surface or extracellular matrix rather than freely diffusible in the medium. Papkoff and Schryver, *Mol. Cell. Biol.*, 10: 2723-2730 (1990); Bradley and Brown, *EMBO J.*, 9: 1569-1575 (1990).

Studies of mutations in *Wnt* genes have indicated a role for Wnts in growth control and tissue patterning. In *Drosophila*, *wingless* (*wg*) encodes a Wnt-related gene (Rijsewijk et al., *Cell*, 50: 649-657 (1987)) and *wg* mutations alter the pattern of embryonic ectoderm, neurogenesis, and imaginal disc outgrowth. Morata and Lawerence, *Dev. Biol.*, 56: 227-240 (1977); Baker, *Dev. Biol.*, 125: 96-108 (1988); Klingensmith and Nusse, *Dev. Biol.*, 166: 396-414 (1994). In *Caenorhabditis elegans*, *lin-44* encodes a Wnt homolog which is required for asymmetric cell divisions. Herman and Horvitz, *Development*, 120: 1035-1047 (1994). Knock-out mutations in mice have shown Wnts to be essential for brain development (McMahon and Bradley, *Cell*, 62: 1073-1085 (1990); Thomas and Cappelli, *Nature*, 346: 847-850 (1990)), and the outgrowth of embryonic primordia for kidney (Stark et al., *Nature*, 372: 679-683 (1994)), tail bud (Takada et al., *Genes Dev.*, 8: 174-189 (1994)), and limb bud. Parr and McMahon, *Nature*, 374: 350-353 (1995). Overexpression of *Wnt-1* in the mammary gland can result in mammary hyperplasia (McMahon, *supra* (1992); Nusse and Varmus, *supra* (1992)), precocious alveolar development (Bradbury et al., *Dev. Biol.*, 170: 553-563 (1995)), and mammary adenocarcinomas (Li et al., *Oncogene*, 19:1002-1009 (2000)).

40 *Wnt-5a* and *Wnt-5b* are expressed in the posterior and lateral mesoderm and the extraembryonic mesoderm of the day 7-8 murine embryo. Gavin et al., *supra* (1990). These embryonic domains

contribute to the AGM region and yolk sac tissues from which multipotent hematopoietic precursors and HSCs are derived. Dzierzak and Medvinsky, Trends Genet., 11: 359-366 (1995); Zon et al., in Gluckman and Coulombel, ed., Colloque, INSERM, 235: 17-22 (1995), 5 presented at the Joint International Workshop on Foetal and Neonatal Hematopoiesis and Mechanism of Bone Marrow Failure, Paris France, April 3-6, 1995; Kanatsu and Nishikawa, Development, 122: 823-830 (1996). Wnt-5a, Wnt-10b, and other Wnts have been detected in limb buds, indicating possible roles in the development and patterning of 10 the early bone microenvironment as shown for Wnt-7b. Gavin et al., *supra* (1990); Christiansen et al., Mech. Devel., 51: 341-350 (1995); Parr and McMahon, *supra* (1995).

For a review on Wnt, see Cadigan and Nusse, Genes & Dev., 11: 3286-3305 (1997).

15 Pennica et al., Proc. Natl. Acad. Sci., 95:14717-14722 (1998) describe the cloning and characterization of two genes, *WISP-1* and *WISP-2*, that are up-regulated in the mouse mammary epithelial cell line C57MG transformed by Wnt-1, and a third related gene, *WISP-3*. (See also, WO 99/21998 published May 6, 1999; WO 99/21999 published 20 May 6, 1999). Pennica et al. report that these *WISP* genes may be downstream of Wnt-1 signaling and that aberrant levels of *WISP* expression in colon cancer may play a role in colon tumorigenesis. *WISP-1* has recently been identified as a  $\beta$ -catenin-regulated gene and the characterization of its oncogenic activity demonstrated that 25 *WISP-1* might contribute to  $\beta$ -catenin-mediated tumorigenesis (Xu et al., Gene & Develop., 14:585-595 (2000)). *WISP-1* overexpression in normal rat kidney cells (NRK-49F) induced morphological transformation, accelerated cell growth and enhanced saturation density. In addition, these cells readily form tumors when injected 30 into nude mice suggesting that *WISP-1* may play some role in tumorigenesis (Xu et al., *supra* 2000). *WISP-1* is also overexpressed in transformed human breast cancer cell lines and in about 47% of primary human breast cancer associated with certain advanced features. Xie et al., Cancer Res., 61:8917-8923 (2001); Saxena et 35 al., Mol. Cell Biochem., 228:99-104 (2001); Michaelson et al., Oncogene, 20:5093-5099 (2001). A particular *WISP-1* variant has also been reported to be overexpressed in about 86% of human scirrhous gastric carcinoma cells. Tanaka et al., Oncogene, 20:5525-5532 (2001).

40 Hurvitz et al., Nature Genetics, 23:94-97 (1999) describe a study involving *WISP3* in which nine different mutations of *WISP3* in

unrelated individuals were found to be associated with the autosomal recessive skeletal disorder, progressive pseudorheumatoid dysplasia (PPD). *WISP3* expression by RT-PCR was observed by Hurvitz et al. in human synoviocytes, articular cartilage chondrocytes, and bone-  
5 marrow-derived mesenchymal progenitor cells.

PCT application WO98/21236 published May 22, 1998 discloses a human connective tissue growth factor gene-3 (CTGF-3) encoding a 26 kD member of the growth factor superfamily. WO98/21236 discloses that the CTGF-3 amino acid sequence was deduced from a human  
10 osteoblast cDNA clone, and that CTGF-3 was expressed in multiple tissues like ovary, testis, heart, lung, skeletal muscle, adrenal medulla, adrenal cortex, thymus, prostate, small intestine and colon.

Hyaluronic acid (also referred to as HA, hyaluronate, or hyaluronan) is recognized in the literature as being an important  
15 component of the extracellular matrix (See, e.g., Hardingham et al., *FASEB J.*, 6:861-870 (1992); Laurent et al., *FASEB J.*, 6:2397-2404 (1992)). HA is a component of skin and mesenchymal tissues where it facilitates cell migration during wound healing, inflammation, and embryonic morphogenesis. (Knudson et al., *FASEB J.*, 7:1233-1241  
20 (1993); Knudson et al., *CIBA Found. Symp.*, 143:150-169 (1989)). HA has also been reported to play a role in certain types of metastases.  
25 (Naor et al., CD44: Structure, Function and Association with the Malignant Process, Advances in Cancer Research, Academic Press (1997), pages 241-319). The largest concentrations of HA are found in the skin and musculo-skeletal system which account for over 50% of total body HA. (Banerji et al., *J. Cell Biol.*, 144:789-801 (1999)).

Various investigators have reported on receptors which bind HA. One of the receptors identified for HA is the CD44 protein. (See, e.g., Culty et al., *J. Cell Biology*, 111:2765-2774 (1990); Aruffo et  
30 al., *Cell*, 61:1303-1313 (1990); Naor et al., CD44: Structure, Function and Association with the Malignant Process, Advances in Cancer Research, Academic Press (1997), pages 241-319); Ropponen et al., *Cancer Res.*, 58:342-347 (1998); Masaki et al., *Cancer*, 92:2539-2546 (2001). CD44 is a family of cell-surface glycoproteins  
35 generated from a single gene by alternative splicing and differential glycosylation. (Wielenga et al., *Am. J. Pathology*, 154:515-523 (1999)). CD44 is believed to function as a cell adhesion receptor, linking extracellular molecules, specifically hyaluronate, to the cell and the cytoskeleton (Wielenga et al., *supra*). CD44 is  
40 expressed on epithelial, mesenchymal and lymphoid cells. (Lesley et al., *Adv. Immunol.*, 54:271-335 (1994)). Wielenga et al. report that

CD44 expression may be regulated by the WNT pathway, based on certain experiments analyzing CD44 expression in the intestinal mucosa of mice and humans with genetic defects in either APC or Tcf-4. (Wielenga et al., supra).

5 Other HA receptors characterized to date include RHAMM (also referred to as receptor for hyaluronic acid mediated motility), a 58 kD intracellular protein expressed transiently on the surface of transformed lymphocytes (Hardwick et al., J. Cell Biol., 117:1343-1350 (1992); Turley et al., Exp. Cell Res., 207:277-282 (1993)). RHAMM expression in 0 fibroblasts was reported to promote metastasis and play an important role in H-Ras transformation (Hall et al., infra).

Another receptor which binds HA was described by Banerji et al. (Banerji et al., supra). Banerji et al. report a receptor on lymph vessel walls, referred to as "LYVE-1", which is a 322-residue type I integral 5 membrane polypeptide which has a 41% similarity to the CD44 receptor. Unlike the CD44 receptor for HA, the LYVE-1 protein is absent in blood vessels. In addition, layilin (Bono et al., Mol. Biol. Cell, 12:891-900 (2001)) and HARE (Zhou et al., J. Biol. Chem., 275:37733-37741 (2000)) were also described as HA receptors.

0 It is to be understood that, if any prior art publication is referred to herein, such reference does not constitute an admission that the publication forms a part of the common general knowledge in the art, in Australia or any other country.

#### 5 Summary of the Invention

A first aspect provides a method of inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, comprising exposing said mammalian cells to an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists 30 of a polypeptide selected from the group consisting of:

- a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and
- b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

35 A second aspect provides a method of treating cancer in a mammal, comprising administering to said mammal an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

- a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

A third aspect provides a method of inhibiting cancer cell motility, comprising exposing a cancer cell to an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

A fourth aspect provides a method of inhibiting cancer cell motility in a mammal comprising administering to said mammal an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

A fifth aspect provides use of a WISP-1-domain 1 polypeptide in the manufacture of a medicament for inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

A sixth aspect provides use of a WISP-1-domain 1 polypeptide in the manufacture of a medicament for inhibiting cancer cell motility, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

A seventh aspect provides use of a WISP-1-domain 1 polypeptide in the manufacture of a medicament for treating cancer, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

5 Applicant has surprisingly found that WISP-1 can induce HAS2 (hyaluronan synthase 2), CD44, and RHAMM mRNA expression, CD44 protein synthesis, and HA secretion. The induction or secretion of such molecules may promote or increase cancer cell growth, motility and/or metastatic potential. Disclosed herein are WISP-1 antagonists and methods of using 10 such antagonists. The antagonists described herein find utility for, among other things, *in vitro*, *in situ*, or *in vivo* diagnosis or treatment of mammalian cancer cells or other pathological conditions associated with the induction or secretion of HAS2, HA, CD44 or RHAMM.

Disclosed herein are isolated WISP-1 antagonists. Such antagonists 15 may comprise antibodies, such as WISP-1 antibodies. In preferred embodiments, the antagonists may block or neutralize WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM. Such antagonistic antibodies may, for example, be monoclonal antibodies, chimeric antibodies, humanized antibodies, or human antibodies. The WISP-1 antagonists contemplated for 20 use according to the disclosure include WISP-1 immunoadhesins, WISP-1 variants, covalently modified forms thereof, or fusion proteins thereof. By way of example, such antagonists may include pegylated WISP-1 or WISP-1 fused to heterologous sequences such as epitope tags or leucine zippers. The methods contemplate the use of a single type of antagonist molecule or 25 a combination of two or more types of antagonists.

The methods of the disclosure include methods to treat pathological 30 conditions or diseases in mammals associated with or resulting from WISP-1, including, induction or secretion of HAS2, HA, CD44 or RHAMM by WISP-1. In the methods of treatment, WISP-1 antagonists may be administered to the mammal suffering from such pathological condition or disease. For example, 35 also disclosed is a method comprising exposing a mammalian cell(s), such as a cancer cell(s), to one or more WISP-1 antagonists in an amount effective to decrease, neutralize or block WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM. The cell may be in cell culture or in a mammal, e.g. a mammal suffering from, for instance, cancer.

Also disclosed are compositions which comprise one or more WISP-1 40 antagonists. Optionally, the compositions of the invention will include pharmaceutically acceptable carriers or diluents. Preferably, the compositions will include one or more WISP-1 antagonists in an amount which is therapeutically effective to treat a pathological condition or disease.

Also disclosed are articles of manufacture and kits which include one or more WISP-1 antagonists.

Also disclosed are methods of conducting screening assays to identify candidate molecules, such as small molecule compounds, polypeptides or antibodies, which act as antagonists with respect to blocking or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM.

Also disclosed are isolated WISP-1 antagonists which inhibit or neutralize the induction or secretion of HAS2, HA, CD44 or RHAMM by native WISP-1 polypeptide in at least one type of mammalian cell, said antagonist being selected from the group consisting of an anti-WISP-1 antibody, a WISP-1 immunoadhesin, a WISP-1 variant, and fusion proteins thereof. The antagonist may comprise an anti-WISP-1 antibody which binds native human WISP-1 polypeptide comprising amino acids 23-367 of Figures 9A-9C or one or more domains of WISP-1 comprising amino acids encoded by the sequences of SEQ ID NO:3; 4; 5; 6; 7; 8; 9; 10; or 11 herein. The anti-WISP-1 antibody may be a chimeric, humanized or human antibody.

Also disclosed are compositions comprising the antagonists described herein and a carrier, optionally the carrier is a pharmaceutically-acceptable carrier.

Also disclosed are methods of inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, comprising exposing said mammalian cells to an effective amount of WISP-1 antagonist, wherein said WISP-1 antagonist is selected from the group consisting of:

- a) a WISP-1 immunoadhesin;
- b) a WISP-1 polypeptide linked to a nonproteinaceous polymer selected from the group consisting of polyethylene glycol, polypropylene glycol, and polyoxyalkylene;
- c) a WISP-1 antibody; and
- d) a WISP-1 variant.

The WISP-1 immunoadhesins employed in the methods may comprise a WISP-1 sequence fused to a Fc region of an immunoglobulin. The anti-WISP-1 antibodies employed in the methods may bind native human WISP-1 comprising amino acids 23-367 of Figures 9A-9C, or one or more of the domains of WISP-1 described in the Examples below. In the methods, the mammalian cells may comprise cancer cells, and optionally the cells comprise colon or colorectal cancer cells, breast cancer cells, lung cancer cells, or brain cancer cells (such as glioma or glioblastoma).

Also disclosed are methods of treating cancer in a mammal, comprising administering to said mammal an effective amount of WISP-1 antagonist.

21 Feb 2012  
2006236649

Optionally, in said methods, the antagonist may inhibit or neutralize induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell and said antagonist is selected from the group consisting of an anti-WISP-1 antibody, a WISP-1 immunoadhesin and a WISP-1 variant. Optionally, the antagonist may act to inhibit cancer cell growth or cancer cell metastasis. The cancer in the mammal may comprise colon or colorectal cancer cells, breast cancer cells, lung cancer cells, or brain cancer cells (such as glioma or glioblastoma). Optionally, the antagonist(s) employed in the methods inhibits or reduces cancer cell growth or metastasis. Also, in the methods, chemotherapy, radiation, prodrug, cytotoxic agent, growth inhibitory agent, or cytokine may also be administered to the mammal.

Also disclosed are antibodies which specifically bind to one or more domains of WISP-1 polypeptide (described further in the Examples below) comprising amino acids encoded by the sequences of SEQ ID NO:3; 4; 5; 6; 7; 8; 9; 10; or 11 herein. Optionally, the antibody is a monoclonal antibody. Optionally, the monoclonal antibody comprises the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody secreted by the hybridoma deposited with ATCC as accession number PTA-4624, PTA-4628, PTA-4626, PTA-4625, or PTA-4627, respectively.

Also disclosed are antibodies which bind to the same epitope as the epitope to which the 3D11, 11C2, 9C10, 5D4, or 9C11 monoclonal antibody produced by the hybridoma cell line deposited as ATCC accession number PTA-4624, PTA-4628, PTA-4626, PTA-4625, or PTA-4627, respectively, binds.

Also disclosed is the hybridoma cell line which produces monoclonal antibody 3D11, 11C2, 9C10, 5D4, or 9C11 and deposited with ATCC as accession number PTA-4624, PTA-4628, PTA-4626, PTA-4625, or PTA-4627, respectively, and the monoclonal antibody 3D11, 11C2, 9C10, 5D4, or 9C11 secreted by the hybridoma deposited with ATCC as accession number PTA-4624, PTA-4628, PTA-4626, PTA-4625, or PTA-4627, respectively.

Also disclosed are isolated anti-WISP-1 monoclonal antibodies, comprising antibodies which bind to WISP-1 polypeptide and competitively inhibit binding of the monoclonal antibody produced by the hybridoma deposited as ATCC accession no. PTA-4624, PTA-4628, PTA-4626, PTA-4625, or PTA-4627 to said WISP-1 polypeptide.

Also disclosed are chimeric or humanized anti-WISP-1 antibodies which specifically bind to WISP-1 polypeptide and comprise (a) a sequence derived from the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody secreted by the hybridoma deposited with ATCC as accession number PTA-4624, PTA-4628, PTA-4626, PTA-4625, or PTA-4627, respectively. Optionally, such antibodies may comprise

a heavy chain, light chain or variable regions derived from the 3D11, 11C2, 9C10, 5D4, or 9C11 antibody.

The anti-WISP-1 antibodies may be linked to one or more non-proteinaceous polymers selected from the group consisting of polyethylene glycol, polypropylene glycol, and polyoxyalkylene, or to a cytotoxic agent or enzyme, or to a radioisotope, fluorescent compound or chemiluminescent compound.

Brief Description of the Drawings

Figures 1A-1F show WISP-1 promotion of hyaluronan production.

5 Particle exclusion assay demonstrating a hyaluronan coat at the surface of NRK/WISP-1H (A) and NRK/WISP-1L (B) but absent from NRK/control (C). NRK/WISP-1H (D) and NRK/control (E) were stained for hyaluronan with bHABP. F, Time course of HA accumulation in NRK/WISP-1H and NRK/control media.

10 Figures 2A-2C show WISP-1 increases HAS2, CD44 and RHAMM mRNA expression and CD44 protein expression. A, Real time RT-PCR analysis of HAS1, HAS2, HAS3, CD44, RHAMM and hyaluronidase (Hyal) mRNA expression in NRK/WISP-1H, NRK/WISP-1L and NRK/control cells. Results are given as fold induction over the expression in

15 NRK/control cells. B, Flow cytometry analysis of CD44 expression in NRK/WISP-1H and NRK/control cells. The shaded area represents the fluorescence intensity of the secondary antibody alone. C, Western blot analysis of CD44 protein in NRK/WISP-1H and NRK/control cells. Actin staining was used as a loading control.

20 Figures 3A-3G show WISP-1 expression increases cell motility and modifies cellular morphology. NRK/control cells (A) formed defined colonies when plated at low density whereas NRK/WISP-1L (B) and NRK/WISP-1H cells (C) scattered. NRK/WISP-1H (E) showed a de-differentiated spindloid morphology with lamelipodia compared to

25 NRK/control cells (D). F, The random migration of NRK/WISP-1H and NRK/control cells was measured over 15 hours using time lapse microscopy. Results represent the typical average migration distance of cells in one field. G, Motility of NRK/WISP-1H and NRK/control cells was evaluated by a cell wound healing assay and measured after

30 15 hours using time lapse microscopy. Data represent the results of a typical experiment.

Figures 4A-4C show WISP-1 addition induces HAS2 mRNA expression and haptotactic migration of NRK cells. A, Real time RT-PCR analysis of HAS2 and CD44 expression in NRK cells seeded on coated surface.

35 In certain cases the coated substrates were further incubated with WISP-1 (treatment). Results are given in fold induction over expression in NRK cells plated on non-coated, non-treated surface. The underside of modified Boyden chamber filters were coated and NRK cells (B) or SW480 cells (C) were added to the upper chamber. In

40 certain cases, additions were made directly to the lower chamber.

Cell motility was evaluated by counting the cells that migrated to the lower side of the insert.

Figures 5A-5E show WISP-1, HAS2 and CD44 over-expression and increased cell motility in C57MG/Wnt-1 cells and MMTV-Wnt-1 transgenic mice mammary tumors. A, Real time RT-PCR analysis of WISP-1, HAS2 and CD44 expression in C57MG/Wnt-1 and C57MG/control cells. Results are given as fold induction over expression in C57MG/control cells. B, Western blot analysis of CD44 content in C57MG/Wnt-1 and C57MG/control cells. Actin staining was used as a loading control. C, C57MG/control cells formed distinct colonies when plated at low density whereas C57MG/Wnt-1 cells scattered. Semi-quantitative RT-PCR analysis of HAS2 (D) and CD44 (E) in mammary tumors of MMTV-Wnt-1 transgenic mice. Results are expressed as relative fold induction over the expression in normal mammary gland tissue.

Figures 6A-6H show WISP-1, HAS2 and CD44 expression and HA accumulation in MMTV-Wnt-1 transgenic mammary tumors. In situ hybridization of WISP-1 (A, B) HAS2 (C, D) and CD44 (E, F), immunohistochemistry of CD44 (G) and bHABP fluorescent staining of hyaluronan (H) in MMTV-Wnt-1 transgenic mouse mammary tumors. t, tumor; s, stroma.

Figures 7A-7L show WISP-1 expression promotes metastatic lung colonization and tumor growth. The effect of WISP-1 was analyzed by injecting NRK/control, NRK/WISP-1L or NRK/WISP-1H cells in the tail vein of nude mice. At various times after injection, lungs were imaged by MRI (A, D, G, J), excised and their gross appearance (B, E, H, K) and histological features (C, F, I, L) were recorded. The severity of lung invasion by tumors was graded as normal (A, B, C); grade I (D, E, F); grade II (G, H, I) or grade III (J, K, L).

Figures 8A-B show CD44 antibody prevents NRK/WISP-1H cell metastasis. NRK/WISP-1H cells ( $2.5 \times 10^5$  cells) were injected in the tail vein of nude mice. Mice were treated twice a week with 10 mg/kg of a CD44 antibody, an isotype control antibody or with buffer only (PBS). The lungs were excised after four weeks for gross anatomical analysis. Picture of a normal lung is shown for comparison.

Figures 9A-9C show the nucleotide (SEQ ID NO:2) and putative amino acid sequence (SEQ ID NO:1) of native human WISP-1.

Figure 10 shows the effects of WISP-1 expression on NRK metastatic potential.

Figures 11A-B show the binding properties of WISP-1 antibodies to various WISP-1 constructs and preparations.

Figures 12A-G show diagrams of various domains of WISP-1 polypeptide and the results of assays conducted to identify the epitopes recognized by the WISP-1 antibodies 11C2, 5D4, 9C11, and 3D11, respectively. Figure 12C illustrates the amino acid sequence 5 provided in SEQ ID NO: 20.

Figure 13 shows the results on an assay conducted to identify the epitope recognized by the WISP-1 antibody 9C10.

Figure 14 shows the results of an ELISA demonstrating detection of WISP-1 by WISP-1 antibodies.

10 Figure 15 shows the results of an ELISA binding assay demonstrating that WISP-1 antibodies recognizing the variable region of WISP-1 can inhibit WISP-1 binding to heparin.

15 Figures 16A-B show the results of an assay detecting inhibition of haptotaxis of NRK cells by WISP-1 antibodies (16A) and a chart summarizing the properties and characteristics of WISP-1 antibodies (16B).

20 Figures 17A-E show the results of an in vivo study of the effects of WISP-1 antibodies. After 3 weeks, the severity of the lesions found in WISP-1 antibody treated animals was greatly attenuated compared to control (Fig. 17 a, b). The number of nodules and the average area of the metastatic foci found in mice treated with WISP-1 antibodies (n=5) were reduced compared to animals treated with a control antibody (Fig. 17 c,d). The total pulmonary area covered by the lesions was reduced by 82-97% compared to animal 25 treated with an isotype control antibody (Fig. 17e).

Figure 18 is a bar diagram showing expression of WISP-1 in 4T1/control, 4T1/WISP-1L, 4T1/WISP-1H, NRK/control, NRK/WISP-1L, and WISP-1H cell lines as measured by semi-quantitative RT-PCR (Taqman).

30 Figure 19 shows the results of an in vitro assay on colony formation of 4T1/control, 4T1/WISP-1L, and 4T1/WISP-1H cell lines.

Figure 20 shows the results of an in vitro assay measuring invasion index using a Matrigel modified Boyden chamber system and 4T1/control, 4T1/WISP-1L, and 4T1/WISP-1H cell lines.

35 Figures 21A-B show effects of WISP-1 expression on mammary epithelial cell tumorigenesis. The effect of WISP-1 was analyzed by injecting 4T1/control 1, 4T1/control 2, 4T1/WISP-1L or 4T1/WISP-1H cells in Balb/C mice. Tumor volume (21A) and tumor weight (21 B) are reported.

40 Figure 22 is a bar diagram illustrating the relative expression of HAS2 and CD44 in tumors formed by the inoculated 4T1/control, 4T1/WISP-1L and 4T1/WISP-1H cells.

Figures 23A-F illustrate the WISP-1 effects on mammary epithelial cells metastasis evaluated in vivo by inoculating 4T1 cells in mice mammary fat pads and examining the extent of the metastatic propagation by micro computer tomography and histology.

5 After 31 days, the mice inoculated with 4T1/WISP-1L or 4T1/WISP-1H cells had extensive lung metastasis (Fig. 23b and 23d) compared to the 4T1/control injected mice (Fig. 23a and 23c). Using immunohistochemistry, it was also observed that the 4T1/WISP-1 pulmonary metastatic foci expressed high levels of CD44 (Fig. 23e).  
10 In these tumors, CD44 was localized at the plasma membrane of the 4T1/WISP-1 cells (Fig. 23f).

Figure 24A shows WISP-1 expression in NRK/control, NRK/WISP-1\_1234, NRK/WISP-1\_134, NRK/WISP-1\_234, NRK/WISP-1\_123 and NRK/WISP-1\_124 cells lines, as measured by semi-quantitative RT-PCR (Taqman);  
15 the results are expressed as fold relative to WISP-1 expression in NRK/WISP-1\_1234. Figure 24B illustrates the effects of these constructs on tumorigenesis in vivo; the results are shown by the effects on tumor volume.

Figure 25 shows the histological analysis of the excised tumors referred to in Figure 24 and revealed that the neoplastic cells from NRK/WISP-1\_234 tumors were phenotypically similar to the neoplastic cells from NRK/WISP-1\_1234 tumors. In these tumors, cells appeared fibroblastic, differentiated and spindloid. The neoplastic cells from NRK/WISP-1\_134 tumors were phenotypically different and appeared less differentiated, frequently multinucleated and had a brisk mitotic rate as demonstrated by the presence of multiple mitotic figures (arrows).  
20

Figure 26 shows the histological analysis of the excised tumors in the NRK/WISP-1\_134 xenograft referred to in Figure 24, and reveals the neoplastic cells invaded several blood vessels adjacent to the tumor.  
30

Figure 27 shows expression as fold invasion compared to a control in which HEK 293 cells transfected with an empty vector were plated in the modified Boyden chamber system. 4T1 cells demonstrated a 4 fold increase invasion when HEK 293 cells expressing constructs WISP-1\_1234, WISP-1\_134 or WISP-1\_123 were seeded in the lower chamber.  
35

Figures 28A-28B illustrate the effects of soluble WISP-1-domain-1 on WISP-1 promoted 4T1 cell invasion, as evaluated using a Matrigel coated modified Boyden chamber system. The results are expressed as relative fold invasion compared to 4T1/ control cells.  
40

21 Feb 2012

2006236649

WISP-1-domain-1-His and WISP-1-domain-1-Fc demonstrated a dose dependent antagonistic activity toward WISP-1 promoted 4T1 invasion.

Figure 29 illustrates an *in situ* hybridization analysis of tissues from the HPAC xenograft model of human pancreatic adenocarcinomas to evaluate WISP-1 expression.

Figure 30 shows a graph illustrating the effects of WISP-1 constructs on tumor volume in a nude mouse model.

Figure 31 shows *in situ* hybridization analyses evaluating WISP-1 expression in primary human pancreatic adenocarcinomas.

Figure 32 shows immunohistochemical staining evaluating WISP-1 expression in primary colon adenocarcinomas.

#### Detailed Description of the Invention

##### I. Definitions

In the claims which follow and in the description of the invention, except where the context requires otherwise due to express language or necessary implication, the word "comprise" or variations such as "comprises" or "comprising" is used in an inclusive sense, i.e. to specify the presence of the stated features but not to preclude the presence or addition of further features in various embodiments of the invention.

The term "WISP polypeptide" refers to the family of native-sequence human and mouse WISP proteins and variants described herein whose genes are induced at least by Wnt-1. This term includes WISP-1 and variants thereof.

Such WISP-1 proteins are described further below and in PCT application WO99/21998 published May 6, 1999, US 6,387,657 B1 issued May 14, 2002, and in Pennica et al., *Proc. Natl. Acad. Sci.*, 95:14717-14722 (1998).

The terms "WISP-1 polypeptide", "WISP-1 homologue" and grammatical variants thereof, as used herein, encompass native- sequence WISP-1 protein and variants (which are further defined herein). The WISP-1 polypeptide may be isolated from a variety of sources, such as from human tissue types or from another source, or prepared by recombinant or synthetic methods, or by any combination of these and similar techniques.

The term "native-sequence WISP-1 polypeptide" comprises a polypeptide having the same amino acid sequence as a WISP-1 polypeptide derived from nature. Such native-sequence WISP-1 polypeptides can be isolated from nature or can be produced by recombinant or synthetic means. The term "native-sequence WISP-1 polypeptide" specifically encompasses naturally occurring truncated or secreted forms of a WISP-1 polypeptide disclosed herein, naturally occurring variant forms (e.g., alternatively spliced

forms or splice variants), and naturally occurring allelic variants of a WISP-1 polypeptide. In one embodiment of the invention, the native-sequence WISP-1 polypeptide is a mature or full-length native-sequence human WISP-1 polypeptide comprising amino acids 23 to 367 of Figure 9 or

amino acids 1 to 367 of Figure 9, respectively, with or without N-terminal methionine.

In another embodiment of the invention, the native-sequence WISP-1 polypeptide is the full-length or mature native-sequence human 5 WISP-1 polypeptide comprising amino acids 23 to 367 or 1 to 367 of Figure 9 wherein the valine residue at position 184 or the alanine residue at position 202 has/have been changed to an isoleucine or serine residue, respectively, with or without N-terminal methionine.

In another embodiment of the invention, the native-sequence WISP-1 10 polypeptide is the full-length or mature native-sequence human WISP-1 polypeptide comprising amino acids 23 to 367 or 1 to 367 of the Figure 9 wherein the valine residue at position 184 and the alanine residue at position 202 has/have been changed to an isoleucine or serine residue, respectively, with or without N-terminal methionine.

15

In another embodiment of the invention, the native-sequence WISP-1 polypeptide is one which is encoded by a nucleotide sequence comprising one of the human WISP-1 splice or other native-sequence variants, including SEQ ID NOS:23, 24, 25, 26, 27, 28, or 29 shown in 20 WO99/21998, with or without an N-terminal methionine.

The term "WISP-1 variant" means an active WISP-1 polypeptide as defined below having at least about 80%, preferably at least about 85%, 86%, 87%, 88%, 89%, more preferably at least about 90%, 91%, 92%, 93%, 94%, most preferably at least about 95%, 96%, 97%, 98%, or 25 99% amino acid sequence identity with human mature WISP-1 having the deduced amino acid sequence shown in Fig. 9, or a soluble fragment thereof. Such variants include, for instance, WISP-1 polypeptides wherein one or more amino acid residues are added to, or deleted from, the N- or C-terminus of the full-length or mature sequences of 30 Figure 9, or WISP-1 polypeptides wherein one or more amino acid residues are inserted or deleted from the internal sequence or domains of the polypeptide, including variants from other species, but excludes a native-sequence WISP-1 polypeptide. Preferably such a variant acts as an antagonist, as defined below.

35 An "extracellular domain", "ECD" or "secreted" protein refers to a form of a polypeptide which is essentially free of any transmembrane and cytoplasmic domains. A "secreted" form of a protein or polypeptide may or may not include a signal sequence.

40 "Stringent conditions" are those that (1) employ low ionic strength and high temperature for washing, for example, 0.015 M sodium chloride/0.0015 M sodium citrate/0.1% sodium dodecyl sulfate

at 50°C; (2) employ during hybridization a denaturing agent, such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM sodium chloride, 75 mM sodium citrate at 42°C; (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 0.075 M sodium citrate), 50 mM sodium phosphate (pH 6.8), 0.1% sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 µg/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS; or (4) employ a buffer of 10% dextran sulfate, 2 x SSC (sodium chloride/sodium citrate), and 50% formamide at 55°C, followed by a high-stringency wash consisting of 0.1 x SSC containing EDTA at 55°C.

"Moderately stringent conditions" are described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989), and include the use of a washing solution and hybridization conditions (e.g., temperature, ionic strength, and percent SDS) less stringent than described above. An example of moderately stringent conditions is a condition such as overnight incubation at 37°C in a solution comprising: 20% formamide, 5 x SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5 x Denhardt's solution, 10% dextran sulfate, and 20 mg/mL denatured sheared salmon sperm DNA, followed by washing the filters in 1 x SSC at about 37-50°C. The skilled artisan will recognize how to adjust the temperature, ionic strength, etc., as necessary to accommodate factors such as probe length and the like.

"Isolated," when used to describe the various polypeptides disclosed herein, means polypeptide that has been identified and separated and/or recovered from a component of its natural environment. Contaminant components of its natural environment are materials that would typically interfere with diagnostic or therapeutic uses for the polypeptide, and may include enzymes, hormones, and other proteinaceous or non-proteinaceous solutes. In preferred embodiments, the polypeptide will be purified (1) to a degree sufficient to obtain at least 15 residues of N-terminal or internal amino acid sequence by use of a spinning cup sequenator, or (2) to homogeneity by SDS-PAGE under non-reducing or reducing conditions using Coomassie blue or, preferably, silver stain. Isolated polypeptide includes polypeptide *in situ* within recombinant cells, since at least one component of the WISP natural environment will not be present. Ordinarily, however, isolated polypeptide will be prepared by at least one purification step.

The term "control sequences" refers to DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotes, for example, include a promoter, optionally an 5 operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

Nucleic acid is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For 10 example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate 15 translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not 20 exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

A "liposome" is a small vesicle composed of various types of lipids, phospholipids and/or surfactant which is useful for delivery 25 of a drug (such as the insulin and insulin variants disclosed herein) to a mammal. The components of the liposome are commonly arranged in a bilayer formation, similar to the lipid arrangement of biological membranes.

The terms "amino acid" and "amino acids" refer to all naturally occurring L-alpha-amino acids. This definition is meant to include 30 norleucine, ornithine, and homocysteine. The amino acids are identified by either the single-letter or three-letter designations:

Asp	D	aspartic acid	Ile	I	isoleucine	
Thr	T	threonine	Leu	L	leucine	
Ser	S	serine	Tyr	Y	tyrosine	
Glu	E	glutamic acid	Phe	F	phenylalanine	
35	Pro	proline	His	H	histidine	
Gly	G	glycine	Lys	K	lysine	
Ala	A	alanine	Arg	R	arginine	
Cys	C	cysteine	Trp	W	tryptophan	
40	Val	V	valine	Gln	Q	glutamine
Met	M	methionine	Asn	N	asparagine	

5        In the Sequence Listing and Figures, certain other single-letter or three-letter designations may be employed to refer to and identify two or more amino acids or nucleotides at a given position in the sequence.

      "Percent (%) amino acid sequence identity" with respect to the polypeptide sequences identified herein is defined as the percentage of amino acid residues in a candidate sequence that are identical with the amino acid residues in such a polypeptide sequence  
10      identified herein, after aligning the sequences and introducing gaps, if necessary, to achieve the maximum percent sequence identity, and not considering any conservative substitutions as part of the sequence identity. Alignment for purposes of determining percent amino acid sequence identity can be achieved in various ways that are  
15      within the skill in the art, for instance, using publicly available computer software such as BLAST, BLAST-2, ALIGN, ALIGN-2 or Megalign (DNASTAR) software. Those skilled in the art can determine appropriate parameters for measuring alignment, including any algorithms needed to achieve maximal alignment over the full-length  
20      of the sequences being compared. For purposes herein, however, % amino acid sequence identity values are obtained as described below by using the sequence comparison computer program ALIGN-2, wherein the complete source code for the ALIGN-2 program is provided in the table below. The ALIGN-2 sequence comparison computer program was  
25      authored by Genentech, Inc. and the source code shown in the table below has been filed with user documentation in the U.S. Copyright Office, Washington D.C., 20559, where it is registered under U.S. Copyright Registration No. TXU510087. The ALIGN-2 program is publicly available through Genentech, Inc., South San Francisco,  
30      California or may be compiled from the source code provided in the table below. The ALIGN-2 program should be compiled for use on a UNIX operating system, preferably digital UNIX V4.0D. All sequence comparison parameters are set by the ALIGN-2 program and do not vary.  
      The term "epitope tagged" when used herein refers to a chimeric  
35      polypeptide comprising a polypeptide fused to a "tag polypeptide". The tag polypeptide has enough residues to provide an epitope against which an antibody can be made. The tag polypeptide preferably also is fairly unique so that the antibody does not substantially cross-react with other epitopes. Suitable tag polypeptides generally have  
40      at least six amino acid residues and usually between about 8 and 50

amino acid residues (preferably, between about 10 and 20 amino acid residues).

As used herein, the term "immunoadhesin" designates antibody-like molecules which combine the binding specificity of a heterologous protein (an "adhesin") with the effector functions of immunoglobulin constant domains. Structurally, the immunoadhesins comprise a fusion of an amino acid sequence with the desired binding specificity which is other than the antigen recognition and binding site of an antibody (i.e., is "heterologous"), and an immunoglobulin constant domain sequence. The adhesin part of an immunoadhesin molecule typically is a contiguous amino acid sequence comprising at least the binding site of a receptor or a ligand. The immunoglobulin constant domain sequence in the immunoadhesin may be obtained from any immunoglobulin, such as IgG-1, IgG-2, IgG-3, or IgG-4 subtypes, IgA (including IgA-1 and IgA-2), IgE, IgD or IgM.

The term "antagonist" is used in the broadest sense, and includes any molecule that partially or fully blocks, inhibits, or neutralizes one or more biological activities of WISP-1 polypeptide, *in vitro*, *in situ*, or *in vivo*. Examples of such biological activities of WISP-1 polypeptides include induction or secretion of HAS2, HA, CD44 or RHAMM in at least one type of mammalian cell. An antagonist may function in a direct or indirect manner. For instance, the antagonist may function to partially or fully block, inhibit or neutralize one or more biological activities of WISP-1 polypeptide, *in vitro*, *in situ*, or *in vivo*, e.g., as a result of its direct binding to WISP-1 polypeptide. The antagonist may also function indirectly to partially or fully block, inhibit or neutralize one or more biological activities of WISP-1 polypeptide, *in vitro*, *in situ*, or *in vivo* as a result of, e.g., blocking or inhibiting another effector molecule.

The term "WISP-1 antagonist" refers to any molecule that partially or fully blocks, inhibits, or neutralizes a biological activity of WISP-1 and include but are not limited to, antibodies, immunoadhesins, WISP-1 immunoadhesins, WISP-1 fusion proteins, covalently modified forms of WISP-1, WISP-1 variants and fusion proteins thereof, WISP-1 antibodies, and higher oligomer forms of WISP-1 (dimers, aggregates) or homo- or heteropolymer forms of WISP-1. To determine whether a WISP-1 antagonist molecule partially or fully blocks, inhibits or neutralizes a biological activity of WISP-1, assays may be conducted to assess the effect(s) of the antagonist molecule on, for example, various cells (as described in the Examples)

or in an *in vivo* murine model of lung cancer metastasis. The various assays may be conducted in known *in vitro* or *in vivo* assay formats. Preferably, the WISP-1 antagonists employed in the methods described herein will be capable of blocking or neutralizing at least one type 5 of WISP-1 activity, which may optionally be determined in assays such as described herein.

The term "antibody" is used in the broadest sense and specifically covers, for example, single monoclonal antibodies, antibody compositions with polyepitopic specificity, single chain 10 antibodies, and fragments of antibodies. "Antibody" as used herein includes intact immunoglobulin or antibody molecules, polyclonal antibodies, multispecific antibodies (i.e., bispecific antibodies formed from at least two intact antibodies) and immunoglobulin fragments (such as Fab, F(ab')<sub>2</sub>, or Fv), so long as they exhibit any of 15 the desired antagonistic properties described herein.

Antibodies are typically proteins or polypeptides which exhibit binding specificity to a specific antigen. Native antibodies are usually heterotetrameric glycoproteins, composed of two identical light (L) chains and two identical heavy (H) chains. Typically, each 20 light chain is linked to a heavy chain by one covalent disulfide bond, while the number of disulfide linkages varies between the heavy chains of different immunoglobulin isotypes. Each heavy and light chain also has regularly spaced intrachain disulfide bridges. Each heavy chain has at one end a variable domain (V<sub>H</sub>) followed by a number of constant 25 domains. Each light chain has a variable domain at one end (V<sub>L</sub>) and a constant domain at its other end; the constant domain of the light chain is aligned with the first constant domain of the heavy chain, and the light chain variable domain is aligned with the variable domain of the heavy chain. Particular amino acid residues are 30 believed to form an interface between the light and heavy chain variable domains [Chothia et al., J. Mol. Biol., 186:651-663 (1985); Novotny and Haber, Proc. Natl. Acad. Sci. USA, 82:4592-4596 (1985)]. The light chains of antibodies from any vertebrate species can be assigned to one of two clearly distinct types, called kappa and 35 lambda, based on the amino acid sequences of their constant domains. Depending on the amino acid sequence of the constant domain of their heavy chains, immunoglobulins can be assigned to different classes. There are five major classes of immunoglobulins: IgA, IgD, IgE, IgG and IgM, and several of these may be further divided into subclasses 40 (isotypes), e.g., IgG-1, IgG-2, IgG-3, and IgG-4; IgA-1 and IgA-2. The heavy chain constant domains that correspond to the different classes

of immunoglobulins are called alpha, delta, epsilon, gamma, and mu, respectively.

"Antibody fragments" comprise a portion of an intact antibody, generally the antigen binding or variable region of the intact 5 antibody. Examples of antibody fragments include Fab, Fab', F(ab')<sub>2</sub>, and Fv fragments, diabodies, single chain antibody molecules, and multispecific antibodies formed from antibody fragments.

The term "variable" is used herein to describe certain portions 10 of the variable domains which differ in sequence among antibodies and are used in the binding and specificity of each particular antibody for its particular antigen. However, the variability is not usually evenly distributed through the variable domains of antibodies. It is typically concentrated in three segments called complementarity 15 determining regions (CDRs) or hypervariable regions both in the light chain and the heavy chain variable domains. The more highly conserved portions of the variable domains are called the framework (FR). The variable domains of native heavy and light chains each comprise four FR regions, largely adopting a  $\beta$ -sheet configuration, connected by three CDRs, which form loops connecting, and in some cases forming 20 part of, the  $\beta$ -sheet structure. The CDRs in each chain are held together in close proximity by the FR regions and, with the CDRs from the other chain, contribute to the formation of the antigen binding site of antibodies [see Kabat, E.A. et al., Sequences of Proteins of Immunological Interest, National Institutes of Health, Bethesda, MD 25 (1987)]. The constant domains are not involved directly in binding an antibody to an antigen, but exhibit various effector functions, such as participation of the antibody in antibody-dependent cellular toxicity.

The term "monoclonal antibody" as used herein refers to an 30 antibody obtained from a population of substantially homogeneous antibodies, i.e., the individual antibodies comprising the population are identical except for possible naturally-occurring mutations that may be present in minor amounts. Monoclonal antibodies are highly specific, being directed against a single antigenic site. 35 Furthermore, in contrast to conventional (polyclonal) antibody preparations which typically include different antibodies directed against different determinants (epitopes), each monoclonal antibody is directed against a single determinant on the antigen.

The monoclonal antibodies herein include chimeric, hybrid and 40 recombinant antibodies produced by splicing a variable (including

hypervariable) domain of the antibody of interest with a constant domain (e.g. "humanized" antibodies), or a light chain with a heavy chain, or a chain from one species with a chain from another species, or fusions with heterologous proteins, regardless of species of origin 5 or immunoglobulin class or subclass designation, as well as antibody fragments (e.g., Fab, F(ab')<sub>2</sub>, and Fv), so long as they exhibit the desired biological activity or properties. See, e.g. U.S. Pat. No. 4,816,567 and Mage et al., in Monoclonal Antibody Production Techniques and Applications, pp.79-97 (Marcel Dekker, Inc.: New York, 10 1987).

Thus, the modifier "monoclonal" indicates the character of the antibody as being obtained from a substantially homogeneous population of antibodies, and is not to be construed as requiring production of the antibody by any particular method. For example, the monoclonal 15 antibodies to be used in accordance with the present invention may be made by the hybridoma method first described by Kohler and Milstein, Nature, 256:495 (1975), or may be made by recombinant DNA methods such as described in U.S. Pat. No. 4,816,567. The "monoclonal antibodies" may also be isolated from phage libraries generated using the 20 techniques described in McCafferty et al., Nature, 348:552-554 (1990), for example.

"Humanized" forms of non-human (e.g. murine) antibodies are specific chimeric immunoglobulins, immunoglobulin chains, or fragments thereof (such as Fv, Fab, Fab', F(ab')<sub>2</sub> or other antigen-binding 25 subsequences of antibodies) which contain minimal sequence derived from non-human immunoglobulin. For the most part, humanized antibodies are human immunoglobulins (recipient antibody) in which residues from a complementary determining region (CDR) of the recipient are replaced by residues from a CDR of a non-human species 30 (donor antibody) such as mouse, rat, or rabbit having the desired specificity, affinity, and capacity. In some instances, Fv framework region (FR) residues of the human immunoglobulin are replaced by corresponding non-human residues. Furthermore, the humanized antibody may comprise residues which are found neither in the recipient 35 antibody nor in the imported CDR or framework sequences. These modifications are made to further refine and optimize antibody performance. In general, the humanized antibody will comprise substantially all of at least one, and typically two, variable domains, in which all or substantially all of the CDR regions 40 correspond to those of a non-human immunoglobulin and all or substantially all of the FR regions are those of a human

immunoglobulin consensus sequence. The humanized antibody optimally also will comprise at least a portion of an immunoglobulin constant region or domain (Fc), typically that of a human immunoglobulin.

A "human antibody" is one which possesses an amino acid sequence which corresponds to that of an antibody produced by a human and/or has been made using any of the techniques for making human antibodies known in the art or as disclosed herein. This definition of a human antibody includes antibodies comprising at least one human heavy chain polypeptide or at least one human light chain polypeptide, for example an antibody comprising murine light chain and human heavy chain polypeptides. Human antibodies can be produced using various techniques known in the art. In one embodiment, the human antibody is selected from a phage library, where that phage library expresses human antibodies (Vaughan et al. Nature Biotechnology, 14:309-314 (1996); Sheets et al. PNAS, (USA) 95:6157-6162 (1998)); Hoogenboom and Winter, J. Mol. Biol., 227:381 (1991); Marks et al., J. Mol. Biol., 222:581 (1991)). Human antibodies can also be made by introducing human immunoglobulin loci into transgenic animals, e.g., mice in which the endogenous immunoglobulin genes have been partially or completely inactivated. Upon challenge, human antibody production is observed, which closely resembles that seen in humans in all respects, including gene rearrangement, assembly, and antibody repertoire. This approach is described, for example, in U.S. Patent Nos. 5,545,807; 5,545,806; 5,569,825; 5,625,126; 5,633,425; 5,661,016, and in the following scientific publications: Marks et al., Bio/Technology, 10: 779-783 (1992); Lonberg et al., Nature, 368: 856-859 (1994); Morrison, Nature, 368:812-13 (1994); Fishwild et al., Nature Biotechnology, 14: 845-51 (1996); Neuberger, Nature Biotechnology, 14: 826 (1996); Lonberg and Huszar, Intern. Rev. Immunol., 13:65-93 (1995). Alternatively, the human antibody may be prepared via immortalization of human B lymphocytes producing an antibody directed against a target antigen (such B lymphocytes may be recovered from an individual or may have been immunized *in vitro*). See, e.g., Cole et al., Monoclonal Antibodies and Cancer Therapy, Alan R. Liss, p. 77 (1985); Boerner et al., J. Immunol., 147 (1):86-95 (1991); and US Pat No. 5,750,373.

The term "Fc region" is used to define the C-terminal region of an immunoglobulin heavy chain which may be generated by papain digestion of an intact antibody. The Fc region may be a native sequence Fc region or a variant Fc region. Although the boundaries of the Fc region of an immunoglobulin heavy chain might vary, the

human IgG heavy chain Fc region is usually defined to stretch from an amino acid residue at about position Cys226, or from about position Pro230, to the carboxyl-terminus of the Fc region (using herein the numbering system according to Kabat et al., *supra*). The Fc region of 5 an immunoglobulin generally comprises two constant domains, a CH2 domain and a CH3 domain, and optionally comprises a CH4 domain.

By "Fc region chain" herein is meant one of the two polypeptide chains of an Fc region.

The "CH2 domain" of a human IgG Fc region (also referred to as 10 "C $\gamma$ 2" domain) usually extends from an amino acid residue at about position 231 to an amino acid residue at about position 340. The CH2 domain is unique in that it is not closely paired with another domain. Rather, two N-linked branched carbohydrate chains are 15 interposed between the two CH2 domains of an intact native IgG molecule. It has been speculated that the carbohydrate may provide a substitute for the domain-domain pairing and help stabilize the CH2 domain. Burton, *Molec. Immunol.* 22:161-206 (1985). The CH2 domain herein may be a native sequence CH2 domain or variant CH2 domain.

The "CH3 domain" comprises the stretch of residues C-terminal 20 to a CH2 domain in an Fc region (i.e. from an amino acid residue at about position 341 to an amino acid residue at about position 447 of an IgG). The CH3 region herein may be a native sequence CH3 domain or a variant CH3 domain (e.g. a CH3 domain with an introduced 25 "protoberance" in one chain thereof and a corresponding introduced "cavity" in the other chain thereof; see US Patent No. 5,821,333). Such variant CH3 domains may be used to make multispecific (e.g. bispecific) antibodies as herein described.

"Hinge region" is generally defined as stretching from about 30 Glu216, or about Cys226, to about Pro230 of human IgG1 (Burton, *Molec. Immunol.* 22:161-206 (1985)). Hinge regions of other IgG isotypes may be aligned with the IgG1 sequence by placing the first 35 and last cysteine residues forming inter-heavy chain S-S bonds in the same positions. The hinge region herein may be a native sequence hinge region or a variant hinge region. The two polypeptide chains of a variant hinge region generally retain at least one cysteine residue per polypeptide chain, so that the two polypeptide chains of the variant hinge region can form a disulfide bond between the two chains. The preferred hinge region herein is a native sequence human hinge region, e.g. a native sequence human IgG1 hinge region.

A "functional Fc region" possesses at least one "effector function" of a native sequence Fc region. Exemplary "effector functions" include C1q binding; complement dependent cytotoxicity (CDC); Fc receptor binding; antibody-dependent cell-mediated cytotoxicity (ADCC); phagocytosis; down regulation of cell surface receptors (e.g. B cell receptor; BCR), etc. Such effector functions generally require the Fc region to be combined with a binding domain (e.g. an antibody variable domain) and can be assessed using various assays known in the art for evaluating such antibody effector functions.

A "native sequence Fc region" comprises an amino acid sequence identical to the amino acid sequence of a Fc region found in nature.

A "variant Fc region" comprises an amino acid sequence which differs from that of a native sequence Fc region by virtue of at least one amino acid modification. Preferably, the variant Fc region has at least one amino acid substitution compared to a native sequence Fc region or to the Fc region of a parent polypeptide, e.g. from about one to about ten amino acid substitutions, and preferably from about one to about five amino acid substitutions in a native sequence Fc region or in the Fc region of the parent polypeptide. The variant Fc region herein will preferably possess at least about 80% sequence identity with a native sequence Fc region and/or with an Fc region of a parent polypeptide, and most preferably at least about 90% sequence identity therewith, more preferably at least about 95% sequence identity therewith.

"Antibody-dependent cell-mediated cytotoxicity" and "ADCC" refer to a cell-mediated reaction in which nonspecific cytotoxic cells that express Fc receptors (FcRs) (e.g. Natural Killer (NK) cells, neutrophils, and macrophages) recognize bound antibody on a target cell and subsequently cause lysis of the target cell. The primary cells for mediating ADCC, NK cells, express Fc $\gamma$ RIII only, whereas monocytes express Fc $\gamma$ RI, Fc $\gamma$ RII and Fc $\gamma$ RIII. FcR expression on hematopoietic cells is summarized in Table 3 on page 464 of Ravetch and Kinet, *Annu. Rev. Immunol.*, 9:457-92 (1991). To assess ADCC activity of a molecule of interest, an *in vitro* ADCC assay, such as that described in US Patent No. 5,500,362 or 5,821,337 may be performed. Useful effector cells for such assays include peripheral blood mononuclear cells (PBMC) and Natural Killer (NK) cells. Alternatively, or additionally, ADCC activity of the molecule of

interest may be assessed *in vivo*, e.g., in a animal model such as that disclosed in Clynes et al. PNAS (USA), 95:652-656 (1998).

"Human effector cells" are leukocytes which express one or more FcRs and perform effector functions. Preferably, the cells express 5 at least Fc $\gamma$ RIII and perform ADCC effector function. Examples of human leukocytes which mediate ADCC include peripheral blood mononuclear cells (PBMC), natural killer (NK) cells, monocytes, cytotoxic T cells and neutrophils; with PBMCs and NK cells being preferred. The effector cells may be isolated from a native source 10 thereof, e.g. from blood or PBMCs as described herein.

The terms "Fc receptor" and "FcR" are used to describe a receptor that binds to the Fc region of an antibody. The preferred FcR is a native sequence human FcR. Moreover, a preferred FcR is one which binds an IgG antibody (a gamma receptor) and includes receptors 15 of the Fc $\gamma$ RI, Fc $\gamma$ RII, and Fc $\gamma$ RIII subclasses, including allelic variants and alternatively spliced forms of these receptors. Fc $\gamma$ RII receptors include Fc $\gamma$ RIIA (an "activating receptor") and Fc $\gamma$ RIIB (an "inhibiting receptor"), which have similar amino acid sequences that differ primarily in the cytoplasmic domains thereof. Activating 20 receptor Fc $\gamma$ RIIA contains an immunoreceptor tyrosine-based activation motif (ITAM) in its cytoplasmic domain. Inhibiting receptor Fc $\gamma$ RIIB contains an immunoreceptor tyrosine-based inhibition motif (ITIM) in its cytoplasmic domain (reviewed in Daëron, Annu. Rev. Immunol., 15:203-234 (1997)). FcRs are reviewed in Ravetch and Kinet, Annu. Rev. Immunol., 25:457-92 (1991); Capel et al., Immunomethods, 4:25-34 (1994); and de Haas et al., J. Lab. Clin. Med., 126:330-41 (1995). Other FcRs, including those to be identified in the future, are 30 encompassed by the term "FcR" herein. The term also includes the neonatal receptor, FcRn, which is responsible for the transfer of maternal IgGs to the fetus (Guyer et al., J. Immunol., 117:587 (1976); and Kim et al., J. Immunol., 24:249 (1994)).

"Complement dependent cytotoxicity" and "CDC" refer to the lysing of a target in the presence of complement. The complement activation pathway is initiated by the binding of the first component 35 of the complement system (C1q) to a molecule (e.g. an antibody) complexed with a cognate antigen. To assess complement activation, a CDC assay, e.g. as described in Gazzano-Santoro et al., J. Immunol. Methods, 202:163 (1996), may be performed.

An "affinity matured" antibody is one with one or more 40 alterations in one or more CDRs thereof which result an improvement

in the affinity of the antibody for antigen, compared to a parent antibody which does not possess those alteration(s). Preferred affinity matured antibodies will have nanomolar or even picomolar affinities for the target antigen. Affinity matured antibodies are

5 produced by procedures known in the art. Marks et al.

Bio/Technology, 10:779-783 (1992) describes affinity maturation by VH and VL domain shuffling. Random mutagenesis of CDR and/or framework residues is described by: Barbas et al. Proc Nat. Acad. Sci., USA 91:3809-3813 (1994); Schier et al. Gene, 169:147-155 (1995); Yelton 10 et al. J. Immunol., 155:1994-2004 (1995); Jackson et al., J. Immunol., 154(7):3310-9 (1995); and Hawkins et al., J. Mol. Biol., 226:889-896 (1992).

The term "immunospecific" as used in "immunospecific binding of antibodies" for example, refers to the antigen specific binding 15 interaction that occurs between the antigen-combining site of an antibody and the specific antigen recognized by that antibody.

The terms "cancer", "cancerous", "metastasis" and "malignant" refer to or describe the physiological condition in mammals that is typically characterized by unregulated cell growth. Examples of

20 cancer include but are not limited to, carcinoma including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, and leukemia.

More particular examples of such cancers include squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non-Hodgkin's lymphoma,

25 pancreatic cancer, glioblastoma, cervical cancer, ovarian cancer, liver cancer such as hepatic carcinoma and hepatoma, bladder cancer,

breast cancer, colon cancer, colorectal cancer, endometrial carcinoma, myeloma (such as multiple myeloma), salivary gland carcinoma, kidney cancer such as renal cell carcinoma and Wilms'

30 tumors, basal cell carcinoma, melanoma, prostate cancer, vulval cancer, thyroid cancer, testicular cancer, esophageal cancer, and various types of head and neck cancer. The preferred cancers for treatment herein include breast cancer, gastric cancer, lung cancer,

colon or colorectal cancer, glioma and glioblastoma.

35 The term "prodrug" as used in this application refers to a precursor or derivative form of a pharmaceutically active substance that is less cytotoxic to cancer cells compared to the parent drug and is capable of being enzymatically activated or converted into the more active parent form. See, e.g., Wilman, "Prodrugs in Cancer

40 Chemotherapy" Biochemical Society Transactions, 14, pp. 375-382,

615th Meeting Belfast (1986) and Stella et al., "Prodrugs: A

Chemical Approach to Targeted Drug Delivery," *Directed Drug Delivery*, Borchardt et al., (ed.), pp. 247-267, Humana Press (1985). The prodrugs of this invention include, but are not limited to, phosphate-containing prodrugs, thiophosphate-containing prodrugs, 5 sulfate-containing prodrugs, peptide-containing prodrugs, D-amino acid-modified prodrugs, glycosylated prodrugs, beta-lactam-containing prodrugs, optionally substituted phenoxyacetamide-containing prodrugs or optionally substituted phenylacetamide-containing prodrugs, 5-fluorocytosine and other 5-fluorouridine prodrugs which can be converted into the more active cytotoxic free drug. Examples of cytotoxic drugs that can be derivatized into a prodrug form for use in this invention include, but are not limited to, those chemotherapeutic agents described below.

The term "cytotoxic agent" as used herein refers to a substance that inhibits or prevents the function of cells and/or causes destruction of cells. The term is intended to include radioactive isotopes (e.g. At<sup>211</sup>, I<sup>131</sup>, I<sup>125</sup>, Y<sup>90</sup>, Re<sup>186</sup>, Re<sup>188</sup>, Sm<sup>153</sup>, Bi<sup>212</sup>, P<sup>32</sup> and radioactive isotopes of Lu), chemotherapeutic agents, and toxins such as small molecule toxins or enzymatically active toxins of bacterial, fungal, plant or animal origin, including fragments and/or variants thereof.

A "chemotherapeutic agent" is a chemical compound useful in the treatment of conditions like cancer. Examples of chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (CYTOXAN<sup>TM</sup>); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylololomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; calystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CBI-TMI); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, ranimustine; antibiotics such as the enediyne

antibiotics (e.g. calicheamicin, especially calicheamicin  $\gamma_1^I$  and calicheamicin  $\theta^I_1$ , see, e.g., *Agnew Chem Intl. Ed. Engl.*, 33:183-186 (1994); dynemicin, including dynemicin A; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne 5 antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabacin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (including morpholino- doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and 10 deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti- metabolites such as methotrexate and 5-fluorouracil (5-FU); folic 15 acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprime, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine, 5-FU; androgens such as 20 calusterone, dromostanolone propionate, epitiostanol, mepitiostane, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolinic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; 25 diaziquone; el fornithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; podophyllinic acid; 2-ethylhydrazide; procarbazine; 30 PSK<sup>®</sup>; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2, 2', 2''-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); 35 cyclophosphamide; thiotepa; taxoids, e.g. paclitaxel (TAXOL<sup>®</sup>, Bristol-Myers Squibb Oncology, Princeton, NJ) and doxetaxel (TAXOTERE<sup>®</sup>, Rhône-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; 40 etoposide (VP-16); ifosfamide; mitomycin C; mitoxantrone; vincristine; vinorelbine; navelbine; novantrone; teniposide;

daunomycin; aminopterin; xeloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoic acid; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in this definition  
5 are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens including for example tamoxifen, raloxifene, aromatase inhibiting 4(5)-imidazoles, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston); and anti-androgens such as flutamide,  
10 nilutamide, bicalutamide, leuprolide, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above.

A "growth inhibitory agent" when used herein refers to a compound or composition which inhibits growth of a cell, either *in vitro* or *in vivo*. Thus, the growth inhibitory agent is one which significantly reduces the percentage of cells overexpressing such genes in S phase. Examples of growth inhibitory agents include agents that block cell cycle progression (at a place other than S phase), such as agents that induce G1 arrest and M-phase arrest.  
15  
20 Classical M-phase blockers include the vincas (vincristine and vinblastine), taxol, and topo II inhibitors such as doxorubicin, epirubicin, daunorubicin, etoposide, and bleomycin. Those agents that arrest G1 also spill over into S-phase arrest, for example, DNA alkylating agents such as tamoxifen, prednisone, dacarbazine, mechlorethamine, cisplatin, methotrexate, 5-fluorouracil, and ara-C.  
25 Further information can be found in *The Molecular Basis of Cancer*, Mendelsohn and Israel, eds., Chapter 1, entitled "Cell cycle regulation, oncogens, and antineoplastic drugs" by Murakami *et al.* (WB Saunders: Philadelphia, 1995), especially p. 13.

30 The term "cytokine" is a generic term for proteins released by one cell population which act on another cell as intercellular mediators. Examples of such cytokines are lymphokines, monokines, and traditional polypeptide hormones. Included among the cytokines are growth hormone such as human growth hormone, N-methionyl human growth hormone, and bovine growth hormone; parathyroid hormone; thyroxine; insulin; proinsulin; relaxin; prorelaxin; glycoprotein hormones such as follicle stimulating hormone (FSH), thyroid stimulating hormone (TSH), and luteinizing hormone (LH); hepatic growth factor; fibroblast growth factor; prolactin; placental  
35 lactogen; tumor necrosis factor- $\alpha$  and - $\beta$ ; Apo-2 ligand (also referred  
40

to as TRAIL); mullerian-inhibiting substance; mouse gonadotropin-associated peptide; inhibin; activin; vascular endothelial growth factor; integrin; thrombopoietin (TPO); nerve growth factors; platelet-growth factor; transforming growth factors (TGFs) such as 5 TGF- $\alpha$  and TGF- $\beta$ ; insulin-like growth factor-I and -II; erythropoietin (EPO); osteoinductive factors; interferons such as interferon- $\alpha$ , - $\beta$ , and -gamma; colony stimulating factors (CSFs) such as macrophage-CSF (M-CSF); granulocyte-macrophage-CSF (GM-CSF); and granulocyte-CSF (G-CSF); interleukins (ILs) such as IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, 10 IL-7, IL-8, IL-9, IL-11, IL-12; and other polypeptide factors including LIF and kit ligand (KL). As used herein, the term cytokine includes proteins from natural sources or from recombinant cell culture and biologically active equivalents of the native sequence cytokines.

15 "Treatment" is an intervention performed with the intention of preventing the development or altering the pathology of a disorder. Accordingly, "treatment" refers to both therapeutic treatment and prophylactic or preventative measures, wherein the object is to prevent or slow down (lessen) the targeted pathological condition or 20 disorder. Those in need of treatment include those already with the disorder as well as those in which the disorder is to be prevented. In treatment of a disorder, a therapeutic agent may directly decrease or increase the magnitude of response of a pathological component of the disorder, or render the disease more susceptible to treatment by 25 other therapeutic agents, e.g. antibiotics, antifungals, anti-inflammatory agents, chemotherapeutics, etc.

The term "effective amount" is the minimum concentration of WISP-1 antagonist which causes, induces or results in either a detectable improvement or reduction of a pathological condition. In 30 a method of treating cancer, an effective amount is one which causes, induces, or results in reduction of cancer cell number or tumor volume. Furthermore a "therapeutically effective amount" is the minimum concentration (amount) of WISP-1 antagonist administered to a mammal which would be effective in at least attenuating a 35 pathological symptom.

"Chronic" administration refers to administration of the factor(s) in a continuous mode as opposed to an acute mode, so as to maintain the initial therapeutic effect (activity) for an extended period of time. "Intermittent" administration is treatment that is

done not consecutively without interruption, but rather is cyclic in nature.

5 "Mammal" for purposes of treatment refers to any animal classified as a mammal, including humans, domestic and farm animals, and zoo, sports, or pet animals, such as dogs, horses, cats, cattle, pigs, hamsters, etc. Preferably, the mammal is human.

Administration "in combination with" one or more further therapeutic agents includes simultaneous (concurrent) and consecutive administration in any order.

10 "Carriers" as used herein include pharmaceutically acceptable carriers, excipients, or stabilizers which are nontoxic to the cell or mammal being exposed thereto at the dosages and concentrations employed. Often the physiologically acceptable carrier is an aqueous pH buffered solution. Examples of physiologically acceptable carriers include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid; low molecular weight (less than about 10 residues) polypeptide; proteins, such as serum albumin, gelatin, or immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, 15 asparagine, arginine or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugar alcohols such as mannitol or sorbitol; salt-forming counterions such as sodium; and/or nonionic surfactants such as TWEEN<sup>®</sup>, polyethylene glycol (PEG), and 20 PLURONICS<sup>®</sup>, hyaluronic acid (HA).

25

**II. Methods and Compositions of the Invention**

As described further in the Examples below, Applicant has surprisingly found that ectopic expression of WISP-1 in fibroblasts 30 triggered HA production, namely increased HA production induced its accumulation at the cell surface and in the culture media. It is believed that cell surface HA can enhance anchorage-independent growth and tumorigenicity (Kosaki et al., Cancer Res., 59:1141-1145 (1999)) and may diminish contact inhibition of cell proliferation, 35 thereby promoting cell migration (Ichikawa et al., J. Invest. Dermatol., 113:935-939 (1999); Itano et al., Proc. Natl. Acad. Sci., 99:3609-3614 (2002)). Hyaluronan coat formation has also been correlated with cellular metastatic potential (Zhang et al., Cancer Res., 55:428-433 (1995); Toole et al., J. Biol. Chem., 277:4593-4596 40 (2002)). Analysis of HA synthases (HAS1, HAS2, and HAS3) expression revealed that WISP-1 promoted HAS2 induction whereas HAS1 and HAS3

mRNA levels remained unchanged. *HAS2* is the main target of growth factor and cytokine up-regulated HA synthesis (Pienimaki et al., J. Biol. Chem., 276:20428-20435 (2001)). Ectopic expression of *HAS2* increases HA secretion and induces hyaluronan coat formation (Kosaki et al., supra). Moreover, *HAS2* is believed to be important for hyaluronan-mediated transformation of epithelium to mesenchyme during cardiac morphogenesis (Camenisch et al., J. Clin. Invest., 106:349-360 (2000)). Because epithelial to mesenchymal transition is also important for tumor invasion and metastasis, *HAS2* is postulated to 10 play an important role in epithelial tumor progression (Boyer et al., Biochem. Pharmacol., 60:1091-1099 (2000); Hay, Acta Anat., 154:8-20 (1995); Arias, Cell, 105:425-431 (2001)). These results indicate that HA secretion induced by WISP-1 may be important for tumor invasion and metastasis.

15 WISP-1 also induced the expression of two HA receptors, CD44 and RHAMM. By inducing HA receptor expression and increasing HA production, WISP-1 may activate an autocrine and/or paracrine loop. Hyaluronan interaction with CD44 and RHAMM promotes cell locomotion and proliferation *in vitro* and tumor growth and metastasis *in vivo* 20 (Turley et al., J. Biol. Chem., 277:4589-4592 (2002); Sy et al., Curr. Top. Microbiol. Immunol., 213:129-153 (1996); Hall et al., J. Neurooncol., 26:221-229 (1995)).

25 The effects of WISP-1 on cell migration were analyzed in a cell wound healing assay and by time lapse microscopy, and the data showed that WISP-1 expression promoted cell motility. Further, isolated cells showed increased migration suggesting that WISP-1 could act through an autocrine mechanism. Purified recombinant WISP-1 promoted HAS2 and CD44 expression, though, when attached to a surface. This induction was further increased when WISP-1 was tethered through its 30 interaction with decorin. Similarly, WISP-1 promoted cell motility when surface bound, suggesting that WISP-1 may also act through a paracrine mechanism when tethered to a substrate. The Examples herein show that WISP-1 induced cell motility was mediated by CD44, by abolishing migration with an anti-CD44 antibody. WISP-1 induced 35 cell motility was also inhibited by WISP-1 antibodies. Moreover, WISP-1 haptotactic activity was not restricted to a single cell type as it induced both normal fibroblast and colon adenocarcinoma cell migration.

40 Because WISP-1 is believed to be a Wnt-1 downstream effector, the C57MG/Wnt-1 cell line was analyzed for phenotypes found in NRK/WISP-1 cells. Consistent with a role for WISP-1 downstream of

Wnt-1, C57MG/Wnt-1 cells overexpressed HAS2 and CD44 and had higher CD44 protein content compared to the parental cell line. In addition, C57/Wnt-1 cells spontaneously scattered in culture and demonstrated a de-differentiated spindloid morphology similar to NRK/WISP-1H cells.

5 Expression analysis was also performed on a group of MMTV-Wnt-1 mammary tumors, and elevated CD44 and HAS2 expression was detected in all mammary tumors from the MMTV-Wnt-1 transgenic mice. In these tumors, WISP-1 expression was localized to the stromal fibroblasts whereas CD44 and HAS2 were expressed by tumor epithelial cells.

10 Although negative for HAS2 expression, the peritumoral stroma contained high levels of HA whereas the tumor parenchyma stained only weakly for HA. Although it is generally accepted that fibroblasts are responsible for HA production, the origin of stromal HA is not definitely known. Because HAS2 expression was found only in the 15 parenchyma, the experimental results disclosed herein suggest that the tumor cells are responsible for the HA synthesis and deposition in the stroma. Hyaluronan accumulation in the peritumoral stroma is frequently encountered in several tumor types and was previously reported for ovary, breast, prostate and colon adenocarcinomas

20 (Ropponen et al., Cancer Res., 58:342-347 (1998); Lippinen et al., Eur. J. Cancer, 37:849-856 (2001); Auvinen et al., Am. J. Pathol., 156:529-536 (2000); Anttila et al., Cancer Res., 60:150-155 (2000)).

Moreover, a high stromal level of HA was associated with poor differentiation, metastatic behavior and unfavorable prognosis.

25 Because HA and CD44 are associated with tumor invasion, the metastatic potential of WISP-1 expressing cells was evaluated in vivo. After tail inoculation, NRK/WISP-1 cells readily colonized the lungs of injected mice and formed invasive tumors. NRK/WISP-1 cells exhibited a significant metastatic potential while NRK cells did not.

30 Histological observation revealed that NRK/WISP-1 cells populated the vasculature and invaded the pulmonary airways. The lung colonization was proportional to the number of cells injected, the time after injection and the WISP-1 expression. In addition, Applicant found the metastatic potential to be proportional to CD44 35 and HAS2 expression levels, and the treatment of mice inoculated with NRK/WISP-1H cells with CD44 antibody or WISP-1 antibody greatly diminished the number and size of the tumors in the lung. Thus, it is likely that levels of WISP-1 promoted lung colonization through a hyaluronan-CD44 mechanism. This is consistent with previous reports 40 demonstrating the importance of CD44 and its interaction with HA for tumor growth, metastasis and the retention of metastatic cells to the

lung vasculature (Sy et al., supra; Kogerman et al., Proc. Natl. Acad. Sci., 94:13233-13238 (1997)).

Although activation of the Wnt pathway by APC or  $\beta$ -catenin mutations may typically be associated with colorectal cancer, several 5 lines of evidence suggest that it also plays a role in other types of cancer including mammary adenocarcinoma (Polakis, Genes Dev., 14:1837-1851 (2000); Brown, Breast Cancer Res., 3:351-355 (2001)). APC truncation and increased  $\beta$ -catenin levels were found in human 10 breast cancer cell lines (Schlosshauer et al., Carcinogenesis, 21:1453-1456 (2000)). Somatic mutations of the APC gene were found in primary breast cancers (Furuuchi et al., Am. J. Pathol., 156:1997-2005 (2000)). In addition,  $\beta$ -catenin activating mutations promote 15 mouse mammary adenocarcinomas Michaelson et al., Oncogene, 20:5525-5532 (2001)). Elevated levels of Wnt-1 and  $\beta$ -catenin were found in invasive ductal breast carcinomas and correlated with poor prognosis 20 (Lin et al., Proc. Natl. Acad. Sci., 97:4262-4266 (2000)). It is therefore possible that WISP-1 expression found in certain breast adenocarcinomas resulted from Wnt pathway activation. Applicants have also found, by Taqman analysis, that WISP-1 is overexpressed in cancers such as breast and glial tumors.

For at least these reasons, it is believed that WISP-1 antagonists will be particularly useful in treating and diagnosing various pathological disorders, such as cancer. The present invention accordingly provides methods for modulating, blocking or neutralizing 25 WISP-1 activity in mammalian cells which comprise exposing the cells to a desired amount of WISP-1 antagonist. Preferably, the amount of WISP-1 antagonist employed will be an amount effective to reduce or inhibit cancer cell growth, metastasis or motility. This can be accomplished *in vivo* or *ex vivo* in accordance, for instance, with the 30 methods described below and in the Examples. Exemplary conditions or disorders to be treated with such WISP-1 antagonists include conditions in mammals clinically referred to as cancer.

Diagnostic methods are also provided herein. For instance, the antagonists may be employed to detect invasive or metastatic cancers. 35 The antagonist molecule may be used, e.g., in assays to detect or quantitate metastatic cancer cells in a sample. A sample, such as cells obtained from a mammal, can be incubated in the presence of a labeled antagonist, and detection of the labeled antagonist bound in the sample can be performed.

The antagonists which can be employed in the methods include, but are not limited to, WISP-1 immunoadhesins, fusion proteins comprising WISP-1, covalently modified forms of WISP-1, WISP-1 variants, fusion proteins thereof, and WISP-1 antibodies. Various 5 techniques that can be employed for making the antagonists are described herein. For instance, methods and techniques for preparing WISP-1 polypeptides are described. Further modifications of the polypeptides, and antibodies to WISP-1 are also described.

In addition to the full-length native sequence WISP-1 10 polypeptide described herein, it is contemplated that WISP-1 polypeptide variants can be prepared. WISP-1 variants can be prepared by introducing appropriate nucleotide changes into the encoding DNA, and/or by synthesis of the desired polypeptide. Those skilled in the art will appreciate that amino acid changes may alter 15 post-translational processes of the WISP-1 polypeptide, such as changing the number or position of glycosylation sites or altering the membrane anchoring characteristics.

Variations in the WISP-1 polypeptides described herein, can be made, for example, using any of the techniques and guidelines for 20 conservative and non-conservative mutations set forth, for instance, in U.S. Patent No. 5,364,934. Variations may be a substitution, deletion or insertion of one or more codons encoding the polypeptide that results in a change in the amino acid sequence as compared with the native sequence polypeptide. Optionally the variation is by 25 substitution of at least one amino acid with any other amino acid in one or more of the domains of the WISP-1 polypeptide. Guidance in determining which amino acid residue may be inserted, substituted or deleted without adversely affecting the desired activity may be found by comparing the sequence of the WISP-1 polypeptide with that of 30 homologous known protein molecules and minimizing the number of amino acid sequence changes made in regions of high homology. Amino acid substitutions can be the result of replacing one amino acid with another amino acid having similar structural and/or chemical properties, such as the replacement of a leucine with a serine, i.e., 35 conservative amino acid replacements. Insertions or deletions may optionally be in the range of about 1 to 5 amino acids. The variation allowed may be determined by systematically making insertions, deletions or substitutions of amino acids in the sequence and testing the resulting variants for activity exhibited by the 40 full-length or mature native sequence.

WISP-1 polypeptide fragments are provided herein. Such fragments may be truncated at the N-terminus or C-terminus, or may lack internal residues, for example, when compared with a full length native protein. Certain fragments lack amino acid residues that are 5 not essential for a desired biological activity of the WISP-1 polypeptide.

WISP-1 polypeptide fragments may be prepared by any of a number of conventional techniques. Desired peptide fragments may be chemically synthesized. An alternative approach involves generating 10 polypeptide fragments by enzymatic digestion, e.g., by treating the protein with an enzyme known to cleave proteins at sites defined by particular amino acid residues, or by digesting the DNA with suitable restriction enzymes and isolating the desired fragment. Yet another suitable technique involves isolating and amplifying a DNA fragment 15 encoding a desired polypeptide fragment, by polymerase chain reaction (PCR). Oligonucleotides that define the desired termini of the DNA fragment are employed at the 5' and 3' primers in the PCR.

In particular embodiments, conservative substitutions of interest are shown in the Table below under the heading of preferred 20 substitutions. If such substitutions result in a change in biological activity, then more substantial changes, denominated exemplary substitutions in the Table, or as further described below in reference to amino acid classes, are introduced and the products screened.

25 Table

	Original	Exemplary	Preferred
	Residue	Substitutions	
		Substitutions	
30	Ala (A)	val; leu; ile	val
	Arg (R)	lys; gln; asn	lys
	Asn (N)	gln; his; lys; arg	gln
	Asp (D)	glu	glu
35	Cys (C)	ser	ser
	Gln (Q)	asn	asn
	Glu (E)	asp	asp
	Gly (G)	pro; ala	ala
	His (H)	asn; gln; lys; arg	arg
40	Ile (I)	leu; val; met; ala; phe; norleucine	leu

Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
5 Phe (F)	leu; val; ile; ala; tyr	leu
Pro (P)	ala	ala
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr; phe	tyr
10 Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

Substantial modifications in function or immunological identity  
 15 of the WISP-1 polypeptide are accomplished by selecting substitutions  
 that differ significantly in their effect on maintaining (a) the  
 structure of the polypeptide backbone in the area of the  
 substitution, for example, as a sheet or helical conformation, (b)  
 the charge or hydrophobicity of the molecule at the target site, or  
 20 (c) the bulk of the side chain. Naturally occurring residues are  
 divided into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- (3) acidic: asp, glu;
- 25 (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic: trp, tyr, phe.

Non-conservative substitutions will entail exchanging a member  
 of one of these classes for another class. Such substituted residues  
 30 also may be introduced into the conservative substitution sites or,  
 more preferably, into the remaining (non-conserved) sites.

The variations can be made using methods known in the art such  
 as oligonucleotide-mediated (site-directed) mutagenesis, alanine  
 scanning, and PCR mutagenesis. Site-directed mutagenesis [Carter et  
 35 al., Nucl. Acids Res., 13:4331 (1986); Zoller et al., Nucl. Acids Res., 10:6487 (1987)], cassette mutagenesis [Wells et al., Gene, 34:315 (1985)], restriction selection mutagenesis [Wells et al., Philos. Trans. R. Soc. London SerA, 317:415 (1986)] or other known  
 techniques can be performed on the cloned DNA to produce the WISP-1  
 40 polypeptide variant DNA.

Scanning amino acid analysis can also be employed to identify one or more amino acids along a contiguous sequence. Among the preferred scanning amino acids are relatively small, neutral amino acids. Such amino acids include alanine, glycine, serine, and 5 cysteine. Alanine is typically a preferred scanning amino acid among this group because it eliminates the side-chain beyond the beta-carbon and is less likely to alter the main-chain conformation of the variant [Cunningham and Wells, Science, 244:1081-1085 (1989)]. Alanine is also typically preferred because it is the most common 10 amino acid. Further, it is frequently found in both buried and exposed positions [Creighton, The Proteins, (W.H. Freeman & Co., N.Y.); Chothia, J. Mol. Biol., 150:1 (1976)]. If alanine substitution does not yield adequate amounts of variant, an isoteric amino acid can be used.

15 Any cysteine residue not involved in maintaining the proper conformation of the WISP-1 polypeptide also may be substituted, generally with serine, to improve the oxidative stability of the molecule and prevent aberrant crosslinking. Conversely, cysteine bond(s) may be added to the WISP-1 polypeptide to improve its 20 stability.

The description below relates primarily to production of WISP-1 polypeptides by culturing cells transformed or transfected with a vector containing WISP-1 polypeptide-encoding nucleic acid. It is, of course, contemplated that alternative methods, which are well 25 known in the art, may be employed to prepare WISP-1 polypeptides. For instance, the appropriate amino acid sequence, or portions thereof, may be produced by direct peptide synthesis using solid-phase techniques [see, e.g., Stewart et al., Solid-Phase Peptide Synthesis, W.H. Freeman Co., San Francisco, CA (1969); Merrifield, J. Am. Chem. Soc., 85:2149-2154 (1963)]. *In vitro* protein synthesis may 30 be performed using manual techniques or by automation. Automated synthesis may be accomplished, for instance, using an Applied Biosystems Peptide Synthesizer (Foster City, CA) using manufacturer's instructions. Various portions of the WISP-1 polypeptide may be 35 chemically synthesized separately and combined using chemical or enzymatic methods to produce the desired WISP-1 polypeptide. The methods and techniques described are similarly applicable to production of WISP-1 variants, modified forms of WISP-1 and WISP-1 antibodies.

1. Isolation of DNA Encoding WISP-1 Polypeptide

DNA encoding WISP-1 polypeptide may be obtained from a cDNA library prepared from tissue believed to possess the WISP-1 polypeptide mRNA and to express it at a detectable level.

5 Accordingly, human WISP-1 polypeptide DNA can be conveniently obtained from a cDNA library prepared from human tissue. The WISP-1 polypeptide-encoding gene may also be obtained from a genomic library or by known synthetic procedures (e.g., automated nucleic acid synthesis).

10 Libraries can be screened with probes (such as oligonucleotides of at least about 20-80 bases) designed to identify the gene of interest or the protein encoded by it. Screening the cDNA or genomic library with the selected probe may be conducted using standard procedures, such as described in Sambrook et al., Molecular Cloning: A Laboratory Manual (New York: Cold Spring Harbor Laboratory Press, 1989). An alternative means to isolate the gene encoding WISP-1 polypeptide is to use PCR methodology [Sambrook et al., supra; Dieffenbach et al., PCR Primer: A Laboratory Manual (Cold Spring Harbor Laboratory Press, 1995)].

20 Techniques for screening a cDNA library are well known in the art. The oligonucleotide sequences selected as probes should be of sufficient length and sufficiently unambiguous that false positives are minimized. The oligonucleotide is preferably labeled such that it can be detected upon hybridization to DNA in the library being screened. Methods of labeling are well known in the art, and include the use of radiolabels like  $^{32}\text{P}$ -labeled ATP, biotinylation or enzyme labeling. Hybridization conditions, including moderate stringency and high stringency, are provided in Sambrook et al., supra.

30 Sequences identified in such library screening methods can be compared and aligned to other known sequences deposited and available in public databases such as GenBank or other private sequence databases. Sequence identity (at either the amino acid or nucleotide level) within defined regions of the molecule or across the full-length sequence can be determined using methods known in the art and as described herein.

35 Nucleic acid having protein coding sequence may be obtained by screening selected cDNA or genomic libraries using the deduced amino acid sequence disclosed herein for the first time, and, if necessary, using conventional primer extension procedures as described in 40 Sambrook et al., supra, to detect precursors and processing

intermediates of mRNA that may not have been reverse-transcribed into cDNA.

2. Selection and Transformation of Host Cells

Host cells are transfected or transformed with expression or 5 cloning vectors described herein for WISP-1 polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding the desired sequences. The culture conditions, such as media, temperature, pH and the like, can be selected by the 10 skilled artisan without undue experimentation. In general, principles, protocols, and practical techniques for maximizing the productivity of cell cultures can be found in Mammalian Cell Biotechnology: a Practical Approach, M. Butler, ed. (IRL Press, 1991) and Sambrook et al., supra.

15 Methods of eukaryotic cell transfection and prokaryotic cell transformation are known to the ordinarily skilled artisan, for example,  $\text{CaCl}_2$ ,  $\text{CaPO}_4$ , liposome-mediated and electroporation. Depending on the host cell used, transformation is performed using 20 standard techniques appropriate to such cells. The calcium treatment employing calcium chloride, as described in Sambrook et al., supra, or electroporation is generally used for prokaryotes. Infection with *Agrobacterium tumefaciens* is used for transformation of certain plant cells, as described by Shaw et al., Gene, 23:315 (1983) and WO 89/05859 published 29 June 1989. For mammalian cells without such 25 cell walls, the calcium phosphate precipitation method of Graham and van der Eb, Virology, 52:456-457 (1978) can be employed. General aspects of mammalian cell host system transfections have been described in U.S. Patent No. 4,399,216. Transformations into yeast are typically carried out according to the method of Van Solingen et 30 al., J. Bact., 130:946 (1977) and Hsiao et al., Proc. Natl. Acad. Sci. (USA), 76:3829 (1979). However, other methods for introducing DNA into cells, such as by nuclear microinjection, electroporation, bacterial protoplast fusion with intact cells, or polycations, e.g., polybrene, polyornithine, may also be used. For various techniques 35 for transforming mammalian cells, see Keown et al., Methods in Enzymology, 185:527-537 (1990) and Mansour et al., Nature, 336:348-352 (1988).

Suitable host cells for cloning or expressing the DNA in the vectors herein include prokaryote, yeast, or higher eukaryote cells.

40 Suitable prokaryotes include but are not limited to eubacteria, such as Gram-negative or Gram-positive organisms, for example,

Enterobacteriaceae such as *E. coli*. Various *E. coli* strains are publicly available, such as *E. coli* K12 strain MM294 (ATCC 31,446); *E. coli* X1776 (ATCC 31,537); *E. coli* strain W3110 (ATCC 27,325) and K5 772 (ATCC 53,635). Other suitable prokaryotic host cells include 5 *Enterobacteriaceae* such as *Escherichia*, e.g., *E. coli*, *Enterobacter*, *Erwinia*, *Klebsiella*, *Proteus*, *Salmonella*, e.g., *Salmonella typhimurium*, *Serratia*, e.g., *Serratia marcescans*, and *Shigella*, as well as *Bacilli* such as *B. subtilis* and *B. licheniformis* (e.g., *B. licheniformis* 41P disclosed in DD 266,710 published 12 April 1989), 10 *Pseudomonas* such as *P. aeruginosa*, and *Streptomyces*. These examples are illustrative rather than limiting. Strain W3110 is one particularly preferred host or parent host because it is a common host strain for recombinant DNA product fermentations. Preferably, the host cell secretes minimal amounts of proteolytic enzymes. For 15 example, strain W3110 may be modified to effect a genetic mutation in the genes encoding proteins endogenous to the host, with examples of such hosts including *E. coli* W3110 strain 1A2, which has the complete genotype *tonA*; *E. coli* W3110 strain 9E4, which has the complete genotype *tonA ptr3*; *E. coli* W3110 strain 27C7 (ATCC 55,244), which 20 has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT kan<sup>r</sup>*; *E. coli* W3110 strain 37D6, which has the complete genotype *tonA ptr3 phoA E15 (argF-lac)169 degP ompT rbs7 ilvG kan<sup>r</sup>*; *E. coli* W3110 strain 40B4, which is strain 37D6 with a non-kanamycin 25 resistant *degP* deletion mutation; and an *E. coli* strain having mutant periplasmic protease disclosed in U.S. Patent No. 4,946,783 issued 7 August 1990. Alternatively, *in vitro* methods of cloning, e.g., PCR or other nucleic acid polymerase reactions, are suitable.

In addition to prokaryotes, eukaryotic microbes such as filamentous fungi or yeast are suitable cloning or expression hosts 30 for WISP-1 polypeptide-encoding vectors. *Saccharomyces cerevisiae* is a commonly used lower eukaryotic host microorganism. Others include *Schizosaccharomyces pombe* (Beach and Nurse, *Nature*, 290: 140 [1981]; EP 139,383 published 2 May 1985); *Kluyveromyces* hosts (U.S. Patent No. 4,943,529; Fleer et al., *Bio/Technology*, 9:968-975 (1991)) such 35 as, e.g., *K. lactis* (MW98-8C, CBS683, CBS4574; Louvencourt et al., *J. Bacteriol.*, 154(2):737-742 [1983]), *K. fragilis* (ATCC 12,424), *K. bulgaricus* (ATCC 16,045), *K. wickeramii* (ATCC 24,178), *K. waltii* (ATCC 56,500), *K. drosophilarum* (ATCC 36,906; Van den Berg et al., *Bio/Technology*, 8:135 (1990)), *K. thermotolerans*, and *K. marxianus*; 40 *yarrowiae* (EP 402,226); *Pichia pastoris* (EP 183,070; Sreekrishna et al., *J. Basic Microbiol.*, 28:265-278 [1988]); *Candida*; *Trichoderma*

reesia (EP 244,234); *Neurospora crassa* (Case et al., Proc. Natl. Acad. Sci. USA, 76:5259-5263 [1979]); *Schwanniomyces* such as *Schwanniomyces occidentalis* (EP 394,538 published 31 October 1990); and filamentous fungi such as, e.g., *Neurospora*, *Penicillium*, 5 *Tolypocladium* (WO 91/00357 published 10 January 1991), and *Aspergillus* hosts such as *A. nidulans* (Ballance et al., Biochem. Biophys. Res. Commun., 112:284-289 [1983]; Tilburn et al., Gene, 26:205-221 [1983]; Yelton et al., Proc. Natl. Acad. Sci. USA, 81: 1470-1474 [1984]) and *A. niger* (Kelly and Hynes, EMBO J., 4:475-479 10 [1985]). Methylotrophic yeasts are suitable herein and include, but are not limited to, yeast capable of growth on methanol selected from the genera consisting of *Hansenula*, *Candida*, *Kloeckera*, *Pichia*, *Saccharomyces*, *Torulopsis*, and *Rhodotorula*. A list of specific species that are exemplary of this class of yeasts may be found in C. 15 Anthony, The Biochemistry of Methylotrophs, 269 (1982).

Suitable host cells for the expression of glycosylated WISP-1 polypeptide are derived from multicellular organisms. Examples of invertebrate cells include insect cells such as *Drosophila S2* and *Spodoptera Sf9*, as well as plant cells, such as cell cultures of 20 cotton, corn, potato, soybean, petunia, tomato, and tobacco. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts such as *Spodoptera frugiperda* (caterpillar), *Aedes aegypti* (mosquito), *Aedes albopictus* (mosquito), *Drosophila melanogaster* (fruitfly), and *Bombyx mori* have been 25 identified. A variety of viral strains for transfection are publicly available, e.g., the L-1 variant of *Autographa californica* NPV and the Bm-5 strain of *Bombyx mori* NPV, and such viruses may be used as the virus herein according to the present invention, particularly for transfection of *Spodoptera frugiperda* cells.

30 However, interest has been greatest in vertebrate cells, and propagation of vertebrate cells in culture (tissue culture) has become a routine procedure. Examples of useful mammalian host cell lines are monkey kidney CV1 line transformed by SV40 (COS-7, ATCC CRL 1651); human embryonic kidney line (293 or 293 cells subcloned for 35 growth in suspension culture, Graham et al., J. Gen Virol. 36:59 (1977)); baby hamster kidney cells (BHK, ATCC CCL 10); Chinese hamster ovary cells/-DHFR (CHO, Urlaub et al., Proc. Natl. Acad. Sci. USA 77:4216 (1980)); mouse sertoli cells (TM4, Mather, Biol. Reprod. 23:243-251 (1980)); monkey kidney cells (CV1 ATCC CCL 70); African 40 green monkey kidney cells (VERO-76, ATCC CRL-1587); human cervical carcinoma cells (HELA, ATCC CCL 2); canine kidney cells (MDCK, ATCC

CCL 34); buffalo rat liver cells (BRL 3A, ATCC CRL 1442); human lung cells (W138, ATCC CCL 75); human liver cells (Hep G2, HB 8065); mouse mammary tumor (MMT 060562, ATCC CCL51); TRI cells (Mather et al., *Annals N.Y. Acad. Sci.* 383:44-68 (1982)); MRC 5 cells; FS4 cells; and 5 a human hepatoma line (Hep G2).

Host cells are transformed with the above-described expression or cloning vectors for WISP-1 polypeptide production and cultured in conventional nutrient media modified as appropriate for inducing promoters, selecting transformants, or amplifying the genes encoding 10 the desired sequences.

### 3. Selection and Use of a Replicable Vector

The nucleic acid (e.g., cDNA or genomic DNA) encoding WISP-1 polypeptide may be inserted into a replicable vector for cloning (amplification of the DNA) or for expression. Various vectors are 15 publicly available. The vector may, for example, be in the form of a plasmid, cosmid, viral particle, or phage. The appropriate nucleic acid sequence may be inserted into the vector by a variety of procedures. In general, DNA is inserted into an appropriate restriction endonuclease site(s) using techniques known in the art. 20 Vector components generally include, but are not limited to, one or more of a signal sequence, an origin of replication, one or more marker genes, an enhancer element, a promoter, and a transcription termination sequence. Construction of suitable vectors containing one or more of these components employs standard ligation techniques 25 which are known to the skilled artisan.

The WISP-1 may be produced recombinantly not only directly, but also as a fusion polypeptide with a heterologous polypeptide, which may be a signal sequence or other polypeptide having a specific cleavage site at the N-terminus of the mature protein or polypeptide. 30 In general, the signal sequence may be a component of the vector, or it may be a part of the WISP-1 polypeptide-encoding DNA that is inserted into the vector. The signal sequence may be a prokaryotic signal sequence selected, for example, from the group of the alkaline phosphatase, penicillinase, lpp, or heat-stable enterotoxin II leaders. For yeast secretion the signal sequence may be, e.g., the yeast invertase leader, alpha factor leader (including *Saccharomyces* 35 and *Kluyveromyces*  $\alpha$ -factor leaders, the latter described in U.S. Patent No. 5,010,182), or acid phosphatase leader, the *C. albicans* glucoamylase leader (EP 362,179 published 4 April 1990), or the 40 signal described in WO 90/13646 published 15 November 1990. In

mammalian cell expression, mammalian signal sequences may be used to direct secretion of the protein, such as signal sequences from secreted polypeptides of the same or related species, as well as viral secretory leaders.

5 Both expression and cloning vectors contain a nucleic acid sequence that enables the vector to replicate in one or more selected host cells. Such sequences are well known for a variety of bacteria, yeast, and viruses. The origin of replication from the plasmid pBR322 is suitable for most Gram-negative bacteria, the 2 $\mu$  plasmid 10 origin is suitable for yeast, and various viral origins (SV40, polyoma, adenovirus, VSV or BPV) are useful for cloning vectors in mammalian cells.

15 Expression and cloning vectors will typically contain a selection gene, also termed a selectable marker. Typical selection genes encode proteins that (a) confer resistance to antibiotics or other toxins, e.g., ampicillin, neomycin, methotrexate, or tetracycline, (b) complement auxotrophic deficiencies, or (c) supply critical nutrients not available from complex media, e.g., the gene encoding D-alanine racemase for *Bacilli*.

20 An example of suitable selectable markers for mammalian cells are those that enable the identification of cells competent to take up the WISP-1 polypeptide-encoding nucleic acid, such as DHFR or thymidine kinase. An appropriate host cell when wild-type DHFR is employed is the CHO cell line deficient in DHFR activity, prepared 25 and propagated as described by Urlaub et al., Proc. Natl. Acad. Sci. USA, 77:4216 (1980). A suitable selection gene for use in yeast is the *trp1* gene present in the yeast plasmid YRp7 [Stinchcomb et al., Nature, 282:39 (1979); Kingsman et al., Gene, 7:141 (1979); Tschemper et al., Gene, 10:157 (1980)]. The *trp1* gene provides a selection 30 marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example, ATCC No. 44076 or PEP4-1 [Jones, Genetics, 85:12 (1977)].

35 Expression and cloning vectors usually contain a promoter operably linked to the WISP-1 polypeptide-encoding nucleic acid sequence to direct mRNA synthesis. Promoters recognized by a variety of potential host cells are well known. Promoters suitable for use with prokaryotic hosts include the  $\beta$ -lactamase and lactose promoter systems [Chang et al., Nature, 275:615 (1978); Goeddel et al., Nature, 281:544 (1979)], alkaline phosphatase, a tryptophan (trp) 40 promoter system [Goeddel, Nucleic Acids Res., 8:4057 (1980); EP

36,776], and hybrid promoters such as the tac promoter [deBoer et al., Proc. Natl. Acad. Sci. USA, 80:21-25 (1983)]. Promoters for use in bacterial systems also will contain a Shine-Dalgarno (S.D.) sequence operably linked to the DNA encoding WISP polypeptide.

5 Examples of suitable promoting sequences for use with yeast hosts include the promoters for 3-phosphoglycerate kinase [Hitzeman et al., J. Biol. Chem., 255:2073 (1980)] or other glycolytic enzymes [Hess et al., J. Adv. Enzyme Reg., 7:149 (1968); Holland, Biochemistry, 17:4900 (1978)], such as enolase, glyceraldehyde-3-  
10 phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase.

15 Other yeast promoters, which are inducible promoters having the additional advantage of transcription controlled by growth conditions, are the promoter regions for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, metallothionein, glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose  
20 utilization. Suitable vectors and promoters for use in yeast expression are further described in EP 73,657.

25 WISP polypeptide transcription from vectors in mammalian host cells is controlled, for example, by promoters obtained from the genomes of viruses such as polyoma virus, fowlpox virus (UK 2,211,504 published 5 July 1989), adenovirus (such as Adenovirus 2), bovine papilloma virus, avian sarcoma virus, cytomegalovirus, a retrovirus, hepatitis-B virus and Simian Virus 40 (SV40), from heterologous mammalian promoters, e.g., the actin promoter or an immunoglobulin promoter, and from heat-shock promoters, provided such promoters are  
30 compatible with the host cell systems.

35 Transcription of a DNA encoding the WISP-1 polypeptide by higher eukaryotes may be increased by inserting an enhancer sequence into the vector. Enhancers are *cis*-acting elements of DNA, usually about from 10 to 300 bp, that act on a promoter to increase its transcription. Many enhancer sequences are now known from mammalian genes (globin, elastase, albumin,  $\beta$ -fetoprotein, and insulin). Typically, however, one will use an enhancer from a eukaryotic cell virus. Examples include the SV40 enhancer on the late side of the replication origin (bp 100-270), the cytomegalovirus early promoter  
40 enhancer, the polyoma enhancer on the late side of the replication origin, and adenovirus enhancers. The enhancer may be spliced into

the vector at a position 5' or 3' to the WISP-1 polypeptide coding sequence, but is preferably located at a site 5' from the promoter.

Expression vectors used in eukaryotic host cells (yeast, fungi, insect, plant, animal, human, or nucleated cells from other 5 multicellular organisms) will also contain sequences necessary for the termination of transcription and for stabilizing the mRNA. Such sequences are commonly available from the 5' and, occasionally 3', untranslated regions of eukaryotic or viral DNAs or cDNAs. These regions contain nucleotide segments transcribed as polyadenylated 10 fragments in the untranslated portion of the mRNA encoding WISP-1 polypeptide.

Still other methods, vectors, and host cells suitable for adaptation to the synthesis of WISP polypeptide in recombinant vertebrate cell culture are described in Gething et al., Nature, 15 293:620-625 (1981); Mantei et al., Nature, 281:40-46 (1979); EP 117,060; and EP 117,058.

#### 4. Culturing the Host Cells

The host cells used to produce the WISP polypeptide of this invention may be cultured in a variety of media. Commercially 20 available media such as Ham's F10 (Sigma), Minimal Essential Medium ((MEM), (Sigma), RPMI-1640 (Sigma), and Dulbecco's Modified Eagle's Medium ((DMEM), Sigma) are suitable for culturing the host cells. In addition, any of the media described in Ham et al., Meth. Enz. 58:44 (1979), Barnes et al., Anal. Biochem. 102:255 (1980), U.S. Pat. Nos. 25 4,767,704; 4,657,866; 4,927,762; 4,560,655; or 5,122,469; WO 90/03430; WO 87/00195; or U.S. Patent Re. 30,985 may be used as culture media for the host cells. Any of these media may be supplemented as necessary with hormones and/or other growth factors (such as insulin, transferrin, or epidermal growth factor), salts 30 (such as sodium chloride, calcium, magnesium, and phosphate), buffers (such as HEPES), nucleotides (such as adenosine and thymidine), antibiotics (such as GENTAMYCIN™ drug), trace elements (defined as inorganic compounds usually present at final concentrations in the micromolar range), and glucose or an equivalent energy source. Any 35 other necessary supplements may also be included at appropriate concentrations that would be known to those skilled in the art. The culture conditions, such as temperature, pH, and the like, are those previously used with the host cell selected for expression, and will be apparent to the ordinarily skilled artisan.

5. Detecting Gene Amplification/Expression

Gene amplification and/or expression may be measured in a sample directly, for example, by conventional Southern blotting, Northern blotting to quantitate the transcription of mRNA [Thomas, 5 Proc. Natl. Acad. Sci. USA, 77:5201-5205 (1980)], dot blotting (DNA analysis), or *in situ* hybridization, using an appropriately labeled probe, based on the sequences provided herein. Alternatively, antibodies may be employed that can recognize specific duplexes, including DNA duplexes, RNA duplexes, and DNA-RNA hybrid duplexes or 10 DNA-protein duplexes. The antibodies in turn may be labeled and the assay may be carried out where the duplex is bound to a surface, so that upon the formation of duplex on the surface, the presence of antibody bound to the duplex can be detected.

Gene expression, alternatively, may be measured by 15 immunological methods, such as immunohistochemical staining of cells or tissue sections and assay of cell culture or body fluids, to quantitate directly the expression of gene product. Antibodies useful for immunohistochemical staining and/or assay of sample fluids may be either monoclonal or polyclonal, and may be prepared in any 20 mammal. Conveniently, the antibodies may be prepared against a native sequence WISP polypeptide or against a synthetic peptide based on the DNA sequences provided herein or against exogenous sequence fused to WISP DNA and encoding a specific antibody epitope.

6. Purification of WISP Polypeptide

25 Forms of WISP polypeptide may be recovered from culture medium or from host cell lysates. If membrane-bound, it can be released from the membrane using a suitable detergent solution (e.g. Triton-X 100) or by enzymatic cleavage. Cells employed in expression of WISP-1 polypeptide can be disrupted by various physical or chemical means, 30 such as freeze-thaw cycling, sonication, mechanical disruption, or cell lysing agents.

It may be desired to purify WISP-1 polypeptide from recombinant cell proteins or polypeptides. The following procedures are exemplary of suitable purification procedures: by fractionation on an 35 ion-exchange column; ethanol precipitation; reverse phase HPLC; chromatography on silica or on a cation-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; protein A Sepharose columns to remove contaminants such as IgG; and metal chelating 40 columns to bind epitope-tagged forms of the WISP-1 polypeptide. Various methods of protein purification may be employed and such

methods are known in the art and described for example in Deutscher, Methods in Enzymology, 182 (1990); Scopes, Protein Purification: Principles and Practice, Springer-Verlag, New York (1982). The purification step(s) selected will depend, for example, on the nature 5 of the production process used and the particular WISP-1 polypeptide produced.

Soluble forms of WISP-1 may be employed as antagonists in the methods of the invention. Such soluble forms of WISP-1 may comprise modifications, as described below (such as by fusing to an 10 immunoglobulin, epitope tag or leucine zipper). Immunoadhesin molecules are further contemplated for use in the methods herein. WISP-1 immunoadhesins may comprise various forms of WISP-1, such as the full length polypeptide as well as soluble forms of the WISP-1 or a fragment thereof. In particular embodiments, the molecule may 15 comprise a fusion of the WISP-1 polypeptide with an immunoglobulin or a particular region of an immunoglobulin. For a bivalent form of the immunoadhesin, such a fusion could be to the Fc region of an IgG molecule. The Ig fusions preferably include the substitution of a soluble (transmembrane domain deleted or inactivated) form of the 20 polypeptide in place of at least one variable region within an Ig molecule. In a particularly preferred embodiment, the immunoglobulin fusion includes the hinge, CH2 and CH3, or the hinge, CH1, CH2 and CH3 regions of an IgG1 molecule. For the production of immunoglobulin fusions, see also US Patent No. 5,428,130 issued June 25 27, 1995 and Chamow et al., TIBTECH, 14:52-60 (1996).

The simplest and most straightforward immunoadhesin design combines the binding domain(s) of the adhesin (e.g. the WISP-1) with the Fc region of an immunoglobulin heavy chain. Ordinarily, when preparing the immunoadhesins of the present invention, nucleic acid 30 encoding the binding domain of the adhesin will be fused C-terminally to nucleic acid encoding the N-terminus of an immunoglobulin constant domain sequence, however N-terminal fusions are also possible.

Typically, in such fusions the encoded chimeric polypeptide will retain at least functionally active hinge, C<sub>H</sub>2 and C<sub>H</sub>3 domains 35 of the constant region of an immunoglobulin heavy chain. Fusions are also made to the C-terminus of the Fc portion of a constant domain, or immediately N-terminal to the C<sub>H</sub>1 of the heavy chain or the corresponding region of the light chain. The precise site at which the fusion is made is not critical; particular sites are well known 40 and may be selected in order to optimize the biological activity, secretion, or binding characteristics of the immunoadhesin.

In a preferred embodiment, the adhesin sequence is fused to the N-terminus of the Fc region of immunoglobulin G<sub>1</sub> (IgG<sub>1</sub>). It is possible to fuse the entire heavy chain constant region to the adhesin sequence. However, more preferably, a sequence beginning in 5 the hinge region just upstream of the papain cleavage site which defines IgG Fc chemically (i.e. residue 216, taking the first residue of heavy chain constant region to be 114), or analogous sites of other immunoglobulins is used in the fusion. In a particularly preferred embodiment, the adhesin amino acid sequence is fused to (a) 10 the hinge region and C<sub>H</sub>2 and C<sub>H</sub>3 or (b) the C<sub>H</sub>1, hinge, C<sub>H</sub>2 and C<sub>H</sub>3 domains, of an IgG heavy chain.

For bispecific immunoadhesins, the immunoadhesins are assembled as multimers, and particularly as heterodimers or heterotetramers. 15 Generally, these assembled immunoglobulins will have known unit structures. A basic four chain structural unit is the form in which IgG, IgD, and IgE exist. A four chain unit is repeated in the higher molecular weight immunoglobulins; IgM generally exists as a pentamer of four basic units held together by disulfide bonds. IgA globulin, and occasionally IgG globulin, may also exist in multimeric form in 20 serum. In the case of multimer, each of the four units may be the same or different.

Various exemplary assembled immunoadhesins within the scope herein are schematically diagrammed below:

- (a) AC<sub>L</sub>-AC<sub>L</sub>;
- 25 (b) AC<sub>H</sub>- (AC<sub>H</sub>, AC<sub>L</sub>-AC<sub>H</sub>, AC<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>, or V<sub>L</sub>C<sub>L</sub>-AC<sub>H</sub>) ;
- (c) AC<sub>L</sub>-AC<sub>H</sub>- (AC<sub>L</sub>-AC<sub>H</sub>, AC<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>, V<sub>L</sub>C<sub>L</sub>-AC<sub>H</sub>, or V<sub>L</sub>C<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>)
- (d) AC<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>- (AC<sub>H</sub>, or AC<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>, or V<sub>L</sub>C<sub>L</sub>-AC<sub>H</sub>) ;
- (e) V<sub>L</sub>C<sub>L</sub>-AC<sub>H</sub>- (AC<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>, or V<sub>L</sub>C<sub>L</sub>-AC<sub>H</sub>) ; and
- (f) (A-Y)<sub>n</sub>- (V<sub>L</sub>C<sub>L</sub>-V<sub>H</sub>C<sub>H</sub>)<sub>2</sub>,

30 wherein each A represents identical or different adhesin amino acid sequences;

V<sub>L</sub> is an immunoglobulin light chain variable domain;

V<sub>H</sub> is an immunoglobulin heavy chain variable domain;

C<sub>L</sub> is an immunoglobulin light chain constant domain;

35 C<sub>H</sub> is an immunoglobulin heavy chain constant domain;

n is an integer greater than 1;

Y designates the residue of a covalent cross-linking agent.

In the interests of brevity, the foregoing structures only show key features; they do not indicate joining (J) or other domains of 40 the immunoglobulins, nor are disulfide bonds shown. However, where such domains are required for binding activity, they shall be

constructed to be present in the ordinary locations which they occupy in the immunoglobulin molecules.

Alternatively, the adhesin sequences can be inserted between immunoglobulin heavy chain and light chain sequences, such that an 5 immunoglobulin comprising a chimeric heavy chain is obtained. In this embodiment, the adhesin sequences are fused to the 3' end of an immunoglobulin heavy chain in each arm of an immunoglobulin, either between the hinge and the C<sub>H</sub>2 domain, or between the C<sub>H</sub>2 and C<sub>H</sub>3 domains. Similar constructs have been reported by Hoogenboom et al., 10 Mol. Immunol., 28:1027-1037 (1991).

Although the presence of an immunoglobulin light chain is not required in the immunoadhesins of the present invention, an immunoglobulin light chain might be present either covalently associated to an adhesin-immunoglobulin heavy chain fusion 15 polypeptide, or directly fused to the adhesin. In the former case, DNA encoding an immunoglobulin light chain is typically coexpressed with the DNA encoding the adhesin-immunoglobulin heavy chain fusion protein. Upon secretion, the hybrid heavy chain and the light chain will be covalently associated to provide an immunoglobulin-like 20 structure comprising two disulfide-linked immunoglobulin heavy chain-light chain pairs. Methods suitable for the preparation of such structures are, for example, disclosed in U.S. Patent No. 4,816,567, issued 28 March 1989. Immunoadhesins are most conveniently 25 constructed by fusing the cDNA sequence encoding the adhesin portion in-frame to an immunoglobulin cDNA sequence. However, fusion to genomic immunoglobulin fragments can also be used (see, e.g. Aruffo et al., Cell, 61:1303-1313 (1990); and Stamenkovic et al., Cell, 66:1133-1144 (1991)). The latter type of fusion requires the presence of Ig regulatory sequences for expression. cDNAs encoding 30 IgG heavy-chain constant regions can be isolated based on published sequences from cDNA libraries derived from spleen or peripheral blood lymphocytes, by hybridization or by polymerase chain reaction (PCR) techniques. The cDNAs encoding the "adhesin" and the immunoglobulin parts of the immunoadhesin are inserted in tandem into a plasmid 35 vector that directs efficient expression in the chosen host cells.

In another embodiment, the WISP-1 or WISP-1 antagonist may be covalently modified by linking the receptor polypeptide to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol (PEG), polypropylene glycol, or polyoxyalkylenes, in the manner set 40 forth in U.S. Patent Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337, or other like molecules such as

polyglutamate. Such pegylated forms may be prepared using techniques known in the art.

Leucine zipper forms of these molecules are also contemplated by the invention. "Leucine zipper" is a term in the art used to refer to a leucine rich sequence that enhances, promotes, or drives dimerization or trimerization of its fusion partner (e.g., the sequence or molecule to which the leucine zipper is fused or linked to). Various leucine zipper polypeptides have been described in the art. See, e.g., Landschulz et al., Science, 240:1759 (1988); US Patent 5,716,805; WO 94/10308; Hoppe et al., FEBS Letters, 344:1991 (1994); Maniatis et al., Nature, 341:24 (1989). Those skilled in the art will appreciate that a leucine zipper sequence may be fused at either the 5' or 3' end of the WISP-1 or WISP-1 antagonist molecule.

The WISP-1 polypeptides of the present invention may also be modified in a way to form chimeric molecules by fusing the polypeptide to another, heterologous polypeptide or amino acid sequence. Preferably, such heterologous polypeptide or amino acid sequence is one which acts to oligomerize the chimeric molecule. In one embodiment, such a chimeric molecule comprises a fusion of the WISP-1 polypeptide with a tag polypeptide which provides an epitope to which an anti-tag antibody can selectively bind. The epitope tag is generally placed at the amino- or carboxyl- terminus of the polypeptide. The presence of such epitope-tagged forms of the polypeptide can be detected using an antibody against the tag polypeptide. Also, provision of the epitope tag enables the polypeptide to be readily purified by affinity purification using an anti-tag antibody or another type of affinity matrix that binds to the epitope tag. Various tag polypeptides and their respective antibodies are well known in the art. Examples include poly-histidine (poly-his) or poly-histidine-glycine (poly-his-gly) tags; the flu HA tag polypeptide and its antibody 12CA5 [Field et al., Mol. Cell. Biol., 8:2159-2165 (1988)]; the c-myc tag and the 8F9, 3C7, 6E10, G4, B7 and 9E10 antibodies thereto [Evan et al., Molecular and Cellular Biology, 5:3610-3616 (1985)]; and the Herpes Simplex virus glycoprotein D (gD) tag and its antibody [Paborsky et al., Protein Engineering, 3(6):547-553 (1990)]. Other tag polypeptides include the Flag-peptide [Hopp et al., BioTechnology, 6:1204-1210 (1988)]; the KT3 epitope peptide [Martin et al., Science, 255:192-194 (1992)]; an  $\alpha$ -tubulin epitope peptide [Skinner et al., J. Biol. Chem., 266:15163-15166 (1991)]; and the T7 gene 10 protein peptide tag

[Lutz-Freyermuth et al., Proc. Natl. Acad. Sci. USA, 87:6393-6397 (1990)].

It is contemplated that anti-WISP-1 antibodies may also be employed in the presently disclosed methods. Examples of such 5 molecules include neutralizing or blocking antibodies which can reduce cancer cell growth, metastasis or motility. The anti-WISP-1 may be monoclonal antibodies.

Monoclonal antibodies may be prepared using hybridoma methods, such as those described by Kohler and Milstein, Nature, 256:495 10 (1975). In a hybridoma method, a mouse, hamster, or other appropriate host animal, is typically immunized with an immunizing agent to elicit lymphocytes that produce or are capable of producing antibodies that will specifically bind to the immunizing agent. Alternatively, the lymphocytes may be immunized *in vitro*.

15 The immunizing agent will typically include a WISP-1 polypeptide or a fusion protein thereof, such as a WISP-1-IgG fusion protein.

Generally, either peripheral blood lymphocytes ("PBLs") are used if cells of human origin are desired, or spleen cells or lymph node cells are used if non-human mammalian sources are desired. The 20 lymphocytes are then fused with an immortalized cell line using a suitable fusing agent, such as polyethylene glycol, to form a hybridoma cell [Goding, Monoclonal Antibodies: Principles and Practice, Academic Press, (1986) pp. 59-103]. Immortalized cell lines are usually transformed mammalian cells, particularly myeloma cells of 25 rodent, bovine and human origin. Usually, rat or mouse myeloma cell lines are employed. The hybridoma cells may be cultured in a suitable culture medium that preferably contains one or more substances that inhibit the growth or survival of the unfused, immortalized cells. For example, if the parental cells lack the enzyme hypoxanthine 30 guanine phosphoribosyl transferase (HGPRT or HPRT), the culture medium for the hybridomas typically will include hypoxanthine, aminopterin, and thymidine ("HAT medium"), which substances prevent the growth of HGPRT-deficient cells.

Preferred immortalized cell lines are those that fuse 35 efficiently, support stable high level expression of antibody by the selected antibody-producing cells, and are sensitive to a medium such as HAT medium. More preferred immortalized cell lines are murine myeloma lines, which can be obtained, for instance, from the Salk Institute Cell Distribution Center, San Diego, California and the 40 American Type Culture Collection, Manassas, Virginia. Human myeloma and mouse-human heteromyeloma cell lines also have been described for

the production of human monoclonal antibodies [Kozbor, J. Immunol., 133:3001 (1984); Brodeur et al., Monoclonal Antibody Production Techniques and Applications, Marcel Dekker, Inc., New York, (1987) pp. 51-63].

5 The culture medium in which the hybridoma cells are cultured can then be assayed for the presence of monoclonal antibodies directed against WISP-1. Preferably, the binding specificity of monoclonal antibodies produced by the hybridoma cells is determined by immunoprecipitation or by an *in vitro* binding assay, such as 10 radioimmunoassay (RIA) or enzyme-linked immunoabsorbent assay (ELISA). Such techniques and assays are known in the art. The binding affinity of the monoclonal antibody can, for example, be determined by the Scatchard analysis of Munson and Pollard, Anal. Biochem., 107:220 (1980).

15 After the desired hybridoma cells are identified, the clones may be subcloned by limiting dilution procedures and grown by standard methods [Goding, supra]. Suitable culture media for this purpose include, for example, Dulbecco's Modified Eagle's Medium or RPMI-1640 medium. Alternatively, the hybridoma cells may be grown *in vivo* as 20 ascites in a mammal.

25 The monoclonal antibodies secreted by the subclones may be isolated or purified from the culture medium or ascites fluid by conventional immunoglobulin purification procedures such as, for example, protein A-Sepharose, hydroxylapatite chromatography, gel electrophoresis, dialysis, or affinity chromatography.

The monoclonal antibodies may also be made by recombinant DNA methods, such as those described in U.S. Patent No. 4,816,567. DNA encoding the monoclonal antibodies is readily isolated and sequenced using conventional procedures (e.g., by using oligonucleotide probes 30 that are capable of binding specifically to genes encoding the heavy and light chains of the monoclonal antibodies). The hybridoma cells serve as a preferred source of such DNA. Once isolated, the DNA may be placed into expression vectors, which are then transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster 35 ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, to obtain the synthesis of monoclonal antibodies in the recombinant host cells. The DNA also may be modified, for example, by substituting the coding sequence for human heavy and light chain constant domains in place of the homologous 40 murine sequences, Morrison, et al., Proc. Nat. Acad. Sci. 81, 6851 (1984), or by covalently joining to the immunoglobulin coding

sequence all or part of the coding sequence for a non-immunoglobulin polypeptide.

Typically such non-immunoglobulin polypeptides are substituted for the constant domains of an antibody of the invention, or they are substituted for the variable domains of one antigen-combining site of an antibody of the invention to create a chimeric bivalent antibody comprising one antigen-combining site having specificity for WISP-1 and another antigen-combining site having specificity for a different antigen.

10 Chimeric or hybrid antibodies also may be prepared in vitro using known methods in synthetic protein chemistry, including those involving crosslinking agents. For example, immunotoxins may be constructed using a disulfide exchange reaction or by forming a thioether bond. Examples of suitable reagents for this purpose 15 include iminothiolate and methyl-4-mercaptopbutyrimidate.

Single chain Fv fragments may also be produced, such as described in Iliades et al., FEBS Letters, 409:437-441 (1997). Coupling of such single chain fragments using various linkers is described in Kortt et al., Protein Engineering, 10:423-433 (1997). A 20 variety of techniques for the recombinant production and manipulation of antibodies are well known in the art. Illustrative examples of such techniques that are typically utilized by skilled artisans are described in greater detail below.

(i) *Humanized antibodies*

25 Generally, a humanized antibody has one or more amino acid residues introduced into it from a non-human source. These non-human amino acid residues are often referred to as "import" residues, which are typically taken from an "import" variable domain. Humanization can be essentially performed following the method of Winter and 30 co-workers [Jones et al., Nature, 321:522-525 (1986); Riechmann et al., Nature, 332:323-327 (1988); Verhoeyen et al., Science, 239:1534-1536 (1988)], by substituting rodent CDRs or CDR sequences for the corresponding sequences of a human antibody.

Accordingly, such "humanized" antibodies are chimeric 35 antibodies wherein substantially less than an intact human variable domain has been substituted by the corresponding sequence from a non-human species. In practice, humanized antibodies are typically human antibodies in which some CDR residues and possibly some FR residues are substituted by residues from analogous sites in rodent 40 antibodies.

It is important that antibodies be humanized with retention of high affinity for the antigen and other favorable biological properties. To achieve this goal, according to a preferred method, humanized antibodies are prepared by a process of analysis of the 5 parental sequences and various conceptual humanized products using three dimensional models of the parental and humanized sequences. Three dimensional immunoglobulin models are commonly available and are familiar to those skilled in the art. Computer programs are available which illustrate and display probable three-dimensional 10 conformational structures of selected candidate immunoglobulin sequences. Inspection of these displays permits analysis of the likely role of the residues in the functioning of the candidate immunoglobulin sequence, i.e. the analysis of residues that influence the ability of the candidate immunoglobulin to bind its antigen. In 15 this way, FR residues can be selected and combined from the consensus and import sequence so that the desired antibody characteristic, such as increased affinity for the target antigen(s), is achieved. In general, the CDR residues are directly and most substantially involved in influencing antigen binding.

20 (ii) *Human antibodies*

Human monoclonal antibodies can be made by the hybridoma method. Human myeloma and mouse-human heteromyeloma cell lines for the production of human monoclonal antibodies have been described, for example, by Kozbor, J. Immunol. 133, 3001 (1984), and Brodeur, et 25 al., Monoclonal Antibody Production Techniques and Applications, pp.51-63 (Marcel Dekker, Inc., New York, 1987).

It is now possible to produce transgenic animals (e.g. mice) that are capable, upon immunization, of producing a repertoire of human antibodies in the absence of endogenous immunoglobulin production. 30 For example, it has been described that the homozygous deletion of the antibody heavy chain joining region ( $J_H$ ) gene in chimeric and germ-line mutant mice results in complete inhibition of endogenous antibody production. Transfer of the human germ-line immunoglobulin gene array in such germ-line mutant mice will result in the 35 production of human antibodies upon antigen challenge. See, e.g. Jakobovits et al., Proc. Natl. Acad. Sci. USA 90, 2551-255 (1993); Jakobovits et al., Nature 362, 255-258 (1993).

Mendez et al. (Nature Genetics 15: 146-156 [1997]) have further 40 improved the technology and have generated a line of transgenic mice designated as "Xenomouse II" that, when challenged with an antigen, generates high affinity fully human antibodies. This was achieved by

germ-line integration of megabase human heavy chain and light chain loci into mice with deletion into endogenous  $J_H$  segment as described above. The Xenomouse II harbors 1,020 kb of human heavy chain locus containing approximately 66  $V_H$  genes, complete  $D_H$  and  $J_H$  regions and 5 three different constant regions ( $\mu$ ,  $\delta$  and  $\gamma$ ), and also harbors 800 kb of human  $\kappa$  locus containing 32  $V_K$  genes,  $J_K$  segments and  $C_K$  genes. The antibodies produced in these mice closely resemble that seen in humans in all respects, including gene rearrangement, assembly, and 10 repertoire. The human antibodies are preferentially expressed over endogenous antibodies due to deletion in endogenous  $J_H$  segment that prevents gene rearrangement in the murine locus.

Alternatively, the phage display technology (McCafferty et al., Nature 348, 552-553 [1990]) can be used to produce human antibodies and antibody fragments *in vitro*, from immunoglobulin variable (V) 15 domain gene repertoires from unimmunized donors. According to this technique, antibody V domain genes are cloned in-frame into either a major or minor coat protein gene of a filamentous bacteriophage, such as M13 or fd, and displayed as functional antibody fragments on the surface of the phage particle. Because the filamentous particle 20 contains a single-stranded DNA copy of the phage genome, selections based on the functional properties of the antibody also result in selection of the gene encoding the antibody exhibiting those properties. Thus, the phage mimicks some of the properties of the B-cell. Phage display can be performed in a variety of formats; for 25 their review see, e.g. Johnson, Kevin S. and Chiswell, David J., Current Opinion in Structural Biology 3, 564-571 (1993). Several sources of V-gene segments can be used for phage display. Clackson et al., Nature 352, 624-628 (1991) isolated a diverse array of anti-oxazolone antibodies from a small random combinatorial library of V 30 genes derived from the spleens of immunized mice. A repertoire of V genes from unimmunized human donors can be constructed and antibodies to a diverse array of antigens (including self-antigens) can be isolated essentially following the techniques described by Marks et al., J. Mol. Biol. 222, 581-597 (1991), or Griffith et al., EMBO J. 35, 12, 725-734 (1993). In a natural immune response, antibody genes 35 accumulate mutations at a high rate (somatic hypermutation). Some of the changes introduced will confer higher affinity, and B cells displaying high-affinity surface immunoglobulin are preferentially replicated and differentiated during subsequent antigen challenge.

40 This natural process can be mimicked by employing the technique known

as "chain shuffling" (Marks et al., Bio/Technol. 10, 779-783 [1992]).

In this method, the affinity of "primary" human antibodies obtained by phage display can be improved by sequentially replacing the heavy and light chain V region genes with repertoires of naturally

5 occurring variants (repertoires) of V domain genes obtained from unimmunized donors. This technique allows the production of antibodies and antibody fragments with affinities in the nM range. A strategy for making very large phage antibody repertoires (also known as "the mother-of-all libraries") has been described by Waterhouse et 10 al., Nucl. Acids Res. 21, 2265-2266 (1993). Gene shuffling can also be used to derive human antibodies from rodent antibodies, where the human antibody has similar affinities and specificities to the starting rodent antibody. According to this method, which is also referred to as "epitope imprinting", the heavy or light chain V 15 domain gene of rodent antibodies obtained by phage display technique is replaced with a repertoire of human V domain genes, creating rodent-human chimeras. Selection on antigen results in isolation of human variable capable of restoring a functional antigen-binding site, i.e. the epitope governs (imprints) the choice of partner.

20 When the process is repeated in order to replace the remaining rodent V domain, a human antibody is obtained (see PCT patent application WO 93/06213, published 1 April 1993). Unlike traditional humanization of rodent antibodies by CDR grafting, this technique provides completely human antibodies, which have no framework or CDR residues 25 of rodent origin.

As discussed below, the antibodies of the invention may optionally comprise monomeric, antibodies, dimeric antibodies, as well as multivalent forms of antibodies. Those skilled in the art may construct such dimers or multivalent forms by techniques known in the 30 art. Methods for preparing monovalent antibodies are also well known in the art. For example, one method involves recombinant expression of immunoglobulin light chain and modified heavy chain. The heavy chain is truncated generally at any point in the Fc region so as to prevent heavy chain crosslinking. Alternatively, the relevant 35 cysteine residues are substituted with another amino acid residue or are deleted so as to prevent crosslinking.

*(iii) Bispecific antibodies*

Bispecific antibodies are monoclonal, preferably human or humanized, antibodies that have binding specificities for at least 40 two different antigens. In the present case, one of the binding specificities is for WISP-1. For example, bispecific antibodies

specifically binding WISP-1 or WISP-1 variants and another CNN family member (e.g., WISP-2, WISP-3, CTGF, Cyr61, or Nov) or other molecules such as CD44 are within the scope of the present invention.

Methods for making bispecific antibodies are known in the art. 5 Traditionally, the recombinant production of bispecific antibodies is based on the coexpression of two immunoglobulin heavy chain-light chain pairs, where the two heavy chains have different specificities (Millstein and Cuello, Nature 305, 537-539 (1983)). Because of the random assortment of immunoglobulin heavy and light chains, these 10 hybridomas (quadromas) produce a potential mixture of 10 different antibody molecules, of which only one has the correct bispecific structure. The purification of the correct molecule, which is usually done by affinity chromatography steps, is rather cumbersome, and the product yields are low. Similar procedures are disclosed in 15 PCT application publication No. WO 93/08829 (published 13 May 1993), and in Traunecker et al., EMBO 10, 3655-3659 (1991).

According to a different and more preferred approach, antibody variable domains with the desired binding specificities (antibody-antigen combining sites) are fused to immunoglobulin constant domain 20 sequences. The fusion preferably is with an immunoglobulin heavy chain constant domain, comprising at least part of the hinge, CH2 and CH3 regions. It is preferred to have the first heavy chain constant region (CH1) containing the site necessary for light chain binding, present in at least one of the fusions. DNAs encoding the 25 immunoglobulin heavy chain fusions and, if desired, the immunoglobulin light chain, are inserted into separate expression vectors, and are cotransfected into a suitable host organism. This provides for great flexibility in adjusting the mutual proportions of the three polypeptide fragments in embodiments when unequal ratios of 30 the three polypeptide chains used in the construction provide the optimum yields. It is, however, possible to insert the coding sequences for two or all three polypeptide chains in one expression vector when the expression of at least two polypeptide chains in equal ratios results in high yields or when the ratios are of no 35 particular significance. In a preferred embodiment of this approach, the bispecific antibodies are composed of a hybrid immunoglobulin heavy chain with a first binding specificity in one arm, and a hybrid immunoglobulin heavy chain-light chain pair (providing a second binding specificity) in the other arm. It was found that this 40 asymmetric structure facilitates the separation of the desired bispecific compound from unwanted immunoglobulin chain combinations,

as the presence of an immunoglobulin light chain in only one half of the bispecific molecule provides for a facile way of separation. This approach is disclosed in PCT Publication No. WO 94/04690, published on March 3, 1994.

5 For further details of generating bispecific antibodies see, for example, Suresh et al., Methods in Enzymology 121, 210 (1986).

(iv) *Heteroconjugate antibodies*

10 Heteroconjugate antibodies are also within the scope of the present invention. Heteroconjugate antibodies are composed of two covalently joined antibodies. Such antibodies have, for example, been proposed to target immune system cells to unwanted cells (U.S. Patent No. 4,676,980), and for treatment of HIV infection (PCT application publication Nos. WO 91/00360 and WO 92/200373; EP 03089).

15 Heteroconjugate antibodies may be made using any convenient cross-linking methods. Suitable cross-linking agents are well known in the art, and are disclosed in U.S. Patent No. 4,676,980, along with a number of cross-linking techniques.

(v) *Antibody fragments*

20 In certain embodiments, the anti-WISP-1 antibody (including murine, human and humanized antibodies, and antibody variants) is an antibody fragment. Various techniques have been developed for the production of antibody fragments. Traditionally, these fragments were derived via proteolytic digestion of intact antibodies (see, e.g., Morimoto et al., *J. Biochem. Biophys. Methods* 24:107-117 (1992) and Brennan et al., *Science* 229:81 (1985)). However, these fragments can now be produced directly by recombinant host cells. For example, Fab'-SH fragments can be directly recovered from *E. coli* and chemically coupled to form F(ab')<sub>2</sub> fragments (Carter et al., *Bio/Technology* 10:163-167 (1992)). In another embodiment, the F(ab')<sub>2</sub> 25 is formed using the leucine zipper GCN4 to promote assembly of the F(ab')<sub>2</sub> molecule. According to another approach, Fv, Fab or F(ab')<sub>2</sub> fragments can be isolated directly from recombinant host cell culture. A variety of techniques for the production of antibody fragments will be apparent to the skilled practitioner. For 30 instance, digestion can be performed using papain. Examples of papain digestion are described in WO 94/29348 published 12/22/94 and U.S. Patent No. 4,342,566. Papain digestion of antibodies typically produces two identical antigen binding fragments, called Fab fragments, each with a single antigen binding site, and a residual Fc 35 fragment. Pepsin treatment yields an F(ab')<sub>2</sub> fragment that has two 40

antigen combining sites and is still capable of cross-linking antigen.

The Fab fragments produced in the antibody digestion also contain the constant domains of the light chain and the first constant 5 domain (CH<sub>1</sub>) of the heavy chain. Fab' fragments differ from Fab fragments by the addition of a few residues at the carboxy terminus of the heavy chain CH<sub>1</sub> domain including one or more cysteines from the antibody hinge region. Fab'-SH is the designation herein for Fab' in which the cysteine residue(s) of the constant domains bear a free 10 thiol group. F(ab')<sub>2</sub> antibody fragments originally were produced as pairs of Fab' fragments which have hinge cysteines between them. Other chemical couplings of antibody fragments are also known.

Antibodies are glycosylated at conserved positions in their constant regions (Jefferis and Lund, Chem. Immunol. 65:111-128 15 [1997]; Wright and Morrison, TibTECH 15:26-32 [1997]). The oligosaccharide side chains of the immunoglobulins affect the protein's function (Boyd et al., Mol. Immunol. 32:1311-1318 [1996]; Wittwe and Howard, Biochem. 29:4175-4180 [1990]), and the intramolecular interaction between portions of the glycoprotein 20 which can affect the conformation and presented three-dimensional surface of the glycoprotein (Hefferis and Lund, *supra*; Wyss and Wagner, Current Opin. Biotech. 7:409-416 [1996]). Oligosaccharides may also serve to target a given glycoprotein to certain molecules based upon specific recognition structures. For example, it has 25 been reported that in agalactosylated IgG, the oligosaccharide moiety 'flips' out of the inter-CH<sub>2</sub> space and terminal N-acetylglucosamine residues become available to bind mannose binding protein (Malhotra et al., Nature Med. 1:237-243 [1995]). Removal by glycopeptidase of the oligosaccharides from CAMPATH-1H (a 30 recombinant humanized murine monoclonal IgG1 antibody which recognizes the CDw52 antigen of human lymphocytes) produced in Chinese Hamster Ovary (CHO) cells resulted in a complete reduction in complement mediated lysis (CMCL) (Boyd et al., Mol. Immunol. 32:1311-1318 [1996]), while selective removal of sialic acid 35 residues using neuraminidase resulted in no loss of DMCL. Glycosylation of antibodies has also been reported to affect antibody-dependent cellular cytotoxicity (ADCC). In particular, CHO cells with tetracycline-regulated expression of  $\beta(1,4)$ -N-acetylglucosaminyltransferase III (GnTIII), a glycosyltransferase 40 catalyzing formation of bisecting GlcNAc, was reported to have

improved ADCC activity (Umana *et al.*, Mature Biotech. **17**:176-180 [1999]).

Glycosylation variants of antibodies are variants in which the glycosylation pattern of an antibody is altered. By altering is 5 meant deleting one or more carbohydrate moieties found in the antibody, adding one or more carbohydrate moieties to the antibody, changing the composition of glycosylation (glycosylation pattern), the extent of glycosylation, etc. Glycosylation variants may, for example, be prepared by removing, changing and/or adding one or more 10 glycosylation sites in the nucleic acid sequence encoding the antibody.

Glycosylation of antibodies is typically either N-linked or O-linked. N-linked refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide 15 sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. 20 O-linked glycosylation refers to the attachment of one of the sugars N-acetylglactosamine, galactose, or xylose to a hydroxyamino acid, most commonly serine or threonine, although 5-hydroxyproline or 5-hydroxylysine may also be used.

Addition of glycosylation sites to the antibody is 25 conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). The alteration may also be made by the addition of, or substitution by, one or more serine or threonine residues to the sequence of the original 30 antibody (for O-linked glycosylation sites).

The glycosylation (including glycosylation pattern) of antibodies may also be altered without altering the underlying nucleotide sequence. Glycosylation largely depends on the host cell used to express the antibody. Since the cell type used for 35 expression of recombinant glycoproteins, e.g. antibodies, as potential therapeutics is rarely the native cell, significant variations in the glycosylation pattern of the antibodies can be expected (see, e.g. Hse *et al.*, J. Biol. Chem. **272**:9062-9070 [1997]). In addition to the choice of host cells, factors which 40 affect glycosylation during recombinant production of antibodies include growth mode, media formulation, culture density,

oxygenation, pH, purification schemes and the like. Various methods have been proposed to alter the glycosylation pattern achieved in a particular host organism including introducing or overexpressing certain enzymes involved in oligosaccharide production (U. S. Patent Nos. 5,047,335; 5,510,261 and 5,278,299). Glycosylation, or certain types of glycosylation, can be enzymatically removed from the glycoprotein, for example using endoglycosidase H (Endo H). In addition, the recombinant host cell can be genetically engineered, e.g. make defective in processing certain types of polysaccharides.

10 These and similar techniques are well known in the art.

The glycosylation structure of antibodies can be readily analyzed by conventional techniques of carbohydrate analysis, including lectin chromatography, NMR, Mass spectrometry, HPLC, GPC, monosaccharide compositional analysis, sequential enzymatic digestion, and HPAEC-PAD, which uses high pH anion exchange chromatography to separate oligosaccharides based on charge. Methods for releasing oligosaccharides for analytical purposes are also known, and include, without limitation, enzymatic treatment (commonly performed using peptide-N-glycosidase F/endo- $\beta$ -galactosidase), elimination using harsh alkaline environment to release mainly O-linked structures, and chemical methods using anhydrous hydrazine to release both N- and O-linked oligosaccharides.

20 Triabodies are also within the scope of the invention. Such antibodies are described for instance in Iliades et al., supra and Kortt et al., supra.

The antibodies of the present invention may be modified by conjugating the antibody to a cytotoxic agent (like a toxin molecule) or a prodrug-activating enzyme which converts a prodrug (e.g. a peptidyl chemotherapeutic agent, see WO81/01145) to an active anti-30 cancer drug. See, for example, WO 88/07378 and U.S. Patent No. 4,975,278. This technology is also referred to as "Antibody Dependent Enzyme Mediated Prodrug Therapy" (ADEPT).

35 The enzyme component of the immunoconjugate useful for ADEPT includes any enzyme capable of acting on a prodrug in such a way so as to convert it into its more active, cytotoxic form. Enzymes that are useful in the method of this invention include, but are not limited to, alkaline phosphatase useful for converting phosphate-containing prodrugs into free drugs; arylsulfatase useful for converting sulfate-containing prodrugs into free drugs; cytosine 40 deaminase useful for converting non-toxic 5-fluorocytosine into the

anti-cancer drug, 5-fluorouracil; proteases, such as serratia protease, thermolysin, subtilisin, carboxypeptidases and cathepsins (such as cathepsins B and L), that are useful for converting peptide-containing prodrugs into free drugs; caspases such as caspase-3; D-5 alanylcarboxypeptidases, useful for converting prodrugs that contain D-amino acid substituents; carbohydrate-cleaving enzymes such as beta-galactosidase and neuraminidase useful for converting glycosylated prodrugs into free drugs; beta-lactamase useful for converting drugs derivatized with beta-lactams into free drugs; and 10 penicillin amidases, such as penicillin V amidase or penicillin G amidase, useful for converting drugs derivatized at their amine nitrogens with phenoxyacetyl or phenylacetyl groups, respectively, into free drugs. Alternatively, antibodies with enzymatic activity, also known in the art as "abzymes", can be used to convert the 15 prodrugs of the invention into free active drugs (see, e.g., Massey, *Nature* 328: 457-458 (1987)). Antibody-abzyme conjugates can be prepared as described herein for delivery of the abzyme to a tumor cell population.

The enzymes can be covalently bound to the antibodies by 20 techniques well known in the art such as the use of heterobifunctional crosslinking reagents. Alternatively, fusion proteins comprising at least the antigen binding region of an antibody of the invention linked to at least a functionally active portion of an enzyme of the invention can be constructed using 25 recombinant DNA techniques well known in the art (see, e.g., Neuberger *et al.*, *Nature*, 312: 604-608 (1984)).

Further antibody modifications are contemplated. For example, the antibody may be linked to one of a variety of nonproteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, 30 polyoxyalkylenes, or copolymers of polyethylene glycol and polypropylene glycol, or other molecules such as polyglutamate. The antibody also may be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization (for example, hydroxymethylcellulose or gelatin-microcapsules, and 35 poly-(methylmethacrylate) microcapsules, respectively), in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules), or in macroemulsions. Such techniques are disclosed in Remington's *Pharmaceutical Sciences*, 16th edition, Osol, A., Ed., (1980). To 40 increase the serum half life of the antibody, one may incorporate a salvage receptor binding epitope into the antibody (especially an

antibody fragment) as described in U.S. Patent 5,739,277, for example. As used herein, the term "salvage receptor binding epitope" refers to an epitope of the Fc region of an IgG molecule (e.g., IgG<sub>1</sub>, IgG<sub>2</sub>, IgG<sub>3</sub>, or IgG<sub>4</sub>) that is responsible for increasing the *in vivo* 5 serum half-life of the IgG molecule.

#### FORMULATIONS

The WISP-1 antagonists described herein, are optionally employed in a carrier. Suitable carriers and their formulations are described in Remington's Pharmaceutical Sciences, 16th ed., 1980, Mack 10 Publishing Co., edited by Osol et al. Typically, an appropriate amount of a pharmaceutically-acceptable salt is used in the carrier to render the formulation isotonic. Examples of the carrier include saline, Ringer's solution and dextrose solution. The pH of the carrier is preferably from about 5 to about 8, and more preferably 15 from about 7.4 to about 7.8. It will be apparent to those persons skilled in the art that certain carriers may be more preferable depending upon, for instance, the route of administration and concentration of active agent being administered. The carrier may be in the form of a lyophilized formulation or aqueous solution.

Acceptable carriers, excipients, or stabilizers are preferably nontoxic to cells and/or recipients at the dosages and concentrations employed, and include buffers such as phosphate, citrate, and other organic acids; antioxidants including ascorbic acid and methionine; preservatives (such as octadecyldimethylbenzyl ammonium chloride; 25 hexamethonium chloride; benzalkonium chloride, benzethonium chloride; phenol, butyl or benzyl alcohol; alkyl parabens such as methyl or propyl paraben; catechol; resorcinol; cyclohexanol; 3-pentanol; and m-cresol); low molecular weight (less than about 10 residues) polypeptides; proteins, such as serum albumin, gelatin, or 30 immunoglobulins; hydrophilic polymers such as polyvinylpyrrolidone; amino acids such as glycine, glutamine, asparagine, histidine, arginine, or lysine; monosaccharides, disaccharides, and other carbohydrates including glucose, mannose, or dextrins; chelating agents such as EDTA; sugars such as sucrose, mannitol, trehalose or 35 sorbitol; salt-forming counter-ions such as sodium; and/or non-ionic surfactants such as TWEEN<sup>TM</sup>, PLURONICS<sup>TM</sup> or polyethylene glycol (PEG).

The formulation may also contain more than one active compound as necessary for the particular indication being treated, preferably those with complementary activities that do not adversely affect each 40 other.

The antagonists described herein may also be entrapped in microcapsules prepared, for example, by coacervation techniques or by interfacial polymerization, for example, hydroxymethylcellulose or gelatin-microcapsules and poly-(methylmethacrylate) microcapsules, 5 respectively, in colloidal drug delivery systems (for example, liposomes, albumin microspheres, microemulsions, nano-particles and nanocapsules) or in macroemulsions. Such techniques are disclosed in Remington's Pharmaceutical Sciences 16th edition, Osol, A. Ed. (1980).

10 The formulations to be used for *in vivo* administration should be sterile. This is readily accomplished by filtration through sterile filtration membranes.

Sustained-release preparations may be prepared. Suitable examples of sustained-release preparations include semipermeable 15 matrices of solid hydrophobic polymers containing the active agent, which matrices are in the form of shaped articles, e.g. films, or microcapsules. Examples of sustained-release matrices include polyesters, hydrogels (for example, poly(2-hydroxyethyl-methacrylate), or poly(vinylalcohol)), polylactides (U.S. Pat. No. 20 3,773,919), copolymers of L-glutamic acid and  $\gamma$  ethyl-L-glutamate, non-degradable ethylene-vinyl acetate, degradable lactic acid-glycolic acid copolymers such as the LUPRON DEPOT<sup>TM</sup> (injectable microspheres composed of lactic acid-glycolic acid copolymer and leuprolide acetate), and poly-D-(-)-3-hydroxybutyric acid. While 25 polymers such as ethylene-vinyl acetate and lactic acid-glycolic acid enable release of molecules for over 100 days, certain hydrogels release proteins for shorter time periods.

#### MODES OF THERAPY

30 The molecules described herein are useful in treating various pathological conditions, such as cancer. These conditions can be treated by administration of one or more antagonists described herein.

35 Diagnosis in mammals of the various pathological conditions described herein can be made by the skilled practitioner. Diagnostic techniques are available in the art which allow, e.g., for the diagnosis or detection of cancer or immune related disease in a mammal. For instance, cancers may be identified through techniques, including but not limited to, palpation, blood analysis, x-ray, NMR and the like.

40 The antagonist(s) can be administered in accord with known methods, such as intravenous administration as a bolus or by

continuous infusion over a period of time, by intramuscular, intraperitoneal, intracerebrospinal, subcutaneous, intra-articular, intrasynovial, intrathecal, oral, topical, or inhalation routes. Optionally, administration may be performed through mini-pump 5 infusion using various commercially available devices. The antagonists may also be employed using gene therapy techniques which have been described in the art.

Effective dosages and schedules for administering antagonists may be determined empirically, and making such determinations is 10 within the skill in the art. Single or multiple dosages may be employed. It is presently believed that an effective dosage or amount of antagonist used alone may range from about 1  $\mu$ g/kg to about 100 mg/kg of body weight or more per day. Interspecies scaling of dosages can be performed in a manner known in the art, e.g., as disclosed in 15 Mordenti et al., Pharmaceut. Res., 8:1351 (1991).

When *in vivo* administration of an antagonist thereof is employed, normal dosage amounts may vary from about 10 ng/kg to up to 100 mg/kg of mammal body weight or more per day, preferably about 1  $\mu$ g/kg/day to 10 mg/kg/day, depending upon the route of 20 administration. Guidance as to particular dosages and methods of delivery is provided in the literature; see, for example, U.S. Pat. Nos. 4,657,760; 5,206,344; or 5,225,212. It is anticipated that different formulations will be effective for different treatment 25 compounds and different disorders, that administration targeting one organ or tissue, for example, may necessitate delivery in a manner different from that to another organ or tissue. Those skilled in the art will understand that the dosage of antagonist that must be administered will vary depending on, for example, the mammal which will receive the agonist or antagonist, the route of administration, 30 and other drugs or therapies being administered to the mammal.

Depending on the type of cells and/or severity of the disease, about 1  $\mu$ g/kg to 15 mg/kg (*e.g.* 0.1-20 mg/kg) of antagonist antibody is an initial candidate dosage for administration, whether, for example, by one or more separate administrations, or by continuous 35 infusion. A typical daily dosage might range from about 1  $\mu$ g/kg to 100 mg/kg or more, depending on the factors mentioned above. For repeated administrations over several days or longer, depending on the condition, the treatment is sustained until a desired suppression of disease symptoms occurs. However, other dosage regimens may be 40 useful.

A single type of antagonist may be used in the methods of the invention. For example, a WISP-1 immunoadhesin molecule may be administered. Alternatively, the skilled practitioner may opt to employ a combination of antagonists in the methods, e.g., a 5 combination of a WISP-1 immunoadhesin and WISP-1 antibody. It is contemplated that yet additional therapies may be employed in the methods. The one or more other therapies may include but are not limited to, administration of radiation therapy, cytokine(s), growth inhibitory agent(s), chemotherapeutic agent(s), cytotoxic agent(s), 10 tyrosine kinase inhibitors, ras farnesyl transferase inhibitors, angiogenesis inhibitors, and cyclin-dependent kinase inhibitors which are known in the art and defined further with particularity in Section I above. In addition, therapies based on therapeutic antibodies that target tumor antigens such as Rituxan™ or Herceptin™ as well as anti- 15 angiogenic antibodies such as anti-VEGF.

Preparation and dosing schedules for chemotherapeutic agents may be used according to manufacturers' instructions or as determined empirically by the skilled practitioner. Preparation and dosing schedules for such chemotherapy are also described in *Chemotherapy 20 Service Ed.*, M.C. Perry, Williams & Wilkins, Baltimore, MD (1992). The chemotherapeutic agent may precede, or follow administration of, e.g. an antagonist, or may be given simultaneously therewith. The antagonist, for instance, may also be combined with an anti-oestrogen compound such as tamoxifen or an anti-progesterone such as 25 onapristone (see, EP 616812) in dosages known for such molecules.

It may be desirable to also administer antibodies against other antigens, such as antibodies which bind to CD20, CD11a, CD18, CD40, CD44, ErbB2, EGFR, ErbB3, ErbB4, vascular endothelial factor (VEGF), or other TNFR family members (such as DR4, DR5, OPG, TNFR1, TNFR2). 30 Alternatively, or in addition, two or more antibodies binding the same or two or more different antigens disclosed herein may be co-administered to the patient. Sometimes, it may be beneficial to also administer one or more cytokines to the patient. In one embodiment, the antagonists herein are co-administered with a growth inhibitory 35 agent. For example, the growth inhibitory agent may be administered first, followed by an antagonist of the present invention.

The antagonist and one or more other therapies may be administered concurrently or sequentially. Following administration of antagonist, treated cells *in vitro* can be analyzed. Where there 40 has been *in vivo* treatment, a treated mammal can be monitored in various ways well known to the skilled practitioner.

ARTICLES OF MANUFACTURE

In another embodiment of the invention, an article of manufacture containing materials useful for the treatment of the disorders described above is provided. The article of manufacture 5 comprises a container and a label. Suitable containers include, for example, bottles, vials, syringes, and test tubes. The containers may be formed from a variety of materials such as glass or plastic. The container holds a composition which is effective for treating the condition and may have a sterile access port (for example the 10 container may be an intravenous solution bag or a vial having a stopper pierceable by a hypodermic injection needle). The active agents in the composition may comprise antagonist(s). The label on, or associated with, the container indicates that the composition is used for treating the condition of choice. The article of 15 manufacture may further comprise a second container comprising a pharmaceutically-acceptable buffer, such as phosphate-buffered saline, Ringer's solution and dextrose solution. It may further include other materials desirable from a commercial and user standpoint, including other buffers, diluents, filters, needles, 20 syringes, and package inserts with instructions for use.

The following examples are offered for illustrative purposes only, and are not intended to limit the scope of the present invention in any way.

All patent and literature references cited in the present 25 specification are hereby incorporated by reference in their entirety.

EXAMPLES

Commercially available reagents referred to in the examples were used according to manufacturer's instructions unless otherwise 30 indicated. The source of those cells identified in the following examples, and throughout the specification, by ATCC accession numbers is the American Type Culture Collection, Manassas, VA. Unless otherwise noted, the present invention uses standard procedures of recombinant DNA technology, such as those described hereinabove and 35 in the following textbooks: Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press N.Y., 1989; Ausubel et al., *Current Protocols in Molecular Biology*, Green Publishing Associates and Wiley Interscience, N.Y., 1989; Innis et al., *PCR Protocols: A Guide to Methods and Applications*, Academic Press, Inc., 40 N.Y., 1990; Harlow et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, 1988; Gait, M.J.,

*Oligonucleotide Synthesis*, IRL Press, Oxford, 1984; R.I. Freshney, *Animal Cell Culture*, 1987; Coligan et al., *Current Protocols in Immunology*, 1991.

5 A description of the assays and materials referred to in the following Examples is provided below:

Proteins and Antibodies: Full length human WISP-1-Fc fusion protein was expressed and purified as described previously (Desnoyers et al., *J. Biol. Chem.*, 276:47599-47607 (2001)). Purified mouse anti-rat CD44 antibody (clone OX49) was from BD Bioscience (Bedford 10 MA). The mouse IgG2a isotype control antibody was from Pharmingen. The FITC conjugated mouse anti-rat CD44 was from Serotec (Raleigh, NC) and the FITC conjugated mouse IgG1 isotype control antibody was from DAKO (Denmark). The rat anti-mouse CD44 (clone KM114) was from Pharmingen. The actin antibody (clone C4) was from ICN biomedicals 15 (Aurora, OH). Murine anti- human WISP-1 monoclonal antibodies were generated and selected using WISP-1-Fc as an immunogen and the protocols and reagents described in the Examples of US Patent No. 6,387,657 issued May 14, 2002. Five such monoclonal antibodies have been deposited by Applicants with the ATCC, as noted below.

20 Cells and Tissue Specimens: NRK-49F and SW480 cell lines (ATCC, Manassas, VA) were maintained in High glucose Dulbecco's Modified Eagle's Medium (HGDMEM) supplemented with 10% fetal calf serum (FBS). Parental C57MG, C57MG expressing Wnt-1 (C57MG/Wnt-1), NRK-49F expressing high levels of hWISP-1 (NRK/WISP-1H), NRK-49F 25 expressing lower levels of hWISP-1 (NRK/WISP-1L) and NRK-49F containing an empty vector (NRK/control) were obtained from Arnold Levine (Princeton University, Princeton, NJ) and maintained in media containing 2.5  $\mu$ g ml<sup>-1</sup> puromycin (Xu et al., *Genes Dev.*, 14:585-595 (2000)). Wnt-1 transgenic mice obtained from Harold Varmus (National 30 Cancer Institute, National Institutes of Health, Bethesda, MD) were bred with p53 null mice obtained from Jackson Laboratories (Bar Harbor, MA) to generate the Wnt-1 transgenic/p53 knock out mice. Specimens of breast tumor and mammary duct epithelium were excised from those mice for analysis.

35 Particle Exclusion Assay: The erythrocyte sedimentation assay was used as described previously, with minor modifications. (Knudson et al., *J. Cell Sci.*, 99:227-235 (1991)). NRK/WISP-1 ((normal rat kidney fibroblasts transfected with WISP-1; see Xu et al., *Genes Dev.*, 14:585-595 (2000)) and NRK/control cells (normal rat kidney 40 fibroblasts; obtained from Arnold Levine, Princeton University, Princeton, NJ) (1 X 10<sup>5</sup> cells/well) were seeded in Falcon 6 well

plates (Becton Dickinson, Franklin Lanes, NJ) and maintained in high glucose Dulbecco's modified Eagle's Medium supplemented with 10% fetal bovine serum (FBS) (Life Technologies, Gaithersburg, MD) overnight at 37° C, 5% CO<sub>2</sub>. The next day, the media was removed and 5 10<sup>8</sup> sheep RBCs (Inter-Cell Technologies, Hopewell, NJ) in 750µl PBS/0.1% BSA (Boehringer Mannheim, Indianapolis, IN) were added and allowed to settle for 15 minutes. The cells were observed using a Nikon Diaphot 300 inverted microscope, and digital images were acquired using a SONY DSC-5000 digital photo camera (Japan).

10 Hyaluronan Staining of Cultured Cells: Hyaluronan Staining was performed as previously described, with minor modifications (Pienimaki et al., J. Biol. Chem., 276:20428-20435 (2001)). NRK/WISP-1 and NRK/control cells (8-10 X 10<sup>3</sup>) were plated in 8 well plastic chamber slides and maintained (as described above) 15 overnight at 37°C, 5% CO<sub>2</sub>. The next day the cells were washed with phosphate buffered saline (PBS) and fixed with 4% paraformaldehyde in PBS for 20 minutes at room temperature. The cells were washed with PBS and the non-specific binding sites were saturated for 30 minutes with distilled water containing 1% bovine serum albumin (BSA) and 20 0.1% Triton X-100 (Calbiochem, LaJolla, CA). Biotinylated HA binding protein (HA-BP) (Seikagaku America, Falmouth, MA) (5µg/ml in PBS/3% BSA) was added and incubated overnight at room temperature. The next day the cells were washed with PBS and incubated for 30 minutes with 1:1000 FITC-conjugated streptavidin (DAKO, Carpinteria, CA) in TBS 25 and 1%/BSA. The cells were washed in PBS, mounted using Vectashield (Burlingame, CA) containing 1 µg/ml Hoechst 33342 (Molecular Probes, Eugene, OR) and visualized under a Nikon Eclipse 800 fluorescent microscope. Images were acquired with a Photometrics 300 CCD camera operated at -40°C interfaced with an Apple G3 computer. The same 30 procedure was also used to stain frozen sections of mammary tumors from Wnt-1 transgenic mice).

Real time RT-PCR: Relative RNA expression level was determined using real-time RT-PCR as previously reported (Winer et al., Anal. Biochem., 270:41-49 (1999)). Total RNA was extracted from cells with 35 Tri-Reagent using the manufacturer's protocol (Molecular Research Center). Specific primers and fluorogenic probes were designed and used to amplify and quantitate gene expression level. The genes' specific signals were normalized to that of the glyceraldehyde-3-phosphate dehydrogenase housekeeping gene. Triplicate sets of data 40 were averaged for each condition. All TaqMan RT-PCR reagents were purchased from Applied Biosystems (Foster City, CA). To test the

effect of recombinant WISP-1 on gene expression, petri dishes (35 mm) were coated overnight at 4°C with 75 µg ml<sup>-1</sup> hWISP-1-Fc in PBS. The next day, cells (3 X 10<sup>5</sup> cells/dish) were plated on the coated surface and the expression analysis was performed after 18 hours. In 5 certain cases the coated surface was washed and treated for an additional 2 hours at room temperature before cell were seeded.

FACS Analysis: Cells were harvested by trypsinization, washed with PBS and incubated in PBS 1% BSA/ 20% glucose containing 1 µg ml<sup>-1</sup> FITC conjugated anti-CD44 (Serotec, Raleigh, NC) or an isotype 10 control Ab (DAKO, Denmark) for 30 minutes at room temperature. Cells were washed with the same buffer and analyzed on an EPICS XL-MCL (Coulter, Miami, FL).

Western Blot: NRK/C and NRK/W1H were cultured to confluence in 10 cm<sup>2</sup> plates. The cells were washed with PBS and extracted with 300 15 µl of 1% Triton-X100 in TBS for 5 minutes. Proteins were precipitated by adding 10 volumes of acetone to the extract. Samples were reduced, denatured and loaded on a 4-20% gradient SDS polyacrylamide gel. The gel was transferred to PVDF membranes and probed with a anti-rat CD44 antibody (OX49) or an anti-mouse CD44 20 (KM114) at a 1 µg/ml concentration and detected using a HRP-conjugated anti-mouse IgG secondary antibody and the Amersham ECL detection reagents. Actin detection in the samples was used as loading control.

Scatter Assay: The cells were plated in 24 well plates at low 25 concentration (~1000 cells/well) and allowed to form colonies over 3 days. The cells were stained using Diff-Quik (Dade Behring, Newark, DE) and colonies were observed under the microscope.

Cell Wound Healing Assay: Cell migration was evaluated using a cell wound healing assay previously described with minor 30 modifications (Pienimaki et al., J. Biol. Chem., 276:20428-20435 (2001)). Briefly, cells were cultured until they reached confluence. One line (~1 mm) was drawn in the cell layer using a micropipette equipped with a disposable yellow tip. Cell migration over the cleared area was monitored by time lapse microscopy over 15 hours. 35 The area newly occupied by the cells after this period was measured using NIH image software calibrated with a stage micrometer.

Time Lapse Microscopy: Cells were plated in HGDMEM/10% FBS and their random migration was observed for 15 hours using time lapse 40 microscopy. The distance covered by the cells was measured using NIH image software calibrated with a stage micrometer.

Migration Assay: Haptotaxis was measured using a modified Boyden chamber system as described previously (Bourguignon et al., J. Biol. Chem., 275:1829-1838 (2000)). The underside of 8  $\mu$ m porosity 24-well format PET membrane filters (Falcon) were coated overnight at 5  $^{\circ}$ C with 50  $\mu$ l of protein (50  $\mu$ g  $ml^{-1}$ ) in PBS. The coated inserts were rinsed in serum-free media and 0.5 X  $10^5$  NRK cells in 0.5 ml HGDMEM/10% FBS were added to the upper chamber. The lower chamber was filled with 0.75 ml of the same media and the plates were incubated overnight at 37 $^{\circ}$ C. The next day the upper chamber was wiped with a 10 cotton swab and the cells that migrated to the lower side of the insert were stained with the Diff-Quik Stain Kit (Dade Behring Inc.) and counted under the microscope. Triplicate sets of data were averaged for each condition.

15 In certain cases, the coated inserts were washed and treated for an additional 2 hours at room temperature.

In situ Hybridization: PCR primers were designed to amplify either a 740 bp fragment of murine WISP-1 spanning from nt 440-1180 of NM\_018865 (upper- 5' GGCTGCCATCTGTGACCCA (SEQ ID NO:12) and lower- 5' CATAGGACCT GCCGGGAGAA A (SEQ ID NO:13)) or a 706 bp fragment of 20 murine WISP-1 spanning from nt 204-910 of NM\_018865 (upper- 5' GCCGTGGCAGTCCTGAGGG (SEQ ID NO:14) and lower- 5' CAGCACCGGG CATTGACGTT A (SEQ ID NO:15)) or a 464 bp fragment of murine CD44 spanning from nt 144-608 of M27129 (upper- 5' TGGAGAAAAATGGCCGCTAC A (SEQ ID NO:16)) and lower-5' TGGGGTGCTC TTCTCGATGG (SEQ ID NO:17)) or 25 a 630 bp fragment of murine HAS2 spanning from nt 927-1557 of NM\_008216 (upper- 5' GGACAAATCGGCCACGTACA T (SEQ ID NO:18) and lower- 5' CTTGCTCCAT CGGGTCTGC (SEQ ID NO:19)) . Primers included 30 extensions encoding 27-nucleotide T7 or T3 RNA polymerase initiation sites to allow in vitro transcription of sense or antisense probes, respectively, from the amplified products. (Lu et al., Cell Vision, 1:169-176 (1994)). All tissues were fixed in 4% formalin and paraffin-embedded. Sections 3-5 microns thick were deparaffinized, deproteinated in 4 mg/ml of proteinase K for 30 minutes at 37  $^{\circ}$ C, and further processed for in situ hybridization as previously described. 35 (Holcomb et al., EMBO J., Aug 2000 1;19(15):4046-55).  $^{33}P$ -UTP labeled sense and antisense probes were hybridized to the sections at 55 $^{\circ}$ C overnight. Unhybridized probe was removed by incubation in 20mg/ml RNase A for 30 minutes at 37 $^{\circ}$ C, followed by a high stringency wash at 55  $^{\circ}$ C in 0.1 X SSC for 2 hours and dehydration through graded 40 ethanols. The slides were dipped in NBT2 nuclear track emulsion,

exposed in sealed plastic slide boxes containing dessicant for 4 weeks at 4 °C, developed and counterstained with hematoxylin and eosin.

5

EXAMPLE 1

Stable fibroblast cell lines expressing high (NRK/WISP-1H) or low (NRK/WISP-1L) levels of WISP-1 and a control cell line containing an empty vector (NRK/control) were used to evaluate the effect of WISP-1 on HA production. As demonstrated by the particle exclusion assay, NRK/WISP-1H cells accumulated a large hyaluronan pericellular coat capable of excluding sedimenting erythrocytes (Fig. 1a). NRK/WISP-1L accumulated a smaller pericellular matrix (Fig. 1b) whereas no matrix surrounded NRK/control (Fig. 1c) or NRK/WISP-1H cells treated with hyaluronidase (data not shown).  
15 Accumulation of cell surface associated HA was also evaluated by fluorescent staining using the biotinylated HA binding protein (bHABP). The staining revealed HA accumulation at the NRK/WISP-1H cell surface (Fig. 1d) whereas no staining was detected on NRK control cells (Fig. 1e).

20 The effect of WISP-1 on HA secretion was evaluated by comparing the accumulation of HA in the culture media of NRK/WISP-1H to NRK/control cells over time. After 24 hours, HA concentration in the NRK/WISP-1H media was 3.5 fold greater in NRK/control cell media and gradually increased to 8 fold after 144 h (fig. 1f). Together these 25 results show that WISP-1 expression in NRK cells promotes HA accumulation at the cell surface and in the culture media.

EXAMPLE 2

To identify the enzyme responsible for the WISP-1 triggered HA production, the expression of the known HA synthases (HAS1, HAS2 and HAS3) in NRK/WISP-1H, NRK/WISP-1L and NRK/control cells was analyzed. HAS2 expression was increased up to 10 fold in WISP-1 producing cells whereas HAS1 and HAS3 mRNA levels were identical to the control (Fig. 2a). CD44 and RHAMM mRNA levels were also increased in 35 NRK/WISP-1 cell lines whereas hyaluronidase expression remained unchanged (Fig 2a). Moreover, HAS2, CD44 and RHAMM mRNA levels were proportional to WISP-1 expression. As demonstrated by FACS analysis (Fig. 2b) and Western blot (Fig. 2c), the increase in CD44 mRNA expression in the NRK/WISP-1H cells resulted in a 4- fold increase in 40 CD44 protein level.

EXAMPLE 3

HA was shown to stimulate cell migration by interacting with two cell surface receptors, CD44 and RHAMM (Hall et al., supra; Bourguignon et al., supra). Because WISP-1 increased both HA 5 production and CD44 and RHAMM expression, the motility of WISP-1 expressing cells was evaluated. When plated at a low density, NRK/control cells proliferated and formed well defined colonies (Fig 3a). Proliferating NRK/WISP-1L cells formed less defined groups as some cells departed from the growing colonies (Fig 3b). No NRK/WISP- 10 1H colonies were seen, cells scattered in a random pattern (Fig 3c). Contrary to the control (Fig. 3d), WISP-1 expressing cells also revealed a hyper elongated morphology with extended lamellipodia characteristic of high motility (Fig. 3e). Using time lapse microscopy, a 4 fold increase in the migration distance of NRK/WISP- 15 1H was observed compared to control cells (Fig. 3f). NRK/WISP-1H also showed a 2.5 fold increase in migration area in a cell wound healing assay (Fig. 3g). Together these results demonstrate that WISP-1 expression may promote cell migration.

20

EXAMPLE 4

To determine whether the ectopic addition of WISP-1 could promote HAS2 expression and cell migration, an assay was conducted using purified recombinant WISP-1-IgG (Desnoyers et al., supra). WISP-1 addition to the culture media did not promote HAS2 expression 25 (data not shown). When NRK cells were plated onto a WISP-1-IgG coated surface, HAS2 expression increased compared to cells plated onto uncoated plastic or onto an irrelevant IgG chimeric protein (TNFR-IgG) coated surface (Fig. 4a). Although decorin alone promoted HAS2 expression, this induction was further increased three fold 30 after the coated surface was incubated with WISP-1. The effects of ectopic WISP-1 addition on NRK cell migration were also examined. In a transwell assay, WISP-1-IgG induced the haptotactic migration of NRK cells when coated on the filter's lower surface (Fig. 4b). Coating of an irrelevant IgG chimeric protein (TNFR-IgG) or addition 35 of WISP-1-IgG to the lower chamber did not promote migration. A CD44 or a WISP-1 antibody (Fig. 4b) inhibited the haptotactic migration induced by WISP-1. Moreover, WISP-1 haptotactic activity was not limited to fibroblasts as it also promoted the migration of SW480 cells, a colon adenocarcinoma cell line (Fig. 4c). Together these 40 results demonstrate that ectopic WISP-1 addition increases HAS2 expression and promotes cell migration through a CD44 mediated

mechanism. It is believed, though not fully understood, that WISP-1 presentation may be important for migration as it may need to be tethered to a substrate to elicit this activity.

5

EXAMPLE 5

Because WISP-1 is believed to be a downstream effector of Wnt-1, WISP-1, HAS2 and CD44 expression in a mammary epithelial cell line stably transfected with Wnt-1 (C57MG/Wnt-1) were analyzed. When compared to the control cell line (C57MG), C57MG/Wnt-1 cells showed a 10 2.7, 5.8 and 3 fold increase in WISP-1, HAS2 and CD44 mRNA expression respectively (Fig. 5a). As demonstrated by Western blot, C57/Wnt-1 cells also showed a 6-fold increase in CD44 protein content (Fig. 5b). Moreover, unlike the control cell line, Wnt-1 expressing cells failed to form distinct colonies and scattered when put in culture at 15 low density (Fig. 5c). These results show that HAS2 and CD44 expression and cell motility are elevated in cell lines where WISP-1 expression is triggered by Wnt-1.

20

EXAMPLE 6

Wnt-1 expression in mammary epithelium promotes tumor development in transgenic mice (Li et al., *Oncogene*, 19:1002-1009 (2000); Tsukamoto et al., *Cell*, 55:619-625 (1988)). Because WISP-1 is a putative downstream effector of Wnt-1, the mRNA expression of HAS2 and CD44 in spontaneous mammary tumors from MMTV-Wnt-1 transgenic 25 mice was measured. When compared to mammary duct epithelium, HAS2 mRNA expression was increased between 2.5 and 5.5 fold in all mammary tumors analyzed (n=5; Fig. 5d). Similarly, CD44 mRNA expression was induced between 2.2 and 4.2 fold in all tumors (n=5; Fig 5e). These results show that HAS2 and CD44 are overexpressed in MMTV-Wnt-1 30 transgenic mice mammary tumors expressing WISP-1.

35

EXAMPLE 7

In situ hybridization demonstrated elevated WISP-1 expression in the peritumoral stroma of MMTV-Wnt-1 transgenic mouse mammary tumors (Fig. 6a, b). In contrast, HAS2 (Fig. 6c-d) and CD44 (Fig. 6 e-f) expression was found only in tumoral epithelial cells. The localization of CD44 in the tumor parenchyma was confirmed by immunohistochemistry (Fig. 6g). Link protein staining revealed an accumulation of hyaluronan associated with the tumor stroma (Fig. 40 6h). The highest staining intensity was found in the immediate vicinity of the tumor lobule whereas weaker staining was localized to

the normal mammary duct epithelium. These findings are consistent with the suggestion that WISP-1 may regulate interactions between tumor and stromal cells that involve hyaluronic acid and CD44.

5

EXAMPLE 8

The metastasis and growth potential of transformed lines was evaluated as described previously (Welch et al., Cancer Res., 60:1552-1556 (2000)). Nine-week-old Swiss nu/nu female mice were used. Briefly, cells were harvested by trypsinization and washed 10 twice with PBS. Each mouse was injected into the lateral tail vein with 100  $\mu$ l of a suspension containing  $5 \times 10^4$  or  $2.5 \times 10^5$  cells. At 2, 3 and 4 week postinjection, mice were examined using cine-magnetic resonance imaging (MRI) for apparition of pulmonary lesions and necropsies were performed. The lungs were perfused in Bouin's 15 fixative, excised and H&E stained sections were generated for evaluation of pulmonary tumor colonization.

Longitudinal sections of the left lung, and a single transverse section of the cranial, medial, caudal, and accessory lobes of the right lung were evaluated. Histologically, affected lungs had 20 spindloid neoplastic cells that infiltrated the pulmonary interstitium. However, because the severity of the changes was variable the following grading system was established.

I = Minimal involvement of the pulmonary interstitium - 10-20% of the lung is affected. Nests and clusters of spindloid neoplastic 25 cells multifocally disrupt the pulmonary interstitium. Affected foci are commonly along the pleural surface, extending into the interstitium. No vessels or large airways are affected.

II = Moderate involvement of the pulmonary interstitium - 20 - 50% of the lung is affected. Nests and clusters of spindloid neoplastic 30 cells multifocally disrupt the pulmonary interstitium. Some vessels or large airways are filled with spindloid cells. Spindloid cells frequently form a broad band or large mass subjacent to the pleura and extending into the interstitium. Overlying mesothelium is plump (reactive).

35 III = Severe involvement of the pulmonary interstitium - 50- 100% of the lung is affected. Nests and clusters of spindloid neoplastic cells multifocally disrupt the pulmonary interstitium. Blood vessels and large airways are filled with spindloid cells. Spindloid cells form a broad band or large mass subjacent to the pleura and extending 40 into the interstitium. Spindloid cells are often embedded in a pale basophilic to amphophilic acellular material. Overlying mesothelium

is plump (reactive). Minimal unaffected area for gas exchange remains.

The results of the lung colonization of nude mice inoculated with NRK/WISP-1H, NRK/WISP-1L and NRK/control cells are summarized in Figures 7 and 10. In mice inoculated with NRK/WISP-1H or NRK/WISP-1L, cells formed lung masses in a time- and dose-dependent manner. The most severe lesions were seen in mice injected with  $2.5 \times 10^5$  NRK/WISP-1H and necropsied at 4 weeks post-injection. At 2 and 3 weeks post-injection, the mice had less severe but progressive tumor formation (grade I - II; see grading system above). Mice injected with  $2.5 \times 10^5$  NRK/WISP-1L had only minimal lung tumor formation by 4 weeks post-injection. After 4 weeks post-injection, mice inoculated with  $0.5 \times 10^5$  NRK/WISP-1L cells were either normal or had minimal neoplastic infiltration (grade I), whereas mice injected with  $0.5 \times 10^5$  NRK/WISP-1H cells showed increased neoplastic infiltration (grade I-II). Histological observation revealed that the lungs of animals inoculated with NRK/control cells were normal (Fig. 7a-c). In NRK/WISP-1 injected animals, neoplastic spindloid cells initially formed small clusters within, protruding from, or subjacent to the pleural surface of the lungs (grade I; Fig. 7d-f) often embedded in a pale basophilic to amphophilic acellular material. In more severely affected animals, the neoplastic cells formed a broad confluent band subjacent to the pleura (grade II; Fig. 7g-i). In the most severely affected animals, blood vessels and large airways were filled with spindloid cells. (grade III; Fig. 7j-l). MRI analysis showed increased density (hypersignal) outlining the right and left lungs along the pleura in animals with histologic grade II and grade III invasion. Severity of the lesion on MRI was consistent with the histologic score (Figure 7a, d, g, j). These results indicate that WISP-1 expression promotes cellular metastatic growth potential.

#### EXAMPLE 9

NRK/WISP-1H cells ( $2.5 \times 10^5$  cells) were inoculated in the tail vein of nude mice. Starting on the day of the cell inoculation, the mice were injected intraperitoneally twice a week with 10 mg/kg of a CD44 antibody, an isotype control antibody or with buffer only (PBS). The lungs were fixed and excised after four weeks for gross anatomical analysis.

After 4 weeks, the severity of the lesions found in CD44 antibody treated animals (n=5) varied from normal to grade I whereas all (n=10) animals treated with a control antibody or saline had

grade III lesions (Fig. 8a). The average area of metastatic foci in the lungs of animals treated with CD44 antibody was reduced by 99% ( $P < 0.00003$ ) compared to animals treated with a control antibody (Fig. 8b).

5

EXAMPLE 10

An assay was conducted to examine binding specificity of certain WISP-1 antibodies. Full length mouse WISP-1 (GenBank accession number NM\_018865) and full length human WISP-1 (GenBank 10 accession number AF100779; Fig. 9) were cloned into an expression vector encoding the human IgG<sub>1</sub> Fc region downstream of the WISP-1 sequence. The resulting recombinant fusion protein (WISP-1-Fc) was synthesized in a baculovirus expression system using Sf9 insect cells and purified to homogeneity from serum-free conditioned medium by 15 affinity chromatography on a Protein A-Sepharose 4 Fast Flow (Amersham Pharmacia Biotech). Full length human WISP-1 was also expressed with an amino terminal hexa-histidine tag (WISP-1-His) in an *E. coli* strain. The cell lysate was subjected to chromatography 20 on a Ni<sup>2+</sup>-NTA agarose column (Qiagen). WISP-1-His was eluted with a 0 to 500 mM imidazole gradient. Fractions containing the eluted WISP-1-His were then pooled and dialyzed. Human WISP-1 from a 25 mammalian expression system was obtained by lysing NRK cells stably transfected with human WISP-1 (Arnold Levine; Princeton University, Princeton, NJ) with SDS-PAGE sample buffer. A control cell lysate was generated with NRK cells stably transfected with an empty vector.

WISP-1 (50 ng) from various expression systems was electrophoresed on a SDS polyacrylamide gel and electro-transferred onto polyvinylidifluoride (PVDF) membranes and probed with different WISP-1 monoclonal antibodies.

30 WISP-1 antibodies 3D11.D7 (also referred to herein as "3D11"), 11C2.C10 (also referred to herein as "11C2"), 9C11.C7 (also referred to herein as "9C11") and 5D4.F6 (also referred to herein as "5D4") bound specifically to WISP-1 generated from baculovirus, bacterial and mammalian expression systems (Fig. 11a). These antibodies did 35 not bind to the murine WISP-1 from baculovirus and did not recognize any protein from the control lysate. The WISP-1 antibodies 6F8, 3A7, 10H12, 3A11, 6E3, 3H10, 5G1, and 10B1 recognized both human and murine WISP-1 only when generated with the baculovirus expression system (Fig. 11b). These antibodies did not recognize human WISP-1 40 when produced in a bacterial or mammalian expression system. The

antibody from clone 9C10 did not bind to any protein after Western blot.

These results suggest that WISP-1 antibodies 3D11, 11C2, 9C11 and 5D4 specifically recognize human WISP-1 and can be used for WISP-5 1 detection by Western blot.

EXAMPLE 11

An assay was conducted to identify the epitopes recognized by the WISP-1 antibodies 11C2, 9C11, 5D4 and 3D11.

10 Full length human WISP-1 (GenBank accession number AF100779) was cloned into a pIRESpuro2 expression vector (Clontech Laboratories, Palo Alto, CA) encoding 6 histidines downstream of the WISP-1 sequence. Deletion mutants were also generated by removing one, two or three domains of human WISP-1. The resulting constructs 15 were also cloned into the pIRESpuro2 expression vector. The nomenclature used to identify the different WISP-1 constructs refer to the domains they contain. (see Fig. 12B) Domain 1 is the insulin-like growth factor binding protein domain (IFGBP), domain 2 is the von Willebrand factor C (VWFc) domain, domain 3 is the 20 thrombospondin (TSP) domain, and the domain 4 is the C-terminal (CT) domain. The variable region resides between domain 2 and 3. These regions and domains of WISP-1 are illustrated in Fig. 12A.

The sequences encoding these domains of WISP-1 are as follows:  
Sequences of WISP-1 Constructs

25 Domain 1:

GAATTCAACATGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCAGCACCGTC  
CTGGCCACGGCCCTCTCTCCAGCCCCCTACGACCATGGACTTACTCCAGCTCCACTGGAGGACACCTCC  
TCACGCCCCAATTCTGCAAGTGGCCATGTGAGTGCCCCGCCATCCCCACCCCGCTGCCGCTGGGGTC  
AGCCTCATCACAGATGGCTGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAACTGACGGAG  
30 GCTGCCATCTGTGACCCCCACCGGGGCCTACTGTGACTACAGCGGGACCGCCCGAGGTACGCAATA  
GGAGTGTGTGCACAGGCGGCCGCACACCACCATCACCATCACCATACTAAGTGAGGCCGCATAGATAA  
CTGATCCAGTGTGCTGGAATTAATTC (SEQ ID NO:3)

Domain 2:

35 GAATTCAACATGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCAGCACCGTC  
CTGGCCACTGCAGTGGTCGGTGTGGCTGCGCTGGATGGGTGCGCTACAACAACGGCCAGTCCTTC  
CAGCTTAACGTCAAGTACAACGTGCACGTGCATCGACGGCGGGTGGCTGCACACCACTGTGCCCTCGA  
GTGCGCCCCCGCGTCTGGTGCCCCCACCGCGCGTGTGAGCATACTGGCCACTGCTGTGAGCAG  
TGGGTATGTGCGGCCGCACACCACCATCACCATACTAAGTGAGGCCGCATAGATAAC (SEQ  
40 ID NO:4)

## Domain 3:

GAATTCAACCAGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTACAGCAGCAGCCAGCACCGTC  
 CTGGCCACTGCAGCATGGCACAGGAACATGCATAGCCTACACAAGCCCCTGGAGCCCTGCTCCACCAGC  
 TCGGCCCTGGGGCTCCACTCGGATCTCAATGTTAACGCCAGTGCTGGCTGAGCAAGAGAGCCGC  
 5 CTCTGCAACTTGGGCCATGCGATGTGGACATCCATACACTCATTAAGGCCGCACACCACCATCAC  
 CATCACCATCACTAAGTGAGGCCGCATAGATAACTGATCCAGTGT (SEQ ID NO:5)

## Domain 4:

GAATTCAACCAGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTACAGCAGCAGCCAGCACCGTC  
 10 CTGGCCACTGCAGGGAAAGAAGTGTCTGGCTGTGTACAGCCAGAGGCATCCATGAACCTCACACTTGCG  
 GGCTGCATCAGCACACGCTCCTATCAACCCAAGTACTGTGGAGTTGCATGGACAATAGGTGCTGCATC  
 CCCTACAAGTCTAAGACTATCGACGTGTCCTCCAGTGTCTGATGGGCTTGGCTCTCCGCCAGGTC  
 CTATGGATTAATGCCTGCTCTGTAACCTGAGCTGTAGGAATCCAATGACATCTTGCTGACTTGGAA  
 15 TCCTACCTGACTTCTCAGAAATTGCCAACGCCGCACACCACCATCACCACCATCACTAAGTG  
 AGGCCGCATAGATAACTGATCCAGTGTG (SEQ ID NO:6)

## Domain 1,2:

GAATTCAACCAGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTACAGCAGCAGCCAGCACCGTC  
 CTGGCCACGGCCCTCTCCAGCCCCTACGACCATGGACTTACTCCAGCTCCACTGGAGGACACCTCC  
 20 TCACGCCCCAATTCTGCAAGTGGCCATGTGAGTGCCCACATCCCCACCCCGCTGCCGCTGGGGTC  
 AGCCTCATCACAGATGGCTGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAATGCA CGGAG  
 GCTGCCATCTGTGACCCCCACCGGGGCTCTACTGTGACTACAGCAGGGACCGCCCGAGGTACGCAATA  
 GGAGTGTGTGCACAGGTGGTGGTGGCTGCGCTGGATGGGGTGCCTACAAACACGGCCAGTCC  
 TTCCAGCCTAACTGCAAGTACAACGTGACGTGCATCGACGGCGGGTGGCTGCACACCAACTTGCC  
 25 CGAGTGCACGGCCCGCGTCTGGTGGCCCCACCCCGCGCGTGAGCATACCTGCCACTGCTGTGAG  
 CAGTGGGTATGTGCGGCCGCACACCACCATCACCACCATCACTAAGTGAGGCCGCATAGATAAC  
 (SEQ ID NO:7)

## Domain 1,2,3:

GAATTCAACCAGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTACAGCAGCAGCCAGCACCGTC  
 CTGGCCACGGCCCTCTCCAGCCCCTACGACCATGGACTTACTCCAGCTCCACTGGAGGACACCTCC  
 TCACGCCCCAATTCTGCAAGTGGCCATGTGAGTGCCCACATCCCCACCCCGCTGCCGCTGGGGTC  
 AGCCTCATCACAGATGGCTGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTTGGGGACAATGCA CGGAG  
 GCTGCCATCTGTGACCCCCACCGGGGCTCTACTGTGACTACAGCAGGGACCGCCCGAGGTACGCAATA  
 35 GGAGTGTGTGCACAGGTGGTGGTGGCTGCGCTGGATGGGGTGCCTACAAACACGGCCAGTCC  
 TTCCAGCCTAACTGCAAGTACAACGTGACGTGCATCGACGGCGGGTGGCTGCACACCAACTTGCC  
 CGAGTGCACGGCCCGCGTCTGGTGGCCCCACCCCGCGCGTGAGCATACCTGCCACTGCTGTGAG  
 CAGTGGGTATGTGAGGACGACGCCAGCAAGAGGCCACCGCAAGACCGCACCCGTGACACAGGAGCCTCGAT  
 GCTGTGGGTGAGGTGGAGGCATGGCACAGGAACATGCATAGCCTACACAAGCCCCTGGAGCCCTGCTCC  
 40 ACCAGCTGCGGCCCTGGGGCTCCACTCGGATCTCAATGTTAACGCCAGTGCTGGCTGAGCAAGAG  
 AGCCGCCTGCAACTTGGGCCATGCGATGTGGACATCCATACACTCATTAAGGCGGCCACACCAC

CATCACCATCACCATCACTAAGTGAGGCCGCATAGATAACTGATCCAGTGTGCTGGA (SEQ ID NO: 8)

Domain 1,2,4:

5 GAATTCACCATGAGGTGGTTCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTC  
 CTGGCCACGGCCCTCTCTCCAGCCCCTACGACCATGGACTTACTCCAGCTCCACTGGAGGACACCTCC  
 TCACGCCCCCAATTCTGCAAGTGGCCATGTGAGTGCCCCCATCCCCACCCGCTGCCGCTGGGGGTC  
 AGCCTCATCACAGATGGCTGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTGGGGACAACGTGACGGAG  
 GCTGCCATCTGTGACCCCCACGGGGCCTCTACTGTGACTACAGCGGGGACCGCCCGAGGTACGCAATA  
 10 GGAGTGTGTGCACAGGTGGTCGGTGTGGCTGCGTCCTGGATGGGTGCGCTACAACAACGGCCAGTCC  
 TTCCAGCCTAACTGCAAGTACAACACTGCACGTCATCGACGGCGGGTGGCTGCACACCACTGTGCCTC  
 CGAGTGCAGCCCCCGCGTCTCTGGTCCCCCACCAGCGGGCGGTGAGCATACTGCCACTGCTGTGAG  
 CAGTGGGTATGTCTGCAAGGAGGGAAAGAAGTGTCTGGCTGTGACTACAGCCAGAGGCATCCATGAACTTC  
 ACACATTGCGGGCTGCATCAGCACACGCTCTATCAACCCAAGTACTGTGGAGTTGCATGGACAATAGG  
 15 TGCTGCATCCCCATAAGTCTAAAGACTATCGACGTGCTCCAGTGTCCCTGATGGCTTGGCTTC  
 CGCCAGGTCTATGGATTAATGCCTGCTCTGTAACCTGAGCTGTAGGAATCCAATGACATTTGCT  
 GACTTGGAAATCCTACCCCTGACTTCTCAGAAATTGCCAACCGGGCCGACACCACCATCACCACCAT  
 CACTAAGTGAGGCCGCATAGATAACTGATCCAGTGTGCTGGAATTAATCGCTGTGCGAGGGCCAGC  
 TGTTGGGGTGAGTACTCCCTCTAAAAGCGGGCATGACTCTGCGCTA (SEQ ID NO: 9)

20

Domain 1,3,4:

GAATTCACCATGAGGTGGTTCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGCACCGTC  
 CTGGCCACGGCCCTCTCTCCAGCCCCTACGACCATGGACTTACTCCAGCTCCACTGGAGGACACCTCC  
 TCACGCCCCCAATTCTGCAAGTGGCCATGTGAGTGCCCCCATCCCCACCCGCTGCCGCTGGGGGTC  
 25 AGCCTCATCACAGATGGCTGTGAGTGCTGTAAGATGTGCGCTCAGCAGCTGGGGACAACGTGACGGAG  
 GCTGCCATCTGTGACCCCCACGGGGCCTCTACTGTGACTACAGCGGGGACCGCCCGAGGTACGCAATA  
 GGAGTGTGCGCATGCTGGGTGAGGTGGAGGCATGGCACAGGAACGTGCATAGCCTACACAAGCCCC  
 TGGAGCCCTGCTCCACCAGCTGCGGCTGGGGCTCCACTCGGATCTCAATGTTAACGCCAGTGC  
 TGGCCTGAGCAAGAGAGCCGCTCTGCAACTTGGCCATGCGATGTGGACATCCATACACTCATTAAG  
 30 GCAGGGAAGAAGTGTCTGGCTGTGACCGCCAGAGGCATCCATGAACTTCACACTTGCGGGCTGCATC  
 AGCACACGCTCTATCAACCCAAGTACTGTGGAGTTGCATGGACAATAGGTGCTGCATCCCCTACAAG  
 TCTAAGACTATCGACGTGCTCCAGTGTCTGATGGCTTGGCTCTCCAGGTCTATGGATT  
 AATGCCCTGCTCTGTAACCTGAGCTGTAGGAATCCAATGACATCTTGCTGACTTGGAAATCCTACCC  
 GACTTCTCAGAAATTGCCAACCGGGCCGACACCACCATCACCACCATCACTAAGTGAGGCCAGC  
 35 AGATAAC (SEQ ID NO: 10)

Domain 2,3,4:

GAATTCACCATGAGGTGGTTCTGCCCTGGACGCTGGCAGCAGTGACAGCAGCAGCCGCCAGC  
 ACCGTCCTGGCCACTGCAGTGGTCGGTGTGGCTGCGCTCTGGATGGGTGCGCTACAACAACGGCCAG  
 40 TCCTTCCAGCCTAACTGCAAGTACAACACTGCACGTGCATCGACGGCGCGGTGGCTGCACACCACTGTGC  
 CTCCGAGTGCAGCCCCCGCGTCTCTGGTCCCCCACCGCGCGTGAACATACCTGCCACTGCTGT

GAGCAGTGGGTATGTGAGGACGACGCCAAGAGGCCACGCAAGACCGCACCCGTGACACAGGAGCCTTC  
GATGCTGTGGGTGAGGTGGAGGCATGGCACAGGAACACTGCATAGCCTACACAAGCCCTGGAGCCCTTGC  
TCCACCAGCTGCGGCCTGGGGTCTCCACTCGGATCTCAATGTTAACGCCAGTGCTGGCTGAGCAA  
GAGAGCCGCCTCTGCAACTTGCAGGCCATGCGATGTGGACATCCATACACTCATTAAGGCAGGGAAAGAAG  
5 TGTCTGGCTGTGTACAGGCCAGAGGCATCCATGAACCTCACACTTGCAGGCCATCAGCACACGCTCC  
TATCAACCCAAGTACTGTGGAGTTGCATGGACAATAGGTGCTGCATCCCTACAAGTCTAAGACTATC  
GACGTGCTTCCAGTGTCTGATGGGCTTGGCTCTCCGCCAGGTCTATGGATTAAATGCCTGCTTC  
TGTAAACCTGAGCTGTAGGAATCCAATGACATCTTGCTGACTTGGAAATCCTACCCTGACTTCTCAGAA  
ATTGCCAACGCGGCCGCACACCACCATCACCATCACCATCAACTAAGTGAGGCCGCATAGATAACTGATC  
10 CAGTGTGCTGGAATTAATTGCTGTGCGA (SEQ ID NO:11)

Cells (HEK 293T) were transfected with the different constructs, and the culture media was collected after 48 hours. One milliliter of culture media was incubated with 20  $\mu$ l of cobalt-agarose for 1 hour, centrifuged and washed. The adsorbed proteins were eluted by heating the pellet at 100 °C for 5 minutes in 20  $\mu$ l of SDS-PAGE sample buffer. The samples were electrophoresed, electro-transferred onto PVDF and probed with the different WISP-1 antibodies.

20 Antibodies 11C2, 9C11 and 5D4 recognized only WISP-1 constructs containing the 19 first amino acids of the variable region located between domain 2 and 3 (Fig. 12C; 12E; 12G). The WISP-1 antibody 3D11 recognized only WISP-1 constructs containing the domain 1 (amino acids 24 to 117; Fig. 12D; 12F).

25 These results indicate that the antibodies 11C2, 9C11 and 5D4 recognize specifically the variable region of WISP-1 whereas the antibody 3D11 recognizes specifically the domain 1 of WISP-1.

#### EXAMPLE 12

30 An assay was conducted to identify the epitope recognized by the WISP-1 antibody 9C10.F5 (also referred to herein as "9C10").

Culture media from HEK 293T cells transfected with the various WISP-1 deletion constructs (as described above in Example 11) was incubated with 1  $\mu$ g of WISP-1 antibody 9C10 and 20  $\mu$ l of protein 35 A-agarose for 1 hour at room temperature. The immunocomplex was precipitated by centrifugation and eluted by heating the pellet at 100 °C for 5 minutes in 20  $\mu$ l of SDS-PAGE sample buffer. The samples were electrophoresed, electro-transferred onto PVDF and probed with WISP-1 antibody 11C2.

40 The antibody 9C10 immunoprecipitated only constructs containing the domain 1 of WISP-1 (Fig. 13). These results demonstrate that the

antibody 9C10 specifically recognizes the domain 1 of WISP-1 and can be used for immunoprecipitation.

EXAMPLE 13

5 WISP-1 antibody 9C10 (100  $\mu$ l of 2  $\mu$ g/ml in carbonate buffer, pH 9.6) was coated to Maxisorb plates overnight at 4 °C. The plates were blocked with 200  $\mu$ l of PBS/3% BSA for 1 hour. A standard curve was made of serial dilutions of WISP-1-Fc (100  $\mu$ l in PBS/3% BSA) and incubated for 1 hour. After the incubation, the plates were washed 10 with 100  $\mu$ l PBS/0.05% Tween and WISP-1 antibodies (100  $\mu$ l of 2  $\mu$ g/ml) in PBS/3% BSA (biotinylated 11C2 or 55B) were incubated for 1 hour. For biotinylated 11C2, the plates were further incubated with 2  $\mu$ g/ml HRP-conjugated streptavidin. For 55B, the plates are washed and incubated with HRP-conjugated donkey anti-rabbit IgG for 1 hour. At 15 the end of the incubation, the wells were washed 6 times with 200  $\mu$ l of PBS containing 0.05% Tween-20, and the signal was visualized using 100  $\mu$ l of the horseradish peroxidase chromogenic substrate TMB (Kirkegaard & Perry Laboratories). The reaction was stopped with 100  $\mu$ l of 1 M phosphoric acid, and the OD at 450 nm was measured. 20 Non-specific binding was determined in parallel incubations by omitting microtiter well coating. No signal was generated when WISP-1-Fc or a WISP-1 antibody was omitted.

Using the antibody 9C10 for capture and the antibodies 11C2 and 55B for detection, an ELISA was conducted capable of detecting 25 concentration of WISP-1 as low as 0.4  $\mu$ g/ml (Fig. 14). This ELISA may be useful for detecting WISP-1 protein in biological fluids such as serum.

EXAMPLE 14

30 Maxisorb plates were coated overnight at 4 °C with 50  $\mu$ l/well of 10  $\mu$ g/ml heparin (Sigma). The non specific binding sites were blocked with 200  $\mu$ l of PBS/3% BSA for 1 hour. The plates were then incubated for 1 hour with 50  $\mu$ l of 6  $\mu$ g/ml hWISP-1-Fc in PBS/3% BSA in the presence of serial dilutions of WISP-1 antibodies. The plates 35 were washed with PBS/0.05% Tween and further incubated 1 hour with 50  $\mu$ l of 2  $\mu$ g/ml HRP conjugated anti-human IgG-Fc in PBS/3% BSA. The plates were washed, and 100  $\mu$ l of HRP substrate (TMB) was added. The color development was stopped with 100  $\mu$ l of 1 M phosphoric acid and the OD at 450 nm was measured.

The WISP-1 antibodies 11C2, 5D4 and 9C11 inhibited WISP-1 binding to heparin with an IC<sub>50</sub> of 1.9, 2.5 and 3.7 µg/ml, respectively (Fig. 15). The antibody 3D11 moderately reduced WISP-1 binding to heparin with a maximal inhibition of 62% at the highest 5 concentration tested (40 µg/ml). The antibody 9C10 did not attenuate WISP-1 heparin binding, showing an inhibition curve similar to the irrelevant antibody control.

These results demonstrate that antibodies recognizing the variable region can inhibit WISP-1 binding to heparin. Because the 10 two WISP-1 antibodies recognizing domain 1 have little or no effect on WISP-1 binding to heparin, it is presently believed that the domain 1 is less likely to participate in this interaction.

#### EXAMPLE 15

15 Haptotaxis was measured using a modified Boyden chamber system. The underside of 8 µm porosity 24-well format PET membrane filters (Becton Dickinson, Franklin Lakes, New Jersey) were coated overnight at 4 °C with 50 µl of protein (50 µg/ml) in PBS. Normal rat kidney cells (NRK; 5 x 10<sup>4</sup>/0.5 ml HGDMEM/10% FBS) were added to the upper 20 chamber, the lower chamber was filled with the same media and the plates were incubated at 37 °C. The next day, the upper chamber was wiped with a cotton swab, and the cells that migrated to the lower side of the insert were stained and counted under a microscope. Triplicate sets of data were averaged for each condition. In certain 25 cases, the coated inserts were washed and treated for an additional 2 hours at room temperature.

In a transwell assay, WISP-1-Fc coated on the filters' lower surface induced the haptotactic migration of NRK cells (Fig. 16A). Coating of an irrelevant IgG chimeric protein (TNFR-Fc) or addition 30 of WISP-1-Fc to the lower chamber did not promote migration. In the presence of coated WISP-1-Fc, five different WISP-1 antibodies (9C10, 11C2, 3D11, 9C11, 5D4) markedly inhibited cell migration. In the absence of coated WISP-1-Fc, these antibodies did not show any effect on cell migration. These results demonstrate that WISP-1 antibodies 35 can modulate cell migration in the presence of WISP-1. By blocking cell migration, WISP-1 antibodies may play an important therapeutic role in preventing cancer progression.

A summary of the characteristics and properties of the 3D11, 9C10, 11C2, 5D4, and 9C11 antibodies discussed in the Examples above 40 is provided in Fig. 16B.

EXAMPLE 16

Nine-week-old Swiss nu/nu female mice were injected into the lateral tail vein with 100  $\mu$ l of a suspension containing  $2.5 \times 10^5$  cells. Starting on the day of the inoculation, the mice were treated semiweekly by intraperitoneal injection (10 mg/kg) of WISP-1 antibodies (9C11, 11C2, 5D4, 9C10, 3D11) or isotype control antibody.

After 3 weeks, lungs were perfused with 4% neutral buffered formalin, excised and H&E stained sections were generated.

Longitudinal sections of the left lung, and a transverse section of the cranial, medial, caudal, and accessory lobes of the right lung were evaluated. For each slide the number of metastatic foci was counted and the average area of metastatic foci was measured using the SPOT RT software (Diagnostic Instruments Incorporated, Burlingame, California). The area ( $\mu\text{m}^2$ ) was determined for at least five individual metastatic foci in four sections of lung.

After 3 weeks, the severity of the lesions found in WISP-1 antibody treated animals was greatly attenuated compared to control (Fig. 17 a, b). The number of nodules and the average area of the metastatic foci found in mice treated with WISP-1 antibodies (n=5) were reduced compared to animals treated with a control antibody (Fig. 17 c,d). Furthermore, upon WISP-1 antibody treatment, the total pulmonary area covered by the lesions was reduced by 82-97% compared to animal treated with an isotype control antibody (Fig. 17e). These results demonstrate the *in vivo* efficacy of WISP-1 antibodies at reducing the tumor burden related to metastasis. Although the mechanism of action of the WISP-1 antibodies is not fully understood, it is believed their efficacy may be mediated by a growth reducing ability and/or a capacity at inhibiting the motility, invasion and seeding of cancer cells at a tissue site.

EXAMPLE 17

Full length mouse WISP-1 (GenBank accession number NM\_018865) was cloned into the pRK mammalian expression vector. The resulting construct (pRK-WISP-1; 18ug) was co-transfected with 2ug pSVi-puromycin plasmid in a 4T1 mouse mammary adenocarcinoma cell line (obtained from Dr. Fred Miller, Barbara Ann Karmanos Cancer Institute, Detroit, MI) using Fugene6 (Roche) according to manufacturer's instructions. After 2 days, cells were selected in 2  $\mu$ g/ml puromycin. After 2 weeks, clones were isolated and WISP-1

expression was evaluated by immunofluorescence. The same procedure was used to generate control cell lines using an empty vector. The resulting 4T1/control and 4T1/WISP-1 cell lines were maintained in Iscove's Media containing 10% FBS and 3 µg/ml puromycin.

5 WISP-1 expression in 4T1/control, 4T1/WISP-1L, 4T1/WISP-1H, NRK/control, NRK/WISP-1L and NRK/WISP-1H cells lines was measured by semi-quantitative RT-PCR (Taqman) using primers and probes that do not distinguish between human and mouse WISP-1 gene.

10 WISP-1 was not expressed in 4T1/control cell lines. WISP-1 expression in 4T1/WISP-1H cell line was 2 fold higher than in 4T1/WISP-1L but 700 fold lower than in NRK/WISP-1H (Figure 18).

EXAMPLE 18

15 To evaluate WISP-1 effect on 4T1 cell scattering, 100,000 4T1/control, 4T1/WISP-1L or 4T1/WISP-1H cells were seeded in 6 well plate in Iscove's Media containing 10% fetal bovine serum. When plated at a low density, 4T1/control cells proliferated and formed well defined colonies (Fig. 19). Proliferating 4T1/WISP-1L cells formed less defined groups as some cells departed from the growing 20 colonies (Fig. 19). No 4T1/WISP-1H colonies were seen, and cells scattered in a random pattern (Fig. 19). Together these results suggest that WISP-1 expression promotes cell migration.

EXAMPLE 19

25 WISP-1 effect on 4T1 cell invasion was also evaluated using a Matrigel coated modified Boyden chamber system of 8 µm porosity 24-well format PET membrane filters (Falcon). 4T1/control, 4T1/WISP-1L or 4T1/WISP-1H cells (100,000 cells) were added to the upper chamber in 0.5 ml Iscove's Media. The lower chamber was filled with 0.75 ml 30 of the same media containing 5% fetal bovine serum and the plates were incubated overnight at 37°C. The next day the upper chamber was wiped with a cotton swab and the cells that migrated to the lower side of the insert were stained with the Diff-Quik Stain Kit (Dade Behring Inc.) and counted under the microscope. Triplicate sets of 35 data were averaged for each condition. 4T1/WISP-1H and 4T1/WISP-1L cells demonstrated a 12 and 6 fold increase invasion compared to 4T1/control, respectively (Fig. 20). These results suggest that WISP-1 promotes tumorigenic mammary epithelial cell invasion and may play a role in metastasis.

EXAMPLE 20

WISP-1 effects on mammary epithelial cells tumorigenesis were evaluated by injecting  $1.5 \times 10^5$  cells (4T1/control 1, 4T1/control 2, 4T1/WISP-1L or 4T1/WISP-1H) into the fourth mammary fat pad of 6-8 week old female BALB/c mice (6 mice/group). Tumor volumes were measured three times per week. Thirty one days after injection, the mice were sacrificed, the tumors were excised and weighted.

Inoculation of 4T1/WISP-1L cells (Fig. 21a; empty squares) and 4T1/WISP-1H cells (Fig. 21a; filled squares) generated faster growing tumors compared to 4T1/control 1 and 4T1/control 2 cells (Fig. 21a; empty and filled circles). After 31 days, the tumors formed by 4T1/WISP-1L and 4T1/WISP-1H inoculated were 4 fold larger (Fig. 21a) and 3 fold heavier (Fig. 21b) than the tumors formed by 4T1/control cells. These results suggest that WISP-1 may increase proliferation of mammary epithelial tumor cells.

EXAMPLE 21

The expression of HAS2 and CD44 in tumors formed by the inoculated 4T1/control, 4T1/WISP-1L and 4T1/WISP-1H cells was measured. When compared to 4T1/control cells tumor, CD44 expression was increased between 5 and 23 fold in 4T1/WISP-1 cells tumors (Fig. 22). On the other hand, HAS2 expression remained identical in all tumors analyzed. These results suggest that WISP-1 increases expression of CD44 in mice inoculated with 4T1 cells. The overexpression of CD44 in these tumors may contribute to the promotion of metastasis.

EXAMPLE 22

WISP-1 effects on mammary epithelial cells metastasis were evaluated by inoculating 4T1 cells in mice mammary fat pads (see Example 20) and examining the extent of the metastatic propagation by micro computer tomography and histology. After 31 days, the mice inoculated with 4T1/WISP-1L or 4T1/WISP-1H cells had extensive lung metastasis (Fig. 23b and 23d) compared to the 4T1/control injected mice (Fig. 23a and 23c). No significant differences were seen between the metastatic potential of 4T1/WISP-1L and 4T1/WISP-1H cells. The mice inoculated with 4T1/control cells had an average of 2.11 pulmonary foci with an average mass of 0.68 grams whereas mice inoculated with 4T1/WISP-1 cells had an average of 20.25 pulmonary foci with an average mass of 12.46 grams. Also, the average

histology score for the lung of 4T1/WISP-1 cells injected mice was 0.92 compared to 0.11 for the NRK/control cells injected mice.

Using immunohistochemistry, it was also observed that the 4T1/WISP-1 pulmonary metastatic foci expressed high levels of CD44 (Fig. 23e). In these tumors, CD44 was localized at the plasma membrane of the 4T1/WISP-1 cells (Fig. 23f). Together these results demonstrate that WISP-1 promotes the metastatic potential of 4T1 cells, increasing the number of pulmonary metastatic foci (10 fold) and size (18 fold). Also, CD44 increased expression was maintained 10 after the 4T1/WISP-1 had metastasized to the lungs.

Clinical observations of mice inoculated with 4T1/control 1, or 4T1/WISP-1 cells demonstrated that WISP-1 expression promoted metastasis at additional secondary sites. Two mice inoculated with 4T1/WISP-1H had a discolored white mass on the kidney whereas no 15 other mice had evidence of kidney tumors.

#### EXAMPLE 23

The human WISP-1 full length and deletion mutant constructs 20 described above were transfected in normal rat kidney cells (NRK-49F). After 48 hours, the culture media was supplemented with 2 µg/ml puromycin. After 2 weeks, the pools of expressing cells were analyzed and the resulting cell lines were maintained in High glucose Dulbecco's Modified Eagle's Medium (HGDMEM) supplemented with 10% 25 fetal calf serum (FBS) and 2 µg/ml puromycin.

WISP-1 expression in NRK/control, NRK/WISP-1\_1234, NRK/WISP-1\_134, NRK/WISP-1\_234, NRK/WISP-1\_123 and NRK/WISP-1\_124 cells lines was measured by semi-quantitative RT-PCR (Taqman). Results are expressed as fold relative to WISP-1 expression in NRK/WISP-1\_1234 30 (Figure 24A). WISP-1 expression in NRK/WISP-1\_134 and NRK/WISP-1\_234 is 0.035 and 0.09 fold respectively compared to NRK/WISP-1\_1234. NRK/WISP-1\_123 expresses 16.1 fold and NRK/WISP-1\_124 7.6 fold compared to NRK/WISP-1\_1234.

The contribution of particular domains to WISP-1 promoted 35 tumorigenesis was evaluated by injecting  $5 \times 10^6$  cells (NRK/control, NRK/WISP-1\_1234, NRK/WISP-1\_123, NRK/WISP-1\_124, NRK/WISP-1\_134, NRK/WISP-1\_234 cells) subcutaneously in 6-8 week old female nude mice (10 mice/group). Tumor volumes were measured weekly. After 35 days the mice were sacrificed, the tumors were excised and analyzed 40 histologically.

Inoculation of NRK/WISP-1\_1234 (Figure 24B; filled triangles), NRK/WISP-1\_134 (Figure 24B; filled circles) and NRK/WISP-1\_234 cells (Figure 24B; filled squares) generated tumors. However the inoculation of NRK/WISP-1\_123 (Figure 24B; empty squares), NRK/WISP-1\_124 (Figure 24B; empty circles) and NRK/control (Figure 24B; empty triangles) did not generate tumors. After 35 days, the tumors formed by NRK/WISP-1\_234 cells were comparable in size to the tumors formed by NRK/WISP-1\_1234 cells. However the tumors formed by NRK/WISP-1\_134 cells were 2.5 fold larger than the tumors formed by NRK/WISP-1\_1234 cells. These results suggest that the domains 1 and 2 are not required for WISP-1 promoted NRK tumorigenesis but that the domains 3 and 4 are required for WISP-1 promoted NRK tumorigenesis. It is believed that WISP-1 antagonists of the invention may preferably bind to and/or block domains 3 and/or 4 of WISP-1.

15 In addition, results suggest the deletion of the domain 2 of WISP-1 may increase its tumorigenic potential.

20 Histological analysis of the excised tumors (Figure 25) revealed that the neoplastic cells from NRK/WISP-1\_234 tumors were phenotypically similar to the neoplastic cells from NRK/WISP-1\_1234 tumors. In these tumors, cells appeared fibroblastic, differentiated and spindloid. However the neoplastic cells from NRK/WISP-1\_134 tumors were phenotypically different and appeared less differentiated, frequently multinucleated and had a brisk mitotic rate as demonstrated by the presence of multiple mitotic figures (arrows). In addition, neoplastic cells invaded several blood vessels adjacent to NRK/WISP-1\_134 tumors (Figure 26). These results indicate that WISP-1\_134 expression may confer tumorigenicity to NRK cells by promoting their *in vivo* growth and invasive potential. Moreover the expression of this construct induces a phenotypic 25 transformation characteristic of aggressive neoplastic cells.

#### EXAMPLE 24

30 The contribution of the particular WISP-1 domains to the promoted 4T1 cell invasion was evaluated using a Matrigel coated modified Boyden chamber system of 8  $\mu$ m porosity 24-well format PET membrane filters (BD Biosciences). HEK 293 cells ( $20 \times 10^6$ ) stably expressing the previously mentioned WISP-1 constructs (WISP-1\_1234, WISP-1\_134, WISP-1\_123, WISP-1\_12, WISP-1\_23, WISP-1\_3, WISP-1\_1 and WISP-1\_4) were seeded in the lower chamber with in 0.75 ml of 50:50 Media (1:1 mixture of DME and Ham's F-12 medium) containing 5% fetal

bovine serum and the plates were kept overnight at 37°C. The next day, 4T1 cells (100,000 cells) were added to the upper chamber in 0.5 ml of serum free 50:50 media and the plates were incubated overnight at 37°C. The following day the upper chamber was wiped with a cotton swab and the cells that migrated to the lower side of the insert were stained with the Diff-Quik Stain Kit (Dade Behring Inc.) and counted under the microscope. Triplicate sets of data were averaged for each condition and results are expressed as fold invasion compared to a control in which HEK 293 cells transfected with an empty vector were plated in the lower chamber. 4T1 cells demonstrated a 4 fold increase invasion when HEK 293 cells expressing constructs WISP-1\_1234, WISP-1\_134 or WISP-123 were seeded in the lower chamber (Fig. 27). HEK 293 cells expressing the construct WISP-1\_12 increased 4T1 cells invasion by only 2.5 fold. The increase in 4T1 cell invasion was marginal when HEK 293 cells expressing WISP-1\_23, WISP-1\_3, WISP-1\_1 or WISP-1\_4 were plated in the lower chamber. These results suggest that the domain 1 is required for WISP-1 promoted 4T1 cell invasion. In addition, the data shows that WISP-1's domain 1 alone is likely not sufficient to promote 4T1 cell invasion.

20

EXAMPLE 25

An assay was conducted to evaluate the antagonistic activity of WISP-1's domain 1 on WISP-1 promoted 4T1 cell invasion. The domain 1 of WISP-1 was cloned into an expression vector immediately upstream of the human IgG<sub>1</sub> Fc region sequence. The resulting recombinant fusion protein (WISP-1-domain 1-Fc) was synthesized in a mammalian expression system using Chinese hamster ovary (CHO) cells and purified to homogeneity from conditioned medium by affinity chromatography on a Protein A-Sepharose 4 Fast Flow (Amersham 25 Pharmacia Biotech). WISP-1's domain 1 was also expressed with an amino terminal hexa-histidine tag (WISP-1-domain-1-His) in a CHO expression system and purified to homogeneity from culture supernatant by affinity chromatography on a Ni<sup>2+</sup>-NTA agarose column (Qiagen). Fractions containing the eluted WISP-1-domain-1-His were 30 then pooled and dialyzed.

The antagonistic effect of purified soluble WISP-1-domain-1 on WISP-1 promoted 4T1 cell invasion was evaluated using a Matrigel coated modified Boyden chamber system of 8 µm porosity 24-well format PET membrane filters (BD Biosciences). 4T1/control or 4T1/WISP-1 35 cells (100,000 cells) were added to the upper chamber in 0.5 ml

Iscove's Media. The lower chamber was filled with 0.75 ml of the same media containing 5% fetal bovine serum. Various concentrations of WISP-1-domain-1-Fc or WISP-1-domain-1-His were added to the upper and lower chambers and the plates were incubated overnight at 37°C.

5 The next day the upper chamber was wiped with a cotton swab and the cells that migrated to the lower side of the insert were stained with the Diff-Quik Stain Kit (Dade Behring Inc.) and counted under the microscope. Triplicate sets of data were averaged for each condition. The results are expressed as relative fold invasion

10 compared to 4T1/ control cells. WISP-1-domain-1-His and WISP-1-domain-1-Fc demonstrated a dose dependent antagonistic activity toward WISP-1 promoted 4T1 invasion (Figure 28). Both WISP-1-domain-1-His and WISP-1-domain-1-Fc showed an apparent IC<sub>50</sub> of 0.01 µg/ml. These results suggest that purified WISP-1-domain-1 antagonizes WISP-1 promoted cell invasion.

15

#### EXAMPLE 26

An animal study was conducted to evaluate WISP-1 expression in the HPAC xenograft model of human pancreatic adenocarcinomas. 5 x 20 10<sup>6</sup> human pancreatic adenocarcinomas cells (HPAC cell line, cat. no. CRL-2119; ATCC, Manassas, VA, USA;) were injected subcutaneously in 6-8 week old female nude mice. After 21 days the mice were sacrificed, the tumors were excised and WISP-1 expression was analyzed by *in situ* hybridization using a human WISP-1 specific probe 25 and a mouse WISP-1 specific probe (Figure 29). The results demonstrate a strong mouse WISP-1 expression in areas corresponding to tumor stroma. On the other hand, no human WISP-1 expression was found in the excised tumors. In addition, the tumors revealed a strong α-smooth muscle actin and a weak vimentin immunohistochemical 30 staining over the tumoral stroma. These results indicate that HPAC cells inoculation in nude mice induces the recruitment of a mouse myofibroblastic stroma in which WISP-1 is expressed.

#### EXAMPLE 27

35 An animal study was conducted to evaluate the *in vivo* efficacy of WISP-1-domain-1 at retarding tumor growth. The human WISP-1\_1, WISP-1\_12 constructs cloned into an expression vector (pIRES puro2; Clontech Laboratories, Palo Alto, CA) were transfected in the HPAC human pancreatic adenocarcinomas cell line. After 48 hours, the 40 culture media was supplemented with 2 µg/ml puromycin. After 2

weeks, the pools of expressing cells were analyzed and the resulting cell lines were maintained in 50:50 media (1:1 mixture of DME and Ham's F-12 medium) supplemented with 10% fetal calf serum (FBS) and 2  $\mu$ g/ml puromycin.

5 The *in vivo* efficacy of WISP-1-domain-1 at retarding tumor growth was evaluated by injecting  $5 \times 10^6$  cells (HPAC/WISP-1\_12, HPAC/WISP-1\_1 cells) subcutaneously in 6-8 week old female nude mice (10 mice/group). Tumor volumes were measured twice weekly until day 25.

10 The results demonstrate that the expression of WISP-1\_1 or WISP-1\_12 in HPAC cells retards their *in vivo* growth. The tumor volume of HPAC/WISP-1\_12 and HPAC/WISP-1\_1 cells were 12% and 54% smaller respectively compared to HPAC control cells tumors (Figure 30). These results indicates that WISP-1-domain-1 and WISP-1-domain-12 expressed by HPAC cells retard their tumor growth possibly by 15 antagonizing mouse WISP-1 tumor promoting activities.

#### EXAMPLE 28

20 *In situ* hybridization studies were conducted to evaluate WISP-1 expression in primary human pancreatic adenocarcinomas. Using a tumor microarray (TMA), it was shown that 68% (125/184) of the samples cores were positive for WISP-1 expression. In addition, in these samples WISP-1 expression was restricted to the stromal compartment (Figure 31). These results indicate a high incidence of WISP-1 expression in the stromal compartment of human primary 25 pancreatic tumors.

#### EXAMPLE 29

30 Immunohistochemical staining was conducted to evaluate WISP-1 expression in primary colon adenocarcinomas. A panel of 15 samples of primary human adenocarcinomas were stained for WISP-1 using a WISP-1 specific polyclonal goat antibody (R&D systems; cat. No. AF1627). WISP-1 was detected in 100% (15/15) of the samples analyzed. In all cases WISP-1 was located in the stromal compartment surrounding the tumors (Figure 32). These results indicate that 35 WISP-1 is prevalent in the stromal compartment of human colon adenocarcinomas.

WISP-1 constructs:

Domain 2,3:

40 GAATTCAACATGAGGTGGTTCCTGCCCTGGACGCTGGCAGCAGTGACAGCAGGCCAGCACCGTC CTGGCCACTGCAGTGGTCGGTGTGGGCTGCGTCCTGGATGGGGTGCCTACAACAAACGGCCAGTCCTTC

CAGCCTAACTGCAAGTACAAC TGACCGTGCATCGACGGCGGGTGGCTGCACACCCTGTGCCCTCGA  
 GTGCGCCCCCGCGTCTCTGGTGCCCCCACCGCGCGTGAGCATACTGCCACTGCTGTGAGCAG  
 TGGGTATGTGAGGACGACGCCAAGAGGCCACGCAAGACCGCACCCGTGACACAGGAGCCTTCGATGCT  
 5 GTGGGTGAGGTGGAGGCATGGCACAGGAAC TGCA TAGCCTACACAAGCCCCTGGAGCCCTTGCTCCACC  
 AGCTCGGGCTGGGGTCTCCACTCGGATCTCCAATGTTAACGCCAGTGTGCTGCCCTGAGCAAGAGAGC  
 CGCCCTGCAACTTGCGGCCATGCGATGTGGACATCCATACTCATTAAAGCGGCCGACACCACCAT  
 CACCATCACCATCACTAAGTGAGGCCATAGATAACTGATCCAGTGTGCTGGAATTAAATCGCTGTCT  
 GCGA (SEQ ID NO:21)

10

## Domain 1-IgG:

CCTCGGTTCTATCGATTGAATTCCACCATGAGGTGGTTCCCTGCCCTGGACGCTGGCAGCAGTG  
 ACAGCAGCAGCCGCCAGCACCGTCCGGCCACGCCCTCTCTCCAGGCCCTACGACCATGGACTTTACT  
 CCAGCTCCACTGGAGGACACCTCCTCACGCCCCAATTCTGCAAGTGGCCATGTGAGTGCCGCCATCC  
 15 CCACCCCGCTGCCCGCTGGGGTCAGCCTCATCACAGATGGCTGTGAGTGTGTAAGATGTGCGCTCAG  
 CAGCTTGGGGACAAC TGCA CGGAGGCTGCCATCTGTGACCCCCACGGGGCTACTGTGACTACAGC  
 GGGGACCGCCCGAGGTACGCAATAGGAGTGTGTCACAGGTGGTCGGTGGGGTACCGACAAAAC  
 CACACATGCCACCGTGCCAGCACCTGAACTCCTGGGGGACCGTCAGTCTCCTTCCCCCCCCAAA  
 CCCAAGGACACCCCTCATGATCTCCGGACCCCTGAGGTACATGCGTGGTGGACGTGAGCCACGAA  
 20 GACCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGATAATGCCAAGACAAAGCCGCGG  
 GAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCCTACCGTCTGCACCAGGACTGGCTGAAT  
 GGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTCCCAGCCCCATCGAGAAAACCATCTCCAAA  
 GCCAAAGGGCAGCCCCGAGAACCCACAGGTGTACACCCCTGCCCTACCCGGGAAGAGATGACCAAGAAC  
 CAGGTCA CGCCTGACCTGCCCTGGTCAAAGGCTTCTATCCAGCGACATGCCGTGGAGTGGAGAGCAAT  
 25 GGGCAGCCGGAGAACAAACTACAAGACCCACGCCCTCCGTGCTGGACTCCGACGGCTCCTTCCCTAC  
 AGCAAGCTCACCGTGGACAAGAGCAGGTGGCAGCAGGGAACGTCTCTCATGCTCCGTGATGCATGAG  
 GCTCTGCACAACCACTACACCGAGAAGAGCAGCTCCCTGTCTCCGGTAAATGAGTGCAGCGACGGCCCTAG  
 AGTCGACCTGCAGAAGCTTGGCCGCCATGGCC (SEQ ID NO:22)

30

## Deposit of Material

The following materials have been deposited with the  
 American Type Culture Collection, 10801 University Blvd., Manassas,  
 VA 20110-2209, USA (ATCC):

35	<u>Material</u>	<u>ATCC Dep. No.</u>	<u>Deposit Date</u>
	3D11.D7	PTA-4624	Sept. 4, 2002
	11C2.C10	PTA-4628	Sept. 4, 2002
	9C10.F5	PTA-4626	Sept. 4, 2002
	5D4.F6	PTA-4625	Sept. 4, 2002
40	9C11.C7	PTA-4627	Sept. 4, 2002

This deposit was made under the provisions of the Budapest Treaty on the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure and the 5 Regulations thereunder (Budapest Treaty). This assures maintenance of a viable culture of the deposit for 30 years from the date of deposit. The deposit will be made available by ATCC under the terms of the Budapest Treaty, and subject to an agreement between Genentech, Inc. and ATCC, which assures permanent and unrestricted 10 availability of the progeny of the culture of the deposit to the public upon issuance of the pertinent U.S. patent or upon laying open to the public of any U.S. or foreign patent application, whichever comes first, and assures availability of the progeny to one determined by the U.S. Commissioner of Patents and Trademarks to be 15 entitled thereto according to 35 USC '122 and the Commissioner's rules pursuant thereto (including 37 CFR '1.14 with particular reference to 886 OG 638).

The assignee of the present application has agreed that if a culture of the materials on deposit should die or be lost or 20 destroyed when cultivated under suitable conditions, the materials will be promptly replaced on notification with another of the same. Availability of the deposited material is not to be construed as a license to practice the invention in contravention of the rights granted under the authority of any government in accordance with its 25 patent laws.

The foregoing written description is considered to be sufficient to enable one skilled in the art to practice the invention. The present invention is not to be limited in scope by the example presented herein. Indeed, various modifications of the 30 invention in addition to those shown and described herein will become apparent to those skilled in the art from the foregoing description and fall within the scope of the appended claims.

THE CLAIMS DEFINING THE INVENTION ARE AS FOLLOWS:

1. A method of inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, comprising exposing said mammalian cells to an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

- a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and
- b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

2. The method according to claim 1, wherein said WISP-1-domain 1 polypeptide is fused to a Fc region of an immunoglobulin.

3. The method according to claim 1, wherein said mammalian cells comprise cancer cells.

4. The method according to claim 3, wherein said cancer cells comprise pancreatic cancer cells, colon or colorectal cancer cells, breast cancer cells, lung cancer cells, or brain cancer cells.

5. The method according to claim 4, wherein said cancer cells comprise pancreatic adenocarcinoma cells or colonic adenocarcinoma cells.

6. A method of treating cancer in a mammal, comprising administering to said mammal an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

- a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and
- b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

35 7. The method according to claim 6, wherein said WISP-1-domain 1 polypeptide is fused to a Fc region of an immunoglobulin.

40 8. The method according to claim 6, wherein said cancer is a pancreatic cancer, colon or colorectal cancer, breast cancer, lung cancer or brain cancer.

9. The method according to claim 8, wherein said cancer is pancreatic adenocarcinoma or colonic adenocarcinoma.

5 10. The method according to claim 6, wherein said polypeptide inhibits or reduces cancer cell growth or metastasis.

11. The method according to claim 8, wherein said polypeptide inhibits or reduces lung cancer cell metastasis at a site in the mammal secondary or  
10 different from the primary lung tumor site in the mammal.

12. The method according to claim 6, wherein chemotherapy, radiation, a prodrug, a cytotoxic agent, a growth inhibitory agent, or a cytokine is also administered to said mammal.

15 13. The method according to claim 6, wherein said polypeptide inhibits or neutralizes induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell.

20 14. A method of inhibiting cancer cell motility, comprising exposing a cancer cell to an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117

25 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

30 15. The method according to claim 14, wherein said WISP-1-domain 1 polypeptide is fused to a Fc region of an immunoglobulin.

16. The method according to claim 14, wherein said cancer cell is a pancreatic cancer cell, colon or colorectal cancer cell, breast cancer cell, lung cancer cell, or brain cancer cell.

35 17. A method of inhibiting cancer cell motility in a mammal comprising administering to said mammal an effective amount of a WISP-1-domain 1 polypeptide, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

40 a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117

of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

5 18. The method according to claim 17, wherein said WISP-1-domain 1 polypeptide is fused to a Fc region of an immunoglobulin.

10 19. The method according to claim 17, wherein said cancer cell is a pancreatic cancer cell, colon or colorectal cancer cell, breast cancer cell, lung cancer cell, or brain cancer cell.

20. The method according to claim 17, wherein said polypeptide inhibits or neutralizes induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell.

5 21. Use of a WISP-1-domain 1 polypeptide in the manufacture of a medicament for inhibiting or neutralizing WISP-1 induction or secretion of HAS2, HA, CD44 or RHAMM in mammalian cells, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

10 a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

15 22. Use of a WISP-1-domain 1 polypeptide in the manufacture of a medicament for inhibiting cancer cell motility, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

30 a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.

35 23. Use of a WISP-1-domain 1 polypeptide in the manufacture of a medicament for treating cancer, wherein said WISP-1-domain 1 polypeptide consists of a polypeptide selected from the group consisting of:

a) a WISP-1-domain 1 polypeptide consisting of amino acids 24 - 117 of SEQ ID NO:1; and

40 b) a WISP-1-domain 1 polypeptide consisting of a polypeptide with at



21 Feb 2012




2006236649








least about 95% identity to amino acids 24 - 117 of SEQ ID NO:1.




24. The use according to claim 23, wherein said polypeptide inhibits or neutralizes induction or secretion of HAS2, HA, CD44 or RHAMM by native human WISP-1 polypeptide in at least one type of mammalian cell.

25. A method according to any one of claims 1, 6, 14 or 17, or a use according to any one of claims 21 to 23, substantially as herein described with reference to any one of the examples or drawings.

***FIG.\_ 1A***   ***FIG.\_ 1B***   ***FIG.\_ 1C******FIG.\_ 1D******FIG.\_ 1E******FIG.\_ 1F***

**FIG.\_2A****FIG.\_2B**

**FIG.\_3A****FIG.\_3B****FIG.\_3C****FIG.\_3D****FIG.\_3E****FIG.\_3F****FIG.\_3G**

**FIG.\_4A****FIG.\_4B****FIG.\_4C**




FIG.\_5C




FIG.\_5B




FIG.\_5A









CD44 mRNA Expression  
(Relative to Normal Mucosa)HAS2 mRNA Expression  
(Relative to Normal Mucosa)

FIG.\_5E


***FIG.\_6A******FIG.\_6B******FIG.\_6C******FIG.\_6D***

**FIG.\_6E****FIG.\_6F****FIG.\_6G****FIG.\_6H**

**FIG. - 7A**



**FIG. - 7C**




**FIG. - 7E**




**FIG. - 7D**



*FIG. 7G*



*FIG. 7H*



*FIG. 7I*

9/36

*FIG. 7J*


9/36

*FIG. 7K*


9/36

*FIG. 7L*

9/36



***FIG.\_8A***



***FIG.\_8B***

11/36

1 CCCACGGCTC CGCTGGGCC AGCTCCCCC AGAGCTCTC GGATCTCTG GCTCTCTCG TCGATCTCG TGCCAGCC GTCCACTGAC GTCCAGCC GAGGTGGTTC  
GGTGGCAG GCGACCGGG TCGAGGGGC CCTCCACCAAG TCTCCACCAAG CCTAGAGAC CGCACAGGC AGCTACGGAC AGGTGACTG CAGGTGACTG CAGGTGACTG CAGGTGACTG  
M R W F

101 CTGCCCTGGA CGCTGGAGC AGTGACAGCA GCAGGGCCA GCACCGTCCCT GCACCGGGC CTCTCTCCAG CCCCTAGCAC CATGGACTT ACTCCAGCTC  
GACGGACCT GCGACCGTGT CGTACTGTCGT CGTCGGGGT CGTGGAGGA CGGGTGGGG GAGGAGGTC GGGGATGCTG GTACCTGAAA TGAGGTCGAG  
M R W F

5 L P W T L A A V T A A A S T V L A T A L S P A P T T M D F T P A P

201 CACTGAGGA CACCTCCCTCA CGCCCCAAT TCTGCAAGGTG GCCATGTGAG TGCCCGCCAT CCCCCACCCCG CTGCCCCGCTG GGGGTCAGCC TCATCACAGA  
GTGACCTCCCT GTGGAGGGAGT GGGGGTTA AGACGTTACAC CGGTACACTC ACGGGTTACCA GGGGGGGGC GACGGGGGAC CCCAGTCGG AGTAGTGTCT  
39 L E D T S S R P Q F C K W P C E C P P S P P R C P L G V S L I T D

301 TGGCTGTGAG TGTGTAAAGA TGTGGCTCA GACAAGTGG GAGGCTTGGG CGAGGCTGAC CATCTGTGAC CCCACCGGG GCCTCTACTG TGACTACAGC  
ACCGACACTC ACGACATTCT ACACGGAGT CTCGAAACCC CTGTTGACGT GCCTCCGACG GTAGACACTG GGGTGGCCC CGGAGATGAC ACTGATGTCG  
72 G C E C C K M C A Q D N C T E A A I C D P H R G L Y C D Y S

401 GGGGACCGCC CGAGGTACGC AATAGGAGTG TGTGGCACAGG TGGTGGGTG GGGCTGGTC CTGGATGGGG TGCGCTACAA CAACGGCCAG TCCTTCCAGC  
CCCTGGGG GCTCCATGCG TTATCCTCAC ACACGTTGTC ACCAGCCACA CCCGACGGCAG GACCTAACCC ACGGATGTTT GTGGCCGGTC AGGAAGGTCG  
105 G D R P R Y A I G V C A Q V V G V G C V L D G V R Y N N G Q S F Q P

501 CTAACTGCAA GTACAAACTGC ACGTGCATCG ACGGGGGGT GGGCTGCACA CCCGAGTGTGCG TCCGAGTGGCC CCCACCCGGT CTCTGGTGC  
GATTGACGTT CATGGTGACG TGCACGGTAGC TGCCGGGCCA CCCGAGTGTGCG AGGCTACGG GGTGACACGG GAGACACGG GGGGGGGGG  
139 N C K Y N C T C I D G A V G C T P L C L R V R P P R I W C P H P R

601 GCGGGTGGCC ATACCTGGCC ACTGCTGTGA CGAGTGGGTA TGTGAGGAGC AGGCCAAAGG GCCACGCCAG ACGGCACCCC GTGACACAGG AGCCTTCGAT  
CGCGCACTCG TAGGGACCCG TGACGACACT GTACCTCTGC TGGGGTTCTC CGGTGGGG CACTGTTGC TCGGAAGCTA  
172 R V S I P G H C C E Q W V C E D D A K R P R K T A P R D T G A F D

701 GCTGTGGGTG AGGTGGAGG ATGGCACAGG AACTGCAATAG CCTACACAAG CCCCTGGAGC CCTTGCTCCA CGAGCTGGG CCGGGGGTC TCACACTGGGA  
CGACACCCAC TCCACCTCG TACCGTGTCC TGACGATTC GGATGTTTC GGGGACCTCG GGAACGAGGT GGTCGAGGCC GGACCCCCAG AGGTGAGGCT  
205 A V G E V E A W H R N C I A Y T S P W S P C S T S C G L G V S T R I

801 TCTCCAAATGT TAACGCCAG TGCTGGCTG AGCAAGAGAG CGGCCCTCTGC AACTTGCGGC CATGGATGT GGACATCCAT ACACCTTAAAGGCAGGGAA  
AGAGGTTACA ATTGCGGGTC CGGACGGGAC TTGAACGGCG GTACGGTACA CCTGTTAGGTA TGAGGAAAT TCCGGTCCCT  
239 S N V N A Q C W P E Q E S R L C N L R P C D V D I H T L I K A G K

**FIG.-9A**

901 GAAGTGTCTG GCTGTGTACCGCCAGCAAGGGC ATCCATGAACTTCACAGTCGAT CCGGGCTGCAAT CAGGCACACGC TOCTATCAAC CCAAGTACTG TGAGAGTTGCG  
272 K C L A V Y Q . P E A S M N F T L A G C I S T R S Y Q P K Y C G V C  
CTTCACAGAC CGACACATGG TCGGTCTCCG TAGGTACTTG AAGTGTGAAAC GCCCAGCTA GTCGGTGTGCG AGGATAGTTG GGTTCATGAC ACTCTAAACG

1001 ATGGGAAATA GGTGCTGCAAT CCCCTACAAG TCTAAAGACTA TCGAGCTGTC CTTCCAGTGT GACTCTGGGT CCTGATGGGC TTGGCTTCTC CCGCCAGGTC CTATGGATTA  
TACCTGTTAT CCACGACGGTA GGGGATGTTCA AGATTCTGAT AGCTGACAG GAAGGTCAACAA GGACTAACCCG AACCGAAGAG GGGGGTCCAG GATACCTAAT  
305 M D N R C C I P Y K S K T I D V S F Q C P D G L G F S R Q Y L W I N

1101 ATGCCCTGCTT CTGTAACCTG AGCTGTAGGA ATCCCAATGAA CATCTTGT GACTTGGAA CCTACCCCTGA CTCTCTAGAA ATGGCCAACT AGGCAGGGCAC  
TACGGACGAA GACATTGGAC TCGACATCCT TAGGGTTACT GTAGAAACGA CTGAAACCTTA GGATGGACT GAAGAGTCTT TAACGGTTGA TCGGTCCGTG  
339 A C F C N L S C R N P N D I F A D L E S Y P D F S E I A N O

1201 AAATCTTGGG TCTTGGGGAC TAACCCAATG CTCGTGAAGC AGTCAGCCCT TATGGCCAAT AACTTTTCAC CAATGAGCCCT TAGTTACCCCT GATCTGGACC  
TTTAGGAAACCC AGAACCCCTG ATTGGTTAC GAGACACTTCG TCAGTGGGA ATACCGGTTA TTGAAAAGTG GTTAACCTCGGA ATCAATGGGA CTAGACCTGG

1301 CTTGGCCTCC ATTTCTGCTT CTAACCATTCAATGACGCC TGATGGCT GCTCAGGGCC ATGCTATGAG TTTTCTCTT GATATCATT AGCATCTACT  
GAACGGAGG TAAAGACAGA GATGGTAAG TTTACTGCGG ACTACACGA CGAGTCGGG TAGGATACTC AAAGAGGAA CTATAGTAAG TCGTAGATGA

1401 CTTAAAGAAA ATGCGCTGCTT CTAAGCTGTC TGGACTACAC CCAAGCTGTA TCCAGGCCCTT CCAAGTCACT AGAAGTCCCTG CTGGATCTTG CCTAAATCCC  
GATTCTTCTT TAGGGACAGA GATGGACAGA GATGGACAAAG AGCTGATGTG GGTTCGGACT AGGTGGAA GGTTCACTGAA GACCTAGAAC GGATTCTAGGG

1501 AAGAAATGGA ATCAGGTAGA CTTTTAAT CACTAATTCTT CTTCTTAGAT GCCAAACCA AAGACTCTT GGTCAGGCC ATGAAATAG ATGGAATTG  
TTCTTACCT TAGCCATCTT GAAATTATA GTGATTAAG AAGAATCTA CGGTGGGTG TTCTGAGAA CCAGGTAG AGTACTATC TACCTTAAC

1601 GAACAAATAGA ATAATCTTATT ATTGGAGCC TGCCAAGAGG TACTGTAATG GGTAAATTCTG ACGTCAAGGC ACAAAACAA TCCTGTATCC AAATATGTAT  
CTTGTATCTT TATTAGATAA TAAACCTCGG ACGGTTCTCC ATGACATTAC CCATTAAGAC TGCAGTCGCG TGTTTTGAT AGGACTAAGG TTATACATA

1701 GCACCTCAAG GTCATCAAC ATTGGCAAG TGAGTGTAA AGTGTGTTAA TTTTGATT TAATGAAAG TTGTATCCAT TAACCTGGG ATGTTGAGG  
CGTGGAGTTC CAGTAGTTG TAAACGGTTCACTCAACTTA TCAACGAATT AAAACTAAA ATTACCTTC AACATAGGTAA ATTGGACCCG TAACAACCTCC

1801 TTAAGTTCTT CTTCACCCCT ACACGTGAA GGGTACAGAT TAGGTGAACTT CCAAGTGTCAAT ATCCAAACAG GTTCAGTCTT TATTAAAC TATTGTAAAG GACAACCTAC  
AATTCAAAGA GAAGTGGGA TGTGACACTT CCCATGTCAAT ATCCAAACAG GTTCAGTCTT TATTAAAC TATTGTAAAG GACAACCTAC CTTTTGGGG


1901 CAGTTAAATAC TCCAGAGAGCA GGGAAAGGTC AGCCCAATTTC AGAAGGACCA ATTGACTCTC ACACGTAAATC AGTGTGCTGAC TGGCAGGGCT TGGGCAAGT  
GTCAAAATTATG AGGTCTCTGT CCCTTCCAG TCGGTAAAG TCTTCTCTGT TAACTGAGAG TGTGACTTAG TGACGACTG ACCGTCGGAA AACCCGGTCAA


**FIG.-9B**


2001 GGCCAGGCTC TTCCCTTGAAT CTTCTCCCTT GTCCCTGCTG GGTTCATAGG ATTGGTAAG GCCTCTGGAC TGGCCTGCTT GGCCCCCTGAG AGTGGTGGCC  
 CCGGTCCGAG AAGGAACCTTA GAAGGGAA CAGGACCAAC CCAAGTATCC TTAACCCTTC CCGAGACCTG ACCGGACAGA CCGGGGACTC TCACCAACGGG  
  
 2101 TGGAAACACTC CTCTACTCTT ACAGAGCCTT GAGAGACCA GCTGCAGACC ATGCCAGACC CACTGAAATG ACCAAGACAG GTTCAGGTAG GGGTGTGGGT  
 ACCCTGTGAG GAGATGAGAA TGTCTCGAA CTCCTCGGGT CGACCGTCGG TAGGGTCTGG GTGACTTTAC TGGTCTGTGCA CAAGTCCATC CCACACCCCA  
  
 2201 CAAACCAAGA ACTGGGTGCC CTTGGTAGCA CCCTGGGGTG ACCCTCTAGAG CTGGAGGTG TGGGACTCCA GGGGGCCCG TGTTCAGGAC ACATCTATG  
 GTTGGTTCT TCACCCACGG GAACCATCGT CGGACCCAC TGGAGATCTC GACCTCCGAC ACCCTGAGGT CCCGGGGGC ACAAGTCCCTG TGAGATAAC  
  
 2301 CAGAGACTCA TTTCACAGCC TTTCGTTCTG CTGACCAAAT GGCCAGTTT CTGGTAGGA GATGGAGGT TACCAAGTTT TTAGAACAG AAATAGACTT  
 GTCTCTGAGT AAAGTGTGG AAAGCAAGAC GACTGGTTA CCGGTCAAAA GACCATCCCTT CTACCTCCAA ATGGTCAACA AATCTTGTC TTATATCTGAA  
  
 2401 AATAAAGGTT TAAAGCTGAA GAGGGTGAAG CTAAAAGGAA AAGGGTGTGG TTAATGAATA TCAGGCTATT ATTATTTGTA TTAGGAAAT ATAATATTTA  
 TTATTCCTAA ATTTCGACTT CTCCAACTTC GATTTTCCTT TTCCAAACAC ATTACTTAT AGTCCGATAA TAAATAACAT AATCCCTTTA TATTATAATT  
  
 2501 CTGTTAGAAT TCTTTTATT AGGGCTTT CTGGCCAGA CATTGCTCTC AGTGGCTTGC ATGTTATTAGC TCACTGAATC TTCAAGACAA TTGTTGAGAAG  
 GACAATCTTA AGAAATAAA TCCCCGAAA GACACGGTCT GTAACGAGAG TACATAATCG AGTGACTTAG AAGTGGTGTGTT ACAACTCTTC  
  
 2601 TTCCCATTT TATTCTGTT CTTACAAATG TGAAACGGAA GCTCATAGAG GTGAGAAAC TCAACAGAG TCACCCAGT GGTGACTGGG AAAGTTAGGA  
 AAGGTAATA ATAAAGACAA GAATGTTTAC ACTTTGCCTT CGAGTATCTC CACTCTTTG AGTGGTCTC AGTGGGTCAA CCACTGACCC TTCAATCTCT  
  
 2701 TTCAAGATCGA ATTGGACTG TCTTATAAC CCATATTTC CCCCTGTTT TAGAGCTTCC AAATGTGCA GAATAGGAAA ACATTTGCAAT AAATGGCTTG  
 AAGTCTAGCT TTAACCTGAC AGAAATATTG GGTATAAAAG GGGGACAAAA ATCTCGAAGG TTTACACAGT CTATCCCTT TGTAACGTTA TTACCGAAC  
  
 2801 ATTTTTAAA AAAAAAAA AAAAAAAA  
 TAAAAAATTT TTTTTTTTTT TTTTTTTT


**FIG.-9C**


Effect of WISP-1 Expression on NRK Metastatic Potential				
No. Cells Injected	Weeks after Innoculation	Incidence of Lung Metastasis		
		NRK/Control	NRK/WISP-1L	NRK/WISP-1H
$2.5 \times 10^5$	2	0/1	0/1	2/2 (Grade I)
	3	0/4	3/3 (Grade I)	4/4 (Grade I-II)
	4	0/4	4/4 (Grade I)	3/3 (Grade III)
$0.5 \times 10^5$	2	0/2	0/2	1/2 (Grade I)
	3	0/4	2/4 (Grade I)	3/4 (Grade I)
	4	0/4	4/4 (Grade I)	3/4 (Grade I-II)


**FIG.\_10**















**FIG. 12E**





**FIG. 12F**













**FIG. 14**




**FIG.\_15****FIG.\_16A**

WISP-1 Antibodies Characterization								
Clones	Isotype	Epitope		IP	WB	Blocking Activity		
		Struct.	Cross Compet.			Heparin Binding	Cell Migrat.	Lung Met.
3D11	2b	Domain 1	A	—	+	+/-	+	+
9C10	2b	Domain 1	B	+	—	—	+	+
11C2	2b	Variable Region	C	+	+	+	+	+
5D4	2a	Variable Region	C	+	+	+	+	+
9C11	2a	Variable Region	C	+	+	+	+	+

**FIG.\_ 16B****FIG.\_ 17A****FIG.\_ 17B**

**FIG.\_ 17C****FIG.\_ 17D****FIG.\_ 17E**


**FIG.\_18****FIG.\_19****FIG.\_20**



*FIG.\_23B*




*FIG.\_23A*



*FIG.\_23D*



*FIG.\_23C*



*FIG.\_23F*



*FIG.\_23E*






Figure 24




Figure 25

Vascular Invasion in NRK/WISP-1\_134 Xenograft




Figure 26




Figure 27




Figure 28




Figure 29






Figure 30




H&E Staining



WISP-1 *in situ* Hybridization

Figure 31

HP9278



HP9300





Figure 32

P1918R1P1.txt  
Sequence Listing

<110> GENENTECH, INC.  
DESNOYERS, LUC  
FILVAROFF, ELLEN

<120> Methods and Compositions for Modulating and Detecting  
WISP Activitiy

<130> P1918R1P1

<150> US 11/105,876

<151> 2005-04-14

<160> 22

<210> 1  
<211> 367  
<212> PRT  
<213> Homo sapiens

<400> 1

Met	Arg	Trp	Phe	Leu	Pro	Trp	Thr	Leu	Ala	Ala	Val	Thr	Ala	Ala
1				5				10					15	
Ala	Ala	Ser	Thr	Val	Leu	Ala	Thr	Ala	Leu	Ser	Pro	Ala	Pro	Thr
				20				25					30	
Thr	Met	Asp	Phe	Thr	Pro	Ala	Pro	Leu	Glu	Asp	Thr	Ser	Ser	Arg
	35							40					45	
Pro	Gln	Phe	Cys	Lys	Trp	Pro	Cys	Glu	Cys	Pro	Pro	Ser	Pro	Pro
	50							55					60	
Arg	Cys	Pro	Leu	Gly	Val	Ser	Leu	Ile	Thr	Asp	Gly	Cys	Glu	Cys
	65							70					75	
Cys	Lys	Met	Cys	Ala	Gln	Gln	Leu	Gly	Asp	Asn	Cys	Thr	Glu	Ala
	80							85					90	
Ala	Ile	Cys	Asp	Pro	His	Arg	Gly	Leu	Tyr	Cys	Asp	Tyr	Ser	Gly
	95							100					105	
Asp	Arg	Pro	Arg	Tyr	Ala	Ile	Gly	Val	Cys	Ala	Gln	Val	Val	Gly
	110								115				120	
Val	Gly	Cys	Val	Leu	Asp	Gly	Val	Arg	Tyr	Asn	Asn	Gly	Gln	Ser
	125								130				135	
Phe	Gln	Pro	Asn	Cys	Lys	Tyr	Asn	Cys	Thr	Cys	Ile	Asp	Gly	Ala
	140								145				150	
Val	Gly	Cys	Thr	Pro	Leu	Cys	Leu	Arg	Val	Arg	Pro	Pro	Arg	Leu
	155								160				165	
Trp	Cys	Pro	His	Pro	Arg	Arg	Val	Ser	Ile	Pro	Gly	His	Cys	Cys
	170								175				180	
Glu	Gln	Trp	Val	Cys	Glu	Asp	Asp	Ala	Lys	Arg	Pro	Arg	Lys	Thr
	185								190				195	
Ala	Pro	Arg	Asp	Thr	Gly	Ala	Phe	Asp	Ala	Val	Gly	Glu	Val	Glu
	200									205			210	
Ala	Trp	His	Arg	Asn	Cys	Ile	Ala	Tyr	Thr	Ser	Pro	Trp	Ser	Pro
	215								220				225	

P1918R1P1.txt

Cys	Ser	Thr	Ser	Cys	Gly	Leu	Gly	Val	Ser	Thr	Arg	Ile	Ser	Asn
230														240
Val	Asn	Ala	Gln	Cys	Trp	Pro	Glu	Gln	Glu	Ser	Arg	Leu	Cys	Asn
245														255
Leu	Arg	Pro	Cys	Asp	Val	Asp	Ile	His	Thr	Leu	Ile	Lys	Ala	Gly
260														270
Lys	Lys	Cys	Leu	Ala	Val	Tyr	Gln	Pro	Glu	Ala	Ser	Met	Asn	Phe
275														285
Thr	Leu	Ala	Gly	Cys	Ile	Ser	Thr	Arg	Ser	Tyr	Gln	Pro	Lys	Tyr
290														300
Cys	Gly	Val	Cys	Met	Asp	Asn	Arg	Cys	Ile	Pro	Tyr	Lys	Ser	
305														315
Lys	Thr	Ile	Asp	Val	Ser	Phe	Gln	Cys	Pro	Asp	Gly	Leu	Gly	Phe
320														330
Ser	Arg	Gln	Val	Leu	Trp	Ile	Asn	Ala	Cys	Phe	Cys	Asn	Leu	Ser
335														345
Cys	Arg	Asn	Pro	Asn	Asp	Ile	Phe	Ala	Asp	Leu	Glu	Ser	Tyr	Pro
350														360
Asp	Phe	Ser	Glu	Ile	Ala	Asn								
365														

&lt;210&gt; 2

&lt;211&gt; 2830

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 2

```

cccacgcgtc cgctgggccc agctccccc agaggtggtc ggatcctctg 50
ggctgctcgg tcgatgcctg tgccactgac gtccaggcat gaggtggttc 100
ctgcccctgga cgctggcagc agtgcacagca gcagccgcca gcaccgtcct 150
ggccacggcc ctctctccag cccctacgac catggacttt actccagctc 200
caactggagga cacctcctca cgcccccata tctgcaagtg gccatgtgag 250
tgcccgccat cccccaccccg ctgcccgcgt ggggtcagcc tcacacaga 300
tggctgtgag tgctgttaaga tgtgcgtca gcagcttggg gacaactgca 350
cgaggctgc catctgtgac ccccacccgg gcctctactg tgactacagc 400
ggggaccgccc cgaggtacgc aataggagtg tgtgcacagg tggtcggtgt 450
gggctgcgtc ctggatgggg tgcgtacaa caacggccag tccttccagc 500
ctaactgcaa gtacaactgc acgtgcacatcg acggcgcggg gggctgcaca 550
ccactgtgcc tccgagtgcg ccccccgcgt ctctggtgcc cccacccgcg 600
gcgcgtgagc atacctggcc actgctgtga gcagtgggta tgtgaggacg 650
acgccaagag gccacgcaga accgcacccccc gtgacacagg agccttcgat 700
gctgtgggtg aggtggaggc atggcacacagg aactgcatacg cctacacaag 750
ccccctggagc ctttgctcca ccagctgcgg cctgggggtc tccactcgga 800

```

P1918R1P1.txt  
tctccaatgt taacgcccag tgctggcctg agcaagagag ccgcctctgc 850  
aacttgcggc catgcgatgt ggacatccat acactcatta aggcaggaa 900  
gaagtgtctg gctgtgtacc agccagaggc atccatgaac ttcacacttg 950  
cgggctgcat cagcacacgc tcctatcaac ccaagtactg tggagttgc 1000  
atggacaata ggtgctgcat cccctacaag tctaagacta tcgacgtgtc 1050  
cttccagtgt cctgatgggc ttggcttctc ccgcaggc ctagggatta 1100  
atgcctgctt ctgtaacctg agctgttagga atcccaatga catcttgct 1150  
gacttggaat cctaccctga cttctcagaa attgccaact aggcaggcac 1200  
aaatcttggg tcttggggac taacccaatg cctgtgaagc agtcagccct 1250  
tatggccaat aactttcac caatgagcct tagttaccct gatctggacc 1300  
cttggcctcc atttctgtct ctaaccatc aaatgacgcc ttagtggct 1350  
gctcaggccc atgctatgag ttttccctt gatatcattc agcatctact 1400  
ctaaagaaaa atgcctgtct ctagctgttc tggactacac ccaagccctga 1450  
tccagccctt ccaagtcact agaagtcctg ctggatcttg cctaaatccc 1500  
aagaaatgga atcaggtaga ctttaatat cactaattc ttcttttagat 1550  
gccaaaccac aagactcttt gggtccattc agatgaatag atgaaatttg 1600  
gaacaataga ataatctatt atttggagcc tgccaagagg tactgtaatg 1650  
ggtaattctg acgtcagcgc accaaaacta tcctgattcc aaatatgtat 1700  
gcaccccaag gtcataaac atttgccaag tgagttgaat agttgctaa 1750  
ttttgatttt taatggaaag ttgtatccat taacctggc attgttgagg 1800  
ttaagttct cttcacccct acactgtgaa gggtacagat taggtttgtc 1850  
ccagtcagaa ataaaatttg ataaacattc ctgttgcattt gaaaagcccc 1900  
cagttataac tccagagaca gggaaaggc agcccatattc agaaggacca 1950  
attgactctc acactgaatc agctgctgac tggcaggcgt ttggcagtt 2000  
ggccaggcgtc ttccctgaat cttctccctt gtccctgctt ggttcatagg 2050  
aattggtaag gcctctggac tggcctgtct ggcccctgag agtgggtgcc 2100  
tggaaacactc ctctactctt acagagcctt gagagacca gctgcagacc 2150  
atgccagacc cactgaaatg accaagacag gttcaggtag gggtgtgggt 2200  
caaaccaaga agtgggtgcc cttggtagca gcctgggtg acctcttagag 2250  
ctggaggcgtg tggactcca ggggcccccg tgttcaggac acatctattg 2300  
cagagactca tttcacagcc tttcgttctg ctgaccaat ggccagtttt 2350  
ctggtaggaa gatggaggtt taccagttgt ttagaaacag aaatagactt 2400  
aataaagggtt taaagctgaa gaggttgaag ctaaaaggaa aaggttgg 2450  
ttaatgaata tcaggctatt atttattgtt ttaggaaaat ataatattt 2500

P1918R1P1.txt  
ctgttagaat tcttttattt agggccttt ctgtgccaga cattgctctc 2550  
agtgcattgc atgtatttagc tcactgaatc ttacgacaa tggtgagaag 2600  
ttcccattat tatttctgtt cttacaaatg tgaaacggaa gctcatagag 2650  
gtgagaaaac tcaaccagag tcacccagtt ggtgactggg aaagtttagga 2700  
ttcagatcga aattggactg tcttataaac ccatatttc cccctgttt 2750  
tagagcttcc aaatgtgtca gaataggaaa acattgcaat aaatggcttg 2800  
atttttaaa aaaaaaaaaaaa aaaaaaaaaa 2830  
<210> 3  
<211> 440  
<212> DNA  
<213> Homo sapiens  
  
<400> 3  
gaattcacca tgaggtggtt cctgcccctgg acgctggcag cagtgacagc 50  
agcagccgccc agcaccgtcc tggccacggc cctctctcca gcccctacga 100  
ccatggactt tactccagct ccactggagg acaccccttc acgcccccaa 150  
ttctgcaagt ggccatgtga gtgcccggca tccccacccc gctgcccgt 200  
gggggtcagc ctcatcacag atggctgtga gtgctgtaag atgtgcgtc 250  
agcagcttgg ggacaactgc acggaggctg ccatctgtga cccccaccgg 300  
ggcctctact gtgactacag cggggaccgc ccgaggtacg caataggagt 350  
gtgtgcacag gcggccgcac accaccatca ccatcaccat cactaagtga 400  
ggccgcatacg ataaactgatc cagtgtgtcga attaatttc 440  
<210> 4  
<211> 340  
<212> DNA  
<213> Homo sapiens  
  
<400> 4  
gaattcacca tgaggtggtt cctgcccctgg acgctggcag cagtgacagc 50  
agcagccgccc agcaccgtcc tggccactgac agtggtcgggt gtgggctgcg 100  
tcctggatgg ggtgcgtac aacaacggcc agtccttcca gcctaactgc 150  
aagtacaact gcacgtgcat cgacggcgcg gtgggctgca caccactgtg 200  
cctccgagtgcg cggcccccgc gtctctggcgtt ccccccacccg cggcgcgtga 250  
gcataacctgg ccactgctgt gagcagtggg tatgtgcggc cgcacaccac 300  
catcaccatc accatcacta agtgaggccg catagataac 340  
<210> 5  
<211> 321  
<212> DNA  
<213> Homo sapiens  
  
<400> 5  
gaattcacca tgaggtggtt cctgcccctgg acgctggcag cagtgacagc 50  
agcagccgccc agcaccgtcc tggccactgac agcatggcac aggaactgca 100

P1918R1P1.txt  
tagcctacac aagccctgg agcccttgct ccaccagctg cggcctgggg 150  
gtctccactc gatatctccaa tgttaacgcc cagtgtggc ctgagcaaga 200  
gagccgcctc tgcaacttgc ggccatgcga tgtggacatc catacactca 250  
ttaaggcggc cgcacaccac catcaccatc accatcacta agtgaggccg 300  
catagataac tgatccagtg t 321

<210> 6  
<211> 442  
<212> DNA  
<213> Homo sapiens

<400> 6  
gaattcacca tgaggtggtt cctgcccggc acgctggcag cagtgcacgc 50  
agcagccgccc agcaccgtcc tggccactgc agggaaagaag tgtctggctg 100  
tgttaccagcc agaggcatcc atgaacttca cacttgcggg ctgcacatcagc 150  
acacgctcct atcaacccaa gtactgtgga gtttgcattgg acaataggtg 200  
ctgcacatcccc tacaagtcta agactatcga cgtgtcccttc cagtgtccctg 250  
atgggcttgg cttctcccgcc caggtccatgg ggattatgc ctgcttctgt 300  
aacctgagct gtaggaatcc caatgacatc tttgctgact tggaaatccta 350  
ccctgacttc tcagaaatttgc ccaacgcggc cgcacaccac catcaccatc 400  
accatcacta agtgaggccg catagataac tgatccagtg tg 442

<210> 7  
<211> 619  
<212> DNA  
<213> Homo sapiens

<400> 7  
gaattcacca tgaggtggtt cctgcccggc acgctggcag cagtgcacgc 50  
agcagccgccc agcaccgtcc tggccacggc cctctctcca gcccctacga 100  
ccatggactt tactccagct ccactggagg acaccccttc acgcccccaa 150  
ttctgcaagt ggccatgtga gtgcccggca tccccacccc gctgcccgt 200  
gggggtcagc ctcacatcacag atggctgtga gtgctgtaaatgtgcgc 250  
agcagcttgg ggacaactgc acggaggctg ccatctgtga ccccccacccgg 300  
ggcctctact gtgactacag cggggaccgc ccgaggtacg caataggagt 350  
gtgtgcacag gtggctgggt tggtctgcgt cctggatggg gtgcgttaca 400  
acaacggcca gtccttccag cctaactgca agtacaactg cacgtgcac 450  
gacggcgcgg tgggctgcac accactgtgc ctccgagtgc gcccccccg 500  
tctctgggtc ccccacccgc ggcgcgtgag catacctggc cactgctgtg 550  
agcagttgggt atgtgcggcc gcacaccacc atcaccatca ccatcactaa 600  
gtgaggccgc atagataac 619

<210> 8  
<211> 885

## P1918R1P1.txt

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 8

gaattcacca tgaggtggtt cctgcccctgg acgctggcag cagtgacagc 50  
agcagccgccc agcaccgtcc tggccacggc cctctctcca gcccctacga 100  
ccatggactt tactccagct ccactggagg acacccctc acgcccccaa 150  
ttctgcaagt ggccatgtga gtgcccggca tccccacccc gctgcccgt 200  
gggggtcagc ctcacacag atggctgtga gtgctgtaa atgtgcgtc 250  
agcagcttgg ggacaactgc acggaggctg ccacatgtga cccccaccgg 300  
ggcctctact gtgactacag cggggaccgc ccgaggtacg caataggagt 350  
gtgtgcacag gtggctgggt tgggctgcgt cctggatggg gtgcgtaca 400  
acaacggcca gtccttccag cctaactgca agtacaactg cacgtgcac 450  
gacggcgcgg tgggctgcac accactgtgc ctccgagtgc gccccccgcg 500  
tctctgggtc ccccacccgc ggccgtgag catacctggc cactgctgtg 550  
agcagtgggt atgtgaggac gacgccaaga ggccacgcaa gaccgcaccc 600  
cgtgacacag gagccttcga tgctgtgggt gaggtggagg catggcacag 650  
gaactgcata gcctacacaa gcccctggag cccttgctcc accagctgcg 700  
gcctgggggt ctccactcggt atctccaatg ttaacgcccgt 750  
gagcaagaga gcccctctg caacttgcgg ccatgcgtatg tggacatcca 800  
tacactcatt aaggcggccg cacaccacca tcaccatcac catcactaag 850  
tgaggccgca tagataactg atccagtgtg ctgga 885

&lt;210&gt; 9

&lt;211&gt; 1014

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

gaattcacca tgaggtggtt cctgcccctgg acgctggcag cagtgacagc 50  
agcagccgccc agcaccgtcc tggccacggc cctctctcca gcccctacga 100  
ccatggactt tactccagct ccactggagg acacccctc acgcccccaa 150  
ttctgcaagt ggccatgtga gtgcccggca tccccacccc gctgcccgt 200  
gggggtcagc ctcacacag atggctgtga gtgctgtaa atgtgcgtc 250  
agcagcttgg ggacaactgc acggaggctg ccacatgtga cccccaccgg 300  
ggcctctact gtgactacag cggggaccgc ccgaggtacg caataggagt 350  
gtgtgcacag gtggctgggt tgggctgcgt cctggatggg gtgcgtaca 400  
acaacggcca gtccttccag cctaactgca agtacaactg cacgtgcac 450  
gacggcgcgg tgggctgcac accactgtgc ctccgagtgc gccccccgcg 500  
tctctgggtc ccccacccgc ggccgtgag catacctggc cactgctgtg 550

P1918R1P1.txt  
agcagtgggt atgtctgcag gcagggaaaga agtgtctggc tgtgtaccag 600  
ccagaggcat ccatgaactt cacacttgcg ggctgcatca gcacacgctc 650  
ctatcaaccc aagtactgtg gagttgcat ggacaatagg tgctgcatcc 700  
cctacaagtc taagactatc gacgtgtcct tccagtgtcc tcatgggctt 750  
ggcttctccc gccaggtcct atggattaat gcctgcttct gtaacctgag 800  
ctgttaggaat cccaatgaca tctttgctga cttgaaatcc taccctgact 850  
tctcagaaat tgccaacgcg gccgcacacc accatcacca tcaccatcac 900  
taagtgaggc cgcatagata actgatccag tgtgctggaa ttaattcgct 950  
gtctgcgagg gccagctgtt ggggtgagta ctccctctca aaagcgggca 1000  
tgacttctgc gcta 1014  
<210> 10  
<211> 904  
<212> DNA  
<213> Homo sapiens  
<400> 10  
gaattcacca tgaggtggtt cctgcccgg acgctggcag cagtgacagc 50  
agcagccgcc agcaccgtcc tggccacggc cctctctcca gcccctacga 100  
ccatggactt tactccagct ccactggagg acaccccttc acgcccccaa 150  
ttctgcaagt ggccatgtga gtgcccggca tccccacccc gctgcccgt 200  
gggggtcagc ctcatcacag atggctgtga gtgctgtaa atgtgcgctc 250  
agcagcttgg ggacaactgc acggaggctg ccacatgtga cccccaccgg 300  
ggcctctact gtgactacag cggggaccgc ccgaggtacg caataggagt 350  
gtgtgcgcat gctgtgggtg aggtggaggc atggcacagg aactgcata 400  
cctacacaag cccctggagc cttgctcca ccagctgcgg cctgggggtc 450  
tccactcgga tctccaatgt taacgcccag tgctggcctg agcaagagag 500  
ccgcctctgc aacttgcggc catgcgttgt ggacatccat acactcatta 550  
aggcaggaa gaagtgtctg gctgtgtacc agccagaggc atccatgaac 600  
ttcacacttgc cgggctgcat cagcacacgc tcctatcaac ccaagtactg 650  
tggagttgc atggacaata ggtgctgcat cccctacaag tctaagacta 700  
tcgacgtgtc cttccagtgt cctgatgggc ttggcttctc cggccaggtc 750  
ctatggatta atgcctgctt ctgtAACCTG agctgttagga atcccaatga 800  
catcttgct gacttggaaat cctaccctga cttctcagaa attgccaacg 850  
cggccgcaca ccaccatcac catcaccatc actaagttag gcccata 900  
taac 904  
<210> 11  
<211> 922  
<212> DNA  
<213> Homo sapiens

## P1918R1P1.txt

<400> 11  
gaattcacca tgagggtggtt cctgcccctgg acgctggcag cagtacacgc 50  
agcagccgccc agcaccgtcc tggccactgc agtggtcggt gtgggctgcg 100  
tcctggatgg ggtgcgctac aacaacggcc agtccttcca gcctaactgc 150  
aagtacaact gcacgtgcat cgacggcgcg gtgggctgca caccactgtg 200  
cctccgagtg cgccccccgc gtctctgggtg ccccccacccg cggcgcggtga 250  
gcataccctgg ccactgctgt gagcagtggg tatgtgagga cgacgccaag 300  
aggccacgca agaccgcacc ccgtgacaca ggagccttcg atgctgtggg 350  
tgagggtggag gcatggcaca ggaactgcat agcctacaca agccccttgg 400  
gcccttgctc caccagctgc ggcctggggg tctccactcg gatctccaaat 450  
gttaacgccc agtgctggcc tgagcaagag agccgcctct gcaacttgcg 500  
gccatgcgat gtggacatcc atacactcat taaggcaggg aagaagtgtc 550  
tggctgtgta ccagccagag gcatccatga acttcacact tgcgggctgc 600  
atcagcacac gctcctatca acccaagtac tgtggagttt gcatggacaa 650  
taggtgctgc atccccatac agtctaagac tatcgacgtg tccttccagt 700  
gtcctgatgg gcttggcttc tcccgccagg tcctatggat taatgcctgc 750  
ttctgttaacc tgagctgttag gaatccaaat gacatctttg ctgacttgg 800  
atcctaccct gacttctcag aaattgccaa cgcggccgca caccaccatc 850  
accatcacca tcactaagtg aggccgcata gataactgat ccagtgtgct 900  
ggaattaaatt cgctgtctgc ga 922

<210> 12

<211> 19

<212> DNA

<213> Mus musculus

<400> 12  
ggctgccatc tgtgaccca 19

<210> 13

<211> 21

<212> DNA

<213> Mus musculus

<400> 13  
cataggacct gcccggagaa a 21

<210> 14

<211> 19

<212> DNA

<213> Mus musculus

<400> 14  
gccgtggcag tcctgaggg 19

<210> 15

<211> 21

<212> DNA

<213> Mus musculus

## P1918R1P1.txt

<400> 15  
cagcaccggg cattgacgtt a 21

<210> 16  
<211> 21  
<212> DNA  
<213> Mus musculus

<400> 16  
tggagaaaaa tggccgctac a 21

<210> 17  
<211> 20  
<212> DNA  
<213> Mus musculus

<400> 17  
tggggtgctc ttctcgatgg 20

<210> 18  
<211> 21  
<212> DNA  
<213> Mus musculus

<400> 18  
ggacaaatcg gccacgtaca t 21

<210> 19  
<211> 19  
<212> DNA  
<213> Mus musculus

<400> 19  
cttgctccat cgggtctgc 19

<210> 20  
<211> 42  
<212> PRT  
<213> Homo sapiens

<400> 20  
Glu Gln Trp Val Cys Glu Asp Asp Ala Lys Arg Pro Arg Lys Thr  
1 5 10 15

Ala Pro Arg Asp Thr Gly Ala Phe Asp Ala Val Gly Glu Val Glu  
20 25 30

Ala Trp His Arg Asn Cys Ile Ala Tyr Thr Ser Pro  
35 40

<210> 21  
<211> 625  
<212> DNA  
<213> Homo sapiens

<400> 21  
gaattcacca tgaggtggtt cctgccctgg acgctggcag cagtgacagc 50  
agcagccgcc agcaccgtcc tggccactgc agtggtcgggt gtgggctgcg 100  
tcctggatgg ggtgcgtac aacaacggcc agtccttcca gcctaactgc 150  
aagtacaact gcacgtgcat cgacggcgcg gtgggctgca caccactgtg 200  
cctccgagtg cgccccccgc gtctctggtg cccccacccg cggcgcgtga 250  
gcataacctgg ccactgctgt gagcagtggg tatgtgagga cgacgccaag 300

## P1918R1P1.txt

aggccacgca agaccgcacc ccgtgacaca ggagccttcg atgctgtgg 350  
tgaggtggag gcatggcaca ggaactgcat agcctacaca agcccctgga 400  
gcccttgctc caccagctgc ggcctgggg tctccactcg gatctccaaat 450  
gttaacgccc agtgctggcc tgagcaagag agccgcctct gcaacttgcg 500  
gccccatgcgt gtggacatcc atacactcat taaggcggcc gcacaccacc 550  
atcaccatca ccatcactaa gtgaggccgc atagataact gatccagtgt 600  
gctggaatta attcgctgtc tgcga 625

<210> 22  
<211> 1131  
<212> DNA  
<213> Homo sapiens

<400> 22  
cctcggttct atcgattgaa ttccaccatg aggtgggttcc tgccctggac 50  
gctggcagca gtgacagcag cagccgcag caccgtcctg gccacggccc 100  
tctctccagc ccctacgacc atggacttta ctccagctcc actggaggac 150  
acctccctcac gcccccaatt ctgcaagtgg ccatgtgagt gcccgcacatc 200  
cccacccgc tgcccgctgg gggtcagcct catcacagat ggctgtgagt 250  
gctgttaagat gtgcgctcag cagcttgggg acaactgcac ggaggctgcc 300  
atctgtgacc cccaccgggg cctctactgt gactacagcg gggaccgc 350  
gaggtacgca ataggagtgt gtgcacaggt ggtcggtgtg ggggtcaccg 400  
acaactca cacatgccc cctgtccag cacctgaact cctgggggga 450  
ccgtcagtct tccttttccc cccaaaaccc aaggacaccc tcatgatctc 500  
ccggaccctt gaggtcacat gcgtgggtt ggacgtgagc cacgaagacc 550  
ctgaggtcaa gttcaactgg tacgtggacg gcgtggaggt gcataatgcc 600  
aagacaaagc cgcgggagga gcagtacaac agcacgtacc gtgtggtcag 650  
cgtcctcacc gtcctgcacc aggactggct gaatggcaag gagtacaagt 700  
gcaaggcttc caacaaagcc ctcccaagccc ccatcgagaa aaccatctcc 750  
aaagccaaag ggcagccccc agaaccacag gtgtacaccc tgccccatc 800  
ccggaaagag atgaccaaga accaggtcag cctgacctgc ctggtaaag 850  
gcttctatcc cagcgacatc gccgtggagt gggagagcaa tggcagccg 900  
gagaacaact acaagaccac gcctccgtg ctggactccg acggctccctt 950  
cttcctctac agcaagctca ccgtggacaa gagcaggtgg cagcaggggga 1000  
acgtcttc atgctccgtg atgcatgagg ctctgcacaa ccactacacg 1050  
cagaagagcc tctccctgtc tccgggtaaa tgagtgcac gcccctagag 1100  
tcgacctgca gaagcttggc cgccatggcc c 1131