

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
22 June 2006 (22.06.2006)

PCT

(10) International Publication Number  
WO 2006/063737 A1

(51) International Patent Classification:  
A61K 31/4184 (2006.01) A61P 9/12 (2006.01)  
A61K 31/5415 (2006.01)

Helmut E. [NL/DE]; Albrecht Dürer Strasse 19, 55218 Ingelheim am Rhein (DE). BÖHM, Peter [DE/DE]; Ledderhoser Weg 50, 55543 Bad Kreuznach (DE). RIEDEL, Axel [DE/DE]; Auhaldenstr. 8, 88437 Maselheim (DE).

(21) International Application Number:

PCT/EP2005/013224

(74) Agents: HAMMANN, Heinz et al.; Boehringer Ingelheim Pharma GMBH & CO. KG, Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(22) International Filing Date:

9 December 2005 (09.12.2005)

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MC, MD, MG, MK, ML, MN, MR, MW, MX, MZ, NA, NE, NG, NI, NL, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/637,062 17 December 2004 (17.12.2004) US

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

(71) Applicant (for AE, AG, AL, AM, AT, AU, AZ, BA, BB, BE, BF, BG, BJ, BR, BW, BY, BZ, CA, CF, CG, CH, CI, CM, CN, CO, CR, CU, CY, CZ, DK, DM, DZ, EC, EE, EG, ES, FI, FR, GA, GB, GD, GE, GH, GM, GN, GQ, GR, GW, HR, HU, ID, IE, IL, IN, IS, IT, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MC, MD, MG, MK, ML, MN, MR, MW, MX, MZ, NA, NE, NG, NI, NL, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD only): BOEHRINGER INGELHEIM INTERNATIONAL GMBH [DE/DE]; Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(71) Applicant (for DE only): BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG [DE/DE]; Binger Strasse 173, 55216 Ingelheim am Rhein (DE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHUMACHER,

Published:

— with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: COMBINATION THERAPY COMPRISING TELMISARTAN AND HYDROCHLOROTHIAZIDE


(57) Abstract: A pharmaceutical composition comprising about 80 mg telmisartan and about 25 mg hydrochlorothiazide or about 160 mg telmisartan and about 50 mg hydrochlorothiazide for the treatment of hypertension in patients with an insufficient blood pressure reduction upon treatment either with an angiotensin II receptor antagonist, or a pharmaceutical composition of an angiotensin II receptor antagonist and a low dose of hydrochlorothiazide.

## COMBINATION THERAPY COMPRISING TELMISARTAN AND HYDROCHLOROTHIAZIDE

The present invention relates to a pharmaceutical composition comprising as active ingredients about 80 mg of the angiotensin II receptor antagonist (ARB) telmisartan and about 25 mg of the diuretic hydrochlorothiazide (HCTZ) and a pharmaceutical composition which comprises about 160 mg telmisartan and about 50 mg hydrochlorothiazide and can be split into halves. The composition is used to treat hypertension in patients with an insufficient blood pressure reduction upon treatment either with an angiotensin II receptor antagonist, due to low plasma levels of renin or with a pharmaceutical composition of an angiotensin II receptor antagonist and lower doses of hydrochlorothiazide.

### Background of the invention

Telmisartan, a white to slightly yellowish solid, is an angiotensin II receptor antagonist developed for the treatment of hypertension and other medical indications as disclosed in EP-A-502314. It is a nonpeptide molecule chemically described as 4'-[{(1,4'-dimethyl-2'-propyl[2,6'-bi-1Hbenzimidazol]-1'-yl)methyl]-[1,1'-biphenyl]-2-carboxylic acid or 4'-[2-n-propyl-4-methyl-6-(1-methylbenzimidazol-2-yl)-benzimidazol-1-ylmethyl]-biphenyl-2-carboxylic acid. Its empirical formula is  $C_{33}H_{30}N_4O_2$ , its molecular weight is 514.63, and its structural formula is:



Telmisartan is manufactured and supplied in the free acid form. It is characterized by its very poor solubility in aqueous systems at the physiological pH range of the


- 2 -

gastro-intestinal tract of between pH 1 to 7. As disclosed in WO 00/43370, crystalline telmisartan exists in two polymorphic forms having different melting points. Under the influence of heat and humidity, the lower melting polymorph B transforms irreversibly into the higher melting polymorph A.

5

Hydrochlorothiazide (HCTZ), a white, odorless, crystalline powder with a molecular weight of 297.74, is a diuretic used in the treatment of edema and hypertension. It is chemically described as 6-chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide-1,1-dioxide. Its empirical formula is  $C_7H_8ClN_3O_4S_2$ , and its structural

10 formula is:



15 Object of the invention

There exists a striking relationship between high blood pressure and cardiovascular morbidity and mortality, i.e. an increased risk of myocardial infarction, heart failure, stroke, or kidney disease in case of increased blood pressure. An incremental increase of 20 mmHg in systolic or 10 mmHg in diastolic blood pressure in individuals

20 between 40 and 70 years of age doubles the risk of cardiovascular diseases.

Therefore a target blood pressure of <140/90 mmHg is recommended and an even lower target of <130/80 mmHg for patients with co-morbidities such as diabetes or chronic kidney disease.

25 Many patients require two or more antihypertensive drugs to achieve this goal. One of the possible combination partners are thiazide-diuretics which are able to facilitate salt and water excretion. For a combination of an angiotensin-II-receptor antagonist (ARB) with HCTZ a synergistic BP-lowering effect has been reported while additional

- 3 -

ARB treatment results in almost no additional side effects. Available combination products combine the ARB with either a low dose of HCTZ or a high dose of HCTZ, wherein low dose means less than 15 mg, preferably 12.5 mg of HCTZ and high dose means more than 15 mg, preferably 25 mg of HCTZ. Unfortunately a subgroup 5 of hypertensive patients does still not adequately respond to treatment with an ARB or a combination therapy of an ARB plus a low dose diuretic meaning that not in all patients the target blood pressure levels are achieved as suggested by the most recent guidelines, especially for patients with co-morbidities. In patients affected low 10 plasma renin activity (PRA) is frequently observed. Renin is an enzyme released by the kidney to help control the body's sodium-potassium balance, fluid volume, and blood pressure. Renin itself is not actually measured in the PRA test, because it is difficult to measure renin in routine lab assay. In the most commonly used renin assay, the test actually determines, by a procedure called radioimmunoassay, the rate of angiotensin I generation per unit time, while the plasma renin concentration 15 (PRC) measures the maximum renin effect. Both the PRA and the PRC are difficult to measure. Not only is renin itself unstable, but the patient's body position and the time of day affect the results. Also, the sample must be collected properly: drawn into a chilled syringe and collection tube, placed on ice, and sent to the performing laboratory immediately. Even if all these procedures are followed, results can vary 20 significantly.

The current invention is based on the surprising finding, that administering a daily dose of 80 mg of telmisartan in combination with 25 mg instead of 12.5 mg of hydrochlorothiazide, results in an unexpected strong increase of the responder rate 25 compared to patients treated with telmisartan or another ARB such as candesartan cilexetil, eprosartan, irbesartan, losartan, olmesartan, pratosartan, ripisartan, telmisartan, valsartan or zolasartan or a combination of those, including telmisartan, with a low dose of the diuretic HCTZ. Therefore it is the object of the present invention to provide a further treatment option for patients whose blood pressure has 30 not been adequately controlled with an ARB or a combination of an ARB with the low dose of the diuretic HCTZ. Said option comprises the manufacture of a pharmaceutical composition comprising 80 mg telmisartan and 25 mg

- 4 -

hydrochlorothiazide or a pharmaceutical composition which comprises about 160 mg telmisartan and about 50 mg hydrochlorothiazide and can be split into halves.

#### Definitions

5 As used herein, the term "substantially amorphous" refers to a product comprising amorphous constituents in a proportion of at least 90%, preferably at least 95%, as determined by X-ray powder diffraction measurement.

The term "dissolving tablet matrix" refers to a pharmaceutical tablet base formulation having instant release (fast dissolution) characteristics that readily dissolves in a 10 physiological aqueous medium.

The term "disintegrating tablet matrix" refers to a pharmaceutical tablet base formulation having instant release characteristics that readily disintegrates in a physiological aqueous medium.

15 Description of the invention

The present invention comprises a pharmaceutical composition which can be used in a method for the treatment of hypertension comprising as active ingredients about 80 mg of the angiotensin II receptor antagonist telmisartan and about 25 mg of the diuretic hydrochlorothiazide and another pharmaceutical composition for the 20 treatment of hypertension, which can be split into halves, comprising as active ingredients about 160 mg of telmisartan and about 50 mg hydrochlorothiazide.

The active ingredient telmisartan is generally supplied in its free acid form, although pharmaceutically acceptable salts such as the sodium salt may also be used. Since 25 during subsequent processing telmisartan is normally dissolved and transformed into a substantially amorphous form, its initial crystal morphology and particle size are frequently of little importance for the physical and biopharmaceutical properties of the pharmaceutical formulation. It is, however, preferred to remove agglomerates from the starting material, e.g. by sieving, in order to facilitate wetting and dissolution 30 during further processing.

- 5 -

The diuretic is usually used as a fine-crystalline powder, optionally in fine-milled, peg-milled or micronized form. For instance, the particle size distribution of hydrochlorothiazide, as determined by the method of laser light scattering in a dry dispersion system (Sympatec Helos/Rodos, focal length 100 mm) is preferably as follows:

- 5  $d_{10}$  :  $\leq 20 \mu\text{m}$ , preferably 2 to 10  $\mu\text{m}$
- 10  $d_{50}$  : 5 to 50  $\mu\text{m}$ , preferably 10 to 30  $\mu\text{m}$
- 15  $d_{90}$  : 20 to 100  $\mu\text{m}$ , preferably 40 to 80  $\mu\text{m}$

Preferred embodiments of the above pharmaceutical composition are tablets or capsules. Particularly preferred are bilayer tablets consisting of a first tablet layer comprising telmisartan in a dissolving tablet matrix having instant release (fast dissolution) characteristics and a separate second tablet layer comprising the active ingredient HCTZ in a disintegrating tablet matrix. The dissolving tablet matrix may have neutral or basic properties, although a basic tablet matrix is preferred.

Conveniently the composition of the present invention comprises telmisartan in substantially amorphous form which may be produced by any suitable method known to those skilled in the art, for instance, by freeze drying of aqueous solutions, coating of carrier particles in a fluidized bed, and solvent deposition on sugar pellets or other carriers. Preferably, however, the substantially amorphous telmisartan is prepared by the specific spray-drying method described in WO03/059327.

- 20 25 Additionally, the composition of the present invention preferably comprises as inactive ingredients sodium hydroxide, meglumine, povidone, sorbitol, magnesium stearate, lactose monohydrate, microcrystalline cellulose, maize starch and sodium starch glycolate.
- 30 A dissolving matrix of a telmisartan tablet layer may comprise a basic agent, a water-soluble diluent and, optionally, other excipients and adjuvants.

- 6 -

Specific examples of suitable basic agents are alkali metal hydroxides such as NaOH and KOH; basic amino acids such as arginine and lysine; and meglumine (N-methyl-D-glucamine), NaOH and meglumine being preferred.

- 5 Specific examples of suitable water-soluble diluents are carbohydrates such as monosaccharides like glucose; oligosaccharides like sucrose, anhydrous lactose and lactose monohydrate; and sugar alcohols like sorbitol, mannitol, erythrol and xylitol. Sorbitol is a preferred diluent.
- 10 The other excipients and/or adjuvants are, for instance, selected from binders, carriers, fillers, lubricants, flow control agents, crystallization retarders, solubilizers, coloring agents, pH control agents, surfactants and emulsifiers, specific examples of which are given below in connection with the second tablet layer composition. The excipients and/or adjuvants for a telmisartan tablet layer composition are preferably chosen such that a non-acidic, fast dissolving tablet matrix is obtained.
- 15

Such a first tablet layer composition generally comprises 3 to 50 wt.%, preferably 5 to 35 wt.%, of active ingredient; 0.25 to 20 wt.%, preferably 0.40 to 15 wt.%, of basic agent; and 30 to 95 wt.%, preferably 60 to 80 wt.% of water-soluble diluent (filler).

- 20 Other (optional) constituents may, for instance, be chosen from one or more of the following excipients and/or adjuvants in the amounts indicated:

- 10 to 30 wt.%, preferably 15 to 25 wt.%, of binders, carriers and fillers, thereby replacing the water-soluble diluent;
- 25 0.1 to 5 wt.%, preferably 0.5 to 3 wt.%, of lubricants;  
0.1 to 5 wt.%, preferably 0.3 to 2 wt.%, of flow control agents;  
1 to 10 wt.%, preferably 2 to 8 wt.%, of crystallization retarders;  
1 to 10 wt.%, preferably 2 to 8 wt.%, of solubilizers;  
0.05 to 1.5 wt.%, preferably 0.1 to 0.8 wt.%, of coloring agents;
- 30 0.5 to 10 wt.%, preferably 2 to 8 wt.%, of pH control agents;  
0.01 to 5 wt.%, preferably 0.05 to 1 wt.%, of surfactants and emulsifiers.

Such a telmisartan tablet layer may be produced by spray-drying an aqueous solution comprising telmisartan and a basic agent to obtain a spray-dried granulate, mixing said spray-dried granulate with a water-soluble diluent to obtain a premix, mixing said premix with a lubricant to obtain a final blend and compressing the final

5 blend to form the first tablet layer.

A separate second tablet layer comprising HCTZ in a fast disintegrating tablet matrix preferably comprises one or more fillers, a binder or polymer, a disintegrant, a lubricant and, optionally, other excipients and adjuvants.

10 Preferred fillers are selected from the group consisting of pregelatinized starch, microcristalline cellulose, low-substituted hydroxypropylcellulose, cellulose, mannitol, erythritol, lactose, saccharose, calciumhydrogenphosphate, sorbitol, and xylitol. Particularly preferred are pregelatinized starch, microcristalline cellulose, mannitol and lactose monohydrate. Particularly preferred are anhydrous lactose, spray-dried

15 lactose and lactose monohydrate.

Preferred disintegrants are selected from the group consisting of croscarmellose sodium salt (cellulose carboxymethylether sodium salt, crosslinked), sodium starch glycolate, crosslinked polyvinylpyrrolidone (crospovidone), corn starch and low-substituted hydroxypropylcellulose. Particularly preferred are sodium starch glycolate and

20 croscarmellose sodium salt.

Preferred binders are selected from the group consisting of polyvinyl pyrrolidone (Povidone), copolymers of vinylpyrrolidone with other vinylderivatives (Copovidone), hydroxypropylmethylcellulose, methylcellulose, hydroxypropyl-cellulose and low-substituted hydroxypropyl-cellulose. Particularly preferred are hydroxypropyl-

25 methylcellulose and Copovidone.

Preferred lubricants are sodium stearyl fumarate and magnesium stearate.

The other excipients and adjuvants, if used, are preferably selected from

- diluents and carriers such as cellulose powder, microcrystalline cellulose, cellulose

30 derivatives like hydroxymethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose and hydroxy-propylmethylcellulose, dibasic calcium phosphate, corn starch, pregelatinized starch, polyvinyl pyrrolidone (Povidone) etc.;

- 8 -

- lubricants such as stearic acid, magnesium stearate, sodium stearyl fumarate, glycerol tribehenate, etc.;
- flow control agents such as colloidal silica, talc, etc.;
- crystallization retarders such as Povidone, etc.;

5 - solubilizers such as Pluronic, Povidone, etc.;

- coloring agents, including dyes and pigments such as Iron Oxide Red or Yellow, titanium dioxide, talc, etc.;
- pH control agents such as citric acid, tartaric acid, fumaric acid, sodium citrate, dibasic calcium phosphate, dibasic sodium phosphate, etc.;

10 - surfactants and emulsifiers such as Pluronic, polyethylene glycols, sodium carboxymethyl cellulose, polyethoxylated and hydrogenated castor oil, etc.;

- antioxidants;

and mixtures of two or more of these excipients and/or adjuvants.

15 The layers can be differentiated by using different colors.

A separate second tablet layer comprising HCTZ generally comprises 1.5 to 35 wt.%, preferably 2 to 15 wt.%, of active ingredient; 25 to 75 wt.%, preferably 35 to 65 wt.%, of filler; 10 to 40 wt.%, preferably 15 to 35 wt.%, of dry binder; 0.5 to 5 wt.%, 20 preferably 1 to 4 wt.%, of wet granulation binder; and 1 to 10 wt.%, preferably 2 to 8 wt.%, of disintegrant. The other excipients and adjuvants are generally employed in the same amount as in the first tablet layer composition.

25 For preparing a bilayer tablet according to the present invention, the first and second tablet layer compositions may be compressed in the usual manner in a bilayer tablet press, e.g. a high-speed rotary press in a bilayer tableting mode. However, care should be taken not to employ an excessive compression force for the first tablet layer. Preferably, the ratio of the compression force applied during compression of the first tablet layer to the compression force applied during compression of both the 30 first and second tablet layers is in the range of from 1:10 to 1:2. For instance, the first tablet layer may be compressed at moderate force of 4 to 8 kN, whereas the main compression of first plus second layer is performed at a force of 10 to 20 kN.

- 9 -

During bilayer tablet compression adequate bond formation between the two layers is achieved by virtue of distance attraction forces (intermolecular forces) and mechanical interlocking between the particles.

5 The tablets obtained release the active ingredients rapidly and in a largely pH-independent fashion, with complete release occurring within less than 60 min and release of the major fraction occurring within less than 15 min. The dissolution/disintegration kinetics of the multilayer tablet may be controlled in different ways. For instance, the layers may dissolve/disintegrate simultaneously. Preferably, the tablet

10 layer comprising HCTZ disintegrates first whereas the layer comprising telmisartan dissolves subsequently.

In accordance with the present invention, at least 70% and typically at least 90% of the active ingredients are dissolved after 30 min.

15 Bilayer tablets according to the present invention tend to be slightly hygroscopic and are therefore preferably packaged using a moisture-proof packaging material such as aluminium foil blister packs, or polypropylene tubes and HDPE bottles which preferably contain a desiccant.

20 A preferred method of producing the bilayer tablet according to the present invention comprises

(i) providing a first tablet layer composition by

25 a) preparing an aqueous solution of telmisartan, at least one basic agent and, optionally, a solubilizer and/or a crystallization retarder;

b) spray-drying said aqueous solution to obtain a spray-dried granulate;

c) mixing said spray-dried granulate with a water-soluble diluent to obtain a premix;

d) mixing said premix with a lubricant to obtain a final blend for the first layer;

30 e) optionally, adding other excipients and/or adjuvants in any of steps a) to d);

(ii) providing a second tablet comprising HCTZ;

- 10 -

- (iii) compressing both the first and the second tablet layer composition to form a tablet layer; and
- (v) compressing the separate tablet layers to form a biilayer tablet.

5 To provide a first tablet layer composition an aqueous alkaline solution of telmisartan is prepared by dissolving the active ingredient in purified water with the help of one or more basic agents like sodium hydroxide and meglumine. Optionally, a solubilizer and/or a recrystallization retarder may be added. The dry matter content of the starting aqueous solution is generally 10 to 40 wt.%, preferably 20 to 30 wt.%.

10 The aqueous solution is then spray-dried at room temperature or preferably at increased temperatures of, for instance, between 50 and 100°C in a co-current or countercurrent spray-drier at a spray pressure of, for instance, 1 to 4 bar. Generally speaking, the spray-drying conditions are preferably chosen in such a manner that a spray-dried granulate having a residual humidity of  $\leq$  5 wt.%, preferably  $\leq$  3.5 wt.%, is obtained in the separation cyclone. To that end, the outlet air temperature of the spray-drier is preferably kept at a value of between about 80 and 90°C while the other process parameters such as spray pressure, spraying rate, inlet air temperature, etc. are adjusted accordingly.

15 The spray-dried granulate obtained is preferably a fine powder having the following particle size distribution:

20  $d_{10}$  :  $\leq$  20  $\mu\text{m}$ , preferably  $\leq$  10  $\mu\text{m}$   
 $d_{50}$  :  $\leq$  80  $\mu\text{m}$ , preferably 20 to 55  $\mu\text{m}$   
 $d_{90}$  :  $\leq$  350  $\mu\text{m}$ , preferably 50 to 150  $\mu\text{m}$

25 After spray-drying, the active ingredient telmisartan as well as the excipients contained in the spray-dried granulate are in a substantially amorphous state with no crystallinity being detectable. From a physical point of view, the spray-dried granulate is a solidified solution or glass having a glass transition temperature  $T_g$  of preferably

30  $> 50^\circ\text{C}$ , more preferably  $> 80^\circ\text{C}$ .

Based on 100 parts by weight of active ingredient telmisartan, the spray-dried granulate preferably contains 5 to 200 parts by weight of basic agent and, optionally, solubilizer and/or crystallization retarder.

The water-soluble diluent is generally employed in an amount of 30 to 95 wt.%,

5 preferably 60 to 80 wt.%, based on the weight of the first tablet layer composition.

The lubricant is generally added to the premix in an amount of 0.1 to 5 wt.%, preferably 0.3 to 2 wt.%, based on the weight of the first tablet layer composition.

Mixing is carried out in two stages, i.e. in a first mixing step the spray-dried granulate and the diluent are admixed using, e.g., a high-shear mixer or a free-fall blender,

10 and in a second mixing step the lubricant is blended with the premix, preferably also under conditions of high shear. The method of the invention is however not limited to these mixing procedures and, generally, alternative mixing procedures may be employed in steps c), d), and also in the subsequent steps f) and g), such as, e.g., container mixing with intermediate screening.

15

To provide a second tablet layer composition comprising HCTZ the constituent components may be prepared by dry-mixing, e.g. by means of a high-intensity mixer or a free-fall blender. Alternatively and preferably, the second tablet layer composition is prepared using a wet granulation technique wherein an aqueous

20 solution of a wet granulation binder is added to a premix and subsequently the wet granulate obtained is dried, e.g. in a fluidized-bed dryer or drying chamber. The dried mixture is screened and then a lubricant is admixed, e.g. using a tumbling mixer or free-fall blender.

25 First and second tablet layer compositions as described above can be compressed into bilayer tablets of the target tablet weight with appropriate size and crushing strength, using an appropriate tablet press, e.g. a rotary press in the bilayer tableting mode. Optional an appropriate external lubricant spray system for dies and punches can be used during manufacturing of tablets in order to improve lubrication. In order 30 to avoid any cross-contamination between the tablet layers (which could lead to decomposition of HCTZ), any granulate residues should be carefully removed during tableting by intense suction of the die table within the tableting chamber.

- 12 -

In addition to the treatment of hypertension the composition according to the present invention can also be used to treat or prevent a condition selected from the group consisting of stroke, myocardial infarction, transient ischaemic attack, congestive

5 heart failure, cardiovascular disease, insulin resistance, impaired glucose tolerance, pre-diabetes, type 2 diabetes mellitus, metabolic syndrome (syndrome X), obesity, hypertriglyceridemia, elevated serum concentrations of C-reactive protein, elevated serum concentrations of lipoprotein(a), elevated serum concentration of homocysteine, elevated serum concentration of low-density lipoprotein (LDL)-

10 cholesterol, elevated serum concentration of lipoprotein-associated phospholipase (A2), reduced serum concentration of high density lipoprotein (HDL)-cholesterol, reduced serum concentration of HDL(2b)-cholesterol, reduced serum concentration of adiponectin, cognitive decline and dementia.

15 In order to further illustrate the present invention, the following non-limiting examples are given.

## Examples

### Example 1: Surprising responder rate using 80mg telmisartan + 25 mg HCTZ

5 An important goal of up-titrating antihypertensive medication is to increase the number of patients adequately responding to treatment. Surprisingly improved responder rates have become obvious when the current clinical database of telmisartan was analysed, in particular with respect to compare the responder rates upon treatment with telmisartan 80 mg /HCTZ 25 mg to those with telmisartan 80 mg

10 /HCTZ 12.5 mg. Responders are defined as having DBP <90 mm Hg or a reduction of at least 10 mm Hg. Regarding SBP an adequate response is defined as SBP <140 or a reduction of at least 10 mm Hg. When comparing telmisartan 80 mg/HCTZ 25 mg with telmisartan 80 mg/HCTZ 12.5 mg and applying the above mentioned definitions the diastolic (DBP) response rates increased by 6.6 % for patients in

15 controlled studies and by 15.4 % for patients from follow-up studies. Systolic (SBP) response rates improved by 7.8% for patients from controlled and from follow-up studies.

The tables below show the detailed blood pressure response data for telmisartan

20 80 mg/HCTZ 25 mg in comparison to telmisartan 80 /HCTZ 12.5 mg from the project database:

- 14 -

**TABLE 1: Diastolic Responder Rates**

<b>Diastolic Blood Pressure</b>			
<b>Patients from Controlled Studies</b>			
<b>Treatment</b>	<b>N</b>	<b>Rate of Non-Responder</b>	<b>Rate of Responder</b>
T80/H12.5	528	35.0%	65.0%
T80/H25	134	28.4%	71.6%
<b>Patients from Follow-up Studies</b>			
<b>Treatment</b>	<b>N</b>	<b>Rate of Non-Responder</b>	<b>Rate of Responder</b>
T80/H12.5	799	41.9%	58.1%
T80/H25	442	26.5%	73.5%

**TABLE 2: Systolic Responder Rates**

5

<b>Systolic Blood Pressure</b>			
<b>Patients from Controlled Studies</b>			
<b>Treatment</b>	<b>N</b>	<b>Rate of Non-Responder</b>	<b>Rate of Responder</b>
T80/H12.5	528	24.2%	75.8%
T80/H25	134	16.4%	83.6%
<b>Patients from Follow-up Studies</b>			
<b>Treatment</b>	<b>N</b>	<b>Rate of Non-Responder</b>	<b>Rate of Responder</b>
T80/H12.5	799	27.3%	72.7%
T80/H25	442	19.5%	80.5%

- 15 -

When comparing telmisartan 80 mg/HCTZ 25 mg with telmisartan 80 mg/HCTZ 12.5 mg and applying the more recent response criteria for systolic blood pressure of SBP <140 or a reduction of at least 20 mm Hg, responder rate increase by 5.7 % (from 61.5 % to 67.2%) for patients from controlled studies and 9.4 % (from 56.2 % to, 5 65.6%) for patients from follow-up studies.

The analysis of existing evidence from the clinical database of telmisartan suggests that the telmisartan 80 mg/HCTZ 25 mg combination does consistently provide higher clinical efficacy in terms of blood pressure lowering and, especially, responder 10 rates. Despite an expected increase in responder rates due to the increased dose of HCTZ the extent of the observed effect for the combination of telmisartan 80 mg and HCTZ 25 mg goes far beyond what was to be expected.

15 **Example 2: Composition of a 680 mg total weight bilayer tablet comprising 80mg telmisartan and 25 mg HCTZ**

**Telmisartan Layer**

Ingredient	mg per tablet
Telmisartan	80.000
Sodium hydroxide	6.720
Polyvidone K25	24.000
Meglumine	24.000
Purified water	(400.000)
Sorbitol	337.280
Magnesium stearate	8.000
<b>Total Telmisartan Layer</b>	<b>480.000</b>

### Hydrochlorothiazide Layer

Ingredient	mg per tablet
Hydrochlorothiazide	25.000
Lactose monohydrate	99.100
Microcrystalline cellulose	64.000
Maize starch	6.000
Ferric oxide	0.900
Sodium starch glycollate	4.000
Purified water	(64.000)
Magnesium stearate	1.000
<b>Total Hydrochlorothiazide Layer</b>	<b>200.000</b>

Example 3: Confirmation of the surprising high responder rate using 80mg

5 **telmisartan + 25 mg HCTZ in a separate clinical trial**

To confirm the surprising responder rate of 80mg telmisartan + 25 mg HCTZ found upon analyzing the available clinical telmisartan database, the corresponding responder rate was determined for a clinical trial actually designed to compare the safety and efficacy of a 80 mg 10 telmisartan + 25 mg hydrochlorothiazide combination with the currently available 160 mg valsartan + 25 mg hydrochlorothiazide combination in patients with Stage 1 and Stage 2 hypertension (confirmation study). This study was a randomized, double-blind, double-dummy, placebo-controlled, forced-titration trial, with a total duration of up to 12 weeks (eight weeks active treatment). The target population included both male and female hypertensive 15 patients, at least 18 years of age.

The primary objective of this study was to show that the combination of MICARDIS® HCT (telmisartan 80 mg / hydrochlorothiazide 25 mg) is superior to placebo in lowering DBP and SBP, at least as effective as DIOVAN® HCT (valsartan 160 mg / hydrochlorothiazide 25 mg) in lowering DBP, and possibly superior to DIOVAN® HCT in lowering DBP and SBP in 20 patients with Stage 1 and Stage 2 hypertension as measured by seated trough cuff blood pressure.

The primary endpoints were the change from baseline (Visit 2) for in-clinic mean seated trough cuff diastolic (DBP) and systolic blood pressure (SBP) at the end of an eight week

- 17 -

(Visit 6) treatment period (i.e., two weeks treatment with MICARDIS® 80 mg or DIOVAN® 160 followed by six weeks treatment MICARDIS® HCT 80/25 mg or DIOVAN® HCT 160/25 mg respectively, or placebo for the entire eight weeks).

5 Measurements of blood pressure were made at trough (i.e., within 23 - 26 hours after most recent intake of the study medication).

Secondary efficacy endpoints in this study, as measured by in-clinic trough cuff blood pressure at the end of an eight week treatment period included:

10 1.) The percentage of patients responding to treatment based on mean seated trough cuff measurements defined as:  
DBP Control: Mean seated DBP <90 mmHg at trough  
DBP Response: Mean seated DBP <90 mmHg at trough and/or a change from baseline of  $\geq 10$  mmHg

15 SBP Response: Mean seated SBP <140 mmHg at trough and/or a change from baseline of  $\geq 10$  mmHg  
Normal BP: Mean seated SBP <130 mmHg at trough and mean seated DBP <85 mmHg at trough  
High Normal: Mean seated SBP  $\geq 130$  and <140 mmHg at trough and a mean seated DBP  $\geq 85$  and <90 mmHg at trough

20 2.) The percentage of patients with uncontrolled HTN defined as systolic BP  $\geq 180$  mmHg and/or diastolic BP  $\geq 120$  mmHg at the end of study.

25 Safety was evaluated through the review of adverse events and through the measurement of changes from the baseline in physical examinations, laboratory parameters and vital signs (mean SBP, mean DBP) and pulse rate. At any time during the study, patients with a mean in-clinic SBP  $\geq 180$  mmHg and/or DBP  $\geq 120$  mmHg were to be withdrawn from the study for safety reasons. The mean value was calculated from three successive in-clinic blood pressure measurements taken two minutes apart after resting quietly in the seated position

30 for five minutes.

Patient inclusion criteria were

35 1. Ability to provide written informed consent.

- 18 -

2. Age 18 years or older.
3. Ability to stop current antihypertensive therapy without unacceptable risk to the patient (investigator's discretion).
4. Seated cuff DBP of  $\geq 95$  mmHg at Visit 2 (baseline).

5

Patient exclusion criteria were

1. Pre-menopausal women (last menstruation  $\leq 1$  year prior to start of run-in period) who:
  - a. were not surgically sterile; and/or
  - b. were nursing or pregnant.
- 10 c. were of child-bearing potential and were NOT practicing acceptable means of birth control, did NOT plan to continue using this method throughout the study and did NOT agree to submit to periodic pregnancy testing during participation in studies of >three-months duration. Acceptable methods of birth control included oral, implantable or injectable contraceptives.
- 15 2. Known or suspected secondary hypertension.
3. Mean sitting SBP  $>180$  mmHg or mean sitting DBP  $>120$  mmHg at any time during the study.
4. Hepatic and/or renal dysfunction as defined by the following laboratory parameters:
  - a. SGPT (ALT) or SGOT (AST)  $>$ two times the upper limit of normal range, or
  - 20 b. Serum creatinine  $>3.0$  mg/dL or creatinine clearance  $< 0.6$  ml/sec.
5. Bilateral renal artery stenosis, renal artery stenosis in a solitary kidney, post-renal transplant or with only one kidney.
6. Clinically relevant hypokalaemia or hyperkalaemia.
7. Uncorrected volume depletion.
- 25 8. Uncorrected sodium depletion.
9. Primary aldosteronism.
10. Hereditary fructose intolerance.
11. Biliary obstructive disorders, cholestasis or moderate to severe hepatic insufficiency.
12. Patients who had previously experienced symptoms characteristic of angioedema during
- 30 treatment with ACE inhibitors or angiotensin II receptor antagonists.
13. History of drug or alcohol dependency within six months prior to start of run-in period.
14. Chronic administration of any medications known to affect blood pressure, except medication allowed by the protocol.
15. Any investigational drug therapy within one month of start of run-in period.

- 19 -

16. Known hypersensitivity to any component of the formulation study drugs (telmisartan, valsartan or hydrochlorothiazide).
17. Contra-indication to a placebo run-in period (e.g. stroke within the past six months, MI, cardiac surgery, PTCA or angina within the past three months prior to start of run-in period).
- 5 18. Any other clinical condition which, in the opinion of the principal investigator, did not allow safe completion of the protocol and safe administration of telmisartan, valsartan or hydrochlorothiazide.
19. Night shift workers.
- 10 20. Clinically significant ventricular tachycardia, atrial fibrillation, atrial flutter or other clinically relevant cardiac arrhythmias as determined by the investigator.
21. NYHA functional class CHF III-IV.
22. Hypertrophic obstructive cardiomyopathy, aortic stenosis, hemodynamically relevant stenosis of aortic or mitral valve.
- 15 23. Patients whose diabetes was unstable and uncontrolled for at least the past three months as defined by a HbA1C  $\geq 10\%$ .
24. Concomitant use of lithium or cholestyramine or colestipol resins (potential drug interactions with hydrochlorothiazide).
25. History of non-compliance with prescribed medication or protocol procedures.

20 The analysis of the data of this study reveals, that the responder rate of 80mg telmisartan + 25mg HCTZ (T80/H25) are even higher than after the analysis of the clinical telmisartan database. Additionally, this responder rate is higher than the responder rate of 160mg valsartan + 25mg HCTZ (Val160/H25), the difference, however, cannot be interpreted as 25 statistically significant.

- 20 -

The detailed responder rate values for 80mg telmisartan + 25 mg HCTZ (T80/H25) and 160mg valsartan + 25 mg HCTZ (Val160/H25) are:

	T80/H25 (confirmation study)	T80/H25 (controlled studies of example 1)	T80/H25 (follow-up studies of example 1)	Val160/H25 (confirmation study)
DBP Response	82.4 %	71.6 %	73.5 %	78.7 %
SBP Response	87.6 %	83.6 %	80.5 %	84.8 %
New EMEA SBP Response Criteria	75.2 % *	67.2 %	65.6 %	68.7 % **

\* Overall value of the responder rates for non-black (76.6 %) and black patients (70.6 %)

\*\* Overall value of the responder rates for non-black (71.6 %) and black patients (59.5 %)

**Claims**

1. A pharmaceutical composition for the treatment of hypertension comprising as active ingredients about 80 mg of the angiotensin II receptor antagonist 5 telmisartan and about 25 mg of the diuretic hydrochlorothiazide.
2. A pharmaceutical composition for the treatment of hypertension comprising as active ingredients about 160 mg of the angiotensin II receptor antagonist telmisartan and about 50 mg of the diuretic hydrochlorothiazide which can be 10 split into halves.
3. The composition of claim 1 or 2, wherein the pharmaceutical composition is a tablet or a capsule.
- 15 4. The composition of claim 3, wherein the pharmaceutical composition contains the component telmisartan in a dissolving tablet matrix having instant release characteristics.
5. The composition of claim 3, wherein the diuretic hydrochlorothiazide forms a 20 separate layer in a disintegrating pharmaceutical matrix.
6. The composition of claim 3, wherein telmisartan or a salt thereof is in a substantially amorphous form.
- 25 7. The composition of claim 3 comprising as inactive ingredients sodium hydroxide, meglumine, povidone, sorbitol, magnesium stearate, lactose monohydrate, microcristalline cellulose, maize starch and sodium starch glycolate.
- 30 8. The composition of claim 4, wherein the dissolving tablet matrix comprises a basic agent, a water-soluble diluent and, optionally, other excipients and adjuvants.

- 22 -

9. The composition of claim 8, wherein the basic agent is selected from alkali metal hydroxides, basic amino acids and meglumine.
- 5 10. The composition of claim 8, wherein the water-soluble diluent is selected from monosaccharides like glucose; oligosaccharides like sucrose and lactose; and sugar alcohols like sorbitol, mannitol, erythrol and xylitol.
- 10 11. The composition of claim 8, wherein the other excipients and adjuvants are selected from binders, carriers, fillers, lubricants, flow control agents, crystallization retarders, solubilizers, coloring agents, pH control agents, surfactants and emulsifiers.
- 15 12. The composition of claim 8, wherein the tablet matrix is produced by spray-drying an aqueous solution comprising telmisartan and a basic agent to obtain a spray-dried granulate, mixing said spray-dried granulate with a water-soluble diluent to obtain a premix, mixing said premix with a lubricant to obtain a final blend and compressing the final blend to form the first tablet layer.
- 20 13. The composition of claim 5, wherein the disintegrating tablet matrix comprises a filler, a binder, a disintegrant and, optionally, other excipients and adjuvants.
14. The composition of claim 13, wherein the other excipients and adjuvants are selected from carriers, diluents, lubricants, flow control agents, solubilizers, 25 antioxidants, coloring agents, pH control agents, surfactants and emulsifiers.
15. The composition of claim 1, wherein the pharmaceutical composition is packaged in a moisture proof packaging material such as aluminium foil blister packs or polypropylene tubes and HDPE bottles.

- 23 -

16. A method of treating hypertension in a patient comprising the administration of a pharmaceutical composition combining 25 mg of the diuretic hydrochlorothiazide with 80 mg of the angiotensin II receptor antagonist telmisartan.
- 5 17. The method of claim 16, wherein the patient's blood pressure is not adequately controlled by therapy with an angiotensin II receptor antagonist or by combination of said angiotensin II receptor antagonist and a lower dose of HCTZ.
- 10 18. The method of claim 17, wherein the patient has a low plasma renin activity or plasma renin concentration.
19. The method of claim 16, additionally treating or preventing a condition selected from the group consisting of stroke, myocardial infarction, transient ischaemic attack, congestive heart failure, cardiovascular disease, insulin resistance, impaired glucose tolerance, pre-diabetes, type 2 diabetes mellitus, metabolic syndrome (syndrome X), obesity, hypertriglyceridemia, elevated serum concentrations of C-reactive protein, elevated serum concentrations of lipoprotein(a), elevated serum concentration of homocysteine, elevated serum concentration of low-density lipoprotein (LDL)-cholesterol, elevated serum concentration of lipoprotein-associated phospholipase (A2), reduced serum concentration of high density lipoprotein (HDL)-cholesterol, reduced serum concentration of HDL(2b)-cholesterol, reduced serum concentration of adiponectin, cognitive decline and dementia.
- 20 25 20. A method for the manufacture of a pharmaceutical unit dosage form comprising the angiotensin II receptor antagonist telmisartan and the diuretic hydrochlorothiazide to treat hypertension in a patient whose blood pressure is not adequately controlled by therapy with an angiotensin II receptor antagonist or by combination of said angiotensin II receptor antagonist and a low dose of the diuretic HCTZ, wherein the weight ratio of telmisartan and

- 24 -

hydrochlorothiazide allows to treat the patient with a daily dose of about 80 mg telmisartan and about 25 mg hydrochlorothiazide.

21. The method of claim 20, wherein the dosage form comprises about 80 mg  
5 telmisartan and about 25 mg hydrochlorothiazide.
22. The method of claim 20, wherein the dosage form comprises about 40 mg  
telmisartan and about 12.5 mg hydrochlorothiazide.
- 10 23. The method of claim 20, wherein the dosage form comprises about 160 mg  
telmisartan and about 50 mg hydrochlorothiazide and can be split into halves.
24. Use of telmisartan and HCTZ for the manufacture of the pharmaceutical  
composition of claim 1 or 2.

15

# INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2005/013224

A. CLASSIFICATION OF SUBJECT MATTER A61K31/4184 A61K31/5415 A61P9/12		
According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols) A61K A61P		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
Electronic data base consulted during the international search (name of data base and, where practical, search terms used) EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, EMBASE, CHEM ABS Data		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 03/059327 A (BOEHRINGER INGELHEIM PHARMA GMBH & CO. KG; FRIEDL, THOMAS; SCHEPKY, GO) 24 July 2003 (2003-07-24) page 5, line 5 – page 7, line 9; page 9, line 1-23 ----- -/-	1-24
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C.		<input checked="" type="checkbox"/> See patent family annex.
* Special categories of cited documents :		
*A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier document but published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed		
*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art. *&* document member of the same patent family		
Date of the actual completion of the international search  30 January 2006		Date of mailing of the international search report  28/02/2006
Name and mailing address of the ISA/ European Patent Office, P.B. 5818 Patentlaan 2 NL – 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl, Fax: (+31-70) 340-3016		Authorized officer  Borst, M

## INTERNATIONAL SEARCH REPORT

International application No PCT/EP2005/013224
---

## C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	ANONYMOUS: "Micardis HCT (telmisartan and hydrochlorothiazide) Tablets, 40mg/12.5 mg 80 mg/12.5 mg and 80 mg/25 mg" BOEHRINGER INGELHEIM, 'Online! 19 April 2004 (2004-04-19), XP002365096 Retrieved from the Internet: URL: <a href="http://www.fda.gov/medwatch/SAFETY/2004/apr_PI/MicardisHCT_PI.pdf">http://www.fda.gov/medwatch/SAFETY/2004/apr_PI/MicardisHCT_PI.pdf</a> > 'retrieved on 2006-01-30! paragraphs entitled "Description"; "Clinical Trials"; "Indications and Usage"; "Dosage and Administration"; "Dose Titration by Clinical Effect" -----	1-24
A	WO 94/05264 A (BOEHRINGER MANNHEIM GMBH) 17 March 1994 (1994-03-17) claims -----	2
X	FREYTAG FRANK ET AL: "Long-term exposure to telmisartan as monotherapy or combination therapy: Efficacy and safety" BLOOD PRESSURE, vol. 11, no. 3, 2002, pages 173-181, XP009060475 ISSN: 0803-7051 page 177, right hand column, last full paragraph -----	1-24
X	NEUTEL JOEL M ET AL: "Long-term efficacy and tolerability of telmisartan as monotherapy and in combination with other antihypertensive medications." BLOOD PRESSURE. 2002, vol. 11, no. 5, 2002, pages 302-309, XP009060476 ISSN: 0803-7051 table III on page 306 -----	1-24
A	RAM C VENTIKA S: "Antihypertensive efficacy of angiotensin receptor blockers in combination with hydrochlorothiazide: a review of the factorial-design studies." JOURNAL OF CLINICAL HYPERTENSION (GREENWICH, CONN.) OCT 2004, vol. 6, no. 10, October 2004 (2004-10), pages 569-577, XP009060495 ISSN: 1524-6175 paragraph bridging pages 571-572 ----- -/-	1-24

## INTERNATIONAL SEARCH REPORT

International application No  
PCT/EP2005/013224

## C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>WHITE WILLIAM B ET AL: "Effects of the angiotensin II receptor blockers telmisartan versus valsartan on the circadian variation of blood pressure Impact on the early morning period" AMERICAN JOURNAL OF HYPERTENSION, vol. 17, no. 4, April 2004 (2004-04), pages 347-353, XP002365097 ISSN: 0895-7061 page 352-353, paragraph entitled "Discussion"</p> <p>-----</p>	1-24
P, A	<p>WHITE W ET AL: "Telmisartan plus hydrochlorothiazide (80/25 mg) has a greater antihypertensive effect than valsartan plus hydrochlorothiazide (160/25 mg) in patients with stage 1 and 2 hypertension" AMERICAN JOURNAL OF HYPERTENSION, ELSEVIER, vol. 18, no. 5, May 2005 (2005-05), page A97, XP004895902 ISSN: 0895-7061 abstract</p> <p>-----</p>	1-24

## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/EP2005/013224

### Box II Observations where certain claims were found unsearchable (Continuation of item 2 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1.  Claims Nos.: - because they relate to subject matter not required to be searched by this Authority, namely:

Rule 39.1(iv) PCT : Although claims 16-19 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effects of the compound/composition.

2.  Claims Nos.: because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:

3.  Claims Nos.: because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

### Box III Observations where unity of invention is lacking (Continuation of item 3 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1.  As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2.  As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3.  As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4.  No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

#### Remark on Protest

The additional search fees were accompanied by the applicant's protest.

No protest accompanied the payment of additional search fees.

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International application No

PCT/EP2005/013224

Patent document cited in search report		Publication date		Patent family member(s)		Publication date
WO 03059327	A	24-07-2003	AU	2002242676 A1		30-07-2003
			BR	0215514 A		21-12-2004
			CA	2472392 A1		24-07-2003
			CZ	20040939 A3		16-02-2005
			EP	1467712 A1		20-10-2004
			JP	2005514439 T		19-05-2005
			SK	2902004 A3		03-11-2004
			US	2005089575 A1		28-04-2005
-----	-----	-----	-----	-----	-----	-----
WO 9405264	A	17-03-1994	AT	144417 T		15-11-1996
			AU	4950093 A		29-03-1994
			DE	4229085 A1		03-03-1994
			DK	658104 T3		17-03-1997
			EP	0658104 A1		21-06-1995
			ES	2095070 T3		01-02-1997
			GR	3022182 T3		31-03-1997
			JP	8500599 T		23-01-1996
			SG	66741 A1		21-05-2002
			US	5562920 A		08-10-1996
-----	-----	-----	-----	-----	-----	-----