

(12) STANDARD PATENT
(19) AUSTRALIAN PATENT OFFICE

(11) Application No. **AU 2012212127 B2**

(54) Title
Digital sequence analysis of DNA methylation

(51) International Patent Classification(s)
C12Q 1/68 (2006.01)

(21) Application No: **2012212127** (22) Date of Filing: **2012.02.02**

(87) WIPO No: **WO12/106525**

(30) Priority Data

(31) Number	(32) Date	(33) Country
61/438,649	2011.02.02	US

(43) Publication Date: **2012.08.09**

(44) Accepted Journal Date: **2016.06.02**

(71) Applicant(s)
Mayo Foundation for Medical Education and Research;Exact Sciences Corporation

(72) Inventor(s)
Ahlquist, David A.;Taylor, William R.;Zou, Hongzhi;Lidgard, Graham P.

(74) Agent / Attorney
Davies Collison Cave, Level 15 1 Nicholson Street, MELBOURNE, VIC, 3000

(56) Related Art
WO 2003/044232
WO 2003/064701
ZOU, H. et al., AACCC Annual Meeting 2010, Abstract No. D-144, Clinical Chemistry, 2010, vol. 56, no. 6, Supplement, page A199
AHLQUIST, D. A., Gastroenterology, 2010, vol. 138, pages 2127-2139
WO 2010/118016

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
9 August 2012 (09.08.2012)

(10) International Publication Number
WO 2012/106525 A2

- (51) International Patent Classification:
C40B 20/00 (2006.01) *C12Q 1/68* (2006.01)
- (21) International Application Number:
PCT/US2012/023646
- (22) International Filing Date:
2 February 2012 (02.02.2012)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:
61/438,649 2 February 2011 (02.02.2011) US
- (71) Applicants (for all designated States except US): **EXACT SCIENCES CORPORATION** [US/US]; 441 Charmany Drive, Madison, Wisconsin 53719 (US). **MAYO FOUNDATION FOR MEDICAL EDUCATION AND RESEARCH** [US/US]; 200 First Street SW, Rochester, Minnesota 55905 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **AHLQUIST, David A.** [US/US]; 6576 Buckridge Court NE, Rochester, Minnesota 55906 (US). **TAYLOR, William R.** [US/US]; 218 N. Garden St., Lake City, Minnesota 55041 (US). **ZOU, Hongzhi** [CN/US]; 8570 Greenway Blvd. #406, Middleton, Wisconsin 53562 (US). **LIDGARD, Graham P.** [US/US]; 8921 Timber Wolf Lane, Madison, Wisconsin 53717 (US).
- (74) Agent: **BROW, Mary Ann D.**; Casimir Jones, S.C., 2275 Deming Way, Suite 310, Middleton, Wisconsin 53562 (US).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.
- (84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:
- without international search report and to be republished upon receipt of that report (Rule 48.2(g))
 - with sequence listing part of description (Rule 5.2(a))



WO 2012/106525 A2

(54) Title: DIGITAL SEQUENCE ANALYSIS OF DNA METHYLATION

(57) Abstract: The present invention relates to methods and compositions for determination of and uses of specific methylation patterns indicative of adenoma and carcinoma. In particular, the invention relates to analysis of defined CpG loci that are coordinately methylated in DNAs from cancer and adenoma samples, methods for identifying coordinately methylated loci, and methods of using analysis of coordinately methylated loci in one or more marker regions in the design of assays for adenoma and cancer.

DIGITAL SEQUENCE ANALYSIS OF DNA METHYLATION

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application claims the benefit of U.S. Provisional Patent Application Serial Number 61/438,649, filed February 2, 2011, which is incorporated by reference in its
5 entirety.

FIELD OF THE INVENTION

The present invention relates to methods and compositions for determination of and uses of specific methylation patterns indicative of adenoma and carcinoma. In particular, the invention relates to analysis of defined CpG loci that are coordinately methylated in DNAs
10 from cancer and adenoma samples, methods for identifying coordinately methylated loci, and methods of using analysis of coordinately methylated loci in one or more marker regions in the design of assays for adenoma and cancer having improved sensitivity and specificity.

BACKGROUND OF THE INVENTION

In higher order eukaryotes, DNA may be methylated at cytosines located 5' to
15 guanosine in CpG dinucleotides. This modification has important regulatory effects on gene expression, especially when involving CpG rich areas, known as CpG islands, often found in the promoter regions of genes. While approximately 75% of the CpG sites throughout the human genome are methylated, CpG sites within CpG islands are normally unmethylated, and aberrant methylation of CpG islands has been associated with certain diseases, including
20 cancers. For example, CpG island hypermethylation is associated with transcriptional inactivation of defined tumor suppressor genes in human cancers, *e.g.*, colorectal cancer. Therefore, detection of hypermethylated nucleic acid could indicate susceptibility or onset of various forms of cancers.

Despite indications suggesting a link between CpG island methylator phenotype
25 (CIMP) and cancers (see, *e.g.*, Baylin SB, *et al.*, *Adv Cancer Res* 1998;72:141–196 and Jones PA, *et al.*, *Nat Rev Genet* 2002;3:415–428), the idea that analysis of methylation status alone could be a useful diagnostic or prognostic tool has been controversial. As discussed by Issa, *et al.* in an editorial in *Gastroenterology* 179(3):2005, researchers had mixed results in confirming the link between CIMP and cancers. Although CIMP was reportedly
30 demonstrated in multiple other malignancies (Shen, I., *et al.* *J Natl Cancer Inst* 2002;94:755–761; Garcia-Manero G, *et al.*, *Clin Cancer Res* 2002;8:2217–2224; Toyota M, *et al.*, *Blood*

2001;97:2823–2829; Ueki T, *et al.*, *Cancer Res* 2000;60:1835–1839; Toyota M, *et al.*,
Cancer Res 1999;59:5438–5442; Strathdee G, *et al.*, *Am J Pathol* 2001;158:1121–1127; Abe
M, *et al.*, *Cancer Res* 2005;65:828–834) and several groups confirmed the original findings
using similar markers and technology (Whitehall VL, *et al.*, *Cancer Res* 2002;62:6011–6014;
5 van Rijnsoever M, *et al.*, *Gut* 2002;51:797–802) other groups were not able to establish such
links (Eads CA, *et al.*, *Cancer Res* 2001;61:3410–3418; Esteller M, *et al.*, *Cancer Res*
2000;60:129–133). As late as 2003, a publication concluded that all methylation events in
colorectal cancer were related to aging rather than neoplasia (Yamashita K, *et al.*, *Cancer*
Cell 2003;4:121–131).

10 The discrepant results have been attributed in part to the fact that it has been
demonstrated that 70% to 80% of aberrant DNA methylation events in colorectal cancer are
age-related (Toyota M, *et al.*, *Proc Natl Acad Sci U S A* 1999;96:8681–8686) and that
cancer-linked phenotypes are only clear when these are filtered out. It has also been noted
that overly sensitive, non-quantitative methods can overestimate methylation and mask the
15 distinctions between methylation that is associated with cancer and that which is not. Issa
states that "methylation events (alone) may not provide the ideal universal cancer marker they
were once thought to be because CIMP target genes will not be useful to screen for all
colorectal cancers (many false negatives are predicted), and non-CIMP target genes will
likely yield a high rate of false-positives because they are also methylated in normal
20 appearing mucosa of older individuals without tumors" (Issa, *et al.*, *supra*).

One approach to increase the clinical specificity of methylation analyses in cancer
detection is to look at multiple marker genes. For example, Zou, *et al.*, examined the
methylation status of BMP3, EYA2, ALX4, and vimentin in cancer samples. While
methylation levels were significantly higher in both cancer and adenoma than in normal
25 epithelium, for each of the four genes, the sensitivity as determined by receiver operating
curves was not significantly improved by combining any or all markers compared with the
best single marker. (Zou, *et al.*, *Cancer Epidemiol Biomarkers Prev* 2007;16(12):2686).

Zou also looked at neoplasms showing methylation in more than one of the marker
genes and found that co-methylation was frequent, with 72% of the cancers and 84% of the
30 adenomas tested showing hypermethylation in two or more of the genes. Zou reported that
methylation of one or more of four (at least one), two or more of four, three or more of four,
or four of four of these marker genes was noted in 88%, 72%, 53%, and 41% of 74 cancers
and 98%, 84%, 60%, and 39% of 62 adenomas, compared with 24%, 7%, 3%, and 0% of 70

normal epithelia, respectively, demonstrating that although the assay gets progressively more specific as when more genes are included in the comethylation set, the sensitivity declines precipitously.

SUMMARY OF THE INVENTION

5 The present invention relates to the methods of identifying regions of specific genes and specific regions of genomic nucleic acid useful in the detection of methylation associated with colorectal cancer. Methods comprise, *e.g.*, detecting methylated sequences in, for example, tissue biopsy, stool extract, or other body fluids with improved sensitivity and specificity. In preferred embodiments, the present invention provides methods of
10 methylation analysis comprising identifying methylation loci showing advantageous methylation ratios when methylation in non-normal cells, *e.g.*, cancer or adenoma cells is compared to background methylation in normal cells. In some embodiments, the present invention relates to methods of analyzing methylation at each of several loci in a set of possible methylation sites within a marker sequence, wherein the presence of methylation at
15 all of the loci within the defined set of sites occurs more frequently in cancer and adenoma cells than in normal cells, such that a finding of methylation at all of the loci in the defined subset of loci in a sample is indicative of adenoma or cancer.

In some embodiments, the present invention provides a method of identifying a set of methylated CpG loci in a marker nucleic acid wherein methylation is indicative of adenoma,
20 comprising:

- a) determining the methylation status of a defined set of CpG loci in each of a plurality of individual copies of a marker nucleic acid from a plurality of normal samples;
- b) determining the methylation status of said defined set of CpG loci in each of a plurality of individual copies of said marker nucleic acid from a plurality of non-normal (*e.g.*,
25 adenoma or cancer) samples to identify a defined subset of CpG loci from within said defined set,

wherein the percentage of individual copies of said marker nucleic acid from said plurality of normal samples that are methylated at all of said CpG loci in said defined subset is less than the percentage of individual copies of said marker nucleic acid from said plurality
30 of non-normal samples that are methylated at all of said CpG loci in said defined subset, and wherein methylation at all of said CpG loci in said defined subset in said marker nucleic acid is indicative of a non-normal state, *e.g.*, adenoma and/or cancer. In certain embodiments, the

mean percentage of individual copies of the marker nucleic acid methylated at all loci in said defined set of CpG loci in said plurality of non-normal samples is greater than the mean percentage of individual copies of the marker nucleic acid methylated at all loci in said defined set of CpG loci in the plurality of normal samples. In preferred embodiments, the
5 mean percentage of individual copies of the marker nucleic acid methylated at all loci in said defined set of CpG loci in the plurality of non-normal samples is at least one standard deviation, preferably two standard deviations, more preferably three standard deviations greater than the mean percentage of individual copies of said marker nucleic acid methylated at all loci in said defined set of CpG loci in said plurality of normal samples.

10 In some embodiments, the defined subset of CpG loci consists of the same loci in the defined set of CpG loci.

Determination of the methylation status of the set of CpG loci may be accomplished by any method known to those of skill in the art. In some embodiments, the method comprises treating DNA from the samples with bisulfite. Bisulfite modification treatment is
15 described, *e.g.*, in U.S. Pat. No. 6,017,704, the entire disclosure of which is incorporated herein by reference. In some embodiments, determining the methylation status of the defined set of CpG loci comprises digital analysis of each of a plurality of CpG loci in a plurality of individual copies of a marker nucleic acid. In some preferred embodiments, digital analysis comprise digital sequencing, and/or digital PCR.

20 In certain preferred embodiments, non-normal sample comprises an adenoma sample, and in particular preferred embodiments, comprises a colorectal adenoma sample. In some preferred embodiments, a non-normal sampled comprises a cancer sample, and in certain preferred embodiments, comprises a colorectal cancer sample.

The present invention provides methods of detecting cancer or adenoma in a sample,
25 *e.g.*, from a subject. In some embodiments, the present invention provides methods comprising determining the methylation status of each CpG locus in a defined subset of CpG loci in at least one cancer or adenoma marker nucleic acid molecule, wherein methylation at each of the CpG loci in the defined subset of CpG loci in the cancer or adenoma marker nucleic acid molecule is indicative of cancer or adenoma in the sample. In certain preferred
30 embodiments, the defined subset comprises at least three CpG loci, while in some preferred embodiments, the defined subset comprises at least four CpG loci or at least five CpG loci.

In certain embodiments, the determining comprises analysis of the CpG loci in a nucleic acid detection assay configured to determine the methylation status of each of the loci in a single nucleic acid detection assay. In some preferred embodiments, the determining comprises analysis of the CpG loci in a nucleic acid detection assay configured to determine the methylation status of each of said loci in a single reaction mixture. In some 5 embodiments, the nucleic acid detection assay comprises a primer extension assay. In certain preferred embodiments, the nucleic acid detection assay may comprise one or more of a nucleic acid amplification assay, a nucleic acid sequencing assay, a structure-specific cleavage assay, a 5' nuclease cleavage assay, an invasive cleavage assay and/or a ligation 10 assay.

The methods of the present invention are not limited to the analysis of a single cancer or adenoma marker nucleic acid. For example, in some embodiments, the methylation status of each CpG locus in a defined subset of CpG loci in at least one cancer or adenoma marker nucleic acid molecule comprises analysis of nucleic acid molecules from a plurality of cancer 15 or adenoma markers. In some embodiments, the plurality of cancer or adenoma markers comprises at least three cancer or adenoma markers, while in some embodiments, the plurality comprises at least four cancer or adenoma markers. In some preferred embodiments, the cancer or adenoma markers and nucleic acid molecules are selected from the group comprising Vimentin, BMP3, Septin 9, TFPI2, 2 regions of LRAT, and EYA4 20 markers and nucleic acid molecules. In some embodiments, the assay methods of the present invention are combined with the analysis of one or more other cancer markers, such as fecal occult blood markers (*e.g.*, hemoglobin, alpha-defensin, calprotectin, α 1-antitrypsin, albumin, MCM2, transferrin, lactoferrin, and lysozyme).

In certain preferred embodiments of the method described herein, a cancer or 25 adenoma marker nucleic acid molecule comprises a vimentin nucleic acid molecule, and in some particularly preferred embodiments, the defined subset of CpG loci in the vimentin nucleic acid molecule comprises loci 37, 40, and 45.

In certain preferred embodiments of the method described herein, a cancer or adenoma marker nucleic acid molecule comprises a BMP3 nucleic acid molecule, and in 30 some particularly preferred embodiments, the defined subset of CpG loci in the BMP3 nucleic acid molecule comprises loci 34, 53, and 61.

In certain preferred embodiments of the method described herein, a cancer or adenoma marker nucleic acid molecule comprises a Septin9 nucleic acid molecule, and in some particularly preferred embodiments, the defined subset of CpG loci in said Septin9 nucleic acid molecule comprises loci 59, 61, 68, and 70.

5 In certain preferred embodiments of the method described herein, a cancer or adenoma marker nucleic acid molecule comprises a TFPI2 nucleic acid molecule and in some particularly preferred embodiments, the defined subset of CpG loci in said TFPI2 nucleic acid molecule comprises loci 55, 59, 63, and 67.

10 In certain preferred embodiments of the method described herein, a cancer or adenoma marker nucleic acid molecule comprises an EYA4 nucleic acid molecule, and in some particularly preferred embodiments, the defined subset of CpG loci in said EYA4 nucleic acid molecule comprises loci 31, 34, 37, and 44.

In certain preferred embodiments of the method described herein, the at least one cancer or adenoma marker or nucleic acid molecule comprises a plurality markers or nucleic acid molecules comprising Vimentin, BMP3, Septin9, and TFPI2 markers or nucleic acid molecules.

The present invention further provides methods of selecting a defined set of CpG loci in a marker nucleic acid wherein methylation is indicative of non-normal status, *e.g.*, adenoma or cancer, comprising a) determining the methylation status of a plurality of CpG loci in each of a plurality of individual copies of a marker nucleic acid from a plurality of normal samples; b) determining the methylation status of the plurality of CpG loci in each of a plurality of individual copies of said marker nucleic acid from a plurality of non-normal (*e.g.*, adenoma or cancer) samples; c) determining methylation ratios for each locus in the plurality of said CpG loci in the marker nucleic acid; and d) selecting a defined set of CpG loci in the marker nucleic acid, wherein the defined set of CpG loci comprises a plurality of CpG loci having advantageous methylation ratios correlating with non-normal status (*e.g.*, adenoma or cancer).

In some embodiments, determining the methylation ratios comprises determining the ratio between the mean methylation at each of the plurality of CpG loci in the normal samples to the mean methylation at each corresponding CpG locus in said plurality of CpG loci in the non-normal samples. In preferred embodiments, the plurality of individual copies of a marker nucleic acid analyzed in the normal and non-normal (*e.g.*, adenoma or cancer)

5 samples comprises at least 10, preferably at least 100, more preferably at least 1000, still more preferably at least 10,000 and still more preferably at least 100,000 copies. The number of copies analyzed is not limited to these whole numbers, but may be any integer above about 10. The number of copies from different sample types, *e.g.*, normal and non-normal need not be equal.

10 In certain preferred embodiments of the methods of selecting a defined set of CpG loci in a marker nucleic acid described herein, the plurality of normal and non-normal (*e.g.*, adenoma or cancer) samples compared comprises at least 10, preferably at least 25, still more preferably at least 100 samples. The number of samples analyzed is not limited to these whole numbers, but may be any integer above about 10. The number of different samples of the different sample types, *e.g.*, normal and non-normal, need not be equal.

In certain embodiments, the defined set of CpG loci comprises at least three CpG loci, preferably at least four CpG loci, more preferably at least five CpG loci.

15 Determination of the methylation status of the plurality of CpG loci may be accomplished by any method known to those of skill in the art, including those described in more detail, below. In some embodiments, the method comprises treating DNA from the samples with bisulfite. In some embodiments, determining the methylation status of the defined set of CpG loci comprises digital analysis of each of a plurality of CpG loci in a plurality of individual copies of a marker nucleic acid. In some preferred embodiments, 20 digital analysis comprises digital sequencing, and/or digital PCR. Methods of preparing samples, *e.g.*, stool samples, for analysis are also known in the art. See, *e.g.*, US7005266; 6,303,304; 5,741,650; 5,952,178; and 6,268,136, each incorporated herein by reference.

DEFINITIONS

25 To facilitate an understanding of the present invention, a number of terms and phrases are defined below.

As used herein, the terms "digital sequencing" and "single molecule sequencing" are used interchangeably and refer to determining the nucleotide sequence of individual nucleic acid molecules. Systems for individual molecule sequencing include but are not limited to the 454 FLX™ or 454 TITANIUM™ (Roche), the SOLEXA™/ Illumina Genome Analyzer 30 (Illumina), the HELISCOPE™ Single Molecule Sequencer (Helicos Biosciences), and the SOLID™ DNA Sequencer (Life Technologies/Applied Biosystems) instruments), as well as

other platforms still under development by companies such as Intelligent Biosystems and Pacific Biosystems.

As used herein, the term "background" as used in reference to methylation of a locus or region refers to methylation observed in a normal cell or sample at a nucleic acid locus or region that is generally unmethylated in normal cells. For example, CpG islands are generally considered unmethylated in normal human cells but methylation is not completely absent in the CpG islands of normal cells.

As used herein, "methylation" or "methylated," as used in reference to the methylation status of a cytosine, *e.g.*, in a CpG locus, generally refers to the presence or absence of a methyl group at position 5 of the cytosine residue (*i.e.*, whether a particular cytosine is 5-methylcytosine). Methylation may be determined directly, *e.g.*, as evidenced by routine methods for analysis of methylation status of cytosines, *e.g.*, by determining the sensitivity (or lack thereof) of a particular C-residue to conversion to uracil by treatment with bisulfite. For example, a cytosine residue in a sample that is not converted to uracil when the sample is treated with bisulfite in a manner that would be expected to convert that residue if non-methylated (*e.g.*, under conditions in which a majority or all of the non-methylated cytosines in the sample are converted to uracils) may generally be deemed "methylated".

As used herein, the terms "digital PCR", "single molecule PCR" and "single molecule amplification" refer to PCR and other nucleic acid amplification methods that are configured to provide amplification product or signal from a single starting molecule. Typically, samples are divided, *e.g.*, by serial dilution or by partition into small enough portions (*e.g.*, in microchambers or in emulsions) such that each portion or dilution has, on average, no more than a single copy of the target nucleic acid. Methods of single molecule PCR are described, *e.g.*, in US 6,143,496, which relates to a method comprising dividing a sample into multiple chambers such that at least one chamber has at least one target, and amplifying the target to determine how many chambers had a target molecule; US 6,391,559; which relates to an assembly for containing and portioning fluid; and US 7,459,315, which relates to a method of dividing a sample into an assembly with sample chambers where the samples are partitioned by surface affinity to the chambers, then sealing the chambers with a curable "displacing fluid." See also US 6,440,706 and US 6,753,147, and Vogelstein, *et al.*, Proc. Natl. Acad. Sci. USA Vol. 96, pp. 9236–9241, August 1999. See also US 20080254474, describing a combination of digital PCR combined with methylation detection.

As used herein, "sensitivity" as used in reference to a diagnostic assay, *e.g.*, a methylation assay, refers to clinical sensitivity - the proportion of positive samples that give a positive result using a diagnostic assay. Sensitivity is generally calculated as the number of true positives identified by the assay, divided by the sum of the number of true positives and the number of false negatives determined by the assay on known positive samples. Similarly, the term "specificity" refers to the proportion or number of true negatives determined by the assay divided by the sum of the number of true negatives and the number of false positives determined by the assay on known negative sample(s).

As used herein in reference to diagnostic or analysis assays, the term "complementary" refers to different assays that, when used together, provide a more sensitive and/or specific result than can be provided by any one of the different assays used alone.

As used herein, the term "informative" or "informativeness" refers to a quality of a marker or panel of markers, and specifically to the likelihood of finding a marker (or panel of markers) in a positive sample.

The term "sample" as used herein is used in its broadest sense. For example, a sample suspected of containing a human gene or chromosome or sequences associated with a human chromosome may comprise a cell, chromosomes isolated from a cell (*e.g.*, a spread of metaphase chromosomes), genomic DNA (in solution or bound to a solid support such as for Southern blot analysis), RNA (in solution or bound to a solid support such as for Northern blot analysis), cDNA (in solution or bound to a solid support) and the like.

As used herein, the term "CpG island" refers to a genomic DNA region that contains a high percentage of CpG sites relative to the average genomic CpG incidence (per same species, per same individual, or per subpopulation (*e.g.*, strain, ethnic subpopulation, or the like). Various parameters and definitions for CpG islands exist; for example, in some embodiments, CpG islands are defined as having a GC percentage that is greater than 50% and with an observed/expected CpG ratio that is greater than 60% (Gardiner-Garden *et al.* (1987) *J Mol. Biol.* 196:261-282; Baylin *et al.* (2006) *Nat. Rev. Cancer* 6:107-116; Irizarry *et al.* (2009) *Nat. Genetics* 41:178-186; each herein incorporated by reference in its entirety). In some embodiments, CpG islands may have a GC content >55% and observed CpG/expected CpG of 0.65 (Takai *et al.* (2007) *PNAS* 99:3740-3745; herein incorporated by reference in its entirety). Various parameters also exist regarding the length of CpG islands. As used herein, CpG islands may be less than 100 bp; 100-200 bp, 200-300 bp, 300-500 bp, 500-750 bp; 750-1000 bp; 1000 or more bp in length. In some embodiments, CpG islands show altered

methylation patterns relative to controls (*e.g.*, altered methylation in cancer subjects relative to subjects without cancer; tissue-specific altered methylation patterns; altered methylation in stool from subjects with colorectal neoplasia (*e.g.*, colorectal cancer, colorectal adenoma) relative to subjects without colorectal neoplasia). In some embodiments, altered methylation involves hypermethylation. In some embodiments, altered methylation involves hypomethylation.

As used herein, the term "CpG shore" or "CpG island shore" refers to a genomic region external to a CpG island that is or that has potential to have altered methylation patterns (see, *e.g.*, Irizarry *et al.* (2009) Nat. Genetics 41:178-186; herein incorporated by reference in its entirety). CpG island shores may show altered methylation patterns relative to controls (*e.g.*, altered methylation in cancer subjects relative to subjects without cancer; tissue-specific altered methylation patterns; altered methylation in stool from subjects with colorectal neoplasia (*e.g.*, colorectal cancer, colorectal adenoma) relative to subjects without colorectal neoplasia). In some embodiments, altered methylation involves hypermethylation. In some embodiments, altered methylation involves hypomethylation. CpG island shores may be located in various regions relative to CpG islands (see, *e.g.*, Irizarry *et al.* (2009) Nat. Genetics 41:178-186; herein incorporated by reference in its entirety). Accordingly, in some embodiments, CpG island shores are located less than 100 bp; 100-250 bp; 250-500 bp; 500-1000 bp; 1000-1500 bp; 1500-2000 bp; 2000-3000 bp; 3000 bp or more away from a CpG island.

The term "target," when used in reference to a nucleic acid detection or analysis method, refers to a nucleic acid having a particular sequence of nucleotides to be detected or analyzed, *e.g.*, in a sample suspected of containing the target nucleic acid. In some embodiments, a target is a nucleic acid having a particular sequence for which it is desirable to determine a methylation status. When used in reference to the polymerase chain reaction, "target" generally refers to the region of nucleic acid bounded by the primers used for polymerase chain reaction. Thus, the "target" is sought to be sorted out from other nucleic acid sequences that may be present in a sample. A "segment" is defined as a region of nucleic acid within the target sequence. The term "sample template" refers to nucleic acid originating from a sample that is analyzed for the presence of a target.

As used herein, the term "locus" refers to a particular position, *e.g.*, of a mutation, polymorphism, or a C residue in a CpG dinucleotide, within a defined region or segment of nucleic acid, such as a gene or any other characterized sequence on a chromosome or RNA

molecule. A locus is not limited to any particular size or length, and may refer to a portion of a chromosome, a gene, functional genetic element, or a single nucleotide or basepair. As used herein in reference to CpG sites that may be methylated, a locus refers to the C residue in the CpG dinucleotide.

5 As used herein, the term "methylation ratio" refers to the amount or degree of methylation observed for particular methylation region or locus (*e.g.*, a CpG locus in a marker gene or region) in a plurality of non-normal cells (*e.g.*, cells in a particular disease state, such as cancerous or pre-cancerous cells) compared to the amount or degree of methylation observed for the same region or locus in a plurality of normal cells (*e.g.*, cells
10 that are not in the particular disease state of interest). For example, for a CpG locus showing mean methylation of 8.39889% in a sampling of normal cells and a mean methylation of 74.0771% in a sampling of a plurality of adenoma cells, a methylation ratio may be expressed as the ratio of the means determined for normal cells:adenoma cells, or 0.11348. A methylation ratio need not be expressed in any particular manner or by any particular
15 calculation. By way of example and not limitation, the methylation ratio above may alternatively be expressed, *e.g.*, as 8.39889:74.0771; 8.39889/74.0771; 74.0771:8.39889; as a calculated "fold methylation over background" 8.81987, *etc.*

 As used herein, the term "advantageous methylation ratio" refers to a methylation ratio for a locus at which methylation correlates with a cellular status, *e.g.*, a particular
20 disease state (for example, normal, precancerous, cancerous) that, when compared to other methylation loci correlated with the same disease state, displays a higher percentage methylation in a population of non-normal cells compared to background levels of methylation at the same locus in a population of normal cells. In some instances, certain CpG loci *e.g.*, within a methylation marker sequence, display a much greater signal-to-noise, *i.e.*,
25 degree in methylation compared to background than other loci in the same marker sequence. In other instances, certain disease-associated marker genes or regions display advantageous methylation ratios at some or all loci compared to the methylation ratios observed for some or all loci within another marker sequence.

 As used herein, the term "coordinately methylated" is used in reference to methylation
30 loci, *e.g.*, CpG loci in a marker sequence, that exhibit a particular pattern of methylation that correlates with a cellular status, *e.g.*, a particular disease state (for example, normal, precancerous, cancerous). In preferred embodiments, methylation loci that are all methylated in a manner correlated with a disease state may be deemed to be coordinately methylated in

cells having that disease state. "Coordinate methylation" is not limited to situations in which all of the coordinated loci are methylated. Any pattern of methylation among a particular set of loci that correlates with a cellular status, including patterns in which all of the coordinate loci are methylated, patterns in which the loci exhibit a reproducible pattern of methylation and non-methylation, and patterns in which none of the loci within the set are methylated are all included within the meaning of "coordinate methylation."

As used herein, the term "coordinate methylation analysis" is used interchangeably with "multimethylation analysis" and refers to an assay in which the methylation statuses of a plurality of individual methylation loci in a marker sequence, *e.g.*, CpG loci, are determined together. In preferred embodiments, coordinate methylation analysis is performed using a digital/single copy method (*e.g.*, digital sequencing) or an assay method configured to interrogate all of the selected CpG loci on each molecule tested, such that the methylation pattern in each single molecule tested is revealed.

As used herein, the term "defined set" of CpG loci (or other methylation loci) refers to the set of CpG loci in a marker gene or region selected for methylation analysis. A defined set of CpG loci in a marker gene or region may comprise all CpG loci in the gene or region, or it may comprise fewer than all of the loci in that gene or region.

As used herein the term "defined subset" of CpG loci (or other methylation loci) refers to a subset of the defined set of CpG loci in a marker gene or region, the methylation of which has been determined to be indicative of a non-normal state, *e.g.*, adenoma or cancer. For example, in coordinate methylation analysis to determine the presence of colorectal cancer, the methylation status of a defined subset of CpG loci in at least one cancer marker nucleic acid molecule is determined, with simultaneous methylation at all of said CpG loci in the defined subset being indicative of cancer in the sample. A defined subset of CpG loci in a marker gene or region may comprise all CpG loci in the defined set, or it may comprise fewer than all of the loci in the defined set of loci in that gene or region.

As used herein, the term "colorectal cancer" is meant to include the well-accepted medical definition that defines colorectal cancer as a medical condition characterized by cancer of cells of the intestinal tract below the small intestine (*e.g.*, the large intestine (colon), including the cecum, ascending colon, transverse colon, descending colon, and sigmoid colon, and rectum). Additionally, as used herein, the term "colorectal cancer" is meant to further include medical conditions which are characterized by cancer of cells of the duodenum and small intestine (jejunum and ileum).

As used herein, the term "metastasis" is meant to refer to the process in which cancer cells originating in one organ or part of the body relocate to another part of the body and continue to replicate. Metastasized cells subsequently form tumors which may further metastasize. Metastasis thus refers to the spread of cancer from the part of the body where it originally occurs to other parts of the body. As used herein, the term "metastasized colorectal cancer cells" is meant to refer to colorectal cancer cells which have metastasized; colorectal cancer cells localized in a part of the body other than the duodenum, small intestine (jejunum and ileum), large intestine (colon), including the cecum, ascending colon, transverse colon, descending colon, and sigmoid colon, and rectum.

As used herein, "an individual is suspected of being susceptible to metastasized colorectal cancer" is meant to refer to an individual who is at an above-average risk of developing metastasized colorectal cancer. Examples of individuals at a particular risk of developing metastasized colorectal cancer are those whose family medical history indicates above average incidence of colorectal cancer among family members and/or those who have already developed colorectal cancer and have been effectively treated who therefore face a risk of relapse and recurrence. Other factors which may contribute to an above-average risk of developing metastasized colorectal cancer which would thereby lead to the classification of an individual as being suspected of being susceptible to metastasized colorectal cancer may be based upon an individual's specific genetic, medical and/or behavioral background and characteristics.

The term "neoplasm" as used herein refers to any new and abnormal growth of tissue. Thus, a neoplasm can be a premalignant neoplasm or a malignant neoplasm.

The term "neoplasm-specific marker" refers to any biological material that can be used to indicate the presence of a neoplasm. Examples of biological materials include, without limitation, nucleic acids, polypeptides, carbohydrates, fatty acids, cellular components (*e.g.*, cell membranes and mitochondria), and whole cells. In some instances, markers are particular nucleic acid regions, *e.g.*, genes, intragenic regions, *etc.* Regions of nucleic acid that are markers may be referred to, *e.g.*, as "marker genes," "marker regions," "marker sequences," *etc.*

The term "colorectal neoplasm-specific marker" refers to any biological material that can be used to indicate the presence of a colorectal neoplasm (*e.g.*, a premalignant colorectal neoplasm; a malignant colorectal neoplasm). Examples of colorectal neoplasm-specific markers include, but are not limited to, exfoliated epithelial markers (*e.g.*, bmp-3, bmp-4, SFRP2, vimentin, septin9, ALX4, EYA4, TFPI2, NDRG4, FOXE1, long DNA, BAT-26, K-

- ras, APC, melanoma antigen gene, p53, BRAF, and PIK3CA) and fecal occult blood markers (e.g., hemoglobin, alpha-defensin, calprotectin, α 1-antitrypsin, albumin, MCM2, transferrin, lactoferrin, and lysozyme). See also US 7485420; US7432050; US5352775; US5648212; USRE36713; US5527676; US5955263; US6090566; US6245515; US6677312; 5 US6800617; US7087583; and US7267955, each incorporated herein by reference.

Additional markers include but are not limited those in Table 1, below:

Table 1

Accession	Symbol	GeneID	Reference
NM_000038	APC	324	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000044	AR	367	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
AB033043	KIAA1217	56243	www.methdb.de/ and/or www.mdanderson.org/
AK055404	KIAA0984	23329	www.methdb.de/ and/or www.mdanderson.org/
AK090480			www.methdb.de/ and/or www.mdanderson.org/
BC041476			www.methdb.de/ and/or www.mdanderson.org/
BX648962	DKFZp686K1684	440034	www.methdb.de/ and/or www.mdanderson.org/
NM_000017	ACADS	35	www.methdb.de/ and/or www.mdanderson.org/
NM_000022	ADA	100; 79015	www.methdb.de/ and/or www.mdanderson.org/
NM_000038	APC	324	www.methdb.de/ and/or www.mdanderson.org/
NM_000038	APC	324	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000043	FAS	355; 819114	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000044	AR	367	www.methdb.de/ and/or www.mdanderson.org/
NM_000044	AR	367	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000076	CDKN1C	1028	www.methdb.de/ and/or www.mdanderson.org/
NM_000077	CDKN2A	1029; 51198	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000077	CDKN2A	1029; 51198	www.methdb.de/ and/or www.mdanderson.org/
NM_000077	CDKN2A	1029; 51198	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000088	COL1A1	1277	www.methdb.de/ and/or www.mdanderson.org/
NM_000095	COMP	1311	www.methdb.de/ and/or www.mdanderson.org/
NM_000104	CYP1B1	1545	www.methdb.de/ and/or www.mdanderson.org/
NM_000115	EDNRB	1910	www.methdb.de/ and/or www.mdanderson.org/
NM_000115	EDNRB	1910	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000115	EDNRB	1910	www.methdb.de/ and/or www.mdanderson.org/
NM_000125	ESR1	2099	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000125	ESR1	2099	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000125	ESR1	2099	www.methdb.de/ and/or www.mdanderson.org/
NM_000125	ESR1	2099	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000182	HADHA	3030	www.methdb.de/ and/or www.mdanderson.org/
NM_000193	SHH	6469	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000249	MLH1	4292	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000249	MLH1	4292	www.methdb.de/ and/or www.mdanderson.org/
NM_000280	PAX6	5080	www.methdb.de/ and/or www.mdanderson.org/

NM_000280	PAX6	5080	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000280	PAX6	5080	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000308	PPGB	5476	www.methdb.de/ and/or www.mdanderson.org/
NM_000314	PTEN	5728	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000321	RB1	5925	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000321	RB1	5925	www.methdb.de/ and/or www.mdanderson.org/
NM_000336	SCNN1B	6338	www.methdb.de/ and/or www.mdanderson.org/
NM_000362	TIMP3	7078	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000362	TIMP3	7078	www.methdb.de/ and/or www.mdanderson.org/
NM_000362	TIMP3	7078	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000378	WT1	7490	www.methdb.de/ and/or www.mdanderson.org/
NM_000402	G6PD	2539	www.methdb.de/ and/or www.mdanderson.org/
NM_000438	PAX3	5077	www.methdb.de/ and/or www.mdanderson.org/
NM_000443	ABCB4	5244	www.methdb.de/ and/or www.mdanderson.org/
NM_000453	SLC5A5	6528	www.methdb.de/ and/or www.mdanderson.org/
NM_000453	SLC5A5	6528	www.methdb.de/ and/or www.mdanderson.org/
NM_000475	NR0B1	190	www.methdb.de/ and/or www.mdanderson.org/
NM_000492	CFTR	1080	www.methdb.de/ and/or www.mdanderson.org/
NM_000492	CFTR	1080	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000514	GDNF	2668	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000517	HBA2	3040	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000520	HEXA	3073; 80072	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000524	HTR1A	3350	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000551	VHL	7428	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000551	VHL	7428	www.methdb.de/ and/or www.mdanderson.org/
NM_000551	VHL	7428	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000610	CD44	960	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000610	CD44	960	www.methdb.de/ and/or www.mdanderson.org/
NM_000612	IGF2	3481; 492304	www.methdb.de/ and/or www.mdanderson.org/
NM_000620	NOS1	4842	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000680	ADRA1A	148	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000717	CA4	762	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000721	CACNA1E	777	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000782	CYP24A1	1591	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000799	EPO	2056	www.methdb.de/ and/or www.mdanderson.org/
NM_000813	GABRB2	2561	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000818	GAD2	2572	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000829	GRIA4	2893	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000830	GRIK1	2897	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000834	GRIN2B	2904	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000843	GRM6	2916	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000852	GSTP1	2950	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000852	GSTP1	2950	www.methdb.de/ and/or www.mdanderson.org/
NM_000852	GSTP1	2950	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000857	GUCY1B3	2983	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000863	HTR1B	3351	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000902	MME	4311	www.methdb.de/ and/or www.mdanderson.org/
NM_000914	OPRM1	4988	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000915	OXT	5020	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_000926	PGR	5241	www.methdb.de/ and/or www.mdanderson.org/
NM_000927	ABCB1	5243	www.methdb.de/ and/or www.mdanderson.org/
NM_000959	PTGFR	5737	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_000965	RARB	5915	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_000965	RARB	5915	www.methdb.de/ and/or www.mdanderson.org/
NM_000965	RARB	5915	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM_000997	RPL37	6167	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001001336	CYB5R2	51700	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001001723	TMEM1	7109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001002295	GATA3	2625	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001003689	L3MBTL2	83746	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001003891	PCQAP	51586	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_001007792	NTRK1	4914	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001008503	OPRM1	4988	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001008504	OPRM1	4988	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001008505	OPRM1	4988	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001009598	RXRG	6258	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001011545	BACH1	571	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001012331	NTRK1	4914	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001013464	LOC401363	401363; 441242; 402532	www.methdb.de/ and/or www.mdanderson.org/
NM_001018084	SLC26A10	65012	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001020658	PUM1	9698; 28997	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001024844	CD82	3732	www.methdb.de/ and/or www.mdanderson.org/
NM_001025205	AP2M1	1173	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001025604	ARRDC2	27106	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001033044	GLUL	2752	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001033056	GLUL	2752	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001033518	WIPI2	26100	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001033519	WIPI2	26100	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001033520	WIPI2	26100	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001033952	CALCA	796	www.methdb.de/ and/or www.mdanderson.org/
NM_001036	RYR3	6263	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001043	SLC6A2	6530	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001053	SSTR5	6755	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001059	TACR3	6870	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001063	TF	7018	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001100	ACTA1	58	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001109	ADAM8	101	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001176	ARHGDI3	398	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001186	BACH1	571	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001204	BMP2	659	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001228	CASP8	841	www.methdb.de/ and/or www.mdanderson.org/
NM_001250	CD40	958	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001257	CDH13	1012	www.methdb.de/ and/or www.mdanderson.org/
NM_001319	CSNK1G2	1455	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001325	CSTF2	1478	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001385	DPYS	1807; 55412	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001451	FOXF1	2294	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001454	FOXJ1	2302	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001458	FLNC	2318	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001480	GALR1	2587	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001538	HSF4	3299	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001553	IGFBP7	3490; 818325	www.methdb.de/ and/or www.mdanderson.org/
NM_001572	IRF7	3665	www.methdb.de/ and/or www.mdanderson.org/
NM_001628	AKR1B1	231	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001635	AMPH	273	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_001651	AQP5	362	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001718	BMP6	654	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_001753	CAV1	857	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_001753	CAV1	857	www.methdb.de/ and/or www.mdanderson.org/
NM_001768	CD8A	925	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001801	CDO1	1036	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001851	COL9A1	1297	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001883	CRHR2	1395	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001884	HAPLN1	1404	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001927	DES	1674	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001954	DDR1	780	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_001958	EEF1A2	1917	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001972	ELA2	1991	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001975	ENO2	2026	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_001989	EVX1	2128	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002007	FGF4	2249	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002012	FHIT	2272; 246734	www.methdb.de/ and/or www.mdanderson.org/
NM_002024	FMR1	2332	www.methdb.de/ and/or www.mdanderson.org/
NM_002065	GLUL	2752	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002110	HCK	3055	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002127	HLA-G	3135	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002148	HOXD10	3236	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002155	HSPA6	3310	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002191	INHA	3623	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_002212	ITGB4BP	3692	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002221	ITPKB	3707	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002235	KCNA6	3742	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002253	KDR	3791	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002344	LTK	4058	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002412	MGMT	4255	www.methdb.de/ and/or www.mdanderson.org/
NM_002457	MUC2	4583	www.methdb.de/ and/or www.mdanderson.org/
NM_002478	MYOD1	4654	www.methdb.de/ and/or www.mdanderson.org/
NM_002529	NTRK1	4914	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002588	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002658	PLAU	5328; 414236	www.methdb.de/ and/or www.mdanderson.org/
NM_002700	POU4F3	5459	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002807	PSMD1	5707; 7410	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002848	PTPRO	5800	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002873	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_002923	RGS2	5997	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003027	SH3GL3	6457	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003088	FSCN1	6624	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003097	SNRPN	6638; 8926; 145624; 8123; 63968; 3653	www.methdb.de/ and/or www.mdanderson.org/
NM_003149	STAC	6769	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003204	NFE2L1	4779	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_003219	TERT	7015	www.methdb.de/ and/or www.mdanderson.org/
NM_003238	TGFB2	7042	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003238	TGFB2	7042	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003246	THBS1	7057	www.methdb.de/ and/or www.mdanderson.org/
NM_003274	TMEM1	7109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003277	CLDN5	7122	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		7178; 51447;	
NM_003295	TPT1	2982	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003300	TRAF3	7187	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003391	WNT2	7472	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003392	WNT5A	7474	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003408	ZFP37	7539; 7551	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003417	ZNF264	9422	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003426	ZNF74	7625	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003435	ZNF134	7693	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003451	ZNF177	7730	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003474	ADAM12	8038	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003508	FZD9	8326	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003539	HIST1H4D	8360	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003540	HIST1H4F	8361	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003541	HIST1H4K	8362	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003546	HIST1H4L	8368	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003666	BLZF1	8548	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		5098; 26025; 56108; 56112; 9708;	
NM_003735	PCDHGC3	56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		5098; 26025; 56108; 56112; 9708;	
NM_003736	PCDHGC3	56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003775	EDG6	8698	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003777	DNAH11	8701; 9026	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003806	HRK	8739	www.methdb.de/ and/or www.mdanderson.org/
		8771; 51750;	
NM_003823	TNFRSF6B	10139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003888	ALDH1A2	8854	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003914	CCNA1	8900	www.methdb.de/ and/or www.mdanderson.org/
NM_003923	FOXH1	8928	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003984	SLC13A2	9058	www.methdb.de/ and/or www.mdanderson.org/
NM_003991	EDNRB	1910	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_003999	OSMR	9180	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004004	GJB2	2706	www.methdb.de/ and/or www.mdanderson.org/
NM_004064	CDKN1B	1027	www.methdb.de/ and/or www.mdanderson.org/
NM_004068	AP2M1	1173	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004102	FABP3	2170	www.methdb.de/ and/or www.mdanderson.org/
NM_004113	FGF12	2257	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004122	GHSR	2693	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004135	IDH3G	3421	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004181	UCHL1	7345	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004230	EDG5	9294	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004248	PRLHR	2834	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004267	CHST2	9435	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004291	CART	9607	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004297	GNA14	9630	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_004327	BCR	613; 26226	www.methdb.de/ and/or www.mdanderson.org/ DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004360	CDH1	999	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004360	CDH1	999	www.methdb.de/ and/or www.mdanderson.org/ DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004360	CDH1	999	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004378	CRABP1	1381	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004385	CSPG2	1462	www.methdb.de/ and/or www.mdanderson.org/
NM_004387	NKX2-5	1482	www.methdb.de/ and/or www.mdanderson.org/
NM_004394	DAP	1611	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004411	DYNC111	1780	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004441	EPHB1	2047	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004464	FGF5	2250	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004477	FRG1	2483	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004480	FUT8	2530	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004484	GPC3	2719	www.methdb.de/ and/or www.mdanderson.org/
NM_004525	LRP2	4036	www.methdb.de/ and/or www.mdanderson.org/
NM_004530	MMP2	4313	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004612	TGFBR1	7046	www.methdb.de/ and/or www.mdanderson.org/
NM_004621	TRPC6	7225	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004714	DYRK1B	9149	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004737	LARGE	9215	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004787	SLIT2	9353	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004817	TJP2	9414	www.methdb.de/ and/or www.mdanderson.org/
NM_004865	TBPL1	9519	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004887	CXCL14	9547	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004929	CALB1	793	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004936	CDKN2B	1030	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004936	CDKN2B	1030	www.methdb.de/ and/or www.mdanderson.org/ DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004938	DAPK1	1612	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_004975	KCNB1	3745	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004976	KCNC1	3746	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_004988	MAGEA1	4100	www.methdb.de/ and/or www.mdanderson.org/
NM_005032	PLS3	5358	www.methdb.de/ and/or www.mdanderson.org/
NM_005048	PTHR2	5746	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005073	SLC15A1	6564	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005100	AKAP12	9590	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005117	FGF19	9965	www.methdb.de/ and/or www.mdanderson.org/
NM_005117	FGF19	9965	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005157	ABL1	25	www.methdb.de/ and/or www.mdanderson.org/
NM_005159	ACTC	70	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005181	CA3	761	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005233	EPHA3	2042	www.methdb.de/ and/or www.mdanderson.org/
NM_005284	GPR6	2830	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005285	NPBWR1	2831	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005286	NPBWR2	2832	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005346	HSPA1B	3304	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005382	NEF3	4741	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005386	NNAT	4826	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005398	PPP1R3C	5507	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005427	TP73	7161	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_005437	NCOA4	8031	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_005523	HOXA11	3207	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005556	KRT7	3855	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005584	MAB21L1	4081	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_005638	SYBL1	6845	www.methdb.de/ and/or www.mdanderson.org/
NM_005668	ST8SIA4	7903	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005806	OLIG2	10215	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005825	RASGRP2	10235	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_005946	MT1A	4489	www.methdb.de/ and/or www.mdanderson.org/
NM_005959	MTNR1B	4544	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006000	TUBA1	7277; 84854	www.methdb.de/ and/or www.mdanderson.org/
NM_006019	TCIRG1	10312	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006041	HS3ST3B1	9953; 84815	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006043	HS3ST2	9956	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006053	TCIRG1	10312	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006142	SFN	2810	www.methdb.de/ and/or www.mdanderson.org/
NM_006158	NEFL	4747	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006158	NEFL	4747	www.methdb.de/ and/or www.mdanderson.org/
NM_006161	NEUROG1	4762	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006194	PAX9	5083	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006211	PENK	5179	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006306	SMC1L1	8243	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006307	SRPX	8406	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006463	STAMPB	10617	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006483	DYRK1B	9149	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006484	DYRK1B	9149	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006497	HIC1	3090	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_006497	HIC1	3090	www.methdb.de/ and/or www.mdanderson.org/
NM_006539	CACNG3	10368	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006587	CORIN	10699	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006614	CHL1	10752	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006735	HOXA2	3199	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006765	TUSC3	7991	www.methdb.de/ and/or www.mdanderson.org/
NM_006788	RALBP1	10928	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006874	ELF2	1998; 26472	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006898	HOXD3	3232	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_006917	RXRG	6258	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_007117	TRH	7200	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_007181	MAP4K1	11184	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_007182	RASSF1	11186	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_007182	RASSF1	11186	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_007182	RASSF1	11186	www.methdb.de/ and/or www.mdanderson.org/
NM_007197	FZD10	11211	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_007294	BRCA1	672	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_007294	BRCA1	672	www.methdb.de/ and/or www.mdanderson.org/
NM_007332	TRPA1	8989	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_007345	ZNF236	7776	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_007361	NID2	22795	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012200	B3GAT3	26229	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012202	GNG3	2785	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012261	C20orf103	24141	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012295	CABIN1	23523	www.methdb.de/ and/or www.mdanderson.org/
NM_012295	CABIN1	23523	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012301	MAGI2	9863	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012309	SHANK2	22941	www.methdb.de/ and/or www.mdanderson.org/
NM_012309	SHANK2	22941	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012399	PITPNB	23760	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_012444	SPO11	23626	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_012458	TIMM13	26517	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_013250	ZNF215	7762	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_013291	CPSF1	29894	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_013381	TRHDE	29953	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_013435	RAX	30062	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_013942	PAX3	5077	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014020	LR8	28959	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014080	DUOX2	50506	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014155	BTBD15	29068	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014228	SLC6A7	6534	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014234	HSD17B8	7923	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014325	CORO1C	23603	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014379	KCNV1	27012	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014386	PKD2L2	27039	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014459	PCDH17	27253; 144997	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014468	VENTX	27287	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014522	PCDH11X	27328; 83259	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014574	STRN3	29966	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014587	SOX8	30812	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014588	VSX1	30813	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014618	DBC1	1620	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014618	DBC1	1620	DNA Methylation And Cancer Therapy, Landes Bioscience 2005, ed. Moshe Szyf
NM_014631	SH3PXD2A	9644	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014653	KIAA0789	9671	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014676	PUM1	9698; 28997	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014710	GPRASP1	9737	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014724	ZNF96	9753	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014786	ARHGEF17	9828	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014817	KIAA0644	9865	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_014979	SV2C	22987	www.methdb.de/ and/or www.mdanderson.org/
NM_015002	FBXO21	23014	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015094	HIC2	23119	www.methdb.de/ and/or www.mdanderson.org/
NM_015101	GLT25D2	23127	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015163	TRIM9	114088	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015472	WWTR1	25937	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015507	EGFL6	25975	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015610	WIPI2	26100	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015641	TES	26136	www.methdb.de/ and/or www.mdanderson.org/
NM_015683	ARRDC2	27106	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015722	DRD1IP	50632	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_015920	RPS27L	51065	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016003	WIPI2	26100	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016135	ETV7	51513	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016157	TRO	7216	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016162	ING4	51147	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016179	TRPC4	7223	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016192	TMEFF2	23671	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016192	TMEFF2	23671	www.methdb.de/ and/or www.mdanderson.org/
NM_016223	PACSIN3	29763	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016229	CYB5R2	51700	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016301	ATPBD1C	51184	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016442	ARTS-1	51752	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016521	TFDP3	51270	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016535	ZNF581	51545	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016538	SIRT7	51547	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_016540	GPR83	10888	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016552	ANKMY1	51281	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016568	RLN3R1	51289	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016605	FAM53C	51307; 995	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016931	NOX4	50507	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016950	SPOCK3	50859	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_016954	TBX22	50945	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_017514	PLXNA3	55558	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_017649	CNNM2	54805	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_017729	EPS8L1	54869	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_017798	YTHDF1	54915	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_017844	ANKMY1	51281	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_017847	C1orf27	54953	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018061	PRPF38B	55119	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018074	FLJ10374	55702	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018129	PNPO	55163	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018135	MRPS18A	55168	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018197	ZFP64	55734	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018310	BRF2	55290	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018354	C20orf46	55321	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018401	STK32B	55351	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018431	DOK5	55816	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018898	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018899	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018901	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018906	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018911	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018920	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018925	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_018926	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018927	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018928	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018950	HLA-F	3134	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018976	SLC38A2	54407	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_018997	MRPS21	54460	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_019043	APBB1IP	54518	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_019102	HOXA5	3202	www.methdb.de/ and/or www.mdanderson.org/
NM_020166	MCCC1	56922	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020201	NT5M	56953	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020208	SLC6A20	54716	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020226	PRDM8	56978	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020230	PPAN	56342	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020348	CNNM1	26507	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020469	ABO	28	www.methdb.de/ and/or www.mdanderson.org/
NM_020549	CHAT	1103	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020630	RET	5979	www.methdb.de/ and/or www.mdanderson.org/
NM_020650	RCN3	57333	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020657	ZNF304	57343	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020660	CX36	57369	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020685	C3orf14	57415	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020815	PCDH10	57575	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020873	LRRN1	57633	www.methdb.de/ and/or www.mdanderson.org/
NM_020984	CHAT	1103	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020985	CHAT	1103	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020986	CHAT	1103	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_020999	NEUROG3	50674	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021032	FGF12	2257	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021101	CLDN1	9076	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021179	C1orf114	57821	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021193	HOXD12	3238	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021216	ZNF71	58491	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021257	NGB	58157	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021614	KCNN2	3781	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021911	GABRB2	2561	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021926	ALX4	60529	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_021956	GRIK2	2898	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022076	DUSP21	63904	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022088	ZFP64	55734	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022169	ABCG4	64137	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022405	SLC6A20	54716	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022443	MLF1	4291	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022468	MMP25	64386; 4328	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022469	GREM2	64388	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_022718	MMP25	64386;	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

		4328	
NM_022750	PARP12	64761	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_023926	ZNF447	65982	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024012	HTR5A	3361	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024046	CAMKV	79012	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024101	MLPH	79083	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024306	FA2H	79152	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024409	NPPC	4880	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024504	PRDM14	63978	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024593	EFCAB1	79645	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024600	C16orf30	79652	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024826	ASAP	79884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024882	C6orf155	79940	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024893	C20orf39	79953	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_024944	CHODL	140578	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025019	TUBA4	80086	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025058	TRIM46	80128	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025061	LRRC8E	80131	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025087	FLJ21511	80157	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025197	CDK5RAP3	80279	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025204	RP3-402G11.12	80305	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		80310; 414301	
NM_025208	PDGFD	414301	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025218	ULBP1	80329	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_025263	PRR3	80742	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030577	MGC10993	80775	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030667	PTPRO	5800	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030668	PTPRO	5800	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030669	PTPRO	5800	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030670	PTPRO	5800	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030671	PTPRO	5800	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030760	EDG8	53637	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		81563; 116492	
NM_030806	C1orf21	116492	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_030920	ANP32E	81611	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031277	RNF17	56163	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031283	TCF7L1	83439	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031424	C20orf55	83541	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031466	NIBP	83696	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031488	L3MBTL2	83746	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		56142; 56145; 56134; 56147; 56146; 56139	
NM_031497	PCDHA6	56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		56142; 56145; 56134; 56147; 56146; 56139	
NM_031856	PCDHA6	56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		56142; 56145; 56134; 56147; 56146; 56139	
NM_031859	PCDHA6	56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
		56142; 56145; 56134; 56147;	
NM_031860	PCDHA6	56147;	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

		56146; 56139	
NM_031882	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031883	PCDHA6	56142; 56145; 56134; 56147; 56146; 56139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031901	MRPS21	54460	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031912	SYT15	83849	www.methdb.de/ and/or www.mdanderson.org/
NM_031922	REPS1	85021	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031934	RAB34	83871	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_031994	RNF17	56163	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032034	SLC4A11	83959	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032087	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032094	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032098	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032099	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032100	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032101	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032109	OTP	23440	www.methdb.de/ and/or www.mdanderson.org/
NM_032134	QRICH2	84074	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032140	C16orf48	84080	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032192	PPP1R1B	84152	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032256	TMEM117	84216	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032303	HSDL2	84263	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032391	PRAC	84366	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM_032402	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032403	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032406	PCDHGC3	5098; 26025; 56108; 56112; 9708; 56109	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032411	ECRG4	84417	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032412	ORF1-FL49	84418	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032603	LOXL3	84695	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032625	C7orf13	129790	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032803	SLC7A3	84889	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032825	ZNF382	84911	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032838	ZNF566	84924	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032883	C20orf100	84969	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032918	RERG	85004	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032945	TNFRSF6B	8771; 51750; 10139	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032961	PCDH10	57575	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032967	PCDH11X	27328; 83259	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032968	PCDH11X	27328; 83259	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_032969	PCDH11X	27328; 83259	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033126	PSKH2	85481	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033135	PDGFD	80310; 414301	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033143	FGF5	2250	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033224	PURB	5814	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033302	ADRA1A	148	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033303	ADRA1A	148	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033304	ADRA1A	148	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033445	HIST3H2A	92815	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_033624	FBXO21	23014	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_052902	STK11IP	114790	www.methdb.de/ and/or www.mdanderson.org/
NM_052954	CYYR1	116159	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_052961	SLC26A8	116369	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_052978	TRIM9	114088	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_054021	GPR101	83550	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_054108	HRASLS5	117245	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_058165	MOGAT1	116255	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_078485	COL9A1	1297	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_080552	SLC32A1	140679	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_080617	CBLN4	140689	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_080671	KCNE4	23704	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_080742	B3GAT2	135152	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_130773	CNTNAP5	129684	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_130900	RAET1L	154064	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM_133180	EPS8L1	54869	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM 133266	SHANK2	22941	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133338	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133339	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133340	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133341	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133342	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133343	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133344	RAD17	5884	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133489	SLC26A10	65012	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133493	CD109	135228	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 133642	LARGE	9215	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 138290	RPIB9	154661	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 138718	SLC26A8	116369	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 138996	CNTNAP5	129684	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 139204	EPS8L1	54869	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 139316	AMPH	273	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 144497	AKAP12	9590	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 144725	FLJ25439	153657	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM 145725	TRAF3	7187	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 145726	TRAF3	7187	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 152562	CDCA2	157313	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM 152854	CD40	958	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 153819	RASGRP2	10235	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 170696	ALDH1A2	8854	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 170697	ALDH1A2	8854	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 170775	KCNN2	3781	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 171827	CD8A	925	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 172337	OTX2	5015	www.methdb.de/ and/or www.mdanderson.org/
NM 173479	LOC126248	126248	www.methdb.de/ and/or www.mdanderson.org/
NM 174869	IDH3G	3421	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 175052	ST8SIA4	7903	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 175611	GRIK1	2897	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 175709	CBX7	23492	www.methdb.de/ and/or www.mdanderson.org/
NM 175768	GRIK2	2898	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 176095	CDK5RAP3	80279	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 176096	CDK5RAP3	80279	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 177555	TRO	7216	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 177556	TRO	7216	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 177557	TRO	7216	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 177558	TRO	7216	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 177959	DOK5	55816	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 178154	FUT8	2530	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 178155	FUT8	2530	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 178156	FUT8	2530	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 178157	FUT8	2530	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181457	PAX3	5077	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181458	PAX3	5077	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181459	PAX3	5077	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181460	PAX3	5077	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181461	PAX3	5077	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181466	ITGB4BP	3692	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181467	ITGB4BP	3692	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181468	ITGB4BP	3692	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181469	ITGB4BP	3692	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181505	PPP1R1B	84152	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 181657	LTB4R	1241	www.methdb.de/ and/or www.mdanderson.org/
NM 181689	NNAT	4826	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 182609	ZNF677	342926	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM 198265	SPO11	23626	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153

NM 198287	ING4	51147	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 198407	GHSR	2693	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 198570	UNQ739	375567	www.methdb.de/ and/or www.mdanderson.org/
NM 198849	LOC283514	283514	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM 199051	FAM5C	339479	www.methdb.de/ and/or www.mdanderson.org/
NM 199076	CNNM2	54805	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 199077	CNNM2	54805	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 199231	GDNF	2668	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 199234	GDNF	2668	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 199425	VSX1	30813	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 199426	ZFP64	55734	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 199427	ZFP64	55734	www.methdb.de/ and/or www.mdanderson.org/
NM 199427	ZFP64	55734	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 201647	STAMPB	10617	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 201999	ELF2	1998; 26472	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 206827	RASL11A	387496	Weber <i>et al.</i> Nature Genetics 37(8), 2005, 853-862
NM 206866	BACH1	571	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 206961	LTK	4058	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 207121	C20orf55	83541	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NM 213622	STAMPB	10617	Keshet <i>et al.</i> Nature Genetics 38(2), 2006, 149-153
NP 536846			www.methdb.de/ and/or www.mdanderson.org/
NR 002196			www.methdb.de/ and/or www.mdanderson.org/

See also Ilana Keshet, *et al.*, Nature Genetics 38, 149–153 (1 February 2006) and Gerd P Pfeifer, *et al.*, Expert Opinion on Medical Diagnostics, September 2007, Vol. 1, No. 1, Pages 99-108, each of which is incorporated herein by reference.

- 5 As used herein, the term "adenoma" refers to a benign tumor of glandular origin. Although these growths are benign, over time they may progress to become malignant. As used herein the term "colorectal adenoma" refers to a benign colorectal tumor in which the cells form recognizable glandular structures or in which the cells are clearly derived from glandular epithelium.
- 10 The term "amplifying" or "amplification" in the context of nucleic acids refers to the production of multiple copies of a polynucleotide, or a portion of the polynucleotide, typically starting from a small amount of the polynucleotide (*e.g.*, a single polynucleotide molecule), where the amplification products or amplicons are generally detectable. Amplification of polynucleotides encompasses a variety of chemical and enzymatic
- 15 processes. The generation of multiple DNA copies from one or a few copies of a target or template DNA molecule during a polymerase chain reaction (PCR) or a ligase chain reaction (LCR; see, *e.g.*, U.S. Patent No. 5,494,810; herein incorporated by reference in its entirety) are forms of amplification. Additional types of amplification include, but are not limited to, allele-specific PCR (see, *e.g.*, U.S. Patent No. 5,639,611; herein incorporated by reference in
- 20 its entirety), assembly PCR (see, *e.g.*, U.S. Patent No. 5,965,408; herein incorporated by reference in its entirety), helicase-dependent amplification (see, *e.g.*, U.S. Patent No.

7,662,594; herein incorporated by reference in its entirety), Hot-start PCR (see, *e.g.*, U.S. Patent Nos. 5,773,258 and 5,338,671; each herein incorporated by reference in their entireties), intersequence-specific PCR, inverse PCR (see, *e.g.*, Triglia, *et al.* (1988) *Nucleic Acids Res.*, 16:8186; herein incorporated by reference in its entirety), ligation-mediated PCR (see, *e.g.*, Guilfoyle, R. *et al.*, *Nucleic Acids Research*, 25:1854-1858 (1997); U.S. Patent No. 5,508,169; each of which are herein incorporated by reference in their entireties), methylation-specific PCR (see, *e.g.*, Herman, *et al.*, (1996) *PNAS* 93(13) 9821-9826; herein incorporated by reference in its entirety), miniprimer PCR, multiplex ligation-dependent probe amplification (see, *e.g.*, Schouten, *et al.*, (2002) *Nucleic Acids Research* 30(12): e57; herein incorporated by reference in its entirety), multiplex PCR (see, *e.g.*, Chamberlain, *et al.*, (1988) *Nucleic Acids Research* 16(23) 11141-11156; Ballabio, *et al.*, (1990) *Human Genetics* 84(6) 571-573; Hayden, *et al.*, (2008) *BMC Genetics* 9:80; each of which are herein incorporated by reference in their entireties), nested PCR, overlap-extension PCR (see, *e.g.*, Higuchi, *et al.*, (1988) *Nucleic Acids Research* 16(15) 7351-7367; herein incorporated by reference in its entirety), real time PCR (see, *e.g.*, Higuchi, *et al.*, (1992) *Biotechnology* 10:413-417; Higuchi, *et al.*, (1993) *Biotechnology* 11:1026-1030; each of which are herein incorporated by reference in their entireties), reverse transcription PCR (see, *e.g.*, Bustin, S.A. (2000) *J. Molecular Endocrinology* 25:169-193; herein incorporated by reference in its entirety), solid phase PCR, thermal asymmetric interlaced PCR, and Touchdown PCR (see, *e.g.*, Don, *et al.*, *Nucleic Acids Research* (1991) 19(14) 4008; Roux, K. (1994) *Biotechniques* 16(5) 812-814; Hecker, *et al.*, (1996) *Biotechniques* 20(3) 478-485; each of which are herein incorporated by reference in their entireties). Polynucleotide amplification also can be accomplished using digital PCR (see, *e.g.*, Kalinina, *et al.*, *Nucleic Acids Research*. 25; 1999-2004, (1997); Vogelstein and Kinzler, *Proc Natl Acad Sci USA*. 96; 9236-41, (1999); International Patent Publication No. WO05023091A2; US Patent Application Publication No. 20070202525; each of which are incorporated herein by reference in their entireties).

The term "polymerase chain reaction" ("PCR") refers to the method of K.B. Mullis U.S. Patent Nos. 4,683,195, 4,683,202, and 4,965,188, that describe a method for increasing the concentration of a segment of a target sequence in a mixture of genomic DNA without cloning or purification. This process for amplifying the target sequence consists of introducing a large excess of two oligonucleotide primers to the DNA mixture containing the desired target sequence, followed by a precise sequence of thermal cycling in the presence of a DNA polymerase. The two primers are complementary to their respective strands of the

double stranded target sequence. To effect amplification, the mixture is denatured and the primers then annealed to their complementary sequences within the target molecule. Following annealing, the primers are extended with a polymerase so as to form a new pair of complementary strands. The steps of denaturation, primer annealing, and polymerase extension can be repeated many times (*i.e.*, denaturation, annealing and extension constitute one "cycle"; there can be numerous "cycles") to obtain a high concentration of an amplified segment of the desired target sequence. The length of the amplified segment of the desired target sequence is determined by the relative positions of the primers with respect to each other, and therefore, this length is a controllable parameter. By virtue of the repeating aspect of the process, the method is referred to as the "polymerase chain reaction" ("PCR"). Because the desired amplified segments of the target sequence become the predominant sequences (in terms of concentration) in the mixture, they are said to be "PCR amplified" and are "PCR products" or "amplicons."

As used herein, the term "nucleic acid detection assay" refers to any method of determining the nucleotide composition of a nucleic acid of interest. Nucleic acid detection assay include but are not limited to, DNA sequencing methods, probe hybridization methods, structure specific cleavage assays (*e.g.*, the INVADER assay, (Hologic, Inc.) and are described, *e.g.*, in U.S. Patent Nos. 5,846,717, 5,985,557, 5,994,069, 6,001,567, 6,090,543, and 6,872,816; Lyamichev et al., *Nat. Biotech.*, 17:292 (1999), Hall et al., *PNAS, USA*, 97:8272 (2000), and US 2009/0253142, each of which is herein incorporated by reference in its entirety for all purposes); enzyme mismatch cleavage methods (*e.g.*, Variagenics, U.S. Pat. Nos. 6,110,684, 5,958,692, 5,851,770, herein incorporated by reference in their entireties); polymerase chain reaction; branched hybridization methods (*e.g.*, Chiron, U.S. Pat. Nos. 5,849,481, 5,710,264, 5,124,246, and 5,624,802, herein incorporated by reference in their entireties); rolling circle replication (*e.g.*, U.S. Pat. Nos. 6,210,884, 6,183,960 and 6,235,502, herein incorporated by reference in their entireties); NASBA (*e.g.*, U.S. Pat. No. 5,409,818, herein incorporated by reference in its entirety); molecular beacon technology (*e.g.*, U.S. Pat. No. 6,150,097, herein incorporated by reference in its entirety); E-sensor technology (Motorola, U.S. Pat. Nos. 6,248,229, 6,221,583, 6,013,170, and 6,063,573, herein incorporated by reference in their entireties); cycling probe technology (*e.g.*, U.S. Pat. Nos. 5,403,711, 5,011,769, and 5,660,988, herein incorporated by reference in their entireties); Dade Behring signal amplification methods (*e.g.*, U.S. Pat. Nos. 6,121,001, 6,110,677, 5,914,230, 5,882,867, and 5,792,614, herein incorporated by reference in their entireties);

ligase chain reaction (*e.g.*, Barnay Proc. Natl. Acad. Sci USA 88, 189-93 (1991)); and sandwich hybridization methods (*e.g.*, U.S. Pat. No. 5,288,609, herein incorporated by reference in its entirety).

As used herein, the terms "complementary" or "complementarity" used in reference to
5 polynucleotides (*i.e.*, a sequence of nucleotides) refers to polynucleotides related by the base-pairing rules. For example, the sequence "5'-A-G-T-3'," is complementary to the sequence "3'-T-C-A-5'." Complementarity may be "partial," in which only some of the nucleic acids' bases are matched according to the base pairing rules. Or, there may be "complete" or "total" complementarity between the nucleic acids. The degree of complementarity between nucleic
10 acid strands has significant effects on the efficiency and strength of hybridization between nucleic acid strands. This is of particular importance in amplification reactions, as well as detection methods that depend upon binding between nucleic acids.

As used herein, the term "primer" refers to an oligonucleotide, whether occurring naturally, as in a purified restriction digest, or produced synthetically, that is capable of
15 acting as a point of initiation of synthesis when placed under conditions in which synthesis of a primer extension product that is complementary to a nucleic acid strand is induced (*e.g.*, in the presence of nucleotides and an inducing agent such as a biocatalyst (*e.g.*, a DNA polymerase or the like). The primer is typically single stranded for maximum efficiency in amplification, but may alternatively be partially or completely double stranded. The portion
20 of the primer that hybridizes to a template nucleic acid is sufficiently long to prime the synthesis of extension products in the presence of the inducing agent. The exact lengths of the primers will depend on many factors, including temperature, source of primer and the use of the method. Primers may comprise labels, tags, capture moieties, *etc.*

As used herein, the term "nucleic acid molecule" refers to any nucleic acid containing
25 molecule, including but not limited to, DNA or RNA. The term encompasses sequences that include any of the known base analogs of DNA and RNA including, but not limited to, 4 acetylcytosine, 8-hydroxy-N6-methyladenosine, aziridinylcytosine, pseudoisocytosine, 5-(carboxyhydroxyl-methyl) uracil, 5-fluorouracil, 5-bromouracil, 5-carboxymethylaminomethyl-2-thiouracil, 5-carboxymethyl-aminomethyluracil,
30 dihydrouracil, inosine, N6-isopentenyladenine, 1-methyladenine, 1-methylpseudo-uracil, 1-methylguanine, 1-methylinosine, 2,2-dimethyl-guanine, 2-methyladenine, 2-methylguanine, 3-methyl-cytosine, 5-methylcytosine, N6-methyladenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxy-amino-methyl-2-thiouracil, beta-D-mannosylqueosine,

5'-methoxycarbonylmethyluracil, 5-methoxyuracil, 2-methylthio-N- isopentenyladenine, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, oxybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, N-uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid, pseudouracil, queosine, 2-thiocytosine, and 2,6-diaminopurine.

As used herein, the term "nucleobase" is synonymous with other terms in use in the art including "nucleotide," "deoxynucleotide," "nucleotide residue," "deoxynucleotide residue," "nucleotide triphosphate (NTP)," or deoxynucleotide triphosphate (dNTP).

An "oligonucleotide" refers to a nucleic acid that includes at least two nucleic acid monomer units (*e.g.*, nucleotides), typically more than three monomer units, and more typically greater than ten monomer units. The exact size of an oligonucleotide generally depends on various factors, including the ultimate function or use of the oligonucleotide. To further illustrate, oligonucleotides are typically less than 200 residues long (*e.g.*, between 15 and 100), however, as used herein, the term is also intended to encompass longer polynucleotide chains. Oligonucleotides are often referred to by their length. For example a 24 residue oligonucleotide is referred to as a "24-mer". Typically, the nucleoside monomers are linked by phosphodiester bonds or analogs thereof, including phosphorothioate, phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoranilidate, phosphoramidate, and the like, including associated counterions, *e.g.*, H⁺, NH₄⁺, Na⁺, and the like, if such counterions are present. Further, oligonucleotides are typically single-stranded. Oligonucleotides are optionally prepared by any suitable method, including, but not limited to, isolation of an existing or natural sequence, DNA replication or amplification, reverse transcription, cloning and restriction digestion of appropriate sequences, or direct chemical synthesis by a method such as the phosphotriester method of Narang *et al.* (1979) *Meth Enzymol.* 68: 90-99; the phosphodiester method of Brown *et al.* (1979) *Meth Enzymol.* 68: 109-151; the diethylphosphoramidite method of Beaucage *et al.* (1981) *Tetrahedron Lett.* 22: 1859-1862; the triester method of Matteucci *et al.* (1981) *J Am Chem Soc.* 103:3185-3191; automated synthesis methods; or the solid support method of U.S. Pat. No. 4,458,066, entitled "PROCESS FOR PREPARING POLYNUCLEOTIDES," issued Jul. 3, 1984 to Caruthers *et al.*, or other methods known to those skilled in the art. All of these references are incorporated by reference.

A "sequence" of a biopolymer refers to the order and identity of monomer units (*e.g.*, nucleotides, amino acids, *etc.*) in the biopolymer. The sequence (*e.g.*, base sequence) of a nucleic acid is typically read in the 5' to 3' direction.

The term "wild-type" refers to a gene or gene product that has the characteristics of that gene or gene product when isolated from a naturally occurring source. A wild-type gene is that which is most frequently observed in a population and is thus arbitrarily designed the "normal" or "wild-type" form of the gene. In contrast, the terms "modified," "mutant," and "variant" refer to a gene or gene product that displays modifications in sequence and or functional properties (*i.e.*, altered characteristics) when compared to the wild-type gene or gene product. It is noted that naturally occurring mutants can be isolated; these are identified by the fact that they have altered characteristics when compared to the wild-type gene or gene product.

As used herein, the term "gene" refers to a nucleic acid (*e.g.*, DNA) sequence that comprises coding sequences necessary for the production of a polypeptide, precursor, or RNA (*e.g.*, rRNA, tRNA). The polypeptide can be encoded by a full length coding sequence or by any portion of the coding sequence so long as the desired activity or functional properties (*e.g.*, enzymatic activity, ligand binding, signal transduction, immunogenicity, *etc.*) of the full-length or fragment polypeptide are retained. The term also encompasses the coding region of a structural gene and the sequences located adjacent to the coding region on both the 5' and 3' ends for a distance of about 1 kb or more on either end such that the gene corresponds to the length of the full-length mRNA. Sequences located 5' of the coding region and present on the mRNA are referred to as 5' non-translated sequences. Sequences located 3' or downstream of the coding region and present on the mRNA are referred to as 3' non-translated sequences. The term "gene" encompasses both cDNA and genomic forms of a gene. A genomic form or clone of a gene contains the coding region interrupted with non-coding sequences termed "introns" or "intervening regions" or "intervening sequences." Introns are segments of a gene that are transcribed into nuclear RNA (*e.g.*, hnRNA); introns may contain regulatory elements (*e.g.*, enhancers). Introns are removed or "spliced out" from the nuclear or primary transcript; introns therefore are absent in the messenger RNA (mRNA) transcript. The mRNA functions during translation to specify the sequence or order of amino acids in a nascent polypeptide.

In addition to containing introns, genomic forms of a gene may also include sequences located on both the 5' and 3' end of the sequences that are present on the RNA transcript. These sequences are referred to as "flanking" sequences or regions (these flanking sequences are located 5' or 3' to the non-translated sequences present on the mRNA transcript). The 5' flanking region may contain regulatory sequences such as promoters and

enhancers that control or influence the transcription of the gene. The 3' flanking region may contain sequences that direct the termination of transcription, post-transcriptional cleavage and polyadenylation.

As used herein, the terms "multimethylation," "series methylation" and "specific
5 methylation" are used interchangeably to refer to defined combinations of CpG sites or loci in a marker sequence must be methylated to call that sequence methylated in a coordinate or multimethylation assay. For example, a specific methylation assay of the CpG sites for BMP3 might require that the CpG positions at 23, 34, 53, 61, 70, and 74, numbered by reference to Figures 1A and 1B, all be methylated in order for a sample to be classified as
10 methylated at the BMP3 marker. Specific methylation of BMP3 is not limited to this set of particular loci, but may include more, fewer, or a different collection of CpG loci. The CpG loci selected for co-analysis in a multimethylation assay are preferably selected. *e.g.*, by analysis of normal (non-adenoma, non-cancer) samples to identify CpG methylation combinations that are less frequently represented in normal samples. In preferred
15 embodiments, combinations of methylation sites are selected to produce good signal-to-noise in cancer and adenoma samples (*i.e.*, the mean multimethylation at a particular combination of loci in cancer samples divided by the mean multimethylation in at those loci in normal samples is high).

As used herein, the terms "individual" and "average" methylation are used
20 interchangeably to refer to analyses in which each CpG locus is analyzed individually, such that all molecules in which that base is methylated are included in a count, regardless of the methylation status of other loci, *e.g.*, in the same marker. Generally, the methylation percentages of all the loci in a marker/region are then averaged, to produce a percent methylation figure for that marker.

As used herein, the term "kit" refers to any delivery system for delivering materials.
25 In the context of reaction assays, such delivery systems include systems that allow for the storage, transport, or delivery of reaction reagents (*e.g.*, oligonucleotides, enzymes, *etc.* in the appropriate containers) and/or supporting materials (*e.g.*, buffers, written instructions for performing the assay *etc.*) from one location to another. For example, kits include one or
30 more enclosures (*e.g.*, boxes) containing the relevant reaction reagents and/or supporting materials. As used herein, the term "fragmented kit" refers to a delivery systems comprising two or more separate containers that each contain a subportion of the total kit components. The containers may be delivered to the intended recipient together or separately. For

example, a first container may contain an enzyme for use in an assay, while a second container contains oligonucleotides. The term "fragmented kit" is intended to encompass kits containing Analyte specific reagents (ASR's) regulated under section 520(e) of the Federal Food, Drug, and Cosmetic Act, but are not limited thereto. Indeed, any delivery system comprising two or more separate containers that each contains a subportion of the total kit components are included in the term "fragmented kit." In contrast, a "combined kit" refers to a delivery system containing all of the components of a reaction assay in a single container (*e.g.*, in a single box housing each of the desired components). The term "kit" includes both fragmented and combined kits.

As used herein, the term "information" refers to any collection of facts or data. In reference to information stored or processed using a computer system(s), including but not limited to internets, the term refers to any data stored in any format (*e.g.*, analog, digital, optical, *etc.*). As used herein, the term "information related to a subject" refers to facts or data pertaining to a subject (*e.g.*, a human, plant, or animal). The term "genomic information" refers to information pertaining to a genome including, but not limited to, nucleic acid sequences, genes, allele frequencies, RNA expression levels, protein expression, phenotypes correlating to genotypes, *etc.* "Allele frequency information" refers to facts or data pertaining to allele frequencies, including, but not limited to, allele identities, statistical correlations between the presence of an allele and a characteristic of a subject (*e.g.*, a human subject), the presence or absence of an allele in an individual or population, the percentage likelihood of an allele being present in an individual having one or more particular characteristics, *etc.*

DESCRIPTION OF THE DRAWINGS

Figures 1A and 1B provide sequence and CpG information for exemplary marker regions used in the present analysis. For each target gene, the native sequence of the region is shown in the top line. Unmethylated C-residues that would be converted by bisulfite and amplification to Ts are shown as T residues. Candidate methylation positions are shown boxed. Reference numbering for base and CpG positions is shown above each native sequence. Primer locations for amplification are shown as a row of underlined base positions.

Figures 2 A-J provide tables showing analyses of normal, adenoma, and cancer samples in which the average methylation was determined at each of the indicated CpG positions, in the indicated marker regions. For each marker, the numbered CpG positions are as indicated by the reference numbers in Figures 1A and 1B. The Mean methylation at each specific locus is shown at the bottom of each column for normal, adenoma, and cancer samples. The ratio of normal/mutant methylation for each locus (a methylation ratio at each locus) is shown at the bottom of each column of the Adenoma and Cancer sample data. The Mean columns on the right of each table indicate the average of methylation across all indicated CpG loci for each of the samples. The Mean and SD values across all normal samples at all loci are as indicated below each table of values from normal samples.

Figures 3 A-I provide tables showing the analyses of normal, adenoma, and cancer samples in which average methylation across all of the CpG loci indicated in Figures 2A-J were calculated for each marker in each sample. For the normal samples in Fig 3A, the average, standard deviation and the mean plus 2 or 3 standard deviations for each marker are indicated. For the adenoma and cancer samples, shaded cells in Figs. 3B and 3C indicate a positive result, reflected as an average methylation value for that marker that is greater than the mean methylation + 3 standard deviations determined for that marker in the normal samples.

Figures 3D and 3E show the calculated effect of a 20-fold dilution of adenoma and cancer DNA into normal DNA, Figs. 3F and 3G show a calculated 10-fold dilution, and Figs. 3H and 3I show a calculated 5-fold dilution. In each of the calculated dilutions, the average methylation for a marker is divided by the 20, 10, or 5, added to the mean methylation of the normal DNA for that marker. Shaded cells in Figs. 3D-3I indicate an average methylation value for that marker that is greater than the mean methylation + 2 standard deviations (specificity of 97.5%) determined for that marker in the normal samples.

Below each of Figures 3B-3I, the percentage of positive values for each marker in the sample type and dilution for that panel are indicated. The percentage of samples giving a positive signal for at least one of the Vimentin, BMP3, Septin 9 and TFPI2 markers are indicated at the bottom of each panel.

Figures 4A and 4B provide sequence and CpG information for exemplary genes used in the present analysis. The CpG loci in each marker gene included in the defined subsets of

CpG loci for coordinate methylation analysis in colorectal adenoma and cancer samples are shown with a black background and in white typeface.

Figures 5 A-I provide tables showing the analyses of normal, adenoma, and cancer samples in which methylation was determined at each of the indicated CpG positions in the indicated marker regions (*i.e.*, samples were assayed for the percentage of DNA copies that displayed methylation at *all* of the CpG loci in the defined subset). Each marker was tested at each of the CpG loci in the defined subsets indicated in Figures 4A and 4B and the percentage methylation data reflects the percentage of marker copies having methylation at all of the tested CpG loci (coordinate methylation or "multimethylation" analysis). For the normal samples in Fig. 5A, the mean multimethylation, standard deviation and the mean plus 2 or 3 standard deviations for each marker are indicated. For the adenoma and cancer samples, shaded cells in Figs. 5B and 5C indicate a positive result, reflected as multimethylation value for that marker that is greater than the mean multimethylation + 3 standard deviations determined for that marker in the normal samples.

Figures 5D and 5E show the calculated effect of a 20-fold dilution of adenoma and cancer DNA into normal DNA, Figs. 5F and 5G show a calculated 10-fold dilution, and 5H and 5I show a calculated 5-fold dilution. In each of the calculated dilutions, the average multimethylation for a marker is divide by the 20, 10, or 5, added to the mean multimethylation of the normal DNA for that marker. Shaded cells in Figs. 5D-5I indicate an average multimethylation value for that marker that is greater than the mean multimethylation + 2 standard deviations (specificity of 97.5%) determined for that marker in the normal samples.

Below each of Figs. 5B-5I, the percentage of positive values for each marker in the sample type and dilution for that panel are indicated. The percentage of samples giving a positive signal for at least one of the vimentin, BMP3, Septin 9 and TFPI2 markers are indicated at the bottom of each panel.

Figure 6 shows a table and graph comparing the percent positive values calculated for each marker in adenoma and cancer samples, as indicated, using either individual/average methylation or multimethylation analysis methods to test each of the indicated markers, at each of the indicated calculated dilutions.

Figure 7 shows a table and graph comparing the percent positive values determined in adenoma and cancer samples, determined using the four markers with the lowest mean

background in these samples (vimentin, BMP3, Septin9, TFPI2), using either the individual/average methylation or the multimethylation analysis method, at each of the indicated calculated dilutions into normal DNA.

DETAILED DESCRIPTION OF THE INVENTION

5 Embodiments of the invention are described in this summary, and in the Summary of the Invention, above, which is incorporated here by reference. Although the invention has been described in connection with specific embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments.

10 The present invention relates to methods and compositions for determination of, and uses of, specific methylation patterns indicative of adenoma and carcinoma. In particular, the invention relates to analysis of defined subsets of CpG loci that are coordinately methylated in DNAs from cancer and adenoma samples, methods for identifying coordinately methylated loci, and methods of using analysis of coordinately methylated loci in one or more markers or regions in the design of assays for adenoma and cancer having improved sensitivity and
15 specificity.

 The present invention relates to the observation that, within marker nucleic acids for which methylation status is indicative of cellular status, *e.g.*, cancerous, pre-cancerous, normal, *etc.*, a subset of the individual methylation loci, *e.g.*, CpG loci, in non-normal cells generally displays a greater degree of methylation relative to the background levels of
20 methylation observed at corresponding loci in normal cells, while other methylation loci in the non-normal cells may exhibit levels of methylation that are closer to background levels. In some embodiments, the degree of methylation observed for a particular locus in plurality of cancerous or pre-cancerous cells relative to normal cells is expressed as a methylation ratio.

25 Some embodiments of the present invention relate to screening known or suspected marker genes to identify specific methylation loci that exhibit greater ratios of disease-associated methylation relative to background methylation, as compared to other marker genes or other loci in the same marker gene. In some preferred embodiments, the present invention relates to coordinate methylation analysis, to measure the degree to which a marker
30 molecule or a sample exhibits methylation at all of a plurality of selected loci.

 The present invention relates to analyzing methylation statuses of a defined set of individual CpG loci in methylation markers (or target regions within such markers) in a

significant enough number of individual DNA molecules in adenoma samples or cancer samples to identify defined subsets of CpG loci that have advantageous methylation ratios compared to other loci in the same adenoma or cancer samples. A defined subset of CpG loci that have advantageous methylation ratios in a sample may comprise the entirety of a set of
5 CpG loci in a particular marker or target region of a marker, or it may be fewer than all of the CpG loci in the characterized region of the marker.

Conventional methods of analyzing methylation status of a marker generally involve analysis of a mixed population of molecules. For example, amplification of a marker nucleic acid from a sample generally produces a mixture of amplicons coming from many copies of a
10 target molecule. If the amplification conditions are not selective for a gene variant, the amplicon product contains a mixture of the variant and the normal or wildtype DNA. Even if primers are specific for a mutation or for a particular methylation site, when DNA is amplified from many copies of target DNA derived from many cells, there can be heterogeneity in other base positions in the resulting amplicon. If these mixed amplicons are
15 sequenced directly, the resulting sequence traces reveal the consensus sequence of the mixed population, and particular sequences or mutations present in a small portion of the population are essentially undetectable. Although some researchers have sequenced individual clones from such amplifications to examine sequence information for individual molecules from the mixture, only small numbers of molecules have been analyzed and the data gathered did not
20 suggest any that any specific loci within the markers predictably exhibited advantageous methylation ratios compared to other methylated loci within the same target, or that coordinated analysis of loci having advantageous methylation ratios could be useful in improving the specificity and sensitivity of assays in detection of neoplasms. An aspect of the present invention is based on the observation that collecting methylation ratio information
25 from a very large number of individual molecules in both normal and non-normal samples reveals that some methylation loci in marker regions or sequences exhibit a greater degree of methylation in non-normal cells compared to background (methylation at the same loci in normal cells) than do other individual loci in the same marker region or gene. These loci in non-normal sequences that have a greater level of methylation compared to background can
30 be viewed as being particularly advantageous in that they are easier to identify over the background level of methylation observed in normal cells. One aspect of this advantage is that analysis of these particular loci permits identification of cancer-associated methylation with more sensitivity, and in a greater background of normal cells.

The present invention also relates to the observation that coordinated analysis of multiple loci provides a significantly enhanced level of sensitivity in the identification of cancerous or precancerous cells, especially in samples that may also comprise a significant number of normal cells. For example, Figure 6 compares the sensitivity of detecting adenoma and cancer cells. For each of the indicated marker genes, the methylation was either determined as an average across the marker region (*e.g.*, the mean methylation in the vimentin marker across all of loci 26, 37, 40, 45, 52, 54, 59, 63, and 74; see Figures 3A-I), indicated as "Individual" average methylation, or as a percentage of molecules displaying methylation at *all* of a subset of selected loci (*e.g.*, methylation in the vimentin marker at all three of loci 37, 40 and 45; see Figures 5A-I), *i.e.*, coordinate methylation analysis of multiple individual loci, indicated as "multi". The sensitivities for the same samples are also shown in calculated 5, 10 or 20-fold dilutions into normal DNA. Figure 6 shows that, while assay sensitivities may be similar in DNA analyzed directly from tissue without dilution, as the amount background from normal DNA is increased at the larger dilutions, the coordinate methylation analysis is shown to be far more sensitive than the average methylation analysis. For example, in the analysis of Septin9, the adenoma and cancer samples can be detected above background only in the undiluted and 5-fold dilution profiles when average methylation across the marker is analyzed, while these same samples can be detected with over about 69-74% sensitivity at 20-fold dilution, and 90-93% sensitivity at 10-fold dilution, when coordinate methylation analysis of loci 37, 40 and 45 is used.

In some embodiments, the present invention provides a method for designing a methylation assay to identify a disease state, comprising I) selecting at least one sequence for analysis; II) determining the methylation status of a plurality of loci in the sequence in a population of normal cells and a population of non-normal cells to determine an average rate of methylation for each of the plurality of loci each both normal and non-normal cells; and III) identifying at least two loci in said plurality of loci having advantageous methylation ratios.

I. Selection of sequence(s). Methylation markers associated with particular disease states have been identified for a number of disease states. For example, colorectal neoplasm-specific marker include, *e.g.*, bmp-3, bmp-4, SFRP2, vimentin, septin9, ALX4, EYA4, TFPI2, NDRG4, FOXE1, long DNA, BAT-26, K-ras, APC, melanoma antigen gene, p53, BRAF, and PIK3CA. Additional markers include but are not limited those in Table 1, above. Analysis of candidate methylation loci to identify those with advantageous methylation ratios

may comprise analysis of every locus in a target sequence (*e.g.*, every CpG) or it may comprise analysis of a subset of the methylation loci. In some embodiments, CpGs are selected for analysis by their location in particular methylation hotspots, while in other embodiments, loci for analysis may be conveniently located with respect to primer binding sites or other sequence features. Figure 1A provides an exemplary selection of marker neoplasm-associated markers with each of the C residues in the CpG loci indicated by a box. For each target gene, the native sequence of the region is shown in the top line and the sequences for unmethylated and methylated DNA as they would appear following bisulfite conversion and amplification are shown below. Unmethylated C-residues that would be converted by bisulfite and amplification to T residues are shown as Ts.

In some embodiments, the present invention provides use of a nucleic acid detection assay to coordinately analyze a plurality of the advantageous loci in a sample, thereby determining the disease state of cells in sample.

II. Determining methylation ratios for loci in the selected sequence(s). As discussed above, determining a methylation ratio for a locus comprises determining an average rate of methylation for that locus in a population of normal cells and determining the average rate of methylation at the same locus in a population of non-normal cells. As noted above, commonly used methods of methylation analysis of marker genes are performed on mixed nucleic acids, *e.g.*, amplicons produced from unfractionated DNA from a mixed population of cells (such as DNA purified from a multicellular tissue sample). While some studies have analyzed individual clones of amplicons made from unfractionated sample DNA, the numbers of clones analyzed has typically been too small to reveal significant or reproducible differences in methylation ratios at individual CpG loci within the sequences. For example, in their comparison of highly methylated genes in colorectal cancer, Zou, *et al.*, analyzed only six clones from each sample. (Zou, *et al.*, *Cancer Epidemiol Biomarkers Prev* 2007;16(12):2686), while Weisenberg, *et al.* used The present invention comprises large-scale analysis of individual DNA molecules, for example, by direct sequencing of individual DNA molecules, or by sequencing of clonally amplified DNA.

While not limiting the present invention to any particular methods, methods of clonally amplifying individual copies of nucleic acids (*e.g.*, using PCR) can be used in the rapid analysis of large numbers of individual markers from normal and non-normal samples. Single-molecule amplification methods may comprise use of microchambers, emulsion reactions, "bridge PCR" on solid supports, or any of a number of established methods of

segregating the amplification products arising from individual target molecules. Following single molecule amplification, amplicons can be sequenced.

Improved methods of sequencing individual molecules directly obviate the need to clone molecules into cells or, in some methods, the need to clonally amplify prior to sequencing. Elimination of cloning into cells makes analysis of much larger collections of molecules dramatically more efficient. Platforms for individual molecule sequencing include the 454 FLX™ or 454 TITANIUM™ (Roche), the SOLEXA™ / Illumina Genome Analyzer (Illumina), the HELISCOPE™ Single Molecule Sequencer (Helicos Biosciences), the Ion Personal Genome Machine (Ion Torrent), and the SOLID™ DNA Sequencer (Life Technologies/Applied Biosystems) instruments, as well as other platforms still under development by companies such as Intelligent Biosystems and Pacific Biosystems. Although the chemistry by which sequence information is generated varies for the different next-generation sequencing platforms, all of them share the common feature of generating sequence data from a very large number of individual sequencing templates, in sequencing reactions that are run simultaneously. Data from the reactions are collected using, *e.g.*, a flow cell, a chemical or optical sensor, and/or scanner, and sequences are assembled and analyzed using bioinformatics software.

In certain preferred embodiments, the present invention provides methods of analysis of methylation markers using digital sequencing to identify neoplasm-associated methylation loci that have methylation ratios that are statistically significantly advantageous compared to other loci in the same markers. In preferred embodiments, digital sequencing is done in a highly or massively parallel fashion, providing higher precision in identifying CpG methylation sites having advantageous methylation ratios.

For the massively parallel digital sequencing methods mentioned above, each molecule is analyzed for methylation at each CpG locus, so the percentage of DNA copies having methylation at any combination of the CpG loci can be analyzed after the experimental run. Further, each particular marker sequence, *e.g.*, each target nucleic acid molecule, or clonal amplicon may be interrogated many, many times, *e.g.*, at least 100 times, sometimes over 1000 times, and in some instances over 100,000 times, or as many as 500,000 times. Thus, patterns of coordinate methylation indicative of cancer or adenoma that would be undetectable in analysis of a handful of individual target molecules may be revealed.

III. Selection of a subset of methylation loci for coordinate analysis

As noted above, determination of the methylation status of a set of CpG loci in a large number of copies marker DNA from both normal samples and non-normal samples (*e.g.*, adenoma or cancer samples) reveals that certain certain CpG loci in marker genes or regions
5 may tend to be coordinately methylated. Further, design of nucleic acid detection assays to interrogate a plurality of CpG loci for which coordinate methylation is indicative of adenoma or cancer in a sample can provide an assay that has improved signal-to-noise compared to assays that survey average percent methylation across entire marker genes.

One aspect of selecting a subset of CpG loci comprises selecting loci that have been
10 determined to be coordinately methylated by use, *e.g.*, of digital analysis methods. Another aspect comprises selecting CpG loci determined to have advantageous methylation ratios when normal DNA is compared to adenoma or cancer DNA. Assay designs may, but need not, make use of a CpG locus having the most advantageous methylation ratio compared to other loci in the same marker. In some embodiments, selection of a plurality of CpG loci as a
15 subset comprises selecting the plurality of loci having the most advantageous methylation ratios. In other embodiments, selection of a plurality of CpG loci as a subset comprises selecting the locus having the most advantageous methylation ratio, then selecting at least additional CpG loci that are conveniently situated with respect to the first selected locus for the configuration of a particular nucleic acid detection assay (*e.g.*, the selection of CpG loci
20 having particular proximity to each other for configuring an invasive cleavage assay, ligation assay, amplification assay, *etc.*) in order to interrogate all of the selected CpG loci on copies of the target DNA in a single assay. In some embodiments, a candidate subset of CpG loci is further analyzed to determine the percentage of copies of marker DNA from non-normal samples that are coordinately methylated at those candidate loci, and that have little or no
25 coordinated methylation in normal samples.

Analysis of samples for detection of adenoma or cancer

Conventional methods of methylation analysis, (*e.g.*, conventional methylation-specific PCR, real time methylation-specific PCR, see, *e.g.*, US 5,786,146, 6,017,704,
30 6,200,756, 6,265,171.), typically analyze in a non-digital fashion, *e.g.*, analyzing a mixture of co-amplified molecules derived from a mixture of DNA target nucleic acids, so that analysis of the amplified products provides sequence information that reflects that aggregate or

average methylation status in the amplicon population, but does not provide information on the percentage of starting molecules having coordinated methylation at all of a plurality of CpG loci. In some instances researchers have analyzed a number of cloned amplicons, which can reveal the diversity in methylation in the CpG loci within a target marker gene.

- 5 However, sequencing individual clones has not provided enough data to reveal statistically significant coordinate methylation of specific subsets of CpG loci.

In contrast to conventional methods, we sought to analyze methylation marker genes in a massively parallel digital sequencing fashion to identify statistically significant coordinate methylation of specific CpG loci associated with neoplasms (adenoma and carcinoma). This method of analysis allowed us to:

1. Analyze samples for coordinate methylation in a marker gene as a means of detecting neoplasms without the need for testing any genetic (mutation) markers
2. Analyze samples for coordinate methylation in a plurality of marker genes as a means of detecting neoplasms without the need for testing any genetic (mutation) markers

15 We decided to use "digital" sequencing on a larger number of tissue samples obtained by biopsy from colorectal adenomas, colorectal cancers normal colorectal epithelia and other GI cancers and sequence a number of specific regions within several genes. This type of sequencing provides a methylation pattern for each individual methylated gene. For the first run we have 9 normal tissues, 38 adenomas and 36 cancer samples with the following

20 markers – Vimentin, BMP3, Septin 9, TFPI2, 2 regions of LRAT, and EYA4.

Surprisingly we found that in some of the genes, the background observed as methylation in normal samples is randomly distributed in the sequences, while the methylation associated with cancer and adenoma is not. Thus, if certain rules are applied *e.g.* if all of C residues a, b, and c have to be methylated in a diagnostic assay, then the number of

25 DNA copies presenting methylation at all three positions in that sequence is reduced compared to the number of DNA copies displaying methylation at a subset of the positions. In some of the marker genes or regions tested, the reduction in number of DNA copies in normal DNA displaying methylation at all of the selected sites drops to a greater degree than it does in the DNA from cancer and/or adenoma sample, resulting in an significantly

30 enhanced ratio of specific signal to background noise. For certain genes, the background from normal DNA is dramatically reduced by using multimethylation (coordinate methylation) analysis, while no equivalent reduction in signal from cancer and adenoma

DNA is seen. For other genes the background in normal samples is less reduced and/or the signal from cancer DNA also decreases with multimethylation analysis, such that there is less or no net improvement in the signal-to-noise and the advantages of using a multimethylation analysis approached are less. Genes having favorable signal to noise in multimethylation analyses are readily determined empirically.

From the multimethylation data presented herein (see, *e.g.*, Figures 2A-J) it is possible to:

- a. Identify regions within the gene sequences that give higher discrimination between normal and non-normal (*e.g.*, cancer and adenoma) cells;
- 10 b. Identify particular genes that have greater signal-to-noise (non-normal cell signal compared to normal cell background);
- c. Identify particular methylation loci that have greater signal-to-noise;
- d. Identify particular methylation loci that are coordinately methylation in adenoma and cancer samples but not in normal samples, such that detection coordinate methylation at these loci is a sensitive indicator of adenoma or cancer;
- 15 e. Identify genes with very low background methylation allowing for greater dilution of methylated DNA in normal DNA with less decrease in assay sensitivity;
- 20 f. Identify genes that are diagnostically complementary to each other, *i.e.*, that when analyzed in combination produce diagnostic information of elevated sensitivity and/or elevated specificity compared to the genes analyzed alone.
- g. Identify combinations of genes that give elevated sensitivity at elevated specificity, *e.g.*, 100% sensitivity for cancer and adenoma at 100% specificity.

25

EXPERIMENTAL EXAMPLES**EXAMPLE 1****Use of digital PCR and sequencing for the identification of specific subsets of CpG loci methylated in cancer and adenoma samples**

5 DNA extracted from frozen tissue samples was treated with an EPITECT bisulfite conversion kit (Qiagen) to convert non-methylated cytosines to uracil. Methylated cytosines remain unconverted. Primers for each gene region were designed for each sequence such that the composition of the amplification products remained the same as the original target sequences and methylated and non methylated sequences were amplified with equal
10 efficiencies. Amplification of the dU-containing converted DNA produced amplicons having T-residues in place of the dU residues. The amplicons were then prepared for sequencing on the Illumina instrument. For each tissue sample, the amplification reaction for each target was prepared from the same sample of bisulfite-treated DNA.

After sequencing, the data was analyzed quantitatively as an average methylation
15 similar to Sanger sequencing, but at a higher precision and resolution in that the combined signal at each position is calculated from individual molecules. For each amplicon sequence, a set of CpG loci was evaluated for percent methylation in the different tissues, to identify a subset the loci that were co-methylated more frequently in cancer and/or adenoma samples than in normal tissues.

20 Illumina Sequencing Protocol:

Sequencing was conducted according to the procedure recommended for the Illumina Genome Analyzer IIX, GAIIx, Data collection software ver. 2.5, and Pipeline analysis software ver 1.5. Briefly, the Illumina procedure comprises a) preparation of a library from sample DNA by attachment of known sequence tags that permit indexing, flow cell
25 attachment, amplification, and sequencing; b) attachment of the library to a flow cell surface; c) bridge amplification to produce clusters of DNA fragments derived from single molecules, and d) sequencing in using iterative primer extension reactions using labeled reversible terminators to determine the nucleotide sequence of each cluster of amplicons. See, *e.g.*, Bentley, *et al*, Nature 456, 53-59 (6 Nov.r 2008)/ doi:10.1038/nature07517 with
30 supplementary methods and data, incorporated herein by reference. Use of unique tag

sequences for indexing permits analysis of multiple samples in a single flow cell. See, *e.g.*, Craig, *et al.*, Nat. Methods Nat Methods. 2008 Oct;5(10):887-93 (Epub 14 Sept. 2008), incorporated herein by reference.

- 5 **Sample Set:** N=82, composed of tissue DNA extracted from 42 colorectal cancers, 31 pre-cancerous adenomas and 9 normal colonic mucosa.

Flow Cell Configuration

Samples were indexed at 12 per lane for a total of 7 lanes. A flow cell is composed of
10 8 lanes, one of which is dedicated to a phiX quality control.

Library Preparation:

Tissue-extracted DNA from patients was bisulfite-treated and a 2-step amplification using approximately 10,000 genome copies of initial material was carried out. The first
15 round used tailed (T1)(Illumina) primers specific for marker sequences. These tails were Illumina-derived sequences needed for round two. The second round (T2)(Illumina) PCR uses primers specific for the Illumina tails added in T1, and incorporates the index, sequencing primer, and flow cell attachment sequences. During the library preparation, multiple qPCR checks were run on the samples to ensure equimolar representations of all
20 amplicons in the libraries.

Primer Design:

Forward and reverse primers specific for regions with converted, non-CpG cytosines are designed (using, *e.g.*, MethPrimer software) to amplify each of the specific biomarker
25 sites in a non-methylation specific manner. When CpG cytosines cannot be avoided in the primer design, degenerate mixtures (C/T; G/A) are used those sites in the primers. If additional sequences outside the primary amplicon need to be queried, additional primers may be designed. If CpGs in the target sequence cannot be avoided, the primers may incorporate degenerate bases at CpG sites (BiSearch software).

30 Primers for second round PCR comprise sequences for Illumina flow cell attachment (bridge amplification sites), sequencing primer sites (for the sample read), index sites, and sequencing primers sites (for the indexing read). Each of the primer sets (x) has 12 different index tags, for a total of 12x sets. Index-independent primer sets (n=x) are optimized on

converted non-methylated DNA (e.g., human DNA) and converted methylated DNA. For example, the control DNAs are amplified, purified (e.g, using AMPURE treatment (Agencourt)), and run on an Agilent 2100 Bioanalyzer to assess the size and quantity of the amplified nucleic acids.

5

Experimental Steps:DNA Isolation and Bisulfite Conversion:

- 1) DNA was extracted and purified from tissue using either DNAZOL (Invitrogen), or QIAAMP kits (Qiagen) and the concentration and purity was measured by absorbance (A230/A260/A280) using a Nanodrop ND-1000 spectrophotometer (Thermo Scientific). PICOGREEN fluorescence (Molecular Probes) was used in conjunction with a TECAN F-200 (Tecan) plate-reader for high samples exhibiting high A230 values.
- 2) Samples were adjusted to a concentration of at least 200ng/uL using a Speedvac evaporation concentrator (Thermo Scientific), as necessary.
- 3) For each sample, 2ug of DNA was bisulfite-treated using EPITECT 96-well plates (Qiagen).
- 4) Recovery was assessed by absorbance and OLIGOGREEN fluorescence (Molecular Probes) and the conversion efficiency was assessed with quantitative PCR using cytosine-containing, non-CpG primers specific for unconverted DNA. Conversion efficiency was determined to be greater than 99%

10
15
20First-Round PCR:

- 5) The 84 samples were amplified in reactions with 30ng DNA using marker-specific (T1) primer sets. The number of cycles used was specific for each marker set and was empirically determined by the initial control reactions on both methylated and unmethylated DNA. The number of cycles is approximately set at the mean calculated Ct value. For example, the following numbers of cycles were used for the indicated markers:

25
30

TFPI12; 26 cycles
SEPT9; 27 cycles
BMP3; 28 cycles
VIM; 28 cycles
EYA4; 29 cycles

- 6) The amplified product from each reaction was purified using AMPURE beads (Agencourt), with elution in EB buffer (Qiagen).
- 5 7) The product for each marker was quantified by qPCR as described above, using a T2 primer set. Master plates were prepared containing equal amounts of each biomarker for each sample.

Second-Round PCR:

- 10 8) The first-round samples were then amplified with the 12 T2 indexed primers.
- 9) The product for each reaction was again purified and concentrations measured with qPCR, this time with flow-cell-specific primers and a standard curve created using serial dilution of PhiX control DNA.

15 Final Library Prep:

- 10) The 12 columns of each plate were combined into 1 master column in equimolar proportions. 1uL of each library was loaded on a high sensitivity DNA chip (Agilent) and run on the Bioanalyzer. A final qPCR was also performed on the 480 LightCycler (Roche) with the PhiX standards.
- 20 11) The libraries were sequenced on the Illumina Instrument and the sequence data obtained for each marker for each sample.
- 12) Average methylation at each CpG site was calculated for each marker for each sample. See Figures 2A-J.
- 13) The percentage of molecules methylated at all of the CpG loci in a defined subset of
25 the CpG loci in each marker was calculated for each marker for each sample. See Figures 5A-I.

All publications and patents mentioned in the above specification are herein
30 incorporated by reference. Various modifications and variations of the described methods and systems of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various

2012212127 17 May 2016

60 modifications of the described modes for carrying out the invention that are obvious to those skilled in the relevant fields are intended to be within the scope of the following claims.

5 Throughout this specification, unless the context requires otherwise, the word "comprise", or variations such as "comprises" or "comprising", will be understood to imply the inclusion of a stated element or integer or group of elements or integers but not the exclusion of any other element or integer or group of elements or integers.

10 The reference in this specification to any prior publication (or information derived from it), or to any matter which is known, is not, and should be taken as an acknowledgement or admission or any form of suggestion that the prior publication (or information derived from it) or known matter forms part of the common general knowledge in the field of endeavour to which this specification relates.

CLAIMS

1. A method of selecting a defined subset of CpG loci in a marker nucleic acid wherein coordinate methylation of the defined subset of CpG loci is indicative of adenoma or cancer, the method comprising:
 - a) determining the methylation status of a plurality of CpG loci in each of a plurality of individual copies of a marker nucleic acid from a plurality of normal samples;
 - b) determining the methylation status of said plurality of CpG loci in each of a plurality of individual copies of said marker nucleic acid from a plurality of adenoma or cancer samples;
 - c) determining methylation ratios for each locus in said plurality of said CpG loci in said marker nucleic acid, wherein said determining said methylation ratios comprises determining the ratio between the mean methylation at each of said plurality of CpG loci in said normal samples to the mean methylation at each corresponding CpG locus in said plurality of CpG loci in said adenoma or cancer samples;
 - d) selecting a defined set of CpG loci in said marker nucleic acid, wherein said defined set of CpG loci comprises at least three of CpG loci, each having an advantageous methylation ratio correlating with adenoma or cancer; and
 - e) selecting a defined subset of CpG loci from within said defined set wherein said defined subset of CpG loci comprises at least three CpG loci, and wherein said defined subset is selected such that the percentage of individual copies of said marker nucleic acid from said plurality of normal samples that are methylated at all of said at least three CpG loci in said defined subset is less than the percentage of individual copies of said marker nucleic acid from said plurality of adenoma or cancer samples that are methylated at all of said at least three CpG loci in said defined subset.

2. The method of claim 1, wherein said adenoma or cancer samples comprise colorectal samples.

3. The method of Claim 1 or Claim 2, wherein the plurality of individual copies of a marker nucleic acid analyzed in said normal, adenoma or cancer samples comprises at least 100, preferably 1000, preferably at least 10,000.
4. The method of Claim 1 or Claim 2, wherein the plurality of individual copies of a marker nucleic acid analyzed in said normal, adenoma or cancer samples comprises at least 100,000.
5. The method of any one of Claims 1 to 4 , wherein the plurality of normal, adenoma or cancer samples comprises at least 10.
6. The method of any one of Claims 1 to 4, wherein the plurality of normal, adenoma or cancer samples comprises at least 25.
7. The method of any one of Claims 1 to 4, wherein the plurality of normal, adenoma or cancer samples comprises at least 100.
8. The method of any one of Claims 1 to 7, wherein said defined set of CpG loci comprises at least four CpG loci.
9. The method of any one of Claims 1 to 7, wherein said defined set of CpG loci comprises at least five CpG loci.
10. The method of any one of Claims 1 to 9, wherein said determining the methylation status of said defined set of CpG loci comprises treating DNA from said samples with bisulfite.
11. The method of any one of Claims 1 to 10, wherein said determining the methylation status of said defined set of CpG loci comprises digital analysis of each of a plurality of CpG loci in said marker nucleic acid.
12. The method of Claim 11, wherein said digital analysis comprise digital sequencing and/or digital PCR.

13. A method of detecting cancer or adenoma in a sample, comprising determining the methylation status of each CpG locus in a defined subset of CpG loci in at least one cancer or adenoma marker nucleic acid molecule, wherein said defined subset of CpG loci comprises at least three CpG loci, and wherein methylation at all of said CpG loci in said defined subset of CpG loci in said cancer or adenoma marker nucleic acid molecule is indicative of cancer or adenoma in said sample.
14. The method of Claim 13, wherein said defined subset of CpG loci comprises at least four CpG loci.
15. The method of Claim 13, wherein said defined subset of CpG loci comprises at least five CpG loci.
16. The method of any one of Claims 13 to 15, wherein said determining comprises analysis of said CpG loci in a nucleic acid detection assay configured to determine the methylation status of each of said loci in a single nucleic acid detection assay.
17. The method of Claim 16, wherein said determining comprises analysis of said CpG loci in a nucleic acid detection assay configured to determine the methylation status of each of said loci in a single reaction mixture.
18. The method of Claim 16, wherein said nucleic acid detection assay comprises a primer extension assay, a nucleic acid amplification assay, a nucleic acid sequencing assay, a structure-specific cleavage assay, a 5' nuclease cleavage assay, an invasive cleavage assay, and/or a ligation assay.
19. The method of any one of Claims 13 to 18, wherein said cancer or adenoma marker nucleic acid molecule is selected from the group consisting of Vimentin, BMP3, Septin 9, TFPI2, 2 regions of LRAT, and EYA4 nucleic acid molecules.

20. The method of Claim 13, wherein said defined subset of CpG loci comprises at least one defined subset of CpG loci selected from the group consisting of: loci 37, 40, and 45 in a vimentin nucleic acid molecule; loci 34, 53, and 61 in a BMP3 nucleic acid molecule; loci 59, 61, 68, and 70 in a septin 9 nucleic acid molecule; loci 55, 59, 63, and 67 in a TFPI2 nucleic acid molecule; and loci 31, 34, 37, and 44 in an EYA4 nucleic acid molecule.

FIG. 2A

Vimentin										
CpG	26	37	40	45	52	54	59	63	74	
NORMAL										Mean
8-4	11.63	6.08	6.62	3.52	3.65	2.31	3.44	3.66	3.88	4.98
8-5	6.92	3.4	2.97	1.26	1.19	0.5	2.61	0.97	1.14	2.33
8-6	12.25	8.92	5.06	4.01	2	2.45	2.27	3.44	5.21	5.07
8-7	15.74	9.41	5.7	6.72	3.73	5.26	5.92	4.29	4.46	6.80
8-8	6.48	2.52	2.71	2.72	4.26	3.14	1.67	2.67	2.84	3.22
8-9	5.65	2.48	1.27	1.48	1.5	0.87	1.7	2.02	3.18	2.24
8-10	5.36	2.77	1.09	2.84	2.09	0.85	1.89	1.97	2.55	2.38
8-11	7.5	2.65	2.34	1.66	4.9	4.84	1.71	3.5	3.04	3.57
8-12	5.91	5.12	3.52	1.91	2.53	2.27	1.84	1.86	1.58	2.95
Mean	8.60444	4.81667	3.47556	2.90222	2.87222	2.49889	2.56111	2.70889	3.09778	3.73
										SD all
										2.68
										M+2SD
										9.09
										M+3SD
										11.77
ADENOMA										
CpG	26	37	40	45	52	54	59	63	74	
ADENOMA										Mean
3-4	50.2	52.6	47	24.66	21.46	16.34	20.96	27.91	36.05	33.02
3-5	79.04	70.33	65.69	56.54	33.15	25.6	35.34	32.14	40.64	48.72
3-6	37.08	32.91	32.94	25.05	23.78	25.64	23.57	23.22	29.4	28.18
3-7	47.73	45.01	44.47	42.42	41.78	41.24	41.83	41.75	40.02	42.92
3-8	56.11	48.88	46.52	44.12	41.26	41.34	41.29	40.82	43.15	44.83
3-9	6.49	3.18	4.26	2.92	3.28	2.87	2.55	2.17	3.46	3.46
3-10	5.7	2.61	4.61	2.02	2.77	2.46	2.2	2.15	3.01	3.06
3-11	84.54	84.38	83.07	41.43	43.76	51.8	77.13	75.74	48.14	65.55
3-12	90.13	90.15	88.53	88.89	88.89	85.22	89.03	83.48	83.35	86.85
5-1	71.75	71.01	70.12	65.83	72.49	71.45	69.97	61.69	65.53	68.87
5-2	39.07	35.75	33.57	31.84	31.98	31.84	30.25	30.1	30.97	32.82
5-4	40.85	32.71	17.84	21.28	18.47	20.19	20.5	23.79	40.53	26.24
5-11	70.93	70.52	69.58	64.06	70.39	70.52	68.51	60.3	63.66	67.51
5-12	84.36	84.75	84.36	77.41	73.36	57.53	49.03	68.92	79.15	73.21
6-1	87.8	86.44	87.45	86.38	85.89	82.93	84.12	80.86	80.54	84.71
6-2	50.08	46.61	40.99	37.87	36.69	32.92	32.63	39.06	39.44	39.59
6-3	68.72	67.13	59.29	60.01	55.34	55.21	49.64	54.92	55.63	58.43
6-4	59.32	59.36	53.62	43.22	42.23	39.54	44.19	45.19	38.26	47.21
6-5	14.51	8.39	8.32	4.44	3.24	2.17	3.35	2.1	4	5.61
6-6	55.32	54.47	53.14	43.9	36.87	39.01	34.56	36.08	39.69	43.89
7-5	71.06	70.46	71.36	70.54	70.74	70.65	68.07	63.9	64.36	69.02
7-6	58.16	55.35	49.14	45.4	42.42	45.59	52.06	44.81	47.71	48.96
7-7	20.5	16.52	15.05	12.88	12	11.04	12.63	11.37	10.61	13.62
7-8	28.05	20.31	22.19	13.34	15.5	10.57	19.98	18.19	17.36	18.39
7-9	79.93	79.58	78.49	78.5	78.21	76.44	75.6	72.93	68.5	76.46
7-10	80.88	81.15	80.46	80.04	79.32	77.6	76.5	74.03	70.63	77.85
7-11	22.15	12.4	10.05	5.74	7.02	2.31	6.61	6.09	9.06	9.05
7-12	33.99	32.48	33.46	31.53	32.34	31.36	31.44	29.17	30.14	31.77
8-1	48.74	49.22	50.37	48.32	50.05	50.83	50.9	49.21	47.73	49.49
8-2	82.35	82.4	82.33	81.75	81.4	81.89	80.65	79.84	78.95	81.28
8-3	63.16	63.53	63.87	65.06	64.59	64.34	62.43	62.09	61.29	63.37
Mean	54.4742	51.9545	50.0690	45.0771	43.9571	42.5303	43.5974	43.3555	44.2245	
N/M	0.15795	0.09271	0.06942	0.06438	0.06594	0.05876	0.05874	0.06248	0.07005	

FIG. 2B

Vimentin											
CpG	26	37	40	45	52	54	59	63	74		
CANCER										Mean	
1-1	55.95	46.38	39.19	39.35	39.25	38.19	30.93	26.81	33.13	38.80	
1-2	46.07	48.21	39.2	26.64	33.32	21.46	34.99	27.12	35.62	34.74	
1-3	36.4	23.79	25.7	5.9	6.22	4.63	4.54	4.83	6.72	13.19	
1-4	80.52	66.55	81.59	80.93	80.51	81.01	79.67	70.78	76.98	77.62	
1-5	86.12	85.72	85.71	85.99	85.66	86.21	84.61	81.46	82.86	84.93	
1-6	76.07	75.26	76.89	75.46	74.46	75.74	74.47	70.81	57.16	72.92	
1-7	11.26	13.06	26.2	7.2	5.52	2.47	3.57	2.68	3.34	8.37	
1-8	3.24	2.4	3.04	3.05	3.85	2.63	4.58	3.05	2.85	3.19	
1-9	91.73	91.67	90.09	90.69	91.4	87.87	88.51	72.29	75.96	86.69	
1-10	91.81	91.44	90.57	91.05	91.41	88.35	88.74	71.95	75.29	86.72	
1-11	96.17	95.74	94.21	85.46	66.31	55.84	45.36	53.05	55.95	72.01	
1-12	81.46	70.09	58.61	49.52	40.5	39.07	40.96	34.12	29.02	49.26	
2-1	3.2	2.71	2.37	2.63	4.47	1.65	3.08	2.32	2.89	2.81	
2-2	27.81	27.74	28.16	24.33	27.01	26.02	25.55	19.5	21.84	25.33	
2-3	50.74	50.46	50.99	50.06	50.1	50.5	48.51	47.2	46.41	49.44	
2-4	49.58	48.5	49.13	42.19	48.89	40.01	47.42	35.7	43.83	45.03	
2-5	5.4	3.66	3.69	3.1	3.06	2.56	2.54	2.58	4.21	3.42	
2-6	39.3	39.04	39.7	38.65	39.38	38.32	37.14	36.43	36.53	38.28	
2-7	55.16	54.9	54.89	54.26	55.2	55.07	54.09	51.47	51.4	54.05	
2-8	62.61	62.07	63.43	61.37	62.91	62.71	62.75	58.15	57.97	61.55	
2-9	5.64	4.28	1.97	2.06	3.81	2.26	6.47	2.86	2.97	3.59	
2-10	4.96	3.92	1.36	1.52	3.48	2.24	5.74	1.94	2.31	3.05	
2-11	68.03	67.01	59.74	66.53	68.15	64.62	65.4	57.5	61.65	64.29	
2-12	60.01	47.56	57	32.01	44.13	40.93	33.52	30.48	45.93	43.51	
3-2	45.95	45.94	46.16	45.91	45.76	46.43	44.81	43.34	42.49	45.20	
5-3	30.38	29.55	29.08	28.84	30.73	30.97	28.13	27.42	28.37	29.27	
5-5	29.85	17.44	6.57	7.31	3.29	3.22	2.62	2.62	5.57	8.72	
5-6	9.19	15.6	3.9	4.11	10.38	3.48	2.23	9.96	13.16	8.09	
5-7	52.6	51.71	39.08	34.92	25.56	28.83	38.34	45.32	48.14	40.50	
5-8	21.65	20.69	20.69	19.08	22.4	21.76	21.76	19.51	19.4	20.77	
5-9	44.3	42.83	43.32	39.74	44.14	41.53	28.83	39.41	40.88	40.55	
5-10	43.44	41.51	42.53	40.36	42.76	40.94	28.62	37.06	39.34	39.62	
6-7	91.08	90.06	90.06	90.28	89.75	90.85	88.49	64.97	85.04	86.95	
6-8	65.53	62.59	59.24	59.03	51.16	50.98	53.92	58.67	60.01	57.90	
6-9	7.35	4.62	4.49	2.5	3.93	4.24	5.55	4.66	8.91	5.14	
6-10	5.56	3.22	4.2	1.36	2.97	2.07	3.03	2.9	7.64	3.66	
6-11	66.81	65.15	64.79	64.41	67.47	63.55	64.79	60.5	61.13	64.29	
6-12	68.38	63.74	45	34.86	35.79	25.68	31.84	25.16	33.91	40.48	
7-1	63.71	61.07	60.24	55.62	56.07	57.06	45.3	48.45	49.69	55.25	
7-2	49.63	41.8	31.28	30.46	23.63	26.56	15.68	23.23	24.1	29.60	
7-3	61.72	60.06	60.77	60.47	60.53	60.58	59.07	57.15	55.66	59.56	
7-4	83.11	82.56	77.6	81.19	67.36	56.23	74.45	56.05	69.23	71.98	
Mean	48.3210	45.7690	44.1055	40.9619	40.7781	38.6957	38.3476	35.9871	38.2260		
N/M	0.17807	0.10524	0.07880	0.07085	0.07044	0.06458	0.06679	0.07527	0.08104		

FIG. 2C

BMP3						
CpG	34	53	61	70	74	
NORMAL						Mean
8-4	3.25	3.26	2.63	3.22	2.92	3.06
8-5	2.37	2.3	1.4	1.88	2.21	2.03
8-6	3.18	1.59	2.69	2.09	2.37	2.38
8-7	3.78	4.15	0.84	2.35	1.84	2.59
8-8	3.99	2.95	2.9	2.91	4.26	3.40
8-9	5.49	3.69	1.71	3.03	3.65	3.51
8-10	5.29	3.44	0.02	0.06	0.11	1.78
8-11	4.22	2.92	2.82	2.65	2.51	3.02
8-12	2.98	3.23	2.71	1.64	1.98	2.51
Mean	3.83889	3.05889	1.96889	2.20333	2.42778	2.70
						SD all
						M+2SD
						M+3SD
ADENOMA						
CpG	34	53	61	70	74	
ADENOMA						Mean
3-4	2.33	3.14	3.61	2.69	4.13	3.18
3-5	19.45	8	4	2.88	4.25	7.72
3-6	17.33	14.18	13.48	15.42	17.16	15.51
3-7	45.19	46.07	41.96	41.15	41.65	43.21
3-8	9.43	4.4	3.57	6.31	6.61	6.06
3-9	5.04	4.75	3.42	3.29	3.1	3.92
3-10	5.32	4.02	3.39	2.71	1.73	3.43
3-11	4.77	3.54	3.99	12.56	3.79	5.73
3-12	5.63	3.77	4.42	6.87	5.14	5.17
5-1	87.27	85.28	84.85	83.63	86.85	85.58
5-2	34.33	32.92	25.52	31.5	28.84	30.62
5-4	8.5	2.83	2.75	4.72	3.95	4.55
5-11	30.56	23.64	15.75	20.94	18.2	21.82
5-12	9.12	15.55	22.11	23.91	10.78	16.29
6-1	90.01	88.79	86.46	88.95	88.02	88.85
6-2	3.22	3.38	3.03	2.23	3.28	3.03
6-3	82.39	80.24	80.21	79.47	80.87	80.64
6-4	2.98	2.51	2.52	2.34	3.17	2.70
6-5	2.8	2.51	2.02	2.46	2.61	2.48
6-6	26.76	25.01	23.53	24.17	24.84	24.86
7-5	41.72	40.49	40.45	39.64	40.58	40.58
7-6	72.74	70.9	66.24	68.78	70.47	69.83
7-7	14.78	14.42	13.32	13	12.37	13.58
7-8	18.74	15.98	11.73	6.85	6.02	11.86
7-9	20.44	21.31	20.56	19.66	20.77	20.55
7-10	16.24	13.64	12.34	16.49	15.25	14.79
7-11	5.49	3.8	3.88	8.94	3.31	5.08
7-12	27.12	27.51	27.26	26.17	26.11	26.83
8-1	56.1	55.54	55.86	56.41	55.4	55.86
8-2	82.63	81.89	81.36	82.11	80.68	81.73
8-3	1.36	2.08	0.67	2.7	2.4	1.84
Mean	27.4126	25.8742	24.7184	25.7726	24.9142	
N/M	0.14004	0.11822	0.07965	0.08549	0.09745	

FIG. 2D

BMP3							
CpG	34	53	61	70	74		
CANCER						Mean	
1-1	4.1	3.27	2.75	2.48	2.97	3.11	
1-2	2.51	2.26	1.79	1.69	1.79	2.01	
1-3	3.93	2.97	2.67	2.33	3.2	3.02	
1-4	3.03	5.87	6.21	6.31	6.12	5.51	
1-5	84.92	84.93	84.66	84.32	84.63	84.69	
1-6	69.86	68.52	68.6	68.14	69.27	68.88	
1-7	4.12	2.89	2.73	3.84	3.19	3.35	
1-8	3.57	2.38	3	1.77	2.7	2.68	
1-9	46.41	30.05	14.12	5.5	5.4	20.30	
1-10	45.95	28.29	12.83	4.28	3.9	19.05	
1-11	3.71	6.67	4.29	4.63	16.61	7.18	
1-12	3.68	3.03	2.44	2.2	2.58	2.79	
2-1	2.8	2.42	1.17	1.05	2.37	1.96	
2-2	3.54	2.31	2.53	1.19	2.97	2.51	
2-3	47.85	47.99	46.97	46.63	46.65	47.22	
2-4	21.33	4.65	3.75	2.92	3.23	7.18	
2-5	74.6	74.29	73.31	74.06	74.61	74.17	
2-6	43.55	43.4	42.74	42.2	42.68	42.91	
2-7	57.93	57.12	56.09	56.29	56.34	56.75	
2-8	20.14	6	3.12	4.79	4.2	7.65	
2-9	84.2	83.37	82.82	82.49	83.03	83.18	
2-10	83.9	82.79	82.46	82.14	82.69	82.80	
2-11	60.49	57.84	63.95	61.56	63.87	61.54	
2-12	4.53	2.96	2.4	2.63	3.37	3.18	
3-2	52.98	52.28	52.39	52.08	52.32	52.41	
5-3	27.71	31.07	25.97	27.23	21.08	26.61	
5-5	21.21	17.74	17	13.13	13.84	16.58	
5-6	4.02	2.82	2.56	1.47	2.25	2.62	
5-7	3.07	5.48	5.09	3.07	2.63	3.87	
5-8	37.56	36.21	34.38	35.91	36.51	36.11	
5-9	9.87	9.23	3.47	5.94	4.84	6.67	
5-10	8.23	8.55	3.47	3.54	4.44	5.65	
6-7	87.32	85.77	86.05	85.91	86.14	86.24	
6-8	48.14	52.49	48.07	49.61	48.67	49.40	
6-9	3.81	2.96	2.49	1.99	2.75	2.80	
6-10	3.32	2.1	1.89	1.48	2.68	2.29	
6-11	36.92	55.67	54.4	54.85	53.99	51.17	
6-12	15.79	14.37	13.98	13.37	14.73	14.45	
7-1	2.37	2.63	2.21	2.07	2.63	2.38	
7-2	3.5	2.31	3.04	3.24	3.77	3.17	
7-3	64.11	63.08	63.22	62.45	63.17	63.21	
7-4	4.82	2.5	3.03	2.53	2.49	3.07	
Mean	28.9381	27.4650	25.9548	25.3645	25.8881		
N/M	0.13266	0.11137	0.07586	0.08687	0.09378		

FIG. 2E

Septin9							
CpG	31	38	59	61	68	70	
NORMAL							Mean
8-4	4.33	4.68	3.65	3.4	3.05	2.55	3.61
8-5	5.81	5.22	3.88	2.55	2.56	2.62	3.77
8-6	9.7	8.48	6.03	5.02	5.4	3.5	6.36
8-7	25.07	23.83	15.73	15.38	12.7	7.3	16.67
8-8	9.6	9.34	7.07	7.68	7.06	5.31	7.68
8-9	2.93	3.79	2.85	2.19	2.67	2	2.74
8-10	2.95	3.94	2.22	3.17	3.03	2.33	2.94
8-11	9.91	10.36	7.98	6.95	6.22	4.78	7.70
8-12	7.37	8.04	5.18	3.9	3.49	3.71	5.28
Mean	8.63000	8.63111	6.06556	5.58222	5.13111	3.78889	6.30
						SD all	5.06
						M+2SD	16.42
						M+3SD	21.47
CpG	31	38	59	61	68	70	
ADENOMA							Mean
3-4	71.21	71.25	67.4	57.63	70.36	70.48	69.72
3-5	78.85	78.62	78.74	79.41	78.69	77.51	78.64
3-6	47.45	47.25	47.36	46.33	45.52	43.7	46.27
3-7	54.28	54.19	53.89	54.5	52.93	51.27	53.51
3-8	82.73	80.42	82.52	83.81	81.98	83.04	82.42
3-9	59.76	59.11	59.97	59.88	59.03	59.65	59.57
3-10	59.64	59.13	60.14	60.14	59.08	59.62	59.63
3-11	81.43	80.79	81.85	82.38	81.68	81.7	81.64
3-12	87.2	87.14	87.78	85.16	87.49	86.78	86.93
5-1	81.86	81.66	82.4	81.83	81.39	82.63	81.96
5-2	58.22	58.48	57.93	56.61	55.53	52.77	56.59
5-4	90.56	89.44	91.06	90.32	89.35	89.37	90.02
5-11	82.52	81.57	82.17	82.04	80.93	80.54	81.63
5-12	87.76	87.67	88.6	87.5	87.52	83.04	87.02
6-1	88.48	88.58	88.91	88.89	88.14	87.99	88.50
6-2	54.02	53.55	52.96	51.62	51.85	49.39	52.23
6-3	75.28	74.77	75.14	74.75	73.85	70.73	74.09
6-4	66.97	67.68	66.53	55.84	46.89	43.19	57.83
6-5	39.2	39.97	29.71	29.79	24.9	18.71	30.38
6-6	46.52	46.87	47.65	46.69	46.41	43.43	46.26
7-5	71.84	71.15	71.81	71.76	71.65	70.46	71.45
7-6	67.38	67.19	66.38	67.37	67.74	65.9	66.99
7-7	39.96	39.74	40.16	38.96	38.06	35.38	38.71
7-8	76.27	73.19	78.23	77.87	70.99	55.35	71.98
7-9	82.36	81.58	82.87	80.95	81.01	79.39	81.36
7-10	82.1	81.35	83.09	81.42	80.71	79.28	81.33
7-11	79.96	79.14	76.53	75.27	74.93	63.99	74.97
7-12	28.03	27.75	28.58	28.03	27.64	27.52	27.93
8-1	47.19	46.29	48.75	47.99	47.07	47.31	47.43
8-2	78.05	78.32	78.79	79.02	77.53	78.26	78.33
8-3	73.6	73.1	72.21	73.63	72.92	73.09	73.26
Mean	68.4058	67.9658	68.1006	67.3352	66.2506	64.2410	67.0498
N/M	0.12616	0.12699	0.08907	0.08290	0.07745	0.05898	0.09403

FIG. 2F

CpG	Septin9						Mean
	31	38	59	61	66	70	
CANCER							
1-1	74.24	74.37	74.74	69.16	73.67	69.78	72.66
1-2	64.3	63.59	64.48	63.97	64.03	64.55	64.15
1-3	78.49	77.13	77.94	75.03	76.7	75.33	76.77
1-4	76.88	76.83	61.04	77.26	77.25	76.34	74.27
1-5	83.22	82.77	82.91	89.1	78.22	81.8	82.00
1-6	72.72	72.6	73.26	73.34	72.55	72.53	72.83
1-7	58.64	58.9	59.5	59.02	57.95	58.03	58.66
1-8	41.04	41.59	42.34	39.9	39.31	31.59	39.30
1-9	87.69	87.79	87.9	85.22	87.34	86.76	87.12
1-10	87.52	88.04	88.3	85.78	87.66	87.37	87.45
1-11	93.45	93.07	83.24	92.82	92.33	92.82	91.29
1-12	54.85	51.7	55.46	53.56	51.61	48.38	52.59
2-1	59.5	59.92	61.38	60.56	66.72	60.88	58.49
2-2	52.9	57.7	53.1	57.04	57.76	46.02	54.09
2-3	49.88	49.48	50.03	50.05	48.88	49.32	49.61
2-4	45.75	46.15	46.94	47	46.06	29.99	43.65
2-5	65.13	65.64	67.05	67.26	66.23	66.47	66.30
2-6	20.94	20.51	20.35	19.89	20.44	9.66	18.62
2-7	47.93	46.04	41.52	41.4	36.64	39.59	42.19
2-8	68.51	68.14	69.55	69.26	68.23	68.64	68.72
2-9	75.02	73.17	56.65	75.24	71.97	75.17	71.20
2-10	74.22	72.24	56.32	74.72	71	74.53	70.51
2-11	37.38	38.85	24.02	14.56	10.66	5.93	21.90
2-12	59.19	58.71	59.1	58.62	55.6	54.69	57.65
3-2	46.5	46.36	47.19	48.3	45.89	45.97	46.70
3-3	47.82	48.26	49.38	48.05	48.32	48.3	48.36
3-5	30.03	28.6	30.72	27.46	28.69	29.34	29.14
3-6	64.74	65.65	65.21	60.89	64.87	65.19	64.43
3-7	57.61	58.27	59.96	58.38	58.15	54.59	57.83
3-8	30.81	30.44	32.62	31.11	31.36	31.13	31.25
3-9	57.09	57.39	58.38	57.84	56.92	57.5	57.52
3-10	56.43	56.72	58.5	57.27	56.35	56.66	56.99
6-7	86.96	86.4	81.86	86.91	86.55	84.06	85.46
6-8	61.99	60.83	61.43	62.11	62.02	62.04	61.73
6-9	55.17	54.87	55.62	55.64	55.6	55.37	55.38
6-10	54.31	54.02	54.91	54.51	54.35	54.33	54.41
6-11	2.04	2.13	3.89	3.2	2.29	2.48	2.67
6-12	75.22	75.08	76.44	76.09	75.94	76.3	75.85
7-1	63.39	63.35	63.77	63.97	62.75	59.07	62.71
7-2	63.88	62.96	64.86	63.63	63.35	62.23	63.49
7-3	73.04	73.06	73.73	74.34	73.65	73.43	73.54
7-4	75.73	77.28	76.63	79.06	79.23	79.13	77.84
Mean	60.2864	60.1548	58.8624	59.5824	58.9786	57.4593	
N/M	0.14315	0.14348	0.10305	0.09369	0.08700	0.06594	

FIG. 2G

	TFPI2									
CpG	28	33	41	50	55	59	63	67	74	
NORMAL										Mean
8-4	16.8	11.72	18.34	15.92	14.43	12.65	20.44	11.45	10.89	14.74
8-5	10.4	6.72	10.5	9.92	8.66	8.57	14.69	7.26	6.75	9.27
8-6	5.78	3.74	5.21	5.57	4.66	5.04	7.19	2.99	4.17	4.93
8-7	7.42	3.99	6.42	6.28	5.29	4.82	8.15	3.56	3.07	5.44
8-8	5.19	3.55	3.84	3.46	2.28	2.69	5.08	2.4	2.54	3.45
8-9	9.18	6.49	10.3	7.05	6.96	6.93	11.24	4.28	4.52	7.44
8-10	9.2	5.94	10.48	7.21	6.49	6.71	11.22	4.83	4.22	7.37
8-11	8.32	3.01	5.69	6.61	3.66	4.03	6.71	3.07	3.07	4.91
8-12	5.23	3.81	5.83	4.96	3.17	2.96	7.42	3.75	2.28	4.38
Mean	8.61333	5.44111	8.51222	7.44222	6.17778	6.04444	10.23778	4.84444	4.61111	6.88
										SD all
										3.86
										M+2SD
										14.61
										M+3SD
										18.47
CpG	28	33	41	50	55	59	63	67	74	
ADENOMA										Mean
3-4	66.66	49.94	65.74	66.29	59.94	62.71	64.28	48.29	30.53	57.15
3-5	77.23	76.12	77.55	76.96	75.45	73.61	76.39	72.7	71	75.22
3-6	41.89	36.35	40.76	40.97	40.56	38.13	40.11	37.77	34.48	39.06
3-7	52.43	48.78	52.34	53.57	50.9	49.98	52.74	47.83	44.76	50.37
3-8	80.17	79.31	79.2	79.83	79.76	77.16	76.28	78.09	64.06	77.10
3-9	41.24	39.22	40.04	40.87	40.69	40.06	39.84	39.23	39.04	40.03
3-10	39.75	37.16	38.59	39.38	39.06	38.61	38.26	37.68	36.9	38.38
3-11	84.09	81.71	83.36	83.58	83.66	82.34	81.57	82.53	80.87	82.63
3-12	86.28	75.8	86.62	87.69	78.28	79.03	86.59	79.89	58.94	79.90
5-1	78.42	77.48	78.38	73.72	77.69	76.86	72.05	75.7	70.17	75.61
5-2	59.47	53.62	57.42	53.97	55.91	55.91	54.51	52.46	50.29	54.84
5-4	91.15	91.11	92.16	87.49	91.68	86.88	86	90.22	89.68	89.53
5-11	73.27	70.21	70.36	68.17	69.56	68.83	66.21	69.63	69.41	69.52
5-12	78.05	75.39	84.77	80.16	80.59	83.95	78.32	83.67	83.63	80.95
6-1	86.13	84.43	85.44	85.9	85.74	83.62	83.43	84.01	82.29	84.55
6-2	52.6	48.51	50.23	51.37	50.91	48.11	50.74	48.74	47.62	49.87
6-3	71.67	68.75	70.6	70.04	70.4	67.48	69.53	67.59	64.92	69.00
6-4	53.89	27.64	52.9	53.58	47.65	29	48.93	48.09	39.87	44.62
6-5	18.96	10.51	19.35	14.32	13.37	12.03	19.15	7.52	6.76	13.55
6-6	52.19	48.46	50.11	50.18	50.03	47.91	49.4	47.35	40.79	48.49
7-5	57.72	56.03	57.33	57.64	57.01	55.73	56.81	55.51	53.79	56.40
7-6	52.68	49.32	51	50.35	50.47	48.95	48.43	49.85	49.02	50.01
7-7	25.22	18.22	21.12	20.96	19.79	18.46	22.15	16.57	15.59	19.79
7-8	33.85	15.37	23.72	19.92	27.03	37.42	54.82	30.88	33.69	30.74
7-9	73.9	59.02	70.99	65.87	66.99	61.66	71.3	62.11	56.11	65.33
7-10	71.51	56.26	68.66	63.4	64.27	58.32	69.59	60.46	55.45	63.10
7-11	63.86	52.6	68.02	62.26	55.02	53.43	67.1	48.71	39.43	56.71
7-12	26.9	21.82	25.75	26.12	26.72	24.49	27.01	22.98	22.07	24.87
8-1	39.67	36.94	38.87	39.82	38.12	37.04	39.47	36.65	35.48	38.01
8-2	4.39	2.77	3.43	9.4	3.36	4.15	5.24	5.01	4.29	4.67
8-3	18.38	6.67	3.77	8.39	7.12	8.05	22.38	3.59	7.32	9.52
Mean	56.5684	50.1781	55.1155	54.2635	53.4752	51.9326	55.4397	51.3326	47.6661	
N/M	0.15226	0.10844	0.15444	0.13715	0.11553	0.11639	0.18467	0.09437	0.09674	

FIG. 2H

CoG	28	33	41	50	55	59	63	67	74	Mean
CANCER										
1-1	75.92	73.75	76.91	76.98	76.89	75.12	74.58	75.44	71.99	75.29
1-2	60.7	58.48	59.83	60.11	59.24	58.56	58.47	58.39	58.02	59.09
1-3	69.4	67.39	69.05	69.3	69.58	66.17	67.34	67.9	66.92	68.12
1-4	84.99	82.51	85.32	85.75	83.16	80.99	83.51	84.26	86.55	77.45
1-5	80.24	78.4	79.98	77.16	79.59	79.17	78.2	72.3	76.71	77.97
1-6	72.82	70.18	72.1	72.24	72.46	71.33	70.86	69.43	69.23	71.18
1-7	56.49	13.25	31.78	55.62	33.25	30.17	54.09	28.21	32.74	37.29
1-8	32.13	20.84	30.98	32.68	33.26	30.28	30.56	29.42	29.77	30.00
1-9	89.35	86.99	84.73	89.13	88.76	87.47	86.92	87.1	60.92	84.60
1-10	88.63	86.02	83.65	88.27	87.98	86.9	86.05	86.28	60.23	83.78
1-11	70.4	12.47	93.04	93.91	75.62	91.84	90.9	64.22	18.62	67.89
1-12	67	65.56	66.77	66.96	66.98	65.79	64.81	64.2	64.96	65.89
2-1	15.11	11.81	12.62	12.17	13.99	11.93	12.78	12.63	11.69	12.75
2-2	28.93	6.35	10.73	33.33	17.12	15.95	30.68	8.02	7.15	17.58
2-3	45.41	43.39	44.33	44.06	44.47	43.25	42.52	43.4	42.42	43.69
2-4	45.2	42.77	44.85	45.04	45.13	43.42	43.83	41.69	35.59	43.06
2-5	63.22	51.49	65.06	65.35	65.29	63.52	63.19	62.25	52.05	61.27
2-6	31.68	28.87	29.77	30.31	30.84	29.71	29.63	29.67	29.19	29.96
2-7	45.62	43.81	44.36	44.25	41.67	43.54	43.71	42.12	41.39	43.39
2-8	64.7	62.61	63.61	63.79	57.58	60.84	60.93	62.33	58.59	61.66
2-9	75.87	71.86	75.47	74.9	75.49	71.68	71.42	74.03	73.19	73.77
2-10	72.47	68.98	72.08	71.33	72.4	68.8	67.93	70.49	69.81	70.48
2-11	59.41	50.01	64.27	64.23	56.2	58.22	60.85	58.34	39.96	56.83
2-12	50.9	45.83	56.05	55.83	52.73	51.94	54.45	52.23	48.45	52.05
3-2	43.65	40.73	43.07	42.99	42.94	43.01	42.86	40.76	40.71	42.30
5-3	38.56	30.45	42.56	40.53	35.54	34.59	40.19	36.63	28.62	36.41
5-5	61.12	56.82	57.86	54.72	56.43	56.68	54.83	50.83	54.89	56.02
5-6	53.8	48.22	59.5	38.46	17.73	19.02	28.45	18.28	13.88	33.04
5-7	63.5	60.54	61.93	58.59	62.75	60.84	58.35	58.15	58.39	60.34
5-8	22.3	18.7	19.36	18.14	20.25	18.93	19.06	19.85	19.06	19.52
5-9	52.64	51.06	52.01	48.3	50.75	49.65	47.16	46.77	49.25	49.73
5-10	49.62	48.7	48.35	45.2	48.18	47.48	46.33	44.56	46.76	47.24
6-7	64.49	83.96	84.44	83.97	84.53	79.72	81.88	82.5	81.27	82.97
6-8	52.41	45.12	51.33	50.98	50.21	49.08	50.3	45.16	39.78	48.26
6-9	34.12	12.99	37.31	25.11	14.81	25.65	37.63	14.79	31.82	26.03
6-10	34.11	12.42	36.31	23.71	13.2	24.85	37.09	14.15	31.6	25.27
6-11	61.62	30.74	60.7	61.09	59.77	59.31	58.74	59.47	58.63	56.67
6-12	57.77	42.67	65.73	64.01	62.69	63.07	64.1	61.68	49.37	59.01
7-1	62.23	60.07	61.28	61.27	62.01	57.54	59.89	60.22	59.01	60.39
7-2	38.32	36.07	41.16	40.3	38.66	33.41	39.09	37.17	35.3	37.72
7-3	55.98	53.42	54.54	54.6	54.66	53.01	53.42	53.37	51.96	53.88
7-4	76.63	56.91	76.83	76.78	57.28	71.59	73.28	65.43	54.48	67.69
Mean	56.8919	48.4098	56.4669	56.2250	53.1431	53.1910	55.2610	50.5743	47.1645	
N/M	0.15140	0.11240	0.15075	0.13237	0.11625	0.11364	0.18526	0.09579	0.09777	

FIG. 21

EYA4		27	29	31	34	37	44	46	55	65	74			
CpG												Mean		
NORMAL														
8-4		7.95	3.92	5.81	15.44	11.76	10	11.72	14.57	6.23	6.67	9.41		
8-5		0.52	0.72	5.76	3.59	0.23	2.35	3.48	0.41	2.95	0.38	2.04		
8-6		2.46	2.56	2.74	2.51	2.84	0.1	2.47	3.88	2.24	2.18	2.40		
8-7		27.16	27.12	7.42	26.64	26.48	4	4.06	10.23	23.1	25.77	18.20		
8-8		3.46	6.25	0.63	5.31	5.58	6.13	9.73	4.37	4.21	10.11	5.58		
8-9		1.55	1.81	0.51	1.55	1.55	0.23	0.26	1.66	6.07	8.5	2.37		
8-10		0.18	0.7	0.37	0.2	0.18	0.17	0.42	0.18	1.71	7.13	1.12		
8-11		14.45	9.34	8.79	11.49	11.46	8.65	3.08	8.6	2.32	10.83	8.90		
8-12		3.31	3.92	2.34	8.86	2.6	4.66	2.22	9.79	0.06	1.56	3.93		
Mean		6.78222	6.26000	3.81889	8.39889	6.96444	4.03222	4.16000	5.96556	5.43222	8.12556	5.99		
												SD all	6.64	
													M+2SD	19.27
													M+3SD	25.90
CpG		27	29	31	34	37	44	46	55	65	74	Mean		
ADENOMA														
3-4		42.34	7.24	13.95	15.1	34.99	13.74	20.19	23.02	14.87	25.63	21.11		
3-5		75.62	55.47	57.53	79.82	76.73	47.75	54.51	49.29	29.16	62.02	58.79		
3-6		94.03	94.1	93.31	90.55	87.59	92.01	88.36	66.23	38.32	65.91	81.04		
3-7		95.43	96.07	95.59	95.71	95.27	92.45	93.84	89.14	68.37	72.86	89.47		
3-8		95.86	86.62	49.24	95.35	79.12	96.8	94.79	92.04	63.24	70.73	82.38		
3-9		14.55	23.73	32.45	24.76	18.23	11.68	13.59	23.87	8.92	47.61	21.94		
3-10		13.57	23.23	29.82	21.68	16.82	10.17	13.21	21.92	7.26	44.63	20.23		
3-11		97.79	98.59	97.25	97.61	82.96	94.67	96.87	43.46	62.52	84.91	85.67		
3-12		65.73	63.48	64.89	71.19	67.65	69.9	72.69	81.74	39.26	68.67	66.52		
5-1		90.16	93.09	89.89	95.48	81.65	33.24	91.22	76.6	56.12	76.33	78.38		
5-2		83.81	71.43	69.95	87.62	79.52	27.62	79.52	58.1	45.71	66.18	66.05		
5-4		28.96	38.8	27.69	64.66	61.75	13.3	38.62	14.57	12.93	48.45	34.97		
5-11		94.41	94.41	92.31	97.2	90.91	28.32	76.92	66.08	39.86	80.07	76.05		
5-12		98.79	96.76	91.09	97.17	91.9	11.74	56.3	25.51	36.44	86.64	69.43		
6-1		95.63	96.8	96.26	96.51	94.84	94.08	96.72	89.66	67	71.73	89.94		
6-2		89.62	87.37	86.95	88.6	83.53	85.34	83.06	83.31	52.22	62.18	80.32		
6-3		97	96.79	95.1	96.68	95.73	95.3	94.9	82.6	59.33	71.11	88.45		
6-4		47.71	28.77	29.19	89.09	38.35	14.73	24.06	15.42	5.76	29.73	31.68		
6-5		15.49	7.45	9.53	20.3	13.44	8.9	5.48	15.51	6.9	11.4	11.44		
6-6		85.06	85.26	84.76	85.88	84.14	85.08	84.83	79.57	64.17	63.75	80.25		
7-5		93.98	94.73	95.05	92.77	94.15	92.06	90.95	87.51	60.32	65.65	86.72		
7-6		92.4	93	92.47	92.76	83.21	91.36	87.67	86.05	63.1	64.39	84.64		
7-7		61.1	61.34	59.78	65.15	62.99	59.16	60.7	52.74	37.07	44.69	56.47		
7-8		17	11.63	8.37	25.03	17.31	5.2	4.95	12.33	3.31	50.91	15.60		
7-9		80.91	74.06	77.26	82.16	76.42	76.69	77.13	72.52	52.25	65.57	73.50		
7-10		77	71.06	70.81	78.33	72.76	78.15	74.08	69.86	51.78	64.53	70.84		
7-11		0.53	0.84	9.01	2.12	16.04	6.46	9.41	8.97	12.98	22.93	8.93		
7-12		73.35	76.12	76.18	71.6	75.34	74.97	76.41	64.9	42.94	51.83	68.36		
8-1		89.74	89.94	90.16	89.41	89.4	89.69	88.68	74.41	70.39	76.14	84.80		
8-2		96.69	96.5	98.29	97.88	92.95	97.53	97.08	79.52	75.3	83.15	91.49		
8-3		89.79	93.92	94.97	94.28	71.33	59.43	61.38	34.46	42.65	55.09	69.73		
Mean		70.7758	68.0194	66.7774	74.0771	68.6142	56.7265	64.8426	56.1645	41.6274	59.8526			
N/M		0.09583	0.09203	0.05719	0.11338	0.10150	0.07108	0.06416	0.10622	0.13050	0.13576			

FIG. 2J

CoG	EYA4										
	27	39	31	34	37	44	46	55	65	74	Mean
CANCER											
1-1	95.77	95.43	95.77	96.12	90.28	94.76	94.95	72.84	67.49	82.09	88.55
1-2	91.26	92.07	92.65	93.03	91.62	91.94	91.19	69.82	62.06	79.13	85.48
1-3	95.81	96.96	96.22	95.96	94.14	96.56	94.2	80.24	62.51	77.93	89.05
1-4	96.31	95.85	97.08	96.12	87.07	94.71	76.53	36.85	36.16	69.45	78.61
1-5	97.36	97.86	97.2	97.58	97.13	97.41	97.29	84.8	65.65	81.89	91.42
1-6	94.33	95.64	95.97	95.08	92.67	95.52	93.72	69.51	72.14	82.97	88.76
1-7	18.72	9.23	9.21	53.23	36.24	14.01	44.41	58.09	7.14	13.19	26.35
1-8	46.66	30.42	29.64	59.15	24.97	19.9	15.52	10.52	31.39	67.42	33.56
1-9	96.11	94.22	93.91	97.09	90.15	93.97	93.78	73.93	73.3	67.96	87.44
1-10	96.21	94.28	93.99	97.24	91.01	94.45	94.19	73.8	72.18	70.03	87.74
1-11	98.77	97.62	97.6	96.75	97.25	98.23	96.89	93.56	41.81	78.11	89.66
1-12	91.41	91.71	92.15	92.58	91.12	91.65	69.38	67.07	55.93	78.28	82.13
2-1	19.91	21.85	20.02	19.4	20.66	18.45	19.59	16.34	12.97	12.04	18.12
2-2	33.9	37.54	33.6	35.41	26.72	32.37	30.33	25.63	12.73	21.64	28.99
2-3	87.09	85.4	86.41	87.06	70.66	82.49	87.01	52.21	38.48	67.66	74.45
2-4	81.28	84.94	84.36	84.62	81.79	75.22	67.07	18.47	8.36	20.2	60.63
2-5	95.51	84.65	89.45	97.75	96.96	89.4	93.64	71.6	60.1	71.7	85.68
2-6	89.64	90.39	92	89.4	86.3	86.95	89.9	78.41	50.38	66.56	81.99
2-7	87.95	83.38	87.88	86.33	79.7	62.37	74.51	58.32	41.49	63.64	72.56
2-8	88.93	79.13	74.75	91.49	78.51	87.98	88.76	67.95	49.06	66.23	77.28
2-9	96.63	96.37	96.75	95.33	94.76	91.66	91.7	82.93	62.73	69.07	87.79
2-10	96.23	96.18	96.45	95.35	94.09	91.68	92.03	84.12	62.29	68.48	87.69
2-11	78.21	91.62	93.38	82.83	80.84	89.21	86.42	47.49	19.68	55.01	72.41
2-12	78.92	77.21	65.25	88.2	87.17	48.02	48.4	43.86	32.71	55.41	62.52
3-2	83.19	83.33	85.51	87.83	84.07	85.4	83.93	72.44	38.71	62.51	76.69
5-3	75	77.63	73.03	76.97	65.13	21.71	66.45	14.47	8.55	28.29	50.72
5-5	76.69	89.83	90.89	79.24	49.15	28.39	67.37	31.78	19.07	40.04	57.25
5-6	84.14	66.9	82.76	80.69	60.69	18.62	65.52	26.9	13.79	33.1	53.31
5-7	83.85	83.08	85	83.46	76.54	29.23	84.23	41.92	30.38	70	66.77
5-8	59.34	63.74	62.09	68.13	56.59	23.08	62.09	39.01	27.47	46.7	50.82
5-9	82.26	85.35	86.38	84.32	80.21	29.56	81.23	18.25	26.74	74.29	64.86
5-10	78.22	81.24	81.41	80.57	78.06	25.46	79.23	18.76	23.95	71.36	61.83
6-7	96.25	97.94	97.18	96.54	94.24	95.22	96.77	90.47	68.78	71.86	80.53
6-8	75.42	60.15	66.23	85.35	70.66	76.79	71.49	67.79	49.75	59.12	68.28
6-9	74.63	72.85	75.4	77.97	75.44	66.67	58.5	43.07	8.53	12.99	56.61
6-10	72.13	68.52	70.34	73.37	69.14	63.44	53.88	42.34	6.23	11.98	53.14
6-11	90.57	91.22	90.96	88.64	90.51	87.87	90.7	78.51	60.18	62.03	83.12
6-12	81.63	94.2	95.03	93.02	79.24	89.94	88.49	58.09	32.59	53.01	76.52
7-1	95.77	93.81	90.2	96.4	94.38	94.6	93.33	85.21	55.69	65.68	86.51
7-2	69.02	62.6	67.89	67.45	67.11	63.19	61.07	42.99	27.53	29.84	55.87
7-3	87.53	87.41	85.13	87.34	86.47	85.59	86.17	77.97	56.43	57.76	79.78
7-4	96.06	96.57	94.2	94.28	78.02	15.36	26.08	27.32	17.28	53.07	59.82
Mean	81.3005	80.3886	80.7457	83.9207	77.0824	67.5360	74.9510	55.1343	39.7569	56.8981	
N/M	0.08342	0.07787	0.04730	0.10008	0.09035	0.05965	0.05550	0.10820	0.13664	0.14281	

FIG. 3A

Average of all individual base percent methylation					
Normal	Vim	BMP3	Septin9	TFPI2	EYA4
8-4	4.98	3.06	3.61	14.74	9.41
8-5	2.33	2.03	3.77	9.27	2.04
8-6	5.07	2.38	6.36	4.93	2.40
8-7	6.80	2.59	16.67	5.44	18.20
8-8	3.22	3.40	7.68	3.45	5.58
8-9	2.24	3.51	2.74	7.44	2.37
8-10	2.38	1.78	2.94	7.37	1.12
8-11	3.57	3.02	7.70	4.91	8.90
8-12	2.95	2.51	5.28	4.38	3.93
AVG	3.73	2.70	6.30	6.88	5.99
SD	2.68	1.18	5.06	3.86	6.64
M+2SD	9.09	5.05	16.42	14.61	19.27
M+3SD	11.77	6.23	21.47	18.47	25.90

FIG. 3C

Cancer	Vim	BMP3	Septin9	TFPI2	EYA4
1-1	38.80	3.11	72.66	75.29	88.55
1-2	34.74	2.01	64.15	59.09	85.48
1-3	13.19	3.02	76.77	68.12	89.05
1-4	77.62	5.51	74.27	77.45	78.61
1-5	84.93	84.69	82.00	77.97	91.42
1-6	72.92	68.88	72.83	71.18	88.76
1-7	8.37	3.35	58.66	37.29	26.35
1-8	3.19	2.68	39.30	30.00	33.56
1-9	86.69	20.30	87.12	84.60	87.44
1-10	86.72	19.05	87.45	83.78	87.74
1-11	72.01	7.18	91.29	67.89	89.66
1-12	49.26	2.79	52.59	65.89	82.13
2-1	2.81	1.96	58.49	12.75	18.12
2-2	25.33	2.51	54.09	17.58	28.99
2-3	49.44	47.22	49.61	43.69	74.45
2-4	45.03	7.18	43.65	43.06	60.63
2-5	3.42	74.17	66.30	61.27	85.08
2-6	38.28	42.91	18.62	29.96	81.99
2-7	54.05	56.75	42.19	43.39	72.56
2-8	61.55	7.65	68.72	61.66	77.28
2-9	3.59	83.18	71.20	73.77	87.79
2-10	3.05	82.80	70.51	70.48	87.69
2-11	64.29	61.54	21.90	56.83	72.41
2-12	43.51	3.18	57.65	52.05	62.52
3-2	45.20	52.41	46.70	42.30	76.69
5-3	29.27	26.61	48.36	36.41	50.72
5-5	8.72	16.58	29.14	56.02	57.25
5-6	8.00	2.62	64.43	33.04	53.31
5-7	40.50	3.87	57.83	60.34	66.77
5-8	20.77	36.11	31.25	19.52	50.82
5-9	40.55	6.67	57.52	49.73	64.86
5-10	39.62	5.65	56.99	47.24	61.83
6-7	88.95	86.24	85.46	82.97	90.53
6-8	57.90	49.40	61.73	48.26	68.28
6-9	5.14	2.80	55.38	26.03	56.61
6-10	3.66	2.29	54.41	25.27	53.14
6-11	64.29	51.17	2.67	56.67	83.12
6-12	40.48	14.45	75.85	59.01	76.52
7-1	55.25	2.38	62.71	60.39	86.51
7-2	29.60	3.17	63.49	37.72	55.87
7-3	59.56	63.21	73.54	53.88	79.78
7-4	71.98	3.07	77.84	67.69	59.82
	76.19	57.14	95.24	95.24	97.62
Vim, BMP, Sep TFPI2				100	

FIG. 3D

	Twenty fold dilution at 97.5% specificity (+2SD of mean)				
Cut off	9.09	5.05	15.73	14.61	19.27
Adenom.	Vim	BMP3	Septin9	TFPI2	EYA4
3-4	5.38	2.86	9.79	9.74	7.05
3-5	6.16	3.09	10.24	10.64	8.93
3-6	5.14	3.48	8.62	8.83	10.05
3-7	5.87	4.86	8.98	9.40	10.47
3-8	5.97	3.00	10.43	10.74	10.11
3-9	3.90	2.90	9.28	8.88	7.09
3-10	3.88	2.87	9.29	8.80	7.01
3-11	7.00	2.99	10.39	11.01	10.28
3-12	8.07	2.96	10.65	10.88	9.32
5-1	7.17	6.98	10.40	10.66	9.91
5-2	5.37	4.23	9.13	9.62	9.30
5-4	5.04	2.93	10.81	11.36	7.74
5-11	7.11	3.79	10.39	10.36	9.80
5-12	7.39	3.51	10.66	10.93	9.47
6-1	7.96	7.14	10.73	11.11	10.49
6-2	5.71	2.85	8.92	9.37	10.01
6-3	6.65	6.73	10.01	10.33	10.42
6-4	6.09	2.83	9.20	9.11	7.58
6-5	4.01	2.82	7.82	7.56	6.57
6-6	5.92	3.94	8.62	9.31	10.01
7-5	7.18	4.73	9.88	9.70	10.33
7-6	6.17	6.19	9.65	9.38	10.23
7-7	4.41	3.38	8.24	7.87	8.82
7-8	4.65	3.29	9.90	8.42	6.77
7-9	7.55	3.73	10.37	10.15	9.67
7-10	7.62	3.44	10.37	10.04	9.54
7-11	4.18	2.95	10.05	9.72	6.44
7-12	5.31	4.04	7.70	8.12	9.41
8-1	6.20	5.49	8.68	8.78	10.23
8-2	7.79	6.79	10.22	7.11	10.57
8-3	6.90	2.79	9.97	7.36	9.48
	0.00	12.90	0.00	0.00	0.00
	Vim, BMP, Sep TFPI2			12.90	

FIG. 3F

	Ten Fold dilution at 97.5% specificity (+2SD of mean)				
Cut off	9.09	5.05	15.73	14.61	19.27
Adenom.	Vim	BMP3	Septin9	TFPI2	EYA4
3-4	7.03	3.02	13.28	12.60	8.10
3-5	8.60	3.47	14.17	14.40	11.87
3-6	6.54	4.25	10.93	10.78	14.10
3-7	8.02	7.02	11.66	11.92	14.94
3-8	8.21	3.31	14.55	14.59	14.23
3-9	4.07	3.09	12.26	10.88	8.19
3-10	4.03	3.04	12.27	10.72	8.02
3-11	10.28	3.27	14.47	15.14	14.56
3-12	12.41	3.22	15.00	14.87	12.65
5-1	10.61	11.26	14.50	14.44	13.83
5-2	7.01	5.76	11.96	12.36	12.60
5-4	6.35	3.15	15.31	15.83	9.49
5-11	10.49	4.88	14.47	13.83	13.60
5-12	11.05	4.33	15.01	14.98	12.94
6-1	12.20	11.58	15.15	15.34	14.99
6-2	7.69	3.00	11.53	11.87	14.03
6-3	9.57	10.76	13.71	13.78	14.84
6-4	8.45	2.97	12.09	11.34	9.16
6-5	4.29	2.95	9.34	8.24	7.14
6-6	8.12	5.19	10.93	11.73	14.02
7-5	10.63	6.76	13.45	12.52	14.67
7-6	8.62	9.68	13.00	11.88	14.46
7-7	5.09	4.06	10.18	8.86	11.64
7-8	5.57	3.89	13.50	9.95	7.55
7-9	11.37	4.75	14.44	13.41	13.34
7-10	11.51	4.18	14.44	13.19	13.08
7-11	4.63	3.21	13.80	12.55	6.89
7-12	6.90	5.38	9.10	9.37	12.83
8-1	8.67	8.29	11.05	10.68	14.47
8-2	11.85	10.87	14.14	7.35	15.14
8-3	10.06	2.88	13.63	7.83	12.97
	38.71	35.48	0.00	16.13	0.00
	Vim, BMP, Sep TFPI2			61.29	

FIG. 5A

Methylation for specific sequence identified						
		Vim	BMP3	SEPTIN9	TFPI2	EYA4
	CpG loci:	37,40,45	34,53,61	59,61,68,71	55,59,63,6	31,34,37,44
Normal						
8-4		1.20%	0.18%	0.56%	5.46%	3.17%
8-5		0.20%	0.38%	0.71%	3.28%	0.03%
8-6		1.89%	0.02%	0.78%	0.74%	0.00%
8-7		1.62%	0.02%	3.77%	0.84%	3.85%
8-8		0.14%	0.01%	1.19%	0.13%	0.39%
8-9		0.01%	0.01%	0.07%	0.82%	0.09%
8-10		0.04%	0.01%	0.13%	0.91%	0.09%
8-11		0.30%	0.12%	1.52%	0.72%	8.49%
8-12		0.47%	0.12%	0.29%	0.05%	0.06%
Mean (M)		0.65%	0.10%	1.00%	1.44%	1.80%
Std Dev		0.72%	0.12%	1.14%	1.77%	2.92%
M+2SD		2.10%	0.34%	3.29%	4.99%	7.63%
M+3SD		2.82%	0.46%	4.43%	6.76%	10.55%

FIG. 5B

Cut off	2.82%	0.46%	4.43%	6.76%	10.55%
Adenoma	Vim	BMP3	Septin9	TFPI2	EYA4
3-4	19.12%	0.24%	60.96%	39.99%	2.07%
3-5	47.82%	0.90%	72.27%	63.44%	32.24%
3-6	21.85%	9.52%	39.62%	30.89%	84.77%
3-7	39.45%	37.77%	47.14%	39.66%	89.11%
3-8	39.08%	0.01%	77.07%	67.72%	43.89%
3-9	0.39%	0.28%	55.75%	32.78%	1.43%
3-10	0.03%	0.01%	55.92%	32.13%	0.38%
3-11	39.51%	0.03%	77.15%	76.22%	77.21%
3-12	85.15%	0.01%	78.97%	58.54%	54.28%
5-1	61.69%	81.06%	76.07%	64.28%	27.66%
5-2	26.34%	17.95%	47.65%	43.47%	16.19%
5-4	11.27%	0.00%	82.45%	74.81%	3.10%
5-11	61.64%	10.99%	74.59%	56.30%	24.83%
5-12	71.24%	2.97%	79.10%	68.40%	9.72%
6-1	83.10%	85.08%	84.48%	76.31%	87.64%
6-2	34.40%	0.42%	44.25%	41.48%	76.82%
6-3	52.37%	77.20%	65.30%	60.22%	89.49%
6-4	37.43%	0.07%	34.06%	24.61%	2.53%
6-5	1.08%	0.04%	10.43%	2.12%	0.69%
6-6	41.40%	21.23%	39.83%	41.53%	80.47%
7-5	57.63%	37.63%	65.54%	48.68%	87.49%
7-6	40.21%	64.61%	61.84%	44.09%	80.73%
7-7	9.25%	10.03%	31.53%	11.86%	51.79%
7-8	11.07%	6.96%	50.61%	11.99%	0.34%
7-9	75.10%	14.42%	72.15%	46.46%	59.02%
7-10	76.96%	7.90%	70.54%	42.87%	53.10%
7-11	3.41%	0.08%	52.03%	30.58%	0.39%
7-12	29.62%	23.60%	23.02%	17.75%	69.49%
8-1	45.04%	52.53%	42.61%	32.02%	88.68%
8-2	80.42%	78.85%	73.17%	0.36%	88.99%
8-3	61.90%	0.01%	69.09%	2.59%	52.72%
Sensitivity	90.32	58.06	100.00	90.32	70.97
vim,BMP3, Sep,TFPI2				100.00%	

FIG. 5C

Cancer	Vim	BMP3	Septin9	TFPI2	EYA4
1-1	33.41%	0.13%	61.92%	67.12%	85.88%
1-2	17.84%	0.06%	59.03%	52.16%	88.01%
1-3	1.73%	0.04%	63.33%	59.72%	88.62%
1-4	64.08%	0.31%	57.10%	45.72%	82.91%
1-5	82.75%	81.13%	71.85%	66.21%	94.37%
1-6	72.60%	65.31%	67.97%	64.34%	90.24%
1-7	1.69%	0.09%	53.46%	18.01%	1.48%
1-8	0.56%	0.27%	26.22%	23.61%	5.03%
1-9	87.66%	10.27%	80.69%	81.23%	84.91%
1-10	88.29%	8.62%	81.50%	80.11%	85.49%
1-11	82.28%	0.03%	78.50%	43.16%	92.99%
1-12	42.32%	0.10%	38.16%	58.35%	88.08%
2-1	0.49%	0.14%	45.78%	8.88%	17.68%
2-2	22.24%	0.13%	36.09%	1.25%	24.55%
2-3	48.22%	25.78%	45.63%	38.24%	64.23%
2-4	40.22%	0.78%	26.66%	35.78%	66.08%
2-5	1.11%	69.66%	62.58%	55.66%	80.10%
2-6	37.83%	41.05%	6.66%	25.54%	78.18%
2-7	52.55%	53.57%	28.93%	34.27%	55.18%
2-8	58.40%	0.88%	62.42%	48.36%	62.57%
2-9	0.59%	80.67%	51.00%	64.49%	85.96%
2-10	0.05%	79.46%	53.89%	61.26%	85.68%
2-11	56.80%	50.96%	0.75%	41.21%	65.12%
2-12	28.56%	0.15%	49.26%	39.41%	38.51%
3-2	43.57%	49.35%	43.20%	36.53%	80.82%
5-3	26.24%	15.25%	45.03%	20.39%	17.11%
5-5	1.95%	13.60%	19.90%	40.01%	12.29%
5-6	1.18%	0.04%	56.10%	2.94%	15.86%
5-7	27.04%	0.22%	51.50%	49.43%	23.85%
5-8	17.47%	31.72%	26.88%	12.76%	18.68%
5-9	38.11%	0.64%	53.60%	38.45%	26.48%
5-10	38.54%	0.32%	52.75%	36.60%	23.79%
6-7	86.89%	83.13%	74.34%	72.65%	89.23%
6-8	51.90%	36.83%	56.44%	38.21%	52.14%
6-9	1.38%	0.48%	52.25%	4.93%	60.58%
6-10	0.67%	0.09%	51.43%	4.31%	56.33%
6-11	57.37%	32.79%	0.50%	51.88%	84.25%
6-12	28.81%	11.47%	72.27%	51.57%	72.63%
7-1	48.16%	0.22%	53.59%	52.32%	86.00%
7-2	22.43%	0.05%	56.56%	26.31%	48.93%
7-3	58.13%	60.11%	70.26%	47.15%	78.31%
7-4	73.59%	0.21%	72.36%	42.14%	8.20%
Sensitivity	73.81	57.14	97.62	90.48	92.86
vim bmp3, sep, TFPI2				100.00%	

FIG. 5D

		Twenty fold dilution at 97.5% specificity (+2SD of mean)			
Cut off					
	2.10%	0.34%	3.29%	4.99%	7.63%
Adenoma	Vim	BMP3	Septin9	TFPI2	EYA4
3-4	1.61%	0.11%	4.05%	3.44%	1.90%
3-5	3.04%	0.14%	4.61%	4.61%	3.41%
3-6	1.74%	0.57%	2.98%	2.98%	6.04%
3-7	2.62%	1.98%	3.36%	3.42%	6.25%
3-8	2.60%	0.10%	4.85%	4.82%	3.99%
3-9	0.67%	0.11%	3.79%	3.08%	1.87%
3-10	0.65%	0.10%	3.80%	3.04%	1.82%
3-11	2.63%	0.10%	4.86%	5.25%	5.66%
3-12	4.91%	0.10%	4.95%	4.36%	4.51%
5-1	3.73%	4.15%	4.80%	4.65%	3.18%
5-2	1.97%	0.99%	3.38%	3.61%	2.61%
5-4	1.21%	0.10%	5.12%	5.18%	1.95%
5-11	3.73%	0.65%	4.73%	4.25%	3.04%
5-12	4.21%	0.25%	4.96%	4.86%	2.28%
6-1	4.81%	4.35%	5.23%	5.25%	6.18%
6-2	2.37%	0.12%	3.21%	3.51%	5.64%
6-3	3.27%	3.96%	4.27%	4.45%	6.27%
6-4	2.52%	0.10%	2.70%	2.67%	1.93%
6-5	0.70%	0.10%	1.52%	1.54%	1.83%
6-6	2.72%	1.16%	2.99%	3.51%	5.82%
7-5	3.53%	1.98%	4.28%	3.87%	6.17%
7-6	2.66%	3.33%	4.09%	3.64%	5.83%
7-7	1.11%	0.60%	2.58%	2.03%	4.39%
7-8	1.20%	0.44%	3.53%	2.04%	1.82%
7-9	4.41%	0.82%	4.61%	3.76%	4.75%
7-10	4.50%	0.49%	4.53%	3.58%	4.45%
7-11	0.82%	0.10%	3.60%	2.97%	1.82%
7-12	2.13%	1.28%	2.15%	2.32%	5.27%
8-1	2.90%	2.72%	3.13%	3.04%	6.23%
8-2	4.67%	4.04%	4.66%	1.46%	6.25%
8-3	3.75%	0.10%	4.46%	1.57%	4.43%
	Sensitivity				
	67.74	54.84	74.19	6.45	0.00
	v+B+S+TRPI2			96.77%	

FIG. 5E

Cancer	Twenty fold dilution at 97.5% specificity (+2SD of mean)				
	Vim	BMP3	Septin9	TFPI2	EYA4
1-1	2.32%	0.10%	4.10%	4.79%	6.09%
1-2	1.54%	0.10%	3.95%	4.05%	6.20%
1-3	0.74%	0.10%	4.17%	4.42%	6.28%
1-4	3.85%	0.11%	3.86%	3.72%	5.94%
1-5	4.79%	4.15%	4.59%	4.75%	6.52%
1-6	4.28%	3.36%	4.40%	4.65%	6.31%
1-7	0.73%	0.10%	3.67%	2.34%	1.87%
1-8	0.68%	0.11%	2.31%	2.62%	2.05%
1-9	5.03%	0.61%	5.04%	5.50%	6.04%
1-10	5.06%	0.53%	5.08%	5.44%	6.07%
1-11	4.76%	0.10%	4.93%	3.59%	6.45%
1-12	2.77%	0.10%	2.91%	4.35%	6.20%
2-1	0.67%	0.10%	3.29%	1.88%	2.68%
2-2	1.76%	0.10%	2.81%	1.50%	3.03%
2-3	3.06%	1.39%	3.28%	3.35%	5.01%
2-4	2.66%	0.14%	2.33%	3.23%	5.10%
2-5	0.71%	3.58%	4.13%	4.22%	5.80%
2-6	2.54%	2.15%	1.33%	2.71%	5.71%
2-7	3.28%	2.78%	2.45%	3.15%	4.56%
2-8	3.57%	0.14%	4.12%	3.86%	4.93%
2-9	0.68%	4.13%	3.55%	4.66%	6.10%
2-10	0.65%	4.07%	3.70%	4.50%	6.08%
2-11	3.49%	2.64%	1.04%	3.50%	5.05%
2-12	2.08%	0.10%	3.46%	3.41%	3.72%
3-2	2.83%	2.56%	3.16%	3.26%	5.84%
5-3	1.96%	0.86%	3.25%	2.46%	2.65%
5-5	0.75%	0.78%	2.00%	3.44%	2.41%
5-6	0.71%	0.10%	3.81%	1.58%	2.59%
5-7	2.00%	0.11%	3.58%	3.91%	2.99%
5-8	1.52%	1.68%	2.34%	2.08%	2.73%
5-9	2.56%	0.13%	3.68%	3.36%	3.12%
5-10	2.58%	0.11%	3.64%	3.27%	2.99%
6-7	4.99%	4.25%	4.72%	5.07%	6.26%
6-8	3.25%	1.94%	3.82%	3.35%	4.41%
6-9	0.72%	0.12%	3.61%	1.68%	4.83%
6-10	0.68%	0.10%	3.57%	1.65%	4.62%
6-11	3.52%	1.74%	1.03%	4.03%	6.01%
6-12	2.09%	0.67%	4.61%	4.02%	5.43%
7-1	3.06%	0.11%	3.68%	4.05%	6.10%
7-2	1.77%	0.10%	3.83%	2.75%	4.24%
7-3	3.56%	3.10%	4.51%	3.79%	5.71%
7-4	4.33%	0.11%	4.62%	3.54%	2.21%
Sensitivity					
	59.52	45.24	69.05	4.76	0.00
vim bmp3, sep, TFPI2				97.62%	

FIG. 5F

Ten Fold dilution at 97.5% specificity (+2SD of mean)					
Cut off	2.10%	0.34%	3.29%	4.99%	7.63%
Adenoma	Vim	BMP3	Septin9	TFF12	EYA4
3-4	2.56%	0.12%	7.10%	5.44%	2.01%
3-5	5.43%	0.19%	8.23%	7.78%	5.02%
3-6	2.84%	1.05%	4.96%	4.53%	10.28%
3-7	4.60%	3.87%	5.71%	5.40%	10.71%
3-8	4.56%	0.10%	8.71%	8.21%	6.19%
3-9	0.69%	0.12%	6.58%	4.72%	1.94%
3-10	0.65%	0.10%	6.59%	4.65%	1.84%
3-11	4.60%	0.10%	8.72%	9.06%	9.52%
3-12	9.16%	0.10%	8.90%	7.29%	7.23%
5-1	6.82%	8.20%	8.61%	7.87%	4.56%
5-2	3.28%	1.89%	5.77%	5.78%	3.42%
5-4	1.78%	0.10%	9.25%	8.92%	2.11%
5-11	6.81%	1.20%	8.46%	7.07%	4.28%
5-12	7.77%	0.39%	8.91%	8.28%	2.77%
6-1	8.96%	8.60%	9.45%	9.07%	10.56%
6-2	4.09%	0.14%	5.43%	5.59%	9.48%
6-3	5.89%	7.82%	7.53%	7.46%	10.75%
6-4	4.39%	0.10%	4.41%	3.90%	2.05%
6-5	0.76%	0.10%	2.04%	1.65%	1.87%
6-6	4.79%	2.22%	4.98%	5.59%	9.85%
7-5	6.41%	3.86%	7.55%	6.30%	10.55%
7-6	4.67%	6.56%	7.18%	5.85%	9.87%
7-7	1.57%	1.10%	4.15%	2.62%	6.98%
7-8	1.76%	0.79%	6.06%	2.64%	1.83%
7-9	8.16%	1.54%	8.22%	6.08%	7.70%
7-10	8.35%	0.89%	8.05%	5.72%	7.11%
7-11	0.99%	0.10%	6.20%	4.50%	1.84%
7-12	3.61%	2.46%	3.30%	3.21%	8.75%
8-1	5.15%	5.35%	5.26%	4.64%	10.67%
8-2	8.69%	7.98%	8.32%	1.47%	10.70%
8-3	6.84%	0.10%	7.91%	1.70%	7.07%
Sensitivity					
	77.42	58.06	93.55	61.29	38.71
v+B+S+TRP12				96.77%	


FIG. 5H

Cut off	Five Fold dilution at 97.5% specificity (+2SD of mean)				
	Vim	BMP3	Septin9	TFPI2	EYA4
3-4	4.47%	0.14%	13.19%	9.43%	2.21%
3-5	10.22%	0.28%	15.45%	14.13%	8.25%
3-6	5.02%	2.00%	8.92%	7.61%	18.75%
3-7	8.54%	7.65%	10.43%	9.37%	19.62%
3-8	8.47%	0.10%	16.41%	14.98%	10.58%
3-9	0.73%	0.15%	12.15%	7.99%	2.08%
3-10	0.66%	0.10%	12.18%	7.86%	1.87%
3-11	8.55%	0.10%	16.43%	16.68%	17.24%
3-12	17.68%	0.10%	16.80%	13.14%	12.65%
5-1	12.99%	16.31%	16.21%	14.29%	7.33%
5-2	5.92%	3.69%	10.53%	10.13%	5.04%
5-4	2.90%	0.10%	17.49%	16.40%	2.42%
5-11	12.98%	2.29%	15.92%	12.70%	6.76%
5-12	14.90%	0.69%	16.82%	15.12%	3.74%
6-1	17.27%	17.11%	17.90%	16.70%	19.33%
6-2	7.53%	0.18%	9.85%	9.73%	17.16%
6-3	11.13%	15.54%	14.06%	13.48%	19.70%
6-4	8.14%	0.11%	7.81%	6.36%	2.30%
6-5	0.87%	0.10%	3.09%	1.86%	1.94%
6-6	8.93%	4.34%	8.97%	9.74%	17.89%
7-5	12.18%	7.62%	14.11%	11.17%	19.30%
7-6	8.69%	13.02%	13.37%	10.25%	17.94%
7-7	2.50%	2.10%	7.31%	3.81%	12.16%
7-8	2.86%	1.49%	11.12%	3.83%	1.87%
7-9	15.67%	2.98%	15.43%	10.73%	13.60%
7-10	16.04%	1.68%	15.11%	10.01%	12.42%
7-11	1.33%	0.11%	11.41%	7.55%	1.88%
7-12	6.57%	4.82%	5.60%	4.99%	15.70%
8-1	9.66%	10.60%	9.52%	7.84%	19.54%
8-2	16.74%	15.87%	15.63%	1.51%	19.60%
8-3	13.03%	0.10%	14.82%	1.96%	12.34%
Sensitivity					
	87.10	58.06	96.77	80.65	61.29
	v+B+S+TRPI2		96.77%		

FIG. 7

Combination Vimentin, BMP3, Septin9, TFPI2

	Individual		Multi	
	Adenoma	Cancer	Adenoma	Cancer
Tissue @3SD	100	100	100	100
5 fold Dil:	96.8	97.6	96.8	100
10 fold Dil:	61.3	54.8	96.8	100
20 fold Dil:	12.9	23.8	96.8	97.6



31446-WO-1-ORD_ST25
SEQUENCE LISTING

<110> Ahlquist, David A.
Taylor, William R.
Zou, Hongzhi
Lidgard, Graham P.

<120> DIGITAL SEQUENCE ANALYSIS OF DNA METHYLATION

<130> EXCT-31446/WO-1/ORD

<150> US 61/438,649

<151> 2011-02-02

<160> 15

<170> PatentIn version 3.5

<210> 1

<211> 121

<212> DNA

<213> Homo sapiens

<400> 1

gaagagccca cctgtcaggc tgcgctgggt cagcgcagca agtggggctg gccgctatct 60

cgctgcaccc ggccgcgtcc cgggctccgt gcgccctcgc cccagctggt ttggagttca 120

a 121

<210> 2

<211> 121

<212> DNA

<213> Artificial

<220>

<223> Synthetic

<400> 2

gaagagtta tttgttaggt tgtgttgggt tagttagta agtggggttg gttgttattt 60

tgttgattt gttgtgttt tgggttttgt gtgtttttgt tttagttggt ttggagttta 120

a 121

<210> 3

<211> 121

<212> DNA

<213> Artificial

<220>

<223> Synthetic

<400> 3

gaagagtta tttgttaggt tgcgttgggt tagcgtagta agtggggttg gtcgttattt 60

cgttgtattc ggtcgcgttt cgggtttcgt gcgttttcgt tttagttggt ttggagttta 120

a 121

<210> 4

<211> 129

<212> DNA

<213> Homo sapiens

31446-W0-1-ORD_ST25

<400> 4
 aaataaaaac ggagtgggtc ccccgcgcgc cgccgcccc cgcgccctg gcggcctccc 60
 ccgaggcccc cggcggcctc acgagcccgc agtagccggt ggcgacgtcg cccccgcccc 120
 acctccccg 129

<210> 5
 <211> 132
 <212> DNA
 <213> Arti fi ci al

<220>
 <223> Syntheti c

<400> 5
 gttaaataaa aatggagtgg gttttttgtg tgttgttgtt ttttgtgttt ttggtggttt 60
 tttttgaggt ttttgggtgt tttatgagtt tgtagtagtt ggtggtgatg ttgtttttgt 120
 tttatttttt tg 132

<210> 6
 <211> 132
 <212> DNA
 <213> Arti fi ci al

<220>
 <223> Syntheti c

<400> 6
 gttaaataaa aacggagtgg gtttttcgcg cgtcgctggt tttcgcgttt ttggcggttt 60
 ttttcgaggt tttcggcgggt tttacgagtt cgtagtagtc ggtggcgacg tcgttttcgt 120
 tttatttttt tg 132

<210> 7
 <211> 99
 <212> DNA
 <213> Homo sapi ens

<400> 7
 ggccgcagca gccagcccag caccacctt cgaagtccga aatgatccca tccagctgcg 60
 cgttgaccgc ggggtccgac atgatggctg gtgggcagc 99

<210> 8
 <211> 99
 <212> DNA
 <213> Arti fi ci al

<220>
 <223> Syntheti c

<400> 8
 ggttgtagta gttagtttag ttttttttt tgaagtttga aatgatttta tttagttgtg 60
 tgttgattgt ggggtttgat atgatggtg gtgggtagt 99

<210> 9
 <211> 99
 <212> DNA

<213> Arti fi ci al
 <220>
 <223> Syntheti c
 <400> 9
 ggtcgtagta gttagtttag tatttatttt cgaagttcga aatgatttta tttagttgcg 60
 cgttgatcgc ggggttcgat atgatggtt gtgggtagc 99

<210> 10
 <211> 116
 <212> DNA
 <213> Homo sapi ens
 <400> 10
 aggaggtgcg cggctttctg ctccaggcgg cccgggtgcc cgctttatgc ggggagcgcg 60
 tccggccgac ccccgccggg gcggagcctg aggggtggct gattcatgca cgggga 116

<210> 11
 <211> 116
 <212> DNA
 <213> Arti fi ci al
 <220>
 <223> Syntheti c
 <400> 11
 aggaggtgtg tggttttttg ttttaggtgg tttgggtggt tgttttatgt ggggtgagtg 60
 tttggttgat tttgttggg gtggagttt aggggtggtt gatttatgta tgggga 116

<210> 12
 <211> 116
 <212> DNA
 <213> Arti fi ci al
 <220>
 <223> Syntheti c
 <400> 12
 aggaggtgcg cggttttttg ttttaggcgg ttcgggtggt cgctttatgc ggggagcgcg 60
 ttcggtcgat tttcgtcggg gcggagttt aggggtggtt gatttatgta cgggga 116

<210> 13
 <211> 117
 <212> DNA
 <213> Homo sapi ens
 <400> 13
 caggctccgtg tcctcgtcct cctaccgag gatgttcggc ggcccgggca ccgagagccg 60
 gccgagctcc agccggagct acgtgactac gtccaccgc acctacagcc tgggca 117

<210> 14
 <211> 117
 <212> DNA
 <213> Arti fi ci al
 <220>
 <223> Syntheti c

31446-W0-1-ORD_ST25

<400> 14
taggtttgtg tttttgtttt tttattgtag gatgtttggt ggtttgggta ttgtgagttg 60
gttgagtttt agttggagtt atgtgattat gtttatttgt atttatagtt tgggtag 117

<210> 15
<211> 117
<212> DNA
<213> Arti fi ci al

<220>
<223> Synthetic

<400> 15
taggttcgtg ttttcgtttt tttatcgtag gatgttcggc ggttcgggta tgcgagtcg 60
gtcgagtttt agtcggagtt acgtgattac gtttattcgt atttatagtt tgggtag 117