



US00PP14355P29

(12) **United States Plant Patent**
Hoogendoorn

(10) **Patent No.:** **US PP14,355 P2**

(45) **Date of Patent:** **Dec. 9, 2003**

(54) **ALSTROEMERIA PLANT NAMED 'STALIDI'**

(22) Filed: **Nov. 4, 2002**

(50) Latin Name: *Alstroemeria hybrida*
Varietal Denomination: **Stalidi**

(51) **Int. Cl.**⁷ **A01H 5/00**

(52) **U.S. Cl.** **Plt./309**

(58) **Field of Search** **Plt./309**

(75) Inventor: **Cornelis Arie Hoogendoorn,**
Rijsenhout (NL)

Primary Examiner—Kent Bell

(74) *Attorney, Agent, or Firm*—C. A. Whealy

(73) Assignee: **Van Zanten Plants B.V.,** Hillegom
(NL)

(57) **ABSTRACT**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

A new and distinct cultivar of Alstroemeria plant named 'Stalidi', characterized by its erect flowering stems; light and pale yellow bi-colored flowers with dark purple spots and stripes; and excellent postproduction longevity.

(21) Appl. No.: **10/287,083**

1 Drawing Sheet

1

2

Botanical classification/cultivar designation: Alstroemeria hybrida cultivar 'Stalidi'.

BACKGROUND OF THE INVENTION

The present Invention relates to a new and distinct cultivar of Alstroemeria plant, botanically known as Alstroemeria hybrida, commercially used as a cut flower Alstroemeria, and hereinafter referred to by the name 'Stalidi'.

The new Alstroemeria is a product of a planned breeding program conducted by the Inventor in Rijsenhout and Aalsmeer, The Netherlands. The objective of the breeding program was to develop new cut flower Alstroemeria cultivars with strong plant growth, attractive flower colors and excellent postproduction longevity.

The new Alstroemeria originated from a cross made by the Inventor in April, 1998 in Rijsenhout, The Netherlands, of a proprietary Alstroemeria hybrida selection identified as 94T319-2, not patented, as the female, or seed, parent with a proprietary Alstroemeria hybrida selection identified as 87G1069-2, not patented, as the male, or pollen, parent. The new Alstroemeria was discovered and selected by the Inventor as a flowering plant within the progeny of the stated cross in a controlled environment in Aalsmeer, The Netherlands in June, 1999.

Asexual reproduction of the new cultivar by root divisions taken in a controlled environment in Rijsenhout, The Netherlands, since June, 1999, has shown that the unique features of this new Alstroemeria are stable and reproduced true to type in successive generations of asexual propagation.

SUMMARY OF THE INVENTION

Plants of the cultivar Stalidi have not been observed under all possible environmental conditions. The phenotype may vary somewhat with variations in environment such as temperature and light intensity without, however, any variance in genotype.

The following traits have been repeatedly observed and are determined to be the unique characteristics of 'Stalidi'. These characteristics in combination distinguish 'Stalidi' as a new and distinct cultivar:

1. Erect flowering stems.
2. Light and pale yellow bi-colored flowers with dark purple-colored spots and stripes.
3. Excellent postproduction longevity.

Plants of the new Alstroemeria are most similar to plants of the parent selections. However, plants of the new Alstroemeria differ from plants of the parents in flower coloration as plants of the female parent have red-colored flowers and plants of the male parent have darker yellow-colored flowers.

Plants of the new Alstroemeria can be compared to plants of the cultivar Stabelin, disclosed in U.S. Plant Pat. No. 10,254. In side-by-side comparisons conducted in Rijsenhout, The Netherlands, plants of the new Alstroemeria differed primarily from plants of the cultivar Stabelin in flower color as plants of the cultivar Stabelin had darker yellow-colored flowers than plants of the new Alstroemeria.

BRIEF DESCRIPTION OF THE PHOTOGRAPH

The accompanying colored photograph illustrates the overall appearance of the new Alstroemeria, showing the colors as true as it is reasonably possible to obtain in colored reproductions of this type. Colors in the photograph may differ slightly from the color values cited in the detailed botanical description which accurately describe the colors of the new Alstroemeria. The photograph comprises a side perspective view of typical flowers of 'Stalidi'.

DETAILED BOTANICAL DESCRIPTION

The aforementioned photograph, following observations and measurements describe plants of the new Alstroemeria grown in Rijsenhout, The Netherlands in a glass-covered greenhouse in ground beds. During the production of the plants, day temperatures ranged from 15 to 20° C., night temperatures ranged from 10 to 15° C., soil temperatures of 14° C. and light levels were about 4,000 lux. Plants used for the photograph and description were about 12 months from planting root divisions. The photograph and the description were taken during March and April, 2002.

Color references are made to The Royal Horticultural Society Colour Chart, 1995 Edition, except where general terms of ordinary dictionary significance are used.

Botanical classification: *Alstroemeria hybrida* cultivar Stalidi.

Parentage:

Female parent.—Proprietary *Alstroemeria hybrida* selection identified as 94T319-2, not patented.

Male parent.—Proprietary selection of *Alstroemeria hybrida* identified as 87G1069-2, not patented.

Propagation:

Type.—By root divisions.

Root description.—Fibrous, fleshy, thick; white, close to 155D, in color.

Rooting habit.—Freely branching.

Rhizomes.—Shape: Elongate; rounded. Length: About 10 to 30 cm. Diameter: About 0.3 to 1 cm. Texture: Smooth. Color: Close to 155D.

Plant description:

Plant habit.—Upright; freely basal-branching, bushy appearance. Time from planting to harvest of cut flowers: About 80 to 90 days.

Number of flowering stems produced per year.—About 184 to 200.

Plant height.—About 124 to 180 cm.

Plant diameter (spread).—About 25 to 30 cm.

Flowering stem description.—Aspect: Erect. Length: About 130 cm. Diameter: About 4.75 to 10 mm. Internode length: About 1 to 7 cm. Strength: Strong. Texture: Glabrous. Color: Close to 144A to 144B.

Foliage description.—Leaves asymmetrical; sessile. Length: About 14 to 20 cm. Width: About 2.5 to 3.5 cm. Shape: Lanceolate. Apex: Acute. Base: Attenuate. Margin: Entire. Texture, upper and lower surfaces: Glabrous. Venation pattern: Parallel. Color: Young and fully developed foliage, upper surface: Close to 137A; moderately glossy. Young and fully developed foliage, lower surface: Close to 137B. Venation: Upper surface, close to 137A; lower surface, close to 137B.

Flower description:

Flower type and habit.—Single cup-shaped flowers arranged in compound umbels. Perianth segments separate. Freely and continuously flowering. Flowers not persistent.

Natural flowering season.—Flowering continuous during the spring in The Netherlands.

Fragrance.—None detected.

Flower longevity on the plant.—About four weeks.

Flower longevity as a cut flower.—About 20 to 25 days.

Flower buds (showing color).—Length: About 2.5 to 3.5 cm. Diameter: About 1.5 to 2 cm. Shape: Roughly ovoid. Color: Close to 16C.

Umbel height.—About 12 to 18 cm.

Umbel diameter.—About 20 to 30 cm.

Number of flowers per umbel.—About 15 to 28.

Flower height (length).—About 7.5 to 8.5 cm.

Flower diameter.—About 7.5 to 8.5 cm.

Flower depth.—About 5 to 6 cm.

Perianth.—Arrangement: Six arranged in two whorls, each whorl with two lateral and one median segments. Size: Inner perianth: Length: Laterals, about 5.2 to 6 cm; median, 5.5 to 6 cm. Width, laterals and median: About 2 to 2.5 cm. Outer perianth: Length, laterals and median: About 6 to 7 cm. Width, laterals and median: About 3.5 to 4 cm. Shape: Inner perianth, all segments: Oblanceolate. Outer perianth, all segments: Obovate. Apex: Inner perianth, all segments: Acute. Outer perianth, all segments: Bracket-shaped. Base, inner and outer perianths, all segments: Attenuate. Margin, inner and outer perianths, all segments: Entire; slightly undulate. Texture, inner and outer perianths, all segments: Smooth, glabrous; velvety. Color: Inner perianth: When opening and fully opened, upper surface: Laterals: At apex, close to 146A; towards apex, close to 16A; mid-section, close to 7A; towards base, close to 16C; spots and stripes, close to 187A. Median: At apex, close to 146A; towards apex, close to 61A; towards base, close to 61C; spots and stripes, close to 187A. When opening and fully opened, lower surface: Laterals: Close to 16C; at apex, close to 146A. Median: Close to 61C; at apex, close to 146A. Outer perianth: When opening and fully opened, upper surface, laterals and median: At apex, close to 146A; towards apex, close to 16A; towards base, close to 16B. When opening and fully opened, lower surface: Laterals: Close to 16C; at apex, close to 146A. Median: Close to 33A; towards margins, close to 28B; at apex, close to 146A.

Pedicels.—Length: About 1 to 3 cm. Diameter: About 2 to 4 mm. Strength: Strong. Angle: About 20 to 50° from vertical. Texture: Smooth, glabrous. Color: Close to 137A.

Reproductive organs.—Stamens: Quantity per flower: Six. Anther shape: Elliptical. Anther length: About 8 mm. Anther diameter: About 3 mm. Anther color: Close to 177B. Pollen amount: Scarce. Pollen color: Close to 185A. Pistils: Quantity per flower: One. Style length: About 3 to 4 cm. Stigma color: Orange red. Ovary color: Close to 144B.

Fruit.—Shape: Globular. Color: Brown.

Disease/pest resistance: Plants of the new *Alstroemeria* have not been observed to be resistant to pathogens and pests common to *Alstroemerias*.

Temperature tolerance: Plants of the new *alstroemeria* have been observed to tolerate temperatures from -5 to 40° C. It is claimed:

1. A new and distinct cultivar of *Alstroemeria* plant named 'Stalidi', as illustrated and described.

* * * * *

